

Title (en)
METHOD FOR ORTHO-ALKYLATION OF HYDROXYAROMATIC COMPOUNDS

Publication
EP 0129065 B1 19920930 (EN)

Application
EP 84105611 A 19840517

Priority
US 50068483 A 19830603

Abstract (en)
[origin: EP0129065A2] Hydroxyaromatic compounds such as phenol are alkylated with alkanols such as methanol in the presence of a catalyst comprising magnesium oxide with up to about 0.1% by weight of copper deposited thereon in a layer of submicroscopic thickness. The catalyst is prepared by preparing a slurry of a substantially water-insoluble magnesium reagent such as magnesium hydroxide or magnesium carbonate in an aqueous solution of at least one copper salt. removing and drying the solid phase, and calcining the same at a temperature within the range of about 350-550°C. The catalyst promotes ortho-alkylation of the hydroxyaromatic compound in high yield, minimizes decomposition of the alkanol, and has long life.

IPC 1-7
B01J 23/74; C07C 37/16

IPC 8 full level
B01J 23/72 (2006.01); **B01J 23/74** (2006.01); **B01J 23/78** (2006.01); **C07B 61/00** (2006.01); **C07C 27/00** (2006.01); **C07C 37/11** (2006.01); **C07C 37/16** (2006.01); **C07C 39/06** (2006.01); **C07C 67/00** (2006.01)

CPC (source: EP US)
B01J 23/74 (2013.01 - EP US); **C07C 37/16** (2013.01 - EP US)

Cited by
US5594123A; EP0208988A1; US4753913A; US11007508B2; WO2018023015A3

Designated contracting state (EPC)
DE FR GB IT NL

DOCDB simple family (publication)
EP 0129065 A2 19841227; EP 0129065 A3 19850731; EP 0129065 B1 19920930; AU 2876284 A 19841206; AU 574680 B2 19880714; BR 8402749 A 19850514; DE 3485946 D1 19921105; DE 3485946 T2 19930422; JP H0341213 B2 19910621; JP S59228940 A 19841222; US 4554267 A 19851119

DOCDB simple family (application)
EP 84105611 A 19840517; AU 2876284 A 19840528; BR 8402749 A 19840604; DE 3485946 T 19840517; JP 10931684 A 19840529; US 50068483 A 19830603