

Title (en)

Improved synthesis of molybdenum/alkylene glycol complexes useful as epoxidation catalysts.

Title (de)

Synthese von Molybdän-Alkylenglycol-Komplexen verwendbar als Epoxydationskatalysatoren.

Title (fr)

Synthèse de complexes molybdène/alkylène-glycol utilisables comme catalyseurs d'époxydation.

Publication

EP 0193682 A1 19860910 (EN)

Application

EP 85309344 A 19851220

Priority

US 68770184 A 19841231

Abstract (en)

Catalytically-active complexes, useful inter alia for olefin epoxidation can be obtained by reacting an ammonium compound containing molybdenum and a glycol, in a mol ratio from 7:1 to 20:1 at an elevated temperature, and if necessary stripping water to give a final water content from 0.5 to 6.0 wt.%. <??>The reaction is carried out in the substantial absence of solvents or diluents, and the glycol has a molecular weight up to 200 and contains only primary or secondary hydroxy groups, on adjacent carbon atoms, or on carbon atoms separated by an even number of other carbon atoms.

IPC 1-7

B01J 23/22; **C07D 301/19**

IPC 8 full level

B01J 31/22 (2006.01); **C07D 301/19** (2006.01); **C07F 11/00** (2006.01)

CPC (source: EP US)

B01J 31/2208 (2013.01 - EP US); **B01J 31/2226** (2013.01 - EP US); **C07D 301/19** (2013.01 - EP US); **B01J 2231/72** (2013.01 - EP US); **B01J 2531/64** (2013.01 - EP US)

Citation (search report)

- [X] US 4009122 A 19770222 - LINES ELLWOOD L, et al
- [A] FR 1550166 A 19681220

Cited by

US5620938A; EP0524816A3; FR2781169A1; EP0321612A1; EP0343293A1; EP0264184A1; US6642170B1; WO0003802A1; WO9404268A1

Designated contracting state (EPC)

BE DE FR GB NL

DOCDB simple family (publication)

EP 0193682 A1 19860910; **EP 0193682 B1 19900307**; CA 1266676 A 19900313; DE 3576336 D1 19900412; JP H0440358 B2 19920702; JP S61161294 A 19860721; US 4626596 A 19861202

DOCDB simple family (application)

EP 85309344 A 19851220; CA 483634 A 19850611; DE 3576336 T 19851220; JP 29091485 A 19851225; US 68770184 A 19841231