

Title (en)

Protein-protein interactions and methods for identifying them

Title (de)

Verfahren zur Identifizierung von Modulatoren wechselwirkender Proteine

Title (fr)

Procédé d'identification de modulateurs de protéines interagissantes

Publication

EP 1684077 A1 20060726 (EN)

Application

EP 06000197 A 20000512

Priority

- EP 00929188 A 20000512
- US 13425999 P 19990514

Abstract (en)

The invention relates to protein-protein interactions and methods for identifying interacting proteins and the amino acid sequence at the site of interaction. Using overlapping hexapeptides that encode for the entire amino acid sequences of the linker domains of human P-glycoprotein gene 1 and 3 (HPgp1 and HP-gp3), a direct and specific binding between P-gpl and 3 linker domains and intracellular proteins was demonstrated. Three different stretches (617 EKGIFYFKLVTM 627 , 658 SRSSLIRKRSTRRSVRGSOA 677 and 694 PVSFWRIMKLNLT 706 for P-gpl and 618 LMKKEGVYFKLVNM 631 , 648 KAA-TRMAPNGWKSRLFRHSTQKNLKNS 674 and 695 PVSFLKVLKLNKT 677 for P-gp3) in linker domains bound to proteins with apparent molecular masses of ~80 kDa, 57 kDa and 30 kDa. The binding of the 57 kDa protein was further characterized. Purification and partial N-terminal amino acid sequencing of the 57 kDa protein showed that it encodes the N-terminal amino acids of alpha and beta-tubulins. The method of the present invention was further validated with Annexin. The present invention thus demonstrates a novel concept whereby the interactions between two proteins are mediated by strings of few amino acids with high and repulsive binding energies, enabling the identification of high-affinity binding sites between any interacting proteins.

IPC 8 full level

G01N 33/68 (2006.01)

CPC (source: EP)

C40B 30/04 (2013.01)

Citation (search report)

- [Y] WO 8403564 A1 19840913 - COMMW SERUM LAB COMMISSION [AU]
- [Y] WO 9815833 A1 19980416 - UNIV UTRECHT [NL], et al
- [Y] EP 0818467 A2 19980114 - NEC CORP [JP]
- [Y] RIVERO-LEZCANO O M ET AL: "Specificity of protein interactions with highly related SRC homology (SH) domains of FGR and FYN protein-tyrosine kinases.", FEBS LETTERS, 31 JAN 1994, vol. 338, no. 2, 31 January 1994 (1994-01-31), pages 183 - 186, XP002384213, ISSN: 0014-5793
- [Y] KITAY MARY KATE ET AL: "Protein-protein interactions between Epstein-Barr virus nuclear antigen-iP and cellular gene products: Binding of 70 kilodalton heat shock proteins", VIROLOGY, vol. 220, no. 1, 1996, pages 91 - 99, XP002384214, ISSN: 0042-6822
- [Y] CIANFRIGLIA M. ET AL.: "P-glycoprotein epitope mapping: II. The murine monoclonal antibody MM6.15 to human multidrug-resistant cells binds with three distinct loops in the MDR1-P-glycoprotein extracellular domain", INTERN. J. CANCER, vol. 61, 1995, pages 142 - 147, XP000957487
- [Y] ELLGAARD L ET AL: "Nested sets of protein fragments and their use in epitope mapping: characterization of the epitope for the S4D5 monoclonal antibody binding to receptor associated protein", JOURNAL OF IMMUNOLOGICAL METHODS, ELSEVIER SCIENCE PUBLISHERS B.V.,AMSTERDAM, NL, vol. 180, no. 1, 13 March 1995 (1995-03-13), pages 53 - 61, XP004021070, ISSN: 0022-1759
- [DY] GEORGES E. ET AL.: "Topology of P-glycoprotein as determined by epitope mapping of MRK-16 monoclonal antibody", J. BIOL. CHEM., vol. 268, no. 3, 1993, pages 1792 - 1798, XP002150421
- [DY] GEORGES E. ET AL.: "Detection of P-glycoprotein isoforms by gene-specific monoclonal antibodies", PROC. NATL. ACAD. SCI. USA, vol. 87, 1990, pages 152 - 156, XP002150436

Designated contracting state (EPC)

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE

DOCDB simple family (publication)

EP 1684077 A1 20060726

DOCDB simple family (application)

EP 06000197 A 20000512