



(11) **EP 1 313 508 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
18.04.2007 Bulletin 2007/16

(51) Int Cl.:
A61K 45/06^(2006.01) A61P 35/00^(2006.01)
A61K 31/505^(2006.01) A61K 31/70^(2006.01)

(21) Application number: **01948214.0**

(86) International application number:
PCT/US2001/014860

(22) Date of filing: **15.06.2001**

(87) International publication number:
WO 2002/002093 (10.01.2002 Gazette 2002/02)

(54) **COMBINATION CONTAINING AN ANTIFOLATE AND METHYLMALONIC ACID LOWERING AGENT**

ZUSAMMENSETZUNG, WELCHE EIN ANTIFOLATE UND EIN METHYLMALONSÄURESENKENDES MITTEL ENTHÄLT

COMPOSITION COMPRENANT UN ANTIFOLATE ET UN AGENT REDUCTEUR D'ACIDE METHYLMALONIQUE

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE TR
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **30.06.2000 US 215310 P**
27.09.2000 US 235859 P
18.04.2001 US 284448 P

(43) Date of publication of application:
28.05.2003 Bulletin 2003/22

(73) Proprietor: **ELI LILLY AND COMPANY**
Indianapolis, Indiana 46285 (US)

(72) Inventors:
• **NIYIKIZA, Clet**
Indianapolis, IN 46278 (US)
• **PAOLETTI, Paolo**
Indianapolis, IN 46240 (US)
• **RUSTHOVEN, James, Jacob**
Ancaster, Ontario L9G 1G4 (CA)

(74) Representative: **Burnside, Ivan John et al**
Eli Lilly and Company Limited
European Patent Operations
Lilly Research Centre
Erl Wood Manor, Sunnighill Road
Windlesham, Surrey GU20 6PH (GB)

(56) References cited:
EP-A- 0 546 870 **US-A- 5 405 839**

- **CALVERT H.: "Folate status and the safety profile of antifolates." SEMINARS IN ONCOLOGY, (2002) 29/2 SUPPL. 5 (3-7). , XP008005755**
- **CALVERT H.: "Future directions in the development of pemetrexed" SEMINARS IN ONCOLOGY, (2002) 29/2 SUPPL. 5 (54-61). , XP008005744**
- **WESTERHOF, G. ROBBIN ET AL: "Carrier- and receptor-mediated transport of folate antagonists targeting folate -dependent enzymes: correlates of molecular-structure and biological activity" MOL. PHARMACOL. (1995), 48 (3), 459-71 , XP008005762**
- **WORZALLA, JOHN F. ET AL: "Role of folic acid in modulating the toxicity and efficacy of the multitargeted antifolate, LY231514" ANTICANCER RESEARCH (1998), 18(5A), 3235-3239 , XP008005757**
- **HANAUSSKE A-R ET AL: "PEMETREXED DISODIUM: A NOVEL ANTIFOLATE CLINICALLY ACTIVE AGAINST MULTIPLE SOLID TUMORS" ONCOLOGIST, ALPHAMED PRESS, US, vol. 4, no. 6, 2001, pages 363-373, XP008005751 ISSN: 1083-7159**
- **BUNN P ET AL: "VITAMIN B12 AND FOLATE REDUCE TOXICITY OF ALIMTA (PEMETREXED DISODIUM, LY231514, MTA), A NOVEL ANTIFOLATE/ANTIMETABOLITE" PROGRAM/ PROCEEDINGS - AMERICAN SOCIETY OF CLINICAL ONCOLOGY, THE SOCIETY, US, vol. 76A, no. 20, 2001, page 300 XP008005885 ISSN: 1081-0641**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 313 508 B1

Description

[0001] Potentially, life-threatening toxicity remains a major limitation to the optimal administration of antifolates. (see, generally, *Antifolate Drugs in Cancer Therapy*, edited by Jackman, Ann L., Humana Press, Totowa, NJ, 1999.) In some cases, a supportive intervention is routinely used to permit safe, maximal dosing. For example, steroids, such as dexamethone, can be used to prevent the formation of skin rashes caused by the antifolate. (*Antifolate*, pg 197.)

[0002] Antifolates represent one of the most thoroughly studied classes of antineoplastic agents, with aminopterin initially demonstrating clinical activity approximately 50 years ago. Methotrexate was developed shortly thereafter, and today is a standard component of effective chemotherapeutic regimens for malignancies such as lymphoma, breast cancer, and head and neck cancer. (Bonnadonna G, Zambetti M, Valagussa P. Sequential or alternating doxorubicin and CMF regimens in breast cancer with more than three positive nodes: Ten year results. *JAMA* 1995;273(7):542-547; Bonnadonna G, Valagussa P, Moliterni A, Zambetti M, Brambilla C. Adjuvant cyclophosphamide, methotrexate, and fluorouracil in node-positive breast cancer: The results of 20 years of follow-up. *N Engl J Med* 1995;332(14):901-906; and Hong WK, Schaefer S, Issell B, et al A prospective randomized trial of methotrexate versus cisplatin in the treatment of recurrent squamous cell carcinoma of the head and neck. *Cancer* 1983;52:206-210.) Antifolates inhibit one or several key folate-requiring enzymes of the thymidine and purine biosynthetic pathways, in particular, thymidylate synthase (TS), dihydrofolate reductase (DHFR), and glycinamide ribonucleotide formyltransferase (GARFT), by competing with reduced folates for binding sites of these enzymes. (Shih C, Habeck LL, Mendelsohn LG, Chen VJ, Schultz RM. Multiple folate enzyme inhibition: Mechanism of a novel pyrolopyrimidine-based antifolate LY231514 (MTA). *Advan Enzyme Regul*, 1998; 38:135-152 and Shih C, Chen VJ, Gossett LS, et al. LY231514, a pyrrolo[2,3-d]pyrimidine-based antifolate that inhibits multiple folate-requiring enzymes. *Cancer Res* 1997;57:1116-1123.) Several antifolate drugs are currently in development. Examples of antifolates that have thymidylate synthase inhibiting ("TSI") characteristics include 5-fluorouracil and Tomudex®. An example of an antifolate that has dihydrofolate reductase inhibiting ("DHFR") characteristic is Methotrexate®. An example of an antifolate that has glycinamide ribonucleotide formyltransferase inhibiting ("GARFT") characteristics is Lometrexol. Many of these antifolate drugs inhibit more than one biosynthetic pathway. For example Lometrexol is also an inhibitor of dihydrofolate reductase and pemetrexed disodium (Alimta®, Eli Lilly and Company, Indianapolis, IN) has demonstrated thymidylate synthase, dihydrofolate reductase, and glycinamide ribonucleotide formyltransferase inhibition.

[0003] A limitation to the development of these drugs is that the cytotoxic activity and subsequent effectiveness of antifolates may be associated with substantial toxicity for some patients. Additionally antifolates as a class are associated with sporadic severe myelosuppression with gastrointestinal toxicity which, though infrequent, carries a high risk of mortality. The inability to control these toxicities led to the abandonment of clinical development of some antifolates and has complicated the clinical development of others, such as Lometrexol and raltitrexed. (Jackman AL, Calvert AH Folate-Based Thymidylate Synthase Inhibitors as Anticancer Drugs. *Ann Oncol* 1995;6(9):871-881; Laohavinij S, Wedge SR, Lind MJ, et al. A phase I clinical study of the antipurine antifolate Lometrexol (DDATHF) given with oral folic acid. *Invest New Drugs* 1996;14:325-335; and Maughan TS, James RD, Kerr D, et al., on behalf of the British MRC Colorectal Cancer Working Party. Preliminary results of a multicenter randomized trial comparing 3 chemotherapy regimens (deGramont, Lokich, and raltitrexed) in metastatic colorectal cancer. *Proc ASCO* 1999;18:Abst 1007.)

[0004] Initially, folic acid was used as a treatment for toxicities associated with GARFTI see, e.g. U.S. Pat. No. 5,217,974. Folic acid has been shown to lower homocysteine levels (see e.g. Homocysteine Lowering Trialist's Collaboration. Lowering blood homocysteine with folic acid based supplements: meta-analysis of randomized trials. (*BMJ* 1998; 316:894-898 and Naurath HJ, Joosten E, Riezler R, Stabler SP, Allen RH', Lindenbaum J). The role of folic acid in modulating the toxicity and efficacy of the multitargeted antifolate LY 231514 (pemetrexed) was discussed in Worzalla *et al.* (*Anticancer Research* 18: 3235-3240 (1998) Worzalla JF, Chuan S and Schultz RM). EP-A-0546870 relates to nutrient compositions which are intended to prevent and cure infectious diseases and which are intended to be administered to patients being administered with anticancer drugs to prevent and treat infectious diseases due to immunosuppression induced by the anticancer drug therapy. The compositions of EP-A-0546870 are characterized in that they comprise a certain amount of retinoid compound(s) such as vitamin A which is indicated as being responsible for the immunoreactivity. Effects of vitamin B12, folate and vitamin B6 supplements in elderly people with normal serum vitamin concentrations (*Lancet* 1995;346:85-89), and homocysteine levels have been shown to be a predictor of cytotoxic events related to the use of GARFT inhibitors, see e.g. U.S. Pat. No. 5,217,974. However, even with this treatment, cytotoxic activity of GARFT inhibitors and antifolates as a class remains a serious concern in the development of antifolates as pharmaceutical drugs. The ability to lower cytotoxic activity would represent an important advance in the use of these agents.

[0005] Surprisingly and unexpectedly, we have now discovered that certain toxic effects such as mortality and non-hematologic events, such as skin rashes and fatigue, caused by antifolates, as a class, can be significantly reduced by the presence of a methylmalonic acid lowering agent as vitamin B12, without adversely affecting therapeutic efficacy. The present invention thus generally relates to a use in the manufacture of a medicament for improving the

therapeutic utility of antifolate drugs by administering to the host undergoing treatment with a methylmalonic acid lowering agent as vitamin B12. We have discovered that increased levels of methylmalonic acid is a predictor of toxic events in patients that receive an antifolate drug and that treatment for the increased methylmalonic acid, such as treatment with vitamin B12, reduces mortality and nonhematologic events, such as skin rashes and fatigue events previously associated with the antifolate drugs. Thus, the present invention generally relates to a use in the manufacture of a medicament for reducing the toxicity associated with the administration of an antifolate to a mammal by administering to said mammal an effective amount of said antifolate in combination with a methylmalonic acid lowering agent as vitamin B12.

[0006] Additionally, we have discovered that the combination of a methylmalonic acid lowering agent as vitamin B12 and folic acid synergistically reduces the toxic events associated with the administration of antifolate drugs. Although, the treatment and prevention of cardiovascular disease with folic acid in combination with vitamin B12 is known, the use of the combination for the treatment of toxicity associated with the administration of antifolate drugs was unknown heretofore.

[0007] Furthermore, the present invention generally relates to a use in the manufacture of a medicament for inhibiting tumor growth in mammals by administering to said mammals an effective amount of an antifolate in combination with a methylmalonic acid lowering agent.

[0008] Furthermore, the present invention generally relates to a use in the manufacture of a medicament for inhibiting tumor growth in mammals by administering to said mammals an effective amount of an antifolate in combination with a methylmalonic acid lowering agent and a FBP binding agent. A preferred FBP binding agent is folic acid.

[0009] Furthermore the present invention generally relates to the use of a methylmalonic acid lowering agent as vitamin B12, alone or in combination with a FBP binding agent, for lowering the mammalian toxicity of an antifolate. A preferred FBP binding agent is folic acid.

[0010] The invention specifically provides the use of the antifolate pemetrexed disodium in the manufacture of a medicament for use in combination therapy for inhibiting tumor growth in mammals wherein said medicament is to be administered in combination with a methylmalonic acid lowering agent selected from vitamin B12 and pharmaceutical derivatives thereof.

[0011] The invention also specifically provides the use of the antifolate pemetrexed disodium in the manufacture of a medicament for use in combination therapy for inhibiting tumor growth in mammals wherein said medicament is to be administered in combination with a methylmalonic acid lowering agent selected from vitamin B12 and pharmaceutical derivatives thereof and a FBP binding agent selected from folic acid, (6R)-5-methyl-5,6,7,8-tetrahydrofolic acid and (6R)-5-formyl-5,6,7,8-tetrahydrofolic acid or a physiologically available salt or ester thereof.

[0012] Furthermore, the present invention specifically relates to the use of a methylmalonic acid lowering agent selected from vitamin B12 and pharmaceutical derivatives thereof in the preparation of a medicament useful in lowering the mammalian toxicity associated with the antifolate pemetrexed disodium, and the medicament is administered in combination with the antifolate pemetrexed disodium.

[0013] Furthermore, the present invention specifically relates to the use of a methylmalonic acid lowering agent selected from vitamin B12 and pharmaceutical derivatives thereof in the preparation of a medicament useful in lowering the mammalian toxicity associated with the antifolate pemetrexed, disodium and the medicament is administered in combination with the antifolate pemetrexed disodium and a FBP binding agent selected from folic acid, (6R)-5-methyl-5,6,7,8-tetrahydrofolic acid and (6R)-5-formyl-5,6,7,8-tetrahydrofolic acid, or a physiologically available salt thereof.

[0014] Furthermore, the present invention specifically relates to the use of a methylmalonic acid lowering agent selected from vitamin B12 and pharmaceutical derivatives thereof in the manufacture of a medicament for use in a method of inhibiting tumor growth in mammals, which method comprises administering said methylmalonic acid lowering agent selected from vitamin B12 and pharmaceutical derivatives thereof in combination with the antifolate pemetrexed disodium,

[0015] Furthermore, the present invention specifically provides a product containing a methylmalonic acid lowering agent selected from vitamin B 12 and pharmaceutical derivatives thereof , the antifolate pemetrexed disodium and optionally a FBP binding agent selected from folic acid, (6R)-5-methyl-5,6,7,8-tetrahydrofolic acid and (6R)-5-formyl-5,6,7,8-tetrahydrofolic acid, or a physiologically available salt as a combined preparation for the simultaneous, separate or sequential use in inhibiting tumor growth.

[0016] The current invention concerns the discovery that administration of a methylmalonic acid lowering agent such as vitamin B 12 or a pharmaceutical derivative thereof, in combination with an antifolate drug such as pemetrexed disodium reduces the toxicity of the said antifolate drug.

[0017] The term "inhibit" as it relates to antifolate drugs refers to prohibiting, alleviating, ameliorating, halting, restraining, slowing or reversing the progression of, or reducing tumor growth.

[0018] As used herein, the term "effective amount" refers to an amount of a compound or drug, which is capable of performing the intended result. For example, an effective amount of an antifolate drug that is administered in an effort to reduce tumor growth is that amount which is required to reduce tumor growth.

[0019] As used herein, the term "toxicity" refers to a toxic event associated with the administration on an antifolate. Such events include, but are not limited to, neutropenia, thrombopenia, toxic death, fatigue, anorexia, nausea, skin rash,

infection, diarrhea, mucositis, and anemia. For further explanation of the types of toxicity experienced by patients receiving antifolates, see, generally, Antifolate Drugs in Cancer Therapy. Preferably, toxicity refers to toxic death, fatigue, neutropenia, thrombopenia, and mucositis.

[0020] As used herein, the term "nonhematologic event" refers to the occurrence of skin rash or fatigue due to the administration of an antifolate.

[0021] As used herein, the term "in combination with" refers to the administration of the vitamin B12 or pharmaceutical derivative, pemetrexed disodium, and optionally the folic acid; in any order such that sufficient levels of methylmalonic acid lowering agent and optionally folic acid are present to reduce the toxicity of an antifolate in a mammal. The administration of the compounds maybe simultaneous as a single composition or as two separate compositions or can be administered sequentially as separate compositions such that an effective amount of the agent first administered is in the patient's body when the second and/or third agent is administered. The antifolate drug pemetrexed disodium may be administered to the mammal first, followed by treatment with the methylmalonic acid lowering agent vitamin B12. Alternatively, the mammal may be administered the antifolate drug simultaneously with the methylmalonic acid lowering agent vitamin B12. Preferably, the mammal is pretreated methylmalonic acid lowering agent vitamin B12 and then treated with the antifolate pemetrexed disodium. If folic acid is to be administered in addition to the methylmalonic acid lowering agent vitamin B12, the folic acid may be administered at any time prior, post, or simultaneously to the administration of either the methylmalonic acid lowering agent vitamin B12 or the antifolate. Preferably, the mammal is pretreated with the methylmalonic acid lowering agent vitamin B12, and then treated with folic acid, followed by treatment with the antifolate pemetrexed disodium.

[0022] The terms "antifolate" and "antifolate drug" generally refer to a chemical compound which inhibits at least one key folate-requiring enzyme of the thymidine or purine biosynthetic pathways, preferably thymidylate synthase ("TS"), dihydrofolate reductase ("DHFR"), or glycinamide ribonucleotide formyltransferase ("GARFT"), by competing with reduced folates for binding sites of these enzymes. The "antifolate" or "antifolate drug" for use in this invention is Pemetrexed Disodium (ALIMTA®), as manufactured by Eli Lilly & Co.

[0023] The terms "methylmalonic acid" and "MMA" refer to a structural isomer of succinic acid present in minute amounts in healthy human urine.

[0024] The term "methylmalonic acid lowering agent" generally refers to a substrate, which lowers the concentration of methylmalonic acid in a mammal. A preferred example of such a substrate for use in the provided invention is vitamin B12. For methods of determining methylmalonic acid and substrates therefore, see, e.g., Matchar DB, Feussner JR, Millington DS, et al. Isotope dilution assay for urinary methylmalonic acid in the diagnosis of vitamin B 12 deficiency. A prospective clinical evaluation. *Ann Intern Med* 1987; 106: 707-710; Norman EJ, Morrison JA. Screening elderly populations for cobalamin (vitamin B12) deficiency using the urinary methylmalonic acid assay by gas chromatography mass spectrometry. *Am J Med* 1993; 94: 589-594; Norman EJ. Gas Chromatography mass spectrometry screening of urinary methylmalonic acid: early detection of vitamin B 12 (cobalamin) deficiency to prevent permanent neurologic disability. *GC/MS News* 1984; 12:120-129; Martin DC, Francis J, Protetch J, Huff FJ. Time dependency of cognitive recovery with cobalamin replacement: report of a pilot study. *JAGS* 1992; 40: 168-172; Norman EJ, Cronin C. Cobalamin deficiency. *Neurol* 1996; 47: 310-311; Rasmussen K, Moelby I, Jensen MK. Studies on methylmalonic acid in humans; Savage DG, Lindenbaum J, Stabler SP, Allen RH. Sensitivity of methylmalonic acid and total homocysteine determination for diagnosing cobalamin and folate deficiency. *Am J Med* 1994; 96: 239-246.

[0025] The "methylmalonic acid lowering agent" for use in this invention is selected from vitamin B12 (cyanocobalamin) and its pharmaceutical derivatives, such as hydroxocobalamin, cyano-10-chlorocobalamin, aquocobalamin perchlorate, aquo-10-chlorocobalamin perchlorate, azidocobalamin, chlorocobalamin, and cobalamin. Preferably the term refers to vitamin B 12, cobalamin, and chlorocobalamin.

[0026] The dosage used in the invention generally will be provided in the form of a vitamin supplement, namely as a tablet administered orally, such as a sustained release formulation, as an aqueous solution added to drinking water, or as an aqueous parenteral formulation. Preferably the methylmalonic acid lowering agent is administered as an intramuscular injection formulation. Such formulations are known in the art and are commercially available.

[0027] The skilled artisan will appreciate that the methylmalonic lowering agents for use in the invention are effective over a wide dosage range. For example, when cobalamin is used as the methylmalonic lowering agent, the dosage of cobalamin may fall within the range of 0.2 µg to 3000 µg of cobalamin from once daily for a month to once every nine weeks for a year. Preferably, cobalamin will be dosed as an intramuscular injection of 500 µg to 1500 µg administered from every 24 hours to every 1680 hours. Preferably, it is an intramuscular injection of about 1000 µg administered initially from 1 to 3 weeks prior to administration of the antifolate and repeated from every 24 hours to every 1680 hours, regardless of when treatment with the antifolate is started and continued until the administration of the antifolate is discontinued. Most preferred is an intramuscular injection of about 1000 µg administered initially from 1 to 3 weeks prior to the first administration of the antifolate and repeated every 6 to 12 weeks, preferably about every 9 weeks, and continued until the discontinuation of the antifolate administrations. However, it will be understood that the amount of the methylmalonic acid lowering agent actually administered will be determined by a physician, in the light of the relevant

circumstances, including the condition to be treated, the chosen route of administration, the actual agent administered, the age, weight, and response of the individual patient, and the severity of the patient's symptoms, and therefore the above dosage ranges are not intended to limit the scope of the invention in any way. In some instances dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effect.

[0028] The term "FBP binding agent" as used herein refers to a folic binding protein binding agent which includes folic acid, (6R)-5-methyl-5,6,7,8-tetrahydrofolic acid, and (6R)-5-formyl-5,6,7,8-tetrahydrofolic acid, or a physiologically-available salt or ester thereof. This latter compound is the (6R)-isomer of leucovorin as disclosed in J. Am. Chem. Soc., 74, 4215 (1952). Both of the tetrahydrofolic acid compounds are in the unnatural configuration at the 6-position. They are 10-20 fold more efficient in binding the folate binding protein compared with their respective (6S)-isomer, see Ratnam, et. al., Folate and Antifolate Transport in Mammalian Cells Symposium, Mar. 21-22, 1991, Bethesda, MD. These compounds are usually prepared as a mixture with their natural form (6S) of diastereomers by non-stereoselective reduction from the corresponding dehydro precursors followed by separation through chromatographic or enzymatic techniques. See e.g. PCT Patent Application Publication WO 880844 (also Derwent Abstract 88-368464/S1) and Canadian Patent 1093554. See, e.g. Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline (2000), 8 Folate, pp. 196-305.

[0029] "Physiologically-available salt" refers to potassium, sodium, lithium, magnesium, or preferably a calcium salt of the FBP binding agent. "Physiologically-available...ester" refers to esters which are easily hydrolyzed upon administration to a mammal to provide the corresponding FBP binding agent free acid, such as for example C₁-C₄ alkyl esters, and mixed anhydrides.

[0030] The FBP binding agent to be utilized according to this invention can be in its free acid form, or can be in the form of a physiologically-acceptable salt or ester which is converted to the parent acid in a biological system. The dosage generally will be provided in the form of a vitamin supplement, namely as a tablet administered orally, preferably as sustained release formulation, as an aqueous solution added to drinking water, an aqueous parenteral formulation, e.g., an intravenous formulation.

[0031] The FBP binding agent is, usually administered to the subject mammal prior to treatment with the antifolate pemetrexed disodium. Pretreatment with the suitable amount of FBP binding agent from 1 to 24 hours is usually sufficient to substantially bind to and block the folate binding protein prior to administration of the antifolate. Although one single dose of the FBP binding agent, preferably an oral administration of folic acid, should be sufficient to load the folate binding protein, multiple dosing of the FBP binding agent can be employed for periods up to weeks before treatment with the active agent to ensure that the folate binding protein is sufficiently bound in order to maximize the benefit derived from such pretreatment.

[0032] In the especially preferred embodiment of this invention, 0.1 mg to 30 mg, most preferably 0.3 mg to 5 mg, of folic acid is administered orally to a mammal about 1 to 3 weeks post administration of the methylmalonic acid lowering agent as vitamin B12 and 1 to 24 hours prior to the parenteral administration of the amount of antifolate pemetrexed disodium. However, it will be understood that the amount of the methylmalonic acid lowering agent as vitamin B12 actually administered will be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual agent administered, the age, weight, and response of the individual patient, and the severity of the patient's symptoms, and therefore the above dosage ranges are not intended to limit the scope of the invention in any way. In some instances dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effect.

[0033] In general, the term "pharmaceutical" when used as an adjective means substantially non-toxic to living organisms.

Methods

[0034] To assess the effect of a methylmalonic acid lowering agent as vitamin B12, alone or in combination with folic acid on the antitumor efficacy of antifolate pemetrexed disodium in a human tumor xenograft model, female nude mice bearing human MX-1 breast carcinoma were treated with pemetrexed disodium (ALIMTA[®]) alone or along with super-physiologic doses of folic acid or vitamin B12 (cobalamin).

[0035] The animals were maintained on sterilized standard lab chow ad libitum and sterilized water ad libitum. The human MX-1 tumor cells (5 x 10⁶) obtained from donor tumors were implanted subcutaneously in a thigh of female nude mice 8- to 10-weeks old. Beginning on day 7 post tumor cell implantation, the animals were treated with pemetrexed disodium (ALIMTA[®]) (100 mg/kg or 150 mg/kg) once daily on days 7 through 11 and 14 through 18 by intraperitoneal injection alone or along with folic acid (6 mg/kg or 60 mg/kg) and/or vitamin B12 (165 mg/kg) by intraperitoneal injection on the same schedule.

[0036] Tumor response was monitored by tumor volume measurements twice weekly over the course of the experiment.

Toxicity was monitored by body weight measurements made at the same time as the tumor volume measurements. Tumor growth delay was the difference in days for the treated and the controls tumors to reach 1000 mm³.

[0037] The human MX-1 breast carcinoma xenograft was responsive to treatment with pemetrexed disodium (ALIMTA[®]) with doses of 100 mg/kg and 150 mg/kg producing tumor growth delays of 17 days and 21 days, respectively. Folic acid was administered to the animals alone at two doses 6 mg/kg and 60 mg/kg on the same schedule as pemetrexed disodium (ALIMTA[®]) and produced tumor growth delays of 7 days and 12 days, respectively. Vitamin B12 administered alone at a dose of 165 mg/kg resulted in a tumor growth delay of 12 days.

Combinations of pemetrexed disodium (ALIMTA[®]) at each of the two doses were administered along with each of the vitamins as simultaneous combination regimens. Administration of folic acid (6 mg/kg) along with pemetrexed disodium (ALIMTA[®]) did not alter the tumor growth delay produced from that obtained with pemetrexed disodium (ALIMTA[®]) alone. The addition of folic acid at the higher dose (60 mg/kg) along with each dose of pemetrexed disodium (ALIMTA[®]) resulted in small increases in tumor growth delay to 22 days and 23 days at the pemetrexed disodium (ALIMTA[®]) doses of 100 mg/kg and 150 mg/kg, respectively. The tumor growth delays with pemetrexed disodium (ALIMTA[®]) and vitamin B12 (165 mg/kg) treatment were 22 days and 24 days at pemetrexed disodium (ALIMTA[®]) doses of 100 mg/kg and 150 mg/kg, respectively.

Body weight was used as a general measure of toxicity for each of the treatment regimens. The body weight loss pattern reflected the treatment regimens with weight decrease during the treatment times of days 7 through 11 and 14 through 18 with some weight recovery during the intervening two days. The weight loss due to pemetrexed disodium (ALIMTA[®]) was dose dependent but overall minor (3%).

Folic acid alone at either 6 mg/kg or 60 mg/kg did not cause weight loss, in fact folic acid treated animals maintained weight and gained weight over the course of the experiment better than the control animals. The animals treated with pemetrexed disodium (ALIMTA[®]) (100 mg/kg) and folic acid (60 mg/kg) gained weight (about 20%) over the course of the experiment.

[0038] Administration of vitamin B12 did not prevent weight gain in the animals over the time course of the experiment. The animals treated with pemetrexed disodium (ALIMTA[®]) (100 mg/kg) along with vitamin B12 gained weight while those treated with pemetrexed disodium (ALIMTA[®]) (150 mg/kg) along with vitamin B12 maintained weight over the course of the experiment.

[0039] In conclusion, administration of super-physiologic but non-toxic doses of the vitamins, folic acid and vitamin B12, did not alter the antitumor activity of pemetrexed disodium (ALIMTA[®]) in the human MX-1 breast carcinoma xenograft tumor in nude mice and did not increase the toxicity of pemetrexed disodium (ALIMTA[®]) as determined by body weight measurements of the animals.

[0040] The effect of vitamin B12, alone or in combination with folic acid, on antifolates can be demonstrated in standard tests commonly utilized to determine the antitumor activity and toxic effects of the antifolates themselves. In one such test, mice are inoculated with the C3H strain of mammary adenocarcinoma by inserting a 2 mm by 2 mm section of tumor into the axillary region of the mice by trocar. The timing of administering the methylmalonic acid lowering agent, alone or in combination with the folic acid, and the antifolate may be varied. Ten animals are used at each dosage level. Antitumor activity is assessed on day ten (when day one is first dosage of antifolate) by measuring the length and width of the tumor growth using vernier calipers, and the activity is expressed as a percent inhibition of tumor growth.

[0041] When the antifolate is administered to infected mice which are maintained on a diet totally free of vitamin B 12 and optionally folic acid for two weeks prior to and during treatment, it exhibits moderated antitumor activity at very low doses, but also causes severe toxicity at a very low dose (measured as death of mice).

[0042] A test group of mice are maintained on a vitamin B12 and optionally folic acid free diet for two weeks before treatment. Vitamin B 12 and optionally folic acid is then administered during the treatment by intramuscular injection of 0.0003% vitamin B12 (weight/volume) and optionally providing the animals drinking water containing 0.0003% folic acid (weight/volume). This concentration translates to about 1.75 mg of vitamin B12 and optionally folic acid per square meter of body surface per day. As the foregoing results indicate, addition of the indicated level of vitamin B12 to the diet of a subject receiving an antifolate results in excellent antitumor activity at low doses, with little or no toxic effects.

[0043] The foregoing tests establish that for tumor bearing mice maintained on a vitamin B12 and optionally folic acid free diet prior to and during treatment with an antifolate, the toxicity of the antifolate is very large, with 1 mg/kg/day being lethal to the majority of the mice, and lower antitumor activity is observed at non-toxic drug doses. Very low doses of vitamin B12 partially reverses drug toxicity and improved antitumor activity. Larger doses of vitamin B 12 reduce antifolate toxicity even more significantly. Pretreatment of the mouse with vitamin B12 and then administering folic acid prior to administering the antifolate demonstrates a striking reduction in toxicity, almost eliminating the antifolate toxicity completely. Thus, the use of vitamin B 12 in combination with an antifolate reduces drug toxicity without adversely affecting antitumor activity, and the use of vitamin B12 in conjunction with folic acid synergistically reduces drug toxicity.

[0044] In a typical clinical evaluation involving cancer patients, all of whom have histologically or cytologically confirmed diagnosis of cancer, an antifolate is administered in combination with vitamin B12. Vitamin B12 is administered as a 1000 µg intramuscular injection 1-3 weeks prior to treatment with the antifolate, and 1000 µg intramuscular injection of

EP 1 313 508 B1

vitamin B12 is made approximately every 9 weeks until the patient discontinues from therapy. The antifolate is administered in four doses over a two week period by rapid intravenous injection, followed by two weeks of non-therapy. Dosing is made on days 1, 4, 8 and 11 of any two week period. Patients will have an initial course of therapy at a dose of 5 mg/m²/dose, and depending upon the toxic effects observed in the initial course, their subsequent courses may be at the same dose, or may be escalated to 6 mg/m², or may be attenuated to 4 mg/m².

[0045] In preparation for the foregoing clinical study, pilot studies in humans have established that vitamin B12 given to patients receiving pemetrexed disodium (ALIMTA[®]) has effected reduced side effects due to the pemetrexed disodium (ALIMTA[®]). One to two weeks prior to administration of pemetrexed disodium (ALIMTA[®]) urine is collected and blood is drawn from a human subject; and vitamin metabolite levels, methylmalonic acid and homocysteine, are determined. Homocysteine levels are determined in blood by a fluorescent polarization immunoassay kit manufactured by Abbot Laboratories. Methylmalonic acid levels are determined by urine levels using a 24 hour urine collection kit available from Biolab Medical Unit (a United Kingdom company). Additionally urine and blood may be collected one week prior to administration of pemetrexed disodium (ALIMTA[®]) (after at least 5 days of folic acid supplementation and at least 1 week vitamin B 12 supplementation), and up to 4 days prior to every cycle.

Method of administration and dosing procedures :

1. Folic Acid:

[0046] Folic acid will be supplied as one of the following options, with preference in order from option #1 to option #3:

1. 350-600 µg folic acid.
2. A multivitamin containing folic acid in the range of 350 µg to 600 µg is acceptable if option #1 is not available.
3. A dose of folic acid between 350 µg and 1000 µg is acceptable if neither option #1 or option # 2 is available.

[0047] For purposes of this study, patients should take oral folic acid daily beginning approximately 1 to 3 weeks before treatment with pemetrexed disodium (ALIMTA[®]) plus cisplatin or cisplatin alone and continuing daily until discontinuation from study therapy.

2. Vitamin B12

[0048] Vitamin B12 will be obtained and administered as a 1000 µg intramuscular injection. A vitamin B12 injection must be administered approximately 1 to 3 weeks before treatment with pemetrexed disodium (ALIMTA[®]) and should be repeated approximately every 9 weeks until the patient discontinues from study therapy.

[0049] Folic acid supplementation, 350 - 600 µg or equivalent should be taken orally daily beginning approximately 1 to 3 weeks prior to the first dose of pemetrexed disodium (ALIMTA[®]) plus cisplatin and continue daily until the patient discontinues from study therapy. A vitamin B12 injection, 1000 µg, must be given intramuscularly approximately 1 to 3 weeks prior to the first dose of pemetrexed disodium (ALIMTA[®]) and should be repeated approximately every 9 weeks until the patient discontinues from study therapy.

[0050] Compare presupplementation homocysteine and methylmalonic acid levels to a) the level immediately prior to the initial dose of study drug, and b) to the level immediately prior to the second dose of study drug (i.e., after a full cycle of supplementation), and compare the prevalence of specific toxicities experienced in up to the first seven cycles of therapy in patients who have been supplemented from baseline to the prevalence seen in the earlier patients (n = 246) who were not supplemented (Farber et al.)

[0051] Toxicity may be compared in specific patients in non-supplemented cycles versus supplemented cycles (cross-over patients).

[0052] The data to be compared are:

- 1) Patient numbers and baseline demographic data for those supplemented from baseline.
- 2) Homocysteine and methylmalonic acid levels, levels at baseline, prior to first dose, prior to second dose, and prior to each therapy cycle depending of the type of cancer under study.
- 3) Grade 3 and 4 hematologic toxicity in these fully supplemented patients.
- 4) Grade 3 and 4 nonhematologic toxicity in these fully supplemented patients.

[0053] The grading of toxicities in chemotherapeutic clinical trials is well known to a person of skill in the art. Examples of fatigue and skin rash grading are provided below.

EP 1 313 508 B1

Fatigue Grading --

Neuromotor

- Grade 0 none or no change
- Grade 1 subjective weakness; no objective findings
- Grade 2 mild objective weakness without significant impairment of function
- Grade 3 objective weakness with impairment of function
- Grade 4 paralysis

Rash Grading --

Skin

- Grade 0 none or no change
- Grade 1 scattered macular or papular eruption or erythema that is asymptomatic
- Grade 2 scattered macular or papular eruption or erythema with pruritus or other associated eruption symptoms
- Grade 3 generalized symptomatic macular, papular, or vesicular eruption
- Grade 4 exfoliative dermatitis or ulcerating dermatitis

[0054] The vitamins (both folic acid and B12) to be used in the following studies may be obtained from Zenith Gold Line, Centrum, Folvite, or in Canada Apo-Folic. Cyanocobalamin is used as the methylmalonic acid lowering agent in these studies.

[0055] Current and past clinical trials show a 4% drug-related death total, 50% grade 3/4 neutropenia, 7% grade 4 thrombocytopenia, and 10% grade 3/4 diarrheas and mucositis in patients administered pemetrexed disodium (ALIMTA®) and folic acid as described in U.S. Pat. No. 5,217,974. Vitamin B12 supplementation with pemetrexed disodium (ALIMTA®) has a moderate effect on drug related toxicity, lowering drug related deaths to 3% and severe toxicities by about 25%. The combination of vitamin B12 and folic acid with pemetrexed disodium (ALIMTA®) has lowered the drug related deaths to <1% in over 480 so treated. The combination of vitamin B 12 and folic acid has lowered the drug related grade 3/4 toxic events, see Table 1.

Table 1

	Percent of occurrences prior to B 12/folic acid treatment (N=246)	Percent of occurrences post B 12/folic acid treatment (N=78)
Hematologic Toxicity/Non-Hematologic Toxicity	37%	6.4%
Neutropenia	32%	2.6%
Mucositis	5%	1,3%
Diarrhea	6%	2.6%
Neutropenia and Mucositis	3%	0%
Neutropenia and Diarrhea	3%	0%
Neutropenia and Infection	2%	0%

[0056] Additionally, sixty-two chemo-naïve patients requiring chemotherapeutic treatment were divided into two groups. Seventeen of these patients received pemetrexed disodium (ALIMTA®), but did not receive vitamin B12 or folic acid, as described *supra*. The remaining patients received treatment with vitamin B12, folic acid, and pemetrexed disodium (ALIMTA®), as described *supra*. Of patients who received the combination treatment, 8 out of 45 responded to the chemotherapy, Of patients who did not receive the combination treatment, but rather, received only treatment with pemetrexed disodium (ALIMTA®), only 1 out of 17 patients responded.

Claims

1. Use of pemetrexed disodium in the manufacture of a medicament for use in combination therapy for inhibiting tumor growth in mammals wherein said medicament is to be administered in combination with vitamin B12 or a pharmaceutical derivative thereof, said pharmaceutical derivative of vitamin B12 being hydroxocobalamin, cyano-10-chlorocobalamin, aquocobalamin perchlorate, aquo-10-chlorocobalamin perchlorate, azidocobalamin, chlorocobalamin or cobalamin.
2. Use according to claim 1 wherein said medicament is to be administered in combination with vitamin B12 or a pharmaceutical derivative thereof, said pharmaceutical derivative of vitamin B12 being hydroxocobalamin, cyano-10-chlorocobalamin, aquocobalamin perchlorate, aquo-10-chlorocobalamin perchlorate, azidocobalamin, chlorocobalamin or cobalamin, and a folic binding protein binding agent selected from folic acid, (6R)-5-methyl-5,6,7,8-tetrahydrofolic acid and (6R)-5-formyl-5,6,7,8-tetrahydrofolic acid or a physiologically available salt or ester thereof.
3. Use according to claim 2 wherein the folic binding protein binding agent is folic acid.
4. Use according to any one of claims 1 to 3 wherein the vitamin B12 or pharmaceutical derivative thereof is vitamin B12, cobalamin or chlorocobalamin.
5. Use according to any one of claims 1 to 3 wherein the vitamin B 12 or pharmaceutical derivative thereof is selected from vitamin B12 or hydroxocobalamin.
6. Use according to any one of claims 1 to 5 wherein the medicament, the vitamin B12 or pharmaceutical derivative thereof and optionally the folic binding protein binding agent are to be administered simultaneously, separately or sequentially.
7. Use according to any one of claims 1 to 6 wherein the medicament is to be administered after administration of the vitamin B12 or pharmaceutical derivative thereof.
8. Use according to any one of claims 1 to 7 wherein the medicament is to be administered after the folic binding protein binding agent
9. Use according to any one of claims 2 to 8 wherein the medicament is to be administered after pretreatment with the vitamin B12 or pharmaceutical derivative thereof followed by folic acid.
10. Use according to any one of claims 1 to 9 wherein vitamin B12 or pharmaceutical derivative thereof is to be administered as an intramuscular injection.
11. Use according to any one of claims 2 to 10 wherein the folic binding protein binding agent is to be administered orally as a tablet.
12. A product containing pemetrexed disodium, vitamin B 12 or a pharmaceutical derivative thereof said pharmaceutical derivative of vitamin B12 being hydroxocobalamin, cyano-10-chlorocobalamin, aquocobalamin perchlorate, aquo-10-chlorocobalamin perchlorate, azidocobalamin, chlorocobalamin or cobalamin, and, optionally, a folic binding protein binding agent selected from the group consisting of folic acid, (6R)-5-methyl-5,6,7,8-tetrahydrofolic acid and (6R)-5-formyl-5,6,7,8-tetrahydrofolic acid, or a physiologically available salt or ester thereof, as a combined preparation for the simultaneous, separate or sequential use in inhibiting tumor growth.
13. A product according to claim 12 wherein the vitamin B12 or pharmaceutical derivative thereof is vitamin B12, cobalamin or chlorocobalamin and, if present, the folic binding protein binding agent is folic acid.
14. A product according to claim 12 wherein the vitamin B 12 or pharmaceutical derivative thereof is vitamin B12 or hydroxocobalamin and, if present, the folic binding protein binding agent is folic acid.

Patentansprüche

1. Verwendung von Pemetrexednatrium zur Herstellung eines Arzneimittels zur Verwendung in einer Kombinations-

EP 1 313 508 B1

therapie zur Hemmung eines Tumorwachstums bei Säugern, worin das Arzneimittel in Kombination mit Vitamin B12 oder einem pharmazeutischen Derivat hiervon verabreicht werden soll, wobei das pharmazeutische Derivat von Vitamin B12 Hydroxocobalamin, Cyano-10-chlorcobalamin, Aquocobalaminperchlorat, Aquo-10-chlorcobalaminperchlorat, Azidocobalamin, Chlorcobalamin oder Cobalamin ist.

- 5 2. Verwendung nach Anspruch 1, worin das Arzneimittel verabreicht werden soll in Kombination mit Vitamin B12 oder einem pharmazeutischen Derivat hiervon, wobei das pharmazeutische Derivat von Vitamin B12 Hydroxocobalamin, Cyano-10-chlorcobalamin, Aquocobalaminperchlorat, Aquo-10-chlorcobalamin-perchlorat, Azidocobalamin, Chlorcobalamin oder Cobalamin ist, und mit einem Folsäurebindeproteinbindemittel, das ausgewählt ist aus Folsäure, (6R)-5-Methyl-5,6,7,8-tetrahydrofolsäure und (6R)-5-Formyl-5,6,7,8-tetrahydrofolsäure oder einem physiologisch annehmbaren Salz oder Ester hiervon.
- 10 3. Verwendung nach Anspruch 2, worin das Folsäurebindeproteinbindemittel Folsäure ist.
- 15 4. Verwendung nach einem der Ansprüche 1 bis 3, worin das Vitamin B12 oder das pharmazeutische Derivat hiervon Vitamin B12, Cobalamin oder Chlorcobalamin ist.
- 20 5. Verwendung nach einem der Ansprüche 1 bis 3, worin das Vitamin B12 oder das pharmazeutische Derivat hiervon aus Vitamin B12 oder Hydroxocobalamin ausgewählt ist.
- 25 6. Verwendung nach einem der Ansprüche 1 bis 5, worin das Arzneimittel, das Vitamin B12 oder ein pharmazeutisches Derivat hiervon und optional das Folsäurebindeproteinbindemittel simultan, separat oder sequenziell verabreicht werden sollen.
- 30 7. Verwendung nach einem der Ansprüche 1 bis 6, worin das Arzneimittel nach der Verabreichung des Vitamin B12 oder eines pharmazeutischen Derivats hiervon verabreicht werden soll.
- 35 8. Verwendung nach einem der Ansprüche 1 bis 7, worin das Arzneimittel nach dem Folsäurebindeproteinbindemittel verabreicht werden soll.
- 40 9. Verwendung nach einem der Ansprüche 2 bis 8, worin das Arzneimittel nach einer Vorbehandlung mit dem Vitamin B12 oder einem pharmazeutischen Derivat hiervon und gefolgt von Folsäure verabreicht werden soll.
- 45 10. Verwendung nach einem der Ansprüche 1 bis 9, worin das Vitamin B12 oder ein pharmazeutisches Derivat hiervon als eine intramuskuläre Injektion verabreicht werden soll.
- 50 11. Verwendung nach einem der Ansprüche 2 bis 10, worin das Folsäurebindeproteinbindemittel oral als eine Tablette verabreicht werden soll.
- 55 12. Produkt, das Pemetrexednatrium, Vitamin B12 oder ein pharmazeutisches Derivat hiervon enthält, wobei das pharmazeutische Derivat von Vitamin B12 Hydroxocobalamin, Cyano-10-chlorcobalamin, Aquocobalaminperchlorat, Aquo-10-chlorcobalaminperchlorat, Azidocobalamin, Chlorcobalamin oder Cobalamin ist, und das optional ein Folsäurebindeproteinbindemittel enthält, das aus der Gruppe ausgewählt ist, die besteht aus Folsäure, (6R)-5-Methyl-5,6,7,8-tetrahydrofolsäure und (6R)-5-Formyl-5,6,7,8-tetrahydrofolsäure oder einem physiologisch verfügbaren Salz oder Ester hiervon, als ein Kombinationspräparat zur simultanen, separaten oder sequenziellen Verwendung bei der Hemmung eines Tumorwachstums.
13. Produkt nach Anspruch 12, worin das Vitamin B12 oder ein pharmazeutisches Derivat hiervon Vitamin B12, Cobalamin oder Chlorcobalamin ist und, falls vorhanden, das Folsäurebindeproteinbindemittel Folsäure ist.
14. Produkt nach Anspruch 12, worin das Vitamin B12 oder ein pharmazeutisches Derivat hiervon Vitamin B12 oder Hydroxocobalamin ist und, falls vorhanden, das Folsäurebindeproteinbindemittel Folsäure ist.

Revendications

1. Utilisation du pemetrexed disodique dans la fabrication d'un médicament pour une utilisation dans une thérapie en combinaison destinée à inhiber la croissance tumorale chez des mammifères auxquels ledit médicament doit être

EP 1 313 508 B1

administré en combinaison avec la vitamine B12 ou un dérivé pharmaceutique de celle-ci, ledit dérivé pharmaceutique de la vitamine B12 étant l'hydroxocobalamine, la cyano-10-chlorocobalamine, le perchlorate d'aquocobalamine, le perchlorate d'aquo-10-chlorocobalamine, l'azidocobalamine, la chlorocobalamine ou la cobalamine.

- 5 **2.** Utilisation selon la revendication 1, dans laquelle ledit médicament doit être administré en combinaison avec la vitamine B12 ou un dérivé pharmaceutique de celle-ci, ledit dérivé pharmaceutique de la vitamine B12 étant l'hydroxocobalamine, la cyano-10-chlorocobalamine, le perchlorate d'aquocobalamine, le perchlorate d'aquo-10-chlorocobalamine, l'azidocobalamine, la chlorocobalamine ou la cobalamine, et un agent de liaison à la protéine fixant les folates choisi parmi l'acide folique, l'acide (6R)-5-méthyl-5,6,7,8,-tétrahydrofolique et l'acide (6R)-5-formyl-10 5,6,7,8,-tétrahydrofolique ou un sel ou un ester de ceux-ci physiologiquement acceptable.
- 3.** Utilisation selon la revendication 2, dans laquelle l'agent de liaison à la protéine fixant les folates est l'acide folique.
- 4.** Utilisation selon l'une quelconque des revendications 1 à 3, dans laquelle la vitamine B12 ou le dérivé pharmaceutique de celle-ci est la vitamine B12, la cobalamine ou la chlorocobalamine.
15
- 5.** Utilisation selon l'une quelconque des revendications 1 à 3, dans laquelle la vitamine B12 ou le dérivé pharmaceutique de celle-ci est choisi parmi la vitamine B12 ou l'hydroxocobalamine.
- 20 **6.** Utilisation selon l'une quelconque des revendications 1 à 5, dans laquelle le médicament, la vitamine B12 ou le dérivé pharmaceutique de celle-ci et, éventuellement l'agent de liaison à la protéine fixant les folates doivent être administrés simultanément, séparément ou successivement.
- 7.** Utilisation selon l'une quelconque des revendications 1 à 6, dans laquelle le médicament doit être administré après l'administration de la vitamine B12 ou du dérivé pharmaceutique de celle-ci.
25
- 8.** Utilisation selon l'une quelconque des revendications 1 à 7, dans laquelle le médicament doit être administré après l'agent de liaison à la protéine fixant les folates.
- 30 **9.** Utilisation selon l'une quelconque des revendications 2 à 8, dans laquelle le médicament doit être administré après un prétraitement avec la vitamine B12 ou un dérivé pharmaceutique de celle-ci, suivi par l'acide folique.
- 10.** Utilisation selon l'une quelconque des revendications 1 à 9, dans laquelle la vitamine B12 ou le dérivé pharmaceutique de celle-ci doit être administré sous la forme d'une injection intramusculaire.
35
- 11.** Utilisation selon l'une quelconque des revendications 2 à 10, dans laquelle l'agent de liaison à la protéine fixant les folates doit être administré oralement sous la forme d'un comprimé.
- 40 **12.** Produit contenant du pemetrexed disodique, de la vitamine B12 ou un dérivé pharmaceutique de celle-ci, ledit dérivé pharmaceutique de la vitamine B12 étant l'hydroxocobalamine, la cyano-10-chlorocobalamine, le perchlorate d'aquocobalamine, le perchlorate d'aquo-10-chlorocobalamine, l'azidocobalamine, la chlorocobalamine ou la cobalamine, et, éventuellement, un agent de liaison à la protéine fixant les folates choisi dans le groupe constitué de l'acide folique, l'acide (6R)-5-méthyl-5,6,7,8,-tétrahydrofolique et l'acide (6R)-5-formyl-5,6,7,8,-tétrahydrofolique, ou un sel ou un ester de ceux-ci physiologiquement acceptable, sous forme d'une préparation combinée pour l'utilisation
45 simultanée, séparée ou successive dans l'inhibition de la croissance tumorale.
- 13.** Produit selon la revendication 12, dans lequel la vitamine B12 ou le dérivé pharmaceutique de celle-ci est la vitamine B12, la cobalamine ou la chlorocobalamine et, le cas échéant, l'agent de liaison à la protéine fixant les folates est l'acide folique.
50
- 14.** Produit selon la revendication 12, dans lequel la vitamine B12 ou le dérivé pharmaceutique de celle-ci est la vitamine B12 ou l'hydroxocobalamine et, le cas échéant, l'agent de liaison à la protéine fixant les folates est l'acide folique.
55