

(19)



(11)

EP 2 921 180 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:

14.08.2019 Bulletin 2019/33

(51) Int Cl.:

A61K 47/10 ^(2017.01) **A61K 47/18** ^(2017.01)
A61K 47/02 ^(2006.01) **A61K 9/19** ^(2006.01)
A61P 7/04 ^(2006.01) **A61K 47/38** ^(2006.01)
A61K 38/37 ^(2006.01)

(21) Application number: **15154020.0**

(22) Date of filing: **22.02.2000**

(54) **Albumin-free factor VIII formulations**

Albuminfreie Faktor-VIII-Formulierungen

Formulations du facteur VIII sans albumine

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE

(30) Priority: **22.02.1999 US 255279**

01.12.1999 US 452752

(43) Date of publication of application:

23.09.2015 Bulletin 2015/39

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:

10075019.9 / 2 193 809
06077229.0 / 1 820 516
00907322.2 / 1 154 796

(73) Proprietors:

- **University of Connecticut**
Farmington, CT 06030-6207 (US)
- **Baxalta GmbH**
6300 Zug (CH)
- **Baxalta Incorporated**
Bannockburn, IL 60015 (US)

(72) Inventors:

- **Besman, Marc**
San Carlos, California 94070 (US)
- **Jameel, Feroz**
Newbury Park, CA California 91320 (US)
- **Bjornson, Erik**
Sherman Oaks
California 91423 (US)

• **Kashi, Ramesh**

Walnut, CA California 91798 (US)

• **Pikal, Michael**

Mansfield Center, CT Connecticut 06250 (US)

• **Tchessalov, Serguei**

U-92 Storrs, CT Connecticut 06269-2092 (US)

• **Carpenter, John**

Littleton, CO Colorado 80127 (US)

(74) Representative: **Hoffmann Eitle**

Patent- und Rechtsanwälte PartmbB
Arabellastraße 30
81925 München (DE)

(56) References cited:

EP-A1- 0 314 095 WO-A1-96/30041

• **ANGELICA FATOUROS ET AL:**

PHARMACEUTICAL RESEARCH, vol. 14, no. 12,
1 January 1997 (1997-01-01), pages 1679-1684,
XP055104627, ISSN: 0724-8741, DOI:
10.1023/A:1012163309468

• **OESTERBERG T ET AL: "DEVELOPMENT OF A**
FREEZE-DRIED ALBUMIN-FREE FORMULATION
OF RECOMBINANT FACTOR VIII SQ",
PHARMACEUTICAL RESEARCH, KLUWER
ACADEMIC PUBLISHERS, NEW YORK, NY, US,
vol. 14, no. 7, 1 January 1997 (1997-01-01), pages
892-898, XP009011546, ISSN: 0724-8741

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 921 180 B1

Description

BACKGROUND OF THE INVENTION

5 [0001] Factor VIII is a protein found in blood plasma which acts as a cofactor in the cascade of reactions leading to blood coagulation. A deficiency in the amount of Factor VIII activity in the blood results in the clotting disorder known as hemophilia A, an inherited condition primarily affecting males. Hemophilia A is currently treated with therapeutic preparations of Factor VIII derived from human plasma or manufactured using recombinant DNA technology. Such preparations are administered either in response to a bleeding episode (on-demand therapy) or at frequent, regular intervals to prevent uncontrolled bleeding (prophylaxis).

10 [0002] Factor VIII is known to be relatively unstable in therapeutic preparations. In blood plasma, Factor VIII is usually complexed with another plasma protein, von Willebrand factor (vWF), which is present in plasma in a large molar excess to Factor VIII and is believed to protect Factor VIII from premature degradation. Another circulating plasma protein, albumin, may also play a role in stabilizing Factor VIII *in vivo*. Currently marketed Factor VIII preparations therefore primarily rely on the use of albumin and/or vWF to stabilize Factor VIII during the manufacturing process and during storage.

15 [0003] The albumin and vWF used in currently marketed Factor VIII preparations is derived from human blood plasma, however, and the use of such material has certain drawbacks. Because a large molar excess of albumin compared to Factor VIII is generally added in order to increase the stability of the Factor VIII in such preparations, it is difficult to characterize the Factor VIII protein itself in these preparations. The addition of human-derived albumin to Factor VIII is also perceived as being a disadvantage with respect to recombinantly-produced Factor VIII preparations. This is because recombinantly-derived Factor VIII preparations, in the absence of such added albumin, would otherwise contain no human-derived proteins, and the theoretical risk of transmitting a virus would be reduced.

20 [0004] Several attempts to formulate Factor VIII without albumin or vWF (or with relatively low levels of these excipients) have been described. For example, U.S. Patent No. 5,565,427 (EP 508 194) to Freudenberg (assigned to Behringwerke) describes Factor VIII preparations which contain particular combinations of detergent and amino acids, specifically arginine and glycine, in addition to excipients such as sodium chloride and sucrose. The detergent, polysorbate 20 or polysorbate 80, is described as being present in amounts of between 0.001 to 0.5% (v/v), while arginine and glycine are present in amounts of between 0.01 to 1 mol/l. Sucrose is described as being present in amounts of between 0.1 and 10%. Example 2 of this patent asserts that solutions of (1) 0.75% sucrose, 0.4 M glycine, and 0.15M NaCl, and (2) 0.01 M sodium citrate, 0.08 M glycine, 0.016M lysine, 0.0025 M calcium chloride, and 0.4 M sodium chloride were not stable in solution over 16 hours, whereas solutions of (3) 1% sucrose, 0.14 M arginine, 0.1 M sodium chloride and (4) 1% sucrose, 0.4 M glycine, 0.14 M arginine, 0.1 M sodium chloride, and 0.05% Tween 80 exhibited stability.

25 [0005] U.S. Patent No. 5,763,401 (EP 818 204) to Nayer (assigned to Bayer) also describes a therapeutic Factor VIII formulation without albumin, comprising 15-60 mM sucrose, up to 50 mM NaCl, up to 5 mM calcium chloride, 65-400 mM glycine, and up to 50 mM histidine. The following specific formulations were identified as being stable: (1) 150 mM NaCl, 2.5 mM calcium chloride, and 165 mM mannitol; and (2) 1% sucrose, 30 mM sodium chloride, 2.5 mM calcium chloride, 20 mM histidine, and 290 mM glycine. A formulation containing higher amounts of sugar (10% maltose, 50 mM NaCl, 2.5 mM calcium chloride, and 5 mM histidine) was found to exhibit poor stability in the lyophilized state compared with formulation (2).

30 [0006] U.S. Patent No. 5,733,873 (EP 627 924) to Osterberg (assigned to Pharmacia & Upjohn) discloses formulations which include between 0.01 - 1 mg/ml of a surfactant. This patent discloses formulations having the following ranges of excipients: polysorbate 20 or 80 in an amount of at least 0.01 mg/ml, preferably 0.02 - 1.0 mg/ml; at least 0.1 M NaCl; at least 0.5mM calcium salt; and at least 1 mM histidine. More particularly, the following specific formulations are disclosed: (1) 14.7 - 50 - 65 mM histidine, 0.31 - 0.6 M NaCl, 4 mM calcium chloride, 0.001 - 0.02 - 0.025% polysorbate 80, with or without 0.1% PEG 4000 or 19.9 mM sucrose; and (2) 20 mg/ml mannitol, 2.67 mg/ml histidine, 18 mg/ml NaCl, 3.7 mM calcium chloride, and 0.23 mg/ml polysorbate 80.

35 [0007] Other attempts to use low or high concentrations of sodium chloride have also been described. U.S. Patent No. 4,877,608 (EP 315 968) to Lee (assigned to Rhone-Poulenc Rorer) teaches formulations with relatively low concentrations of sodium chloride, namely formulations comprising 0.5 mM - 15 mM NaCl, 5 mM calcium chloride, 0.2 mM - 5 mM histidine, 0.01 - 10 mM lysine hydrochloride and up to 10% sugar. The "sugar" can be up to 10% maltose, 10% sucrose, or 5% mannitol.

40 [0008] US 5,605,884 (EP 0 314 095) to Lee (assigned to Rhone-Poulenc Rorer) teaches the use of formulations with relatively high concentrations of sodium chloride. These formulations include 0.35 M -1.2 M NaCl, 1.5 - 40 mM calcium chloride, 1 mM - 50 mM histidine, and up to 10% of a "sugar" such as mannitol, sucrose, or maltose. A formulation comprising 0.45 M NaCl, 2.3 mM calcium chloride, and 1.4 mM histidine is exemplified.

45 [0009] International Patent Application WO 96/22107 to Roser (assigned to Quadrant Holdings Cambridge Limited) describes formulations which include the sugar trehalose. These formulations comprise: (1) 0.1 M NaCl, 15 mM calcium

chloride, 15 mM histidine, and 1.27 M (48%) trehalose; or (2) 0.011% calcium chloride, 0.12% histidine, 0.002% Tris, 0.002% Tween 80, 0.004% PEG 3350, 7.5% trehalose, and either 0.13% or 1.03% NaCl.

[0010] WO 9630041 A1 relates to a protein formulation comprising coagulation factor VIII or factor IX in an aqueous solution.

[0011] Other therapeutic Factor VIII formulations of the prior art generally include albumin and/or vWF for the purpose of stabilizing Factor VIII and are therefore not relevant to the present invention. For example, U.S. Patent No. 5,328,694 (EP 511 234) to Schwinn (assigned to Octapharma AG) describes a formulation which includes 100 - 650 mM disaccharide and 100 mM- 1.0 M amino acid. Specifically, the following formulations are disclosed: (1) 0.9 M sucrose, 0.25 M glycine, 0.25 M lysine, and 3 mM calcium chloride; and (2) 0.7 M sucrose, 0.5 M glycine, and 5 mM calcium chloride.

[0012] While several attempts have been made to formulate Factor VIII without albumin or vWF, there remains a need for therapeutic Factor VIII formulations which are stable in the absence of albumin or other proteins.

SUMMARY OF THE INVENTION

[0013] The present invention is defined by the appended claims. Also disclosed is the following:

The present disclosure relates to therapeutic Factor VIII compositions which are stable in the absence of albumin. In particular, the present disclosure comprises a Factor VIII composition comprising, in addition to Factor VIII: 4% to 10 % of a bulking agent selected from the group consisting of mannitol, glycine and alanine; 1% to 4% of a stabilizing agent selected from the group consisting of sucrose, trehalose, raffinose, arginine; 1 mM to 5 mM calcium salt; 100 mM to 300 mM NaCl; and a buffering agent for maintaining a pH of approximately between 6 and 8. This composition can additionally comprise a surfactant such as polysorbate 20, polysorbate 80, Pluronic F68, or Brij 35. When the surfactant is polysorbate 80, it should be present in an amount of less than 0.1%.

[0014] The buffer of the Factor VIII compositions according to the present disclosure is preferably present in a concentration of from 10 mM to 50 mM, and is preferably selected from the group consisting of histidine, Tris, BIS-Tris Propane, PIPES, MOPS, HEPES, MES and ACES. Advantageously, the buffering agent is either histidine or Tris. The Factor VIII composition of the present disclosure can further comprise an antioxidant.

[0015] The Factor VIII compositions of the present disclosure include both a bulking agent and a stabilizer. The bulking agent can be present in an amount of from about 6% to about 8%, preferably about 8%. The stabilizing agent is preferably present in an amount of about 2%. Sodium chloride is also present in these compositions, preferably in an amount of from 150 to 350 mM, and more preferably in an amount of about 225 mM. The calcium salt of the composition is also preferably calcium chloride, and the composition itself is preferably in lyophilized form.

[0016] In another embodiment, the present disclosure can comprise a Factor VIII composition formulated without adding albumin which includes the following excipients in addition to Factor VIII: 2% to 6 % hydroxyethyl starch; 1% to 4% of a stabilizing agent selected from the group consisting of sucrose, trehalose, raffinose, arginine; 1 mM to 5 mM calcium salt; 100 mM to 300 mM NaCl; and a buffering agent for maintaining a pH of approximately between 6 and 8.

[0017] Preferably, such a composition comprises about 4% hydroxyethyl starch, and the NaCl is present in an amount of 200 mM. The stabilizing agent is also preferably present in an amount of about 2%.

[0018] In a further embodiment, the present disclosure includes a Factor VIII composition, formulated without albumin, comprising: 300 mM to 500 mM NaCl; 1% to 4% of a stabilizing agent selected from the group consisting of sucrose, trehalose, raffinose, arginine; 1 mM to 5 mM calcium salt; and a buffering agent for maintaining a pH of approximately between 6 and 8. Preferably, the NaCl is present in a concentration of about 400 mM.

[0019] In yet another embodiment, the present disclosure comprises a process for lyophilizing an aqueous Factor VIII composition in a container using a lyophilizer, wherein the process comprises an initial freezing step, and the initial freezing step further comprises the steps of: (a) lowering the temperature of the lyophilizer chamber to at least about -45°C; (b) raising the temperature of the chamber to between about -15°C and -25°C; and subsequently (c) lowering the temperature of the chamber to at least about -45°C. In this process, the temperature of the chamber is preferably lowered or raised at a rate of between about 0.5°C and about 1.0°C per minute. In step (a), the temperature is preferably maintained for about 1 hour, and is lowered to about -55°C. In step (b) the temperature is preferably maintained between -15°C and -25°C for between 1 and 3 hours, and more preferably is at -22°C, and the temperature in step (c) is preferably maintained for about 1 hour. The Factor VIII composition used in this process preferably comprises between 4% and 10 % of an agent selected from the group consisting of mannitol, glycine and alanine, and also preferably comprises between 1% and 4% of an agent selected from the group consisting of sucrose, trehalose, raffinose, and arginine. In addition, the Factor VIII composition used in this process also preferably comprises between 100 mM and 300 mM NaCl.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

5 **[0020]** As used herein, the terms below and variations thereof shall be defined as follows, unless otherwise indicated:
Factor VIII - The Factor VIII molecule exists naturally and in therapeutic preparations as a heterogeneous distribution
of polypeptides arising from a single gene product (see, e.g., Andersson et al., Proc. Natl. Acad. Sci. USA, 83, 2979-2983,
May 1986). The term "Factor VIII" as used herein refers to all such polypeptides, whether derived from blood plasma or
10 produced through the use of recombinant DNA techniques. Commercially available examples of therapeutic preparations
containing Factor VIII include those sold under the trade names of HEMOFIL M and RECOMBINATE (available from
Baxter Healthcare Corporation, Deerfield, Illinois, U.S.A.). Other preparations currently in development comprise primarily
a single subpopulation of Factor VIII molecules which lack the B domain portion of the molecule.

[0021] *International Unit, IU* - International Unit, or IU, is a unit of measurement of the blood coagulation activity
15 (potency) of Factor VIII as measured by a standard assay, such as one of the following:

One stage assay. One stage assays are known to the art, such as that described in Lee, Martin L, et al., An Effect
of Predilution on Potency Assays of Factor VIII Concentrates, Thrombosis Research (Pergamon Press Ltd.) 30,
511-519 (1983).

Chromogenic assay. Chromogenic assays may be purchased commercially, such as the Coatest Factor VIII, avail-
20 able from Chromogenix AB, Molndal, Sweden.

[0022] *Anneal* -The term anneal shall be used to indicate a step in the lyophilization process of a pharmaceutical
preparation undergoing lyophilization, prior to the freeze-drying of the preparation, in which the temperature of the
preparation is raised from a lower temperature to a higher temperature and then cooled again after a period of time.

25 **[0023]** *Bulking Agent* - For the purposes of this application, bulking agents are those chemical entities which provide
structure to the "cake" or residual solid mass of a pharmaceutical preparation after it has been lyophilized and which
protect it against collapse. A crystallizable bulking agent shall mean a bulking agent as described herein which can be
crystallized during lyophilization, other than sodium chloride. HES is not included in this group of crystallizable bulking
agents.

30 **[0024]** *Freeze-drying, freezing, lyophilizing* - "Freeze-drying," unless otherwise indicated by the context in which it
appears, shall be used to denote the portion of a lyophilization process in which the temperature of a pharmaceutical
preparation is raised in order to drive water out of the preparation. The "freezing" steps of a lyophilization process are
those steps which occur prior to the freeze-drying stage. "Lyophilizing," unless otherwise indicated, shall refer to the
entire process of lyophilization, including both the freezing steps and the freeze-drying steps.

35 **[0025]** Unless otherwise noted, percentage terms express weight/volume percentages and temperatures are in the
Celsius scale.

Formulation Components

40 **[0026]** The liquid Factor VIII compositions of the present invention include bulking agents, stabilizing agents, buffering
agents, sodium chloride, calcium salts, and, advantageously, other excipients. These excipients have been chosen in
order to maximize the stability of Factor VIII in lyophilized preparations. However, the Factor VIII compositions of the
present invention exhibit stability in the liquid state as well.

[0027] The bulking agents used in the present formulations, which form the crystalline portion of the lyophilized product
45 (except in the case of HES), are selected from the group consisting of mannitol, glycine, alanine, and hydroxyethyl starch
(HES). Mannitol, glycine, or alanine are present in an amount of 4 - 10%, preferably 6 - 9%, and more preferably about
8%. When HES is used as a bulking agent, it is present in an amount of 2 - 6%, preferably 3 - 5%, and more preferably
about 4%.

[0028] The stabilizing agents used in the liquid formulations of the present invention are selected from the group
50 consisting of sucrose, trehalose, raffinose, and arginine. These agents are present in the liquid formulations of the
present invention in an amount of between 1 - 4%, preferably 2 - 3%, more preferably about 2%. Sorbitol and glycerol
were evaluated as possible stabilizers but were found to be poor stabilizers in the present formulations.

[0029] Sodium chloride is included in the present formulations in an amount of 100 - 300 mM, preferably 150 - 250
mM, and most preferably about 225 mM. In one embodiment of the present invention, sodium chloride itself can be used
55 without any of the aforementioned bulking agents, in which case it would be included in the liquid formulation in an
amount of between 300 mM and 500 mM NaCl, preferably 350 to 450 mM NaCl, and more preferably about 400 mM NaCl.

[0030] In addition, buffers are present in these formulations, because it is believed that the Factor VIII molecule can
be adversely affected by pH shifts during lyophilization. The pH should preferably be maintained in the range of between

EP 2 921 180 B1

6 and 8 during lyophilization, and more preferably at a pH of about 7. The buffering agent can be any physiologically acceptable chemical entity or combination of chemical entities which have the capacity to act as buffers, including histidine, Tris, BIS-Tris Propane, PIPES, MOPS, HEPES, MES and ACES. The full chemical designations of these buffering agents is listed in Table 1 below. Typically, the buffering agent is included in a concentration of 10 - 50 mM. When histidine is added to the formulations, concentrations of over 20 mM and preferably about 25 mM are used, alone or in combination with other buffers such as Tris. Histidine is especially preferred for use in the liquid compositions of the present invention, as described in greater detail below.

Table 1 - Buffering Agents

Tris	tris-(hydroxymethyl)-aminomethane
BIS-Tris Propane	1,3-bis-[tris-(hydroxy-methyl)methylamino]-propane
PIPES	piperazine-N,N'-bis-(2-ethanesulfonic acid)
MOPS	3-(N-morpholino) propanesulfonic acid
HEPES	N-2-hydroxyethyl-piperazine-N'-2-ethanesulfonic acid
MES	2-(N-morpholino) ethanesulfonic acid
ACES	N-2-acetamido-2-aminoethanesulfonic acid
In order to preserve the activity of Factor VIII, it is important that the	

liquid formulations of the present invention also include calcium or another divalent cation able to interact with Factor VIII and maintain its activity, presumably by maintaining the association of the heavy and light chains of Factor VIII. Between 1 mM and 5 mM of a calcium salt can be used, more preferably 3 - 4 mM, and most preferably about 4 mM. The calcium salt is preferably calcium chloride, but can also be other calcium salts such as calcium gluconate, calcium gluconate, or calcium gluceptate.

[0031] The liquid Factor VIII compositions of the present invention also preferably include a surfactant, preferably in an amount of 0.1 % or less, and more preferably in an amount of about 0.03%. The surfactant can, for example, be chosen from the group consisting of polysorbate 20, polysorbate 80, pluronic polyols, and Brij 35 (polyoxyethylene 23 lauryl ether). Several grades of pluronic polyols (sold under the trade name Pluronic, manufactured by the BASF Wyandotte Corporation) are available. These polyols, of diversified molecular weight (from 1,000 to over 16,000) and physicochemical properties have been used as surfactants. Pluronic F-38, of a molecular weight of 5,000 and Pluronic F-68, molecular weight 9,000, both contain (by weight) 80 per cent hydrophilic polyoxyethylene groups and 20 percent hydrophobic polyoxypropylene groups. Tween-80, a commercial polysorbate, however, is preferred in the present formulations, in particular vegetable-derived Tween-80.

[0032] The liquid Factor VIII formulations of the present invention also preferably include an antioxidant. The addition of antioxidants to the lyophilized formulations of the invention has been found to improve the stability of these formulations, and thus extend their shelf lives. The antioxidants used must be compatible for use with a pharmaceutical preparation, and in addition are preferably water soluble. When adding antioxidants to a formulation, it is preferable to add such antioxidants as late in the process prior to lyophilization as possible, in order to avoid spontaneous oxidation of the antioxidant. Table 2 below lists suitable antioxidants, which are available commercially through companies such as Calbiochem and Sigma.

Table 2 - Antioxidants

N-Acetyl-L-Cysteine / Homocysteine
Glutathione
6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox)
Lipoic acid
Methionine
Sodium Thiosulfate
Platinum
Glycine-glycine-histidine (tripeptide)

(continued)

Butylatedhydroxytoluene (BHT)

5 [0033] Of the foregoing antioxidants, glutathione is preferred. Concentrations in the range of about 0.05 mg/ml to more than 1.0 mg/ml have all been found to enhance the stability of Factor VIII compositions, and it is believed that higher concentrations would also be useful (up to the point of any toxic effects or adverse manufacturing effects, such as a depression of the glass transition temperature of the lyophilized product).

10 [0034] It has been found in particular that the combination of histidine and glutathione produces synergistically beneficial effects on the stability of Factor VIII compositions. Histidine, while acting as a buffer, can also act as a metal chelator. To the extent that Factor VIII inactivation is caused by metal-induced oxidation, histidine can therefore act to stabilize Factor VIII by binding such oxidizing metal ions. It is believed that by binding these metals, the glutathione (or indeed any other antioxidant present) is thereby able to provide further antioxidative protection, since the oxidative effect of the metal ions bound by the histidine has been contained.

15 [0035] Other chelating agents might also be used in the liquid compositions of the present invention. Such agents should preferably bind metals such as copper and iron with greater affinity than calcium, if a calcium salt is being used in the composition. One such chelator is deferoxamine, a chelating agent that facilitates the removal of Al⁺⁺ and iron. Deferoxamine Mesylate, C₂₅H₄₈N₆O₈·CH₄O₃S, can be obtained from Sigma (Sigma Prod, No. D9533). It is an aluminum and iron(II) chelator which chelates iron (as a 1:1 chelate complex) only in the +3 oxidation state, not +2 oxidation state, and can also bind manganese ion and other metals. Deferoxamine can be used advantageously in an amount of 0.25 mg/l.

20 [0036] The Factor VIII used in the present formulations can be either highly purified human plasma-derived Factor VIII or more preferably can be recombinantly produced Factor VIII. Recombinant Factor VIII can be produced by Chinese hamster ovary (CHO) cells transfected with a vector carrying a DNA sequence coding for the Factor VIII molecule. Methods for creating such transfected CHO cells are described, *inter alia*, in U.S. Patent No. 4,757,006 to Toole, Jr., though alternative methods are also known to the art (see, e.g., U.S. Patent No. 4,868,112, also to Toole, Jr., and PCT International Application WO-A-91/09122). The methods used to culture such CHO cells to produce Factor VIII are also known to the art, for example in European Patent Application No. 0 362 218 to Genetics Institute, entitled "Improved method for producing Factor VIII:C-type proteins." Recombinant Factor VIII can, however, also be produced in other cell lines, such as baby hamster kidney (BHK) cells. The Factor VIII molecule itself, if recombinantly produced, can be either full-length Factor VIII or a deletion derivative thereof, such as a B domain-deleted Factor VIII molecule.

25 [0037] While the Factor VIII compositions described in this application can be lyophilized and reconstituted in the indicated concentrations, one of skill in the art will understand that these preparations can also be reconstituted in more dilute form. For example, a liquid preparation according the present invention which is lyophilized and/or normally reconstituted in 2 ml of solution can also be reconstituted in a larger volume of diluent, such as 5 ml. This is particularly appropriate when the Factor VIII preparation is being injected into a patient immediately, since in this case the Factor VIII is less likely to lose activity, which may occur more rapidly in more dilute solutions of Factor VIII.

40 Formulation and Lyophilization Development

[0038] In order to achieve maximal stability, the liquid Factor VIII compositions of the present invention are preferably lyophilized. During lyophilization, Factor VIII is converted from being in an aqueous phase to being in an amorphous solid phase, which is thought to protect the protein from chemical and/or conformational instability. The lyophilized preparation not only contains an amorphous phase, but also includes a component which crystallizes during lyophilization. This is thought to allow the rapid lyophilization of the Factor VIII composition and the formation of a more elegant cake (that is, a cake with minimal shrinkage from the sides of the container in which it was lyophilized). In the liquid formulations of the present invention, the stabilizing agents have been selected to exist primarily in an amorphous phase of the lyophilized product, while the bulking agents (except HES) have been selected to crystallize during freezing.

45 [0039] Both the Factor VIII and the stabilizer are preferably dispersed in the amorphous phase of the lyophilized cake. The mass of the stabilizer is also preferably large compared to the other excipients in the amorphous form. In addition, the apparent glass transition temperature (T_g') of the amorphous phase is preferably relatively high during freeze-drying, and the glass transition temperature (T_g) of the solid is likewise preferably high during storage. Crystallization of sodium chloride in the product was found to be desirable, since amorphous sodium chloride will depress the T_g' of the amorphous phase.

50 [0040] In order to avoid the collapse of the cake of a particular composition, primary drying is preferably carried out at a product temperature below the apparent glass transition temperature of the freeze concentrate. An increase in drying time may also be required to offset a decrease in T_g' . Further information on lyophilization may be found in

EP 2 921 180 B1

Carpenter, J.F. and Chang, B.S., Lyophilization of Protein Pharmaceuticals, Biotechnology and Biopharmaceutical Manufacturing, Processing and Preservation, K.E. Avis and V.L. Wu, eds. (Buffalo Grove, IL: Interpharm Press, Inc.), pp. 199-264 (1996).

5 **Example 1**

10 **[0041]** The effects of the concentration of Factor VIII and of the addition of a stabilizer on the recovery of Factor VIII were investigated in several studies. These studies were performed using mannitol as a model bulking agent and sucrose as a model stabilizer. The three sample formulations described in Table 3 below were used in these studies. All formulations used in these studies included 10 mM Tris, 200 mM NaCl, 8% mannitol, 4 mM CaCl₂, and 0.02% Tween-80 and were conducted at pH 7.0.

15 **Table 3**

Sample I.D.	Initial Factor VIII (IU/ml)	Sucrose %
IA	600	-
IB	60	-
IC	60	2

20 **[0042]** These samples were lyophilized using the freeze-drying cycle shown in Table 4 below in order to maintain a product temperature below the apparent glass transition temperature (T_g). Differential scanning calorimetric (DSC) studies indicated the presence of a transition at approximately -40°C in the mannitol formulations. In order to maintain a product temperature below this value, the shelf temperature was set to -32°C during primary drying. Primary drying under these conditions was performed for about 55 hours, with a total cycle time of about 80 hours.

25 **Table 4**

Freezing/Processing Method	Description
I (Freezing)	Cool to +5°C; Cool to -5°C at 1°C/minute, hold for 20 minutes; Cool to -20 ±5°C at 1°C/minute, hold for 1 hour (up to 3 hours); Cool to -45°C at 0.5°C/minute, hold for 1 hour.
II	Freeze per method I Hold at -35°C for 48 hours.
III	Freeze per method I Hold at -35°C for 48 hours; Hold at -20°C for 48 hours.
IV (Freeze-drying)	Shelf -32°C during primary drying for about 55 hours (up to 100 hours); Product < -40°C during primary drying; Ramp from -32°C to +40°C at 0.2°C/minute; Shelf +40°C during secondary drying for 3 hours.

45 **[0043]** The Factor VIII activity of these samples, as determined by the one-stage clotting assay, was compared against a control held at -45°C. The assay results are shown in Table 5 below.

50 **Table 5**

Processing Method	% Loss in Factor VIII Activity During Each Step		
	Formulation IA (600 IU/ml)	Formulation IB (60 IU/ml)	Formulation IC (60 IU/ml, 2% Sucrose)
I	6.7	37.5	41.7
II	2.0	9.3	3.9
III	7.3	11.6	5.0

EP 2 921 180 B1

(continued)

Processing Method	% Loss in Factor VIII Activity During Each Step		
IV (Lyophilization)	20.0	24.2	18.3

[0044] These results indicate that protein concentration has an effect on the recovery of Factor VIII during freezing. Formulations containing 60 IU/ml lost approximately 37%-42% of the initial Factor VIII activity during the freezing step, while 6.7% of Factor VIII activity was lost for the formulation containing 600 IU/ml. These results indicate that a higher protein concentration has a protective effect during freezing. Although sucrose provided some protection to the Factor VIII during the intermediate temperature holds as well as during freeze-drying, it failed to protect the protein during the initial freezing step.

Example 2

[0045] Following the development of the lyophilization process outlined in Example 1, further optimization of this process was undertaken. It has been found that a lyophilized composition having a higher glass transition temperature, (and, theoretically, better Factor VIII stability) can be produced by: (1) lowering the freezing temperature initially to -45°C or lower (such as down to about -50°C or -55°C); (2) raising the temperature to -20°C or -22°C (±5°C); and then (3) lowering the temperature again to 45°C or lower. The temperature is lowered or raised, as the case may be, at a rate of between about 0.5°C and about 1.0°C per minute. Once the desired temperature is reached, the composition is held at that temperature for between 1 and 3 hours. This improved freezing cycle is shown in Table 6 below.

Table 6

Freezing Method	Description
I	Cool to +5°C; Cool to -5°C at 0.5-1°C/minute, hold for 20 minutes; Cool to between -55°C and -45°C at 0.5-1°C/minute, hold for about 1 hour; Warm to -22°C (±5°C) at 0.5-1°C/minute, hold for 1 to 3 hours; Cool to -45°C at 0.5-1°C/minute, hold for about 1 hour.

[0046] Unless otherwise indicated, the temperatures referred to in this example and in other examples refer to the shelf temperature of the lyophilizer and not to the temperature of the product per se. Following the improved freezing cycle, the remainder of the lyophilization process can be conducted as outlined in Example 1 above, or otherwise as described further herein or as determined by one of skill in the art.

[0047] This improved lyophilization process was found to be useful for formulations which include glycine as the bulking agent as well as those which use mannitol. It is further believed to have applicability to liquid formulations which make use of the other bulking agents of the present invention as well.

Example 3

[0048] It is believed that in order to produce a freeze-dried product with acceptable cake appearance and glass transition temperature, the bulking agent of lyophilized pharmaceutical preparations which contain sodium chloride, such as glycine or mannitol, may need to be crystallized. The following improved lyophilization process for crystallizable bulking agents was therefore developed.

Table 7a - Freezing Steps

Process Step	Temperature	Duration of Step
Initial freezing	-40°C or less	1 hour
First annealing	between -23°C and -27°C	3 hours
Second freezing	-55°C	1 hour
Second annealing	-36°C	4 hours
Third freezing	-50°C	1 hour

Table 7b - Freeze-Drying Steps

Process Step	Temperature	Duration of Step
Primary Drying	-35°C	up to 100 hours
Secondary Drying: First step	40°C	3 hours
Secondary Drying: Second step	45°C	3 hours
Secondary Drying: Third step	50°C	3 hours

[0049] In the freezing steps, changes in the temperatures occurred at a rate of between about 0.5°C/minute and 1°C/minute. It is believed that steps of longer duration would also be effective.

[0050] Prior to the first freezing step, the temperature is brought to between about 2°C and 8°C for about one hour for the purpose of bringing all the vials to approximately the same temperature. After this the lyophilizer is cooled to -5°C. The first freezing step should be performed at a temperature less than -30°C, preferably below -35°C, and more preferably at about -40°C. Following this, the first annealing step should occur at a temperature of between -30°C and -19°C, more preferably either between about -25°C and -28°C (if glycine is the bulking agent) or between -21°C and -24°C (if mannitol is the bulking agent), with the temperatures of -23°C and -26°C being most preferred, at which temperatures it is believed that the crystallizable bulking agents crystallize, at least in part. However, the lower range around -27°C is not recommended for formulations containing mannitol and arginine. This step is preferably carried out for about 3 hours.

[0051] Following the first annealing step, the temperature is lowered, preferably to less than about -50°C and more preferably to less than -55°C, for about 1 hour. It is believed that the sodium chloride in the preparation nucleates at this time.

[0052] During the second annealing step, the temperature of the pharmaceutical preparation is raised to between about -30°C and -39°C, and preferably to about -33°C for mannitol-containing compositions and -36°C for glycine-containing compositions. It is believed that NaCl crystal growth occurs at this time, at least in part. This step is preferably conducted for about 4 hours. Following this, the temperature of the lyophilizer is reduced to about -50°C, preferably for about 1 hour in order to reduce the temperature of the preparation.

[0053] In the freeze-drying steps which follow, changes in temperature occurred at a rate of between about 0.1°C/minute and 0.5°C/minute. After reducing the pressure in the lyophilizer to about 65 mTorr, the temperature is raised to between about -32°C and -35°C for primary drying. Ice crystals in the preparation will sublime at this temperature. This step is performed for up to about 100 hours, or until most of the ice has been sublimated from the preparation. The point at which most of the ice has sublimated can be determined, for example, using a dewpoint sensor, which indicates the end of the sublimation of ice when the readings decrease (the point of inflection).

[0054] Following primary drying, the temperature is raised to +40°C, preferably at a rate of 0.2°C/minute, to initiate secondary drying to remove further water from the preparation. This temperature is preferably maintained for about three hours. Second and third secondary drying steps follow this first step, where the temperature is raised to about +45°C for about three hours and then to about +50°C for three more hours in order to reduce the moisture in the lyophilized cake to less than 2% (w/w).

Example 4

[0055] Further studies were performed to examine specifically the effect of histidine on lyophilized Factor VIII compositions containing glycine or mannitol as bulking agents. Non-reversing heat flow (Modulated DSC, mDSC) was used to detect the crystallization of these bulking agents during cooling. Both the temperature of crystallization and the total heat of crystallization were determined from the crystallization exotherm. The appearance of the NaCl eutectic melt endotherm during warming was used to detect NaCl crystallization. In mDSC, the extent of crystallization was determined as the ratio of the enthalpy of melting of the formulation to the enthalpy of melting of pure NaCl solution by using the total heat flow signal. In addition, X-ray diffraction analyses were performed in order to determine the extent of crystallization in the lyophilized formulations. While histidine concentrations less than 20 mM did not significantly impact the crystallization of glycine, 50 mM histidine reduced the extent of glycine crystallization. Well-defined NaCl crystallization exotherms were not observed during cooling of formulations containing glycine. However, eutectic melting endotherms during heating indicated that NaCl was crystallized (> 50%) after cooling lower than -50°C and annealing at -30°C, -35°C and -40°C. The inclusion of 50 mM histidine in the glycine-containing formulation retarded NaCl crystallization. Consequently, the annealing time was increased 3-fold for such formulations in order to achieve an equivalent crystallinity.

[0056] However, the effect of 20 mM histidine on the crystallization of NaCl in the glycine-containing formulations was

minimal. In freeze-drying studies, collapse of the lyophilized cake was observed visually in glycine-containing formulations containing 50 mM histidine. X-ray powder diffraction data indicated a decrease in the crystallinity of NaCl in samples containing histidine. In mannitol-containing formulations, typically 83% - 90% of the sodium chloride crystallized during cooling between -40°C and -50°C without the need for annealing. While inclusion of 20 mM histidine to the formulation suppressed NaCl crystallization during cooling, annealing resulted in approximately 40% crystallization of the NaCl.

[0057] Therefore, in formulations containing a crystallizable bulking agent, such as glycine or mannitol, and NaCl, the inclusion of histidine may decrease the extent of crystallization of NaCl. Although this could in some cases lead to the collapse of the cake which is formed during lyophilization, the use of relatively lower concentrations of histidine in such formulations can mitigate this effect. Nonetheless, acceptable cakes have been formed with concentrations of histidine of 35 mM and 50 mM. Histidine may also be preferable to HEPES as a buffer in mannitol- and glycine-based formulations, as the use of HEPES has been observed to lower the T_g to a greater extent than a similar amount of histidine.

Example 5

[0058] The physical characteristics of a number of potential Factor VIII formulations, including seven candidate stabilizers and five bulking agents, were evaluated in another study. In addition to a bulking agent and stabilizer, all formulations listed in Table 8 below (except for formulation 11) contained 10 mM Tris•HCl, 200 mM NaCl, 0.02% Tween-80, 4 mM CaCl₂ and were at pH 7.0. Formulation 11 contained 10 mM Tris•HCl, 0.02% Tween-80, and 4 mM CaCl₂, also at pH 7.0. All pH measurements were performed at ambient temperature.

Table 8

Sample I.D.	Bulking Agent	Protein Stabilizer
1	8% Mannitol	2% Sucrose
2	8% Mannitol	2% Trehalose
3	8% Mannitol	2% Raffinose
4	8% Mannitol	2% Arginine
5	8% Mannitol	2% Lysine
6	8% Mannitol	2% Sorbitol
7	8% Mannitol	2% Glycerol
8	4% Hydroxyethyl Starch	2% Sucrose
9	8% Glycine	2% Sucrose
10	8% Glycine	2% Trehalose
11	400 mM NaCl	2% Sucrose
12	8% Alanine	2% Sucrose

[0059] Collapse temperature measurements by freeze-dry microscopy and thermal transition measurements by DSC were used to predict freeze-drying behavior. DSC, X-ray powder diffraction and polarized light microscopy were also used to determine the crystallinity of the lyophilized samples. The reconstitution time and the appearance of the samples were also evaluated. The results of all of these measurements are summarized in Table 9 below.

Table 9

Sample I.D.	T_{pc} (°C)	T_c (°C)	T_g (°C)	Reconstitution (seconds)	Water Content (%)	Appearance
1	-14	-10	54	64	n/c	Elegant
2	-20	-15	53	62	1.4	Top partially collapsed
3	-15	-10	54	77	1.7	Elegant
4	-	-	-	-	-	Partial collapse
5	-	-	-	-	-	Collapsed
6	n/c	n/c	< 10°C*	63	0.6	Elegant
7	-	-	< 10°C*	-	-	Elegant
8	-	-	86	49	0.7	Elegant but shrunk from sides
9	-	-	54	22	0.8	Elegant

EP 2 921 180 B1

(continued)

Sample I.D.	T _{pc} (°C)	T _c (°C)	T _g (°C)	Reconstitution (seconds)	Water Content (%)	Appearance
10	-	-	63	18	-	Elegant
11	-	-	66	11	0.4	Elegant (layer on bottom)
12	-	-	-	57	0.5	Elegant

Sorbitol and Glycerol have glass transitions at < 10°C. The DSC scan range did not include temperatures in this range.
n/c = not clear
T_{pc} = Temperature at which partial collapse occurs in the freeze-dry microscope
T_c = Temperature at which total collapse occurs in the freeze-dry microscope
T_g = Glass transition temperature

[0060] With the exception of mannitol:lysine, all of the formulations appeared to have adequate physical appearance. Lysine interfered with the crystallization of both mannitol and glycine, which caused a depression in the glass transition temperature and a collapse of the lyophilized cake.

Example 6

[0061] The Factor VIII compositions described in Table 8 above were placed in storage at -70°C, 25°C, 40°C, and 50°C for varying lengths of time in order to evaluate their stability. Factor VIII activity levels were evaluated after 2 weeks, 1 month, 2 months, and 3 months, and the results are summarized in Table 10 below. Two of the samples, one employing mannitol as the bulking agent and sorbitol as the stabilizer, and the other employing mannitol as the bulking agent and glycerol as the stabilizer, exhibited poor stability. The remaining formulations all exhibited the ability to stabilize Factor VIII.

Table 10

Formulation Description	Temperature (°C)	% of initial at month				
		0	0.5	1	2	3
Glycine: Sucrose	-70	100.00	97.43	101.71	99.89	97.97
	25	100.00				85.44
	40	100.00		79.87	71.52	63.06
	50	100.00	76.34	67.99	52.14	47.64
Glycine:Trehalose	-70	100.00	89.22	96.00	95.90	94.64
	25	100.00				83.17
	40	100.00		79.93	72.42	68.03
	50	100.00	80.97	64.28	57.60	50.92
Mannitol:Trehalose	-70	100.00	91.32	97.72	96.10	98.26
	25	100.00				85.79
	40	100.00		82.54	70.72	59.44
	50	100.00	66.16	65.51	48.81	52.06
Mannitol: Sucrose	-70	100.00	100.45	100.56	105.47	99.22
	25	100.00				87.04
	40	100.00		85.59	80.78	55.42
	50	100.00	81.68	75.53	57.88	43.46
Mannitol:Arginine	-70	100.00	102.26	105.53	103.72	105.08
	25	100.00				95.15
	40	100.00		91.53	80.93	69.19

EP 2 921 180 B1

(continued)

Formulation Description	Temperature (°C)	% of initial at month				
			50	100.00	82.28	68.06
Mannitol:Raffinose	-70	100.00	93.88	98.41	100.68	103.62
	25	100.00				83.13
	40	100.00		81.09	73.61	67.16
	50	100.00	71.69	68.52	54.25	47.11
Mannitol: Glycerol	-70					
	25					
	40					
	50					
Mannitol: Sorbitol	-70	100.00	104.06			
	25	100.00				
	40	100.00				
	50	100.00	32.73			
HES:Sucrose	-70	100.00	102.74	103.03	100.90	
	25	100.00				
	40	100.00		76.89	77.47	
	50	100.00	71.47	67.40	30.02	
NaCl:Sucrose	-70	100.00	88.54	88.44	95.58	
	25	100.00				
	40	100.00		71.56	58.30	
	50	100.00	52.71	37.90	30.34	
Alanine:Sucrose	-70	100.00	109.78	109.67	108.96	
	25	100.00				
	40	100.00		92.99	73.03	
	50	100.00	83.25	74.91	57.65	
Glycine: Raffinose	-70	100.00	111.57	114.51	105.25	
	25	100.00				
	40	100.00		89.20	82.10	
	50	100.00	93.21	72.22	53.24	

Example 7

[0062] Based on the information developed during the studies described in Examples 5 and 6, it was decided that candidate formulations having the excipients shown in Table 11 below would be further developed.

Table 11

Excipient	Concentration
mannitol or glycine	6-9%
arginine or trehalose	1-3%
tween 80	0.005-0.04%

(continued)

Excipient	Concentration
NaCl	200-250 mM
CaCl ₂	3-5 mM
TRIS	20-30 mM
histidine or HEPES	10-50 mM
glutathione	0.15-0.25 mg/ml

[0063] Based on these parameters, the following specific formulations were developed:

Table 12

Formulation #1	Formulation #2	Formulation #3
10mM HEPES	10mM HEPES	10mM HEPES
20mM Tris	20mM Tris	20mM Tris
225mM NaCl	225mM NaCl	225mM NaCl
0.03% (v/v) Tween-80	0.03% (v/v) Tween-80	0.03% (v/v) Tween-80
8% (w/v) mannitol	8% (w/v) glycine	8% (w/v) mannitol
2% (w/v) trehalose	2% (w/v) trehalose	2% (w/v) arginine
0.2 mg/ml reduced glutathione	0.2 mg/ml reduced glutathione	0.2 mg/ml reduced glutathione
4 mM CaCl ₂	4 mM CaCl ₂	4 mM CaCl ₂

Formulation #4	Formulation #5
25mM histidine	25mM histidine
20mM Tris	20mM Tris
225mM NaCl	225mM NaCl
0.03% (v/v) Tween-80	0.03% (v/v) Tween-80
8% (w/v) mannitol	8% (w/v) glycine
2% (w/v) trehalose	2% (w/v) trehalose
0.2 mg/ml reduced glutathione	0.2 mg/ml reduced glutathione
4 mM CaCl ₂	4 mM CaCl ₂

[0064] The disclosure also provides:

(1) A Factor VIII composition formulated without adding albumin to said composition, comprising the following formulation excipients in addition to Factor VIII:

- 4% to 10% of a bulking agent selected from the group consisting of mannitol, glycine and alanine;
- 1% to 4% of a stabilizing agent selected from the group consisting of sucrose, trehalose, raffinose, and arginine;
- 1 mM to 5 mM calcium salt;
- 100 mM to 300 mM NaCl; and
- a buffering agent for maintaining a pH of approximately between 6 and 8.

(2) The Factor VIII composition of (1), additionally comprising a surfactant.

EP 2 921 180 B1

(3) The Factor VIII composition of (2), wherein said surfactant is selected from the group consisting of polysorbate 20, polysorbate 80, Pluronic F68, and Brij 35.

5 (4) The Factor VIII composition of (3), wherein said surfactant is polysorbate 80, and wherein said polysorbate 80 is present in an amount of less than 0.1%.

(5) The Factor VIII composition according to (1)-(4), wherein said surfactant is present in an amount of about 0.03%.

10 (6) The Factor VIII composition according to (1)-(5), wherein said buffering agent is selected from the group consisting of Tris, BIS-Tris Propane, histidine, PIPES, MOPS, HEPES, MES and ACES.

(7) The Factor VIII composition of (6), wherein said buffering agent comprises Tris.

15 (8) The Factor VIII composition of (7), wherein the Tris is present in an amount of about 20 mM.

(9) The Factor VIII composition of (6), wherein said buffering agent comprises between about 10 mM and about 50 mM histidine.

20 (10) The Factor VIII composition of (9), wherein the histidine is present in an amount of about 25 mM.

(11) The Factor VIII composition according to (1)-(10), further comprising an antioxidant.

(12) The Factor VIII composition of (11), wherein said antioxidant is glutathione.

25 (13) The Factor VIII composition of (12), wherein said glutathione is present in an amount of between about 0.05 mg/ml and about 1.0 mg/ml.

(14) The Factor VIII composition according to (1)-(13), wherein said bulking agent is present in an amount of about 8%.

30 (15) The Factor VIII composition according to (1)-(14), wherein said bulking agent is mannitol.

(16) The Factor VIII composition according to (1)-(14), wherein said bulking agent is glycine.

35 (17) The Factor VIII composition according to (1)-(16), wherein said stabilizing agent is present in an amount of about 2%.

(18) The Factor VIII composition according to (1)-(17), wherein said stabilizing agent is sucrose.

40 (19) The Factor VIII composition according to (1)-(17), wherein said stabilizing agent is arginine.

(20) The Factor VIII composition according to (1)-(17), wherein said stabilizing agent is trehalose.

45 (21) The Factor VIII composition according to (1)-(20), wherein said NaCl is present in an amount of from about 200 mM to about 250 mM.

(22) The Factor VIII composition of (21), wherein said NaCl is present in an amount of about 225 mM.

(23) The Factor VIII composition of (1)-(22), wherein said calcium salt is calcium chloride.

50 (24) The Factor VIII composition according to (1)-(23), wherein said composition is in lyophilized form.

(25) A Factor VIII composition formulated without adding albumin to said composition, comprising the following formulation excipients in addition to Factor VIII:

55 2% to 6% hydroxyethyl starch;
1% to 4% of a stabilizing agent selected from the group consisting of sucrose, trehalose, raffinose, and arginine;
1 mM to 5 mM calcium salt;
100 mM to 300 mM NaCl; and

a buffering agent for maintaining a pH of approximately between 6 and 8.

(26) The Factor VIII composition of (25), comprising about 4% hydroxyethyl starch.

5 (27) The Factor VIII composition according to (25)-(26), comprising about 200 mM NaCl.

(28) The Factor VIII composition according to (25)-(27), wherein said stabilizing agent is present in an amount of about 2%.

10 (29) The Factor VIII composition according to (25)-(28), wherein said stabilizing agent is sucrose.

(30) The Factor VIII composition according to (25)-(28), wherein said stabilizing agent is arginine.

(31) The Factor VIII composition according to (25)-(28), wherein said stabilizing agent is trehalose.

15 (32) The Factor VIII composition formulated without adding albumin to said composition, comprising the following formulation excipients in addition to Factor VIII:

20 300 mM to 500 mM NaCl;
1% to 4% of a stabilizing agent selected from the group consisting of sucrose, trehalose, raffinose, and arginine;
1 mM to 5 mM calcium salt; and
a buffering agent for maintaining a pH of approximately between 6 and 8.

(33) The composition of (32), wherein the NaCl is present in an amount of about 400 mM.

25 (34) Use of Factor VIII composition according to any of (1)-(33) for the preparation of a medicament for the treatment of hemophilia.

(35) An improved method of lyophilizing an aqueous pharmaceutical formulation containing a crystallizable bulking agent and NaCl, wherein said method comprises the steps of:

30 (a) freezing the aqueous pharmaceutical formulation at a temperature of less than about -35 °C;
(b) annealing the pharmaceutical formulation at between about -30 °C and - 19 °C;
(c) lowering the temperature of the pharmaceutical formulation to less than about -50 °C;
35 (d) annealing the pharmaceutical formulation at between about -30 °C and - 39 °C; and then
(e) freeze-drying the pharmaceutical formulation.

[0065] Although the invention has been described in connection with specific embodiments, it should be understood that the invention is not limited to such specific embodiments, but is as defined in the following claims.

40

Claims

1. A Factor VIII composition for use in treating a patient by reconstituting a lyophilised Factor VIII composition in a volume of a diluent to prepare a reconstituted Factor VIII composition and administering the reconstituted Factor VIII composition to the patient, wherein the lyophilised Factor VIII composition is obtainable by lyophilizing a volume of a liquid Factor VIII composition formulated without adding albumin to said composition, wherein the liquid Factor VIII composition comprises the following formulation excipients in addition to Factor VIII:

45 50 4% to 10% of a bulking agent selected from the group consisting of mannitol, glycine and alanine;
1% to 4% of a stabilizing agent selected from the group consisting of sucrose, trehalose, raffinose, and arginine;
1 mM to 5 mM calcium salt;
100 mM to 300 mM NaCl; and
55 a buffering agent for maintaining a pH of approximately between 6 and 8;
and wherein the volume of the diluent is equal to or greater than the volume of the liquid Factor VIII composition.

2. A method for preparing a reconstituted Factor VIII composition comprising the steps of:

(i) providing a lyophilised Factor VIII composition obtainable by lyophilizing a volume of a liquid Factor VIII composition formulated without adding albumin to said composition, wherein the liquid Factor VIII composition comprises the following formulation excipients in addition to Factor VIII:

- 5 4% to 10% of a bulking agent selected from the group consisting of mannitol, glycine and alanine;
 1% to 4% of a stabilizing agent selected from the group consisting of sucrose, trehalose, raffinose, and
 arginine;
 1 mM to 5 mM calcium salt;
 100 mM to 300 mM NaCl; and
 10 a buffering agent for maintaining a pH of approximately between 6 and 8; and

(ii) reconstituting the lyophilised Factor VIII composition in a volume of diluent wherein the volume of the diluent is equal to or greater than the volume of the liquid Factor VIII composition.

15 **3.** A reconstituted Factor VIII composition comprising:

(i) a lyophilised Factor VIII composition obtainable by lyophilizing a volume of a liquid Factor VIII composition formulated without adding albumin to said composition, wherein the liquid Factor VIII composition comprises the following formulation excipients in addition to Factor VIII:

- 20 4% to 10% of a bulking agent selected from the group consisting of mannitol, glycine and alanine;
 1% to 4% of a stabilizing agent selected from the group consisting of sucrose, trehalose, raffinose, and
 arginine;
 1 mM to 5 mM calcium salt;
 100 mM to 300 mM NaCl; and
 25 a buffering agent for maintaining a pH of approximately between 6 and 8; and

(ii) a volume of a diluent which is equal to or greater than the volume of the liquid Factor VIII composition.

30 **4.** The Factor VIII composition for use according to Claim 1 or the method for preparing a reconstituted Factor VIII composition of Claim 2 or the reconstituted Factor VIII composition of Claim 3 wherein the volume of the liquid Factor VIII composition is 2 mL and the volume of the diluent is 2 mL or 5 mL.

35 **5.** The Factor VIII composition for use according to Claim 1 or 4 or the method for preparing a reconstituted Factor VIII composition of Claim 2 or 4 or the reconstituted Factor VIII composition of Claim 3 or 4 wherein the diluent is water.

40 **6.** The Factor VIII composition for use according to Claim 1, 4 or 5 or the method for preparing a reconstituted Factor VIII composition of Claim 2, 4 or 5 or the reconstituted Factor VIII composition of any one of Claims 3 to 5 wherein the liquid Factor VIII composition additionally comprises a surfactant, optionally selected from the group consisting of polysorbate 20, polysorbate 80, Pluronic F68 and Brij 35, and optionally in an amount of about 0.03%.

45 **7.** The Factor VIII composition for use according to Claim 1, 4, 5 or 6 or the method for preparing a reconstituted Factor VIII composition of Claim 2, 4, 5 or 6 or the reconstituted Factor VIII composition of any one of Claims 3 to 6 wherein the buffering agent of the liquid Factor VIII composition comprises Tris, optionally in an amount of about 20 mM.

50 **8.** The Factor VIII composition for use according to Claim 1, 4, 5 or 6 or the method for preparing a reconstituted Factor VIII composition of Claim 2, 4, 5 or 6 or the reconstituted Factor VIII composition of any one of Claims 3 to 6 wherein the buffering agent of the liquid Factor VIII composition comprises histidine, optionally at a concentration of about 25 mM.

55 **9.** The Factor VIII composition for use according to Claim 1, 4, 5, 6, 7 or 8 or the method for preparing a reconstituted Factor VIII composition of Claim 2, 4, 5, 6, 7 or 8 or the reconstituted Factor VIII composition of any one of Claims 3 to 8 wherein the liquid Factor VIII composition optionally further comprises an antioxidant, optionally glutathione, optionally glutathione in an amount of between 0.05 mg/mL and 1.0 mg/mL.

10. The Factor VIII composition for use according to Claim 1, 4, 5, 6, 7, 8 or 9 or the method for preparing a reconstituted Factor VIII composition of Claim 2, 4, 5, 6, 7, 8 or 9 the reconstituted Factor VIII composition of any one of Claims 3 to 9 wherein the bulking agent of the liquid Factor VIII composition is mannitol, optionally in an amount of about 8%.

11. The Factor VIII composition for use according to Claim 1, 4, 5, 6, 7, 8, 9 or 10 or the method for preparing a reconstituted Factor VIII composition of Claim 2, 4, 5, 6, 7, 8, 9 or 10 or the reconstituted Factor VIII composition of any one of Claims 3 to 10 wherein the stabilizing agent of the liquid Factor VIII composition is trehalose, optionally in an amount of about 2%.

5
12. The Factor VIII composition for use according to Claim 1, 4, 5, 6, 7, 8, 9, 10 or 11 or the method for preparing a reconstituted Factor VIII composition of Claim 2, 4, 5, 6, 7, 8, 9, 10 or 11 or the reconstituted Factor VIII composition of any one of Claims 3 to 11 wherein the NaCl of the liquid Factor VIII composition is present in an amount of from 200 to 250 mM, such as in an amount of about 225 mM.

10
13. The Factor VIII composition for use according to Claim 1, 4, 5, 6, 7, 8, 9, 10, 11 or 12 or the method for preparing a reconstituted Factor VIII composition of Claim 2, 4, 5, 6, 7, 8, 9, 10, 11 or 12 or the reconstituted Factor VIII composition of any one of Claims 3 to 12 wherein the calcium salt of the liquid Factor VIII composition is calcium chloride.

15

Patentansprüche

20
1. Faktor VIII-Zusammensetzung zur Verwendung bei der Behandlung eines Patienten durch das Rekonstituieren einer lyophilisierten Faktor VIII-Zusammensetzung in einem Volumen eines Verdünnungsmittels, um eine rekonstituierte Faktor VIII-Zusammensetzung herzustellen, und das Verabreichen der rekonstituierten Faktor VIII-Zusammensetzung an den Patienten,
wobei die lyophilisierte Faktor VIII-Zusammensetzung durch das Lyophilisieren eines Volumens einer flüssigen Faktor VIII-Zusammensetzung erhältlich ist, die ohne Zugabe von Albumin zu der Zusammensetzung formuliert ist,
25 wobei die flüssige Faktor VIII-Zusammensetzung die folgenden Formulierungshilfsstoffe zusätzlich zu Faktor VIII umfasst:

30
4% bis 10% eines Füllmittels, das ausgewählt ist aus der Gruppe, bestehend aus Mannitol, Glycin und Alanin;
1% bis 4% eines Stabilisierungsmittels, das ausgewählt ist aus der Gruppe, bestehend aus Saccharose, Trehalose, Raffinose und Arginin;
1 mM bis 5 mM Calciumsalz;
100 mM bis 300 mM NaCl; und
ein Puffermittel zum Beibehalten eines pH-Werts von ungefähr zwischen 6 und 8;
35 und wobei das Volumen des Verdünnungsmittels gleich zum oder größer als das Volumen der flüssigen Faktor VIII-Zusammensetzung ist.

2. Verfahren zur Herstellung einer rekonstituierten Faktor VIII-Zusammensetzung, umfassend die Schritte:

40
(i) des Bereitstellens einer lyophilisierten Faktor VIII-Zusammensetzung, die durch das Lyophilisieren eines Volumens einer flüssigen Faktor VIII-Zusammensetzung erhältlich ist, die ohne Zugabe von Albumin zu der Zusammensetzung formuliert ist, wobei die flüssige Faktor VIII-Zusammensetzung die folgenden Formulierungshilfsstoffe zusätzlich zu Faktor VIII umfasst:

45
4% bis 10% eines Füllmittels, das ausgewählt ist aus der Gruppe, bestehend aus Mannitol, Glycin und Alanin;
1% bis 4% eines Stabilisierungsmittels, das ausgewählt ist aus der Gruppe, bestehend aus Saccharose, Trehalose, Raffinose und Arginin;
1 mM bis 5 mM Calciumsalz;
100 mM bis 300 mM NaCl; und
ein Puffermittel zum Beibehalten eines pH-Werts von ungefähr zwischen 6 und 8; und

50
(ii) des Rekonstituierens der lyophilisierten Faktor VIII-Zusammensetzung in einem Volumen eines Verdünnungsmittels, wobei das Volumen des Verdünnungsmittels gleich zum oder größer als das Volumen der flüssigen Faktor VIII-Zusammensetzung ist.

55
3. Rekonstituierte Faktor VIII-Zusammensetzung, umfassend:

(i) eine lyophilisierte Faktor VIII-Zusammensetzung, die durch das Lyophilisieren eines Volumens einer flüssigen Faktor VIII-Zusammensetzung erhältlich ist, die ohne Zugabe von Albumin zu der Zusammensetzung formuliert

EP 2 921 180 B1

ist, wobei die flüssige Faktor VIII-Zusammensetzung die folgenden Formulierungshilfsstoffe zusätzlich zu Faktor VIII umfasst:

5 4% bis 10% eines Füllmittels, das ausgewählt ist aus der Gruppe, bestehend aus Mannitol, Glycin und Alanin;
1% bis 4% eines Stabilisierungsmittels, das ausgewählt ist aus der Gruppe, bestehend aus Saccharose,
Trehalose, Raffinose und Arginin;
1 mM bis 5 mM Calciumsalz;
100 mM bis 300 mM NaCl; und
10 ein Puffermittel zum Beibehalten eines pH-Werts von ungefähr zwischen 6 und 8; und

(ii) ein Volumen eines Verdünnungsmittels, das gleich zum oder größer als das Volumen der flüssigen Faktor VIII-Zusammensetzung ist.

- 15 **4.** Faktor VIII-Zusammensetzung zur Verwendung gemäß Anspruch 1, oder Verfahren zur Herstellung einer rekonstituierten Faktor VIII-Zusammensetzung gemäß Anspruch 2, oder rekonstituierte Faktor VIII-Zusammensetzung gemäß Anspruch 3, wobei das Volumen der flüssigen Faktor VIII-Zusammensetzung 2 mL beträgt und das Volumen des Verdünnungsmittels 2 mL oder 5 mL beträgt.
- 20 **5.** Faktor VIII-Zusammensetzung zur Verwendung gemäß Anspruch 1 oder 4, oder Verfahren zur Herstellung einer rekonstituierten Faktor VIII-Zusammensetzung gemäß Anspruch 2 oder 4, oder rekonstituierte Faktor VIII-Zusammensetzung gemäß Anspruch 3 oder 4, wobei das Verdünnungsmittel Wasser ist.
- 25 **6.** Faktor VIII-Zusammensetzung zur Verwendung gemäß Anspruch 1, 4 oder 5, oder Verfahren zur Herstellung einer rekonstituierten Faktor VIII-Zusammensetzung gemäß Anspruch 2, 4 oder 5, oder rekonstituierte Faktor VIII-Zusammensetzung gemäß irgendeinem der Ansprüche 3 bis 5, wobei die flüssige Faktor VIII-Zusammensetzung zusätzlich ein Tensid umfasst, das optional ausgewählt ist aus der Gruppe, bestehend aus Polysorbat 20, Polysorbat 80, Pluronic F68 und Brij 35, und optional in einer Menge von etwa 0,03%.
- 30 **7.** Faktor VIII-Zusammensetzung zur Verwendung gemäß Anspruch 1, 4, 5 oder 6, oder Verfahren zur Herstellung einer rekonstituierten Faktor VIII-Zusammensetzung gemäß Anspruch 2, 4, 5 oder 6, oder rekonstituierte Faktor VIII-Zusammensetzung gemäß irgendeinem der Ansprüche 3 bis 6, wobei das Puffermittel der flüssigen Faktor VIII-Zusammensetzung Tris umfasst, optional in einer Menge von etwa 20 mM.
- 35 **8.** Faktor VIII-Zusammensetzung zur Verwendung gemäß Anspruch 1, 4, 5 oder 6, oder Verfahren zur Herstellung einer rekonstituierten Faktor VIII-Zusammensetzung gemäß Anspruch 2, 4, 5 oder 6, oder rekonstituierte Faktor VIII-Zusammensetzung gemäß irgendeinem der Ansprüche 3 bis 6, wobei das Puffermittel der flüssigen Faktor VIII-Zusammensetzung Histidin umfasst, optional in einer Konzentration von etwa 25 mM.
- 40 **9.** Faktor VIII-Zusammensetzung zur Verwendung gemäß Anspruch 1, 4, 5, 6, 7 oder 8, oder Verfahren zur Herstellung einer rekonstituierten Faktor VIII-Zusammensetzung gemäß Anspruch 2, 4, 5, 6, 7 oder 8, oder rekonstituierte Faktor VIII-Zusammensetzung gemäß irgendeinem der Ansprüche 3 bis 8, wobei die flüssige Faktor VIII-Zusammensetzung optional weiter ein Antioxidans umfasst, optional Glutathion, optional Glutathion in einer Menge von zwischen 0,05 mg/mL und 1,0 mg/mL.
- 45 **10.** Faktor VIII-Zusammensetzung zur Verwendung gemäß Anspruch 1, 4, 5, 6, 7, 8 oder 9, oder Verfahren zur Herstellung einer rekonstituierten Faktor VIII-Zusammensetzung gemäß Anspruch 2, 4, 5, 6, 7, 8 oder 9, oder rekonstituierte Faktor VIII-Zusammensetzung gemäß irgendeinem der Ansprüche 3 bis 9, wobei das Füllmittel der flüssigen Faktor VIII-Zusammensetzung Mannitol ist, optional in einer Menge von etwa 8%.
- 50 **11.** Faktor VIII-Zusammensetzung zur Verwendung gemäß Anspruch 1, 4, 5, 6, 7, 8, 9 oder 10, oder Verfahren zur Herstellung einer rekonstituierten Faktor VIII-Zusammensetzung gemäß Anspruch 2, 4, 5, 6, 7, 8, 9 oder 10, oder rekonstituierte Faktor VIII-Zusammensetzung gemäß irgendeinem der Ansprüche 3 bis 10, wobei das Stabilisierungsmittel der flüssigen Faktor VIII-Zusammensetzung Trehalose ist, optional in einer Menge von etwa 2%.
- 55 **12.** Faktor VIII-Zusammensetzung zur Verwendung gemäß Anspruch 1, 4, 5, 6, 7, 8, 9, 10 oder 11, oder Verfahren zur Herstellung einer rekonstituierten Faktor VIII-Zusammensetzung gemäß Anspruch 2, 4, 5, 6, 7, 8, 9, 10 oder 11, oder rekonstituierte Faktor VIII-Zusammensetzung gemäß irgendeinem der Ansprüche 3 bis 11, wobei das NaCl der flüssigen Faktor VIII-Zusammensetzung in einer Menge von 200 bis 250 mM vorliegt, beispielsweise in einer

Menge von etwa 225 mM.

- 5 13. Faktor VIII-Zusammensetzung zur Verwendung gemäß Anspruch 1, 4, 5, 6, 7, 8, 9, 10, 11 oder 12, oder Verfahren zur Herstellung einer rekonstituierten Faktor VIII-Zusammensetzung gemäß Anspruch 2, 4, 5, 6, 7, 8, 9, 10, 11 oder 12, oder rekonstituierte Faktor VIII-Zusammensetzung gemäß irgendeinem der Ansprüche 3 bis 12, wobei das Calciumsalz der flüssigen Faktor VIII-Zusammensetzung Calciumchlorid ist.

10 Revendications

- 15 1. Composition de facteur VIII pour utilisation dans le traitement d'un patient en reconstituant une composition de facteur VIII lyophilisée dans un volume d'un diluant pour préparer une composition de facteur VIII reconstituée et en administrant la composition de facteur VIII reconstituée au patient, dans laquelle la composition de facteur VIII lyophilisée peut être obtenue en lyophilisant un volume d'une composition de facteur VIII liquide formulée sans ajouter d'albumine à ladite composition, dans laquelle la composition de facteur VIII liquide comprend les excipients de formulation suivants en plus du facteur VIII :

20 4 % à 10 % d'un agent gonflant choisi dans le groupe constitué du mannitol, de la glycine et de l'alanine ;
1 % à 4 % d'un agent stabilisant choisi dans le groupe constitué du saccharose, du tréhalose, de la raffinose et de l'arginine ;
1 mM à 5 mM de sel de calcium ;
100 mM à 300 mM de NaCl ; et
un agent tampon pour maintenir un pH d'environ 6 à 8 ; et
dans laquelle le volume du diluant est égal ou supérieur au volume de la composition de facteur VIII liquide.

- 25 2. Procédé de préparation d'une composition de facteur VIII reconstituée comprenant les étapes consistant à :

30 (i) fournir une composition de facteur VIII lyophilisée qui peut être obtenue en lyophilisant un volume d'une composition de facteur VIII liquide formulée sans ajouter d'albumine à ladite composition, dans laquelle la composition de facteur VIII liquide comprend les excipients de formulation suivants en plus du facteur VIII :

35 4 % à 10 % d'un agent gonflant choisi dans le groupe constitué du mannitol, de la glycine et de l'alanine ;
1 % à 4 % d'un agent stabilisant choisi dans le groupe constitué du saccharose, du tréhalose, de la raffinose et de l'arginine ;
1 mM à 5 mM de sel de calcium ;
100 mM à 300 mM de NaCl ; et
un agent tampon pour maintenir un pH d'environ 6 à 8 ; et

40 (ii) reconstituer la composition de facteur VIII lyophilisée dans un volume de diluant, dans laquelle le volume du diluant est égal ou supérieur au volume de la composition de facteur VIII liquide.

3. Composition de facteur VIII reconstituée comprenant :

45 (i) une composition de facteur VIII lyophilisée qui peut être obtenue en lyophilisant un volume d'une composition de facteur VIII liquide formulée sans ajouter d'albumine à ladite composition, dans laquelle la composition de facteur VIII liquide comprend les excipients de formulation suivants en plus du facteur VIII :

50 4 % à 10 % d'un agent gonflant choisi dans le groupe constitué du mannitol, de la glycine et de l'alanine ;
1 % à 4 % d'un agent stabilisant choisi dans le groupe constitué du saccharose, du tréhalose, de la raffinose et de l'arginine ;
1 mM à 5 mM de sel de calcium ;
100 mM à 300 mM de NaCl ; et
un agent tampon pour maintenir un pH d'environ 6 à 8 ; et

55 (ii) un volume d'un diluant qui est égal ou supérieur au volume de la composition de facteur VIII liquide.

4. Composition de facteur VIII pour utilisation selon la revendication 1 ou procédé de préparation d'une composition de facteur VIII reconstituée selon la revendication 2 ou composition de facteur VIII reconstituée selon la revendication

EP 2 921 180 B1

3, dans lequel ou laquelle le volume de la composition de facteur VIII liquide est de 2 mL et le volume du diluant est de 2 mL ou de 5 mL.

- 5
5. Composition de facteur VIII pour utilisation selon la revendication 1 ou 4 ou procédé de préparation d'une composition de facteur VIII reconstituée selon la revendication 2 ou 4 ou composition de facteur VIII reconstituée selon la revendication 3 ou 4, dans lequel ou laquelle le diluant est de l'eau.
- 10
6. Composition de facteur VIII pour utilisation selon la revendication 1, 4 ou 5 ou procédé de préparation d'une composition de facteur VIII reconstituée selon la revendication 2, 4 ou 5 ou composition de facteur VIII reconstituée selon l'une quelconque des revendications 3 à 5, dans lequel ou laquelle la composition de facteur VIII liquide comprend en outre un tensioactif éventuellement choisi dans le groupe constitué du polysorbate 20, du polysorbate 80, du Pluronic F68 et du Brij 35 et éventuellement en quantité d'environ 0,03 %.
- 15
7. Composition de facteur VIII pour utilisation selon la revendication 1, 4, 5 ou 6 ou procédé de préparation d'une composition de facteur VIII reconstituée selon la revendication 2, 4, 5 ou 6 ou composition de facteur VIII reconstituée selon l'une quelconque des revendications 3 à 6, dans lequel ou laquelle l'agent tampon de la composition de facteur VIII liquide comprend du Tris, éventuellement en quantité d'environ 20 mM.
- 20
8. Composition de facteur VIII pour utilisation selon la revendication 1, 4, 5 ou 6 ou procédé de préparation d'une composition de facteur VIII reconstituée selon la revendication 2, 4, 5 ou 6 ou composition de facteur VIII reconstituée selon l'une quelconque des revendications 3 à 6, dans lequel ou laquelle l'agent tampon de la composition de facteur VIII liquide comprend de l'histidine, éventuellement en concentration d'environ 25 mM.
- 25
9. Composition de facteur VIII pour utilisation selon la revendication 1, 4, 5, 6, 7 ou 8 ou procédé de préparation d'une composition de facteur VIII reconstituée selon la revendication 2, 4, 5, 6, 7 ou 8 ou composition de facteur VIII reconstituée selon l'une quelconque des revendications 3 à 8, dans lequel ou laquelle la composition de facteur VIII liquide comprend en outre éventuellement un antioxydant, éventuellement de la glutathione, éventuellement de la glutathione en quantité comprise entre 0,05 mg/mL et 1,0 mg/mL.
- 30
10. Composition de facteur VIII pour utilisation selon la revendication 1, 4, 5, 6, 7, 8 ou 9 ou procédé de préparation d'une composition de facteur VIII reconstituée selon la revendication 2, 4, 5, 6, 7, 8 ou 9 ou composition de facteur VIII reconstituée selon l'une quelconque des revendications 3 à 9, dans lequel ou laquelle l'agent gonflant de la composition de facteur VIII liquide est le mannitol, éventuellement en quantité d'environ 8%.
- 35
11. Composition de facteur VIII pour utilisation selon la revendication 1, 4, 5, 6, 7, 8, 9 ou 10 ou procédé de préparation d'une composition de facteur VIII reconstituée selon la revendication 2, 4, 5, 6, 7, 8, 9 ou 10 ou composition de facteur VIII reconstituée selon l'une quelconque des revendications 3 à 10, dans lequel ou laquelle l'agent stabilisant de la composition de facteur VIII liquide est la tréhalose, éventuellement en quantité d'environ 2 %.
- 40
12. Composition de facteur VIII pour utilisation selon la revendication 1, 4, 5, 6, 7, 8, 9, 10 ou 11 ou procédé de préparation d'une composition de facteur VIII reconstituée selon la revendication 2, 4, 5, 6, 7, 8, 9, 10 ou 11 ou composition de facteur VIII reconstituée selon l'une quelconque des revendications 3 à 11, dans lequel ou laquelle le NaCl de la composition de facteur VIII liquide est présent en quantité de 200 à 250 mM, notamment en quantité d'environ 225 mM.
- 45
13. Composition de facteur VIII pour utilisation selon la revendication 1, 4, 5, 6, 7, 8, 9, 10, 11 ou 12 ou procédé de préparation d'une composition de facteur VIII reconstituée selon la revendication 2, 4, 5, 6, 7, 8, 9, 10, 11 ou 12 ou composition de facteur VIII reconstituée selon l'une quelconque des revendications 3 à 12, dans lequel ou laquelle le sel de calcium de la composition de facteur VIII liquide est le chlorure de calcium.
- 50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 5565427 A [0004]
- EP 508194 A, Freudenberg [0004]
- US 5763401 A [0005]
- EP 818204 A, Nayer [0005]
- US 5733873 A [0006]
- EP 627924 A, Osterberg [0006]
- US 4877608 A [0007]
- EP 315968 A, Lee [0007]
- US 5605884 A [0008]
- EP 0314095 A, Lee [0008]
- WO 9622107 A, Roser [0009]
- WO 9630041 A1 [0010]
- US 5328694 A [0011]
- EP 511234 A, Schwinn [0011]
- US 4757006 A [0036]
- US 4868112 A, Toole, Jr. [0036]
- WO A9109122 A [0036]
- EP 0362218 A [0036]

Non-patent literature cited in the description

- **ANDERSSON et al.** *Proc. Natl. Acad. Sci. USA*, May 1986, vol. 83, 2979-2983 [0020]
- An Effect of Predilution on Potency Assays of Factor VIII Concentrates. **LEE, MARTIN L et al.** *Thrombosis Research*. Pergamon Press Ltd, 1983, vol. 30, 511-519 [0021]
- Lyophilization of Protein Pharmaceuticals. **CARPENTER, J.F. ; CHANG, B.S.** *Biotechnology and Biopharmaceutical Manufacturing, Processing and Preservation*. Interpharm Press, Inc, 1996, 199-264 [0040]