



(11)

EP 3 272 748 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
19.06.2019 Bulletin 2019/25

(51) Int Cl.:
C07D 453/02 ^(2006.01) **A61K 31/439** ^(2006.01)
A61P 25/00 ^(2006.01)

(21) Application number: **17186513.2**

(22) Date of filing: **17.05.2011**

(54) **PHARMACEUTICAL FORMULATIONS COMPRISING CRYSTALLINE FORMS OF
(R)-7-CHLORO-N-(QUINUCLIDIN-3-YL)BENZO(B)THIOPHENE-2-CARBOXAMIDE
HYDROCHLORIDE MONOHYDRATE**

PHARMAZEUTISCHE FORMULIERUNGEN MIT KRISTALLINE FORMEN VOM
(R)-7-CHLORO-N-(CHINUCLIDIN-3-YL)BENZO(B)THIOPHEN-2-CARBOXAMID-HYDROCHLORID-
MONOHYDRAT

FORMULATIONS PHARMACEUTIQUES COMPRENANT DES FORMES CRISTALLINES DE
MONOHYDRATE D'HYDROCHLORURE DE
(R)-7-CHLORO-N-(QUINUCLIDIN-3-YL)BENZO(B)THIOPHENE-2-CARBOXAMIDE

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **17.05.2010 US 345363 P**
07.06.2010 US 352092 P

(43) Date of publication of application:
24.01.2018 Bulletin 2018/04

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
15202184.6 / 3 029 039
11721413.0 / 2 571 874

(73) Proprietor: **Forum Pharmaceuticals Inc.**
Waltham, MA 02451 (US)

(72) Inventors:
• **OLIVER-SHAFFER, Patricia**
Acton, MA Massachusetts 01720 (US)
• **SHAPIRO, Gideon**
Gainesville, FL Florida 32653 (US)
• **CHESWORTH, Richard**
Concord, MA Massachusetts 01742 (US)
• **KISHIDA, Muneki**
Osaka 541-8505 (JP)
• **ISHIGE, Takayuki**
Osaka 541-8505 (JP)

(74) Representative: **Holzwarth-Rochford, Andreas**
Jones Day
Nextower
Thurn-und-Taxis-Platz 6
60313 Frankfurt am Main (DE)

(56) References cited:
US-A1- 2005 119 325

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 3 272 748 B1

Description**FIELD**

[0001] The present disclosure relates to a pharmaceutical formulation comprising crystalline forms of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate for use in methods of therapeutic treatments.

BACKGROUND OF THE INVENTION

[0002] The endogenous neurotransmitter acetylcholine (ACh) mediates diverse physiological functions in the peripheral and central nervous systems (CNS) via muscarinic and nicotinic subclasses of acetylcholine receptors (AChRs). The nicotinic acetylcholine receptors (nAChRs) are ligand-gated cell surface ion channels that are selectively activated by the natural product nicotine. The diverse molecular subtypes or variants of nicotinic acetylcholine receptor are based on the pentameric structure of the receptor. The nAChR subtypes are formed from diverse pentameric combinations of nine molecularly distinct alpha subunits and four molecularly distinct beta subunits. A particularly interesting molecular target for therapeutic intervention is the alpha-7 nicotinic receptor subtype, which is comprised of five alpha-7 monomeric subunits. Thus, agonists which are selective for the alpha-7 receptor have potential to treat a range of diseases. Alpha-7 agonists are expected to be especially useful for the treatment of CNS disorders associated with cognitive deficits. This expectation is based on beneficial effects of alpha-7 receptor activation on cognition, learning and memory. At the same time, selective alpha-7 agonists are expected to cause fewer or less severe undesirable side effects, e.g. nausea, vomiting, tachycardia, which are mediated by the activation of certain other nicotinic receptor subtypes as for example by the non-selective agonist nicotine.

[0003] As such, there is a need for additional selective alpha-7 agonists for the treatment of CNS disorders associated with cognitive deficits.

SUMMARY OF THE INVENTION

[0004] Accordingly, the present invention is directed to a pharmaceutical formulation for use in a method of treatment for improving cognition or treating cognitive loss in a subject in need thereof, wherein the pharmaceutical formulation comprises a therapeutically effective amount of a crystalline Form I of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate, wherein the pharmaceutical formulation is a tablet or capsule; wherein the crystalline Form I of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate is characterized by an x-ray powder diffraction pattern having peaks expressed as 2θ at one or both of 17.48 and 20.58 ± 0.20 degrees when measured against an internal silicon standard and characterized by an x-ray powder diffraction pattern further having at least four peaks, six peaks, eight peaks or all peaks expressed as 2θ at 4.50, 9.04, 14.60, 15.14, 15.80, 16.60, 18.16, 18.44, 19.48, 21.74, and 25.46 ± 0.20 degrees when measured against an internal silicon standard; and wherein the subject is suffering from a disorder selected from the group consisting of: attention deficit disorder, attention deficit hyperactivity disorder, and Parkinson's Disease.

[0005] In a further aspect, the present invention is directed to a pharmaceutical formulation for use in a method of treatment for improving cognition or treating cognitive loss in a subject in need thereof, wherein the pharmaceutical formulation comprises a therapeutically effective amount of a crystalline Form II of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate, wherein the pharmaceutical formulation is a tablet or capsule; wherein the crystalline Form II of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate is characterized by an x-ray powder diffraction pattern having peaks expressed as 2θ at one or both of 21.16 and 21.38 ± 0.20 degrees when measured against an internal silicon standard and characterized by an x-ray powder diffraction pattern further having at least four peaks, six peaks, eight peaks or all peaks expressed as 2θ at 4.48, 9.00, 13.58, 15.62, 16.48, 19.02, 19.44, 22.46 and 25.00 ± 0.20 degrees when measured against an internal silicon standard; and wherein the subject is suffering from a disorder selected from the group consisting of: attention deficit disorder, attention deficit hyperactivity disorder, and Parkinson's Disease.

[0006] In another aspect the present invention is directed to a pharmaceutical formulation for oral administration, comprising a therapeutically effective amount of a crystalline Form I of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate and a pharmaceutically acceptable carrier, wherein the pharmaceutical formulation is a tablet or capsule; wherein the crystalline Form I of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate is characterized by an x-ray powder diffraction pattern having peaks expressed as 2θ at one or both of 17.48 and 20.58 ± 0.20 degrees when measured against an internal silicon standard and characterized by an x-ray powder diffraction pattern further having at least four peaks, six peaks, eight peaks or all peaks expressed as 2θ at 4.50, 9.04, 14.60, 15.14, 15.80, 16.60, 18.16, 18.44, 19.48, 21.74, and 25.46 ± 0.20 degrees when

measured against an internal silicon standard.

[0007] In a still further aspect the present invention is directed to a pharmaceutical formulation for oral administration comprising a therapeutically effective amount of a crystalline Form II of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate and a pharmaceutically acceptable carrier, wherein the pharmaceutical formulation is a tablet or capsule; wherein the crystalline Form II of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate is characterized by an x-ray powder diffraction pattern having peaks expressed as 2θ at one or both of 21.16 and 21.38 ± 0.20 degrees when measured against an internal silicon standard and characterized by an x-ray powder diffraction pattern further having at least four peaks, six peaks, eight peaks or all peaks expressed as 2θ at 4.48 , 9.00 , 13.58 , 15.62 , 16.48 , 19.02 , 19.44 , 22.46 and 25.00 ± 0.20 degrees when measured against an internal silicon standard.

BRIEF DESCRIPTION OF THE DRAWINGS

[0008]

Figure 1 is a phase diagram for Form I and Form II.

Figure 2 depicts an X-ray powder diffraction (XRPD) spectrum for Form X.

Figure 3 is a graph depicting the relationship between water activity and volume fraction of water in acetonitrile/water systems at various temperatures.

Figure 4 depicts a diagram of Form I and Form II using plots of temperature against the value of water activity.

Figure 5 depicts an X-ray powder diffraction (XRPD) spectrum for Form I.

Figure 6 depicts an X-ray powder diffraction (XRPD) spectrum for Form II.

DETAILED DESCRIPTION OF THE INVENTION

[0009] The present invention will be described with reference to the following definitions that, for convenience, are set forth below. Unless otherwise specified, the below terms used herein are defined as follows:

I. Definitions

[0010] As used herein and unless otherwise specified, the term "crystal forms," "crystalline forms" and related terms herein refer to solid forms that are crystalline. Crystal forms include single-component crystal forms and multiple-component crystal forms, and include, but are not limited to, polymorphs, solvates, hydrates, and/or other molecular complexes. In one embodiment, the crystalline forms of the invention are monohydrates. In certain embodiments, a crystalline form is substantially pure, isolated or enriched in one crystalline form, and/or is substantially free of amorphous forms and/or other crystal forms.

[0011] As used herein and unless otherwise specified, the term "crystalline" and related terms used herein, when used to describe a compound, substance, modification, material, component or product, unless otherwise specified, mean that the compound, substance, modification, material, component or product is substantially crystalline as determined by X-ray diffraction. See, e.g., Remington: The Science and Practice of Pharmacy, 21st edition, Lippincott, Williams and Wilkins, Baltimore, Md. (2005); The United States Pharmacopeia, 23rd ed., 1843-1844 (1995).

[0012] Moreover, more detailed characterizations techniques for characterizing crystal forms and amorphous forms may include, but are not limited to, thermal gravimetric analysis (TGA), differential scanning calorimetry (DSC), X-ray powder diffractometry (XRPD), single-crystal X-ray diffractometry, vibrational spectroscopy, e.g., infrared (IR) and Raman spectroscopy, solid-state and solution nuclear magnetic resonance (NMR) spectroscopy, optical microscopy, hot stage optical microscopy, scanning electron microscopy (SEM), electron crystallography and quantitative analysis, particle size analysis (PSA), surface area analysis, solubility measurements, dissolution measurements, elemental analysis and Karl Fischer analysis. Characteristic unit cell parameters may be determined using one or more techniques such as, but not limited to, X-ray diffraction and neutron diffraction, including single-crystal diffraction and powder diffraction. Techniques useful for analyzing powder diffraction data include profile refinement, such as Rietveld refinement, which may be used, e.g., to analyze diffraction peaks associated with a single phase in a sample comprising more than one solid phase. Other methods useful for analyzing powder diffraction data include unit cell indexing, which allows one of skill in the art to determine unit cell parameters from a sample comprising crystalline powder. Furthermore, it would be understood by the ordinarily skilled artisan that identification of a crystal may be made using one of these techniques, e.g., X-ray powder diffractometry, and may be confirmed using additional noted characterization techniques.

[0013] As used herein and unless otherwise specified, a sample comprising a particular crystal form or amorphous form that is "substantially pure," contains the particular crystal form or amorphous form in a chemical and/or physical purity greater than about 75%, e.g., 80%, e.g., 85%, e.g., 90%, e.g., 91%, e.g., 92%, e.g., 93%, e.g., 94%, e.g., 95%,

e.g., 96%, e.g., 97%, e.g., 98%, e.g., 99%, e.g., 99.25%, e.g., 99.50%, e.g., 99.75%, e.g., 99.9%, e.g., 100% physically and/or chemically pure. In certain embodiments, the particular crystal form or amorphous form is greater than about 90%, e.g., 91%, e.g., 92%, e.g., 93%, e.g., 94%, e.g., 95%, e.g., 96%, e.g., 97%, e.g., 98%, e.g., 99%, e.g., 99.25%, e.g., 99.50%, e.g., 99.75%, e.g., 99.9%, e.g., 100% physically and/or chemically pure. In particular embodiments, the particular crystal form or amorphous form is greater than about 95%, e.g., 96%, e.g., 97%, e.g., 98%, e.g., 99%, e.g., 99.25%, e.g., 99.50%, e.g., 99.75%, e.g., 99.9%, e.g., 100% physically and/or chemically pure. In specific embodiments, the particular crystal form or amorphous form is greater than about 99%, e.g., 99.25%, e.g., 99.50%, e.g., 99.75%, e.g., 99.9%, e.g., 100% physically and/or chemically pure.

[0014] As used herein and unless otherwise specified, a sample or composition that is "substantially free" of one or more other solid forms and/or other chemical compounds means that the composition contains, in particular embodiments, less than about 25%, e.g., 20%, e.g., 15%, e.g., 10%, e.g., 9%, e.g., 8%, e.g., 7%, e.g., 6%, e.g., 5%, e.g., 4%, e.g., 3%, e.g., 2%, e.g., 1%, e.g., 0.75%, e.g., 0.5%, e.g., 0.25%, e.g., or 0.1% percent by weight of one or more amorphous forms and/or other crystal forms. In certain embodiments, the composition contains less than 10%, e.g., 9%, e.g., 8%, e.g., 7%, e.g., 6%, e.g., 5%, e.g., 4%, e.g., 3%, e.g., 2%, e.g., 1%, e.g., 0.75%, e.g., 0.5%, e.g., 0.25%, e.g., or 0.1% percent by weight of one or more amorphous forms and/or other crystal forms. In particular embodiments, the composition contains less than 5%, e.g., 4%, e.g., 3%, e.g., 2%, e.g., 1%, e.g., 0.75%, e.g., 0.5%, e.g., 0.25%, e.g., or 0.1% percent by weight of one or more amorphous forms and/or other crystal forms. In specific embodiments, the composition contains less than 1%, e.g., 0.75%, e.g., 0.5%, e.g., 0.25%, e.g., or 0.1% percent by weight of one or more amorphous forms and/or other crystal forms. In certain embodiments, a crystal form of a substance may be physically and/or chemically pure.

[0015] As used herein and unless otherwise specified, the terms "polymorphs," "polymorphic forms" and related terms herein, refer to two or more crystal forms that consist essentially of the same molecule, molecules, and/or ions. Different polymorphs may have different physical properties such as, e.g., melting temperature, heat of fusion, solubility, dissolution properties and/or vibrational spectra, as a result of the arrangement or conformation of the molecules and/or ions in the crystal lattice. The differences in physical properties may affect pharmaceutical parameters such as storage stability, compressibility and density (important in formulation and product manufacturing), and dissolution rate (an important factor in bioavailability). Differences in stability can result from changes in chemical reactivity (e.g., differential oxidation, such that a dosage form discolors more rapidly when comprised of one polymorph than when comprised of another polymorph) or mechanical changes (e.g., tablets crumble on storage as a kinetically favored polymorph converts to thermodynamically more stable polymorph) or both (e.g., tablets of one polymorph are more susceptible to breakdown at high humidity). As a result of solubility/dissolution differences, in the extreme case, some solid-state transitions may result in lack of potency or, at the other extreme, toxicity. In addition, the physical properties may be important in processing (e.g., one polymorph might be more likely to form solvates or might be difficult to filter and wash free of impurities, and particle shape and size distribution might be different between polymorphs).

[0016] The terms "hydrate" and "hydrated" refer to a solvate wherein the solvent comprises water. "Polymorphs of solvates" refers to the existence of more than one crystal form for a particular solvate composition. Similarly, "polymorphs of hydrates" refers to the existence of more than one crystal form for a particular hydrate composition.

[0017] As used herein and unless otherwise specified, the term "amorphous," "amorphous form," and related terms used herein, describe that the substance, component or product in question is not substantially crystalline as determined by X-ray diffraction. In particular, the term "amorphous form" describes a disordered solid form, i.e., a solid form lacking long range crystalline order. In certain embodiments, an amorphous form of a substance may be substantially free of other amorphous forms and/or crystal forms. In other embodiments, an amorphous form of a substance may contain less than about 1%, 2%, 3%, 4%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45% or 50% of one or more other amorphous forms and/or crystal forms on a weight basis. In certain embodiments, an amorphous form of a substance may be physically and/or chemically pure. In certain embodiments, an amorphous form of a substance be about 99%, 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91% or 90% physically and/or chemically pure.

[0018] As used herein and unless otherwise specified, the terms "about" and "approximately," when used in connection with a numeric value or a range of values which is provided to characterize a particular solid form, e.g., a specific temperature or temperature range, such as, e.g., that describing a DSC or TGA thermal event, including, e.g., melting, dehydration, desolvation or glass transition events; a mass change, such as, e.g., a mass change as a function of temperature or humidity; a solvent or water content, in terms of, e.g., mass or a percentage; or a peak position, such as, e.g., in analysis by IR or Raman spectroscopy or XRPD; indicate that the value or range of values may deviate to an extent deemed reasonable to one of ordinary skill in the art while still describing the particular solid form. For example, in particular embodiments, the terms "about" and "approximately," when used in this context and unless otherwise specified, indicate that the numeric value or range of values may vary within 25%, 20%, 15%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1.5%, 1%, 0.5%, or 0.25% of the recited value or range of values.

[0019] As used herein, and unless otherwise specified, the terms "treat," "treating" and "treatment" refer to the eradication or amelioration of a disease or disorder, or of one or more symptoms associated with the disease or disorder. In certain embodiments, the terms refer to minimizing the advancement or worsening of the disease or disorder resulting

from the administration of a compound of the invention to a patient with such a disease or disorder. In some embodiments, the terms refer to the administration of a compound provided herein, with or without other additional active agents, after the onset of symptoms of the particular disease. The terms "treating", "treatment", or the like, as used herein covers the treatment of a disease-state in a subject, e.g., a mammal, and includes at least one of: (i) inhibiting the disease-state, i.e., partially or completely arresting its development; (ii) relieving the disease-state, i.e., causing regression of symptoms of the disease-state, or ameliorating a symptom of the disease; and (iii) reversal or regression of the disease-state, preferably eliminating or curing of the disease. In a particular embodiment the terms "treating", "treatment", or the like, covers the treatment of a disease-state in a mammal, e.g., a primate, e.g., a human, and includes at least one of (i), (ii) and (iii) above. As is known in the art, adjustments for systemic versus localized delivery, age, body weight, general health, sex, diet, time of administration, drug interaction and the severity of the condition may be necessary, and will be ascertainable with routine experimentation by one of ordinary skill in the art.

[0020] As used herein, and unless otherwise specified, the terms "prevent", "preventing" and "prevention" refer to the prevention of the onset, recurrence or spread of a disease or disorder, or of one or more symptoms thereof. In certain embodiments, the terms refer to the administration of a compound provided herein to a subject, with or without other additional active compound, prior to the onset of symptoms, particularly to patients at risk of diseases or disorders provided herein. The terms encompass the inhibition or reduction of a symptom of the particular disease. Subjects with familial history of a disease in particular are candidates for preventive regimens in certain embodiments. In addition, subjects who have a history of recurring symptoms are also potential candidates for the prevention. In this regard, the term "prevention" may be interchangeably used with the term "prophylactic treatment." In certain embodiments, the prevention is achieved by administration of a prophylactically effective amount of a compound of the invention.

[0021] As used herein, and unless otherwise specified, a "therapeutically effective amount" of a compound is an amount sufficient to provide a therapeutic benefit in the treatment or management of a disease or disorder, or to delay or minimize one or more symptoms associated with the disease or disorder. A therapeutically effective amount of a compound means an amount of therapeutic agent, alone or in combination with other therapies, which provides a therapeutic benefit in the treatment or management of the disease or disorder. The term "therapeutically effective amount" can encompass an amount that improves overall therapy, reduces or avoids symptoms or causes of disease or disorder, or enhances the therapeutic efficacy of another therapeutic agent.


[0022] As used herein, and unless otherwise specified, the terms "manage", "managing" and "management" refer to preventing or slowing the progression, spread or worsening of a disease or disorder, or of one or more symptoms thereof. Often, the beneficial effects that a subject derives from a prophylactic and/or therapeutic agent do not result in a cure of the disease or disorder. In this regard, the term "managing" encompasses treating a subject who had suffered from the particular disease in an attempt to prevent or minimize the recurrence of the disease.

[0023] As used herein, and unless otherwise specified, a "prophylactically effective amount" of a compound is an amount sufficient to prevent a disease or disorder, or prevent its recurrence. A prophylactically effective amount of a compound means an amount of therapeutic agent, alone or in combination with other agents, which provides a prophylactic benefit in the prevention of the disease. The term "prophylactically effective amount" can encompass an amount that improves overall prophylaxis or enhances the prophylactic efficacy of another prophylactic agent.

[0024] The term "composition" as used herein is intended to encompass a product comprising the specified ingredients (and in the specified amounts, if indicated), as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts. The term "pharmaceutical composition" encompasses compositions containing a compound of the invention, e.g., crystal Form I or II, and a pharmaceutically acceptable carrier. A "pharmaceutically acceptable carrier" is a diluent, excipient or carrier that is compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

II. Compounds of the Invention

[0025] In one embodiment, the invention provides a pharmaceutical formulation for oral administration of crystalline Form I and Form II of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate having the following formula.



[0026] For clarity, the alpha-7 receptor agonist compound (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride was disclosed in United States Patent Application Publication No. US 2005-0119325. However, in contrast to the present invention, such disclosure did not disclose or suggest the present invention, (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate, nor did it disclose any crystal forms of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate. These crystalline forms comprised in the inventive pharmaceutical formulations for oral administration are defined below.

[0027] In one embodiment, the invention provides a crystalline Form I of (R)-7-chloro-N-(quinuclidin-3-yl)benzo [b] thiophene-2-carboxamide hydrochloride monohydrate, characterized by an x-ray powder diffraction pattern having peaks expressed as 2θ at one or both of 17.48 and 20.58 ± 0.20 degrees when measured against an internal silicon standard.

[0028] In another embodiment, the invention provides the crystalline Form I as defined above, characterized by an x-ray powder diffraction pattern further having at least one peak expressed as 2θ at $4.50, 9.04, 14.60, 15.14, 15.80, 16.60, 18.16, 18.44, 19.48, 21.74,$ and 25.46 ± 0.20 degrees when measured against an internal silicon standard.

[0029] In another embodiment, the invention includes the crystalline Form I as defined above, characterized by an x-ray powder diffraction pattern further having at least two peaks expressed as 2θ at $4.50, 9.04, 14.60, 15.14, 15.80, 16.60, 18.16, 18.44, 19.48, 21.74$ and 25.46 ± 0.20 degrees when measured against an internal silicon standard.

[0030] In yet another embodiment, the invention provides the crystalline Form I as defined above, characterized by an x-ray powder diffraction pattern further having at least four peaks expressed as 2θ at $4.50, 9.04, 14.60, 15.14, 15.80, 16.60, 18.16, 18.44, 19.48, 21.74$ and 25.46 ± 0.20 degrees when measured against an internal silicon standard.

[0031] The invention further includes the crystalline Form I as defined above, characterized by an x-ray powder diffraction pattern further having at least six peaks expressed as 2θ at $4.50, 9.04, 14.60, 15.14, 15.80, 16.60, 18.16, 18.44, 19.48, 21.74$ and 25.46 ± 0.20 degrees when measured against an internal silicon standard.

[0032] The invention further includes the crystalline Form I as defined above, characterized by an x-ray powder diffraction pattern further having at least eight peaks expressed as 2θ at $4.50, 9.04, 14.60, 15.14, 15.80, 16.60, 18.16, 18.44, 19.48, 21.74$ and 25.46 ± 0.20 degrees when measured against an internal silicon standard.

[0033] The invention further provides the crystalline Form I as defined above, characterized by an x-ray powder diffraction pattern further having peaks expressed as 2θ at $4.50, 9.04, 14.60, 15.14, 15.80, 16.60, 18.16, 18.44, 19.48, 21.74$ and 25.46 ± 0.20 degrees when measured against an internal silicon standard.

[0034] In another embodiment, the present invention provides a crystalline Form II of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate, characterized by an x-ray powder diffraction pattern having peaks expressed as 2θ at one or both of 21.16 and 21.38 ± 0.20 degrees when measured against an internal silicon standard.

[0035] In another embodiment, the present invention provides the crystalline Form II as defined above, characterized by an x-ray powder diffraction pattern further having at least one peak expressed as 2θ at $4.48, 9.00, 13.58, 15.62, 16.48, 19.02, 19.44, 22.46$ and 25.00 ± 0.20 degrees when measured against an internal silicon standard.

[0036] In yet another embodiment, provided herein is the crystalline Form II as defined above, characterized by an x-ray powder diffraction pattern further having at least two peaks expressed as 2θ at $4.48, 9.00, 13.58, 15.62, 16.48, 19.02, 19.44, 22.46$ and 25.00 ± 0.2 degrees when measured against an internal silicon standard.

[0037] In another embodiment, the present invention includes the crystalline Form II as defined above, characterized by an x-ray powder diffraction pattern further having at least four peaks expressed as 2θ at $4.48, 9.00, 13.58, 15.62, 16.48, 19.02, 19.44, 22.46$ and 25.00 ± 0.2 degrees when measured against an internal silicon standard.

[0038] Further provided herein is the crystalline Form II as defined above, characterized by an x-ray powder diffraction pattern further having at least six peaks expressed as 2θ at $4.48, 9.00, 13.58, 15.62, 16.48, 19.02, 19.44, 22.46$ and 25.00 ± 0.2 degrees when measured against an internal silicon standard.

[0039] Also provided here is the crystalline Form II as defined above, characterized by an x-ray powder diffraction pattern further having at least eight peaks expressed as 2θ at $4.48, 9.00, 13.58, 15.62, 16.48, 19.02, 19.44, 22.46$ and 25.00 ± 0.2 degrees when measured against an internal silicon standard.

[0040] In another embodiment, provided herein is the crystalline Form II as defined above, characterized by an x-ray powder diffraction pattern further having peaks expressed as 2θ at $4.48, 9.00, 13.58, 15.62, 16.48, 19.02, 19.44, 22.46$ and 25.00 ± 0.2 degrees when measured against an internal silicon standard.

III. Methods of the Invention

A. Pharmaceutical formulation for use in methods of treatment

[0041] Described herein is a pharmaceutical formulation of the crystalline Form I for the treatment and/or prophylaxis of a disease which can be treated or prevented by alpha-7 receptor activation. Also described herein is a pharmaceutical formulation of crystalline Form II for the treatment and/or prophylaxis of a disease which can be treated or prevented by alpha-7 receptor activation.

[0042] Also described herein is a pharmaceutical formulation for use in a method of treating or preventing a disease which can be treated or prevented by alpha-7 receptor activation comprising administering to a subject the pharmaceutical formulation comprising crystalline Form I. Also described herein is a pharmaceutical formulation for use in a method of treating or preventing a disease which can be treated or prevented by alpha-7 receptor activation comprising administering to a subject the pharmaceutical formulation comprising crystalline Form II.

[0043] In one embodiment, the present invention provides a pharmaceutical formulation provided herein for use in a method for improving cognition or treating cognitive loss in a subject comprising administering to a subject the pharmaceutical formulation comprising crystalline Form I. In another embodiment, the present invention provides the pharmaceutical formulation provided herein for use in a method for improving cognition or treating cognitive loss by administering to a subject the pharmaceutical formulation of crystalline Form I wherein the subject is suffering from a disorder selected from: attention deficit disorder, attention deficit hyperactivity disorder, and Parkinson's Disease. In another embodiment, the present invention provides the pharmaceutical formulation for use in a method for improving cognition or treating cognitive loss by administering the pharmaceutical formulation comprising crystalline Form I wherein the subject is suffering from a disorder selected from Alzheimer's Disease, pre-senile dementia and schizophrenia.

[0044] In another embodiment, the present invention provides the pharmaceutical formulation provided herein for use in a method of treating a disorder selected from attention deficit disorder, attention deficit hyperactivity disorder, Parkinson's Disease, Alzheimer's Disease and schizophrenia, the method comprising administering to a subject the pharmaceutical formulation comprising crystalline Form I. In another embodiment, the present invention includes the pharmaceutical formulation provided herein for use in a method of treating a subject that is at risk for developing a disorder selected from: Alzheimer's disease, Parkinson's Disease and schizophrenia, the method comprising administering to the subject the pharmaceutical formulation comprising crystalline Form I. In yet another embodiment, the present invention includes the pharmaceutical formulation provided herein for use in a method of treating a subject over age 60, comprising administering to the subject the pharmaceutical formulation comprising crystalline Form I. In a further embodiment, the present invention includes the pharmaceutical formulation provided herein for use in a method of treating a subject for age-related memory loss, comprising administering to the subject the pharmaceutical formulation comprising crystalline Form I. In another embodiment the present invention includes the pharmaceutical formulation provided herein for use in a method of treating a subject for age-related memory loss, comprising administering to the subject the pharmaceutical formulation comprising crystalline Form I wherein the subject is over age 60.

[0045] In another embodiment, the present invention provides the pharmaceutical formulation provided herein for use in a method for improving cognition or treating cognitive loss in a subject comprising administering to a subject the pharmaceutical formulation comprising crystalline Form II. In another embodiment, the present invention provides the pharmaceutical formulation provided herein for use in a method for improving cognition or treating cognitive loss by administering the pharmaceutical formulation comprising crystalline Form II to a subject, wherein the subject is suffering from a disorder selected from: attention deficit disorder, attention deficit hyperactivity disorder, and Parkinson's Disease. In another embodiment, the present invention provides the pharmaceutical formulation provided herein for use in a method for improving cognition or treating cognitive loss by administering the pharmaceutical formulation comprising crystalline Form II to a subject, wherein the subject is suffering from a disorder selected from Alzheimer's Disease, pre-senile dementia and schizophrenia.

[0046] In another embodiment, the present invention provides the pharmaceutical formulation provided herein for use in a method of treating a disorder selected from attention deficit disorder, attention deficit hyperactivity disorder, Parkinson's Disease, Alzheimer's Disease and schizophrenia, the method comprising administering to a subject the pharmaceutical formulation comprising crystalline Form II. In another embodiment, the present invention includes the pharmaceutical formulation provided herein for use in a method of treating a subject that is at risk for developing a disorder selected from: Alzheimer's disease, Parkinson's Disease and schizophrenia, the method comprising administering to the subject the pharmaceutical formulation comprising crystalline Form II. In yet another embodiment, the present invention includes the pharmaceutical formulation provided herein for use in a method of treating a subject over age 60, comprising administering to the subject the pharmaceutical formulation comprising crystalline Form II. In a further embodiment, the present invention includes the pharmaceutical formulation provided herein for use in a method of treating a subject for age-related memory loss, comprising administering to the subject the pharmaceutical formulation comprising crystalline Form II. In another embodiment the present invention includes the pharmaceutical formulation provided herein for use in a method of treating a subject for age-related memory loss, comprising administering to the subject the pharmaceutical formulation comprising crystalline Form II wherein the subject is over age 60.

B. Methods of Preparation

[0047] It would be beneficial to provide (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate in a stable crystalline form. After extensive studies, two types of stable crystalline forms were identified: Form I and Form II. One crystalline form does not convert to the other readily in solid condition because each of two

crystalline forms is stable. On the other hand, it was found that when one crystalline form was dissolved in an aqueous solvent and the crystallization was carried out from the solution, it was difficult to predict which crystalline form was preferentially produced. In addition, one form might be converted to the other or a mixture of two forms quite easily in the solution under certain conditions. Therefore the mechanism of crystallization was unclear, and it was difficult to design methods for producing each form at high purity. After extensive investigation, the inventors arrived at the methods for selectively manufacturing each pure crystalline form. The methods can be carried out using a variety of different solvents.

[0048] A crystal form of a compound is usually obtained by: 1) dissolving the compound in a solvent at a high temperature, where the solubility of the product is high, 2) lowering the temperature of the solution to cause crystallization of the compound, and 3) isolating the resulting crystals.

[0049] However, solid state investigations revealed that there are two enantiotropic crystalline forms of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate, and the present inventors discovered that the usual crystallization procedure is apt to produce a mixture of the two crystal forms of the compound. This is because at higher temperatures, one crystal form of the compound is a little more stable in a solvent system, whereas in at lower temperatures, the other crystal form of the compound is a little more stable in the same solvent system. This effect can be seen in Examples 5, 6, 7 and 8. Furthermore, the boundary temperature for converting from one form to the other was found to vary depending on the solvent system in which the compound is dissolved.

[0050] The inventors extensively investigated the stable crystal forms at various temperatures and in various solvent systems and recognized that Form I and Form II of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate can be separately prepared based on the relationship of temperature and water activity of the solvent independently of the particular organic solvent. This recognition led to the creation of novel methods for separately manufacturing pure Form I and pure Form II.

[0051] Described herein is a method for preparing a pharmaceutical composition, the method comprising combining the crystalline form I with an excipient or pharmaceutically acceptable carrier. The method may further include combining the crystalline form I with a liquid. The method may include filling a capsule with a composition comprising the crystalline form I.

[0052] Described herein is a method for preparing a pharmaceutical composition comprising combining the crystalline form II with an excipient or pharmaceutically acceptable carrier. The method may further include combining the crystalline form II with a liquid. The method may include filling a capsule with a composition comprising the crystalline form II.

[0053] A method of manufacturing a crystalline Form I comprises: (1) stirring (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride (hereinafter, "(R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride" includes any form of anhydrate, hydrate and solvate, preferably anhydrate and hydrate) in an aqueous organic solvent within the temperature-water activity range I, if required with decreasing the temperature and/or water activity thereof, to form a substantially pure crystalline Form I; and (2) isolating the resulting crystalline Form I, wherein the temperature-water activity range I is defined by the following relationship of the temperature and the water activity of the aqueous organic solvent, the water activity (x) of the aqueous organic solvent is from 0.16 to 0.73; and the temperature (T) of the aqueous organic solvent is higher than $(183x - 64.2)$ and lower than the boiling temperature of the aqueous organic solvent.

[0054] The method of manufacturing crystalline Form I may include a method where, in step (1) the initial state of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride in an aqueous organic solvent is a solution, and the solution is stirred within the temperature-water activity range I with decreasing the temperature and/ or water activity thereof. The method of manufacturing crystalline Form I may include a method where in step (1) the initial state of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride in an aqueous organic solvent is a suspension, and the suspension is stirred within the temperature-water activity range I, if required with decreasing the temperature and/ or water activity thereof. The method of manufacturing crystalline Form I may include a method where in step (1) the seed crystals of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride is added to the solution. The method of manufacturing crystalline Form I may also include a method where the seed crystals are Form I of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate.

[0055] As described herein, the method of manufacturing crystalline Form I may include a method where the water activity is from 0.29 to 0.59. The method of manufacturing crystalline Form I may include a method where the temperature is between -10°C and 60°C and higher than the following $T: T = 183x - 57.6$; wherein x is water activity of the aqueous organic solvent and T is a temperature ($^{\circ}\text{C}$). The method of manufacturing crystalline Form I may also include a method where the end point temperature in step (1) is between 0°C and 35°C .

[0056] As described herein, the method of manufacturing crystalline Form I may include a method where the aqueous organic solvent is mixture of water and one or more of organic solvents which are miscible with water and are selected from alcohols, ketones, nitriles and ethers. The method of manufacturing crystalline Form I may include a method where the aqueous organic solvent is mixture of water and one or more of organic solvents which are miscible with water and are selected from propanol, butanol, butanone and acetonitrile.

[0057] Also described herein is a method of manufacturing a crystalline Form II; comprising: (1) stirring (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride in an aqueous organic solvent within the temperature-water activity range II, if required with decreasing the temperature and/or water activity thereof, to form a crystalline Form II, and (2) isolating the resulting crystalline Form II, wherein the temperature-water activity range II is defined by the following relationship of the temperature and the water activity of the aqueous organic solvent, the water activity (x) of the aqueous organic solvent is from 0.16 to 0.73; and the temperature (T) of the aqueous organic solvent is lower than $(183x - 64.2)$ and higher than the freezing-point temperature of the aqueous organic solvent.

[0058] The method of manufacturing crystalline Form II may include the method where in step (1) the initial state of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride in an aqueous organic solvent is a solution, and the solution is stirred within the temperature-water activity range II with decreasing the temperature and/or water activity thereof. The method of manufacturing crystalline Form II may include a method wherein in step (1) the initial state of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride in an aqueous organic solvent is a suspension, and the suspension is stirred within the temperature-water activity range II, optionally decreasing the temperature and/or water activity thereof.

[0059] As described herein, the method of manufacturing crystalline Form II may include a method wherein in step (1) the seed crystals of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride is added to the solution. The method of manufacturing crystalline Form II may include a method wherein the seed crystals are Form II of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate. The method of manufacturing crystalline Form II may include a method wherein the water activity is from 0.29 to 0.59. The method of manufacturing crystalline Form II may also include a method wherein the temperature is between -10°C and 60°C and lower than the following $T: T = 183x - 70.8$; wherein x is water activity of the aqueous organic solvent and T is a temperature ($^{\circ}\text{C}$). The method of manufacturing crystalline Form II may include a method wherein the end point temperature in step (1) is between 0°C and 35°C . The method of manufacturing crystalline Form II may include a method wherein the aqueous organic solvent is mixture of water and one or more of organic solvents which are miscible with water and are selected from alcohols, ketones, nitriles and ethers.

[0060] As described herein, the method of manufacturing crystalline Form II may include a method wherein the aqueous organic solvent is mixture of water and one or more of organic solvents which are miscible with water and are selected from propanol, butanol, butanone and acetonitrile.

[0061] Also described herein is another method for preparing the crystalline Form I, comprising: (a) heating 10 - 30% by weight of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride in acetonitrile or an aqueous acetonitrile to between 60°C and the boiling point of the solution, (b) optionally adding water to the mixture to fully dissolve the (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride; (c) cooling the solution until crystals are just visible; (d) if the water content is greater than 3% volume/volume when crystals are just visible, adding acetonitrile to the mixture so that the water content is less than 3% volume/volume; (e) cooling the resulting mixture to below 15°C ; and (f) isolating crystalline (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate.

[0062] The method of preparing the crystalline Form I may include the method as defined in steps (a) through (f) above, wherein the water added in step (a) does not bring the water content of the mixture above 30% volume/volume. The method of preparing the crystalline Form I may include the method as defined in steps (a) through (f) above, wherein (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride is present at 15-25% by weight in step (a). The method of preparing the crystalline Form I may include the method as defined in steps (a) through (f) above, wherein (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride is present at 16-20% by weight in step (a). The method of preparing the crystalline Form I may include the method as defined in steps (a) through (f) above, wherein (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride is present at 17-19% by weight in step (a). The method of preparing the crystalline Form I may include the method as defined in steps (a) through (f) above, further comprising adding Form I (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate to the mixture after crystals are just visible.

[0063] As described herein, the method of preparing the crystalline Form I may include the method as defined in steps (a) through (f) above, wherein step (c) comprises cooling the solution to below 55°C . The method of preparing the crystalline Form I may include the method as defined in steps (a) through (f) above, wherein step (c) comprises cooling the solution to below 50°C .

[0064] Also described herein is a method for preparing the crystalline Form II, the method comprising: (a) heating 5 - 15 % by weight of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride in 2-butanol or an aqueous 2-butanol to between 60°C and the boiling point of the solution; (b) if the water content is smaller than 5% volume/volume, adding water to the mixture so that the water content is not less than 5% volume/volume; (c) cooling the solution to below 10°C ; (d) keeping the resulting mixture to below 10°C ; and (e) isolating crystalline (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate.

[0065] Further described herein is a method for preparing the crystalline Form II, the method comprising: (a) adding

(R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride to (i) acetonitrile or (ii) aqueous acetonitrile to create a composition that is 10-20% by weight of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride; (b) optionally adding water to the composition to make the water content 6-10%; (c) optionally cooling the solution at below 10 °C; (d) allowing crystals to form; and (e) isolating crystalline (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate.

[0066] The crystalline Forms I and II are very stable in many aspects. Both forms are stable under storage conditions. No degradation products from either form were detected under the storage conditions: 11 %RH at 40 °C, 75 %RH at 40 °C, 11 %RH at 60°C and 75 %RH at 60 °C after 2 weeks, and no degradation products from either form were detected under the photo storage conditions: exposure to light (D65 lamp) of 1.2 million Lux hours at 25 °C. Both forms are also stable under physical stress. The XRD charts of both forms were not changed after compression experiments with a planar pestle (1000 kgf/cm²).

[0067] The pure crystalline Forms I and II can be manufactured by the special methods, which are described herein.

[0068] Pure crystalline Form I can be manufactured by a method which comprises: (1) stirring (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride in an aqueous organic solvent within the temperature-water activity range I (as depicted in Figure 4), if required with decreasing the temperature and/ or water activity thereof, to form a substantially pure crystalline Form I, and (2) isolating the resulting crystalline Form I.

[0069] Pure crystalline Form II can be manufactured by the method which comprises: (1) stirring (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride in an aqueous organic solvent within the temperature-water activity range II (as depicted in Figure 4), if required with decreasing the temperature and/ or water activity thereof, to form a crystalline Form II, and (2) isolating the resulting crystalline Form II.

[0070] Water activity is a thermophysical coefficient used to represent the energy status of the water in a system and is defined as the vapor pressure of water above a sample divided by that of pure water at the same temperature. It can be measured with a capacitance hygrometer or a dew point hygrometer. It can be also predicted by COSMO-RS method (Fluid Phase Equilibria, 172 (2000) 43-72).

[0071] (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride was prepared, for example, by the method described in WO03/55878. 7-Chloro-benzo[b]thiophene-2-carboxylic acid was reacted with carbonyldiimidazole to give 7-chloro-2-imidazolyl-carbonylbenzo[b]thiophene, followed by reacting with (R)-3-aminoquinuclidine dihydrochloride to give (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride.

[0072] (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride used for the above manufacturing methods can be for example crystals (e.g., Forms I, II, and mixtures thereof), amorphous products, an oil or a solution, and preferably a solution. The crystallization can be performed in the same vessel after hydrochlorination. An aqueous organic solvent is a mixture of water and one or more organic solvents. The preferable organic solvents are water-miscible organic solvents, and more preferable are for example alcohols (e.g., C₁₋₆alkanol such as methanol, ethanol, 1-propanol, 2-propanol, 1-butanol, 2-butanol, and C₂₋₆alkanediol such as ethylene glycol, propylene glycol), ketones (e.g., C₃₋₆alkanone such as acetone, butanone), nitriles (e.g., acetonitrile, propanonitrile) and ethers (e.g. dimethoxyethane, tetrahydrofuran). Preferable solvents are alcohols, nitriles and ketones, and more preferable are propanols, butanols, butanone and acetonitrile.

[0073] The solution may be supersaturated prior to the formation of crystals. The boundary between the temperature-water activity ranges for Forms I and II is shown in Figure 4 as a line which divides the domains of Forms I and II.

[0074] Crystal forms may be monitored during the production method. Any analytical methods can be used for monitoring as long as it can distinguish crystal forms, and XRD is one of most preferable methods. In order to manufacture a pure form, stirring of the mixture is continued until undesired form completely converts to the desired form.

[0075] In the method for manufacturing Form I, Form X which is different from both Forms I and II may appear temporally, but Form X can be converted to Form I and disappears if stirring of the mixture is continued.

IV. Pharmaceutical Formulations of the Invention

[0076] Also provided herein in the present invention is a pharmaceutical formulation for oral administration comprising the crystalline Form I. Also provided herein is a pharmaceutical formulation for oral administration comprising the crystalline Form II.

[0077] The crystalline Forms I and II may be used to prepare a medicament to treat disease or condition in a mammal in need thereof, wherein mammal would receive symptomatic relief from the administration of a therapeutically effective amount of a crystalline Form I or II. The crystalline Forms I and II may be administered in combination with other medications for additive or synergistic therapeutic benefit for a given disease. Diseases include, but are not limited to, those described below. Medications include, but are not limited to, drugs which are approved for a given indication e.g. acetylcholinesterase inhibitors for Alzheimer's Disease.

[0078] Because Form I is very stable and can be stored for a considerable length of time prior to its use in the preparation of a drug product, Form I is useful in the manufacture of drug product even when the manufacturing process, i.e., the

formulation of the active ingredient, causes some or all of the Form I to convert to another form.

[0079] The crystalline Forms I and II may be formulated as solutions or suspensions, in the form of tablets, capsules (each including timed release and sustained release formulations), pills, oils, powders, granules, elixirs, tinctures, suspensions, syrups, emulsions, microemulsions, or with excipients. Likewise, they may also be administered by any conventional route, for example in intravenous (both bolus and infusion), intraperitoneal, intraocularly, subcutaneous, intramuscular form, enterally, preferably orally (e.g., in the form of tablets or capsules), or in a nasal, buccal, sub-lingual, transdermal, or a suppository form, using well known formulations to those of ordinary skill in the pharmaceutical arts. In addition, the crystalline Forms I and II can be administered in the form of liposomes or the like. Disintegrators include, without limitation, delivery systems such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines. According to the present invention, the pharmaceutical formulation is a tablet or capsule.

[0080] For oral administration in the form of a tablet or capsule, the crystalline Forms I and II can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Suitable lubricants used in these dosage forms include, for example, sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Suitable disintegrating agents are, for example, starches, carboxymethylstarch sodium, crosscarmellose sodium and the like. Examples of the suitable coloring agents are iron sesquioxide, yellow iron sesquioxide, amaranth, erythrosine, tartrazine, Sunset Yellow FCF and the like.

[0081] The dosage regimen for the pharmaceutical formulations comprising crystalline Forms I and II is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient. An ordinarily skilled physician or veterinarian can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition.

[0082] In one embodiment satisfactory results in animals are indicated to be obtained at a daily dosage of from about 0.1 to about 600 mg or from about 0.01 to about 5 mg/kg animal body weight.

[0083] Injected intravenous, subcutaneous or intramuscular dosages of the crystalline Forms I and II, when used for the indicated effects, will range between about 0.001 to 1.0 mg/kg. Furthermore, the crystalline Forms I and II can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration can be continuous rather than intermittent throughout the dosage regimen. Transdermal delivery can also be achieved using approaches known to those skilled in the art.

[0084] Diseases that may be treated using the pharmaceutical formulations comprising crystalline Forms I and II include, but are not limited to: condition cognitive and attention deficit symptoms of Alzheimer's, neurodegeneration associated with diseases such as Alzheimer's disease, pre-senile dementia (mild cognitive impairment), senile dementia, schizophrenia, psychosis attention deficit disorder, attention deficit hyperactivity disorder, mood and affective disorders, amyotrophic lateral sclerosis, borderline personality disorder, traumatic brain injury, behavioral and cognitive problems associated with brain tumors, AIDS dementia complex, dementia associated with Down's syndrome, dementia associated with Lewy Bodies, Huntingdon disease, depression, general anxiety disorder, age related macular degeneration, Parkinson's disease, tardive dyskinesia, Pick's disease, post traumatic stress disorder, dysregulation of food intake including bulimia and anorexia nervosa, withdrawal symptoms associated with smoking cessation and dependant drug cessation, Gilles de la Tourette's Syndrome, glaucoma, neurodegeneration associated with glaucoma or symptoms associated with pain or is the treatment and/or prophylaxis for the improvement of perception, concentration, learning and/or memory.

EXEMPLIFICATION

[0085] The present invention is illustrated by the following examples, which are not intended to be limiting in any way.

Example 1

Preparation of the crystalline Form I

[0086] (R)- 7-Chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride was synthesized by the procedure described in US 2005-0119325. To prepare Form I, 1.0 Kg of the compound was dissolved in acetonitrile (5 L) and heated to 72 - 78 °C. Once at this temperature, water (0.5 L) was added. The mixture was cooled to 50 - 60 °C, wherein crystals are just visible and seed with Form I seed crystals. The mixture was held for a minimum of 2 hours,

and then acetonitrile (20 L) was added while maintaining an internal temperature of 50 - 60 °C. The material was cooled to 5 - 10 °C. Crystals were isolated by vacuum filtration and washed with acetonitrile (2 L). The material was dried at 40 °C in a vacuum oven with humidity control to provide 0.8 kg of pure Form I.

Example 2

Preparation of the crystalline Form I

[0087] Acetonitrile (90 mL) and water (10 mL) were mixed at room temperature. 1.0ml of this solution was added to 100.7 mg of the crystalline Form I of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate. This suspension was stirred at 80 °C until solid component was dissolved, then the temperature was decreased to 40°C for 80 minutes. During the cooling, spontaneous crystallization was observed round 52°C. To the suspension, 2.40 ml of acetonitrile was dropped slowly, and then the temperature was decreased to 10 °C for 60 minutes. The suspension was stirred at same temperature for 15 hours, and then the solid was filtered and washed with 0.20 ml of acetonitrile. After vacuum drying, 81.1mg of the crystalline Form I was recovered.

Example 3

Preparation of the crystalline Form II

[0088] Acetonitrile (90 mL) and water (10 mL) were mixed at room temperature. 1.0ml of this solution was added to 100.9 mg of the crystalline Form I of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate. This suspension was stirred at 80 °C until solid component was dissolved, then the temperature was decreased to 10 °C for 140 minutes. During the cooling, spontaneous crystallization was observed round 51 °C. The suspension was stirred at same temperature for 15 hours, then the solid was filtered and washed with 0.20 ml of acetonitrile. After vacuum drying, 48.7mg of the crystalline Form II was recovered.

Example 4

Preparation of the crystalline Form II

[0089] (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate (462 g) was triturated in 2308.5 mL acetonitrile and 230.85 mL of water at ambient temperature for 4.75 hours. The product was isolated by filtration and dried to afford 314 g of isolated pure Form II.

Example 5

Phase diagram in aqueous acetonitrile

(1) Solubility measurements of Forms I and II

[0090] Solubility of Forms I and II was measured at various temperatures between 5 °C and 45 °C in aqueous acetonitrile in which water concentration was from 0 to 10 v/v % respectively.

[0091] Solubility was measured according to the following procedure. Form I or II crystals and an aqueous acetonitrile were added to a glass vessel. The mixture was stirred with a Teflon-coated magnetic stirrer bar at a defined temperature controlled with aluminum block. The liquid phase was sampled periodically, and the concentration of the compound was measured with high performance liquid chromatography (HPLC). Solid material was also collected at the same time to identify the crystal form using XRPD. In analyzing the time course of the change in concentration, that plateau zone was identified as an equilibrium condition, and the mean value of these concentrations was defined as "solubility" under that condition. Measured solubility is summarized in Table 1.

Table 1 Results of the solubility measurement

Conditions		Solubility			Stable form (Estimated from solubility)
Water concentration [v/v %]	Temp [°C]	Form I	Form II	Form X	
0	5	1.0	*	**	I
	25	1.3	*	**	I

(continued)

Conditions		Solubility			Stable form (Estimated from solubility)
Water concentration [v/v %]	Temp [°C]	Form I	Form II	Form X	
2	5	2.4	2.6	**	I
	25	4.1	4.7	4.2	I
	45	8.7	*	9.1	I
3.5	12.5	5.4	5.7	**	I
	37.5	10.0	*	10.3	I
6	5	12.8	11.2	**	II
	25	17.1	16.7	**	II
	45	27.2	29.6	**	I
8.5	12.5	27.6	25.1	**	II
	37.5	44.4	44.4	**	I and II
10	5	*	29.7	**	II
	25	47.5	44.6	**	II
	45	71.6	74.5	**	I
* Could not measure ** Did not conduct					

[0092] (2) Solubility equations were modeled using JMP 6 (SAS Institute). With the response surface method, measured solubility values were applied to equation (1) to obtain solubility models as a function of temperature and water concentration. W' and T' were defined as equations (2) and (3) respectively. Here, C_x^* , W and T mean solubility of a certain crystal form, water concentration in aqueous acetonitrile (v/v %) and temperature (°C) respectively. Constant values from a, b, c, d and f were determined by applying measured solubility values with least squares fitting method within the range of 2 to 10 v/v % water concentration and within the range of 5 to 45 °C. Perspective of defects or accuracy, measured solubility values at 2 % - 45 °C of both forms, 3.5% - 37.5 °C of form II and 10 % - 10.0 °C of form I are excluded to build mathematical formulas.

$$C_x^* = \text{Exp}(a + b W + c T' + d W T' + e W^2 + f T'^2) \quad (1)$$

$$W = (W - 6) / 4 \quad (2)$$

$$T' = (T - 25) / 20 \quad (3)$$

[0093] Fitting results showed that constants were determined successfully, and these models can describe solubility within the range with only minor deviation. Obtained solubility equations of Forms I and II are shown in equation (4) and (5) respectively.

$$C_{I}^* = \text{Exp}(2.8448 + 1.2517 W + 0.4185 T' - 0.1086 W T' - 0.2249 W^2 + 0.0681 T'^2) \quad (4)$$

$$C_{II}^* = \text{Exp}(2.8389 + 1.1503 W + 0.5101 T' - 0.0638 W T' - 0.1888 W^2 + 0.0488 T'^2) \quad (5)$$

(3) Development of phase diagram

[0094] Thermodynamic relationship between polymorphic crystal phases is consistent with solubility. Based on the

results of the solubility measurements, it is apparent that thermodynamic relationship between forms I and II crystals is enantiotropy. Boundary of stable crystal form should exist in the range that solubility measurement was carried out. At boundary condition, solubility of forms I and II should be same. Hence, boundary condition can be induced from equation (4) and (5) and simplified as described in equation (6).

$$0.0059 + 0.1014 W - 0.0916 T - 0.0448 W T - 0.0361 W^2 + 0.0193 T^2 = 0 \quad (6)$$

[0095] By solving boundary equation (6), boundary condition can be determined. Solved values are shown in Table 2. By plotting the results, phase diagram was described in Figure 1. For convenience, this boundary line which obtained from equation (6) was fitted by fourth degree equation of water concentration. This approximation formula and its solved values are shown in equation (7) and Table 2 respectively.

$$T = 0.0056W^4 - 0.1305W^3 + 0.2831W^2 + 11.3942W - 31.3235 \quad (7)$$

Table 2 Solved values of boundary equations

Water concentration [%v/v]	Temperature[°C]	
	Equation (6)	Equation (7)
3.0	2.34	2.34
3.5	7.26	7.27
4.0	11.86	11.86
4.5	16.08	16.09
5.0	19.91	19.91
5.5	23.33	23.32
6.0	26.31	26.30
6.5	28.85	28.86
7.0	30.97	30.99
7.5	32.69	32.72
8.0	34.03	34.07
8.5	35.03	35.07
9.0	35.73	35.76
9.5	36.16	36.20
10.0	36.35	36.43

Example 6

Inter-conversion tests

[0096] Inter-conversion tests were also carried out in order to confirm reliability of obtained phase diagram of Example 5 (Figure 1).

[0097] Solvent was added to a glass vessel and temperature was controlled with aluminum block. Identical amounts of Forms I and II crystals were added to the vessel. The solutions were stirred for 13 to 40 hr with Teflon-coated magnetic stirrer bar. Solid component was sampled and analyzed with XRPD to determine its crystal form.

[0098] The results were summarized in Table 3. These results were consistent with phase diagram of Example 5.

Table 3 Experimental Results of Inter-conversion tests

run	Temperature [°C]	Water concentration [v/v %]	Results
1	10	3.0	I
2		3.5	I
3		4.0	II
4		4.5	II
5	20	4.0	I
6		4.5	I
7		5.0	I
8		5.5	II
9		6.0	II
10	30	5.5	I
11		6.0	I
12		6.5	I
13		7.0	I
14		7.5	II

Example 7***Crystallization behavior in aqueous acetonitrile*****(1) Form X**

[0099] Form X was found as another solid form in the solubility study. A typical XRPD pattern of form X is shown in Figure 2.

(2) Crystallization behavior in 98 v/v % aqueous acetonitrile

[0100] Form I crystals were added to 98 v/v % aqueous acetonitrile in a glass vessel. Next, 98 v/v % aqueous acetonitrile was added to make the mixture 40 v/w times relative to the Form I crystal. The mixture was stirred with Teflon-coated magnetic stirrer bar and heated to 80°C with aluminum block. After the crystals were dissolved, the mixture was cooled to a determined temperature at the rate of 30 °C per hour. After a determined holding time, Form I crystals were added as seed crystals according to object of an experiment. Precipitates were sampled periodically and analyzed by XRPD.

[0101] Initial precipitates in 98 v/v % aqueous acetonitrile system were confirmed as Form X. Spontaneous transformation from Form X to Form I was not observed within 16 hours. From the results of the seeded experiments, it was estimated that Form I is more stable than Form X regardless of temperature. Form X could be recovered in 84.7% yield.

Table 4 Results of crystallization experiments in 98 v/v % aqueous acetonitrile

Run	Temperature [°C]	Holding time [hr]	Crystal form
1	60	0	X
	40	0	X
	10	0	X
		15	X
		Seeding*	
		2	X
		8	X
		27	X(I)**
2	60	0	X
	40	0	X
		16	X
		Seeding*	
		2	I+X
		8	I
		27	I
3	60	0	X
	40	0	X
	10	0	X
		15	X
	-	-	X
* ca. 4 % w/w Form I crystals			
** Slight amount of Form I was detected			

(3) Transformation behavior in 98 v/v % aqueous acetonitrile

[0102] Form I crystals and 40 v/w times volume of 98 v/v % aqueous acetonitrile were added to a glass vessel. The mixture was stirred with Teflon-coated magnetic stirrer bar and heated to 80 °C with aluminum block. After dissolved, solution was cooled to 5 °C at the rate of 30 °C per hour. To the slurry of Form X, 10 w/w % of Form I crystals were added as seed crystals at 5 °C, then controlled to a certain temperature. Samples of solid material were analyzed by XRPD on a periodic basis.

[0103] Transformation from Form X to Form I was observed above ambient temperature. This tendency to transform was also observed at 5 °C. In 98 v/v % aqueous acetonitrile system, it was estimated that Form I is more stable than Form X regardless of temperature though transformation kinetics was extremely slow at a low temperature.

Table 5 Results of transformation experiments in 98 v/v % aqueous acetonitrile

Run	Temperature [°C]	Time [hr]	Crystal form
1	5	0*	X
		16	X
		40	X(I)**

EP 3 272 748 B1

(continued)

Run	Temperature [°C]	Time [hr]	Crystal form
2	25	0*	X
		16	I+X
		40	I
3	40	0*	X
		16	I
		40	I
*Right after the seeding at 5 °C			
**Slight amount of Form I was detected			

(4) Transformation behavior in 97 v/v % aqueous acetonitrile

[0104] Form X crystals and 97 v/v % aqueous acetonitrile were added to a glass vessel. The mixture was stirred with Teflon-coated magnetic stirrer bar at the temperature controlled with aluminum block. Solid component was sampled periodically and analyzed by XRPD to determine the crystal form.

[0105] Spontaneous transformation from Form X to Form I was observed in all experiments. It was estimated that Form I is more stable than Form X regardless of temperature in 97 v/v % aqueous acetonitrile.

Table 6 Results of transformation experiments in 97 v/v % aqueous acetonitrile

Run	Temperature [°C]	Time [hr]	Crystal form
1	10	1	X
		3	X
		21	I
2	20	1	X
		3	X
		21	I
3	30	1	X
		3	I
		21	I
4	40	1	X
		3	I
		21	I
5	50	1	I
		3	I
		21	I

Example 8

Inter-conversion tests in various solvents

[0106] A mixture of crystalline Forms I and II (25mg/25mg) was stirred in 0.5 mL of each organic solvent/water shown in Table 7 for 3 days at 5, 25, 40 and 60 °C, respectively, and the precipitates were filtrated, then the crystal forms were confirmed by XRPD. The results were shown in the Table 7.

Table 7 Results of inter-conversion tests in various solvents

Solvent	Water Concentration [v/v%]	Temperature[°C]			
		5	25	40	60
2-Propanol	0	I(II)	I(II)	I(II)	I
2-Propanol	2	I(II)	I	I	X
2-Propanol	5	I	I	I	I
2-Propanol	10	II	I	I	sol
1-Propanol	0	I/X	I/X	I/II	I/II
1-Propanol	2	I	X	I	sol
1-Propanol	5	I	I	I(II)	sol
1-Propanol	10	II	sol	sol	sol
Acetone	0	I	I/II	I/X	-
Acetone	2	I	I	I	-
Acetone	5	I	I	I	-
Acetone	10	II	I	I	-
1-Butanol	0	I	I	I	I
1-Butanol	2	I	I	I	I
1-Butanol	5	II	I	I	I
1-Butanol	10	II	sol	Sol	sol
2-Butanol	0	I	I	I	I
2-Butanol	2	I	I	I	I
2-Butanol	5	II	I	I	I
2-Butanol	10	II	II	II	sol
Acetonitrile	0	I/II	I	I	I
Acetonitrile	2	I/II	I	I	I
Acetonitrile	5	II	I	I	I
Acetonitrile	10	II	II	I/II	sol
2-Butanone	0	I	I/X	I/X	I/X
2-Butanone	2	II	I	I	I
2-Butanone	5	II	II	II	I
2-Butanone	10	II	sol	sol	sol
I: form I II: form II I/II: form - I (small amount of form -II) I/II: form -I/II mixture I/X: form -I/X mixture Sol: solution					

Example 9***Relationship between water concentrations and the values of water activity at various temperatures***

[0107] The values of water activity of various water concentrations (i.e., 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 and 20 v/v%) in various organic solvents (i.e., 1-propanol, 2-propanol, 1-butanol, 2-butanol, acetone, 2-butanone, and acetonitrile) at various temperatures (i.e., 0, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55 and 60 °C), respectively, were calculated using COSMOTHERME version 2.1 based on Cosmo-RS method.

[0108] The values of water concentrations (v/v %) were calculated on the boundary line (equation (6)) between Form I and Form II at various temperatures (i.e., 0, 5, 10, 15, 20, 25, 30, and 35 °C), and converted into the values of water activity using a regression curve of the cubic equation for plots of the values of water activity versus water concentrations at various temperatures (Figure 3). Table 8 indicates the relationship between water contents (v/v %) and the values of

water activity on the boundary line of the phase diagram. The values of water activity and the corresponding temperatures were fitted by equation (7) with a good correlation (correlation coefficient: 0.997).

$$T = 183X - 64.2 \quad (7)$$

Table 8 Relationship between water concentrations (v/v %) and the values of water activity at various temperatures

Temperature [°C]	Water Concentration [V/V %]	Water activity
0	2.77	0.349
5	3.27	0.382
10	3.79	0.411
15	4.37	0.437
20	5.01	0.460
25	5.77	0.483
30	6.76	0.509
35	8.48	0.551

[0109] The results of the inter-conversion tests were plotted and the boundary line by equation (7) was drawn on the phase diagram of Form I and Form II as shown in Figure 4. The boundary line approximately separated the Form I and Form II in all experimented solvent systems.

Example 10

Crystallization of crystalline Form I

(1) 1-Propanol

[0110] Crystalline Form I (100.1 mg) was dissolved in 1 mL of 1-propanol/water (9:1(v/v)) at 70°C. The mixture was gradually cooled to 60°C during 20 minutes, and 1 mL of 1-propanol was added thereto. The mixture was again gradually cooled to 5°C during 110 minutes with stirring. Then 3 mL of 1-propanol was added thereto, and the mixture was stirred at 5 °C overnight. The crystals were isolated by filtration in vacuo and dried under air at room temperature to give crystalline Form I (11.1 mg).

(2) 2-Propanol

[0111] Crystalline Form I (100.1 mg) was dissolved in 1 mL of 2-propanol/water (9:1(v/v)) at 90°C. The mixture was gradually cooled to 25°C during 130 minutes, and 1 mL of 2-propanol was added thereto. The mixture was again gradually cooled to 5 °C, and 3 mL of 2-propanol was added. The mixture was stirred at 5°C for 4 days. The crystals were isolated by filtration in vacuo and dried under air at room temperature to give crystalline Form I (48.9 mg).

(3) 1-Butanol

[0112] Crystalline Form I (100.0 mg) was dissolved in 1 mL of 1-butanol/water (9:1(v/v)) at 60°C, and 1 mL of 1-butanol was added thereto. The mixture was gradually cooled to 25°C during 70 minutes, and 3 mL of 1-butanol was added thereto. The mixture was again gradually cooled to 5°C, and stirred at 5°C overnight. The crystals were isolated by filtration in vacuo and dried under air at room temperature to give crystalline Form I (29.0 mg).

(4) 2-Butanol

[0113] Crystalline Form I (100.0 mg) was dissolved in 1 mL of 2-butanol/water (9:1(v/v)) at 90°C. The mixture was gradually cooled to 60°C during 60 minutes, and 1 mL of 2-butanol was added thereto. The mixture was gradually cooled to 25°C during 70 minutes, and 3 mL of 2-butanol was added thereto. The mixture was again gradually cooled to 5 °C,

and stirred at 5°C overnight. The crystals were isolated by filtration in vacuo and dried under air at room temperature to give crystalline Form I (52.1 mg).

(5) Acetone

[0114] Crystalline Form I (100.2 mg) was dissolved in 1.3 mL of acetone/water (9:1(v/v)) under reflux. The mixture was gradually cooled to 25°C during 70 minutes, and 1.3 mL of acetone was added thereto. The mixture was gradually cooled to 5°C, and 3.9 mL of acetone was added thereto. The mixture was stirred for 4 days at 5°C. The crystals were isolated by filtration in vacuo and dried under air at room temperature, to give crystalline Form I (74.5 mg).

(6) 2-Butanone

[0115] Crystalline Form I (100.3 mg) was dissolved in 1 mL of 2-butanone/water (9:1(v/v)) at 60°C. 4 mL of 2-butanone was added thereto. The mixture was gradually cooled to 25°C during 70 minutes, and the mixture was stirred at room temperature for 4 days. The crystals were isolated by filtration in vacuo and dried under air at room temperature to give crystalline Form I (70.0 mg).

Example 11

Crystallization of crystalline Form II

(1) 1-Propanol

[0116] Crystalline Form I (100.0 mg) was dissolved in 1 mL of 1-propanol/water (9:1(v/v)) at 60°C. The mixture was gradually cooled to 5°C during 110 minutes, and was stirred for 4 days at 5°C. The crystals were isolated by filtration in vacuo and dried under air at room temperature to give crystalline Form II (11.0 mg).

(2) 2-Propanol

[0117] Crystalline Form I (100.3 mg) was dissolved in 1 mL of 2-propanol/water (9:1(v/v)) at 90°C. The mixture was gradually cooled to 5°C during 170 minutes, and was stirred for 5 days at 5°C. The crystals were isolated by filtration in vacuo and dried under air at room temperature to give crystalline Form II (40.2 mg).

(3) 1-Butanol

[0118] Crystalline Form I (100.1 mg) was dissolved in 1 mL of 1-butanol/water (9:1(v/v)) at 70°C. The mixture was gradually cooled to 5°C during 130 minutes, and 1 mL of 1-butanol was added thereto. The mixture was stirred for 4 days at 5°C. The crystals were isolated by filtration in vacuo and dried under air at room temperature to give crystalline Form II (31.6 mg).

(4) 2-Butanol

[0119] Crystalline Form I (100.2 mg) was dissolved in 1 mL of 2-butanol/water (9:1(v/v)) at 90°C. The mixture was gradually cooled to 5°C during 170 minutes, and 1 mL of 2-butanol was added thereto. The mixture was stirred for 4 days at 5°C. The crystals were isolated by filtration in vacuo and dried under air at room temperature to give crystalline Form II (54.7 mg).

(5) Acetone

[0120] Crystalline Form I (100.3 mg) was dissolved in 1.2 mL of acetone/water (9:1(v/v)) under reflux. The mixture was gradually cooled to 5°C during 110 minutes, and 1 mL of acetone was added thereto. The mixture was stirred for 4 days at 5°C. The crystals were isolated by filtration in vacuo and dried under air at room temperature to give crystalline Form II (36.4 mg).

(6) 2-Butanone

[0121] Crystalline Form I (100.2 mg) was dissolved in 1 mL of 2-butanone/water (9:1(v/v)) at 60°C. The mixture was cooled to 5°C during 110 minutes. Next, 4 mL of 2-butanone was added thereto. The mixture was stirred for 4 days at

5°C. The crystals were isolated by filtration in vacuo and dried under air at room temperature to give crystalline Form II (76.7 mg).

Example 12

Physical data of Forms I and II

(1) Powder X-ray diffraction (XRD)

[0122] Diffraction patterns were taken at room temperature and humidity using a Rigaku RINT-TTRIII diffractometer with Cu K α radiation. The diffraction angle, 2θ , was scanned from 3 to 40 at rate of 2°/minute at a step size of 0.02°. The results of this analysis are shown in Figures 5 and 6, which are the same as those measured against an internal silicon standard.

(2) Solubility

[0123] Excess amount of the samples, Forms I and II, were suspended in water, and were equilibrated by shaking for 20 minutes at 25 °C or 37 °C, respectively. The amounts of dissolved the compound were determined using a Waters alliance HPLC system 2695, detected by UV 210nm. The results of this analysis are shown in Table 9.

Table 9 Solubility in water of the crystalline Forms I and II

Temperature	Form I	Form II
25 °C	39 mg/mL	30 mg/mL
37 °C	90 mg/mL	54 mg/mL

(3) Hygroscopicity

[0124] The hygroscopicity of Forms I and II were investigated using a Surface measurement systems, DVS-1, between 10 %RH and 90 %RH at 25 °C. There was no hygroscopicity in Form I. On the other hand, there was a hygroscopicity in Form II, and the water value of approximately 4% was increased and decreased between 10 %RH and 90 %RH by absorption and desorption of ca. 1 mol of channel water.

Equivalents

[0125] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific procedures described herein. Such equivalents were considered to be within the scope of this invention and are covered by the following claims. Moreover, any numerical or alphabetical ranges provided herein are intended to include both the upper and lower value of those ranges. In addition, any listing or grouping is intended, at least in one embodiment, to represent a shorthand or convenient manner of listing independent embodiments; as such, each member of the list should be considered a separate embodiment.

Claims

1. A pharmaceutical formulation for use in a method of treatment for improving cognition or treating cognitive loss in a subject in need thereof, wherein the pharmaceutical formulation comprises a therapeutically effective amount of a crystalline Form I of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate, wherein the pharmaceutical formulation is a tablet or capsule; wherein the crystalline Form I of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate is **characterized by** an x-ray powder diffraction pattern having peaks expressed as 2θ at one or both of 17.48 and 20.58 ± 0.20 degrees when measured against an internal silicon standard and **characterized by** an x-ray powder diffraction pattern further having at least four peaks, six peaks, eight peaks or all peaks expressed as 2θ at 4.50, 9.04, 14.60, 15.14, 15.80, 16.60, 18.16, 18.44, 19.48, 21.74, and 25.46 ± 0.20 degrees when measured against an internal silicon standard; and wherein the subject is suffering from a disorder selected from the group consisting of: attention deficit disorder, attention deficit hyperactivity disorder, and Parkinson's Disease.

2. A pharmaceutical formulation for use in a method of treatment for improving cognition or treating cognitive loss in a subject in need thereof, wherein the pharmaceutical formulation comprises a therapeutically effective amount of a crystalline Form II of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate, wherein the pharmaceutical formulation is a tablet or capsule; wherein the crystalline Form II of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate is **characterized by** an x-ray powder diffraction pattern having peaks expressed as 2θ at one or both of 21.16 and 21.38 ± 0.20 degrees when measured against an internal silicon standard and **characterized by** an x-ray powder diffraction pattern further having at least four peaks, six peaks, eight peaks or all peaks expressed as 2θ at 4.48 , 9.00 , 13.58 , 15.62 , 16.48 , 19.02 , 19.44 , 22.46 and 25.00 ± 0.20 degrees when measured against an internal silicon standard; and wherein the subject is suffering from a disorder selected from the group consisting of: attention deficit disorder, attention deficit hyperactivity disorder, and Parkinson's Disease.
3. The pharmaceutical formulation for use of any one of claims 1-2, wherein the pharmaceutical formulation is a tablet formulation.
4. The pharmaceutical formulation for use of any one of claims 1-2, wherein the pharmaceutical formulation is a capsule formulation.
5. The pharmaceutical formulation for use of any one of claims 3-4, wherein the pharmaceutical formulation is a timed release formulation.
6. The pharmaceutical formulation for use of any one of claims 3-4, wherein the pharmaceutical formulation is a sustained release formulation.
7. The pharmaceutical formulation for use of any one of claims 1-6, wherein the subject is suffering from attention deficit disorder.
8. The pharmaceutical formulation for use of any one of claims 1-6, wherein the subject is suffering from attention deficit hyperactivity disorder.
9. The pharmaceutical formulation for use of any one of claims 1-6, wherein the subject is suffering from Parkinson's Disease.
10. A pharmaceutical formulation for oral administration, comprising a therapeutically effective amount of a crystalline Form I of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate and a pharmaceutically acceptable carrier, wherein the pharmaceutical formulation is a tablet or capsule; wherein the crystalline Form I of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate is **characterized by** an x-ray powder diffraction pattern having peaks expressed as 2θ at one or both of 17.48 and 20.58 ± 0.20 degrees when measured against an internal silicon standard and **characterized by** an x-ray powder diffraction pattern further having at least four peaks, six peaks, eight peaks or all peaks expressed as 2θ at 4.50 , 9.04 , 14.60 , 15.14 , 15.80 , 16.60 , 18.16 , 18.44 , 19.48 , 21.74 , and 25.46 ± 0.20 degrees when measured against an internal silicon standard.
11. A pharmaceutical formulation for oral administration comprising a therapeutically effective amount of a crystalline Form II of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate and a pharmaceutically acceptable carrier, wherein the pharmaceutical formulation is a tablet or capsule; wherein the crystalline Form II of (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophene-2-carboxamide hydrochloride monohydrate is **characterized by** an x-ray powder diffraction pattern having peaks expressed as 2θ at one or both of 21.16 and 21.38 ± 0.20 degrees when measured against an internal silicon standard and **characterized by** an x-ray powder diffraction pattern further having at least four peaks, six peaks, eight peaks or all peaks expressed as 2θ at 4.48 , 9.00 , 13.58 , 15.62 , 16.48 , 19.02 , 19.44 , 22.46 and 25.00 ± 0.20 degrees when measured against an internal silicon standard.
12. The pharmaceutical formulation of any one of claims 10-11, wherein the pharmaceutical formulation is a tablet formulation.
13. The pharmaceutical formulation of any one of claims 10-11, wherein the pharmaceutical formulation is a capsule formulation.

14. The pharmaceutical formulation of any one of claims 12-13, wherein the pharmaceutical formulation is a timed release formulation.
15. The pharmaceutical formulation of any one of claims 12-13, wherein the pharmaceutical formulation is a sustained release formulation.
16. The pharmaceutical formulation of any one of claims 10-15 for use in a method of treatment of a subject suffering from Alzheimer's disease, Schizophrenia, or pre-senile dementia (mild cognitive impairment).

Patentansprüche

1. Pharmazeutische Formulierung zur Verwendung in einem Verfahren zur Behandlung zur Verbesserung der Kognition oder Behandlung von kognitivem Verlust bei einem Probanden, der dieser bedarf, wobei die pharmazeutische Formulierung eine therapeutisch wirksame Menge einer kristallinen Form I von (R)-7-Chloro-N-(chinuclidin-3-yl)benzo [b]thiophen-2-carboxamid-Hydrochlorid-Monohydrat umfasst, wobei die pharmazeutische Formulierung eine Tablette oder Kapsel ist; wobei die kristalline Form I von (R)-7-Chloro-N-(chinuclidin-3-yl)benzo[b]thiophen-2-carboxamid-Hydrochlorid-Monohydrat **gekennzeichnet ist durch** ein Röntgenpulverdiffraktogramm mit Peaks, ausgedrückt in 2θ , bei $17,48^\circ$ und/oder $20,58^\circ \pm 0,20^\circ$, wenn gegen einen internen Silicium-Standard gemessen, und **gekennzeichnet ist durch** ein Röntgenpulverdiffraktogramm, das weiterhin mindestens vier Peaks, sechs Peaks, acht Peaks oder alle Peaks, ausgedrückt als 2θ , bei $4,50^\circ$, $9,04^\circ$, $14,60^\circ$, $15,14^\circ$, $15,80^\circ$, $16,60^\circ$, $18,16^\circ$, $18,44^\circ$, $19,48^\circ$, $21,74^\circ$ und $25,46^\circ \pm 0,20^\circ$, wenn gegen einen internen Silicium-Standard gemessen, aufweist; und wobei der Proband an einer Erkrankung leidet, ausgewählt aus der Gruppe umfassend : Aufmerksamkeitsdefizitstörung, Aufmerksamkeitsdefizithyperaktivitätsstörung und Parkinson-Krankheit.
2. Pharmazeutische Formulierung zur Verwendung in einem Verfahren zur Behandlung zur Verbesserung der Kognition oder Behandlung von kognitivem Verlust bei einem Probanden, der dessen bedarf, wobei die pharmazeutische Formulierung eine therapeutisch wirksame Menge einer kristallinen Form II von (R)-7-Chloro-N-(chinuclidin-3-yl)benzo [b]thiophen-2-carboxamid-Hydrochlorid-Monohydrat umfasst, wobei die pharmazeutische Formulierung eine Tablette oder Kapsel ist; wobei die kristalline Form II von (R)-7-Chloro-N-(chinuclidin-3-yl)benzo[b]thiophen-2-carboxamid-Hydrochlorid-Monohydrat **gekennzeichnet ist durch** ein Röntgenpulverdiffraktogramm mit Peaks, ausgedrückt als 2θ , bei $21,16^\circ$ und/oder $21,38^\circ \pm 0,20^\circ$, wenn gegen einen internen Silicium-Standard gemessen, und **gekennzeichnet ist durch** ein Röntgenpulverdiffraktogramm, das weiterhin mindestens vier Peaks, sechs Peaks, acht Peaks oder alle Peaks, ausgedrückt als 2θ , bei $4,48^\circ$, $9,00^\circ$, $13,58^\circ$, $15,62^\circ$, $16,48^\circ$, $19,02^\circ$, $19,44^\circ$, $22,46^\circ$ und $25,00^\circ \pm 0,20^\circ$, wenn gegen einen internen Silicium-Standard gemessen, aufweist; und wobei der Proband an einer Erkrankung leidet, ausgewählt aus der Gruppe umfassend: Aufmerksamkeitsdefizitstörung, Aufmerksamkeitsdefizithyperaktivitätsstörung und Parkinson-Krankheit.
3. Pharmazeutische Formulierung zur Verwendung nach einem der Ansprüche 1-2, wobei die pharmazeutische Formulierung eine Tablettenformulierung ist.
4. Pharmazeutische Formulierung zur Verwendung nach einem der Ansprüche 1-2, wobei die pharmazeutische Formulierung eine Kapselformulierung ist.
5. Pharmazeutische Formulierung zur Verwendung nach einem der Ansprüche 3-4, wobei die pharmazeutische Formulierung eine Formulierung mit zeitlich kontrollierter Freisetzung ist.
6. Pharmazeutische Formulierung zur Verwendung nach einem der Ansprüche 3-4, wobei die pharmazeutische Formulierung eine Retardformulierung ist.
7. Pharmazeutische Formulierung zur Verwendung nach einem der Ansprüche 1-6, wobei der Proband an einer Aufmerksamkeitsdefizitstörung leidet.
8. Pharmazeutische Formulierung zur Verwendung nach einem der Ansprüche 1-6, wobei der Proband an einer Aufmerksamkeitsdefizithyperaktivitätsstörung leidet.
9. Pharmazeutische Formulierung zur Verwendung nach einem der Ansprüche 1-6, wobei der Proband an einer Parkinson-Krankheit leidet.

10. Pharmazeutische Formulierung zur oralen Verabreichung, umfassend eine therapeutisch wirksame Menge einer kristallinen Form I von (R)-7-Chloro-N-(chinuclidin-3-yl)benzo[b]thiophen-2-carboxamid-Hydrochlorid-Monohydrat und einen pharmazeutisch annehmbaren Träger, wobei die pharmazeutische Formulierung eine Tablette oder Kapsel ist; wobei die kristalline Form I von (R)-7-Chloro-N-(chinuclidin-3-yl)benzo[b]thiophen-2-carboxamid-Hydrochlorid-Monohydrat **gekennzeichnet ist durch** ein Röntgenpulverdiffraktogramm mit Peaks, ausgedrückt als 2θ , bei $17,48^\circ$ und/oder $20,58^\circ \pm 0,20^\circ$, wenn gegen einen internen Silicium-Standard gemessen, und **gekennzeichnet ist durch** ein Röntgenpulverdiffraktogramm, das weiterhin mindestens vier Peaks, sechs Peaks, acht Peaks oder alle Peaks, wiedergegeben als 2θ , bei $4,50^\circ$, $9,04^\circ$, $14,60^\circ$, $15,14^\circ$, $15,80^\circ$, $16,60^\circ$, $18,16^\circ$, $18,44^\circ$, $19,48^\circ$, $21,74^\circ$ und $25,46^\circ \pm 0,20^\circ$, wenn gegen einen internen Silicium-Standard gemessen, aufweist.
11. Pharmazeutische Formulierung zur oralen Verabreichung, umfassend eine therapeutisch wirksame Menge einer kristallinen Form II von (R)-7-Chloro-N-(chinuclidin-3-yl)benzo[b]thiophen-2-carboxamid-Hydrochlorid-Monohydrat und einen pharmazeutisch annehmbaren Träger, wobei die pharmazeutische Formulierung eine Tablette oder Kapsel ist; wobei die kristalline Form II von (R)-7-Chloro-N-(chinuclidin-3-yl)benzo[b]thiophen-2-carboxamid-Hydrochlorid-Monohydrat **gekennzeichnet ist durch** ein Röntgenpulverdiffraktogramm mit Peaks, ausgedrückt als 2θ , bei $21,16^\circ$ und/oder $21,38^\circ \pm 0,20^\circ$, wenn gegen einen internen Silicium-Standard gemessen, und **gekennzeichnet ist durch** ein Röntgenpulverdiffraktogramm, das weiterhin mindestens vier Peaks, sechs Peaks, acht Peaks oder alle Peaks, wiedergegeben als 2θ , bei $4,48^\circ$, $9,00^\circ$, $13,58^\circ$, $15,62^\circ$, $16,48^\circ$, $19,02^\circ$, $19,44^\circ$, $22,46^\circ$ und $25,00^\circ \pm 0,20^\circ$, wenn gegen einen internen Silicium-Standard gemessen, aufweist.
12. Pharmazeutische Formulierung nach einem der Ansprüche 10-11, wobei die pharmazeutische Formulierung eine Tablettenformulierung ist.
13. Pharmazeutische Formulierung nach einem der Ansprüche 10-11, wobei die pharmazeutische Formulierung eine Kapselformulierung ist.
14. Pharmazeutische Formulierung nach einem der Ansprüche 12-13, wobei die pharmazeutische Formulierung eine Formulierung mit zeitlich kontrollierter Freisetzung ist.
15. Pharmazeutische Formulierung nach einem der Ansprüche 12-13, wobei die pharmazeutische Formulierung eine Retardformulierung ist.
16. Pharmazeutische Formulierung nach einem der Ansprüche 10-15 zur Verwendung in einem Verfahren zur Behandlung eines Probanden, der an einer Alzheimer-Krankheit, Schizophrenie oder präseniler Demenz (leichter kognitiver Beeinträchtigung) leidet.

Revendications

1. Formulation pharmaceutique pour une utilisation dans une méthode de traitement visant à améliorer la cognition ou à traiter un déclin cognitif chez un sujet qui le requiert, la formulation pharmaceutique comprenant une quantité thérapeutiquement efficace d'une forme cristalline I de l'hydrochlorure de (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophène-2-carboxamide monohydraté, où la formulation pharmaceutique est un comprimé ou une capsule; où la forme cristalline I de l'hydrochlorure de (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophène-2-carboxamide monohydraté est **caractérisée par** un diagramme de diffraction de poudre des rayons X ayant des pics, exprimés en 2θ , à $17,48$ et/ou $20,58 \pm 0,20$ degrés, la mesure étant effectuée par rapport à un étalon interne de silicium, et **caractérisée par** un diagramme de diffraction de poudre des rayons X ayant en outre au moins quatre pics, six pics, huit pics ou tous les pics, exprimés en 2θ , à $4,50$, $9,04$, $14,60$, $15,14$, $15,80$, $16,60$, $18,16$, $18,44$, $19,48$, $21,74$, et $25,46 \pm 0,20$ degrés, la mesure étant effectuée par rapport à un étalon interne de silicium; et où le sujet souffre d'une affection sélectionnée dans le groupe consistant en les suivantes: trouble déficitaire de l'attention, trouble déficitaire de l'attention avec hyperactivité et maladie de Parkinson.
2. Formulation pharmaceutique pour une utilisation dans une méthode de traitement visant à améliorer la cognition ou à traiter un déclin cognitif chez un sujet qui le requiert, la formulation pharmaceutique comprenant une quantité thérapeutiquement efficace d'une forme cristalline II de l'hydrochlorure de (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophène-2-carboxamide monohydraté, où la formulation pharmaceutique est un comprimé ou une capsule; où la forme cristalline II de l'hydrochlorure de (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophène-2-carboxamide monohydraté est **caractérisée par** un diagramme de diffraction de poudre des rayons X ayant des pics, exprimés en 2θ ,

à 21,16 et/ou 21,38 $\pm$ 0,20 degrés, la mesure étant effectuée par rapport à un étalon interne de silicium, et **caractérisée par** un diagramme de diffraction de poudre des rayons X ayant en outre au moins quatre pics, six pics, huit pics ou tous les pics, exprimés en 2 θ , à 4,48, 9,00, 13,58, 15,62, 16,48, 19,02, 19,44, 22,46 et 25,00 $\pm$ 0,20 degrés, la mesure étant effectuée par rapport à un étalon interne de silicium; et où le sujet souffre d'une affection sélectionnée dans le groupe consistant en les suivantes: trouble déficitaire de l'attention, trouble déficitaire de l'attention avec hyperactivité et maladie de Parkinson.

3. Formulation pharmaceutique pour l'utilisation de l'une quelconque des revendications 1-2, où la formulation pharmaceutique est une formulation en comprimé.

4. Formulation pharmaceutique pour l'utilisation de l'une quelconque des revendications 1-2, où la formulation pharmaceutique est une formulation en capsule.

5. Formulation pharmaceutique pour l'utilisation de l'une quelconque des revendications 3-4, où la formulation pharmaceutique est une formulation à libération contrôlée.

6. Formulation pharmaceutique pour l'utilisation de l'une quelconque des revendications 3-4, où la formulation pharmaceutique est une formulation à libération prolongée.

7. Formulation pharmaceutique pour l'utilisation de l'une quelconque des revendications 1-6, où le sujet souffre d'un trouble déficitaire de l'attention.

8. Formulation pharmaceutique pour l'utilisation de l'une quelconque des revendications 1-6, où le sujet souffre d'un trouble déficitaire de l'attention avec hyperactivité.

9. Formulation pharmaceutique pour l'utilisation de l'une quelconque des revendications 1-6, où le sujet souffre de la maladie de Parkinson.

10. Formulation pharmaceutique pour administration orale comprenant une quantité thérapeutiquement efficace d'une forme cristalline I de l'hydrochlorure de (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophène-2-carboxamide monohydraté et un support pharmaceutiquement acceptable, où la formulation pharmaceutique est un comprimé ou une capsule; où la forme cristalline I de l'hydrochlorure de (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophène-2-carboxamide monohydraté est **caractérisée par** un diagramme de diffraction de poudre des rayons X ayant des pics, exprimés en 2 θ , à 17,48 et/ou 20,58 $\pm$ 0,20 degrés, la mesure étant effectuée par rapport à un étalon interne de silicium, et **caractérisée par** un diagramme de diffraction de poudre des rayons X ayant en outre au moins quatre pics, six pics, huit pics ou tous les pics, exprimés en termes de position 2 θ , à 4,50, 9,04, 14,60, 15,14, 15,80, 16,60, 18,16, 18,44, 19,48, 21,74, et 25,46 $\pm$ 0,20 degrés, la mesure étant effectuée par rapport à un étalon interne de silicium.

11. Formulation pharmaceutique pour administration orale comprenant une quantité thérapeutiquement efficace d'une forme cristalline II de l'hydrochlorure de (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophène-2-carboxamide monohydraté et un support pharmaceutiquement acceptable, où la formulation pharmaceutique est un comprimé ou une capsule; où la forme cristalline II de l'hydrochlorure de (R)-7-chloro-N-(quinuclidin-3-yl)benzo[b]thiophène-2-carboxamide monohydraté est **caractérisée par** un diagramme de diffraction de poudre des rayons X ayant des pics, exprimés en 2 θ , à 21,16 et/ou 21,38 $\pm$ 0,20 degrés, la mesure étant effectuée par rapport à un étalon interne de silicium, et **caractérisée par** un diagramme de diffraction de poudre des rayons X ayant en outre au moins quatre pics, six pics, huit pics ou tous les pics, exprimés en 2 θ , à 4,48, 9,00, 13,58, 15,62, 16,48, 19,02, 19,44, 22,46 et 25,00 $\pm$ 0,20 degrés, la mesure étant effectuée par rapport à un étalon interne de silicium.

12. Formulation pharmaceutique pour l'utilisation de l'une quelconque des revendications 10-11, où la formulation pharmaceutique est une formulation en comprimé.

13. Formulation pharmaceutique pour l'utilisation de l'une quelconque des revendications 10-11, où la formulation pharmaceutique est une formulation en capsule.

14. Formulation pharmaceutique pour l'utilisation de l'une quelconque des revendications 12-13, où la formulation pharmaceutique est une formulation à libération contrôlée.

EP 3 272 748 B1

15. Formulation pharmaceutique pour l'utilisation de l'une quelconque des revendications 12-13, où la formulation pharmaceutique est une formulation à libération prolongée.
16. Formulation pharmaceutique de l'une quelconque des revendications 10-15 pour une utilisation dans une méthode de traitement d'un sujet souffrant de la maladie d'Alzheimer, de schizophrénie ou de démence présénile (déficit cognitif léger).

5

10

15

20

25

30

35

40

45

50

55

FIGURE 1




FIGURE 2





Figure 3

FIGURE 4




FIGURE 5 (Crystalline Form I)





FIGURE 6 (Crystalline Form II)



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 20050119325 A [0026] [0086]
- WO 0355878 A [0071]

Non-patent literature cited in the description

- Remington: The Science and Practice of Pharmacy. Lippincott, Williams and Wilkins, 2005 [0011]
- The United States Pharmacopeia. 1995, 1843-1844 [0011]
- *Fluid Phase Equilibria*, 2000, vol. 172, 43-72 [0070]