



(11) **EP 2 139 494 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
11.03.2020 Bulletin 2020/11

(51) Int Cl.:
A61K 31/70 (2006.01) **A61K 9/16** (2006.01)
A61K 9/20 (2006.01) **A61P 3/10** (2006.01)

(21) Application number: **08732695.5**

(86) International application number:
PCT/US2008/057888

(22) Date of filing: **21.03.2008**

(87) International publication number:
WO 2008/116179 (25.09.2008 Gazette 2008/39)

(54) **PHARMACEUTICAL FORMULATIONS CONTAINING DAPAGLIFLOZIN PROPYLENE GLYCOL HYDRATE**

PHARMAZEUTISCHE FORMULIERUNGEN MIT DAPAGLIFLOZIN-PROPYLENGLYCOL-HYDRAT
PRÉPARATIONS PHARMACEUTIQUES CONTENANT DE L'HYDRATE DE PROPYLÈNEGLYCOL DE DAPAGLIFLOZINE

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT RO SE SI SK TR
Designated Extension States:
AL BA MK RS

- **PATEL, Jatin, M.**
New Brunswick, NJ 08903 (US)
- **TAO, Li**
New Brunswick, NJ 08903 (US)
- **TEJWANI, Ravindra, W.**
New Brunswick, NJ 08903 (US)
- **VATSARAJ, Nipa**
New Brunswick, NJ 08903 (US)
- **WU, Yongmei**
New Brunswick, NJ 08903 (US)

(30) Priority: **22.03.2007 US 896286 P**

(43) Date of publication of application:
06.01.2010 Bulletin 2010/01

(60) Divisional application:
12153939.9 / 2 508 188

(73) Proprietor: **AstraZeneca AB**
151 85 Södertälje (SE)

(74) Representative: **AstraZeneca**
Milstein Building
Granta Park
Cambridge CB21 6GH (GB)

- (72) Inventors:
- **BINDRA, Dilbir, S.**
New Brunswick, NJ 08903 (US)
 - **DALI, Mandar, V.**
New Brunswick, NJ 08903 (US)
 - **PARAB, Prakash, V.**
New Brunswick, NJ 08903 (US)

(56) References cited:
WO-A-03/099836 WO-A-2008/002824

- **MENG WEI ET AL: "Discovery of dapagliflozin: A potent, selective renal sodium-dependent glucose cotransporter 2 (SGLT2) inhibitor for the treatment of type 2 diabetes" JOURNAL OF MEDICINAL CHEMISTRY, vol. 51, no. 5, March 2008 (2008-03), pages 1145-1149, XP002491733 ISSN: 0022-2623**

EP 2 139 494 B1

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description


FIELD OF THE INVENTION

5 [0001] The present invention provides an immediate release pharmaceutical formulation in the form of a stock granulation for loading in capsules or forming tablets, or in the form of a capsule or a tablet containing the stock granulation, containing the sodium dependent glucose transporter (SGLT2) inhibitor dapagliflozin (S)-propylene glycol hydrate.

BACKGROUND OF THE INVENTION


10 [0002] At least 171 million people worldwide suffer from type II diabetes (NIDDM), which is characterized by hyperglycemia due to excessive hepatic glucose production and peripheral insulin resistance. Hyperglycemia is considered to be the major risk factor for the development of diabetic complications, and is likely to contribute directly to the impairment of insulin secretion seen in advanced NIDDM. Thus, consistent control of plasma glucose levels in NIDDM patients can offset the development of diabetic complications and beta cell failure seen in advanced disease. Plasma glucose is normally filtered in the kidney in the glomerulus and actively reabsorbed in the proximal tubule. SGLT2 appears to be the major transporter responsible for the reuptake of glucose at this site. A selective inhibitor of the sodium-dependent glucose transporter SGLT2 in the kidney is expected to normalize plasma glucose levels by enhancing the excretion of glucose in the urine, thereby improving insulin sensitivity, and delaying the development of diabetic complications.

20 [0003] The compound of the structure (I)



35 or pharmaceutically acceptable salts or solvates thereof (hereinafter dapagliflozin), an orally active SGLT2 inhibitor is disclosed in U.S. Patent No. 6,515,117.


[0004] U.S. Application Serial No. 11/765,481 filed June 28, 2007, published as US 2008-0004336 A1 discloses dapagliflozin in the form of its (S)-propylene glycol ((S)-PG) hydrate and its (R)-propylene glycol ((R)-PG) hydrate. The (S)-propylene glycol hydrate is referred to as form SC-3 and has the structure shown as (Ia).



55 [0005] The (R)-propylene glycol hydrate is referred to as form SD-3 and has the structure shown as (Ib).

5

10



Ib (SD-3)

15

[0006] Methods for preparing the (S)-PG hydrate and the (R)-PG hydrate of dapagliflozin are provided in U.S. Application Serial No. 11/765,481 filed June 28, 2007, published as US 2008-0004336 A1.


BRIEF DESCRIPTION OF THE INVENTION

20

[0007] In accordance with the present disclosure, pharmaceutical formulations are provided which can be in the form of a capsule formulation or a tablet formulation, for oral use, designed for immediate release, and include as the medicament dapagliflozin which has the structure (I)

25

30



I


35

or a pharmaceutically acceptable salt, solvate, mixed solvate, or complex thereof (which is disclosed in U.S. Patent No. 6,515,117) and a pharmaceutically acceptable carrier thereof.

[0008] The preparation of dapagliflozin in the form of its (S)-propylene glycol ((S)-PG) hydrate (SC-3) which is shown as Compound (Ia)

40

45



Compound Ia

50


is disclosed in U.S. Application Serial No. 11/765,481, filed June 28, 2007, U.S. Publication No. 2008-0004336 A1, and Provisional Application No. 60/817,118, filed June 28, 2006.

55

[0009] The preparation of dapagliflozin in the form of its (R)-propylene glycol ((R)-PG) hydrate (SD-3), which is shown as Compound (Ib)

5

10



Compound Ib

15

is disclosed in U.S. Application Serial No. 11/765,481, filed June 28, 2007, U.S. Publication No. 2008-0004336 A1, and Provisional Application No. 60/817,118, filed June 28, 2006.

[0010] Described herein are immediate release pharmaceutical formulations in the form of a stock granulation (e.g., granules, beads, and/or beadlets), for loading in capsules or forming into tablets, comprising

20

- a) dapagliflozin or dapagliflozin propylene glycol hydrate;
- b) one or more bulking agents;
- c) optionally one or more binders;
- d) optionally one or more disintegrants;
- e) optionally one or more glidants and/or anti-adherents; and
- f) optionally one or more lubricants.

25


[0011] According to the invention there is provided an immediate release pharmaceutical formulation in the form of a stock granulation for loading in capsules or forming tablets, or in the form of a capsule or a tablet containing the stock granulation, the stock granulation comprising:

30

- a) dapagliflozin propylene glycol hydrate having the formula

35

40



45

- b) one or more bulking agents;
- c) one or more binders;
- d) one or more disintegrants;
- e) optionally one or more glidants and/or anti-adherents; and
- f) optionally one or more lubricants.

50

[0012] The stock granulation comprises dapagliflozin (S) propylene glycol hydrate and one or more bulking agents. Suitable bulking agents include, for example, microcrystalline cellulose and/or lactose, as well as others provided herein and known in the art. The stock granulation also comprises one or more of the following compounds: (1) one or more binders; (2) one or more disintegrants; (3) optionally one or more glidants and/or anti-adherents; and (4) optionally one or more lubricants. Suitable binders include, for example, pregelatinized starch, as well as others provided herein and known in the art. Suitable disintegrants include, for example, sodium starch glycolate, croscovidone, and croscamellose sodium, as well as others provided herein and known in the art. Suitable glidants and/or anti-adherents include, for example, silicon dioxide and talc, as well as others provided herein and known in the art. Suitable lubricants include, for example, magnesium stearate, as well as others provided herein and known in the art.

55

[0013] The stock granulation described above, and capsules and tablets containing same, is prepared by mixing together dapagliflozin propylene glycol and one or more bulking agents, in any desired order, to form the stock granulation; and filling the capsules with or forming tablets from desired quantities of the stock granulation. Alternatively, the stock granulation described is prepared by mixing together dapagliflozin propylene glycol hydrate and one or more bulking agents; and optionally one or more of the following compounds: binder(s); disintegrant(s); glidant(s) and/or anti-adherent(s); and lubricant(s) in any desired order, to form the stock granulation and filling the capsules with or forming tablets from desired quantities of stock granulation.

[0014] The tablets described above are prepared by compressing the stock granulation into tablet form. The described tablets may be prepared by compressing the stock granulation having one or more binder(s). The tablets described may be prepared by compressing the stock granulation containing one or more anti-adherent(s) and/or glidant(s). The tablets described may be prepared by compressing the stock granulation comprising one or more of the following compounds: (1) one or more binders; (2) one or more disintegrants; (3) one or more glidants and/or anti-adherents; and (4) one or more lubricants.

[0015] Optionally, the tablets and/or capsules of the invention can include an outer protective coating which comprises a coating polymer, such as, for example, polyvinyl alcohol (PVA), hydroxypropyl methyl cellulose, and hydroxypropyl cellulose, and/or a plasticizer(s) and optional colorant(s). Other optional components of the outer protective coating include anti-adherent(s) and/or glidant(s) and opacifying agent(s).

[0016] The pharmaceutical dapagliflozin propylene glycol hydrate formulations of the invention including the stock granulation, capsules containing same, and tablets of the invention are useful in the treatment of mammals, such as humans, dogs, and cats, for diseases or disorders associated with SGLT2 activity. Thus, the invention provides pharmaceutical dapagliflozin propylene glycol hydrate formulations for use in the treatment of diseases or disorders associated with SGLT2 activity, for example, Type I and Type II diabetes; impaired glucose tolerance; insulin resistance; and diabetic complications, such as nephropathy, retinopathy, neuropathy and cataracts; hyperglycemia; hyperinsulinemia; hypercholesterolemia; dyslipidemia; elevated blood levels of free fatty acids or glycerol; hyperlipidemia; hypertriglyceridemia; obesity; wound healing; tissue ischemia; atherosclerosis; hypertension; and Syndrome X or Metabolic Syndrome.

[0017] In one embodiment, the invention provides the pharmaceutical formulation of the invention for use in the treatment of type II diabetes. In another embodiment, the invention provides the pharmaceutical formulation of the invention for use in delaying the progression or onset of type II diabetes.

[0018] Also described is a method for treating or delaying the progression or onset of diseases or disorders associated with SGLT2 activity comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of a pharmaceutical formulation described herein. Also described is a method for treating type II diabetes comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of a pharmaceutical formulation as described herein. Also described is a method for delaying the progression or onset of type II diabetes comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of a pharmaceutical formulation described herein.

[0019] Other therapeutic agent(s) suitable for combination with the formulations of the present invention include known therapeutic agents useful in the treatment of the aforementioned disorders associated with SGLT2 activity including: anti-diabetic agents; anti-hyperglycemic agents; hypolipidemic or lipid lowering agents; anti-obesity agents; anti-hypertensive agents and appetite suppressants.

[0020] Also described is a method for treating or delaying the progression or onset of diseases or disorders associated with SGLT2 activity comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of a pharmaceutical formulation described herein and one or more of the following: an anti-diabetic agent(s), anti-hyperglycemic agent(s); hypolipidemic or lipid lowering agent(s); anti-obesity agent(s); anti-hypertensive agent(s) and appetite suppressant(s).

[0021] Also described is a method for treating type II diabetes comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of a pharmaceutical formulation described herein and one or more anti-diabetic agent(s). Also described is a method for delaying the progression or onset of type II diabetes comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of a pharmaceutical formulation described herein and one or more anti-diabetic agent(s). Also described is a method for treating or delaying the progression or onset of type II diabetes comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of a pharmaceutical formulation described herein and one or more of the following: an anti-hyperglycemic agent(s); hypolipidemic or lipid lowering agent(s); anti-obesity agent(s); anti-hypertensive agent(s) and appetite suppressant(s).

DETAILED DESCRIPTION OF THE INVENTION

[0022] The invention provides immediate release pharmaceutical formulations, which include, among others, tablet and capsule formulations, containing the sodium dependent glucose transporter (SGLT2) inhibitor dapagliflozin (S)-pro-

propylene glycol hydrate.

[0023] As used herein, the term "dapagliflozin" is intended to mean the structure shown as structure I or Compound I. The term "dapagliflozin propylene glycol hydrate" is meant to refer to and encompass both dapagliflozin (S)-propylene glycol hydrate (structure Ia or Compound Ia) and dapagliflozin (R)-propylene glycol hydrate (structure Ib or Compound Ib). For the avoidance of doubt, the claims relate to dapagliflozin (S)-propylene glycol hydrate (structure Ia or Compound Ia) As used herein, the terms "pharmaceutical formulation", "pharmaceutical formulation of the invention", and "formulation" are meant to refer to formulations containing dapagliflozin (S)-propylene glycol hydrate. Likewise, the term "medicament" is meant in the present application to refer to dapagliflozin (S)-propylene glycol hydrate.

[0024] As used herein, the terms "immediate release" and "immediate release pharmaceutical formulation" are intended to mean that the pharmaceutical formulations of the invention are not produced using excipients that interfere with absorption of the active pharmaceutical ingredient, for example, dapagliflozin or dapagliflozin propylene glycol hydrate, when administered to a mammal or human.

[0025] The pharmaceutical formulation of the invention can be in the form of a capsule, tablet, bead, beadlet, granule or pill, all of the above being collectively referred to as pharmaceutical formulations, and contains medicament, namely dapagliflozin (S)-propylene glycol hydrate. In one embodiment, the medicament is dapagliflozin (S)-propylene glycol hydrate. In one embodiment of the invention, the immediate release pharmaceutical formulation of the invention is in the form of a stock granulation (e.g., granules, beads, and/or beadlets), for loading in capsules or forming into tablets.

[0026] As described herein the dapagliflozin or dapagliflozin propylene glycol hydrate is in an amount within the range of from about 0.1% to about 70% by weight of the stock granulation and preferably in an amount within the range of from about 0.1% to about 30% by weight of the stock granulation.

[0027] The pharmaceutical formulation as described herein can include pharmaceutical excipients as described herein to aid in the formation of a stock granulation suitable in the form of granules, beads, or beadlets for capsule loading and for tablets of the invention. In one embodiment, the pharmaceutical formulation is in the form of a capsule or tablet containing a stock granulation comprising

- a) dapagliflozin (S)-propylene glycol hydrate;
- b) at least one bulking agent or filler;
- c) at least one binder;
- d) at least one disintegrant;
- e) optionally at least one glidant and/or anti-adherent; and
- f) optionally at least one lubricant.

[0028] In the described embodiments of the pharmaceutical formulations of the invention, the amounts of medicament (dapagliflozin (S)-propylene glycol hydrate) and each of the excipient(s) are expressed as a percentage weight of the total weight of the stock granulation, which is equivalent in measurement to the percentage weight of the total weight of the tablet or capsule fill.

[0029] In one embodiment, the dapagliflozin (S)-propylene glycol hydrate is in an amount within the range of from about 0.1% to about 70% by weight of the stock granulation. The dapagliflozin propylene glycol hydrate may be in an amount within the range of from about 0.1% to about 30% by weight of the stock granulation.

[0030] In one embodiment, the bulking agent or filler is present in an amount within the range of from about 1% to about 95% by weight of the stock granulation. In another embodiment, the bulking agent or filler is present in an amount within the range of from about 10% to about 85% by weight of the stock granulation.

[0031] In one embodiment, the binder is present in an amount within the range of up to about 20% by weight of the stock granulation. In another embodiment, the binder is present in an amount within the range of from about 1% to about 10% by weight of the stock granulation. In another embodiment, the binder is present in an amount within the range of from about 2% to about 4% by weight of the stock granulation.

[0032] In one embodiment, the disintegrant is present in an amount within the range of up to about 20% by weight of the stock granulation. In another embodiment, the disintegrant is present in an amount within the range of from about 0.25% to about 10% by weight of the stock granulation.

[0033] In one embodiment, the glidant and/or anti-adherent, if present, is present in an amount within the range of from about 0% to about 20% by weight of the stock granulation. In another embodiment, the glidant and/or anti-adherent, if present, is present in an amount within the range of from about 1% to about 15% by weight of the stock granulation.

[0034] In one embodiment, the lubricant, if present, is present in an amount within the range of from about 0% to about 5% by weight of the stock granulation. In another embodiment, the lubricant, if present, is present in an amount within the range of from about 0.1% to about 5% by weight of the stock granulation. In another embodiment, the lubricant, if present, is present in an amount within the range of from about 0.2% to about 2% by weight of the stock granulation.

[0035] In one embodiment, the pharmaceutical formulation is in the form of a capsule or tablet containing a stock granulation comprising

- a) dapagliflozin (S)-propylene glycol hydrate;
- b) at least one bulking agent or filler;
- c) at least one binder;
- d) at least one disintegrant;
- e) optionally at least one glidant and/or anti-adherent; and
- f) optionally at least one lubricant.

wherein

- a) the dapagliflozin or dapagliflozin propylene glycol hydrate is present in an amount within the range of from about 0.1% to about 70% by weight;
- b) the bulking agent or filler is present in an amount within the range of from about 1% to about 95% by weight;
- c) the binder is present in an amount of up to 20% by weight;
- d) the disintegrant is present in an amount of up to 20% by weight;
- e) the glidant and/or anti-adherent, if present, is present in an amount within the range of from 0% to 20% by weight; and
- f) the lubricant, if present, is present in an amount within the range of from 0% to 5% by weight, all of the above % by weight being based on the weight of the stock granulation.

[0036] In one embodiment, the pharmaceutical formulation is in the form of a capsule or tablet containing a stock granulation comprising

- a) dapagliflozin (S)-propylene glycol hydrate;
- b) at least one bulking agent or filler;
- c) at least one binder;
- d) at least one disintegrant;
- e) optionally at least one glidant and/or anti-adherent; and
- f) optionally at least one lubricant.

wherein

- a) the dapagliflozin or dapagliflozin propylene glycol hydrate is present in an amount within the range of from about 0.1% to about 30% by weight
- b) the bulking agent or filler is present in an amount within the range of from about 10% to about 85% by weight;
- c) the binder, if present, is present in an amount within the range of from about 1% to about 10% by weight;
- d) the disintegrant, if present, is in an amount within the range of from about 0.25% to about 10% by weight;
- e) the glidant and/or anti-adherent, if present, is in an amount within the range of from about 1% to about 15% by weight; and
- f) the lubricant, if present, is in an amount within the range of from about 0.2% to about 2% by weight, all of the above % by weight being based on the weight of the stock granulation.

[0037] In one embodiment, the medicament in the pharmaceutical formulations has 90% of the particles smaller than 200 micrometers. In another embodiment, the medicament has 90% of its particles smaller than 100 micrometers. In another embodiment, the medicament has 90% of its particles smaller than 50 micrometers. Dapagliflozin or dapagliflozin propylene glycol hydrate can be milled or micronized as needed to obtain the above mentioned characteristics.

[0038] Examples of bulking agents or fillers suitable for use herein include, but are not limited to, cellulose derivatives, such as microcrystalline cellulose or wood cellulose, lactose, sucrose, starch, pregelatinized starch, dextrose, mannitol, fructose, xylitol, sorbitol, corn starch, modified corn starch, inorganic salts such as calcium carbonate, calcium phosphate, dicalcium phosphate, calcium sulfate, dextrin/dextrates, maltodextrin, compressible sugars, and other known bulking agents or fillers, and/or mixtures of two or more thereof. Several types of microcrystalline cellulose are suitable for use in the formulations described herein, for example, microcrystalline cellulose selected from the group consisting of Avicel® types: PH101, PH102, PH103, PH105, PH 112, PH113, PH200, PH301, and other types of microcrystalline cellulose, such as silicified microcrystalline cellulose. Several types of lactose are suitable for use in the formulations described herein, for example, lactose selected from the group consisting of anhydrous lactose, lactose monohydrate, lactose fast flo, directly compressible anhydrous lactose, and modified lactose monohydrate. In one embodiment of the invention, the bulking agent of the stock granulation is microcrystalline cellulose and/or lactose. Lactose is particularly useful for tablet formulation.

[0039] Examples of binders suitable for use herein include, but are not limited to, hydroxypropyl cellulose, corn starch,

pregelatinized starch, modified corn starch, polyvinyl pyrrolidone (PVP) (typical molecular weight ranging from about 5,000 to about 1,000,000, preferably about 40,000 to 50,000), hydroxypropyl methylcellulose (HPMC), lactose, gum acacia, ethyl cellulose, cellulose acetate, as well as a wax binder such as carnauba wax, paraffin, spermaceti, polyethylenes or microcrystalline wax, as well as other conventional binding agents and/or mixtures of two or more thereof. In one embodiment of the invention, the binding agent, if present, of the stock granulation is pregelatinized starch.

[0040] Examples of disintegrants suitable for use herein include, but are not limited to, croscarmellose sodium, crospovidone, starch, potato starch, pregelatinized starch, corn starch, sodium starch glycolate, microcrystalline cellulose, low substituted hydroxypropyl cellulose and other known disintegrants. Several specific types of disintegrant are suitable for use in the formulations described herein. For example, any grade of crospovidone can be used, including for example crospovidone XL-10, and includes members selected from the group consisting of Kollidon CL®, Polyplasdone XL®, Kollidon CL-M®, Polyplasdone XL-10®, and Polyplasdone INF-10®. In one embodiment, the disintegrant, if present, of the stock granulation is sodium starch glycolate, croscarmellose sodium and/or crospovidone. In one embodiment, the disintegrant is sodium starch glycolate. In another embodiment, the disintegrant is croscarmellose sodium and/or crospovidone, which are particularly useful for tablet formulation. In one specific embodiment, the disintegrant is crospovidone XL-10 with peroxide levels below 400 parts per million (ppm). These materials are also referred to as insoluble polyvidone, insoluble PVP, crosslinked PVP, and PVPP. The crospovidone can be substituted with croscarmellose sodium, sodium starch glycolate, or pregelatinized starch (at, for example, a 5-10% concentration).

[0041] Examples of lubricants suitable for use herein include, but are not limited to, magnesium stearate, zinc stearate, calcium stearate, talc, carnauba wax, stearic acid, palmitic acid, sodium stearyl fumarate sodium laurel sulfate, glyceryl palmitostearate, palmitic acid, myristic acid and hydrogenated vegetable oils and fats, as well as other known lubricants, and/or mixtures of two or more thereof. In one embodiment, the lubricant, if present, of the stock granulation is magnesium stearate.

[0042] Examples of glidants and/or anti-adherents suitable for use herein include but are not limited to, silicon dioxide (generally), colloidal silicon dioxide, magnesium silicate, magnesium trisilicate, talc, and other forms of silicon dioxide, such as aggregated silicates and hydrated silica.

[0043] In one embodiment of the stock granulation, the bulking agent is microcrystalline cellulose and/or lactose monohydrate, the binder, if present, is pregelatinized starch, the disintegrant, if present, is sodium starch glycolate, croscarmellose sodium and/or crospovidone, the lubricant, if present, is magnesium stearate and the glidant and/or anti-adherent, if present, is silicon dioxide and/or talc.

[0044] In one embodiment, the tablet or capsule has a protective outer layer. The protective outer layer of the tablet or capsule, where present, can include from about 10% to about 95% of polymer based on the weight of the coating layer, and can be prepared employing conventional procedures. In one embodiment, the outer layer of the tablet or capsule includes from about 20% to about 90% of polymer based on the weight of the coating layer. The formulation can contain at least one coating layer polymer and a coating solvent, for example, water, which is used for processing and removed by drying. Suitable examples of polymer for the coating layer include, but are not limited to, hydroxypropyl methylcellulose, polyvinyl alcohol (PVA), ethyl cellulose, methacrylic polymers, hydroxypropyl cellulose, and starch. In one embodiment, the coating layer polymer is PVA. In another embodiment, the coating layer polymer is hydroxypropyl cellulose. Use of PVA allows for enhanced logo definition, film adhesion, and facilitates faster coating of the drug, the latter of which can be important for dapagliflozin formulations due to the temperature sensitivity of the compound.

[0045] The coating can also optionally include a plasticizer of from about 0% to about 30% by weight, based on the weight of the coating layer. In one embodiment, the plasticizer is from about 15% to about 25% by weight of the coating layer. Suitable plasticizers include, but are not limited to, triacetin, diethyl phthalate, tributyl sebacate, polyethylene glycol (PEG), glycerin, triacetin, and triaethyl citrate, for example. In one embodiment, the plasticizer is polyethylene glycol of molecular weight 200 to 20,000. In another embodiment, the plasticizer is polyethylene glycol of molecular weight 400 to 4,000. In another embodiment, the plasticizer is polyethylene glycol of molecular weight 400.

[0046] In another embodiment, the coating can also optionally include an anti-adherent or glidant such as talc, fumed silica, or magnesium stearate, for example. In another embodiment, the coating can also optionally include an opacifying agent, such as titanium dioxide, for example. In yet another embodiment, the coating layer can also optionally include one or more colorants, for example, iron oxide based colorant(s). Examples of commercially available coating material include Opadry® HP and Opadry® II white.

[0047] The pharmaceutical formulations disclosed herein can further comprise antioxidants and chelating agents. For example, the pharmaceutical formulations can comprise butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), propyl gallate (PG), sodium metabisulfite, ascorbyl palmitate, potassium metabisulfite, disodium EDTA (ethylenediamine tetraacetic acid; also known as disodium edentate), EDTA, tartaric acid, citric acid, citric acid monohydrate, and sodium sulfite. In one embodiment, the foregoing compounds are included in the pharmaceutical formulations in amounts in the range of about 0.01% to about 5% w/w. In one specific embodiment, the pharmaceutical formulation includes BHA, BHT, or PG used at a range of about 0.02% to about 1% and disodium EDTA, citric acid, or citric acid monohydrate used at a range of about 2% to about 5%. In a preferred embodiment, the pharmaceutical formulation

EP 2 139 494 B1

includes BHA used at about 0.05% w/w.

[0048] The pharmaceutical formulations described above can be prepared by mixing together dapagliflozin propylene glycol hydrate and one or more of the desired excipients described herein in any desired order, to form the stock granulation; and filling the capsules with or forming tablets from desired quantities of the stock granulation. The stock granulation, capsules and tablets of the invention can be prepared by a variety of processes and order of addition of excipients. The utility of these formulations is not limited to a specific dosage form or manufacturing process. For example, stock formulation tablets can be manufactured by wet granulation, dry granulation, direct blending or any other pharmaceutically acceptable process described herein or otherwise known in the art.

[0049] The pharmaceutical formulations of the invention can be packaged in any packaging that facilitates stability of the drug formulation. For example, sealed high density polyethylene (HDPE) bottles containing silica gel desiccant or aluminum blister lined with PVC can be used. Use of such packaging helps to control unwanted oxidation of the product at room temperature.

[0050] Examples of certain specific embodiments of tablet and capsule formulations in accordance with the invention are set out below.

Table I. Tablet and Capsule Formulations

Material	Possible Range % by weight of tablet or capsule fill	Preferred Range % by weight of tablet or capsule fill
Dapagliflozin or Dapagliflozin propylene glycol hydrate	0.1 to 70%	0.1 to 30%
Bulking Agent/binder	1 to 95%	10 to 85%
Anhydrous Lactose	0 to 95%	20 to 75%
Microcrystalline cellulose	0 to 95%	20 to 75%
Pregelatinized starch	0 to 95%	10 to 75%
Disintegrant	0 to 20%	0.25 to 10%
Croscarmellose sodium	0 to 20%	2 to 10%
Crospovidone	0 to 12%	4 to 10%
Sodium Starch glycolate	0 to 20%	2 to 10%
Lubricant	0.1 to 5%	0.2 to 2%
Magnesium Stearate	0.1 to 5%	0.2 to 2%
Anti adherent/glidant Talc, silicon dioxide	0 to 10%	1 to 10% more preferably 1 to 4%
Outer Protective Coating Layer	% by weight of tablet or capsule fill	% by weight of tablet or capsule fill
Coating polymer, and optional plasticizer(s), glidant (s), anti-tacking agent(s), and colorant(s)	0.5 to 50%	1 to 5%

Table II. Granulation Composition (% w/w) for Tablets and Capsules

Ingredient	Possible Range % by weight	Preferred range % by weight
Dapagliflozin or Dapagliflozin Propylene Glycol Hydrate	0.1-40	0.1-10
Microcrystalline Cellulose	q.s.	q.s.
Anhydrous Lactose	0-50	10-30
Crospovidone	1-15	3-10
Silicon Dioxide	0-6	0.5-4

(continued)

Ingredient	Possible Range % by weight	Preferred range % by weight
Magnesium Stearate	0.0-4.0	0.5-2.0
q.s. refers to the quantity sufficient to make the granulation composition 100% w/w.		

[0051] A film coating for capsules or tablets of Table II comprises, for example, polyvinyl alcohol (PVA), titanium dioxide, polyethylene glycol, talc, and colorant.

[0052] Tablets or capsules of various strengths (0.1 - 50mg) can be prepared using different weights of the stock granulations described herein.

[0053] The pharmaceutical formulation in the form of a tablet can be obtained by a process comprising the steps of:

- a) mixing the inactive ingredients with the medicament (Dapagliflozin or Dapagliflozin propylene glycol hydrate) using a combination of blending and milling processes;
- b) formulating granules;
- c) drying and/or screening the granules;
- d) blending the granules; and
- e) tableting the blend obtained in (d) into tablets.


[0054] In one embodiment, step a) of the process employs impact milling and/or sizing equipment. In one embodiment, the granules in step b) of the process are formulated by dry granulation, wet granulation, or direct compression. In one embodiment, the granules are formulated by dry granulation. In one embodiment, the granules in step d) of the process are blended with a tableting aid or a lubricant and filler.

[0055] The pharmaceutical formulation in the form of a capsule can be obtained by a process comprising the steps of:

- a) mixing the inactive ingredients with the medicament using a combination of blending and milling processes;
- b) formulating granules;
- c) drying and/or screening the granules; and
- d) loading the granules into capsules.

[0056] In one embodiment, step a) of the process employs impact milling and/or sizing equipment. In one embodiment, the granules in step b) of the process are formulated by dry granulation, wet granulation, or direct compression. In one embodiment, the granules are formulated by dry granulation.

[0057] The dapagliflozin propylene glycol hydrate ((S) form and (R) form) can be prepared, for example, by a process as described in U.S. Application Serial No: 11/765,481, filed June 28, 2007, U.S. Publication No. 2008-0004336 A1, and provisional application No. 60/817,118 filed June 28, 2006.




Compound 1a

For example, dapagliflozin (S)-propylene glycol hydrate (Compound 1a) can be prepared by the following steps: providing a compound A (prepared as described in U.S. application Serial No. 10/745,075 filed December 23, 2003, Examples 17 to 20), of the structure (A);

5

10




Compound A

15 treating compound A with an alcohol solvent, such as methanol or ethanol, and aqueous base, such as sodium hydroxide, and water, if necessary, under an inert atmosphere, and elevated temperature, if necessary; adding an acid, such as hydrochloric acid to neutralize the reaction mixture, to form compound I of the structure;

20

25



30


Compound I

35 treating the reaction mixture containing compound I with an organic solvent, such as methyl t-butyl ether, an alkyl acetate, such as ethyl acetate, methyl acetate, isopropyl acetate, or butyl acetate, and (S)-propylene glycol; optionally adding seeds of (S)-propylene glycol compound Ia (SC-3) to the mixture, to form dapagliflozin (S)-propylene glycol compound Ia (SC-3 form).

[0058] In another example, dapagliflozin propylene glycol hydrate can be prepared in a process comprising the step of reducing a compound of the structure (B)

40


45



B

50 to remove the methoxy group; treating compound B with a reducing agent, such as triethylsilyl hydride and an activating group which is a Lewis acid, such as $\text{BF}_3 \cdot \text{Et}_2\text{O}$, and an organic solvent, such as CH_3CN , and water; separating out the compound of the structure (I);

55




and treating compound I with (S)-propylene glycol in the presence of a solvent, such as t-butylmethyl ether, and optionally with seeds of compound Ia (dapagliflozin (S)-propylene glycol), to form a crystal slurry of compound Ia (dapagliflozin (S)-propylene glycol) and separating out compound Ia (dapagliflozin (S)-propylene glycol).

15 **[0059]** The above process of the invention is a one-pot operation which minimizes the production of intermediates, resulting in improved yield and priority of the final crystalline compound Ia dapagliflozin(S)-propylene glycol.

[0060] In carrying out the formation of compound Ia, the (S)-propylene glycol is employed in a molar ratio to compound I with the range of from about 0.9:1 to about 1.5:1. In one embodiment, the (S)-propylene glycol is employed in a molar ratio to compound I with the range of from about 0.98:1 to about 1.2:1.


20 **[0061]** Dapagliflozin (R)-propylene glycol hydrate (Compound Ib) can be prepared by the following steps: providing a compound A (prepared as described in U.S. application Serial No. 10/745,075 filed December 23, 2003, Examples 17 to 20), of the structure (A);



35

treating compound A with an alcohol solvent, such as methanol or ethanol, and aqueous base, such as sodium hydroxide, and water, if necessary, under an inert atmosphere, and elevated temperature, if necessary; adding an acid, such as hydrochloric acid to neutralize the reaction mixture, to form compound I of the structure;


40



55 treating the reaction mixture containing compound I with an organic solvent, such as methyl t-butyl ether, an alkyl acetate, such as ethyl acetate, methyl acetate, isopropyl acetate, or butyl acetate, and (R)-propylene glycol; optionally adding seeds of (R)-propylene glycol Compound Ib (SD-3) to the mixture, to form dapagliflozin (R)-propylene glycol Compound Ib (SD-3 form).

5

10



Compound Ib (SD-3)

[0062] The activity of dapagliflozin or dapagliflozin propylene glycol hydrate can be determined using, for example, the assay system described below or any appropriate assay system known in the art.

[0063] The mRNA sequence for human SGLT2 (GenBank #M95549) is cloned by reverse-transcription and amplification from human kidney mRNA, using standard molecular biology techniques. The cDNA sequence is stably transfected into CHO cells, and clones are assayed for SGLT2 activity essentially as described in Ryan et al., "HK-2: an immortalized proximal tubule epithelial cell line from normal adult human kidney", *Kidney International*, 45:48-57 (1994). Evaluation of inhibition of SGLT2 activity in a clonally selected cell line is performed essentially as described in Ryan et al. (1994), with the following modifications. Cells are grown in 96-well plates for 2-4 days to 75,000 or 30,000 cells per well in F-12 nutrient mixture (Ham's F-12), 10% fetal bovine serum, 300 ug/ml Geneticin and penicillin-streptomycin. At confluence, the cells are washed twice with 10 mM HEPES/Tris, pH 7.4, 137 mM N-methyl-D-glucamine, 5.4 mM KCl, 2.8 mM CaCl₂, 1.2 mM MgSO₄. Cells are then incubated with 10 μM [¹⁴C]AMG, and 10 μM inhibitor (final DMSO = 0.5%) in 10 mM HEPES/Tris, pH 7.4, 137 mM NaCl, 5.4 mM KCl, 2.8 mM CaCl₂, 1.2 mM MgSO₄ at 37°C for 1.5 hours. Uptake assays are quenched with ice cold IX PBS containing 0.5 mM phlorizin, and cells are then lysed with 0.1% NaOH. After addition of MicroScint scintillation fluid, the cells are allowed to shake for 1 hour, and then [¹⁴C]AMG (glucose analog α-methyl-D-glucopyranoside) is quantitated on a TopCount scintillation counter. Controls are performed with and without NaCl. For determination of EC₅₀ values, 10 inhibitor concentrations (dapagliflozin) are used over 2 log intervals in the appropriate response range, and triplicate plates are averaged across plates.

[0064] The pharmaceutical formulations described herein containing dapagliflozin or dapagliflozin propylene glycol hydrate possess activity as an inhibitor of the sodium dependent glucose transporters found in the intestine and kidney of mammals, is a selective inhibitor of renal SGLT2 activity, and therefore can be used in the treatment of diseases or disorders associated with SGLT2 activity.

[0065] Accordingly, the pharmaceutical dapagliflozin and dapagliflozin propylene glycol hydrate formulations described herein can be administered to mammals, preferably humans, for the treatment of a variety of conditions and disorders associated with SGLT2 activity including treating or delaying the progression or onset of diabetes (including Type I and Type II diabetes, impaired glucose tolerance, insulin resistance, and diabetic complications, such as nephropathy, retinopathy, neuropathy and cataracts), hyperglycemia, hyperinsulinemia, hypercholesterolemia, dyslipidemia, elevated blood levels of free fatty acids or glycerol, hyperlipidemia, hypertriglyceridemia, obesity, wound healing, tissue ischemia, atherosclerosis and hypertension. The formulations described herein can also be utilized to increase the blood levels of high density lipoprotein (HDL). In addition, the conditions, diseases, and maladies collectively referenced to as "Syndrome X" or Metabolic Syndrome as detailed in Johannsson, *J. Clin. Endocrinol. Metab.*, 82, 727-34 (1997), can be treated employing the formulations described herein.

[0066] In one embodiment, the invention provides the pharmaceutical dapagliflozin (S)-propylene glycol hydrate formulations of the invention for use in the treatment of type II diabetes. In another embodiment, the invention provides the pharmaceutical dapagliflozin (S)-propylene glycol hydrate formulations of the invention for use in delaying the progression or onset of type II diabetes.

[0067] Also described is a method for treating or delaying the progression or onset of diseases or disorders associated with SGLT2 activity comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of the pharmaceutical dapagliflozin propylene glycol hydrate formulation of the invention. Also described is a method for treating type II diabetes comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of the pharmaceutical dapagliflozin propylene glycol hydrate formulation of the invention. Also described is a method for delaying the progression or onset of type II diabetes comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of the pharmaceutical dapagliflozin propylene glycol hydrate formulation of the invention.

[0068] Also described is the use of the pharmaceutical dapagliflozin or dapagliflozin propylene glycol hydrate formulation (including the stock granulation, capsules containing same, and tablets thereof) in the manufacture of a medicament

for the treatment of type II diabetes. Also described is the use of the pharmaceutical dapagliflozin propylene glycol hydrate formulation in the manufacture of a medicament for delaying the progression or onset of type II diabetes. Also described is the pharmaceutical dapagliflozin propylene glycol hydrate formulation for use in therapy in treating or delaying the progression or onset of type II diabetes.

5 **[0069]** Other therapeutic agent(s) suitable for combination with the formulations of the present invention include, but are not limited to, known therapeutic agents useful in the treatment of the aforementioned disorders associated with SGLT2 activity including: anti-diabetic agents; anti-hyperglycemic agents; hypolipidemic or lipid lowering agents; anti-obesity agents; anti-hypertensive agents and appetite suppressants.

10 **[0070]** Also described is a method for treating or delaying the progression or onset of diseases or disorders associated with SGLT2 activity comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of the pharmaceutical formulation and one or more of the following: anti-diabetic agent(s), anti-hyperglycemic agent(s); hypolipidemic or lipid lowering agent(s); anti-obesity agent(s); anti-hypertensive agent(s) and appetite suppressant(s).

15 **[0071]** Also described is a method for treating type II diabetes comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of the pharmaceutical formulation and one or more anti-diabetic agent(s). Also described is a method for delaying the progression or onset of type II diabetes comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of the pharmaceutical formulation described herein and one or more anti-diabetic agent(s).

20 **[0072]** Also described is a method for treating or delaying the progression or onset of type II diabetes comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of the pharmaceutical formulation described herein and one or more of the following: anti-hyperglycemic agent(s); hypolipidemic or lipid lowering agent(s); anti-obesity agent(s); anti-hypertensive agent(s) and appetite suppressant(s). For example, described is a method for treating or delaying the progression or onset of type II diabetes comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of a pharmaceutical formulation described herein and an anti-hyperglycemic agent(s). Also described is a method for treating or delaying the progression or onset of type II diabetes comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of a pharmaceutical formulation described herein and a hypolipidemic agent(s). Also described is a method for treating or delaying the progression or onset of type II diabetes comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of a pharmaceutical formulation described herein and an anti-obesity agent(s). Also described is a method for treating or delaying the progression or onset of type II diabetes comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of a pharmaceutical formulation described herein and an anti-hypertensive agent(s). Also described is a method for treating or delaying the progression or onset of type II diabetes comprising administering to a mammalian species in need of such treatment a therapeutically effective amount of a pharmaceutical formulation described herein and an appetite suppressant(s).

30 **[0073]** Also described is the use of the combination of the pharmaceutical dapagliflozin propylene glycol hydrate formulations described herein and one or more agents selected from the group consisting of anti-diabetic agents, anti-hyperglycemic agents, hypolipidemic or lipid lowering agents, anti-obesity agents, anti-hypertensive agents, and appetite suppressants as a medicament for the treatment of diseases or disorders associated with SGLT2 activity, for example, Type I and Type II diabetes; impaired glucose tolerance; insulin resistance; and diabetic complications, such as nephropathy, retinopathy, neuropathy and cataracts; hyperglycemia; hyperinsulinemia; hypercholesterolemia; dyslipidemia; elevated blood levels of free fatty acids or glycerol; hyperlipidemia; hypertriglyceridemia; obesity; wound healing; tissue ischemia; atherosclerosis; hypertension; and Syndrome X or Metabolic Syndrome.

40 **[0074]** Also described is the combination of the pharmaceutical dapagliflozin propylene glycol hydrate formulation and one or more anti-diabetic agents as a medicament for the treatment of type II diabetes. Also described is the combination of the pharmaceutical dapagliflozin propylene glycol hydrate formulation and one or more anti-diabetic agents as a medicament for delaying the progression or onset of type II diabetes. Also described is the combination of the pharmaceutical dapagliflozin propylene glycol hydrate formulation and one or more anti-hyperglycemic agents as a medicament for the treatment of type II diabetes. Also described is the combination of the pharmaceutical dapagliflozin propylene glycol hydrate formulation and one or more anti-hyperglycemic agents as a medicament for delaying the progression or onset of type II diabetes. Also described is the combination of the pharmaceutical dapagliflozin propylene glycol hydrate formulation and one or more hypolipidemic agents or lipid-lowering agents as a medicament for the treatment of type II diabetes. Also described is the combination of the pharmaceutical dapagliflozin propylene glycol hydrate formulation and one or more hypolipidemic agents or lipid-lowering agents as a medicament for delaying the progression or onset of type II diabetes. Also described is the combination of the pharmaceutical dapagliflozin or dapagliflozin propylene glycol hydrate formulation of the invention and one or more anti-obesity agents as a medicament for the treatment of type II diabetes. Also described is the combination of the pharmaceutical dapagliflozin propylene glycol hydrate formulation and one or more anti-obesity agents as a medicament for delaying the progression or onset of type II diabetes. Also described is the combination of the pharmaceutical dapagliflozin propylene glycol hydrate formulation and one or

more anti-hypertensive agents as a medicament for the treatment of type II diabetes. Also described is the combination of the pharmaceutical dapagliflozin propylene glycol hydrate formulation and one or more anti-hypertensive agents as a medicament for delaying the progression or onset of type II diabetes. Also described is the combination of the pharmaceutical dapagliflozin propylene glycol hydrate formulation and one or more appetite suppressants as a medicament for delaying the progression or onset of type II diabetes.

[0075] The formulations of the invention in the form of capsules or tablets containing dapagliflozin propylene glycol hydrate can be administered in dosages of about 0.1 mg to about 750 mg per day, in single or divided doses or multiple doses which can be administered 1 to 4 times daily. In one embodiment, the formulations of the invention in the form of capsules or tablets containing dapagliflozin propylene glycol hydrate is administered in dosages of about 0.2 mg to about 600 mg per day, in single or divided doses or multiple doses which can be administered 1 to 4 times daily. In another embodiment, the formulations of the invention in the form of capsules or tablets containing dapagliflozin propylene glycol hydrate is administered in dosages of from about 0.5 mg to about 100 mg per day, in single or divided doses or multiple doses which can be administered 1 to 4 times daily.

[0076] The present invention includes within its scope pharmaceutical formulations containing, as an active ingredient, a therapeutically effective amount of dapagliflozin propylene glycol hydrate, alone or in combination with a pharmaceutical carrier or diluent as described. Optionally, the formulations of the present invention can be utilized as an individual treatment, or utilized in combination with one or more other therapeutic agent(s) in the same dosage form (fixed dosage) or separate dosage forms.

[0077] Other therapeutic agent(s) suitable for combination with the formulations of the present invention include, but are not limited to, known therapeutic agents useful in the treatment of the aforementioned disorders including: anti-diabetic agents; anti-hyperglycemic agents; hypolipidemic/lipid lowering agents; anti-obesity agents; anti-hypertensive agents and appetite suppressants.

[0078] Examples of suitable anti-diabetic agents for use in combination with the formulations of the present invention include, but are not limited to, biguanides (e.g., metformin or phenformin), glucosidase inhibitors (e.g., acarbose or miglitol), insulins (including insulin secretagogues or insulin sensitizers), meglitinides (e.g., repaglinide), sulfonylureas (e.g., glimepiride, glyburide, gliclazide, chlorpropamide and glipizide), biguanide/glyburide combinations (e.g., Glucovance®), thiazolidinediones (e.g., troglitazone, rosiglitazone and pioglitazone), PPAR-alpha agonists, PPAR-gamma agonists, PPAR alpha/gamma dual agonists, glycogen phosphorylase inhibitors, inhibitors of fatty acid binding protein (aP2), glucagon-like peptide-1 (GLP-1) and other agonists of the GLP-1 receptor, and dipeptidyl peptidase IV (DPP4) inhibitors.

[0079] Other suitable thiazolidinediones include, but are not limited to, MCC-555 (disclosed in U.S. Patent No. 5,594,016, Mitsubishi), faraglitazar (GI-262570, Glaxo-Wellcome), englitazone (CP-68722, Pfizer) or darglitazone (CP-86325, Pfizer; isaglitazone, MIT/Johnson & Johnson), reglitazar (JTT-501, (JPNT/Pharmacia & Upjohn), rivoglitazone (R-119702, Sankyo/WL), liraglutide (NN-2344, Dr. Reddy/NN), and (Z)-1,4-bis-4-[(3,5-dioxo-1,2,4-oxadiazolidin-2-ylmethyl)]phenoxybut-2-ene (YM-440, Yamanouchi).

[0080] Examples of PPAR-alpha agonists, PPAR-gamma agonists and PPAR alpha/gamma dual agonists include, but are not limited to, muraglitazar, peliglitazar, tesaglitazar AR-HO39242 (Astra/Zeneca), GW-501516 (Glaxo-Wellcome), KRP297 (Kyorin Merck), as well as those disclosed by Murakami et al, "A Novel Insulin Sensitizer Acts As a Coligand for Peroxisome Proliferation - Activated Receptor Alpha (PPAR alpha) and PPAR gamma. Effect on PPAR alpha Activation on Abnormal Lipid Metabolism in Liver of Zucker Fatty Rats", Diabetes 47, 1841-1847 (1998); WO 01/21602 and in U.S. Patent No. 6,414,002 and U.S. Patent No. 6,653,314.

[0081] Suitable aP2 inhibitors include, but are not limited to, those disclosed in U.S. application Serial No. 09/391,053, filed September 7, 1999, and in U.S. Patent No. 6,548,529.

[0082] Suitable DPP4 inhibitors include, but are not limited to, those disclosed in WO99/38501, WO99/46272, WO99/67279 (PROBIODRUG), WO99/67278 (PROBIODRUG), WO99/61431 (PROBIODRUG), NVP-DPP728A (1-[[[2-[(5-cyanopyridin-2-yl)amino]ethyl]amino]acetyl]-2-cyano-(S)-pyrrolidine] (Novartis) as disclosed by Hughes et al, Biochemistry, 38(36), 11597-11603, 1999, TSL-225 (tryptophyl-1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid (disclosed by Yamada et al, Bioorg. & Med. Chem. Lett. 8 (1998) 1537-1540), 2-cyanopyrrolidides and 4-cyanopyrrolidides, as disclosed by Ashworth et al, Bioorg. & Med. Chem. Lett., Vol. 6, No. 22, pp 1163-1166 and 2745-2748 (1996), the compounds disclosed in U.S. application Serial No. 10/899,641, WO 01/68603 and U.S. patent 6,395,767. In one embodiment, the DPP4 inhibitor is saxagliptin.

[0083] Other suitable meglitinides include nateglinide (Novartis) or KAD1229 (PF/Kissei).

[0084] Examples of suitable anti-hyperglycemic agents for use in combination with the formulations of the present invention include, but are not limited to, glucagon-like peptide-1 (GLP-1) such as GLP-1(1-36) amide, GLP-1(7-36) amide, GLP-1(7-37) (as disclosed in U.S. Patent No. 5,614,492), as well as exenatide (Amylin/Lilly), LY-315902 (Lilly), MK-0431 (Merck), liraglutide (NovoNordisk), ZP-10 (Zealand Pharmaceuticals A/S), CJC-1131 (Conjuchem Inc), and

the compounds disclosed in WO 03/033671.

[0085] Examples of suitable hypolipidemic/lipid lowering agents for use in combination with the formulations of the present invention include one or more MTP inhibitors, HMG CoA reductase inhibitors, squalene synthetase inhibitors, fibric acid derivatives, ACAT inhibitors, lipoxygenase inhibitors, cholesterol absorption inhibitors, ileal Na⁺/bile acid co-transporter inhibitors, up-regulators of LDL receptor activity, bile acid sequestrants, cholesterol ester transfer protein (e.g., CETP inhibitors, such as torcetrapib (CP-529414, Pfizer) and JTT-705 (Akros Pharma)), PPAR agonists (as described above) and/or nicotinic acid and derivatives thereof. The hypolipidemic agent can be an up-regulator of LD2 receptor activity, such as 1(3H)-isobenzofuranone,3-(13-hydroxy-10-oxotetradecyl)-5,7-dimethoxy- (MD-700, Taisho Pharmaceutical Co. Ltd) and cholestan-3-ol,4-(2-propenyl)-(3a,4a,5a)- (LY295427, Eli Lilly). Preferred hypolipidemic agents include pravastatin, lovastatin, simvastatin, atorvastatin, fluvastatin, cerivastatin, atavastatin and rosuvastatin (ZD-4522), for example.

[0086] Examples of MTP inhibitors that can be employed as described above include, but are not limited to, those disclosed in U.S. Patent No. 5,595,872, U.S. Patent No. 5,739,135, U.S. Patent No. 5,712,279, U.S. Patent No. 5,760,246, U.S. Patent No. 5,827,875, U.S. Patent No. 5,885,983 and U.S. Patent No 5,962,440.

[0087] Examples of HMG CoA reductase inhibitors that can be employed in combination with the formulations of the invention include, but are not limited to, mevastatin and related compounds, as disclosed in U.S. Patent No. 3,983,140, lovastatin (mevinolin) and related compounds, as disclosed in U.S. Patent No. 4,231,938, pravastatin and related compounds, such as disclosed in U.S. Patent No. 4,346,227, simvastatin and related compounds, as disclosed in U.S. Patent Nos. 4,448,784 and 4,450,171. Other suitable HMG CoA reductase inhibitors that can be employed herein include, but are not limited to, fluvastatin, disclosed in U.S. Patent No. 5,354,772, cerivastatin, as disclosed in U.S. Patent Nos. 5,006,530 and 5,177,080, atorvastatin, as disclosed in U.S. Patent Nos. 4,681,893, 5,273,995, 5,385,929 and 5,686,104, atavastatin (Nissan/Sankyo's nisvastatin (NK-104)), as disclosed in U.S. Patent No. 5,011,930, rosuvastatin (Shionogi-Astra/Zeneca (ZD-4522)), as disclosed in U.S. Patent No. 5,260,440, and related statin compounds disclosed in U.S. Patent No. 5,753,675, pyrazole analogs of mevalonolactone derivatives, as disclosed in U.S. Patent No. 4,613,610, indene analogs of mevalonolactone derivatives, as disclosed in PCT application WO 86/03488, 6-[2-(substituted-pyrrol-1-yl)-alkyl]pyran-2-ones and derivatives thereof, as disclosed in U.S. Patent No. 4,647,576, Searle's SC-45355 (a 3-substituted pentanedioic acid derivative) dichloroacetate, imidazole analogs of mevalonolactone, as disclosed in PCT application WO 86/07054, 3-carboxy-2-hydroxy-propane-phosphonic acid derivatives, as disclosed in French Patent No. 2,596,393, 2,3-disubstituted pyrrole, furan and thiophene derivatives, as disclosed in European Patent Application No. 0221025, naphthyl analogs of mevalonolactone, as disclosed in U.S. Patent No. 4,686,237, octahydronaphthalenes, such as disclosed in U.S. Patent No. 4,499,289, keto analogs of mevinolin (lovastatin), as disclosed in European Patent Application No.0142146 A2, and quinoline and pyridine derivatives, as disclosed in U.S. Patent No. 5,506,219 and 5,691,322. In addition, phosphinic acid compounds useful in inhibiting HMG CoA reductase, such as those disclosed in GB 2205837, are suitable for use in combination with the formulations of the present invention.

[0088] Examples of squalene synthetase inhibitors suitable for use herein include, but are not limited to, α -phosphono-sulfonates disclosed in U.S. Patent No. 5,712,396, those disclosed by Biller et al., J. Med. Chem., 1988, Vol. 31, No. 10, pp. 1869-1871, including isoprenoid (phosphinyl-methyl)phosphonates, as well as other known squalene synthetase inhibitors, for example, as disclosed in U.S. Patent No. 4,871,721 and 4,924,024 and in Biller, S.A., Neuenschwander, K., Ponpipom, M.M., and Poulter, C.D., Current Pharmaceutical Design, 2, 1-40 (1996). Other squalene synthetase inhibitors suitable for use herein include the terpenoid pyrophosphates disclosed by P. Ortiz de Montellano et al, J. Med. Chem., 1977, 20, 243-249; the farnesyl diphosphate analog A and presqualene pyrophosphate (PSQ-PP) analogs as disclosed by Corey and Volante, J. Am. Chem. Soc., 1976, 98, 1291-1293; phosphinylphosphonates reported by McClard, R.W. et al, J.A.C.S., 1987, 109, 5544; and cyclopropanes reported by Capson, T.L., PhD dissertation, June, 1987, Dept. Med. Chem. U of Utah, Abstract, Table of Contents, pp 16, 17, 40-43, 48-51, Summary.

[0089] Examples of fibric acid derivatives that can be employed in combination the formulations of the invention include, but are not limited to, fenofibrate, gemfibrozil, clofibrate, bezafibrate, ciprofibrate, clinofibrate and the like, probucol, and related compounds, as disclosed in U.S. Patent No. 3,674,836, bile acid sequestrants, such as cholestyramine, colestipol and DEAE-Sephadex (Secholex®, Policexide®), as well as lipostabil (Rhone-Poulenc), Eisai E-5050 (an N-substituted ethanolamine derivative), imanixil (HOE-402), tetrahydrolipstatin (THL), istigmastanylphos-phorylcholine (SPC, Roche), aminocyclodextrin (Tanabe Seiyoku), Ajinomoto AJ-814 (azulene derivative), melinamide (Sumitomo), Sandoz 58-035, American Cyanamid CL-277,082 and CL-283,546 (disubstituted urea derivatives), nicotinic acid, acipimox, acifran, neomycin, p-aminosalicylic acid, aspirin, poly(diallylmethylamine) derivatives, such as disclosed in U.S. Patent No. 4,759,923, quaternary amine poly(diallyldimethylammonium chloride) and ionenes, such as disclosed in U.S. Patent No. 4,027,009, and other known serum cholesterol lowering agents. In one embodiment, the fibric acid derivative is probucol or gemfibrozil.

[0090] Examples of ACAT inhibitors that can be employed in combination with the formulations of the invention include, but are not limited to, those disclosed in Drugs of the Future 24, 9-15 (1999), (Avasimibe); "The ACAT inhibitor, CI-1011 is effective in the prevention and regression of aortic fatty streak area in hamsters", Nicolosi et al, Atherosclerosis

(Shannon, Irel). (1998), 137(1), 77-85; "The pharmacological profile of FCE 27677: a novel ACAT inhibitor with potent hypolipidemic activity mediated by selective suppression of the hepatic secretion of ApoB100-containing lipoprotein", Ghiselli, Giancarlo, *Cardiovasc. Drug Rev.* (1998), 16(1), 16-30; "RP 73163: a bioavailable alkylsulfanyl-diphenylimidazole ACAT inhibitor", Smith, C., et al, *Bioorg. Med. Chem. Lett.* (1996), 6(1), 47-50; "ACAT inhibitors: physiologic mechanisms for hypolipidemic and anti-atherosclerotic activities in experimental animals", Krause et al, Editor(s): Ruffolo, Robert R., Jr.; Hollinger, Manfred A., *Inflammation: Mediators Pathways* (1995), 173-98, Publisher: CRC, Boca Raton, Fla.; "ACAT inhibitors: potential anti-atherosclerotic agents", Sliskovic et al, *Curr. Med. Chem.* (1994), 1(3), 204-25; "Inhibitors of acyl-CoA:cholesterol O-acyl transferase (ACAT) as hypocholesterolemic agents. The first water-soluble ACAT inhibitor with lipid-regulating activity. Inhibitors of acyl-CoA:cholesterol acyltransferase (ACAT). Development of a series of substituted N-phenyl-N'-[(1-phenylcyclopentyl)methyl]ureas with enhanced hypocholesterolemic activity", Stout et al, *Chemtracts: Org. Chem.* (1995), 8(6), 359-62, or TS-962 (Taisho Pharmaceutical Co. Ltd).

[0091] Examples of suitable cholesterol absorption inhibitors for use in combination with the formulations of the invention include, but are not limited to, SCH48461 (Schering-Plough), as well as those disclosed in *Atherosclerosis* 115, 45-63 (1995) and *J. Med. Chem.* 41, 973 (1998).

[0092] Examples of suitable ileal Na⁺/bile acid co-transporter inhibitors for use in combination with the formulations of the invention include, but are not limited to, compounds as disclosed in *Drugs of the Future*, 24, 425-430 (1999).

[0093] Examples of lipoxygenase inhibitors that can be employed in combination with the formulations of the invention include, but are not limited to, 15-lipoxygenase (15-LO) inhibitors, such as benzimidazole derivatives, as disclosed in WO 97/12615, 15-LO inhibitors, as disclosed in WO 97/12613, isothiazolones, as disclosed in WO 96/38144, and 15-LO inhibitors, as disclosed by Sendobry et al "Attenuation of diet-induced atherosclerosis in rabbits with a highly selective 15-lipoxygenase inhibitor lacking significant antioxidant properties", *Brit. J. Pharmacology* (1997) 120, 1199-1206, and Comicelli et al., "15-Lipoxygenase and its Inhibition: A Novel Therapeutic Target for Vascular Disease", *Current Pharmaceutical Design*, 1999, 5, 11-20.

[0094] Examples of suitable anti-hypertensive agents for use in combination with the formulations of the present invention include, but are not limited to, beta adrenergic blockers, calcium channel blockers (L-type and T-type; e.g. diltiazem, verapamil, nifedipine, amlodipine and mibefradil), diuretics (e.g., chlorothiazide, hydrochlorothiazide, flumethiazide, hydroflumethiazide, bendroflumethiazide, methylchlorothiazide, trichloromethiazide, polythiazide, benzthiazide, ethacrynic acid, tricyclics, chlorthalidone, furosemide, musolimine, bumetanide, triamterene, amiloride, spironolactone), renin inhibitors, ACE inhibitors (e.g., captopril, zofenopril, fosinopril, enalapril, ceranopril, cilazapril, delapril, pentopril, quinapril, ramipril, lisinopril), AT-1 receptor antagonists (e.g., losartan, irbesartan, valsartan), ET receptor antagonists (e.g., sitaxsentan, atrisentan and compounds disclosed in U.S. Patent Nos. 5,612,359 and 6,043,265), Dual ET/All antagonist (e.g., compounds disclosed in WO 00/01389), neutral endopeptidase (NEP) inhibitors, vasopeptidase inhibitors (dual NEP-ACE inhibitors) (e.g., omapatrilat and gemopatrilat), and nitrates.

[0095] Examples of suitable anti-obesity agents for use in combination with the formulations of the present invention include, but are not limited to, beta 3 adrenergic agonists, lipase inhibitors, serotonin (and dopamine) reuptake inhibitors, thyroid receptor beta drugs, 5HT_{2C} agonists, (such as Arena APD-356); MCHR1 antagonists, such as Synaptic SNAP-7941 and Takeda T-226926, melanocortin receptor (MC4R) agonists, melanin-concentrating hormone receptor (MCHR) antagonists (such as Synaptic SNAP-7941 and Takeda T-226926), galanin receptor modulators, orexin antagonists, CCK agonists, NPY1 or NPY5 antagonist, NPY2 and NPY4 modulators, corticotropin releasing factor agonists, histamine receptor-3 (H3) modulators, 11-beta-HSD-1 inhibitors, adipopectin receptor modulators, monoamine reuptake inhibitors or releasing agents, ciliary neurotrophic factors (CNTF, such as AXOKINE® by Regeneron), BDNF (brain-derived neurotrophic factor), leptin and leptin receptor modulators, cannabinoid-1 receptor antagonists (such as SR-141716 (Sanofi) or SLV-319 (Solvay)), and anorectic agents.

[0096] Beta 3 adrenergic agonists that can be optionally employed in combination with formulations of the present invention include, but are not limited to, AJ9677 (Takeda/Dainippon), L750355 (Merck), CP331648 (Pfizer,) or other known beta 3 agonists, as disclosed in U.S. Patent Nos. 5,541,204, 5,770,615, 5,491,134, 5,776,983 and 5,488,064.

[0097] Examples of lipase inhibitors that can be employed in combination with formulations of the present invention include, but are not limited to, orlistat and ATL-962 (Alizyme).

[0098] Serotonin (and dopamine) reuptake inhibitors (or serotonin receptor agonists) that can be employed in combination with the formulations of the present invention include, but are not limited to, BVT-933 (Biovitrum), sibutramine, topiramate (Johnson & Johnson) and axokine (Regeneron).

[0099] Examples of thyroid receptor beta compounds that can be employed in combination with formulations of the present invention include, but are not limited to, thyroid receptor ligands, such as those disclosed in WO 97/21993 (U. Cal SF), WO 99/00353 (KaroBio) and WO 00/039077 (KaroBio).

[0100] Examples of monoamine reuptake inhibitors that can be employed in combination with the formulations of the present invention include, but are not limited to, fenfluramine, dexfenfluramine, fluvoxamine, fluoxetine, paroxetine, sertraline, chlorphentermine, cloforex, clortermine, picilorex, sibutramine, dexamphetamine, phentermine, phenylpropanolamine and mazindol.

[0101] Anorectic agents that can be employed in combination with the formulations of the present invention include, but are not limited to, topiramate (Johnson & Johnson), dexamphetamine, phentermine, phenylpropanolamine and mazindol.

[0102] Where any of the formulations of the invention are used in combination with other therapeutic agent(s), the other therapeutic agent(s) can be used, for example, in the amounts indicated in the Physician's Desk Reference, as in the cited patents and patent applications set out above, or as otherwise known and used by one of ordinary skill in the art.

[0103] Where any of the formulations of the invention are used in combination with other therapeutic agent(s), each of the compounds of the combination can be administered simultaneously or sequentially and in any order, and the components can be administered separately or as a fixed combination, in jointly therapeutically effective amounts, for example, in daily dosages as described herein. In one embodiment of the invention, a fixed combination of the invention can be prepared by mixing a dry granulation of the dapagliflozin or dapagliflozin propylene glycol hydrate formulation of the invention and a dry granulation of the other therapeutic agent(s) and filling the mixture into capsules of desired size, shape, color, or other characteristics, or compressed to form tablets.

[0104] The following examples are provided to describe the invention in further detail. These examples, which set forth the best mode presently contemplated for carrying out the invention, are intended to illustrate and not to limit the invention.

EXAMPLES


[0105] The following working Examples are illustrative of the present invention. All temperatures are expressed in degrees Centigrade unless otherwise indicated.

Example 1: Preparation of dapagliflozin (compound I)

[0106] The preparation of compounds of structure I is generally described in U.S. Patent 6,414,126, and specifically described in Scheme 1 and Example 1 of U.S. Patent 5,515,117. Stable forms of compounds of structure (I) can be crystallized as solvates (e.g., hydrates) or complexes.

Example 2A: Preparation of dapagliflozin (S)-propylene glycol hydrate (Ia)

[0107] The preparation of structure Ia is described and depicted schematically below.



[0108] Compound A can be prepared as described in Example 1, Part E of U.S. Patent 6,515,117.

[0109] A 10-L glass reactor equipped with a thermocouple and a nitrogen inlet was charged with MeOH (1.25 L), deionized water (3.6 L) followed by 50% aqueous NaOH (205.9 ml, 3.899 mol). The residual solution of NaOH in the measuring cylinder was transferred with water (94 ml) to the reaction vessel. Compound A (503.11 g, 0.872 mol) was added and the mixture was stirred and heated to ~68°C over 1.5 hour. After 1 hour, the circulation bath temperature was lowered from 80°C to 70°C; internal temperature became 65°C. After a total of 3 hours, HPLC indicated completion of reaction, Compound I AP ~99.5. (HPLC: Column: YMC ODS-A (C-18) S3, 4.6 x 50 mm. Solvent A: 0.2% aq. H₃PO₄. Solvent B: 90% CH₃CN/10%H₂O Start %B = 0, final %B = 100 Gradient time 8 min; hold time 3 minutes. Integration

EP 2 139 494 B1

stop time 11.0 minutes. Flow rate 2.5 ml/minute. UV wave length 220 nm.)

[0110] After the mixture was cooled to 25°C, isopropyl acetate (2.5 L) was added. The mixture was stirred for 10 minutes and then the aqueous layer was separated (pH = 12.5) and organic layer was washed with water (1 L). During this wash the pH of the biphasic system was adjusted to 6.0 with concentrated HCl (5.0 ml) and then the aqueous layer was separated. Neutralization before phase split was done to prevent contamination of the product with NaOH. The (S)-propylene glycol structure prepared without neutralization was slightly basic [pH 8.3 of a suspension sonicated in water (~20 mg/ml)].


[0111] The organic layer was collected in a separate vessel. The reactor was washed with water (2 L), MeOH (2 L) and flushed with nitrogen gas. The wet solution of compound B was recharged into the reactor and (S)-propylene glycol ((S)-PG) (67.03 g, 0.872 mole) was introduced. Optionally, seed crystals of (S)-PG Ia can be added at this stage. Seed crystals can be prepared by dissolving compound I in a solvent such as MTBE and treating the resulting solution with (S)-propylene glycol and proceeding as described above without the use of seeding.

[0112] Instantaneous crystallization produced a thick slurry. After stirring for 1 hour, cyclohexane (2.5 L) was added rapidly over 10 minutes and the stirring was continued for 21 hours. The product was filtered through a filter paper (Whatman #5, Buchner funnel 24" diameter). The filtration was rapid and took about 15 minutes. The filter cake was washed with a mixture (1:1) of MTBE/cyclohexane (2 x 1 L) and dried under suction for 0.5 hour. The solid was transferred to a pyrex tray and dried under vacuum (25 mm Hg) in an oven at 25-30°C for two days till water analysis by K.F. corresponded to monohydrate (3.6 wt.%). The (S)-PG product Ia was obtained (0.425 kg, yield 97%) as a snow white solid, mp 71°C, HPLC AP 99.7. (HPLC method: Mobile Phase A: 0.05% TFA in H₂O. Mobile Phase B: 0.05% TFA in CAN. Column: YMC Hydrosphere 4.6x150 (3μ). Gradient: 30-90%B over 45 minutes, hold 5 minutes; back to 30%B and re-equilibrate for 10 min. Wavelength: 220 nm. Injection Volume: 10μl. Temperature: Ambient).


[0113] The (R) form of dapagliflozin can be prepared using these methods and substituting (S)-propylene glycol with (R)-propylene glycol.

Example 2B: Preparation of dapagliflozin (S)-propylene glycol hydrate (compound Ia)

[0114] The structure Ia can alternatively be prepared as described and depicted schematically below.



$C_{29}H_{33}ClO_{10}$
Exact Mass: 576.18
Mol. Wt.: 577.02
C, 60.36; H, 5.76; Cl, 6.14; O, 27.73



$C_{24}H_{35}ClO_9$
Exact Mass: 502.20
Mol. Wt.: 502.98
C, 57.31; H, 7.01; Cl, 7.05; O, 28.63

[0115] 20g of compound A was charged to a reactor at ambient temperature and pressure. 30mL Methanol and 49.75mL 3N NaOH were added to the reactor and the reaction mixture was heated to 80°C or reflux, and held about 2-3 hours for reaction completion < 0.5 AP. The batch was cooled to 20°C and neutralized to pH 6.0-7.5 using 1N acetic acid (requires ~ 1mL/gm input).

[0116] Extraction: The product was extracted from the reaction mixture into 100mL isopropyl acetate, the aqueous

EP 2 139 494 B1

phase was split away and the organic phase washed with water until conductivity < 10mS (~ 4mL/gm input). The aqueous phase was split away.

[0117] Crystallization: 2.8g (1.05 eq) (S)-(+)-1,2 Propanediol 96%+ was added to the reaction mixture. The batch was seeded with 0.1g compound I seed. 160mL Cyclohexane was added and the batch cooled to 5°C. The batch was allowed to stir at 5°C at least 1 hour before isolation.

[0118] Isolation and Drying: Each load of isolated cake was washed with 50/50 by volume isopropyl acetate/cyclohexane mixture. The cake was dried at 30°C in a vacuum oven under full vacuum. (Cake is dry when KF = 3.6% - 4.1%).

Yield = 84% (uncorrected)

Typical purity = 99.81AP

Typical PG content = 15.1 - 15.8% by GC

[0119] Capsules containing the SGLT2 inhibitor of Formula I (dapagliflozin) or Formula Ia (dapagliflozin (S)- Propylene glycol hydrate) were prepared in strengths of 2.5 mg (Example 3), 10 mg (Example 4) and 100 mg (Example 5) as two-piece, gray opaque size #0 (2.5 mg and 10 mg) and size #00 (for 100 mg) hard gelatin capsules.

Example 3 - Preparation of Dapagliflozin/Dapagliflozin propylene glycol hydrate Capsule, 2.5 mg

[0120] A 25.0 mg of stock granulation was prepared containing 10% dapagliflozin or dapagliflozin propylene glycol hydrate filled in gray, opaque, size #0 capsule shell.

A. Stock Granulation Composition

[0121]

Ingredient	Amount (% w/w)
Dapagliflozin or Dapagliflozin propylene glycol hydrate (Or equivalent amount of Dapagliflozin propylene glycol hydrate)	10.0
Pregelatinized Starch	15.0
Microcrystalline Cellulose	68.75
Sodium Starch Glycolate	3.0
Silicon Dioxide	2.0
Magnesium Stearate	1.25

[0122] The amount of dapagliflozin is theoretically equivalent to 81.29% of dapagliflozin propylene glycol hydrate, either of which can be used. The actual amount of dapagliflozin propylene glycol hydrate will depend on the purity. The microcrystalline cellulose is the compensating excipient whose amount can vary depending on the actual amount of dapagliflozin propylene glycol hydrate and magnesium stearate used. The preferred amount of magnesium stearate is 1.25% (w/w). A useful range is 1.25-1.50% (w/w).

[0123] The stock granulation of Part A and the Example 3 capsules were prepared according to the following procedures.

B. Example 3 Stock granulation procedure

[0124]

1. Screen dapagliflozin or dapagliflozin propylene glycol hydrate.
2. Screen silicon dioxide.
3. Mix silicon dioxide with dapagliflozin or dapagliflozin propylene glycol hydrate in a suitable blender.
4. Screen pregelatinized starch and microcrystalline cellulose, if necessary.
5. Add ingredients from Step 4 to a suitable blender.
6. Add mixture from Step 3 to the blend from Step 5, and mix.
7. Screen sodium starch glycolate.
8. Add ingredient from Step 7 to the blend from Step 6, and mix.

EP 2 139 494 B1

9. Screen the blend from Step 8, and mix.
10. Screen portion of magnesium stearate.
11. Add ingredient from Step 10 to the blend from Step 9, and mix.
12. Densify the blend from Step 11.
13. Reduce the densified blend Step 12.
14. Screen the remaining portion of magnesium stearate.
15. Add ingredient from Step 14 to the granulation from Step 13, and mix.

C. Example 3 Product: Dapagliflozin/Dapagliflozin propylene glycol hydrate Capsule, 2.5 mg

[0125]

1. Fill empty capsule shells with sufficient Example 3 Part A stock granulation for capsules (10.0%) w/w (as the non-solvated form), to provide 2.5 mg capsules.
2. De-dust the capsules.

Example 4-Preparation of Dapagliflozin/Dapagliflozin propylene glycol hydrate Capsule, 10 mg

A. Stock Granulation Composition

[0126] Stock granulation composition was prepared as described in Example 3A.

B. Example 4 Stock granulation procedure

[0127] Stock granulation procedure was performed as described in Example 3B.

C. Example 4 Product: Dapagliflozin Capsule, 10 mg

[0128]

1. Fill empty capsule shells with Example 3 Part A stock granulation for capsules (10.0 % w/w as the non-solvated form), to provide 10 mg capsules.
2. De-dust the capsules.
3. Weight sort the capsules.

Example 5: Preparation of Dapagliflozin/Dapagliflozin propylene glycol hydrate Capsule, 100 mg

[0129] Composition: 438.6 mg of dapagliflozin (Example 5 Part A) Stock Granulation for Capsules (22.8% w/w), filled in Gray, Opaque, Size #0 Capsule Shell was prepared.

A. Stock Granulation Composition

[0130]

Ingredient	Amount (% w/w)
Dapagliflozin or Dapagliflozin propylene glycol hydrate (Or equivalent amount of Dapagliflozin propylene glycol hydrate)	22.8
Pregelatinized Starch	15.0
Microcrystalline Cellulose	55.95
Sodium Starch Glycolate	3.0
Silicon Dioxide	2.0
Magnesium Stearate	1.25

[0131] The amount of dapagliflozin is theoretically equivalent to 81.29% of dapagliflozin propylene glycol hydrate

EP 2 139 494 B1

hydrate, either of which can be used. The actual amount of dapagliflozin propylene glycol hydrate will depend on the purity. The microcrystalline cellulose is the compensating excipient whose amount can vary depending on the actual amount of dapagliflozin propylene glycol hydrate and magnesium stearate used. The preferred amount of magnesium stearate is 1.25% (w/w). A useful range is 1.25-1.50% (w/w).

5 [0132] The stock granulation of Part 5A and the Example 5 capsules were prepared according to the following procedures.

B. Stock Granulation Procedure

10 [0133]

1. Screen silicon dioxide.
2. Mix silicon dioxide with dapagliflozin or dapagliflozin propylene glycol hydrate in a suitable blender.
3. Screen the blend from Step 2, and mix again.
- 15 4. Screen pregelatinized starch and microcrystalline cellulose, if necessary.
5. Add ingredients from Step 4 to the blend from Step 3, and mix.
6. Screen sodium starch glycolate.
7. Add ingredient from Step 6 to the blend from Step 5, and mix.
8. Screen a portion of magnesium stearate.
- 20 9. Add ingredient from Step 8 to the blend from Step 7, and mix.
10. Densify the blend from Step 9.
11. Reduce the densified blend from Step 10.
12. Screen the remaining portion of magnesium stearate.
- 25 13. Add ingredient from Step 12 to the granulation from Step 11, and mix.

C. Example 5 Product: Dapagliflozin/Dapagliflozin propylene glycol hydrate Capsule, 100 mg

[0134]

- 30 1. Fill empty capsule shells with Example 5 stock granulation for capsules (22.8 % w/w as the non-solvated form).
2. De-dust the capsules.
3. Weight sort the capsules.

35 [0135] The formed capsules of Example 3 (2.5 mg), Example 4 (10 mg), and Example 5 (100 mg) are used in treating metabolic disorders, including obesity.

Example 6 -Treatment of Metabolic Disorders

40 [0136] An oral solution (0.5 mg/mL) was prepared by dissolving dapagliflozin or dapagliflozin propylene glycol hydrate in a mixture of polyethylene glycol 400, NF and water (USP or purified water) 30:70% v/v. The oral solution was clear and colorless.

45 [0137] The glucosuric effects of dapagliflozin propylene glycol hydrate results in significant loss of calories in the urine versus a known SGLT2 inhibitor (GSK 869,682). The results of an indirect comparison of two single ascending dose studies of SGLT2 inhibitors is described. The amount of glucose excretion/day in healthy subjects taking 50, 100, 200 or 500 mg of GSK 869,682 was approximately 5g, 6g, 12g, and 16g, respectively. The amount of glucose excretion/day in healthy subjects taking 5, 20, 50 or 100 mg of dapagliflozin propylene glycol hydrate was approximately 30g, 55g, 60g, and 70g, respectively. The results of the dapagliflozin propylene glycol hydrate study was further confirmed in a 14-day multiple ascending dose phase 2a study in subjects with type 2 diabetes. Patients with type 2 diabetes were treated with placebo, 5 mg dapagliflozin propylene glycol hydrate, 25 mg dapagliflozin propylene glycol hydrate, or 100 mg dapagliflozin propylene glycol hydrate. Results from the 24-hour glucose excretion show that subjects taking 5 mg, 25 mg, and 100mg dapagliflozin propylene glycol hydrate had significantly higher urine glucose excretion compared to subjects taking placebo.

Diet-Induced Obesity in Rats

55 [0138] Obesity was induced in male Sprague-Dawley rats (mean baseline weight = 220 g) via ad libitum access to 2 diets: normal diet (Harlan Teklad rat chow; 3.5 kcal/gm, 5% vegetable fat) and high-sucrose/high-fat diet (Research Diets D12327; 4.6 kcal/gm, 40% sucrose and 40% vegetable fat). Rats under these conditions typically consume ap-

EP 2 139 494 B1

proximately 30 g/day of the high-sucrose/high-fat chow and 2 g/day of the normal Harlan Teklad rat chow. A 220-g rat given access to both diets will weigh approximately 750 g after 10 weeks.

Acute Glucosuria Study

[0139] Dapagliflozin propylene glycol hydrate (1, 5, or 10 mg/kg) or placebo (vehicle) was orally administered to diet-induced obese (DIO) rats after 24-hour baseline urine samples were collected. Urine volume and glucose concentration were used to determine total urine glucose loss over 24 hours post dose.

[0140] Total urine glucose was determined after 24 hours administration of dapagliflozin propylene glycol hydrate. Total glucose lost was calculated as volume of urine x glucose concentration. The results showed that the total amount of glucose lost over 24 hours post dose was significantly increased with increasing doses of dapagliflozin propylene glycol hydrate in a dose-dependent manner.

Chronic Weight Loss Study

[0141] DIO rats were sorted into treatment groups based on body weight, total kilocalories consumed, and body composition (via echo MRI). Dapagliflozin propylene glycol hydrate (0.5, 1, or 5 mg/kg) or placebo was orally administered to DIO rats for 28 days. To assess the importance of compensatory over-eating in drug-treated animals, a subgroup of rats that received 5 mg/kg of dapagliflozin propylene glycol hydrate was restricted to the food intake of the placebo group. Body weight and the weight of both diets were determined daily. Respiratory quotient data were obtained on days 2 and 15 of the study, echo MRI was obtained on day 22, and blood was collected for a fasting clinical chemistry tests on day 27.

[0142] Chronically administered dapagliflozin propylene glycol hydrate (administered daily over 25 days) produced significant weight loss ($p < .05$ versus vehicle) in diet-induced obese rats. If the compound-induced overeating was prevented (dapagliflozin propylene glycol hydrate mg/kg pair fed to vehicle group), then the weight loss was greater. Percent weight changes were calculated as $\text{daily weight} - \text{day 0 weight} \times 100$.

Weight Loss in Patients with Type II Diabetes

[0143] Treatment naive type II diabetes mellitus patients, $n=389$, with inadequate blood glycemic control and low mean glucosuria at baseline were given once-daily oral treatments with dapagliflozin propylene glycol hydrate (2.5, 5, 10, 20, or 50 mgs), metformin XR® (750 mg titrated to 1500 mgs), or placebo over 12-weeks.

[0144] Treatment with dapagliflozin propylene glycol hydrate resulted in consistent and sustained increases in urinary glucose excretion, rising to mean glucosuria values between 51.8 g/day to 85.0 g/day at week 12 from baseline means between 5.8 grams/day to 10.9 grams/day. Mean glucosuria with placebo and metformin both remained low, 5.7 grams/day and 5.6 grams/day respectively at week 12. A higher proportion of patients in each of the dapagliflozin propylene glycol hydrate groups achieved a 5% weight reduction over those patients taking placebo. Mean percent reductions for body weight and absolute changes in body mass index (BMI) over 12 weeks are shown in Table III.

Table III

	Dapagliflozin-PGS Dose					Placebo n=54	Metformin n=56
	2.5 mgs n=59	5 mgs n=58	10 mgs n=47	20 mgs n=59	50 mgs n=56		
Baseline weight (kg)	90	89	86	88	91	89	88
Mean reduction in weight (%)	-2.7	-2.5	-2.7	-3.4	-3.4	-1.2	-1.7
Baseline BMI (kg/m²)	31	31	30	31	32	32	32
Mean reduction in BMI	-0.9	-0.8	-0.8	-1.0	-1.1	-0.3	-0.5

Example 7 - Preparation of Dapagliflozin/Dapagliflozin Propylene glycol hydrate Tablet, 2.5 mg

[0145] Tablets containing the SGLT2 inhibitor dapagliflozin or dapagliflozin propylene glycol hydrate were prepared

EP 2 139 494 B1

in strengths of 2.5 mg (Example 7), 10 mg (Example 8) and 50 mg (Example 9) as described below.

Product: Dapagliflozin/Dapagliflozin Propylene glycol hydrate Tablet, 2.5 mg

5 A. Tablet Composition

[0146]

	Ingredient	Amount
10	Dapagliflozin propylene glycol hydrate (Or equivalent amount of Dapagliflozin)	3.08 mg
	Microcrystalline Cellulose	67.11 mg
	Anhydrous Lactose	25.00 mg
15	Crospovidone	8.75 mg
	Croscarmellose Sodium	3.75 mg
	Talc	12.50 mg
	Silicon Dioxide	2.88 mg
20	Magnesium Stearate	1.94 mg

[0147] The amount of dapagliflozin is theoretically equivalent to 81.29% of dapagliflozin propylene glycol hydrate, either of which can be used. The actual amount of dapagliflozin propylene glycol hydrate will depend on the purity. The microcrystalline cellulose is the compensating excipient whose amount can vary depending on the actual amount of dapagliflozin propylene glycol hydrate and magnesium stearate used. The target amount of magnesium stearate is 1.94 mg. An acceptable range is about 1.55 to about 2.33 mg.

[0148] The stock granulation of Part 7A and the Example 7 tablets were prepared according to the following procedures.

B. Stock Granulation Procedure

30 [0149]

1. Deaggregate dapagliflozin propylene glycol hydrate or dapagliflozin and magnesium stearate separately using a suitable screen.
- 35 2. Mix dapagliflozin propylene glycol hydrate or dapagliflozin with a portion of microcrystalline cellulose in a suitable mixer; pass through a mill; and transfer it into a suitable blender.
3. "Dry Rinse" the mixer used for mixing Step 2 with a portion of microcrystalline cellulose.
- 40 4. Add the blend from Step 3 to the blend from Step 2.
5. Mix the mixture from Step 4 with remaining microcrystalline cellulose, portion of crospovidone, portion of croscarmellose sodium, portion of silicon dioxide and Anhydrous Lactose.
- 45 6. Add talc and intragranular magnesium stearate to the mixture from Step 5 and mix.
7. Compact the powder blend from Step 6.
- 50 8. Reduce compact from Step 7 to form granules.
9. Mix the granules from Step 8 with remaining amounts of crospovidone, croscarmellose sodium and silicon dioxide.
10. Mix the granules from Step 9 with remaining amount of magnesium stearate.

55 C. Example 7 Product: Dapagliflozin/Dapagliflozin propylene glycol hydrate Tablet, 2.5 mg

[0150]

EP 2 139 494 B1

1. Setup the tableting equipment.
2. Compress the Example 7 stock granulation into tablets.

5 Example 8- Preparation of Dapagliflozin/Dapagliflozin propylene glycol hydrate Tablet, 10 mg

Product: Dapagliflozin/Dapagliflozin propylene glycol hydrate Tablet, 10 mg

10 A. Tablet Composition

[0151]

	Ingredient	Amount
15	Dapagliflozin propylene glycol hydrate (Or equivalent amount of Dapagliflozin)	12.30 mg
	Microcrystalline Cellulose	57.89 mg
	Anhydrous Lactose	25.00 mg
	Crospovidone	8.75 mg
20	Croscarmellose Sodium	3.75 mg
	Talc	12.50 mg
	Silicon Dioxide	2.88 mg
	Magnesium Stearate	1.94 mg

25 **[0152]** The amount of dapagliflozin is theoretically equivalent to 81.29% of dapagliflozin propylene glycol hydrate, either of which can be used. The actual amount of dapagliflozin propylene glycol hydrate will depend on the purity. The microcrystalline cellulose is the compensating excipient whose amount can vary depending on the actual amount of dapagliflozin propylene glycol hydrate and magnesium stearate used. The target amount of magnesium stearate is 1.94 mg. An acceptable range is about 1.55 to about 2.33 mg.

30 **[0153]** The stock granulation of Part 8A and the Example 8 tablets were prepared according to the following procedures.

B. Stock Granulation Procedure

35 [0154]

1. Deaggregate dapagliflozin propylene glycol hydrate or dapagliflozin and magnesium stearate separately using a suitable screen.
2. Mix microcrystalline cellulose, dapagliflozin propylene glycol hydrate or dapagliflozin, portion of crospovidone, portion of croscarmellose sodium, portion of silicon dioxide and anhydrous lactose in a suitable blender.
3. Add talc and intragranular magnesium stearate to the mixture from Step 2 and mix in a suitable blender.
4. Compact the powder blend from Step 3.
5. Reduce compact from Step 4 to form granules.
6. Mix the granules from Step 5 with remaining amounts of crospovidone, croscarmellose sodium and silicon dioxide.
7. Mix the granules from Step 6 with remaining amount of magnesium stearate.

C. Example 8- Product: Dapagliflozin/Dapagliflozin propylene glycol hydrate Tablet, 10 mg

50 [0155]

1. Setup the tableting equipment.
2. Compress the Example 8 stock granulation into tablets.

55

Example 9- Preparation of Dapagliflozin/Dapagliflozin propylene glycol hydrate Tablet, 50 mg

Product: Dapagliflozin/Dapagliflozin propylene glycol hydrate Tablet, 50 mg

5 **A. Tablet Composition**

[0156]

	Ingredient	Amount
10	Dapagliflozin propylene glycol hydrate (Or equivalent amount of Dapagliflozin)	61.66 mg
	Microcrystalline Cellulose	114.09 mg
	Anhydrous Lactose	62.60 mg
15	Crospovidone	21.91 mg
	Croscarmellose Sodium	9.39 mg
	Talc	31.30 mg
	Silicon Dioxide	7.20 mg
20	Magnesium Stearate	4.85 mg

[0157] The amount of dapagliflozin is theoretically equivalent to 81.29% of dapagliflozin propylene glycol hydrate, either of which can be used. The actual amount of dapagliflozin propylene glycol hydrate will depend on the purity. The microcrystalline cellulose is the compensating excipient whose amount can vary depending on the actual amount of dapagliflozin propylene glycol hydrate and magnesium stearate used. The target amount of magnesium stearate is 4.85 mg. An acceptable range is about 3.76 to about 5.95 mg.

[0158] The stock granulation of Part 9A and the Example 9 tablets were prepared according to the following procedures.

B. Stock Granulation Procedure

30 [0159]

1. Mix dapagliflozin propylene glycol hydrate or dapagliflozin, microcrystalline cellulose, anhydrous lactose, crospovidone, croscarmellose sodium, talc and silicon dioxide in a suitable blender.
- 35 2. Pass the mixture from Step 1 through a suitable mill.
3. Determine the yield from Step 1 and calculate the amount of magnesium stearate required.
- 40 4. Mix the mixture from Step 2 in a suitable blender.
5. Mix the mixture from Step 4 with magnesium stearate.
6. Dry granulate the powder blend from Step 5.
- 45 7. Size the granulation from Step 6.
8. Determine the yield based on Step 7.
- 50 9. Mix the granules from Step 8 with remaining amount of crospovidone, croscarmellose sodium and silicon dioxide.
10. Mix the granules from Step 9 with remaining amount of magnesium stearate.

C. Example 9 Product: Dapagliflozin/Dapagliflozin propylene glycol hydrate Tablet, 50 mg

55 [0160]

1. Setup the tableting equipment.

EP 2 139 494 B1

2. Compress the Example 9 stock granulation into tablets.

Example 10- Preparation of Dapagliflozin/Dapagliflozin propylene glycol hydrate Tablets (10-, 25-, and 40 mg)

Product: Dapagliflozin/Dapagliflozin propylene glycol hydrate Tablets (10 mg, 25 mg, and 40 mg)

A. Granulation composition (% w/w)

[0161]

Ingredient	Formulation % w/w
Dapagliflozin propylene glycol hydrate (Or equivalent amount of Dapagliflozin)	9.84
Microcrystalline Cellulose	63.91
Anhydrous Lactose	20
Crospovidone	4
Silicon Dioxide	1.5
Magnesium Stearate	0.75

B. Stock Granulation Procedure:

[0162]

1. Mix dapagliflozin propylene glycol hydrate or dapagliflozin with microcrystalline cellulose.
2. Pass the mixture from step 1 through a suitable mill.
3. Mix the blend from step 2 with microcrystalline cellulose, lactose anhydrous, crospovidone, and silicon dioxide.
4. Mix the blend from step 3 with magnesium stearate.
5. Dry granulate the powder blend from Step 4.
6. Size the granulation from Step 5 using appropriate sieve(s).
7. Determine the yield based on step 6.
8. Mix the granules from Step 7 with remaining amount of crospovidone, and silicon dioxide.
9. Mix granules from step 8 with remaining amount of magnesium stearate.

[0163] Tablets or capsules of various strengths (8 - 50mg) can be prepared using different weights of this granulation using tableting procedure described above.

[0164] Tableting/capsule filling operations: Same as other formulations provided herein.

[0165] Film coating: Hydroxypropylmethyl cellulose, titanium dioxide, polyethylene glycol, and colorant. Alternative film coating: Polyvinyl alcohol (PVA), titanium dioxide, polyethylene glycol, talc, and colorant.

Example 11- Preparation of Dapagliflozin/Dapagliflozin propylene glycol hydrate Tablets (1, 2.5, 5, 10 mg)

Product: Dapagliflozin/Dapagliflozin propylene glycol hydrate Tablets 1, 2.5, 5, 10 mg

A. Granulation composition

[0166]

Ingredient	1 mg Tablet	2.5 mg Tablet	5 mg Tablet	10 mg Tablet
Dapagliflozin propylene glycol hydrate (Or equivalent amount of Dapagliflozin)	1.23 mg	3.075 mg	6.15 mg	12.30 mg
Microcrystalline Cellulose	50-90 mg	60-115 mg	60-115 mg	120-230 mg
Lactose	10-30 mg	12.5-38 mg	12.5-38 mg	25-75 mg
Crospovidone	2-10 mg	2.5-13 mg	2.5-13 mg	5-25 mg

EP 2 139 494 B1

(continued)

Ingredient	1 mg Tablet	2.5 mg Tablet	5 mg Tablet	10 mg Tablet
Silicon Dioxide	0.5-4 mg	0.6-5 mg	0.6-5 mg	1-10 mg
Magnesium Stearate	0.5-2.0 mg	0.6-2.5 mg	0.6-2.5 mg	1-5 mg
Antioxidant and/or chelating agent	0-0.5 mg	0-0.6 mg	0-0.6 mg	0-1.25 mg

B. Stock granulation procedure

[0167]

1. Mix dapagliflozin propylene glycol hydrate or dapagliflozin with microcrystalline cellulose.
2. Pass the mixture from step 1 through a suitable mill.
3. Mix the blend from step 2 with microcrystalline cellulose, lactose anhydrous, croscopovidone, silicon dioxide.
4. Mix the blend from step 3 with magnesium stearate.
5. Dry granulate the powder blend from Step 4.
6. Size the granulation from Step 5 using appropriate sieve(s).
7. Determine the yield based on step 6.
8. Mix the granules from Step 7 with remaining amount of croscopovidone, and silicon dioxide.
9. Mix granules from step 8 with remaining amount of magnesium stearate.

[0168] Tablets or capsules of various strengths (1 - 20mg) can be prepared using different weights of this granulation.


[0169] Tableting/capsule filling operations: Same as other formulations provided herein.

[0170] Film coating: Polyvinyl alcohol (PVA), titanium dioxide, polyethylene glycol, talc, and colorant.

Claims

1. An immediate release pharmaceutical formulation in the form of a stock granulation for loading in capsules or forming tablets, or in the form of a capsule or a tablet containing the stock granulation, the stock granulation comprising:

a) dapagliflozin propylene glycol hydrate having the formula



- b) one or more bulking agents;
- c) one or more binders;
- d) one or more disintegrants;

- e) optionally one or more glidants and/or anti-adherents; and
- f) optionally one or more lubricants.

2. The formulation as defined in Claim 1, wherein

- a) the dapagliflozin propylene glycol hydrate is present in an amount within the range of from 0.1 to 70% by weight of stock granulation or of tablet or capsule fill;
- b) the bulking agent is present in an amount within the range of from 1 to 95% by weight of the stock granulation or of tablet or capsule fill;
- c) the binder is present in an amount of up to 20% by weight of stock granulation or of tablet or capsule fill;
- d) the disintegrant is present in an amount of up to 20% by weight of the stock granulation or of tablet or capsule fill;
- e) the glidant and/or anti-adherent, if present, is present in an amount within the range of from 0 to 20% by weight of the stock granulation or of tablet or capsule fill; and
- f) the lubricant, if present, is present in an amount within the range of from 0% to 5% by weight of tablet of the stock granulation or of tablet or capsule fill.

3. The formulation according to Claim 1, wherein the bulking agent is microcrystalline cellulose and/or lactose monohydrate, the binder is pregelatinized starch, the disintegrant is sodium starch glycolate, crospovidone, and/or croscarmellose sodium, the glidant and/or antiadherent is silicon dioxide and/or talc, and the lubricant is magnesium stearate.

4. The formulation as defined in Claim 1 in the form of a capsule filled with the stock granulation, the capsule containing an amount of the dapagliflozin propylene glycol hydrate that is equivalent to 2.5 mg or 10 mg dapagliflozin, the stock granulation comprising:

- a) the dapagliflozin propylene glycol hydrate, wherein the dapagliflozin propylene glycol hydrate is present in an amount 10.0% by weight of capsule fill;
- b) microcrystalline cellulose, wherein the microcrystalline cellulose is present in an amount of 68.75% by weight of capsule fill;
- c) pregelatinized starch, wherein the pregelatinized starch is present in an amount of 15.0% by weight of capsule fill;
- d) sodium starch glycolate, wherein the sodium starch glycolate is present in an amount of 3.0% by weight of capsule fill;
- e) silicon dioxide, wherein the silicon dioxide is present in an amount of 2.0% by weight of capsule fill; and
- f) magnesium stearate, wherein the magnesium stearate is present in an amount of 1.25% by weight of capsule fill.

5. The formulation as defined in Claim 1 in the form of a capsule filled with the stock granulation, the capsule containing an amount of the dapagliflozin propylene glycol hydrate that is equivalent to 100 mg dapagliflozin, the stock granulation comprising:

- a) the dapagliflozin propylene glycol hydrate, wherein the dapagliflozin propylene glycol hydrate is present in an amount of 22.8% by weight of capsule fill;
- b) microcrystalline cellulose, wherein the microcrystalline cellulose is present in an amount of 55.95% by weight of capsule fill;
- c) pregelatinized starch, wherein the pregelatinized starch is present in an amount of 15% by weight of capsule fill;
- d) sodium starch glycolate, wherein the sodium starch glycolate is present in an amount of 3.0% by weight of capsule fill;
- e) silicon dioxide, wherein the silicon dioxide is present in an amount of 2.0% by weight of capsule fill; and
- f) magnesium stearate, wherein the magnesium stearate is present in an amount of 1.25% by weight of capsule fill.

6. The formulation as defined in Claim 1 in the form of a tablet comprising:

- a) the dapagliflozin propylene glycol hydrate in an amount of 3.08 mg;
- b) microcrystalline cellulose, wherein the microcrystalline cellulose is present in an amount of 67.11 mg;
- c) anhydrous lactose, wherein the anhydrous lactose is present in an amount of 25.00 mg;
- d) crospovidone, wherein the crospovidone is present in an amount of 8.75 mg; and croscarmellose sodium, wherein the croscarmellose sodium is present in an amount of 3.75 mg;
- e) talc, wherein the talc is present in an amount of 12.5 mg; and silicon dioxide, wherein the silicon dioxide is

present in an amount of 2.88 mg; and

f) magnesium stearate, wherein the magnesium stearate is present in an amount of 1.94 mg.

7. The formulation as defined in Claim 1 in the form of a tablet comprising:

a) the dapagliflozin propylene glycol hydrate in an amount of 12.30 mg;

b) microcrystalline cellulose, wherein the microcrystalline cellulose is present in an amount of 57.89 mg;

c) anhydrous lactose, wherein the anhydrous lactose is present in an amount of 25.00 mg;

d) crospovidone, wherein the crospovidone is present in an amount of 8.75 mg; and croscarmellose sodium, wherein the croscarmellose sodium is present in an amount of 3.75 mg;

e) talc, wherein the talc is present in an amount of 12.50 mg; and silicon dioxide, wherein the silicon dioxide is present in an amount of 2.88 mg; and

f) magnesium stearate, wherein the magnesium stearate is present in an amount of 1.94 mg.

8. The formulation as defined in Claim 1 in the form of a tablet comprising:

a) dapagliflozin propylene glycol hydrate, wherein the dapagliflozin propylene glycol hydrate is present in an amount of 61.66 mg;

b) microcrystalline cellulose, wherein the microcrystalline cellulose is present in an amount of 114.09 mg;

c) anhydrous lactose, wherein the anhydrous lactose is present in an amount of 62.6 mg;

d) crospovidone, wherein the crospovidone is present in an amount of 21.91 mg;

e) croscarmellose sodium, wherein the croscarmellose sodium is present in an amount of 9.39 mg;

f) talc, wherein the talc is present in an amount of 31.3 mg;

g) silicon dioxide, wherein the silicon dioxide is present in an amount of 7.2 mg; and

h) magnesium stearate, wherein the magnesium stearate is present in an amount of 4.85 mg.

9. A formulation as defined in any of Claims 1 to 3, further comprising one or more therapeutic agents selected from the group consisting of anti-diabetic agents, antihyperglycemic agents, hypolipidemic/lipid lowering agents; anti-obesity agents; antihypertensive agents and appetite suppressants.

10. A formulation as defined in any of Claims 1 to 3, further comprising saxagliptin.

11. A formulation as defined in any of Claims 1 to 3, further comprising metformin.

12. The pharmaceutical formulation of any of Claims 1 to 11 for use in treating or delaying the progression or onset of Type I and Type II diabetes, impaired glucose tolerance, insulin resistance, nephropathy, retinopathy, neuropathy, cataracts, hyperglycemia, hyperinsulinemia, hypercholesterolemia, dyslipidemia, elevated blood levels of free fatty acids or glycerol, hyperlipidemia, hypertriglyceridemia, obesity, wound healing, tissue ischemia, atherosclerosis, hypertension, or Syndrome X (Metabolic Syndrome).

13. The pharmaceutical formulation of any one of Claims 1-12 for use in treating or delaying the progression or onset of type II diabetes.


Patentansprüche

1. Pharmazeutische Formulierung mit sofortiger Freisetzung in der Form eines Stammgranulats zum Laden in Kapseln oder zur Herstellung von Tabletten oder in der Form einer Kapsel oder einer Tablette, die das Stammgranulat enthält, wobei das Stammgranulat umfasst:

a) Dapagliflozin-Propylenglycolhydrat mit der Formel

5

10



15

- b) ein oder mehrere Füllstoffe;
- c) ein oder mehrere Bindemittel;
- d) ein oder mehrere Sprengmittel;
- e) gegebenenfalls ein oder mehrere Fließregulierungsmittel und/oder Antihafmittel; und
- f) gegebenenfalls ein oder mehrere Schmiermittel.

20

2. Formulierung gemäß Anspruch 1, wobei

25

- a) das Dapagliflozin-Propylenglycolhydrat in einer Menge in dem Bereich von 0,1 bis 70 Gew.-% des Stammgranulats oder der Tablette oder der Kapselfüllung vorhanden ist;
- b) der Füllstoff in einer Menge in dem Bereich von 1 bis 95 Gew.-% des Stammgranulats oder der Tablette oder der Kapselfüllung vorhanden ist;
- c) das Bindemittel in einer Menge von bis zu 20 Gew.-% des Stammgranulats oder der Tablette oder der Kapselfüllung vorhanden ist;
- d) das Sprengmittel in einer Menge von bis zu 20 Gew.-% des Stammgranulats oder der Tablette oder der Kapselfüllung vorhanden ist;
- e) das Fließregulierungsmittel und/oder Antihafmittel, falls vorhanden, in einer Menge in dem Bereich von 0 bis 20 Gew.-% des Stammgranulats oder der Tablette oder der Kapselfüllung vorhanden ist; und
- f) das Schmiermittel, falls vorhanden, in einer Menge in dem Bereich von 0 Gew.-% bis 5 Gew.-% der Tablette des Stammgranulats oder der Tablette oder der Kapselfüllung vorhanden ist.

30

35

3. Formulierung gemäß Anspruch 1, wobei der Füllstoff mikrokristalline Cellulose und/oder Lactosemonohydrat ist, das Bindemittel vorgelierte Stärke ist, das Sprengmittel Natriumstärkeglycolat, Crospovidon und/oder Croscarmellose-Natrium ist, das Fließregulierungsmittel und/oder Antihafmittel Siliciumdioxid und/oder Talkum ist und das Schmiermittel Magnesiumstearat ist.

40

4. Formulierung gemäß Anspruch 1 in der Form einer mit dem Stammgranulat gefüllten Kapsel, wobei die Kapsel eine Menge des Dapagliflozin-Propylenglycolhydrats enthält, die zu 2,5 mg oder 10 mg Dapagliflozin äquivalent ist, wobei das Stammgranulat umfasst:

45

- a) das Dapagliflozin-Propylenglycolhydrat, wobei das Dapagliflozin-Propylenglycolhydrat in einer Menge von 10 Gew.-% der Kapselfüllung vorhanden ist;
- b) mikrokristalline Cellulose, wobei die mikrokristalline Cellulose in einer Menge von 68,75 Gew.-% der Kapselfüllung vorhanden ist;
- c) vorgelierte Stärke, wobei die vorgelierte Stärke in einer Menge von 15,0 Gew.-% der Kapselfüllung vorhanden ist;
- d) Natriumstärkeglycolat, wobei das Natriumstärkeglycolat in einer Menge von 3,0 Gew.-% der Kapselfüllung vorhanden ist;
- e) Siliciumdioxid, wobei das Siliciumdioxid einer Menge von 2,0 Gew.-% der Kapselfüllung vorhanden ist; und
- f) Magnesiumstearat, wobei das Magnesiumstearat in einer Menge von 1,25 Gew.-% der Kapselfüllung vorhanden ist.

55

5. Formulierung gemäß Anspruch 1 in der Form einer mit dem Stammgranulat gefüllten Kapsel, wobei die Kapsel eine Menge des Dapagliflozin-Propylenglycolhydrats enthält, die zu 100 mg Dapagliflozin äquivalent ist, wobei das Stammgranulat umfasst:

EP 2 139 494 B1

- a) das Dapagliflozin-Propylenglycolhydrat, wobei das Dapagliflozin-Propylenglycolhydrat in einer Menge von 22,8 Gew.-% der Kapselfüllung vorhanden ist;
- b) mikrokristalline Cellulose, wobei die mikrokristalline Cellulose in einer Menge von 55,95 Gew.-% der Kapselfüllung vorhanden ist;
- 5 c) vorgelierte Stärke, wobei die vorgelierte Stärke in einer Menge von 15,0 Gew.-% der Kapselfüllung vorhanden ist;
- d) Natriumstärkeglycolat, wobei das Natriumstärkeglycolat in einer Menge von 3,0 Gew.-% der Kapselfüllung vorhanden ist;
- 10 e) Siliciumdioxid, wobei das Siliciumdioxid einer Menge von 2,0 Gew.-% der Kapselfüllung vorhanden ist; und
- f) Magnesiumstearat, wobei das Magnesiumstearat in einer Menge von 1,25 Gew.-% der Kapselfüllung vorhanden ist.

6. Formulierung gemäß Anspruch 1 in der Form einer Tablette, umfassend:

- 15 a) das Dapagliflozin-Propylenglycolhydrat in einer Menge von 3,08 mg;
- b) mikrokristalline Cellulose, wobei die mikrokristalline Cellulose in einer Menge von 67,11 mg vorhanden ist;
- c) wasserfreie Lactose, wobei die wasserfreie Lactose in einer Menge von 25,00 mg vorhanden ist;
- d) Crospovidon, wobei das Crospovidon in einer Menge von 8,75 mg vorhanden ist; und Croscarmellose-Natrium, wobei das Croscarmellose-Natrium in einer Menge von 3,75 mg vorhanden ist;
- 20 e) Talkum, wobei das Talkum einer Menge von 12,5 mg vorhanden ist; und Siliciumdioxid, wobei das Siliciumdioxid einer Menge von 2,88 mg vorhanden ist; und
- f) Magnesiumstearat, wobei das Magnesiumstearat in einer Menge von 1,94 mg vorhanden ist.

7. Formulierung gemäß Anspruch 1 in der Form einer Tablette, umfassend:

- 25 a) das Dapagliflozin-Propylenglycolhydrat in einer Menge von 12,30 mg;
- b) mikrokristalline Cellulose, wobei die mikrokristalline Cellulose in einer Menge von 57,89 mg vorhanden ist;
- c) wasserfreie Lactose, wobei die wasserfreie Lactose in einer Menge von 25,00 mg vorhanden ist;
- d) Crospovidon, wobei das Crospovidon in einer Menge von 8,75 mg vorhanden ist; und Croscarmellose-Natrium, wobei das Croscarmellose-Natrium in einer Menge von 3,75 mg vorhanden ist;
- 30 e) Talkum, wobei das Talkum einer Menge von 12,50 mg vorhanden ist; und Siliciumdioxid, wobei das Siliciumdioxid einer Menge von 2,88 mg vorhanden ist; und
- f) Magnesiumstearat, wobei das Magnesiumstearat in einer Menge von 1,94 mg vorhanden ist.

8. Formulierung gemäß Anspruch 1 in der Form einer Tablette, umfassend:

- 35 a) Dapagliflozin-Propylenglycolhydrat, wobei das Dapagliflozin-Propylenglycolhydrat in einer Menge von 61,66 mg vorhanden ist;
- b) mikrokristalline Cellulose, wobei die mikrokristalline Cellulose in einer Menge von 114,09 mg vorhanden ist;
- 40 c) wasserfreie Lactose, wobei die wasserfreie Lactose in einer Menge von 62,6 mg vorhanden ist;
- d) Crospovidon, wobei das Crospovidon in einer Menge von 21,91 mg vorhanden ist;
- e) Croscarmellose-Natrium, wobei das Croscarmellose-Natrium einer Menge von 9,39 mg vorhanden ist;
- f) Talkum, wobei das Talkum einer Menge von 31,3 mg vorhanden ist;
- 45 g) Siliciumdioxid, wobei das Siliciumdioxid einer Menge von 7,2 mg vorhanden ist; und
- h) Magnesiumstearat, wobei das Magnesiumstearat in einer Menge von 4,85 mg vorhanden ist.

9. Formulierung gemäß einem der Ansprüche 1 bis 3, ferner umfassend ein oder mehrere therapeutische Mittel ausgewählt aus der Gruppe bestehend aus antidiabetischen Mitteln, antihyperglykämischen Mitteln, hypolipidämischen/lipidsenkenden Mitteln; Antiadipositasmitteln; blutdrucksenkenden Mitteln und Appetithemmern.

50

10. Formulierung gemäß einem der Ansprüche 1 bis 3, ferner umfassend Saxagliptin.

11. Formulierung gemäß einem der Ansprüche 1 bis 3, ferner umfassend Metformin.

- 55 12. Pharmazeutische Formulierung gemäß einem der Ansprüche 1 bis 11 zur Verwendung bei der Behandlung oder Verzögerung der Progression oder des Ausbruchs von Typ-I- und Typ-II-Diabetes, beeinträchtigter Glucosetoleranz, Insulinresistenz, Nephropathie, Retinopathie, Neuropathie, Katarakten, Hyperglykämie, Hyperinsulinämie, Hypercholesterolämie, Dyslipidämie, erhöhten Blutspiegeln von freien Fettsäuren oder Glycerol, Hyperlipidämie, Hyper-


triglyceridämie, Adipositas, Wundheilung, Gewebeischämie, Atherosklerose, Bluthochdruck oder Syndrom X (metabolisches Syndrom).

13. Pharmazeutische Formulierung gemäß einem der Ansprüche 1 bis 12 zur Verwendung bei der Behandlung oder Verzögerung der Progression oder des Ausbruchs von Typ-II-Diabetes.

Revendications

1. Formulation pharmaceutique à libération immédiate sous la forme d'une granulation mère pour chargement dans des capsules ou formation de comprimés, ou sous la forme d'une capsule ou d'un comprimé contenant la granulation mère, la granulation mère comprenant :

a) du dapagliflozine-propylène glycol hydraté ayant la formule



b) un ou plusieurs agents de charge ;

c) un ou plusieurs liants ;

d) un ou plusieurs délitants ;

e) facultativement un ou plusieurs agents glissants et/ou agents antiadhérents ; et

f) facultativement un ou plusieurs lubrifiants.

2. Formulation telle que définie dans la revendication 1, dans laquelle

a) le dapagliflozine-propylène glycol hydraté est présent en une quantité dans la plage de 0,1 à 70 % en poids de granulation mère ou de remplissage de comprimé ou de capsule ;

b) l'agent de charge est présent en une quantité dans la plage de 1 à 95 % en poids de la granulation mère ou de remplissage de comprimé ou de capsule ;

c) le liant est présent en une quantité allant jusqu'à 20 % en poids de granulation mère ou de remplissage de comprimé ou de capsule ;

d) le délitant est présent en une quantité allant jusqu'à 20 % en poids de la granulation mère ou de remplissage de comprimé ou de capsule ;

e) l'agent glissant et/ou un agent antiadhérent, s'il est présent, est présent en une quantité dans la plage de 0 à 20 % en poids de la granulation mère ou de remplissage de comprimé ou de capsule ; et

f) le lubrifiant, s'il est présent, est présent en une quantité dans la plage de 0 % à 5 % en poids de comprimé de la granulation mère ou de remplissage de comprimé ou de capsule.

3. Formulation selon la revendication 1, dans laquelle l'agent de charge est la cellulose microcristalline et/ou le lactose monohydraté, le liant est l'amidon prégélatinisé, le délitant est le glycolate d'amidon sodique, la crospovidone, et/ou la croscarmellose sodique, l'agent glissant et/ou l'agent antiadhérent est le dioxyde de silicium et/ou le talc, et le lubrifiant est le stéarate de magnésium.

4. Formulation telle que définie dans la revendication 1 sous la forme d'une capsule remplie avec la granulation mère, la capsule contenant une quantité du dapagliflozine-propylène glycol hydraté qui est équivalente à 2,5 mg ou 10 mg de dapagliflozine, la granulation mère comprenant :

a) du dapagliflozine-propylène glycol hydraté, le dapagliflozine-propylène glycol hydraté étant présent en une quantité de 10,0 % en poids de remplissage de capsule ;

EP 2 139 494 B1

- b) de la cellulose microcristalline, la cellulose microcristalline étant présente en une quantité de 68,75 % en poids de remplissage de capsule ;
c) de l'amidon prégélatinisé, l'amidon prégélatinisé étant présent en une quantité de 15,0 % en poids de remplissage de capsule ;
d) du glycolate d'amidon sodique, le glycolate d'amidon sodique étant présent en une quantité de 3,0 % en poids de remplissage de capsule ;
e) du dioxyde de silicium, le dioxyde de silicium étant présent en une quantité de 2,0 % en poids de remplissage de capsule ; et
f) du stéarate de magnésium, le stéarate de magnésium étant présent en une quantité de 1,25 % en poids de remplissage de capsule.

5. Formulation telle que définie dans la revendication 1 sous la forme d'une capsule remplie avec la granulation mère, la capsule contenant une quantité du dapagliflozine-propylène glycol hydraté qui est équivalente à 100 mg de dapagliflozine, la granulation mère comprenant :

- a) du dapagliflozine-propylène glycol hydraté, le dapagliflozine-propylène glycol hydraté étant présent en une quantité de 22,8 % en poids de remplissage de capsule ;
b) de la cellulose microcristalline, la cellulose microcristalline étant présente en une quantité de 55,95 % en poids de remplissage de capsule ;
c) de l'amidon prégélatinisé, l'amidon prégélatinisé étant présent en une quantité de 15 % en poids de remplissage de capsule ;
d) du glycolate d'amidon sodique, le glycolate d'amidon sodique étant présent en une quantité de 3,0 % en poids de remplissage de capsule ;
e) du dioxyde de silicium, le dioxyde de silicium étant présent en une quantité de 2,0 % en poids de remplissage de capsule ; et
f) du stéarate de magnésium, le stéarate de magnésium étant présent en une quantité de 1,25 % en poids de remplissage de capsule.

6. Formulation telle que définie dans la revendication 1 sous la forme d'un comprimé comprenant :

- a) du dapagliflozine-propylène glycol hydraté en une quantité de 3,08 mg ;
b) de la cellulose microcristalline, la cellulose microcristalline étant présente en une quantité de 67,11 mg ;
c) du lactose anhydre, le lactose anhydre étant présent en une quantité de 25,00 mg ;
d) de la crospovidone, la crospovidone étant présente en une quantité de 8,75 mg ; et de la croscarmellose sodique, la croscarmellose sodique étant présente en une quantité de 3,75 mg ;
e) du talc, le talc étant présent en une quantité de 12,5 mg ; et du dioxyde de silicium, le dioxyde de silicium étant présent en une quantité de 2,88 mg ; et
f) du stéarate de magnésium, le stéarate de magnésium étant présent en une quantité de 1,94 mg.

7. Formulation telle que définie dans la revendication 1 sous la forme d'un comprimé comprenant :

- a) du dapagliflozine-propylène glycol hydraté en une quantité de 12,30 mg ;
b) de la cellulose microcristalline, la cellulose microcristalline étant présente en une quantité de 57,89 mg ;
c) du lactose anhydre, dans laquelle le lactose anhydre est présent en une quantité de 25,00 mg ;
d) de la crospovidone, la crospovidone étant présente en une quantité de 8,75 mg ; et de la croscarmellose sodique, la croscarmellose sodique étant présente en une quantité de 3,75 mg ;
e) du talc, le talc étant présent en une quantité de 12,50 mg ; et du dioxyde de silicium, le dioxyde de silicium étant présent en une quantité de 2,88 mg ; et
f) du stéarate de magnésium, le stéarate de magnésium étant présent en une quantité de 1,94 mg.

8. Formulation telle que définie dans la revendication 1 sous la forme d'un comprimé comprenant :

- a) du dapagliflozine-propylène glycol hydraté, le dapagliflozine-propylène glycol hydraté étant présent en une quantité de 61,66 mg ;
b) de la cellulose microcristalline, la cellulose microcristalline étant présente en une quantité de 114,09 mg ;
c) du lactose anhydre, le lactose anhydre étant présent en une quantité de 62,6 mg ;
d) de la crospovidone, la crospovidone étant présente en une quantité de 21,91 mg ;
e) de la croscarmellose sodique, la croscarmellose sodique étant présente en une quantité de 9,39 mg ;

EP 2 139 494 B1

f) du talc, le talc étant présent en une quantité de 31,3 mg ;

g) du dioxyde de silicium, le dioxyde de silicium étant présent en une quantité de 7,2 mg ; et

h) du stéarate de magnésium, le stéarate de magnésium étant présent en une quantité de 4,85 mg.

- 5
- 9.** Formulation telle que définie dans l'une quelconque des revendications 1 à 3, comprenant en outre un ou plusieurs agents thérapeutiques choisis dans le groupe constitué d'agents antidiabétiques, agents antihyperglycémiques, agents hypolipidémiques/hypolipidémiants ; agents antiobésité ; agents antihypertenseurs et agents anorexigènes.
- 10
- 10.** Formulation telle que définie dans l'une quelconque des revendications 1 à 3, comprenant en outre de la saxagliptine.
- 11.** Formulation telle que définie dans l'une quelconque des revendications 1 à 3, comprenant en outre de la metformine.
- 15
- 12.** Formulation pharmaceutique selon l'une quelconque des revendications 1 à 11 pour utilisation dans le traitement ou le retardement de la progression ou l'apparition du diabète de type I et de type II, d'une tolérance au glucose diminuée, de l'insulinorésistance, d'une néphropathie, d'une rétinopathie, d'une neuropathie, de cataractes, de l'hyperglycémie, de l'hyperinsulinémie, de l'hypercholestérolémie, de la dyslipidémie, d'une élévation des taux sanguins d'acides gras libres ou de glycérol, de l'hyperlipidémie, de l'hypertriglycéridémie, de l'obésité, de la cicatrisation des plaies, d'une ischémie tissulaire, de l'athérosclérose, de l'hypertension ou du syndrome X (syndrome métabolique).
- 20
- 13.** Formulation pharmaceutique selon l'une quelconque des revendications 1 à 12 pour utilisation dans le traitement ou le retardement de la progression ou l'apparition du diabète de type II.
- 25
- 30
- 35
- 40
- 45
- 50
- 55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 6515117 B [0003] [0007] [0108]
- US 76548107 [0004] [0006] [0008] [0009] [0057]
- US 20080004336 A1 [0004] [0006] [0008] [0009] [0057]
- US 60817118 B [0008] [0009] [0057]
- US 74507503 [0057] [0061]
- US 5594016 A [0079]
- WO 0121602 A [0080]
- US 6414002 B [0080]
- US 6653314 B [0080]
- US 39105399 [0081]
- US 6548529 B [0081]
- WO 9938501 A [0082]
- WO 9946272 A [0082]
- WO 9967279 A [0082]
- WO 9967278 A [0082]
- WO 9961431 A [0082]
- US 899641 [0082]
- WO 0168603 A [0082]
- US 6395767 B [0082]
- US 5614492 A [0084]
- WO 03033671 A [0084]
- US 5595872 A [0086]
- US 5739135 A [0086]
- US 5712279 A [0086]
- US 5760246 A [0086]
- US 5827875 A [0086]
- US 5885983 A [0086]
- US 5962440 A [0086]
- US 3983140 A [0087]
- US 4231938 A [0087]
- US 4346227 A [0087]
- US 4448784 A [0087]
- US 4450171 A [0087]
- US 5354772 A [0087]
- US 5006530 A [0087]
- US 5177080 A [0087]
- US 4681893 A [0087]
- US 5273995 A [0087]
- US 5385929 A [0087]
- US 5686104 A [0087]
- US 5011930 A [0087]
- US 5260440 A [0087]
- US 5753675 A [0087]
- US 4613610 A [0087]
- WO 8603488 A [0087]
- US 4647576 A [0087]
- WO 8607054 A [0087]
- FR 2596393 [0087]
- EP 0221025 A [0087]
- US 4686237 A [0087]
- US 4499289 A [0087]
- EP 0142146 A2 [0087]
- US 5506219 A [0087]
- US 5691322 A [0087]
- GB 2205837 A [0087]
- US 5712396 A [0088]
- US 4871721 A [0088]
- US 4924024 A [0088]
- US 3674836 A [0089]
- US 4759923 A [0089]
- US 4027009 A [0089]
- WO 9712615 A [0093]
- WO 9712613 A [0093]
- WO 9638144 A [0093]
- US 5612359 A [0094]
- US 6043265 A [0094]
- WO 0001389 A [0094]
- US 5541204 A [0096]
- US 5770615 A [0096]
- US 5491134 A [0096]
- US 5776983 A [0096]
- US 5488064 A [0096]
- WO 9721993 U [0099]
- WO 9900353 A [0099]
- WO 00039077 A [0099]
- US 6414126 B [0106]
- US 5515117 A [0106]

Non-patent literature cited in the description

- **RYAN et al.** HK-2: an immortalized proximal tubule epithelial cell line from normal adult human kidney. *Kidney International*, 1994, vol. 45, 48-57 [0063]
- **JOHANSSON.** *J. Clin. Endocrinol. Metab.*, 1997, vol. 82, 727-34 [0065]
- **MURAKAMI et al.** A Novel Insulin Sensitizer Acts As a Coligand for Peroxisome Proliferation - Activated Receptor Alpha (PPAR alpha) and PPAR gamma. Effect on PPAR alpha Activation on Abnormal Lipid Metabolism in Liver of Zucker Fatty Rats. *Diabetes*, 1998, vol. 47, 1841-1847 [0080]

- **HUGHES et al.** *Biochemistry*, 1999, vol. 38 (36), 11597-11603 [0082]
- **YAMADA et al.** *Bioorg. & Med. Chem. Lett.*, 1998, vol. 8, 1537-1540 [0082]
- **ASHWORTH et al.** *Bioorg. & Med. Chem. Lett.*, 1996, vol. 6 (22), 1163-1166, 2745-2748 [0082]
- **BILLER et al.** *J. Med. Chem.*, 1988, vol. 31 (10), 1869-1871 [0088]
- **BILLER, S.A. ; NEUENSCHWANDER, K. ; PON-PIPOM, M.M. ; POULTER, C.D.** *Current Pharmaceutical Design*, 1996, vol. 2, 1-40 [0088]
- **P. ORTIZ DE MONTELLANO et al.** *J. Med. Chem.*, 1977, vol. 20, 243-249 [0088]
- **COREY ; VOLANTE.** *J. Am. Chem. Soc.*, 1976, vol. 98, 1291-1293 [0088]
- **MCCLARD, R.W. et al.** *J.A.C.S.*, 1987, vol. 109, 5544 [0088]
- **CAPSON, T.L.** Dept. Med. Chem. U of Utah, Abstract, Table of Contents. *PhD dissertation*, June 1987, 16, , 17, , 40-43, 48-51 [0088]
- *Drugs of the Future*, 1999, vol. 24, 9-15 [0090]
- **NICOLOSI et al.** The ACAT inhibitor, CI-1011 is effective in the prevention and regression of aortic fatty streak area in hamsters. *Atherosclerosis (Shannon, Irel)*, 1998, vol. 137 (1), 77-85 [0090]
- **GHISELLI ; GIANCARLO.** The pharmacological profile of FCE 27677: a novel ACAT inhibitor with potent hypolipidemic activity mediated by selective suppression of the hepatic secretion of ApoB100-containing lipoprotein. *Cardiovasc. Drug Rev.*, 1998, vol. 16 (1), 16-30 [0090]
- **SMITH, C. et al.** RP 73163: a bioavailable alkylsulfanyl-diphenylimidazole ACAT inhibitor. *Bioorg. Med. Chem. Lett.*, 1996, vol. 6 (1), 47-50 [0090]
- ACAT inhibitors: physiologic mechanisms for hypolipidemic and anti-atherosclerotic activities in experimental animals. **KRAUSE et al.** *Inflammation: Mediators Pathways*. Publisher: CRC, 1995, 173-98 [0090]
- **SLISKOVIC et al.** ACAT inhibitors: potential anti-atherosclerotic agents. *Curr. Med. Chem.*, 1994, vol. 1 (3), 204-25 [0090]
- **STOUT et al.** Inhibitors of acyl-CoA:cholesterol O-acyl transferase (ACAT) as hypocholesterolemic agents. The first water-soluble ACAT inhibitor with lipid-regulating activity. Inhibitors of acyl-CoA:cholesterol acyltransferase (ACAT). Development of a series of substituted N-phenyl-N'-[(1-phenylcyclopentyl)methyl]ureas with enhanced hypocholesterolemic activity. *Chemtracts: Org. Chem.*, 1995, vol. 8 (6), 359-62 [0090]
- *Atherosclerosis*, 1995, vol. 115, 45-63 [0091]
- *J. Med. Chem.*, 1998, vol. 41, 973 [0091]
- *Drugs of the Future*, 1999, vol. 24, 425-430 [0092]
- **SENDOBRY et al.** Attenuation of diet-induced atherosclerosis in rabbits with a highly selective 15-lipoxygenase inhibitor lacking significant antioxidant properties. *Brit. J. Pharmacology*, 1997, vol. 120, 1199-1206 [0093]
- **COMICELLI et al.** 15-Lipoxygenase and its Inhibition: A Novel Therapeutic Target for Vascular Disease. *Current Pharmaceutical Design*, 1999, vol. 5, 11-20 [0093]