



(19) Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number : **0 375 265 B1**

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication of patent specification : **05.05.93 Bulletin 93/18**

(51) Int. Cl.⁵ : **C07D 275/02**

(21) Application number : **89313046.8**

(22) Date of filing : **13.12.89**

(54) Purification of isothiazolones.

(30) Priority : **22.12.88 US 289069**

(73) Proprietor : **ROHM AND HAAS COMPANY**
Independence Mall West
Philadelphia Pennsylvania 19105 (US)

(43) Date of publication of application :
27.06.90 Bulletin 90/26

(72) Inventor : **Petigara, Ramesh Balubhai**
2185 Stewart Drive
Hatfield, Pennsylvania 19440 (US)
Inventor : **Sharp, Douglas Norman**
2248 Rittenhouse Square
Bensalem, Pennsylvania 19020 (US)

(45) Publication of the grant of the patent :
05.05.93 Bulletin 93/18

(84) Designated Contracting States :
AT BE CH DE ES FR GB GR IT LI LU NL SE

(74) Representative : **Angell, David Whilton et al**
ROHM AND HAAS (UK) LTD. European
Operations Patent Department Lennig House
2 Mason's Avenue
Croydon CR9 3NB (GB)


(56) References cited :
EP-A- 0 095 907
EP-A- 0 271 761
US-A- 3 761 488

EP 0 375 265 B1

Note : Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Description

This invention is concerned with the preparation of isothiazolones, more particularly involving the purification of complex reaction mixtures to obtain high purity (>98%) isothiazolones. The invention involves first forming and isolating readily filterable high purity crystalline salts of isothiazolones of the formula I:



wherein R is hydrogen or methyl and Y is a linear or branched alkyl, cycloalkyl or substituted cycloalkyl of from 20 3 to 10 carbon atoms, followed by hydrolysis with water to the desired isothiazolones which are well known biocides.


25 A process for preparing isothiazolone reaction mixtures is described in U.S.-A-3,761,488, where a N,N'-disubstituted 3,3'-dithiodipropionamide is halogenated and cyclized with such reactants as chlorine or sulfonyl chloride to a crude reaction mixture containing 2-substituted-3-isothiazolone hydrohalide salt in solution. The halogenation reactant is maintained at about 3 equivalents of halogen per equivalent of dithiodiamide to obtain a complete reaction which unfortunately contains small to significant amount of unwanted products such as 5-halo-2-substituted isothiazolone, 4-halo-2-substituted isothiazolone, 4,5-dihalo- 2-substituted isothiazolone, 3-halo-N-substituted alkyl propionamide, and other amide impurities. From the reaction mixture the hydrohalide salt is crystallized, isolated and then hydrolyzed with water to the desired isothiazolone of high purity (>98%).

30 An alternate route to isothiazolones is disclosed in EP-A-95,907, wherein a 3-mercaptopropionamide is treated with at least about 2 equivalents of chlorine or other halogenating agent to form and isolate the isothiazolone hydrohalide salt, followed by its hydrolysis to obtain the desired isothiazolone.


Both of the above processes are also useful in preparing the 4-methyl-2-substituted isothiazolones from the corresponding mercaptoisobutyramide or dithiodiisobutyramide intermediate.

35 However, in these processes some amounts of the above mentioned impurities are formed. For improvement of product quality, safety reasons, and for many end uses, it is desirable to remove such impurities.

The materials to be chlorinated and cyclized to form a reaction mixture containing the corresponding isothiazolone hydrochloride have the following formulae:



(II) and



(III)

Some solvents used to conduct the chlorination/cyclization have been lower alkyl esters, such as ethyl acetate,

butyl acetate, and the like. Upon removal of the excess hydrogen chloride, the isothiazolone salt precipitates. There are disadvantages to using such solvents because:


- a. certain impurities precipitate with the desired product and can be removed only with difficulty;
- b. the solvent becomes hydrolyzed during the hydrolysis of the isothiazolone salt, leading to a loss of solvent on recovery or introduction of the hydrolyzed solvent as an impurity in the desired product;
- c. upon recycling the solvent or its hydrolysed products react during the chlorination/cyclization step to lower product yield and have deleterious effects; or
- d. the desired hydrochloride salt may be more soluble in the ester solvent to cause significant yield losses.

We have now surprisingly found a suitable solvent for the chlorination reaction and subsequent isothiazolone purification which is sufficiently inert to chlorination under the reaction conditions, sufficiently inert to the hydrolysis reaction conditions employed to hydrolyse the salt, sufficiently volatile to be removed completely from the isothiazolone when desired, and possesses such solubility characteristics for the isothiazolone salt as to allow quantitative separation of pure salt after removal of excess hydrogen chloride. We have further found processing conditions for the crystallization which can produce pure clustered crystal aggregates of a size and shape suitable for easy filtration and washing, and which avoid the formation of tiny secondary nucleation crystals which can clog filtration apparatus and are difficult to free from mother liquor containing reaction impurities.

EP-A-95,907 teaches that other solvents may be used, such as toluene or perchloroethylene. These solvents are reactive under halogenation conditions or are toxic or create environmental problems.


EP-A-271,761, teaches the purification of crude isothiazolones, by isolation of the isothiazolone (not the hydrochloride) and process impurities (in an unspecified procedure), dissolving this material in an organic solvent immiscible with water, treating this solution with a strong acid, such as hydrogen chloride, to form the salt, followed by separation and hydrolysis of the salt in the presence of the organic solvent. Monochlorobenzene and chlorinated hydrocarbons are mentioned as solvents, but are not used. There is no teaching in this reference which would suggest the process of the present invention to the skilled man.

Accordingly, this invention provides a process for preparing substantially pure (over 98%) isothiazolone by preparing crystalline isothiazolones monohydrochloride salts of the formula:




wherein R is hydrogen or methyl and Y is a linear or branched alkyl group, or an optionally substituted cycloalkyl group, of from 3 to 10 carbon atoms, by the chlorination of compounds of the formula:

45



50 wherein X is - SH or

55



characterised in that:

- a) the chlorination is conducted in a chlorinated hydrocarbon solvent so as to afford a crude reaction mix-

ture containing isothiazolone hydrochloride and excess hydrogen chloride in solution;

5 b) the amount of hydrogen chloride is reduced to below about 120% of that needed to form the salt;

c) a seeded solution is formed;

d) the seeded solution is cooled under temperature/time conditions to form filterable crystals of the salt;

e) the pure crystals are collected.

When Y is a substituted cycloalkyl group the substituents may be halo, C₁-C₄ alkyl or C₁-C₄ alkoxy subject to the upper carbon atom limit of 10.

It is desired to employ a solvent which is substantially or completely unreactive with chlorine during the chlorination/cyclization reaction and is further substantially or completely unreactive with water during the conversion of the isothiazolone hydrochloride to the free isothiazolone. Appropriate solvents satisfying these parameters are chlorinated hydrocarbons, either aromatic or aliphatic such as monochlorobenzene, chlorotoluene, dichlorobenzene, ethylene dichloride or perchloro ethylene and the like. Since an appropriate solvent is selected based on such considerations as least toxicity, moderate volatility, low cost and least environmental problems, monochlorobenzene is a preferred solvent.

We found, however, that monochlorobenzene has a serious deficiency in that the reaction mixture, on degassing, gave the isothiazolone hydrochloride in the form of thin, long needles forming a paste which is impossible to collect. Monochlorobenzene is trapped within the fine crystals, and any impurities dissolved in the monochlorobenzene are difficult to remove. This deficiency is also noted with other chlorinated solvents, such as perchloroethylene.

Thus, to be able to utilize the chlorinated hydrocarbons as a solvent in the chlorination and as a crystallizing media for the isothiazolone salt, we had to overcome this problem by discovering a means for drastically altering the crystal size and shape, so as to produce easily filterable high purity large crystals.

We have found that the isothiazolone hydrochloride salt is much more soluble in the reaction mixture containing an excess of hydrogen chloride generated during the chlorination in the chlorinated hydrocarbons than it is in monochlorobenzene containing little or no hydrogen chloride. In other words, we found that the solubility of the monohydrochloride salts of the 2-substituted isothiazolones is significantly decreased as the concentration of hydrogen chloride in the reaction mixture decreases. Elimination of the large excess of hydrogen chloride from the chlorination mixture enables the formation of a seeded solution of the isothiazolone hydrochloride salt either in the mixture, or the creation of conditions where separately added seed crystals of the salt will neither redissolve nor cause further massive precipitation of small crystals due to secondary nucleation.

In the initial step of the present process, the appropriate compound of formula II or III is reacted with chlorine in the presence of the chlorinated hydrocarbon solvent. Even under the controlled chlorination conditions small to appreciable amounts of 4-chloro-, 5-chloro- and 4,5-dichloroisothiazolones and other by-products are formed.

Other impurities which may be present in the reaction mixture include unreacted compound III and 3-chloro-N-(alkyl or cycloalkyl)propionamide (or isobutyramide). Formation of such by-products detracts from the yield of non-chlorinated isothiazolone salt, and such impurities will remain in the solvent mixture.

The stoichiometry requires that 2 moles of chlorine react with compound II to form the salt and 3 excess moles of hydrogen chloride, and that 3 moles of chlorine react with compound III to form the salt and 5 excess moles of hydrogen chloride. Thus the solvent for the formed isothiazolone is a mixture of appropriate inert solvent, such as chlorobenzene, with excess hydrogen chloride and various impurities as are mentioned generated in the reaction. In such a mixture, the desired isothiazolone salt is fully soluble.

The removal of hydrogen chloride, i.e. degassing, should be carried out without introducing any new chemicals. Thus, it is best removed by an appropriate physical process, such as evaporation. Preferred methods are sparging of the mixture with an appropriate inert gas, such as nitrogen, which will not further react with the isothiazolone or hydrogen chloride, or by applying sufficient vacuum to cause the hydrogen chloride and some solvent to volatilize. The degassing may be carried out at temperatures within a wide range; preferred is a temperature in the range of from about 46° to about 60°C and most preferred is from about 46° to 52°C.

A combination of sparge and vacuum may also be employed. The excess hydrogen chloride removed from the reaction mixture may be neutralized with base or collected in other ways to avoid corrosion or contamination. Some solvent is also removed at this point. If necessary additional solvent may be added to the reaction mixture to maintain a particular solids level.

The level of remaining hydrogen chloride can be measured by titration. The level, which prior to the sparging, vacuum, or other reduction operation was about 220% of that needed to form the salt, should be reduced to below about 120% of that needed to form the salt; that is to say the reaction system should contain less than 0.2 moles of hydrogen chloride beyond that present in 1 mol of the isothiazolone monohydrochloride salt.

At this point, with the temperature preferably high enough to maintain most, if not all, of the salt in solution, and with the concentration of the salt near the concentration at which crystallization could occur, seed crystals are formed or added. If added from an external source, such seed crystals are preferably small, such as below about 100 μm by 30 μm by 30 μm , and may contain entrapped solvent. The amount of added seed crystals is preferably from about 3 to about 6 weight percent of of the batch, although somewhat smaller or larger amounts may be used.

It is preferred to form the seed crystals internally in the degassed reaction mixture by control of temperature and extent of removal of hydrogen chloride. The initial crystals formed are small, such as about 100 μm by 30 μm by 30 μm or smaller, and will consist of from about 25 to about 60 % by weight of the total salt in the partially precipitated mixture. Preferred is from about 25 to about 35% present as seed crystals; this level may be achieved by dissolving some of the initially precipitated small crystals by upward temperature adjustment.

It is well known in crystallization that creation of or introduction of seed crystals can lead to rapid crystallization of the whole mass of dissolved crystallizable material. In the present case, the rapid crystallization can be eliminated by careful control of temperature, optionally including a step where the temperature is raised slightly or cycled to adjust the size and amount of seed crystal present. Then the initial seed crystals are allowed to "ripen", i.e., grow slightly larger, but without significant crystallization of new material from the dissolved phase. The "ripening" phase produces modified seed crystals which are more useful in controlling the subsequent crystallization.

Once appropriate "ripened" seed crystals are available, crystallization to form the filterable large clustered crystals is best conducted by cooling of the reaction medium. The temperature should preferably be lowered at a slow rate, preferably about 1 degree C. or less per 5 minutes. Also it is preferred to cool the batch; initially temperatures of from about 45° to about 55°C, preferably about 52° to about 53°C, through the first 10-degree to 15-degree drop (to about 30 to about 45, preferably to about 42 to about 43 °) at a slower rate than for subsequent cooling; a rate of about 1°C per 15 minutes has been found most effective. Further cooling to room temperature and below may be effected at a somewhat faster rate, but in no case above about 1°C per 5 minutes. Intermediate rates may also be employed during the mid-portion of the cooling cycle. The cooling rate selected is such that it avoids any significant supersaturation which may lead to massive secondary nucleation passing tiny new crystals.

A final temperature of room temperature or below, preferably about 20°C, is utilized.

The clustered crystals formed by the present process may be collected by conventional means, including vacuum draining, mechanically pressing the filter cake, use of a pressure column, and the like. The crystals may be washed with monochlorobenzene to remove residual mother liquor containing soluble impurities; if desired, other appropriate chlorinated solvents which will dissolve the impurities may be used in the wash step.

The isothiazolone salt can be converted to the free isothiazolone by hydrolysis with water by known means, the water phase containing HCl is separated, the organic phase containing product is preferably re-washed with water, the water phase again separated. From the product layer the residual monochlorobenzene or other appropriate solvent can be removed by application of vacuum and distillation or stripping, such as steam stripping.

As noted earlier, the present process is mainly useful for the purification of isothiazolones wherein the 2-substituent radical is an alkyl group, linear or branched, of from three to ten carbon atoms. When the alkyl group is less than three carbon atoms, the solubility of the isothiazolone salt is so small in the organic phase that the present process is not effective. If the substituent group is higher in molecular weight than those shown here, the solubility of the hydrochloride salt will be lessened in monochlorobenzene to the extent that the present process cannot be effectively practiced. Within the present series of isothiazolones, some routine trial and error experimentation may be necessary, such as adjustment of solids, extent of removal of excess hydrogen chloride, and adjustment of temperature, to produce the best results in terms of extent of aggregated crystals and yields of recovered product.

The preferred isothiazolones or isothiazolone hydrochloride salts for purification by the process of the present invention are those where R is H and Y is (C₃-C₁₀) alkyl. Especially preferred is *n*-octylisothiazolone, wherein R is H and Y is *n*-octyl.

The process is not suitable for isothiazolones substituted with chlorine in the four and/or five position, as the hydrochloride salts are too soluble in the appropriate solvents. The process is less suitable for purification of hydrohalide salts other than the hydrochloride, mainly for economic reasons.

The purification process described herein may also be applied to purification of isothiazolones made by other processes, such as by processes taught by Virgilio et al. in U.S.-A-4,281,136 and 4,508,136, wherein 2-alkyl-3-haloisothiazolium salts may be converted to a mixture of the corresponding isothiazolone with by-products related to the salts.

The isothiazolones are known to have many preservative uses including deodorizers, preservative for liq-

uid and powder soaps, food processing, dairy chemicals, food preservatives, animal food preservatives, preservatives for wood, lazures and paints, preservatives for paint films and plastics, hospital and medical anti-septics, metal working fluids, cooling water, air washers, paper treatment, petroleum products, adhesives, pigment slurries, latexes, leather and hide treatment, agricultural formulations, mining, nonwoven fabrics, petroleum storage, rubber, sugar processing, tobacco, cosmetics, chemical toilets, household laundry, detergent, and dishwashing products, diesel fuel additives, waxes and polishes and many other applications where water and organic materials come in contact under conditions which allow the growth of undesired microorganisms.

5 Isothiazolones also are used as disinfectants, in oil field water treatment, as watercooling system micro-biocides, as preservatives for aqueous dispersions or organic polymers, as wood pulp white water slimicides, 10 as cosmetic preservatives, as cutting oil, jet fuel, and heating oil preservatives, and the like. They are also useful in adhesives, agricultural chemical preservation, air washing devices, alcohol stabilization, carpet backing, caulk and sealants, ceramics, cleaners, cement modifiers, diesel and other fuels, electrocoating systems, 15 electronic circuitry, commercial enzymes, fabric softeners, feed preservation, fibers, printing, household and industrial cleaners, inks, laundered goods, laundry wash water, marine antifoulants, such as ship bottoms, propellers, fishnets, and the like, medical devices, membranes, odor control, pasteurization baths, photographic emulsions, pharmaceutical and therapeutic uses, preservation of reagent chemicals, sanitizers, swimming pools, textile manufacture and uses, toiletries, waste treatment, water purification, and the like.

15 Solutions of isothiazolones are also usefully applied to a solid substrate, such as fabric, leather, or wood, as a preservative, or admixed with plastics.

20 The isothiazolones with which this invention is primarily concerned, having organic solubility and limited water solubility, have especial utility in preservatives for starch and dextrin pastes used on wallpaper, as preservatives for plastics, such as plasticized poly(vinyl chloride), and as mildewcides for painted surfaces where the paint is applied as a water-based latex.

25 The isothiazolones prepared by the process of the present invention may also be useful in the various applications mentioned herein when combined with other isothiazolones, such as 5-chloro-2-methylisothiazolone, 2-methylisothiazolone, and mixtures thereof. The present isothiazolones may also be useful in combination with other biocides.

30 The present invention is especially useful in the preparation of 2-n-octylisothiazolone which offers superior resistance to mildew growth, especially on latex paints, but is also effective in solvent based paints. Mildew in paints is noted as disfigurement of exterior house paints and other coatings by a superficial fungal dark-colored growth caused mainly by *Aureobasidium pullulans*, *cladosporium*, and *alertia pullulans*. The n-octylisothiazolone is effective in killing the organism and so preventing discoloration and in killing bacteria responsible for paint spoilage during storage.

35 The invention will now be further illustrated by the following Examples some of which are for comparison purposes only and in which all invention and percentages are by weight unless otherwise specified.

EXAMPLE 1 - 2-n-Octylisothiazolone, hydrochloride salt

40 This example illustrates the formation of clustered readily filterable crystals of high purity hydrochloride salt of 2-n-octyl isothiazolone by controlled removal of excess hydrogen chloride, generation of seed crystals, temperature cycling of seed crystals, and controlled cooling of the seeded mixture.

45 In a 1325 l. reactor equipped with an agitator, means for adding liquids, means for adding gases, means for pumping liquid from the reaction and recirculating it to the reactor, and a vent equipped for applying vacuum and for scrubbing or collecting removed volatiles, is added 273.4 kgs. of monochlorobenzene. Over a period of four hours are fed 729 kgs. of 55% n-octyl-3-mercaptopropionamide in monochlorobenzene and 195.9 kgs. of chlorine gas, while agitating and maintaining the temperature at 46°C. The chlorine feed was then continued to add 63.8 kgs. over a further two hours. The batch is analyzed as follows:

50	2-n-Octylisothiazolone (OIT)	29%
	Hydrogen chloride bound to OIT	5.5%
	Hydrogen chloride free in solution	6.5%
55	Monochlorobenzene	52%
	Process impurities	7%

Analysis is by gas chromatographic analysis of the organic layer after weighing an aliquot, treatment of the solution with water/ethyl acetate, and analysis of a known weight of the ethyl acetate solution. The process impurities include 4-chloro-2-octylisothiazolone, 4,5-dichloro-2-octylisothiazolone 3-chloro-N-octylpropionamide and related amide by-products. The hydrogen chloride content of the reaction mixture is measured by titration of an aliquot of known weight with 0.1 N sodium hydroxide to a phenolphthalein endpoint. The amount bound as the monohydrochloride was calculated based on OIT concentration.

While maintaining agitation and temperature control to 46°C the reaction is sparged with nitrogen (feed rate 566.4 liters/minute at standard conditions) for 8 hours. After about 4 hours, small crystals of the 2-octyl-3(2H)isothiazolone hydrochloride began to precipitate; at the end of the degassing, the batch contained about 22% weight percent of precipitated solids consisting of the monohydrochloride salt. The batch was again analyzed:

15	2-n-Octylisothiazolone (OIT)	35.5
	Hydrogen chloride bound to OIT.....	6.5
	Hydrogen chloride free in solution..	\<0.5
	Monochlorobenzene.....	49
20	Process impurities.....	8.5

Three separate repeats of this process afforded values of 21.8% solids precipitated as monohydrochloride salt from the batch and 107.5 % of HCl present relative to the stoichiometric amount needed to form the mono-hydrochloride salt.

The batch is cooled to 38°C, then re-heated to 46°C. The cycle is repeated two to three times, with agitation maintained throughout. The heating and cooling rates were both 1°C per 5 minutes. The batch is then cooled to 20°C at the same rate, to give clustered crystals representing 36 weight percent of the batch. The crystallized batch is readily filtered on a pressure column filter at 1400-2100 gms/sq.cm. nitrogen pressure. The wet filter cake of crystals was then washed with 455 kgs. of monochlorobenzene at 20°C. An aliquot of the snow-white crystals is dried in a vacuum oven at 45°C/0.013 Bar (10 mm. mercury) for one hour, and is analyzed as OIT 85.3%; HCl 14.5%; total accountability 99.8%.

EXAMPLE 2 - Comparative Example

This example illustrates the formation of long thin needles like crystals which were extremely difficult to filter and to wash, and is presented here to show the difficulties of purification when utilizing an inert solvent without applying the critical steps of the present invention.

The process of Example 1 is followed until the chlorination step is completed. Vacuum of 0.0667 to 0.0267 to 0.0267 Bar (50-20 mm. mercury) is then applied to the reaction mixture with no control of reaction temperature. After four hours, the reaction is at room temperature, and long thin crystals of OIT.HCl (ca. 300 x 10 x 10 µm) precipitates as a paste soaking-up all of the solvent. When filtration of the paste was attempted, crystals free of solvent could not be isolated, either by use of a fritted glass dip stick or of a Buechner funnel under vacuum.

EXAMPLE 3 - 2-n-Octylisothiazolone, hydrochloride salt

This example illustrates the use of variable rate cooling to produce clustered crystals, readily filterable, and of high purity.

The steps of Example 1 are followed through the degassing step. The mixture containing small crystals (21% solids) is heated to 52°C with agitation and allowed to equilibrate until the content of undissolved "seed" crystals is reduced to 12% of the batch weight. The mixture is then cooled from 52°C to 42°C at a rate of 1°C/15 minutes, from 42°C to 32°C at 1°C/10 minutes, and from 32°C to 20°C at a rate of 1°C/5 minutes. The mixture is then held at 20°C prior to filtration. Readily filterable crystals (size, 300x150x100 um free of fines) of high purity are obtained after washing with MCB.

EXAMPLE 4 - 2-n-Octylisothiazolone

This example illustrates the conversion of the purified crystals of OIT hydrochloride to free OIT of high purity.

5 Wet-cake clustered crystals of OIT monohydrochloride (from Example 1) are directly treated in the pressure filtration column with an equal weight of hot (65°C) water. After two hours of recirculation and one hour of separation, the upper aqueous layer is removed, and the organic layer further washed with an equal weight of water to remove any dissolved residual hydrochloric acid. The organic layer is then charged to a reactor equipped with an agitator, means for adding steam, means for applying vacuum, and means for adding and 10 removing liquid, and stripped with steam at 80°C under a vacuum of 0.0667 Bar(50 mm. mercury) to give 306 kgs. of the final product, which analyzed as follows:

15	2-n-Octylisothiazolone (OIT)	98.5%
	hydrogen chloride	≤0.2%
	Monochlorobenzene	1.5%
	Water	0.2%
20	Process impurities	≤0.1%

Crystals from the crystallization procedure of Example 3 gave equivalent results.

EXAMPLE 5 - 2-n-Octylisothiazolone

This example illustrates an alternate procedure for removal of solvent from the purified *n*-octylisothiazolone.

30 Crystals from Example 1 were charged to the reactor of Example 3, hydrolyzed as in Example 4, and stripped under vacuum to a final pressure of 0.0133 Bar(10 mm. mercury) at 55°C. The purity and yield of OIT was equivalent to that from the stripping process of Example 4.

EXAMPLE 6 - Formulation of 2-n-Octylisothiazolone

35 This example illustrates a formulation. *N*-octylisothiazolone (320 kgs.) is dissolved in 1,2-propylene glycol (375 kgs.) to afford a 46% solution. The product may be added to a commercial acrylic latex paint formulation, preferably containing 0.012 kg./l. of zinc oxide, at a level of 0.0048 kgs. solution/l. of paint. The paint will exhibit good storage stability in the can against bacteriocidal attack, and will exhibit excellent resistance to mildew upon application to exterior surfaces.

40 *N*-octylthiazolone may also be dissolved at other concentrations in propylene glycol or other appropriate solvents to prepare solutions appropriate for use as mildewcides for fabric, hides or leather.

EXAMPLE 7 - n-Octylisothiazolone

45 This example illustrates preparation of pure OIT from the dithiodipropionamide starting material.

To a reactor as in Example 1 containing 110 kgs. of monochlorobenzene is co-fed a) a slurry of 394 kgs. of *N,N*'-diocetyl dithiodipropionamide in 483 kgs. of monochlorobenzene and b) 145 kgs. of chlorine, the co-feed being carried out over a four hour period at 45°C with agitation. An additional 49 kgs. of chlorine was further added over two hours. Further reaction steps are carried out as in Examples 3 and 4 to produce 301 kgs. of 50 99.5% pure OIT.

EXAMPLE 8

This example illustrates vacuum degassing to reduce and control the content of excess hydrogen chloride after chlorination is complete.

The reaction conditions of Example 1 are followed through the completion of chlorination. Vacuum is then applied with the temperature maintained at 46°C and agitation of the mixture. The batch is then recirculated to aid in degassing and removal of excess hydrogen chloride under vacuum. The pressure is lowered to 0.06

5 Bar(45 mm. mercury) over a four hour period, at which point the OIT hydrochloride began to precipitate with a mild exotherm. The exotherm is controlled to maintain temperature at 46°C, and vacuum is continued without recirculation for another two hours. The crystals formed at this point could not be filtered readily because of their small size. The temperature was raised to 52°C and held for one hour when the weight of precipitate crystals decreased from 21.6 to 12.8%. Hydrogen chloride present at this point was 112% of the stoichiometric amount needed for complete formation of the hydrochloride salt.

10 The batch was cooled to 20°C with a variable cooling rate (as shown in Example 3) to afford readily filterable crystals. After hydrolysis and separation as in Example 4, the OIT was shown to be 98.8% purity with the remainder being essentially monochlorobenzene and water.

10

EXAMPLE 9

This example illustrates vacuum degassing at a temperature where the salt remains soluble.

15 The conditions of Example 1 are followed through the completion of the chlorination. The batch is then subjected to vacuum degassing at 60°C and a final pressure of 0.099 to 0.107 Bar(75 to 80 mm. mercury). After four hours, the hydrogen chloride content is reduced to 108% of the stoichiometric amount. The mixture remains free of any solids. On cooling to 46°C, exothermic nucleation occurs. After equilibration for three hours at 46°C, the batch contained 19.6 wt.% solids precipitated as small unfilterable crystals. Upon following the 20 temperature cycling and cooling process of Example 1, readily filterable crystals of dimensions 330x100 x50 µm are formed. (After filtration and washing, the dimensions of the crystals are measured from calibrated photographs taken from optical microscopy). Hydrolysis and workup as in Example 4 affords the hydrochloride salt of OIT of high purity.

25

EXAMPLE 10

Use of variable cooling in precipitating clustered crystals after vacuum degassing of a solution.

25 The procedure of Example 9 is followed, except that after formation of the seed crystals, the temperature is raised to 52°C and equilibrated for one hour to provide about 12 wt. % of the batch as seed crystals. The cooling procedure of Example 3 is then employed to produce crystals of the salt which are readily filtered and washed.

30

EXAMPLE 11

35 This example illustrates vacuum degassing under conditions where smaller amounts of seed crystals were formed.

The procedure of Example 8 is followed, except that vacuum degassing is conducted at 52°C at a final pressure of 0.067 Bar(50 mm. mercury), reached over a four-hour period, followed by at least a two-hour equilibration. Small crystals amounting to 10% of the total batch were formed. Cooling and isolation as in Example 3 affords crystals of the salt which are readily filtered and washed.

40

EXAMPLE 12

45 This example illustrates the preparation of filterable crystals through the external introduction of seed crystals. The process of Example 9 is followed to afford a clear solution at 60°C containing 35.5 weight percent of isothiazolone and 6.6 weight percent hydrogen chloride. The batch is cooled to 52°C at 1°C/5 minutes and 3 weight percent OIT.HCl crystals, size 150x30x30 µm are added. The batch is held at 52°C for two hours, whereupon about 13 percent of the batch is present as small crystals. The process of Example 3 is then used to produce easily filterable crystals of high purity.

50

EXAMPLE 13 - n-Propylisothiazolone

55 A stirred slurry of 438 gms. of N,N'-di-n-propyl dithiodipropionamide in 660 gms. of monochlorobenzene is chlorinated at 46°C with 320 gms. of chlorine over a three hour period. The reaction mixture is degassed under vacuum at 46°C, when small crystals of the 2-n-propylisothiazolone monohydrochloride precipitated. The reaction mixture is heated to 52°C and then cooled to 20°C under the variable rate cooling conditions of Example 3. The crystalline slurry is readily filtered on a Buechner funnel, washed with monochlorobenzene, and dried under vacuum to afford 464 gms. (86% yield) 99.5% of pure 2-n-propylisothiazolone hydrochloride.

EXAMPLE 14 - 2-n-Hexylisothiazolone


In a manner similar to that of Example 13, 377 gms. of a 40% solution of N,N'-di-n-hexylidithiopropionamide in monochlorobenzene is reacted with 213 gms. of chlorine fed over three hours at a reaction temperature of 45°C. The reaction mixture is then degassed under vacuum at 46°C, when small crystals of the 2-n-hexylisothiazolone precipitated. The reaction mixture is heated to 52°C and then cooled to 20°C under the variable rate cooling conditions of Example 3. The crystalline slurry is readily filtered on a Buechner funnel, washed with 100 mls. of monochlorobenzene, and dried under vacuum to afford 355 gms. (80% yield) of 99% pure 2-n-hexylisothiazolone hydrochloride.

10

Claims

1. A process for preparing crystalline isothiazolone monohydrochloride salts of the formula:


15



25

wherein R is hydrogen or methyl and Y is linear or branched alkyl, of from 3 to 10 carbon atoms, by the chlorination of a compound of the formula


30



35

where X is -SH or

40



wherein:

45 a) the chlorination is conducted in a chlorinated hydrocarbon solvent to afford a crude reaction mixture containing isothiazolone hydrochloride in solution and to excess hydrogen chloride; characterised in that

b) the excess of hydrogen chloride is reduced to less than 0.2 moles in excess of that present in one mole of the isothiazolone mono-hydrochloride salt; then,

c) the solution is formed into a seeded solution; then,

50 d) the seeded solution is cooled under temperature/time conditions to form filterable crystals of the salt; and then

e) the crystals are collected.

2. A process of Claim 1 wherein step (d) is carried out in such a way as to avoid the formation of crystals too small to be filtered.

55

3. A process of Claim 1 or 2 which further includes, subsequent to formation of the seeded solution, three or four cycles of heating and cooling of the seeded solution at a temperature of from 46°C to 36°C before


final cooling to 20°C.

4. A process of Claim 1 or 2 wherein, after the formation of seeded solution the batch is heated to about 52°C and then subjected to cooling at an increasing rate.
5. A process of any preceding Claim wherein the solvent comprises monochlorobenzene and/or ethylene dichloride perchloro ethylene.
6. A process of any preceding Claim wherein Y is linear C₃ to CH₁₀ alkyl and R is hydrogen.
10. 7. A process of any preceding Claim wherein Y is n-propyl, n-hexyl or n-octyl.
8. A process of any preceding Claim wherein the size of the crystals are at least 200 µm by about at least 100 µm by about at least 50 µm.
15. 9. A process according to any preceding Claim wherein the collected crystals are washed and then hydrolyzed and the resulting isothiazolone is isolated.
10. 10. The process of any preceding Claim wherein Y is n-octyl and R is hydrogen.

20 **Patentansprüche**

1. Verfahren zur Herstellung von kristallinen Isothiazolonmonohydrochloridsalzen der Formel:


25



30


35 worin R Wasserstoff oder Methyl ist und Y ein lineares oder verzweigtes C₃-C₁₀ Alkyl ist, durch Chlorierung einer Verbindung der Formel

40



wo X -SH oder

45



50

ist, bei dem

- (a) die Chlorierung in einem chlorierten Kohlenwasserstofflösungsmittel durchgeführt wird, wodurch ein Rohreaktionsgemisch mit Isothiazolonhydrochlorid in Lösung und Chlorwasserstoff im Überschuß erhalten wird, gekennzeichnet dadurch, daß
- (b) der Überschuß von Chlorwasserstoff auf weniger als 0,2 Mol Überschuß dessen, was in einem Mol des Isothiazolonmonohydrochloridsalzes vorliegt, reduziert wird, dann
- (c) die Lösung zu einer Saatlösung ausgebildet wird, dann
- (d) die Saatlösung unter Temperatur-/Zeitbedingungen abgekühlt wird, wodurch filterbare Kristalle des Salzes ausgebildet werden, und dann

(e) die Kristalle gesammelt werden.

2. Verfahren nach Anspruch 1, worin Schritt (d) in solcher Weise durchgeführt wird, daß die Ausbildung von zur Filtrierung zu kleinen Kristallen vermieden wird.

5 3. Verfahren nach Anspruch 1 oder 2, bei dem ferner nach Ausbildung der Saatlösung drei oder vier Zyklen von Erwärmen und Abkühlen der Saatlösung bei einer Temperatur von 46 °C bis 36 °C vor der Endabkühlung auf 20 °C erfolgen.

10 4. Verfahren nach Anspruch 1 oder 2, worin nach Ausbildung der Saatlösung der Ansatz auf etwa 52 °C erwärmt und dann einem Abkühlen mit erhöhter Rate unterzogen wird.

5. Verfahren nach einem der vorhergehenden Ansprüche, worin das Lösungsmittel Monochlorbenzol und/ oder Ethyldichloridperchlorethylen umfaßt.

15 6. Verfahren nach einem der vorhergehenden Ansprüche, worin Y ein lineares $C_3\text{-}CH_{10}$ Alkyl ist und R Wasserstoff ist.

7. Verfahren nach einem der vorhergehenden Ansprüche, worin Y n-Propyl, n-Hexyl oder n-Octyl ist.


20 8. Verfahren nach einem der vorhergehenden Ansprüche, worin die Größe der Kristalle mindestens etwa 200 μm auf mindestens etwa 100 μm auf mindestens etwa 50 μm ist.

9. Verfahren nach einem der vorhergehenden Ansprüche, worin die gesammelten Kristalle gewaschen und dann hydrolysiert werden und das erhaltene Isothiazolon isoliert wird.


25 10. Verfahren nach einem der vorhergehenden Ansprüche, worin Y n-Octyl ist und R Wasserstoff ist.

Revendications


30 1. Un procédé pour préparer des monochlorhydrates cristallins d'isothiazolone de formule:



dans laquelle R est de l'hydrogène ou un radical méthyle et Y est un radical alkyle linéaire ou ramifié ayant de 3 à 10 atomes de carbone, par chloration d'un composé de formule



50 dans laquelle X est -SH ou



dans lequel:

a) la chloration est mise en oeuvre dans un solvant hydrocarboné chloré pour avoir un mélange de réaction brute contenant du chlorhydrate d'isothiazolone en solution et du chlorure d'hydrogène en excès, caractérisé en ce que:

5 b) l'excès de chlorure d'hydrogène est réduit à une valeur inférieure à 0,2 mole par rapport à celui présent dans une mole du chlorhydrate d'isothiazolone et

c) la solution est formée dans une solution d'ensemencement puis

d) la solution d'ensemencement est refroidie sous des conditions de température/durée pour former des cristaux de sel susceptibles d'être filtrés; et ensuite

e) les cristaux sont recueillis.

10 2. Procédé selon la revendication 1 dans lequel l'étape (d) est mise en oeuvre de façon à éviter la formation de cristaux trop petits pour être filtrés.

15 3. Procédé selon l'une des revendications 1 ou 2 qui comprend, en outre, après la formation de la solution d'ensemencement, trois ou quatre cycles de chauffage et de refroidissement de la solution d'ensemencement à une température comprise dans la gamme d'environ 46°C à 36°C avant un refroidissement final à 20°C.

20 4. Procédé selon la revendication 1 ou 2, dans lequel après la formation de la solution d'ensemencement, le lot est chauffé à environ 52°C et ensuite soumis à un refroidissement à une vitesse croissante.

5. Procédé selon l'une quelconque des revendications précédentes, dans lequel le solvant comprend du monochlorobenzène et/ou du dichlorure d'éthylène et du tétrachloréthylène.

25 6. Procédé selon l'une quelconque des revendications précédentes, dans lequel Y est un radical alkyle linéaire en C₃ à CH₁₀ et R est de l'hydrogène.

7. Procédé selon l'une quelconque des revendications précédentes, dans lequel Y est un radical n-propyle, un radical n-hexyle ou un radical n-octyle.

30 8. Procédé selon l'une quelconque des revendications précédentes, dans lequel la dimension des cristaux est d'au moins 200 µm par au moins 100 µm par au moins 50 µm.

9. Procédé selon l'une quelconque des revendications précédentes, dans lequel les cristaux recueillis sont lavés puis hydrolysés et l'isothiazolone résultante est isolée.

35 10. Procédé selon l'une quelconque des revendications précédentes, dans lequel Y est le radical n-octyle et R est de l'hydrogène.

40

45

50

55