

(18)



Europäisches Patentamt

European Patent Office

Office européen des brevets

(11) Publication number:

0 034 039
B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication of patent specification: **26.09.84**

(21) Application number: **81300459.5**

(22) Date of filing: **04.02.81**

(51) Int. Cl.³: **C 11 D 1/83, C 11 D 1/72,**
C 11 D 1/75, C 11 D 3/32

(54) **Liquid detergent composition.**

(30) Priority: **05.02.80 US 118705**
30.01.81 US 230143

(43) Date of publication of application:
19.08.81 Bulletin 81/33

(45) Publication of the grant of the patent:
26.09.84 Bulletin 84/39

(84) Designated Contracting States:
BE DE FR GB IT NL

(58) References cited:
DE-A-2 424 856
DE-A-2 536 107
DE-A-2 715 351
US-A-3 963 649
US-A-4 144 201

**The file contains technical information
submitted after the application was filed and
not included in this specification**

(78) Proprietor: **THE PROCTER & GAMBLE
COMPANY**
301 East Sixth Street
Cincinnati Ohio 45202 (US)

(72) Inventor: **Welch, John B.**
308 Forest Avenue
Cincinnati Ohio 45215 (US)

(74) Representative: **Gibson, Tony Nicholas et al**
Procter & Gamble (NTC) Limited Whitley Road
Longbenton Newcastle upon Tyne NE12 9TS
(GB)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European patent convention).

Courier Press, Leamington Spa, England.

EP 0 034 039 B1

Description

The invention relates to aqueous high sudsing liquid detergent compositions containing specified amounts and types of surfactants especially useful in the washing of tableware, kitchenware and other
5 hard surfaces.

The compositions of this invention provide more complete drainage of rinse water from surfaces such as glass, ceramics and metal, thereby reducing spotting and filming, particularly in a dishwashing procedure that involves drain drying without towel drying and polishing.

The performance of a detergent composition for cleaning glasses, dishes, and other articles with a
10 normally shiny surface is evaluated by the consumer in terms of shine and the absence of filming, streaking, and spotting. The liquid dishwashing detergent compositions presently on the market are designed to remove the soils from glasses, dishes, and other tableware and kitchen utensils. The detergent solution and redeposited soil residues are normally removed from the washed articles by rinsing and optionally by towel drying the articles when they are still wet. If not rinsed and towel dried,
15 these residues can dry upon the surfaces of the washed articles, leaving films, streaks, or spots.

Even when such articles are entirely clean but rinsed in plain water containing dissolved salts such as water hardness, spots and streaks can appear on the washed and rinsed surfaces upon evaporation of the water.

Towel drying of washed articles, e.g., glasses and dishes, immediately after removal from the
20 washing and rinsing solution, is undesirable from the standpoints of convenience and hygiene. Therefore, it is common practice to put the washed or washed and rinsed articles aside for draining and air-drying. Consequently, the cleaning efficacy of the product used, which the housewife may have visually appreciated at the end of the washing or rinsing cycle, is diminished due to the adherence of redeposited soil, residual dried detergent, and water hardness residues.

U.S. Patent 3,963,649, Spadini et al, discloses liquid detergent compositions containing a
25 nonionic surfactant and a water-soluble gel-forming gelatin. These compositions are said to minimize filming, streaking and spotting of tableware and kitchen utensils. The essential nonionic surfactant may be a tertiary amine or phosphine oxide, an amide or a condensation product of ethylene oxide and an organic hydrophobic compound.

U.S. Patent 3,983,079, Spadini et al, discloses dishwashing detergent compositions said to have
30 good rinse water draining characteristics. The compositions contain a water-soluble quaternary ammonium compound, a nonionic surfactant containing both ethylene oxide and propylene oxide and a sultaine or betaine zwitterionic surfactant.

U.S. Patent 4,144,201, Winterbotham et al, discloses liquid dishwashing detergent compositions
35 containing anionic material; amine oxide or mixtures thereof with nonionic alkylene oxide condensates; and soluble casein to improve drain-dry and mildness properties.

Belgian Patent 845,184 discloses liquid and granular dishwashing detergent compositions containing one or more specified classes of surfactants to ensure rapid drainage and provide a shiny surface.


It is an object of the present invention to provide liquid detergent compositions and a process for
40 dishwashing that promote rapid and relatively complete drainage of rinse water thereby reducing spotting and filming on surfaces such as glass, ceramics and metal.

There is a continuing need for compositions and methods which can be employed during
dishwashing operations to improve the final dry appearance of washed and dried kitchen utensils and
45 articles. If such compositions and methods are intended to be useful for conventional dishwashing soil removal operations, there is a continuing need for a compatible combination of materials which will simultaneously provide the surfactancy, sudsing, and mildness attributes of an acceptable dishwashing detergent composition as well as the anti-spotting and anti-filming benefits described above.

According to the present invention there is provided an aqueous liquid detergent composition
50 containing 10% to 50% by weight of an anionic surfactant, from 2% to 20% by weight of an ethoxylated nonionic surfactant and from 0% to 10% of an organic or inorganic builder salt comprising a drainage-promoting ethoxylated nonionic surfactant selected from:

an aqueous liquid detergent composition containing from 10% to 50% by weight of an anionic
surfactant, from 2% to 20% by weight of an ethoxylated nonionic surfactant, from 1.5% to 10% of a
55 suds stabilizing agent selected from

a) amides of general formula



60 wherein R_1 is a C_7-C_{21} saturated or unsaturated aliphatic hydrocarbon radical, R_2 is methylene or ethylene and m is 1, 2 or 3;

65


b) amine oxides of general formula



wherein R_3 is an alkyl, 2-hydroxyalkyl, 3-hydroxyalkyl or 3-alkoxy 2 hydroxypropyl radical in which the alkyl and alkoxy groups respectively have 8—18 carbon atoms, R_4 and R_5 are C_1 — C_3 alkyl or hydroxy C_2 — C_3 alkyl and n is 0 to 10, and

c) mixtures thereof; from 0% to 10% of an organic or inorganic builder salt; and at least 20% of water; wherein the ethoxylated nonionic surfactant is a drainage-promoting ethoxylated nonionic surfactant selected from

i) an ethoxylated aliphatic alcohol of the formula



wherein R is an aliphatic hydrocarbyl radical containing from 16 to 30 carbon atoms, wherein n is from 16 to 100;

ii) an ethoxylated alkyl phenol of the formula



wherein R is an alkyl phenol radical containing a total of from 18 to 30 carbon atoms and at least one alkyl group containing at least 12 carbon atoms and wherein n is from 16 to 100;

iii) the condensation product of mono C_{16-22} fatty acid esters of polyols with from 13 to 100 moles of ethylene oxide per mole of the mono-ester;

iv) the condensation product of cholesterol and from 13 to 100 moles of ethylene oxide;

v) a material which is a condensate of ethylene oxide, propylene oxide and a compound containing hydroxy or amine groups onto which alkyl oxides can be polymerized, said polymer having a molecular weight of from 500 to 15,000, an ethylene oxide content of from 30% to 70% by weight and a propylene oxide content of from 30% to 70% by weight; and

vi) mixtures thereof;

said composition being free of proteinaceous materials.

Dishware, glassware, and other tableware and kitchenware may be washed in water solutions of the detergent composition, generally at a weight concentration of 0.05% to 0.4% of the composition in water at a temperature of 26°C to 49°C. The tableware and kitchenware are then rinsed, drained, and allowed to dry in a rack or other means of separation.

Detailed description of the invention

The liquid detergent compositions of the present invention contain four essential components:

- a) an anionic surfactant
- b) a drainage promoting ethoxylated nonionic surfactant
- c) a suds stabilizing nonionic surfactant
- d) water.

Optional ingredients can be added to provide various performance and aesthetic characteristics.

Anionic surfactant

The compositions of this invention contain from 10% to 50% by weight of an anionic surfactant or mixtures thereof. Preferred compositions contain from 20% to 35% of anionic surfactant by weight.

Most anionic detergents can be broadly described as the water-soluble salts, particularly the alkali metal, alkaline earth metal, ammonium and amine salts, of organic sulfuric reaction products having in their molecular structure an alkyl radical containing from 8 to 22 carbon atoms and a radical selected from the group consisting of sulfonic acid and sulfuric acid ester radicals. Included in the term alkyl is the alkyl portion of acyl radicals. Examples of the anionic synthetic detergents which can form the surfactant component of the compositions of the present invention are the sodium, ammonium, potassium or magnesium alkyl sulfates, especially those obtained by sulfating the higher alcohols (C_8 — C_{18} carbon atoms) sodium or magnesium alkyl benzene or alkyl toluene sulfonates, in which the alkyl group contains from 9 to 15 carbon atoms, the alkyl radical being either a straight or branched aliphatic

chain; sodium or magnesium paraffin sulfonates and olefin sulfonates in which the alkyl or alkenyl group contains from 10 to 20 carbon atoms; sodium C₁₀₋₂₀ alkyl glyceryl ether sulfonates, especially those ethers of alcohols derived from tallow and coconut oil; sodium coconut oil fatty acid monoglyceride sulfates and sulfonates; sodium, ammonium or magnesium salts of alkyl phenol
 5 ethylene oxide ether sulfates with 1 to 30 units of ethylene oxide per molecule and in which the alkyl radicals contain from 8 to 12 carbon atoms; the reaction products of fatty acids esterified with isethionic acid and neutralized with sodium hydroxide where, for example, the fatty acids are derived from coconut oil; sodium or potassium salts of fatty acid amides of a methyl tauride in which the fatty acids, for example, are derived from coconut oil and sodium or potassium beta-acetoxy or beta-
 10 acetamido-alkanesulfonates where the alkane has from 8 to 22 carbon atoms.

Specific examples of alkyl sulfate salts which can be employed in the instant detergent compositions include sodium lauryl alkyl sulfate, sodium stearyl alkyl sulfate, sodium palmityl alkyl sulfate, sodium decyl sulfate, sodium myristyl alkyl sulfate, potassium lauryl alkyl sulfate, potassium stearyl alkyl sulfate, potassium decyl sulfate, potassium palmityl alkyl sulfate, potassium myristyl alkyl
 15 sulfate, sodium dodecyl sulfate, magnesium dodecyl sulfate, potassium tallow alkyl sulfate, sodium tallow alkyl sulfate, sodium coconut alkyl sulfate, potassium coconut alkyl sulfate, magnesium C₁₂₋₁₅ alkyl sulfate and mixtures of these surfactants. Preferred alkyl sulfates include sodium C₁₂₋₁₅ alkyl sulfates and magnesium C₁₂₋₁₅ alkyl sulfate.

Suitable alkylbenzene or alkyltoluene sulfonates include the alkali metal (lithium, sodium, potassium), alkaline earth (calcium, magnesium) ammonium and alkanolamine salts of straight or
 20 branched-chain alkylbenzene or alkyltoluene sulfonic acids. Alkylbenzene sulfonic acids useful as precursors for these surfactants include decyl benzene sulfonic acid, undecyl benzene sulfonic acid, dodecyl benzene sulfonic acid, tridecyl benzene sulfonic acid, tetrapropylene benzene sulfonic acid and mixtures thereof. Preferred sulfonic acids as precursors of the alkyl-benzene sulfonates useful for
 25 compositions herein are those in which the alkyl chain is linear and averages 11 to 13 carbon atoms in length. Examples of commercially available alkyl benzene sulfonic acids useful in the present invention include Conoco SA 515 and SA 597 marketed by the Continental Oil Company and Calsoft LAS 99 marketed by the Pilot Chemical Company.

Particularly preferred anionic surfactants useful herein are alkyl ether sulfates having the formula
 30 $RO(C_2H_4O)_xSO_3M$ wherein R is alkyl or alkenyl of 10 to 20 carbon atoms, x is 1 to 30, and M is a water-soluble cation. The alkyl ether sulfates useful in the present invention are condensation products of ethylene oxide and monohydric alcohols having from 10 to 20 carbon atoms. Preferably, R has 10 to 16 carbon atoms. The alcohols can be derived from natural fats, e.g., coconut oil or tallow, or can be synthetic. Such alcohols are reacted with 1 to 30, and especially 1 to 12, molar proportions of ethylene
 35 oxide and the resulting mixture of molecular species is sulfated and neutralized.

Specific examples of alkyl ether sulfates of the present invention are sodium coconut alkyl triethylene glycol ether sulfate, magnesium C₁₂₋₁₅ alkyl triethylene glycol ether sulfate, and sodium tallow alkyl hexaoxy ethylene sulfate. Preferred alkyl ether sulfates are those comprising a mixture of individual compounds, said mixture having an average alkyl chain length of from 12 to 16 carbon atoms
 40 and an average degree of ethoxylation of from 1 to 12 moles of ethylene oxide.


Additional examples of anionic surfactants useful herein are the compounds which contain two anionic functional groups. These are referred to as dianionic surfactants. Suitable dianionic surfactants are the disulfonates, disulfates, or mixtures thereof which may be represented by the following formula:



where R is an acyclic aliphatic hydrocarbyl group having 15 to 20 carbon atoms and M is a water-solubilizing cation, for example, the C₁₅ to C₂₀ disodium, 1,2-alkyldisulfates, C₁₅ to C₂₀ dipotassium-1,2-alkyldisulfonates or disulfates, di-sodium 1,9-hexadecyl disulfates, C₁₅ to C₂₀ disodium 1,2-
 50 alkyldisulfonates, disodium 1,9-stearyldisulfates and 6,10-octadecyldisulfates.

Drainage promoting ethoxylated nonionic surfactant

The ethoxylated nonionic surfactants of the present invention are the condensation product of alcohols, alkyl phenols and other specified hydrophobic molecules with ethylene oxide. The materials
 55 hereinafter disclosed have not been used in aqueous liquid detergent compositions having the required formulation characteristics of the present invention. Their ability to improve rinse water drainage characteristics had not been recognized. Suds stabilizing nonionic surfactants hereinafter described have been in general use, but by themselves do not provide the improved drainage characteristics. Preferably, the compositions of the present invention contain from 2% to 20%, more preferably from
 60 3% to 12%, and most preferably from 3% to 8%, of drainage promoting ethoxylated aliphatic alcohols of the formula



wherein R is an aliphatic hydrocarbyl radical containing from 16 to 30 carbon atoms, wherein n is from
 65 16 to 100.

O 034 039

Other ethoxylated nonionic surfactants at a level of from 2% to 20% can provide the drainage promoting characteristics of ethoxylated alcohols, but are less desirable for reasons of biodegradability and effect on sudsing or cleaning performance. Examples of such alternate ethoxylated nonionic surfactants are:

- 1) an ethoxylated alkyl phenol of the formula $R(OC_2H_4)_nOH$ wherein R is an alkyl phenyl radical containing a total of from 18 to 30 carbon atoms and at least one alkyl group containing at least 12 carbon atoms wherein n is from 16 to 100;
- 2) the condensation product of mono C_{16-22} fatty acid esters of polyols with from 13 to 100 moles of ethylene oxide per mole of partial ester;
- 3) the condensation product of cholesterol and from 13 to 100 moles of ethylene oxide;
- 4) a material which is a condensate of ethylene oxide, propylene oxide and a compound containing hydroxy or amine groups onto which the alkylene oxides can be polymerized, said polymer having a molecular weight of from 500 to 15,000, an ethylene oxide content of from 30% to 70% by weight and a propylene oxide content of from 30% to 70% by weight.

In a particularly preferred embodiment an aliphatic alcohol contains from 16 to 22 carbon atoms and is ethoxylated to an average degree of from 18 to 50 moles of ethylene oxide per mole of alcohol.

20 Suds stabilizing nonionic surfactant

The compositions of this invention contain from 1.5% to 10%, preferably from 2% to 8%, of suds stabilizing nonionic surfactant or mixture thereof having a different chemical structure and function than the essential drainage promoting nonionic surfactant.

Suds stabilizing nonionic surfactants operable in the instant compositions are two basic types—amides and the amine oxide semi-polar nonionics.

The amide type of nonionic surface active agent include the ammonia, monoethanol and diethanol amides of fatty acids having an acyl moiety of from 8 to 18 carbon atoms and represented by the general formula



wherein R_1 is a saturated or unsaturated, aliphatic hydrocarbon radical having from 7 to 21, preferably from 11 to 17 carbon atoms; R_2 represents a methylene or ethylene group; and m is 1, 2, or 3, preferably 1 or 2. Specific examples of said amides are mono-ethanol coconut fatty acid amide and diethanol dodecyl fatty acid amide. These acyl moieties may be derived from naturally occurring glycerides, e.g., coconut oil, palm oil, soybean oil and tallow, but can be derived synthetically, e.g., by the oxidation of petroleum, or hydrogenation of carbon monoxide by the Fischer-Tropsch process. The monoethanol amides and diethanolamides of C_{12-14} fatty acids are preferred.

Amine oxide semi-polar nonionic surface active agents comprise compounds and mixtures of compounds having the formula:



wherein R_3 is an alkyl, 2-hydroxyalkyl, 3-hydroxyalkyl, or 3-alkoxy-2-hydroxypropyl radical in which the alkyl and alkoxy, respectively, contain from 8 to 18 carbon atoms, R_4 and R_5 are methyl, ethyl, propyl, isopropyl, 2-hydroxyethyl, 2-hydroxypropyl, or 3-hydroxypropyl and n is from 0 to 10. Particularly preferred are amine oxides of the formula:



wherein R_3 is a C_{10-14} alkyl and R_4 and R_5 are methyl or ethyl.

The preferred sudsing characteristics of the compositions of the invention are those which will provide the user of the product with an indication of cleaning potential in a dishwashing solution. Soils encountered in dishwashing act as suds depressants and the presence or absence of suds from the surface of a dishwashing solution is a convenient guide to product usage. Mixtures of anionic surfactants and suds stabilizing nonionic surfactants are utilized in the compositions of the invention because of their high sudsing characteristics, their suds stability in the presence of food soils and their

ability to indicate accurately an adequate level of product usage in the presence of soil. Additionally, and most importantly, compositions containing the other two essential surfactants of the invention but not the suds stabilizing nonionic surfactants as defined herein, do not provide an optimum draining promoting effect.

5 In preferred embodiments of the invention, the ratio of anionic surfactants to total nonionic surfactants in the composition will be in a molar ratio of from 11:1 to 1:1, and more preferably from 8:1 to 3:1. From the standpoint of sudsing, the suds stabilizing nonionic surfactants are generally preferred, but the essential relatively highly ethoxylated drainage promoting nonionic surfactants of the invention can contribute to sudsing performance and are included in the calculation of ratios of anionic to
10 nonionic surfactant.

Other optional surfactants

The compositions of the invention may contain optional surfactants such as ampholytic, zwitterionic and cationic surfactants.

15 Ampholytic surfactants can be broadly described as derivatives of aliphatic amines which contain a long chain of 8 to 18 carbon atoms and an anionic water-solubilizing group, e.g. carboxy, sulfo or sulfate. Examples of compounds falling within this definition are sodium-3-dodecylamino propane sulfonate, and dodecyl dimethylammonium hexanoate.

Zwitterionic surface active agents operable in the instant composition are broadly described as
20 internally-neutralized derivatives of aliphatic quaternary ammonium and phosphonium and tertiary sulfonium compounds in which the aliphatic radical can be straight chain or branched, and wherein one of the aliphatic substituents contains from 8 to 18 carbon atoms and one contains an anionic water solubilizing group, e.g., carboxy, sulfo, sulfato, phosphato, or phosphono.

Cationic surfactants such as quaternary ammonium compounds can find optional use in the
25 practice of the invention to the extent they are compatible with the other surfactants in the particular composition.

Water

The compositions of this invention contain at least 20%, preferably from 40% to 70%, water.
30

Additional optional ingredients

The compositions of this invention can contain up to 10%, by weight of detergency builders either of the organic or inorganic type. Examples of water-soluble inorganic builders which can be used, alone or in admixture with themselves and organic alkaline sequestrant builder salts, are alkali metal
35 carbonates, phosphates, polyphosphates and silicates. Specific examples of such salts are sodium tripolyphosphate, sodium carbonate, potassium carbonate, sodium pyrophosphate, potassium pyrophosphate, potassium tripolyphosphate, and sodium hexametaphosphate. Examples of organic builder salts which can be used alone, or in admixture with each other or with the preceding inorganic alkaline builder salts, are alkali metal polycarboxylates, e.g., water-soluble citrates such as sodium and
40 potassium citrate, sodium and potassium tartrate, sodium and potassium ethylenediaminetetraacetate, sodium and potassium N-(2-hydroxyethyl)-ethylene diamine triacetates, sodium and potassium nitrilo triacetates (NTA) and sodium and potassium N-(2-hydroxyethyl)-nitrilo diacetates. Other organic detergency builders such as water-soluble phosphonates can find use in the compositions of the invention. In general, however, detergency builders have limited value in
45 dishwashing detergent compositions and use at levels above 10% can restrict formulation flexibility in liquid compositions because of solubility and phase stability considerations.

Alcohols, such as ethyl alcohol, and hydrotropes, such as sodium and potassium toluene sulfonate, sodium and potassium xylene sulfonate, trisodium sulfosuccinate and related compounds (as disclosed in U.S. Patent 3,915,903), and urea, can be utilized in the interest of achieving a desired
50 product phase stability and viscosity. Ethyl alcohol at a level of from 3% to 15% and potassium or sodium toluene, xylene or cumene sulfonate at a level of from 1% to 6% are particularly useful in the compositions of the invention.

The detergent compositions of this invention can contain, if desired, any of the usual adjuvants, diluents and additives, for example, perfumes, enzymes, dyes, anti-tarnishing agents, antimicrobial
55 agents, and the like, without detracting from the advantageous properties of the compositions. Alkalinity sources and pH buffering agents such as monoethanolamine, triethanolamine and alkali metal hydroxides can also be utilized.

The following examples are given to illustrate the compositions of the invention. All percentages are by weight unless otherwise indicated.
60

O 034 039

Example I

The following liquid detergent compositions were prepared.

		A	B	C
5	Ammonium C ₁₂ alkyl sulfate	4.3%	4.3%	12.5%
	Ammonium C ₁₂ alkyl sulfate-ethoxy (1)	19.9	19.9	—
	Ammonium C ₁₂ alkyl sulfate-ethoxy (3)	—	—	13.5
10	Dimethyldodecylamine oxide	2.3	1.7	4.0
	Tallow alcohol-ethoxy (22)	5.0	4.0	—
15	Ethanol	6.2	6.2	6.2
	Ammonium xylene sulfonate	2.5	2.5	2.5
	Magnesium chloride	3.1	3.1	—
20	Potassium chloride	—	—	0.7
	Water	56.2	57.8	60.1
25	Perfume and miscellaneous	0.5	0.5	0.5

Compositions A and B are within the scope of the present invention. Composition C is typical of presently used dishwashing liquid detergent compositions and is outside the scope of the present invention.

30

Spotting and filming performance

"Libby" glasses were soiled with a fatty soil containing milk solids and washed in 46°C water solutions containing 0.2% of Compositions A, B and C. The glasses were rinsed in 46°C water, rack dried, graded on a 1—10 scale (1 poorest, 10 best) for spotting and filming. The glasses were also
35 comparison graded for overall appearance using a scale of 0 to 4 to indicate no difference to a large advantage for one of the glasses in the comparison.

Average Conditions—0.25% soil and 7 grains/gallon water hardness measured as CaCO₃

40

	A	B	C	Least significant Difference—.05
Spotting	7.4	7.3	6.7	0.32
45 Filming	7.7	7.5	7.4	0.22
Appearance	+0.5	+0.2	—1.4	0.7

50 Stress Conditions—0.35% soil and 10 grains/gallon water hardness measured as CaCO₃

	A	B	C	Least significant Difference—0.05
55 Spotting	7.3	7.3	6.4	0.27
Filming	7.4	7.5	7.2	0.25
Appearance	+0.1	—0.1	—1.8	1.1

60

Sudsing

Suds were generated by agitation in dishpans containing 2 gallons of 46°C water using Compositions A, B and C at a 0.2% product concentration. Dinner plates were washed with the introducing of 4.0 ml of a triglyceride-containing soil on each plate. Suds height is measured after washing sets of five plates. This procedure is repeated five times for a total of 25 plates. The suds
65 height after washing each set is expressed in terms of percent of original suds height and an average of

O 034 039

the five values is reported as suds during washing (SDW). The number of plates washed when suds disappear from the surface of the dishwashing solution is recorded as "mileage".

The following sudsing results were obtained:

5	Soil	Water Hardness	A	B	SDW/Mileage C	LSD—.05
	100% fat	2	16.3/—	17.4/—	14.9/—	2.5/—
10		7	17.2/—	17.2/—	17.3/—	1.6/—
		14	13.0/—	12.4/—	11.4/—	6.0/—
15	Mixed fat, protein, carbohydrate and acid	2	14.8/32	14.5/31	13.6/30	4.8/5.3
		7	12.5/21	12.2/21	13.0/21	2.7/4.9
		14	12.4/31	12.6/29	10.5/28	5.0/9.1

20 Compositions A and B had sudsing characteristics equivalent to C and provided a superior appearance and freedom from spotting and filming after washing and rinsing.

The following materials are substituted for the tallow alcohol-ethoxy (22) in Compositions A and B:

- 25
- 1) C₁₃₋₁₅ alkyl phenol-ethoxy (30)
 - 2) cholesterol-ethoxy (24)
 - 3) sorbitan monoleate-ethoxy (80)
 - 4) glyceryl monostearate-ethoxy (20)
 - 5) C₁₈ alkanol-ethoxy (20)

30 Comparable sudsing, appearance, filming and spotting performance relative to Composition C is obtained.

Example II

The following liquid dishwashing detergent compositions were prepared:

35		D	E	F
	Ammonium C ₁₂ alkyl sulfate	12.5%	4.3%	—
40	Ammonium C ₁₂ alkyl sulfate-ethoxy (1)	—	19.9	—
	Ammonium C ₁₂ alkyl sulfate-ethoxy (3)	13.5	—	12.5
	Ammonium C ₁₁₋₁₃ alkylbenzene sulfonate	—	—	18.8
45	Magnesium chloride	—	3.1	—
	Dimethyldodecylamine oxide	4.0	1.7	—
50	C ₁₂ alkyl monoethanolamide	—	—	1.9
	Ammonium xylene sulfonate	2.5	—	2.4
	Ethanol	6.2	2.5	3.0
55	Water	60.3	68.5	61.4
	Perfume and miscellaneous	1.0	—	—

60 For purposes of evaluating spotting and filming performance a variation of Composition D was prepared in which 5% tallow alcohol-ethoxy (22) replaced 5% water. Similarly variations of Composition E were prepared in which 1%, 5% and 20% tallow alcohol-ethoxy (22) replaced water.

65 Paired comparison grading of the two base compositions and four variations containing tallow alcohol-ethoxy (22) for filming and spotting performance on glassware after washing and rinsing showed:

O 034 039

- 1) no difference between base formulas
- 2) only a marginal benefit at a 1% tallow alcohol-ethoxy (22) level
- 3) a substantial and easily noticeable benefit for the 5% tallow alcohol-ethoxy (22) level
- 4) only a marginal benefit in an increase in tallow alcohol-ethoxy (22) from 5% to 20%.

5

Sudsing of Composition D with 5% tallow alcohol-ethoxy (22) was equivalent to sudsing of Composition C. Composition D with 20% tallow alcohol-ethoxy (22) provided an average suds level approximately 50% of that provided by Composition C.

10 C_{12-15} alkylbenzene sulfonate, C_{13-16} paraffin sulfonate and C_{12-16} olefin sulfonate are substituted for the C_{12} alkyl sulfate in Compositions C and D. Comparable results are obtained.

C_{12} monoethanolamide at a 6% level replaces 4% dimethyldodecylamine oxide and 2% water in Composition C. Comparable results are obtained.

Example III

15 Variations of Compositions D of Example II were prepared with the following materials replacing water.

- 1) 5% tallow alcohol-ethoxy (22)
- 2) 5% tallow alcohol-ethoxy (11)
- 20 3) 5% C_{14-15} alkanol-ethoxy (7)
- 4) 5% C_{12} alkanol-ethoxy (12)
- 5) 5% C_{16} alkanol-ethoxy (20)

Grading of glassware for filming and spotting after washing and rinsing in the compositions showed:

25 Compositions #3 and #4 provide no improvement over the base Composition D containing no ethoxylated nonionic surfactant. Compositions #1 and #5 provided an easily noticeable benefit relative to the base Composition D and Compositions #2, #3 and #4. Composition #2 had a slightly lower level of filming and spotting relative to the base Composition D, but was substantially less effective than Compositions #1 and #5.

30 Sudsing of Composition #2 was substantially reduced relative to base Composition D and Composition #1 which was essentially the same as base Composition D.

Example IV

35 A variation of Composition D of Example II was prepared with 17% of Pluridot HA—430 replacing water. Pluridot HA—430 is manufactured by BASF-Wyandotte and is a condensate of ethylene oxide and propylene oxide polymerized on a triol base and having a molecular weight of 3700—4200. The resultant composition provided an advantage of reduced filming and spotting of glassware relative to Composition D.

40

45

50

55

60

65

0 034 039

Example V

The following liquid dishwashing detergent compositions were prepared:

		G	H	I	J
5	Ammonium C ₁₂ alkyl sulfate	4.3%	4.3%	—	12.5%
	Ammonium C ₁₂ alkyl ether sulfate-ethoxy (1)	19.9	19.9	—	—
10	Ammonium C ₁₂ alkyl ether sulfate-ethoxy (3)	—	—	12.5	13.5
	Ammonium C ₁₂₋₁₃ alkylbenzene sulfonate	—	—	18.8	—
	Dimethyldodecylamine oxide	2.8	2.8	—	4.0
15	C ₁₂ alcohol-ethoxy (23)	—	3.6	—	—
	Tallow alcohol-ethoxy (18)	3.3	—	—	—
20	Tallow alcohol-ethoxy (22)	—	—	5.0	—
	Ethanol	6.2	6.2	3.0	6.2
	Ammonium xylene sulfonate	2.5	2.5	2.4	2.5
25	Magnesium chloride	3.1	3.1	—	—
	Potassium chloride	—	—	—	0.7
30	Perfume and miscellaneous	0.5	0.5	0.5	0.5
	Water	—remainder—			

Spotting and filming performance

35 Water glasses were soiled with a fatty soil containing milk solids and washed in 46°C water solutions containing 0.2% of Compositions G, H, I and J. The glasses were rinsed in 46°C water and rack dried. Glasses from each treatment were compared for overall spotting and filming appearance. Graders assigned values of +4 to -4 to indicate their satisfaction or dissatisfaction with the end result appearance of the glasses.

40 Conditions—0.3% soil and 10 grains/gallon water hardness measured as CaCO₃.


45		G	H	I	J	Least significant Difference—0.05
	Average appearance grade	+0.6	−0.2	−0.2	−0.9	0.8

50 Composition G provided a significant appearance advantage relative to the other three compositions. Compositions H and I provided a numerical advantage over Composition J not statistically significant at a 95% confidence level. Only Composition G provided results on the positive side of the satisfaction scale.

Claims

55 1. An aqueous liquid detergent composition containing from 10% to 50% by weight of an anionic surfactant, from 2% to 20% by weight of an ethoxylated nonionic surfactant, from 1.5% to 10% of a suds stabilizing agent selected from

60 a) amides of general formula



65 wherein R₁ is a C₇—C₂₁ saturated or unsaturated aliphatic hydrocarbon radical, R₂ is methylene or ethylene and m is 1, 2 or 3;

b) amine oxides of general formula



wherein R_3 is an alkyl, 2-hydroxyalkyl, 3-hydroxyalkyl or 3-alkoxy-2-hydroxypropyl radical in which the alkyl and alkoxy groups respectively have 8—18 carbon atoms, R_4 and R_5 are C_1 — C_3 alkyl or hydroxy C_2 — C_3 alkyl and n is 0 to 10, and

c) mixtures thereof; from 0% to 10% of an organic or inorganic builder salt; and at least 20% of water;


characterised in that the ethoxylated nonionic surfactant is a drainage-promoting ethoxylated nonionic surfactant selected from

i) an ethoxylated aliphatic alcohol of the formula



wherein R is an aliphatic hydrocarbyl radical containing from 16 to 30 carbon atoms, wherein n is from 16 to 100;

ii) an ethoxylated alkyl phenol of the formula



wherein R is an alkyl phenol radical containing a total of from 18 to 30 carbon atoms and at least one alkyl group containing at least 12 carbon atoms and wherein n is from 16 to 100;

iii) the condensation product of mono C_{16-22} fatty acid esters of polyols with from 13 to 100 moles of ethylene oxide per mole of the mono-ester;

iv) the condensation product of cholesterol and from 13 to 100 moles of ethylene oxide;

v) a material which is a condensate of ethylene oxide, propylene oxide and a compound containing hydroxy or amine groups onto which alkyl oxides can be polymerized, said polymer having a molecular weight of from 500 to 15,000, an ethylene oxide content of from 30% to 70% by weight and a propylene oxide content of from 30% to 70% by weight; and

vi) mixtures thereof;


said composition being free of proteinaceous materials.

2. A composition according to Claim 1 characterised in that the drainage-promoting nonionic surfactant has the formula (i) wherein R is a C_{16} — C_{22} aliphatic hydrocarbyl radical and n has a value of from 18 to 50.

Patentansprüche


1. Wäßriges flüssiges Reinigungsmittel, enthaltend 10 bis 50 Gew.-% eines anionischen oberflächenaktiven Stoffes, 2 bis 20 Gew.-% eines ethoxylierten nichtionischen oberflächenaktiven Stoffes, 1,5 bis 10% eines Schaumstabilisators ausgewählt aus

a) Amiden der allgemeinen Formel



worin R_1 ein C_7 — C_{21} -gesättigter oder ungesättigter aliphatischer Kohlenwasserstoffrest, R_2 Methylen oder Ethylen und m 1, 2 oder 3 sind;

b) Aminoxiden der allgemeinen Formel



worin R_3 ein Alkyl-, 2-Hydroxyalkyl-, 3-Hydroxyalkyl- oder 3-Alkoxy-2-hydroxypropylrest, worin die Alkyl- bzw. Alkoxygruppen 8 bis 18 C-Atome aufweisen, R_4 und R_5 C_{1-3} -Alkyl oder Hydroxy- C_2 — C_3 -alkyl und n 0 bis 10 sind, und

c) Gemischen davon,

0 bis 10% eines organischen oder anorganischen Gerüststoffsalzes und mindestens 20% Wasser, dadurch gekennzeichnet, daß der erthoxylierte nichtionische oberflächenaktive Stoff ein ablauffördernder
5 ethoxylierter nichtionischer Stoff ist ausgewählt aus

- i) einem ethoxylierten aliphatischen Alkohol der Formel $R(OC_2H_4)_nOH$, worin R einen aliphatischen Kohlenwasserstoffrest mit 16 bis 30 C-Atomen und n 16 bis 100 bedeuten;
- 10 ii) einem ethoxylierten Alkylphenol der Formel $R(OC_2H_4)_nOH$, worin R einen Alkylphenolrest mit einer Gesamtmenge von 18 bis 30 c-Atomen und mindestens einer Alkylgruppe mit mindestens 12 C-Atomen und n 16 bis 100 bedeuten;
- iii) dem Kondensationsprodukt von Mono- C_{16-22} -fettsäureestern und Polyolen mit 13 bis 100 Mol Ethylenoxid je Mol des Monoesters;
- iv) dem Kondensationsprodukt von Cholesterol und 13 bis 100 Mol Ethylenoxid;
- 15 v) einem Material, das ein Kondensat aus Ethylenoxid, Propylenoxid und einer Verbindung ist, die Hydroxy- oder Amingruppen enthält, auf die Alkyloxyde aufpolymerisiert sein können, wobei das Polymer ein Molekulargewicht von 500 bis 15 000, einen Ethylenoxidgehalt von 30 bis 70 Gew.-% und einen Propylenoxidgehalt von 30 bis 70 Gew.-% aufweist und
- vi) Gemischen davon,

20 wobei das Mittel frei von proteinhaltigen Materialien ist.

2. Mittel nach Anspruch 1, dadurch gekennzeichnet, daß der ablauffördernde nichtionische oberflächenaktive Stoff die Formel (i) aufweist, worin R einen $C_{16}-C_{22}$ -aliphatischen Kohlenwasserstoffrest bedeutet und n einen Wert von 18 bis 50 aufweist.

25

Revendications


1. Composition détergente liquide aqueuse contenant de 10 à 50% en poids d'un tensio-actif anionique, de 2% à 20% en poids d'un tensio-actif non ionique éthoxylé, de 1,5 à 10% d'un agent de
30 stabilisation des mousses, choisi parmi:

a) les amides de formule générale:



(dans laquelle R_1 représente un radical d'hydrocarbure aliphatique en C_7-C_{21} , saturé ou non saturé; R_2 représente un groupe méthylène ou éthylène et m vaut 1, 2 ou 3);

b) les oxydes d'amines de formule générale:



45


(dans laquelle R_3 représente un radical alkyle, hydroxy-2 alkyle, hydroxy-3 alkyle ou alcoxy-3 hydroxy-2 propyle, dans lequel les groupes alkyles et alcoxy ont respectivement 8 à 18 atomes de carbone; R_4 et R_5 représentent des groupes alkyles en C_1-C_3 ou hydroxyalkyles en C_2-C_3 , et n vaut 0 à 10), et

50 c) leurs mélanges;

de 0% à 10% d'un sel minéral ou organique qui est un adjuvant de détergence; et au moins 20% d'eau; ladite composition étant caractérisée en ce que le tensio-actif non ionique éthoxylé est un tensio-actif non ionique éthoxylé favorisant l'égouttage et qui est choisi parmi:

55


i) un alcool aliphatique éthoxylé de formule:



60

(dans laquelle R représente un radical hydrocarbyle aliphatique contenant de 16 à 30 atomes de carbone, et n vaut de 16 à 100);

ii) un alkyl phénol éthoxylé de formule:



65

O 034 039

(dans laquelle R représente un radical alkyl-phénol contenant au total de 18 à 30 atomes de carbone et au moins un groupe alkyle contenant au moins 12 atomes de carbone, et n vaut de 16 à 100);

- 5 iii) le produit de condensation de monoesters d'acides gras en C_{16} — C_{22} et de polyols avec de 13 à 100 moles d'oxyde d'éthylène par mole de monoester;
- iv) le produit de condensation du cholestérol et de 13 à 100 moles d'oxyde d'éthylène;
- v) une substance qui est un produit de condensation de l'oxyde d'éthylène, de l'oxyde de propylène et d'un composé contenant des groupes hydroxyle ou amine sur lesquels les oxydes d'alkyle peuvent se polymériser, ledit polymère ayant un poids moléculaire de 500 à 15 000, une teneur en oxyde d'éthylène de 30 à 70% en poids et une teneur en oxyde de propylène de 30 à 70% en poids; et
- 10 vi) leurs mélanges;

ladite composition ne comportant pas de matières protéiniques.

2. Composition selon la revendication 1, caractérisée en ce que le tensio-actif non ionique
- 15 favorisant l'égouttage possède la formule (i) dans laquelle R est un radical hydrocarbyle aliphatique en C_{16} — C_{22} et n vaut de 18 à 50.

20

25

30

35

40

45

50

55

60

65