

12

EUROPEAN PATENT APPLICATION

21 Application number: 81200669.0

51 Int. Cl.³: **C 11 D 9/02**
C 11 D 17/08

22 Date of filing: 16.06.81

30 Priority: 20.06.80 GB 8020159

43 Date of publication of application:
30.12.81 Bulletin 81/52

84 Designated Contracting States:
AT BE CH DE FR GB IT LI NL SE

71 Applicant: **UNILEVER NV**
Burgemeester 's Jacobplein 1
NL-3000 DK Rotterdam(NL)

84 Designated Contracting States:
BE CH DE FR IT LI NL SE AT

71 Applicant: **UNILEVER PLC**
Unilever House Blackfriars P O Box 68
London EC4P 4BQ(GB)

84 Designated Contracting States:
GB

72 Inventor: **Brouwer, Hendrik Willem**
Johan Willem Frisolaan 30
NL-3136 BG Vlaardingen(NL)

72 Inventor: **Hemmes, Simon Nico**
Jan van Nassastraat 18
NL-3331 BG Zwijndrecht(NL)

72 Inventor: **van de Griend, Jacob Adriaan**
Eindhovenreef 75
NL-3146 VH Maassluis(NL)

74 Representative: **Van Gent, Jan Paulus et al,**
Unilever N.V. Patent Division P.O. Box 137
NL-3130 AC Vlaardingen(NL)

54 **Aqueous, soap-based liquid detergent composition.**

57 A liquid, soap-based detergent composition having a satisfactory detergency over a broad temperature range, being less sensitive to water hardness, is obtained by using a linoleate soap which is substantially free from lauric soaps. Mixtures of oleate/linolate soaps, optionally with small amounts of stearic soaps, can equally be used. These soap blends are better soluble than coconut/oleate soap blends.

EP 0 042 648 A1

AQUEOUS, SOAP-BASED LIQUID DETERGENT COMPOSITION

The present invention relates to an aqueous liquid detergent composition, in which the detergent-active material is or comprises a soap.

- 5 Soap-based liquid detergent compositions are well-known in the art. Because of the water-solubility limits of the commonly-known soaps, such soap-based liquids contain either a relatively low concentration of soap, when soap is the sole detergent-active material in these liquids, or they contain, next to the soap, another synthetic
10 detergent-active material, by the presence whereof the concentration of soap in the liquid can be made higher.

- Soap-based liquid detergent compositions often suffer from physical drawbacks; thus, for example, it is a problem to prepare a liquid
15 composition which is physically stable, has a satisfactory pourability or viscosity, or which does not gel. Overcoming these problems often creates another one, i.e. that in doing so the washing and laundering efficacy of the composition is impaired.

- 20 Hence it is an object of the present invention to provide an aqueous, soap-based liquid detergent composition which is physically stable, has satisfactory pourability or viscosity characteristics, which does not gel and which has a satisfactory washing, laundering, and cleaning performance.

25

- It is another object of the present invention to provide for such an aqueous, soap-based liquid detergent composition which has a satisfactory washing and laundering performance over a broad temperature range, i.e. at low, medium and high washing or laun-
30 dering temperatures.

These and other objects of the invention have been found to be met by using in the soap component a sodium soap of linoleic acid. It has been found that if the soap component contains a sodium

linoleate (a further definition of the soap component will follow below), the liquid detergent composition has a satisfactory detergency over a broad temperature range and can contain a relatively high concentration of soap. The detergency efficacy is less
5 impaired by the water hardness than with soap systems not containing the linoleate.

As stated above, it is an essential feature that in the soap component a sodium linoleate is used. The soap component may consist
10 solely of sodium linoleate, which may in practice, however, meet with cost and availability problems. Therefore - and this is a preferred embodiment of the invention - it has been found that if the soap component, next to the sodium linoleate, also contains a sodium soap of a C_{16} - C_{24} mono-unsaturated fatty acid, the same
15 benefits are obtained to practically the same degree. Thus, it has been found that if the soap component contains a mixture of sodium linoleate and sodium oleate, in such a relative weight proportion that at least 10% by weight of the sum of these two soaps consists of sodium linoleate, quite satisfactory liquid products are obtained,
20 particularly as regards its detergency over a broad temperature range. Such mixtures can e.g. be obtained from the soapstock fatty acids, obtained from refining and margarine industries. Instead of, or in addition to sodium oleate, a sodium soap of palmitoleic, petroselinic, vaccenic, gadoleic, eicosenoic, cetoleic, erucic and
25 selacholeic acid can also be used.

The soap component may consist solely of the mixture of sodium linoleate and sodium oleate, but it has been found that certain amounts of certain saturated fatty acid sodium soaps can be
30 tolerated without significantly impairing the benefits of the present invention. Thus it has been found that the soap component may contain up to 20% by weight of a saturated C_{16} - C_{24} fatty acid soap. However, this saturated fatty acid soap may not be a sodium laurate or sodium myristate, since the presence thereof significantly detracts from the benefits of the present invention. Small
35 amounts, i.e. up to 5, preferably 3%, can be tolerable, provided there is sufficient linoleate or linoleate plus oleate present, but it is highly preferred that no sodium laurate or

sodium myristate is present in the soap component. Typical examples of sodium soaps of saturated C_{16} and high fatty acids are sodium palmitate, sodium margarate, sodium stearate, sodium nonadecylate, sodium arachidate, sodium heneicosinate, sodium behenate, sodium tricosanate, sodium lignocerate, and sodium soaps of trans-fatty acids of hardened oils and fats.

Mixtures of the above-identified soaps, i.e. sodium linoleate plus sodium oleate plus up to 20% of a soap of a saturated C_{16} - C_{24} fatty acid can be obtained from the separate fatty acids, but may also be obtained from suitable sources which contain these fatty acids in the appropriate proportions. Such sources are grapeseed oil, chufa oil, groundnut oil, maize oil, mustard seed oil, soybean oil, sesame oil, sunflower oil, cottonseed oil, rapeseed oil and tall oil fatty acids. Of these, the last-mentioned five oils and the tall oil fatty acids are preferred. The sodium soaps may be prepared by neutralizing the fatty acids, obtained from these sources and this is preferred when tall oil fatty acids are used.

In the case of the other suitable oils, however, it has been found that the sodium soap can also be prepared in a practical and advantageous manner by carrying out saponification of the oil in the presence of suitable emulsifying agents and neutralization of the fatty acids in situ in the liquid detergent composition. In this manner, an easy way of preparing the compositions of the invention is provided, leading to physically stable products.

It is essential in this process that an emulsifying agent is present in the liquid in which the saponification takes place. Such emulsifying agents are e.g. nonionic detergent surfactants like the ethoxylation and/or propoxylation products of linear or branched-chain primary or secondary, natural or synthetic alcohols having 8-24 carbon atoms in their alkyl chain, of mono- or dialkylphenols having from 8-18 carbon atoms in thier alkyl chain, of C_8 - C_{22} fatty acids, of polyethylene- or polypropylene-glycols, of C_8 - C_{22} fatty amines, -amides or -alkylolamides and so on. Tertiary amine oxides having one C_8 - C_{18} alkyl chain and two C_1 - C_5 alkyl chains are also included in the term nonionic detergent. Amphoteric and zwitterionic detergents may also be used, such as sulphobetaines, amido-betaines,

alkylaminocarboxylic acids etc.

The amount of such emulsifying agent is not critical; in fact it is often desirable to add more than is necessary for the saponification, i.e. in the cases where the presence of a nonionic detergent in the compositions of the invention is required for further improved washing and laundering efficacy, e.g. in underbuilt situations as well as physical stability. In general therefore the compositions of the invention contain from 0-40% by weight of a nonionic detergent material (0% only in the case where the sodium linoleate or sodium linoleate/sodium oleate constitutes the whole of the soap component, and is not prepared by in situ saponification and neutralization) and preferably from 5-30% of a nonionic detergent material is present.

The amount of the soap component in the liquid composition is generally from 2.5-40, preferably 5-35% by weight.

The compositions of the invention may furthermore contain hydro-tropes, buffers, fluorescers, soil-suspending agents, enzymes, stabilizers for enzymes, dyes, perfumes, anti-oxidants and so on, all in amounts usually encountered in products of this type. Sequestrant builders can also be included in an amount of up to 20% by weight.

The invention, as well as its benefits, will further be illustrated by the following Examples.

Example 1

The following liquid detergent compositions were formulated

	% by weight	
Tall oil fatty acids	25	30
C ₁₂ -C ₁₃ p-alcohol, condensed with 6.5 moles of ethylene oxide	25	10
NaOH (33% aqueous solution)	15	18
triethanolamine	10	10
isopropyl alcohol	10	10
enzyme (Alcalase ^(R))	0.3	0.3
dye and perfume	0.25	0.25
boric acid	2	2
water	balance	balance

These liquid compositions were stable: they remained homogeneous after the two weeks' storage at 0°C.

Example 2

5 In the formulations of Example 1, the tall oil fatty acids were replaced by fatty acids, derived from soybean-, sunflower-, rape-seed- and cottonseed oil, and mixtures thereof. This was done in the following way: the mixture of the lye, the triethanolamine, the nonionic detergent and water was heated till 80°C, after which
10 the oil was added. Subsequently, after cooling till 60°C, the isopropyl alcohol was added, and subsequently, after further cooling, the enzyme and the other ingredients.

Similar formulations were prepared in the same manner, with varying amounts of soap and nonionic detergent. These formulations were
15 stored for 2 weeks at 0°C, and their phase stability was assessed.

The following results were obtained:

With soybean oil fatty acids: the liquids were stable at the 10%
20 soap level in the whole range of 10-25% of the nonionic detergent. With sunflower oil fatty acids: the liquids were stable at a 10 and 15% soap level in the whole range of 10-25% nonionic detergent, and also at the 20% soap level with 15% nonionic detergent.

25 With rapeseed oil fatty acids, the liquids were all stable in the whole soap range of 10-25% and in the whole nonionic detergent range of 10-25%. With cottonseed oil fatty acids, stability was obtained at the 10% soap level in the nonionic range of 10-25%, as well as at the 15% soap level and 15% nonionic level. Mix-
30 tures of soybean- and rapeseed oil fatty acids (ratio of 1 to 1) gave products, which were stable at nearly all soap and non-ionic levels within the 10-25% range.

Example 3

35 Using coconut dimethyl amine oxide (5%) in the formulation of Example 1 as the nonionic with 5, 10, 15 and 20% sodium soap of soybean oil fatty acids (saponified and neutralized in situ)

gave stable products.

Example 4

The following liquid compositions C and D were assessed for detergency in a Tergotometer at 60°C, 15 minutes washing, hardness 26° French hardness (only Ca²⁺) at a dosage of 8, 9.5 and 11 g/l. Three types of soiled standard test pieces (WFK, ERTC and VCD) were used, and the number of repeats was 8.

The compositions were as follows:

		% by weight	
		C	D
	sodium oleate	12.5	12.5
	sodium linoleate	12.5	-
15	sodium coconut soap	-	12.5
	triethanolamine	10.0	10.0
	isopropylalcohol	10.0	10.0
	water	balance	balance

The following differences in reflectance (ΔR) were measured:

		concentr.			
Test piece		composition	8	9.5	11
25	WFK	D	5.0	6.9	11.9
		C	7.2	10.3	14.7
	ERTC	D	3.3	11.1	22.7
		C	2.9	19.5	22.7
30	VCD	D	3.3	13.4	16.9
		C	4.6	16.7	18.5

C L A I M S

1. An aqueous soap-based liquid detergent composition, characterized in that it comprises from 2.5 - 40% by weight of a soap component which is or comprises sodium linoleate, the composition not containing more than 5% by weight of sodium laurate or sodium myristate.
5
2. A composition according to claim 1, characterized in that the soap component comprises a mixture of sodium linoleate and a sodium salt of a C_{16} - C_{24} mono-unsaturated fatty acid.
10
3. A composition according to claim 2, characterized in that the soap component comprises a mixture of sodium linoleate and sodium oleate, the sodium linoleate being at least 10% by weight of the mixture.
15
4. A composition according to any one of claims 1 - 3, characterized in that the soap component further comprises up to 20% by weight of a saturated C_{16} - C_{24} fatty acid.
- 20 5. A composition according to any one of claims 1 - 4, characterized in that it further contains up to 40% by weight of a nonionic detergent.
- 25 6. A composition according to any one of claims 1 - 5, further comprising up to 20% by weight of sequestrant builders.
7. A process for the preparation of a composition according to claim 1, in which the soap component is prepared from the corresponding oil by means of saponification of the oil in the presence of an emulsifying agent and neutralization of the fatty acids in situ in the liquid detergent composition.
30



European Patent
Office

EUROPEAN SEARCH REPORT

0042648

Application number
EP 81 20 0669

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
	<u>US - A - 4 065 398</u> (H.W. BROUWER) * Whole document *	1-6	C 11 D 9/02 17/08
	--		
	<u>US - A - 3 503 896</u> (A.E. FISHMAN) * Whole document *	1,2	
	--		
	<u>FR - A - 1 602 731</u> (HENKEL) * Page 1, lines 31-36; examples and claims *	1,5	TECHNICAL FIELDS SEARCHED (Int. Cl.)
	--		
A	<u>FR - A - 1 526 318</u> (C. PELIZZA) * Whole document *	1	C 11 D 9/00 9/02 10/04 17/08
	--		
A	<u>FR - A - 1 077 641</u> (UNILEVER) * Whole document *	1	
	--		
			CATEGORY OF CITED DOCUMENTS
			X: particularly relevant A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention E: conflicting application D: document cited in the application L: citation for other reasons
			&: member of the same patent family. corresponding document
The present search report has been drawn up for all claims			
Place of search	Date of completion of the search	Examiner	
The Hague	13-07-1981	GOLLER	