

12

**EUROPEAN PATENT APPLICATION**

21 Application number: 82304461.5

22 Date of filing: 24.08.82

51 Int. Cl.<sup>3</sup>: **G 03 C 7/32, G 03 C 7/34,**  
**G 03 C 7/26, G 03 C 7/00,**  
**C 07 C 147/10, C 07 C 147/14**  
**// G03C7/16**

30 Priority: 25.08.81 US 296086

71 Applicant: **EASTMAN KODAK COMPANY, 343 State Street, Rochester, New York 14650 (US)**

43 Date of publication of application: 09.03.83  
Bulletin 83/10

72 Inventor: **Lestina, Gregory James, 42 Alta Vista Drive, Rochester New York 14625 (US)**

84 Designated Contr. States: CH DE FR GB LI

74 Representative: **Pepper, John Herbert et al, KODAK LIMITED Patent Department P.O. Box 114 190 High Holborn, London WC1V 7EA (GB)**

54 **Photographic elements containing ballasted couplers.**

57 A photographic element comprising a support, a photographic silver halide emulsion and a nondiffusible photographic coupler, characterized in that said coupler contains attached to a position other than the coupling position a ballast terminated with a hydroxyphenyl-sulfonyl group or a hydroxyphenylsulfinyl group.

**EP 0 073 636 A1**

PHOTOGRAPHIC ELEMENTS  
CONTAINING BALLASTED COUPLERS

5 This invention relates to photographic elements containing ballasted couplers.

Images are commonly obtained in the photographic art by a coupling reaction between the development product of a silver halide developing agent (i.e., oxidized aromatic primary amino developing agent) and a color forming compound commonly referred to as a coupler. The dyes produced by coupling are indoaniline, azomethine, indamine or indophenol dyes, depending upon the chemical composition of the coupler and the developing agent. The subtractive process of color formation is ordinarily employed in multicolor photographic elements and the resulting image dyes are usually cyan, magenta and yellow dyes which are formed in or adjacent silver halide layers sensitive to radiation complementary to the radiation absorbed by the image dye; i.e, silver halide emulsions sensitive to red, green and blue radiation.

15 Since this is a mature art, the patent and technical literature is replete with references to compounds which can be used as couplers for the formation of photographic images. Preferred couplers which form cyan dyes upon reaction with oxidized color developing agents are phenols and naphthols. Representative couplers are described in the following patents and publications: U.S. Patents  
25 2,772,162; 2,895,826; 3,002,836; 3,034,892;  
30 2,474,293; 2,423,730; 2,367,531; 3,041,236 and  
"Farbkuppler-ein Literaturubersicht," published in  
Agfa Mitteilungen, Band II, pp. 156-175 (1961).

35 Preferred couplers which form magenta dyes upon reaction with oxidized color developing agent

are pyrazolones, pyrazolotriazoles, pyrazolobenzimidazoles and indazolones. Representative couplers are described in such patents and publications as U.S. Patents 2,600,788; 2,369,489; 2,343,703; 5 2,311,082; 2,673,801; 3,152,896; 3,519,429; 3,061,432; 3,062,653; 3,725,067; 2,908,573 and "Farbkuppler-eine Literaturubersicht," published in Agfa Mitteilungen, Band II, pp. 126-156 (1961).

10 Couplers which form yellow dyes upon reaction with oxidized color developing agent are acylacetanilides such as benzoylacetanilides and pivalylacetanilides. Representative couplers are described in the following patents and publications: U.S. Patents 2,875,057; 2,407,210; 3,265,506; 15 2,298,443; 3,048,194; 3,447,928 and "Farbkuppler-eine Literaturubersicht," published in Agfa Mitteilungen, Band II, pp. 112-126 (1961).

20 Also known are couplers which form black or neutral dyes upon reaction with oxidized color developing agent. Representative couplers are resorcinols and m-aminophenols such as are described in U.S. Patents 1,939,231; 2,181,944; 2,333,106; 4,126,461; German OLS 2,644,194 and German OLS 2,650,764.

25 Also known are compounds which react with oxidized color developing agent in the same way as couplers but which do not yield a dye. Such compounds are employed to modify the photographic image by competing with dye-forming coupler for oxidized 30 color developing agent or by releasing a photographic reagent, such as a development inhibitor, as a result of the coupling reaction. While many such compounds are not commonly referred to as couplers, it is convenient to consider them as such in view of the 35 similarities in the ways they and couplers react during photographic processing. For the purposes of

the present invention, they are considered couplers. Representative couplers are described in such patents and published patent applications as U.S. Patents 3,632,345; 3,928,041; 3,938,996; 3,958,993; 5 3,961,959; 4,010,035; 4,029,503; 4,046,574; 4,049,455; 4,052,213; 4,063,950; 4,075,021; 4,121,934; 4,157,916; 4,171,223; 4,186,012 and 4,187,110; U.K. Patent Specifications 1,445,797; 1,504,094; 1,536,341 and 2,032,914A; German OLS's 10 2,448,063; 2,552,505; 2,610,546 and 2,617,310; and Belgian Patent 839,083.

When intended for incorporation in photographic elements, couplers are commonly dispersed therein with the aid of a high boiling organic solvent, referred to as a coupler solvent. Couplers are 15 rendered nondiffusible in photographic elements, and compatible with coupler solvents, by including in the coupler molecule a group referred to as a ballast group. This group is located on the coupler in a 20 position other than the coupling position and impart to the coupler sufficient bulk to render the coupler nondiffusible in the element as coated and during processing. It will be appreciated that the size and 25 nature of the ballast group will depend upon the bulk of the unballasted coupler and the presence of other substituents on the coupler.


Although numerous couplers are known in the art, there is a continuing problem to improve, or optimize for particular applications, many properties 30 of the coupler and the resultant dye.

It is an object of this invention to provide novel photographic elements that contain couplers that have improved stability, reactivity and compatibility with other components in the photographic 35 element, the dyes derived from such couplers having efficient light absorption and good stability and hue.

Such can be accomplished by a photographic element comprising a support, a photographic silver halide emulsion and a nondiffusible photographic coupler which reacts with oxidized color developing agent to give a compound which may or may not be an image dye, characterized in that the coupler contains attached to a position other than the coupling position a ballast terminated with a hydroxyphenylsulfonyl group or a hydroxyphenylsulfinyl group.

The coupling group of the couplers used in the photographic elements of the invention can be any coupling group known or used in the art to form a colored or colorless reaction product with oxidized color developing agent. The ballast group of the couplers used in the invention can be any ballast, or portion thereof, which is terminated with a hydroxyphenylsulfonyl or hydroxyphenylsulfinyl group.

Preferred couplers used in the invention have the structural formula:



where:

- COUP represents a coupling group;
- p is 1 or 2;
- q is 1 to 3; and

25 L is a direct linkage (i.e., a covalent bond) or a bivalent linking group.

The coupling group represented by COUP can be any coupling group commonly used in photographic elements. The remainder of the molecule shown in formula I can be joined to the coupling group at any position, other than the coupling position, where ballast groups commonly are joined. The coupling position of the coupling group can be unsubstituted, or substituted with a coupling off group which can

modify the equivalency of the coupler, its reactivity, its dispersibility or which, upon release from the coupler, interacts with other components of the element. The coupling group can include substituents  
5 in other positions.

The bivalent linking group represented by L can be any of the groups found in ballast groups, such as alkylene of 1 to 10 carbon atoms, arylene of 6 to 10 carbon atoms, heterocyclene of 5 to 10 carbon  
10 atoms, oxygen, sulfur, amino, amido, sulfonamido, carbamoyl, sulfamoyl, and combinations of such linking groups, e.g., alkarylene, aralkylene, aminoarylene, aminoalkylene, amidoarylene, amido-  
alkylene, ureido, alkarylamido, amidoarylsulfamoyl,  
15 aminoarylamido and aminoarylsulfamoylalkyl.


Other preferred couplers used in the invention have the structural formula:

20

25

30

35




where:


5

COUP is as defined above;  
 1, m and n are each individually 0 or 1;  
 L<sup>1</sup> represents a bivalent group selected from

10




15




L<sup>2</sup> represents a bivalent group selected from

20



L<sup>3</sup> represents a bivalent group selected from

25



30

R<sup>1</sup> and R<sup>3</sup> are each individually hydrogen, alkyl of 1 to 20 carbon atoms or aryl of 6 to 20 carbon atoms;

R<sup>2</sup> is hydrogen or one or more halogen, alkyl or alkoxy substituents;

35

X is -O- or -S- ;

O


||

Q is -C- or -SO<sub>2</sub> ;

5 r is 0 or 1; and





s is 0 to 10.

Particularly preferred couplers used in the invention have the structural formula:



where:

15 L<sup>4</sup> represents a bivalent group selected from



COUP, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, x, r and s are as defined above.

35 In an especially preferred embodiment, the


hydroxy group in structural formulae II and III is in the para position.

5 In the above structural formulae the alkyl, alkylene, aryl, arylene and heterocyclene groups can be unsubstituted or substituted with one or more groups such as halogen, nitro, amino, carboxy, alkyl, alkoxy, aryl, aryloxy, heterocyclyl, carbamoyl, amido, sulfamoyl and sulfonamido.


10 As indicated above, common yellow dye-forming couplers are acylacetanilides such as pivalylacetanilides and benzoylacetanilides. Common magenta dye-forming couplers are pyrazolones, pyrazolotriazoles, pyrazolobenzimidazoles and indazolones. Common cyan dye-forming couplers are phenols and naphth-  
15 ols, common neutral dye-forming couplers are resorcinols and m-aminophenols. Common non-dye-forming couplers are acyclic and cyclic compounds in which the active position, corresponding to the coupling position, is adjacent to or in conjugation with a  
20 carbonyl group or an imino group, such as  $\alpha$ - or  $\gamma$ -substituted ketones or imines, e.g. cyclopentanones, cyclohexanones, indanones, indanoimines, oxyindoles and oxazolinones. These couplers can form the coupling group COUP in the above formulae.  
25 Structures of representative coupling groups are shown below. In these structures Z represents hydrogen or a coupling-off group and the unsatisfied bond, or bonds, indicates the preferred position, or positions, at which there can be attached the remainder  
30 of the molecule shown in the above structures; it being recognized that the coupling group can contain other substituents. Typical suitable coupling groups that can be used in the photographic elements of the invention are set out below.

35 Cyan dye-forming coupling groups:

5




10




Magenta dye-forming coupling groups:

15




20




25


(B represents a blocking group capable of being removed during processing, e.g., by alkaline cleavage or coupling)

30




35






5

Yellow dye-forming coupling groups:



10




15


Non-dye-forming coupling groups:




20



25




30




35

5




Neutral dye-forming coupling groups:

10




15



Specific couplers used in the invention, which are shown below, contain ballasts of general structures B<sup>1</sup> through B<sup>6</sup>, where Y is -OH.

20




B<sup>1</sup> : R<sup>3</sup> = C<sub>4</sub>H<sub>9</sub>-r

B<sup>2</sup> : R<sup>3</sup> = C<sub>10</sub>H<sub>21</sub>-r


B<sup>3</sup> : R<sup>3</sup> = C<sub>12</sub>H<sub>25</sub>-r

25




B<sup>4</sup>

30



B<sup>5</sup>


35


















B<sup>6</sup>












5 Cyan dye-forming couplers used in the invention include the following; the group Y in B<sup>2</sup>, B<sup>3</sup> and B<sup>4</sup> being -OH:

10



	Coupler No.	R <sub>4</sub>	R <sub>5</sub>	Z
15	C-1	n-C <sub>3</sub> H <sub>7</sub> CO-	-B <sup>2</sup>	-H
	C-2	B <sup>3</sup> -	-COC <sub>3</sub> H <sub>7</sub> -n	-H
	C-3	B <sup>3</sup> -	-COCF <sub>3</sub>	-H
	C-4	B <sup>3</sup> -	-COC <sub>3</sub> F <sub>7</sub> -n	-H
20	C-5	B <sup>2</sup> -	-CO- 	-Cl
25	C-6	B <sup>4</sup> -	-CO- 	-Cl
	C-7	B <sup>2</sup> -	-CONH-  -CN	-H
30	C-8	B <sup>4</sup> -	-CONH-  -CN	-H
	C-9	B <sup>2</sup> -	-CONH-  -SO <sub>2</sub> NHC <sub>4</sub> H <sub>9</sub> -n	-H
35	C-10	B <sup>2</sup> -	-CONH-  -SO <sub>2</sub> NH- 	-H

Coupler No.	R <sub>4</sub>	R <sub>5</sub>	Z
5 C-11	B <sup>2</sup> -		-OC <sub>6</sub> H <sub>5</sub>
10 C-12	B <sup>2</sup> -		-H
15 C-13	B <sup>2</sup> -		-OCH <sub>3</sub>
20 C-14	B <sup>2</sup> -		-OC <sub>6</sub> H <sub>5</sub>
25 C-15	B <sup>2</sup> -		-Cl
30 C-16	B <sup>2</sup> -		-OC <sub>6</sub> H <sub>5</sub>
35 C-17	B <sup>2</sup> -		-Cl
C-18	B <sup>2</sup> -		-OC <sub>6</sub> H <sub>5</sub>


Coupler No.	R <sub>4</sub>	R <sub>5</sub>	Z
5 C-19	B <sup>4</sup> -		-OC <sub>6</sub> H <sub>5</sub>
C-20	B <sup>2</sup> -		-OC <sub>6</sub> H <sub>5</sub>
10 C-21	B <sup>2</sup> -		-OC <sub>6</sub> H <sub>5</sub>
C-22	B <sup>2</sup> -		-OC <sub>6</sub> H <sub>5</sub>
15 C-23	B <sup>2</sup> -		-OC <sub>6</sub> H <sub>5</sub>
20 C-24	B <sup>2</sup> -		
25 C-25	B <sup>2</sup> -		-OC <sub>6</sub> H <sub>5</sub>
C-26	B <sup>2</sup> -		
30 C-27	B <sup>2</sup> -		-OC <sub>6</sub> H <sub>5</sub>
35			

Coupler No.	R <sub>4</sub>	R <sub>5</sub>	Z
-------------	----------------	----------------	---

5


C-28

B<sup>2</sup> -



-OC<sub>6</sub>H<sub>5</sub>

10



Coupler No.	R <sup>6</sup>	R <sup>5</sup>	W	Z
-------------	----------------	----------------	---	---

15

29

CH<sub>3</sub> -

-B<sup>3</sup>

-H

-H

30

CH<sub>3</sub> -

-B<sup>2</sup>

-H

-H

31

C<sub>2</sub>H<sub>5</sub> -

-B<sup>2</sup>

-Cl


-Cl

20

Magenta dye-forming couplers used in the invention include the following, the group Y in B<sup>2</sup> or B<sup>3</sup> being OH:

Coupler No.	Z
-------------	---


M-1




-H

25

M-2



M-3




-H

30

M-4



35



5

10







15










20

25



30

35

Coupler No.	R <sup>7</sup>	R <sup>8</sup>	Z
M-5	CH <sub>3</sub> -	-(CH <sub>2</sub> ) <sub>2</sub> -  -NHB <sup>2</sup>	-Cl
M-6	CH <sub>3</sub> -	-(CH <sub>2</sub> ) <sub>2</sub> -  -NHB <sup>2</sup>	-OC <sub>6</sub> H <sub>5</sub>
M-7	CH <sub>3</sub> -	-SCH <sub>2</sub> CH <sub>2</sub> -  -NHB <sup>2</sup>	-Cl
M-8	CH <sub>3</sub> -	-(CH <sub>2</sub> ) <sub>3</sub> -  -NHB <sup>2</sup>	-H
M-9	CH <sub>3</sub> -	"	-Cl
M-10	CH <sub>3</sub> -	"	-SC <sub>7</sub> H <sub>15</sub> -n
M-11	CH <sub>3</sub> -	"	
M-12	CH <sub>3</sub> OCH <sub>2</sub> -	"	-Cl
M-13	CH <sub>3</sub> -	"	-OC <sub>6</sub> H <sub>5</sub>
M-14	CH <sub>3</sub> -	"	-SCH <sub>2</sub> CH <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub>
M-15	CH <sub>3</sub> -	"	


Coupler No.	R <sup>7</sup>	R <sup>8</sup>	Z	
M-16	CH <sub>3</sub> -	"		
5	M-17	CH <sub>3</sub> -	"	
	M-18	CH <sub>3</sub> -	"	
10	M-19	CH <sub>3</sub> -	"	
	M-20	CH <sub>3</sub> -	"	
15	M-21	CH <sub>3</sub> -	"	
	M-22	CH <sub>3</sub> -	"	
20	M-23	CH <sub>3</sub> -	"	
	M-24	CH <sub>3</sub> -	"	-H
25	M-25	CH <sub>3</sub> -	"	-Cl
	M-25	CH <sub>3</sub> -	"	
30				
35				

M-26



5


10



Coupler No.	R <sup>9</sup>	R <sup>10</sup>	R <sup>11</sup>	Z	
15	M-27	Cl-	-H	-NHB <sup>9</sup>	-H
	M-28	Cl-	-H	-NHB <sup>2</sup>	-H
	M-29	Cl-	-H	-NHB <sup>2</sup>	-SC <sub>7</sub> H <sub>15</sub> <sup>n</sup>
20	M-30	Cl-	-H	-NHB <sup>2</sup>	-SC <sub>6</sub> H <sub>5</sub>
	M-31	Cl-	-H	-NHB <sup>2</sup>	
25	M-32	(CH <sub>3</sub> ) <sub>2</sub> NSO <sub>2</sub> -	-SO <sub>2</sub> B <sup>5</sup>	-H	-H
	M-33	B <sup>2</sup> NH-	-Cl	-H	-H
	M-34	E <sup>2</sup> NH-	-SO <sub>2</sub> NHCH <sub>3</sub>	-H	-H
30	M-35	B <sup>2</sup> NH-	-SO <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub>	-H	-H
	M-36	B <sup>4</sup> NH-	-SO <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub>	-H	-H
	M-37	B <sup>2</sup> ONH-	-SO <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub>	-H	-H
35		or B <sup>2</sup> N(OH)-			


M-38

5



M-39


10



15

Yellow eye-forming couplers used in the invention include the following, the group Y in B<sup>1</sup>, P<sup>2</sup> or B<sup>6</sup> being -OH:

20



Coupler No.

W

R<sup>1 2</sup>


Z

25

Y-1

Cl-


-B<sup>6</sup>



Y-2

Cl-

-P<sup>1</sup>




30

Y-3

H-


-P<sup>1</sup>



Y-4

Cl-

-B<sup>2</sup>



35

Y-5


Cl-

-B<sup>2</sup>

-H

Coupler No.	W	R <sup>1 3</sup>	Z
Y-6	CH <sub>3</sub> O-	-B <sup>2</sup>	
5 Y-7	Cl-	-B <sup>2</sup>	
10 Y-8	CH <sub>3</sub> O-	-B <sup>2</sup>	

Noncolor forming couplers used in the invention include the following:



Coupler No.	R <sup>1 3</sup>	R <sup>1 4</sup>
20 U-1	B <sup>2</sup> NH-	H-
U-2	H-	B <sup>2</sup> NH-

25 Couplers used in the invention can be prepared by attaching a blocked hydroxyphenylsulfonyl or blocked hydroxyphenylsulfinyl group directly to the coupling group or by attaching such a group to the remainder of the ballast group after which the ballast group is attached to the coupling group. Thereafter the blocking group is removed. Conventional condensation reactions can be employed in

30 joining the various groups which ultimately form the coupler. For many of the couplers used in the invention it is convenient to provide the hydroxyphenylsulfonyl group using a 4,4'-sulfonyldiphenol mono

ether (e.g. benzyl ether) or mono ester (e.g. acetyl ester.) Conventional reaction techniques can be employed to attach such a compound to the remainder of the ballast group and the thus formed ballast group to the coupling group. Thereafter, the blocking group can be removed by hydrogenation (in the case of the ether) or alkaline hydrolysis (in the case of an ester). When the coupler, or the remainder of the ballast group, has an amino group available for reaction, it is convenient to react that amino group with a blocked hydroxybenzene-sulfonyl chloride after which the blocking group is removed.

The couplers used in the invention can be used in the ways and for the purposes that nondiffusible couplers are used in the photographic art.

Typically, the couplers are incorporated in silver halide emulsions and the emulsions coated on a support to form the photographic elements of the invention. Alternatively, the couplers can be incorporated in photographic layers adjacent a silver halide emulsion layer where, during development, the coupler will be in reactive association with development products such as oxidized color developing agent. Thus, as used herein, the term "associated therewith" signifies that the coupler is in the silver halide emulsion or in an adjacent location where, during processing, it will come into reactive association with silver halide development products.

The photographic elements of the invention can be single color elements or multicolor elements. Multicolor elements contain dye image-forming units sensitive to each of the three primary regions of the spectrum. Each unit can be comprised of a single emulsion layer or of multiple emulsion layers sensitive to a given region of the spectrum. The layers

of the photographic element, including the layers of the image-forming units, can be arranged in various orders as known in the art. In an alternative format, the emulsions sensitive to each of the three primary regions of the spectrum can be disposed as a single segmented layer, e.g., as by the use of microvessels as described in Belgian Patent 881,513.

A typical multicolor photographic element of the invention comprises a support bearing a cyan dye image-forming unit comprised of at least one red-sensitive silver halide emulsion layer having associated therewith at least one cyan dye-forming coupler, a magenta dye image-forming unit comprising at least one green-sensitive silver halide emulsion layer having associated therewith at least one magenta dye-forming coupler and a yellow dye image-forming unit comprising at least one blue-sensitive silver halide emulsion layer having associated therewith at least one yellow dye-forming coupler, at least one of the couplers in the element being a coupler as described above. The element can contain additional layers, such as filter layers, interlayers, overcoat layers, subbing layers, and the like.

In the following discussion of suitable materials for use in the emulsions used in the invention, reference will be made to Research Disclosure, December 1978, Item 17643. This publication will be identified hereafter by the term "Research Disclosure."

The silver halide emulsions employed in the photographic elements of this invention can be either negative-working or positive-working. Suitable emulsions and their preparation are described in Research Disclosure Sections I and II and the publications

cited therein. Suitable vehicles for the emulsion layers and other layers of elements of this invention are described in Research Disclosure Section IX and the publications cited therein.

5           In addition to the couplers used in the invention, additional couplers as described in Research Disclosure Section VII, paragraphs D, E, F and G and the publications cited therein can be used. These couplers can be incorporated in the elements and  
10 emulsions as described in Research Disclosure Section VII paragraph C and the publications cited therein.

          The photographic elements of this invention or individual layers thereof, can contain brighteners (see Research Disclosure Section V), antifoggants and  
15 stabilizers (see Research Disclosure Section VI), antistain agents and image dye stabilizer (see Research Disclosure Section VII, paragraphs I and J), light absorbing and scattering materials (see Research Disclosure Section VIII), hardeners (see Research Disclosure Section XI), plasticizers and lubricants (see Research Disclosure Section XII), anti-static agents (see Research Disclosure Section XIII), matting agents (see Research Disclosure Section XVI) and development modifiers (see Research Disclosure  
20 Section XXI).

          The photographic elements of the invention can be coated on a variety of supports as described in Research Disclosure Section XVII and the references described therein.

30           Photographic elements of the invention can be exposed to actinic radiation, typically in the visible region of the spectrum, to form a latent image as described in Research Disclosure Section XVIII and then processed to form a visible dye image  
35 as described in Research Disclosure Section XIX. Processing to form a visible dye image includes the

step of contacting the element with a color developing agent to reduce developable silver halide and oxidize the color developing agent. Oxidized color developing agent in turn reacts with the coupler to yield a dye.

Preferred color developing agents are p-phenylene diamines. Especially preferred are 4-amino-N,N-diethyl-aniline hydrochloride, 4-amino-3-methyl-N,N-diethylaniline hydrochloride, 4-amino-3-methyl-N-ethyl-N- $\beta$ -(methanesulfonamido) ethylaniline sulfate hydrate, 4-amino-3-methyl-N-ethyl-N- $\beta$ -hydroxyethylaniline sulfate, 4-amino-3- $\beta$ -(methanesulfonamido)ethyl-N,N-diethyl-aniline hydrochloride and 4-amino-N-ethyl-N-(2-methoxy ethyl)-m-toluidine di-p-toluene sulfonic acid.


With negative working silver halide this processing step leads to a negative image. To obtain a positive (or reversal) image, this step can be preceded by developing exposed silver halide with a non-chromogenic developing agent without forming a dye, and then uniformly fogging the element to render unexposed silver halide developable. Alternatively, a direct positive emulsion can be employed to obtain a positive image.

Development is followed by the conventional steps of bleaching and fixing, or bleach-fixing, to remove silver and silver halide, washing and drying.

The following examples are included for a further understanding of this invention.

Preparative Example 1

Preparation of Ballast Group Intermediate



To a solution of 90 g (0.31 mol) methyl 2-bromododecanoate and 104.4 g (0.31 mol) 4,4'-sulfonyldiphenol monobenzyl ether in 0.35 L dry acetone were added 1 g sodium iodide and 214.2 g (1.55 mol)

potassium carbonate. After refluxing the mixture 20 h, solids were removed by filtration and the filtrate concentrated to a waxy solid. Recrystallization from methanol gave a white solid ( $B^2OCH_3$ ,  $Y = OBz$ ),  
5 mp 73-75°C, with the correct elemental analysis and expected NMR spectrum. A solution of 120 g (0.21 mol) of this product in 0.8 L dimethylformamide was added with stirring to 0.5 L of 2.3 M aqueous potassium hydroxide solution, water was added and the  
10 cloudy solution stirred 0.5 h before pouring into acidic ice-water. The resulting solid was collected, dissolved in dichloromethane, and the solution washed, dried over magnesium sulfate, and concentrated. Recrystallization from acetonitrile yielded  
15 67 g white solid ( $B^2OH$ ,  $Y = OBz$ ), mp 119-121°C, with the expected NMR spectrum and elemental analysis. This acid was converted to the acid chloride by dissolving 67 g (0.12 mol) in 0.4 L thionyl chloride and stirring 5 h. Excess thionyl chloride was re-  
20 moved under vacuum and the product recrystallized from dry acetonitrile to give a white solid ( $B^2Cl$ ,  $Y = OBz$ ), mp 84-85°C, with the expected NMR spectrum and elemental analysis.

Preparative Example 2

25 Preparation of Ballast Group Intermediate

$B^2Cl$ , where  $Y = OAc$

A solution of 455 g (0.82 mol),  
 $B^2OCH_3$ ,  $Y = OBz$  in 1.6 L tetrahydrofuran and  
0.4 L acetic acid was hydrogenated 12 h at 50 psi and  
30 50°C over 45 g 5% palladium on charcoal catalyst. The catalyst was removed by filtration and the concentrated filtrate drowned in water. An ethyl acetate solution of resulting white solid was washed, dried, concentrated, and the product recrystallized  
35 from acetonitrile to give 340 g (0.74 mol) white

solid ( $B^2OCH_3$ ,  $Y = OH$ ), mp 63-65°C with the expected elemental analysis. Hydrolysis of this ester was accomplished by slowly adding an aqueous solution containing 40 g (1 mol) sodium hydroxide to a stirred solution of the ester in 1 L dimethylformamide, stirring 2 h then pouring into acidified ice-water. The resulting gummy solid dissolved in ethyl acetate was washed with dilute hydrochloric acid, dried, and concentrated. Recrystallization from acetonitrile yielded a white solid ( $B^2OH$ ,  $Y = OH$ ), mp 116-117°C. This phenolic acid was acetylated by dissolving in 70 mL acetic anhydride and 7 mL concentrated sulfuric acid, stirring 30 minutes at 20°C., then on a steam bath for 30 minutes, cooling, and pouring into 8 L water. The product was recrystallized from methanol to give a white solid ( $B^2OH$ ,  $Y = OAc$ ), mp 73-75°C. Refluxing 35 g (0.07 mol) of this acid in excess thionyl chloride for 5 h and concentrating yielded a colorless oil, which on trituration in ligroin gave 22 g white solid ( $B^2Cl$ ,  $Y = OAc$ ), mp 66-69°C.

Preparative Example 3:

Preparation of Ballast Group Intermediate

$B^3Cl$ , where  $Y = OBz$

The procedural steps were similar to those for preparation of  $B^2Cl$  in Preparative Example 1, except that ethyl 2-bromotetradecanoate was the starting material. Intermediates included white solids  $B^3OCH_2CH_3$ ,  $Y = OBz$  (mp 55-61°C);  $B^3OH$ ,  $Y = OBz$  (mp 117-118°C); and  $B^3Cl$ ,  $Y = OBz$  (mp 81-84°C).

Preparative Example 4

Preparation of Ballast Group Intermediate

$B^1Cl$  where  $Y = OBz$

The procedural steps were similar to those for preparation of  $B^2Cl$  in Preparative Example 1,

except that ethyl 2-bromobutyrate was the starting material. Intermediates included white solids  $B^1 OCH_2 CH_3$ ,  $Y = OBz$  (mp 102-105°C);  $B^1 OH$ ,  $Y = OBz$  (mp 147.5-148.5°C); and  $B^1 Cl$ ,  $Y = OBz$  (mp 40°C).

Preparative Example 5

Preparation of Ballast Intermediate

$B^4 Cl$  where  $Y = OBz$

A solution of 10.2 g (0.029 mol) methyl 2-p-nitrophenoxy)dodecanoate in 100 mL tetrahydrofuran was shaken 6 h under 40 psi hydrogen in the presence of 0.7 g 10% palladium on charcoal catalyst to reduce the nitro group. Then 6.3 mL (0.04 mol) N,N-dimethylaniline and 8.2 g (0.029 mol) p-benzyloxybenzenesulfonyl chloride were added and the mixture stirred 15 hours at 20°C. The catalyst was removed by filtration and the filtrate poured into cold dilute hydrochloric acid. Ethyl acetate extraction, washing, drying, concentration, and purification through silica gel yielded 14 g colorless oil ( $B^4 OCH_3$ ,  $Y = OBz$ ). This was dissolved in 60 mL tetrahydrofuran and 40 mL methanol, stirred 0.5 h with 20 mL aqueous sodium hydroxide solution and poured into cold dilute hydrochloric acid. Ethyl acetate extraction, washing, drying, concentration, ligroin trituration yielded 12 g white crystals ( $B^4 OH$ ,  $Y = OBz$ ), mp 100-101°C, with the correct elemental analysis. To a stirred solution of 10 g (0.018 mol) of this acid in 50 mL tetrahydrofuran was added 1.8 mL (0.022 mol) oxalyl chloride and 5 drops dimethyl formamide. After 1.5 h concentration gave 0.018 mol of brown oil  $B^4 Cl$ ,  $Y = OBz$ .

Preparative Example 6

Preparation of Ballast Intermediate

B<sup>5</sup>H·HCl where Y = OH

To a solution of 24 g (0.043 mol) acid chloride  
5 B<sup>2</sup> Cl, Y=OBz in 400 mL tetrahydrofuran was added a  
40% aqueous solution containing 10 g (0.125 mol)  
methylamine. After 0.5 h stirring, the mixture was  
poured over acidified ice-water, extracted with  
diethyl ether, and the organic layer washed, dried,  
10 and concentrated to yield, after further purification  
on a 50:50 silica gel/Fluorisil (trade mark) column, a  
clear colorless oil (B<sup>2</sup>NHCH<sub>3</sub>, Y = OBz). Reduction  
was accomplished by refluxing 16 g (0.029 mol) of  
15 this amide product and 16 mL 2 M borane-methyl sul-  
fide complex in 400 mL tetrahydrofuran for 3 h. The  
cooled reaction mixture was slowly acidified with 50%  
hydrochloric acid solution, then extracted with di-  
ethyl ether. Acidification and concentration of the  
washed and dried organic layer gave 14 g of white  
20 solid (B<sup>5</sup>H·HCl, Y = OH).

Final steps in the synthesis of couplers  
used in this invention generally involved the attach-  
ment of the ballast group and the removal of the  
ballast blocking group, if any. For example, in  
25 Scheme I an amino-substituted coupling group,  
COUP-NH<sub>2</sub>, is allowed to react with an acid  
chloride ballast group and the resulting intermediate  
is converted to the desired coupler by hydrogenation  
to remove the benzyl group.


30

35

Scheme I

(n = 0, 1)

5




15

Where the blocking group is acetyl, as in Scheme II, it is removed by alkaline hydrolysis:

Scheme II

20



30


35

Alternatively, if the coupler is substituted with an acidic function, a ballast group containing

an amine function may be attached according to Scheme III:

Scheme III

5



10

Preparative Example 7:

15

Preparation of Coupler C-8 by Scheme I

A suspension of 5.4 g (0.018 mol) 2-(p-cyanophenylureido)-5-nitrophenol in 200 ml tetrahydrofuran was shaken overnight under 40 psi hydrogen with 1.6 g 10% palladium on charcoal catalyst and 0.3 ml acetic acid. Then 0.018 mol of the acid chloride B<sup>1</sup>Cl prepared in Preparative Example 5 and 6.8 ml dimethylaniline were added under nitrogen and the mixture stirred 0.5 h before removing the catalyst by filtration and pouring the filtrate into cold dilute hydrochloric acid. Ethyl acetate extraction, washing, drying, concentration, and crystallization from acetonitrile yielded 10.2 g of the pale white solid benzyl ether of the desired coupler. A solution of this product in 100 ml tetrahydrofuran was shaken for 15 hours under 40 psi hydrogen with 2.5 g 10% palladium on charcoal catalyst and 0.5 ml acetic acid. The catalyst was removed by filtration and the reduction product concentrated and crystallized from acetonitrile to give 6.1 g white solid coupler C-8, mp 103-106°C, with an infrared spectrum and elemental analysis consistent with the desired structure.

35

Preparative Example 8:

Preparation of Coupler M-28 by Scheme II

A solution of 50 g (0.115 mol) 3-(2-chloro-5-nitroanilino)-1-(2,4,6-trichlorophenyl)-2-pyrazolin-5-one in dimethylformamide and tetrahydrofuran was reduced with 35 psi hydrogen and Raney nickel catalyst. Removal of the catalyst by filtration and concentration of the filtrate gave 21 g (0.052 mol) light yellow solid 3-(2-chloro-5-aminoanilino)-1-(2,4,6-trichlorophenyl)-2-pyrazolin-5-one. To an acetic acid solution containing 6.9 g (0.017 mol) of this amine and 9.5 g potassium acetate was added, in small portions, 9.5 g (0.019 mol) of the acid chloride B<sup>2</sup>Cl prepared in Example 2. After stirring 15 hours, the mixture was concentrated to a third of its volume, poured into a large volume of water and extracted with diethyl ether. The organic layers were washed, dried, concentrated, and crystallized from methanol to give 14 g (0.016 mol) buff-colored solid acetate ester of the desired coupler, mp 115-116°C. To a solution of this product in dimethylformamide stirred under nitrogen was added an aqueous solution containing 2 g potassium hydroxide. After 15 minutes the mixture was acidified with hydrochloric acid, poured into dilute hydrochloric acid, extracted with diethyl ether, washed, dried, concentrated, and crystallized from methanol to give 9.2 g white crystalline coupler M-28, mp 127-130°C.

Preparative Example 9:

Preparation of Coupler M-8 by Scheme I

A suspension of 10 g (0.035 mol) 6-methyl-3-[3-(p-nitrophenyl)-propyl]-1H-pyrazole[3,2-c]-5-triazole in 300 mL tetrahydrofuran was shaken about 2 h

at 25°C with 35 psi hydrogen and a palladium on charcoal catalyst. Removal of the catalyst, concentration of the filtrate, and recrystallization from acetonitrile gave a buff-colored solid amine, mp  
5 194-6°C. To a stirred acetic solution of 7.6 g (0.03 mol) of this amine product and 2 g potassium acetate was added, in small portions, 16.7 g (0.03 mol) of the acid chloride B<sup>2</sup>Cl prepared in Example 1. After stirring 15 h, the mixture was poured into a  
10 large volume of water, extracted with diethyl ether, and the combined extracts then washed, dried, and concentrated to give the white solid benzyl ether of the desired coupler, mp 122-124°C. A concentrated solution of 7 g (0.009 mol) of the product in tetra-  
15 hydrofuran was shaken 2 h under 40 psi hydrogen with a mixture of palladium on charcoal catalyst in ethanol. Removal of the catalyst by filtration, concentration, and recrystallization from acetonitrile gave 5.5 g cream-colored solid coupler M-32,  
20 mp 170-172°C, with an infrared spectrum and elemental analysis consistent with the desired compound.

Preparative Example 10:

Preparation of Coupler M-32 by Scheme III

A mixture of 8 g (0.015 mol) 3-(2-chloro-  
25 4-fluorosulfonylanilino)-1-(2,6-dichloro-4-dimethylsulfamoyl phenyl)-2-pyrazolin-5-one and 5.5 g aluminum chloride in 175 mL 1,2-dichloroethane was refluxed 15 minutes. After cooling to 20°C, 7.25 g  
(0.015 mol) of the amine salt B<sup>5</sup>H·HCl prepared in  
30 Example 6 in 25 mL pyridine was added and the mixture refluxed 2 h. Then a mixture of hydrochloric acid, ice, and diethyl ether was added and the organic layer washed, dried, concentrated, and triturated with hexane to give 14.2 g light tan solid coupler M-32  
35 with the correct elemental analysis.

Preparative Example 11:Preparation of Coupler Y-1 by Scheme II

To a solution of 30.3 g (0.05 mol)  $\alpha$ -pivalyl- $\alpha$ -[4-(p-benzyloxyphenylsulfonyl)phenoxy]-2-chloro-5-aminoacetanilide in 150 mL pyridine was added 11.8 g (0.05 mol) p-acetoxybenzenesulfonyl chloride at 20°C. After stirring for 15 hours, the mixture was poured onto 1 L of ice-water containing 50 mL concentrated hydrochloric acid. The resulting solid was collected, dissolved in 250 mL ethanol and treated with an alcoholic solution of potassium hydroxide. After stirring 1 h the mixture was poured over acidified ice-water. The collected solid was then dissolved in boiling benzene and cyclohexane added to give a precipitate. Recrystallization from ethanol-cyclohexane yielded 23 g coupler Y-1, mp 174-175°C, with the correct elemental analysis.

Preparative Example 12:Preparation of Coupler Y-2 by Scheme I

To a solution of 57.8 g (0.095 mol)  $\alpha$ -pivalyl- $\alpha$ -[4-(p-benzyloxyphenylsulfonyl)-phenoxy]-2-chloro-5-amino acetanilide and 13 g quinoline, cooled to 0°C, was added in one portion 45 g (0.095 mol) of the acid chloride B<sup>1</sup>Cl prepared in Preparative Example 4. After stirring 1 h the reaction mixture was poured into 3 L cold water to produce 98 g (after oven drying) crude product. Recrystallization from toluene yielded 84.5 g (0.018 mol) white crystalline dibenzyl ether of the desired coupler. A suspension of 10.4 g (0.01 mol) of this product in 200 mL ethanol and 200 mL tetrahydrofuran was shaken 3 h under 35 psi hydrogen with palladium on charcoal catalyst. Removal of catalyst by filtration, concentration of the filtrate and trituration in hot cyclohexane gave a good yield of coupler Y-2 with the correct elemental analysis.

Examples 1-14

Photographic elements of this invention and control elements were prepared and tested according to the procedures described below.

5 All photographic elements were prepared by coating a cellulose acetate butyrate film support with a photosensitive layer containing a silver bromo-iodide emulsion at 0.91 g Ag/L<sup>2</sup> (when the coupler is 4-equivalent) or 0.46 g Ag/m<sup>2</sup> (when the  
10 coupler is 2-equivalent)., gelatin at 3.78 g/m<sup>2</sup>, and one of the couplers identified in Table I dispersed in one-half its weight of the coupler solvent described and coated at  $1.62 \times 10^{-3}$  moles/m<sup>2</sup>. The photosensitive layer was overcoated  
15 with a layer containing gelatin at 1.08 g/m<sup>2</sup> and bis-vinyl-sulfonylmethyl ether at 1.75 weight percent based on total gelatin.

Samples of each element were imagewise exposed through a graduated-density test object and  
20 processed at 40°C employing one of three color developing solutions identified below then stopped, bleached, fixed and washed.

In each element, well-defined, magenta dye images were produced which were evaluated by plotting  
25 dye density vs. log exposure sensitometric curves and recording the maximum dye density ( $D_{max}$ ) and gamma ( $\gamma$ ) i.e., the contrast determined by the slope of the straight line portion of the curve. Additionally, dye hues were evaluated from spectrophotometric  
30 curves by measuring the maximum absorption peak ( $\lambda_{max}$ ) normalized to a density of 1.0 and the half band width (HBW). Halfband width is the width, in nanometers, of the spectrophotometric curve at one-half the difference between maximum density and  
35 stain. Similarly, the top-band width (TBW) and bottom-band width (BBW) of the curve were measured at

three-fourths and one-fourth, respectively, of the normalized density. Curve shape factor (CSF) equals 100 X TbW/HBW and provides a ratio of the width near the top and bottom of the absorption curve. The greater this ratio, the steeper are the sides of the absorption peak, and the more efficient is the dye's absorption of light in its spectral region.

All results are recorded in Table I.

These results show that couplers of this invention have enhanced activity, which results in increased maximum dye density and gamma. In addition many of the dyes formed from couplers of this invention have absorption maxima at desirably longer wavelengths and have broader half band widths and larger curve shape factors, resulting in more efficient spectral absorption.

20

25

30

35

T A B L E I

Example No.	Coupler No.(1)	Coupler Solvent (2)	Developer (3)	Activity		Hue	
				D-max	Y	$\lambda$ -max	HBW
1	M-3	CS-1	D-1	3.70	1.12	546	96
	CC-1	CS-1	D-1	3.28	1.06	531	94
2	M-8	CS-2	D-3	4.21	2.03	550	86
	CC-3	CS-2	D-3	3.68	1.30	556	88
3	M-27	CS-1	D-1	4.40	2.14	545	91
	CC-1	CS-1	D-1	3.36	1.27	531	94
4	M-28	CS-1	D-1	4.32	2.01	549	92
	CC-1	CS-1	D-1	3.20	1.19	531	94
5	M-28	CS-2	D-2	4.12	1.65	541	93
	CC-2	CS-2	D-2	2.96	1.04	539	81
6	M-28	CS-2	D-3	4.98	1.88	549	94
	CC-3	CS-2	D-3	3.66	1.30	556	88
7	M-29	CS-2	D-2	4.53	2.92	540	88
	CC-2	CS-2	D-2	2.12	0.75	539	79
8	M-29	CS-2	D-3	4.87	3.14	545	90
	CC-2	CS-2	D-3	3.03	1.05	545	79
9	M-32	CS-2	D-2	3.15	1.32	549	82
	CC-2	CS-2	D-2	2.37	0.99	540	78

$\lambda$ -max

D-max

Developer (3)

Coupler Solvent (2)

Coupler No.(1)

Example No.

HBW

CSF


T A B L E 1 (continued)

Example No.	Coupler No. (1)	Coupler Solvent (2)	Developer (3)	Activity		Hue		
				D-max	Y	$\lambda$ -max	HBW	CSF
10	C-31	CS-3	D-2	3.25	1.14	675	160	47.0
	CC-4	CS-3	D-2	2.22	0.78	659	141	45.2
11	Y-1	CS-3	D-2	3.39	1.44	449	88	43.7
	CC-5	CS-3	D-2	3.14	0.98	446	88	43.5
	CC-6	CS-3	D-2	1.78	0.51	441	85	42.0
12	Y-2	CS-3	D-2	2.82	1.13	445	90	44.4
	CC-7	CS-3	D-2	1.92	0.55	445	85	45.2
13	Y-3	CS-3	D-2	1.89	0.62	441	93	44.4
	CC-8	CS-3	D-2	0.57	0.14	439	90	44.3
14	Y-4	CS-3	D-1	3.86	1.90	450	90	44.6
	CC-9	CS-3	D-1	3.69	1.48	450	87	43.8

1) Comparison Couplers:

CC-1


5



10

CC-2


15



20


CC-3

25








30

CC-4



35





(2) Coupler Solvents:

30 CS-1 - 1,4-Cyclohexylenedimethylene bis(2-ethylhexanoate)

CS-2 - Tri-cresyl phosphate

CS-3 - Dibutyl phthalate

CS-4 - 2,4-Di-t-pentylphenol

35

(3) Developer Formulations:

	<u>D-1</u>	<u>D-2</u>	<u>D-3</u>
4-Amino-3-methyl-N,N-di- ethylaniline hydrochloride	2.45 g	--	--
5 4-Amino-3-methyl-N-ethyl- N- $\beta$ -(methanesulfon- amido)ethylaniline sulfate	--	5.0 g	--
10 4-Amino-3-methyl-N-ethyl- N- $\beta$ -hydroxyethyl aniline sulfate	--	--	3.55 g
Potassium sulfite	2.0 g	2.0 g	2.0 g
Potassium carbonate (anhydrous)	30.0 g	30.0 g	30.0 g
15 Potassium bromide	1.25 g	1.25 g	1.25 g
Potassium iodide	0.6 mg	0.6 mg	0.6 mg
1% Solution in methanol of 5-nitro-1H-indazole	4.0 mL	--	--
Water to:	1.0 L	1.0 L	1.0 L
20 pH adjusted to:	10.0	10.0	10.0

Example 15

Photographic elements containing additional couplers were prepared, processed and evaluated as described above in connection with Examples 1-14.

25 The results are reported in Table II below.


TABLE II

<u>Coupler</u>	<u>Coupler Solvent</u>	<u>Dev.</u>	<u>D-max</u>	<u><math>\alpha</math></u>	<u><math>\lambda</math>-max (nm)</u>	<u>HBW (nm)</u>	<u>CSF %</u>
30 C-2	CS-3	D-1	3.78	1.77	655	144	45.7
C-3	CS-3	D-2	3.30	1.28	659	161	47.8
C-13	CS-3	D-3	2.99	1.11	702	135	44.4
M-3	CS-1	D-1	3.70	1.12	546	96	45.7
M-9	CS-4	D-3	4.16	1.81	558	90	44.2
35 M-28	CS-2	D-3	4.58	1.88	549	94	42.6

Example 16

For each of the couplers identified below, photographic elements were prepared as described above in connection with Examples 1-14. Four samples  
5 from each element were exposed as described above. One pair of the exposed elements was developed in developer D-2, described above, and the other pair was developed in this developer to which had been added  
1.5 g/L of the soluble competing coupler citrazinic  
10 acid. The remaining processing for one element from each pair was stopping, bleaching, fixing and washing while for the second element from each pair the bleaching step was omitted so that the developed silver remained in the element. For those elements in  
15 which the silver remained, the amount of developed silver, in  $\text{g/m}^2$ , was determined by x-ray fluorescence analysis and plotted against exposure. For those elements from which the developed silver had been removed, dye density vs exposure curves were  
20 generated. From the plots for pairs of elements developed with the same developer composition there was plotted, for each exposure step, dye density vs developed silver. The slope of the line for the elements developed in the absence of a competing coupler  
25 ( $Y_0$  in Table III, below) is a measure of the efficiency with which the coupler forms dye; the greater the slope the more efficient the coupler. The slope of the line for the elements developed in the presence of the competing coupler ( $Y_c$  in Table III, below)  
30 is a measure of the reactivity of the coupler, the greater the slope, the more reactive the coupler.


The couplers employed had the following structure:



10

Coupler	Q
M-8	-SO <sub>2</sub> -
CC-10	-C(CH <sub>3</sub> ) <sub>2</sub> -

15



25

Coupler	R'	B
M-28	-C <sub>10</sub> H <sub>21</sub> -n	-O-C <sub>6</sub> H <sub>4</sub> -SO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> -OH
CC-11	-C <sub>2</sub> H <sub>5</sub>	-O-C <sub>6</sub> H <sub>4</sub> -C <sub>5</sub> H <sub>11</sub> -t

30

TABLE III

35

Coupler	Coupler Solvent	Y <sub>o</sub>	Y <sub>c</sub>	Y <sub>c</sub> /Y <sub>o</sub>
M-8	CS-4	5.55	1.73	0.312
CC-10	CS-4	1.62	0.21	0.130
M-28	CS-2	6.30	4.65	0.738
CC-11	CS-2	4.50	1.50	0.337

It is apparent from the values for  $Y_o$  and  $Y_c$  in Table III that the couplers used in the invention react more efficiently with oxidized developer to form image dye, in the presence or absence of a competing coupler, than do those couplers with ballasts not used in this invention.

10

15

20

25


30

35

## CLAIMS:

1. A photographic element comprising a support, a photographic silver halide emulsion and a nondiffusible photographic coupler which reacts with  
 5 oxidized color developing agent to give a compound which may or may not be an image dye characterized in that said coupler contains attached to a position other than the coupling position a ballast terminated with a hydroxyphenylsulfonyl group or a hydroxyphenylsulfinyl  
 10 group.


2. A photographic element according to claim 1 wherein the coupler has the structural formula:



where:


15 COUP represents a coupling group;  
 p is 1 or 2;  
 q is 1 to 3; and  
 L is direct linkage or a bivalent linking  
 group.

20 3. A photographic element according to claim 1 wherein the coupler has the structural formula:



where:

25 COUP represents a coupling group;  
 1, m and n are each individually 0 or 1;  
 L<sup>1</sup> represents a bivalent group selected from






where:

COUP represents a coupling group

L<sup>4</sup> represents a bivalent group selected

from



where:

10 R<sup>1</sup> and R<sup>3</sup> are each individually hydrogen, alkyl of 1 to 20 carbon atoms or aryl of 6 to 20 carbon atoms;

R<sup>2</sup> is hydrogen or one or more halogen, alkyl or alkoxy substituents,

15 x is -O- or -S-;

r is 0 or 1; and

s is 1 to 10,

any alkyl, alkylene, aryl, arylene or heterocyclene group possibly being substituted.



20 5. A photographic element according to claim 2, 3, or 4 wherein the coupler is a yellow dye-form-





ing coupler and COUP is an acylacetanilide coupling group.



5 6. A photographic element according to claim 2, 3 or 4 wherein the coupler is a cyan dye-forming coupler and COUP is a phenol or naphthol coupling group.

10 7. A photographic element according to claim 2, 3 or 4 wherein the coupler is a magenta dye-forming coupler and COUP is a pyrazolone, pyrazolo-triazole, pyrazolobenzimidazole or indazolone coupling group.

8. A photographic element according to claim 1 wherein the coupler has the structure:







20

25

30

35



DOCUMENTS CONSIDERED TO BE RELEVANT			EP 82304461.5
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 3)
A	<u>GB - A - 2 029 977 (FUJI)</u> * Claims 1-3, 5; page 2, line 46 - page 3, line 16 * --	1,2	G 03 C 7/32 G 03 C 7/34 G 03 C 7/26 G 03 C 7/00
A	<u>GB - A - 2 038 808 (FUJI)</u> * Claims 1,6-8; page 2, line 58 - page 3, line 41 * --	1,2	C 07 C 147/10 C 07 C 147/14// G 03 C 7/16
A	<u>DE - A1 - 2 529 991 (FUJI)</u> * Claims 1,3-5; page 5, line 28 - page 7, line 11 * ----	1,2	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 3)  G 03 C C 07 C
X The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 03-12-1982	Examiner SCHÄFER
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			