



**EUROPEAN PATENT SPECIFICATION**

Date of publication of patent specification :  
**07.04.93 Bulletin 93/14**

Int. Cl.<sup>5</sup> : **C10L 1/22, C10L 1/14**

Application number : **90301791.1**

Date of filing : **20.02.90**

**Middle distillate fuel having improved storage stability.**

Priority : **02.03.89 US 318748**

Date of publication of application :  
**05.09.90 Bulletin 90/36**

Publication of the grant of the patent :  
**07.04.93 Bulletin 93/14**

Designated Contracting States :  
**BE DE ES FR GB IT NL**

References cited :  
**EP-A- 0 182 940**  
**EP-A- 0 261 795**  
**EP-A- 0 294 045**  
**US-A- 2 928 790**  
**US-A- 3 442 791**  
**US-A- 3 701 641**  
**US-A- 4 025 316**  
**US-A- 4 166 726**  
**US-A- 4 172 707**

Proprietor : **ETHYL PETROLEUM ADDITIVES, INC.**  
**20 South Fourth Street**  
**St. Louis Missouri 63102-1886 (US)**

Inventor : **Bostick, John Gray**  
**742 Klein Drive**  
**Smithtown Illinois 62285 (US)**  
Inventor : **Cunningham, Larry John**  
**1928 Windyhill**  
**Kirkwood Missouri 63122 (US)**  
Inventor : **Hanlon, John Vincent**  
**2610 Bopp Road**  
**St.Louis, Missouri 63131 (US)**

Representative : **Collier, Jeremy Austin Grey et al**  
**J.A.Kemp & Co. 14, South Square Gray's Inn**  
**London WC1R 5EU (GB)**

**EP 0 385 633 B1**

Note : Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).


## Description

This invention relates generally to improving the stability of middle distillate fuels and more particularly to stabilized middle distillate fuel compositions which contain a combination of N,N-dimethylcyclohexyl amine and a Mannich Base.


Middle distillate fuels such as diesel oil, fuel oil, jet fuel and kerosene when stored for long periods of time are subject to the formation of color and solid deposits. The deposits accumulate on filters causing the filters to become plugged. Various additives and combinations of additives have been employed to reduce color and deposit formation. For example: U.S. Patent 2,984,550 discloses the use of Mannich bases derived from phenols, formaldehyde and polyamines for stabilization; U.S. Patent 3,490,882 discloses stabilized petroleum distillate fuel oils containing N,N-dimethylcyclohexylamine antioxidant and a N,N'-di(ortho-hydroxyarylidene)-1,2-alkylenediamine metal deactivator such as N,N'-disalicylidene-1,2-propylenediamine; U.S. Patent 4,166,726 discloses a fuel additive which is a mixture of a polyalkylene amine and a Mannich Base; and U.S. Patents 4,501,595 and 4,533,361 disclose diesel oil which contains a condensate of tetraethylene pentamine, paraformaldehyde, a hindered phenol such as 2,6-di-t-butylphenol and polyisobutenyl succinic anhydride.

The effectiveness of any particular type of additive combination can vary with different fuel stocks and combinations which are more effective at the same total additive concentration reduce treatment cost. We have now discovered novel, synergistic additive combinations which include certain Mannich Bases and provide middle distillate fuels having generally improved storage stability compared to fuels containing the same total concentrations of either N,N-dimethylcyclohexylamine antioxidant alone or N,N-dimethylcyclohexylamine in combination with an N,N'-di(ortho-hydroxyarylidene)-1,2-alkylenediamine metal deactivator.

In accordance with this invention, there is provided a fuel additive concentrate comprising a mixture of N,N-dimethylcyclohexylamine and a Mannich Base which is the reaction product of an aldehyde, an amine and an alkyl phenol selected from (a) hindered phenol having the formula:



where  $R_1$ ,  $R_2$ ,  $R_3$  are independently selected from hydrogen, t-butyl, t-amyl and isopropyl, provided that at least one of  $R_1$ ,  $R_2$  and  $R_3$  is hydrogen and at least one of  $R_1$  and  $R_2$  is t-butyl, t-amyl or isopropyl; and (b) p-alkyl phenol having the formula:



where  $R_4$  is  $C_9$  to  $C_{30}$  alkyl.

The concentrate can also contain a N,N'-di(ortho-hydroxyarylidene)-1,2-alkylenediamine metal deactivator such as N,N'-disalicylidene-1,2-propylenediamine. Also provided is a stabilized fuel containing from 1 to 1400 mg/l of N,N-dimethylcyclohexylamine, from 0.5 to 1100 mg/l of Mannich Base and from 0 to 400 mg/l of an N,N'-di(ortho-hydroxyarylidene)-1,2-alkylenediamine.

The N,N-dimethylcyclohexylamine component of the compositions of the invention is a commercially available fuel antioxidant.

The N,N'-di(ortho-hydroxyarylidene)-1,2-alkylenediamine component, in which, typically, the arylidene radical contains 6-7 carbon atoms and the alkylene radical contains 2-3 carbon atoms, is a metal deactivator

whose presence in combination with the other components provides fuel compositions of the invention having the most improved stability. The preferred metal deactivator is N,N'-disalicylidene-1,2-propylenediamine which is commercially available.


The Mannich Base component of the invention is produced by the Mannich condensation reaction of a hindered or p-alkyl phenol, an aldehyde, such as formaldehyde, ethanal, propanal, and butanal (preferably formaldehyde in its monomeric form or paraformaldehyde) and primary and secondary amines.

The hindered phenols which are useful in preparing the Mannich Base component of the invention are phenols which are characterized by the presence of at least one and preferably two ortho-t-butyl, t-amyl, and/or isopropyl groups. Specific examples of such hindered phenols include: 2,4-di-t-butylphenol, 2,4-diisopropylphenol, 2,6-diisopropylphenol, 2-t-butylphenol, and 2-t-amylphenol with 2,6-di-t-butylphenol being most preferred.

The p-alkyl phenols which are useful in preparing the Mannich Base component of the invention are those which contain from 9 to 30 carbons which can be arranged in either a straight or a branched chain. Preferred phenols are C<sub>9</sub> to C<sub>12</sub> p-alkylphenols such as, for example, p-nonylphenol and p-dodecylphenol.


The amines which are useful in preparing the Mannich Base component of the invention are primary and secondary amines which can be selected from one or more of:

A. alkyl monoamines of the formula;




where R<sub>5</sub> is selected from H and C<sub>1</sub> to C<sub>5</sub> alkyl, and R<sub>6</sub> is selected from C<sub>1</sub> to C<sub>14</sub> alkyl and the group -(CH<sub>2</sub>)<sub>n</sub>-OR<sub>7</sub> where n = 1 to 10 and R<sub>7</sub> is C<sub>1</sub> to C<sub>20</sub> alkyl,

B. alkyl diamines of the formula;




where R<sub>5</sub> is selected from H and C<sub>1</sub> to C<sub>5</sub> alkyl, R<sub>8</sub> is C<sub>1</sub> to C<sub>6</sub> alkylene and A and B are independently selected from H, C<sub>1</sub> to C<sub>5</sub> alkyl, mono-hydroxysubstituted C<sub>1</sub> to C<sub>5</sub> alkyl, and the group (CH<sub>2</sub>)<sub>n</sub>-OR<sub>7</sub> where n = 1 to 10 and R<sub>7</sub> is C<sub>1</sub> to C<sub>20</sub> alkyl,

C. ethylene polyamines of the formula;



where n = 2 to 10, and

D. cyclic amines of the formula;




where n and m are independently integers from 1 to 3, X is selected from CH<sub>2</sub>, O, S and NR<sub>9</sub> where R<sub>9</sub> is H, C<sub>1</sub> to C<sub>10</sub> alkyl, or the group (CH<sub>2</sub>)<sub>n</sub>-NH<sub>2</sub> where n is 1 to 10. The alkyl groups can have a branched chain.

Specific examples of such amines include 1,3-diaminopropane, 1,2 diaminopropane, dimethylamine, die-




thylamine, dipropylamine, dibutylamine, N,N-dimethyl-1,3-diaminopropane, 1,1-dimethyldodecylamine, mixed C<sub>12</sub>-C<sub>14</sub> t-alkyl amines, 2-methyl-1,5-pentadiamine, ethylenediamine; cyclic amines such as piperazine, aminoethylpiperazine, morpholine and thiomorpholine; and ethylene polyamines such as diethylene triamine and triethylene tetraamine.


5 The Mannich Base can be formed by reacting from 1 to 5 moles of aldehyde, from about 1 to 2 moles of amine and from 1 to 4 moles of phenol at a temperature of from 0°C to 150°C for 0.5 to 10 hours. An inert solvent such as isopropanol can be used which is distilled from the product along with water formed in the reaction.

The Mannich Base product is usually a mixture of materials which may contain unreacted ingredients, especially the phenol. The Mannich Bases can be isolated from the product mixture but the product mixture itself  
10 can conveniently be used in forming the compositions of the invention. Examples of Mannich reactions and products are illustrated below:



20 or





where R<sub>1</sub>, R<sub>2</sub>, R<sub>4</sub>, R<sub>5</sub> and R<sub>6</sub> are as defined above.

15 The additive mixtures of the invention are usually prepared and marketed in the form of concentrates for addition to the fuel by the customer although the individual components could be added directly into the fuel. Suitable proportions of additives in the concentrates of the invention, based on the total weight of concentrate, include from 25 to 95 wt% N,N-dimethylcyclohexylamine, from 0 to 25 wt% N,N'-di(ortho-hydroxyarylidene)-1,2-alkylenediamine and, from 5 to 75 wt% Mannich Base.

20 The concentrates are added to the fuel in effective amounts to provide improved stability. Suitable amounts of additive concentrate in the fuel are from 1 to 500 pounds per thousand barrels (Ptbs) (3 to 1500 mg/l, preferred 2.5 to 100 Ptbs or 8 to 300 mg/l). This will provide a stabilized fuel containing from 1 to 1400 mg/liter (preferred 2 to 250 mg/l) N,N-dimethylcyclohexylamine, from 0 to 400 mg/l (preferred 0 to 100 mg/l) N,N'-di(ortho-hydroxyarylidene)-1,2-alkylenediamine metal deactivator and from 1 to 1100 mg/l (preferred 1 to 250 mg/l) of Mannich Base. When used, the metal deactivator is present in amounts of 1.0% to 25 wt% of concentrate or .3 to 400 mg/l of fuel. The concentrates can also contain an inert diluent or solvent which can be, for example, an aliphatic hydrocarbon such as kerosene or an aromatic hydrocarbon such as xylene.

25 The middle distillate fuels whose stability is improved by the invention typically include those boiling within a temperature range of 150°-400°C which may commonly be labeled as kerosene, fuel oil, diesel oil, No. 1-D, or No. 2-D.

30 The compositions of the invention are further illustrated by, but are not intended to be limited to, the following examples wherein parts are parts by weight unless otherwise indicated.

### Example 1

35 A Mannich Base reaction product of formaldehyde, 1,3-diaminopropane and 2,6-di-t-butylphenol is prepared by the following process.

40 Dissolve 103 grams (0.5 mole) of 2,6-di-t-butylphenol in 100 grams of isopropyl alcohol (IPA) in a 500 ml round bottom flask. Add 18.5 grams (0.25 mole) of 1,3-diaminopropane dropwise over 15 minutes while the contents of the flask are stirred. There is an exotherm observed as the amine is added. Cool the contents of the flask to below 30°C and add a 10% excess, (44.6 grams 0.55 mole) of 37% aqueous formaldehyde solution dropwise over 30 minutes while maintaining the temperature below 30°C. Heat the contents of the flask to reflux and continue to reflux for one hour. Switch from reflux to distillation and distill off IPA/water mixture to 105°C. Apply 28 in. Hg vacuum to remove residual materials. The total product yield is 122.2 or 96% of theory which contains compounds of the Structure III and IV.

45 Additive blends of the reaction product were prepared and tested in different fuels using both the D 4625 43°C (110°F) Storage Stability Test, in which the color change (using ASTM D1500) and the total insolubles in the fuel (reported in mg/100 ml) are determined on 400 ml samples stored for 13 weeks in the dark and the F-21-61 149°C (300°F) Accelerated Stability Test in which the color change and insoluble gums are determined on 50 ml samples heated to 149°C for a selected time, which was 90 minutes, allowed to cool in the dark, tested for color (ASTM D1500), and then filtered (using a 4.25 cm Whatman #1 filter paper) and the filtrate discarded. 50 The filter is washed clean of fuel with isooctane and measured for deposits by comparison with a set of reference papers. The blend compositions and test results in comparison to untreated fuel and blends without the Mannich Base product are reported in Table I below.

TABLE I  
Composition in Pounds Per Thousand Barrels

Components	Fuel #1					Fuel #2			Fuel #3			Fuel #4				
	1	2	3	4	5	1	2	3	1	2	3	1	2	3	4	5
DMCA <sup>1</sup>	0.0	5.0	4.0	4.75	3.80	0.0	4.75	3.8	0.0	9.5	7.6	0.0	2.38	1.90	4.75	3.80
MDA <sup>2</sup>	0.0	0.0	0.0	0.25	0.25	0.0	0.25	0.24	0.0	0.5	0.5	0.0	0.12	0.12	0.25	0.24
Mannich Base	0.0	0.0	1.0	0.00	0.95	0.0	0.0	0.96	0.0	0.0	1.9	0.0	0.00	0.48	0.00	0.96
Total																
Additives	0.0	5.0	5.0	5.0	5.0	0.0	5.0	5.0	0.0	10.0	10.0	0.0	2.5	2.5	5.0	5.0

Test Results  
149°C (F-21-61)

Components	Fuel #1					Fuel #2			Fuel #3			Fuel #4				
	1	2	3	4	5	1	2	3	1	2	3	1	2	3	4	5
Color	L7	L3.5	2.5	L3	2	8	3	3	8	L2.5	2	L2.5	2	2	2.5	L2.5
Deposit	13	5	4	4	2	17	10	5	17	5	3	6	4	4	4	4

43°C (D 4625)

Components	Fuel #1					Fuel #2			Fuel #3			Fuel #4				
	1	2	3	4	5	1	2	3	1	2	3	1	2	3	4	5
Color	L5	4	4	4	L4	L3.5	L3.5	L3.5	L3.5	L3.5	L3.5	2.5	L2.5	L2.5	L2.5	L2.5
Deposit	6.8	3.2	2.4	2.5	1.8	7.9	4	4.7	6.7	3	2.1	2.2	2	1.6	1.3	1

L = less than

<sup>1</sup>N,N-dimethylcyclohexylamine

<sup>2</sup>N,N'-disalicylidene-1,2-propylenediamine

Fuel #1 is Midwest Refinery

Fuel #2 is Mid-Continent #2 Diesel (Corning Crude)

Fuel #3 is Mid-Continent #2 Diesel (Ill. Basin Crude)

Fuel #4 is Midwest #2 Diesel (KS/TX Crude)

A significant difference in stability at 149°C is indicated by a color difference of about 1/2 number and/or a deposit difference of 2 numbers and a significant difference in stability at 43°C is indicated by a color difference of about 1/2 number and a deposit difference of 20%. The results in Table I show that the blends of the invention which contain Mannich Base in addition to DMCA or DMCA and MDA gave significantly better overall stability when compared to comparable blends which did not contain the Mannich Base, for example, blend 3 vs blend 2 and blend 5 vs blend 4 of Fuel #1.

#### Example 2

A Mannich Base reaction product of formaldehyde, dimethylamine, and 2,6-di-t-butylphenol is prepared by the following process.

Dissolve 103 grams (0.5 mole) of 2,6-di-t-butylphenol in 100 grams of IPA in a 500 ml round bottom flask and add 72 grams (0.64 mole) of a 40% aqueous dimethylamine solution. Cool the mixture to about 30°C and

add dropwise with stirring 44.6 grams (0.55 mole) of 37% formaldehyde while keeping the mixture at a temperature below 40°C. Heat the mixture to reflux and reflux for 4 hours. Remove IPA/water by distillation and apply vacuum to remove residual materials. The product yield is 113 grams or 86% of theory which contains N,N-dimethyl-3,5-di-t-butyl-4-hydroxybenzylamine.

Additive blends of the above reaction product were prepared and tested in different fuels using the test procedures described in Example 1. The blend compositions and test results in comparison to untreated fuel and blends which did not contain the Mannich Base product are reported in Table II below.

TABLE II  
Composition in Pounds Per Thousand Barrels

Components	Fuel #1					Fuel #2			Fuel #3			Fuel #4				
	1	2	3	4	5	1	2	3	1	2	3	1	2	3	4	5
DMCA	0.0	5.0	4.0	4.75	3.80	0.0	9.5	7.6	0.0	28.5	22.8	0.0	2.38	1.90	4.75	3.80
MDA	0.0	0.0	0.0	0.25	0.25	0.0	0.5	0.5	0.0	1.5	1.5	0.0	0.12	0.12	0.25	0.24
Mannich Base	0.0	0.0	1.0	0.00	0.95	0.0	0.0	1.9	0.0	0.0	5.7	0.0	0.00	0.48	0.00	0.96
Total																
Additives	0.0	5.0	5.0	5.0	5.0	0.0	10.0	10.0	0.0	30.0	30.0	0.0	2.5	2.5	5.0	5.0

Test Results  
149°C (F-21-61)

Components	Fuel #1					Fuel #2			Fuel #3			Fuel #4				
	1	2	3	4	5	1	2	3	1	2	3	1	2	3	4	5
Color	L7	L3.5	3	L3	L3	8	3	3	4.5	L4	3.5	L2.5	2	L2.5	2.5	L2.5
Deposit	13	5	6	4	4	17	10	6	5	3	3	6	4	5	4	3

43°C (D 4625)

Components	Fuel #1					Fuel #2			Fuel #3			Fuel #4				
	1	2	3	4	5	1	2	3	1	2	3	1	2	3	4	5
Color	L5	4	4	4	4	L3.5	L3.5	L3.5	--	--	--	2.5	L2.5	L3	L2.5	L2.5
Deposit	6.8	3.2	2.8	2.5	2	6.7	4	3.9	--	--	--	2.2	2	1.9	1.3	0.9

The fuels were the same as in Example 1 except that Fuel #3 is a fuel containing unhydro-treated residual cracked stock.

The results indicated that the blends containing Mannich Base gave fuels having significantly improved stability except in the case of Fuel #4 where the results were mixed.

### Example 3

A Mannich Base reaction product of formaldehyde, C<sub>12</sub>-C<sub>14</sub> t-alkyl amine mixture (Primene 81R) and 2,6-

di-t-butyl phenol is prepared by the process described in Example 2 using 95.5 grams (0.5 mole) of Primene 81R in place of the dimethylamine. The product yield is 200 grams or 82% of theory which contains N-[3,5-di-t-butyl-4-hydroxybenzyl]-mixed C<sub>12</sub>-C<sub>14</sub> t-alkyl amines.

Additive blends of the above reaction product were prepared and tested in #2 diesel fuel using the test procedures described in Example 1. The blend compositions and test results are reported in Table III below.

TABLE III  
Composition Pounds Per Thousand Barrels

<u>Components</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>6</u>	<u>7</u>
DMCA	0.0	9.5	0.0	4.75	19.0	0.0	9.5
MDA	0.0	0.5	0.5	0.50	1.0	1.0	1.0
Mannich Base	0.0	0.0	9.5	4.75	0.0	19.0	9.5
Total Additives	0.0	10.0	10.0	10.0	20.0	20.0	20.0

Test Results  
149°C (F-21-61)

<u>Components</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>6</u>	<u>7</u>
Color	15.5	14.5	15	14.5	14.5	15	14.5
Deposit	8	6	4	3	4	3	2

43°C (D 4625)

<u>Components</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>6</u>	<u>7</u>
Color	16.5	16	6	16	5.5	16	15.5
Deposit	8.3	3.2	5	2.8	3.3	5	3

The results indicate that blends 4 and 7 according to the invention which contain the Mannich Base in addition to DMCA and MDA have better stability at the same total additive levels compared to blends 2 and 5 containing only DMCA and MDA.

#### Example 4

A Mannich Base reaction product of formaldehyde, 1,2-diaminopropane, and 2,6-di-t-butyl phenol is prepared by the following process.

Dissolve 103 gm (0.5 mole) of 2,6-di-t-butyl phenol in 100 grams of IPA in a 500 ml roundbottom flask and add 18.5 grams (0.25) moles of 1,2-diaminopropane. Cool this mixture to about 30°C and add dropwise with stirring 44.6 grams (0.55 mole) of 37% formaldehyde while keeping the temperature of the mixture below 40°C. Heat the mixture to reflux and reflux for 1 hour. Remove IPA/water by distillation and apply vacuum to remove residual materials.

Additive blends of the above reaction product were prepared and tested in Fuel #1 fuel using the test procedures described in Example 1. The blend compositions and results are reported in Table IV below.



TABLE IV  
Composition Pounds Per Thousand Barrels

<u>Components</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>
DMCA	0.0	5.0	4.75	4.0	3.8
MDA	0.0	0.0	0.25	0.0	0.25
Mannich Base	0.0	0.0	0.00	1.0	0.95
Total Additives	0.0	5.0	5.0	5.0	5.0

Test Results  
149°C (F-21-61)

<u>Components</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>
Color	L7	L3.5	L3	L3	L2.5
Deposit	13	5	4	3	3

43°C (D 4625)

<u>Components</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>
Color	L5	4	4	4	L4
Deposit	6.8	3.2	2.5	2.6	1.7

The results indicate that blend 4 of the invention containing the Mannich Base has improved stability compared to blend 2 which containing DMCA alone. Blend 5 containing the Mannich Base has improved stability over blend 3 which contained DMCA and MDA alone.

#### Example 5

A Mannich Base reaction product of formaldehyde, N,N-dimethyl-1,3-diaminopropane, and p-dodecylphenol was prepared by the following procedure.

Combine 262.4 grams (1.0 mole) of the alkyl (C<sub>12</sub>) phenol with 102.2 grams (1.0 mole) of N,N-dimethyl-1,3-diamino-propane and add 89.2 grams (1.1 mole) of 37% formaldehyde with stirring while keeping the temperature below 40°C. Heat the mixture to 100°C for two hours and then remove water by distillation (100°C - 28 in vacuum). The product yield is 176 grams or 93% of theory which contains N,N-dimethyl-N'-[2 hydroxy-5-dodecylbenzyl]-1,3-diaminopropane.

Additive blends of the above reaction product were prepared and tested in midcontinent #2 diesel fuel using the test procedures described in Example 1. The blend compositions and results are reported on Table V below.

TABLE V  
Composition Pounds Per Thousand Barrels

<u>Components</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>
DMCA	0.0	4.75	2.05	9.5	4.5
MDA	0.0	0.25	0.25	0.5	0.5
Mannich Base	0.0	0.00	2.70	0.0	5.4
Total Additives	0.0	5.0	5.0	10.0	10.0

Test Results  
149°C (F-21-61)

<u>Components</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>
Color	8	3	13	12.5	12.5
Deposit	13	10	5	5	5


43°C (D 4625)

<u>Components</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>
Color	13.5	13.5	3	13.5	13.5
Deposit	6	4	3.5	3	2.4


The results indicate that blends 3 and 5 of the invention have over-all improved stability compared to blends at the same total additive level which did not include the Mannich Base.

### Claims

1. A fuel additive concentrate comprising a mixture of N,N-dimethylcyclohexylamine and a Mannich Base which is the reaction product of an aldehyde, a primary or secondary amine and an alkyl phenol selected from (a) hindered phenol having the formula:




where  $R_1$ ,  $R_2$ ,  $R_3$  are independently selected from hydrogen, t-butyl, t-amyl and isopropyl, provided that at least one of  $R_1$ ,  $R_2$  and  $R_3$  is hydrogen and at least one of  $R_1$  and  $R_2$  is t-butyl, t-amyl or isopropyl; and (b) p-alkyl phenol having the formula:



where  $R_4$  is  $C_9$  to  $C_{30}$  alkyl; said concentrate containing, based on the total weight of concentrate, from 25 to 95 wt% N,N-dimethylcyclohexylamine from 5 to 75 wt% Mannich Base.


2. The concentrate of claim 1 wherein the concentrate also contains from 1 to 25 wt% N,N'-di(ortho-hydroxyarylidene)-1,2-alkylenediamine metal deactivator.
3. The concentrate of claim 1 or 2 wherein the aldehyde is formaldehyde, and the amine is selected from one or more of:

A. alkyl monoamines of the formula;




where  $\text{R}_5$  is selected from H and  $\text{C}_1$  to  $\text{C}_5$  alkyl, and  $\text{R}_6$  is selected from  $\text{C}_1$  to  $\text{C}_{14}$  alkyl and the group  $-(\text{CH}_2)_n\text{-OR}_7$  where  $n = 1$  to  $10$  and  $\text{R}_7$  is  $\text{C}_1$  to  $\text{C}_{20}$  alkyl,

B. alkyl diamines of the formula;




where  $\text{R}_5$  is selected from H and  $\text{C}_1$  to  $\text{C}_5$  alkyl,  $\text{R}_8$  is  $\text{C}_1$  to  $\text{C}_6$  alkylene and A and B are independently selected from H,  $\text{C}_1$  to  $\text{C}_5$  alkyl, mono-hydroxysubstituted  $\text{C}_1$  to  $\text{C}_5$  alkyl, and the group  $(\text{CH}_2)_n\text{-OR}_7$  where  $n = 1$  to  $10$  and  $\text{R}_7$  is  $\text{C}_1$  to  $\text{C}_{20}$  alkyl,

C. ethylene polyamines of the formula;




where  $n = 2$  to  $10$ , and

D. cyclic amines of the formula;




where  $n$  and  $m$  are independently integers from  $1$  to  $3$ , X is selected from  $\text{CH}_2$ , O, S and  $\text{NR}_9$  where  $\text{R}_9$  is H,  $\text{C}_1$  to  $\text{C}_{10}$  alkyl, or the group  $(\text{CH}_2)_n\text{-NH}_2$  where  $n$  is  $1$  to  $10$ .

4. The concentrate of any of the preceding claims wherein the phenol is a hindered phenol, and the Mannich Base comprises N-[3,5-di-t-butyl-4-hydroxybenzyl]-mixed  $\text{C}_{12}$  to  $\text{C}_{14}$  t-alkyl amines, N,N-dimethyl-3,5-di-t-butyl-4-hydroxybenzylamine, a compound of the formula:



where  $\text{R}_1$  and  $\text{R}_2$  are independently selected from hydrogen, t-butyl, t-amyl and isopropyl provided that at least one of  $\text{R}_1$  and  $\text{R}_2$  is t-butyl, t-amyl, or isopropyl, or a compound of the formula:



where  $R_1$  and  $R_2$  are independently selected from hydrogen, t-butyl, t-amyl and isopropyl provided that at least one of  $R_1$  and  $R_2$  is t-butyl, t-amyl, or isopropyl.

- 15
5. The concentrate of any of claims 1-3 wherein the phenol is a p-alkylphenol, and the Mannich Base comprises N,N-dimethyl-N'-[2-hydroxy-5-dodecylbenzyl]-1,3-diaminopropane.
- 20
6. The concentrate of claim 2 wherein the metal deactivator is N,N'-disalicylidene-1,2-propylenediamine and the Mannich Base is the reaction product of formaldehyde, 2,6-di-t-butyl phenol and an amine selected from 1,3-diaminopropane, 1,2-diaminopropane, mixed  $C_{12}$ - $C_{14}$  t-alkyl amines, and dimethylamine.
- 25
7. The concentrate of claim 2 wherein the metal deactivator is N,N'-disalicylidene-1,2-propylenediamine and the Mannich Base is the reaction product of formaldehyde, p-dodecyl phenol, and N,N-dimethyl-1,3-diaminopropane.
- 30
8. A fuel composition comprising middle distillate fuel containing from 3 to 1500 mg/l of the concentrate of any of claims 1-7.
9. A fuel composition comprising a middle distillate fuel and from 1 to 1400 mg/l N,N-dimethylcyclohexylamine, from 0 to 400 mg/l N,N-di(ortho-hydroxyarylidene)-1,2-alkylenediamine and from 1 to 1100 mg/l of a Mannich Base which is the reaction product of an aldehyde, an amine and an alkyl phenol selected from (a) hindered phenol having the formula:



45

where  $R_1$ ,  $R_2$ ,  $R_3$  are independently selected from hydrogen, t-butyl, t-amyl and isopropyl, provided that at least one of  $R_1$ ,  $R_2$  and  $R_3$  is hydrogen and at least one of  $R_1$  and  $R_2$  is t-butyl, t-amyl or isopropyl; and (b) p-alkyl phenol having the formula:




where  $R_4$  is  $C_9$  to  $C_{30}$  alkyl.

10. The fuel composition of claim 9 wherein the aldehyde is formaldehyde and the amine is selected from one or more of:


A. alkyl monoamines of the formula;



10 where  $\text{R}_5$  is selected from H and  $\text{C}_1$  to  $\text{C}_5$  alkyl and  $\text{R}_6$  is selected from  $\text{C}_1$  to  $\text{C}_{14}$  alkyl and the group  $-(\text{CH}_2)_n\text{-OR}_7$  where  $n = 1$  to 10 and  $\text{R}_7$  is  $\text{C}_1$  to  $\text{C}_{20}$  alkyl,  
 B. alkyl diamines of the formula;



25 where  $\text{R}_5$  is selected from H and  $\text{C}_1$  to  $\text{C}_5$  alkyl  $\text{R}_8$  is  $\text{C}_1$  to  $\text{C}_6$  alkylene and A and B are independently selected from H,  $\text{C}_1$  to  $\text{C}_5$  alkyl, monohydroxysubstituted  $\text{C}_1$  to  $\text{C}_5$  alkyl, and the group  $(\text{CH}_2)_n\text{-OR}_7$  where  $n = 1$  to 10 and  $\text{R}_7$  is  $\text{C}_1$  to  $\text{C}_{20}$  alkyl,  
 C. ethylene polyamines of the formula;



40 where  $n$  and  $m$  are independently integers from 1 to 3, X is selected from  $\text{CH}_2$ , O, S and  $\text{NR}_9$  where  $\text{R}_9$  is H,  $\text{C}_1$  to  $\text{C}_{10}$  alkyl, or the group  $(\text{CH}_2)_n\text{-NH}_2$  where  $n$  is 1 to 10.


### Patentansprüche

- 45 1. Treibstoffzusatzkonzentrat umfassend eine Mischung aus N,N-Dimethylcyclohexylamin und einer Mannich-Base, die das Reaktionsprodukt eines Aldehyds, eines primären oder sekundären Amins und eines Alkylphenols ist, das ausgewählt ist aus (a) gehindertem Phenol mit der Formel



in der  $\text{R}_1$ ,  $\text{R}_2$  und  $\text{R}_3$  unabhängig voneinander aus Wasserstoff, t-Butyl, t-Amyl und Isopropyl ausgewählt

sind, mit der Maßgabe, daß mindestens einer der Reste  $R_1$ ,  $R_2$  und  $R_3$  Wasserstoff ist und mindestens einer der Reste  $R_1$  und  $R_2$  t-Butyl, t-Amyl oder Isopropyl ist, und  
(b) p-Alkylphenol mit der Formel




in der  $R_4$   $C_9$ - $C_{30}$ -Alkyl ist, wobei das Konzentrat auf der Basis seines Gesamtgewichts zwischen 25 und 95 Gew.-% N,N-Dimethylcyclohexylamin und 5 bis 75 Gew.-% Mannich-Base enthält.

2. Konzentrat nach Anspruch 1, welches außerdem zwischen 1 und 25 Gew.-% N,N'-Di(ortho-hydroxyary-  
liden)-1,2-alkylendiamin-Metaldeaktivator enthält.


3. Konzentrat nach Anspruch 1 oder 2, in dem das Aldehyd Formaldehyd ist und das Amin ausgewählt ist  
aus einer oder mehreren der folgenden Substanzen:

A. Alkylmonoamine der Formel



in der  $R_5$  ausgewählt ist aus H und  $C_1$ - bis  $C_5$ -Alkyl und  $R_6$  ausgewählt ist aus  $C_1$ - bis  $C_{14}$ -Alkyl und der  
Gruppe  $-(CH_2)_n-OR_7$ , in der  $n = 1$  bis 10 ist und  $R_7$   $C_1$ - $C_{20}$ -Alkyl ist,

B. Alkyldiamine der Formel




in der  $R_5$  ausgewählt ist aus H und  $C_1$ - bis  $C_5$ -Alkyl,  $R_8$   $C_1$ - bis  $C_6$ -Alkyl ist und A und B unabhängig  
voneinander aus H,  $C_1$ - bis  $C_5$ -Alkyl, mono-hydroxysubstituiertem  $C_1$ - bis  $C_5$ -Alkyl und der Gruppe  
 $(CH_2)_n-OR_7$ , in der  $n = 1$  bis 10 ist und  $R_7$   $C_1$ - bis  $C_{20}$ -Alkyl ist, ausgewählt sind.

C. Ethylenpolyamine der Formel


$H_2N\{(CH_2)_2NH\}_nH$ , in der  $n = 2$  bis 10 ist, und

D. cyclische Amine der Formel




in der  $n$  und  $m$  unabhängig ganze Zahlen von 1 bis 3 sind, X ausgewählt ist aus  $CH_2$ , O, S und  $NR_9$ ,  
worin  $R_9$  H,  $C_1$ - bis  $C_{10}$ -Alkyl oder die Gruppe  $(CH_2)_n-NH_2$  ist, in der  $n$  1 bis 10 beträgt.

4. Konzentrat nach einem der vorstehenden Ansprüche, in dem das Phenol ein gehindertes Phenol ist und die Mannich-Base N-[3,5-di-t-Butyl-4-hydroxybenzyl]-vermischte C<sub>12</sub>- bis C<sub>14</sub>-t-Alkylamine, N,N-Dimethyl-3,5-di-t-butyl-4-hydroxybenzylamin, eine Verbindung der Formel:




in der R<sub>1</sub> und R<sub>2</sub> unabhängig voneinander aus Wasserstoff, t-Butyl, t-Amyl und Isopropyl ausgewählt sind, mit der Maßgabe, daß mindestens einer der Reste R<sub>1</sub> und R<sub>2</sub> t-Butyl, t-Amyl oder Isopropyl ist, oder eine Verbindung der Formel



in der R<sub>1</sub> und R<sub>2</sub> unabhängig aus Wasserstoff, t-Butyl, t-Amyl und Isopropyl ausgewählt sind, mit der Maßgabe, daß mindestens einer der Reste R<sub>1</sub> und R<sub>2</sub> t-Butyl, t-Amyl oder Isopropyl ist, umfaßt.


5. Konzentrat nach einem der Ansprüche 1 bis 3, in dem das Phenol ein p-Alkylphenol ist und die Mannich-Base N,N-Dimethyl-N'-[2-hydroxy-5-dodecylbenzyl]-1,3-diaminopropan enthält.
6. Konzentrat nach Anspruch 2, in dem der Metalldeaktivator N,N'-Disalicyliden-1,2-propylendiamin und die Mannich-Base das Reaktionsprodukt aus Formaldehyd, 2,6-Di-t-butylphenol und einem aus der Gruppe 1,3-Diaminopropan, 1,2-Diaminopropan, gemischten C<sub>12</sub>-C<sub>14</sub>-t-Alkylaminen und Dimethylamin ausgewählten Amin ist.
7. Konzentrat nach Anspruch 2, in dem der Metalldeaktivator N,N'-Disalicyliden-1,2-propylendiamin und die Mannich-Base das Reaktionsprodukt aus Formaldehyd, p-Dodecylphenol und N,N-Dimethyl-1,3-diaminopropan ist.
8. Treibstoffzusammensetzung, die Mitteldestillattreibstoff mit 3 bis 1500 mg/l des Konzentrats nach einem der Ansprüche 1 bis 7 enthält.
9. Treibstoffzusammensetzung umfassend einen Mitteldestillattreibstoff und 1 bis 1400 mg/l N,N-Dimethylcyclohexylamin, 0 bis 400 mg/l N,N-Di(ortho-hydroxyaryliden)-1,2-alkylendiamin und 1 bis 1100 mg/l einer Mannich-Base, die das Reaktionsprodukt aus einem Aldehyd, einem Amin und einem Alkylphenol ist, das ausgewählt ist aus (a) einem gehinderten Phenol mit der Formel



5

10

in der  $R_1$ ,  $R_2$  und  $R_3$  unabhängig voneinander aus Wasserstoff, t-Butyl, t-Amyl und Isopropyl ausgewählt sind, mit der Maßgabe, daß mindestens einer der Reste  $R_1$ ,  $R_2$  und  $R_3$  Wasserstoff ist und mindestens einer der Reste  $R_1$  und  $R_2$  t-Butyl, t-Amyl oder Isopropyl ist, und (b) einem p-Alkylphenol mit der Formel



15

20


in der  $R_{\text{bis}}$   $C_{30}$ -Alkyl ist.

25

10. Treibstoffzusammensetzung nach Anspruch 9, in der das Aldehyd Formaldehyd ist und das Amin aus einer oder mehreren der folgenden Substanzen ausgewählt ist:

A. Alkylmonoamine der Formel

30




35

in der  $R_5$  aus H und  $C_1$ - bis  $C_5$ -Alkyl und  $R_6$  aus  $C_1$  bis  $C_{14}$ -Alkyl und der Gruppe  $-(CH_2)_n-OR_7$ , in der  $n$  1 - 10 und  $R_7$   $C_1$ - bis  $C_{20}$ -Alkyl ist, ausgewählt ist;

B. Alkyldiamine der Formel

40



45

in der  $R_5$  aus H und  $C_1$ - bis  $C_5$ -Alkyl ausgewählt ist,  $R_8$   $C_1$ - bis  $C_6$ -Alkyl ist, und A und B unabhängig voneinander aus H,  $C_1$ - bis  $C_5$ -Alkyl, monohydroxysubstituiertem  $C_1$ - bis  $C_5$ -Alkyl und der Gruppe  $(CH_2)_n-OR_7$ , in der  $n$  1 bis 10 ist und  $R_7$   $C_1$ - bis  $C_{20}$ -Alkyl ist, ausgewählt sind;

50


C. Ethylenpolyamine der Formel

$H_2((CH_2)_2NH)_nH$ , in der  $n$  2 bis 10 ist, und

D. Cyclische Amine der Formel

55







in der n und m unabhängig voneinander ganze Zahlen von 1 bis 3 sind und X ausgewählt ist aus der Gruppe CH<sub>2</sub>, O, S und NR<sub>9</sub>, in der R<sub>9</sub> H, C<sub>1</sub>- bis C<sub>10</sub>-Alkyl ist, oder der Gruppe (CH<sub>2</sub>)<sub>n</sub>-NH<sub>2</sub>, in der n 1 bis 10 ist.

### Revendications

1. Concentré d'additifs pour combustible, comprenant un mélange de N,N-diméthylcyclohexylamine et d'une base de Mannich qui est le produit de réaction d'un aldéhyde, d'une amine primaire ou secondaire et d'un alkylphénol choisi entre (a) un phénol à encombrement stérique répondant à la formule :




dans laquelle R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> sont choisis, indépendamment, entre l'hydrogène et les radicaux tertio-butyle, tertio-amyle et isopropyle, sous réserve que l'un au moins de R<sub>1</sub>, R<sub>2</sub> et R<sub>3</sub> soit l'hydrogène et que l'un au moins de R<sub>1</sub> et R<sub>2</sub> soit un radical tertio-butyle, tertio-amyle ou isopropyle ; et (b) un p-alkylphénol répondant à la formule :




dans laquelle R<sub>4</sub> est un groupe alkyle en C<sub>9</sub> à C<sub>30</sub> ; ledit concentré contenant, sur la base de son poids total, 25 à 95 % en poids de N,N-diméthylcyclohexylamine et 5 à 75 % en poids de base de Mannich.

2. Concentré suivant la revendication 1, qui contient aussi 1 à 25 % en poids de N,N'-di(ortho-hydroxy-arylidène)-1,2-alkylènediamine comme désactivateur des métaux.
3. Concentré suivant la revendication 1 ou 2, dans lequel l'aldéhyde est le formaldéhyde et l'amine consiste en une ou plusieurs des amines suivantes :
  - A. des alkylmono-amines de formule :




dans laquelle  $R_5$  est choisi entre H et des radicaux alkyle en  $C_1$  à  $C_5$ , et  $R_6$  est choisi entre des radicaux alkyle en  $C_1$  à  $C_{14}$  et le groupe  $-(CH_2)_n-OR_7$  dans lequel  $n$  a une valeur de 1 à 10 et  $R_7$  est un groupe alkyle en  $C_1$  à  $C_{20}$ ,

B. des alkyldiamines de formule :




dans laquelle  $R_5$  est choisi entre H et un groupe alkyle en  $C_1$  à  $C_5$ ,  $R_8$  est un groupe alkylène en  $C_1$  à  $C_6$  et A et B sont choisis, indépendamment, entre H, un groupe alkyle en  $C_1$  à  $C_5$ , alkyle en  $C_1$  à  $C_5$  portant un substituant hydroxy et le groupe  $(CH_2)_n-OR_7$  dans lequel  $n$  a une valeur de 1 à 10 et  $R_7$  est un groupe alkyle en  $C_1$  à  $C_{20}$ ,

C. des éthylènepolyamines de formule :




dans laquelle  $n$  a une valeur de 2 à 10, et

D. des amines cycliques de formule :




dans laquelle  $n$  et  $m$  sont, indépendamment, des nombres entiers de 1 à 3, X est choisi entre  $CH_2$ , O, S et  $NR_9$ , où  $R_9$  représente H, un radical alkyle en  $C_1$  à  $C_{10}$  ou le groupe  $(CH_2)_n-NH_2$  dans lequel  $n$  a une valeur de 1 à 10.

4. Concentré suivant l'une quelconque des revendications précédentes, dans lequel le phénol est un phénol à encombrement stérique et la base de Mannich comprend des N-[3,5-di-tertio-butyl-4-hydroxybenzyl]-tertio-alkylamines à groupes tertio-alkyle mixtes en  $C_{12}$  à  $C_{14}$ , la N,N-diméthyl-3,5-di-tertio-butyl-4-hydroxybenzylamine, un composé de formule :




dans laquelle  $R_1$  et  $R_2$  sont choisis, indépendamment, entre l'hydrogène, les radicaux tertio-butyle, tertio-amyle et isopropyle sous réserve que l'un au moins de  $R_1$  et  $R_2$  soit un radical tertio-butyle, tertio-amyle ou isopropyle, ou un composé de formule :



dans laquelle R<sub>1</sub> et R<sub>2</sub> sont choisis, indépendamment, entre l'hydrogène, les radicaux tertio-butyle, tertio-amyle et isopropyle, sous réserve que l'un au moins de R<sub>1</sub> et R<sub>2</sub> soit un radical tertio-butyle, tertio-amyle ou isopropyle.

- 15
5. Concentré suivant l'une quelconque des revendications 1 à 3, dans lequel le phénol est un p-alkylphénol et la base de Mannich comprend le N,N-diméthyl-N'-[2-hydroxy-5-dodécylbenzyl]-1,3-diaminopropane.
- 20
6. Concentré suivant la revendication 2, dans lequel le désactivateur de métaux est la N,N'-disalicylidène-1,2-propylènediamine et la base de Mannich est le produit de réaction du formaldéhyde, du 2,6-di-tertio-butylphénol et d'une amine choisie entre le 1,3-diaminopropane, le 1,2-diaminopropane, des tertio-alkylamines à groupes alkyle mixtes en C<sub>12</sub> à C<sub>14</sub> et la diméthylamine.
- 25
7. Concentré suivant la revendication 2, dans lequel le désactivateur de métaux est la N,N'-disalicylidène-1,2-propylènediamine et la base de Mannich est le produit de réaction du formaldéhyde, du p-dodécylphénol et du N,N-diméthyl-1,3-diaminopropane.
- 30
8. Composition de combustible, comprenant un combustible à base de distillat moyen contenant 3 à 1500 mg/l du concentré suivant l'une quelconque des revendications 1 à 7.
- 35
9. Composition de combustible, comprenant un combustible à base de distillat moyen et 1 à 1400 mg/l de N,N-diméthylcyclohexylamine, 0 à 400 mg/l de N,N-di(ortho-hydroxy-arylidène)-1,2-alkylènediamine et 1 à 1100 mg/l d'une base de Mannich qui est le produit de réaction d'un aldéhyde, d'une amine et d'un alkylphénol choisi entre (a) un phénol à encombrement stérique répondant à la formule :



45


dans laquelle R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> sont choisis, indépendamment, entre l'hydrogène, les radicaux tertio-butyle, tertio-amyle et isopropyle, sous réserve que l'un au moins de R<sub>1</sub>, R<sub>2</sub> et R<sub>3</sub> soit l'hydrogène et que l'un au moins de R<sub>1</sub> et R<sub>2</sub> soit un radical tertio-butyle, tertio-amyle ou isopropyle ; et (b) un p-alkylphénol répondant à la formule :



dans laquelle  $R_4$  est un radical alkyle en  $C_9$  à  $C_{30}$ .


10. Composition de combustible suivant la revendication 9, dans laquelle l'aldéhyde est le formaldéhyde et l'amine est une ou plusieurs des amines suivantes :

A. des alkylmono-amines de formule :




où  $R_5$  est choisi entre H et un radical alkyle en  $C_1$  à  $C_5$ , et  $R_6$  est choisi entre un radical alkyle en  $C_1$  à  $C_{14}$  et le groupe  $-(CH_2)_n-OR_7$  dans lequel  $n$  a une valeur de 1 à 10 et  $R_7$  est un radical alkyle en  $C_1$  à  $C_{20}$ ,

B. des alkyldiamines de formule :




dans laquelle  $R_6$  est choisi entre H et un radical alkyle en  $C_1$  à  $C_5$ ,  $R_8$  est un radical alkylène en  $C_1$  à  $C_6$  et A et B sont choisis, indépendamment, entre H, un radical alkyle en  $C_1$  à  $C_5$ , alkyle en  $C_1$  à  $C_5$  portant un substituant hydroxy et le groupe  $(CH_2)_n-OR_7$  dans lequel  $n$  a une valeur de 1 à 10 et  $R_7$  est un radical alkyle en  $C_1$  à  $C_{20}$ ,

C. des éthylènepolyamines de formule :



dans laquelle  $n$  a une valeur de 2 à 10, et

D. des amines cycliques de formule :



dans laquelle  $n$  et  $m$  représentent, indépendamment, des nombres entiers de 1 à 3, X est choisi entre  $CH_2$ , O, S et un groupe  $NR_9$  dans lequel  $R_9$  représente H, un radical alkyle en  $C_1$  à  $C_{10}$  ou le groupe  $(CH_2)_n-NH_2$  dans lequel  $n$  a une valeur de 1 à 10.