



12 **EUROPEAN PATENT APPLICATION**

21 Application number : **92610022.3**

51 Int. Cl.⁵ : **C12N 9/24, C12N 15/56,**
D21C 9/10, A23K 1/165,
A21D 2/00

22 Date of filing : **27.03.92**

30 Priority : **02.04.91 EP 91610027**

43 Date of publication of application :
07.10.92 Bulletin 92/41

84 Designated Contracting States :
PT

71 Applicant : **NOVO NORDISK A/S**
Novo Allé
DK-2880 Bagsvaerd (DK)

72 Inventor : **Schülein, Martin**
Wiedeweltsgade 51
DK-2100 Koebenhavn O (DK)
Inventor : **Halkier, Torben**
Vodroffsvej 6G
DK-1900 Frederiksberg C (DK)
Inventor : **Heldt-Hansen, Hans Peter**
Vangeledet 53
DK-2830 Virum (DK)
Inventor : **Pedersen, Lars Saaby**
Bagsvaerdvej 69B.14
DK-2800 Lyngby (DK)
Inventor : **Dalbøge, Henrik**
Ligustervaenget 63
DK-2830 Virum (DK)

54 **Xylanase, corresponding recombinant DNA sequence, xylanase containing agent, and use of the agent.**

57 The xylanase is characterized by several partial amino acid sequences and is immunoreactive with an antibody raised against a purified xylanase derived from *Humicola insolens*, DSM 1800. This xylanase preparation is practically free of cellulase and is well suited for treatment of paper pulp, as a baking agent and as an additive to fodder.

The invention comprises a xylanase, a corresponding recombinant DNA sequence, a vector, a transformed host, a method for production of the xylanase, an agent containing the xylanase, and a use of the agent.

Xylan, a major component of plant hemicellulose, is a polymer of D-xylose linked by β -1,4-xylosidic bonds. Xylan can be degraded to xylose and xylo-oligomers by acid or enzymatic hydrolysis. Enzymatic hydrolysis of xylan produces free sugars without the by-products formed with acid (e.g. furans).

There are currently five major applications for xylanases; 1) enzymatic breakdown of agricultural wastes for production of alcohol fuels; 2) enzymatic modification of animal feeds or feed components or addition to animal feeds for *in vivo* breakdown of the hemicellulose fraction; 3) use as a baking agent; 4) manufacturing of dissolving pulps yielding cellulose; and 5) bio-bleaching of wood pulp. [Detroy R.W. In: Organic Chemicals from Biomass, (CRC Press, Boca Raton, FL, 1981) 19-41.; Paice, M.G., and L. Jurasek. J. Wood Chem. Technol. 4: 187-198.; Pommier, J.C., J.L. Fuentes, G. Goma. Tappi Journal (1989): 187-191.; Senior, D.J., et al., Biotechnol. Letters 10 (1988):907-912.]

The pulp and paper industry is using xylanase compositions in the bleaching process to enhance the brightness of bleached pulps, to decrease the amount of chemicals used in the bleaching stages, and to increase the freeness of pulps in the recycled paper process [Eriksson, K. E. L., Wood Science and Technology 24 (1990); 79-101.; Paice, M. G., R. Bernier, and L. Jurasek, Biotechnol. and Bioeng. 32 (1988): 235-239.; Pommier, J. C., J. L. Fuentes, and G. Goma, Tappi Journal (1989): 187-191.]

Kraft pulping, a process widely used in the pulp and paper industry, involves the alkaline sulfate cooking of pulp to remove 90-98% of the lignin. The remaining 2-10% of lignin gives the pulp a dark brown color which has the tendency to darken in UV light or by oxidation. In order to obtain a white pulp for high quality paper, the crown color is removed by a multi-stage bleaching process using chemicals, e.g. chlorine, chlorine dioxide, ozone, oxygen or hydrogen peroxide.

Presently, there is much concern about the environmental impact of the chemicals generated from the bleaching process. Enzymes can aid in the removal of lignin from the pulp without any harmful side products. Reports show that lignin in wood is linked to xylan [Eriksson, O., et al., Wood Sci. Technol. 14 (1980); 267.; Takashi, N., and T. Koshijiima, Wood Sci. Technol. 22 (1988); 177-189]. By a limited hydrolysis of the xylan a greater release of lignin occurs during bleaching. Thus, by enzymatically treating the pulp prior to bleaching the amount of bleaching chemicals needed would in turn decrease. [Viikari, L., et al., Proceedings of the 3rd International Symposium on Biotechnology in the Pulp and Paper Industry (1986); 67.]

According to the technical literature, good results have been obtained by means of fungal preparations from *Trichoderma* [Paice, M. G., L. Jurasek, J. Wood Chem. Technol. 4 (1989): 187-198.; Senior, D.J., et al., Biotechnol. Letters. 10 (1988): 907-912], which require pH adjustment of the wood pulps below pH 6.0.

A *Trichoderma* xylanase preparation, Pulpzyme™ HA (commercially available from Novo Nordisk A/S) can be used for delignification of kraft pulps at pH 5-7. At 50°C and pH higher than 7, the enzyme shows only little effect.

Also a *Bacillus pumilus* xylanase preparation, Pulpzyme HB (commercially available from Novo Nordisk A/S) can be used for delignification of Kraft pulps at pH 5-7. At 50°C and pH higher than 7, the enzyme shows only little effect.

Humicola insolens xylanases have been described (Yoshioka, H et al., Agric. Biol. Chem. 43(3) (1981) 579-586). In the crude preparation they have a pH-optimum of 6.0 and a temperature optimum of 60°C. At pH 9 they show 25% of the activity at pH 6. According to the information in the article the enzymes were not purified and will therefore contain significant cellulase activity, as *H. insolens* is known as a good cellulase producer, reference being made to US 4,435,307. Further the enzymes are not characterized with respect to cellulases, M_w , pl or amino acid composition and has not been tested on kraft pulps, in baking or for animal feed.

The prior art xylanase preparation from *Humicola insolens* YH-8 described in Agric. Biol. Chem. and the other prior art xylanase preparations indicated above are not as well suited for use in delignification of kraft pulp, partly due to the relatively high cellulase content.

Thus, it is the purpose of the invention to provide a xylanase, which can be produced as a preparation with very small amounts of other enzyme activities, especially cellulase activities and other xylanase activities, and which is well suited for use in delignification of kraft pulp, as a baking agent and as an additive to animal fodder.

The xylanase according to the invention is characterized by the fact that it has the following partial amino acid sequences

1 Thr-Asn-Thr-Gly-Asn-Phe-Val-Gly-Gly-Lys-Trp-Asn-Pro-Gly-Thr-Gly-Arg-Thr-Lys-Asn-Tyr-,

2 Thr-Ala-Asn-Pro-Leu-Val-Glu-Tyr-Tyr-,

3 Ser-Trp-Trp-Ser-Asp-Gly-Gly-Gln-Val-Gln-Tyr-,

4 Val-Ser-Thr-Arg-Tyr-Asn-Gln-Pro-Ser-Ile-Asp-Gly-Thr-Arg-Thr-Phe-Gln-Gln-Tyr-Trp-Ser-Ile-Arg-Lys-,

5 Tyr-Val-Ile-Glu-Ser-Tyr-Gly-Thr-Tyr-Asn-Pro-Gly-Ser-Gln-Ala-Gln-Tyr-Lys-Gly-Thr-Phe-Tyr-Thr-Asp-Gly-Asp-Gln-Tyr-Asp-,

and

6 Gln-Val-Thr-Pro-Asn-Ala-Glu-Gly-Trp-His-Asn-Gly-Tyr-Phe-Tyr-,
or a partial amino acid sequence with a homology thereto of at least 80%, preferably at least 90%.

Surprisingly, it has been found that is possible to produce the xylanase according to the invention as part
5 of a xylanase preparation, which contains enzymatic activities, e.g. cellulases besides xylanase in very small concentrations. Especially, it has to be noted that the xylanase according to the invention is a special xylanase selected among the several xylanases produced inherently from *Humicola insolens*, DSM 1800, which is excellently suited both as an agent for addition to paper pulp, as a baking agent and as an additive to animal fodder. Also, it has been found that the xylanase according to the invention exhibits a specific activity which is larger
10 than the specific activity of any of the prior art xylanases. It is most surprising that the xylanase according to the invention exhibits superior properties in relation to all these technical areas, which are otherwise unrelated to each other.

The xylanase according to this invention exhibits a significant effect in removal of lignin from pulp at pH 8, as will be demonstrated later. The lignin removing effect is significantly higher than prior art (*B. pumilus*), as
15 will be shown later.

The high performance in regard to lignin removal is surprisingly obtained with xylanase according to the invention alone, even though it is one out of several xylanase components in *H. insolens*. The process for lignin removal is more economical as only one xylanase component is needed, a component which by genetic engineering can be produced in high yields.

Surprisingly it is possible to produce the xylanase according to the invention without significant cellulase activity, thereby avoiding yield loss caused by cellulolytic attack on the cellulose fibres in the pulp during the use of the xylanase for lignin removal.

As will be shown later the xylanase according to the invention surprisingly can remove more lignin from softwood than the *B. pumilus* xylanase even though the two xylanases have very similar pH profiles.

Xylanase (the designation pentosanase is commonly used in the baking industry) is used as a baking agent for wheat bread for several purposes:

- dough development
- improving dough elasticity and stability
- increasing bread volume
- improving crumb structure
- anti-staling

It is believed that some xylanases degrade tire pentosans (arabinoxylans) in such a way that the thereby modified pentosans improve dough elasticity, stability and development. Surprisingly the xylanase according to the invention can modify the pentosans in such a way. The xylanase according to the invention can be produced without other xylanases and xylan modifying enzymes and thereby makes it possible to obtain a controlled modification of the pentosans.

The pH in dough is 6.0 to 5.5 which makes this xylanase ideal for the use as a baking agent, as the pH optimum of the xylanase according to the invention is 5.5 to 7.5.

The xylanase according to the invention can also be used for modification of animal feeds or for addition to animal feeds for in vivo breakdown of the hemicellulose fraction. The xylanase according to the invention can be used as a preparation with practically no side activities, whereas the prior art xylanase, represented by an *H. insolens* xylanase product, would contain several side activities, including other xylanases. It was found that the xylanase according to the invention and the prior art xylanase gave rise to similar effects in regard to weight gain in a chicken feeding trial. This shows that the xylanase according to the invention is the active or one of the active *H. insolens* xylanases in relation to the feed additive application. Furthermore, the xylanase according to the invention exhibits the significant advantage that it contains a very pure xylanase, whereby a reproducible weight gain can be obtained, in contradistinction to the prior art xylanase, which contains many different side activities with varying proportions from batch to batch. Also, use of the xylanase according to the invention, which is a single component xylanase, opens up the possibility of a more economic feed additive

A preferred embodiment of the xylanase according to the invention is characterized by the fact that the xylanase is immunoreactive with an antibody raised against a purified xylanase derived from *Humicola insolens*, DSM 1800.

A preferred embodiment of the xylanase according to the invention is characterized by the fact that the xylanase has an isoelectric point of 7.5 - 9.5, preferably 8.0 - 8.5.

A preferred embodiment of the xylanase according to the invention is characterized by the fact that the xylanase exhibits a specific activity above 330 EXU/mg of protein, preferably above 400 EXU/mg of protein. The EXU xylanase activity unit is defined in AF 293.9/1. This AF publication and other AF publications indicated in this specification are available on request from Novo Nordisk A/S, Novo Allé, DK-2880 Bagsvaerd, Denmark.

EP 0 507 723 A1

Also, the invention comprises a recombinant DNA sequence, which is characterized by encoding for the xylanase according to the invention.

A preferred embodiment of the recombinant DNA sequence according to the invention is characterized by the fact that it comprises the following partial DNA sequence

5

```

1           5           10           15
ATG GTC TCG CTC AAG TCT GTC CTC GCG GCC GCC ACG GCT GTG AGC

10          20          25          30
TCT GCC ATT GCT GCC CCT TTT GAC TTC GTT CCT CGG GAC AAC TCG

          35          40          45
ACG GCC CTT CAG GCT CGC CAG GTG ACC CCC AAC GCC GAG GGC TGG

15          50          55          60
CAC AAC GGC TAC TTC TAC TCG TGG TGG TCC GAC GGC GGA GGC CAG

          65          70          75
GTT CAG TAC ACC AAC CTC GAG GGC AGC CGC TAC CAG GTC AGA TGG

20          80          85          90
NNN AAC ACC GGC AAC TTC GTC GGT GGT AAG GGT TGG AAC CCG GGA

          95          100          105
ACC GGC CCC ACG ATC AAC TAC GGC GGC TAC TTC AAC CCC CAG GGC

          110          115          120
AAC GGC TAC CTG GCC GTC TAC GGC TGG ACC NNN AAC CCG CTC GTC

30          125          130          135
GAG TAC TAT GTC ATC GAG TCG TAC GGC ACG TAC AAT CCC GGC AGC

          140          145          150
CAG GCT CAG TAC AAG GGC ACA TTC TAT ACC GAC GGC GAT CAG TAT

35          155          160          165
GAC ATC TTT GTG AGC ACC CGT NNN AAC CAG CCC AGC ATC ACG GCA

          170
CCC GGA CGT CCA GCT AGT ACT
40

```

A preferred embodiment of the recombinant DNA sequence according to the invention is characterized by the fact that it comprises a DNA sequence selected from

45

- a) the *Humicola insolens* xylanase DNA insert in pH 450
- b) a DNA sequence which hybridizes to the coding region for the mature xylanase DNA comprised by the DNA insert of a) and which comprises a structural gene for a polypeptide with xylanase activity, and optionally a promoter, a coding region for a signal or leader peptide and/or transcriptional terminator
- c) a DNA sequence with a homology sufficient to hybridize to the sequence indicated in Claim 6 under relative stringent conditions (1.0 x SSC, 0.1% SDS, 65°C), reference being made to T. Maniatis, A laboratory Manual (CSH)
- d) a derivative of a DNA sequence defined in a), b) or c), or
- e) a DNA sequence which codes for a mature xylanase or a signal peptide or a leader peptide thereof and which is degenerate within the meaning of the genetic code with respect to a DNA sequence of a), b) or c).

55

Also, the invention comprises a vector, which is characterized by the fact that it comprises the recombinant DNA sequence according to the invention.

A preferred embodiment of the vector according to the invention is characterized by the fact that the pro-

motor is the *Aspergillus oryzae* takaamylase promoter and/or the xylanase gene is isolated from EC 3-2, and/or the terminator is the *Aspergillus oryzae* AMG terminator, preferably pHD450.

Also, the invention comprises a transformed host, which is characterized by the fact that it contains the vector according to the invention.

5 A preferred embodiment of the transformed host according to the invention is characterized by the fact that it is an *Aspergillus* strain.

A preferred embodiment of the transformed host according to the invention is characterized by the fact that it is a strain belonging to the species *Aspergillus aculeatus*, *Aspergillus niger*, *Aspergillus oryzae* or *Aspergillus awamori*.

10 A preferred embodiment of the transformed host according to the invention is characterized by the fact that it is a microorganism, which in its non-transformed condition does not produce xylanase or only produces xylanase in insignificant amounts, preferably *Bacillus sp.*, *E. coli* or *S. cerevisiae*.

Also, the invention comprises a method for production of xylanase, which is characterized by the fact that the method utilizes a transformed host according to the invention.

15 Also, the invention comprises the xylanase, produced by means of the method according to the invention.

Also, the invention comprises an agent containing the xylanase according to the invention, preferably in the form of a non-dusting granulate, a stabilized liquid or a protected enzyme, whereby the xylanase comprises at least 10%, preferably at least 30% of the total enzyme protein.

20 A preferred embodiment of the agent according to the invention is characterized by the fact that the proportion between the xylanase activity and the cellulase activity, as expressed by the ratio between the xylanase activity in EXU/g and the cellulase activity in ECU/g has a value above 10, preferably above 30, most preferably above 100. The cellulase activity unit ECU is defined in AF 302-1.

A preferred embodiment of the agent according to the invention is characterized by the fact that it contains a xylanase activity of at least 10 EXU/mg of enzyme protein, preferably at least 100 EXU/mg of enzyme protein, more preferably at least 300 EXU/mg of enzyme protein.

25 Also, the invention comprises a use of the agent according to the invention, the use being characterized by the fact that the agent is used for xylan degradation.

A preferred embodiment of the use according to the invention is characterized by the fact that the use is related to chemical pulp or recycle paper pulp before or as part of bleaching, preferably at a pH value above 7.

30 A preferred embodiment of the use according to the invention is characterized by the fact that the use is related to production of bread, which contains wheat flour.

A preferred embodiment of the use according to the invention is characterized by the fact that the use is related to animal feed.

35 The xylanase according to the invention can be produced in the following manner.

Xylanase was produced by cultivating *Humicola insolens* DSM 1800, as described in US 4,435,307, Example 6 from the beginning thereof to column 11, line 29. The freeze dried powder was diluted with water to a dry matter content of 10%, and pH was lowered to 2.3 with 10% HCl. The mixture was left alone at 22°C for 50 minutes, and subsequently pH was increased to 8.0 with NaOH. Then a salt precipitation with 250 g of Na₂SO₄ per kg of liquid at pH 5.0 was carried out. The salt cake was redissolved and then concentrated and washed by ultrafiltration. Finally the preparation was frozen down.

In a 12% dry matter solution the xylanase preparation exhibited the following enzyme activities: 444 CSU/ml and 144 EXU/ml. CSU is the cellulase activity unit, vide AF 267.

45 This preparation was further purified by batch ion exchange with DEAE Sephadex A-50 (Pharmacia), concentration by AMICON ultrafiltration, gel filtration on Sephacryl S-200 (Pharmacia), and cation exchange with high load S-Sepharose (Pharmacia).

50 It was found that the xylanase did not exhibit any detectable cellulase side activity. Thus, the purified xylanase product showed a cellulase activity of less than 0.1 CSU/mg protein and a xylanase activity higher than 350 EXU/mg protein. The purified xylanase product shows only one SDS-PAGE band with molecular weight 22 kD. The pI was determined to 8.5 by isoelectric focusing.

The antibody reactive with this purified xylanase product was produced in the following manner. Antiserum against the purified xylanase was raised by immunizing rabbits according to the procedure described by N. Axelsen *et al.*: A Manual of Quantitative Immunoelectrophoresis, Blackwell Scientific Publications, 1973, Chapter 23. A purified immunoglobulin was obtained from the antiserum by salt precipitation ((NH₄)₂SO₄), followed by dialysis and ion exchange chromatography on DEAE-Sephadex. Immunochemical characterization of the purified xylanase was conducted by rocket immunoelectrophoresis (N. Axelsen *et al.*, Chapter 2,). With the antibody indicated above the purified xylanase showed a single arch moving against the cathode in rocket immunoelectrophoresis.

Also, after enzymatic degradation with chymotrypsin three peptides are sequenced, whereby one of these is determined as a side sequence (ss) during the sequencing of another peptide:

Chymo-22: Thr-Asn-Thr-Gly-Asn-Phe-Val-Gly-Gly-Lys-Trp-Asn-Pro-Gly-Thr-Gly-Arg-Thr-Lys-Asn-Tyr-
(SEQ ID NO. 1)

5 Chymo-22: Thr-Ala-Asn-Pro-Leu-Val-Glu-Tyr-Tyr- (ss) (SEQ ID NO. 2)

Chymo-24: Ser-Trp-Trp-Ser-Asp-Gly-Gly-Gly-Gln-Val-Gln-Tyr-(SEQ ID NO. 3)

After chemical degradation with CNBr followed by enzymatic degradation with pepsin two peptides are sequenced.

10 CNBr/Pep-16: Val-Ser-Thr-Arg-Tyr-Asn-Gln-Pro-Ser-Ile-Asp-Gly-Thr-Arg-Thr-Phe-Gln-Gln-Tyr-Trp-Ser-Ile-Arg-Lys- (SEQ ID NO. 4)

CNBr/Pep-23: Tyr-Val-Ile-Glu-Ser-Tyr-Gly-Thr-Tyr-Asn-Pro-Gly-Ser-Gln-Ala-Gln-Tyr-Lys-Gly-Thr-Phe-Tyr-
-Thr-Asp-Gly-Asp-Gln-Tyr-Asp- (SEQ ID NO. 5)

It has been found that the xylanase according to the invention exhibits a weak homology to prior art xylanases, e.g. xylanases producible from *Schizophyllum commune*, *Bacillus pumilus* and *Bacillus subtilis*.

15 The invention will be illustrated by the following examples.

Example 1 illustrates the selection of the xylanase producing gene and production of the xylanase by means of a genetic modified host organism. Example 2 illustrates the production in pilot plant of the xylanase and purification of the xylanase. Example 3 illustrates the use of the xylanase as a bleach booster during paper pulp production and Example 4 illustrates the use of the xylanase as a baking agent.

20

EXAMPLE 1

Media

25 YPD: 10 g yeast extract, 20 g peptone, H₂O to 810 ml. Autoclaved, 90 ml 20% glucose (sterile filtered) added.
10 × Basal salt: 66.8 g yeast nitrogen base, 100 g succinic acid, 60 g NaOH, H₂O ad 1000 ml, sterile filtered.
SC-URA: 90 ml 10 × Basal salt, 22.5 ml 20% casamino acids, 9 ml 1% tryptophane, H₂O ad 806 ml, autoclaved,
3.6 ml 5% threonine and 90 ml 20% glucose added.

30 SC-H agar: 7.5 g/l yeast nitrogen base without amino acids, 11.3 g/l succinic acid, 6.8 g/l NaOH, 5.6 g/l casamino acids without vitamins, 0.1 g/l tryptophan and 20 g/l agar (Bacto). Autoclaved for 20 min. at 121°C. After autoclaving, 55 ml of a 22% galactose solution and 1.8 ml of a 5% threonine solution were added per 450 ml agar.

YNB-1 agar: 3.3 g/l KH₂P0₄, 16.7 g/l agar, pH adjusted to 7. Autoclaved for 20 min. at 121°C. After autoclaving, 25 ml of a 13.6% yeast nitrogen base without amino acids, 25 ml of a 40% glucose solution, 1.5 ml of
35 a 1 % L-leucine solution and 1.5 ml of a 1 % histidine solution were added per 450 ml agar.

YNB-1 broth: Composition as YNB-1 agar, but without the agar.

Oat spelt xylan overlayer gel: 1% agarose, 1% oat spelt xylan (Sigma Chemical Company) in Tris-maleic buffer, pH 7. The gel was boiled and then cooled to 55°C before the overlayer is poured onto agar plates.

40 FG-4-Agar: 35 g/l agar, 30 g/l Soy bean meal, 15 g/l maltodextrin (Glucidex 6), 5 g/l Bacto pepton, pH 7. Autoclaved 40 min at 121°C

FG-4 medium: 30 g/l Soy bean meal, 15 g/l maltodextrin (Glucidex 6), 5 g/l Bacto pepton. Autoclaved 40 min at 121°C.

Construction of a yeast expression plasmid

45

The commercially available plasmid pYES II (Invitrogen) was cut with SpeI, filled in with Klenow DNA polymerase + dNTP and cut with ClaI. The DNA was size fractionated on an agarose gel, and a fragment of about 2000 bp was purified by electroelution. The same plasmid was cut with ClaI/PvuII, and a fragment of about 3400 bp was purified by electroelution. The two fragments were ligated to a blunt-ended SphI/EcoRI fragment
50 containing the yeast TPI promoter. This fragment was isolated from a plasmid in which the TPI promoter from *S. cerevisiae* (cf. T. Albers and G. Kawasaki, *J. Mol. Appl. Genet.* 1, 1982, pp. 419-434) was slightly modified: an internal SphI site was removed by deleting the four bp constituting the core of this site. Furthermore, redundant sequences upstream of the promoter were removed by BalI exonuclease treatment followed by addition of a SphI linker. Finally, an EcoRI linker was added at position -10. After these modifications, the promoter is
55 included in a SphI-EcoRI fragment. Its efficiency compared to the original promoter appears to be unaffected by the modifications. The resulting plasmid pYHD17 is shown in Fig. 1.

Donor organism

H. insolens, DSM 1800, grown in a cellulose-rich fermentation medium with agitation to ensure sufficient aeration.

Isolation of mRNA

Total RNA was isolated from approximately 7 g of mycelium. The mycelium was frozen in liquid nitrogen and ground in a mortar with 1 g of quartz sand to a consistency of flour. The RNA was extracted with guanidinium thiocyanate and centrifuged through CsCl essentially as described in Sambrook et al., 1989, op. cit.. Poly A RNA was isolated from total RNA by chromatography on oligo dT cellulose.

cDNA synthesis

cDNA synthesis was carried out by means of a cDNA synthesis kit from Invitrogen according to the manufacturer's specifications. The DNA was adapted to the expression vectors by addition of a BstXI linker (Invitrogen) and size fractionated on an agarose gel. Only DNA larger than 5-600 bp was used in the library construction. The adapted cDNA was ligated into an appropriate yeast expression vector cut with BstXI. Following test ligations (in order to determine the size of the library) the library was plated onto 50 agar plates amounting to approximately 5000 transformants per plate. To each plate containing approximately 5000 individual clones was added 3 ml of medium. The bacteria were scraped off, 1 ml glycerol was added, and stored at -80°C as 50 pools. The remaining 2 ml were used for DNA isolation. If the amount of DNA was insufficient to give the required number of yeast transformants (see below), large scale DNA was prepared from 500 ml medium (TB) inoculated with 50 µl -80°C bacterial stock propagated over night. DNA was isolated from 20 individual clones from the library and subjected to analysis for cDNA insertion. The insertion frequency was found to be >90% and the average insert size was approximately 1400bp.

Transformation of yeast

The yeast strain used was yNG231. (MAT alpha, leu2, ura3-52, his4-539, pep4-delta 1, cir+). One colony was grown at 30°C overnight in 10 ml of YPD. 10, 30, and 60 µl of this culture were added to 3 shaker flasks containing 100 ml YPD, and incubated with shaking overnight at 30°C. The culture with an OD₆₀₀ closest to 0.3-0.4 was selected. The cells were harvested in 50 ml tubes in a Beckman centrifuge (speed 6, 10 minutes), the cells were resuspended in 2 x 5 ml H₂O, centrifuged as described above, resuspended in 5 ml buffer containing 0.1 M LiAc, 10 mM Tris-Cl, 1 mM EDTA, pH 7.5, and centrifuged again. The cells were resuspended in 500 µl of the above buffer and incubated for 60 minutes at 30°C. 250 µg carrier DNA (sterile salmon sperm DNA 10 mg/ml, see below) was added and aliquots of 100 µl were prepared. The DNA to be transformed (approx. 5 µg) was added to the 100 µl aliquot, mixed gently, and incubated for 30 minutes at 30°C. 700 µl 40% PEG 4000, 0.1 M LiAc, 10 mM Tris-Cl, 1 mM EDTA, pH 7.5 was added, and incubation was continued for 60 minutes at 30°C. The transformation mixture was subjected to heat shock for 5 minutes at 42°C, spun briefly in a micro centrifuge, resuspended in 100-200 µl H₂O, and plated on SC plates without uracil, followed by incubation for three days at 30°C.

Preparation of carrier DNA

100 mg salmon-sperm DNA was weighed out and dissolved overnight in 10 ml 10 mM Tris-Cl, 1 mM EDTA, pH 7.5 (TE). The solution was then sonicated in a plastic container in ice water until it was no longer viscous. The solution was then phenole extracted and EtOH precipitated, and the pellet was washed and resuspended in 5 ml TE. The suspension was EtOH precipitated, and the pellet was washed and resuspended in 5 ml TE. The OD₂₆₀ was measured, and the suspension was diluted with TE to 10 mg/ml.

Screening of yeast

As described above DNA from the *Humicola* library, pools 1-10, was transformed into yeast, and plates containing 20-25,000 colonies were obtained from each pool. The colonies were scraped off and stored in glycerol at -80°C.

Yeast cells from the library were spread onto YNB agar to a total of about 400,000 colonies. The number of colonies per plate varied from 50 to 500. After 4 or 5 days of growth, the agar plates were replica plated onto

a set of SC-H agar plates. These plates were then incubated for 2-4 days at 30°C before the agar plates were overlaid with a oat spelt xylan overlay gel for the detection of xylanase. After incubation overnight at 40°C, enzyme reactions were visualised with Congo Red. 10-15 ml of a 0.1% solution of Congo Red was poured onto the overlay and removed after 10-20 min. The plates were then washed once or twice by pouring 10-15 ml of 2M NaCl onto the plates. The NaCl solution was removed after 15-25 min. Xylanase-positive colonies identified on the plates overlayers as colourless or pale red clearing zones on a red background.

Cells from enzyme-positive colonies were spread for single colony isolation on agar, and an enzyme-producing single colony was selected for each of the xylanase-producing colonies identified.

10 Characterization of xylanase positive clones

Each of the 147 of the xylanase-producing colonies were isolated. Some of these colonies were inoculated into 20 ml YNB-1 broth in a 50 ml glass test tube. The tube was shaken for 2 days at 30°C. The cells were harvested by centrifugation for 10 min. at 3000 rpm.

15 The cells were resuspended in 1 ml 0.9 M sorbitol, 0.1 M EDTA, pH 7.5. The pellet was transferred to an Eppendorf tube, and spun for 30 seconds at full speed. The cells were resuspended in 0.4 ml 0.9 M sorbitol, 0.1 M EDTA, 14 mM β -mercaptoethanol. 100 μ l 2 mg/ml Zymolase was added, and the suspension was incubated at 37°C for 30 minutes and spun for 30 seconds. The pellet (spheroplasts) was resuspended in 0.4 ml TE. 90 μ l of (1.5 ml 0.5 M EDTA pH 8.0, 0.6 ml 2 M Tris-Cl pH 8.0, 0.6 ml 10% SDS) was added, and the suspension was incubated at 65°C for 30 minutes. 80 μ l 5 M KOAc was added, and the suspension was incubated on ice for at least 60 minutes and spun for 15 minutes at full speed. The supernatant was transferred to a fresh tube which was filled with EtOH (room temperature) followed by thorough but gentle mixing and spinning for 30 seconds. The pellet was washed with cold 70% EtOH, spun for 30 seconds and dried at room temperature. The pellet was resuspended in 50 μ l TE and spun for 15 minutes. The supernatant was transferred to a fresh tube. 2.5 μ l 10 mg/ml RNase was added, followed by incubation at 37°C for 30 minutes and addition of 500 μ l isopropanol with gentle mixing. The mixture was spun for 30 seconds, and the supernatant was removed. The pellet was rinsed with cold 96% EtOH and dried at room temperature. The DNA was dissolved in 50 μ l water to a final concentration of approximately 100 μ g/ml.

30 The DNA was transformed into *E. coli* by standard procedures. Two *E. coli* colonies were isolated from each of the transformations and analysed with the restriction enzymes HindIII and XbaI which excised the DNA insert. DNA from one of these clones was retransformed into yeast and rescreened for enzyme activity.

The DNA sequences of several of the positive clones were partially determined. Based on the DNA sequence, 15 clones were classified as the same family, the sequence of this xylanase family showed full homology with the amino acid sequence of the purified xylanase according to the invention. A partial sequence appears in claim 5.

A strain of *E. coli* containing the xylanase HindIII/XbaI cDNA fragment in pYES2 was deposited in DSM on March 18, 1992, as DSM 6995. The xylanase cDNA fragment was isolated from one of the clones by cleavage with HindIII/XbaI. The HindIII/XbaI fragment was purified by agarose gel electrophoresis electroeluted and made ready for ligation reactions. The cDNA fragment is ligated to HindIII/XbaI digested pHD414 (see below) to generate pHD 450 in which the cDNA is under transcriptional control of the TAKA promoter from *Aspergillus oryzae* and the AMG terminator from *Aspergillus niger*.

After amplification of the DNA in *E. coli* the plasmid was transformed into *Aspergillus oryzae* as described as follows.

45 Construction of an *Aspergillus* expression vector

The vector pHD414 (Fig. 2) is a derivative of the plasmid p775 (described in EP 238 023). In contrast to this plasmid, pHD 414 has a string of unique restriction sites between the promoter and the terminator. The plasmid was constructed by removal of an approximately 200 bp long fragment (containing undesirable RE sites) at the 3' end of the terminator, and subsequent removal of an approximately 250 bp long fragment at the 5' end of the promoter, also containing undesirable sites. The 200 bp region was removed by cleavage with NarI (positioned in the pUC vector) and XbaI (just 3' to the terminator), subsequent filling in the generated ends with Klenow DNA polymerase +dNTP, purification of the vector fragment on gel and religation of the vector fragment. This plasmid was called pHD413. pHD413 was cut with StuI (positioned in the 5' end of the promoter) and PvuII (in the pUC vector), fractionated on gel and religated, resulting in pHD414. Fig. 11 is a map of plasmid pHD414, wherein "AMG Terminator" indicates the *A. niger* glucoamylase terminator, and "TAKA Promoter" indicates the *A. oryzae* TAKA amylase promoter.

Transformation of *Aspergillus oryzae*

100 ml of YPD (Sherman et al., Methods in Yeast Genetics, Cold Spring Harbor Laboratory, 1981) is inoculated with spores of *A. oryzae* or and incubated with shaking at 37°C for about 2 days. The mycelium is harvested by filtration through miracloth and washed with 200 ml of 0.6 M MgSO₄. The mycelium is suspended in 15 ml of 1.2 M MgSO₄, 10 mM NaH₂PO₄, pH = 5.8. The suspension is cooled on ice and 1 ml of buffer containing 120 mg of Novozym® 234 is added. After 5 minutes 1 ml of 12 mg/ml BSA (Sigma type H25) is added and incubation with gentle agitation continued for 1.5-2.5 hours at 37°C until a large number of protoplasts is visible in a sample inspected under the microscope.

The suspension is filtered through miracloth, the filtrate transferred to a sterile tube and overlaid with 5 ml of 0.6 M sorbitol, 100 mM Tris-HCl, pH = 7.0. Centrifugation is performed for 15 minutes at 100 g and the protoplasts are collected from the top of the MgSO₄ cushion. 2 volumes of STC (1.2 M sorbitol, 10 mM Tris-HCl, pH = 7.5, 10 mM CaCl₂) are added to the protoplast suspension and the mixture is centrifugated for 5 minutes at 1000 g. The protoplast pellet is resuspended in 3 ml of STC and repelleted. This is repeated. Finally the protoplasts are resuspended in 0.2-1 ml of STC.

100 µl of protoplast suspension is mixed with 5-25 µg of the appropriate DNA in 10 µl of STC. Protoplasts are mixed with p3SR2 (an *A. nidulans* amdS gene carrying plasmid). The mixture is left at room temperature for 25 minutes. 0.2 ml of 60% PEG 4000 (BDH 29576), 10 mM CaCl₂ and 10 mM Tris-HCl, pH = 7.5 is added and carefully mixed (twice) and finally 0.85 ml of the same solution is added and carefully mixed. The mixture is left at room temperature for 25 minutes, spun at 2500 g for 15 minutes and the pellet is resuspended in 2 ml of 1.2 M sorbitol. After one more sedimentation the protoplasts are spread on the appropriate plates. Protoplasts are spread on minimal plates (Cove, see Biochem.Biophys.Acta 113 (1966) 51-56) containing 1.0 M sucrose, pH = 7.0, 10 mM acetamide as nitrogen source and 20 mM CsCl to inhibit background growth. After incubation for 4-7 days at 37°C spores are picked and spread for single colonies. This procedure is repeated and spores of a single colony after the second reisolation is stored as a defined transformant.

Expression of the xylanase in *Aspergillus*

11 transformants were obtained and inoculated and maintained on YPG-agar. Each of the 8 selected transformants were inoculated from YPG-agar slants on 500 ml shake flask with 150 ml FG-4 media. After 4 days of fermentation with sufficient agitation to ensure good aeration, the culture broths were centrifuged for 10 minutes at 2000 g and the supernatants were analyzed. The best yielded 72 EXU/ml. There is no background from the untransformed host strain. This strain is designated Axy40/8.

EXAMPLE 2

The strain Axy40/8 was fermented in a pilot plant scale in the following way.

An agar substrate with the following composition was prepared in a Fernbach flask:

Sucrose	30 g/l
KH ₂ PO ₄	1 g/l
NaNO ₃	3 g/l
MgSO ₄ , 7H ₂ O	0,5 g/l
FeSO ₄ , 7H ₂ O	0,01 g/l
KCl	0,5 g/l
Agar	25 g/l

pH was adjusted to between 6.4 - 6.5, autoclaved for 20 minutes at 121 °C.

The Fernbach flask was inoculated with a spore suspension and cultivated 5 days at 30°C.

A substrate with the following composition was prepared in a 500 litre seed fermenter:

Yeast extract, 50%	6 kg
Potato starch	9 kg
CaCO ₃	0, 15 kg
Pluronic L61	150 ml
Termamyl® 60 (L)*	9 g

*) Termamyl is an alpha-amylase obtainable from Novo Nordisk A/S

Tap water was added to a total volume of around 225 litres. pH was adjusted to around 6.5 with H₃PO₄.

The temperature was raised from 60 to 90°C in 30 minutes, held for 30 minutes at 90°C, before the substrate was sterilized in the seed fermenter for 1.5 hour at 121°C. Final volume before inoculation was around 300 litres.

The Fernbach flask spore suspension was transferred to the seed fermenter. Seed fermentation conditions were:

Fermenter type: Conventional aerated and agitated fermenter with a height/diameter ratio of around 2.3.

Agitation: 250 rpm
 5 Aeration: 300 normal litre air per minute
 Temperature: 34°C
 Time: around 24 hours

Around 35 hours after inoculation 150 litres was transferred from the seed fermenter to the main fermenter.

A substrate with the following composition was prepared in a 2500 litre main fermenter:

10 MgSO₄, 7H₂O 2.6 kg
 KH₂PO₄ 2.6 kg
 K₂SO₄ 3.9 kg
 Trace metal solution 650 ml
 Potato starch 39.0 kg
 15 Yeast extract, 50% 28.6 kg
 Urea 5.2 kg
 Citric acid 1.053 kg
 Pluronic L61 650 ml
 Termamyl® 60 L 39 g

20 The trace metal solution has the following composition:

ZnSO₄, 7H₂O 14.3 g/l
 CuSO₄, 7H₂O 2.5 g/l
 NiCl₂, 6H₂O 0.5 g/l
 FeSO₄, 7H₂O 13.5 g/l
 25 MnSO₄, H₂O 8.5 g/l
 Citric acid, monohydrate 3.0 g/l

Tap water was added to a total volume of around 900 litres. pH was adjusted to 6.5 with H₃PO₄. The temperature was raised from 60 to 90°C in 30 minutes, held for 30 minutes at 90°C. Then pH was adjusted to 4.5 before the substrate was sterilized in the main fermenter for 1.5 hours at 123°C. Final volume before inoculation was around 1300 litres.

Then 150 litres of seed culture was added.

Fermentation conditions were:

Fermenter type: Conventional aerated and agitated fermenter with a height/diameter ratio of around 2.7.

Agitation: 250 rpm (two turbine impellers)
 35 Aeration: 1500 normal litre air per minute, rising to 1700 NI/min at 130 hours.
 Temperature: 34°C
 Time: around 130 hours

After 25 hours into the fermentation a maltodextrin solution was added aseptically to the main fermenter at a rate increasing from 1 l/hour to 4 l/hour in 12 hours. The dextrin solution was prepared with the following composition in a 550 litre feed tank:

40 Potato starch 184 kg
 Biotin 0.04 g
 Thiamin 0.4 g
 Citric acid 0.2 kg
 45 Pluronic L61 200 ml
 Termamyl® 60L 360 ml

Tap water was added to a total volume of around 300 litres. The temperature was held at 90°C for 60 minutes before the substrate was sterilized in the dosing tank for 1.5 hours at 123°C. Final volume before start of dosage was around 400 litres.

50 After 25 hours of fermentation pH was controlled at 7.3-7.4 by NH₃.

After around 130 fermentation hours the fermentation process was stopped. The around 1850 litres of culture broth were cooled to around 5°C and the enzymes were recovered according to the following method.

The culture broth was centrifuged at pH 7 and the centrifugate was filtered on a Seitz filter sheet (type Supra EKS Neu) using Hyflo Super-Cell diatomaceous earth as a filter aid. The pH in the filtrate was adjusted to 4.7 and the filtrate was concentrated to a dry matter content of 5% using ultrafiltration. Further concentration to a dry matter content of 30% was carried out by evaporation. The pH in the concentrate was adjusted to 6.5 and the concentrate was filtered on a frame filter using Hyflo Super-Cell as filter aid and germ filtered on a Seitz filter sheet (type Supra EKS Neu).

This concentrate is identified later in this specification as preparation 1. Preparation 1 exhibits a xylanase activity of 533 EXU/g.

Preparation 1 was purified as follows.

Totally 200,000 EXU was precipitated with 25% ammonium sulfate. The precipitate was solubilized in 500 ml of water and washed on an Amicon ultrafiltration cell with a membrane GR90PP from Dow Danmark A/S. The conductivity was reduced to 1.7 mS. The yield was 150,000 EXU. The pH was adjusted to 5.0 and the sample filtered through a 0.45 micron filter; yield 100,000 EXU. The sample was treated by cation exchange chromatography by means of 200 ml S-Sepharose fast flow column chromatography and a buffer containing 20 mM sodium acetate with pH 5.0. The xylanase bound to the column at this pH. The xylanase was eluted using a linear gradient containing 1 M sodium chloride in the same buffer. Totally 75,000 EXU was recovered in 160 ml with E₂₈₀ 4.1 which corresponds to 114 EXU per E₂₈₀.

For peptide mapping and measurement of the specific activity and characterization the xylanase is obtained in high purity by a final purification step. By means of size chromatography Superdex 75 a small portion of the xylanase has been highly purified. The purified xylanase has a single band in SDS-PAGE with a molecular weight of 22 kD, and a single band in isoelectric focusing by means of Pharmacia IEF gels with a pI of 8.2.

The native and the cloned xylanase according to the invention purified as described above, react equally with rabbit serum raised against purified native xylanase. By use of rocket immunoelectrophoresis both enzymes on an equal EXU basis give rise to the same immune precipitate.

By use of the EXU method the activity of the purified xylanase on colored xylan is 130 EXU per E₂₈₀ or 450 EXU per mg protein. The extinction coefficient is 3.5 E₂₈₀.

EXAMPLE 3

In this example the prior art *B. pumilus* xylanase has been compared with the xylanase according to the invention. Preparation 1 described in Example 2 was used for the comparison. On a softwood pulp, it was seen that the xylanase according to the invention gave a significantly better bleach boosting effect.

Xylanase according to the invention: *H. insolens* xylanase, liquid preparation, 533 EXU/g

Prior art xylanase: *B. pumilus* xylanase liquid preparation, 1070 EXU/g

Pulp

Unbleached softwood from a Swedish mill. The pulp has been washed, air dried, and stored at < 5°C in a cool room.

The pulp was soaked for > 12 hours in water whereafter it was pulped in a laboratory pulper at 1% DS, 10,000 reversions.

Table 1
Data for the pulp measured after storage

Pulp data measured	Softwood
Kappa No.	25.8
pH	7.35
Dry matter (%)	97.0
ISO brightness (%)	28.3

Two bleaching trials were performed, each consisting of the following two stages:

1) Enzyme treatment

2) (D50C50) E delignification

Laboratory bleaching conditions

Enzyme

5 50°C, pH 8.0 (Britton-Robinson buffer). Treatment time and dosage to be described later under the heading "Enzyme treatment".

(D50C50)

10 45 minutes, 40°C, 5% DS, final pH: 1.9 - 2.9.
aCL-multiples: 0. 15, 0. 19, 0.23, 0.27

E

15 60 minutes, 60°C, 12% DS, final pH: 10.6 - 12.2.

The NaOH dosage is calculated from the chlorine chemicals applied in the delignification stage as
[0.5 - (kg Cl₂/ton + ClO₂/ton) + 3] kg/ton

The bleaching stages were carried out in plastic bags heated in water baths. The bags were kneaded by hand at regular intervals.

20

Enzyme treatment

Two samples of pulp were treated in the same enzyme stage: one with the xylanase according to the invention, and the other with the prior art xylanase. The treatment time was 3 hours and the dosage was 1000 EXU/kg. Afterwards, the pulp was washed thoroughly with cold water. A third sample - the control - was given the same treatment, but without enzyme addition.

25

A significant difference between the two enzymes was seen in this experiment.

The Kappa numbers and absorbances measured after enzyme treatment are presented in Table 2.

30

Table 2

Data measured after the enzyme stage

35

	Control	Xylanase according to the invention	Prior art xylanase
Kappa no.	24.3	23.0	23.7
Abs. 280 nm	0.31	1.36	1.32

40

45

(D50C50) E delignification

From each of the three samples, four portions each of about 17 g of pulp were delignified with different dosages of active chlorine. Afterwards, the Kappa nos. and the ISO brightness values were measured. The results are shown in Table 3.

50

From Table 3 it appears that the *H. insolens* xylanase according to the invention provides a markedly lower Kappa no. and a better brightness than the prior art xylanase. Even if it appears from Table 3 that the Kappa no. in relation to the xylanase according to the invention is only slightly smaller than the Kappa no. in relation to the prior art xylanase, this slight difference corresponds to a much larger and significant saving of used active chlorine.

55

Table 3

Kappa nos. and brightness after enz(D50C50)E delignification of the pulp

aCl multiple*		0.15	0.19	0.23	0.27
kg of aCl/ton		36.5	46.2	56.0	65.7
Control	Kappa No.	8.48	6.33	4.55	3.89
	% ISO	36.1	40.8	45.0	47.5
Xylanase according to the invention	Kappa No.	7.22	4.92	3.88	3.03
	% ISO	39.1	44.0	47.9	51.5
Prior art xylanase	Kappa No.	7.54	5.66	4.07	3.28
	% ISO	38.7	42.0	46.5	51.1

* Multiple on Kappa no. of control

For the pulp used in this experiment it can be concluded that the xylanase according to the invention will generate a higher bleach boosting effect than the prior art xylanase.

EXAMPLE 4

This example illustrates the use of the xylanase according to the invention as a baking agent.

Xylanase (the designation pentosanase is commonly used in the baking industry) is used as a baking agent for wheat bread for several purposes:

- dough development
- improving dough elasticity and stability
- increasing bread volume
- improving crumb structure
- anti-staling

Preparation 1 has been tested in wheat bread and it has very good effect on bread quality. This enzyme has sufficient dough softening effect. Addition of 43150 FXU of preparation 1 increases the volume of rolls by 5-20% and crumb structure become more uniform and crumb is softer than bread without enzyme. It has a better effect in comparison to the commercial products. The xylanase activity unit FXU is defined in AF 293.6.

The prior art xylanase baking agents comprise several enzymatic activities, whereas the baking agent according to the invention easily can be produced with a very low content of enzymatic activities other than the xylanase activity. Thus, by use of the baking agent according to the invention bakery products with more constant characteristics from one baking operation to the next baking operation can be obtained.

Recipe and baking process

The basic recipe in this example is:

- wheat flour 1000 g
- salt 16 g
- sugar 16 g
- yeast 50 g
- water 590 g

- enzyme

The ingredients were mixed by a spiral dough-kneading machine. The dough was then kneaded for 2 minutes, at low speed and 5 minutes, at high speed. The dough temperature was approx. 26-28°C. After 10 minutes of resting, the dough was divided and formed to 30 rolls and 1 loaf. After a proofing time of 45 minutes for rolls and 40 minutes for loaf at 33°C, 80% RH, the rolls and the loaf were baked at 220°C for 15 minutes, and 30 minutes, respectively.

Results

	FXU/kg flour	Dough development	V-rolls index	V-loaf index	Crumb structure	Softness	
Reference	0	+	100	100	+	100	100
Prep. 1	43	++	106	105	++	87	66
	130	++	115	106	+	85	69
Prep. 1 without amylase activity	75	++	108	103	++	79	88
	151	++	116	104	+(+)	94	83
Pentopan	122	++	108	101	++	82	94
	427	++	111	105	+	96	82

In prep. 1 without amylase activity the amylase activity produced by the host organism is separated from the xylanase by means of conventional chromatographic techniques.

The roll and loaf volumes are measured using the traditional rape seeds method. The measured values are then recalculated to relative index.

The crumb softness is measured on a SMS-Texture Analyzer. A plunger with a diameter of 25 mm is pressed on the middle of a 11 mm thick slice of bread, the force needed for the plunger to depress the crumb 3 mm with a speed of 3.3 mm/s is recorded, and it is expressed as the crumb softness. The value is then recalculated to a relative index related to the reference sample which by definition has an index of 100. The lower the index for crumb softness, the softer is the crumb.

The characters for dough development and crumb structure are given according to visual evaluation: + means that the dough is normal, and that the crumb is also normal with coarse structure, and ++ means that the dough is softer than the reference and more elastic, and that the crumb structure is uniform and silky.

Pentopan™, a commercial preparation available from Novo Nordisk A/S, is a prior art baking agent which contains different xylanases from *H. insolens* and also other *H. insolens* enzyme activities. It appears from the above that the xylanase according to the invention has a better performance than the prior art baking agent as it provides larger volume and softer crumb. The most important baking advantage in relation to the xylanase according to the invention, in comparison to the prior art xylanase is the fact that the xylanase according to the invention can be used as a baking agent with practically no side activities, and thus is able to generate bread with very uniform properties from batch to batch.

SEQUENCE LISTING

5

(1) GENERAL INFORMATION:

(i) APPLICANT: NOVO NORDISK A/S

10

(ii) TITLE OF INVENTION: XYLANASE, CORRESPONDING RECOMBINANT DNA
SEQUENCE, XYLANASE CONTAINING AGENT, AND USE OF THE AGENT

(iii) NUMBER OF SEQUENCES: 7

(iv) CORRESPONDENCE ADDRESS:

15

(A) ADDRESSEE: NOVO NORDISK A/S

(B) STREET: Novo Allé

(C) CITY: DK-2880 Bagsvaerd

(E) COUNTRY: Denmark

20

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

25

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(viii) ATTORNEY/AGENT INFORMATION:

30

(A) NAME: BACH, Niels et al.

(B) REGISTRATION NUMBER: GA 24307

(C) REFERENCE/DOCKET NUMBER: 3588.204-WO

(ix) TELECOMMUNICATION INFORMATION:

35

(A) TELEPHONE: +45 4444 8888

(B) TELEFAX: +45 4449 3256

(C) TELEX: 37304

(2) INFORMATION FOR SEQ ID NO:1:

40

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21 amino acids

(B) TYPE: amino acid

45

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

50

(A) ORGANISM: Humicola insolens

(B) STRAIN: DSM 1800

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

5
Thr Asn Thr Gly Asn Phe Val Gly Gly Lys Trp Asn Pro Gly Thr Gly
1 5 10 15
Arg Thr Lys Asn Tyr
20

10 (2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

20 (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Humicola insolens
(B) STRAIN: DSM 1800

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

30 Thr Ala Asn Pro Leu Val Glu Tyr Tyr
1 5

(2) INFORMATION FOR SEQ ID NO:3:

35 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
40 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- 45 (A) ORGANISM: Humicola insolens
(B) STRAIN: DSM 1800

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Ser Trp Trp Ser Asp Gly Gly Gly Gln Val Gln Tyr
1 5 10

55

(2) INFORMATION FOR SEQ ID NO:4:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 amino acids
 (B) TYPE: amino acid
10 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: peptide
(vi) ORIGINAL SOURCE:
15 (A) ORGANISM: Humicola insolens
 (B) STRAIN: DSM 1800

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Val Ser Thr Arg Tyr Asn Gln Pro Ser Ile Asp Gly Thr Arg Thr Phe
1 5 10 15
25 Gln Gln Tyr Trp Ser Ile Arg Lys
 20

(2) INFORMATION FOR SEQ ID NO:5:

- 30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 29 amino acids
 (B) TYPE: amino acid
35 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: peptide
(vi) ORIGINAL SOURCE:
40 (A) ORGANISM: Humicola insolens
 (B) STRAIN: DSM 1800

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Tyr Val Ile Glu Ser Tyr Gly Thr Tyr Asn Pro Gly Ser Gln Ala Gln
1 5 10 15
50 Tyr Lys Gly Thr Phe Tyr Thr Asp Gly Asp Gln Tyr Asp
 20 25

55

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 15 amino acids
- (B) TYPE: amino acid, N-terminal
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- 15 (A) ORGANISM: Humicola insolens
- (B) STRAIN: DSM 1800

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

20 Gln Val Thr Pro Asn Ala Glu Gly Trp His Asn Gly Tyr Phe Tyr
1 5 10 15

25 (2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- 30 (A) LENGTH: 516 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: rDNA

(vi) ORIGINAL SOURCE:

- 40 (A) ORGANISM: Humicola insolens
- (B) STRAIN: DSM 1800

45

50

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

5 ATGGTCTCGC TCAAGTCTGT CCTCGCGGCC GCCACGGCTG TGAGCTCTGC CATTGCTGCC 60
CCTTTTGACT TCGTTCCTCG GGACAACTCG ACGGCCCTTC AGGCTCGCCA GGTGACCCCC 120
AACGCCGAGG GCTGGCACAA CGGCTACTTC TACTCGTGGT GGTCCGACGG CGGAGGCCAG 180
10 GTTCAGTACA CCAACCTCGA GGGCAGCCGC TACCAGGTCA GATGGNNNAA CACCGGCAAC 240
TTCGTCGGTG GTAAGGGTTG GAACCCGGGA ACCGGCCCCA CGATCAACTA CGGCGGCTAC 300
15 TTCAACCCCC AGGGCAACGG CTACCTGGCC GTCTACGGCT GGACCNNA A CCGCTCGTC 360
GAGTACTATG TCATCGAGTC GTACGGCAGG TACAATCCCG GCAGCCAGGC TCAGTACAAG 420
GGCACATTCT ATACCGACGG CGATCAGTAT GACATCTTTG TGAGCACCCG TNNNAACCAG 480
20 CCCAGCATCA CGGCACCCGG ACGTCCAGCT AGTACT 516

25

30

35

40

45

50

55

5
10
15
20
25
30
35
40
45
50
55

MICROORGANISMS	
Optional Sheet in connection with the microorganism referred to on page <u>18</u> , line <u>4</u> of the description ¹	
A. IDENTIFICATION OF DEPOSIT ¹	
Further deposits are identified on an additional sheet <input type="checkbox"/> ²	
Name of depositary Institution ⁴	
DEUTSCHE SAMMLUNG VON MIKROORGANISMEN UND ZELL-KULTUREN GmbH	
Address of depositary Institution (including postal code and country) ⁴	
Mascheroder Weg 1b, D-3300 Braunschweig, Federal Republic of Germany	
Date of deposit ⁵	Accession Number ⁶
18 March 1992	DSM 6995
B. ADDITIONAL INDICATIONS ⁷ (leave blank if not applicable). This information is continued on a separate attached sheet <input type="checkbox"/>	
In respect of those designations in which a European and/or Australian patent is sought, during the pendency of the patent application a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (Rule 28(4) EPC / Regulation 3.25 of Australia Statutory Rules 1991 No 71).	
C. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE ⁸ (if the indications are not for all designated States)	
D. SEPARATE FURNISHING OF INDICATIONS ⁹ (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later ⁹ (Specify the general nature of the indications e.g., "Accession Number of Deposit")	
E. <input type="checkbox"/> This sheet was received with the international application when filed (to be checked by the receiving Office)	
..... (Authorized Officer)	
<input type="checkbox"/> The date of receipt (from the applicant) by the International Bureau ¹⁰	
was	
..... (Authorized Officer)	

Claims

- 5 **1.** Xylanase wherein the xylanase has the following partial amino acid sequences
 1 Thr-Asn-Thr-Gly-Asn-Phe-Val-Gly-Gly-Lys-Trp-Asn-Pro-Gly-Thr-Gly-Arg-Thr-Lys-Asn-Tyr-, (SEQ ID
 NO. 1)
 2 Thr-Ala-Asn-Pro-Leu-Val-Glu-Tyr-Tyr-, (SEQ ID NO. 2)
 3 Ser-Trp-Trp-Ser-Asp-Gly-Gly-Gly-Gln-Val-Gln-Tyr-, (SEQ ID NO. 3)
10 4 Val-Ser-Thr-Arg-Tyr-Asn-Gln-Pro-Ser-Ile-Asp-Gly-Thr-Arg-Thr-Phe-Gln-Gln-Tyr-Trp-Ser-Ile-Arg-
 Lys-, (SEQ ID NO. 4)
 5 Tyr-Val-Ile-Glu-Ser-Tyr-Gly-Thr-Tyr-Asn-Pro-Gly-Ser-Gln-Ala-Gln-Tyr-Lys-Gly-Thr-Phe-Tyr-Thr-
 Asp-Gly-Asp-Gln-Tyr-Asp-, (SEQ ID NO. 5)
 and
15 6 Gln-Val-Thr-Pro-Asn-Ala-Glu-Gly-Trp-His-Asn-Gly-Tyr-Phe-Tyr-, (SEQ ID NO. 6)
 or a partial amino acid sequence with a homology thereto of at least 80%, preferably at least 90%.
- 2.** Xylanase according to Claim 1, wherein the xylanase is immunoreactive with an antibody raised against
 a purified xylanase derived from *Humicola insolens*, DSM 1800.
- 20 **3.** Xylanase according to any of Claims 1 - 2, wherein the xylanase has an isoelectric point of 7.5 - 9.5, pref-
 erably 8.0 - 8.5.
- 4.** Xylanase according to Claims 1 - 3, wherein the xylanase exhibits a specific activity above 330 EXU/mg
 of protein, preferably above 400 EXU/mg of protein.
- 25 **5.** Recombinant DNA sequence, characterized by encoding for the xylanase according to Claims 1 - 4.
- 6.** A recombinant DNA sequence according to Claim 5, comprising the following partial DNA sequence

30

35

40

45

50

55

11. Transformed host according to Claim 10, wherein the transformed host is an *Aspergillus* strain.
12. Transformed host according to Claim 10, wherein the transformed host is a strain belonging to the species *Aspergillus aculeatus*, *Aspergillus niger*, *Aspergillus oryzae* or *Aspergillus awamori*.
- 5 13. Transformed host according to Claim 10, wherein the transformed host is a microorganism, which in its non-transformed condition does not produce xylanase or only produces xylanase in insignificant amounts, preferably *Bacillus sp.*, *E. coli* or *S. cerevisiae*.
- 10 14. Method for production of a xylanase by use of a transformed host according to Claims 10 - 13.
- 15 15. Xylanase produced by the method according to Claim 14.
16. Agent containing the xylanase according to any of claims 1 - 4 or 15, preferably in the form of a non-dusting granulate, a stabilized liquid or a protected enzyme, whereby the xylanase comprises at least 10%, preferably at least 30% of the total enzyme protein.
17. Agent according to Claim 16, wherein the proportion between the xylanase activity and the cellulase activity, as expressed by the ratio between the xylanase activity in EXU/g and the cellulase activity in ECU/g has a value above 10, preferably above 30, most preferably above 100.
- 20 18. Agent according to Claims 16 - 17 which contains a xylanase activity of at least 10 EXU/mg of enzyme protein, preferably at least 100 EXU/mg of enzyme protein, more preferably at least 300 EXU/mg of enzyme protein.
- 25 19. Use of the agent according to Claims 16 - 18, characterized by the fact that the agent is used for xylan degradation.
20. Use according to Claim 19, characterized by the fact that the use is related to chemical pulp or recycle paper pulp before or as part of bleaching, preferably at a pH value above 7.
- 30 21. Use according to Claim 19, characterized by the fact that the use is related to production of bread, which contains wheat flour.
22. Use according to Claim 19, characterized by the fact that the use is related to animal feed.

35

40

45

50

55




FIG. 1




FIG. 2



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 92 61 0022
PAGE1

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
Y	WO-A-9 102 839 (NOVO NORDISK A/S)	1, 5, 8, 10, 14, 16, 19	C12N9/24 C12N15/56 D21C9/10 A23K1/165 A21D2/00
A	* the whole document * ---	20	
Y	EP-A-0 305 216 (NOVO INDUSTRI A/S)	1, 5, 8, 10, 14, 16, 19	
A	* abstract; claims 1-24 * ---	12	
A	AGRICULTURAL AND BIOLOGICAL CHEMISTRY vol. 51, no. 11, November 1987, TOKYO JP pages 3133 - 3135; T.HAMAMOTO ET AL.: 'Alkalophilic Bacillus xylanase A, a secretable protein through outer membrane of E.coli' * the whole document * ---	5, 8, 13	
A	JOURNAL OF ANTIBIOTICS vol. 49, no. 7, July 1986, TOKYO pages 985 - 993; A.IWASAKI ET AL.: 'Molecular cloning of a xylanase gene from Streptomyces sp. no. 36a and its expression in Streptomyces' * abstract * ---	5, 8, 13	TECHNICAL FIELDS SEARCHED (Int. Cl.5) C12N
A	CHEMICAL ABSTRACTS, vol. 101, no. 25, 17 December 1984, Columbus, Ohio, US; abstract no. 225520Y, V.KITPREECHAVANICH ET AL.: 'Purification and properties of endo-1,4-beta-xylanase from Humicola lanuginosa' page 332 ; & J.Ferment.Technol.1984, vol.62, no.5, pages 415-420 * abstract * ---	1	
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 13 MAY 1992	Examiner GURDJIAN D.
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			

EPO FORM 1503 01.92 (P0401)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 92 61 0022
PAGE2

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
A	WORLD PATENTS INDEX LATEST Week 8451, Derwent Publications Ltd., London, GB; AN 84-316055 & JP-A-59 198 978 (OKADA K) 10 November 1984 * abstract * ---	1, 5, 8, 10, 14, 16, 19, 21	
A	WORLD PATENTS INDEX LATEST Week 9045, Derwent Publications Ltd., London, GB; AN 90-336620 & EP-A-396 162 (UNILEVER NV) 7 November 1990 * abstract * -----	1, 5, 8, 10, 14, 16, 19, 22	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
Place of search BERLIN		Date of completion of the search 13 MAY 1992	Examiner GURDJIAN D.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03.82 (P0401)