



(11) **EP 0 529 031 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

Note: Bibliography reflects the latest situation

(15) Correction information:

Corrected version no 1 (W1 B1)
Corrections, see
Claims EN

(51) Int Cl.:

C12N 15/15 ^(1990.01) **C07K 5/00** ^(1985.01)
C07K 7/00 ^(1985.01) **A61K 38/55** ^(1995.01)
A61K 51/00 ^(1995.01) **A61L 33/00** ^(1985.01)

(48) Corrigendum issued on:

21.11.2007 Bulletin 2007/47

(86) International application number:

PCT/US1992/000836

(45) Date of publication and mention
of the grant of the patent:

24.05.2000 Bulletin 2000/21

(87) International publication number:

WO 1992/013952 (20.08.1992 Gazette 1992/22)

(21) Application number: **92905748.7**

(22) Date of filing: **03.02.1992**

(54) **IMPROVED INHIBITORS OF THROMBIN**

VERBESSESTE THROMBININHIBITOREN

INHIBITEURS AMELIORES DE THROMBINE

(84) Designated Contracting States:

AT BE CH DE DK ES FR GB GR IT LI LU MC NL SE

(74) Representative: **Vossius & Partner**

Postfach 86 07 67
81634 München (DE)

(30) Priority: **08.02.1991 US 652929**

(43) Date of publication of application:

03.03.1993 Bulletin 1993/09

(56) References cited:

EP-A- 0 019 589 **EP-A- 0 118 280**
WO-A-91/02750

(73) Proprietor: **BIOGEN, INC.**

Cambridge
Massachusetts 02142 (US)

- Biochemistry, vol. 29, 1990, (Easton, PA, US), J.M. MARAGANORE et al.: "Design and characterization of Hirulogs: A novel class of Bivalent peptide inhibitors of thrombin" pages 7095-7101, see abstract, page 7099, left-hand column, lines 21-23
- Scand. J. Haematol., vol. 31, 1983, (Copenhagen, DK), G.F. HANDELAND et al.: "Simplified assay for antithrombin III activity using chromogenic peptide substrate", pages 427-436, see abstract; page 435, left-hand column, paragraphs 2-3

(72) Inventors:


- MARAGANORE, John, M.
Concord, MA 01742 (US)
- JABLONSKI, Jo-Ann, M.
Middleborough, MA 02346 (US)
- BOURDON, Paul, R.
Sommerville, MA 02143 (US)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).


EP 0 529 031 B9

DescriptionTECHNICAL FIELD OF INVENTION

[0001] This invention relates to novel biologically active molecules which bind to and inhibit thrombin. These molecules comprise a catalytic site directed moiety (CSDM) of the formula:



wherein X is hydrogen or is characterized by a backbone chain consisting of from 1 to 100 atoms; R₁ is selected from the group consisting of unsubstituted, mono-substituted, di-substituted and tri-substituted saturated ring structures; R₂ is a bond or is characterized by a backbone chain consisting of from 1 to 5 atoms; R₃ is a bond or is characterized by a backbone chain consisting of from 1 to 3 atoms; R₄ is any amino acid; R₅ is any L-amino acid which comprises a guanidinium- or amino-containing side chain group; R₆ is a non-amide bond; and Y is a bond or is characterized by a backbone chain consisting of from 1 to 9 atoms; or the formula:



wherein **R'₁** is selected from the group consisting of unsubstituted, mono-substituted, di-substituted and tri-substituted ring structures; **R'₄** is any amino acid comprising a side chain group characterized by the capacity to accept a hydrogen bond at a pH of between about 5.5 and 9.5; and X, R₂, R₃, R₅, R₆ and Y are defined as above. Preferred thrombin inhibitors are further characterized by an anion binding exosite associating domain (ABEAM) and a linker portion of between 18Å and 42Å in length which connects Y to the ABEAM. This invention also relates to compositions, combinations and methods which employ these molecules for therapeutic, prophylactic and diagnostic purposes.

BACKGROUND ART

[0002] Acute vascular diseases, such as myocardial infarction, stroke, pulmonary embolism, deep vein thrombosis, peripheral arterial occlusion, and other blood system thromboses constitute major health risks. Such diseases are caused by either partial or total occlusion of a blood vessel by a blood clot, which contains fibrin and platelets.

[0003] Thrombin is the naturally occurring protein which catalyzes the conversion of fibrinogen to fibrin, the final step in blood clot formation. In addition to catalyzing the formation of a fibrin clot, thrombin also activates platelet aggregation and release reactions. This means that thrombin plays a central role in both acute platelet-dependent (arterial) thrombosis [S. R. Hanson and L. A. Harker, "Interruption of Acute Platelet-Dependent Thrombosis by the Synthetic Antithrombin D-Phenylalanyl-L-Prolyl-L-Arginyichloromethylketone", Proc. Natl. Acad. Sci. USA, 85, pp. 3184-88 (1988)] and fibrin-dependent (venous) thrombosis.

[0004] Thrombin has several other bioregulatory roles [J. W. Fenton, II, "Thrombin Bioregulatory Functions", Adv. Clin. Enzymol., 6, pp. 186-93 (1988)]. For example, thrombin can directly activate an inflammatory response by stimulating the synthesis of platelet activating factor (PAF) in endothelial cells [S. Prescott et al., "Human Endothelial Cells in Culture Produce Platelet-Activating Factor (1-alkyl-2-acetyl-sn-glycero-3-phosphocholine) When Stimulated With Thrombin", Proc. Natl. Acad. Sci. USA, 81, pp. 3534-38 (1984)]. PAF is exposed on the surface of endothelial cells and serves as a ligand for neutrophil adhesion and subsequent degranulation [G. M. Vercolletti et al., "Platelet-Activating Factor Primes Neutrophil Responses to Agonists: Role in Promoting Neutrophil-Mediated Endothelial Damage", Blood, 71, pp. 1100-07 (1988)]. Alternatively, thrombin may promote inflammation by increasing vascular permeability which can lead to edema [P. J. Del Vecchio et al., "Endothelial Monolayer Permeability To Macromolecules", Fed. Proc., 46, pp. 2511-15 (1987)]. Reagents which block the active site of thrombin, such as hirudin, interrupt the activation of platelets and endothelial cells [C. L. Knupp, "Effect of Thrombin Inhibitors on Thrombin-Induced Release and Aggregation", Thrombosis Res., 49, pp. 23-36 (1988)].

[0005] Thrombin has also been implicated in promoting cancer, based on the ability of its native digestion product, fibrin, to serve as a substrate for tumor growth [A. Falanga et al., "Isolation and Characterization of Cancer Procoagulant: A Cysteine Proteinase from Malignant Tissue", *Biochemistry*, 24, pp. 5558-67 (1985); S. G. Gordon et al., "Cysteine Proteinase Procoagulant From Amnion-Chorion", *Blood*, 66, pp. 1261-65 (1985); and A. Falanga et al., "A New Procoagulant In Acute Leukemia", *Blood*, 71, pp. 870-75 (1988)]. And thrombin has been implicated in neurodegenerative diseases based on its ability to cause neurite retraction [D. Gurwitz et al., "Thrombin Modulates and Reverses Neuroblastoma Neurite Outgrowth", *Proc. Natl. Acad. Sci. USA*, 85, pp. 3440-44 (1988)]. Therefore, the ability to regulate the *in vivo* activity of thrombin has many important clinical implications.

[0006] One route to the successful treatment or prevention of acute vascular disease is the inhibition of thrombin. Many types of thrombin inhibitors are already known in the art. Heparin, an indirect inhibitor of thrombin, is widely used to treat venous thrombosis. Although effective against fibrin-dependent clot formation, heparin has little efficacy in inhibiting thrombin-induced activation of platelets. Therefore, this drug is not utilized in the treatment of arterial thromboses. Moreover, heparin produces many undesirable side effects, including hemorrhaging and thrombocytopenia.

[0007] Hirudin is a naturally occurring polypeptide which is produced by the blood sucking leech Hirudo medicinalis. This compound, which is synthesized in the salivary gland of the leech, is the most potent natural inhibitor of coagulation known. Hirudin prevents blood from coagulating by binding tightly to thrombin ($K_d = 2 \times 10^{-11}M$) in a 1:1 stoichiometric complex [S. R. Stone and J. Hofsteenge, "Kinetics of the Inhibition of Thrombin by Hirudin", *Biochemistry*, 25, pp. 4622-28 (1986)]. This, in turn, inhibits thrombin from catalyzing the conversion of fibrinogen to fibrin (clot), as well as inhibiting all other thrombin-mediated processes [J. W. Fenton, II, "Regulation of Thrombin Generation and Functions", *Semin. Thromb. Hemost.*, 14, pp. 234-40 (1988)].

[0008] Hirudin inhibits thrombin by binding to the latter at two separate sites. Initially, the C-terminus of hirudin interacts with an "anion-binding exosite" (ABE) in thrombin [J. W. Fenton, II et al., "Thrombin Anion Binding Exosite Interactions with Heparin and Various Polyanions", *Ann. New York Acad. Sci.*, 556, Pp. 158-65 (1989)]. Following this low affinity binding, the hirudin-thrombin complex undergoes a conformational change and amino terminal portion of hirudin is able to bind to the catalytic site of thrombin [S. Kono et al., "Analysis of Secondary Structure of Hirudin and the Conformational Change Upon Interaction with Thrombin", *Arch. Biochem. Biophys.*, 267, pp. 158-66 (1988)].

[0009] The isolation, purification and amino acid sequence of hirudin are known in the art [P. Walsmann and F. Markwardt, "Biochemical and Pharmacological Aspects of the Thrombin Inhibitor Hirudin", *Pharmazie*, 36, pp. 653-60 (1981); J. Dodt et al., "The Complete Covalent Structure of Hirudin: Localization of the Disulfide Bonds", *Biol. Chem. Hoppe-Sevler*, 366, pp. 379-85 (1985); S. J. T. Mao et al., "Rapid Purification and Revised Amino Terminal Sequence of Hirudin: A Specific Thrombin Inhibitor of the Blood-Sucking Leech", *Anal. Biochem.*, 161, pp. 514-18 (1987); and R. P. Harvey et al., "Cloning and Expression of a cDNA Coding for the Anti-Coagulant Hirudin from the Bloodsucking Leech, *Hirudo medicinalis*", *Proc. Natl. Acad. Sci. USA*, 83, pp. 1084-88 (1986)].

[0010] In animal studies, hirudin, purified from leeches, has demonstrated efficacy in preventing venous thrombosis, vascular shunt occlusion and thrombin-induced disseminated intravascular coagulation. In addition, hirudin exhibits low toxicity and a very short clearance time from circulation [F. Markwardt et al., "Pharmacological Studies on the Antithrombotic Action of Hirudin in Experimental Animals", *Thromb. Haemost.*, 47, pp. 226-29 (1982)].

[0011] Hirudin has more recently been cloned and expressed in E. Coli [European patent applications 158,564, 168,342 and 171,024] and yeast [European patent application 200,655]. Despite these advances, hirudin is still moderately expensive to produce and it is not widely available commercially.

[0012] Recently, efforts have been made to identify peptide fragments of native hirudin or derivatives thereof which are also effective in prolonging clotting times. Such compounds are described in European patent application Nos. 276,014, 291,982, 333,356, 341,607 and 372,670. The molecules described in these patent applications demonstrated varying efficacy in inhibiting clot formation, but were all 2 to 4 orders of magnitude less potent than hirudin. Such peptide fragments, therefore, may not be fully satisfactory to dissolve blood clots in on-going therapy regimens.

[0013] More recently, compounds which mimic the action of hirudin by binding to both the anion binding exosite and the catalytic site of thrombin have been described [copending United States patent application Serial Nos. 395,482 and 549,388]. These compounds demonstrate thrombin inhibitory activity equal to or greater than native hirudin. They are also smaller than hirudin and therefore less antigenic. These inhibitors are also produced synthetically, allowing for the production of commercially feasible quantities at reasonable costs.

[0014] Despite the developments to date, there is an ongoing need for even more potent thrombin inhibitors which can be produced inexpensively and in commercially feasible quantities. Such inhibitors would not only be effective in treating and preventing vascular disease, but may also be therapeutically useful in treating cancer, neurodegenerative disease and inflammation.

SUMMARY OF THE INVENTION

[0015] The present invention provides molecules which are potent inhibitors of thrombin. These molecules have been

designed based on the three-dimensional X-ray crystallographic structure of a thrombin-inhibitor complex. Because of this, the inhibitors of the present invention are spatially configured to provide the best fit into the three-dimensional spaces in and around the catalytic site of thrombin. This results in molecules which have optimal thrombin inhibitory activity.

[0016] The invention further provides thrombin inhibitors which additionally comprise a moiety that binds to the anion binding exosite of thrombin. These inhibitors qualitatively mimic the action of hirudin. Because these molecules are designed for optimal spatial configuration, they are more potent than hirudin. The high potency of the inhibitors of this invention allows them to be administered to patients in dosages which are comparatively lower than those required in hirudin-based therapy regimens.

[0017] The molecules of this invention may be utilized in compositions and methods for inhibiting any thrombin-mediated or thrombin-associated function or process. Pharmaceutical compositions containing these molecules, as well as methods of treatment or prophylaxis of vascular diseases, inflammatory responses, carcinomas, and neurodegenerative diseases using these inhibitors are also part of the present invention. These molecules may also be employed in compositions and methods for *ex vivo* imaging, for storing and treating extracorporeal blood and for coating invasive devices. And the molecules of this invention may be administered to a patient in combination with a fibrinolytic agent to increase the efficacy of a given dose of that agent or to lower the dose of that agent required for a given effect, such as dissolving a blood clot.

[0018] Due to the fact that the molecules of the present invention may be prepared by chemical synthesis techniques, commercially feasible amounts may be produced inexpensively. Moreover, because the molecules of the present invention are significantly smaller than the thrombin inhibitors presently employed in medical treatment, they are less likely to stimulate an undesirable immune response in patients treated with them. Accordingly, the use of these thrombin inhibitors is not limited to the treatment of acute disease. These molecules may also be utilized in therapy for chronic thromboembolic diseases, such as atherosclerosis and restenosis following angioplasty. The molecules of the present invention may also be utilized in a variety of other applications in place of known thrombin inhibitors, especially heparin or hirudin.

[0019] As will be appreciated from the disclosure to follow, the molecules, compositions and methods of this invention are useful in the treatment and prevention of various diseases attributed to the undesirable effects of thrombin, as well as for diagnostic purposes.

BRIEF DESCRIPTION OF THE DRAWINGS

[0020]

Figure 1 depicts a space filling model of the Hirulog-8-thrombin complex.

Figure 2 is a schematic depiction of the interaction between the 1-position D-Phe and 2-position proline of Hirulog-8 and the hydrophobic pocket adjacent the catalytic site of thrombin.

Figure 3 depicts the comparative anticoagulant activity of hirugen, recombinant hirudin, hirulog-8 and D-Cha-hirulog.

DETAILED DESCRIPTION OF THE INVENTION

[0021] The following common abbreviations of the amino acids are used throughout the specification and in the claims:


Orn - ornithine	Gly - glycine
Ala - alanine	Val - valine
Leu - leucine	Ile - isoleucine
Pro - proline	Phe - phenylalanine
Trp - tryptophan	Met - methionine
Ser - serine	Thr - threonine
Cys - cysteine	Tyr - tyrosine
Asn - asparagine	Gln - glutamine
Asp - aspartic acid	Glu - glutamic acid
Lys - lysine	Arg - arginine
His - histidine	Nle - norleucine
Npa - naphthylalanine	Cha - cyclohexylalanine
Hyp - hydroxyproline	TPro - thioproline
Ac - acetyl	Suc - succinyl

(continued)


BOC - <u>tert</u> Butoxycarbonyl	Tos - <u>para</u> Toluenesulfonyl
Cbz - Carbobenzyloxy	Inp - isonipecotic acid
3,4,-dehydroPro - 3,4,-dehydroproline	Sar - sarcosine (N-methylglycine)
Tyr(OSO ₃ H) - tyrosine-O-sulfate	Tyr(SO ₃ H) - 3-sulfo-tyrosine
3-,5-diiodoTyr - 3-,5-diiodotyrosine	

[0022] The term "any amino acid" as used herein includes the L-isomers of the naturally occurring amino acids, as well as other "non-protein" α -amino acids commonly utilized by those in the peptide chemistry arts when preparing synthetic analogs of naturally occurring amino peptides. The naturally occurring amino acids are glycine, alanine, valine, leucine, isoleucine, serine, methionine, threonine, phenylalanine, tyrosine, tryptophan, cysteine, proline, histidine, aspartic acid, asparagine, glutamic acid, glutamine, γ -carboxyglutamic acid, arginine, ornithine and lysine. Examples of "non-protein" α -amino acids include norleucine, norvaline, alloisoleucine, homoarginine, thioproline, dehydroproline, hydroxyproline (Hyp), isonipecotic acid (Inp), homoserine, cyclohexylglycine (Chg), α -amino-n-butyric acid (Aba), cyclohexylalanine (Cha), aminophenylbutyric acid (Pba), phenylalanines substituted at the ortho, meta, or para position of the phenyl moiety with one or two of the following: a (C₁-C₄) alkyl, a (C₁-C₄) alkoxy, halogen or nitro groups or substituted with a methylenedioxy group; β -2- and 3-thienylal-alanine, β -2- and 3-furanylalanine, β -2-, 3- and 4-pyridylalanine, β -(benzothienyl-2- and 3-yl)alanine, β -(1- and 2-naphthyl)alanine, O-alkylated derivatives of serine, threonine or tyrosine, S-alkylated cysteine, S-alkylated homocysteine, O-sulfate, O-phosphate and O-carboxylate esters of tyrosine, 3- and 5-sulfo tyrosine, 3- and 5-carboxy-tyrosine, 3- and 5-phospho-tyrosine, 4-methane sulfonic acid ester of tyrosine, 4-methane phosphonic acid ester of tyrosine, 4-phenylacetic acid, 3,5-diiodotyrosine, 3- and 5-nitrotyrosine, ϵ -alkyl lysine, delta-alkyl ornithine, and the D-isomers of any of the above amino acids. Unless specifically indicated, all amino acids referred to in this application are in the L-form.

[0023] The compounds referred to herein as tyrosine-O-sulfate, Tyr(OSO₃H) and O-sulfate ester of tyrosine are identical and have the structural formula:



[0024] The compounds referred to herein as Tyr(SO₃H), 3-sulfo-tyrosine and 5-sulfo-tyrosine are identical and have the structural formula:



[0025] The term "patient" as used in this application refers to any mammal, especially humans.




[0026] The term "anionic amino acid" as used herein means a meta, para or ortho, mono- or di-substituted phenylalanine, cyclohexylalanine or tyrosine containing a negatively charged moiety, as well as S-alkylated cysteine, S-alkylated homocysteine, γ -carboxyglutamic acid, ϵ -alkyl lysine, delta-alkyl ornithine, glutamic acid, and aspartic acid. Examples of anionic amino acids are O-sulfate, O-phosphate and O-carboxylate esters of tyrosine, 3- and 5-sulfo-tyrosine, 3- and 5-carboxy-tyrosine, 3- and 5-phospho-tyrosine, 4-methane sulfonic acid ester of tyrosine, 4-methane phosphonic acid ester of tyrosine, 4-phenylacetic acid, 3,5-diiodotyrosine and 3- and 5-nitrotyrosine.

[0027] The terms "catalytic site", "active site" and "active site pocket" as used herein, each refer to any or all of the following sites in thrombin: the substrate binding or "S₁" site; the hydrophobic binding or "oily" site; and the site where

cleavage of a substrate is actually carried out ("charge relay site").

[0028] The term "backbone chain" as used herein, refers to the portion of a chemical structure that defines the smallest number of consecutive bonds that can be traced from one end of that chemical structure to the other. The atomic components that make up a backbone chain may comprise any atoms that are capable of forming bonds with at least two other atoms.


[0029] For example, each of the following chemical structures is characterized by a backbone chain of 7 atoms (the atoms which comprise the backbone chain are indicated in boldface):



[0030] The term "calculated length" as used in this application, refers to a predicted measurement derived by summing up the bond lengths between the atoms that comprise the backbone chain. Bond lengths between any two given atoms are well known in the art [see, for example, CRC Handbook of Chemistry and Physics, 65th Edition, R. C. Weist, ed., CRC Press, Inc., Boca Raton, FL, pp. F-166-70 (1984)].

[0031] Applicants have analyzed the structure of a thrombin-Hirulog-8 complex by three-dimensional X-ray crystallography. Hirulog-8 is an inhibitor that binds to both the anionic binding exosite and the catalytic site of thrombin. It has the formula: (D-Phe)-Pro-Arg-Pro-(Gly)₄-Asn-Gly-Asp-Phe-Glu-Glu-Ile-Pro-Glu-Glu-Tyr-Leu. The synthesis of this compound is described in copending United States patent application Serial No. 395,482 and in Example 1 of this application. This crystallographic data revealed several structural features in and around the active site of thrombin that were crucial in designing the improved thrombin inhibitors of this invention.

[0032] One of these structural features is a hydrophobic pocket in thrombin adjacent to its catalytic center. In Hirulog-8, the N-terminal D-Phe residue, specifically the phenyl ring of that amino acid, occupies this space. Substituting a saturated ring for the unsaturated ring increases the lipophilic interactions with thrombin, thus increasing inhibitory potency. Therefore, according to one embodiment of this invention, the thrombin inhibitor has the formula:




wherein x is hydrogen or is characterized by a backbone chain consisting of from 1 to 100 atoms; R₁ is selected from the group consisting of unsubstituted, mono-substituted, di-substituted and tri-substituted saturated homocyclic or heterocyclic ring structures; R₂ is a bond or is characterized by a backbone chain consisting of from 1 to 5 atoms; R₃ is a bond or is characterized by a backbone chain consisting of from 1 to 3 atoms; R₄ is any amino acid; R₅ is any L-amino acid which comprises a guanidinium- or amino-containing side chain group; R₆ is a non-amide bond; and Y is characterized by a backbone chain consisting of from 1 to 9 atoms. Examples of L-amino acids which comprise a guanidinium- or amino-containing side chains are arginine, lysine and ornithine.

[0033] Preferably, the saturated homocyclic or heterocyclic ring structure is supplied by a D-cyclohexyl-alanine (D-Cha), a mono-substituted D-Cha, a di-substituted D-Cha or a tri-substituted D-Cha residue (i.e., X is H₂N; R₁ is selected from the group consisting of unsubstituted, mono-substituted, di-substituted and tri-substituted hexane; R₂ is CH₂-CH; and R₃ is C=O). Most preferably, X is H₂N; R₁ is hexane; R₂ is CH₂-CH; R₃ is C=O; R₄ is proline; R₅ is arginine and Y is proline.

[0034] The presence of a non-amide bond at R₆ retards or prevents cleavage of the inhibitor by thrombin. The non-amide bond component may be formed by chemically modifying an amide bond. This may be achieved by methods well known in the art [M. Szelke et. al., "Potent New Inhibitors of Human Renin", Nature, 299, pp. 555-57 (1982); D. H. Coy et al., "Facile Solid Phase Preparation of Proteins containing the CH₂-NH Peptide Bond Isostere and Application to the Synthesis of Somatostatin (SRIF) Octapeptide Analogues", Peptides 1986, D. Theodopoulos, Ed., Walter Gruyter & Co., Berlin, pp. 143-46 (1987)]. When a non-amide bond is formed in this manner, it is preferable that the chemical modification be performed prior to the addition of the portion of the molecule containing this bond to the rest of the thrombin inhibitor. In this manner, the portion containing this non-amide bond can be added en bloc, in a single synthesis step, to the rest of the inhibitor.

[0035] In the most preferred embodiment R₅ is Arg and Y is Pro. In this embodiment R₆ is a naturally occurring imide bond, which is slowly cleaved by thrombin. This avoids the necessity of pre-forming the non-amide bond and allows Y and R₆ to be added to the rest of the inhibitor sequentially rather than en bloc.

[0036] Further analysis of the Hirulog-8-thrombin crystallographic structure revealed that the proline bound to the D-Phe of Hirulog-8 was within 3.46 Å of the hydroxyl group of Tyr76 of thrombin. Because this distance was close enough to form hydrogen bonds, the substitution of an amino acid comprising a side chain group characterized by the capacity to accept a hydrogen bond at a pH of between about 5.5 and 9.5 for Pro at this position should increase the binding affinity of the inhibitor. Accordingly, another embodiment of this invention is a thrombin inhibitor of the formula:



wherein **R₁'** is selected from the group consisting of unsubstituted, mono-substituted, di-substituted and tri-substituted ring structures; R₂ is a bond or is characterized by a backbone chain consisting of from 1 to 5 atoms; R₃ is a bond or is characterized by a backbone chain consisting of from 1 to 3 atoms; **R₄'** is any amino acid comprising a side chain group characterized by the capacity to accept a hydrogen bond at a pH of between about 5.5 and 9.5; R₅ is any L-amino acid which comprises a guanidinium- or amino-containing side chain group; R₆ is a non-amide bond; and Y is characterized by a backbone chain consisting of from 1 to 9 atoms.

[0037] Amino acids comprising a side chain group characterized by the capacity to accept a hydrogen bond at a pH of between about 5.5 and 9.5 are well known in the art. For example, histidine (which contains an imidazolium nitrogen), thioproline (which contains a thiol group) and isonipecotic acid (which contains a carboxylate side group) are each known to be hydrogen bond acceptors at pH 5.5 to 9.5.

[0038] More preferred thrombin inhibitors according to this embodiment comprise an unsubstituted, mono-substituted,


di-substituted and tri-substituted hexane at the R_1' position. Most preferably X is H_2N , R_1' is hexane, R_2 is CH_2-CH and R_3 is $C=O$ and the resulting amino acid formed by X, R_1' , R_2 and R_3 is in the D -configuration (i.e., D -Cha)

[0039] In another preferred embodiment the thrombin inhibitor of this invention additionally consists of an anion binding exosite associating moiety (ABEAM), and a linker bound to Y at one end and the ABEAM at the other. Similar thrombin inhibitors have been described in copending United States application Serial Nos. 549,388, filed July 6, 1990, and 395,482, filed August 18, 1989, both of which are herein incorporated by reference, but the present preferred inhibitors are surprisingly and unexpectedly more potent.

[0040] In this embodiment, the linker region of the inhibitor provides a bridge between the CSDM and the ABEAM. Accordingly, it is the length of the linker, rather than its structure, that is of prime importance. The calculated length of the backbone chain which characterizes the linker must be at least about 18 Å --the distance between the catalytic site and the anion binding exosite of thrombin -- and less than about 42 Å.

[0041] The backbone chain of the linker may comprise any atoms which are capable of bonding to at least two other atoms. Preferably, the backbone chain consists of any chemically feasible combination of atoms selected from oxygen, carbon, nitrogen and sulfur. Those of skill in the art are aware of what combination of the above backbone chain atoms falls within the required length based on known distances between various bonds [see, for example, R. T. Morrison and R. N. Boyd, Organic Chemistry, 3rd Edition, Allyn and Bacon, Inc., Boston, Massachusetts (1977)]. According to a preferred embodiment, the linker is a peptide which comprises the amino acid sequence Gly-Gly-Gly-Asn-Gly-Asp-Phe. Preferably, the amino acid bound to the ABEAM component is Phe.

[0042] The third domain of these preferred thrombin inhibitors is the ABEAM which binds to the anion binding exosite of thrombin. Preferably the ABEAM has the formula:



wherein W is a bond; B_1 is an anionic amino acid; B_2 is any amino acid; B_3 is Ile, Val, Leu, Nle or Phe; B_4 is Pro, Hyp, 3,4-dehydroPro, thiazolidine-4-carboxylate, Sar, any N-methyl amino acid or D-Ala; B_5 is an anionic amino acid; B_6 is an anionic amino acid; B_7 is a lipophilic amino acid selected from the group consisting of Tyr, Trp, Phe, Leu, Nle, Ile, Val, Cha, Pro, or a dipeptide consisting of one of these lipophilic amino acids and any amino acid; B_8 is a bond or a peptide containing from one to five residues of any amino acid; and Z is OH or is characterized by a backbone chain consisting of between 1 and 6 atoms.

[0043] Peptides which are homologous to the carboxy terminal portion of hirudin have been shown to bind to the anion binding exosite on thrombin [copending United States patent application 314,756 and J. M. Maraganore et al., "Anticoagulant Activity of Synthetic Hirudin Peptides", J. Biol. Chem., 264, pp. 8692-98 (1989); both of which are herein incorporated by reference].

[0044] According to a preferred embodiment of this invention, ABEAM is homologous to amino acids 56-64 of hirudin, i.e., B_1 is Glu; B_2 is Glu; B_3 is Ile; B_4 is Pro; B_5 is Glu; B_6 is Glu; B_7 is Tyr-Leu, Tyr(SO_3H)-Leu or Tyr(OSO_3H)-Leu, or (3-,5-diiodoTyr)-Leu; B_8 is a bond; and Z is OH. It should be noted that native hirudin contains Tyr(OSO_3H) at position 63. However, carboxy terminal hirudin peptides which contain Tyr(SO_3H) have identical anticoagulant activity as those which contain the native Tyr(OSO_3H) [see copending United States patent application 314,756].

[0045] Other ABEAM components within the scope of this invention may comprise those portions of any molecule known to bind to the anion binding site of thrombin. These include amino acids 1675-1686 of Factor V, amino acids 272-285 of platelet glycoprotein Ib, amino acids 415-428 of thrombomodulin, amino acids 245-259 of prothrombin Fragment 2 and amino acids 30 to 44 of fibrinogen A α chain. In addition, the ABEAM component may be selected from any of the hirudin peptide analogues described by J.L. Krstenansky et al., "Development of MDL-28,050, A Small Stable Antithrombin Agent Based On A Functional Domain of the Leech Protein, Hirudin", Thromb. Haemostas., 63, pp. 208-14 (1990), particularly those comprising the sequence Asp-Tyr-Glu-Pro-Ile-Pro-Glu-Glu-Ala-Cha-(D -Glu).

[0046] The thrombin inhibitors of the present invention may be synthesized by various techniques which are well known in the art. These include organic chemical synthesis techniques, solid-phase peptide synthesis, solution-phase peptide synthesis or a combination of these techniques. Portions of some of the inhibitors of this invention may also be produced by other methods, such as enzymatic cleavage of natural or recombinant hirudin or recombinant DNA techniques. These portions may then be bound to the synthetically produced portions of the inhibitor to produce the final product according to this invention. The choice of synthesis technique will, of course, depend upon the composition of the particular inhibitor.

[0047] In a preferred embodiment of this invention, the thrombin inhibitor is synthesized via a mixed heterologous/solid phase technique. This technique involves the solid-phase synthesis of all or most of the peptide portion of the molecule, followed by the addition of the non-amino acid components which are synthesized by solution phase techniques. The non-amino acid may be coupled to the peptidic portion via solid-phase or solution-phase methods. Similarly, any

remaining peptidic portions may also be added via solid-phase or solution phase methods. This constitutes the most cost-efficient procedures for producing commercial quantities of these molecules.

[0048] When "non-protein" amino acids are contained in the thrombin inhibitor of this invention, they may be either added directly to the growing chain during peptide synthesis or prepared by chemical modification of the complete synthesized peptide, depending on the nature of the desired "non-protein" amino acid. Those of skill in the chemical synthesis art are well aware of which "non-protein" amino acids may be added directly and which must be synthesized by chemically modifying the complete peptide chain following peptide synthesis.

[0049] The molecules of the present invention display potent anticoagulant activity. This activity may be assayed *in vitro* using any conventional technique. Preferably, an assay for anticoagulant activity involves direct determination of the thrombin-inhibitory activity of the molecule. Such techniques measure the inhibition of thrombin-catalyzed cleavage of colorimetric substrates or, more preferably, the increase in thrombin times or increase in activated partial thromboplastin times of human plasma. The latter assay measures factors in the "intrinsic" pathway of coagulation. Alternatively, the assay employed may use purified thrombin and fibrinogen to measure the inhibition of release of fibrinopeptides A or B by radioimmunoassay or ELISA.

[0050] The antiplatelet activity of the molecules of this invention may also be measured by any of a number of conventional platelet assays. Preferably, the assay will measure a change in the degree of aggregation of platelets or a change in the release of a platelet secretory component in the presence of thrombin. The former may be measured in an aggregometer. The latter may be measured using RIA or ELISA techniques specific for the secreted component.

[0051] The molecules of the present invention are useful in compositions, combinations and methods for the treatment and prophylaxis of various diseases attributed to thrombin-mediated and thrombin-associated functions and processes. These include myocardial infarction, stroke, pulmonary embolism, deep vein thrombosis, peripheral arterial occlusion, restenosis following arterial injury or invasive cardiological procedures, acute or chronic atherosclerosis, edema and inflammation, various cell regulatory processes (e.g. secretion, shape changes, proliferation), cancer and metastasis, and neurodegenerative diseases.

[0052] The thrombin inhibitors of the present invention may be formulated using conventional methods to prepare pharmaceutically useful compositions, such as the addition of a pharmaceutically acceptable carrier. These compositions and the methods employing them may be used for treating or preventing thrombotic diseases in a patient.

[0053] According to an alternate embodiment of the present invention, the thrombin inhibitors may be employed in combinations, compositions, and methods for treating thrombotic disease, and for decreasing the dosage of a thrombolytic agent required to establish reperfusion or prevent reocclusion in a patient. Additionally, the thrombin inhibitors of this invention may be used in combinations, compositions, and methods for decreasing reperfusion time or increasing reocclusion time in a patient treated with a thrombolytic agent. These combinations and compositions comprise a pharmaceutically effective amount of a thrombin inhibitor of the present invention and a pharmaceutically effective amount of a thrombolytic agent.

[0054] In these combinations and compositions, the thrombin inhibitor and the thrombolytic agent work in a complementary fashion to dissolve blood clots, resulting in decreased reperfusion times and increased reocclusion times in patients treated with them. Specifically, the thrombolytic agent dissolves the clot, while the thrombin inhibitor prevents newly exposed, clot-entrapped or clot-bound thrombin from regenerating the clot. The use of the thrombin inhibitor in the combinations and compositions of this invention advantageously allows the administration of a thrombolytic reagent in dosages previously considered too low to result in thrombolytic effects if given alone. This avoids some of the undesirable side effects associated with the use of thrombolytic agents, such as bleeding complications.

[0055] Thrombolytic agents which may be employed in the combinations and compositions of the present invention are those known in the art. Such agents include, but are not limited to, tissue plasminogen activator purified from natural sources, recombinant tissue plasminogen activator, streptokinase, urokinase, prourokinase, anisolated streptokinase plasminogen activator complex (ASPAC), animal salivary gland plasminogen activators and known, biologically active derivatives of any of the above.

[0056] The term "combination" as used herein, includes a single dosage form containing at least one thrombin inhibitor of this invention and at least one thrombolytic agent; a multiple dosage form, wherein the thrombin inhibitor and the thrombolytic agent are administered separately, but concurrently; or a multiple dosage form wherein the two components are administered separately, but sequentially. In sequential administration, the thrombin inhibitor may be given to the patient during the time period ranging from about 5 hours prior to about 5 hours after administration of the thrombolytic agent. Preferably, the thrombin inhibitor is administered to the patient during the period ranging from 2 hours prior to 2 hours following administration of the thrombolytic agent.

[0057] Alternatively, the thrombin inhibitor and the thrombolytic agent may be in the form of a single, conjugated molecule. Conjugation of the two components may be achieved by standard cross-linking techniques well known in the art. The single molecule may also take the form of a recombinant fusion protein, if both the thrombin inhibitor and the thrombolytic agent are peptidic.

[0058] Various dosage forms may be employed to administer the compositions and combinations of this invention.

These include, but are not limited to, parenteral administration, oral administration and topical application. The compositions and combinations of this invention may be administered to the patient in any pharmaceutically acceptable dosage form, including those which may be administered to a patient intravenously as bolus or by continued infusion, intramuscularly -- including paravertebrally and periarticularly -- subcutaneously, intracutaneously, intra-articularly, intrasynovially, intrathecally, intra-lesionally, periostally or by oral, nasal, or topical routes. Such compositions and combinations are preferably adapted for topical, nasal, oral and parenteral administration, but, most preferably, are formulated for parenteral administration.

[0059] Parenteral compositions are most preferably administered intravenously either in a bolus form or as a constant infusion. If the thrombin inhibitor is being used as an antiplatelet compound, constant infusion is preferred. If the thrombin inhibitor is being used as an anticoagulant, a subcutaneous or intravenous bolus injection is preferred. For parenteral administration, fluid unit dose forms are prepared which contain a thrombin inhibitor of the present invention and a sterile vehicle. The thrombin inhibitor may be either suspended or dissolved, depending on the nature of the vehicle and the nature of the particular thrombin inhibitor. Parenteral compositions are normally prepared by dissolving the thrombin inhibitor in a vehicle, optionally together with other components, and filter sterilizing before filling into a suitable vial or ampule and sealing. Preferably, adjuvants such as a local anesthetic, preservatives and buffering agents are also dissolved in the vehicle. The composition may then be frozen and lyophilized to enhance stability.

[0060] Parenteral suspensions are prepared in substantially the same manner, except that the active component is suspended rather than dissolved in the vehicle. Sterilization of the compositions is preferably achieved by exposure to ethylene oxide before suspension in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of its components.

[0061] Tablets and capsules for oral administration may contain conventional excipients, such as binding agents, fillers, diluents, tableting agents, lubricants, disintegrants, and wetting agents. The tablet may be coated according to methods well known in the art. Suitable fillers which may be employed include cellulose, mannitol, lactose and other similar agents. Suitable disintegrants include, but are not limited to, starch, polyvinylpyrrolidone and starch derivatives, such as sodium starch glycolate. Suitable lubricants include, for example, magnesium stearate. Suitable wetting agents include sodium lauryl sulfate.

[0062] Oral liquid preparations may be in the form of aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, or may be presented as a dry product for reconstitution with water or another suitable vehicle before use. Such liquid preparations may contain conventional additives. These include suspending agents; such as sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminum stearate gel or hydrogenated edible fats; emulsifying agents which include lecithin, sorbitan monooleate, polyethylene glycols, or acacia; non-aqueous vehicles, such as almond oil, fractionated coconut oil, and oily esters; and preservatives, such as methyl or propyl p-hydroxybenzoate or sorbic acid.

[0063] Compositions formulated for topical administration may, for example, be in aqueous jelly, oily suspension or emulsified ointment form.

[0064] The dosage and dose rate of the thrombin inhibitor will depend on a variety of factors, such as the size of the patient, the specific pharmaceutical composition used, the object of the treatment, i.e., therapy or prophylaxis, the nature of the thrombotic disease to be treated, and the judgment of the treating physician.

[0065] According to the present invention, a preferred pharmaceutically effective daily dose of the thrombin inhibitor of this invention is between about 0.5 nmoles/kg body weight of the patient to be treated ("body weight") and about 2.5 μ moles/kg body weight. In combinations containing a thrombolytic agent, a pharmaceutically effective daily dose of the thrombolytic is between about 10% and 80% of the conventional dosage range. The "conventional dosage range" of a thrombolytic agent is the daily dosage used when that agent is employed in a monotherapy. [Physician's Desk Reference 1989, 43rd Edition, Edward R. Barnhart, publisher]. That conventional dosage range will, of course, vary depending on the thrombolytic agent employed. Examples of conventional dosage ranges are as follows: urokinase - 500,000 to 6,250,000 units/patient; streptokinase - 140,000 to 2,500,000 units/patient; tPA - 0.5 to 5.0 mg/kg body weight; ASPAC - 0.1 to 10 units/kg body weight.

[0066] Most preferably, the therapeutic and prophylactic compositions of the present invention comprise a dosage of between about 5 nmoles/kg body weight and about 250 nmoles/kg body weight of the thrombin inhibitor. Most preferred combinations comprise the same amount of the thrombin inhibitor and between about 10% and about 70% of the conventional dosage range of a thrombolytic agent. It should also be understood that a daily pharmaceutically effective dose of either the thrombin inhibitors of this invention or the thrombolytic agent present in combinations of the invention, may be less than or greater than the specific ranges cited above.

[0067] Once improvement in the patient's condition has occurred, a maintenance dose of a combination or composition of this invention is administered, if necessary. Subsequently, the dosage or the frequency of administration, or both, may be reduced, as a function of the symptoms, to a level at which the improved condition is retained. When the symptoms have been alleviated to the desired level, treatment should cease. Patients may, however, require intermittent treatment upon any recurrence of disease symptoms.

[0068] According to an alternate embodiment of this invention, thrombin inhibitors may be used in compositions and methods for coating the surfaces of invasive devices, resulting in a lower risk of clot formation or platelet activation in patients receiving such devices. Surfaces that may be coated with the compositions of this invention include, for example, prostheses, artificial valves, vascular grafts, stents and catheters. Methods and compositions for coating these devices are known to those of skill in the art. These include chemical cross-linking or physical adsorption of the thrombin inhibitor-containing compositions to the surfaces of the devices.

[0069] According to a further embodiment of the present invention, thrombin inhibitors may be used *forex vivo* thrombus imaging in a patient. In this embodiment, the thrombin inhibitor is labelled with a radioisotope. The choice of radioisotope is based upon a number of well-known factors, for example, toxicity, biological half-life and detectability. Preferred radioisotopes include, but are not limited to, ^{125}I , ^{123}I and ^{111}In . Techniques for labelling the thrombin inhibitor are well known in the art. Most preferably, the radioisotope is ^{123}I and the labelling is achieved using ^{123}I -Bolton-Hunter Reagent. The labelled thrombin inhibitor is administered to a patient and allowed to bind to the thrombin contained in a clot. The clot is then observed by utilizing well-known detecting means, such as a camera capable of detecting radioactivity coupled to a computer imaging system. This technique also yields images of platelet-bound thrombin and meizothrombin.

[0070] This invention also relates to compositions containing the thrombin inhibitors of this invention and methods for using such compositions in the treatment of tumor metastases. The efficacy of the thrombin inhibitors of this invention for the treatment of tumor metastases is manifested by the inhibition of metastatic growth. This is based upon the presence of a procoagulant enzyme in certain cancer cells. This enzyme activates the conversion of Factor X to Factor Xa in the coagulation cascade, resulting in fibrin deposition which, in turn, serves as a substrate for tumor growth. By inhibiting fibrin deposition through the inhibition of thrombin, the molecules of the present invention serve as effective anti-metastatic tumor agents. Examples of metastatic tumors which may be treated by the thrombin inhibitors of this invention include, but are not limited to, carcinoma of the brain, carcinoma of the liver, carcinoma of the lung, osteocarcinoma and neoplastic plasma cell carcinoma.

[0071] The invention also relates to methods and compositions employing the above-described thrombin inhibitors to inhibit thrombin-induced endothelial cell activation. This inhibition includes the repression of platelet activation factor (PAF) synthesis by endothelial cells. These compositions and methods have important applications in the treatment of diseases characterized by thrombin-induced inflammation and edema, which is thought to be mediated by PAF. Such diseases include, but are not limited to, adult respiratory distress syndrome, septic shock, septicemia and reperfusion damage.

[0072] Early stages of septic shock include discrete, acute inflammatory and coagulopathic responses. It has previously been shown that injection of baboons with a lethal dose of live *E. coli* leads to marked declines in neutrophil count, blood pressure and hematocrit. Changes in blood pressure and hematocrit are due in part to the generation of a disseminated intravascular coagulopathy (DIC) and have been shown to parallel consumption of fibrinogen [F. B. Taylor et al., "Protein C Prevents the Coagulopathic and Lethal Effects of Escherichia coli Infusion in the Baboon", J.Clin.Invest., 79, pp. 918-25 (1987)]. Neutropenia is due to the severe inflammatory response caused by septic shock which results in marked increases in tumor necrosis factor levels. The thrombin inhibitors of this invention may be utilized in compositions and methods for treating or preventing DIC in septicemia and other diseases.

[0073] This invention also relates to the use of the above-described thrombin inhibitors, or compositions comprising them, as anticoagulants for extracorporeal blood. As used herein, the term "extracorporeal blood" includes blood removed in line from a patient, subjected to extracorporeal treatment, and then returned to the patient in such processes as dialysis procedures, blood filtration, or blood bypass during surgery. The term also includes blood products which are stored extracorporeally for eventual administration to a patient and blood collected from a patient to be used for various assays. Such products include whole blood, plasma, or any blood fraction in which inhibition of coagulation is desired.

[0074] The amount or concentration of thrombin inhibitor in these types of compositions is based on the volume of blood to be treated or, more preferably, its thrombin content. Preferably, an effective amount of a thrombin inhibitor of this invention for preventing coagulation in extracorporeal blood is from about 0.5 nmoles/60 ml of extracorporeal blood to about 2.5 μmoles /60 ml of extracorporeal blood.

[0075] The thrombin inhibitors of this invention may also be used to inhibit clot-bound thrombin, which is believed to contribute to clot accretion. This is particularly important because commonly used anti-thrombin agents, such as heparin and low molecular weight heparin, are ineffective against clot-bound thrombin.

[0076] Finally, the thrombin inhibitors of this invention may be employed in compositions and methods for treating neurodegenerative diseases. Thrombin is known to cause neurite retraction, a process suggestive of the rounding in shape changes of brain cells and implicated in neurodegenerative diseases, such as Alzheimer's disease and Parkinson's disease.

[0077] In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

EXAMPLE 1Design Of A Thrombin Inhibitor Capable Of Blocking The Catalytic Site And Binding To The Anion Binding Exosite

[0078] Carboxy terminal hirudin peptides effectively block thrombin-catalyzed fibrinogen hydrolysis, but not chromogenic substrate hydrolysis [J. M. Maraganore et al., J. Biol. Chem., 264, pp. 8692-98 (1989)]. In addition, hirudin peptides do not neutralize thrombin-catalyzed activation of Factors V and VIII [J. W. Fenton, II, et al., "Hirudin Inhibition by Thrombin", *Angio. Archiv. Biol.*, 18, p. 27 (1989)].

[0079] Hirudin peptides, such as Tyr₆₃-O-sulfate-N-acetyl-hirudin₅₃₋₆₄ ("hirugen"), exhibit potent inhibitory effects toward thrombin-induced platelet activation *in vitro* [J. A. Jakubowski and J. M. Maraganore, "Inhibition of Thrombin-Induced Platelet Activities By A Synthetic 12 Amino Acid Residue Sulfated Peptide (Hirugen)", *Blood*, p. 1213 (1989)]. Nevertheless, a thrombin inhibitor capable of blocking the active site may be required for inhibition of platelet thrombosis *in vivo*, if activation of Factors V and VIII are critical and rate-limiting steps. This conclusion is warranted from results obtained with the irreversible thrombin inhibitor (D-Phe)-Pro-Arg-CH₂Cl [S. R. Hanson and L. A. Harker, "Interruption of Acute Platelet-Dependent Thrombosis by the Synthetic Antithrombin D-Phenylalanyl-L-Prolyl-L-Arginyl Chloromethyl Ketone", *Proc. Natl. Acad. Sci. USA*, 85, pp. 3184-88 (1988)] and other reversible thrombin inhibitors [J. F. Eid et al., "Thrombin is an Important Mediator of Platelet Aggregation in Stenosed Canine Coronary Arteries with Endothelial Injury" *J. Clin. Invest.*, 84, pp. 18-27 (1989)].

[0080] Using the above knowledge that the NH₂-terminus of hirudin peptides is proximal to Lys-149, we employed a three-dimensional model of thrombin [B. Furie, et al., "Computer-Generated Models of Blood Coagulation Factor Xa, Factor IXa, and Thrombin Based Upon Structural Homology with Other Serine Proteases", *J. Biol. Chem.*, 257, pp. 3875-82 (1982)] to design an agent which: 1) binds to the anion binding exosite of thrombin; and, 2) is capable of blocking the active site pocket of thrombin and inhibiting the function of catalytic residues contained therein.

[0081] We determined that the minimal distance from the ε-NH₂ of Lys-149 to the β-hydroxylate of Ser-195 is 18-20 Å. Based on a 3 Å/amino acid residue length, we calculated that at least about 4 - 7 amino acids would be required to link a hirudin peptide, such as Tyr₆₃-O-sulfate-hirudin₅₃₋₆₄, to a domain comprising an active-site inhibitor structure. The composition of the linker was designed to be glycine. Glycine was chosen in order to engineer the greatest flexibility of a linker for these preliminary investigations. It should be understood, however, that other, more rigid biopolymer linkers may also be employed.

[0082] We chose the sequence (D-Phe)-Pro-Arg-Pro as the active site inhibitor because thrombin exhibits specificity for Arg as the P₁ amino acid in the cleavage of substrates. A Pro following the Arg (the P_{1'} amino acid) yields a bond that is cleaved very slowly by thrombin. We designed alternate peptides by replacing that Pro with a sarcosyl- or N-methyl-alanine amino acid or by chemical reduction of an Arg-Gly scissile bond.

EXAMPLE 2Synthesis Of Hirulog-8

[0083] Hirulog-8 has the formula: H-(D-Phe)-Pro-Arg-Pro-(Gly)₄-Asn-Gly-Asp-Phe-Glu-Glu-Ile-Pro-Glu-Glu-Tyr-Leu-OH. We synthesized Hirulog-8 by conventional solid-phase peptide synthesis employing an Applied Biosystems 430 A Peptide Synthesizer. This peptide was synthesized using BOC-L-Leucine-O-divinylbenzene resin. Additional t-BOC-amino acids (Peninsula Laboratories, Belmont, CA) used included BOC-O-2,6-dichlorobenzyl tyrosine, BOC-L-glutamic acid (γ-benzyl ester), BOC-L-proline, BOC-L-isoleucine, BOC-L-phenylalanine, BOC-L-aspartic acid (β-benzyl ester), BOC-glycine, BOC-L-asparagine, BOC-D-phenylalanine, and BOC-L-arginine. In order to achieve higher yields in synthesis, the (Gly)₄ linker segment was attached in two cycles of manual addition of BOC-glycylglycine (Beckman Biosciences, Inc., Philadelphia, PA). After completion of synthesis, the peptide was fully deprotected and uncoupled from the divinylbenzene resin by treatment with anhydrous HF: p-cresol: ethylmethyl sulfate (10:1:1, v/v/v). Following removal from the resin, the peptide was lyophilized to dryness.

[0084] Crude Hirulog-8 was purified by reverse-phase HPLC employing an Applied Biosystems 151A liquid chromatographic system and a Vydac C₁₈ column (2.2 x 25 cm). The column was equilibrated in 0.1% TFA/water and developed with a linear gradient of increasing acetonitrile concentration from 0 to 80% over 45 minutes in the 0.1% TFA at a flow-rate of 4.0 ml/min. The effluent stream was monitored for absorbance at 229 nm and fractions were collected manually. We purified 25-30 mg of crude Hirulog-8 by HPLC and recovered 15-20 mg of pure peptide.

[0085] We confirmed the structure of purified Hirulog-8 by amino acid and sequence analyses. Amino acid hydrolysates were prepared by treating the peptide with 6 N HCl, *in vacuo*, at 110°C for 24 hrs. We then analyzed the hydrolysates by ion-exchange chromatography and subsequent ninhydrin derivatization/detection using a Beckman 6300 automated analyzer. We performed sequence analysis using automated Edman degradation on an Applied Biosystems 470A gas-

phase sequencer equipped with a Model 900A data system. Phenylthiohydantoin (PTH) amino acids were analyzed on-line using an Applied Biosystems 120A PTH-analyzer and a PTH-C₁₈ column (2.1 X 220 mm).

EXAMPLE 3

Design Of 1- And 2-Position Substituted Hirulogs

[0086] We obtained the X-ray crystallographic structure of the Hirulog-8:thrombin complex by the following steps. First, we generated Hirulog-8:thrombin complex crystals of suitable quality to obtain a high resolution diffraction pattern. We next collected diffractometer data employing these crystals. Finally, we determined the 3-dimensional structure of the Hirulog-8:thrombin complex using molecular replacement rotation/translation methods employing the coordinates of the PPACK:thrombin [W. Bode et al., "The Refined 1.9 Å Crystal Structure of Human α -Thrombin: Interaction With D-Phe-Pro-Arg-Chloromethylketone and Significance of the Tyr-Pro-Pro-Trp Insertion Segment", EMBO J., 8, pp. 3467-75 (1989)] and hirudin:thrombin [T. J. Rydel et al., "The Structure of a Complex of Recombinant Hirudin and Human α -Thrombin", Science, 249, pp. 277-80, (1990)] complexes. As shown in Figure 1, the structure of Hirulog-8 bound to thrombin was resolved allowing resolution of the D-Phe-Pro-Arg sequence of the CSDM and the Asp-Phe-Glu-Glu-Ile segment of the ABEAM.

[0087] In Figure 1, thrombin is displayed in white, except for its active site, which is shown in dense dots. Hirulog-8 is depicted in sparse dots. The left-hand portion of Hirulog-8 nearest the thrombin active site is the CSDM. The right-hand portion is the ABEAM. Other amino acids of Hirulog-8 are not shown in Figure 1 because electron densities corresponding to them could not be assigned.

[0088] Examination of the CSDM portion of the Hirulog-8:thrombin structure showed the positioning of the 1-position amino acid (D-Phe) in a hydrophobic pocket formed by His57, Tyr60A, Trp60D, Leu99, Ile174, and Trp215 of thrombin. The D-Phe residue formed close van der Waals contacts with Leu99, Ile174, and Trp215 (Figure 2). In Figure 2, thrombin is displayed in solid lines and Hirulog-8 is in dashed lines. The positioning of the D-Phe residue within the pocket suggested that substitutions at the 1-position which enhance lipophilic contacts would lead to an increased binding affinity of the CSDM moiety in the thrombin inhibitors of this invention. Accordingly, we replaced the D-Phe residue of Hirulog-8 with either D-naphthylalanine (D-NPA) or D-cyclohexylalanine (D-Cha) to form D-NPA-Hirulog-8 and D-Cha-Hirulog-8, respectively.

[0089] The binding of the CSDM of Hirulog-8 to the catalytic site of thrombin was also found to include apolar interactions between the first proline of the inhibitor (adjacent to D-Phe) and a pocket defined by His57, Tyr60A and Trp60D of thrombin (Figure 2). Furthermore, that proline was found to be within 3.46 Å of the phenolic hydroxyl group of thrombin Tyr60A (Figure 2). In Figure 2, thrombin is depicted in solid lines and Hirulog-8 in dashed lines.

[0090] The proximity of this proline to Tyr60A of thrombin suggested the potential for hydrogen bond formation between the two. By substituting proline with an amino acid capable of forming hydrogen bonds, the stability of the CSDM binding to the thrombin active site can be increased. This, in turn, would increase the inhibitory activity of such a molecule. Therefore, we replaced the proline of Hirulog-8 with either L-histidine (His₂-Hirulog-8), L-thioprolin (TPro₂-Hirulog-8) or isonipecotic acid (Inp₂-Hirulog-8). Each of these substitutions created a hydrogen bond acceptor at the 2-position

(**R**₄' component) of the thrombin inhibitors of this invention (i.e., an imidazolium nitrogen, a thiol and a carboxylate, respectively).

EXAMPLE 4

Synthesis of 1-Position Substituted Hirulogs

[0091] D-Npa-Hirulog-8 was synthesized in the same manner as Hirulog-8 (Example 2), except that Boc-D-naphthylalanine (Bachem Inc., Torrance, CA) was employed in place of D-Phe at the last cycle of synthesis. D-Cha-Hirulog-8 was similarly prepared using Boc-D-cyclohexylalanine (Bachem Biosciences, Philadelphia, PA) at the last cycle of synthesis.

[0092] Both 1-position substituted peptides were purified as described for Hirulog-8 in Example 2. The purified peptides were characterized by amino acid analysis and by FAB-MS.

EXAMPLE 5Synthesis of 2-Position Substituted Hirulog Derivatives

[0093] Substitutions at the 2-position were designed having the formula: (D-Cha)-X-Arg-HPro-(Gly)₄-Asn-Gly-Asp-Phe-Glu-Glu-Ile-Pro-Glu-Glu-Tyr-Leu or (D-Phe)-X-Arg-HPro-(Gly)₄-Asn-Gly-Asp-Phe-Glu-Glu-Ile-Pro-Glu-Glu-Tyr-Leu, wherein X is either histidine, thioproline or isonipecotic acid. These peptides are synthesized essentially as described in Example 2 and Example 4 except for incorporation of Boc-L-hydroxyproline (Bachem, Inc.) instead of Boc-L-proline at cycle 16 of the synthesis and either Boc-N-im-CBZ-L-histidine, Boc-L-thioproline, or Boc-isonipecotic acid (all obtained from Bachem, Inc.) instead of Boc-L-proline at cycle 18. HPro is used at the 4-position to slow down the rate of cleavage of the inhibitor by thrombin. The peptides are purified and characterized as described in Example 2.

EXAMPLE 6Characterization Of Antithrombin Activities Of 1-Position Substituted Hirulogs

[0094] We compared the inhibition of thrombin-catalyzed hydrolysis of Spectrozyme TH (tosyl-Gly-Pro-Arg-p-nitroanilide; American Diagnostica, New York, NY) of Hirulog-8, D-Cha-Hirulog-8 and D-Npa-Hirulog-8 in an assay. Specifically, we measured the initial rate velocities in the presence or absence of each inhibitor over a range of substrate concentrations from 2.2 to 22 μ M. The thrombin-catalyzed rate was monitored in a Cary 19 spectrophotometer at 405 nm and recorded continuously as a function of time. Kinetics were performed at room temperature ($25 \pm 1^\circ\text{C}$) in a 0.05 M sodium borate buffer, pH 8.4, containing 0.1 M NaCl.

[0095] For a typical enzyme reaction, 1.0 ml of buffer was added to both the sample and reference cuvettes. Thrombin (3.2×10^{-9} M, final concentration) and the hirulog ($0 - 4 \times 10^{-8}$ M) were added to the sample cuvette prior to addition of Spectrozyme TH (2.2 - 22 μ M). Immediately following addition of substrate, the contents of the sample cuvette were mixed by use of a plastic pipette. The reaction was monitored spectrophotometrically for 5 - 15 minutes.

[0096] Initial rate velocities at each substrate concentration were expressed as moles Spectrozyme TH hydrolyzed/sec/mole thrombin. This was determined during the initial linear phase of the reaction ($\leq 15\%$ total hydrolysis of substrate) by measuring the slope of the hydrolytic reaction. Lineweaver-Burke plots were constructed accordingly, by plotting the inverse of the initial velocity against the inverse of the substrate concentration. Shown below are the inhibitory constants obtained for Hirulog-8 and derivatives of the present invention.

Derivative	K_i , nM
Hirulog-8	1.4
D-Cha-Hirulog-8	0.12
D-Npa-Hirulog-8	4.3

[0097] As can be seen from these results, substitution of D-Phe in Hirulog-8 with D-Cha results in a surprising and unexpected decrease in K_i by one order of magnitude. This finding indicates that substitution of D-Phe by D-Cha increases the binding affinity of the CSDM in the inhibitors of this invention. The failure of D-Npa-Hirulog-8 to decrease K_i indicates that the presence of a saturated ring structure at this position causes the increased binding affinity. D-Cha contains such a saturated ring, while D-Npa contains an unsaturated ring.

[0098] Molecules containing the 2-position substitutions described in Example 5 will display similar surprising and unexpected decreases in K_i .

EXAMPLE 7Anticoagulant Activity Of A₁-Substituted Hirulogs

[0099] We compared the anticoagulant activity of Tyr₆₃-O-sulfate-N-acetyl-hirudin₅₃₋₆₄ ("hirugen"), recombinant hirudin (American Diagnostica), Hirulog-8, and the 1-position substituted hirulogs of the present invention using pooled, normal human plasma (George King Biomedical, Overland Park, KA) and a Coag-A-Mate XC instrument (General Diagnostics, Organon Technica, Oklahoma City, OK). Activity was monitored using the activated partial thromboplastin time (APTT) assay with CaCl₂ and phospholipid solutions obtained from the manufacturer. Recombinant hirudin (American Diagnostica), Hirulog-8, D-Cha-Hirulog-8 or hirugen was then added to the APTT determination wells at a final concentrations of 10 to 32,300 ng/ml in a total volume of 25 μ l prior to addition of 100 μ l of plasma.

[0100] As shown in Figure 3, D-Cha-Hirulog-8 prolonged APTT to 470% of control values at a concentration of 1 $\mu\text{g}/\mu\text{l}$. This increase was significantly greater than the increases in APTT caused by hirugen, recombinant hirudin or Hirulog-8 at the same concentration. Thus, in addition to showing enhanced antithrombin activities *in vitro* over Hirulog-8, D-Cha-Hirulog-8 also showed a significantly increased anticoagulant effect in plasma assays over Hirulog-8.


[0101] Molecules containing the 2-position substitutions described in Example 5 will display increases in APTT that are greater than Hirulog-8.

[0102] While we have hereinbefore presented a number of embodiments of this invention, it is apparent that our basic construction can be altered to provide other embodiments which utilize the molecules, compositions, combinations and methods of this invention. Therefore, it will be appreciated that the scope of this invention is to be defined by the claims appended hereto rather than the specific embodiments which have been presented hereinbefore by way of example.

Claims

1. A thrombin inhibitor comprising:

a) a catalytic site-directed moiety comprising the formula:



wherein X is hydrogen or is **characterized by** a backbone chain consisting of from 1 to 100 atoms; R_1 is selected from the group consisting of homocyclic or heterocyclic saturated ring structures of 6 atoms which are unsubstituted, mono-substituted, di-substituted and tri-substituted; R_2 is **characterized by** a backbone chain consisting of from 1 to 5 atoms; R_3 is a bond or is **characterized by** a backbone chain consisting of from 1 to 3 atoms; R_4 is any amino acid; R_5 is any L-amino acid which comprises a guanadinium- or amino-containing side chain group; R_6 is a non-amide bond; and Y is **characterized by** a backbone chain consisting of from 1 to 9 atoms; and b) a linker moiety **characterized by** a backbone chain having a calculated length of between 18 Å and about 42 Å; and

c) an anion binding exosite associating moiety, wherein said catalytic site-directed moiety is bound to said anion binding exosite associating moiety via said linker moiety, and said inhibitor is capable of simultaneously binding to the catalytic site and the anion binding exosite of thrombin; and wherein said inhibitor is distinguished by increased lipophilic interaction with thrombin as compared to the interaction between thrombin and (D-Phe)-Pro-Arg-Pro-(Gly)4-Asn-Gly-Asp-Phe-Glu-Glu-Ile-Pro-Glu-Glu-Tyr-Leu.


2. The thrombin inhibitor of claim 1, wherein X is H_2N , R_1 is selected from the group consisting of unsubstituted, mono-substituted, di-substituted and tri-substituted hexane, R_2 is $\text{CH}_2\text{-CH}$, and R_3 is C=O .

3. The thrombin inhibitor of claim 1 or 2, wherein said catalytic site-directed moiety has the amino acid sequence: D-Cha-Pro-Arg-Pro

4. The thrombin inhibitor of claim 1 or 2, wherein R_4 is any amino acid comprising a side chain group **characterized by** the capacity to accept a hydrogen bond at a pH of between about 5.5 and 9.5.

5. The thrombin inhibitor of claims 1, 2 or 4, wherein R_1 is selected from the group consisting of histidine, thioproline and isonipecotic acid.

6. The thrombin inhibitor of any one of claims 1 to 5, wherein said anion binding exosite associating moiety consists of the formula:



wherein W is a bond, B_1 is an anionic amino acid, B_2 is any amino acid, B_3 is Ile, Val, Leu, Nle or Phe, B_4 is Pro, Hyp, 3,4-dehydroPro, thiazolidine-4-carboxylate, Sar, any N-methyl amino acid or D-Ala, B_5 is an anionic amino acid, B_6 is an anionic amino acid, B_7 is a lipophilic amino acid selected from the group consisting Tyr, Trp, Phe, Leu, Nle, Ile, Val, Cha, Pro, or a dipeptide consisting of one of these lipophilic amino acids and any amino acid, B_8

is a bond or a peptide containing from one to five residues of any amino acid, and Z is OH or is **characterized by** a backbone chain consisting of from 1 to 6 atoms.

7. The thrombin inhibitor of claim 6, wherein B₁ is Glu, B₂ is Glu, B₃ is Ile, B₄ is Pro, B₅ is Glu, B₆ is Glu, B₇ is Tyr-Leu, Tyr(SO₃H)-Leu, Tyr(OSO₃H)-Leu or (3-, 5-diiodoTyr)-Leu, B₈ is a bond, and Z is OH.
8. The thrombin inhibitor of any one of claims 1 to 7, wherein said backbone chain of said linker moiety consists of any combination of atoms selected from the group consisting of carbon, nitrogen, sulfur and oxygen.
9. The thrombin inhibitor of claim 8, wherein said linker comprises the amino acid sequence:
Gly-Gly-Gly-Asn-Gly-Asp-Phe.
10. The thrombin inhibitor of claim 9, wherein said thrombin inhibitor is D-Cha-Hirulog-8.
11. A pharmaceutical composition comprising a pharmaceutically effective amount of a thrombin inhibitor of any one of claims 1 to 10 and optionally a pharmaceutically acceptable carrier.
12. The pharmaceutically acceptable composition of claim 11, wherein said pharmaceutically effective amount is between about 0.5 nmoles/kg body weight/day to about 2.5 μ moles/kg body weight/day.
13. The pharmaceutically acceptable composition of claim 12, wherein said pharmaceutically effective amount is between about 5 nmoles/kg body weight/day to about 250 nmoles/kg body weight/day.
14. The thrombin inhibitor of any one of claims 1 to 10, further comprising a radioisotope.
15. The thrombin inhibitor of claim 14, wherein said radioisotope is selected from the group consisting of ¹²³I, ¹²⁵I and ¹¹¹In.
16. A composition for ex vivo imaging of a fibrin or a platelet thrombus in a patient, said composition comprising a pharmaceutically acceptable buffer and a thrombin inhibitor of claim 14 or 15.
17. A method for ex vivo imaging of a fibrin or a platelet thrombus in a patient comprising the steps
 - (a) administering to said patient the composition of claim 16; and
 - (b) using detecting means to observe the thrombin inhibitor present in said composition.
18. A composition for coating the surface or an invasive device to be inserted into a patient, wherein said composition comprises a suitable buffer and at Least one thrombin inhibitor of any one of claims 1 to 10.
19. A method for coating the surface of an invasive device to be inserted into a patient, said method comprising the step of contacting said surface with the composition of claim 18.
20. A pharmaceutically effective combination comprising a thrombin inhibitor of any one or claims 1 to 10, a thrombolytic agent, and a pharmaceutically acceptable carrier.
21. The pharmaceutically effective combination of claim 20, wherein said thrombin inhibitor is D-Cha-Hirulog-8 and said thrombolytic agent is tPA.
22. The combination of claim 20 or 21, wherein the daily dosage of said thrombin inhibitor is between about 0.5 nmoles/kg body weight and about 2.5 μ moles/kg body weight and wherein the daily dosage of said thrombolytic agent is between about 10% and about 80% of the conventional dosage range of said thrombolytic agent.
23. The combination of any one of claims 20 to 22, wherein the daily dosage of said thrombin inhibitor is between about 5 nmoles/kg body weight and about 250 nmoles/kg body weight and wherein the daily dosage of said thrombolytic agent is between about 10% and about 70% of the conventional dosage range of said thrombolytic agent.
24. Use of a thrombin inhibitor of any one or claims 1-10 for the preparation of a pharmaceutical composition for inhibiting a thrombin-mediated or thrombin-associated function or process in a patient or in extracorporeal blood, thrombus


accretion in a patient caused by clot-bound thrombin, or platelet-dependent thrombosis in a patient, for treating neurodegenerative disease in a patient, or for treating or preventing disseminated intravascular coagulation in a patient.

- 5 **25.** Use of a thrombin inhibitor of any one or claims 1-10 in combination with a thrombolytic agent for the preparation or a pharmaceutical combination for establishing reperfusion or preventing reocclusion in a patient.
- 10 **26.** The use of claim 25, wherein the amount of said thrombolytic agent in said composition is less than the amount required in an effective monotherapy.
- 15 **27.** Use of a thrombin inhibitor of any one of claims 1 to 10 for the preparation of a pharmaceutical composition for decreasing the reperfusion time and increasing the reocclusion time in a patient treated with a thrombolytic agent.
- 20 **28.** The use of a claim 27, wherein said pharmaceutical composition is administerable to said patient during the time period ranging from about 5 hours prior to about 5 hours following the treatment of said patient with said thrombolytic agent.
- 25 **29.** The use of claim 28, wherein said pharmaceutical composition is administerable to said patient during the time period ranging from about 2 hours prior to about 2 hours following said treatment of said patient with said thrombolytic agent.
- 30 **30.** Use of a thrombin inhibitor of any one of claims 1-10 for the preparation of a pharmaceutical composition for inhibiting the growth of a metastatic tumor in a patient.
- 35 **31.** The use of claim 30, wherein said metastatic tumor is selected from the group consisting of carcinoma of the brain, carcinoma of the lung, carcinoma of the liver, osteocarcinoma and neoplastic cell carcinoma.
- 40 **32.** Use of a thrombin inhibitor of any one of claims 1-10 for the preparation of a pharmaceutical composition for treating or preventing thrombin induced inflammation in a patient.
- 45 **33.** The use of claim 32, wherein said thrombin-induced inflammation is caused by a disease selected from the group consisting of adult respiratory distress syndrome, septic shock, septicemia, and reperfusion damage.
- 50 **34.** Use of a thrombin inhibitor of any one of claims 1-10 for the preparation of a pharmaceutical composition for treating or preventing a thrombotic disease in a patient.
- 55 **35.** The use of any one of claims 24 to 33, wherein the amount of thrombin inhibitor is between 0.5 nmoles/kg body weight/day to 2.5 μ moles/kg body weight/day.
- 60 **36.** The use of claim 34, wherein the amount of thrombin inhibitor is between 5 nmoles/kg body weight/day to 250 nmoles/kg body weight/day.
- 65 **37.** A method for producing the thrombin inhibitor of any one of claims 1 to 10, 14, or 15, comprising the steps of:
 - 70 (a) synthesizing the peptidic portion and the non-amino acid component of the thrombin inhibitor;
 - 75 (b) coupling said non-amino acid component to said peptidic portion; and, optionally
 - 80 (c) labelling the thrombin inhibitor with a radioisotope.
- 85 **38.** A method for producing the pharmaceutical composition of any one of claims 11 to 13, or the pharmaceutically effective combination of any one of claims 20 to 23, comprising adding to the thrombin inhibitor of any one of claims 1 to 10 a pharmaceutically acceptable carrier and, optionally, a thrombolytic agent.

Patentansprüche

1. Thrombininhibitor umfassend:

- (a) eine gegen das katalytische Zentrum gerichtete Einheit mit der Formel:



wobei X ein Wasserstoffatom ist oder durch ein Grundgerüst gekennzeichnet ist, bestehend aus 1 bis 100 Atomen; R₁ ausgewählt ist aus der Gruppe bestehend aus homocyclischen oder heterocyclischen gesättigten Ringstrukturen aus 6 Atomen, die unsubstituiert, mono-substituiert, di-substituiert und tri-substituiert sind; R₂ **gekennzeichnet ist durch** ein Grundgerüst, bestehend aus 1 bis 5 Atomen; R₃ eine Bindung ist oder **durch** ein Grundgerüst gekennzeichnet ist, bestehend aus 1 bis 3 Atomen; R₄ eine Aminosäure ist; R₅ eine L-Aminosäure ist, die eine Guanidinium- oder Amino-enhaltende Seitenkettengruppe umfaßt; R₆ eine nicht-Amid-Bindung ist; und Y **durch** ein Grundgerüst gekennzeichnet ist, bestehend aus 1 bis 9 Atomen; und

(b) eine Verknüpfungseinheit, die **durch** ein Grundgerüst gekennzeichnet ist, das eine errechnete Länge zwischen 18 Å und etwa 42 Å aufweist; und

(c) eine Einheit, die an die Anion-bindende Exo-Stelle ("anion binding exosite") bindet, wobei die gegen das katalytische Zentrum gerichtete Einheit an die Einheit, die an die Anion-bindende Exo-Stelle bindet, über die Verknüpfungseinheit gebunden ist, und der Inhibitor in der Lage ist, gleichzeitig an das katalytische Zentrum und die Anionbindende Exo-Stelle von Thrombin zu binden; und

wobei sich der Inhibitor **durch** eine verstärkte lipophile Interaktion mit Thrombin im Vergleich zu der Interaktion zwischen Thrombin und (D-Phe)-Pro-Arg-Pro-(Gly)₄-Asn-Gly-Asp-Phe-Glu-Glu-Ile-Pro-Glu-Glu-Tyr-Leu unterscheidet.

2. Thrombininhibitor nach Anspruch 1, wobei X H₂N ist, R₁ ausgewählt ist aus der Gruppe bestehend aus unsubstituiertem, mono-substituiertem, di-substituiertem und tri-substituiertem Hexan, R₂ CH₂-CH und R₃ C=O ist.
3. Thrombininhibitor nach Anspruch 1 oder 2, wobei die gegen das katalytische Zentrum gerichtete Einheit die Aminosäuresequenz:
D-Cha - Pro - Arg - Pro
aufweist.
4. Thrombininhibitor nach Anspruch 1 oder 2, wobei R₄ eine Aminosäure ist, die eine Seitenkettengruppe umfaßt, die **dadurch gekennzeichnet ist, daß** sie eine Wasserstoffbindung bei einem pH-Wert zwischen ungefähr 5.5 und 9.5 eingehen kann.
5. Thrombininhibitor nach Anspruch 1, 2 oder 4, wobei R₄ ausgewählt ist aus der Gruppe bestehend aus Histidin, Thioprolin und Isonipekotinsäure.
6. Thrombininhibitor nach einem der Ansprüche 1 bis 5, wobei die Einheit, die die Anion-bindende Exo-Stelle bindet, aus der Formel:
W - B₁ - B₂ - B₃ - B₄ - B₅ - B₆ - B₇ - B₈ - Z
besteht, wobei W eine Bindung, B₁ eine anionische Aminosäure, B₂ eine Aminosäure, B₃ Ile, Val, Leu, Nle oder Phe, B₄ Pro, Hyp, 3,4-DehydroPro, Thiazolidin-4-carboxylat, Sar, eine N-Methyl-Aminosäure oder D-Ala, B₅ eine anionische Aminosäure, B₆ eine anionische Aminosäure, B₇ eine lipophile Aminosäure, ausgewählt aus der Gruppe bestehend aus Tyr, Trp, Phe, Leu, Nle, Ile, Val, Cha, Pro, oder ein Dipeptid, das aus einem dieser lipophilen Aminosäuren und einer Aminosäure besteht, B₈ eine Bindung oder ein Peptid, das 1 bis 5 Aminosäurereste enthält, und Z OH ist oder durch ein Grundgerüst gekennzeichnet ist, bestehend aus 1 bis 6 Atomen.
7. Thrombininhibitor nach Anspruch 6, wobei B₁ Glu, B₂ Glu, B₃ Ile, B₄ Pro, B₅ Glu, B₆ Glu, B₇ Tyr-Leu, Tyr(SO₃H)-Leu, Tyr(OSO₃H)-Leu oder (3-,5-diiodTyr)-Leu, B₈ eine Bindung, und Z OH ist.
8. Thrombininhibitor nach einem der Ansprüche 1 bis 7, wobei das Grundgerüst der Verknüpfungseinheit aus einer Kombination von Atomen besteht, die ausgewählt sind aus der Gruppe bestehend aus Kohlenstoff, Stickstoff, Schwefel und Sauerstoff.
9. Thrombininhibitor nach Anspruch 8, wobei die Verknüpfungseinheit die Aminosäuresequenz:

Gly - Gly - Gly - Asn - Gly - Asp - Phe
umfaßt.

- 5 10. Thrombininhibitor nach Anspruch 9, wobei der Thrombininhibitor D-Cha-Hirulog-8 ist.
11. Arzneimittel, umfassend eine pharmazeutisch wirksame Menge eines Thrombininhibitors nach einem der Ansprüche 1 bis 10 und gegebenenfalls einen pharmazeutisch verträglichen Träger.
- 10 12. Pharmazeutisch verträgliche Zusammensetzung nach Anspruch 11, wobei die pharmazeutisch wirksame Menge bei etwa 0,5 nMol/kg Körpergewicht/Tag bis etwa 2,5 µMol/kg Körpergewicht/Tag liegt.
13. Pharmazeutisch verträgliche Zusammensetzung nach Anspruch 12, wobei die pharmazeutisch wirksame Menge bei etwa 5 nMol/kg Körpergewicht/Tag bis etwa 250 nMol/kg Körpergewicht/Tag liegt.
- 15 14. Thrombininhibitor nach einem die Ansprüche 1 bis 10, zusätzlich ein Radioisotop umfassend.
15. Thrombininhibitor nach Anspruch 14, wobei das Radioisotop ausgewählt ist aus der Gruppe bestehend aus ¹²³I, ¹²⁵I und ¹¹¹In.
- 20 16. Zusammensetzung zur ex vivo-Abbildung eines Fibrin- oder Blutplättchen-Thrombus' in einem Patienten, umfassend einen pharmazeutisch verträglichen Puffer und einen Thrombininhibitor nach Anspruch 14 oder 15.
17. Verfahren zur ex-vivo-Abbildung eines Fibrin- oder Blutplättchen- Thrombus' in einem Patienten, umfassend die Schritte:
 - 25 (a) Verabreichung der Zusammensetzung nach Anspruch 16 an den Patienten; und
 - (b) Verwendung eines Nachweismittels zur Beobachtung des Thrombininhibitors, der in der Zusammensetzung vorhanden ist.
- 30 18. Zusammensetzung zur Beschichtung der Oberfläche einer invasiven Vorrichtung zur Einführung in einen Patienten, wobei die Zusammensetzung einen geeigneten Puffer und mindestens einen Thrombininhibitor nach einem der Ansprüche 1 bis 10 umfaßt.
- 35 19. Verfahren zur Beschichtung der Oberfläche einer invasiven Vorrichtung zur Einführung in einen Patienten, umfassend den Schritt der Beschichtung der Oberfläche mit der Zusammensetzung nach Anspruch 18.
20. Pharmazeutisch wirksame Kombination, umfassend einen Thrombininhibitor nach einem der Ansprüche 1 bis 10, ein thrombolytisches Agens und einen pharmazeutisch verträglichen Träger.
- 40 21. Pharmazeutisch wirksame Kombination nach Anspruch 20, wobei der Thrombininhibitor D-Cha-Hirulog-8 und das thrombolytische Agens tPA ist.
22. Kombination nach Anspruch 20 oder 21, wobei die tägliche Dosis des Thrombininhibitors zwischen etwa 0,5 nMol/kg Körpergewicht und etwa 2,5 µMol/kg Körpergewicht liegt, und wobei die tägliche Dosis des thrombolytischen Agens zwischen etwa 10% und etwa 80% des konventionellen Dosierungsbereiches des thrombolytischen Agens liegt.
- 45 23. Kombination nach einem der Ansprüche 20 bis 22, wobei die tägliche Dosis des Thrombininhibitors zwischen etwa 5 nMol/kg Körpergewicht und etwa 250 nMol/kg Körpergewicht liegt, und wobei die tägliche Dosis des thrombolytischen Agens zwischen etwa 10% und etwa 70% des konventionellen Dosierungsbereiches des thrombolytischen Agens liegt.
- 50 24. Verwendung eines Thrombininhibitors nach einem der Ansprüche 1 bis 10 zur Herstellung eines Arzneimittels zur Hemmung einer/eines Thrombinvermittelten oder Thrombin-assoziierten Funktion oder Prozesses in einem Patienten oder in extrakorporalem Blut, einer Thrombus-Zunahme in einem Patienten verursacht durch an Blutgerinnsel gebundenes Thrombin, oder einer Blutplättchen-abhängigen Thrombose in einem Patienten, zur Behandlung einer neurodegenerativen Erkrankung in einen Patienten, oder zur Behandlung oder Vorbeugung disseminierter intravasaler Koagulation in einem Patienten.
- 55

25. Verwendung eines Thrombininhibitors nach einem der Ansprüche 1 bis 10 in Kombination mit einem thrombolytischen Agens zur Herstellung eines Arzneimittels zur Wiederherstellung der Durchblutung oder zur Vorbeugung einer Reokklusion in einem Patienten.

26. Verwendung nach Anspruch 25, wobei die Menge des thrombolytischen Agens in der Zusammensetzung weniger ist als die Menge, die zu einer wirksamen Monotherapie benötigt wird.

27. Verwendung eines Thrombininhibitors nach einem der Ansprüche 1 bis 10 zur Herstellung eines Arzneimittels zur Verringerung der Reperusionszeit und zur Erhöhung der Reokklusionszeit in einem Patienten, der mit einem thrombolytischen Agens behandelt wurde.

28. Verwendung nach Anspruch 27, wobei das Arzneimittel dem Patienten innerhalb der Zeitspanne zwischen etwa 5 Stunden vor und etwa 5 Stunden nach der Behandlung des Patienten mit dem thrombolytischen Agens verabreichbar ist.

29. Verwendung nach Anspruch 28, wobei das Arzneimittel dem Patienten innerhalb der Zeitspanne zwischen etwa 2 Stunden vor und etwa 2 Stunden nach der Behandlung des Patienten mit dem thrombolytischen Agens verabreichbar ist.

30. Verwendung eines Thrombininhibitors nach einem der Ansprüche 1 bis 10 zur Herstellung eines Arzneimittels zur Hemmung des Wachstums eines metastatischen Tumors in einem Patienten.

31. Verwendung nach Anspruch 30, wobei der metastatische Tumor ausgewählt ist aus der Gruppe bestehend aus Hirncarcinom, Lungencarcinom, Lebercarcinom, Osteocarcinom und neoplastischem Zeltcarcinom.

32. Verwendung eines Thrombininhibitors nach einem der Ansprüche 1 bis 10 zur Herstellung eines Arzneimittels zur Behandlung oder Vorbeugung einer Thrombin-induzierten Entzündung in einem Patienten.

33. Verwendung nach Anspruch 32, wobei die Thrombin-induzierte Entzündung durch eine Erkrankung verursacht ist, die ausgewählt ist aus der Gruppe bestehend aus akuter respiratorischer Insuffizienz, septischem Schock, Septikämie, und Reperusionsverletzung.

34. Verwendung eines Thrombininhibitors nach einem der Ansprüche 1 bis 10 zur Herstellung eines Arzneimittels zur Behandlung oder Vorbeugung einer thrombotischen Erkrankung in einem Patienten.

35. Verwendung nach einem der Ansprüche 24 bis 33, wobei die Menge des Thrombininhibitors zwischen 0,5 nMol/kg Körpergewicht/Tag und 2,5 µMol/kg Körpergewicht/Tag liegt.

36. Verwendung nach Anspruch 34, wobei die Menge des Thrombininhibitors zwischen 5 nMol/kg Körpergewicht/Tag und 250 nMol/kg Körpergewicht/Tag liegt.

37. Verfahren zur Herstellung des Thrombininhibitors nach einem der Ansprüche 1 bis 10, 14 oder 15, umfassend die Schritte:


- (a) Synthese des Peptidanteils und der nicht-Aminosäure-Komponente des Thrombininhibitors;
- (b) Verknüpfung der nicht-Aminosäure-Komponente mit dem Peptidanteil; und gegebenenfalls
- (c) Markierung des Thrombininhibitors mit einem Radioisotop.

38. Verfahren zur Herstellung des Arzneimittels nach einem der Ansprüche 11 bis 13, oder der pharmazeutisch wirksamen Kombination nach einem der Ansprüche 20 bis 23, umfassend die Zugabe eines pharmazeutisch verträglichen Trägers und gegebenenfalls eines thrombolytischen Agens zu dem Thrombininhibitor nach einem der Ansprüche 1 bis 10.

Revendications

1. Inhibiteur de la thrombine comprenant:

a) une entité dirigée vers le site catalytique comprenant la formule:



où X est hydrogène ou est **caractérisé par** une chaîne d'épine dorsale consistant en 1 à 100 atomes; R₁ est sélectionné dans le groupe consistant en structures cycliques saturées homocycliques ou hétérocycliques de 6 atomes qui sont non substitués, mono-substitués, di-substitués et tri-substitués; R₂ est **caractérisé par** une chaîne d'épine dorsale consistant en 1 à 5 atomes; R₃ est une liaison ou est **caractérisé par** une chaîne d'épine dorsale consistant en 1 à 3 atomes; R₄ est tout acide aminé; R₅ est tout acide L-aminé qui comprend un groupe de chaîne latérale contenant un guanidinium ou un amino; R₆ est une liaison non amide; et Y est **caractérisé par** une chaîne d'épine dorsale consistant en 1 à 9 atomes; et

b) une entité d'enchaînement **caractérisée par** une chaîne d'épine dorsale ayant une longueur calculée entre 18 Å et environ 42 Å; et

c) une entité associant un exosite liant un anion, où ladite entité dirigée vers le site catalytique est liée à ladite entité associant un exosite liant un anion via ladite entité d'enchaînement et ledit inhibiteur est capable de se lier simultanément au site catalytique et à l'exosite liant un anion ce la thrombine; et où ledit inhibiteur se distingue par une interaction lipophile accrue avec la thrombine en comparaison avec l'interaction entre la thrombine et (D-Phe)-Pro-Arg-Pro-(Gly)4-Asn-Gly-Asp-Phe-Glu-Glu-Ile-Pro-Glu-Glu-Tyr-Leu.


2. Inhibiteur de la thrombine de la revendication 1, où X est H₂N, R₁ est sélectionné dans le groupe consistant en hexane non substitué, mono-substitué, di-substitué et tri-substitué, R₂ est CH₂-CH, et R₃ est C=O.

3. Inhibiteur de la thrombine de la revendication 1 ou 2, où ladite entité dirigée vers le site catalytique a la séquence d'acides aminés:
D-Cha-Pro-Arg-Pro

4. Inhibiteur de la thrombine de la revendication 1 ou 2, où R₄ est tout acide aminé comprenant un groupe de chaîne latérale **caractérisé par** la capacité d'accepter une liaison hydrogène à un pH entre environ 5,5 et 9,5.

5. Inhibiteur de la thrombine de la revendication 1, 2 ou 4, où R₄ est sélectionné dans le groupe consistant en histidine, thioproline et acide isonipécotique.

6. Inhibiteur de la thrombine selon l'une quelconque des revendications 1 à 5, où ladite entité associant un exosite de liant un anion consiste en la formule :



où W est une liaison, B₁ est un acide aminé anionique, B₂ est tout acide aminé, B₃ est Ile, Val, Leu, Nle ou Phe, B₄ est Pro, Hyp, 3,4-déshydroPro, thiazolidine-4-carboxylate, Sar, tout acide aminé N-méthylé ou D-Ala, B₅ est un acide aminé anionique, B₆ est un acide aminé anionique, B₇ est un acide aminé lipophile sélectionné dans le groupe consistant en Tyr, Trp, Phe, Leu, Nle, Ile, Val, Cha, Pro, ou un dipeptide consistant en l'un de ces acides aminés lipophiles et tout acide aminé, B₈ est une liaison ou un peptide contenant de un à cinq résidus de tout acide aminé, et Z est OH ou est **caractérisé par** une chaîne d'épine dorsale consistant an 1 à 6 atomes.

7. Inhibiteur de la thrombine de la revendication 6, où B₁ est Glu, B₂ est Glu, B₃ est Ile, B₄ est Pro, B₅ est Glu, B₆ est Glu, B₇ est Tyr-Leu, Tyr(SO₃H)-Leu, Tyr(OSO₃H)-Leu ou (3-, 5-diiodoTyr)-Leu, B₈ est une liaison, et Z est OH.

8. Inhibiteur de la thrombine selon l'une quelconque des revendications 1 à 7, où ladite chaîne d'épine dorsale de ladite entité d'enchaînement consiste en toute combinaison d'atomes sélectionnés dans le groupe consistant en carbone, nitrogen sulfur et oxygène.

9. Inhibiteur de la thrombine de la revendication 8, où ledit enchaînement comprend la séquence d'acides aminés:
Gly-Gly-Gly-Asn-Gly-Asp-Phe.

10. Inhibiteur de la thrombine de la revendication 9, où ledit inhibiteur de la thrombine est D-Cha-Hirulog-8.

11. Composition pharmaceutique comprenant une quantité pharmaceutiquement efficace d'un inhibiteur de la thrombine selon l'une quelconque des revendications 1 à 10 et facultativement un véhicule pharmaceutiquement acceptable.
- 5 12. Composition pharmaceutiquement acceptable de la revendication 11, où ladite quantité pharmaceutiquement efficace est comprise entre environ 0,5 nmoles/kg de poids corporel/jour et environ 2,5 μ moles/kg poids corporel/jour.
13. Composition pharmaceutiquement acceptable de la revendication 12, où ladite quantité pharmaceutiquement efficace est entre environ 5 nmoles/kg poids corporel/jour et environ 250 nmoles/kg poids corporel/jour.
- 10 14. Inhibiteur de la thrombine selon l'une quelconque des revendications 1 à 10, comprenant de plus un radioisotope.
- 15 15. Inhibiteur de la thrombine de la revendication 14, où ledit radioisotope est sélectionné dans le groupe consistant en ^{123}I , ^{125}I et ^{111}In .
16. Composition pour imagerie ex vivo d'un thrombus de fibrine ou de plaquettes dans un patient, ladite composition comprenant un buffer pharmaceutiquement acceptable et un inhibiteur de la thrombine de la revendication 14 ou 15.
17. Méthode pour imagerie ex vivo d'un thrombus de fibrine ou de plaquettes dans un patient, comprenant les étapes de :
20 (a) administrer audit patient la composition de la revendication 16; et
(b) utiliser un moyen de détection pour observer l'inhibiteur de la thrombine présent dans ladite composition.
18. Composition pour revêtir la surface d'un dispositif invasif à insérer dans un patient, où ladite composition comprend un tampon approprié et au moins un inhibiteur de la thrombine de l'une des revendications 1 à 10.
- 25 19. Méthode pour revêtir la surface d'un dispositif invasif à insérer dans un patient, ladite méthode comprenant l'étape de mettre ladite surface en contact avec la composition de la revendication 18.
- 30 20. Combinaison pharmaceutiquement efficace, comprenant un inhibiteur de la thrombine selon l'une des revendications 1 à 10, un agent thrombolytique, et un véhicule pharmaceutiquement acceptable.
21. Combinaison pharmaceutiquement efficace de la revendication 20, où ledit inhibiteur de la thrombine est D-Cha-Hirulog-8 et ledit agent thrombolytique est tPA.
- 35 22. Combinaison de la revendication 20 ou 21, où le dosage quotidien dudit inhibiteur de la thrombine est entre environ 0,5 nmoles/kg de poids corporel et environ 2,5 μ moles/kg de poids corporel et où le dosage quotidien dudit agent thrombolytique est compris entre environ 10% et environ 80% de la gamme de dosage conventionnel dudit agent thrombolytique.
- 40 23. Combinaison selon l'une quelconque des revendications 20 à 22, où le dosage quotidien dudit inhibiteur de la thrombine est entre environ 5 nmoles/kg de poids corporel et environ 250 nmoles/kg de poids corporel et où le dosage quotidien dudit agent thrombolytique est compris entre environ 10% et environ 70% de la gamme de dosage conventionnel dudit agent thrombolytique.
- 45 24. Utilisation d'un inhibiteur de la thrombine selon l'une quelconque des revendications 1-10 pour la préparation d'une composition pharmaceutique pour inhiber une fonction ou un processus dû à la thrombine ou associé à la thrombine dans un patient ou dans le sang extracorporel, l'accrétion d'un thrombus chez un patient provoquée par de la thrombine liée à un caillot ou une thrombose dépendant des plaquettes dans un patient, pour le traitement d'une maladie neurodégénérative dans un patient ou pour le traitement ou la prévention d'une coagulation intravasculaire disséminée dans un patient.
- 50 25. Utilisation d'un inhibiteur de la thrombine selon l'une des revendications 1-10 en combinaison avec un agent thrombolytique pour la préparation d'une combinaison pharmaceutique pour établir la reperfusion ou prévenir la réocclusion dans un patient.
- 55 26. Utilisation de la revendication 25, où la quantité dudit agent thrombolytique dans ladite composition est inférieure à la quantité requise dans une monothérapie efficace.

27. Utilisation d'un inhibiteur de la thrombine selon l'une quelconque des revendications 1 à 10 pour la préparation d'une composition pharmaceutique pour diminuer le temps de reperfusion et augmenter le temps de réocclusion dans un patient traité avec un agent thrombolytique.
- 5 28. Utilisation de la revendication 27, où ladite composition pharmaceutique est administrable audit patient pendant la période de temps allant d'environ 5 heures avant jusqu'à environ 5 heures après le traitement dudit patient avec ledit agent thrombolytique.
- 10 29. Utilisation de la revendication 28, où ladite composition pharmaceutique est administrable audit patient pendant la période de temps allant d'environ 2 heures avant jusqu'à environ 2 heures après le traitement dudit patient par ledit agent thrombolytique.
- 15 30. Utilisation d'un inhibiteur de la thrombine selon l'une des revendications 1-10 pour la préparation d'une composition pharmaceutique pour l'inhibition de la croissance d'une tumeur métastatique dans un patient.
31. Utilisation de la revendication 30, où ladite tumeur métastatique est sélectionnée dans le groupe consistant en carcinome du cerveau, carcinome du poumon, carcinome du foie, ostéocarcinome et carcinome des cellules néoplastiques.
- 20 32. Utilisation d'un inhibiteur de la thrombine selon l'une des revendications 1-10 pour la préparation d'une composition pharmaceutique pour le traitement ou la prévention d'une inflammation induite par la thrombine dans un patient.
- 25 33. Utilisation de la revendication 32, où ladite inflammation induite par la thrombine est provoquée par une maladie sélectionnée dans le groupe consistant en syndrome du trouble respiratoire dans l'adulte, choc septique, septicémie, et dégradation par reperfusion.
34. Utilisation d'un inhibiteur de la thrombine de l'une des revendications 1-10 pour la préparation d'une composition pharmaceutique pour le traitement ou la prévention d'une maladie thrombotique dans un patient.
- 30 35. Utilisation selon l'une quelconque des revendications 24 à 33, où la quantité de l'inhibiteur de la thrombine est entre 0,5 nmole/kg de poids corporel/jour et environ 2,5 μ moles/kg/poids corporel/jour.
36. Utilisation de la revendication 34, où la quantité de l'inhibiteur de la thrombine est entre 5 nmoles/kg de poids corporel/jour et 250 nmoles/kg de poids corporel/jour.
- 35 37. Méthode pour produire un inhibiteur de la thrombine selon l'une quelconque des revendications 1-10, 14, ou 15, comprenant les étapes de:
- 40 a) synthétiser la partie peptidique et la partie non-acide aminé de l'inhibiteur de la thrombine;
b) coupler ladite partie non-acidé amine audite partie peptidique; et facultativement
c) marquer l'inhibiteur de la thrombine avec un radioisotope.
- 45 38. Méthode pour produire la composition pharmaceutique selon l'une quelconque des revendications 11-13, ou la combinaison pharmaceutiquement efficace selon l'une quelconque des revendications 20-23, comprenant ajouter a l'inhibiteur de la thrombine selon l'une quelconque des revendications 1-10 un véhicule pharmaceutiquement acceptable et facultativement un agent thrombolytique.




FIG. 1




FIG. 2




FIG. 3

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- EP 158564 A [0011]
- EP 168342 A [0011]
- EP 171024 A [0011]
- EP 200655 A [0011]
- EP 276014 A [0012]
- EP 291982 A [0012]
- EP 333356 A [0012]
- EP 341607 A [0012]
- EP 372670 A [0012]
- US 395482 A [0013] [0031] [0039]
- US 549388 A [0013] [0039]
- US 314756 A [0043] [0044]

Non-patent literature cited in the description

- **S. R. HANSON ; L. A. HARKER.** Interruption of Acute Platelet-Dependent Thrombosis by the Synthetic Antithrombin D-Phenylalanyl-L-Prolyl-L-Arginylichloromethylketone. *Proc. Natl. Acad. Sci. USA*, 1988, vol. 85, 3184-88 [0003]
- **J. W. FENTON, II.** Thrombin Bioregulatory Functions. *Adv. Clin. Enzymol.*, 1988, vol. 6, 186-93 [0004]
- **S. PRESCOTT et al.** Human Endothelial Cells in Culture Produce Platelet-Activating Factor (1-alkyl-2-acetyl-sn-glycero-3-phosphocholine) When Stimulated With Thrombin. *Proc. Natl. Acad. Sci. USA*, 1984, vol. 81, 3534-38 [0004]
- **G. M. VERCOLLETTI et al.** Platelet-Activating Factor Primes Neutrophil Responses to Agonists: Role in Promoting Neutrophil-Mediated Endothelial Damage. *Blood*, 1988, vol. 71, 1100-07 [0004]
- **P. J. DEL VECCHIO et al.** Endothelial Monolayer Permeability To Macromolecules. *Fed. Proc.*, 1987, vol. 46, 2511-15 [0004]
- **C. L. KNUPP.** Effect of Thrombin Inhibitors on Thrombin-Induced Release and Aggregation. *Thrombosis Res.*, 1988, vol. 49, 23-36 [0004]
- **A. FALANGA et al.** Isolation and Characterization of Cancer Procoagulant: A Cysteine Proteinase from Malignant Tissue. *Biochemistry*, 1985, vol. 24, 5558-67 [0005]
- **S. G. GORDON et al.** Cysteine Proteinase Procoagulant From Amnion-Chorion. *Blood*, 1985, vol. 66, 1261-65 [0005]
- **A. FALANGA et al.** A New Procoagulant In Acute Leukemia. *Blood*, 1988, vol. 71, 870-75 [0005]
- **D. GURWITZ et al.** Thrombin Modulates and Reverses Neuroblastoma Neurite Outgrowth. *Proc. Natl. Acad. Sci. USA*, 1988, vol. 85, 3440-44 [0005]
- **S. R. STONE ; J. HOFSTEENGE.** Kinetics of the Inhibition of Thrombin by Hirudin. *Biochemistry*, 1986, vol. 25, 4622-28 [0007]
- **J. W. FENTON, II.** Regulation of Thrombin Generation and Functions. *Semin. Thromb. Hemost.*, 1988, vol. 14, 234-40 [0007]
- **J. W. FENTON, II et al.** Thrombin Anion Binding Exosite Interactions with Heparin and Various Polyanions. *Ann. New York Acad. Sci.*, 1989, vol. 556, 158-65 [0008]
- **S. KONO et al.** Analysis of Secondary Structure of Hirudin and the Conformational Change Upon Interaction with Thrombin. *Arch. Biochem. Biophys.*, 1988, vol. 267, 158-66 [0008]
- **P. WALSMANN ; F. MARKWARDT.** Biochemical and Pharmacological Aspects of the Thrombin Inhibitor Hirudin. *Pharmazie*, 1981, vol. 36, 653-60 [0009]
- **J. DODT et al.** The Complete Covalent Structure of Hirudin: Localization of the Disulfide Bonds. *Biol. Chem. Hoppe-Sevler*, 1985, vol. 366, 379-85 [0009]
- **S. J. T. MAO et al.** Rapid Purification and Revised Amino Terminal Sequence of Hirudin: A Specific Thrombin Inhibitor of the Blood-Sucking Leech. *Anal. Biochem*, 1987, vol. 161, 514-18 [0009]
- **R. P. HARVEY et al.** Cloning and Expression of a cDNA Coding for the Anti-Coagulant Hirudin from the Bloodsucking Leech, *Hirudo medicinalis*. *Proc. Natl. Acad. Sci. USA*, 1986, vol. 83, 1084-88 [0009]
- **F. MARKWARDT et al.** Pharmacological Studies on the Antithrombotic Action of Hirudin in Experimental Animals. *Thromb. Haemost.*, 1982, vol. 47, 226-29 [0010]
- CRC Handbook of Chemistry and Physics. CRC Press, Inc, 1984, F-166-70 [0030]
- **M. SZELKE.** Potent New Inhibitors of Human Renin. *Nature*, 1982, vol. 299, 555-57 [0034]
- Facile Solid Phase Preparation of Proteins containing the CH₂-NH Peptide Bond Isostere and Application to the Synthesis of Somatostatin (SRIF) Octapeptide Analogues. **D. H. COY et al.** Peptides. Walter Gruyter & Co, 1987, vol. 1986, 143-46 [0034]

- **R. T. MORRISON ; R. N. BOYD.** Organic Chemistry. Allyn and Bacon, Inc, 1977 [0041]
- **J. M. MARAGANORE et al.** Anticoagulant Activity of Synthetic Hirudin Peptides. *J. Biol. Chem.*, 1989, vol. 264, 8692-98 [0043]
- **J.L. KRSTENANSKY et al.** Development of MDL-28,050, A Small Stable Antithrombin Agent Based On A Functional Domain of the Leech Protein, Hirudin. *Thromb. Haemostas.*, 1990, vol. 63, 208-14 [0045]
- Physician's Desk Reference. Edward R. Barnhart, 1989 [0065]
- **F. B. TAYLOR et al.** Protein C Prevents the Coagulopathic and Lethal Effects of Escherichia coli Infusion in the Baboon. *J.Clin.Invest.*, 1987, vol. 79, 918-25 [0072]
- **J. M. MARAGANORE et al.** *J. Biol. Chem.*, 1989, vol. 264, 8692-98 [0078]
- **J. W. FENTON, II et al.** Hirudin Inhibition by Thrombin. *Angio. Archiv. Biol.*, 1989, vol. 18, 27 [0078]
- **S. R. HANSON ; L. A. HARKER.** Interruption of Acute Platelet-Dependent Thrombosis by the Synthetic Antithrombin D-Phenylalanyl-L-Prolyl-L-Arginyl Chloromethyl Ketone. *Proc. Natl. Acad. Sci. USA*, 1988, vol. 85, 3184-88 [0079]
- **J. F. EIDT et al.** Thrombin is an Important Mediator of Platelet Aggregation in Stenosed Canine Coronary Arteries with Endothelial Injury. *J. Clin. Invest.*, 1989, vol. 84, 18-27 [0079]
- **B. FURIE et al.** Computer-Generated Models of Blood Coagulation Factor Xa, Factor IXa, and Thrombin Based Upon Structural Homology with Other Serine Proteases. *J. Biol. Chem.*, 1982, vol. 257, 3875-82 [0080]
- **W. BODE et al.** The Refined 1.9 Å Crystal Structure of Human α -Thrombin: Interaction With D-Phe-Pro-Arg-Chloromethylketone and Significance of the Tyr-Pro-Pro-Trp Insertion Segment. *EMBO J.*, 1989, vol. 8, 3467-75 [0086]
- **T. J. RYDEL et al.** The Structure of a Complex of Recombinant Hirudin and Human α -Thrombin. *Science*, 1990, vol. 249, 277-80 [0086]