

19



Europäisches Patentamt
European Patent Office
Office européen des brevets



11 Publication number:

0 555 798 A1

12

EUROPEAN PATENT APPLICATION

21 Application number: **93101923.6**

51 Int. Cl.⁵: **C07H 1/08, C12N 15/10**

22 Date of filing: **08.02.93**

30 Priority: **13.02.92 US 835179**

43 Date of publication of application:
18.08.93 Bulletin 93/33

84 Designated Contracting States:
BE DE ES FR GB IT SE

71 Applicant: **Becton Dickinson and Company**
One Becton Drive
Franklin Lakes New Jersey 07417-1880(US)


72 Inventor: **Woodard, Daniel Lee**
1800-203, Avent Ridge Road

Raleigh, North Carolina 27606(US)
Inventor: **Howard, Adriann Jeanelle**
1321 Nez Castle Road, Apartment A-4
Durham, North Carolina 27704(US)
Inventor: **Down, James Arthur**
101 Charter Oaks Circle
Cary, North Carolina 27511(US)


74 Representative: **Rambelli, Paolo et al**
Jacobacci-Casetta & Perani S.p.A. Via Alfieri
17
I-10121 Torino (IT)

54 **Hydrated celite and purification of DNA.**


57 The invention discloses hydrated celite, how to make hydrated celite, and processes for purifying DNA with hydrated celite. The new composition is of the formula:



wherein R' is OH, R is SiO₂, and R'' is OH, and repeating units of the composition comprising the formula:




wherein R' is independently OH or



EP 0 555 798 A1

R is SiO₂, and R'' is independently OH or



FIELD OF THE INVENTION

The invention is in the field of molecular biology. In particular, the invention is in the area of deoxyribonucleic acid purification.

5

BACKGROUND OF THE INVENTION

The continued advances in molecular biology and related disciplines present continued needs for improvements in tools associated with fully appreciating and developing the advanced technology.

10

A wide range of technologies involve the use of deoxyribonucleic acids (DNA) in a variety of forms. For example, advances in the area of recombinant DNA technology continually require the use of DNA in the form of probes, genomic DNA, and plasmid DNA.

15

Advances in the area of diagnostics also continue to utilize DNA in a variety of ways. For example, DNA probes are routinely used in the detection and diagnosis of human pathogens. Likewise, DNA is used in the detection of genetic disorders. DNA is also used in the detection of food contaminants. And, DNA probes are routinely used in locating, identifying and isolating target DNA of interest for a variety of reasons ranging from genetic mapping to cloning and recombinant expression.

20

In many instances DNA is available in extremely small amounts, and isolation and purification procedures can be laborious and time consuming. The often time consuming and laborious procedures can lead to loss of DNA. In the purification of DNA from specimens obtained from serum, urine, and bacterial cultures, there is the added risk of contamination and false-positive results.

Typical DNA purification protocols involve the use of caustic and poisonous compositions. The typical DNA purification protocol uses high concentrations of chaotropic salts such as sodium iodine and sodium perchlorate.

25

There are numerous protocols for purifying DNA. As evidenced by recent activity in the area of DNA purification, there is a continued pursuit for optimal DNA purification protocols. U. S. Patent 4,923,978 discloses a process for purifying DNA in which a solution of protein and DNA is passed over a hydroxylated support and the protein is bound and the DNA is eluted. U.S. Patent 4,935,342 discloses purification of DNA by selective binding of DNA to anion exchangers and subsequent elution. U. S. Patent 4,946,952 discloses DNA isolation by precipitation with water-soluble ketones. A DNA purification procedure using chaotropes and dialyzed DNA is disclosed in in. S. Patent 4,900,677.

30


While the present protocols for purifying DNA are able to accomplish their goal, it is desirable to purify DNA without the use of such caustic and poisonous compounds such as the most often used chaotropes in addition to obtaining increased amounts of DNA.

35

SUMMARY OF THE INVENTION

The invention provides the monomer unit composition:


40



45


wherein R' is OH, R is SiO₂, and R'' is OH and repeating units of the composition comprising the formula:

50



55


wherein R' is independently OH or



5

R is SiO₂, and R'' is independently OH or

10



15

and compositions comprising the above monomer and repeating units thereof. Repeating units described above can include from about 2 up to infinity. Ranges include about 2 to about 100,000,000, and about 2 to about 100,000.

20


The invention can be used to purify DNA from a variety of sources and from a variety of forms. The process uses the composition of the invention and renders the use of binding buffers, such as chaotropes, optional. The DNA can be bound in aqueous solution such as TE buffer (10mM Tris, 1mM EDTA) at room temperature. In addition, the DNA can be eluted into water from the compositions of the invention by heating, or generally used elution buffers such as TE or 1 X TAE can be employed. Sources of DNA for purification include bacteria, bacteriophage, specimens, plants, animals, and the like. DNA can be found in a variety of forms and includes single-stranded, double-stranded, circular, and linear. The invention can be practiced with DNA from any source in any form.

25

DETAILED DESCRIPTION


30

The invention provides the monomer unit composition:



35


wherein R' is OH, R is SiO₂, and R'' is OH and repeating units of the composition comprising the formula:



45


wherein R' is independently OH or

50



55

R is SiO₂, and R'' is independently OH or



5


and compositions comprising the above monomer and repeating units thereof. Repeating units described above can include from about 2 up to infinity. Ranges include about 2 to 100,000,000, and about 2 to 100,000.

10

The surface provides for bonding of DNA while also allowing easy recovery of DNA from the surface.

Also provided is a process for purifying DNA which comprises contacting DNA with a composition of the formula:


15



20

wherein R' is OH, R is SiO₂, and R'' is OH or repeating units of the composition comprising the formula:


25



30

wherein R' is independently OH or


35



40

R is SiO₂, and R'' is independently OH or

45




Reaction products of hydrating reagents and SiO₂ are also provided.

50

The invention also provides a method for making the composition of the formula:


55



wherein R' is OH, R is SiO₂, and R'' is



10 and repeating units of the composition comprising the formula:



20 wherein R' is independently OH or



30 R is SiO₂, and R'' is independently OH or



Repeating units described above can include from about 2 up to infinity. Ranges include about 2 to 100,000,000, and about 2 to 100,000.

40 Generally, reaction products of hydrating reagents and SiO₂ result in a bead like structure comprising repeating units of the above referenced monomer unit.

45 It is possible that the electronic nature of this polymer is such that surface modifications can be made that are of a more conventional nature but are changed electronically due to the presence of this polymer being at the center of the bead (making it a more efficient surface for the purposes described in this disclosure). For example, the surface could be modified with SiCl₄ followed by hydration which would result in a silanol coating on the surface. The exposure of the repeating unit is what interacts with the DNA, and thus surfaces comprising the repeating unit are also suitable for practicing the invention. Surfaces which can be designed to comprise compositions of the invention include dipstick configurations, tubes, vials, filtration devices, and the like.

50 The procedure for obtaining the compositions of the invention generally comprises adding a hydrating reagent like NaOH(aq) to SiO₂ and refluxing.

The invention also provides a process for purifying DNA which comprises contacting DNA with compositions of the invention.

55 The process for producing the compositions of the invention and the reaction products of NaOH and SiO₂ comprises the addition of NaOH to SiO₂. Any counter ion, not just Na⁺, can be used in any concentration and in any ratio compared to SiO₂. For example, potassium hydroxide can also be used. The ratio of NaOH to SiO₂ is about .1:1 - 10:1, preferably about 2:1. The resultant product is filtered then washed and dried. Suitable washing reagents include acetone and the like. The product is now ready for use in purifying DNA.

The start of any DNA purification or isolation procedure requires obtaining the desired DNA from its source. Typical protocols for obtaining DNA from specimens such as serum, urine and bacterial cultures are well known and routinely carried out. Likewise, the ability to obtain DNA from genomic libraries and the like are routine. The key to the invention is the ability to purify DNA, once obtained from its source. Typical procedures for obtaining DNA end with a suspension of the DNA in solution. References include those for isolation of DNA from biological samples, Harding, J.D., Gebeyehu, G., Bebee, R., Simms, D., Ktevan, L., *Nucleic Acids Research*, **17**:6947 (1989), and Marko, M.A., Chipperfield, A., and Birnboim, H.C., *Analytical Biochemistry*, **121**:382 (1982). Procedures for isolation of plasmid DNA can be found in Lutze, L.H., Winegar, R.A., *Nucleic Acids Research* **20**:6150 (1990). Extraction of double-stranded DNA from biological samples can be found in Yamada, O., Matsumoto, T., Nakashima, M., Hagri, S., Kamahora, T., Ueyama, H., Kishi, Y., Uemura H., Kurimura, T., *Journal of Virological Methods* **27**:203 (1990). Most DNA solutions comprise the DNA in a suitable buffer such as TE (Tris-EDTA (10mM:1mM)), TEA (40mm Tris-acetate, 1mm EDTA) buffer, or a lysate.

Once the DNA is obtained in a suitable solution, a binding matrix is typically added to the solution. Generally used binding matrixes are silica in the form of glass or diatoms. However, procedures using silica require high concentrations of chaotropes or alcohols for the DNA to bind to the surfaces. Currently used chaotropes include sodium iodide (NaI), urea, guanidinium Hydrochloride, sodium perchlorate (NaClO₄), and potassium bromide (KBr). Chaotropes and alcohols can be toxic, caustic, flammable and/or expensive. The process of the present invention does not require the presence of chaotropes or alcohols for binding to surfaces of the invention. Processes of the invention bind DNA in an aqueous solution at room temperature and elute the DNA in water at 37°C. However, if desired, chaotropes, alcohols and the like can be used with the process of the invention.

Typical procedures for using the process of the invention include the addition of the composition of the invention to a solution of DNA, which is generally followed by the addition of a binding buffer. At this point, it is advantageous that the process of the invention does not require a binding buffer. The solution can be incubated for a brief period at room temperature. After spinning, the supernatant can be discarded and the pellet washed. The DNA can then be eluted.

The composition of the invention is typically used in weight ranges from about 1:10 to 1:1 composition weight water. Preferably excess amounts of water are avoided and buffers such as TE can be used in place of water.

Next, a binding buffer is added, if used. After a brief incubation period at room temperature from about 1 to 20 minutes, preferably about 10, the container can be spun to obtain a pellet and supernatant fractions. The supernatant is separated and the pellet is washed with a reagent such as ethanol diluted with 50mM Tris. A preferred wash reagent concentration is 80% ethanol. DNA can then be eluted from the compositions of the invention by using elution buffers such as TE buffer, 1 X TAE buffer, and 1 X TBE buffer. More importantly, the use of elution buffers can be eliminated altogether, and DNA eluted in water by heating. For maximum yields the elution step can be repeated.

The chemical compositions of the invention can be conveniently assembled into a kit. A kit comprising the composition of the invention can include the composition in a container, such as a vial, with a suitable buffer, such as TE buffer and TAE buffer and optionally include a container of a binding buffer such as chaotropes, a container of wash buffer, such as a solution of ethanol diluted with 50mM tris or 1 X TAE, and a container of elution buffer, such as TE buffer, 1 X TAE buffer, and 1 X TBE buffer. Such a kit would allow convenient purification of DNA.

The following examples illustrate the specific embodiments of the invention described in this document. As would be apparent to skilled artisans, various changes and modifications are possible and are contemplated within the scope of the invention described.

EXAMPLES

EXAMPLE 1

The purpose of this experiment is to increase the surface hydroxyl groups on celite 545, acid washed, by refluxing in various solutions of NaOH. The increased presence of hydroxyls allows for easier modifications of the surface by chemical reactions.

EP 0 555 798 A1

Materials

Celite 545, acid washed (Alltech, Deerfield, IL, stock #9043, QC 243)

NaOH (Aldrich Lot #04027EP)

5 Eight experiments were set up exactly the same way except that the amount of NaOH was varied from one experiment to the next.

Experiment	Celite 545		NaOH		
	g	mMol	mg	mMol	eq celite
1	.5	8.33	16.8	.42	.05
2	.5	8.33	33.2	.83	.1
3	.5	8.33	100.0	2.50	.3
4	.5	8.33	168.0	4.20	.5
5	.5	8.33	233.0	5.83	.7
6	.5	8.33	333.0	8.33	1
7	.5	8.33	500.0	12.50	1.5
8	.5	8.33	666.0	16.66	2

20 In a typical experiment, the celite 545 is added to a round bottom flask followed by the NaOH dissolved in 10ml H₂O. Reflux 48 hours with stirring. Filter, wash with water and acetone and air dry. Store in a desiccator.

25 Results and Conclusions for Example 1:

FTIR analysis was done showing increase in the -OH signal of hydrated surfaces compared to the starting material.

30 EXAMPLE 2

This experiment describes how the DNA binding capacity of SUPER FINE SUPER FLOSS CELITE (Manville) was determined and how that compares to the Prep-A-Gene DNA purification kit.

35 Materials:

Super Fine Super Floss (SFSF) (Sample from Manville, Denver, CO (1:5 w/w in H₂O))

λDNA (BRL Cat. 56125A, Lot AJU702)

50mM Tris pH7.0 (Dilute from 1M stock) BRL Cat. 5505UA, Lot 60926

40 (PREP-A-GENE KIT (Bio-Rad, Richmond, CA))

Binding Buffers (Diluted from 6M stock) NaClO₄ Fisher Cat. 5490-500, Lot 914199

Wash Buffer 80% Ethanol in 50mM Tris, pH7.0

Elution Buffer Milli Q H₂O

Ethidium Bromide (10mg/ml) Sigma Cat. E-8751, Lot 99F3722 1% agarose BAL Cat. 5510UA, Lot 9N2204

45 1X TAE (from 50X stock) Tris Base-Sigma CAT T-1503, Lot 80H5633 Acetic Acid - Fisher A38-500 EDTA - Sigma CAT ED255, Lot 117F-0026

Type II Loading Dye (25% Ficoll 400, 0.25% Bromophenol Blue, 0.25% xylene cyanol Ficoll 400 - Sigma CAT F4375,

Bromophenol Blue - BIO-RAD CAT 161-0404, Lot M 1264, Xylene

50 Cyanole - Sigma CAT X-4126, Lot 8043740)

Type 57 and 55 POLAROID Film

METHODS

- 55 1. Two groups of reactions are set up, one for each surface type. Each surface has 8 tubes containing 50μl of the DNA solution. This solution is 0.5μl λDNA in 50μl 50mM Tris, pH7.0 for 31μg DNA/reaction. The titration ranges from 0M NaClO₄ to 6M NaClO₄.
2. Add 20μl of each surface to the reaction mixes.

EP 0 555 798 A1

3. Add 400 μ l Binding Buffer according to the titration. For Prep-A-Gene this was 0M, 2M, 2.5M, 3M, 3.5M, 4M, 4.5M, and 6M NaClO₄. For SFSF, the titration is 0M, 1M, 1.5M, 2M, 2.5M, 3M, 3.5M, and 4M NaClO₄.
4. Incubate for 10 minutes, with rocking, at room temperature.
5. Spin and discard supernatant.
6. Wash pellet 2 times with 80% ethanol/50mM Tris, pH7.0.
7. Elute DNA in 20 μ l H₂O, 37°C, 10 minutes.
8. Spin and remove supernatant to a separate tube. Repeat elution step and combine supernatants for ~40 μ l total.
9. Add 2 μ l, Type II loading dye to each tube.
10. Load onto a 1% agarose, 1 X TAE gel. Run for ~25 minutes at 100-130 volts in 1 X TAE buffer.
11. Stain with ethidium bromide in H₂O (~1:1000) for ~15 minutes. Destain for ~20-30 minutes.
12. Photograph over UV light with Type 57 Polaroid film. If possible, take negatives with Type 55 film.

15 Results and Conclusions

SFSF celite binds DNA strongly at a binding buffer concentration of 2.5m NaClO₄ compared to Prep-a-Gene's matrix which requires 3M NaCl₄ for binding DNA. For this reason SFSF will be used as the standard for comparison to the hydrated surfaces.

20

EXAMPLE 3

The purpose of the following experiment is to determine at what concentration of binding buffer will the hydrated SiO₂ surfaces allow recovery of DNA from a sample. The results will be compared to Super Fine Super Floss celite.

25

Surfaces

1. Acid washed Celite 545 + .05 eq NaOH.
2. Acid washed Celite 545 + .1 eq NaOH.
3. Acid washed Celite 545 + .3 eq NaOH.
4. Acid washed Celite 545 + .5 eq NaOH.
5. Acid washed Celite 545 + .7 eq NaOH.
6. Acid washed Celite 545 + 1 eq NaOH.
7. Acid washed Celite 545 + 1.5 eq NaOH.
8. Acid washed Celite 545 + 2.0 eq NaOH.
9. Super Fine Super Floss celite.

35

The remainder of materials and methods are set forth and carried out in substantial accordance with the teachings of Example 2.

40

45

50

55

Results:	
Celite + NaOH	Concentration of Binding Buffer Required for Strong DNA Binding to the Test Surface DNA Binding 1M-4M [NaClO ₄]
.05 eq NaOH	+
.10 eq NaOH	+
.30 eq NaOH	+ + 1.5M
.50 eq NaOH	+ + 2.0M
.70 eq NaOH	+ + 2.0M
1.0 eq NaOH	+ + 2.0M
1.5 eq NaOH	+ + 1.5M
2.0 eq NaOH	+ + + *
SFSF celite	+ + 3.0M

+ Trace amounts of DNA elute across the titration
 + + Near complete elution of DNA from 1.5, 2.0 or 3.0 M NaClO₄.
 + + + Near complete elution of DNA across the titration.

* Eluted DNA under native conditions. DNA binds in water at room temperature and eluted at 37 °C in water.


Conclusion

The results of gel electrophoresis demonstrates that as the amount of NaOH was increased in proportion to celite, the recovery of DNA from the resulting surface increased as well. As The mole ratio of NaOH:celite reaches 2.0, DNA was recovered under native conditions (no binding buffer required). Most of the surfaces isolated the DNA better than SFSF celite, without requiring concentrated amounts of binding buffer generally required to bring this about.


Although the invention has been described with respect to specific modifications, the details thereof are not to be construed as limitations, for it will be apparent that various equivalents, changes and modifications may be resorted to without departing from the spirit and scope thereof, and it is understood that such equivalent embodiments are to be included therein.

Claims


1. A process for purifying DNA which comprises contacting DNA with a composition of the formula:



wherein R' is OH, R is SiO₂, and R'' is OH or repeating units of the composition comprising the formula:




wherein R' is independently OH or



5

R is SiO₂, and R'' is independently OH or

10




15

or with a composition comprising the above monomer and repeating units thereof

2. The process of Claim 1 which further comprises the use of chaotropes.
3. The process of Claim 1 which further comprises elution of DNA from the hydrated SiO₂.
4. The process of Claim 3 in which elution is obtained by heat and water.
5. A kit for purifying DNA comprising a composition of the formula:


25



30

wherein R' is OH, R is SiO₂, and R'' is OH, or repeating units of the composition comprising the formula:


35



40


wherein R' is independently OH or

45




R is SiO₂, and R'' is independently OH or

50




55

6. A compound of formula:



5


wherein R' is OH, R is SiO₂, and R'' is OH, and repeating units of the composition comprising the formula:



10

15


wherein R' is independently OH or



20

R is SiO₂, and R'' is independently OH or

25



30

or compositions comprising the above monomer and repeating units thereof

7. The reaction products of SiO₂ and hydrating reagents.

35

8. A process for producing reaction products of hydrating reagents and SiO₂ which comprises reacting hydrating reagents with SiO₂.

9. The reaction products of Claim 7 in which the hydrating reagent is NaOH.

40

10. The process of Claim 8 in which the hydrating reagent is NaOH.

45

50

55



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 93 10 1923

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
A	GB-A-2 074 892 (VARIAN ASSOCIATES) * page 5 - page 6; examples 1-2 * * claims * ---	1	C07H1/08 C12N15/10
A	US-A-5 075 430 (MICHAEL C. LITTLE) * claims 1-3 * ---	1-2	
A	WO-A-9 100 924 (UNIVERSITY OF KANSAS) * claims * ---	1	
A	EP-A-0 235 526 (VEB LEIPZIGER ARZNEIMITTELWERK) * page 6, line 30 - line 36 * * claims * -----	1	
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			C07H C12N
The present search report has been drawn up for all claims			
Place of search	Date of completion of the search	Examiner	
THE HAGUE	24 MAY 1993	SIEMENS T.	
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			

EPO FORM 1503 03.82 (P/901)