



Publication number : **0 566 199 A1**

EUROPEAN PATENT APPLICATION

Application number : **93201058.0**

Int. Cl.⁵ : **C07D 413/12, A61K 31/42**

Date of filing : **13.04.93**

Priority : **15.04.92 US 869287**

Date of publication of application :
20.10.93 Bulletin 93/42

Designated Contracting States :
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC
NL PT SE**


Applicant : **STERLING WINTHROP INC.**
90 Park Avenue
New York, NY 10016 (US)

Inventor : **Diana, Guy Dominic, c/o Sterling
Winthrop Inc.**
Patent Department, 90 Park Avenue
New York, New York 10016 (US)
Inventor : **Nitz, Theodore John, c/o Sterling
Winthrop Inc.**
Patent Department, 90 Park Avenue
New York, New York 10016 (US)

Representative : **Haile, Helen Cynthia et al**
Kodak Limited Patent Department Headstone
Drive
Harrow, Middlesex HA1 4TY (GB)

1,2,4-Oxadiazolyl-phenoxyalkylisoxazoles and their use as antiviral agents.

Compounds of the formula



wherein :

R₁ is alkyl, alkoxy, hydroxy, cycloalkyl, hydroxyalkyl, alkoxyalkyl or hydroxyalkoxy ;

Y is alkylene of 3 to 9 carbon atoms,

R₂ and R₃ independently are hydrogen, alkyl, alkoxy, halo, trifluoromethyl or nitro ;


R₄ is alkoxy, hydroxy, halomethyl, dihalomethyl, trihalomethyl, cycloalkyl, alkoxyalkyl, alkoxyalkyl, alkanecarbonyloxyalkyl, cyano, 2,2,2-trifluoroethyl, (4-methylphenyl)sulfonyloxymethyl, N=Q or CON=Q, where N=Q is amino, alkylamino or dialkylamino ; or pharmaceutically acceptable acid-addition salts thereof are useful as antiviral agents.

Preferred compounds are those wherein Y is alkylene of 3 to 5 carbon atoms and especially when R₂ and R₃ are in the 2- and 6- positions of the phenyl ring.

This invention relates to novel 1,2,4-oxadiazolylphenoxyalkylisoxazoles, to methods for the preparation thereof, and compositions and methods for the use thereof as antiviral agents.

Diana U.S. Patent 4,843,087 discloses heterylphenoxyalkylisoxazoles, wherein the heteryl moiety is an oxazole or an oxazine, which exhibit antiviral activity.

5 Diana et al. U.S. Patent 4,857,53915 discloses antivirally active compounds of the formula



15 wherein:

Y is an alkylene bridge of 3 to 9 carbon atoms;

Z is N or HC:


R is hydrogen or lower-alkyl of 1 to 5 carbon atoms, with the proviso that when Z is N, R is lower-alkyl;

20 R₁ and R₂ are hydrogen, halogen, lower-alkyl, lower-alkoxy, nitro, lower-alkoxycarbonyl or trifluoromethyl; and

Het is selected from specified heterocyclic groups.

Included in the definition of Het is unsubstituted 1,3,4-oxadiazol-2-yl and unsubstituted 1,2,4-oxadiazol-5-yl.

25 Diana et al. U.S. Patent 4,861,79129 discloses antivirally active compounds of the formula, inter alia,



35 wherein:

Y is an alkylene bridge of 3 to 9 carbon atoms optionally interrupted by one or two oxygen atoms, by cyclohexyl or by an olefinic linkage;

X is O, S, SO or SO₂;


Z is N or R₈C, where R₈ is hydrogen or lower-alkanoyl;

40 R₁ and R₂ are selected from the group consisting of hydrogen, lower-alkyl, lower-alkenyl, halogen, nitro, lower-alkoxy, lower-alkylthio, difluoromethyl, trifluoromethyl, amino, lower-alkanoylamino, di-lower-alkylamino, hydroxy, lower-alkanoyl, lower-alkanoyl, hydroxymethyl and carboxy;


R and R₃ are each hydrogen or alkyl of 1 to 3 carbon atoms optionally substituted by a member of the group consisting of hydroxy, lower-alkanoyloxy, lower-alkoxy, halo or N=Z', wherein N=Z' is amino, lower-alkanoylamino, lower-alkylamino, di-lower-alkylamino, 1-pyrrolidyl, 1-piperidinyl or 4-morpholinyl; with the proviso that when Z is N, R is other than hydrogen; and

Het is selected from specified heterocyclic groups including unsubstituted 1,3,4-oxadiazol-2-yl.

Diana et al. U.S. Patent 4,942,241 discloses antivirally active compounds of the formulae



and



wherein:


Y is an alkylene bridge of 1 to 9 carbon atoms;

R' is lower-alkyl or hydroxy-lower-alkyl of 1 to 5 carbon atoms;

R₁ and R₂ are hydrogen, halogen, lower-alkyl, lower-alkoxy, nitro, lower-alkoxycarbonyl or trifluoromethyl; and

R₈ is hydrogen or lower-alkyl of 1 to 5 carbon atoms.

Diana U. S. Patent 4,945,164 discloses antivirally active compounds of the formula, *inter alia*,



wherein:


Y is an alkylene bridge of 3 to 9 carbon atoms;

R' is lower-alkyl or hydroxy-lower-alkyl of 1 to 5 carbon atoms;

R₁ and R₂ are hydrogen, halogen, lower-alkyl, lower-alkoxy, nitro, lower-alkoxycarbonyl or trifluoromethyl; and

R₈ is hydrogen or lower-alkyl of 1 to 5 carbon atoms.

G.D. Diana and T.R. Bailey in U.S. Patent Application Serial No. 07/731,569 disclose compounds of the formula



wherein:

Y is alkylene of 3 to 9 carbon atoms;


R₁ is lower-alkyl, lower-alkoxy-(C₁₋₃-alkyl), lower-alkoxycarbonyl, cyclopropyl or trifluoromethyl;

R₂ and R₃ independently are hydrogen, lower-alkyl, halogen, lower-alkoxy, nitro, trifluoromethyl or hydroxy; and

R₄ is hydrogen or lower-alkyl; where lower-alkyl and lower-alkoxy, each occurrence, have from 1 to 5 carbon atoms;

with the proviso that when R₁ is lower-alkyl, at least one of R₂ and R₃ is hydroxy.

The present invention provides a compound of the formula



I

wherein:


R₁ is alkyl, alkoxy, hydroxy, cycloalkyl, hydroxyalkyl, alkoxyalkyl or hydroxyalkoxy;

Y is alkylene of 3 to 9 carbon atoms,

R₂ and R₃ independently are hydrogen, alkyl, alkoxy, halo, trifluoromethyl or nitro; and

R₄ is alkoxy, hydroxy, halomethyl, dihalomethyl, trihalomethyl, cycloalkyl, alkoxyalkyl, alkoxyalkyl, alkanecarbonyloxyalkyl, cyano, 2,2,2-trifluoroethyl, (4-methylphenyl)sulfonyloxymethyl, N=Q or CON=Q, where N=Q is amino, alkylamino or dialkylamino; or a pharmaceutically-acceptable acid-addition salt thereof.


The present invention also provides a compound of the formula



III

wherein Y, R₂ and R₃ are as defined above and R₆ is alkoxy, fluoromethyl, difluoromethyl, trihalomethyl, cycloalkyl or alkoxyalkyl.

The present invention also provides a compound of the formula



IV

wherein R₂ and R₃ are as defined above and R₇ is alkoxy, fluoromethyl, difluoromethyl, trifluoromethyl, cycloalkyl, alkoxyalkyl or cyano.

The present invention also provides compounds of formulae XVII and XXI hereinafter.

The present invention also provides a composition for combatting picornaviruses which comprises an antivirally effective amount of a compound of formula I in admixture with a suitable carrier or diluent and to methods for combatting picornaviruses therewith including combatting a picornaviral infection in a mammalian host.

The compounds of formula I are useful as antipicornaviral agents.

The compounds of formulas III, IV, XVII and XXI are useful as intermediates for the preparation of the compounds of formula I.

Preferred compounds of formula I are those wherein


R₁ is C₁₋₃-alkyl, C₁₋₃-alkoxy, hydroxy, cyclopropyl, hydroxy-C₁₋₃-alkyl, C₁₋₃-alkoxy-C₁₋₃-alkyl or hydroxy-C₁₋₃-alkoxy;

Y is alkylene of 3 to 9 carbon atoms, especially 3 to 5 carbon atoms;

R₂ and R₃ independently are hydrogen, C₁₋₃-alkyl, C₁₋₃-alkoxy or halo; and

R₄ is C₁₋₃-alkoxy, hydroxy, halomethyl, dihalomethyl, trihalomethyl, cyclopropyl, C₁₋₃-alkoxycarbonyl, hydroxy-C₁₋₃-alkyl, C₁₋₃-alkoxy-C₁₋₃-alkyl, (C₁₋₃-alkane)carbonyloxy-C₁₋₃-alkyl, cyano, 2,2,2-trifluoroethyl, 4-(methylphenyl)-sulfonyloxymethyl, N=Q or CON=Q, where N=Q is amino, C₁₋₃-alkylamino or di-(C₁₋₃-alkyl)amino.

More preferred compounds of formula I are compounds of the formula




I A

wherein R₁, Y, R₂, R₃ and R₄ are as defined above for formula I and especially wherein R₁, Y, R₂, R₃ and R₄ are as defined in the previous paragraph for the preferred compounds of formula I.

Especially preferred are the compounds of formula I or IA wherein R₄ is C₁₋₃-alkoxy, fluoromethyl, dihalomethyl, trihalomethyl, cycloalkyl or C₁₋₃-alkoxy-C₁₋₃-alkyl, especially trifluoromethyl.

It should be understood that in the compounds of the invention, when the 1,2,4-oxadiazole ring is substituted by hydroxy, amino or alkylamino, they may exist in any of three possible tautomeric forms as follows:



wherein R₄ is hydroxy, amino or alkylamino and T is O, NH or N-alkyl, and such tautomers are within the purview of the invention.

As used herein, unless otherwise specifically defined, alkyl, alkane, alkoxy, cycloalkyl and halo each has the following meaning:

alkyl and alkoxy mean aliphatic radicals, including branched radicals, of from one to five carbon atoms. Thus the alkyl moiety of such radicals include, for example methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, t-butyl and pentyl;

alkane means a monovalent aliphatic radical, including branched radicals of from one to four carbon atoms. Thus the alkane moiety of such radical includes, for example, methyl, ethyl, propyl, isopropyl, n-butyl and sec-butyl;


cycloalkyl means an alicyclic radical having from three to six carbon atoms as illustrated by cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl; and

halo means bromo, chloro, iodo or fluoro.

As used herein, in hydroxyalkyl and alkoxyalkyl, the hydroxy and alkoxy groups can occur at any available position of alkyl. Thus hydroxyalkyl and alkoxyalkyl include, for example, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2-hydroxypropyl, 2-hydroxyisopropyl, 2, 3, 4 and 5-hydroxypentyl and the like and corresponding alkyl ethers thereof.

As used herein, in hydroxyalkoxy, the hydroxy group can occur at any available position of alkoxy other than the C-1 position. Thus hydroxyalkoxy includes, for example, 2-hydroxyethoxy, 2-hydroxypropoxy, 2-hydroxyisopropoxy, 2-and 5-hydroxypentoxy and the like.

The compounds of formula I wherein R_1 is alkyl, alkoxy, cycloalkyl or alkoxyalkyl, Y , R_2 and R_3 are as defined hereinbefore, and R_4 is hydroxy, halomethyl, dihalomethyl, trihalomethyl, cycloalkyl, alkoxyalkyl, alkoxyalkyl, alkanecarbonyloxyalkyl or 2,2,2-trifluoroethyl, can be prepared by a process which comprises reacting an amidoxime (N-hydroxycarboximidamide) of the formula




V

with an acid halide, R_4COX , an alkyl haloformate, $ROCOX$ (in the case where R_4 in formula I is hydroxy), where R is methyl or ethyl, or an acid anhydride, $(R_4CO)_2O$, where R_1 , Y , R_2 , R_3 and R_4 are as defined above in this paragraph and X is bromo, chloro, fluoro or iodo, under anhydrous conditions to form the corresponding compound of formula I. The process involves the following methods. In one method, the amidoxime V is reacted with the acid halide or the acid anhydride in the presence of an organic or inorganic base, e.g., pyridine, triethylamine or potassium carbonate, in an inert solvent, e.g., acetone, methylene chloride, chloroform, toluene or tetrahydrofuran, or in a base which also functions as the solvent, e.g., pyridine, at an elevated temperature (about 40-130°C) or at a reduced temperature (about 0-15°C). In the latter case an intermediate O-acyl derivative $[C(NH_2)=NOC(=O)-(R_4 \text{ or } OR)]$ is isolated and heated at a temperature in the range of about 100-130°C for a time sufficient for cyclization to the oxadiazole of formula I to occur, generally about 5 minutes to 4 hours. In another method, the amidoxime V is reacted with the acid halide or acid anhydride in an acid which corresponds to the acid halide or acid anhydride at an elevated temperature (about 70-100°C).

The compounds of formula I where R_1 is alkyl, alkoxy, cycloalkyl or alkoxyalkyl, Y , R_2 and R_3 are as defined hereinbefore, and R_4 is dihalomethyl, trihalomethyl, cycloalkyl, alkoxyalkyl, alkanecarbonyloxyalkyl or 2,2,2-trifluoroethyl can be prepared by a process which comprises reacting amidoxime V with the product obtained by reaction of a carboxylic acid, R_4CO_2H , where R_1 , Y , R_2 , R_3 and R_4 are as defined above in this paragraph, with the coupling agent N,N'-carbonyldiimidazole, prepared as described in the examples, in an inert solvent, e.g., acetone, tetrahydrofuran, chloroform, methylene chloride or toluene, at an elevated temperature (about 40-80°C) to form the corresponding compound of formula I.

The compounds of formula I wherein R_1 is alkyl, alkoxy, cycloalkyl or alkoxyalkyl, Y , R_2 and R_3 are as defined hereinbefore, and R_4 is amino can be prepared by a process which comprises reacting amidoxime V, where R_1 , Y , R_2 and R_3 are as defined above in this paragraph, with cyanogen halide, CNX_1 , where X_1 is bromo, chloro or iodo, in the presence of a base, e.g., potassium or sodium bicarbonate, in an alcoholic solvent, e.g., ethyl alcohol, at about room temperature to give the compound of formula I where R_4 is amino.

The compounds of formula I where R_1 is alkyl, alkoxy, cycloalkyl or alkoxyalkyl, Y , R_2 and R_3 are as defined hereinbefore and R_4 is CH_2CF_3 can be prepared by reacting amidoxime V where R_1 is as defined above in this paragraph and Y , R_2 and R_3 are as defined hereinbefore, with a ketene 1,3-propanedithiol acetal of the formula




to give the corresponding compound of formula I.

The amidoxime V and ketene 1,3-propanedithiol acetal are reacted in the presence of silver trifluoroacetate in an inert solvent, e.g., tetrahydrofuran, dioxane, dimethylformamide or N-methylpyrrolidinone, at a temper-

ature in the range of from about 60 to about 100°C. Preferably the reaction is carried out in the dark.

The intermediate amidoxime V is prepared according to the following flow sheet:




The bromophenol VI reacts with the cuprous cyanide in an inert solvent at an elevated temperature, e.g., in dimethylformamide at reflux temperature, to give the cyanophenol VII. The latter is reacted with haloisoxazole VIII, where X_2 is chloro, bromo or iodo, in a dry inert solvent, e.g., acetonitrile or N-methylpyrrolidinone, in the presence of a base, e.g., potassium carbonate or sodium hydroxide, optionally in the presence of a catalytic amount of potassium or sodium iodide, at an elevated temperature (50-120°C) to give the cyano compound IX. The cyano compound IX reacts with the hydroxylamine hydrochloride in the presence of a base, e.g., potassium or sodium carbonate, sodium acetate or sodium hydroxide, in an alcoholic solvent, e.g., ethyl alcohol, at an elevated temperature (50-150°C) to give the amidoxime V.

Certain intermediate compounds of formula IX wherein R_1 is alkyl, cycloalkyl or alkoxyalkyl and Y, R_2 and R_3 are as defined hereinbefore can be prepared by reacting the ethynyl compound XII described hereinafter with a nitrile oxide, $R_1C\equiv N\rightarrow O$, where R_1 is as defined above in this paragraph, using a procedure similar to that described hereinafter for the preparation of compound I from the ethynyl compound III.

The intermediate bromophenols of formula VI and cyanophenols of formula VII belong to generically known classes of compounds and are readily prepared by known procedures.

The intermediate haloisoxazoles of formula VIII can be prepared by the procedure described in U.S. Patent 4,843,087, i.e., by reacting an alkali metal derivative of an isoxazole of the formula






X


wherein R_1 is alkyl, alkoxy, cycloalkyl or alkoxyalkyl, with a dihalide, $\text{X}_2\text{-Y'-X}_2$, where Y' is alkylene of 2 to 8 carbon atoms and X_2 is as defined above. The alkali metal derivative is prepared *in situ* by treating isoxazole X with an organo-alkali metal base such as butyllithium or lithium diisopropylamide under anhydrous conditions.

The compounds of formula I wherein R_1 is alkyl, cycloalkyl or alkoxyalkyl, Y , R_2 and R_3 are as defined hereinbefore, and R_4 is alkoxy, trihalomethyl, cycloalkyl, alkoxycarbonyl, alkoxyalkyl or 2,2,2-trifluoroethyl, can be prepared by a process which comprises reacting an ethynyl compound of formula III hereinabove, wherein R_6 has the meaning defined above in this paragraph for R_4 , with a nitrile oxide of the formula $\text{R}_1\text{C}\equiv\text{N}\rightarrow\text{O}$ which is prepared *in situ* from a hydroxyimino halide of the formula $\text{R}_1\text{C}(\text{X}_3)=\text{NOH}$, where X_3 is chlorine or bromine, in the presence of an amine base, e.g., triethylamine, pyridine or N-methylpyrrolidine. The hydroxyimino halides, which may also be prepared *in situ*, belong to a generically known class of compounds and are readily prepared by conventional procedures, e.g., by reacting the corresponding aldehyde oxime ($\text{R}_1\text{C}=\text{NOH}$) with a halogenating agent, e.g., N-chlorosuccinimide or bromine. The process for preparing the compounds of formula I by reacting the ethynyl compound of formula III takes place by heating the reactants in an inert polar solvent, e.g., dimethylformamide or N-methylpyrrolidone, at a temperature in the range of about 20 to about 120°C.

The intermediate ethynyl compounds of formula III are prepared according to the following flow sheet:



The cyanophenol VII is reacted with haloalkyne XI, where X_2 is as defined hereinbefore, using a procedure similar to that described above for the preparation of the cyano compound IX from compounds VII and VIII, to give the ethynyl compound of formula XII. Ethynyl compound XII is reacted with the hydroxylamine hydrochloride, using a procedure similar to that described above for the preparation of amidoxime V from cyano compound IX, to give the amidoxime of formula XIII. The amidoxime XIII is reacted with the acid halide R_4COX , acid anhydride $(R_4CO)_2O$, carboxylic acid R_4CO_2H or




using procedures similar to those described hereinbefore for the preparation of compounds of formula I from amidoxime V.

The haloalkynes of formula XI belong to a generically known class of compounds.

The compounds of formula I where R_1 is alkyl, alkoxy, cycloalkyl or alkoxyalkyl, Y, R_2 and R_3 are as defined hereinbefore, and R_4 is alkoxy, trihalomethyl, cycloalkyl, alkoxycarbonyl, alkoxyalkyl or 2,2,2-trifluoroethyl, can be prepared by a process which comprises reacting a phenol of the formula IV above wherein R_2 and R_3 are as defined hereinbefore and R_7 is as defined above in this paragraph for R_4 , with a haloisoxazole of formula VIII above where R_1 is as defined above in this paragraph and Y and X_2 are as defined hereinbefore, to give the corresponding compound of formula I. The procedure used is similar to that described above for the preparation of cyano compound IX by reaction of cyanophenol VII and haloisoxazole VIII.


The intermediate haloisoxazole VIII can be prepared as described hereinbefore.

The intermediate phenols of formula IV can be prepared by reacting cyanophenol VII with hydroxylamine hydrochloride, using a procedure similar to that described hereinbefore for the preparation of amidoxime V from cyano compound IX, to give an amidoxime of the formula




XIV

Amidoxime XIV is reacted with R_4COX , $(R_4CO)_2O$, R_4CO_2H or



using procedures similar to those described hereinbefore for the preparation of compounds of formula I from amidoxime V, to give the corresponding phenol of formula IV.

The compounds of formula I wherein R_1 is hydroxyalkyl, Y, R_2 and R_3 are as defined hereinbefore, and R_4 is dihalomethyl, trihalomethyl, cycloalkyl, alkoxyalkyl, 2,2,2-trifluoroethyl or amino can be prepared from a compound of the formula




XVII

wherein R_1' is tert-butyldimethylsilyloxyalkyl $[(CH_3)_3CSi(Me)_2-O-alkyl]$ and Y, R_2 , R_3 and R_4 are as defined above in this paragraph, by cleaving the tert-butyldimethylsilyl ether.

Cleavage of the tert-butyldimethylsilyl ether is carried out by treating compound XVII with a strong organic acid, e.g., acetic acid or trifluoroacetic acid, or inorganic acid, e.g., hydrochloric acid or sulfuric acid, in an inert solvent, e.g., tetrahydrofuran or dioxane in the presence of water at a temperature in the range of from about 20 to about 60°C.

The compound of formula XVII where R_4 is dihalomethyl, trihalomethyl, cycloalkyl, alkoxyalkyl or 2,2,2-trifluoroethyl, can be prepared by a process which comprises reacting phenol IV wherein R_2 and R_3 are as defined hereinbefore and R_7 is as defined above in this paragraph for R_4 , with an isoxazole of the formula



XV

XVI


wherein R_1' , Y and X_2 are as defined hereinbefore.

The phenol IV is reacted with haloisoxazole XVI using a procedure similar to that described hereinbefore for the preparation of cyano compound IX from cyanophenol VII and haloisoxazole VIII.

The phenol IV is reacted with isoxazole XV in the presence of diethyl azodicarboxylate (DEAD) and triphenylphosphine in an inert solvent, e.g., tetrahydrofuran, chloroform, dimethylformamide or N-methylpyrrolidinone, at a temperature in the range of from about -20 to about 20°C.

The intermediate phenol IV can be prepared by the procedure described hereinbefore.

The intermediate isoxazoles XV and XVI can be prepared by reacting isoxazole X, wherein R_1 is hydroxyalkyl, with tert-butyldimethylsilyl chloride to give the corresponding tert-butyldimethylsilyl ether of formula




XVIII

where R_1' is as defined above, and reaction of an alkali metal derivative of compound XVIII with ethylene oxide or $X_2-Y'-X_2$ respectively.

Isoxazole X, wherein R_1 is hydroxyalkyl, is reacted with tert-butyl(dimethyl)silyl chloride in the presence of 4-(dimethylamino)pyridine and a base, e.g., triethylamine, pyridine or imidazole, in a dry inert solvent, e.g., methylene chloride, chloroform or tetrahydrofuran, at room temperature to give compound XVIII. Isoxazole XV is prepared by reacting an alkali metal derivative of compound XVIII with ethylene oxide, preferably in the presence of a chelating agent, e.g., N,N,N',N'-tetramethylethylenediamine or hexamethyl phosphoric triamide, in a dry inert solvent, e.g., tetrahydrofuran, at a temperature in the range of from about -78 to about 20°C. The


alkali metal derivative is prepared *in situ* by reacting compound XVIII with an organo-alkali metal base, e.g., butyllithium or lithium diisopropylamide, under anhydrous conditions.

The compound of formula XVII, where R_4 is dihalomethyl, trihalomethyl, cycloalkyl, alkoxyalkyl or 2,2,2-trifluoroethyl, can also be prepared, as can the compound of that formula where R_4 is amino, according to the following flow sheet:



The reaction of compound XV or XVI with cyanophenol VII to give compound XIX is carried out by procedures similar to those described hereinbefore for preparing compound XVII by reacting phenol IV with isoxazole XV or haloisoxazole XVI respectively. The reaction of cyano compound XIX with hydroxylamine hydrochloride to give amidoxime XX, and the latter with the acid halide, acid anhydride, carboxylic acid, cyanogen halide or ketene 1,3-propanedithiol acetal to give compound XVII can be carried out by procedures similar to those described hereinbefore for preparing amidoxime V from cyano compound IX and for preparing the compound of formula I from amidoxime V.

The compounds of formula I wherein R_1 is hydroxy, Y, R_2 and R_3 are as defined hereinbefore, and R_4 is cycloalkyl or alkoxyalkyl can be prepared by reacting a compound of the formula



XXI

where R_8 is alkyl and Y, R_2 , R_3 and R_4 are as defined above in this paragraph, with hydroxylamine hydrochloride to give the compound of formula I where R_1 is hydroxy.

Compound XXI is reacted with hydroxylamine hydrochloride in the presence of a base, e.g., sodium hydroxide, and water in an alcoholic solvent, e.g., methyl or ethyl alcohol, at a temperature in the range of from about 0 to about 25°C.

The intermediate compounds of formula XXI can be prepared by reacting an alkali metal derivative of compound III, wherein R_6 is as defined for R_4 of compound XXI, with an alkyl haloformate, R_8OCOX , where X is as defined hereinbefore. The reaction takes place in a dry inert solvent, e.g., tetrahydrofuran or dioxane, at an initial temperature of about -78 to about -20°C with subsequent warming to about 20 to about 25°C. The alkali metal derivative can be prepared *in situ* by reacting compound III with an organo-alkali metal, e.g., butyllithium or lithium diisopropylamide, under anhydrous conditions.

Certain compounds of formula I are intermediates for other compounds of formula I as described herein-after.

The acid halides, alkyl haloformates and acid anhydrides used in the hereinbefore described processes for preparing the compounds of formula I and intermediates therefor, belong to well known classes of compounds and can be readily prepared by known procedures.

The compound of formula I wherein R_1 is alkyl, cycloalkyl or alkoxyalkyl, Y, R_2 , R_3 are as defined hereinabove, and R_4 is alkoxy or $\text{N}=\text{Q}$, where $\text{N}=\text{Q}$ is alkylamino or dialkylamino, can be prepared from the corresponding compound of formula I wherein R_4 is trichloromethyl. In the case where R_4 is alkoxy, the trichloromethyl compound is reacted with an alkali metal alkoxide, e.g., sodium methoxide or sodium ethoxide, and in the case where R_4 is $\text{N}=\text{Q}$, with an amine ($\text{HN}=\text{Q}$), in a suitable solvent, e.g., dimethylformamide or N-methylpyrrolidinone, at room temperature to give the corresponding compound of formula I where R_4 is alkoxy, alkylamino or dialkylamino.

The compounds of formula I wherein R_1 is hydroxyalkyl, Y, R_2 and R_3 are as defined hereinbefore, and R_4 is hydroxy, dihalomethyl, trihalomethyl, cycloalkyl, hydroxyalkyl, 2,2,2-trifluoroethyl or amino, can be prepared from the corresponding compound wherein R_1 is alkoxyalkyl by ether cleavage of the alkoxyalkyl moiety. The alkoxyalkyl compound is treated with trimethylsilyl iodide in a dry inert solvent, e.g., 1,2-dichloroethane, chloroform or acetonitrile, at a temperature in the range of from about 60 to about 80°C to give the corresponding hydroxyalkyl compound.

The compounds of formula I wherein R_1 is alkyl, alkoxy, cycloalkyl or alkoxyalkyl, Y, R_2 and R_3 are as defined hereinbefore, and R_4 is $\text{CON}=\text{Q}$, where $\text{N}=\text{Q}$ is amino, alkylamino or dialkylamino, can be prepared by reacting the corresponding compound of formula I wherein R_4 is alkoxycarbonyl with amine $\text{HN}=\text{Q}$ in a polar solvent, e.g., ethyl alcohol or N-methylpyrrolidinone, at room temperature to give the corresponding compound where R_4 is $\text{CON}=\text{Q}$.

The compound of formula I where R_1 is alkyl, alkoxy, cycloalkyl or alkoxyalkyl, Y, R_2 and R_3 are as defined hereinbefore, and R_4 is cyano, can be prepared from the corresponding compound wherein R_4 is $\text{CON}=\text{Q}$, where $\text{N}=\text{Q}$ is amino, by treating the latter with trifluoroacetic anhydride in the presence of a base, e.g., pyridine or triethylamine, in a dry inert solvent, e.g., tetrahydrofuran, chloroform or 1,2-dichloroethane, at a temperature in the range of from about 0 to about 20°C.

The compounds of formula I wherein R_1 is alkoxy or hydroxyalkoxy, Y, R_2 and R_3 are as defined above, and R_4 is alkoxy, trihalomethyl, cycloalkyl, alkoxyalkyl, 2,2,2-trifluoroethyl or dialkylamino, can be prepared by etherification of the corresponding compound of formula I wherein R_1 is hydroxy. The etherification takes place by reacting the hydroxy compound with an alkyl halide or hydroxyalkyl halide, where halide is bromide, chloride or iodide, in the presence of a base, e.g., potassium carbonate or sodium carbonate, in an inert dry solvent, e.g., acetone, butanone or acetonitrile, at a temperature in the range of from about 50 to about 90°C.

The compounds of formula I wherein R_1 is alkyl, cycloalkyl, hydroxyalkyl, alkoxyalkyl or hydroxyalkoxy, Y, R_2 and R_3 are as defined hereinbefore, and R_4 is hydroxyalkyl can be prepared by transesterification of the

corresponding compound of formula I wherein R_4 is alkanecarbonyloxyalkyl. The transesterification is carried out by treating the alkanecarbonyloxyalkyl compound with an inorganic or organic base, e.g., potassium carbonate, sodium bicarbonate or triethylamine, in an alcoholic solvent, e.g., methyl or ethyl alcohol, at room temperature.

The compounds of formula I wherein R_1 is alkyl, cycloalkyl or hydroxyalkyl, Y , R_2 and R_3 are as defined hereinbefore, and R_4 is hydroxyalkyl, can also be prepared by ether cleavage of the corresponding compound of formula I wherein R_4 is alkoxyalkyl. The ether cleavage can be carried out by treating the alkoxy compound with trimethylsilyl iodide using a procedure similar to that described hereinbefore for preparing the compound of formula I wherein R_1 is hydroxyalkyl.

The compound of formula I wherein R_1 is alkyl, alkoxy, cycloalkyl, hydroxyalkyl, alkoxyalkyl or hydroxyalkoxy, Y , R_2 and R_3 are as defined hereinbefore, and R_4 is iodomethyl, can be prepared from the corresponding compound of formula I wherein R_4 is chloromethyl by reaction with alkali metal iodide, e.g., sodium iodide. The reaction takes place by treating the chloromethyl compound with the alkali metal iodide, e.g., sodium or potassium iodide, in an inert solvent, e.g., acetone or butanone, at about 20°C.

The compounds of formula I wherein R_1 is alkyl, alkoxy, cycloalkyl or alkoxyalkyl, Y , R_2 and R_3 are as defined hereinbefore, and R_4 is (4-methylphenyl)sulfonyloxymethyl, are prepared from the corresponding compound of formula I wherein R_4 is hydroxymethyl by reaction with (4-methylphenyl)sulfonyl halide, where halide is bromide, chloride or iodide, in the presence of an inorganic base, e.g., potassium carbonate or sodium bicarbonate. The reaction takes place by reacting the reactants in an inert solvent, e.g., methylene chloride, chloroform or 1,2-dichloroethane, at about 20°C.

The compounds of formula I are sufficiently basic to form stable acid-addition salts with strong acids and such salts are within the purview of the invention. The nature of the acid-addition salt is immaterial, provided it is derived from an acid the anion of which is essentially non-toxic to animal organisms. Appropriate acid-addition salts include hydrochloride, hydrobromide, sulfate, acid sulfate, maleate, citrate, tartrate, methanesulfonate, p-toluenesulfonate, dodecyl sulfate and cyclohexanesulfonate. The acid-addition salts are prepared by conventional methods known in the art.

In the various processes described hereinabove for the preparation of the compounds of the invention, it will be appreciated that the reactions should be carried out for a time sufficient to provide the desired product and that for any specific reaction type, the time of the reaction will depend upon one or more factors such as, e.g., the nature of the reactants, the solvent employed and/or the temperature at which the reaction is carried out.

The antiviral compounds of the invention are formulated for use by preparing a dilute solution or suspension in a pharmaceutically acceptable aqueous, organic or aqueous-organic medium for topical or parenteral administration by intravenous or intramuscular injection, or for intranasal or ophthalmic application; or are prepared in tablet, capsule, or aqueous suspension form with conventional excipients for oral administration.

The structures of the compounds of the invention were established by modes of synthesis and elementary analysis, and by infrared, nuclear magnetic resonance and/or mass spectra.

The invention will now be described with reference to the following examples but the scope is in no way to be limited thereto.

Example 1

a) 3-(3-Methylisoxazol-5-yl)propyl alcohol.

3,5-Dimethylisoxazole (X) (220 g, 2.27 moles) in 2.2 l tetrahydrofuran under nitrogen was cooled with stirring to -75°C and 908 ml 2.5M n-butyllithium (2.27 moles) in hexanes were added over 1 hour keeping the temperature at or less than 65°C. The chilled solution was stirred for thirty minutes after addition was complete and was then treated at about -70°C with a solution of 112 g (2.54 moles) of ethylene oxide in 390 ml tetrahydrofuran over a period of 1.5 hours, keeping the temperature at about -65 °C and stirred overnight. The mixture at 8°C was quenched with continued cooling in an 8°C bath by adding 1.2 l 2.5M hydrochloric acid over a period of 20 minutes, during which time the temperature rose to 23°C, and was stirred for 10 minutes. The organic phase was separated, washed with 500 ml of water and concentrated to give 147 g of title compound as a brown oil. The combined aqueous phases (original + wash phase) were extracted with methyl tert-butyl ether (3 x 200 ml) and the combined organic extracts were concentrated to give an additional 125 g of title compound as a brown oil.

b) 3-(3-Methylisoxazol-5-yl)propyl chloride. (VIII)

To the product from part (a) (125 g, 0.885 mole) in 1225 ml methylene chloride was added 192 ml (2.63 moles) of thionyl chloride over a period of 1 hour during which time the temperature rose to 40°C to a gentle reflux. Heating at reflux was continued for 3 hours, the reaction mixture was allowed to stand over-

night, and then heating at reflux was continued for 1 hour. The reaction mixture was added as a steady stream to 3 kg of ice water with vigorous stirring, stirring was continued for 1 hour and the aqueous phase was separated. Water (1 l) was added to the organic phase followed by 161 g solid sodium bicarbonate in portions with vigorous stirring. The organic phase was separated and concentrated *in vacuo* to give a black oil which was purified by wiped-film distillation to give 94 g of the title compound as a yellow oil, bp 65°C/0.09 mm.

c) 3,5-Dimethyl-4-[3-(3-methylisoxazol-5-yl)propyloxy]-benzonitrile. (IX)

A mixture of 3,5 -dimethyl-4-hydroxybenzonitrile (VII) (7.36 g, 50.0 mmol), dry N-methylpyrrolidinone (100 ml), milled potassium carbonate (13.8 g, 100 mmol), potassium iodide (0.84 g, 5.0 mmol), and the product from part (b) (12.0 g, 75.0 mmol) was stirred at 60°C for 18 hours. After cooling to room temperature, the mixture was partitioned between 200 ml water and 100 ml ethyl acetate. The aqueous layer was extracted twice with 50 ml portions of ethyl acetate. The combined organic extracts were washed with water, brine, dried (MgSO₄), and concentrated *in vacuo* to provide 18.3 g of a yellow oil. MPLC (Silica Gel 60 50x460 mm, 25% ethyl acetate in hexanes) provided 12.7 g (94.1%) of pure title compound as a white solid, m.p. 46-48°C (methanol).

d) 3,5-Dimethyl-4-[3-(3-methylisoxazol-5-yl)propyloxy]-N-hydroxybenzenecarboximidamide. (V)

A mixture of the product prepared according to part (c) (18.4 g, 68.1 mmol), absolute ethanol (200 ml), milled potassium carbonate (46.9 g, 0.340 mol), and hydroxylamine hydrochloride (23.6 g, 0.340 mol) was refluxed for 18 hours. The hot mixture was filtered and the solids remaining washed with hot ethanol. The combined filtrates were concentrated *in vacuo* to provide 19.4 g (93.9%) of the title compound as a white powder which was of sufficient purity to be used in subsequent steps. A sample was recrystallized from ethanol to give a white solid, m.p. 129-130.5°C.

e) 5-{3-[2,6-Dimethyl-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-methylisoxazole [I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = CF₃].

To a solution of the product from part (d) (4.38 g, 14.4 mmol) in 8.0 ml dry pyridine was added 4.07 ml (28.8 mmol) trifluoroacetic anhydride at a rate to maintain a gentle reflux. After addition was complete, the mixture was allowed to cool to room temperature, and diluted with water. The solids obtained were washed with water, dried *in vacuo*, and purified by chromatography (Silica Gel 60, 15-40% ethyl acetate in hexanes), to give 4.76 g of pure title compound as a white solid, m.p. 61-62°C.

Example 2

a) 3,5-Difluoro-4-hydroxybenzonitrile. (VII)

A mixture of 4-bromo-2,6-difluorophenol (VI) (4.00 g, 19.0 mmol), copper (I) cyanide (1.72 g, 19.0 mmol), and dimethylformamide (40 ml) was refluxed for 6 hours, cooled to room temperature, diluted with water (150 ml), and filtered. The tan solids obtained were washed with water and retained. The combined filtrates were acidified (1N HCl) and extracted with ethyl acetate. The combined organic extracts were dried (MgSO₄), concentrated *in vacuo*, and purified by flash chromatography (Silica Gel 60, 20% ethyl acetate in hexanes) to give 1.03 g of pure title compound as an off-white solid, mp 195-197°C.

The tan solid was suspended in ethyl acetate with a small amount of acetone, filtered, and concentrated *in vacuo*. The residue obtained was partitioned between ethyl acetate and 1 N HCl. The aqueous phase was extracted with ethyl acetate and the combined organic phases purified as above to provide an additional 0.43 g (49% combined yield) of pure title compound.

The following compounds were prepared by a procedure similar to that of Example 1(c):

ExampleCompound

2b

3,5-Difluoro-4-[3-(3-methylisoxazol-5-yl)propyl-oxy]benzonitrile, (IX) mp 23-24.5°C (ether/-hexanes) - prepared from 3,5-difluoro-4-hydroxybenzonitrile (VII) and the product of Example 1b (VIII); yield 49.1%.

3a

3,5-Dichloro-4-[3-(3-methylisoxazol-5-yl)propyl-oxy]benzonitrile (IX) mp 69.5-70.5°C (methanol) (white solid) - prepared from 3,5-dichloro-4-hydroxybenzonitrile (VII) and the product of Example 1b (VIII); yield 80.7%.

The following compounds were prepared by a procedure similar to that of Example 1d:

ExampleCompound

2c

3,5-Difluoro-4-[3-(3-methylisoxazol-5-yl)-propyl-oxy]-N-hydroxybenzenecarboximidamide (V) mp 122-124°C - prepared from the product of Example 2b (IX); yield 86%. The crude product was purified by suspension in 10% ethanol in chloroform, filtration, concentration in vacuo and trituration of the resulting white solid in cold chloroform.

3b

3,5-Dichloro-4-[3-(3-methylisoxazol-5-yl)propyl-oxy]-N-hydroxybenzenecarboximidamide (V) prepared from the product of Example 3a (IX) (0.5 g). The product (0.78 g), obtained on concentration of the filtrates as an oily solid, was used in the next step.

The following compounds were prepared by a procedure similar to that of Example 1e:

<u>Example</u>	<u>Compound</u>
2d	<p><u>5-{3-[2,6-Difluoro-4-(5-trifluoromethyl-1,2,4-oxa-</u> <u>diazol-3-yl)phenoxy]propyl}-3-methylisoxazole</u> [I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(F)₂, R₄ = CF₃], mp 36-37°C (hexanes) (white solid) - from the product of Example 2c (V) and trifluoroacetic anhydride; yield 44.5%.</p>
3c	<p><u>5-{3-[2,6-Dichloro-4-(5-trifluoromethyl-1,2,4-oxa-</u> <u>diazol-3-yl)phenoxy]propyl}-3-methylisoxazole</u> [I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(Cl)₂, R₄ = CF₃], mp 65-67°C (hexanes) (white solid) - from the product of Example 3b (V) and trifluoroacetic anhydride; yield 80.5%.</p>

- 4 5-{3-[4-(5-Cyclopropyl-1,2,4-oxadiazol-3-yl)-
 2,6-dimethylphenoxy]propyl}-3-methylisoxazole
 [I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂,
 R₄ = cyclopropyl], mp 85-88°C (methanol) (white
 solid) - from the product of Example 1d (V) and
 cyclopropanecarbonyl chloride; yield 71.0%.
- 5 5-{3-[2,6-Dimethyl-4-(5-methoxymethyl-
 1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-methyl-
 isoxazole [I; R₁ = CH₃, Y = (CH₂)₃,
 R₂ and R₃ = 2,6-(CH₃)₂, R₄ = CH₂OCH₃], mp 63-64°C
 (ether/hexane) (white solid) - from the product of
 Example 1d (V) and methoxyacetyl chloride;
 yield 76.1%.
- 6 5-{3-[2,6-Dimethyl-4-(5-fluoromethyl-1,2,4-oxa-
 diazol-3-yl)phenoxy]propyl}-3-methylisoxazole
 [I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂,
 R₄ = CH₂F], mp 80-80.5°C (methanol) (white solid) -
 from the product of Example 1d (V) and fluoro-
 acetyl chloride; yield 45.6%.
- 7 5-{3-[2,6-Dimethyl-4-(5-ethoxycarbonyl-1,2,4-oxa
 diazol-3-yl)phenoxy]propyl}-3-methylisoxazole
 [I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂,
 R₄ = CO₂CH₂CH₃], mp 105-106°C (ethyl acetate/-
 hexane) (white solid) - from the product of
 Example 1d (V) and ethyl oxalyl chloride;
 yield 67.8%.

Example 8

5-{3-[2,6-Dimethyl-4-(5-oxo-4,5-dihydro-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-methylisoxazole [tautomer
 of I where R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = OH].

To a chilled (0°C) suspension of the product from Example 1d (V) (3.03 g, 10.0 mmol), dry acetone (30 ml) and finely divided potassium carbonate (1.52 g, 11 mmol) was added dropwise a solution of ethyl chloroformate (1.05 ml, 11.0 mmol) in acetone (5.5 ml). After stirring at 0°C for 1 hour, the reaction mixture was diluted with water (100 ml) and extracted with methylene chloride (3x25 ml). The combined organic extracts were washed with brine, dried (MgSO₄), filtered through a short column of Florisil, and concentrated *in vacuo* to give the crude intermediate O-acyl derivative as an off-white solid which was then heated at 120-130°C for 45 minutes to give the title compound (2.38 g, 75.4%), mp 194-195°C (methanol) (white needles).

The following compounds were prepared by a procedure similar to that of Example 8:

ExampleCompound

9

5-{3-[2,6-Dimethyl-4-(5-methylcarbonyloxy-methyl-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-methylisoxazole [I; $R_1 = \text{CH}_3$, $Y = (\text{CH}_2)_3$, R_2 and $R_3 = 2,6-(\text{CH}_3)_2$, $R_4 = \text{CH}_2\text{OCOCH}_3$], mp 71-73°C (ether/hexanes) (white solid) - from the product of Example 1d (V) and acetoxyacetyl chloride; yield 71.3%. The crude product was purified by chromatography (Silica Gel 60, 35% ethyl acetate in hexanes).

10

5-{3-[4-(5-Chloromethyl-1,2,4-oxadiazol-3-yl)-2,6-dimethylphenoxy]propyl}-3-methylisoxazole [I; $R_1 = \text{CH}_3$, $Y = (\text{CH}_2)_3$, R_2 and $R_3 = 2,6-(\text{CH}_3)_2$, $R_4 = \text{CH}_2\text{Cl}$], mp 75-76°C (methanol) (white solid) - from the product of Example 1d (V) and chloroacetyl chloride. The crude product was purified by chromatography (Silica Gel 60, 20% ethyl acetate in hexanes); yield 76.2%.

11

5-{3-[2,6-Dimethyl-4-(5-(1-methylcarbonyloxyethyl)-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-methylisoxazole [I; $R_1 = \text{CH}_3$, $Y = (\text{CH}_2)_3$, R_2 and $R_3 = 2,6-(\text{CH}_3)_2$, $R_4 = \text{CH}(\text{CH}_3)\text{OCOCH}_3$], mp 77-77.5°C (white solid) - from the product of Example 1d (V) and 2-acetoxypropionyl chloride; yield 64.6%.

Example 12

5-{3-[2,6-Dimethyl-4-(5-trichloromethyl-1,2,4-oxadiazol-3-yl)-phenoxy]propyl}-3-methylisoxazole [I; $R_1 = \text{CH}_3$, $Y = (\text{CH}_2)_3$, R_2 and $R_3 = 2,6-(\text{CH}_3)_2$, $R_4 = \text{CCl}_3$].

Trichloroacetic acid (22.8 g, 140 mmol) was added to the product of Example 1d (V) (10.6 g, 34.8 mmol) and heated at 85°C until a thick solution was obtained. Trichloroacetyl chloride (14.5 ml, 69.6 mmol) was added in three equal portions. A vigorous reaction ensued after addition of the first portion. The mixture was heated an additional hour at 94°C. The cooled mixture was diluted with water and extracted with ethyl acetate (3x25 ml). The combined organic phases were washed with saturated sodium bicarbonate, brine, dried (MgSO_4) and concentrated *in vacuo* to give 10.1 g of orange oil. Chromatography (Silica Gel 60, methylene chloride) provided 6.94 g of yellow oil which was crystallized from methanol to give 5.03 g of pure title compound as white needles, mp 77-77.5°C.

Example 13

5-{3-[4-(5-Dichloromethyl-1,2,4-oxadiazol-3-yl)-2,6-dimethylphenoxy]propyl}-3-methylisoxazole [I; $R_1 = \text{CH}_3$, $Y = (\text{CH}_2)_3$, R_2 and $R_3 = 2,6-(\text{CH}_3)_2$, $R_4 = \text{CHCl}_2$].

Dichloroacetic acid (1.24 ml, 15.0 mmol) was added to the product of Example 1d (V) (1.14 g, 3.76 mmol) and heated at 85°C until a solution was obtained. Dichloroacetic anhydride (1.14 ml, 7.52 mmol) was added dropwise rapidly and stirred at 85°C for an additional hour. Work-up as described for Example 12 provided 1.51

g of yellow-brown oil which was purified by chromatography (Silica Gel 60, 25% ethyl acetate in hexanes) to give 1.37 g (91.3%) of pure title compound as a pale yellow oil which solidified upon standing, mp 52-3°C (ethanol).

Example 14

5-{3-[4-(5-Difluoromethyl-1,2,4-oxadiazol-3-yl)-2,6-dimethylphenoxy]propyl}-3-methylisoxazole [I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = CHF₂].

Difluoroacetic acid (0.31 ml, 5.0 mmol) was added to a cold (-25°C) solution of N,N'-carbonyldiimidazole (0.80 g, 5.0 mmol) in dry tetrahydrofuran (5.0 ml). After 5 minutes, the resulting suspension was added dropwise rapidly to a solution of the product of Example 1d (V) in dry tetrahydrofuran (20 ml). The mixture was refluxed for 2 hours, cooled, diluted with water, and extracted with ethyl acetate (3x). The combined organic phases were washed with water, brine, dried (MgSO₄), and concentrated *in vacuo* to give 0.78 g of a pale yellow solid. Chromatography (Silica Gel 60, 30% ethyl acetate in hexanes) provided 0.55 g of pure title compound as a pale yellow oil which solidified upon standing, mp 70.5-71°C (methanol).

Example 15

5-{3-[4-(5-Imino-4,5-dihydro-1,2,4-oxadiazol-3-yl)-2,6-dimethylphenoxy]propyl}-3-methylisoxazole [tautomer of I where R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = NH₂].

Cyanogen bromide (1.17 g, 11.0 mmol) was added in portions to a mixture of the product of Example 1d (V) (3.03 g, 10.0 mmol) and potassium bicarbonate (1.10 g, 11.0 mmol) in 50% aqueous ethanol (8.0 ml). After 15 minutes, the thick yellow suspension was diluted with water and filtered. The yellow solid obtained was washed with water and ether to give 1.48 g (45.1%) of pure title compound as a yellow powder, mp 175-183°C.

Example 16

5-{3-[2,6-Dimethyl-4-(5-methoxy-1,2,4-oxadiazol-3-yl)-phenoxy]propyl}-3-methylisoxazole [I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = OCH₃].

The product of Example 12 (I) (627 mg, 1.46 mmol) was added to a freshly prepared solution of sodium methoxide in methanol (1.5 equivalents sodium in 5 ml methanol) in dry dimethylformamide (3-5 ml) and the mixture was stirred at room temperature for 15-30 minutes. The reaction mixture was diluted with water and extracted with ethyl acetate (3x). The combined organic extracts were washed with water, brine, dried (MgSO₄) and concentrated *in vacuo*. The crude residue (0.64 g) was purified by chromatography (Silica Gel 60, first with 2% methanol in methylene chloride followed by 5% ethyl acetate in methylene chloride) to give pure title compound (308 mg) as a colorless oil which crystallized from methanol, mp 64.5-65.5°C (white solid).

Example 17

5-{3-[2,6-Dimethyl-4-(5-ethoxy-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-methylisoxazole [I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = OCH₂CH₃].

Following the procedure of Example 16 but using sodium ethoxide in ethanol in place of sodium methoxide in methanol there was obtained from the product of Example 12 (I) (905 mg, 2.10 mmol) a crude residue (0.82 g) which was purified by chromatography (Silica Gel 60, 2% ethyl acetate in methylene chloride) to give 0.52 g (69%) of pure title compound as a yellow solid, mp 70-72.5°C (ethanol).

Example 18

5-{3-[2,6-Dimethyl-4-(5(methylimino-4,5-dihydro-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-methylisoxazole [tautomer of I where R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = NHCH₃].

The product of Example 12 (I) (1.00 g, 2.32 mmol) was added to 5 ml 40% aqueous methylamine in dimethylformamide (3-5 ml) and the mixture was stirred at room temperature for 18 hours. The reaction mixture was diluted with water and extracted with ethyl acetate (3x). The combined organic extracts were washed with water, brine, dried (MgSO₄) and concentrated *in vacuo*. The crude residue (0.54 g) was purified by chromatography (Silica Gel 60, first with 2% methanol in methylene chloride and then with 50% ethyl acetate in hexanes) to give 300 mg (37.5%) of pure title compound as a yellow solid, mp 126.5-127°C (ethanol).

Example 19

5 5-{3-[2,6-Dimethyl-4-(5-dimethylamino-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-methylisoxazole [I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = N(CH₃)₂].

Following the procedure of Example 18 but using 40% aqueous dimethylamine in place of 40% aqueous methylamine and reducing the reaction time to 15-30 minutes, there was obtained from the product of Example 12 (I) (0.97 g, 2.2 mmol) a crude residue (0.75 g) which was purified by chromatography (Silica Gel 60, 50% ethyl acetate/hexanes) to give 0.70 g (84%) of pure title compound as a pale yellow solid, mp 123-124°C (ethanol).

Example 20

a) 3,5-Dimethyl-4-(3-ethynylpropoxy)benzonitrile. (XII)

Following the procedure of Example 1c and using 14.7 g (100 mmol) 3,5-dimethyl-4-hydroxybenzonitrile (VII) and substituting 5-chloro-1-pentyne (XI) (12.7 ml 120 mmol) for the product of Example 1b (VIII), there was obtained a red-brown oil which was purified by chromatography (Silica Gel 60, 15% ethyl acetate in hexanes) to give pure title compound (21.2 g, 99.4%) as a pale yellow oil.

b) 3,5-Dimethyl-4-(3-ethynylpropoxy)-N-hydroxybenzenecarboximidamide. (XIII)

Following the procedure of Example 1d and using 13.0 g (61.0 mmol) of the product from part (a), there was obtained the title compound (14.9 g, 99.3%) as a white solid which was sufficiently pure for use in the next step.

c) 3-[3,5-Dimethyl-4-(3-ethynylpropoxy)phenyl]-5-trifluoromethyl-1,2,4-oxadiazole. (III)

Following the procedure of Example 1e and using 7.40 g (30.0 mmol) of the product of part (b), 9.0 ml dry pyridine and 8.50 ml trifluoroacetic anhydride there was obtained pure title compound (6.42 g, 65.9%) as a pale yellow oil which crystallized from methanol to give the title compound as a white solid, mp 45.5-48°C.

Procedure 1 - General procedure for preparing the compounds of Examples 21, 22, 23, 28a and 29a below

To a solution of N-chlorosuccinimide (NCS, 1.8-2.5 equivalents) in dry N,N-dimethylformamide or N-methylpyrrolidinone (1.6-3.0 ml per mmol NCS) and 1-2 drops of pyridine was added dropwise a solution of oxime (1.8-2.5 equivalents) in the same solvent (0.40-0.80 ml per mmol oxime). The internal temperature was maintained at 25-30°C with a 25°C water bath. After 1 hour at room temperature, a solution of the appropriate ethynyl compound (formula III or XII) (1 equivalent) in the same solvent (0.80 ml per mmol the ethynyl compound) was added. The reaction mixture was heated to 85-90°C and a solution of triethylamine (TEA, 1.8-2.5 equivalents) in the same solvent (0.80-1.6 ml per mmol TEA) was added dropwise over 45-90 minutes. After an additional hour at 85-90°C, the mixture was cooled to room temperature, diluted with water, and extracted with ethyl acetate (3x). The combined organic phases were washed with 10% KHSO₄, water, brine, dried (MgSO₄ or Na₂SO₄) and concentrated in vacuo. The crude product was purified by chromatography (Silica Gel 60, 15-40% ethyl acetate in hexanes).

The following compounds were prepared by Procedure 1:

Example**Compound**

21

5-{3-[2,6-Dimethyl-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-(methoxymethyl)isoxazole [I; $R_1 = \text{CH}_2\text{OCH}_3$, $Y = (\text{CH}_2)_3$, R_2 and $R_3 = 2,6-(\text{CH}_3)_2$, $R_4 = \text{CF}_3$], colorless oil (yield 70.1%) - from the product of Example 20c (III) (2.00 g, 6.17 mmol) and methoxyacetaldehyde oxime (1.10 g, 12.3 mmol).

22

5-{3-[2,6-Dimethyl-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-(ethoxymethyl)isoxazole [I; $R_1 = \text{CH}_2\text{OCH}_2\text{CH}_3$, $Y = (\text{CH}_2)_3$, R_2 and $R_3 = 2,6-(\text{CH}_3)_2$, $R_4 = \text{CF}_3$], mp 24-25°C (methanol) (white powder) (yield 35.3%) - from the product of Example 20c (III) (2.00 g, 6.17 mmol) and 2-ethoxyacetaldehyde oxime (1.27 g, 12.3 mmol).

23

3-Cyclopropyl-5-{3-[2,6-dimethyl-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-isoxazole [I; $R_1 = \text{cyclopropyl}$, $Y = (\text{CH}_2)_3$, R_2 and $R_3 = 2,6-(\text{CH}_3)_2$, $R_4 = \text{CF}_3$], mp 63.5-65°C (ethanol) (white needles) (yield 82%) - from the product of Example 20c (III) (0.92 g, 2.8 mmol) and cyclopropylcarboxaldehyde oxime (0.48 g, 5.6 mmol).

2-Ethoxyacetaldehyde oxime (used in Example 22 above)

A solution of hydroxylamine hydrochloride (18.8 g, 0.270 mol), ethanol (25 ml), water (40 ml) and 1,1,2-triethoxyethane was warmed at 45°C for 30 minutes, cooled to room temperature, and extracted with ether (3x). The combined organic phases were dried (MgSO_4), concentrated in vacuo, and filtered through a small plug of cotton to give 10.1 g of title compound as a pale yellow oil which was used as is.

Example 245-Cyclopropyl-3-[3,5-dimethyl-4-(3-ethynylpropoxy)phenyl]-1,2,4-oxadiazole. (III)

Following the procedure of Example 1e and using 5.00 g (20.3 mmol) of the product of Example 20b (XIII), 75 ml dry pyridine and 2.77 ml (30.5 mmol) cyclopropylcarbonyl chloride there was obtained pure title compound (3.98 g, 66.2%) as a nearly colorless oil which solidified on standing, mp 45-46°C (methanol).

Procedure 2 - general procedure for preparing the compounds of Examples 25, 26 and 27 below

To a chilled (0°C) solution of the appropriate aldehyde oxime (2.5 equivalents) in dry dimethylformamide (DMF) (15 ml) was added in 1 portion N-chloro-succinimide (NCS) (2.5 equivalents). After 1-2 hours, the product from Example 24 (1 equivalent) was added and the whole heated to 80°C. A solution of triethylamine (2.5 equivalents) in dry DMF (5 ml) was added dropwise over 90 minutes. The mixture was heated an additional 18 hours. Work-up and purification as described for Example 21 provided the pure product.

The following compounds were prepared by Procedure 2:

ExampleCompound

25

5-{3-[5-(Cyclopropyl-1,2,4-oxadiazol-3-yl)-2,6-dimethylphenoxy]propyl}-3-ethylisoxazole
 [I; R₁ = CH₂CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = cyclopropyl], colorless oil - from the product of Example 24 (III) and propionaldehyde oxime; yield 67%.

26

5-{3-[5-(Cyclopropyl-1,2,4-oxadiazol-3-yl)-2,6-dimethylphenoxy]propyl}-3-(methoxymethyl)-isoxazole [I; R₁ = CH₂OCH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = cyclopropyl], mp 44-45°C (methanol) (white solid) - from the product of Example 24 (III) and methoxy-acetaldehyde oxime; yield 26.1% (from combination of two runs).

27

5-{3-[5-(Cyclopropyl-1,2,4-oxadiazol-3-yl)-2,6-dimethylphenoxy]propyl}-3-cyclopropylisoxazole
 [I; R₁ = R₄ = cyclopropyl, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂], mp 59-60°C (methanol) (white solid) - from the product of Example 24 (III) and cyclopropylcarboxaldehyde oxime; yield 60.4%.

Example 28

a) 3,5-Dimethyl-4-[3-(3-ethylisoxazol-5-yl)propyloxy]benzonitrile. (IX)

Following Procedure 1 above but omitting the pyridine and using propionaldehyde oxime (8.6 g, 118 mmol) and the product of Example 20a (XII) (10.1 g, 47.0 mmol) there was obtained 4.90 g (36.7%) of pure title compound, mp 53.5-54.5°C (ethanol).

b) 3,5-Dimethyl-4-[3-(3-ethylisoxazol-5-yl)propyloxy]N-hydroxybenzenecarboximidamide. (V)

A mixture of the product from part (a) (2.01 g, 7.50 mmol), ethanol (20 ml), hydroxylamine hydrochloride (2.61 g, 37.5 mmol), and finely divided potassium carbonate (5.20 g, 37.5 mmol) was refluxed for 18 hours. The mixture was filtered hot, the filter cake washed with ethanol, and the combined filtrates concentrated in vacuo to give 2.57 g of crude title compound as a pasty yellow solid, which was used as such in the next step.

c) 5-{3-[2,6-Dimethyl-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-ethylisoxazole [I; R₁ = CH₂CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = CF₃].

All the product from part (b) was dissolved in pyridine (2.3 ml) and trifluoroacetic anhydride (2.1 ml, 15 mmol) was added dropwise. The mixture was refluxed for 1 hour, cooled to room temperature, diluted with water, and extracted with methylene chloride (3x). The combined organic phases were washed with 1N HCl, water, brine, dried (MgSO₄), and concentrated in vacuo. The pale yellow oil obtained (2.15 g) was chromatographed Silica Gel 60, methylene chloride) to give 2.10 g (70.7%) of pure title compound as a white solid, mp 157-158°C (methanol).

Example 29

a) 3,5-Dimethyl-4-{3-[3-(2-methoxyethyl)isoxazol-5-yl]propyloxy}benzonitrile. (IX)

Following Procedure 1 above and using 3-methoxypropionaldehyde oxime (1.94 g, 18.8 mmol) and the product of Example 20a (XII) (2.20 g, 10.3 mmol) there was obtained 1.51 g (46.5%) of pure title compound as a colorless oil which crystallized from ethanol as fine white needles, mp 64-64.5°C. There was recovered 0.89 g (40.4%) of starting product of Example 20a.

b) 5-{3-[2,6-Dimethyl-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-(methoxyethyl)isoxazole

[I; R₁ = CH₂CH₂OCH₃, R₂ and R₃ = 2,6-(CH₃)₂, Y = (CH₂)₃, R₄ = CF₃].

Sodium (442 mg, 19.2 mg-atom) was dissolved in dry methanol (20 ml) contained in an addition funnel. This solution was added dropwise to a solution of hydroxylamine hydrochloride (1.34 g, 19.2 mmol) in dry methanol (10 ml). A fine white precipitate formed. After 1 hour, a solution of the product from part (a) (1.21 g, 3.85 mmol) in dry methanol (5 ml) was added and the mixture heated at reflux for 2.5 hours. The hot reaction mixture was filtered, the filter cake washed with methanol, and the combined filtrates concentrated in vacuo. The white oily solid obtained was dissolved in pyridine (4 ml) and trifluoroacetic anhydride (1.63 ml, 11.6 mmol) was added at a rate to maintain a gentle reflux. The mixture was heated at reflux for an additional 30 minutes, cooled to room temperature, diluted with water, and extracted with ethyl acetate (3x). The combined organic phases were washed with 10% KHSO₄, water, brine, dried (MgSO₄), and concentrated in vacuo to give 2.27 g of yellow oil. Chromatography (Silica Gel 60, 30% ethyl acetate in hexanes) provided 1.28 g (78.0%) of pure title compound as a colorless oil. Crystallization from methanol gave a white solid, mp 36.5-37°C.

3-Methoxypropionaldehyde oxime (used in Example 29a above)

To a solution of hydroxylamine hydrochloride (2.80 g, 40.2 mmol), 10% aqueous sodium acetate (4.0 ml) and water (6 ml) was added 1,1,3-trimethoxypropane (2.12 ml, 14.9 mmol) and heated at 40-50°C for 30 minutes. After cooling to room temperature, the solution was saturated with sodium chloride and extracted with ether (3x) and methylene chloride (3x). The combined organic phases were dried (MgSO₄), filtered through a pad of Florisil, and concentrated in vacuo to provide 1.6 g of title compound as a colorless oil which was used as is.

Procedure 3 - General procedure for the preparation of the compounds of Example 30a and b, 31a and b, and 32a and b

A mixture of the appropriate 4-hydroxybenzonitrile (VII) (1 equivalent), dry ethanol (3.7-8.9 ml per mmol of the 4-hydroxybenzonitrile), hydroxylamine hydrochloride (5 equivalents), and finely divided potassium carbonate (5 equivalents) was refluxed with efficient stirring for 18 hours. The hot reaction mixture was filtered and the filter cake washed with ethanol. The combined filtrates were concentrated in vacuo to give the crude amidoximes (XIV) which were dissolved into pyridine (1-2 ml per mmol of the 4-hydroxybenzonitrile). Trifluoroacetic anhydride (5 equivalents) was added at a rate to maintain a gentle reflux. After heating an additional 0.5-3 hours, the cooled reaction mixture was diluted with ethyl acetate and water (4:1) until homogeneous. The organic phase was extracted with cold 1N KOH (3x). The basic extracts were acidified with concentrated HCl and extracted with ethyl acetate (3x). The combined organic phases were washed with brine, dried (MgSO₄), concentrated in vacuo. Chromatography (Silica Gel 60, ethyl acetate in hexanes or Florisil, methylene chloride) provided the pure 4-hydroxyphenyl-5-trifluoromethyl-1,2,4-oxadiazole.

Following Procedure 3 there were prepared the following crude intermediate amidoximes and corresponding 4-hydroxy-5-trifluoromethyl-1,2,4-oxadiazoles:

Example	Compound
30a	<u>3,5-Dimethyl-4,N-dihydroxybenzene-carboximidamide (XIV)- from 3,5-dimethyl-4-hydroxybenzonitrile.</u>
30b	<u>3-(3,5-Dimethyl-4-hydroxyphenyl)-5-trifluoromethyl-1,2,4-oxadiazole, (IV) mp 114-115°C (hexane) (white needles) - from the product of Example 30a; yield 75.2%.</u>
31a	<u>3,5-Dichloro-4,N-dihydroxybenzene-carboximidamide (XIV)- from 3,5-dichloro-4-hydroxybenzonitrile.</u>
31b	<u>3-(3,5-Dichloro-4-hydroxyphenyl)-5-trifluoromethyl-1,2,4-oxadiazole, (IV) mp 96-98°C (hexane) (white needles) - from the product of Example 31a; yield 52.0%.</u>
32a	<u>4,N-Dihydroxybenzenecarboximidamide (XIV)- from 4-hydroxybenzonitrile.</u>
32b	<u>3-(4-Hydroxyphenyl)-5-trifluoromethyl-1,2,4-oxadiazole, (IV) mp 74-75°C (hexanes) (white needles) - from the product of Example 32a; yield 56.4%.</u>

Example 30c

5-{5-[2,6-Dimethyl-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)-phenoxy]pentyl}-3-methylisoxazole [I; R₁ = CH₃, Y = (CH₂)₅, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = CF₃].

Following a procedure similar to that of Example 1c but substituting the product from Example 30b (IV) (1.0 g, 3.9 mmol) for 3,5 -dimethyl-4-hydroxybenzonitrile and 5-(3-methylisoxazol-5-yl)pentyl bromide (VIII) (1.0 g, 4.3 mmol) for 3-(3-methylisoxazol-5-yl)propyl chloride and using 0.72 g (4.3 mmol) of potassium iodide there was obtained 0.25 g (16%) of pure title compound as a white solid, mp 41-42°C (methanol).

Example 31c

5-{5-[2,6-Dichloro-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)phenoxy]pentyl}-3-methylisoxazole [I; R₁ = CH₃, Y = (CH₂)₅, R₂ and R₃ = 2,6-(Cl)₂, R₄ = CF₃].

Following a procedure similar to that of Example 1c but substituting the product from Example 31b (IV) (0.93 g, 3.1 mmol) for 3,5-dimethyl-4-hydroxybenzonitrile and 5-(3-methylisoxazol-5-yl)pentyl bromide (VIII) (1.0 g, 4.3 mmol) for 3-(3-methylisoxazol-5-yl)propyl chloride and using 0.72 g (4.3 mmol) of potassium iodide there was obtained 0.83 g (60%) of pure title compound as a white solid, mp 42-43°C (hexanes).

Example 32c

3-Methyl-5-{3-[4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)phenoxy]propyl}isoxazole [I; R₁ = CH₃, Y = (CH₂)₃, R₂ = R₃ = H, R₄ = CF₃].

Following a procedure similar to that of Example 1c but substituting the product of Example 32b (IV) (0.42 g, 1.8 mmol) for 3,5-dimethyl-4-hydroxybenzonitrile and using 0.63 g (4.0 mmol) of the product of Example 1b (VIII) and 0.67 g (4.0 mmol) of potassium iodide there was obtained, after trituration in cold methanol, 0.48 g (76%) of pure title compound as a white powder, mp 68-69°C (methylene chloride-hexanes).

Example 33

5 5-{3-[2,6-Dimethyl-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)-phenoxy]propyl}-3-(2-hydroxyethyl)isoxazole
 [I; R₁ = CH₂CH₂OH, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = CF₃].

A solution of the product of Example 29b (I) (1.28 g, 3.00 mmol), dry 1,2-dichloroethane (9 ml), and trimethylsilyl iodide (1.71 ml, 12.0 mmol) was refluxed for 4 hours. To the cooled reaction mixture was added methanol (8 ml). The mixture was diluted with water and extracted with ethyl acetate (3x). The combined organic phases were washed with 10% NaHSO₃, saturated NaHCO₃, brine, dried (MgSO₄), and concentrated *in vacuo*.
 10 Chromatography (Silica Gel 60, 50% ethyl acetate in hexanes) provided 1.11 g (90.2%) of pure title compound as a colorless oil which solidified upon standing, mp 74.5-75°C (methanol) (white solid).

Example 34

15 a) 3-(tert-Butyldimethylsilyloxymethyl)-5-methylisoxazole. (XVIII)

To a chilled (5°C) solution of 3-hydroxymethyl-5-methylisoxazole (X) (16.8 g, 148 mmol) and tert-butyldimethylsilyl chloride (24.6 g, 163 mmol) in dry methylene chloride (100 ml) was added over 15 minutes a solution of triethylamine (22.7 ml, 163 mmol) in methylene chloride (25 ml). 4-Dimethylaminopyridine (1.81 g, 14.8 mmol) was added and the thick reaction mixture was stirred at room temperature for 48 hours.
 20 Water (100 ml) was added and the aqueous layer extracted with methylene chloride (3x). The combined organic phases were washed with brine, dried (MgSO₄), filtered through a pad composed of a layer of Florisil and a layer of Silica Gel 60, and concentrated *in vacuo*. The yellow oil obtained (36.6 g) was purified by chromatography (Silica Gel 60, 2% ethyl acetate in hexanes) to give 27.7 g (81.9%) of pure title compound as a pale yellow oil.

25 b) 3-[3-(tert-Butyldimethylsilyloxymethyl)isoxazol-5-yl]propyl alcohol. (XV)

To a cold (-78°C) solution of the product from part (a) (13.0 g, 57.0 mmol) and N,N,N',N'-tetramethylethylenediamine (1.2 ml, 7.9 mmol) in dry tetrahydrofuran (THF) (150 ml) was added over 5 minutes n-butyllithium (31.3 ml, 2.0 M in hexane). The bright orange-yellow anion solution was stirred for 25 minutes. Ethylene oxide (50.0 ml of 7.6 M solution in dry THF) was added over 10 minutes. After 1.5 hours, saturated
 30 NH₄Cl (30 ml) was added. The mixture was allowed to warm to room temperature and diluted with water. The aqueous layer was extracted with ethyl acetate (3x). The combined organic phases were washed with brine, dried (MgSO₄), filtered through a pad of Silica Gel 60, and concentrated *in vacuo*. Chromatography (Silica Gel 60, 20% ethyl acetate in hexanes) gave 3.44 g of recovered product from part (a) and 8.18 g (52.7%) of pure title compound as a colorless oil.

35 c) 3-(tert-Butyldimethylsilyloxymethyl)-5-{3-[2,6-dimethyl-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)phenoxy]propyl}isoxazole. (XVII)

A solution of the product from part (b) (XV) (1.00 g, 3.67 mmol), the product of Example 30b (IV) (1.04 g, 4.04 mmol), and triphenylphosphine (1.06 g, 4.04 mmol) in dry tetrahydrofuran (THF) (10 ml) was chilled to 0°C. A solution of diethyl azodicarboxylate (DEAD) (0.61 ml, 1.04 mmol) in dry THF (15 ml) was added dropwise over 20 minutes. The solution was stirred for 30 minutes at 0°C and 18 hours at room temperature, diluted with water, and extracted with ethyl acetate (2x). The combined organic phases were washed with
 40 10% NaOH, brine, dried (MgSO₄), filtered through a pad of Silica Gel 60, and concentrated *in vacuo* to give 3.44 g of yellow oil. Chromatography (Silica Gel 60, 10% ethyl acetate in hexanes) provided 1.73 g (83.6%) of pure title compound as a colorless oil.

45 d) 5-{3-[2,6-Dimethyl-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-(hydroxymethyl)isoxazole

[I; R₁ = CH₂OH, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = CF₃].

A solution of the product from part (c) (XVII) (0.75 g, 1.5 mmol), tetrahydrofuran (60 ml), and 1N HCl (7.5 ml) was stirred at room temperature for 18 hours and diluted with water (100 ml). The pH was adjusted to pH 7 (pH paper) with solid NaHCO₃ and extracted with ethyl acetate (3x). The combined organic phases were washed with brine, dried (MgSO₄), and concentrated *in vacuo* to give 0.73 g of yellow oil which was purified by chromatography (Silica Gel 60, 50% ethyl acetate in hexanes) to provide 0.58 g (100%) of pure
 50 title compound as a white solid, mp 92-3°C (white needles from ethanol).

Example 35

55 a) 3,5-Dimethyl-4-{3-[3-(tert-butyldimethylsilyloxymethyl)-isoxazol-5-yl]propyloxy}benzonitrile. (XIX)

To a chilled (0°C) methylene chloride (25 ml) solution of 3,5-dimethyl-4-hydroxybenzonitrile (VII) (773 mg, 5.26 mmol), the product from Example 34b (XV) (1.43 g, 5.26 mmol), and triphenylphosphine (1.38 g,

5.26 mmol) was added dropwise over 30 minutes a solution of diethyl azodicarboxylate (DEAD) (915 mg, 5.26 mmol) in methylene chloride (5 ml). The solution was stirred at 0°C for 30 minutes and at room temperature for 18 hours, after which it was washed with water, 2.5 M NaOH, brine, dried (Na₂SO₄) and concentrated *in vacuo*. The residue was triturated in ether to remove the bulk of the triphenylphosphine oxide, the filtrate concentrated *in vacuo*, and the residue purified by chromatography (Silica Gel 60, 15% ethyl acetate in hexanes) to give 1.73 g (82.2%) of pure title compound as a colorless oil.

b) 3,5-Dimethyl-4-{3-[3-(tert-butyl)dimethylsilyloxymethyl]isoxazol-5-yl}propyloxy}-N-hydroxybenzene-carboximidamide. (XX)

A mixture of the product from part (a) XIX (1.22 g, 3.05 mmol), ethanol (30 ml), hydroxylamine hydrochloride (1.06 g, 15.2 mmol), and finely divided potassium carbonate (2.10 g, 15.2 mmol) was refluxed for 5 hours and filtered. The filter cake was washed with ethanol and the combined filtrates concentrated *in vacuo* to give 1.30 g of white solid. A portion of this material (0.78 g) was purified by chromatography (reverse phase silica gel, 17% water in methanol) to give 0.47 g of title compound which contained approximately 5 % (NMR analysis) of desilylated material.

c) 5-{3-[4-(5-Cyclopropyl-1,2,4-oxadiazol-3-yl)-2,6-dimethyl-phenoxy]propyl}-3-(hydroxymethyl)isoxazole [I; R₁ = CH₂OH, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = cyclopropyl].

To a solution of the purified product from part (b) (XX) (0.47 g, 1.1 mmol) in pyridine (20 ml) was added cyclopropylcarbonyl chloride (0.15 ml, 1.6 mmol). The mixture was heated at 90°C for 26 hours. The pyridine was removed *in vacuo* and the residue partitioned between water and ethyl acetate. The aqueous phase was extracted with ethyl acetate (3x). The combined organic phases were washed with 3N HCl (2x), brine, dried (Na₂SO₄), and concentrated *in vacuo* to yield 0.61 g of a yellow oil. Chromatography (Silica Gel 60, 35% ethyl acetate in hexanes) provided 0.25 g (62%) of pure title compound as a colorless oil. Crystallization from methylene chloride and hexanes provided the title compound as a white solid, mp 80-1°C.

Example 36

a) 5-Cyclopropyl-3-[4-(5-ethoxycarbonyl-4-pentynyloxy)-3,5-dimethylphenyl]-1,2,4-oxadiazole. (XXI)

To a cold (-78°C) dry tetrahydrofuran solution (20 ml) of the product from Example 24 (III) (1.30 g, 4.41 mmol) was added dropwise n-butyllithium (2.30 ml, 2.3M in hexane) over 15 minutes. After an additional 30 minutes at -78°C, ethyl chloroformate (0.63 ml, 6.6 mmol) was added and the mixture warmed gradually to 0°C over 2 hours. The reaction was quenched with saturated NH₄Cl and extracted with ethyl acetate (3x). The combined organic phases were washed with brine, dried (Na₂SO₄), and concentrated *in vacuo* to give a colorless oil (2.05 g). Chromatography (Silica Gel 60, 10-20% ethyl acetate in hexanes) provided 1.38 g (85.0%) of pure title compound as a colorless oil.

b) 5-{3-[4-(5-Cyclopropyl-1,2,4-oxadiazol-3-yl)-2,6-dimethylphenoxy]propyl}-3-hydroxyisoxazole [I; R₁ = OH, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = cyclopropyl].

A mixture of the product from part (a) XXI (810 mg, 2.20 mmol), ethanol (15 ml), hydroxylamine hydrochloride (400 mg, 5.76 mmol), and 10% NaOH (5 ml) was stirred at room temperature for 24 hours (after 8 hours, a solution was obtained). Water (6 ml) was added, the mixture acidified with concentrated HCl to pH 2 (pH paper), and extracted with ether (4x). The combined organic phases were washed with brine, dried (Na₂SO₄), and concentrated *in vacuo* to a white solid. Chromatography (Silica Gel 60, 50% ethyl acetate in hexanes) provided 0.55 g (70%) of pure title compound as a white solid, mp 155-6°C (ethyl acetate and hexanes).

Example 37

5-{3-[4-(5-Cyclopropyl-1,2,4-oxadiazol-3-yl)-2,6-dimethylphenoxy]propyl}-3-ethoxyisoxazole [I; R₁ = OCH₂CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = cyclopropyl].

A mixture of the product of Example 36b (I) (0.30 g, 0.85 mmol), dry acetone (25 ml), finely divided potassium carbonate (0.24 g, 1.7 mmol), and ethyl iodide (0.18 ml, 2.2 mmol) was heated at 50°C for 18 hours, filtered, and concentrated *in vacuo* to give a pinkish solid. Chromatography (Silica Gel, 50% ethyl acetate in hexanes) provided 0.19 g of slightly impure title compound and 0.12 g (37%) of a pure side product (the corresponding 2,3-dihydro-2-ethyl-3-oxoisoxazole compound) as a colorless oil. Pure title compound was obtained by chromatography (reverse silica gel, 20% water in methanol); yield 0.14 g (43%), mp 70-1°C (methanol).

Example 385-{3-[4-(5-Aminocarbonyl-1,2,4-oxadiazol-3-yl)-2,6-dimethylphenoxy]propyl}-3-methylisoxazole

[I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = CONH₂].

Finely divided product of Example 7 (I) (3.08 g, 8.00 mmol) was added to 10% ethanolic ammonia (80 ml). After 15 minutes, a solution was obtained and a fine precipitate started to form. After 4 hours, the mixture was filtered and the solids obtained washed with cold ethanol to give 2.35 g (82.5%) of pure title compound as a fine white powder, mp 177-8°C (isopropyl acetate).

Example 395-{3-[4-(5-Cyano-1,2,4-oxadiazol-3-yl)-2,6-dimethylphenoxy]propyl}-3-methylisoxazole

[I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = CN].

To a chilled (0°C) suspension of the product of Example 38 (I) (1.60g, 4.50 mmol) and dry pyridine (11.2 ml) in dry tetrahydrofuran (27 ml) was added trifluoroacetic anhydride (1.90 ml, 13.5 mmol). The mixture was stirred at 0°C for 4 hours and at room temperature for 18 hours, diluted with water (100 ml), and extracted with ethyl acetate (2x25 ml). The combined organic phases were washed with 1N HCl (3x), brine, dried (MgSO₄), and concentrated in vacuo. The red solid obtained (1.67 g) was chromatographed (Silica Gel 60, 20% ethyl acetate in hexanes) to give 1.38 g (90.8%) of pure title compound as a white solid, mp 93-4°C (ethyl acetate and hexanes).

Example 405-{3-[2,6-Dimethyl-4-(5-(hydroxymethyl)-1,3,4-oxadiazol-3-yl)phenoxy]propyl}-3-methylisoxazole

[I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = CH₂OH].

A mixture of the product of Example 9 (I) (4.12 g, 10.7 mmol) and finely divided potassium carbonate (1.48 g, 10.7 mmol) in dry methanol (40 ml) was stirred at room temperature for 15 minutes and partitioned between water (50 ml) and ethyl acetate (50 ml). The aqueous phase was extracted with ethyl acetate (1x25 ml) and the combined organic phases washed with brine, dried (MgSO₄), and concentrated in vacuo. Chromatography (Silica Gel 60, 50% ethyl acetate in hexanes) provided 3.35 g (91.2%) of pure title compound as a white solid, mp 116.5-117°C (ether).

Example 415-{3-[2,6-Dimethyl-4-(5-(iodomethyl)-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-methylisoxazole

[I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = CH₂I].

A solution of sodium iodide (0.45 g, 3.0 mmol) in dry acetone (5 ml) was added dropwise to a solution of the product of Example 10 (I) (905 mg, 2.50 mmol) in dry acetone (5 ml). After 4 hours, the yellow suspension was poured into water (50 ml) and extracted with methylene chloride (3x25 ml). The combined organic phases were washed with brine, dried (MgSO₄), and concentrated in vacuo to give a brown oil (1.56 g). Filtration through Florisil (methylene chloride) provided a green-yellow oil (1.43 g) which solidified upon standing at 0°C. Chromatography (Silica Gel 60, 25% ethyl acetate in hexanes) provided 1.06 g (93.8%) of pure title compound as a pale yellow solid, mp 89-90°C (white needles from ether-pentane).

Example 425-{3-[2,6-Dimethyl-4-(5-(4-methylphenylsulfonyloxymethyl)-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-methylisoxazole

[I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂, R₄ = 4-CH₃C₆H₄SO₂OCH₂].

To a chilled (0°C) mixture of the product of Example 40 (I) (343 mg, 1.00 mmol) and finely divided potassium carbonate (0.28 g, 2.0 mmol) in dry methylene chloride (5 ml) was added dropwise a filtered solution of (4-methylphenyl)sulfonyl chloride (0.23 g, 1.2 mmol) in methylene chloride (2 ml). The mixture was stirred at room temperature for 72 hours, after which an additional 0.40 mmol of potassium carbonate and (4-methylphenyl)sulfonyl chloride was added. After 24 hours, the mixture was partitioned between water (10 ml) and ethyl acetate (10 ml). The organic phase was washed with 1N NaOH (1x5 ml), brine, dried (Mg SO₄), and concentrated in vacuo. Chromatography (Silica Gel 60, 40% ethyl acetate in hexanes) provided 478 mg (96.1%) of pure title compound as a white solid, mp 97-8°C (ether).

Example 43

5-{3-[2,6-Dimethyl-4-(5-(2,2,2-trifluoroethyl)-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-methylisoxazole
 [I; $R_1 = \text{CH}_3$, $Y = (\text{CH}_2)_3$, R_2 and $R_3 = 2,6-(\text{CH}_3)_2$, $R_4 = \text{CH}_2\text{CF}_3$].

A mixture of the product of Example 1d (V) (4.55 g, 15.0 mmol), dry tetrahydrofuran (45 ml), 2-trifluoroethylidene-1,3-dithiane (3.60 g, 18.0 mmol), and silver trifluoroacetate (7.3 g, 33 mmol) was refluxed in the dark for 22 hours, cooled to room temperature, and filtered. The green filter cake was washed with ethyl acetate (4x20 ml). The combined filtrates were concentrated *in vacuo*. The residue obtained was dissolved in methylene chloride (50 ml) and washed with water (3x25 ml), 0.1 M NaHCO_3 (freshly prepared, 25 ml), brine, dried (MgSO_4), filtered through a pad of Florisil, and concentrated *in vacuo* to give 5.39 g of a yellow paste. Purification by chromatography (Silica Gel 60, 15% ethyl acetate in hexanes) provided 2.22 g (37.5%) of pure title compound as a white solid, mp 84-85°C (methanol) (white plates).

Example 44

5-{3-[4-(5-Cyclopropyl-1,2,4-oxadiazol-3-yl)-2,6-dimethylphenoxy]propyl}-3-(2-hydroxyethoxy)isoxazole
 [I; $R_1 = \text{HOCH}_2\text{CH}_2\text{O}$, $Y = (\text{CH}_2)_3$, R_2 and $R_3 = 2,6-(\text{CH}_3)_2$, $R_4 = \text{cyclopropyl}$].

A mixture of the product of Example 36b (I) (0.75 g, 2.1 mmol), dry acetone (25 ml), finely divided potassium carbonate (0.32 g, 2.3 mmol), and 2-bromoethanol (0.19 ml, 2.7 mmol) was refluxed for 5 hours, filtered, and concentrated *in vacuo* to give a pinkish oil. Chromatography (Silica Gel 60, 50% ethyl acetate in hexanes) provided 0.51 g of impure title compound and 0.48 g (57%) of a pure side product (the corresponding 2,3-dihydro-2-(2-hydroxyethyl)-3-oxoisoxazole compound) as a white solid. Pure title compound (0.31 g, 37%) was obtained by gradient chromatography (Silica Gel 60, hexanes to 50% ethyl acetate in hexanes), mp 64-65 °C (methylene chloride and hexanes).

Following a procedure similar to that of Example 1c but substituting for 3,5-dimethyl-4-hydroxybenzonitrile (VII) an equivalent amount of the following:

- 4-hydroxy-3-nitrobenzonitrile
- 4-hydroxy-3,5-dimethoxybenzonitrile
- 4-hydroxy-3-trifluoromethylbenzonitrile

there can be obtained respectively the following compounds of formula IX:

- 4-[3-(3-methylisoxazol-5-yl)propyloxy]-3-nitrobenzonitrile
- 3,5-dimethoxy-4-[3-(3-methylisoxazol-5-yl)propyloxy]benzonitrile
- 4-[3-(3-methylisoxazol-5-yl)propyloxy]-3-(trifluoromethyl)benzonitrile.

Following a procedure similar to that of Example 1d but substituting for the product from Example 1c (IX) an equivalent amount of the above compounds of formula IX there can be obtained respectively the following compounds of formula V:

- 4-[3-(3-methylisoxazol-5-yl)propyloxy]-3-nitro-N-hydroxybenzenecarboximidamide
- 3,5-dimethoxy-4-[3-(3-methylisoxazol-5-yl)propyloxy]-N-hydroxybenzenecarboximidamide
- 4-[3-(3-methylisoxazol-5-yl)propyloxy]-3-trifluoromethyl-N-hydroxybenzenecarboximidamide.

Following a procedure similar to that of Example 1e but substituting for the product of Example 1d (V) an equivalent amount of the above compounds of formula V there can be obtained respectively the following compounds of formula I:

- 3-methyl-5-{3-[2-nitro-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)phenoxy]propyl}isoxazole

[I; $R_1 = \text{CH}_3$, $Y = (\text{CH}_2)_3$, $R_2 = 2-\text{NO}_2$, $R_3 = \text{H}$, $R_4 = \text{CF}_3$]

- 5-{3-[2,6-dimethoxy-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-methylisoxazole

[I; $R_1 = \text{CH}_3$, $Y = (\text{CH}_2)_3$, R_2 and $R_3 = 2,6-(\text{OCH}_3)_2$, $R_4 = \text{CF}_3$]

- 3-methyl-5-{3-[2-trifluoromethyl-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)phenoxy]propyl}isoxazole

[I; $R_1 = \text{CH}_3$, $Y = (\text{CH}_2)_3$, $R_2 = 2-\text{CF}_3$, $R_3 = \text{H}$, $R_4 = \text{CF}_3$].

Following the procedures of Example 20a, b and c and using equivalent amounts of reactants in each case but substituting in Example 20a 11-chloro-1-undecyne for 5-chloro-1-pentyne there can be obtained successively the following:

- 3,5-dimethyl-4-(9-ethynynonyloxy)benzonitrile (XII);
- 3,5-dimethyl-4-(9-ethynynonyloxy)-N-hydroxybenzenecarboximidamide (XIII); and
- 3-[3,5-dimethyl-4-(9-ethynynonyloxy)phenyl]-5-trifluoromethyl-1,2,4-oxadiazole. (III)

Following Procedure 1 and using equivalent amounts of acetaldehyde oxime and 3-[3,5-dimethyl-4-(9-ethynynonyloxy)phenyl]-5-trifluoromethyl-1,2,4-oxadiazole (III), there can be obtained 5-{9-[2,6-dimethyl-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)phenoxy]nonyl}-3-methylisoxazole [I; $R_1 = \text{CH}_3$, $Y = (\text{CH}_2)_9$, R_2 and $R_3 = 2,6-(\text{CH}_3)_2$, $R_4 = \text{CF}_3$].

Following Procedure 1 and using equivalent amounts of n-hexyl aldehyde oxime and the product of Example 20c (III), there can be obtained 5-{3-[2,6-dimethyl-4-(5-trifluoromethyl)phenoxy]propyl}-3-(n-pentyl)isoxazole

[I; R₁ = (CH₂)₄CH₃, Y = (CH₂)₃, R₁ and R₂ = 2,6-(CH₃)₂, R₄ = CF₃].

Following the procedure of Example 37 but substituting an equivalent amount of n-pentyl bromide for the ethyl iodide, there can be obtained 5-{3-[4-(5-cyclopropyl-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-pentyloxyisoxazole

[I; R₁ = O(CH₂)₄CH₃, Y = (CH₂)₃, R₁ and R₂ = 2,6-(CH₃)₂.

R₄ = cyclopropyl].

Following the procedure of Example 37 but substituting equivalent amounts of the product of Example 40 (I) and n-pentyl bromide for the product of Example 36b (I) and ethyl iodide respectively, there can be obtained 5-{3-[2,6-di-methyl-4-(5-(n-pentyloxymethyl)-1,2,4-oxadiazol-3-yl)-phenoxy]propyl}-3-methylisoxazole

[I; R₁ = CH₃, Y = (CH₂)₃, R₂ and R₃ = 2,6-(CH₃)₂,

R₄ = CH₂O(CH₂)₄CH₃].

Following the procedure of Example 1e but substituting an equivalent amount of cyclohexanecarbonyl chloride for the trifluoroacetic anhydride, there can be obtained 5-{3-[4-(5-cyclohexyl-1,2,4-oxadiazol-3-yl)-2,6-dimethylphenoxy]propyl}-3-methylisoxazole

[I; R₁ = CH₃, Y = (CH₂)₃, R₁ and R₂ = 2,6-(CH₃)₂, R₄ = cyclohexyl].

Biological evaluation of representative compounds of formula I has shown that they possess antiviral activity. They are useful in inhibiting virus replication *in vitro* and are primarily active against picornaviruses, especially rhino-viruses. The *in vitro* testing of the representative compounds of the invention against picornaviruses showed that viral replication was inhibited at minimum inhibitory concentrations (MIC) ranging from 0.002 to 9.608 micrograms per milliliter. The test procedure used was as follows:

TEST PROCEDURE

The MIC values were determined by an automated tissue culture infectious dose 50% (TCID-50) assay. HeLa (Wisconsin) cells in 96-well cluster plates were infected with a dilution of virus which had been shown empirically to produce 80% to 100% cytopathic effect (CPE) in 3 days in the absence of drug. The compound to be tested was serially diluted through 10, 2-fold cycles and added to the infected cells. After a 3-day incubation at 33°C and 2.5% carbon dioxide, the cells were fixed with a 5% solution of glutaraldehyde followed by staining with a 0.25% solution of crystal violet in water. The plates were then rinsed, dried, and the amount of stain remaining in the well (a measure of intact cells) was quantitated with an optical density reader. The MIC was determined to be the concentration of compound which protected 50% of the cells from virus-induced CPE relative to an untreated virus control.


In the above test procedures, representative compounds of formula I were tested against a panel of fifteen human rhinovirus (HRV) serotypes, namely, HRV-2, -14, -1A, -1B, -6, -21, -22, -15, -25, -30, -50, -67, -89, -86 and -41 and the MIC value, expressed in micrograms per milliliter (μg/ml), for each rhinovirus serotype was determined. The compounds tested were found to exhibit antiviral activity against one or more of these serotypes.

The MIC values (μg/ml) obtained for the compound of Example 1e in the above-described antiviral test procedure were as follows:

HRV Serotype (MIC)	HRV Serotype (MIC)	HRV Serotype (MIC)
-2 (0.027)	-21 (0.015)	-50 (0.154)
-14 (0.022)	-22 (0.011)	-67 (0.070)
-1A (0.119)	-15 (0.147)	-89 (0.015)
-1B (0.054)	-25 (0.036)	-86 (0.029)
-6 not tested	-30 (0.047)	-41 (0.338)

Claims

1. A compound of the formula




wherein

- R_1 is alkyl, alkoxy, hydroxy, cycloalkyl, hydroxyalkyl, alkoxyalkyl or hydroxyalkoxy;
 Y is alkylene of 3 to 9 carbon atoms,
 R_2 and R_3 independently are hydrogen, alkyl, alkoxy, halo, trifluoromethyl or nitro;
 R_4 is alkoxy, hydroxy, halomethyl, dihalomethyl, trihalomethyl, cycloalkyl, alkoxycarbonyl, hydroxyalkyl, alkoxyalkyl, alkanecarbonyloxyalkyl, cyano, 2,2,2-trifluoroethyl, (4-methylphenyl)sulfonyloxymethyl, N=Q or CON=Q, where N=Q is amino, alkylamino or dialkylamino; or a pharmaceutically-acceptable acid-addition salt thereof.

2. A compound as claimed in claim 1 wherein Y is alkylene of 3 to 5 carbon atoms.

3. A compound as claimed in either of the preceding claims of the formula



4. A compound as claimed in claim 3 wherein R_4 is C_{1-3} -alkoxy, fluoromethyl, dihalomethyl, trihalomethyl, cycloalkyl or C_{1-3} -alkoxy- C_{1-3} -alkyl.

5. A compound as claimed in Claim 4 wherein R_4 is trifluoromethyl.

6. 5-{3-[2,6-Dimethyl-4-(5-trifluoromethyl-1,2,4-oxadiazol-3-yl)phenoxy]propyl}-3-methylisoxazole

7. A composition for combatting picornaviruses which comprises an antivirally effective amount of a compound as claimed in claim 1 in admixture with a suitable pharmaceutical carrier or diluent.

8. A composition for combatting picornaviruses which comprises an antivirally effective amount of a compound as claimed in Claim 3 in admixture with a suitable pharmaceutical carrier or diluent.

9. A composition for combatting picornaviruses which comprises an antivirally effective amount of a compound as claimed in Claim 6 in admixture with a suitable pharmaceutical carrier or diluent.

10. The use of a compound as claimed in Claim 1 for the preparation of a medicament for combatting a picornaviral infection in a mammalian host.

11. The use of a compound as claimed in Claim 3 for the preparation of a medicament for combatting a picornaviral infection in a mammalian host.

12. The use of a compound as claimed in Claim 6 for the preparation of a medicament for combatting a picornaviral infection in a mammalian host.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 93 20 1058

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	EP-A-0 413 289 (STERLING DRUG INC.)	1-3,7,8,10,11	C07D413/12 A61K31/42
D	* claims 1,19-21 * & US-A-4 942 241		
A	EP-A-0 207 453 (STERLING DRUG INC.)	1-3,7,8,10,11	
P,X	EP-A-0 523 803 (STERLING WINTHROP INC)	1-3,7,8,10,11	
	* claims 1-3,5-7,9-10 *		
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			C07D
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 28 JUNE 1993	Examiner HENRY J.C.
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 (3.92) (P0401)