

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 593 406 A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **93870200.8**

(51) Int. Cl.⁵: **C11D 17/00, C11D 1/52,
C11D 1/72, C11D 1/68,
C11D 1/66**

(22) Date of filing: **13.10.93**

(30) Priority: **13.10.92 EP 92870164
06.11.92 EP 92870182**

(43) Date of publication of application:
20.04.94 Bulletin 94/16

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU NL
PT SE**

(71) Applicant: **THE PROCTER & GAMBLE
COMPANY
One Procter & Gamble Plaza
Cincinnati Ohio 45202(US)**

(72) Inventor: **Boutique, Jean-Pol
Rue Emile Labarre
B-5800 Genbloux (-Ernage)(BE)**

(74) Representative: **Canonici, Jean-Jacques et al
Procter & Gamble European Technical
Center N.V.
Temselaan 100
B-1853 Strombeek-Bever (BE)**

(54) **Non-aqueous liquid detergent compositions.**

(57) The present invention is a non-aqueous detergent composition comprising a surfactant comprising 2 vicinal hydroxyl groups selected from poly hydroxy fatty acid amides, alkyl mono and di(alkoxy) glyceryl ethers, alkane 1,2-diols, alkyl glycerates and N-alkyl-lactobionamides, and a boron comprising compound. The molar ratio of said surfactant to said boron comprising compound are equal to or lower than 2:1.

EP 0 593 406 A1

Field of the Invention

The present invention is related to non-aqueous liquid detergent compositions.

5 Background of the Invention

Detergent compositions have traditionally contained a number of non-active ingredients. This has resulted in the increase in volume of the compositions and thus the packaging therefore. It is now considered highly desirable to reduce the volume of detergent compositions due to environmental
10 considerations and produce more compact detergent formulations. One means to achieve this is to reduce the amount of non active ingredients such as water in detergent compositions. Reducing and preferably removing non active ingredients, particularly water results in a more compact composition. This in turn results in the need for less packaging material and smaller more manageable containers.

Thus, it is an object of the present invention to formulate non-aqueous detergent compositions which
15 exhibit at least the same cleaning performance of an equivalent aqueous detergent composition.

In order to achieve the required cleaning performance, it is desirable to incorporate surfactants comprising 2 vicinal hydroxyl groups selected from poly hydroxy fatty acid amides, alkyl mono or di(alkoxy) glyceryl ethers, alkane 1,2-diols, alkyl glycerates and N-alkyl-lactobionamides. This is because such surfactants are especially effective with respect to their greasy stain removal performance.

Unfortunately, it has been found that said surfactants comprising 2 vicinal hydroxyl groups are not
20 soluble in non-aqueous compositions and form precipitates.

Thus, it is a further object of the present invention to incorporate these surfactants comprising 2 vicinal hydroxyl groups in non-aqueous compositions without the precipitation of the surfactant in the composition.

In response to this object it has now been found that these surfactants comprising 2 vicinal hydroxyl
25 groups can indeed be solubilized in non-aqueous detergent compositions. This is achieved by the addition of a boron comprising compound such as boric acid.

Without wanting to be bound by theory it is believed that the boron comprising compound interacts with the hydroxyl groups of the surfactant to form complexes. These complexes disrupt the crystallinity of the surfactant and thereby prevent its precipitation.

Another advantage of the present invention is that certain ingredients such as dye transfer inhibition systems which are not compatible with an aqueous environment can be incorporated into non-aqueous liquid detergent compositions. This contributes to overall cleaning performance of liquid detergent compositions.

35 Summary of the Invention

The present invention is a non-aqueous liquid detergent composition comprising a surfactant comprising 2 vicinal hydroxyl groups selected from poly hydroxy fatty acid amides, alkyl mono- or di(alkoxy)-glyceryl ethers, alkane 1,2-diols, alkyl glycerates and N-alkyl-lactobionamides characterized in that said
40 composition further comprises a boron comprising compound wherein the molar ratio of said surfactant to said boron comprising compound is equal or lower than 2:1.

All amounts, parts and percentages are given as a % weight of the total composition unless otherwise stated.

45 Detailed description of the Invention

The present invention is a non-aqueous liquid detergent composition comprising a surfactant comprising 2 vicinal hydroxyl groups. Said surfactant is solubilized in the non-aqueous composition by the incorporation of a boron comprising compound.

As used herein the term non aqueous compositions refers to compositions which are substantially non-aqueous and comprise less than 5% water, preferably from 0.1% to 3% water.

The term detergent composition as used herein refers to complete formulations, premixes and liquid anhydrous bases in which other solid ingredients such as builders and bleaches may be suspended.

55 Surfactants comprising 2 vicinal hydroxyl groups

According to the present invention the compositions comprise a surfactant comprising 2 vicinal hydroxyl groups and mixtures thereof. Suitable surfactants comprising 2 vicinal hydroxyl groups include; surfactants

derived from poly hydroxy fatty acid amides, alkyl mono or di(alkoxy) glyceryl ethers, alkane 1,2 diols, alkyl glycerates and N-alkyl lactobionamides.

The compositions of the present invention comprise from 2% to 80%, preferably from 5% to 50% , most preferably from - 5% to 15% of said surfactant comprising 2 vicinal hydroxyl groups.

5 According to the present invention highly preferred surfactants comprising 2 vicinal hydroxyl groups are polyhydroxy fatty acid amide surfactants and mixtures thereof. Suitable poly hydroxy fatty acid amides for use herein are according to the formula:




15 wherein R¹ is H, C₁-C₄ hydrocarbyl, 2-hydroxy ethyl, 2-hydroxy propyl or a mixture thereof, preferably C₁-C₄ alkyl, more preferably C₁ or C₂ alkyl most preferably C₂ alkyl, R₂ is C₅₋₃₁ hydrocarbyl, preferably straight chain C_{7-C19} alkyl or alkenyl, more preferably straight chain C₁₁₋₁₇ alkyl or alkenyl or mixture thereof and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 2 hydroxyls directly connected to the chain, or an alkoxyated derivative (preferably ethoxylated or propoxylated) thereof. Z preferably is derived from a reducing sugar in a reductive amination reaction; more preferably Z is a glycityl. Suitable reducing sugars include glucose, fructose, maltose, lactose, galactose mannose and xylose. Z is preferably selected from the group consisting of -CH₂-(CHOH)_n-CH₂OH, CH(CH₂OH)-(CHOH)_{n-1}-CH₂OH, -CH₂-(CHOH)₂(CHOR')(CHOH)-CH₂OH, where n is an integer from 3 to 5, inclusive, and R' is H or a cyclic or aliphatic monosaccharide and alkoxyated derivatives thereof. Most preferred are glycityls wherein n is 4, particularly -CH₂-(CHOH)₄-CH₂OH. Z can be 1-deoxylactityl, 1-deoxyglucityl, 2-deoxyfructityl, 1-deoxymaltityl, 1-deoxylactityl, 1-deoxygalactityl, 1-deoxymannityl, 1-deoxymaltotriosityl.

20 R₁ can be for example N-methyl, N-ethyl, N-propyl, N-isopropyl, N-butyl, N-2-hydroxy ethyl, or N-2-hydroxy propyl. R₂ is a straight C₁₁₋₁₅ alkyl or alkenyl chain or mixtures thereof. The compositions according to the present invention comprise from 2% to 80%, preferably from 5% to 50%, most preferably from 5% to 15% of poly hydroxy fatty acid amide surfactants.


Other suitable surfactants comprising 2 vicinal hydroxy groups are selected from the group consisting of alkyl mono or di(alkoxy)glyceryl ethers, alkane 1,2- diols, alkyl glycerates and N-alkyl lactobionamides.

Suitable alkyl mono or di (alkoxy) glyceryl ethers are C₈₋₁₈ alkyl mono (alkoxy) glyceryl ethers according to the formula :




wherein R is a C₈₋₁₈ alkyl or alkenyl group and n is from 0 to 6.

40 Suitable alkyl di(alkoxy) glyceryl ethers are C₈₋₁₈ alkyl di(alkoxy) glyceryl ethers according to the formula :



45 where R is a C₈₋₁₈ alkyl or alkenyl group and n is from 0 to 6.


Suitable alkane diols are C₈₋₁₈ alkyl 1,2 diols, preferably C₁₀₋₁₂ alkyl 1,2 diols. Suitable alkylglycerates are C₈₋₁₈ alkylglycerates according to the formula :



where R is preferably C₁₀₋₁₂.

Suitable N-alkyl lactobionamides are N-C₈₋₁₈ alkyl lactobionamides according to the formula :

55



wherein R is a C₈₋₁₈ alkyl or alkenyl group.

The surfactant comprising 2 vicinal hydroxyl groups of the present invention is solubilized in the non-aqueous composition by the incorporation of a boron comprising compound.

Boron comprising compound

The boron comprising compound employed herein can be boric acid or one of its salts, or mixtures thereof. Preferred salts are the alkanolamine salts of tetraborate or metaborate. Most preferred are boric acid and monoethanolamine borate. According to the present invention the compositions comprise from 0.1% to 10%, preferably from 0.5% to 5%, most preferably from 2% to 4% of the boron comprising compound.

According to the present invention the molar ratio of said surfactant comprising 2 vicinal hydroxyl groups to said boron comprising compound material is equal or lower than 2:1, preferably equal or lower than 1.5:1, most preferably equal or lower than 1:1.

The rest of the liquid detergent composition according to the present invention is made of conventional detergency ingredients, i.e. other surfactants, builders, bleaches, enzymes, polymers, brighteners and others.

The liquid detergent compositions herein may additionally comprise as an optional ingredient from 2% to 90%, preferably from 5% to 50% of an organic surface-active agent selected from anionic, nonionic, cationic and zwitterionic surface active agents and mixtures thereof.


According to the present invention the compositions may further comprise a dye transfer inhibition system. Suitable dye transfer inhibiting systems for use herein include DTI systems comprising a non-iron metallo catalyst. Suitable non-iron metallo catalysts may be selected from:

- non-iron metallo porphin and water-soluble or water-dispersible derivatives thereof;
- non-iron metallo porphyrin and water-soluble or water-dispersible derivatives thereof;
- non-iron metallo phthalocyanine and water-soluble or water-dispersible derivatives thereof;

The preferred usage range of the catalyst in the wash is 10⁻⁸ molar to 10⁻³ molar, more preferred 10⁻⁶ - 10⁻⁴ molar.

The essential metallo porphin structure may be visualized as indicated in Formula I in the accompanying drawings. In Formula I the atom positions of the porphin structure are numbered conventionally and the double bonds are put in conventionally. In other formula, the double bonds have been omitted in the drawings, but are actually present as in I.

Preferred metallo porphin structures are those substituted at one or more of the 5, 10, 15 and 20 carbon positions of Formula I (Meso positions), with a phenyl or pyridyl substituent selected from the group consisting of




wherein n and m may be 0 or 1; A is selected from water-solubilizing group, e.g., sulfate, sulfonate, phosphate, and carboxylate groups; and B is selected from the group consisting of C₁-C₁₀ alkyl, C₁-C₁₀ polyethoxy alkyl or C₁-C₁₀ hydroxy alkyl.

Preferred molecules are those in which the substituents on the phenyl or pyridyl groups are selected from the group consisting of


-CH₃, -C₂H₅, -CH₂CH₂CH₂SO₃⁻, -CH₂--, and -CH₂CH(OH)CH₂SO₃⁻, -SO₃

A particularly preferred metallo porphin is one in which the molecule is substituted at the 5, 10, 15, and 20 carbon positions with the substituent



This preferred compound is known as metallo tetrasulfonated tetraphenylporphin. The symbol X¹ is (=CY-) wherein each Y, independently, is hydrogen, chlorine, bromine, fluorine or meso substituted alkyl, cycloalkyl, aralkyl, aryl, alkaryl or heteroaryl.


The symbol X² of Formula I represents an anion, preferably OH⁻ or Cl⁻. The compound of Formula I may be substituted at one or more of the remaining carbon positions with C₁-C₁₀ alkyl, hydroxyalkyl or oxyalkyl groups.



(I)

Porphin derivatives also include chlorophylls, chlorines, i.e. isobacterio chlorines and bacteriochlorines.

Metallo porphyrin and water-soluble or water-dispersible derivatives thereof have a structure given in formula II.



(II)


where X can be alkyl, alkyl carboxy, alkyl hydroxyl, vinyl, alkenyl, alkyl sulfate, alkylsulfonate, sulfate, sulfonate, aryl.

The symbol X² of Formula II represents an anion, preferably OH⁻ or Cl⁻.

The symbol X_i can be alkyl, alkylcarboxy, alkylhydroxyl, vinyl, alkenyl, alkylsulfate, alkylsulfonate, sulfate, sulfonate.

Metallo phthalocyanine and derivatives have the structure indicated in Formula III, wherein the atom positions of the phthalocyanine structure are numbered conventionally. The anionic groups in the above

structures contain cations selected from the group consisting of sodium and potassium cations or other non-interfering cations which leave the structures water-soluble. Preferred phthalocyanine derivatives are metallo phthalocyanine trisulfonate and metallo phthalocyanine tetrasulfonate.




Still a number of considerations are significant in selecting variants of or substituents in the basic porphrin or azaporphin structure. In the first place, one would choose compounds which are available or can be readily synthesized.

Beyond this, the choice of the substituent groups can be used to control the solubility of the catalyst in water or in detergent solutions. Yet again, especially where it is desired to avoid attacking dyes attached to solid surfaces, the substituents can control the affinity of the catalyst compound for the surface. Thus, strongly negatively charged substituted compounds, for instance the tetrasulfonated porphyrin, may be repelled by negatively charged stained surfaces and are therefore most likely not to cause attack on fixed dyes, whereas the cationic or zwitterionic compounds may be attracted to, or at least not repelled by such stained surfaces.

Another dye transfer inhibition system for use herein comprises polyamine N-oxide containing polymers and metallo catalysts.


Polyamine N-oxide polymers which contain units having the following structure formula :



wherein

P is a polymerisable unit, whereto the R-N-O group can be attached to or wherein the R-N-O group forms part of the polymerisable unit or a combination of both.

A is




-O-, -S-,



x is or O or 1;

R are aliphatic, ethoxylated aliphatics, aromatic, heterocyclic or alicyclic groups or any combination thereof whereto the nitrogen of the N-O group can be attached or wherein the nitrogen of the N-O group is part of these groups.

The N-O group can be represented by the following general structures :



wherein R1, R2, R3 are aliphatic groups, aromatic, heterocyclic or alicyclic groups or combinations thereof, x or/and y or/and z is 0 or 1 and wherein the nitrogen of the N-O group can be attached or wherein the nitrogen of the N-O group forms part of these groups

The N-O group can be part of the polymerisable unit (P) or can be attached to the polymeric backbone or a combination of both. Suitable polyamine N-oxides wherein the N-O group forms part of the polymerisable unit comprise polyamine N-oxides wherein R is selected from aliphatic, aromatic, alicyclic or heterocyclic groups.

One class of said polyamine N-oxides comprises the group of polyamine N-oxides wherein the nitrogen of the N-O group forms part of the R-group. Preferred polyamine N-oxides are those wherein R is a heterocyclic group such as pyridine, pyrrole, imidazole, pyrrolidine, piperidine and derivatives thereof. Another class of said polyamine N-oxides comprises the group of polyamine N-oxides wherein the nitrogen of the N-O group is attached to the R-group.

Other suitable polyamine N-oxides are the polyamine oxides whereto the N-O group is attached to the polymerisable unit.

Preferred class of these polyamine N-oxides are the polyamine N-oxides having the general formula (I) wherein R is an aromatic, heterocyclic or alicyclic groups wherein the nitrogen of the N-O functional group is part of said R group.

Examples of these classes are polyamine oxides wherein R is a heterocyclic compound such as pyridine, pyrrole, imidazole and derivatives thereof.

Another preferred class of polyamine N-oxides are the polyamine oxides having the general formula (I) wherein R are aromatic, heterocyclic or alicyclic groups wherein the nitrogen of the N-O functional group is attached to said R groups. Examples of these classes are polyamine oxides wherein R groups can be aromatic such as phenyl.

Any polymer backbone can be used as long as the amine oxide polymer formed is water-soluble and has dye transfer inhibiting properties. Examples of suitable polymeric backbones are polyvinyls, polyalkylenes, polyesters, polyethers, polyamide, polyimides, polyacrylates and mixtures thereof.

The amine N-oxide polymers of the present invention typically have a ratio of amine to the amine N-oxide of 10 : 1 to 1 : 1000000. However the amount of amine oxide groups present in the polyamine oxide polymer can be varied by appropriate copolymerization or by appropriate degree of N-oxidation. Preferably, the ratio of amine to amine N-oxide is from 3:1 to 1:1000000. The polymers of the present invention actually encompass random or block copolymers where one monomer type is an amine N-oxide and the other monomer type is an N-oxide or not.


The amine oxide unit of the polyamine N-oxides has a pKa < 10, preferably pKa < 7, more preferred pKa < 6. The polyamine oxides can be obtained in almost any degree of polymerisation. The degree of polymerisation is not critical provided the material has the desired water-solubility and dye-suspending power. Typically, the average molecular weight is within the range of 500 to 1000,000; more preferred 1000 to 500,000; most preferred 5000 to 100,000.

The polyamine N-oxides of the present invention are typically present from 0.01 to 10%, more preferably from 0.05 to 1%, most preferred from 0.05 to 0.5 % by weight of the dye transfer inhibiting composition.


Dye transfer inhibition systems comprising said polyamine N-oxide containing polymers may additionally comprise a metallo catalyst.

The preferred usage range of the catalyst in the wash is 10^{-8} molar to 10^{-3} molar, more preferred 10^{-6} - 10^{-4} molar. The essential metallo porphin structure may be visualized as indicated in Formula I in the accompanying drawings. In Formula I the atom positions of the porphin structure are numbered conventionally and the double bonds are put in conventionally. In other formula, the double bonds have been omitted in the drawings, but are actually present as in I.

Preferred metallo porphin structures are those substituted at one or more of the 5, 10, 15 and 20 carbon positions of Formula I (Meso positions), with a phenyl or pyridyl substituent selected from the group consisting of



and




wherein n and m may be 0 or 1; A is selected from water-solubilizing group, e.g., sulfate, sulfonate, phosphate or carboxylate groups; and B is selected from the group consisting of C_1 - C_{10} alkyl, C_1 - C_{10} polyethoxy alkyl and C_1 - C_{10} hydroxy alkyl.

Preferred molecules are those in which the substituents on the phenyl or pyridyl groups are selected from the group consisting of


$-CH_3$, $-C_2H_5$, $-CH_2CH_2CH_2SO_3^-$, $-CH_2-$, and $-CH_2CH(OH)CH_2SO_3^-$, $-SO_3$

A particularly preferred metallo porphin is one in which the molecule is substituted at the 5, 10, 15, and 20 carbon positions with the substituent



This preferred compound is known as metallo tetrasulfonated tetraphenylporphin. The symbol X^1 is (=CY-) wherein each Y, independently, is hydrogen, chlorine, bromine, fluorine or meso substituted alkyl, cycloalkyl, aralkyl, aryl, alkaryl or heteroaryl.


The symbol X^2 of Formula I represents an anion, preferably OH^- or Cl^- . The compound of Formula I may be substituted at one or more of the remaining carbon positions with C_1 - C_{10} alkyl, hydroxyalkyl or oxyalkyl groups.



(I)

Porphin derivatives also include chlorophylls, chlorines, i.e. isobacterio chlorines and bacteriochlorines.

Metallo porphyrin and water-soluble or water-dispersible derivatives thereof have a structure given in formula II.




where X can be alkyl, alkyl carboxy, alkyl hydroxyl, vinyl, alkenyl, alkyl sulfate, alkylsulfonate, sulfate, sulfonate, aryl.

The symbol X^2 of Formula II represents an anion, preferably OH^- or Cl^- .

The symbol X can be alkyl, alkylcarboxy, alkylhydroxyl, vinyl, alkenyl, alkylsulfate, alkylsulfonate, sulfate, sulfonate.

Metallo phthalocyanine and derivatives have the structure indicated in Formula III, wherein the atom positions of the phthalocyanine structure are numbered conventionally. The anionic groups in the above structures contain cations selected from the group consisting of sodium and potassium cations or other non-interfering cations which leave the structures water-soluble. Preferred phthalocyanine derivatives are metallo phthalocyanine trisulfonate and metallo phthalocyanine tetrasulfonate.



Another form of substitution possible for the present invention is substitution of the central metal by Fe, Mn, Co Rh, Cr, Ru, Mo or other transition metals.

Still a number of considerations are significant in selecting variants of or substituents in the basic porphin or azaporphin structure. In the first place, one would choose compounds which are available or can be readily synthesized.

Beyond this, where it is desired to avoid attacking dyes attached to solid surfaces, the substituents can control the affinity of the catalyst compound for the surface. Thus, strongly negatively charged substituted compounds, for instance the tetrasulfonated porphin, may be repelled by negatively charged stained surfaces and are therefore most likely not to cause attack on fixed dyes, whereas the cationic or zwitterionic compounds may be attracted to, or at least not repelled by such stained surfaces.

According to the present invention the composition may comprise a number of additional compounds commonly used in liquid detergent compositions such solvents, chelants, enzymes, perfumes, dyes which are well known and described in the art.

The present invention may be illustrated by the following examples wherein all the ingredients listed below are combined.

EXAMPLES

Example	I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII	XIII	XIV	XV	XVI	XVII	XVIII
Polyethyleneglycol 200	35	35	35	35	35	35	35	35	30	0	0	35	35	50	50	50	50	0
C12-14 EO 3	15	15	15	15	15	15	15	15	15	15	15	15	15	21.4	21.4	21.4	21.4	15
C12-14 EO 7	15	15	15	15	15	15	15	15	15	15	15	15	15	21.4	21.4	21.4	21.4	15
Propanediol	0.6	0.6	0.6	0.6	1.1	1.1	0	0	0	35	35	0	0	0	0	0	0	35
C12 N - methyl glucamide	5	5	5	10	10	10	0	0	6	10	10	0	0	0	0	0	0	10
C12 N - butyl glucamide	0	0	0	0	0	0	10	10	0	0	0	0	0	0	0	0	0	0
C12 N - methoxypropyl glucamide	0	0	0	0	0	0	0	0	0	0	0	10	10	0	0	0	0	0
C12 N - methyl glycerolamide	0	0	0	0	0	0	0	0	0	0	0	0	0	7.2	7.2	0	0	0
C12 N - methyl xylose amide	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	7.2	7.2	0
Na C 12-14 Alkyl sulphate	0	0	0	0	0	0	0	0	5.1	0	0	0	0	0	0	0	0	0
Boric acid	0	0.82	1.67	0	1.67	3.28	0	1	1.7	0	1.2	0	1.2	0	1	0	1	1.2
Poly Vinyl Pyrrolidone N - oxide	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.05
Total parts	70.6	71.4	72.3	75.6	77.8	79.4	75.0	76.0	72.8	75.0	76.2	75.0	76.2	100.0	101.0	100.0	101.0	76.3
Polyol surfactant/ boric acid molar ratio	oo	1	0.49	oo	1	0.5	oo	1.48	0.58	oo	1.37	oo	1.19	oo	1.56	oo	1.29	1.37

Examples 1, IV, VII, X, XII, XIV and XVI, outside the scope of the invention, show poor physical stability. The other examples where the molar ratio of the polyol surfactant/ boric acid is equal or lower than 2 show good physical stability.


Claims

1. A non-aqueous liquid detergent composition comprising a surfactant comprising 2 vicinal hydroxyl groups selected from poly hydroxy fatty acid amides, alkyl mono or di(alkoxy) glyceryl ethers, alkane
5 1,2-diols, alkyl glycerates and N-alkyl-lactobionamides characterized in that said composition further comprises a boron comprising compound wherein the molar ratio of said surfactant to said boron comprising compound is equal to or lower than 2:1.

2. A non-aqueous liquid detergent composition according to claim 1, comprising from 2% to 80% of said
10 surfactant.

3. A non aqueous liquid detergent composition according to claims 1 and 2, comprising from 0.1% to 10% of said boron comprising compound.

4. A non-aqueous liquid detergent composition according to any of the preceding claims, wherein said
15 surfactant is a poly hydroxy fatty acid amides, according to the formula:



wherein R¹ is H, C₁₋₄ hydrocarbyl, 2-hydroxy ethyl, 2-hydroxy propyl or a mixture thereof, R₂ is C₅₋₃₁
25 hydrocarbyl, and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxylated derivative thereof.

5. A non-aqueous liquid detergent composition according to any of the preceding claims, wherein said
30 boron comprising compound is selected from boric acid and the alkanolamine salts of tetraborate and metaborate.

6. A non aqueous liquid detergent composition according to claim 5, wherein said boron comprising
35 compound is boric acid.

7. A non-aqueous liquid detergent composition according to any of the preceding claims, further
40 comprising a dye transfer inhibition system.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 93 87 0200

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.5)
A	WO-A-92 06159 (THE PROCTER & GAMBLE COMPANY) * page 5, line 28 - page 8, line 3; example 12 * * page 27, line 22 - page 28, line 9 * ---	1-6	C11D17/00 C11D1/52 C11D1/72 C11D1/68 C11D1/66
A	WO-A-92 06160 (THE PROCTER & GAMBLE COMPANY) * page 5, line 27 - page 7, line 31 * * page 28, line 36 - page 29, line 24 * ---	1,2,4	
P,A	WO-A-93 01267 (THE PROCTER & GAMBLE COMPANY) * page 4, paragraph 1 - page 5, paragraph 4 * * page 26, paragraph 5 - page 27, paragraph 2; example 1 * ---	1-4	
A	EP-A-0 385 526 (UNILEVER NV) * page 2, line 1 - page 2, line 35; examples 1-3 * ---	1-3,5,6	
P,A	EP-A-0 550 278 (UNILEVER PLC) * page 3, line 20 - page 5, line 21; claim 9 * ---	1-3,5,6	TECHNICAL FIELDS SEARCHED (Int.Cl.5) C11D
A	EP-A-0 491 531 (UNILEVER PLC) * page 2, line 1 - page 2, line 57 * * page 4, line 32 - page 4, line 46; example 1 * ---	1-3,5	
P,A	EP-A-0 553 608 (THE PROCTER & GAMBLE COMPANY) * page 2, line 50 - page 5, line 31 * -----	7	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 2 February 1994	Examiner Doolan, G
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			