



(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

Note: Bibliography reflects the latest situation

- (15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see page(s) 5, 49-51
- (48) Corrigendum issued on:
27.10.2004 Bulletin 2004/44
- (45) Date of publication and mention
of the grant of the patent:
07.03.2001 Bulletin 2001/10
- (21) Application number: **94919288.4**
- (22) Date of filing: **27.05.1994**
- (51) Int Cl.7: **C12N 15/55**, C12N 9/16,
C07K 16/40, C07K 16/46,
C12N 15/11
- (86) International application number:
PCT/US1994/006066
- (87) International publication number:
WO 1994/028144 (08.12.1994 Gazette 1994/27)

(54) **CYCLIC GMP-BINDING, CYCLIC GMP-SPECIFIC PHOSPHODIESTERASE MATERIALS AND METHODS**

CYCLISCH GMP BINDENDE CYCLISCH GMP SPEZIFISCHE PHOSPHODIESTERASE MATERIALIEN UND VERFAHREN.

PHOSPHODIESTERASE DE FIXATION DU MONOPHOSPHATE DE GUANOSINE CYCLIQUE ET AVEC SPECIFICITE POUR CELUI-CI ET PROCEDE D'OBTENTION

- (84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL PT SE
- (30) Priority: **27.05.1993 US 68051**
- (43) Date of publication of application:
17.05.1995 Bulletin 1995/20
- (83) **Declaration under Rule 28(4) EPC (expert solution)**
- (60) Divisional application:
00112074.0 / 1 038 963
- (73) Proprietors:
• **BOARD OF REGENTS OF THE UNIVERSITY OF WASHINGTON**
Seattle, WA 98105 (US)
• **ICOS CORPORATION**
Bothell, WA 98021 (US)
- (72) Inventors:
• **BEAVO, Joseph, A.**
Seattle, WA 98155 (US)
• **CORBIN, Jackie, D.**
Nashville, TN 37215 (US)
- **FERGUSON, Kenneth, M.**
Bothell, WA 98021 (US)
• **FRANCIS, Sharron, H.**
Nashville, TN 37215 (US)
• **KADLECEK, Ann**
Seattle, WA 98103 (US)
• **LOUGHNEY, Kate**
Seattle, WA 98115 (US)
• **McALLISTER-LUCAS, Linda, M.**
Nashville, TN 37212 (US)
• **SONNENBURG, William, K.**
Mountlake Terrace, WA 98043 (US)
• **THOMAS, Melissa, K.**
Boston, MA 02114 (US)
- (74) Representative: **Richardson, Kate et al**
Forrester & Boehmert,
Pettenkoferstrasse 20-22
80336 München (DE)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

(56) References cited:

- JOURNAL OF BIOLOGICAL CHEMISTRY., vol.265, no.25, 5 September 1990, BALTIMORE US pages 14971 - 14978 THOMAS ETAL. 'Substrate- and Kinase-directed regulation of Phosphorylation of a cGMP-binding Phosphodiesterase by cGMP.' cited in the application
- JOURNAL OF BIOLOGICAL CHEMISTRY., vol.265, no.25, 5 September 1990, BALTIMORE, MD US pages 14964 - 14970 THOMAS ET AL. 'Characterization of a Purified Bovine Lung cGMP-binding cGMP Phosphodiesterase' cited in the application
- JOURNAL OF BIOLOGICAL CHEMISTRY., vol.267, no.26, 15 September 1992, BALTIMORE US pages 18683 - 18688 REPASKE ET AL. 'A Polymerase Chain Reaction Strategy to Identify and Clone Cylic Nucleotide Phosphodiesterase cDNAs'
- JOURNAL OF BIOLOGICAL CHEMISTRY., vol.266, no.26, 15 September 1991, BALTIMORE US pages 17655 - 17661 SONNENBURG ET AL. 'Molecular Cloning of a Cyclic GMP-stimulated Cyclic Nucleotide Phosphodiesterase cDNA'
- GENOMICS, vol.13, no.3, July 1992 pages 698 - 704 COLLINS ET AL. 'The human Beta-Subunit of Rod Photoreceptor cGMP Phosphodiesterase: Complete Retinal cDNA Sequence and Evidence for Expression in Brain' cited in the application
- JOURNAL OF BIOLOGICAL CHEMISTRY., vol.268, no.30, 25 October 1993, BALTIMORE US pages 22863 - 22873 MCALLISTER-LUCAS ET AL. 'The Structure of a Bovine Lung cGMP-binding, cGMP-specific Phosphodiesterase Deduced from a cDNA Clone'

Description

[0001] Experimental work described herein was supported in part by Research Grants GM15731, DK21723, DK40029 and GM41269 and the Medical Scientist Training Program Grant GM07347 awarded by the National Institutes of Health. The United States government has certain rights in the invention.

FIELD OF THE INVENTION

[0002] The present invention relates generally to a cyclic guanosine monophosphate-binding, cyclic guanosine monophosphate-specific phosphodiesterase designated cGB-PDE and more particularly to novel purified and isolated polynucleotides encoding cGB-PDE polypeptides, to methods and materials for recombinant production of cGB-PDE polypeptides, and to methods for identifying modulators of cGB-PDE activity.

BACKGROUND

[0003] Cyclic nucleotide phosphodiesterases (PDEs) that catalyze the hydrolysis of 3'5' cyclic nucleotides such as cyclic guanosine monophosphate (cGMP) and cyclic adenosine monophosphate (cAMP) to the corresponding nucleoside 5' monophosphates constitute a complex family of enzymes. By mediating the intracellular concentration of the cyclic nucleotides, the PDE isoenzymes function in signal transduction pathways involving cyclic nucleotide second messengers.

[0004] A variety of PDEs have been isolated from different tissue sources and many of the PDEs characterized to date exhibit differences in biological properties including physicochemical properties, substrate specificity, sensitivity to inhibitors, immunological reactivity and mode of regulation. [See Beavo *et al.*, *Cyclic Nucleotide Phosphodiesterases: Structure, Regulation and Drug Action*, John Wiley & Sons, Chichester, U.K. (1990)] Comparison of the known amino acid sequences of various PDEs indicates that most PDEs are chimeric multidomain proteins that have distinct catalytic and regulatory domains. [See Charbonneau, pp. 267-296 in Beavo *et al.*, *supra*] All mammalian PDEs characterized to date share a sequence of approximately 250 amino acid residues in length that appears to comprise the catalytic site and is located in the carboxyl terminal region of the enzyme. PDE domains that interact with allosteric or regulatory molecules are thought to be located within the amino-terminal regions of the isoenzymes. Based on their biological properties, the PDEs may be classified into six general families: the Ca²⁺/calmodulin-stimulated PDEs (Type I), the cGMP-stimulated PDEs (Type II), the cGMP-inhibited PDEs (Type III), the cAMP-specific PDEs (Type IV), the cGMP-specific phosphodiesterase cGB-PDE (Type V) which is the subject of the present invention and the cGMP-specific photoreceptor PDEs (Type VI).

[0005] The cGMP-binding PDEs (Type II, Type V and Type VI PDEs), in addition to having a homologous catalytic domain near their carboxyl terminus, have a second conserved sequence which is located closer to their amino terminus and which may comprise an allosteric cGMP-binding domain. See Charbonneau *et al.*, *Proc. Natl. Acad. Sci. USA*, 87: 288-292 (1990).

[0006] The Type II cGMP-stimulated PDEs (cGs-PDEs) are widely distributed in different tissue types and are thought to exist as homodimers of 100-105 kDa subunits. The cGs-PDEs respond under physiological conditions to elevated cGMP concentrations by increasing the rate of cAMP hydrolysis. The amino acid sequence of a bovine heart cGs-PDE and a partial cDNA sequence of a bovine adrenal cortex cGs-PDE are reported in LeTrong *et al.*, *Biochemistry*, 29: 10280-10288 (1990) and full length bovine adrenal and human fetal brain cGs-PDE cDNA sequences are described in Patent Cooperation Treaty International Publication No. WO 92/18541 published on October 29, 1992. The full length bovine adrenal cDNA sequence is also described in Sonnenburg *et al.*, *J. Biol. Chem.*, 266: 17655-17661 (1991).

[0007] The photoreceptor PDEs and the cGB-PDE have been described as cGMP-specific PDEs because they exhibit a 50-fold or greater selectivity for hydrolyzing cGMP over cAMP.

[0008] The photoreceptor PDEs are the rod outer segment PDE (ROS-PDE) and the cone PDE (COS-PDE). The holoenzyme structure of the ROS-PDE consists of two large subunits α (88 kDa) and β (84 kDa) which are both catalytically active and two smaller γ regulatory subunits (both 11 kDa). A soluble form of the ROS-PDE has also been identified which includes α , β , and γ subunits and a δ subunit (15 kDa) that appears to be identical to the COS-PDE 15 kDa subunit. A full-length cDNA corresponding to the bovine membrane-associated ROS-PDE α subunit is described in Ovchinnikov *et al.*, *FEBS Lett.*, 223: 169-173 (1987) and a full length cDNA corresponding to the bovine rod outer segment PDE β subunit is described in Lipkin *et al.*, *J. Biol. Chem.*, 265: 12955-12959 (1990). Ovchinnikov *et al.*, *FEBS Lett.*, 204: 169-173 (1986) presents a full-length cDNA corresponding to the bovine ROS-PDE γ subunit and the amino acid sequence of the δ subunit. Expression of the ROS-PDE has also been reported in brain in Collins *et al.*, *Genomics*, 13: 698-704 (1992). The COS-PDE is composed of two identical α' (94 kDa) subunits and three smaller subunits of 11 kDa, 13 kDa and 15 kDa. A full-length cDNA corresponding to the bovine COS-PDE α' subunit is reported in Li *et al.*, *Proc. Natl. Acad. Sci. USA*, 87: 293-297 (1990).

[0009] cGB-PDE has been purified to homogeneity from rat [Francis *et al.*, *Methods Enzymol.*, 159: 722-729 (1988)] and bovine lung tissue [Thomas *et al.*, *J. Biol. Chem.*, 265: 14964-14970 (1990), hereinafter "Thomas I"]. The presence of this or similar enzymes has been reported in a variety of tissues and species including rat and human platelets [Hamet *et al.*, *Adv. Cyclic Nucleotide Protein Phosphorylation Res.*, 16: 119-136 (1984)], rat spleen [Coquil *et al.*, *Biochem. Biophys. Res. Commun.*, 127: 226-231 (1985)], guinea pig lung [Davis *et al.*, *J. Biol. Chem.*, 252: 4078-4084 (1977)], vascular smooth muscle [Coquil *et al.*, *Biochim. Biophys. Acta*, 631: 148-165 (1980)], and sea urchin sperm [Francis *et al.*, *J. Biol. Chem.*, 255: 620-626 (1979)]. cGB-PDE may be a homodimer comprised of two 93 kDa subunits. [See Thomas I, *supra*] cGB-PDE has been shown to contain a single site not found in other known cGMP-binding PDEs which is phosphorylated by cGMP-dependent protein kinase (cGK) and, with a lower affinity, by cAMP-dependent protein kinase (cAK). [See Thomas *et al.*, *J. Biol. Chem.*, 265: 14971-14978 (1990), hereinafter "Thomas II"] The primary amino acid sequence of the phosphorylation site and of the amino-terminal end of a fragment generated by chymotryptic digestion of cGB-PDE are described in Thomas II, *supra*, and Thomas I, *supra*, respectively. However, the majority of the amino acid sequence of cGB-PDE has not previously been described.

[0010] Various inhibitors of different types of PDEs have been described in the literature. Two inhibitors that exhibit some specificity for Type V PDEs are zaprinast and dipyridamole. See Francis *et al.*, pp. 117-140 in Beavo *et al.*, *supra*.

[0011] Elucidation of the DNA and amino acid sequences encoding the cGB-PDE and production of cGB-PDE polypeptide by recombinant methods would provide information and material to allow the identification of novel agents that selectively modulate the activity of the cGB-PDEs. The recognition that there are distinct types or families of PDE isoenzymes and that different tissues express different complements of PDEs has led to an interest in the development of PDE modulators which may have therapeutic indications for disease states that involve signal transduction pathways utilizing cyclic nucleotides as second messengers. Various selective and non-selective inhibitors of PDE activity are discussed in Murray *et al.*, *Biochem. Soc. Trans.*, 20(2): 460-464 (1992). Development of PDE modulators without the ability to produce a specific PDE by recombinant DNA techniques is difficult because all PDEs catalyze the same basic reaction, have overlapping substrate specificities and occur only in trace amounts. As a result, purification to homogeneity of many PDEs is a tedious and difficult process.

[0012] There thus continues to exist a need in the art for DNA and amino acid sequence information for the cGB-PDE, for methods and materials for the recombinant production of cGB-PDE polypeptides and for methods for identifying specific modulators of cGB-PDE activity.

SUMMARY OF THE INVENTION

[0013] The present invention provides novel purified and isolated polynucleotides (e.g., DNA sequences and RNA transcripts, both sense and antisense strands, including splice variants thereof) encoding the human cGMP-binding, cGMP-specific PDE (cGB-PDE) polypeptide set out in SEQ ID NO:23. Also provided by the present invention is a purified and isolated polynucleotide encoding a human allelic variant of the cGB-PDE polypeptide set out in SEQ ID NO: 23, wherein said polynucleotide hybridizes at about 65°C in 3 x SSC, 20 mM sodium phosphate pH 6.8, with washing at about 65°C in 2 x SSC, to the non-coding strand of the DNA set out in SEQ ID NO: 22. According to the present invention, there is further provided a purified and isolated polynucleotide comprising the DNA sequence set out in SEQ ID NO: 22. Preferred DNA sequences of the invention include genomic and cDNA sequences as well as wholly or partially chemically synthesized DNA sequences. DNA sequences encoding cGB-PDE that are set out in SEQ ID NO: 9 or 20 and DNA sequences which hybridize thereto under stringent conditions or DNA sequences which would hybridize thereto but for the redundancy of the genetic code are disclosed herein. Also contemplated are biological replicas (i.e., copies of isolated DNA sequences made *in vivo* or *in vitro*) of DNA sequences of the invention. Autonomously replicating recombinant constructions such as plasmid and viral DNA vectors incorporating cGB-PDE sequences of the invention and especially vectors wherein DNA encoding cGB-PDE of the invention is operatively linked to an endogenous or exogenous expression control DNA sequence and a transcriptional terminator are also provided. Specifically illustrating expression plasmids of the invention is the plasmid hcgbmet156-2 6n in *E. coli* strain JM109 which was deposited with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852, on May 4, 1993 as Accession No. 69296.

[0014] According to another aspect of the invention, host cells including procaryotic and eucaryotic cells, are stably transformed with DNA sequences of the invention in a manner allowing the desired polypeptides to be expressed therein. Host cells expressing cGB-PDE products can serve a variety of useful purposes. Such cells constitute a valuable source of immunogen for the development of antibody substances specifically immunoreactive with cGB-PDE. Host cells of the invention are conspicuously useful in methods for the large scale production of cGB-PDE polypeptides wherein the cells are grown in a suitable culture medium and the desired polypeptide products are isolated from the cells or from the medium in which the cells are grown by, for example, immunoaffinity purification.

[0015] cGB-PDE products may be obtained as isolates from natural cell sources or may be chemically synthesized, but are preferably produced by recombinant procedures involving host cells of the invention. Use of mammalian host

cells is expected to provide for such post-translational modifications (e.g., glycosylation, truncation, lipidation and tyrosine, serine or threonine phosphorylation) as may be needed to confer optimal biological activity on recombinant expression products of the invention. cGB-PDE products of the invention may be full length polypeptides, fragments or variants. Variants may comprise cGB-PDE polypeptide analogs wherein one or more of the specified (i.e., naturally encoded) amino acids is deleted or replaced or wherein one or more nonspecified amino acids are added: (1) without loss of one or more of the biological activities or immunological characteristics specific for cGB-PDE; or (2) with specific disablement of a particular biological activity of cGB-PDE.

[0016] According to the present invention, there is further provided a fragment of human cGB-PDE consisting of the amino acids 516 to 875 of SEQ ID NO: 23, a fragment of human cGB-PDE consisting of the amino acids 1-494 of SEQ ID NO: 23, a fragment of the human cGB-PDE consisting of the amino acids 1-549 of SEQ ID NO: 23 and a fragment of the human cGB-PDE consisting of the amino acids 515-819 of SEQ ID NO: 23.

[0017] Also provided by the present invention are antibody substances (e.g., monoclonal and polyclonal antibodies, single chain antibodies, chimeric antibodies, CDR-grafted antibodies and the like) specifically immunoreactive with the cGB-PDE as set forth in SEQ ID NO: 23. Other binding proteins specific for cGB-PDE are also disclosed. Specific binding proteins can be developed using isolated or recombinant cGB-PDE or cGB-PDE variants or cells expressing such products. Binding proteins are useful, in turn, in compositions for immunization as well as for purifying cGB-PDE polypeptides and detection or quantification of cGB-PDE polypeptides in fluid and tissue samples by known immunological procedures. They are also manifestly useful in modulating (i.e., blocking, inhibiting or stimulating) biochemical activities of cGB-PDE, especially those activities involved in signal transduction. Anti-idiotypic antibodies specific for anti-cGB-PDE antibody substances are also contemplated.

[0018] The scientific value of the information contributed through the disclosures of DNA and amino acid sequences of the present invention is manifest. As one series of examples, knowledge of the sequence of a cDNA for cGB-PDE makes possible the isolation by DNA/DNA hybridization of genomic DNA sequences encoding cGB-PDE and specifying cGB-PDE expression control regulatory sequences such as promoters, operators and the like. DNA/DNA hybridization procedures carried out with DNA sequences of the invention under stringent conditions are likewise expected to allow the isolation of DNAs encoding allelic variants of cGB-PDE, other structurally related proteins sharing one or more of the biochemical and/or immunological properties specific to cGB-PDE, and non-human species proteins homologous to cGB-PDE. Polynucleotides of the invention when suitably labelled are useful in hybridization assays to detect the capacity of cells to synthesize cGB-PDE. Polynucleotides of the invention may also be the basis for diagnostic methods useful for identifying a genetic alteration(s) in the cGB-PDE locus that underlies a disease state or states. Also made available by the invention are anti-sense polynucleotides relevant to regulating expression of cGB-PDE by those cells which ordinarily express the same.

[0019] The DNA and amino acid sequence information provided by the present invention also makes possible the systematic analysis of the structure and function of cGB-PDE and definition of those molecules with which it will interact. Agents that modulate cGB-PDE activity may be identified by incubating a putative modulator with lysate from eucaryotic cells expressing recombinant cGB-PDE and determining the effect of the putative modulator on cGB-PDE phosphodiesterase activity. In a preferred embodiment the eucaryotic cell lacks endogenous cyclic nucleotide phosphodiesterase activity. Specifically illustrating such a eucaryotic cell is the yeast strain YKS45 which was deposited with the ATCC on May 19, 1993 as Accession No. 74225. The selectivity of a compound that modulates the activity of the cGB-PDE can be evaluated by comparing its activity on the cGB-PDE to its activity on other PDE isozymes. The combination of the recombinant cGB-PDE products of the invention with other recombinant PDE products in a series of independent assays provides a system for developing selective modulators of cGB-PDE.

[0020] Selective modulators may include, for example, antibodies and other proteins or peptides which specifically bind to the cGB-PDE or cGB-PDE nucleic acid, oligonucleotides which specifically bind to the cGB-PDE or cGB-PDE nucleic acid and other non-peptide compounds (e.g., isolated or synthetic organic molecules) which specifically react with cGB-PDE or cGB-PDE nucleic acid. Mutant forms of cGB-PDE which affect the enzymatic activity or cellular localization of the wild-type cGB-PDE are also contemplated. Presently preferred targets for the development of selective modulators include, for example: (1) the regions of the cGB-PDE which contact other proteins and/or localize the cGB-PDE within a cell, (2) the regions of the cGB-PDE which bind substrate, (3) the allosteric cGMP-binding site(s) of cGB-PDE, (4) the phosphorylation site(s) of cGB-PDE and (5) the regions of the cGB-PDE which are involved in dimerization of cGB-PDE subunits. Modulators of cGB-PDE activity may be therapeutically useful in treatment of a wide range of diseases and physiological conditions.

BRIEF DESCRIPTION OF THE DRAWINGS

[0021] Numerous other aspects and advantages of the present invention will be apparent upon consideration of the following detailed description thereof, reference being made to the drawing wherein:

EP 0 652 960 B9 (W1B1)

FIGURE 1A to 1C is an alignment of the conserved catalytic domains of several PDE isoenzymes wherein residues which are identical in all PDEs listed are indicated by their one letter amino acid abbreviation in the "conserved" line, residues which are identical in the cGB-PDE and photoreceptor PDEs only are indicated by a star in the "conserved" line and gaps introduced for optimum alignment are indicated by periods;

FIGURE 2A to 2C is an alignment of the cGMP-binding domains of several PDE isoenzymes wherein residues which are identical in all PDEs listed are indicated by their one letter amino acid abbreviation in the "conserved" line and gaps introduced for optimum alignment are indicated by periods;

FIGURE 3 is an alignment of internally homologous repeats from several PDE isoenzymes wherein residues identical in each repeat A and B from all cGMP-binding PDEs listed are indicated by their one letter amino acid abbreviation in the "conserved" line and stars in the "conserved" line represent positions in which all residues are chemically conserved;

FIGURE 4 schematically depicts the domain organization of cGB-PDE;

FIGURE 5 is a bar graph representing the results of experiments in which extracts of COS cells transfected with bovine cGB-PDE sequences or extracts of untransfected COS cells were assayed for phosphodiesterase activity using either 20 μ M cGMP or 20 μ M cAMP as the substrate;

FIGURE 6 is a graph depicting results of assays of extracts from cells transfected with bovine cGB-PDE sequences for cGMP phosphodiesterase activity in the presence of a series of concentrations of phosphodiesterase inhibitors including dipyridamole (closed squares), zaprinast (closed circles), methoxymethylxanthine (closed triangles) and rolipram (open circles);

FIGURE 7 is a bar graph presenting results of experiments in which cell extracts from COS cells transfected with bovine cGB-PDE sequences or control untransfected COS cells were assayed for [3 H]cGMP-binding activity in the absence (-) or presence (+) of 0.2 mM IBMX; and

FIGURE 8 is a graph of the results of assays in which extracts from cells transfected with bovine cGB-PDE sequences were assayed for [3 H]cGMP-binding activity in the presence of excess unlabelled cAMP (open circles) or cGMP (closed circles) at the concentrations indicated.

DETAILED DESCRIPTION

[0022] The following examples illustrate the invention. Example 1 describes the isolation of a bovine cGB-PDE cDNA fragment by PCR and subsequent isolation of a full length cGB-PDE cDNA using the PCR fragment as a probe. Example 2 presents an analysis of the relationship of the bovine cGB-PDE amino acid sequence to sequences reported for various other PDEs. Northern blot analysis of cGB-PDE mRNA in various bovine tissues is presented in Example 3. Expression of the bovine cGB-PDE cDNA in COS cells is described in Example 4. Example 5 presents results of assays of the cGB-PDE COS cell expression product for phosphodiesterase activity, cGMP-binding activity and Zn²⁺ hydrolase activity. Example 6 describes the isolation of human cDNAs homologous to the bovine cGB-PDE cDNA. The expression of a human cGB-PDE cDNA in yeast cells is presented in Example 7. RNase protection assays to detect cGB-PDE in human tissues are described in Example 8. Example 9 describes the bacterial expression of human cGB-PDE cDNA and the development of antibodies reactive with the bacterial cGB-PDE expression product. Example 10 describes cGB-PDE analogs and fragments. The generation of monoclonal antibodies that recognize cGB-PDE is described in Example 11. Example 12 relates to utilizing recombinant cGB-PDE products of the invention to develop agents that selectively modulate the biological activities of cGB-PDE.

Example 1

[0023] The polymerase chain reaction (PCR) was utilized to isolate a cDNA fragment encoding a portion of cGB-PDE from bovine lung first strand cDNA. Fully degenerate sense and antisense PCR primers were designed based on the partial cGB-PDE amino acid sequence described in Thomas I, *supra*, and novel partial amino acid sequence information.

A. Purification of cGB-PDE Protein

[0024] cGB-PDE was purified as described in Thomas I, *supra*, or by a modification of that method as described below.

[0025] Fresh bovine lungs (5-10 kg) were obtained from a slaughterhouse and immediately placed on ice. The tissue was ground and combined with cold PEM buffer (20mM sodium phosphate, pH 6.8, containing 2mM EDTA and 25mM β -mercaptoethanol). After homogenization and centrifugation, the resulting supernatant was incubated with 4-7 liters of DEAE-cellulose (Whatman, UK) for 3-4 hours. The DEAE slurry was then filtered under vacuum and rinsed with multiple volumes of cold PEM. The resin was poured into a glass column and washed with three to four volumes of PEM. The protein was eluted with 100mM NaCl in PEM and twelve 1-liter fractions were collected. Fractions were

EP 0 652 960 B9 (W1B1)

assayed for IBMX-stimulated cGMP binding and cGMP phosphodiesterase activities by standard procedures described in Thomas *et al.*, *supra*. Appropriate fractions were pooled, diluted 2-fold with cold, deionized water and subjected to Blue Sepharose® CL-6B (Pharmacia LKB Biotechnology Inc., Piscataway, NJ) chromatography. Zinc chelate affinity adsorbent chromatography was then performed using either an agarose or Sepharose-based gel matrix. The resulting protein pool from the zinc chelation step treated as described in the Thomas I, *supra*, or was subjected to a modified purification procedure.

[0026] As described in Thomas I, *supra*, the protein pool was applied in multiple loads to an HPLC Bio-Sil TSK-545 DEAE column (150 x 21.5 mm) (BioRad Laboratories, Hercules, CA) equilibrated in PEM at 4°C. After an equilibration period, a 120-ml wash of 50mM NaCl in PEM was followed by a 120-ml linear gradient (50-200mM NaCl in PEM) elution at a flow rate of 2 ml/minute. Appropriate fractions were pooled and concentrated in dialysis tubing against Sephadex G-200 (Boehringer Mannheim Biochemicals, UK) to a final volume of 1.5 ml. The concentrated cGB-PDE pool was applied to an HPLC gel filtration column (Bio-Sil TSK-250, 500 x 21.5 mm) equilibrated in 100mM sodium phosphate, pH 6.8, 2mM EDTA, 25mM β-mercaptoethanol and eluted with a flow rate of 2 ml/minute at 4°C.

[0027] If the modified, less cumbersome procedure was performed, the protein pool was dialyzed against PEM for 2 hours and loaded onto a 10 ml preparative DEAE Sephacel column (Pharmacia) equilibrated in PEM buffer. The protein was eluted batchwise with 0.5M NaCl in PEM, resulting in an approximately 10-15 fold concentration of protein. The concentrated protein sample was loaded onto an 800 ml (2.5 cm x 154 cm) Sephacryl S400 gel filtration column (Boehringer) equilibrated in 0.1M NaCl in PEM, and eluted at a flow rate of 1.7 ml/minute.

[0028] The purity of the protein was assessed by Coomassie staining after sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). Approximately 0.5-3.0 mg of pure cGB-PDE were obtained per 10 kg bovine lung.

[0029] Rabbit polyclonal antibodies specific for the purified bovine cGB-PDE were generated by standard procedures.

B. Amino Acid Sequencing of cGB-PDE

[0030] cGB-PDE phosphorylated with [³²P]ATP and was then digested with protease to yield ³²P-labelled phosphopeptides. Approximately 100 µg of purified cGB-PDE was phosphorylated in a reaction mixture containing 9mM MgCl₂, 9µM [³²P]ATP, 10µM cGMP, and 4.2 µg purified bovine catalytic subunit of cAMP-dependent protein kinase (cAK) in a final volume of 900 µl. Catalytic subunit of cAK was prepared according to the method of Flockhart *et al.*, pp. 209-215 in Marangos *et al.*, *Brain Receptor Methodologies, Part A*, Academic Press, Orlando, Florida (1984). The reaction was incubated for 30 minutes at 30°C, and stopped by addition of 60 µl of 200mM EDTA.

[0031] To obtain a first peptide sequence from cGB-PDE, 3.7 µl of a 1 mg/ml solution of a α-chymotrypsin in KPE buffer (10mM potassium phosphate, pH 6.8, with 2mM EDTA) was added to 100 µg purified, phosphorylated cGB-PDE and the mixture was incubated for 30 minutes at 30°C. Proteolysis was stopped by addition of 50 µl of 10% SDS and 25 µl of β-mercaptoethanol. The sample was boiled until the volume was reduced to less than 400 µl, and was loaded onto an 8% preparative SDS-polyacrylamide gel and subjected to electrophoresis at 50mAmps. The separated digestion products were electroblotted onto Immobilon polyvinylidene difluoride (Millipore, Bedford, MA), according to the method of Matsudaira, *J. Biol. Chem.*, 262: 10035-10038 (1987). Transferred protein was identified by Coomassie Blue staining, and a 50 kDa band was excised from the membrane for automated gas-phase amino acid sequencing. The sequence of the peptide obtained by the α-chymotryptic digestion procedure is set out below as SEQ ID NO: 1.

SEQ ID NO: 1

REXDANRINYMYAQYVKNTM

[0032] A second sequence was obtained from a cGB-PDE peptide fragment generated by V8 proteolysis. Approximately 200 µg of purified cGB-PDE was added to 10mM MgCl₂, 10µM [³²P]ATP, 100µM cGMP, and 1 µg/ml purified catalytic subunit of cAK in a final volume of 1.4 ml. The reaction was incubated for 30 minutes at 30°C, and was terminated by the addition of 160 µl of 0.2M EDTA. Next, 9 µl of 1 mg/ml *Staphylococcal aureus* V8 protease (International Chemical Nuclear Biomedicals, Costa Mesa, CA) diluted in KPE was added, followed by a 15 minute incubation at 30°C. Proteolysis was stopped by addition of 88 µl of 10% SDS and 45 µl β-mercaptoethanol. The digestion products were separated by electrophoresis on a preparative 10% SDS-polyacrylamide gel run at 25 mAmps for 4.5 hours. Proteins were electroblotted and stained as described above. A 28 kDa protein band was excised from the membrane and subjected to automated gas-phase amino acid sequencing. The sequence obtained is set out below as SEQ ID NO: 2.

SEQ ID NO: 2

QSLAAAVVP

5

C. PCR Amplification of Bovine cDNA

[0033] The partial amino acid sequences utilized to design primers (SEQ ID NO: 3, below, and amino acids 9-20 of SEQ ID NO: 1) and the sequences of the corresponding PCR primers (in IUPAC nomenclature) are set below wherein SEQ ID NO: 3 is the sequence reported in Thomas I, *supra*.

10

SEQ ID NO: 3

15

F D N D E G E Q

5' TTY GAY AAY GAY GAR GGN GAR CA 3' (SEQ ID NO: 4)

20

3' AAR CTR TTR CTR CTY CCN CTY GT 5' (SEQ ID NO: 5)

25

SEQ ID NO: 1, Amino acids 9-20

N Y M Y A Q Y V K N T M

30

5' AAY TAY ATG TAY GCN CAR TAY GT 3' (SEQ ID NO: 6)

35

3' TTR ATR TAC ATR CGN GTY ATR CA 5' (SEQ ID NO: 7)

3' TTR ATR TAC ATR CGN GTY ATR CAN TTY TTR TGN TAC 5'
(SEQ ID NO: 8)

40

The sense and antisense primers, synthesized using an Applied Biosystems Model 380A DNA Synthesizer (Foster City, CA), were used in all possible combinations to amplify cGB-PDE-specific sequences from bovine lung first strand cDNA as described below.

[0034] After ethanol precipitation, pairs of oligonucleotides were combined (SEQ ID NO: 4 or 5 combined with SEQ ID NOs: 6, 7 or 8) at 400nM each in a PCR reaction. The reaction was run using 50 ng first strand bovine lung cDNA (generated using AMV reverse transcriptase and random primers on oligo dT selected bovine lung mRNA), 200µM dNTPs, and 2 units of Taq polymerase. The initial denaturation step was carried out at 94°C for 5 minutes, followed by 30 cycles of a 1 minute denaturation step at 94°C, a two minute annealing step at 50°C, and a 2 minute extension step at 72°C. PCR was performed using a Hybaid Thermal Reactor (ENK Scientific Products, Saratoga, CA) and products were separated by gel electrophoresis on a 1% low melting point agarose gel run in 40mM Tris-acetate, 2mM EDTA. A weak band of about 800-840 bp was seen with the primers set out in SEQ ID NOs: 4 and 7 and with primers set out in SEQ ID NOs: 4 and 8. None of the other primer pairs yielded visible bands. The PCR product generated by amplification with the primers set out in SEQ ID NOs: 4 and 7 was isolated using the Gene Clean® (Bio101, La Jolla, CA) DNA purification kit according to the manufacturer's protocol. The PCR product (20 ng) was ligated into 200 ng of linearized pBluescript KS(+) (Stratagene, La Jolla, CA), and the resulting plasmid construct was used to transform *E. coli* XL1 Blue cells (Stratagene Cloning Systems, La Jolla, CA). Putative transformation positives were screened by sequencing. The sequences obtained were not homologous to any known PDE sequence or to the known partial cGB-PDE sequences.

55

EP 0 652 960 B9 (W1B1)

5 [0035] PCR was performed again on bovine lung first strand cDNA using the primers set out in SEQ ID NOs: 4 and 7. A clone containing a 0.8 Kb insert with a single large open reading frame was identified. The open reading frame encoded a polypeptide that included the amino acids KNTM (amino acids 17-20 of SEQ ID NO: 1 which were not utilized to design the primer sequence which is set out in SEQ ID NO: 7) and that possessed a high degree of homology to the deduced amino acid sequences of the cGs-, ROS- and COS-PDEs. The clone identified corresponds to nucleotides 489-1312 of SEQ ID NO: 9.

D. Construction and Hybridization Screening of a Bovine cDNA Library

10 [0036] In order to obtain a cDNA encoding a full-length cGB-PDE, a bovine lung cDNA library was screened using the ³²P-labelled PCR-generated cDNA insert as a probe.

15 [0037] Polyadenylated RNA was prepared from bovine lung as described Sonnenburg et al., *J. Biol. Chem.*, 266: 17655-17661 (1991). First strand cDNA was synthesized using AMV reverse transcriptase (Life Sciences, St. Petersburg, FL) with random hexanucleotide primers as described in Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, New York (1987). Second strand cDNA was synthesized using *E. coli* DNA polymerase I in the presence of *E. coli* DNA ligase and *E. coli* RNase H. Selection of cDNAs larger than 500 bp was performed by Sepharose® CL-4B (Millipore) chromatography. EcoRI adaptors (Promega, Madison, WI) were ligated to the cDNA using T4 DNA ligase. Following heat inactivation of the ligase, the cDNA was phosphorylated using T4 polynucleotide kinase. Unligated adaptors were removed by Sepharose® CL-4B chromatography (Pharmacia, Piscataway, NJ). The cDNA was ligated into EcoRI-digested, dephosphorylated lambda Zap® II arms (Stratagene) and packaged with Gigapack® Gold (Stratagene) extracts according to the manufacturer's protocol. The titer of the unamplified library was 9.9 x 10⁵ with 18% nonrecombinants. The library was amplified by plating 50,000 plaque forming units (pfu) on to twenty 150 mm plates, resulting in a final titer of 5.95 x 10⁶ pfu/ml with 21 % nonrecombinants.

20 [0038] The library was plated on twenty-four 150 mm plates at 50,000 pfu/plate, and screened with the ³²P-labelled cDNA clone. The probe was prepared using the method of Feinberg et al., *Anal. Biochem.*, 137: 266-267 (1984), and the ³²P-labelled DNA was purified using Elutip-D® columns (Schleicher and Schuell Inc., Keene, NH) using the manufacturer's protocol. Plaque-lifts were performed using 15 cm nitrocellulose filters. Following denaturation and neutralization, DNA was fixed onto the filters by baking at 80°C for 2 hours. Hybridization was carried out at 42°C overnight in a solution containing 50% formamide, 5X SSC (0.75M NaCl, 0.75M sodium citrate, pH 7), 25mM sodium phosphate (pH 7.0), 2X Denhardt's solution, 10% dextran sulfate, 90 µg/ml yeast tRNA, and approximately 106 cpm/ml ³²P-labelled probe (5x10⁸ cpm/µg). The filters were washed twice in 0.1X SSC, 0.1% SDS at room temperature for 15 minutes per wash, followed by a single 20 minute wash in 0.1X SSC, 1 % SDS at 45°C. The filters were then exposed to X-ray film at -70°C for several days.

25 [0039] Plaques that hybridized with the labelled probe were purified by several rounds of replating and rescreening. Insert cDNAs were subcloned into the pBluescript SK(-) vector (Stratagene) by the *in vivo* excision method described by the manufacturer's protocol. Southern blots were performed in order to verify that the rescued cDNA hybridized to the PCR probe. Putative cGB-PDE cDNAs were sequenced using Sequenase® Version 2.0 (United States Biochemical Corporation, Cleveland, Ohio) or TaqTrack® kits (Promega).

30 [0040] Three distinct cDNA clones designated cGB-2, cGB-8 and cGB-10 were isolated. The DNA and deduced amino acid sequences of clone cGB-8 are set out in SEQ ID NOs: 9 and 10. The DNA sequence downstream of nucleotide 2686 may represent a cloning artifact. The DNA sequence of cGB-10 is identical to the sequence of cGB-8 with the exception of one nucleotide. The DNA sequence of clone cGB-2 diverges from that of clone cGB-8 5' to nucleotide 219 of clone cgb-8 (see SEQ ID NO: 9) and could encode a protein with a different amino terminus.

35 [0041] The cGB-8 cDNA clone is 4474 bp in length and contains a large open reading frame of 2625 bp. The triplet ATG at position 99-101 in the nucleotide sequence is predicted to be the translation initiation site of the cGB-PDE gene because it is preceded by an in-frame stop codon and the surrounding bases are compatible with the Kozak consensus initiation site for eucaryotic mRNAs. The stop codon TAG is located at positions 2724-2726, and is followed by 1748 bp of 3' untranslated sequence. The sequence of cGB-8 does not contain a transcription termination consensus sequence, therefore the clone may not represent the entire 3' untranslated region of the corresponding mRNA.

40 [0042] The open reading frame of the cGB-8 cDNA encodes an 875 amino acid polypeptide with a calculated molecular mass of 99.5 kD. This calculated molecular mass is only slightly larger than the reported molecular mass of purified cGB-PDE, estimated by SDS-PAGE analysis to be approximately 93 kDa. The deduced amino acid sequence of cGB-8 corresponded exactly to all peptide sequences obtained from purified bovine lung cGB-PDE providing strong evidence that cGB-8 encodes cGB-PDE.

Example 2

45 [0043] A search of the SWISS-PROT and GEnEmbl data banks (Release of February, 1992) conducted using the

FASTA program supplied with the Genetics Computer Group (GCG) Software Package (Madison, Wisconsin) revealed that only DNA and amino acid sequences reported for other PDEs displayed significant similarity to the DNA and deduced amino acid of clone cGB-8.

[0044] Pairwise comparisons of the cGB-PDE deduced amino acid sequence with the sequences of eight other PDEs were conducted using the ALIGN [Dayhoff *et al.*, *Methods Enzymol.*, 92: 524-545 (1983)] and BESTFIT [Wilbur *et al.*, *Proc. Natl. Acad. Sci. USA*, 80: 726-730 (1983)] programs. Like all mammalian phosphodiesterases sequenced to date, cGB-PDE contains a conserved catalytic domain sequence of approximately 250 amino acids in the carboxyl-terminal half of the protein that is thought to be essential for catalytic activity. This segment comprises amino acids 578-812 of SEQ ID NO: 9 and exhibits sequence conservation with the corresponding regions of other PDEs. Table 1 below sets out the specific identity values obtained in pairwise comparisons of other PDEs with amino acids 578-812 of cGB-PDE, wherein "ratdunce" is the rat cAMP-specific PDE; "61 kCaM" is the bovine 61 kDa calcium/calmodulin-dependent PDE; "63 kCaM" is the bovine 63 kDa calcium/calmodulin-dependent PDE; "drosdunce" is the drosophila cAMP-specific dunce PDE; "ROS- α " is the bovine ROS-PDE α -subunit; "ROS- β " is the bovine ROS-PDE β -subunit; "COS- α " is the bovine COS-PDE α ' subunit; and "cGs" is the bovine cGs-PDE (612-844).

Table 1

Phosphodiesterase	Catalytic Domain Residues	% Identity
Ratdunce	77-316	31
61 kCaM	193-422	29
63 kcam	195-424	29
drosdunce	1-239	28
ROS- α	535-778	45
ROS- β	533-776	46
COS- α '	533-776	48
cGs	612-844	40

[0045] Multiple sequence alignments were performed using the Progressive Alignment Algorithm [Feng *et al.*, *Methods Enzymol.*, 183: 375-387 (1990)] implemented in the PILEUP program (GCG Software). FIGURE 1A to 1C shows a multiple sequence alignment of the proposed catalytic domain of cGB-PDE with the all the corresponding regions of the PDEs of Table 1. Twenty-eight residues (see residues indicated by one letter amino acid abbreviations in the "conserved" line on FIGURE 1A to 1C) are invariant among the isoenzymes including several conserved histidine residues predicted to play a functional role in catalysis. See Charbonneau *et al.*, *Proc. Natl. Acad. Sci. USA*, *supra*. The catalytic domain of cGB-PDE more closely resembles the catalytic domains of the ROS-PDEs and COS-PDEs than the corresponding regions of other PDE isoenzymes. There are several conserved regions among the photoreceptor PDEs and cGB-PDE that are not shared by other PDEs. Amino acid positions in these regions that are invariant in the photoreceptor PDE and cGB-PDE sequences are indicated by stars in the "conserved" line of FIGURE 1A to 1C. Regions of homology among cGB-PDE and the ROS- and COS-PDEs may serve important roles in conferring specificity for cGMP hydrolysis relative to cAMP hydrolysis or for sensitivity to specific pharmacological agents.

[0046] Sequence similarity between cGB-PDE, cGs-PDE and the photoreceptor PDEs, is not limited to the conserved catalytic domain but also includes the noncatalytic cGMP binding domain in the amino-terminal half of the protein. Optimization of the alignment between cGB-PDE, cGs-PDE and the photoreceptor PDEs indicates that an amino-terminal conserved segment may exist including amino acids 142-526 of SEQ ID NO: 9. Pairwise analysis of the sequence of the proposed cGMP-binding domain of cGB-PDE with the corresponding regions of the photoreceptor PDEs and cGs-PDE revealed 26-28% sequence identity. Multiple sequence alignment of the proposed cGMP-binding domains with the cGMP-binding PDEs is shown in FIGURE 2A to 2C wherein abbreviations are the same as indicated for Table 1. Thirty-eight positions in this non-catalytic domain appear to be invariant among all cGMP-binding PDEs (see positions indicated by one letter amino acid abbreviations in the "conserved" line of FIGURE 2A to 2C).

[0047] The cGMP-binding domain of the cGMP-binding PDEs contains internally homologous repeats which may form two similar but distinct inter- or intra-subunit cGMP-binding sites. FIGURE 3 shows a multiple sequence alignment of the repeats a (corresponding to amino acids 228-311 of cGB-PDE) and b (corresponding to amino acids 410-500 of cGB-PDE) of the cGMP-binding PDEs. Seven residues are invariant in each A and B regions (see residues indicated by one letter amino acid abbreviations in the "conserved" line of FIGURE 3). Residues that are chemically conserved in the A and B regions are indicated by stars in the "conserved" line of FIGURE 3. cGMP analog studies of cGB-PDE support the existence of a hydrogen bond between the cyclic nucleotide binding site on cGB-PDE and the 2'OH of cGMP.

[0048] Three regions of cGB-PDE have no significant sequence similarity to other PDE isoenzymes. These regions include the sequence flanking the carboxyl-terminal end of the catalytic domain (amino acids 812-875), the sequence

EP 0 652 960 B9 (W1B1)

separating the cGMP-binding and catalytic domains (amino acids 527-577) and the amino-terminal sequence spanning amino acids 1-141. The site (the serine at position 92 of SEQ ID NO: 10) of phosphorylation of cGB-PDE by cGK is located in this amino-terminal region of sequence. Binding of cGMP to the allosteric site on cGB-PDE is required for its phosphorylation.

5 [0049] A proposed domain structure of cGB-PDE based on the foregoing comparisons with other PDE isoenzymes is presented in FIGURE 4. This domain structure is supported by the biochemical studies of cGB-PDE purified from bovine lung.

Example 3

10 [0050] The presence of cGB-PDE mRNA in various bovine tissues was examined by Northern blot hybridization.

[0051] Polyadenylated RNA was purified from total RNA preparations using the Poly(A) Quick® mRNA purification kit (Stratagene) according to the manufacturer's protocol. RNA samples (5 µg) were loaded onto a 1.2% agarose, 6.7% formaldehyde gel. Electrophoresis and RNA transfer were performed as previously described in Sonnenburg *et al.*, *supra*. Prehybridization of the RNA blot was carried out for 4 hours at 45°C in a solution containing 50% formamide, 5X SSC, 25mM sodium phosphate, pH 7, 2X Denhardt's solution, 10% dextran sulfate, and 0.1 mg/ml yeast tRNA. A random hexanucleotide-primer-labelled probe (5 X 10⁸ cpm/µg) was prepared as described in Feinberg *et al.*, *supra*, using the 4.7 kb cGB-8 cDNA clone of Example 2 excised by digestion with *Accl* and *SacII*. The probe was heat denatured and injected into a blotting bag (6 X 10⁵ cpm/ml) following prehybridization. The Northern blot was hybridized overnight at 45°C, followed by one 15 minute wash with 2X SSC, 0.1% SDS at room temperature, and three 20 minute washes with 0.1X SSC, 0.1% SDS at 45°C. The blot was exposed to X-ray film for 24 hours at -70°C. The size of the RNA that hybridized with the cGB-PDE probe was estimated using a 0.24-9.5 kb RNA ladder that was stained with ethidium bromide and visualized with UV light.

20 [0052] The ³²P-labelled cGB-PDE cDNA hybridized to a single 6.8 kb bovine lung RNA species. A mRNA band of the identical size was also detected in polyadenylated RNA isolated from bovine trachea, aorta, kidney and spleen.

Example 4

30 [0053] The cGB-PDE cDNA in clone cGB-8 of Example 2 was expressed in COS-7 cells (ATCC CRL1651).

[0054] A portion of the cGB-8 cDNA was isolated following digestion with the restriction enzyme *XbaI*. *XbaI* cut at a position in the pBluescript polylinker sequence located 30 bp upstream of the 5' end of the cGB-8 insert and at position 3359 within the cGB-8 insert. The resulting 3389 bp fragment, which contains the entire coding region of cGB-8, was then ligated into the unique *XbaI* cloning site of the expression vector pCDM8 (Invitrogen, San Diego, CA). The pCDM8 plasmid is a 4.5 kb eucaryotic expression vector containing a cytomegalovirus promoter and enhancer, an SV40-derived origin of replication, a polyadenylation signal, a procaryotic origin of replication (derived from pBR322) and a procaryotic genetic marker (*supF*). *E. coli* MC1061/P3 cells (Invitrogen) were transformed with the resulting ligation products, and transformation positive colonies were screened for proper orientation of the cGB-8 insert using PCR and restriction enzyme analysis. The resulting expression construct containing the cGB-8 insert in the proper orientation is referred to as pCDM8-cGB-PDE.

35 [0055] The pCDM8-cGB-PDE DNA was purified from large-scale plasmid preparations using Qiagen pack-500 columns (Chatsworth, CA) according to the manufacturer's protocol. COS-7 cells were cultured in Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum, 50 µg/ml penicillin and 50 µg/ml streptomycin at 37°C in a humidified 5 % CO₂ atmosphere. Approximately 24 hours prior to transfection, confluent 100 mm dishes of cells were replated at one-fourth or one-fifth the original density. In a typical transfection experiment, cells were washed with buffer containing 137mM NaCl, 2.7mM KCl, 1.1mM potassium phosphate, and 8.1mM sodium phosphate, pH 7.2 (PBS). Then 4-5 ml of DMEM containing 10% NuSerum (Collaborative Biomedical Products, Bedford, MA) was added to each plate. Transfection with 10 µg pCDM8-cGB-PDE DNA or pCDM8 vector DNA mixed with 400 µg DEAE-dextran (Pharmacia) in 60 µl TBS [Tris-buffered saline: 25mM Tris-HCl (pH 7.4), 137mM NaCl, 5mM KCl, 0.6mM Na₂HPO₄, 0.7mM CaCl₂, and 0.5mM MgCl₂] was carried out by dropwise addition of the mixture to each plate. The cells were incubated at 37°C, 5% CO₂ for 4 hours, and then treated with 10% dimethyl sulfoxide in PBS for 1 minute. After 2 minutes, the dimethyl sulfoxide was removed, the cells were washed with PBS and incubated in complete medium. After 48 hours, cells were suspended in 0.5-1 ml of cold homogenization buffer [40mM Tris-HCl (pH 7.5), 15mM benzamidine, 15mM β-mercaptoethanol, 0.7 µg/ml pepstatin A, 0.5 µg/ml leupeptin, and 5µM EDTA] per plate of cells, and disrupted using a Dounce homogenizer. The resulting whole-cell extracts were assayed for phosphodiesterase activity, cGMP-binding activity, and total protein concentration as described below in Example 5.

Example 5

[0056] Phosphodiesterase activity in extracts of the transfected COS cells of Example 4 or in extracts of mock transfected COS cells was measured using a modification of the assay procedure described for the cGs-PDE in Martins *et al.*, *J. Biol. Chem.*, 257: 1973-1979 (1982). Cells were harvested and extracts prepared 48 hours after transfection. Incubation mixtures contained 40mM MOPS buffer (pH 7), 0.8mM EDTA, 15mM magnesium acetate, 2 mg/ml bovine serum albumin, 20 μ M [³H]cGMP or [³H]cAMP (100,000-200,000 cpm/assay) and COS-7 cell extract in a total volume of 250 μ l. The reaction mixture was incubated for 10 minutes at 30°C, and then stopped by boiling. Next, 10 μ l of 10mg/ml *Crotalus atrox* venom (Sigma) was added followed by a 10 minute incubation at 30°C. Nucleoside products were separated from unreacted nucleotides as described in Martins *et al.*, *supra*. In all studies, less than 15% of the total [³H]cyclic nucleotide was hydrolyzed during the reaction.

[0057] The results of the assays are presented in FIGURE 5 wherein the results shown are averages of three separate transfections. Transfection of COS-7 cells with pCDM8-cGB-PDE DNA resulted in the expression of approximately 15-fold higher levels of cGMP phosphodiesterase activity than in mock-transfected cells or in cells transfected with pCDM8 vector alone. No increase in cAMP phosphodiesterase activity over mock or vector-only transfected cells was detected in extracts from cells transfected with pCDM8-cGB-PDE DNA. These results confirm that the cGB-PDE bovine cDNA encodes a cGMP-specific phosphodiesterase.

[0058] Extracts from the transfected COS cells of Example 4 were also assayed for cGMP PDE activity in the presence of a series of concentrations of the PDE inhibitors zaprinast, dipyridamole (Sigma), isobutyl-1-methyl-8-methoxymethylxanthine (MeOxMeMIX) and rolipram.

[0059] The results of the assays are presented in FIGURE 6 wherein PDE activity in the absence of inhibitor is taken as 100% and each data point represents the average of two separate determinations. The relative potencies of PDE inhibitors for inhibition of cGMP hydrolysis by the expressed cGB-PDE cDNA protein product were identical to those relative potencies reported for native cGB-PDE purified from bovine lung (Thomas I, *supra*). IC₅₀ values calculated from the curves in FIGURE 6 are as follows: zaprinast (closed circles), 2 μ M; dipyridamole (closed squares), 3.5 μ M; MeOxMeMIX (closed triangles), 30 μ M; and rolipram (open circles), >300 μ M. The IC₅₀ value of zaprinast, a relatively specific inhibitor of cGMP-specific phosphodiesterases, was at least two orders of magnitude lower than that reported for inhibition of phosphodiesterase activity of the cGs-PDE or of the cGMP-inhibited phosphodiesterase (cGi-PDEs) (Reeves *et al.*, pp. 300-316 in Beavo *et al.*, *supra*). Dipyridamole, an effective inhibitor of selected cAMP- and cGMP-specific phosphodiesterases, was also a potent inhibitor of the expressed cGB-PDE. The relatively selective inhibitor of calcium/calmodulin-stimulated phosphodiesterase (CaM-PDEs), MeOxMeMIX, was approximately 10-fold less potent than zaprinast and dipyridamole, in agreement with results using cGB-PDE activity purified from bovine lung. Rolipram, a potent inhibitor of low K_m cAMP phosphodiesterases, was a poor inhibitor of expressed cGB-PDE cDNA protein product. These results show that the cGB-PDE cDNA encodes a phosphodiesterase that possesses catalytic activity characteristic of cGB-PDE isolated from bovine tissue, thus verifying the identity of the cGB-8 cDNA clone as a cGB-PDE.

[0060] It is of interest to note that although the relative potencies of the PDE inhibitors for inhibition of cGMP hydrolysis were identical for the recombinant and bovine isolate cGB-PDE, the absolute IC₅₀ values for all inhibitors tested were 2-7 fold higher for the recombinant cGB-PDE. This difference could not be attributed to the effects of any factors present in COS-7 cell extracts on cGMP hydrolytic activity, since cGB-PDE isolated from bovine tissue exhibited identical kinetics of inhibition as a pure enzyme, or when added back to extracts of mock-transfected COS-7 cells. This apparent difference in pharmacological sensitivity may be due to a subtle difference in the structure of the recombinant cGB-PDE cDNA protein product and bovine lung cGB-PDE, such as a difference in post-translational modification at or near the catalytic site. Alternatively, this difference may be due to an alteration of the catalytic activity of bovine lung cGB-PDE over several purification steps.

[0061] Cell extracts were assayed for [³H]cGMP-binding activity in the absence or presence of 0.2mM 3-isobutyl-1-methylxanthine (IBMX) (Sigma), a competitive inhibitor of cGMP hydrolysis. The cGMP binding assay, modified from the assay described in Thomas I, *supra*, was conducted in a total volume of 80 μ l. Sixty μ l of cell extract was combined with 20 μ l of a binding cocktail such that the final concentration of components of the mixture were 1 μ M [³H]cGMP, 5 μ M cAMP, and 10 μ M 8-bromo-cGMP. The cAMP and 8-bromo-cGMP were added to block [³H]cGMP binding to cAK and cGK, respectively. Assays were carried out in the absence and presence of 0.2mM IBMX. The reaction was initiated by the addition of the cell extract, and was incubated for 60 minutes at 0°C. Filtration of the reaction mixtures was carried out as described in Thomas I, *supra*. Blanks were determined by parallel incubations with homogenization buffer replacing cell extracts, or with a 100-fold excess of unlabelled cGMP. Similar results were obtained with both methods. Total protein concentration of the cell extracts was determined by the method of Bradford, *Anal. Biochem.*, 72:248-254 (1976) using bovine serum albumin as the standard.

[0062] Results of the assay are set out in FIGURE 7. When measured at 1 μ M [³H]cGMP in the presence of 0.2mM IBMX, extracts from COS-7 cells transfected with pCDM8-cGB-PDE exhibited 8-fold higher cGMP-binding activity than

extracts from mock-transfected cells. No IBMX stimulation of background cGMP binding was observed suggesting that little or no endogenous cGB-PDE was present in the COS-7 cell extracts. In extracts of pCDM8-cGB-PDE transfected cells cGMP-specific activity was stimulated approximately 1.8-fold by the addition of 0.2mM IBMX. The ability of IBMX to stimulate cGMP binding 2-5 fold is a distinctive property of the cGMP-binding phosphodiesterases.

[0063] Cell extracts were assayed as described above for [³H]cGMP-binding activity (wherein concentration of [³H]cGMP was 2.5μM) in the presence of excess unlabelled cAMP or cGMP. Results are presented in FIGURE 8 wherein cGMP binding in the absence of unlabelled competitor was taken as 100% and each data point represents the average of three separate determinations. The binding activity of the protein product encoded by the cGB-PDE cDNA was specific for cGMP relative to cAMP. Less than 10-fold higher concentrations of unlabelled cGMP were required to inhibit [³H]cGMP binding activity by 50% whereas approximately 100-fold higher concentrations of cAMP were required for the same degree of inhibition.

[0064] The results presented in this example show that the cGB-PDE cDNA encodes a phosphodiesterase which possesses biochemical activities characteristic of native cGB-PDE.

[0065] The catalytic domains of mammalian PDEs and a *Drosophila* PDE contain two tandem conserved sequences (HX₃HX₂₄₋₂₆E) that are typical Zn²⁺-binding motifs in Zn²⁺ hydrolases such as thermolysin [Vallee and Auld, *Biochem.*, 29: 5647-5659 (1990)]. cGB-PDE binds Zn²⁺ in the presence of large excesses of Mg²⁺, Mn²⁺, Fe²⁺, Fe³⁺, Ca²⁺ or Cd²⁺. In the absence of added metal, cGB-PDE has a PDE activity that is approximately 20% of the maximum activity that occurs in the presence of 40 mM Mg²⁺, and this basal activity is inhibited by 1,10-phenanthroline or EDTA. This suggests that a trace metal(s) accounts for the basal PDE activity despite exhaustive treatments to remove metals. PDE activity is stimulated by addition of Zn²⁺ (0.02-1 μM) or Co²⁺ (1-20 μM), but not by Fe²⁺, Fe³⁺, Ca²⁺, Cd²⁺, or Cu²⁺. Zn²⁺ increases the basal PDE activity up to 70% of the maximum stimulation produced by 40mM Mg²⁺. The stimulatory effect of Zn²⁺ in these assays may be compromised by an inhibitory effect that is caused by Zn²⁺ concentrations > 1 μM. The Zn²⁺-supported PDE activity and Zn²⁺ binding by cGB-PDE occur at similar concentrations of Zn²⁺. cGB-PDE thus appears to be a Zn²⁺ hydrolase and Zn²⁺ appears to play a critical role in the activity of the enzyme. See, Colbran *et al.*, *The FASEB J.*, 8: Abstract 2148 (March 15, 1994).

Example 6

[0066] Several human cDNA clones, homologous to the bovine cDNA clone encoding cGB-PDE, were isolated by hybridization under stringent conditions using a nucleic acid probe corresponding to a portion of the bovine cGB-8 clone (nucleotides 489-1312 of SEQ ID NO: 9).

Isolation of cDNA Fragments Encoding Human cGB-PDE

[0067] Three human cDNA libraries (two glioblastoma and one lung) in the vector lambda Zap were probed with the bovine cGB-PDE sequence. The PCR-generated clone corresponding to nucleotides 484-1312 of SEQ ID NO: 9 which is described in Example 1 was digested with *EcoRI* and *SaI* and the resulting 0.8 kb cDNA insert was isolated and purified by agarose gel electrophoresis. The fragment was labelled with radioactive nucleotides using a random primed DNA labelling kit (Boehringer).

[0068] The cDNA libraries were plated on 150 mm petri plates at a density of approximately 50,000 plaques per plate. Duplicate nitrocellulose filter replicas were prepared. The prehybridization buffer was 3X SSC, 0.1% sarkosyl, 10X Denhardt's, 20mM sodium phosphate (pH 6.8) and 50 μg/ml salmon testes DNA. Prehybridization was carried out at 65°C for a minimum of 30 minutes. Hybridization was carried out at 65°C overnight in buffer of the same composition with the addition of 1-5x10⁵ cpm/ml of probe. The filters were washed at 65°C in 2X SSC, 0.1% SDS. Hybridizing plaques were detected by autoradiography. The number of cDNAs that hybridized to the bovine probe and the number of cDNAs screened are indicated in Table 2 below.

Table 2

cDNA Library	Type	Positive Plaques	Plaques Screened
Human SW 1088 glioblastoma	dT-primed	1	1.5x10 ⁶
Human lung	dT-primed	2	1.5x10 ⁶
Human SW 1088 glioblastoma	dT-primed	4	1.5x10 ⁶

Plasmids designated cgbS2.1, cgbS3.1, cgbL23.1, cgbL27.1 and cgbS27.1 were excised *in vivo* from the lambda Zap

EP 0 652 960 B9 (W1B1)

clones and sequenced.

[0069] Clone cgbS3.1 contains 2060 bp of a PDE open reading frame followed by a putative intron. Analysis of clone cgbS2.1 reveals that it corresponds to clone cgbS3.1 positions 664 to 2060 and extends the PDE open reading frame an additional 585 bp before reading into a putative intron. The sequences of the putative 5' untranslated region and the protein encoding portions of the cgbS2.1 and cgbS3.1 clones are set out in SEQ ID NOs: 11 and 12, respectively. Combining the two cDNAs yields a sequence containing approximately 2.7 kb of an open reading encoding a PDE. The three other cDNAs did not extend any further 5' or 3' than cDNA cgbS3.1 or cDNA cgbS2.1.

[0070] To isolate additional cDNAs, probes specific for the 5' end of clone cgbS3.1 and the 3' end of clone cgbS2.1 were prepared and used to screen a SW1088 glioblastoma cDNA library and a human aorta cDNA library. A 5' probe was derived from clone cgbS3.1 by PCR using the primers cgbS3.1S311 and cgbL23.1A1286 whose sequences are set out in SEQ ID NOs: 8 and 9, respectively, and below.

Primer cgbS3.1S311 (SEQ ID NO: 13)

5' GCCACCAGAGAAATGGTC 3'

Primer cgbL23.1A1286 (SEQ ID NO: 14)

5' ACAATGGGTCTAAGAGGC 3'

The PCR reaction was carried out in a 50 ul reaction volume containing 50 pg cgbS3.1 cDNA, 0.2mM dNTP, 10 ug/ml each primer, 50 mM KCl, 10mM Tris-HCl pH 8.2, 1.5mM MgCl₂ and Taq polymerase. After an initial four minute denaturation at 94°C, 30 cycles of one minute at 94°C, two minutes at 50°C and four minutes at 72°C were carried out. An approximately 0.2 kb fragment was generated by the PCR reaction which corresponded to nucleotides 300-496 of clone cgbS3.1.

[0071] A 3' probe was derived from cDNA cgbS2.1 by PCR using the oligos cgbL23.1S1190 and cgbS2.1A231 whose sequences are set out below.

Primer cgbL23.1S1190 (SEQ ID NO: 15)

5' TCAGTGCATGTTTGCTGC 3'

Primer cgbS2.1A231 (SEQ ID NO: 16)

5' TACAAACATGTTCATCAG 3'

The PCR reaction as carried out similarly to that described above for generating the 5' probe, and yielded a fragment of approximately 0.8kb corresponding to nucleotides 1358-2139 of cDNA cgbS2.1. The 3' 157 nucleotides of the PCR fragment (not shown in SEQ ID NO: 12) are within the presumptive intron.

[0072] The two PCR fragments were purified and isolated by agarose gel electrophoresis, and were labelled with radioactive nucleotides by random priming. A random-primed SW1088 glioblastoma cDNA library (1.5x10⁶ plaques) was screened with the labelled fragments as described above, and 19 hybridizing plaques were isolated. An additional 50 hybridizing plaques were isolated from a human aorta cDNA library (dT and random primed, Clontech, Palo Alto, CA).

[0073] Plasmids were excised *in vivo* from some of the positive lambda Zap clones and sequenced. A clone designated cgbS53.2, the sequence of which is set out in SEQ ID NO: 17, contains an approximately 1.1 kb insert whose sequence overlaps the last 61 bp of cgbS3.1 and extends the open reading frame an additional 135 bp beyond that found in cgbS2.1. The clone contains a termination codon and approximately 0.3 kB of putative 3' untranslated sequence.

Generation of a Composite cDNA Encoding Human cGB-PDE

[0074] Clones cgbS3.1, cgbS2.1 and cgbS53.2 were used as described in the following paragraphs to build a composite cDNA that contained a complete human cGB-PDE opening reading frame. The composite cDNA is designated cgbmet156-2 and was inserted in the yeast ADH1 expression vector pBNY6N.

EP 0 652 960 B9 (W1B1)

[0075] First, a plasmid designated cgb stop-2 was generated that contained the 3' end of the cGB-PDE open reading frame. A portion of the insert of the plasmid was generated by PCR using clone cgbS53.2 as a template. The PCR primers utilized were cgbS2.1S1700 and cgbstop-2.

5

Primer cgbS2.1S1700 (SEQ ID NO: 18)

5' TTTGGAAGATCCTCATCA 3'

10

Primer cgbstop-2 (SEQ ID NO: 19)

5' ATGTCTCGAGTCAGTTCGCTTGGCCTG 3'

15 The PCR reaction was carried out in 50 ul containing 50 pg template DNA, 0.2mM dNTPs, 20mM Tris-HCl pH 8.2, 10mM KCl, 6mM (NH₄)₂SO₄, 1.5mM MgCl₂, 0.1% Triton-X-100, 500ng each primer and 0.5 units of Pfu polymerase (Stratagene). The reaction was heated to 94°C for 4 minutes and then 30 cycles of 1 minute at 94°C, 2 minutes at 50°C and four minutes at 72°C were performed. The polymerase was added during the first cycle at 50°C. The resulting PCR product was phenol/chloroform extracted, chloroform extracted, ethanol precipitated and cut with the restriction enzymes *Bcl*I and *Xho*I. The restriction fragment was purified on an agarose gel and eluted.

20

[0076] This fragment was ligated to the cDNA cgbS2.1 that had been grown in dam *E. coli*, cut with the restriction enzymes *Bcl*I and *Xho*I, and gel-purified using the Promega magic PCR kit. The resulting plasmid was sequenced to verify that cgbstop-2 contains the 3' portion of the cGB-PDE open reading frame.

25

[0077] Second, a plasmid carrying the 5' end of the human cGB-PDE open reading frame was generated. Its insert was generated by PCR using clone cgbS3.1 as a template. PCR was performed as described above using primers cgbmet156 and cgbS2.1A2150.

30

Primer cgbmet156 (SEQ ID NO: 20)

5' TACAGAATTCTGACCATGGAGCGGGCCGGC 3'

35

Primer cgbS2.1A2150 (SEQ ID NO: 21)

5' CATTCTAAGCGGATACAG 3'

40

The resulting PCR fragment was phenol/chloroform extracted, chloroform extracted, ethanol precipitated and purified on a Sepharose CL-6B column. The fragment was cut with the restriction enzymes *Eco*RV and *Eco*RI, run on an agarose gel and purified by spinning through glass wool. Following phenol/chloroform extraction, chloroform extraction and ethanol precipitation, the fragment was ligated into *Eco*RI/*Eco*RV digested BluescriptII SK(+) to generate plasmid cgbmet156. The DNA sequence of the insert and junctions was determined. The insert contains a new *Eco*RI site and an additional 5 nucleotides that together replace the original 155 nucleotides 5' of the initiation codon. The insert extends to an *Eco*RV site beginning 531 nucleotides from the initiation codon.

45

[0078] The 5' and 3' portions of the cGB-PDE open reading frame were then assembled in vector pBNI6a. The vector pBNI6a was cut with *Eco*RI and *Xho*I, isolated from a gel and combined with the agarose gel purified *Eco*RI/*Eco*RV fragment from cgbmet156 and the agarose gel purified *Eco*RV/*Xho*I fragment from cgbstop-2. The junctions of the insert were sequenced and the construct was named hcbgmet156-2 6a.

50

[0079] The cGB-PDE insert from hcbgmet156-2 6a was then moved into the expression vector pBNI6n. Expression of DNA inserted in this vector is directed from the yeast ADHI promoter and terminator. The vector contains the yeast 2 micron origin of replication, the pUC19 origin of replication and an ampicillin resistance gene. Vector pBNI6n was cut with *Eco*RI and *Xho*I and gel-purified. The *Eco*RI/*Xho*I insert from hcbgmet156-2 6a was gel purified using Promega magic PCR columns and ligated into the cut pBNI6n. All new junctions in the resulting construct, hcbgmet156-2 6n, were sequenced. The DNA and deduced amino acid sequences of the insert of hcbgmet156-2 6n which encodes a composite human cGB-PDE is set out in SEQ ID NOs: 22 and 23. The insert extends from the first methionine in clone cgbS3.1 (nucleotide 156) to the stop codon (nucleotide 2781) in the composite cDNA. Because the methionine is the most 5' methionine in clone cgbS3.1 and because there are no stop codons in frame with the methionine and upstream of it, the insert in pBNI6n may represent a truncated form of the open reading frame.

55

Variant cDNAs

[0080] Four human cGB-PDE cDNAs that are different from the hcgbmet156-2 6n composite cDNA have been isolated. One cDNA, cgbL23.1, is missing an internal region of hcgbmet156-2 6n (nucleotides 997-1000 to 1444-1447). The exact end points of the deletion cannot be determined from the cDNA sequence at those positions. Three of the four variant cDNAs have 5' end sequences that diverge from the hcgbmet156-2 6n sequence upstream of nucleotide 151 (cDNAs cgbA7f, cgbA5C, cgb12). These cDNAs presumably represent alternatively spliced or unspliced mRNAs.

Example 7

[0081] The composite human cGB-PDE cDNA construct, hcgbmet156-2 6n, was transformed into the yeast strain YKS45 (ATCC 74225) (MAT α his3 trp1 ura3 leu3 pde1::HIS3 pde2::TRP1) in which two endogenous PDE genes are deleted. Transformants complementing the leu⁻ deficiency of the YKS45 strain were selected and assayed for cGB-PDE activity. Extracts from cells bearing the plasmid hcgbmet156-2 6n were determined to display cyclic GMP-specific phosphodiesterase activity by the assay described below.

[0082] One liter of YKS45 cells transformed with the plasmid cgbmet156-2 6n and grown in SC-leu medium to a density of $1-2 \times 10^7$ cells/ml was harvested by centrifugation, washed once with deionized water, frozen in dry ice/ethanol and stored at -70°C . Cell pellets (1-1.5 ml) were thawed on ice in the presence of an equal volume of 25mM Tris-Cl (pH 8.0)/5mM EDTA/5mM EGTA/1mM o-phenanthroline/0.5mM AEBSF (Calbiochem)/0.1 % β -mercaptoethanol and 10 ug/ml each of aprotinin, leupeptin, and pepstatin A. The thawed cells were added to 2 ml of acid-washed glass beads (425-600 μM , Sigma) in 15 ml Corex tube. Cells were broken with 4 cycles consisting of a 30 second vortexing on setting 1 followed by a 60 second incubation on ice. The cell lysate was centrifuged at 12,000 x g for 10 minutes and the supernatant was passed through a 0.8 μ filter. The supernatant was assayed for cGMP PDE activity as follows. Samples were incubated for 20 minutes at 30°C in the presence of 45mM Tris-Cl (pH 8.0), 2mM EGTA, 1mM EDTA, 0.2mg/ml BSA, 5mM MgCl₂, 0.2mM o-phenanthroline, 2ug/ml each of pepstatin A, leupeptin, and aprotinin, 0.1mM AEBSF, 0.02% β -mercaptoethanol and 0.1mM [³H]cGMP as substrate. [¹⁴C]-AMP (0.5 nCi/assay) was added as a recovery standard. The reaction was terminated with stop buffer (0.1M ethanolamine pH 9.0, 0.5M ammonium sulfate, 10mM EDTA, 0.05 % SDS final concentration). The product was separated from the cyclic nucleotide substrate by chromatography on BioRad Affi-Gel 601. The sample was applied to a column containing approximately 0.25 ml of Affi-Gel 601 equilibrated in column buffer (0.1M ethanolamine pH 9.0 containing 0.5M ammonium sulfate). The column was washed five times with 0.5 ml of column buffer. The product was eluted with four 0.5 ml aliquots of 0.25 acetic acid and mixed with 5 ml Ecolume (ICN Biochemicals). The radioactive product was measured by scintillation counting.

Example 8

[0083] Analysis of expression of cGB-PDE mRNA in human tissues was carried out by RNase protection assay.

[0084] A probe corresponding to a portion of the putative cGMP binding domain of cGB-PDE (402 bp corresponding to nucleotides 1450 through 1851 of SEQ ID NO: 13) was generated by PCR. The PCR fragment was inserted into the EcoRI site of the plasmid pBSII SK(-) to generate the plasmid RP3. RP3 plasmid DNA was linearized with XbaI and antisense probes were generated by a modification of the Stratagene T7 RNA polymerase kit. Twenty-five ng of linearized plasmid was combined with 20 microcuries of alpha ³²P rUTP (800 Ci/mmol, 10 mCi/ml), 1X transcription buffer (40mM TrisCl, pH 8, 8mM MgCl₂, 2mM spermidine, 50mM NaCl), 0.25mM each rATP, rGTP and rCTP, 0.1 units of RNase Block II, 5mM DTT, 8 μM rUTP and 5 units of T7 RNA Polymerase in a total volume of 5 μl . The reaction was allowed to proceed 1 hour at room temperature and then the DNA template was removed by digestion with RNase free DNase. The reaction was diluted into 100 μl of 40mM TrisCl, pH 8, 6mM MgCl₂ and 10mM NaCl. Five units of RNase-free DNase were added and the reaction was allowed to continue another 15 minutes at 37°C . The reaction was stopped by a phenol extraction followed by a phenol chloroform extraction. One half volume of 7.5M NH₄OAc was added and the probe was ethanol precipitated.

[0085] The RNase protection assays were carried out using the Ambion RNase Protection kit (Austin, TX) and 10 μg RNA isolated from human tissues by an acid guanidinium extraction method. Expression of cGB-PDE mRNA was easily detected in RNA extracted from skeletal muscle, uterus, bronchus, skin, right saphenous vein, aorta and SW1088 glioblastoma cells. Barely detectable expression was found in RNA extracted from right atrium, right ventricle, kidney cortex, and kidney medulla. Only complete protection of the RP3 probe was seen. The lack of partial protection argues against the cDNA cgbL23.1 (a variant cDNA described in Example 7) representing a major transcript, at least in these RNA samples.

Example 9

[0086] Polyclonal antisera was raised to *E. coli*-produced fragments of the human cGB-PDE.

[0087] A portion of the human cGB-PDE cDNA (nucleotides 1668-2612 of SEQ ID NO: 22, amino acids 515-819 of SEQ ID NO: 23) was amplified by PCR and inserted into the *E. coli* expression vector pGEX2T (Pharmacia) as a *Bam*HI/*Eco*RI fragment. The pGEX2T plasmid carries an ampicillin resistance gene, an *E. coli* *lacI^q* gene and a portion of the *Schistosoma japonicum* glutathione-S-transferase (GST) gene. DNA inserted in the plasmid can be expressed as a fusion protein with GST and can then be cleaved from the GST portion of the protein with thrombin. The resulting plasmid, designated cgbPE3, was transformed into *E. coli* strain LE392 (Stratagene). Transformed cells were grown at 37°C to an OD600 of 0.6. IPTG (isopropylthioalactopyranoside) was added to 0.1mM and the cells were grown at 37°C for an additional 2 hours. The cells were collected by centrifugation and lysed by sonication. Cell debris was removed by centrifugation and the supernatant was fractionated by SDS-PAGE. The gel was stained with cold 0.4M KCl and the GST-cgb fusion protein band was excised and electroeluted. The PDE portion of the protein was separated from the GST portion by digestion with thrombin. The digest was fractionated by SDS-PAGE, the PDE protein was electroeluted and injected subcutaneously into a rabbit. The resultant antisera recognizes both the bovine cGB-PDE fragment that was utilized as antigen and the full length human cGB-PDE protein expressed in yeast (see Example 8).

Example 10

[0088] Polynucleotides encoding various cGB-PDE analogs and cGB-PDE fragments were generated by standard methods.

A. cGB-PDE Analogs

[0089] All known cGMP-binding PDEs contain two internally homologous tandem repeats within their putative cGMP-binding domains. In the bovine cGB-PDE of the invention, the repeats span at least residues 228-311 (repeat A) and 410-500 (repeat B) of SEQ ID NO: 10. Site-directed mutagenesis of an aspartic acid that is conserved in repeats A and B of all known cGMP-binding PDEs was used to create analogs of cGB-PDE having either Asp-289 replaced with Ala (D289A) or Asp-478 replaced with Ala (D478A). Recombinant wild type (WT) bovine and mutant bovine cGB-PDEs were expressed in COS-7 cells. cGB-PDE purified from bovine lung (native cGB-PDE) and WT cGB-PDE displayed identical cGMP-binding kinetics with a K_d of approximately 2 μ M and a curvilinear dissociation profile ($t_{1/2}$ = 1.3 hours at 4°C). This curvilinearity may have been due to the presence of distinct high affinity (slow) and low affinity (fast) sites of cGMP binding. The D289A mutant had significantly decreased affinity for cGMP (K_d > 20 μ M) and a single rate of cGMP-association ($t_{1/2}$ = 0.5 hours), that was similar to that calculated for the fast site of WT and native cGB-PDE. This suggested the loss of a slow cGMP-binding site in repeat A of this mutant. Conversely, the D478A mutant showed higher affinity for cGMP (K_d of approximately 0.5 μ M) and a single cGMP-dissociation rate ($t_{1/2}$ = 2.8 hours) that was similar to the calculated rate of the slow site of WT and native cGB-PDE. This suggested the loss of a fast site when repeat B was modified. These results indicate that dimeric cGB-PDE possesses two homologous but kinetically distinct cGMP-binding sites, with the conserved aspartic acid being critical for interaction with cGMP at each site. See, Colbran *et al.*, *FASEB J.*, 8: Abstract 2149 (May 15, 1994).

B. Amino-Terminal Truncated cGB-PDE Polypeptides

[0090] A truncated human cGB-PDE polypeptide including amino acids 516-875 of SEQ ID NO: 23 was expressed in yeast. A cDNA insert extending from the *Nco*I site at nucleotide 1555 of SEQ ID NO: 22 through the *Xho*I site at the 3' end of SEQ ID NO: 22 was inserted into the ADH2 yeast expression vector YEpC-PADH2d [Price *et al.*, *Meth. Enzymol.*, 185: 308-318 (1990)] that had been digested with *Nco*I and *Sal*I to generate plasmid YEpC-PADH2d HcGB. The plasmid was transformed into spheroplasts of the yeast strain yBJ2-54 (*prc1-407 prb1-1122 pep4-3 leu2 trp1 ura3-52 Δ pde1::URA3, HIS3 Δ pde2::TRP1 cir^o*). The endogenous PDE genes are deleted in this strain. Cells were grown in SC-leu media with 2% glucose to 10⁷ cells/ml, collected by filtration and grown 24 hours in YEP media containing 3% glycerol. Cells were pelleted by centrifugation and stored frozen. Cells were disrupted with glass beads and the cell homogenate was assayed for phosphodiesterase activity essentially as described in Prpic *et al.*, *Anal. Biochem.*, 208: 155-160 (1993). The truncated human cGB-PDE polypeptide exhibited phosphodiesterase activity.

C. Carboxy-Terminal Truncated cGB-PDE Polypeptides

[0091] Two different plasmids encoding carboxy-terminal truncated human cGB-PDE polypeptides were constructed.

[0092] Plasmid pBJ6-84Hin contains a cDNA encoding amino acids 1-494 of SEQ ID NO: 23 inserted into the *Nco*I

EP 0 652 960 B9 (W1B1)

and Sall sites of vector YEpC-PADH2d. The cDNA insert extends from the NcoI site at nucleotide position 10 of SEQ ID NO: 22 through the HindIII site at nucleotide position 1494 of SEQ ID NO: 22 followed by a linker and the Sall site of YEpC-PADH2d.

5 [0093] Plasmid pBJ6-84Ban contains a cDNA encoding amino acids 1-549 of SEQ ID NO: 23 inserted into the NcoI and Sall sites of vector YEpC-PADH2d. The cDNA insert extends from the NcoI site at nucleotide position 10 of SEQ ID NO: 22 through the BanI site at nucleotide position 1657 of SEQ ID NO: 22 followed by a linker and the Sall site of YEpC-PADH2d.

[0094] The truncated cGB-PDE polypeptides are useful for screening for modulators of cGB-PDE activity.

10 Example 11

[0095] Monoclonal antibodies reactive with human cGB-PDE were generated.

15 [0096] Yeast yBJ2-54 containing the plasmid YEpADH2 HcGB (Example 10B) were fermented in a New Brunswick Scientific 10 liter Microferm. The cGB-PDE cDNA insert in plasmid YEpADH2 HcGB extends from the NcoI site at nucleotide 12 of SEQ ID NO: 22 to the XhoI site at the 3' end of SEQ ID NO: 22. An inoculum of 4×10^9 cells was added to 8 liters of media containing SC-leu, 5% glucose, trace metals, and trace vitamins. Fermentation was maintained at 26°C, agitated at 600 rpm with the standard microbial impeller, and aerated with compressed air at 10 volumes per minute. When glucose decreased to 0.3% at 24 hours post-inoculation the culture was infused with 2 liters of 5X YEP media containing 15% glycerol. At 66 hours post-inoculation the yeast from the ferment was harvested by centrifugation at 4,000 x g for 30 minutes at 4°C. Total yield of biomass from this fermentation approached 350 g wet weight.

20 [0097] Human cGB-PDE enzyme was purified from the yeast cell pellet. Assays for PDE activity using 1 mM cGMP as substrate was employed to follow the chromatography of the enzyme. All chromatographic manipulations were performed at 4°C.

25 [0098] Yeast (29g wet weight) were resuspended in 70ml of buffer A (25mM Tris pH 8.0, 0.25mM DTT, 5mM MgCl₂, 10μM ZnSO₄, 1mM benzamidine) and lysed by passing through a microfluidizer at 22-24,000 psi. The lysate was centrifuged at 10,000 x g for 30 minutes and the supernatant was applied to a 2.6 x 28 cm column containing Pharmacia Fast Flow Q anion exchange resin equilibrated with buffer B containing 20mM BisTris-propane pH 6.8, 0.25mM DTT, 1mM MgCl₂, and 10μM ZnSO₄. The column was washed with 5 column volumes of buffer B containing 0.125M NaCl and then developed with a linear gradient from 0.125 to 1.0M NaCl. Fractions containing the enzyme were pooled and applied directly to a 5 x 20 cm column of ceramic hydroxyapatite (BioRad) equilibrated in buffer C containing 20mM BisTris-propane pH 6.8, 0.25mM DTT, 0.25MKCl, 1mM MgCl₂, and 10μM ZnSO₄. The column was washed with 5 column volumes of buffer C and eluted with a linear gradient from 0 to 250mM potassium phosphate in buffer C. The pooled enzyme was concentrated 8-fold by ultrafiltration (YM30 membrane, Amicon). The concentrated enzyme was chromatographed on a 2.6 x 90 cm column of Pharmacia Sephacryl S300 (Piscataway, NJ) equilibrated in 25mM BisTris-propane pH 6.8, 0.25mM DTT, 0.25M NaCl, 1mM MgCl₂, and 20μM ZnSO₄. Approximately 4 mg of protein was obtained. The recombinant human cGB-PDE enzyme accounted for approximately 90% of protein obtained as judged by SDS polyacrylamide gel electrophoresis followed by Coomassie blue staining.

30 [0099] The purified protein was used as an antigen to raise monoclonal antibodies. Each of 19 week old Balb/c mice (Charles River Biotechnical Services, Inc., Wilmington, Mass.) was immunized sub-cutaneously with 50 ug purified human cGB-PDE enzyme in a 200 ul emulsion consisting of 50% Freund's complete adjuvant (Sigma Chemical Co.). Subsequent boosts on day 20 and day 43 were administered in incomplete Freund's adjuvant. A pre-fusion boost was done on day 86 using 50 ug enzyme in PBS. The fusion was performed on day 90.

35 [0100] The spleen from mouse #1817 was removed sterilely and placed in 10ml serum free RPMI 1640. A single-cell suspension was formed and filtered through sterile 70-mesh Nitex cell strainer (Becton Dickinson, Parsippany, New Jersey), and washed twice by centrifuging at 200 g for 5 minutes and resuspending the pellet in 20 ml serum free RPMI. Thymocytes taken from 3 naive Balb/c mice were prepared in a similar manner.

40 [0101] NS-1 myeloma cells, kept in log phase in RPMI with 11% Fetalclone (FBS) (Hyclone Laboratories, Inc., Logan, Utah) for three days prior to fusion, were centrifuged at 200 g for 5 minutes, and the pellet was washed twice as described in the foregoing paragraph. After washing, each cell suspension was brought to a final volume of 10 ml in serum free RPMI, and 20 μl was diluted 1:50 in 1 ml serum free RPMI. 20 μl of each dilution was removed, mixed with 20 μl 0.4% trypan blue stain in 0.85% saline (Gibco), loaded onto a hemocytometer (Baxter Healthcare Corp., Deerfield, Illinois) and counted.

45 [0102] Two x 10⁸ spleen cells were combined with 4.0 x 10⁷ NS-1 cells, centrifuged and the supernatant was aspirated. The cell pellet was dislodged by tapping the tube and 2 ml of 37°C PEG 1500 (50% in 75 mM Hepes, pH 8.0) (Boehringer Mannheim) was added with stirring over the course of 1 minute, followed by adding 14 ml of serum free RPMI over 7 minutes. An additional 16 ml RPMI was added and the cells were centrifuged at 200 g for 10 minutes. After discarding the supernatant, the pellet was resuspended in 200 ml RPMI containing 15% FBS, 100 μM sodium hypoxanthine, 0.4 μM aminopterin, 16 μM thymidine (HAT) (Gibco), 25 units/ml IL-6 (Boehringer Mannheim) and 1.5

EP 0 652 960 B9 (W1B1)

x 10⁶ thymocytes/ml. The suspension was first placed in a T225 flask (Corning, United Kingdom) at 37°C for two hours before being dispensed into ten 96-well flat bottom tissue culture plates (Corning, United Kingdom) at 200 µl/well. Cells in plates were fed on days 3, 4, 5 post fusion day by aspirating approximately 100 µl from each well with an 20 G needle (Becton Dickinson), and adding 100 µl/well plating medium described above except containing 10 units/ml IL-6 and lacking thymocytes.

[0103] The fusion was screened initially by ELISA. Immulon 4 plates (Dynatech) were coated at 4°C overnight with purified recombinant human cGB-PDE enzyme (100ng/well in 50mM carbonate buffer pH9.6). The plates were washed 3X with PBS containing 0.05 % Tween 20 (PBST). The supernatants from the individual hybridoma wells were added to the enzyme coated wells (50 µl/well). After incubation at 37°C for 30 minutes, and washing as above, 50 µl of horseradish peroxidase conjugated goat anti-mouse IgG(fc) (Jackson ImmunoResearch, West Grove, Pennsylvania) diluted 1:3500 in PBST was added. Plates were incubated as above, washed 4X with PBST and 100 µl substrate consisting of 1 mg/ml o-phenylene diamine (Sigma) and 0.1 µl/ml 30% H₂O₂ in 100 mM citrate, pH 4.5, was added. The color reaction was stopped in 5 minutes with the addition of 50 µl of 15% H₂SO₄. A₄₉₀ was read on a plate reader (Dynatech).

[0104] Wells CSG, E4D, F1G, F9H, F11G, J4A, and JSD were picked and renamed 102A, 102B, 102C, 102D, 102E, 102F, and 102G respectively, cloned two or three times, successively, by doubling dilution in RPMI, 15% FBS, 100µM sodium hypoxanthine, 16µM thymidine, and 10 units/ml IL-6. Wells of clone plates were scored visually after 4 days and the number of colonies in the least dense wells were recorded. Selected wells of the each cloning were tested by ELISA.

[0105] The monoclonal antibodies produced by above hybridomas were isotyped in an ELISA assay. Results showed that monoclonal antibodies 102A to 102E were IgG1, 102F was IgG2b and 102G was IgG2a.

[0106] All seven monoclonal antibodies reacted with human cGS-PDE as determined by Western analysis.

Example 12

[0107] Developing modulators of the biological activities of specific PDEs requires differentiating PDE isozymes present in a particular assay preparation. The classical enzymological approach of isolating PDEs from natural tissue sources and studying each new isozyme is hampered by the limits of purification techniques and the inability to definitively assess whether complete resolution of a isozyme has been achieved. Another approach has been to identify assay conditions which might favor the contribution of one isozyme and minimize the contribution of others in a preparation. Still another approach has been the separation of PDEs by immunological means. Each of the foregoing approaches for differentiating PDE isozymes is time consuming and technically difficult. As a result many attempts to develop selective PDE modulators have been performed with preparations containing more than one isozyme. Moreover, PDE preparations from natural tissue sources are susceptible to limited proteolysis and may contain mixtures of active proteolytic products that have different kinetic, regulatory and physiological properties than the full length PDEs.

[0108] Recombinant cGB-PDE polypeptide products of the invention greatly facilitate the development of new and specific cGB-PDE modulators. The use of human recombinant enzymes for screening for modulators has many inherent advantages. The need for purification of an isozyme can be avoided by expressing it recombinantly in a host cell that lacks endogenous phosphodiesterase activity (e.g., yeast strain YKS45 deposited as ATCC 74225). Screening compounds against human protein avoids complications that often arise from screening against non-human protein where a compound optimized on a non-human protein may fail to be specific for or react with the human protein. For example, a single amino acid difference between the human and rodent 5HT_{1B} serotonin receptors accounts for the difference in binding of a compound to the receptors. [See Oskenberg *et al.*, *Nature*, 360: 161-163 (1992)]. Once a compound that modulates the activity of the cGB-PDE is discovered, its selectivity can be evaluated by comparing its activity on the cGB-PDE to its activity on other PDE isozymes. Thus, the combination of the recombinant cGB-PDE products of the invention with other recombinant PDE products in a series of independent assays provides a system for developing selective modulators of cGB-PDE. Selective modulators may include, for example, antibodies and other proteins or peptides which specifically bind to the cGB-PDE or cGB-PDE nucleic acid, oligonucleotides which specifically bind to the cGB-PDE (see Patent Cooperation Treaty International Publication No. WO93/05182 published March 18, 1993 which describes methods for selecting oligonucleotides which selectively bind to target biomolecules) or cGB-PDE nucleic acid (e.g., antisense oligonucleotides) and other non-peptide natural or synthetic compounds which specifically bind to the cGB-PDE or cGB-PDE nucleic acid. Mutant forms of the cGB-PDE which alter the enzymatic activity of the cGB-PDE or its localization in a cell are also contemplated. Crystallization of recombinant cGB-PDE alone and bound to a modulator, analysis of atomic structure by X-ray crystallography, and computer modelling of those structures are methods useful for designing and optimizing non-peptide selective modulators. See, for example, Erickson *et al.*, *Ann. Rep. Med. Chem.*, 27: 271-289 (1992) for a general review of structure-based drug design.

[0109] Targets for the development of selective modulators include, for example: (1) the regions of the cGB-PDE which contact other proteins and/or localize the cGB-PDE within a cell, (2) the regions of the cGB-PDE which bind

EP 0 652 960 B9 (W1B1)

substrate, (3) the allosteric cGMP-binding site(s) of cGB-PDE, (4) the metal-binding regions of the cGB-PDE, (5) the phosphorylation site(s) of cGB-PDE and (6) the regions of the cGB-PDE which are involved in dimerization of cGB-PDE subunits.

5 SEQUENCE LISTING

(1) GENERAL INFORMATION:

[0110]

10

(i) APPLICANT: The Board of Regents of the University of Washington and ICOS Corporation

(ii) TITLE OF INVENTION: Cyclic GMP-Binding, Cyclic GMP-Specific Phosphodiesterase Materials and Methods

15

(iii) NUMBER OF SEQUENCES: 23

(iv) CORRESPONDENCE ADDRESS:

20

- (A) ADDRESSEE: Marshall, O'Toole, Gerstein, Murray & Borun
- (B) STREET: 6300 Sears Tower, 233 S. Wacker Drive
- (C) CITY: Chicago
- (D) STATE: Illinois
- (E) COUNTRY: USA
- (F) ZIP: 60606

25

(v) COMPUTER READABLE FORM:

30

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

35

- (A) APPLICATION NUMBER:
- (B) FILING DATE:
- (C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

40

- (A) APPLICATION NUMBER: US 08/068,051
- (B) FILING DATE: 27-MAY-1993

(viii) ATTORNEY/AGENT INFORMATION:

45

- (A) NAME: Noland, Greta E.
- (B) REGISTRATION NUMBER: 35,302
- (C) REFERENCE/DOCKET NUMBER: 32083

(ix) TELECOMMUNICATION INFORMATION:

50

- (A) TELEPHONE: (312) 474-6300
- (B) TELEFAX: (312) 474-0448
- (C) TELEX: 25-3856

55

EP 0 652 960 B9 (W1B1)

(2) INFORMATION FOR SEQ ID NO:1:

[0111]

5 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

15

Arg Glu Xaa Asp Ala Asn Arg Ile Asn Tyr Met Tyr Ala Gln Tyr Val
1 5 10 15
Lys Asn Thr Met
20

20

(2) INFORMATION FOR SEQ ID NO:2:

[0112]

25 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

35

Gln Ser Leu Ala Ala Ala Val Val Pro
1 5

(2) INFORMATION FOR SEQ ID NO:3:

40

[0113]

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: peptide

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

55

Phe Asp Asn Asp Glu Gly Glu Gln
1 5

(2) INFORMATION FOR SEQ ID NO:4:

[0114]

5 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

TTYGAYAAYG AYGARGGNGA RCA

23

(2) INFORMATION FOR SEQ ID NO:5:

[0115]

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

AARCTRTRRC TRCTYCCNCT YGT

23

(2) INFORMATION FOR SEQ ID NO:6

[0116]

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6

AAYTAYATGT AYGNCARTA YGT

23

(2) INFORMATION FOR SEQ ID NO:7

[0117]

- 5 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 23 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - 10 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (iv) ANTI-SENSE: YES
- 15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7

20 **TTRATRTACA TRCGNGTYAT RCA**

23

(2) INFORMATION FOR SEQ ID NO:8

[0118]

- 25 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 36 base pairs
 - (B) TYPE: nucleic acid
 - 30 (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (iv) ANTI-SENSE: YES
- 35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8

40 **TTRATRTACA TRCGNGTYAT RCANTTYTTR TGNTAC**

36

(2) INFORMATION FOR SEQ ID NO:9

[0119]

- 45 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 4474 base pairs
 - (B) TYPE: nucleic acid
 - 50 (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- 55 (ix) FEATURE:
- (A) NAME/KEY: CDS
 - (B) LOCATION: 99..2723

EP 0 652 960 B9 (W1B1)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9

	GGGAGGGTCT CGAGGCGAGT TCTGCTCCTC GGAGGGAGGG ACCCCAGCTG GAGTGAAAA	60
5	CCAGCACCAG CTGACCGCAG AGACACGCCG CGCTGATC ATG GAG AGG GCC GGC	113
	Met Glu Arg Ala Gly	
	1 5	
	CCC GGC TGC CGC GCG GCC GCA ACA GCA ATG GGA CCA GGA CTC GGT CGA	161
10	Pro Gly Cys Arg Ala Ala Ala Thr Ala Met Gly Pro Gly Leu Gly Arg	
	10 15 20	
	AGC GTG GCT GGA CGA TCA CTG GGA CTT TAC CTT CTC TAC TTT GTT AGG	209
	Ser Val Ala Gly Arg Ser Leu Gly Leu Tyr Leu Leu Tyr Phe Val Arg	
	25 30 35	
15	AAA GGC ACC AGA GAA ATG GTC AAC GCA TGG TTT GCT GAG AGA GTT CAC	257
	Lys Gly Thr Arg Glu Met Val Asn Ala Trp Phe Ala Glu Arg Val His	
	40 45 50	
	ACC ATT CCT GTG TGC AAG GAA GGA ATC AAG GGC CAC ACG GAA TCC TGC	305
20	Thr Ile Pro Val Cys Lys Glu Gly Ile Lys Gly His Thr Glu Ser Cys	
	55 60 65	
	TCT TGC CCC TTG CAG CCA AGT CCC CGT GCA GAG AGC AGT GTC CCT GGA	353
	Ser Cys Pro Leu Gln Pro Ser Pro Arg Ala Glu Ser Ser Val Pro Gly	
	70 75 80 85	
25	ACA CCA ACC AGG AAG ATC TCT GCC TCT GAA TTC GAT CGG CCG CTT AGA	401
	Thr Pro Thr Arg Lys Ile Ser Ala Ser Glu Phe Asp Arg Pro Leu Arg	
	90 95 100	
	CCC ATC GTT ATC AAG GAT TCT GAG GGA ACT GTG AGC TTC CTC TCT GAC	449
30	Pro Ile Val Ile Lys Asp Ser Glu Gly Thr Val Ser Phe Leu Ser Asp	
	105 110 115	
	TCA GAC AAG AAG GAA CAG ATG CCT CTA ACC TCC CCA CGG TTT GAT AAT	497
	Ser Asp Lys Lys Glu Gln Met Pro Leu Thr Ser Pro Arg Phe Asp Asn	
	120 125 130	
35	GAT GAA GGG GAC CAG TGC TCG AGA CTC TTG GAA TTA GTG AAA GAT ATT	545
	Asp Glu Gly Asp Gln Cys Ser Arg Leu Leu Glu Leu Val Lys Asp Ile	
	135 140 145	
	TCT AGT CAC TTG GAT GTC ACA GCC TTA TGT CAC AAA ATT TTC TTG CAC	593
	Ser Ser His Leu Asp Val Thr Ala Leu Cys His Lys Ile Phe Leu His	
	150 155 160 165	
40	ATC CAT GGA CTC ATC TCC GCC GAC CGC TAC TCC TTA TTC CTC GTC TGT	641
	Ile His Gly Leu Ile Ser Ala Asp Arg Tyr Ser Leu Phe Leu Val Cys	
	170 175 180	
	GAG GAC AGC TCC AAC GAC AAG TTT CTT ATC AGC CGC CTC TTT GAT GTT	689
45	Glu Asp Ser Ser Asn Asp Lys Phe Leu Ile Ser Arg Leu Phe Asp Val	
	185 190 195	
	GCA GAA GGT TCA ACA CTG GAA GAA GCT TCA AAC AAC TGC ATC CGC TTA	737
	Ala Glu Gly Ser Thr Leu Glu Glu Ala Ser Asn Asn Cys Ile Arg Leu	
	200 205 210	

EP 0 652 960 B9 (W1B1)

	GAG	TGG	AAC	AAA	GGC	ATC	GTG	GGA	CAC	GTG	GCC	GCT	TTT	GGC	GAG	CCC	785
	Glu	Trp	Asn	Lys	Gly	Ile	Val	Gly	His	Val	Ala	Ala	Phe	Gly	Glu	Pro	
		215					220					225					
5	TTG	AAC	ATC	AAA	GAC	GCC	TAT	GAG	GAT	CCT	CGA	TTC	AAT	GCA	GAA	GTT	833
	Leu	Asn	Ile	Lys	Asp	Ala	Tyr	Glu	Asp	Pro	Arg	Phe	Asn	Ala	Glu	Val	
	230				235					240						245	
10	GAC	CAA	ATT	ACA	GGC	TAC	AAG	ACA	CAA	AGT	ATT	CTT	TGT	ATG	CCA	ATT	881
	Asp	Gln	Ile	Thr	Gly	Tyr	Lys	Thr	Gln	Ser	Ile	Leu	Cys	Met	Pro	Ile	
					250					255					260		
15	AAG	AAT	CAT	AGG	GAA	GAG	GTT	GTT	GGT	GTA	GCC	CAG	GCC	ATC	AAC	AAG	929
	Lys	Asn	His	Arg	Glu	Glu	Val	Val	Gly	Val	Ala	Gln	Ala	Ile	Asn	Lys	
				265					270					275			
20	AAA	TCA	GGA	AAT	GGT	GGG	ACA	TTC	ACT	GAA	AAA	GAC	GAA	AAG	GAC	TTT	977
	Lys	Ser	Gly	Asn	Gly	Gly	Thr	Phe	Thr	Glu	Lys	Asp	Glu	Lys	Asp	Phe	
			280				285						290				
25	GCT	GCT	TAC	TTG	GCA	TTT	TGT	GGA	ATT	GTT	CTT	CAT	AAT	GCT	CAA	CTC	1025
	Ala	Ala	Tyr	Leu	Ala	Phe	Cys	Gly	Ile	Val	Leu	His	Asn	Ala	Gln	Leu	
			295				300					305					
30	TAT	GAG	ACT	TCA	CTG	CTG	GAG	AAC	AAG	AGA	AAT	CAG	GTG	CTG	CTT	GAC	1073
	Tyr	Glu	Thr	Ser	Leu	Leu	Glu	Asn	Lys	Arg	Asn	Gln	Val	Leu	Leu	Asp	
	310				315						320					325	
35	CTT	GCT	AGC	TTA	ATT	TTT	GAA	GAA	CAA	CAA	TCA	TTA	GAA	GTA	ATT	CTA	1121
	Leu	Ala	Ser	Leu	Ile	Phe	Glu	Glu	Gln	Ser	Leu	Glu	Val	Ile	Leu		
					330					335					340		
40	AGG	AAA	ATA	GCT	GCC	ACT	ATT	ATC	TCT	CCC	ATG	CAG	GTG	CAG	AAA	TGC	1169
	Arg	Lys	Ile	Ala	Ala	Thr	Ile	Ile	Ser	Pro	Met	Gln	Val	Gln	Lys	Cys	
				345					350					355			
45	ACC	ATT	TTC	ATA	GTG	GAT	GAA	GAT	TGC	TCC	GAT	TCT	TTT	TCT	AGT	GTG	1217
	Thr	Ile	Phe	Ile	Val	Asp	Glu	Asp	Cys	Ser	Asp	Ser	Phe	Ser	Ser	Val	
				360				365					370				
50	TTT	CAC	ATG	GAG	TGT	GAG	GAA	TTA	GAA	AAA	TCG	TCA	GAT	ACT	TTA	ACA	1265
	Phe	His	Met	Glu	Cys	Glu	Glu	Leu	Glu	Lys	Ser	Ser	Asp	Thr	Leu	Thr	
				375			380					385					
55	CGG	GAA	CGT	GAT	GCA	ACC	AGA	ATC	AAT	TAC	ATG	TAT	GCT	CAG	TAT	GTC	1313
	Arg	Glu	Arg	Asp	Ala	Thr	Arg	Ile	Asn	Tyr	Met	Tyr	Ala	Gln	Tyr	Val	
	390				395					400						405	
60	AAA	AAT	ACC	ATG	GAA	CCA	CTT	AAT	ATC	CCA	GAC	GTC	AGT	AAG	GAC	AAA	1361
	Lys	Asn	Thr	Met	Glu	Pro	Leu	Asn	Ile	Pro	Asp	Val	Ser	Lys	Asp	Lys	
					410					415					420		
65	AGA	TTT	CCC	TGG	ACA	AAT	GAA	AAC	ATG	GGA	AAT	ATA	AAC	CAG	CAG	TGC	1409
	Arg	Phe	Pro	Trp	Thr	Asn	Glu	Asn	Met	Gly	Asn	Ile	Asn	Gln	Gln	Cys	
				425					430					435			
70	ATT	AGA	AGT	TTG	CTT	TGT	ACA	CCT	ATA	AAA	AAT	GGA	AAG	AAG	AAC	AAA	1457
	Ile	Arg	Ser	Leu	Leu	Cys	Thr	Pro	Ile	Lys	Asn	Gly	Lys	Lys	Asn	Lys	
				440				445					450				
75	GTG	ATA	GGG	GTT	TGC	CAA	CTT	GTT	AAT	AAG	ATG	GAG	GAA	ACC	ACT	GGC	1505
	Val	Ile	Gly	Val	Cys	Gln	Leu	Val	Asn	Lys	Met	Glu	Glu	Thr	Thr	Gly	
				455			460					465					

EP 0 652 960 B9 (W1B1)

	AAA	GTT	AAG	GCT	TTC	AAC	CGC	AAC	GAT	GAA	CAG	TTT	CTG	GAA	GCT	TTC	1553
	Lys	Val	Lys	Ala	Phe	Asn	Arg	Asn	Asp	Glu	Gln	Phe	Leu	Glu	Ala	Phe	
	470					475					480					485	
5	GTC	ATC	TTT	TGT	GGC	TTG	GGG	ATC	CAG	AAC	ACA	CAG	ATG	TAC	GAA	GCA	1601
	Val	Ile	Phe	Cys	Gly	Leu	Gly	Ile	Gln	Asn	Thr	Gln	Met	Tyr	Glu	Ala	
					490					495					500		
	GTG	GAG	AGA	GCC	ATG	GCC	AAG	CAA	ATG	GTC	ACG	TTA	GAG	GTT	CTG	TCT	1649
	Val	Glu	Arg	Ala	Met	Ala	Lys	Gln	Met	Val	Thr	Leu	Glu	Val	Leu	Ser	
10				505					510					515			
	TAT	CAT	GCT	TCA	GCT	GCA	GAG	GAA	GAA	ACC	AGA	GAG	CTG	CAG	TCC	TTA	1697
	Tyr	His	Ala	Ser	Ala	Ala	Glu	Glu	Glu	Thr	Arg	Glu	Leu	Gln	Ser	Leu	
				520				525					530				
	GCG	GCT	GCT	GTG	GTA	CCA	TCT	GCC	CAG	ACC	CTT	AAA	ATC	ACT	GAC	TTC	1745
	Ala	Ala	Ala	Val	Val	Pro	Ser	Ala	Gln	Thr	Leu	Lys	Ile	Thr	Asp	Phe	
15				535			540					545					
	AGC	TTC	AGC	GAC	TTT	GAG	CTG	TCT	GAC	CTG	GAA	ACA	GCA	CTG	TGC	ACA	1793
	Ser	Phe	Ser	Asp	Phe	Glu	Leu	Ser	Asp	Leu	Glu	Thr	Ala	Leu	Cys	Thr	
20				555							560					565	
	ATC	CGG	ATG	TTC	ACT	GAC	CTC	AAC	CTT	GTG	CAG	AAC	TTC	CAG	ATG	AAA	1841
	Ile	Arg	Met	Phe	Thr	Asp	Leu	Asn	Leu	Val	Gln	Asn	Phe	Gln	Met	Lys	
					570					575					580		
	CAT	GAG	GTC	CTT	TGC	AAG	TGG	ATT	TTA	AGT	GTG	AAG	AAG	AAC	TAT	CGG	1889
	His	Glu	Val	Leu	Cys	Lys	Trp	Ile	Leu	Ser	Val	Lys	Lys	Asn	Tyr	Arg	
25				585					590					595			
	AAG	AAC	GTC	GCC	TAT	CAT	AAT	TGG	AGA	CAT	GCC	TTT	AAT	ACA	GCT	CAG	1937
	Lys	Asn	Val	Ala	Tyr	His	Asn	Trp	Arg	His	Ala	Phe	Asn	Thr	Ala	Gln	
30				600				605					610				
	TGC	ATG	TTT	GCG	GCA	CTA	AAA	GCA	GGC	AAA	ATT	CAG	AAG	AGG	CTG	ACG	1985
	Cys	Met	Phe	Ala	Ala	Leu	Lys	Ala	Gly	Lys	Ile	Gln	Lys	Arg	Leu	Thr	
				615			620					625					
	GAC	CTG	GAG	ATA	CTT	GCA	CTG	CTG	ATT	GCT	GCC	TTA	AGC	CAT	GAT	CTG	2033
	Asp	Leu	Glu	Ile	Leu	Ala	Leu	Leu	Ile	Ala	Ala	Leu	Ser	His	Asp	Leu	
35						635					640					645	
	GAT	CAC	CGT	GGT	GTC	AAT	AAC	TCA	TAC	ATA	CAG	CGA	AGT	GAA	CAC	CCA	2081
	Asp	His	Arg	Gly	Val	Asn	Asn	Ser	Tyr	Ile	Gln	Arg	Ser	Glu	His	Pro	
					650					655					660		
	CTT	GCT	CAG	CTC	TAC	TGC	CAT	TCA	ATC	ATG	GAG	CAT	CAT	CAT	TTT	GAT	2129
	Leu	Ala	Gln	Leu	Tyr	Cys	His	Ser	Ile	Met	Glu	His	His	His	Phe	Asp	
					665					670					675		
	CAG	TGC	CTG	ATG	ATC	CTT	AAT	AGT	CCT	GGC	AAT	CAG	ATT	CTC	AGT	GGC	2177
	Gln	Cys	Leu	Met	Ile	Leu	Asn	Ser	Pro	Gly	Asn	Gln	Ile	Leu	Ser	Gly	
45				680				685						690			
	CTC	TCC	ATT	GAA	GAG	TAT	AAG	ACC	ACC	CTG	AAG	ATC	ATC	AAG	CAA	GCT	2225
	Leu	Ser	Ile	Glu	Glu	Tyr	Lys	Thr	Thr	Leu	Lys	Ile	Ile	Lys	Gln	Ala	
							700					705					
	ATT	TTA	GCC	ACA	GAC	CTA	GCA	CTG	TAC	ATA	AAG	AGA	CGA	GGA	GAA	TTT	2273
	Ile	Leu	Ala	Thr	Asp	Leu	Ala	Leu	Tyr	Ile	Lys	Arg	Arg	Gly	Glu	Phe	
50						715					720					725	
	TTT	GAA	CTT	ATA	ATG	AAA	AAT	CAA	TTC	AAT	TTG	GAA	GAT	CCT	CAT	CAA	2321
	Phe	Glu	Leu	Ile	Met	Lys	Asn	Gln	Phe	Asn	Leu	Glu	Asp	Pro	His	Gln	
55						730					735					740	

EP 0 652 960 B9 (W1B1)

	AAG GAG TTG TTT TTA GCG ATG CTG ATG ACA GCT TGT GAT CTT TCT GCA	2369
	Lys Glu Leu Phe Leu Ala Met Leu Met Thr Ala Cys Asp Leu Ser Ala	
	745 750 755	
5	ATT ACA AAA CCC TGG CCT ATT CAA CAA CGG ATA GCA GAA CTT GTT GCC	2417
	Ile Thr Lys Pro Trp Pro Ile Gln Gln Arg Ile Ala Glu Leu Val Ala	
	760 765 770	
	ACT GAA TTT TTT GAC CAA GGA GAT AGA GAG AGG AAA GAA CTC AAC ATA	2465
	Thr Glu Phe Phe Asp Gln Gly Asp Arg Glu Arg Lys Glu Leu Asn Ile	
10	775 780 785	
	GAG CCC GCT GAT CTA ATG AAC CGG GAG AAG AAA AAC AAA ATC CCA AGT	2513
	Glu Pro Ala Asp Leu Met Asn Arg Glu Lys Lys Asn Lys Ile Pro Ser	
	790 795 800 805	
15	ATG CAA GTT GGA TTC ATA GAT GCC ATC TGC TTG CAA CTG TAT GAG GCC	2561
	Met Gln Val Gly Phe Ile Asp Ala Ile Cys Leu Gln Leu Tyr Glu Ala	
	810 815 820	
	TTG ACC CAT GTG TCG GAG GAC TGT TTC CCT TTG CTG GAC GGC TGC AGA	2609
	Leu Thr His Val Ser Glu Asp Cys Phe Pro Leu Leu Asp Gly Cys Arg	
20	825 830 835	
	AAG AAC AGG CAG AAA TGG CAG GCT CTT GCA GAA CAG CAG GAG AAG ACA	2657
	Lys Asn Arg Gln Lys Trp Gln Ala Leu Ala Glu Gln Gln Glu Lys Thr	
	840 845 850	
25	CTG ATC AAT GGT GAA AGC AGC CAG ACC AAC CGA CAG CAA CGG AAT TCC	2705
	Leu Ile Asn Gly Glu Ser Ser Gln Thr Asn Arg Gln Gln Arg Asn Ser	
	855 860 865	
	GTT GCT GTC GGG ACA GTG TAGCCAGGTG TATCAGATGA GTGAGTGTGT	2753
	Val Ala Val Gly Thr Val	
30	870 875	
	GCTCAGCTCA GTCCTCTGCA ACACCATGAA GCTAGGCATT CCAGCTTAAT TCCTGCAGTT	2813
	GACTTTAAAA AACTGGCATA AAGCACTAGT CAGCATCTAG TTCTAGCTTG ACCAGTGAAG	2873
	AGTAGAACAC CACCACAGTC AGGGTGCAGA GCAGTTGGCA GTCTCCTTTC CAACCCAGAC	2933
35	TGGTGAATTT AAAGAAGAGC AGTCGTCGTT TATATCTCTG TCTTTTCCTA AGCGGGGTGT	2993
	GGAATCTCTA AGAGGAGAGA GAGATCTGGA CCACAGGTCC AATGCGCTCT GTCCTCTCAG	3053
	CTGCTTCCCC CACTGTGCTG TGACCTCTCA ATCTGAGAAA CGTGTAAGGA AGGTTTCAGC	3113
40	GAATTCCCTT TAAATGTGT CAGACAGTAG CTTCTGGGC CGGGTTGTTC CCGCAGCTCC	3173
	CCATCTGTTT GTTGTCTATC TTGGCTGAAA GAGGCTTTGC TGTACCTGCC ACACTCTCCT	3233
	GGATCCCTGT CCAGTAGCTG ATCAAAAAA AGGATGTGAA ATTCTCGTGT GACTTTTTAG	3293
45	AAAAGGAAAG TGACCCCGAG GATCGGTGTG GATTCAGTAG TTGTCCACAG ATGATCTGTT	3353
	TAGTTTCTAG AATTTTCCAA GATGATACAC TCCTCCCTAG TCTAGGGGTC AGACCCTGTA	3413
	TGGTGGCTGT GACCCTGAG GAACTTCTCT CTTTGCATGA CATTAGCCAT AGAACTGTTC	3473
50	TTGTCCAAAT ACACAGCTCA TATGCAGCTT GCAGGAAACA CTTTAAAAAC ACAACTATCA	3533
	CCTATGTTAT TCTGATTACA GAAGTTATCC CTACTIONTAAACATAAA CAAAGCCCCC	3593
	CAAACCTCAA ATAGTTGTGT GTGGTGAGAA ACTGCAAGTT TTCATCTCCA GAGATAGCTA	3653
55	TAGGTAATAA GTGGGATGTT TCTGAAACTT TAAAAATAA TCTTTTACAT ATATGTTAAC	3713

EP 0 652 960 B9 (W1B1)

TGTTTTCTTA TGAGCACTAT GGTTTGTTTT TTTTTTTTTT TGCTCTGCTT TGACTTGCCC 3773
 TTTTCACTCA ATTATCTTGG CAGTTTTTCT AAATGACTTG CACAGACTTC TCCTGTACTT 3833
 5 CATGGCTGTG CAGTGTTCCTA TGCTGTGAAG GCACCATCGT GTATTAAATC AGTTCCCTGG 3893
 TCACACATAG GTGAGCTGGT TGGAAATTTT TACCATTAAT AAACCACTTT CCCACATTGA 3953
 TGCTTTCTAA TCTGGCACAG GATGCTTCTT TTTTCCCCT TTTTCTCTGT TTAATTATTG 4013
 10 GAAATGGGAT CTGTGGGATC CTCGTCCCT GGCACCTAGC TGCTCTCAAC GTGGCCTGTG 4073
 GCCAGCAGCA TTGGCTAGAC CTGGGGGCTT GTTGGGAACG GAGACCCTCT GCCCTGCCCC 4133
 TGGCCTGCTG ACAAGGACCT GCATTTTGCT GAGCTCCCAG TGACCCTGGT GTTTAATTGT 4193
 15 TAACCATTGA AAAAAATCAA ACTATAGTTT ATTTACAATG TTGTGTTAAT TTCGGGTGTA 4253
 CAGCAAAGTG ACTCAGTGGT CAAGTACATT TAAAACACTG GGCATACTCT CTCCCTCTCC 4313
 TTGTGTACCT GGTGTTGATT TCCAGAAACC ATGCTCTTGT CTGTCCTGTA GTTTTGGAAG 4373
 20 CGCTTTCTCT TTGAAGACTG CCTTCTCTCC TGTGTCTGCC CTACATGGAC TAGTTCGTTT 4433
 ATTGTCCTAC ATGGCTTTGC TTCCATGTTT CTCTCAACTT T 4474

(2) INFORMATION FOR SEQ ID NO:10:

25

[0120]

(i) SEQUENCE CHARACTERISTICS:

30

- (A) LENGTH: 875 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

40

45

50

55

EP 0 652 960 B9 (W1B1)

Met Glu Arg Ala Gly Pro Gly Cys Arg Ala Ala Ala Thr Ala Met Gly
 1 5 10 15
 5 Pro Gly Leu Gly Arg Ser Val Ala Gly Arg Ser Leu Gly Leu Tyr Leu
 20 25 30
 Leu Tyr Phe Val Arg Lys Gly Thr Arg Glu Met Val Asn Ala Trp Phe
 35 40 45
 10 Ala Glu Arg Val His Thr Ile Pro Val Cys Lys Glu Gly Ile Lys Gly
 50 55 60
 His Thr Glu Ser Cys Ser Cys Pro Leu Gln Pro Ser Pro Arg Ala Glu
 65 70 75 80
 15 Ser Ser Val Pro Gly Thr Pro Thr Arg Lys Ile Ser Ala Ser Glu Phe
 85 90 95
 Asp Arg Pro Leu Arg Pro Ile Val Ile Lys Asp Ser Glu Gly Thr Val
 100 105 110
 20 Ser Phe Leu Ser Asp Ser Asp Lys Lys Glu Gln Met Pro Leu Thr Ser
 115 120 125
 Pro Arg Phe Asp Asn Asp Glu Gly Asp Gln Cys Ser Arg Leu Leu Glu
 130 135 140
 25 Leu Val Lys Asp Ile Ser Ser His Leu Asp Val Thr Ala Leu Cys His
 145 150 155 160

30

35

40

45

50

55

EP 0 652 960 B9 (W1B1)

Lys Ile Phe Leu His Ile His Gly Leu Ile Ser Ala Asp Arg Tyr Ser
 165 170 175
 5 Leu Phe Leu Val Cys Glu Asp Ser Ser Asn Asp Lys Phe Leu Ile Ser
 180 185 190
 Arg Leu Phe Asp Val Ala Glu Gly Ser Thr Leu Glu Glu Ala Ser Asn
 195 200 205
 10 Asn Cys Ile Arg Leu Glu Trp Asn Lys Gly Ile Val Gly His Val Ala
 210 215 220
 Ala Phe Gly Glu Pro Leu Asn Ile Lys Asp Ala Tyr Glu Asp Pro Arg
 225 230 235 240
 15 Phe Asn Ala Glu Val Asp Gln Ile Thr Gly Tyr Lys Thr Gln Ser Ile
 245 250 255
 Leu Cys Met Pro Ile Lys Asn His Arg Glu Glu Val Val Gly Val Ala
 260 265 270
 20 Gln Ala Ile Asn Lys Lys Ser Gly Asn Gly Gly Thr Phe Thr Glu Lys
 275 280 285
 Asp Glu Lys Asp Phe Ala Ala Tyr Leu Ala Phe Cys Gly Ile Val Leu
 290 295 300
 25 His Asn Ala Gln Leu Tyr Glu Thr Ser Leu Leu Glu Asn Lys Arg Asn
 305 310 315 320
 Gln Val Leu Leu Asp Leu Ala Ser Leu Ile Phe Glu Glu Gln Gln Ser
 325 330 335
 30 Leu Glu Val Ile Leu Arg Lys Ile Ala Ala Thr Ile Ile Ser Pro Met
 340 345 350
 Gln Val Gln Lys Cys Thr Ile Phe Ile Val Asp Glu Asp Cys Ser Asp
 355 360 365
 35 Ser Phe Ser Ser Val Phe His Met Glu Cys Glu Glu Leu Glu Lys Ser
 370 375 380
 Ser Asp Thr Leu Thr Arg Glu Arg Asp Ala Thr Arg Ile Asn Tyr Met
 385 390 395 400
 Tyr Ala Gln Tyr Val Lys Asn Thr Met Glu Pro Leu Asn Ile Pro Asp
 405 410 415
 40 Val Ser Lys Asp Lys Arg Phe Pro Trp Thr Asn Glu Asn Met Gly Asn
 420 425 430
 Ile Asn Gln Gln Cys Ile Arg Ser Leu Leu Cys Thr Pro Ile Lys Asn
 435 440 445
 45 Gly Lys Lys Asn Lys Val Ile Gly Val Cys Gln Leu Val Asn Lys Met
 450 455 460
 Glu Glu Thr Thr Gly Lys Val Lys Ala Phe Asn Arg Asn Asp Glu Gln
 465 470 475 480
 50 Phe Leu Glu Ala Phe Val Ile Phe Cys Gly Leu Gly Ile Gln Asn Thr
 485 490 495
 Gln Met Tyr Glu Ala Val Glu Arg Ala Met Ala Lys Gln Met Val Thr
 500 505 510

55

EP 0 652 960 B9 (W1B1)

Leu Glu Val Leu Ser Tyr His Ala Ser Ala Ala Glu Glu Glu Thr Arg
 515 520 525
 5
 Glu Leu Gln Ser Leu Ala Ala Ala Val Val Pro Ser Ala Gln Thr Leu
 530 535 540
 Lys Ile Thr Asp Phe Ser Phe Ser Asp Phe Glu Leu Ser Asp Leu Glu
 545 550 555 560
 10
 Thr Ala Leu Cys Thr Ile Arg Met Phe Thr Asp Leu Asn Leu Val Gln
 565 570 575
 Asn Phe Gln Met Lys His Glu Val Leu Cys Lys Trp Ile Leu Ser Val
 580 585 590
 15
 Lys Lys Asn Tyr Arg Lys Asn Val Ala Tyr His Asn Trp Arg His Ala
 595 600 605
 Phe Asn Thr Ala Gln Cys Met Phe Ala Ala Leu Lys Ala Gly Lys Ile
 610 615 620
 20
 Gln Lys Arg Leu Thr Asp Leu Glu Ile Leu Ala Leu Leu Ile Ala Ala
 625 630 635 640
 Leu Ser His Asp Leu Asp His Arg Gly Val Asn Asn Ser Tyr Ile Gln
 645 650 655
 25
 Arg Ser Glu His Pro Leu Ala Gln Leu Tyr Cys His Ser Ile Met Glu
 660 665 670
 His His His Phe Asp Gln Cys Leu Met Ile Leu Asn Ser Pro Gly Asn
 675 680 685
 30
 Gln Ile Leu Ser Gly Leu Ser Ile Glu Glu Tyr Lys Thr Thr Leu Lys
 690 695 700
 Ile Ile Lys Gln Ala Ile Leu Ala Thr Asp Leu Ala Leu Tyr Ile Lys
 705 710 715 720
 35
 Arg Arg Gly Glu Phe Phe Glu Leu Ile Met Lys Asn Gln Phe Asn Leu
 725 730 735
 Glu Asp Pro His Gln Lys Glu Leu Phe Leu Ala Met Leu Met Thr Ala
 740 745 750
 40
 Cys Asp Leu Ser Ala Ile Thr Lys Pro Trp Pro Ile Gln Gln Arg Ile
 755 760 765
 Ala Glu Leu Val Ala Thr Glu Phe Phe Asp Gln Gly Asp Arg Glu Arg
 770 775 780
 45
 Lys Glu Leu Asn Ile Glu Pro Ala Asp Leu Met Asn Arg Glu Lys Lys
 785 790 795 800
 Asn Lys Ile Pro Ser Met Gln Val Gly Phe Ile Asp Ala Ile Cys Leu
 805 810 815
 50
 Gln Leu Tyr Glu Ala Leu Thr His Val Ser Glu Asp Cys Phe Pro Leu
 820 825 830
 Leu Asp Gly Cys Arg Lys Asn Arg Gln Lys Trp Gln Ala Leu Ala Glu
 835 840 845
 55
 Gln Gln Glu Lys Thr Leu Ile Asn Gly Glu Ser Ser Gln Thr Asn Arg
 850 855 860

Gln Gln Arg Asn Ser Val Ala Val Gly Thr Val
865 870 875

5 (2) INFORMATION FOR SEQ ID NO:11:

[0121]

(i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 2060 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

20

25

30

35

40

45

50

55

EP 0 652 960 B9 (W1B1)

GCGGCCGCGC TCCGGCCGCT TTGTGAAAG CCGGCCCGAC TGGAGCAGGA CGAAGGGGGA 60
 GGGTCTCGAG GCCGAGTCCT GTTCTTCTGA GGGACGGACC CCAGCTGGGG TGGAAAAGCA 120
 5 GTACCAGAGA GCCTCCGAGG CGCGCGGTGC CAACCATGGA GCGGGCCGGC CCCAGCTTCG 180
 GGCAGCAGCG ACAGCAGCAG CAGCCCCAGC AGCAGAAGCA GCAGCAGAGG GATCAGGACT 240
 CGGTCGAAGC ATGGCTGGAC GATCACTGGG ACTTTACCTT CTCATACTTT GTTAGAAAAG 300
 10 CCACCAGAGA AATGGTCAAT GCATGGTTTG CTGAGAGAGT TCACACCATC CCTGTGTGCA 360
 AGGAAGGTAT CAGAGGCCAC ACCGAATCTT GCTCTTGTCC CTTGCAGCAG AGTCCTCGTG 420
 CAGATAACAG TGTCCCTGGA ACACCAACCA GAAAATCTC TGCCTCTGAA TTTGACCGGC 480
 15 CTCTTAGACC CATTGTTGTC AAGGATTCTG AGGGAAGTGT GAGCTTCCTC TCTGACTCAG 540
 AAAAGAAGGA ACAGATGCCT CTAACCCCTC CAAGTTTGA TCATGATGAA GGGGACCAGT 600
 GCTCAAGACT CTTGGAATTA GTGAAGGATA TTTCTAGTCA TTTGGATGTC ACAGCCTTAT 660
 20 GTCACAAAAT TTTCTTGCAT ATCCATGGAC TGATATCTGC TGACCGCTAT TCCCTGTTCC 720
 TTGTCTGTGA AGACAGCTCC AATGACAAGT TTCTTATCAG CCGCCTCTTT GATGTTGCTG 780
 AAGGTTCAAC ACTGGAAGAA GTTTCAAATA ACTGTATCCG CTTAGAATGG AACAAAGGCA 840
 25 TTGTGGGACA TGTGGCAGCG CTTGGTGAGC CCTTGAACAT CAAAGATGCA TATGAGGATC 900
 CTCGGTTCAA TGCAGAAGTT GACCAAATTA CAGGCTACAA GACACAAAGC ATTCTTTGTA 960
 TGCCAATTAA GAATCATAGG GAAGAGGTTG TTGGTGTAGC CCAGGCCATC AACAAAGAAAT 1020
 30 CAGGAAACGG TGGGACATTT ACTGAAAAAG ATGAAAAGGA CTTTGCTGCT TATTTGGCAT 1080
 TTTGTGGTAT TGTTCTTCAT AATGCTCAGC TCTATGAGAC TTCACTGCTG GAGAACAAGA 1140
 GAAATCAGGT GCTGCTTGAC CTTGCTAGTT TAATTTTGA AGAACAAACA TCATTAGAAG 1200
 35 TAATTTTGAA GAAAATAGCT GCCACTATTA TCTCTTTCAT GCAAGTGCAG AAATGCACCA 1260
 TTTTCATAGT GGATGAAGAT TGCTCCGATT CTTTTCTAG TGTGTTTAC ATGGAGTGTG 1320
 AGGAATTAGA AAAATCATCT GATACATTA CAAGGGAACA TGATGCAAAC AAAATCAATT 1380
 40 ACATGTATGC TCAGTATGTC AAAAATACTA TGAACCACT TTATATCCCA GATGTCAGTA 1440
 AGGATAAAAG ATTTCCCTGG ACAACTGAAA ATACAGGAAA TGTAAACCAG CAGTGCATTA 1500

45

50

55

EP 0 652 960 B9 (W1B1)

GAAGTTTGCT TTGTACACCT ATAAAAAATG GAAAGAAGAA TAAAGTTATA GGGGTTTGCC 1560
AACTTGTTAA TAAGATGGAG GAGAATACTG GCAAGGTAA GCCTTTCAAC CGAAATGACG 1620
5 AACAGTTTCT GGAAGCTTTT GTCATCTTTT GTGGCTTGGG GATCCAGAAC ACGCAGATGT 1680
ATGAAGCAGT GGAGAGAGCC ATGGCCAAGC AAATGGTCAC ATTGGAGGTT CTGTCGTATC 1740
ATGCTTCAGC AGCAGAGGAA GAAACAAGAG AGCTACAGTC GTTAGCGGCT GCTGTGGTGC 1800
10 CATCTGCCCA GACCCTTAAA ATTACTGACT TTAGCTTCAG TGACTTTGAG CTGTCTGATC 1860
TGGAAACAGC ACTGTGTACA ATTCCGATGT TTACTIONACT CAACCTTGTG CAGAACTTCC 1920
AGATGAAACA TGAGGTTCTT TGCAGATGGA TTTTAAGTGT TAAGAAGAAT TATCGGAAGA 1980
15 ATGTTGCCTA TCATAATTGG AGACATGCCT TTAATACAGC TCAGTGCATG TTTGCTGCTC 2040
TAAAAGCAGG CAAAATTCAG 2060

(2) INFORMATION FOR SEQ ID NO:12:

20

[0122]

(i) SEQUENCE CHARACTERISTICS:

25

- (A) LENGTH: 1982 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

35

40

45

50

55

EP 0 652 960 B9 (W1B1)

AAAAAATTTT CTTGCATATC CATGGACTGA TATCTGCTGA CCGCTATTCC CTGTTCCCTTG 60
 TCTGTGAAGA CAGCTCCAAT GACAAGTTTC TTATCAGCCG CCTCTTTGAT GTTGCTGAAG 120
 5 GTTCAACACT GGAAGAAGTT TCAAATAACT GTATCCGCTT AGAATGGAAC AAAGGCATTG 180
 TGGGACATGT GGCAGCGCTT GGTGAGCCCT TGAACATCAA AGATGCATAT GAGGATCCTC 240
 GGTTCAATGC AGAAGTTGAC CAAATTACAG GCTACAAGAC ACAAAGCATT CTTTGTATGC 300
 10 CAATTAAGAA TCATAGGGAA GAGGTTGTTG GTGTAGCCCA GGCCATCAAC AAGAAATCAG 360
 GAAACGGTGG GACATTTACT GAAAAAGATG AAAAGGACTT TGCTGCTTAT TTGGCATTTT 420
 GTGGTATTGT TCTTCATAAT GCTCAGCTCT ATGAGACTTC ACTGCTGGAG AACAGAGAA 480
 15 ATCAGGTGCT GCTTGACCTT GCTAGTTTAA TTTTGAAGA ACAACAATCA TTAGAAGTAA 540
 TTTTGAAGAA AATAGCTGCC ACTATTATCT CTTTCATGCA AGTCAGAAA TGCACCATTT 600
 TCATAGTGGG TGAAGATTGC TCCGATTCTT TTTCTAGTGT GTTTCACATG GAGTGTGAGG 660
 20 AATTAGAAAA ATCATCTGAT ACATTAACAA GGGAACATGA TGCAAAACAAA ATCAATTACA 720
 TGTATGCTCA GTATGTCAA AATACTATGG AACCACTTAA TATCCCAGAT GTCAGTAAGG 780
 ATAAAAGATT TCCCTGGACA ACTGAAAATA CAGGAAATGT AAACCAGCAG TGCATTAGAA 840
 25 GTTTGCTTTG TACACCTATA AAAAATGGAA AGAAGAATAA AGTTATAGGG GTTTGCCAAC 900
 TTGTTAATAA GATGGAGGAG AATACTGGCA AGGTTAAGCC TTTCAACCGA AATGACGAAC 960
 AGTTTCTGGA AGCTTTTGTG ATCTTTTGTG GCTTGGGGAT CCAGAACACG CAGATGTATG 1020

30

35

40

45

50

55

EP 0 652 960 B9 (W1B1)

	AAGCAGTGGA GAGAGCCATG GCCAAGCAAA TGGTCACATT GGAGGTTCTG TCGTATCATG	1080
	CTTCAGCAGC AGAGGAAGAA ACAAGAGAGC TACAGTCGTT AGCGGCTGCT GTGGTGCCAT	1140
5	CTGCCAGAC CCTTAAAATT ACTGACTTTA GCTTCAGTGA CTTTGAGCTG TCTGATCTGG	1200
	AAACAGCACT GTGTACAATT CGGATGTTTA CTGACCTCAA CCTTGTGCAG AACTTCCAGA	1260
	TGAAACATGA GGTTCCTTTC AGATGGATTT TAAGTGTTAA GAAGAATTAT CGGAAGAATG	1320
10	TTGCCTATCA TAATTGGAGA CATGCCTTTA ATACAGCTCA GTGCATGTTT GCTGCTCTAA	1380
	AAGCAGGCAA AATTCAGAAC AAGCTGACTG ACCTGGAGAT ACTTGCATTG CTGATTGCTG	1440
	CACTAAGCCA CGATTTGGAT CACCGTGGTG TGAATAACTC TTACATACAG CGAAGTGAAC	1500
15	ATCCACTTGC CCAGCTTTAC TGCCATTCAA TCATGGAACA CCATCATTTT GACCAGTGCC	1560
	TGATGATTCT TAATAGTCCA GGCAATCAGA TTCTCAGTGG CCTCTCCATT GAAGAATATA	1620
	AGACCACGTT GAAAATAATC AAGCAAGCTA TTTTAGCTAC AGACCTAGCA CTGTACATTA	1680
20	AGAGGCGAGG AGAATTTTTT GAACTTATAA GAAAAAATCA ATTCAATTTG GAAGATCCTC	1740
	ATCAAAAGGA GTTGTTTTTG GCAATGCTGA TGACAGCTTG TGATCTTTCT GCAATTACAA	1800
	AACCCTGGCC TATTCAACAA CGGATAGCAG AACTTGTAGC AACTGAATTT TTTGATCAAG	1860
25	GAGACAGAGA GAGAAAAGAA CTCAACATAG AACCCACTGA TCTAATGAAC AGGGAGAAGA	1920
	AAAACAAAAT CCCAAGTATG CAAGTTGGGT TCATAGATGC CATCTGCTTG CAACTGTATG	1980
30	AG	1982

(2) INFORMATION FOR SEQ ID NO:13:

[0123]

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GCCACCAGAG AAATGGTC

18

(2) INFORMATION FOR SEQ ID NO:14:

[0124]

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

ACAATGGGTC TAAGAGGC

18

10 (2) INFORMATION FOR SEQ ID NO:15:

[0125]

(i) SEQUENCE CHARACTERISTICS:

15

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

25

TCAGTGCATG TTTGCTGC

18

(2) INFORMATION FOR SEQ ID NO:16:

30 **[0126]**

(i) SEQUENCE CHARACTERISTICS:

35

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

45

TACAAACATG TTCATCAG

18

(2) INFORMATION FOR SEQ ID NO:17:

[0127]

50 (i) SEQUENCE CHARACTERISTICS:

55

- (A) LENGTH: 1107 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

EP 0 652 960 B9 (W1B1)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

	GAGACATGCC TTTAATACAG CTCAGTGCAT GTTTGCTGCT CTAAAAGCAG GCAAAATTCA	60
5	GAACAAGCTG ACTGACCTGG AGATACTTGC ATTGCTGATT GCTGCACTAA GCCACGATTT	120
	GGATCACCGT GGTGTGAATA ACTCTTACAT ACAGCGAAGT GAACATCCAC TTGCCCAGCT	180
	TTACTGCCAT TCAATCATGG AACACCATCA TTTTGACCAG TGCCTGATGA TTCTTAATAG	240
10	TCCAGGCAAT CAGATTCTCA GTGGCCTCTC CATTGAAGAA TATAAGACCA CGTTGAAAAT	300
	AATCAAGCAA GCTATTTTGTAG CTACAGACCT AGCACTGTAC ATTAAGAGGC GAGGAGAATT	360
	TTTTGAACTT ATAAGAAAAA ATCAATTCAA TTTGGAAGAT CCTCATCAA AGGAGTTGTT	420
15	TTTGGAATG CTGATGACAG CTTGTGATCT TTCTGCAATT ACAAACCCT GGCCTATTCA	480
	ACAACGGATA GCAGAACTTG TAGCAACTGA ATTTTTTGAT CAAGGAGACA GAGAGAGAAA	540
	AGAACTCAAC ATAGAACCCA CTGATCTAAT GAACAGGGAG AAGAAAAACA AAATCCCAAG	600
20	TATGCAAGTT GGGTTCATAG ATGCCATCTG CTTGCAACTG TATGAGGCC TGACCCACGT	660
25	GTCAGAGGAC TGTTTCCCTT TGCTAGATGG CTGCAGAAAG AACAGGCAGA AATGGCAGGC	720
	CCTTGCAGAA CAGCAGGAGA AGATGCTGAT TAATGGGGAA AGCGGCCAGG CCAAGCGGAA	780
	CTGAGTGGCC TATTTTCATGC AGAGTTGAAG TTTACAGAGA TGGTGTGTTT TGCAATATGC	840
30	CTAGTTTCTT ACACACTGTC TGTATAGTGT CTGTATTTGG TATATACTTT GCCACTGCTG	900
	TATTTTTTATT TTTGCACAAC TTTTGAGAGT ATAGCATGAA TGTTTTTAGA GGA CTATTAC	960
	ATATTTTTTG TATATTTGTT TTATGCTACT GAACTGAAAG GATCAACAAC ATCCACTGTT	1020
35	AGCACATTGA TAAAAGCATT GTTTGTGATA TTTCTGTGAC TGCAAAGTGT ATGCAGTATT	1080
	CTTGCACTGA GGTTTTTTTG CTTGGGG	1107

(2) INFORMATION FOR SEQ ID NO:18:

40
[0128]

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

50 (ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

55 **TTTGGAAGAT CCTCATCA** **18**

(2) INFORMATION FOR SEQ ID NO:19:

[0129]

5 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

15

ATGTCTCGAG TCAGTTCCGC TTGGCCTG

28

(2) INFORMATION FOR SEQ ID NO:20:

20

[0130]

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

30

TACAGAATTC TGACCATGGA GCGGGCCGGC

30

(2) INFORMATION FOR SEQ ID NO:21:

[0131]

40

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

50

CATTCTAAGC GCATACAG

18

55

EP 0 652 960 B9 (W1B1)

(2) INFORMATION FOR SEQ ID NO:22:

[0132]

5 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2645 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 12..2636

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

20

25	GAATTCTGAC C ATG GAG CGG GCC GGC CCC AGC TTC GGG CAG CAG CGA CAG Met Glu Arg Ala Gly Pro Ser Phe Gly Gln Gln Arg Gln 1 5 10	50
30	CAG CAG CAG CCC CAG CAG CAG AAG CAG CAG CAG AGG GAT CAG GAC TCG Gln Gln Gln Pro Gln Gln Gln Lys Gln Gln Gln Arg Asp Gln Asp Ser 15 20 25	98
35	GTC GAA GCA TGG CTG GAC GAT CAC TGG GAC TTT ACC TTC TCA TAC TTT Val Glu Ala Trp Leu Asp Asp His Trp Asp Phe Thr Phe Ser Tyr Phe 30 35 40 45	146
40	GTT AGA AAA GCC ACC AGA GAA ATG GTC AAT GCA TGG TTT GCT GAG AGA Val Arg Lys Ala Thr Arg Glu Met Val Asn Ala Trp Phe Ala Glu Arg 50 55 60	194
45	GTT CAC ACC ATC CCT GTG TGC AAG GAA GGT ATC AGA GGC CAC ACC GAA Val His Thr Ile Pro Val Cys Lys Glu Gly Ile Arg Gly His Thr Glu 65 70 75	242
50	TCT TGC TCT TGT CCC TTG CAG CAG AGT CCT CGT GCA GAT AAC AGT GTC Ser Cys Ser Cys Pro Leu Gln Gln Ser Pro Arg Ala Asp Asn Ser Val 80 85 90	290
55	CCT GGA ACA CCA ACC AGG AAA ATC TCT GCC TCT GAA TTT GAC CGG CCT Pro Gly Thr Pro Thr Arg Lys Ile Ser Ala Ser Glu Phe Asp Arg Pro 95 100 105	338
60	CTT AGA CCC ATT GTT GTC AAG GAT TCT GAG GGA ACT GTG AGC TTC CTC Leu Arg Pro Ile Val Val Lys Asp Ser Glu Gly Thr Val Ser Phe Leu 110 115 120 125	386

EP 0 652 960 B9 (W1B1)

	TCT	GAC	TCA	GAA	AAG	AAG	GAA	CAG	ATG	CCT	CTA	ACC	CCT	CCA	AGG	TTT	434
	Ser	Asp	Ser	Glu	Lys	Lys	Glu	Gln	Met	Pro	Leu	Thr	Pro	Pro	Arg	Phe	
					130					135					140		
5	GAT	CAT	GAT	GAA	GGG	GAC	CAG	TGC	TCA	AGA	CTC	TTG	GAA	TTA	GTG	AAG	482
	Asp	His	Asp	Glu	Gly	Asp	Gln	Cys	Ser	Arg	Leu	Leu	Glu	Leu	Val	Lys	
				145					150					155			
	GAT	ATT	TCT	AGT	CAT	TTG	GAT	GTC	ACA	GCC	TTA	TGT	CAC	AAA	ATT	TTC	530
	Asp	Ile	Ser	Ser	His	Leu	Asp	Val	Thr	Ala	Leu	Cys	His	Lys	Ile	Phe	
10				160				165					170				
	TTG	CAT	ATC	CAT	GGA	CTG	ATA	TCT	GCT	GAC	CGC	TAT	TCC	CTG	TTC	CTT	578
	Leu	His	Ile	His	Gly	Leu	Ile	Ser	Ala	Asp	Arg	Tyr	Ser	Leu	Phe	Leu	
		175					180					185					
15	GTC	TGT	GAA	GAC	AGC	TCC	AAT	GAC	AAG	TTT	CTT	ATC	AGC	CGC	CTC	TTT	626
	Val	Cys	Glu	Asp	Ser	Ser	Asn	Asp	Lys	Phe	Leu	Ile	Ser	Arg	Leu	Phe	
		190				195					200					205	
	GAT	GTT	GCT	GAA	GGT	TCA	ACA	CTG	GAA	GAA	GTT	TCA	AAT	AAC	TGT	ATC	674
	Asp	Val	Ala	Glu	Gly	Ser	Thr	Leu	Glu	Glu	Val	Ser	Asn	Asn	Cys	Ile	
20					210				215						220		
	CGC	TTA	GAA	TGG	AAC	AAA	GGC	ATT	GTG	GGA	CAT	GTG	GCA	GCG	CTT	GGT	722
	Arg	Leu	Glu	Trp	Asn	Lys	Gly	Ile	Val	Gly	His	Val	Ala	Ala	Leu	Gly	
				225					230					235			
25	GAG	CCC	TTG	AAC	ATC	AAA	GAT	GCA	TAT	GAG	GAT	CCT	CGG	TTC	AAT	GCA	770
	Glu	Pro	Leu	Asn	Ile	Lys	Asp	Ala	Tyr	Glu	Asp	Pro	Arg	Phe	Asn	Ala	
			240					245					250				
	GAA	GTT	GAC	CAA	ATT	ACA	GGC	TAC	AAG	ACA	CAA	AGC	ATT	CTT	TGT	ATG	818
	Glu	Val	Asp	Gln	Ile	Thr	Gly	Tyr	Lys	Thr	Gln	Ser	Ile	Leu	Cys	Met	
30			255				260					265					
	CCA	ATT	AAG	AAT	CAT	AGG	GAA	GAG	GTT	GTT	GGT	GTA	GCC	CAG	GCC	ATC	866
	Pro	Ile	Lys	Asn	His	Arg	Glu	Glu	Val	Val	Gly	Val	Ala	Gln	Ala	Ile	
						275					280					285	
35	AAC	AAG	AAA	TCA	GGA	AAC	GGT	GGG	ACA	TTT	ACT	GAA	AAA	GAT	GAA	AAG	914
	Asn	Lys	Lys	Ser	Gly	Asn	Gly	Gly	Thr	Phe	Thr	Glu	Lys	Asp	Glu	Lys	
					290					295					300		
	GAC	TTT	GCT	GCT	TAT	TTG	GCA	TTT	TGT	GGT	ATT	GTT	CTT	CAT	AAT	GCT	962
	Asp	Phe	Ala	Ala	Tyr	Leu	Ala	Phe	Cys	Gly	Ile	Val	Leu	His	Asn	Ala	
				305					310					315			
40	CAG	CTC	TAT	GAG	ACT	TCA	CTG	CTG	GAG	AAC	AAG	AGA	AAT	CAG	GTG	CTG	1010
	Gln	Leu	Tyr	Glu	Thr	Ser	Leu	Leu	Glu	Asn	Lys	Arg	Asn	Gln	Val	Leu	
				320				325					330				
	CTT	GAC	CTT	GCT	AGT	TTA	ATT	TTT	GAA	GAA	CAA	CAA	TCA	TTA	GAA	GTA	1058
	Leu	Asp	Leu	Ala	Ser	Leu	Ile	Phe	Glu	Glu	Gln	Gln	Ser	Leu	Glu	Val	
45			335				340					345					
	ATT	TTG	AAG	AAA	ATA	GCT	GCC	ACT	ATT	ATC	TCT	TTC	ATG	CAA	GTG	CAG	1106
	Ile	Leu	Lys	Lys	Ile	Ala	Ala	Thr	Ile	Ile	Ser	Phe	Met	Gln	Val	Gln	
						355					360					365	
50	AAA	TGC	ACC	ATT	TTC	ATA	GTG	GAT	GAA	GAT	TGC	TCC	GAT	TCT	TTT	TCT	1154
	Lys	Cys	Thr	Ile	Phe	Ile	Val	Asp	Glu	Asp	Cys	Ser	Asp	Ser	Phe	Ser	
					370					375					380		
	AGT	GTG	TTT	CAC	ATG	GAG	TGT	GAG	GAA	TTA	GAA	AAA	TCA	TCT	GAT	ACA	1202
	Ser	Val	Phe	His	Met	Glu	Cys	Glu	Glu	Leu	Glu	Lys	Ser	Ser	Asp	Thr	
55					385				390						395		

EP 0 652 960 B9 (W1B1)

	TTA	ACA	AGG	GAA	CAT	GAT	GCA	AAC	AAA	ATC	AAT	TAC	ATG	TAT	GCT	CAG	1250
	Leu	Thr	Arg	Glu	His	Asp	Ala	Asn	Lys	Ile	Asn	Tyr	Met	Tyr	Ala	Gln	
			400					405					410				
5	TAT	GTC	AAA	AAT	ACT	ATG	GAA	CCA	CTT	AAT	ATC	CCA	GAT	GTC	AGT	AAG	1298
	Tyr	Val	Lys	Asn	Thr	Met	Glu	Pro	Leu	Asn	Ile	Pro	Asp	Val	Ser	Lys	
			415				420					425					
	GAT	AAA	AGA	TTT	CCC	TGG	ACA	ACT	GAA	AAT	ACA	GGA	AAT	GTA	AAC	CAG	1346
	Asp	Lys	Arg	Phe	Pro	Trp	Thr	Thr	Glu	Asn	Thr	Gly	Asn	Val	Asn	Gln	
10						435					440					445	
	CAG	TGC	ATT	AGA	AGT	TTG	CTT	TGT	ACA	CCT	ATA	AAA	AAT	GGA	AAG	AAG	1394
	Gln	Cys	Ile	Arg	Ser	Leu	Leu	Cys	Thr	Pro	Ile	Lys	Asn	Gly	Lys	Lys	
					450					455					460		
	AAT	AAA	GTT	ATA	GGG	GTT	TGC	CAA	CTT	GTT	AAT	AAG	ATG	GAG	GAG	AAT	1442
	Asn	Lys	Val	Ile	Gly	Val	Cys	Gln	Leu	Val	Asn	Lys	Met	Glu	Glu	Asn	
				465					470					475			
	ACT	GGC	AAG	GTT	AAG	CCT	TTC	AAC	CGA	AAT	GAC	GAA	CAG	TTT	CTG	GAA	1490
	Thr	Gly	Lys	Val	Lys	Pro	Phe	Asn	Arg	Asn	Asp	Glu	Gln	Phe	Leu	Glu	
20			480					485					490				
	GCT	TTT	GTC	ATC	TTT	TGT	GGC	TTG	GGG	ATC	CAG	AAC	ACG	CAG	ATG	TAT	1538
	Ala	Phe	Val	Ile	Phe	Cys	Gly	Leu	Gly	Ile	Gln	Asn	Thr	Gln	Met	Tyr	
			495				500					505					
	GAA	GCA	GTG	GAG	AGA	GCC	ATG	GCC	AAG	CAA	ATG	GTC	ACA	TTG	GAG	GTT	1586
	Glu	Ala	Val	Glu	Arg	Ala	Met	Ala	Lys	Gln	Met	Val	Thr	Leu	Glu	Val	
25						515					520					525	
	CTG	TCG	TAT	CAT	GCT	TCA	GCA	GCA	GAG	GAA	GAA	ACA	AGA	GAG	CTA	CAG	1634
	Leu	Ser	Tyr	His	Ala	Ser	Ala	Ala	Glu	Glu	Glu	Thr	Arg	Glu	Leu	Gln	
					530					535					540		
	TCG	TTA	GCG	GCT	GCT	GTG	GTG	CCA	TCT	GCC	CAG	ACC	CTT	AAA	ATT	ACT	1682
	Ser	Leu	Ala	Ala	Ala	Val	Val	Pro	Ser	Ala	Gln	Thr	Leu	Lys	Ile	Thr	
				545				550						555			
	GAC	TTT	AGC	TTC	AGT	GAC	TTT	GAG	CTG	TCT	GAT	CTG	GAA	ACA	GCA	CTG	1730
	Asp	Phe	Ser	Phe	Ser	Asp	Phe	Glu	Leu	Ser	Asp	Leu	Glu	Thr	Ala	Leu	
35			560					565					570				
	TGT	ACA	ATT	CGG	ATG	TTT	ACT	GAC	CTC	AAC	CTT	GTG	CAG	AAC	TTC	CAG	1778
	Cys	Thr	Ile	Arg	Met	Phe	Thr	Asp	Leu	Asn	Leu	Val	Gln	Asn	Phe	Gln	
			575				580					585					
	ATG	AAA	CAT	GAG	GTT	CTT	TGC	AGA	TGG	ATT	TTA	AGT	GTT	AAG	AAG	AAT	1826
	Met	Lys	His	Glu	Val	Leu	Cys	Arg	Trp	Ile	Leu	Ser	Val	Lys	Lys	Asn	
						595					600					605	
	TAT	CGG	AAG	AAT	GTT	GCC	TAT	CAT	AAT	TGG	AGA	CAT	GCC	TTT	AAT	ACA	1874
	Tyr	Arg	Lys	Asn	Val	Ala	Tyr	His	Asn	Trp	Arg	His	Ala	Phe	Asn	Thr	
45					610					615					620		
	GCT	CAG	TGC	ATG	TTT	GCT	GCT	CTA	AAA	GCA	GGC	AAA	ATT	CAG	AAC	AAG	1922
	Ala	Gln	Cys	Met	Phe	Ala	Ala	Leu	Lys	Ala	Gly	Lys	Ile	Gln	Asn	Lys	
				625					630					635			
	CTG	ACT	GAC	CTG	GAG	ATA	CTT	GCA	TTG	CTG	ATT	GCT	GCA	CTA	AGC	CAC	1970
	Leu	Thr	Asp	Leu	Glu	Ile	Leu	Ala	Leu	Leu	Ile	Ala	Ala	Leu	Ser	His	
50			640					645					650				
	GAT	TTG	GAT	CAC	CGT	GGT	GTG	AAT	AAC	TCT	TAC	ATA	CAG	CGA	AGT	GAA	2018
	Asp	Leu	Asp	His	Arg	Gly	Val	Asn	Asn	Ser	Tyr	Ile	Gln	Arg	Ser	Glu	
55			655				660					665					

EP 0 652 960 B9 (W1B1)

	CAT CCA CTT GCC CAG CTT TAC TGC CAT TCA ATC ATG GAA CAC CAT CAT	2066
	His Pro Leu Ala Gln Leu Tyr Cys His Ser Ile Met Glu His His His	
	670 675 680 685	
5	TTT GAC CAG TGC CTG ATG ATT CTT AAT AGT CCA GGC AAT CAG ATT CTC	2114
	Phe Asp Gln Cys Leu Met Ile Leu Asn Ser Pro Gly Asn Gln Ile Leu	
	690 695 700	
	AGT GGC CTC TCC ATT GAA GAA TAT AAG ACC ACG TTG AAA ATA ATC AAG	2162
10	Ser Gly Leu Ser Ile Glu Glu Tyr Lys Thr Thr Leu Lys Ile Ile Lys	
	705 710 715	
	CAA GCT ATT TTA GCT ACA GAC CTA GCA CTG TAC ATT AAG AGG CGA GGA	2210
	Gln Ala Ile Leu Ala Thr Asp Leu Ala Leu Tyr Ile Lys Arg Arg Gly	
	720 725 730	
15	GAA TTT TTT GAA CTT ATA AGA AAA AAT CAA TTC AAT TTG GAA GAT CCT	2258
	Glu Phe Phe Glu Leu Ile Arg Lys Asn Gln Phe Asn Leu Glu Asp Pro	
	735 740 745	
	CAT CAA AAG GAG TTG TTT TTG GCA ATG CTG ATG ACA GCT TGT GAT CTT	2306
20	His Gln Lys Glu Leu Phe Leu Ala Met Leu Met Thr Ala Cys Asp Leu	
	750 755 760 765	
	TCT GCA ATT ACA AAA CCC TGG CCT ATT CAA CAA CGG ATA GCA GAA CTT	2354
	Ser Ala Ile Thr Lys Pro Trp Pro Ile Gln Gln Arg Ile Ala Glu Leu	
	770 775 780	
25	GTA GCA ACT GAA TTT TTT GAT CAA GGA GAC AGA GAG AGA AAA GAA CTC	2402
	Val Ala Thr Glu Phe Phe Asp Gln Gly Asp Arg Glu Arg Lys Glu Leu	
	785 790 795	
	AAC ATA GAA CCC ACT GAT CTA ATG AAC AGG GAG AAG AAA AAC AAA ATC	2450
30	Asn Ile Glu Pro Thr Asp Leu Met Asn Arg Glu Lys Lys Asn Lys Ile	
	800 805 810	
	CCA AGT ATG CAA GTT GGG TTC ATA GAT GCC ATC TGC TTG CAA CTG TAT	2498
	Pro Ser Met Gln Val Gly Phe Ile Asp Ala Ile Cys Leu Gln Leu Tyr	
	815 820 825	
35	GAG GCC CTG ACC CAC GTG TCA GAG GAC TGT TTC CCT TTG CTA GAT GGC	2546
	Glu Ala Leu Thr His Val Ser Glu Asp Cys Phe Pro Leu Leu Asp Gly	
	830 835 840 845	
	TGC AGA AAG AAC AGG CAG AAA TGG CAG GCC CTT GCA GAA CAG CAG GAG	2594
	Cys Arg Lys Asn Arg Gln Lys Trp Gln Ala Leu Ala Glu Gln Gln Glu	
	850 855 860	
40	AAG ATG CTG ATT AAT GGG GAA AGC GGC CAG GCC AAG CGG AAC	2636
	Lys Met Leu Ile Asn Gly Glu Ser Gly Gln Ala Lys Arg Asn	
	865 870 875	
45	TGACTCGAG	2645

(2) INFORMATION FOR SEQ ID NO:23:

[0133]

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 875 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

EP 0 652 960 B9 (W1B1)

Met Glu Arg Ala Gly Pro Ser Phe Gly Gln Gln Arg Gln Gln Gln Gln
1 5 10 15

5

10

15

20

25

30

35

40

45

50

55

EP 0 652 960 B9 (W1B1)

Pro Gln Gln Gln Lys Gln Gln Gln Arg Asp Gln Asp Ser Val Glu Ala
 20 25 30
 Trp Leu Asp Asp His Trp Asp Phe Thr Phe Ser Tyr Phe Val Arg Lys
 35 40 45
 Ala Thr Arg Glu Met Val Asn Ala Trp Phe Ala Glu Arg Val His Thr
 50 55 60
 Ile Pro Val Cys Lys Glu Gly Ile Arg Gly His Thr Glu Ser Cys Ser
 65 70 75 80
 Cys Pro Leu Gln Gln Ser Pro Arg Ala Asp Asn Ser Val Pro Gly Thr
 85 90 95
 Pro Thr Arg Lys Ile Ser Ala Ser Glu Phe Asp Arg Pro Leu Arg Pro
 100 105 110
 Ile Val Val Lys Asp Ser Glu Gly Thr Val Ser Phe Leu Ser Asp Ser
 115 120 125
 Glu Lys Lys Glu Gln Met Pro Leu Thr Pro Pro Arg Phe Asp His Asp
 130 135 140
 Glu Gly Asp Gln Cys Ser Arg Leu Leu Glu Leu Val Lys Asp Ile Ser
 145 150 155 160
 Ser His Leu Asp Val Thr Ala Leu Cys His Lys Ile Phe Leu His Ile
 165 170 175
 His Gly Leu Ile Ser Ala Asp Arg Tyr Ser Leu Phe Leu Val Cys Glu
 180 185 190
 Asp Ser Ser Asn Asp Lys Phe Leu Ile Ser Arg Leu Phe Asp Val Ala
 195 200 205
 Glu Gly Ser Thr Leu Glu Glu Val Ser Asn Asn Cys Ile Arg Leu Glu
 210 215 220
 Trp Asn Lys Gly Ile Val Gly His Val Ala Ala Leu Gly Glu Pro Leu
 225 230 235 240
 Asn Ile Lys Asp Ala Tyr Glu Asp Pro Arg Phe Asn Ala Glu Val Asp
 245 250 255
 Gln Ile Thr Gly Tyr Lys Thr Gln Ser Ile Leu Cys Met Pro Ile Lys
 260 265 270
 Asn His Arg Glu Glu Val Val Gly Val Ala Gln Ala Ile Asn Lys Lys
 275 280 285
 Ser Gly Asn Gly Gly Thr Phe Thr Glu Lys Asp Glu Lys Asp Phe Ala
 290 295 300
 Ala Tyr Leu Ala Phe Cys Gly Ile Val Leu His Asn Ala Gln Leu Tyr
 305 310 315 320
 Glu Thr Ser Leu Leu Glu Asn Lys Arg Asn Gln Val Leu Leu Asp Leu
 325 330 335
 Ala Ser Leu Ile Phe Glu Glu Gln Gln Ser Leu Glu Val Ile Leu Lys
 340 345 350
 Lys Ile Ala Ala Thr Ile Ile Ser Phe Met Gln Val Gln Lys Cys Thr
 355 360 365

EP 0 652 960 B9 (W1B1)

Ile Phe Ile Val Asp Glu Asp Cys Ser Asp Ser Phe Ser Ser Val Phe
 370 375 380
 5 His Met Glu Cys Glu Glu Leu Glu Lys Ser Ser Asp Thr Leu Thr Arg
 385 390 395 400
 Glu His Asp Ala Asn Lys Ile Asn Tyr Met Tyr Ala Gln Tyr Val Lys
 405 410 415
 10 Asn Thr Met Glu Pro Leu Asn Ile Pro Asp Val Ser Lys Asp Lys Arg
 420 425 430
 Phe Pro Trp Thr Thr Glu Asn Thr Gly Asn Val Asn Gln Gln Cys Ile
 435 440 445
 15 Arg Ser Leu Leu Cys Thr Pro Ile Lys Asn Gly Lys Lys Asn Lys Val
 450 455 460
 Ile Gly Val Cys Gln Leu Val Asn Lys Met Glu Glu Asn Thr Gly Lys
 465 470 475 480
 20 Val Lys Pro Phe Asn Arg Asn Asp Glu Gln Phe Leu Glu Ala Phe Val
 485 490 495
 Ile Phe Cys Gly Leu Gly Ile Gln Asn Thr Gln Met Tyr Glu Ala Val
 500 505 510
 25 Glu Arg Ala Met Ala Lys Gln Met Val Thr Leu Glu Val Leu Ser Tyr
 515 520 525
 His Ala Ser Ala Ala Glu Glu Glu Thr Arg Glu Leu Gln Ser Leu Ala
 530 535 540
 30 Ala Ala Val Val Pro Ser Ala Gln Thr Leu Lys Ile Thr Asp Phe Ser
 545 550 555 560
 Phe Ser Asp Phe Glu Leu Ser Asp Leu Glu Thr Ala Leu Cys Thr Ile
 565 570 575
 35 Arg Met Phe Thr Asp Leu Asn Leu Val Gln Asn Phe Gln Met Lys His
 580 585 590
 Glu Val Leu Cys Arg Trp Ile Leu Ser Val Lys Lys Asn Tyr Arg Lys
 595 600 605
 40 Asn Val Ala Tyr His Asn Trp Arg His Ala Phe Asn Thr Ala Gln Cys
 610 615 620
 Met Phe Ala Ala Leu Lys Ala Gly Lys Ile Gln Asn Lys Leu Thr Asp
 625 630 635 640
 45 Leu Glu Ile Leu Ala Leu Leu Ile Ala Ala Leu Ser His Asp Leu Asp
 645 650 655
 His Arg Gly Val Asn Asn Ser Tyr Ile Gln Arg Ser Glu His Pro Leu
 660 665 670
 50 Ala Gln Leu Tyr Cys His Ser Ile Met Glu His His His Phe Asp Gln
 675 680 685
 Cys Leu Met Ile Leu Asn Ser Pro Gly Asn Gln Ile Leu Ser Gly Leu
 690 695 700
 55 Ser Ile Glu Glu Tyr Lys Thr Thr Leu Lys Ile Ile Lys Gln Ala Ile
 705 710 715 720

EP 0 652 960 B9 (W1B1)

Leu Ala Thr Asp Leu Ala Leu Tyr Ile Lys Arg Arg Gly Glu Phe Phe
 725 730 735
 5 Glu Leu Ile Arg Lys Asn Gln Phe Asn Leu Glu Asp Pro His Gln Lys
 740 745 750
 Glu Leu Phe Leu Ala Met Leu Met Thr Ala Cys Asp Leu Ser Ala Ile
 755 760 765
 10 Thr Lys Pro Trp Pro Ile Gln Gln Arg Ile Ala Glu Leu Val Ala Thr
 770 775 780
 Glu Phe Phe Asp Gln Gly Asp Arg Glu Arg Lys Glu Leu Asn Ile Glu
 785 790 795 800
 15 Pro Thr Asp Leu Met Asn Arg Glu Lys Lys Asn Lys Ile Pro Ser Met
 805 810 815
 Gln Val Gly Phe Ile Asp Ala Ile Cys Leu Gln Leu Tyr Glu Ala Leu
 820 825 830
 20 Thr His Val Ser Glu Asp Cys Phe Pro Leu Leu Asp Gly Cys Arg Lys
 835 840 845
 Asn Arg Gln Lys Trp Gln Ala Leu Ala Glu Gln Gln Glu Lys Met Leu
 850 855 860
 25 Ile Asn Gly Glu Ser Gly Gln Ala Lys Arg Asn
 865 870 875

30

35

40

45

50

55

Applicant's or agent's file reference number	32083	International application No.
--	-------	-------------------------------

5

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

10

A. The indications made below relate to the microorganism referred to in the description on page 5 lines 6-9

15

B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet

Name of depositary institution
American Type Culture Collection

20

Address of depositary institution (including postal code and country)
12301 Parklawn Drive
Rockville, Maryland 20852
US

25

Date of deposit 4 May 1993	Accession Number ATCC 69296
-------------------------------	--------------------------------

30

C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet

"In respect of those designations in which a European patent is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 23(4) EPC)."

35

D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)

EP

40

E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)

The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")

45

50

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on
Authorized officer Doris L. Brock <i>DLB</i> EPC International Division	Authorized officer

55

Claims

1. A purified and isolated polynucleotide encoding the human cyclic GMP-binding, cyclic GMP-specific phosphodiesterase (cGB-PDE) polypeptide set out in SEQ ID NO: 23.
2. A purified and isolated polynucleotide encoding a human allelic variant of the cGB-PDE polypeptide set out in SEQ ID NO: 23, wherein said polynucleotide hybridizes at about 65°C in 3X SSC, 20 mM sodium phosphate pH 6.8, with washing at about 65°C in 2X SSC, to the non-coding strand of the DNA set out in SEQ ID NO: 22.
3. A polynucleotide according to Claim 1 or 2, which is a DNA.
4. A DNA according to Claim 3, which is a cDNA.
5. A DNA according to Claim 3, which is a genomic DNA.
6. An RNA transcript of a DNA according to Claim 5.
7. A DNA according to Claim 5, further comprising an endogenous expression control DNA sequence.
8. A vector comprising a DNA according to Claim 3.
9. A vector according to Claim 8, wherein the DNA is operatively linked to an expression control sequence.
10. A host cell stably transformed or transfected with a DNA sequence according to Claim 3 in a manner allowing the expression in said host cell of the cGB-PDE of SEQ ID NO: 23.
11. A method of producing a cGB-PDE polypeptide, which method comprises growing a host cell according to Claim 10 in a suitable nutrient medium and isolating cGB-PDE polypeptide from the cell or the medium of its growth.
12. A cGB-PDE polypeptide produced by the method of Claim 11.
13. A purified and isolated polynucleotide comprising the DNA sequence set out SEQ ID NO: 22.
14. A fragment of human cGB-PDE consisting of the amino acids 516 to 875 of SEQ ID NO: 23.
15. A fragment of human cGB-PDE consisting of the amino acids 1-494 of SEQ ID NO: 23.
16. A fragment of the human cGB-PDE consisting of the amino acids 1-549 of SEQ ID NO: 23.
17. A fragment of the human cGB-PDE consisting of the amino acids 515-819 of SEQ ID NO: 23.
18. An antibody specifically immunoreactive with the cGB-PDE as set forth in SEQ ID NO: 23.
19. An antibody according to Claim 18, which is a monoclonal antibody.
20. A hybridoma cell line producing an antibody according to Claim 19.
21. An antibody according to Claim 18, which is a humanized antibody.

Patentansprüche

1. Ein aufgereinigtes und isoliertes Polynukleotid, das für das humane, zyklisches GMP-bindende, für zyklisches GMP spezifische Phosphodiesterase(cGB-PDE)-Polypeptid kodiert, das in SEQ ID NO:23 dargestellt ist.
2. Ein aufgereinigtes und isoliertes Polynukleotid, das für eine humane, allelische Variante des in SEQ ID NO:23 dargestellten cGB-PDE-Polypeptids kodiert, wobei besagtes Polynukleotid bei ungefähr 65 °C in 3X SSC, 20 mM Natriumphosphat pH 6,8, unter Waschen bei ungefähr 65 °C in 2X SSC, an den nicht-kodierenden Strang der in

EP 0 652 960 B9 (W1B1)

SEQ ID NO:22 dargestellten DNA hybridisiert.

3. Ein Polynukleotid nach Anspruch 1 oder 2, das eine DNA ist.
- 5 4. Eine DNA nach Anspruch 3, die eine cDNA ist.
5. Eine DNA nach Anspruch 3, die eine genomische DNA ist.
6. Ein RNA-Transkript einer DNA nach Anspruch 5.
- 10 7. Eine DNA nach Anspruch 5, die weiter eine endogene Expressionskontroll-DNA-Sequenz umfaßt.
8. Ein Vektor, der eine DNA nach Anspruch 3 umfaßt.
- 15 9. Ein Vektor nach Anspruch 8, wobei die DNA an eine Expressionskontroll-Sequenz operativ geknüpft ist.
10. Eine Wirtszelle, die mit einer DNA-Sequenz nach Anspruch 3 auf eine Weise stabil transformiert oder transfiziert ist, die die Expression in besagter Wirtszelle der cGB-PDE von SEQ ID NO:23 ermöglicht.
- 20 11. Ein Verfahren zum Herstellen eines cGB-PDE-Polypeptids, wobei das Verfahren umfaßt: Wachsenlassen einer Wirtszelle nach Anspruch 10 in einem geeigneten Nährmedium und Isolieren von cGB-PDE-Polypeptid aus der Zelle oder dem Medium seines Wachstums.
- 25 12. Ein cGB-PDE-Polypeptid, erzeugt durch das Verfahren von Anspruch 11.
13. Ein aufgereinigtes und isoliertes Polynukleotid, das die in SEQ ID NO:22 dargestellte DNA-Sequenz umfaßt.
14. Ein Fragment humaner cGB-PDE, das die Aminosäuren 516 bis 875 von SEQ ID NO: 23 umfaßt.
- 30 15. Ein Fragment humaner cGB-PDE, das die Aminosäuren 1-494 von SEQ ID NO:23 umfaßt.
16. Ein Fragment der humanen cGB-PDE, das die Aminosäuren 1-549 von SEQ ID NO:23 umfaßt.
17. Ein Fragment der humanen cGB-PDE, das die Aminosäuren 515-819 von SEQ ID NO:23 umfaßt.
- 35 18. Ein Antikörper, der spezifisch immunreaktiv ist mit der cGB-PDE, wie dargestellt in SEQ ID NO:23.
19. Ein Antikörper nach Anspruch 18, der ein monoklonaler Antikörper ist.
- 40 20. Eine Hybridom-Zelllinie, die einen Antikörper nach Anspruch 19 produziert.
21. Ein Antikörper nach Anspruch 18, der ein humanisierter Antikörper ist.

45 **Revendications**

1. Polynucléotide purifié et isolé, codant le polypeptide, phosphodiesterase (cGB-PDE), de liaison au monophosphate de guanosine (GMP) cyclique et spécifique au GMP cyclique humain, indiqué dans la séquence n° 23.
- 50 2. Polypeptide purifié et isolé codant un variant allélique humain du polypeptide cGB-PDE indiqué dans la séquence n° 23, dans lequel ledit polypeptide s'hybride, à environ 65°C dans 3X SSC, du phosphate de sodium 20 mM, pH 6,8, avec lavage à environ 65°C dans 2X SSC, au brin non codant de l'ADN indiqué dans la séquence n° 22.
3. Polynucléotide selon la revendication 1 ou 2, qui est un ADN.
- 55 4. ADN selon la revendication 3, qui est un ADNc.
5. ADN selon la revendication 3, qui est un ADN génomique.

EP 0 652 960 B9 (W1B1)

6. Produit de transcription d'ARN d'un ADN selon la revendication 5.
7. ADN selon la revendication 5, comprenant de plus une séquence d'ADN de commande d'expression endogène.
- 5 8. Vecteur comprenant un ADN selon la revendication 3.
9. Vecteur selon la revendication 8, dans lequel l'ADN est lié de façon opératoire à une séquence de commande d'expression.
- 10 10. Cellule hôte transformée ou transfectée de façon stable par une séquence d'ADN selon la revendication 3 d'une manière permettant l'expression, dans ladite cellule hôte, de cGB-PDE de la séquence n° 23.
11. Procédé pour produire un polypeptide cGB-PDE, lequel procédé comprend la croissance d'une cellule hôte selon la revendication 10 dans un milieu nutritif approprié et l'isolation du polypeptide cGB-PDE à partir de la cellule ou
15 du milieu de sa croissance.
12. Polypeptide cGB-PDE produit par le procédé de la revendication 11.
13. Polynucléotide purifié et isolé comprenant la séquence d'ADN indiquée dans la séquence n° 22.
20
14. Fragment de cGB-PDE humaine, constitué des acides aminés 516 à 875 de la séquence n° 23.
15. Fragment de cGB-PDE humaine, constitué des acides aminés 1-494 de la séquence n° 23.
- 25 16. Fragment de cGB-PDE humaine, constitué des acides aminés 1-549 de la séquence n° 23.
17. Fragment de cGB-PDE humaine, constitué des acides aminés 515-819 de la séquence n° 23.
18. Anticorps spécifiquement immunoréactif avec la cGB-PDE comme indiqué dans la séquence n° 23.
30
19. Anticorps selon la revendication 18, qui est un anticorps monoclonal.
20. Lignée cellulaire hybridome produisant un anticorps selon la revendication 19.
- 35 21. Anticorps selon la revendication 18, qui est un anticorps humanisé.

40

45

50

55

CGB-PDE	FOMKHEVLCK	WILSVKKNYR	K..NVAYHNWR	HAFNTAQCMF	AALKAGKIQK	626
ROS- α	FHIPQEALVR	FMYSLSKGYR	R..ITYHNWR	HGFNVGQTMF	SLLVTGKLR	582
ROS- β	FOIPQEVLR	FLFSVSKGYR	R..ITYHNWR	HGFNVAQTMF	TLLMTGKLS	580
CONE- α '	FKVPVEVLR	WMTYVRKGYR	A..VTYHNWR	HGFNVGQTMF	TLLMTGRLKK	580
CGS	YKIDCPTLAR	FCLMVKKGYR	D.P.PYHNWM	HAFSVSHFCY	LLYKNLELTN	659
61 kCAM	FKIPVSCLIA	FAEALEVGYS	KYKNPYHNI	HAADVTQTVH	YIMLHTGIMH	242
63 kCAM	FKIPTVFLMT	FLDALETGYG	KYKNPYHNI	HAADVTQTVH	CFLLRRTGMVH	244
RATDUNCE	FOIPADTLR	YLLTLEGHYH	S.NVAYHNSI	HAADVOSAH	VLLGTPALEA	125
DROSDUNCE	.MIPPKTFLN	FMSTLEDHYV	K.DNPFHNSL	HAADVTQSTN	VLLNTPALEG	48
CONSERVED	-----*	-----Y-	-----HN-*	H-----**	-----*	
CGB-PDE	RLTDLEILAL	LIAALSHDL	HRGVNNSYIQ	RSEHPLAQLY	CH..SIMEHH	674
ROS- α	YFTDLEALAM	VTAAFCHDID	HRGTNNLYQM	KSQNPLAKLH	GS..SILERH	630
ROS- β	YYTDLEAFAM	VTAGLCHDID	HRGTNNLYQM	KSQNPLAKLH	GS..SILERH	628
CONE- α '	YYTDLEAFAM	LAAAFCHDID	HRGTNNLYQM	KSTSPARLH	GS..SILERH	628
CGS	YLEDMEIFAL	FISCMCHDLD	HRGTNNSFOV	ASKSVLAALY	SSEGSMERH	709
61 kCAM	WLTELEILAM	VFAAIIHDYE	HIGTTNFI	QTRSDVAILY	.NDRSVLENH	291
63 kCAM	CLSEIEVLAI	IFAAIIHDYE	HIGTTNSFI	QTKSEQAILY	.NDRSVLENH	293
RATDUNCE	VFTDLEVLA	IFACAIHDVD	HPGVSNOFLI	NTNSELALMY	.NDSSVLENH	174
DROSDUNCE	VFTPLEVGG	LFAACIHDVD	HPGLTNOFLV	NSSSELALMY	.NDESVLENH	97
CONSERVED	-----E	-----HD--	H-G--N-*	-----*-A	-----S-E-H	

FIGURE 1A

CGB-PDE	HFDOCLMILN	SPGNQILSGL	SIEEYKTTLK	IIKQAILATD	LALYIKRRGE	714
ROS- α	HLEFGKTLR	DESLNIFQNL	NRRQHEHAIH	MMDIAIIATD	LALYCKKRTM	680
ROS- β	HLEFGKFLS	EETLNIYQNL	NRRQHEHVIH	LMDIAIIATD	LALYFKKRTM	678
CONE- α'	HLEYSKTLQ	DESLNIFQNL	NKRQYETVIH	LFEVAIIATD	LALYFKKRTM	678
CGS	HFAQAIAILN	THGCNIFDHF	SRKDYORMLD	LMRDIILATD	LAHHLRIFKD	748
61 KCAM	HVSAAYRLMQ	EEEMNVLINL	SKDDWRDLRN	LVIEMVLSTD	MSGHFQQIKN	326
63 KCAM	HISVFRMMQ	DDEMNIIFINL	TKDEFVELRA	LVIEMVLATD	MSCHFQQVKS	328
RATDUNCE	HLAVGFKLLQ	GENCDIFQNL	STKQKLSLRR	MVIDMVLATD	MSKHMSLLAD	220
DROSDUNCE	HLAVAFKLLQ	NOGCCIFCNM	OKKOROTLRK	MVIDIVLSTD	MSKHMSLLAD	143
CONSERVED	H-----	-----	-----	-----TD	---*--*--*--	
CGB-PDE	FFELIMKN..QF	NLEDPHQKEL	FLAMLMTACD	LSAITKPWPI	764
ROS- α	FOKIVDOSKT	YETQQEWTQY	MMLDQTRKEI	VMAMMMTACD	LSAITKPWEV	730
ROS- β	FOKIVDESKN	YEDRKSWEY	LSLETTTRKEI	VMAMMMTACD	LSAITKPWEV	728
CONE- α'	FOKIVDACEK	METEEEEAIKY	VTIDPTKKEI	IMAMMMTACD	LSAITKPWEV	728
CGS	LQKMAE....VGY	DRTNKHHS	LLCLLMTSCD	LSDQTKGWKT	798
61 KCAM	IRNSLQOPEG	L.....DKAK	TMSLILHAAD	ISHPAKSWKL	376
63 KCAM	MKTALQOLER	I.....DKSK	ALSLLLHAAD	ISHPTKQWSV	378
RATDUNCE	LKTMVETKKV	T.....SLGVL	LLDNYSDRIQ	VLOS LVHCAD	LSNPAKPLPL	270
DROSDUNCE	LKTMVETKKV	A.....GSGVL	LLDNYTDRIQ	VLENLVHCAD	LSNPTKPLPL	193
CONSERVED	*-----	-----	-----*	-----D	-S**--K----	

FIGURE 1B

CGB-PDE	QORIAELVAT	EFFDQGDRE	KELNIEPADL	MNREKKNKIP	SMQVGFID..	812
ROS- α	QSKVALLVAA	EFWEQGDLE	TVLQQNP	MDRNKADEL	KLOVGFID..	778
ROS- β	QSKVALLVAA	EFWEQGDLE	TVLQQQPIP	MDRNKAAEL	KLOVGFID..	776
CONE- α'	QSOVALLVAN	EFWEQGDLE	TVLQQQPIP	MDRNKKDEL	KLOVGFID..	776
CGS	TRKIAELIYK	EFFSQGDLE	A.MGNRPMEM	MDREKAY.IP	ELQISFME..	844
61 kCAM	HHRWTMALME	EFFLOGDKEA	EL..GLPFSP	LCDRKSTMVA	OSQIGFID..	422
63 kCAM	HSRWTKALME	EFFRQGDKEA	EL..GLPFSP	LCDRTSTLVA	OSQIGFID..	424
RATDUNCE	YROWTERIMA	EFFQQGDRE	ES..GLDISP	MCDKHTASVE	KSQVGFID..	316
DROSDUNCE	YKRWVALLME	EFFLQGDKER	ES..GMDISP	MCDRHNATIE	KSQVGFID..	239
CONSERVED	*-----*-	EF--QGD-E-	-----	-----	--Q--F----	

FIGURE 1C

CGB-PDE	LLELVKDISS	HLDVTALCHK	IFLHIHGLIS	ADRYSLFLVC	EDSSNDKFLI	188
CGS	ILQLCGELYD	.LDASSLQLK	VLOYLQOQETQ	ASRCCLLLVS	EDN..LQ.LS	245
CONE- α '	LLEVL..LEE	AGSVELAAHR	ALORLAQLLO	ADRCMFLCR	ARNGTPE.VA	106
ROS- β	LFELVQDMQE	NVNMERVVFK	ILRRLCSILH	ADRCSLFMYR	QRNGVAE.LA	107
ROS- α	...LLRDFOD	NLOAEKCVFN	VMKKLCFLLQ	ADRMSLFMYR	ARNGIAE.LA	109
CONSERVED	-----	-----	-----	A-R-----	-----	
CGB-PDE	SRLFDVAEGS	TLEE...ASN	NCIRLEWNGK	IVGHVAAFGE	PLNIKDAYED	237
CGS	CKVIG...DK	VLEE.....	.EISFPLTTG	RLGOVVEDKK	SIQLKDLTSE	292
CONE- α '	SKLLDVTPTS	KFEDNLVVPD	REAVFPLDVG	IVGWAHTKK	TFNVDPVKKN	154
ROS- β	TRLFSVQPS	VLEDCLVPPD	SEIVFPLDIG	VVGHVAQTKK	MVNVQDVMEC	155
ROS- α	TRLFNVHKDA	VLEECLVAPD	SEIVFPLDMG	VVGHVALSKK	IVNVPNTEED	157
CONSERVED	-----	---E-----	-----G	--G-V-----	-----	
CGB-PDE	PRFNAEVDQI	TGYKTQSILC	MPIKNHR.EE	VVGVAQAINK	KSGNGGTFTE	287
CGS	DM..QQLOSM	LGCEVOAMLC	VPVISRATDQ	VVALACAFNK	..LGGDLFTD	342
CONE- α '	SHFSDFMDKQ	TGYVTRNLLA	TPIV..MGKE	VLAVFMAVNK	..VDASEFSK	204
ROS- β	PHFSSFADL	TDYVTRNILA	TPIM..NGKD	VVAVIMAVNK	..LDGPCFTS	205
ROS- α	EHFCDFVDTL	TEYQTKNILA	SPIM..NGKD	VVAIIMAVNK	..VDGPHFTE	207
CONSERVED	-----	-----L-	-P-----	V-----A-NK	-----F--	

FIGURE 2A

CGB-PDE	KDEKDFAAAYL	AFCGIVLHNA	QLYETSLEN	KRNQVLLDLA	SLIFEEQOQL	337
CGS	QDEHVIOHCF	HYTSTVLTST	LAFQKEQKLK	CECOALLOVA	KNLFTHLDDV	390
CONE- α'	QDEEVFSKYL	SFVSIILKLH	HTNYLYNIES	RRSQILMWSA	NKVFEELTDV	252
ROS- β	EDEDVFLKYL	NFGTLNLKIY	HYSYLHNCET	RRGOVLLWSA	NKVFEELTDI	253
ROS- α	NDEEILLKYL	NFANLIMKVF	HLSYLHNCET	RRGOILLWSG	SKVFEELTDI	255
CONSERVED	-DE-----	-----	-----	---Q-L----	---F-----	
CGB-PDE	EVILKkiaat	IISFMQVQKC	TIFIVD.EDC	SDSFSSVFHM	ECEELEKSSD	361
CGS	SVLLQEIITE	ARNLSNAEIC	SVFLID...Q	NELVAKVFDG	GVLEDESY..	409
CONE- α'	ERQFHkALYT	VRTYlNCERY	SIGLLDMTKE	KEYF.DEWPV	KPGEVEPYKG	301
ROS- β	ERQFHkAFYT	VRAylNCDRY	SVGLLDMTKE	KEFF.DVWPV	LMGEAQAYSg	302
ROS- α	ERQFHkALYT	VRAFlNCDRY	SVGLLDMTKQ	KEFF.DVWPV	LMGEAPPYAG	304
CONSERVED	-----	-----	---D-----	-----	---E-----	
CGB-PDE	TLTRE.....RDAMRINY	MYAQYVKNTM	411
CGSEIRI...	.PADQ.....	GIAGHVATTG	459
CONE- α'	PKTPDGREVI	FYKIIDYILH	GKEEIKVIPT	PPMDHWTLIS	GLPTYVAENG	351
ROS- β	PRTPDGREIL	FYKVIDYILH	GKEDIKVIPT	PPADHWALAS	GLPTYVAESG	352
ROS- α	PRTPDGREIN	FYKVIDYILH	GKEDIKVIPT	PPPDHWALVS	GLPTYVAONG	354
CONSERVED	-----	-----	-----	---D-----	---V-----	

FIGURE 2B

cGB-PDE	EPLNIPDVSK	DKRFPWTNEN	MGNINQQCIR	SLLCTPIKNG	KKNKIVGVCCQ	459
cGS	OILNIPDAYA	HPLFY..RGV	DDSTGRF.TR	NILCFPIKN.	ENQEVIGVAE	499
CONE- α '	FICNMLNAPA	DEYFTFOKGP	VDETGWV.IK	NVLSLPIVN.	KKEDIVGVAT	399
ROS- β	FICNIMNAPA	DEMNFQOEGP	LDDSGWI.VK	NVLSMPIVN.	KKEEIVGVAT	400
ROS- α	LICNIMNAPS	EDFFAFOKEP	LDESGMM.IK	NVLSMPIVN.	KKEEIVGVAT	402
CONSERVED	---N---	---F---	-----	--L--PI-N-	-----GV--	
cGB-PDE	LVNKMEETG	KVKAFNRNDE	QFLEAFVIFC	GLGIONTOMY	EAVRAMAKQ	506
cGS	LVNKG... .	PWFSKFDE	DLATAFSIYC	GISIAHSLLY	KKVNEAQYRS	541
CONE- α '	FYNRKDG... .	KPFDEYDE	HIAETLTOFL	GWSLLNTDTY	EKMNKLENRK	441
ROS- β	FYNRKDG... .	KPFDEODE	VLMESLTOFL	GWSVLNTDTY	DKMNKLENRK	442
ROS- α	FYNRKDG... .	KPFDEMDDE	TLMESLAQFL	GWSVLNPDTY	ELMNKLENRK	444
CONSERVED	---N---	---F---DE	-----	G-----Y	-----	
cGB-PDE	MVTLEVLSYH	ASAAAAE				526
cGS	HLANEMMYH	MKVSDDE				561
CONE- α '	DIAQEMLMNH	TKATPDE				461
ROS- β	DIAODMVLVYH	VRCDREE				462
ROS- α	DIFODMVKYH	VKCDNEE				464
CONSERVED	-----H	-----E				

FIGURE 2C

CGB-PDE A EPLNIKDAYEDPRF... NAEVDQITGYKTOSILCMPIKMH. REEVVGAOAIN. KKSGN
ROS- α A KIVNVPNTEDEHF... CDFVDTLLEYQTKNILASPIMNG. K. DVVAIIMAVN. KVDGP
ROS- β A KMNVDVMECPHF... SSFADELTDYVTRNILATPIMNG. K. DVVAVIMAVN. KLDGP
CONE- α ' A KTFNVPDVKKNSHF... SDFMDKQTGYVTRNILATPIVMG. K. EVLAVFMAVN. KVDAS
CGS A KSIQLKDLTSEDM... QQLOSMGLGCEVOAMLCVPPVISRATDQVVALACAFN. KLGGD
CGB-PDE B EPLNIPDVSKDKRFPWTNENMGNIQQCIRSLLCTPIKNGKKNKVIQVQLVN. KMEET
ROS- α B LICNIMNAPSEDFFAFQKEPLDE. SGMKIKNVLSMPIVNK. KEEIVGVATFYNRKDGKP
ROS- β B FICNIMNAPADEMFNFOEGPLDD. SGWIVKNVLSMPIVNK. KEEIVGVATFYNRKDGKP
CONE- α ' B FICMMLNAPADEYFTFOKGPVDE. TGWIKNVLSLPIVNK. KEDIVGVATFYNRKDGKP
CGS B QILNIPDAYAHLF... YRGVDDSTGFRTRNIIICFPPIKNE. NOEVIGVAELVN. KINGP
CONSERVED B -----L--P*-----*****-----N-K-----

CGB-PDE A GG... TFTEKDEKFAAYLAFCGIVLHMAQL. YE
ROS- α A HFTENDEEILLKYLNFANLIMKVFHLSY.
ROS- β A CFTSEDEDEVLKYLNFGLNLKLYHLSY.
CONE- α ' A EFSKODEEVFSKYLSEFVSIILKLHHTNY.
CGS A LFTDODEHVIOHCFHYTSTVL. TSTLAFQ
CGB-PDE B TGKVKAFNRNDEQFLEAFVIFCGLGIQNTQM. YE
ROS- α B FDEMETLMESLAQFLGWSV. LNPDTYE
ROS- β B FVEQDEVLMESLTQFLGWSV. LNTDTYD
CONE- α ' B FDEYDEHIAETLTQFLGWSL. LNTDTYE
CGS B WFSKFDEDLATAFSIYCGISI. AHSLLYK
CONSERVED B -----F---DE-----*-----*-----

FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8