

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 703 216 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
07.04.1999 Bulletin 1999/14

(51) Int Cl.⁶: **C07C 257/18**, C07C 257/14,
C07D 211/14, A61K 31/155

(21) Application number: **95306596.8**

(22) Date of filing: **19.09.1995**

(54) **Amidinophenol derivatives as protease inhibitors**

Amidinophenol-Derivate als Protease-hemmende Verbindungen

Dérivés d'amidinophénol comme inhibiteurs de protéase

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU NL PT
SE**

(30) Priority: **20.09.1994 JP 250158/94**

(43) Date of publication of application:
27.03.1996 Bulletin 1996/13

(73) Proprietor: **ONO PHARMACEUTICAL CO., LTD.**
Chuo-ku Osaka 541 (JP)

(72) Inventors:
• **Nakai, Hisao, c/o Minase Research Institute
Mishima-gun, Osaka (JP)**
• **Kamiyasu, Koumei,
c/o Minase Research Institute
Mishima-gun, Osaka (JP)**

(74) Representative: **Bentham, Stephen**
J.A. KEMP & CO.
14 South Square
Gray's Inn
London WC1R 5LX (GB)

(56) References cited:
EP-A- 0 048 433 EP-A- 0 588 655
EP-A- 0 656 349 CH-A- 520 657
FR-A- 2 556 720

- **BIOMEDICAL RESEARCH**, vol.15, no.1, January 1994, TOKYO, JP pages 55 - 8 A. **MATSUSHIMA ET. AL.** 'Inhibition of a mite protease (Df protease) by synthetic inhibitors'
- **JOURNAL OF MEDICINAL CHEMISTRY.**, vol.38, no.22, 27 October 1995, WASHINGTON US pages 4508 - 17 K. **SENOKUCHI ET. AL.** 'New Orally Active Serine Protease Inhibitors: Structural Requirements for Their Good Oral Activity'

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 703 216 B1

Description

[0001] The present invention relates to amidinophenol derivatives, processes for the preparation thereof and leukotriene (LT)₄ antagonist, phospholipase A₂ and/or trypsin inhibitor containing amidinophenol derivatives as active ingredient.

[0002] The metabolic pathways by which various compounds are biosynthesized, in vivo, from arachidonic acid as a common starting material is called the arachidonic acid cascade.

[0003] Lipoxygenase, for example, 5-lipoxygenase, 12-lipoxygenase or 15-lipoxygenase, respectively, acts on arachidonic acid to produce 5-HPETE, 12-HPETE or 15-HPETE from arachidonic acid.

[0004] The above mentioned HPETEs are converted into 5-HETE, 12-HETE or 15-HETE, by converting a peroxy group into a hydroxy group by the action of peroxidase, and 5-HPETE is also converted into LTA₄.

[0005] LTA₄ is converted into LTB₄ or LTC₄ by enzymatic reaction (see Biochem. Biophys. Res. Commun., 91, 1266 (1979), Prostaglands, 19(5), 645).

[0006] Recently a number of properties of LTB₄ have been revealed.

[0007] It is understood that LTB₄ has strong chemotactic and adhesive activity and degranulation activity on leukocytes (see Nature, 286, 264 (1980), Proc. Nat. Acad. Sci. USA, 78, 3887 (1981))

[0008] LTB₄ also has strong calcium ionophore action, and attacks various cells, and it is considered to accelerate release of metabolites of arachidonic acid from these cells (see J. Biol. Chem, 257, 4746 (1982)).

[0009] High levels of LTB₄ have also been found at sites of various inflammations, for example, rheumatism, spondyl arthritis, gout, psoriasis, ulcerative colitis and respiratory tract diseases, thereby demonstrating that LTB₄ is closely associated with various inflammations (see J. Clin. Invest., 66, 1166 (1980); Lancet II 1122-1123 (1982); J. Invest. Dermatol., 82, 477-479 (1984); Gastroenterology 86, 453-460 (1984)).

[0010] It is therefore considered that LTB₄ antagonists are useful as anti-inflammatory agents or anti-allergic agents.

[0011] It is known that LTB₄ antagonists are also useful for the treatment of rheumatoid arthritis, inflammatory bowel diseases, psoriasis, nonsteroidal anti-inflammatory agent-induced stomach diseases, adult respiratory distress syndrome, cardiac infarction, allergic rhinitis, hemodialysis-induced neutropenia, anaphase asthma (see the specification of the Japanese Patent Kokai No. 5-239008).

[0012] Phospholipase A₂ (PLA₂) is an enzyme which acts on phospholipids existing in cell membranes. It hydrolyzes an ester bond at the second position of the phospholipids. There are two known kinds of PLA₂, membrane-associated PLA₂ and pancreatic PLA₂.

[0013] Membrane-associated PLA₂ acts on phospholipids to release arachidonic acid (AA) from the phospholipids. The AA is converted into prostaglandins, thromboxanes and leukotrienes, which are physiologically active substances inducing various inflammatory diseases and allergic diseases.

[0014] On the other hand, pancreatic PLA₂ degrades phospholipids and destroys cell membrane, thereby producing lysolecithin having strong cytotoxicity. Recently, much importance has been attached to pancreatitis, severity in pancreatitis and multiple organ failure induced by this destructive activity on cell membrane.

[0015] It is also reported that membrane-associated PLA₂ is also concerned with these diseases.


[0016] Accordingly, the inhibition of PLA₂ leads to the suppression of the release of AA, a precursor of various physiologically active substances, and therefore, it is considered to be useful for the prevention and/or the treatment of various inflammatory and allergic diseases. Furthermore, it is considered to be useful for the prevention and/or the treatment of pancreatitis, severity in pancreatitis and multiple organ failure due to the inhibition of the destructive activity on cell membranes.

[0017] It is also known that the inhibition of various proteases such as trypsin, plasmin, thrombin and kallikrein, especially trypsin is useful for the prevention and/or the treatment of disseminated intravascular coagulation, pancreatitis, severity in pancreatitis and multiple organ failure.

[0018] In the specifications of EP-A-588655 and 656349, it is disclosed that certain amidinophenol compounds of the formula (IA) depicted hereinafter have an inhibitory activity on PLA₂ and an inhibitory activity on trypsin and are useful for the prevention and/or the treatment of various inflammatory or allergic diseases, disseminated intravascular coagulation, pancreatitis, severity in pancreatitis and multiple organ failure.

[0019] Several amidinophenol derivatives are already known as LTB₄ antagonists. They are disclosed in WO 94/11341, the specification of Japanese Patent Kokai No. 5-239008 and EP-518819. In these applications, it is disclosed that amidinophenylthioalkoxy(thio)benzamide is useful as LTB₄ antagonist.

[0020] For example, it is described in the specification of EP-518819 that the compound of the formula (A) :



10 wherein R^{1a} is amino which is mono- or disubstituted by a substituent selected from an aliphatic hydrocarbon radical, an araliphatic hydrocarbon radical, an aromatic radical, and a cycloaliphatic hydrocarbon radical or is amino which is disubstituted by a divalent aliphatic hydrocarbon radical; R^{2a} is hydrogen, halogen, trifluoromethyl, an aliphatic hydrocarbon radical, or is hydroxy which is etherified by an aliphatic alcohol, araliphatic alcohol, or aromatic alcohol or which is esterified by an aliphatic or araliphatic carboxylic acid,

15 R^{3a} is hydrogen or an acyl radical which is derived from an organic carbonic acid, an organic carboxylic acid, a sulfonic acid, or a carbamic acid, X_{1a} and X_{3a} , independently of one another, are oxygen (-O-) or sulphur (-S-), X_{2a} is a divalent aliphatic hydrocarbon radical which may be interrupted by an aromatic radical, wherein the phenyl rings of formula (A) may be, independently of one another, further substituted by one or more
 20 substituents selected from halogen, trifluoromethyl, an aliphatic hydrocarbon radical, hydroxy, and hydroxy which is etherified by an aliphatic alcohol or which is esterified by an aliphatic or araliphatic carboxylic acid; wherein aryl moieties in the above definitions may be, independently of one another, further substituted by one or more substituents selected from halogen, trifluoromethyl, an aliphatic hydrocarbon radical, hydroxy, and hydroxy which is etherified by an aliphatic alcohol or which is esterified by an aliphatic or araliphatic carboxylic acid; and
 25 wherein a cycloaliphatic hydrocarbon radical may be substituted by an aliphatic radical, and pharmaceutically acceptable salts thereof are useful as LTB_4 antagonist.

[0021] Biomedical Research 15(1) 55-58, 1994 discloses synthetic guanidine and amidine derivatives which are inhibitors of a house dust mite protease.


30 [0022] In the amidinophenyl(oxy)(thio)alkoxy(thio)benzamide compounds represented by EP-518819 as prior art, it can be seen that the existence of -O(or S)-alkylene-O(or S)- (with the proviso that the alkylene may be interrupted by an aromatic group) is essential in the prior art compounds.

[0023] It has now been discovered that compounds in which it is essential that the amidinophenyl is bonded to the phenyl group via an ester or amide group possess useful properties as LTB_4 antagonists, phospholipase A_2 and/or trypsin inhibitors. The amidinophenol derivatives of the invention possess a strong antagonistic activity on LTB_4 .


35 [0024] That is to say, as mentioned above, compounds of the formula (IA) and processes for the preparation thereof are disclosed in EP-A-588655 and EP-A-656349.

[0025] That is to say, the compound of the present invention, of the formula (IB) has an inhibitory activity on phospholipase A_2 and an inhibitory activity on trypsin and it is useful for the prevention and the treatment of various inflammatory, allergic diseases, disseminated intravascular coagulation, pancreatitis, severity in pancreatitis and multiple organ failure.



[0026] The present invention accordingly provides a new amidinophenol derivative of the formula (IB):




50 wherein:



is a group of the formula:




or








(in which

R^0 is hydrogen, or C1-4 alkyl, or C1-4 alkoxy),
T is NH or oxygen,
E is a single bond, or
a group of the formula:




A_0 is a single bond, C1-4 alkylene, -oxy-(C1-4)alkylene-, -thio-(C1-4)alkylene-, C2-8 alkenylene, or C2-8 alkenylene which is substituted by carboxy or by C1-4 alkoxy carbonyl, R^{100} , R^{200} , R^{300} and R^{400} each independently, is hydrogen or C1-4 alkyl,
R is a group of the formula:





(in which the grouping:



is a 4-10 membered hetero ring containing one or two nitrogen atoms),
 R^{50} , R^{60} and R^{70} each independently, is,


- (i) hydrogen,
- (ii) C1-8 alkyl,
- (iii) C2-8 alkenyl
- (iv) $-\text{COOR}^{110}$ (in which R^{110} is hydrogen or C1-4 alkyl unsubstituted or substituted by phenyl),
- (v) $-(\text{C1-8 alkylene})-\text{COOR}^{110}$ (in which R^{110} has the same meaning as hereinbefore defined),

- (vi) -(C2-8 alkenylene)-COOR¹¹⁰ (in which R¹¹⁰ has the same meaning as hereinbefore defined),
- (vii) C4-7 cycloalkyl,
- (viii) -(C1-4 alkylene)-(4-7 membered hetero ring containing one oxygen),
- (ix) -(C1-4 alkylene)-(4-7 membered hetero ring containing one nitrogen),
- (x) phenyl,
- (xi) C1-8 alkyl which is substituted by one or two phenyl,
- (xii) -(C1-4 alkylene)-O-benzoyl,
- (xiii) -(C1-4 alkylene)-CONH-(C1-4 alkylene)-NR¹²⁰R¹³⁰,
- (xiv) -(C1-4 alkylene)-COO-(C1-4 alkylene)-NR¹²⁰R¹³⁰,
- (xv) -(C1-4 alkylene)-COO-amidinophenyl,
- (xvi) -(C1-4 alkylene)-CONH-(C1-4 alkyl which is substituted by one or two -COOR¹¹⁰) (in which R¹¹⁰ has the same meaning as hereinbefore defined),
- (xvii) -(C1-4 alkylene)-CONR¹²⁰R¹³⁰, or
- (xviii) (C1-4)alkoxy(C1-4)alkyl,

R⁸⁰ and R⁹⁰ each independently, is C1-4 alkyl or -(C1-4 alkylene)-phenyl,
R¹²⁰ and R¹³⁰ each independently, is hydrogen, C1-4 alkyl, or C2-8 alkenyl, with the provisos that:

- (1) R⁵⁰ and R⁶⁰ in the formulae (i) and (iii), and R⁵⁰, R⁶⁰ and R⁷⁰ in the formulae (ii) and (iv), do not represent hydrogen at the same time,
- (2) when at least one substituent in R⁵⁰, R⁶⁰, R⁷⁰ and A₀ represent substituent containing -COO-t-Bu, the other groups do not represent groups containing carboxy,
- (3) R¹²⁰ and R¹³⁰ do not represent hydrogen at the same time,
- (4) when

T is oxygen,
the grouping:




is the formula (i) as hereinbefore described,
E is a single bond,
A₀ is a single bond, C1-4 alkylene or vinylen which is optionally substituted by one or two C1-4 alkyl, and
R is the formula (i) as hereinbefore described, then at least one group in R⁵⁰, and R⁷⁰ is

- (viii) -(C1-4 alkylene)-(4-7 membered hetero ring containing one oxygen),
- (ix) -(C1-4 alkylene)-(4-7 membered hetero ring containing one nitrogen),
- (xi-1) C1-8 alkyl which is substituted by two phenyl,
- (xii) -(C1-4 alkylene)-O-benzoyl,
- (xiii) -(C1-4 alkylene)-CONH-(C1-4 alkylene) -NR¹²⁰R¹³⁰,
- (xiv) -(C1-4 alkylene)-COO-(C1-4 alkylene) -NR¹²⁰R¹³⁰,
- (xv) -(C1-4 alkylene)-COO-amidinophenyl,
- (xvi) -(C1-4 alkylene)-CONH-(C1-4 alkyl which is substituted by one or two -COOR¹¹⁰) (in which R¹¹⁰ has the same meaning as hereinbefore defined), or
- (xvii) -(C1-4 alkylene)-CONR¹²⁰R¹³⁰ (in which R¹²⁰ and R¹³⁰ each independently is C1-4 alkyl or C2-8 alkenyl);

(5) when

T is oxygen,
the grouping:



is the formula (i) as hereinbefore defined,

E is a single bond,

A₀ is a single bond, C1-4 alkylene or vinylene which may be optionally substituted by one or two C1-4 alkyl, and R is the formula (ii) as hereinbefore defined,


then R⁵⁰, R⁶⁰ and R⁷⁰ do not represent hydrogen; and

(6) the compound of formula (IB) is not p-amidinophenyl p-[(2-isopropyl-1-imidazolyl)methyl]benzoate;

or 4-(4-amidinophenoxy)carbonyl-α-methylcinnamic acid N-1-(S)-ethoxycarbonyl-2-benzoyloxyethylamide; and non-toxic salts thereof or non-toxic acid addition salts thereof.

[0027] The compounds of the invention may form hydrates: it is to be understood that such hydrates form part of the present invention and that references to the compounds in this specification including the accompanying claims are to be understood as embracing the hydrates.

[0028] The invention also provides the use of a compound of the formula (IA):



wherein R¹ and R² each independently, is:

(i) hydrogen, or

(ii) -COOR⁴ (in which R⁴ is C1-3 alkyl);

[0029] A is

(i) a single bond,

(ii) C1-4 alkylene, or


(iii) -C(R⁵)=C(R⁶)- (in which R⁵ and R⁶ each independently, is hydrogen or C1-4 alkyl);

[0030] R³ is

(i) -CON(R⁷)R⁸,

(ii) -CONR⁹-CH(R⁷)R⁸, or

(iii)



(wherein R^7 and R^8 each independently, is

- (1) hydrogen,
- (2) phenyl,
- (3) -(C1-4 alkylene)-phenyl,
- (4) -(C1-4 alkylene)-phenyl which is substituted by one or two $-R^{11}-COOR^{12}$
(in which R^{11} is a single bond or C1-8 alkylene, and R^{12} is hydrogen or C1-4 alkyl),
- (5) C1-5 alkyl,
- (6) C2-10 alkenyl containing one to three double bonds,
- (7) $-R^{11a}-COOR^{12}$
(in which R^{11a} is


- (a) a single bond,
- (b) C1-8 alkylene,
- (c) C2-8 alkenylene, or
- (d) C4-8 alkenylene in which one or two carbon atoms in the main chain are replaced by sulfur, and R^{12} has the same meaning as hereinbefore defined), or

- (8) C3-7 cycloalkyl;

[0031] R_9 is

- (1) hydrogen,
- (2) $-R^{11}-COOR^{12}$ (in which the various symbols have the same meanings as hereinbefore defined), or
- (3) C2-6 alkoxyalkyl;

the grouping:



is a 4-7 membered mono hetero ring contain one or two nitrogen;

[0032] R^{10} is

- (1) hydrogen, or
 - (2) -(C1-4 alkylene)-phenyl,
- with the proviso that:

- (1) both R^7 and R^8 do not represent hydrogen at the same time,
- (2) when at least one group in R^7 , R^8 , and R^9 represent the group containing $-COO-t-Bu$, the other groups do not represent the groups containing carboxy;

non-toxic salts thereof and non-toxic acid-addition salts thereof, in the manufacture of a medicament for the prevention and/or treatment of rheumatoid arthritis, inflammatory bowel diseases, psoriasis, nonsteroidal anti-inflammatory agent-induced stomach diseases, adult respiratory distress syndrome, cardiac infarction, allergic rhinitis, hemodialysis-induced neutropenia or anaphase asthma.

[0033] The invention also provides the use of a compound of formula (IB) or a non-toxic salt thereof or non-toxic acid addition salt thereof in the manufacture of a medicament for the prevention and/or treatment of a condition which can be ameliorated by the administration of a leukotriene B_4 antagonist or a phospholipase A_2 or trypsin inhibitor.

[0034] Throughout the specification, it will be understood by those skilled in the art, that all isomers are included in the present invention. For example, the alkyl, alkoxy, alkylene, alkenylene and alkynylene groups include straight-chain and also branched-chain ones and the double bonds in the alkenylene group include E, Z and EZ mixtures. Accordingly, all isomers produced by the existence of asymmetric carbon atoms are included in the present invention when

branched-chain alkyl, alkoxy, alkylene, alkenylene and alkynylene are present.

[0035] Explanation of various symbols in the formula (IB) is shown below.

[0036] The C1-3 alkyl group means methyl, ethyl, propyl and the isomers thereof. C1-4 alkyl group means butyl in addition to the groups above mentioned, and the isomers thereof. C1-5 alkyl group means pentyl in addition to the groups above mentioned, and the isomers thereof.

[0037] C1-4 alkylene group means methylene, ethylene, trimethylene, tetramethylene and the isomers thereof. C1-8 alkylene group means pentamethylene, hexamethylene, heptamethylene, octamethylene in addition to the groups above mentioned, and the isomers thereof.

[0038] C2-8 alkenylene group containing one to three double bonds means ethylene, trimethylene, tetramethylene, pentamethylene, hexamethylene, heptamethylene or octamethylene in which one to three groupings $-CH_2-CH_2-$ (except those at each end of the group) are replaced by double bonds.

[0039] C3-7 cycloalkyl group means cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl.

[0040] The 4-7 membered hetero ring containing one or two nitrogen means, for example, pyrrolyl, pyrrolidinyl, imidazolyl, imidazolidinyl, pyridinyl, piperidinyl, pyrazinyl, piperazinyl or pyrimidinyl. Explanation of various symbols in the formula (IB), is shown below.

[0041] In the formula (IB), C1-4 alkyl represented by R^0 , R^{100} , R^{200} , R^{300} , R^{400} , R^{50} , R^{60} , R^{70} , R^{80} , R^{90} , R^{120} and R^{130} , and that in R^0 , R^{100} , R^{200} , R^{300} , R^{400} , R^{50} , R^{60} , R^{70} , R^{80} , R^{90} , R^{120} and R^{130} , means methyl, ethyl, propyl, butyl and the isomers thereof.

[0042] In the formula (IB), C1-4 alkyl represented by R^0 and A_0 , and that in R^0 and A_0 means methoxy, ethoxy, propoxy, butoxy and the isomers thereof.

[0043] In the formula (IB), C1-4 alkylene represented by A_0 , and that in A_0 , means methylene, ethylene, trimethylene, tetramethylene and the isomers thereof.

[0044] In the formula (IB), C2-8 alkenylene represented by A_0 , and that in A_0 , means ethylene, trimethylene, tetramethylene, pentamethylene, hexamethylene, heptamethylene, octamethylene and the isomers thereof, having one, two or three double bonds.

[0045] In the formula (IB), C1-8 alkyl represented by R^{50} , R^{60} and R^{70} , and that in R^{50} , R^{60} and R^{70} , means methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl and the isomers thereof.

[0046] In the formula (IB), C2-8 alkenyl represented by R^{50} , R^{60} and R^{70} , and that in R^{50} , R^{60} and R^{70} , mean methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl and the isomers thereof, having one, two or three double bonds.

[0047] In the formula (IB), 4-7 cycloalkyl represented by R^{50} , R^{60} and R^{70} , and that in R^{50} , R^{60} , and R^{70} , mean cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.


[0048] In the formula (IB), examples of the 4-7 membered hetero ring containing one oxygen (which may be partially or fully saturated) represented by R^{50} , R^{60} , and R^{70} , and that in R^{50} , R^{60} , and R^{70} , are furyl, pyranyl, dihydrofuryl, dihydropyranyl, tetrahydrofuryl and tetrahydropyranyl.

[0049] In the formula (IB), examples of the 4-7 membered hetero ring containing one nitrogen (which may be partially or fully saturated) represented by R^{50} , R^{60} , and R^{70} , and that in R^{50} , R^{60} , and R^{70} , are pyrrolyl, pyridinyl, piperidinyl, pyrrolinyl, pyrrolidinyl and dihydropyridinyl.

[0050] In the formula (IB), when R is the formula represented by (vi), examples of the 4-10 membered hetero ring containing one or two nitrogen, (which may be partially or fully saturated) are pyrrolyl, pyridinyl, pyrrolinyl, pyrrolidinyl, dihydropyridinyl, imidazolyl, piperidinyl, imidazolyl, imidazolidinyl, pyrimidinyl, pyridazinyl, pyrazinyl, indolyl and tetrahydroindolyl.

[0051] Preferred compounds of the present invention of the formula (IB) are those described in the Examples and the following compounds.

Table 1



Preferable groups as R




Table 1 continued











Preferable groups as R			
			
			
			

Table 2



Preferable groups as R




Table 2 continued











Preferable groups as R			
			
			
			

Table 3

Preferable groups as R














Table 3 continued








Preferable groups as R	
	
	
	

Table 4

Preferable groups as R














Table 4 continued











Preferable groups as R			
			
			
			

Table 5



 Preferable groups as R














Table 5 continued











Preferable groups as R			
			
			
			

Table 6



Preferable groups as R













Table 6 continued











Preferable groups as R			
			
			
			

Table 7



Preferable groups as R













Table 7 continued











Preferable groups as R			
			
			
			

Table 8



 Preferable groups as R















Table 8 continued

Preferable groups as R			

Table 9



Preferable groups as R




Table 9 continued











Preferable groups as R			
			
			
			

Table 10



 Preferable groups as R














Table 10 continued











Preferable groups as R			
			
			
			

Table 11



 Preferable groups as R















Table 11 continued

Preferable groups as R			

Table 12



 Preferable groups as R




Table 12 continued











Preferable groups as R			
			
			
			

Table 13



 Preferable groups as R














Table 13 continued











Preferable groups as R			
			
			
			

Table 14



 Preferable groups as R














Table 14 continued











Preferable groups as R			
			
			
			

Table 15



Preferable groups as R




Table 15 continued











Preferable groups as R			
			
			
			

Table 16



Preferable groups as R















Table 16 continued



[0052] Pharmaceutical compositions of the present invention can be prepared using one active ingredient or two or more active ingredients.


[0053] The compounds of the formula (IA) and (IB), of the present invention may be converted into the corresponding salts and acid-addition salts by known methods. Nontoxic and water-soluble salts are preferred.

[0054] Suitable salts include the salts of alkali metals (sodium, potassium etc), alkaline-earth metal (calcium, magnesium etc.), ammonium salts, salts of pharmacologically acceptable organic amines (tetramethyl ammonium, triethylamine, methylamine, dimethylamine, cyclopentylamine, benzylamine, phenethylamine, piperidine, monoethanolamine, diethanolamine, tris (hydroxymethyl)aminomethane, lysine, arginine, N-methyl-D-glucane etc).


[0055] Suitable acid-addition salts include the salts with inorganic acids such as hydrochloric acid, and the salts with organic acids such as acetic acid, trifluoroacetic acid, lactic acid, tartaric acid, oxalic acid, fumaric acid, maleic acid, citric acid, benzoic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, toluenesulfonic acid, isethionic acid, glucuronic acid and gluconic acid. Preferred salts include the salts with acids such as hydrochloric acid, methanesulfonic, acetic acid and trifluoroacetic acid.

[0056] The compounds of the present invention, of the formula (IA), may be prepared by methods known per se, as hereinbefore defined (see the specification of EP-A-588655 and EP-A-656349).

[0057] The compounds of the present invention, of the formula (IB) may be prepared by the reaction to form an ester or amide bond of a compound of the formula (II):



(wherein the various symbols have the same meanings as hereinbefore defined) with a compound of the formula:



(wherein the various symbols have the same meanings as hereinbefore defined). The esterification and the reaction to form an amide are known and can be carried out by known methods for example:

- (1) using an acid halide,
- (2) using a mixed acid anhydride or
- (3) using a condensing agent.

[0058] Esterification can be carried out, for example, as follows:

(1) the method using an acid halide may be carried out, for example, by reacting a carboxylic acid with an acid halide (e. g., oxalyl chloride, thionyl chloride etc.) in an inert organic solvent (e.g., chloroform, methylene chloride, diethyl ether, tetrahydrofuran etc.) or without a solvent at from -20°C to the reflux temperature of the solvent, and then by reacting the acid halide obtained with a corresponding alcohol in the presence of a tertiary amine (e.g., pyridine, triethylamine, diethylaniline, diethylaminopyridine etc.) in an inert organic solvent (e. g., chloroform, methylene chloride, diethyl ether, tetrahydrofuran etc.) at a temperature of from 0°C to 40°C,

(2) the method using a mixed acid anhydride may be carried out, for example, by reacting a carboxylic acid and an acid halide (e. g., pivaloyl chloride, tosyl chloride, mesyl chloride etc.) or an acid derivative (e. g., ethyl chloroformate, isobutyl chloroformate etc.) in the presence of a tertiary amine (e.g. pyridine, triethylamine, dimethylaniline, dimethylaminopyridine etc.) in an inert organic solvent (e. g., chloroform, methylene chloride, diethyl ether, tetrahydrofuran etc.) or without a solvent at a temperature of from 0°C to 40°C, and then by reacting the mixture of acid anhydride obtained with a corresponding alcohol in an inert organic solvent (e. g., chloroform, methylene chloride, diethyl ether, tetrahydrofuran etc.), at a temperature of from 0°C to 40°C,

(3) the method using a condensing agent (e. g., 1,3-dicyclohexylcarbodiimide (DCC), 1-ethyl-3-[(dimethylamino) propyl] carbodiimide(EDC), 2-chloro-1-methylpyridinium iodide etc.) may be carried out, for example, by reacting a carboxylic acid with a corresponding alcohol using a condensing agent in the presence or absence of a tertiary amine (e.g. pyridine, triethylamine, dimethylaniline, dimethylaminopyridine etc.) in an inert organic solvent (e. g., chloroform, methylene chloride, dimethyl formamide, diethyl ether etc.) or without a solvent at a temperature of from 0°C to 40°C.




[0059] The reaction to form an amide may be carried out by the same reaction as hereinbefore defined, converting alcohol to a corresponding amine.

[0060] The reactions (1), (2) and (3) hereinbefore described may be preferably carried out in an atmosphere of inert gas (e. g., argon, nitrogen etc.) under anhydrous conditions.


[0061] The compounds of the formula (III) may be prepared by the series of reactions depicted in the following Scheme A.

Scheme (A)

when R is (i) or (ii),




MsOH or CF₃COOH
anisole
or
palladium carbon / H₂



(IIIa)


Scheme (A) continued

when R is (iii),







Scheme (A) continued

when R is (iv),




Scheme (A) continued

when R is (v),




MsOH or CF₃COOH
anisole
or
palladium carbon / H₂



(IIIId)

Scheme (A) continued

when R is (v),



In the Scheme A,

R^P is t-butyl or benzyloxycarbonyl,

X¹⁰, X²⁰ and X³⁰ each independently, is halogen,

Ms is methanesulfonic acid,

A₀₀ is bond, C1-3 alkylene, oxy-(C1-3) alkylene, thio-(C1-3)alkylene, C2-7 alkenylene, C2-7 alkenylene which is substituted by carboxy or C1-4 alkoxy carbonyl, and the other symbols have the same meaning as hereinbefore described.

[0062] The reactions in Schemes hereinbefore depicted may be carried out by methods known per se. The compounds of the formulae (II), (IV), (V) and (VI) used as starting materials in the Schemes hereinbefore depicted, are known per se or may be easily prepared by methods known per se.

[0063] Other starting materials and each reagents are known per se or may be prepared by known methods.

[0064] In each reaction in the present specification, products may be purified by conventional manner. For example, it may be carried out by distillation at atmospheric or reduced pressure, high performance liquid chromatography, thin layer chromatography or column chromatography using silica gel or magnesium silicate, washing or recrystallization. Purification may be carried out after each reaction, or after a series of reactions.

[Effect]

[0065] As mentioned above, it is considered that LTB₄ antagonist is useful for anti-inflammatory and anti-allergic agent.

[0066] Therefore, the compounds of the present invention of the formula (IA) and (IB), having LTB₄ antagonistic activity, may be used for the treatment of an animal, preferably a human, as anti-inflammatory and anti-allergic agent.

[0067] It is known that an LTB₄ antagonist is also useful for the treatment of e.g. rheumatoid arthritis, inflammatory bowel diseases, psoriasis, nonsteroidal anti-inflammatory agent-induced stomach diseases, adult respiratory distress syndrome, cardiac infarction, allergic rhinitis, hemodialysis-induced neutropenia, anaphase asthma in an animal, preferably a human; an LTB₄ antagonist may be used for their prevention and/or treatment.

[0068] The compounds of the formula (IB) also have inhibitory activity on phospholipase and inhibitory activity on trypsin, in an animal, preferably a human; therefore the compounds of the formula (IB) are useful for the prevention and/or the treatment of e.g. various inflammatory, allergic diseases, disseminated intravascular coagulation, pancreatitis, severity in pancreatitis and multiple organ failure.

[0069] It is confirmed that the toxicity of the active ingredients and non-toxic salts thereof and non-toxic acid addition salts thereof in the present invention is very weak. For example, LD₅₀ of Compound 1 was 117mg/kg when administered intravenously to male mice. Accordingly, the active substances in the present invention may be considered to be sufficiently safe and suitable for pharmaceutical use.

[0070] For the purpose hereinbefore described, the active ingredient in the present invention and non-toxic salts thereof and non-toxic acid addition salts thereof may be normally administered systemically or partially, usually by oral or parenteral administration.

[0071] The doses to be administered are determined depending upon age, weight, symptom, the desired therapeutic effect, the route of administration, and the duration of the treatment etc. In the human adult, the doses per person per dose are generally between 1 mg and 1000mg, by oral administration, up to several times per day, or between 100μg and 100mg, by parenteral administration (preferably, intravenously) up to several times per day. As mentioned above, the doses to be used depend upon various conditions. Therefore, there are cases in which doses lower than or greater than the ranges specified above may be used.

[0072] When administering of the compounds of the present invention, it is used in the form of solid compositions, liquid compositions or other compositions for oral administration, as injections, liniments or suppositories etc. for parenteral administration.

[0073] Solid compositions for oral administration include compressed tablets, pills, capsules, dispersible powders, and granules.

[0074] In such compositions, one or more of the active compound(s) is or are admixed with at least one inert diluent (such as lactose, mannitol, glucose, hydroxypropyl cellulose, microcrystalline cellulose, starch, polyvinylpyrrolidone, magnesium metasilicate aluminate etc.).

[0075] These compositions may also comprise, as in normal practice, additional substances other than inert diluents: e. g. lubricating agents (such as magnesium stearate, etc.), disintegrating agents (such as cellulose calcium glycolate, etc.), assisting agents for dissolving (such as arginine, glutamic acid, asparaginic acid, etc.) and stabilizer (human serum albumin, lactose, etc.).

[0076] The tablets or pills may, if desired, be coated with a film of gastric or enteric material (such as sugar, gelatin, hydroxypropyl cellulose, hydroxypropylmethyl cellulose phthalate etc.).

[0077] Capsules include hard capsules and soft capsules.

[0078] Liquid compositions for oral administration include solutions, emulsions, suspensions, syrups and elixirs.

[0079] These liquid compositions may comprise inert diluents commonly used in the art (purified water, ethanol etc.).

[0080] Besides inert diluents, such compositions may also comprise adjuvants (such as wetting agents, suspending agents, etc.), sweetening agents, flavoring agents and preserving agents.

[0081] Other compositions for oral administration include spray compositions which may be prepared by known methods and which comprise one or more of the active compound(s). Spray compositions may comprise additional substances other than inert diluents: e. g. stabilizing agents (sodium sulfate etc.), isotonic stabilizing agents (sodium

chloride, sodium citrate, citric acid, etc.). For preparation of such spray compositions, for example, the method described in the United States Patent No. 2,868,691 or 3,095,355 may be used.

[0082] Injections for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions and emulsions.

[0083] In such compositions, one or more of active compound(s) is or are admixed with at least one of inert aqueous diluent(s) (distilled water for injection, physiological salt solution etc.) or inert non-aqueous diluent(s) (propylene glycol, polyethylene glycol, olive oil, ethanol, POLYSORBATE80 (registered trade mark) etc.).

[0084] Injections may comprise furthermore assisting agents such as preserving agents, wetting agents, emulsifying agents, dispersing agents, stabilizing agents (such as human serum albumine, lactose, etc.) and assisting agents for dissolving (arginine, glutamic acid, asparaginic acid, polyvinylpyrrolidone etc.).

[0085] Usually, they may be sterilized by filtration (a bacteria-retaining filter, etc), by incorporation of sterilizing agents in the compositions or by irradiation, or after treated, they may also manufactured in the form of sterile solid compositions, for example, by freeze-drying, and which may be dissolved in sterile water or some other sterile diluent(s) for injection immediately before used, and which may be used.

[Example]

[0086] The following reference Examples and Examples illustrate the present invention.

[0087] The solvents in parentheses show the developing or eluting solvents and the ratios of the solvents used are by volume in chromatographic separations.

Example 1 (A): Binding inhibition against $^3\text{H-LTB}_4$ on the human polymorphonuclear leukocyte (PMN)

[0088] Polypropylene tubes were added 0.049 ml Hanks balanced salt solution (HBSS), 0.001 ml test compound and 0.05 ml $^3\text{H-LTB}_4$ (4nM), and mixed. The reaction was started by addition of thoroughly mixed cell suspension (1.6×10^6 cells), followed by incubation at 0°C for 20 min. The reaction was terminated by the addition of ice-cold HBSS (2.5 ml). PMNs were harvested by vacuum filtration through Whatman GF/C glass fiber filters on Brandel cell harvester (BRANDEL, M-24R). The filters were then washed 2 times to remove free $^3\text{H-LTB}_4$ with 2.5 ml of the ice-cold PBS (-) solution. The filters were transferred to each vial, and equilibrated after adding 8 ml ACS II cocktail (Amersham). The radioactivity was measured by liquid scintillation counter (Aloka, LSC-5100).

[0089] Specific binding of $^3\text{H-LTB}_4$ to receptor was defined as total binding subtracting nonspecific binding. Nonspecific binding was determined as binding in the presence of $1.5\mu\text{M}$ LTB_4 instead of test compound.

The inhibitory effect of test compound was calculated from the following equation.

[0090] The percentage of inhibition (%) = $100 - (B_1/B_0 \times 100)$

B_1 : Specific $^3\text{H-LTB}_4$ binding in presence of test compound

B_0 : Specific $^3\text{H-LTB}_4$ binding in absence of test compound

[Results]

[0091] The results are shown in the following Table 17.

Table 17


Compound No.	European Patent Publication No. 588655 Compound (Example No.)	binding activity (%)
1	1 (i)	91.5
2	1 (m)	76.6
3	1 (p)	75.0
4	1 (aa)	63.7
5	1 (ii)	94.3
6	1 (pp)	71.6
7	1 (qq)	78.0
8	1 (hhh)	82.7
9	1 (III)	91.6
10	1 (mmm)	86.5
11	2 (g)	76.8

Table 17 (continued)


Compound No.	European Patent Publication No. 588655 Compound (Example No.)	binding activity (%)
12	2 (p)	95.2
13	2 (u)	100.2
14	2 (w)	96.5
15	2 (cc)	89.1
16	2 (gg)	83.6
17	2 (kk)	93.9
18	3 (f)	87.0
19	4	74.0
20	4 (a)	83.5
21	5 (r)	90.8
22	5 (w)	89.7
23	5 (ff)	78.0
24	European Patent Publication No. 656349 Example 1 (b)	61.2
25		85.5
26		92.8
27		94.4
28		87.3

The structure of compounds used in the present invention are shown below.


Compound No. 1




Compound No. 2




Compound No.3




Compound No.4




Compound No.5




Compound No.6




Compound No.7




Compound No.8




Compound No.9




Compound No.10




Compound No.11




Compound No.12




Compound No.13




Compound No.14




Compound No.15




Compound No.16




Compound No.17




Compound No.18




Compound No.19




Compound No.20




Compound No.21




Compound No.22




Compound No.23




Compound No.24




Compound No. 25




Compound No. 26



Compound No. 27



Compound No. 28



Example 1(B):

[0092] The compounds of the formula (IB), of the present invention have an antagonistic activity on LTB₄. The results which are measured by method as hereinbefore described in Example 1(A), are shown the following Table 18.

Table 18

Compound (Example No.)	binding activity (%)
2	79.7
2(a)	92.0
2(b)	97.9
2(c)	103.2
2(d)	99.3
2(e)	94.5
2(f)	91.8
2(g)	89.6
2(h)	85.4
2(i)	69.6
2(j)	55.4
2(k)	97.7
2(l)	81.0
2(m)	89.2
2(n)	82.8
2(o)	85.8
2(p)	95.2
2(q)	98.0
2(r)	80.1

Table 18 continued

Compound (Example No.)	binding activity (%)
2(s)	83.0
2(t)	51.5
2(u)	67.6
2(v)	92.0
2(w)	76.7
2(x)	94.1
2(cc)	76.7
2(dd)	50.8
2(ee)	65.3
2(ff)	82.4
3	96.8
4	73.1
4(a)	52.0
5	89.7
5(a)	62.5
5(b)	90.2
6	67.8

Example 1 (C) : inhibitory activity on phospholipase A₂ and on trypsin

[0093] It has been confirmed that the compounds of the formula (IB), of the present invention have inhibitory activities on phospholipase_{A2} (PLA₂) and on trypsin.

[0094] For example, in laboratory tests the following results were obtained.

[Method]

[0095]

(1) Inhibitory activity on PLA₂

A reaction solution including 50 mM tris-HCl buffer (pH7.5, 874μl; containing 100mM sodium chloride, 1mM EDTA), 1M calciumchloride (6μl), 1% bovine serum albumin (10μl) and 2.5mM 10PY-PC (10μl), was prepared. To the solution were added a test compound in various concentration or water (50μl), and a solution of 10mU/ml PLA₂ (derived from hog pancreas) (50μl). The appearance of fluorescence was measured (Ex=345 nm, Em=396 nm). Percentage (%) of the strength of fluorescence in the presence of a test compound was calculated when the strength of that in the absence thereof was regarded as 100%, and therefrom IC₅₀ value was calculated. The results are shown in the following Table19.

(2) Inhibitory activity on trypsin

To a mixture of a 0.2 M HEPES • sodium hydroxide buffer solution (pH 8.0, 100μl) and distilled water (640μl), were added a test compound in various concentration or water (10μl), and a solution of 80 mU/ml trypsin (derived from bovine pancreas) (50μl) and then the mixture was preincubated for one minute at 30°C. To the solution thus obtained was added 2.5mM BAPNA (200μl) and the mixture was incubated at 30°C. The absorbance at 405 nm was measured. Percentage (%) of the absorbance in the presence of a test compound was calculated when the absorbance in the absence thereof was regarded as 100%, and therefrom IC₅₀ value was calculated. The results are shown in the following Table 19.

Table 19

Compound (Example No.)	inhibitory activity on PLA ₂ IC ₅₀ (μM)	inhibitory activity on trypsin IC ₅₀ (μM)
2	--	0.19
2(a)	2.6	0.4
2(b)	3.8	0.56
2(c)	8.1	0.26

Table 19 (continued)

Compound (Example No.)	inhibitory activity on PLA ₂ IC ₅₀ (μM)	inhibitory activity on trypsin IC ₅₀ (μM)
2(d)	8.7	0.14
2(e)	8.5	0.34
2(f)	70	0.10
2(g)	53	0.16
2(h)	11	0.15
2(i)	59	0.14
2(j)	--	0.12
2(k)	20	0.10
2(l)	94	0.12
2(m)	18	0.17
2(n)	10	0.16
2(o)	12	0.14
2(p)	29	0.13
2(q)	34	0.16
2(r)	46	0.16
2(s)	44	0.16
3	4.7	0.12
4	41	0.16
4(a)	--	0.14
5	--	0.13
5(a)	--	0.15
6	4.5	0.17

In the methods hereinbefore described,

10PY-PC represents 3'-palmitoyl-2-(1-pyrenedecanoyl)-L-α-phosphatidylcholine,

HEPES represents 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid, and


BAPNA represents α-N-benzoyl-DL-arginine-p-nitroanilide hydrochloride.

[Preparation of new compounds]

[0096] The following reference Examples and Examples illustrate new compounds of the formula (IB).

Reference Example 1

[0097] N-(2-Propenyl)-N-ethoxycarbonylmethyl-4-benzyloxycarbonylphenoxyacetamide.




[0098] A solution of 4-benzyloxycarbonylphenoxyacetic acid (4.29g) in thionyl chloride (10ml) was refluxed for 15min. After an excess amount of solvent was distilled off, product was dissolved in dichloromethane. And this solution was added dropwise to a solution of N-(2-propenyl)-N-ethoxycarbonylmethylamine (2.14g) in pyridine under cooling with ice. After the solution was stirred for 30min at room temperature, the solution was poured into ice water. The mixture was extracted with ethyl acetate. The extract was washed with a solution of 1N hydrochloric acid, water and a saturated aqueous solution of sodium chloride, successively, evaporated. The residue purified by silica gel column chromatography

to give the title compound (5.96g) having the following physical data:

TLC : Rf 0.43 (hexane : ethyl acetate=3 : 2)

Reference Example 2

[0099] N-(2-Propenyl)-N-ethoxycarbonylmethyl-4-carboxyphenoxyacetamide




[0100] Methanesulfonic acid (28ml) was added to the compound prepared in Reference Example 1 (5.69g) under cooling at 0°C. After reaction solution was stirred for one hour at room temperature, poured into ice water and extracted with ethyl acetate. Organic layer was washed with water, and a saturated aqueous solution of sodium chloride, successively, evaporated. The residue was purified by silica gel column chromatography to give the title compound (4.31g) having the following physical data.

TLC : Rf 0.35 (hexane : ethyl acetate=1 : 1)

Example 2

[0101] N-(2-propenyl)-N-ethoxycarbonylmethyl-4-(4-amidinophenoxycarbonyl)phenoxyacetamide acetate



[0102] To a solution of amidinophenol (1.72g) and the compound prepared in Reference Example 2 (3.21g) in pyridine was added DCC (3.09g) and stirred overnight at room temperature. Reaction solution was filtered and the filtrate was evaporated. The residue was purified by silica gel column chromatography and was formed into acetate by conventional manner to obtain the title compound having the following physical data.

TLC : Rf 0.41 (chloroform : methanol : acetic acid = 10 : 2 : 1),


NMR (CD₃OD) : δ 8.14(2H, d, J=9.0Hz), 7.90(2H, d, J=9.0Hz), 7.49(2H, d, J=9.0Hz), 7.08(2H, d, J=9.0Hz), 5.68-6.07 (1H, m), 5.17-5.37(2H, m), 4.93 and 5.02(2H, s, ratio=7 : 10), 4.03-4.28(6H, m), 1.26 and 1.29(3H, t, J=7.0Hz).

Example 2 (a)~2 (ff)

[0103] By the same procedure as Reference Example 1, 2 and Example 2, the compound having the following physical data was given.

Example 2 (a)


[0104]



TLC : Rf 0.57 (chloroform : methanol : acetic acid = 10 : 2 : 1),
NMR (CD₃OD) : δ 2.60(2H, t, J=8.0Hz), 2.98(2H, t, J=8.0Hz), 5.17(2H, s), 6.99-7.02(2H, m), 7.09-7.16(5H, m), 7.30(5H, s), 7.38(1H, d, J=9.0Hz), 7.43(1H, s), 7.48(2H, d, J=8.0Hz), 7.98(2H, d, J=8.0Hz).

Example 2 (b)


[0105]



TLC : Rf 0.60 (chloroform : methanol : acetic acid = 10 : 2 : 1),
NMR (CD₃OD) : δ 2.41 (2H, t, J=7.0Hz), 3.00(2H, t, J=7.0Hz), 4.69(2H, s), 5.23(2H, s), 7.09-7.42(14H, m), 7.43(2H, d, J=8.0Hz), 7.98(2H, d, J=8.0Hz).

Example 2 (c)

[0106]




TLC : Rf 0.53 (chloroform : methanol : acetic acid = 10 : 2 : 1),
NMR (CD₃OD) : δ 8.0(2H, d, J=8.0Hz), 7.50(2H, d, J=8.0Hz), 7.46(1H, s), 7.40(1H, d, J=8.0Hz), 7.24(5H, s), 7.12(1H, s), 7.10(1H, d, J=8.0Hz), 4.61(2H, s), 4.22(2H, q, J=8.0Hz), 3.00(2H, t, J=9.0Hz), 2.61(2H, t, J=9.0Hz), 1.30(3H, t, J=8.0Hz).

Example 2 (d)

[0107]

5

10



• AcOH

15

TLC : Rf 0.45 (chloroform : methanol : acetic acid = 10 : 2 : 1),
 NMR (CD₃OD) : δ 8.00(2H, d, J=8Hz), 7.80(1H, d, J=16Hz), 7.75(2H, d, J=8Hz), 7.50(2H, d, J=8Hz), 7.35(2H, d, J=8Hz), 7.30-7.20(5H, m), 6.70(1H, d, J=16Hz), 4.65(2H, s), 4.25(2H, q, J=7Hz), 1.30(3H, t, J=7Hz).


Example 2 (e)

20

[0108]

25

30



• AcOH

35

TLC : Rf 0.45 (chloroform : methanol : acetic acid = 10 : 2 : 1),
 NMR (CD₃OD) : δ 1.30(3H, t, J=7.0Hz), 2.18(3H, s), 4.31(2H, q, J=7.0Hz), 4.77(2H, m), 5.02(1H, t, J=4.0Hz), 7.39-7.61 (8H, m), 7.89(2H, d, J=9.0Hz), 8.02(2H, d, J=9.0Hz), 8.22(2H, d, J=9.0Hz).


Example 2 (f)

40

[0109]

45

50




• HCl

TLC : Rf 0.43 (chloroform : methanol : acetic acid = 10 : 2 : 1),
 NMR (CD₃OD) : δ 8.00-7.80(7H, m), 7.50(2H, d, J=8.5Hz), 6.90(1H, d, J=16Hz), 4.60(1H, dd, J=4.5, 4.5Hz), 4.20(2H, q, J=6.5Hz), 4.15(2H, q, J=6.5Hz), 2.50(2H, t, J=7.5Hz), 2.30(1H, m), 2.10(1H, m), 1.30(3H, t, J=6.5Hz), 1.25(3H, t, J=6.5Hz).

Example 2 (g)

[0110]




• HCl

TLC : Rf 0.46 (chloroform : methanol : acetic acid = 10 : 2 : 1),

NMR (CD₃OD) : δ 8.00-7.90(5H, m), 7.65(2H, d, J=8Hz), 7.50(2H, d, J=8Hz), 4.65(1H, dd, J=4.5, 4.5Hz), 4.20(2H, q, J=6.5Hz), 4.15(2H, q, J=6.5Hz), 2.50(2H, t, J=7.5Hz), 2.30(1H, m), 2.25(3H, m), 2.10(1H, m), 1.30(3H, t, J=6.5Hz), 1.25(3H, t, J=6.5Hz).

Example 2 (h)

[0111]




• HCl

TLC : Rf 0.48 (chloroform : methanol : acetic acid = 15 : 2 : 1),

NMR (CD₃OD) : δ 8.24(2H, d, J=8.5Hz), 7.95(2H, d, J=8.5Hz), 7.62(2H, d, J=8.0Hz), 7.55(2H, d, J=8.0Hz), 7.35(1H, s), 6.85(1H, dt, J=7.5, 15.0Hz), 5.93(1H, d, J=15.0Hz), 4.28(4H, q, J=7.5Hz), 4.18(2H, d, J=7.5Hz), 3.23(2H, d, J=7.5Hz), 2.14(3H, s), 1.26(6H, t, J=7.5Hz), 1.23(3H, t, J=7.5Hz).

Example 2 (i)

[0112]



• AcOH


TLC : Rf 0.43 (chloroform : methanol : acetic acid = 10 : 2 : 1),

EP 0 703 216 B1

NMR (CD₃OD) : δ 8.24 and 8.26(2H, d, J=9.0Hz), 7.81(1H, d, J=18.0Hz), 7.75(2H, d, J=9.0Hz), 7.58 and 7.66(2H, d, J=9.0Hz), 7.37(2H, d, J=9.0Hz), 6.73(1H, d, J=18.0Hz), 5.77-5.96(1H, m), 5.22-5.34(2H, m), 4.12-4.28(4H, m), 3.96-4.00(2H, m), 1.20 and 1.30(3H, t, J=7.0Hz).

5 Example 2 (j)

[0113]




TLC : R_f 0.44 (chloroform : methanol : acetic acid = 10 : 2 : 1),

NMR (CD₃OD) : δ 8.18(2H, d, J=9.0Hz), 7.90(2H, d, J=9.0Hz), 7.50(2H, d, J=9.0Hz), 7.17(2H, d, J=9.0Hz), 4.70(2H, s), 4.55(1H, dd, J=9.5, 5.0Hz), 4.18(2H, q, J=7.0Hz), 4.11(2H, q, J=7.0Hz), 2.40(2H, t, J=7.0Hz), 1.97-2.32(2H, m), 1.27(3H, t, J=7.0Hz), 1.23(3H, t, J=7.0Hz).

Example 2 (k)

[0114]




TLC : R_f 0.48 (chloroform : methanol : acetic acid = 15 : 2 : 1),

NMR (CD₃OD) : δ 8.22(2H, d, J=8.0Hz), 7.92(2H, d, J=8.0Hz), 7.60(2H, d, J=8.0Hz), 7.56(2H, d, J=8.0Hz), 7.37(1H, brs), 4.27(4H, q, J=7.5Hz), 4.13(2H, q, J=7.5Hz), 3.47(2H, s), 2.16(3H, s), 1.25(6H, t, J=7.5Hz), 1.22(3H, t, J=7.5Hz).

Example 2 (l)

[0115]



TLC : R_f 0.49 (chloroform : methanol : acetic acid = 10 : 2 : 1),


NMR (CD₃OD) : δ 7.98(1H, s), 7.90(2H, d, J=9.0Hz), 7.58(4H, m), 7.48(2H, d, J=9.0Hz), 5.78-5.96(1H, m), 5.23-5.32

EP 0 703 216 B1

(2H, m), 4.22(2H, q, J=7.0Hz), 4.20(2H, s), 3.98-4.03(2H, m), 2.24(3H, s), 1.30(3H, t, J=7.0Hz).

Example 2 (m)


[0116]



TLC : R_f 0.38 (chloroform : methanol : acetic acid = 10 : 1 : 1),
NMR (CD₃OD) : δ 8.20(2H, d, J=8.4Hz), 7.92(2H, d, J=8.8Hz), 7.74(2H, d, J=8.4Hz), 7.55(2H, d, J=8.8Hz), 7.25(3H, m), 6.32(1H, d, J=14.6Hz), 4.55(1H, m), 4.20(2H, q, J=7.2Hz), 4.14(2H, q, J=7.0Hz), 2.72(3H, s), 2.45(2H, t, J=7.4Hz), 2.36-1.90(2H, m), 1.29(3H, t, J=7.2Hz), 1.25(3H, t, J=7.0Hz).

Example 2 (n)


[0117]



TLC : R_f 0.39 (chloroform : methanol : acetic acid = 10 : 1 : 1),
NMR (CD₃OD) : δ 8.18(2H, d, J=8.4Hz), 7.92(2H, d, J=8.8Hz), 7.73(2H, d, J=8.4Hz), 7.53(2H, d, J=8.8Hz), 7.50-7.15 (2H, m), 7.05(1H, d, J=14.5Hz), 6.75-6.55(1H, m), 6.03-5.81(1H, m), 5.32-5.14(2H, m), 4.20(2H, q, J=7.2Hz), 4.30-4.10 (4H, m), 1.94(3H, s), 1.28(3H, t, J=7.2Hz).

Example 2 (o)

[0118]




EP 0 703 216 B1

TLC : Rf 0.50 (chloroform : methanol : acetic acid = 10 : 2 : 1),

NMR (CD₃OD) : δ 8.20(2H, d, J=8.5Hz), 7.90(2H, d, J=11.5Hz), 7.60(2H, d, J=8.5Hz), 7.55(2H, d, J=11.5Hz), 7.35(1H, br.s), 5.70(1H, m), 5.15(2H, m), 4.25(4H, q, J=7Hz), 3.10(2H, d, J=7Hz), 2.15(3H, s), 1.95(3H, s), 1.25(6H, t, J=7Hz) .

5 Example 2 (p)

[0119]




• MsOH

TLC : Rf 0.50 (chloroform : methanol : acetic acid = 10 : 1 : 1),

NMR (CD₃OD) : δ 7.94(2H, d, J=8.0Hz), 7.89(2H, d, J=8.5Hz), 7.72(2H, d, J=8.5Hz), 7.44(2H, d, J=8.0Hz), 6.49(1H, s), 4.64(1H, m), 4.23(2H, q, J=7.5Hz), 4.14(2H, q, J=7.0Hz), 2.74(3H, s), 2.66(3H, s), 2.52(2H, t, J=7.0Hz), 2.32(2H, m), 2.14(2H, m), 1.30(3H, t, J=7.0Hz), 1.25(3H, t, J=7.5Hz).

Example 2 (q)

[0120]




• AcOH

TLC : Rf 0.50 (chloroform : methanol : acetic acid = 10 : 1 : 1),

NMR (CD₃OD) : δ 7.89(2H, d, J=8.8Hz), 7.73(2H, d, J=8.4Hz), 7.56(2H, d, J=8.4Hz), 7.44(2H, d, J=8.8Hz), 6.49(1H, s), 5.88(1H, m), 5.35-5.20(2H, m), 4.30-4.10(4H, m), 4.00(2H, m), 2.65(3H, s), 1.93(3H, s), 1.31(3H, t, J=7.2Hz).

Example 2 (r)

[0121]



• AcOH

TLC : Rf 0.46 (chloroform : methanol : acetic acid = 10 : 2 : 1),

NMR (CD₃OD) : δ 8.18(2H, d, J=9.0Hz), 7.93(2H, d, J=9.0Hz), 7.82(2H, d, J=9.0Hz), 7.80(1H, s), 7.52(2H, d, J=9.0Hz), 4.66(1H, dd, J=8.5Hz, 4.0Hz), 4.33(2H, q, J=7.0Hz), 4.20(2H, q, J=7.0Hz), 4.12(2H, q, J=7.0Hz), 2.39(2H, t, J=7.0Hz),

5

[0122]



20

[0123]



40

[0124]




55

EP 0 703 216 B1

NMR (CD₃OD) : δ 8.18(2H, d, J=9.0Hz), 7.93(2H, d, J=9.0Hz), 7.82(2H, d, J=9.0Hz), 7.80(1H, s), 7.52(2H, d, J=9.0Hz), 4.66(1H, dd, J=8.5Hz,4.0Hz), 4.33(2H, q, J=7.0Hz), 4.20(2H, q, J=7.0Hz), 4.12(2H, q, J=7.0Hz), 2.39(2H, t, J=7.0Hz), 2.11-2.31(1H, m), 1.82-2.00(1H, m), 1.36(3H, t, J=7.0Hz), 1.24(3H, t, J=7.0Hz), 1.21(3H, t, J=7.0Hz).

5 Example 2 (v)


[0125]



TLC : R_f 0.22 (chloroform : methanol : acetic acid = 10 : 2 : 1),
NMR (CD₃OD) : δ 8.21 (2H, d, J=8.0Hz), 7.95(2H, d, J=8.0Hz), 7.89(2H, d, J=8.0Hz), 7.59(2H, d, J=8.0Hz), 7.55(2H, d, J=8.0Hz), 7.43(2H, d, J=8.0Hz), 6.78(1H, s), 6.15-5.80(1H, m), 5.47-5.28(2H, m), 4.42(2H, s), 4.25(2H, d, J=5.0Hz), 2.68(3H, s, CH₃SO₃H), 2.18(3H, s).

Example 2 (w)


[0126]



TLC : R_f 0.27 (chloroform : methanol : acetic acid = 10 : 2 : 1),
NMR (CD₃OD) : δ 8.20(2H, d, J=8Hz), 7.91(2H, d, J=8Hz), 7.57(2H, d, J=8Hz), 7.53(2H, d, J=8Hz), 6.73(1H, s), 5.8-6.0 (1H, br), 5.2-5.35(2H, m), 4.8-4.9(1H, m), 4.0-4.3(8H, m), 2.12(3H, s), 1.91(3H, s), 1.27(6H, t, J=7Hz).

Example 2 (x)

[0127]



EP 0 703 216 B1

TLC : Rf 0.25 (chloroform : methanol : acetic acid = 10 : 2 : 1),


NMR (CD₃OD) : δ 8.22(2H, d, J=8Hz), 7.91(2H, d, J=8Hz), 7.52 and 7.67(4H, d, J=8Hz, rotamer), 6.65 and 6.78(1H, s, rotamer), 5.6-6.0(3H, br), 5.0-5.3(6H, m), 3.9-4.4(8H, m), 2.11 and 2.16(3H, s, rotamer), 1.92(3H, s).

5 Example 2 (cc)

[0128]

10

15



• MsOH

20 TLC : Rf 0.30 (chloroform : methanol : acetic acid = 10 : 2 : 1),


NMR (CD₃OD) : δ 8.21(2H, d, J=8Hz), 7.92(2H, d, J=8Hz), 7.60-7.45(4H, m), 6.73 and 6.65(1H, s, rotamer), 4.5-4.3(1H, m), 4.3-4.0(2H, br), 4.0-3.7(3H, m), 3.7-3.5(1H, br), 2.70(3H, s), 2.17 and 2.10(3H, s, rotamer), 2.2-1.8(3H, m), 1.8-1.4(1H, m).

25 Example 2 (dd)

[0129]

30

35



• 2CF₃COOH

40

TLC : Rf 0.10 (ethyl acetate : acetic acid : H₂O = 3 : 1 : 1),

NMR (CD₃OD) : δ 8.22(2H, d, J=8Hz), 7.92(2H, d, J=8Hz), 7.7-7.4(4H, m), 6.70(1H, s), 4.5-4.0(3H, br), 3.6-3.4(2H, m), 3.2-3.0(2H, m), 2.3-1.9(7H, br).

45


50

55

Example 2 (ee)

[0130]

5




15 TLC : Rf 0.43 (chloroform : methanol : acetic acid = 3 : 1 : 1),
NMR (CD₃OD) : δ 9.20(1H, br. s), 8.70(1H, br. s), 8.05-7.95(4H, m), 7.85(2H, d, J=9Hz), 7.75(2H, J=8Hz), 6.75(1H, m), 5.95(1H, m), 5.30(2H, m), 4.20(4H, m), 2.75(3H, s, CH₃SO₃H), 2.20(3H, s).

Example 2 (ff)

20


[0131]



TLC : Rf 0.40 (chloroform : methanol : acetic acid = 10 : 2 : 1),
NMR (CDCl₃) : δ 8.02(1H, d, J=9Hz), 7.90(1H, d, J=9Hz), 7.64(1H, s), 7.50(1H, d, J=9Hz), 7.40-7.00(14H, m), 6.95-6.80(2H, m), 6.80-6.72(1H, m), 6.48(1H, d, J=9Hz), 4.00-3.80(1H, m), 3.88(3H, s), 3.70-3.30,(2H, m), 3.10-2.90(1H, m), 2.90-2.70(2H, m), 2.70-2.30(2H, m), 2.30-2.00(2H, m), 1.00-1.24(1H, m).

Reference Example 3

45 **[0132]** 2-(N-Benzyl-N-methylamino)-2-(4-t-butoxycarbonylphenylmethylimino)acetic acid ethyl ester.




To a solution of 2-(N-benzyl-N-methylamino)-2-thioxoacetic acid ethyl ester (4.98g) in dichloromethane under cooling

with ice, was added dropwise $\text{BF}_4 \cdot \text{Et}_2\text{O}$ (72ml). Reaction solution was stirred for 30min at room temperature, extracted with dichloromethane. The extract was evaporated. The residue was purified by silica gel column chromatography to give the title compound having the following the physical data.

TLC : Rf 0.45 (hexane : ethyl acetate = 3 : 1).

Reference Example 4

[0133] 2-(N-Benzyl-N-methylamino)-2-(4-carboxyphenylmethylimino)acetic acid ethyl ester




To a solution of the compound prepared in Reference Example 3 (3.77g) in anisole (10ml) under cooling with ice bath, was added trifluoroacetic acid (20ml) and stirred for two hours at room temperature. Reaction solution was evaporated, neutralized by adding 1N aqueous solution of sodium hydroxide and extracted with ethyl acetate. The extract was evaporated. The residue was purified by silica gel column chromatography to give the title compound (1.87g) having the following physical data.

TLC : Rf 0.36 (hexane : ethyl acetate = 1 : 2).

Example 3

[0134] 2-[4-(4-Amidinophenoxy)phenylmethylimino]-2-(N-benzyl-N-methylamino)acetic acid ethyl ester hydrochloride




[0135] By the same procedure as Example 2, the title compound having the following physical data was given.

TLC : Rf 0.34 (chloroform : methanol : acetic acid = 10 : 2 : 1),

NMR (CD_3OD) : δ 1.26(3H, t, $J=7.0\text{Hz}$), 2.88(3H, s), 4.36(2H, q, $J=7.0\text{Hz}$), 4.49(2H, s), 4.50(2H, s), 7.27-7.35(5H, m), 7.48(2H, d, $J=9.0\text{Hz}$), 7.52(2H, d, $J=9.0\text{Hz}$), 7.92(2H, d, $J=9.0\text{Hz}$), 8.12(2H, d, $J=9.0\text{Hz}$).

Reference Example 5


[0136] Ethyl 1-(3-phenylpropyl)-1-(4-benzoyloxycarbonylphenylmethyl)phosphinate.



[0137] A solution of ethyl phenylpropylphosphinate (1.2g) and triethylamine (2.4ml) in chloroform (30ml) was cooled to 0°C, and a solution of trimethylsilylchloride (1.46ml) and 4-bromomethylbenzoic acid benzyl ester (1.75g) in chloroform (10ml) was added thereof, and stirred at room temperature for 1.5 day. To the reaction mixture was added ice water and extracted with ethyl acetate. Organic layer was washed with water and a saturated aqueous solution of sodium chloride, successively evaporated. The residue was purified by silica gel column chromatography to give the title compound (900mg).

Reference Example 6


[0138] Ethyl 1-(3-phenylpropyl)-1-(4-carboxyphenylmethyl)phosphinate



[0139] A mixture of the compound prepared in Reference Example 5 (900mg), palladium carbon (180mg, 10%) and ethanol (20ml) was stirred for two hours under an atmosphere of hydrogen at room temperature. Reaction mixture was filtered. The filtrate was evaporated and the title compound (815mg) was given.

Example 4

[0140] Ethyl 1-(4-amidinophenoxyacetylphenylmethyl)-1-(3-phenylpropyl)phosphinate acetate




[0141] By the same procedure as Reference Example 5,6 and Example 2, the title compound (805mg) having the following physical data was given.

TLC : Rf 0.62 (chloroform : methanol : acetic acid = 10 : 2 : 1),

NMR (CD₃OD) : δ 8.10(2H, d, J=8Hz), 7.95(2H, d, J=9Hz), 7.55(2H, d, J=9Hz), 7.60-7.40(2H, m), 7.30-7.10(3H, m), 7.20(2H, d, J=8Hz), 4.00(2H, m), 3.40(2H, d, J=24Hz), 2.70(2H, t, J=6.5Hz), 2.00-1.60(4H, m), 1.30(3H, t, J=7.5Hz).

Example 4(a)

[0142]




[0143] By the same procedure as Example 4, the compound having the following physical data was given.

TLC : R_f 0.60 (chloroform : methanol : acetic acid = 10 : 2 : 1),

NMR (CD₃OD) : δ 1.36(6H, t, J=7.0Hz), 4.15(4H, quin, J=7.0Hz), 6.68(1H, t, J=18.0Hz), 7.54(2H, d, J=9.0Hz), 7.56(1H, dd, J=23.0Hz, 18.0Hz), 7.82(2H, d, J=9.0Hz), 7.93(2H, d, J=9.0Hz), 8.22(2H, d, J=9.0Hz).

Reference Example 7


[0144] 4-Phenylpiperidine-1-ylmethylbenzoic acid methyl ester



[0145] A solution of 4-formylbenzoic acid (3.5g) and 4-phenylpiperidine (6.9g) in methanol (35ml) was stirred for one hour at room temperature. After the solution was cooled with ice bath, sodium borohydride (1.63g) was added thereto and reaction solution was stirred. After reaction finished, reaction solution was poured into ice water and extracted with ethyl acetate. The organic layer was washed with water and saturated aqueous solution of sodium chloride, successively, dried over and evaporated. The residue was washed with methanol to give the title compound (4.70g).

Reference Example 8


[0146] 4-(4-Phenylpiperidine-1-ylmethyl)benzoic acid



[0147] A solution of the compound prepared in Reference Example 7 (4.8g) in dioxane (50ml) was cooled with ice bath and 2N aqueous solution of sodium hydroxide (10ml) was added thereof and stirred at 60°C for two hours. The reaction mixture was cooled with ice bath and neutralized by adding 2N hydrochloric acid. Depositing solid was filtered and washed with water, ether successively, dried over. The title compound (4.29g) was given.

Example 5

[0148] 4-(4-Phenylpiperidine-1-ylmethyl)benzoic acid amidinophenol ester 2 hydrochloride



[0149] By the same procedure as Example 2, the title compound having the following physical data was given.

TLC : Rf 0.33 (chloroform : methanol : acetic acid = 5 : 1 : 1),


NMR (CD₃OD) : δ 8.32(2H, d, J=8.0Hz), 7.95(2H, d, J=8.8Hz), 7.88(2H, d, J=8.0Hz), 7.55(2H, d, J=8.8Hz), 7.28(5H, m), 4.52(2H, s), 3.62(2H, br.d), 3.25(2H, br.d), 2.94(1H, m), 2.12(4H, m).

Example 5(a)-5(b)

[0150] By the same procedure as Reference Example 7,8 and Example 5, the compound having the following physical data was given.

Example 5 (a)

[0151]




TLC : Rf 0.3 (chloroform : methanol : acetic acid = 50 : 10 : 1),

NMR (CD₃OD) : δ 8.20(2H, d, J=8.0Hz), 7.95(2H, d, J=8.0Hz), 7.81(1H, d, J=2.0Hz), 7.79(1H, d, J=2.0Hz), 7.69(5H, brs), 7.55(2H, d, J=8.5Hz), 7.39(2H, d, J=8.5Hz), 5.63(2H, s), 2.72(6H, s).

Example 5 (b)

[0152]




TLC : Rf 0.48 (chloroform : methanol : acetic acid = 10 : 1 : 1),

NMR (CD₃OD + CDCl₃) : δ 8.05(2H, d, J=8.4Hz), 7.89(2H, d, J=8.8Hz), 7.71(1H, d, J=8.0Hz), 7.46(2H, d, J=8.8Hz), 7.40(1H, s), 7.37-7.30(2H, m), 7.17(1H, d, J=8.0Hz), 7.16(2H, d, J=8.4Hz), 5.95(2H, s), 4.30(2H, q, J=7.4Hz), 2.73(3H, s), 1.33(3H, t, J=7.4Hz).

Reference Example 9


[0153] 4-(N-Benzyl-N-ethoxycarbonylaminoethyl)benzoic acid benzyl ester



[0154] A solution of 4-(N-benzylaminomethyl)benzoic acid benzyl ester (5.21g) and bromoacetic acid benzyl ester (1.7ml) in DMF (10ml) was stirred for two hours at 80° C and ice water was added thereto. The reaction solution was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous solution of sodium hydrogen carbonate, water and a saturated aqueous solution of sodium chloride, successively. The organic layer was dried over and evaporated. The residue was purified by silica gel column chromatography to give the title compound (2.26g).

Reference Example 10


[0155] 4-(N-Benzyl-N-ethoxycarbonylaminoethyl)benzoic acid hydrochloride



[0156] A mixture solution of the compound prepared in Reference Example 9 (2.26g), methanesulfonic acid (10.5ml) and anisole (25ml) was stirred for one hour at room temperature. To the reaction solution was added ice water and extracted with chloroform. The organic layer was washed with water, a saturated aqueous solution of sodium chloride, dried over and evaporated. The residue was purified by silica gel column chromatography to give amine. 4N hydrochloric acid-dioxane was added to amine and the mixture was evaporated to give the title compound (1.76g).

Example 6

[0157] N-(4-(4-Amidino-phenoxy-carbonyl)phenylmethyl)-N-benzylaminoacetic acid ethyl ester 2 hydrochloride



[0158] By the same procedure as Example 2, the title compound having the following physical data was given.
 TLC : R_f 0.42 (chloroform : methanol : acetic acid = 10 : 2 : 1),
 NMR (CD₃OD) : δ 8.25(2H, d, J=8Hz), 7.90(2H, d, J=8Hz), 7.60(2H, d, J=8Hz), 7.50(2H, d, J=8Hz), 7.40-7.20(5H, m), 4.15(2H, q, J=7Hz), 3.90(2H, s), 3.80(2H, s), 3.30(2H, s), 1.25(3H, t, J=7Hz).

Formulation Example 1

[0159] The following components were admixed in conventional manner and punched out to obtain 100 tablets each

containing 100mg of active ingredient.

Compound number 1	10g
Cellulose calcium glycolate (disintegrating agent)	0.2g
Magnesium stearate (Lubricating agent)	0.1g
Microcrystalline cellulose	1.7g

Formation Example 2

[0160] The following components were admixed conventional method and punched out to obtain 100 tables each containing 100mg of active ingredient.

Compound number 2	10g
Cellulose calcium glycolate (disintegrating agent)	0.2g
Magnesium stearate (Lubricating agent)	0.1g
Microcrystalline cellulose	1.7g

Formation Example 3

[0161] The following components were admixed in conventional manner. The solution was sterilized conventional manner, placed 5ml portions into 10ml ampoules and obtained 100 ampoules each containing 10mg of the active ingredient.

Compound number 1	1g
Citric acid	0.2g
distilled water	500ml


Formation Example 4

[0162] The following components were admixed in conventional manner. The solution was sterilized in conventional manner, placed 5ml portions into 10ml ampoules and obtained 100 ampoules each containing 10mg of the active ingredient.


Compound number 2	1g
Citric acid	0.2g
distilled water	500ml

Claims



1. An amidinophenol derivative of the formula (IB):




wherein:



is a group of the formula :




or








(in which R^0 is hydrogen, or C1-4 alkyl, or C1-4 alkoxy),


T is NH or oxygen,
E is a single bond, or
a group of the formula :




A_0 is a single bond, C1-4 alkylene, -oxy-(C1-4)alkylene-, -thio-(C1-4)alkylene-, C2-8 alkenylene, or C2-8 alkenylene which is substituted by carboxy or by C1-4 alkoxy carbonyl, R^{100} , R^{200} , R^{300} and R^{400} each independently, is hydrogen or C1-4 alkyl, R is a group of the formula:



, or



(in which the grouping:




is a 4-10 membered hetero ring containing one or two nitrogen atoms),
 R^{50} , R^{60} and R^{70} each independently, is,

- (i) hydrogen,
- (ii) C1-8 alkyl,
- (iii) C2-8 alkenyl
- (iv) $-\text{COOR}^{110}$ (in which R^{110} is hydrogen or C1-4 alkyl unsubstituted or substituted by phenyl),
- (v) $-(\text{C1-8 alkenylene})-\text{COOR}^{110}$ (in which R^{110} has the same meaning as hereinbefore defined),
- (vi) $-(\text{C2-8 alkenylene})-\text{COOR}^{110}$ (in which R^{110} has the same meaning as hereinbefore defined),
- (vii) C4-7 cycloalkyl,
- (viii) $-(\text{C1-4 alkenylene})-(4-7 \text{ membered hetero ring containing one oxygen})$,
- (ix) $-(\text{C1-4 alkenylene})-(4-7 \text{ membered hetero ring containing one nitrogen})$,
- (x) phenyl,
- (xi) C1-8 alkyl which is substituted by one or two phenyl,
- (xii) $-(\text{C1-4 alkenylene})-\text{O}-\text{benzoyl}$,
- (xiii) $-(\text{C1-4 alkenylene})-\text{CONH}-(\text{C1-4 alkenylene})-\text{NR}^{120}\text{H}^{130}$,
- (xiv) $-(\text{C1-4 alkenylene})-\text{COO}-(\text{C1-4 alkenylene})-\text{NR}^{120}\text{R}^{130}$,
- (xv) $-(\text{C1-4 alkenylene})-\text{COO}-\text{amidinophenyl}$,
- (xvi) $-(\text{C1-4 alkenylene})-\text{CONH}-(\text{C1-4 alkyl which is substituted by one or two } -\text{COOR}^{110})$ (in which R^{110} has the same meaning as hereinbefore defined),
- (xvii) $-(\text{C1-4 alkenylene})-\text{CONR}^{120}\text{R}^{130}$, or
- (xviii) (C1-4) alkoxy (C1-4) alkyl;

R^{80} and R^{90} each independently, is C1-4 alkyl or $-(\text{C1-4 alkenylene})-\text{phenyl}$,
 R^{120} and R^{130} each independently, is hydrogen, C1-4 alkyl, or C2-8 alkenyl, with the provisos that:

- (1) R^{50} and R^{60} in the formulae (i) and (iii), and R^{50} , R^{60} and R^{70} in the formulae (ii) and (iv), do not represent hydrogen at the same time,
- (2) when at least one substituent in R^{50} , R^{60} , R^{70} and A_0 represent substituent containing $-\text{COO}-t\text{-Bu}$, the other groups do not represent groups containing carboxy,
- (3) R^{120} and R^{130} do not represent hydrogen at the same time,
- (4) when

T is oxygen,
 the grouping:




is the formula (i) as hereinbefore described,
 E is a single bond,
 A_0 is a single bond, C1-4 alkenylene or vinylene which is optionally substituted by one or two C1-4 alkyl,
 and
 R is the formula (i) as hereinbefore described, then at least one group in R^{50} and R^{60} is

- (viii) $-(\text{C1-4 alkenylene})-(4-7 \text{ membered hetero ring containing one oxygen})$,
- (ix) $-(\text{C1-4 alkenylene})-(4-7 \text{ membered hetero ring containing one nitrogen})$,
- (xi-1) C1-8 alkyl which is substituted by two phenyl,

- (xii) -(C1-4 alkylene)-O-benzoyl,
 (xiii) -(C1-4 alkylene)-CONH-(C1-4 alkylene) -NR¹²⁰R¹³⁰,
 (xiv) -(C1-4 alkylene)-COO- (C1-4 alkylene) -NR¹²⁰R¹³⁰,
 (xv) -(C1-4 alkylene) -COO-amidinophenyl,
 (xvi) -(C1-4 alkylene)-CONH-(C1-4 alkyl which is substituted by one or two -COOR¹¹⁰) (in which R¹¹⁰ has the same meaning as hereinbefore defined), or
 (xvii) -(C1-4 alkylene) -CONR¹²⁰R¹³⁰ (in which R¹²⁰ and R¹³⁰ each independently is C1-4 alkyl or C2-8 alkenyl) ;

(5) when

T is oxygen,
 the formula:



is the formula (i) as hereinbefore defined,


E is bond,

A₀ is bond, C1-4 alkylene or vinylene which may be optionally substituted by one or two C1-4 alkyl,
 and R is the formula (ii) as hereinbefore defined,
 then R⁵⁰, R⁶⁰ and R⁷⁰ do not represent hydrogen; and


(6) the compound of formula (IB) is not p-amidinophenyl p-[(2-isopropyl-l-imidazolyl)methyl]benzoate;

or 4-(4-amidinophenoxy-carbonyl)-α-methylcinnamic acid N-1-(S)-ethoxycarbonyl-2-benzoyloxyethylamide; or
 a non-toxic salt thereof or a non-toxic acid addition salt thereof.

2. A compound according to claim 1, wherein R is



or




in which the various symbols are as defined in claim 1.

3. A compound according to claim 1, wherein R is




or




in which the various symbols are as defined in claim 1.

4. A compound according to claim 1, wherein R is




in which the various symbols are as defined in claim 1.

5. A compound according to claim 1, wherein R is




in which the various symbols are as defined in claim 1.

6. A compound according to claim 1, wherein



is a 4-10 membered hetero ring containing one nitrogen.

7. A compound according to claim 1, wherein



is a 4-10 membered hetero ring containing two nitrogen.

8. A compound according to claim 1, wherein R^{50} , R^{60} and R^{70} each independently, is

- (i) hydrogen,
- (ii) C1-8 alkyl,
- (iii) C2-8 alkenyl
- (iv) $-COOR^{110}$,
- (v) $-(C1-8 \text{ alkylene})-COOR^{110}$,
- (vi) $-(C2-8 \text{ alkenylene})-COOR^{110}$,
- (vii) C4-7 cycloalkyl,
- (x) phenyl,
- (xi) C1-8 alkyl which is substituted by one or two phenyl,
- (xii) $-(C1-4 \text{ alkylene})-O\text{-benzoyl}$,
- (xiii) $-(C1-4 \text{ alkylene})-CONH-(C1-4 \text{ alkylene})-NR^{120}R^{130}$,
- (xiv) $-(C1-4 \text{ alkylene})-COO-(C1-4 \text{ alkylene})-NR^{120}R^{130}$,
- (xv) $-(C1-4 \text{ alkylene})-OCO\text{-amidinophenyl}$,
- (xvi) $-(C1-4 \text{ alkylene})-CONH-(C1-4 \text{ alkyl which is substituted by one or two } -COOR^{110})$,
- (xvii) $-(C1-4 \text{ alkylene})-CONR^{120}R^{130}$, or
- (xviii) $-(C1-4)\text{alkoxy}(C1-4)\text{alkyl}$,

in which the various symbols are defined in claim 1.

9. A compound according to claim 1, wherein R^{50} , R^{60} and R^{70} each independently, is

- (viii) $-(C1-4 \text{ alkylene})-(4-7 \text{ membered hetero ring containing one oxygen})$.

10. A compound according to claim 1, wherein R^{50} , R^{60} and R^{70} each independently, is

- (ix) $-(C1-4 \text{ alkylene})-(4-7 \text{ membered hetero ring containing one nitrogen})$.

11. A compound according to claim 1, which is

- 4-(4-amidinophenoxycarbonyl)phenoxyacetic acid N-2-propenyl-N-ethoxycarbonylmethylamide,
- 4-(6-amidino-7,8-dihydro-2-naphthyloxycarbonyl)benzoic acid N-phenyl-N-phenylmethylamide,
- 4-(6-amidino-7,8-dihydro-2-naphthyloxycarbonyl)benzoic acid N-phenyl-N-benzoyloxycarbonylmethylamide,
- 4-(6-amidino-7,8-dihydro-2-naphthyloxycarbonyl)benzoic acid N-phenyl-N-ethoxycarbonylmethylamide,
- 4-[4-(2-amidinoethenyl)phenoxy carbonyl]benzoic acid N-phenyl-N-ethoxycarbonylmethylamide,

4-(4-amidinophenoxy carbonyl)- α -methylcinnamic acid N-1-(S)-ethoxycarbonyl-2-benzoyloxyethylamide,
 4-[2-(4-amidinophenoxy carbonyl)ethenyl]benzoic acid N-1-(S), 3-bis(ethoxycarbonyl)propylamide,
 4-[2-(4-amidinophenoxy carbonyl)-1-propenyl]benzoic acid N-1-(S), 3-bis(ethoxycarbonyl)propylamide,
 4-(4-amidinophenoxy carbonyl)- α -methylcinnamic acid N-[1,1,4-tris(ethoxycarbonyl)-3-butenyl]amide,
 4-[4-(2-amidinoethenyl)phenoxy carbonyl]benzoic acid N-ethoxycarbonylmethyl-N-allylamide,
 4-(4-amidinophenoxy carbonyl)phenoxyacetic acid N-1-(S), 3-bis(ethoxycarbonyl)propylamide,
 4-(4-amidinophenoxy carbonyl)- α -methylcinnamic acid N-1, 1, 2-tris(ethoxycarbonyl)ethylamide,
 4-[2-(4-amidinophenoxy carbonyl)-1-propenyl]benzoic acid N-ethoxycarbonylmethyl-N-allylamide,
 5-[4-(4-amidinophenoxy carbonyl)phenyl]-2, 4-pentadienoic acid N-1-(S), 3-bis(ethoxycarbonyl)propylamide,
 5-[4-(4-amidinophenoxy carbonyl)phenyl]-2, 4-pentadienoic acid N-ethoxycarbonylmethyl-N-allylamide,
 4-(4-amidinophenoxy carbonyl)- α -methylcinnamic acid N-[1,1-bis(ethoxycarbonyl)-3-butenyl]amide,
 4-[2-(4-amidinophenoxy carbonyl)-1-methylethenyl]benzoic acid N-1-(S), 3-bis(ethoxycarbonyl)propylamide,
 4-[2-(4-amidinophenoxy carbonyl)-1-methylethenyl] benzoic acid N-ethoxycarbonylmethyl-N-allylamide,
 3-[4-(4-amidinophenoxy carbonyl)phenyl]-2-ethoxycarbonyl-2-propenoic acid N-1-(S), 3-bis(ethoxycarbonyl)
 propylamide,
 3-[4-(4-amidinophenoxy carbonyl)phenyl]-2-ethoxycarbonyl-2-propenoic acid N-ethoxycarbonylmethyl-N-ally-
 lamide,
 4-(4-amidinophenoxy carbonyl)- α -methylcinnamic acid N-2-dimethylaminoethylcarbamoylmethyl-N-allylamide,
 4-(4-amidinophenoxy carbonyl)- α -methylcinnamic acid N-2-dimethylaminoethoxycarbonylmethyl-N-allylamide,
 4-(4-amidinophenoxy carbonyl)- α -methylcinnamic acid N-4-amidinophenoxy carbonylmethyl-N-allylamide,
 4-(4-amidinophenoxy carbonyl)- α -methylcinnamic acid N-1, 1-bis(ethoxycarbonyl)methylcarbamoylmethyl-N-
 allylamide,
 4-(4-amidinophenoxy carbonyl)- α -methylcinnamic acid N-(diallylcarbamoyl)methyl-N-allylamide,
 4-(4-amidinophenylcarbamoyl)- α -methylcinnamic acid N-allyl-N-carboxymethylamide, or
 4-(4-amidino-1-methoxyphenoxy carbonyl)benzoic acid N-1, 1-bis (phenylmethyl)methyl-N-3-phenylpropylamide.

12. A compound according to claim 1, which is

2-[4-(4-amidinophenoxy carbonyl) phenylmethylimino]-2-(N-benzyl-N-methylamino)acetic acid ethylester, or
 2-[N [4-(4-amidinophenoxy carbonyl)phenyl]methyl-N-benzylamino}acetic acid ethyl ester.

13. A compound according to claim 1, which is

ethyl 1-[4-(4-amidinophenoxy carbonyl)phenylmethyl]-1-(3-phenylpropyl)phosphinate, or
 diethyl 2-[4-(4-amidinophenoxy carbonyl)phenyl]-ethenylphosphonate.

14. A compound according to claim 1, which is

4-(4-amidinophenoxy carbonyl)- α -methylcinnamic acid N-(2,3,4,5-tetrahydro-(furan-2-yl)methyl-N-carboxy-
methylamide.

15. A compound according to claim 1, which is

4-(4-amidinophenoxy carbonyl)- α -methylcinnamic acid N-(4-piperidinylmethyl)-N-carboxymethylamide.

16. A compound according to claim 1, which is


4-(4-phenylpiperidinylmethyl)benzoic acid 4-amidinophenyl ester, or
 4-(2-ethoxycarbonylindolylmethyl)benzoic acid 4-amidinophenyl ester.

17. A compound according to claim 1, which is


4-(2-phenylimidazolylmethyl)benzoic acid 4-amidinophenyl ester.

18. A compound according to any one of the preceding claims in the form of a non-toxic acid addition salt.


19. A process for the preparation of a compound of formula (IB):



10 which comprises the reaction to form an ester or amide bond, of a compound of the formula:



20 (wherein the various symbols are as defined in claim 1)
with a compound of the formula :




30 (wherein the various symbols are as defined in claim 1); and optionally converting the amidinophenol derivative thus obtained into a salt or acid addition salt thereof.

35 **20.** A pharmaceutical composition which comprises, as active ingredient, an amidinophenol derivative of the formula (IB) depicted in claim 1, or a non-toxic salt thereof, or non-toxic acid addition salt thereof, with a carrier or coating.

21. Use of a compound of formula (IB) according to any one of claims 1 to 19 or a non-toxic salt thereof or non-toxic acid addition salt thereof in the manufacture of a medicament for the prevention and/or treatment of a condition which can be ameliorated by the administration of a leukotriene B₄ antagonist or by an inhibitor of phospholipase A₂ or trypsin.

40 **22.** Use of a compound of the formula (IA):



wherein

55 R¹ and R² each independently, is:


- (i) hydrogen, or
- (ii) -COOR⁴ (in which R⁴ is C1-3 alkyl);

A is

- (i) a single bond,
- (ii) C1-4 alkylene, or
- (iii) $-C(R^5)=C(R^6)-$ (in which R^5 and R^6 each independently, is hydrogen or C1-4 alkyl) ;

R^3 is

- (i) $-\text{CON}(R^7)R^8$,
- (ii) $-\text{CONR}^9-\text{CH}(R^7)R^8$, or
- (iii)



(wherein R^7 and R^8 each independently, is


- (1) hydrogen,
- (2) phenyl,
- (3) $-(\text{C1-4 alkylene})-\text{phenyl}$,
- (4) $-(\text{C1-4 alkylene})-\text{phenyl}$ which is substituted by one or two $-\text{R}^{11}-\text{COOR}^{12}$ (in which R^{11} is a single bond or C1-8 alkylene, and R^{12} is hydrogen or C1-4 alkyl),
- (5) C1-5 alkyl,
- (6) C2-10 alkenyl containing one to three double bonds,
- (7) $-\text{R}^{11a}-\text{COOR}^{12}$ (in which R^{11a} is

- (a) a single bond,
- (b) C1-8 alkylene,
- (c) C2-8 alkenylene, or
- (d) C4-8 alkenylene in which one or two carbon atoms in the main chain are replaced by sulfur, and R^{12} has the same meaning as hereinbefore defined), or

- (8) C3-7 cycloalkyl;

R^9 is

- (1) hydrogen,
 - (2) $-\text{R}^{11}-\text{COOR}^{12}$ (in which the various symbols have the same meanings as hereinbefore defined), or
 - (3) C2-6 alkoxyalkyl;
- the grouping:



is a 4-7 membered mono hetero ring contain one or two nitrogen;

R^{10} is


(1) hydrogen, or
(2) -(C1-4 alkylene)-phenyl,
with the proviso that:

- (1) both R^7 and R^8 do not represent hydrogen at the same time,
(2) when at least one group in R^7 , R^8 , and R^9 represent the group containing -COO-t-Bu, the other groups do not represent the groups containing carboxy;


or a non-toxic salt thereof or non-toxic acid addition salt thereof in the manufacture of a medicament for the prevention and/or treatment of rheumatoid arthritis, inflammatory bowel diseases, psoriasis, nonsteroidal anti-inflammatory agent-induced stomach diseases, adult respiratory distress syndrome, cardiac infarction, allergic rhinitis, hemodialysis-induced neutropenia or anaphase asthma.

Patentansprüche

1. Amidinophenolderivat der Formel [IB] :




worin bedeuten:




eine Gruppe der Formeln:

(i)




(ii)



oder


(iii)



(worin R⁰ für Wasserstoff, C1-4-Alkyl oder C1-4-Alkoxy steht);

T NH oder Sauerstoff;

E eine Einfachbindung oder eine Gruppe der Formel:




A₀ eine Einfachbindung, C1-4-Alkylen, -Oxy-(C1-4)-Alkylen-, -Thio -(C1-4) -Alkylen-, C2-8-Alkenylen oder durch Carboxy oder C1-4-Alkoxy-carbonyl substituiertes C2-8-Alkenylen;


R¹⁰⁰, R²⁰⁰, R³⁰⁰ und R⁴⁰⁰ jeweils unabhängig voneinander Wasserstoff oder C1-4-Alkyl;

R eine Gruppe der Formeln:


(i)




(ii)




(iii)




(iv)



(v)




oder




oder

(vi)



(worin die Gruppierung:



für einen 4-10-gliedrigen Heteroring mit einem oder zwei Stickstoffatom(en) steht);
 R^{50} , R^{60} und R^{70} jeweils unabhängig voneinander

(i) Wasserstoff,

(ii) C1-8-Alkyl,

(iii) C2-8-Alkenyl

(iv) ---COOR^{110} (mit R^{110} gleich Wasserstoff oder ggf. mit Phenyl substituiertem C1-4-Alkyl),(v) $\text{---(C1-8-Alkylen)---COOR}^{110}$ (mit R^{110} in der zuvor angegebenen Bedeutung),(vi) $\text{---(C2-8-Alkenylen)---COOR}^{110}$ (mit R^{110} in der zuvor angegebenen Bedeutung),

(vii) C4-7-Cycloalkyl,

(viii) $\text{---(C1-4-Alkylen)---(4-7-gliedriger Heteroring mit einem Sauerstoff)}$,(ix) $\text{---(C1-4-Alkylen)---(4-7-gliedriger Heteroring mit einem Stickstoff)}$,

(x) Phenyl,


(xi) 1- oder 2-fach phenylsubstituiertes C1-8-Alkyl,

(xii) $\text{---(C1-4-Alkylen)---O-Benzoyl}$,(xiii) $\text{---(C1-4-Alkylen)---CONH---(C1-4-Alkylen)---NR}^{120}\text{R}^{130}$,(xiv) $\text{---(C1-4-Alkylen)---COO---(C1-4-Alkylen)---NR}^{120}\text{R}^{130}$,(xv) $\text{---(C1-4-Alkylen)---COO---Amidinophenyl}$,

- (xvi) -(C1-4-Alkylen)-CONH-(1- oder 2-fach durch-COOR¹¹⁰ substituiertes C1-4-Alkyl) (mit R¹¹⁰ in der zuvor angegebenen Bedeutung),
 (xvii) -(C1-4-Alkylen)-CONR¹²⁰R¹³⁰ oder
 (xviii) (C1-4) Alkoxy (C1-4) Alkyl;

R⁸⁰ und R⁹⁰ jeweils unabhängig voneinander C1-4-Alkyl oder (C1-4-Alkylen)-Phenyl,
 R¹²⁰ und R¹³⁰ jeweils unabhängig voneinander Wasserstoff, C1-4-Alkyl oder C2-8-Alkenyl, wobei gilt, daß

- (1) R⁵⁰ und R⁶⁰ in den Formeln (i) und (iii), und R⁵⁰, R⁶⁰ und R⁷⁰ in den Formeln (ii) und (iv) nicht gleichzeitig Wasserstoff bedeuten dürfen,
 (2) im Falle, daß mindestens ein Substituent in R⁵⁰, R⁶⁰, R⁷⁰ und A₀ einem Substituenten mit -COO-tert.-Bu entspricht, die anderen Gruppen nicht für carboxyhaltige Gruppen stehen dürfen,
 (3) R¹²⁰ und R¹³⁰ nicht gleichzeitig Wasserstoff bedeuten dürfen,
 (4) im Falle, daß T für Sauerstoff steht, die Gruppe




der zuvor angegebenen Formel (i) entspricht,
 E eine Einfachbindung bedeutet,

A₀ für eine Einfachbindung, C1-4-Alkylen oder ggf. 1- oder 2-fach durch C1-4-Alkyl substituiertes Vinylen steht, und
 R der zuvor beschriebenen Formel (i) entspricht, mindestens eine Gruppe in R⁵⁰ und R⁶⁰

- (viii) -(C1-4-Alkylen)-(4-7-gliedriger Heteroring mit einem Sauerstoff),
 (ix) -(C1-4-Alkylen)-(4-7-gliedriger Heteroring mit einem Stickstoff),
 (xi-1) 2-fach phenylsubstituiertes C1-8-Alkyl,
 (xii) -(C1-4-Alkylen)-O-Benzoyl,
 (xiii) -(C1-4-Alkylen)-CONH-(C1-4-Alkylen)-NR¹²⁰R¹³⁰,
 (xiv) -(C1-4-Alkylen)-COO-(C1-4-Alkylen)-NR¹²⁰R¹³⁰,
 (xv) - (C1-4-Alkylen)-COO-Amidinophenyl,
 (xvi) -(C1-4-Alkylen)-CONH-(1- oder 2-fach durch -COOR¹¹⁰ substituiertes C1-4-Alkyl) (worin R¹¹⁰ die zuvor angegebenen Bedeutung besitzt), oder
 (xvii) -(C1-4-Alkylen)-CONR¹²⁰R¹³⁰ (mit R¹²⁰ und R¹³⁰ jeweils unabhängig voneinander gleich C1-4-Alkyl oder C2-8-Alkenyl)
 ist;

(5) im Falle, daß
 T Sauerstoff darstellt,
 die Formel



der zuvor angegebenen Formel (i) entspricht,

E eine Bindung bedeutet,
 A₀ für eine Bindung, C1-4-Alkylen oder ggf. 1- oder 2-fach durch C1-4-Alkyl substituiertes Vinylen steht, und
 R der zuvor angegebenen Formel (ii) entspricht, R⁵⁰, R⁶⁰ und R⁷⁰ nicht für Wasserstoff stehen dürfen, und


(6) die Verbindung der Formel (IB) nicht aus p-Amidinophenyl p-[(2-isopropyl-1-imidazolyl)methyl]benzoat;

oder 4-(4-Amidinophenoxy-carbonyl)- α -methylzimtsäure N-1-(S)-ethoxycarbonyl-2-benzoyloxyethylamid besteht, oder ein nicht-toxisches Salz oder ein nicht-toxisches Säureadditionssalz hiervon.

2. Verbindung nach Anspruch 1, wobei R für




oder




worin die verschiedenen Symbole die in Anspruch 1 angegebene Bedeutung besitzen, steht.

3. Verbindung nach Anspruch 1, wobei R für




oder




worin die verschiedenen Symbole die in Anspruch 1 angegebene Bedeutung besitzen, steht.

4. Verbindung nach Anspruch 1, wobei R für




worin die verschiedenen Symbole die in Anspruch 1 angegebene Bedeutung besitzen, steht.

5. Verbindung nach Anspruch 1, wobei R für




worin die verschiedenen Symbole die in Anspruch 1 angegebene Bedeutung besitzen, steht.

6. Verbindung nach Anspruch 1, wobei



für einen 4-10-gliedrigen Heteroring mit einem Stickstoff steht.

7. Verbindung nach Anspruch 1, worin



für einen 4-10-gliedrigen Heteroring mit zwei Stick-stoffen steht.

8. Verbindung nach Anspruch 1, worin R⁵⁰, R⁶⁰ und R⁷⁰ jeweils unabhängig voneinander bedeuten:

- (i) Wasserstoff,
- (ii) C1-8-Alkyl,
- (iii) C2-8-Alkenyl
- (iv) -COOR¹¹⁰,
- (v) -(C1-8-Alkyl)-COOR¹¹⁰,
- (vi) -(C2-8-Alkenyl)-COOR¹¹⁰,
- (vii) C4-7-Cycloalkyl,
- (x) Phenyl,

- (xi) 1- oder 2-fach phenylsubstituiertes C1-8-Alkyl,
- (xii) -(C1-4-Alkyl)-O-Benzoyl,
- (xiii) -(C1-4-Alkyl)-CONH-(C1-4-Alkyl)-NR¹²⁰R¹³⁰,
- (xiv) -(C1-4-Alkyl)-COO-(C1-4-Alkyl)-NR¹²⁰R¹³⁰,
- (xv) -(C1-4-Alkyl)-OCO-Amidinophenyl,
- (xvi) -(C1-4-Alkyl)-CONH-(1- oder 2-fach durch -COOR¹¹⁰ substituiertes C1-4-Alkyl),
- (xvii) -(C1-4-Alkyl)-CONR¹²⁰R¹³⁰ oder
- (xviii) (C1-4) Alkoxy (C1-4) Alkyl,

worin die verschiedenen Symbole die in Anspruch 1 angegebenen Bedeutung besitzen.

9. Verbindung nach Anspruch 1, wobei R⁵⁰, R⁶⁰ und R⁷⁰ jeweils unabhängig voneinander für

(viii) -(C1-4-Alkyl)-(4-7-gliedriger Heteroring mit einem Sauerstoff) stehen.

10. Verbindung nach Anspruch 1, wobei R⁵⁰, R⁶⁰ und R⁷⁰ jeweils unabhängig voneinander für

(ix) -(C1-4-Alkyl)-(4-7-gliedriger Heteroring mit einem Stickstoff) stehen.

11. Verbindung nach Anspruch 1, nämlich

4-(4-Amidinophenoxy-carbonyl)phenoxyessigsäure N-2-propenyl-N-ethoxycarbonylmethylamid,
 4-(6-Amidino-7,8-dihydro-2-naphthyloxy-carbonyl)benzoesäure N-phenyl-N-phenylmethylamid,
 4-(6-Amidino-7,8-dihydro-2-naphthyloxy-carbonyl)benzoesäure N-phenyl-N-benzyloxy-carbonylmethylamid,
 4-(6-Amidino-7,8-dihydro-2-naphthyloxy-carbonyl)benzoesäure N-phenyl-N-ethoxycarbonylmethylamid,
 4-[4-(2-Amidinoethenyl)phenoxy-carbonyl]benzoesäure N-phenyl-N-ethoxycarbonylmethylamid,
 4-(4-Amidinophenoxy-carbonyl)- α -methylzimtsäure N-1-(S)-ethoxycarbonyl-2-benzoyloxyethylamid,
 4-[2-(4-Amidinophenoxy-carbonyl)ethenyl]benzoesäure N-1-(S), 3-bis(ethoxycarbonyl)propylamid,
 4-[2-(4-Amidinophenoxy-carbonyl)-1-propenyl]benzoesäure N-1-(S), 3-bis(ethoxycarbonyl)propylamid,
 4-(4-Amidinophenoxy-carbonyl)- α -methylzimtsäure N-[1,1,4-tris(ethoxycarbonyl)-3-butenyl]amid,
 4-[4-(2-Amidinoethenyl)phenoxy-carbonyl]benzoesäure N-ethoxycarbonylmethyl-N-allylamid,
 4-(4-Amidinophenoxy-carbonyl)phenoxyessigsäure N-1-(S), 3-bis(ethoxycarbonyl)propylamid,
 4-(4-Amidinophenoxy-carbonyl)- α -methylzimtsäure N-1,1,2-tris(ethoxycarbonyl)ethylamid,
 4-[2-(4-Amidinophenoxy-carbonyl)-1-propenyl]benzoesäure N-ethoxycarbonylmethyl-N-allylamid,
 5-[4-(4-Amidinophenoxy-carbonyl)phenyl]-2,4-pentadiensäure N-1-(S), 3-bis(ethoxycarbonyl)propylamid,
 5-[4-(4-Amidinophenoxy-carbonyl)phenyl]-2,4-pentadiensäure N-ethoxycarbonylmethyl-N-allylamid,
 4-(4-Amidinophenoxy-carbonyl)- α -methylzimtsäure N-[1,1-bis(ethoxycarbonyl)-3-butenyl]amid,
 4-[2-(4-Amidinophenoxy-carbonyl)-1-methylethenyl]benzoesäure N-1-(S), 3-bis(ethoxycarbonyl)propylamid,
 4-[2-(4-Amidinophenoxy-carbonyl)-1-methylethenyl]benzoesäure N-ethoxycarbonylmethyl-N-allylamid,
 3-[4-(4-Amidinophenoxy-carbonyl)phenyl]-2-ethoxycarbonyl-2-propensäure N-1-(S), 3-bis(ethoxycarbonyl)propylamid,
 3-[4-(4-Amidinophenoxy-carbonyl)phenyl]-2-ethoxycarbonyl-2-propensäure N-ethoxycarbonylmethyl-N-allylamid,
 4-(4-Amidinophenoxy-carbonyl)- α -methylzimtsäure N-2-dimethylaminoethylcarbamoylmethyl-N-allylamid,
 4-(4-Amidinophenoxy-carbonyl)- α -methylzimtsäure N-2-dimethylaminoethoxycarbonylmethyl-N-allylamid,
 4-(4-Amidinophenoxy-carbonyl)- α -methylzimtsäure N-4-amidinophenoxy-carbonylmethyl-N-allylamid,
 4-(4-Amidinophenoxy-carbonyl)- α -methylzimtsäure N-1,1-bis(ethoxycarbonyl)methylcarbamoylmethyl-N-allylamid,
 4-(4-Amidinophenoxy-carbonyl)- α -methylzimtsäure N-(diallylcarbamoyl)methyl-N-allylamid,
 4-(4-Amidinophenylcarbamoyl)- α -methylzimtsäure N-allyl-N-carboxymethylamid oder
 4-(4-Amidino-1-methoxyphenoxy-carbonyl)benzoesäure N-1,1-bis(phenylmethyl)methyl-N-3-phenylpropylamid.

12. Verbindung nach Anspruch 1, nämlich

2-[4-(4-Amidinophenoxy-carbonyl)phenylmethylimino]-2-(N-benzyl-N-methylamino)essigsäureethylester oder
 2-(N-[4-(4-Amidinophenoxy-carbonyl)phenyl]methyl-N-benzylamino)essigsäureethylester.

13. Verbindung nach Anspruch 1, nämlich

Ethyl 1-[4-(4-amidinophenoxy)carbonyl]phenylmethyl]-1-(3-phenylpropyl)phosphinat oder
Diethyl 2-[4-(4-amidinophenoxy)carbonyl]phenyl]ethenylphosphonat.

14. Verbindung nach Anspruch 1, nämlich

4-(4-Amidinophenoxy)carbonyl)- α -methylzimtsäure N-(2,3,4,5-tetrahydro-furan-2-yl)methyl-N-carboxymethyl-
amid.

15. Verbindung nach Anspruch 1, nämlich

4-(4-Amidinophenoxy)carbonyl)- α -methylzimtsäure N-(4-piperidinylmethyl)-N-carboxymethylamid.

16. Verbindung nach Anspruch 1, nämlich


4-(4-Phenylpiperidinylmethyl)benzoesäure 4-amidinophenylester oder
4-(2-Ethoxycarbonylindolylmethyl)benzoesäure 4-amidinophenylester.

17. Verbindung nach Anspruch 1, nämlich


4-(2-Phenylimidazolylmethyl)benzoesäure 4-amidinophenylester.

18. Verbindung nach einem der vorhergehenden Ansprüche in Form eines nicht-toxischen Säureadditionssalzes.


19. Verfahren zur Herstellung einer Verbindung der Formel (IB) :



durch Umsetzen einer Verbindung der Formel:



(worin die verschiedenen Symbole die in Anspruch 1 angegebene Bedeutung besitzen)
mit einer Verbindung der Formel:




(worin die verschiedenen Symbole die in Anspruch 1 angegebene Bedeutung besitzen) zur Bildung einer Ester-
oder Amidbindung und ggf. Umwandeln des hierbei erhaltenen Amidinophenolderivats in ein Salz oder Säuread-
ditionssalz desselben.

20. Arzneimittelzubereitung, umfassend als aktiven Bestandteil ein Amidinophenolderivat der Formel (IB) gemäß Anspruch 1 oder ein nicht-toxisches Salz oder nicht-toxisches Säureadditionssalz desselben sowie einen Träger oder einen Überzug.

21. Verwendung einer Verbindung der Formel (IB) nach einem der Ansprüche 1 bis 19 oder eines nicht-toxischen Salzes oder nicht-toxischen Säureadditionssalzes derselben bei der Herstellung eines Medikaments zur Verhinderung und/oder Behandlung eines Zustands, der durch Verabreichen eines Leukotriens B₄-Antagonisten oder durch einen Phospholipase A₂- oder Trypsin-Inhibitor gelindert werden kann.

22. Verwendung einer Verbindung der Formel (IA):



worin bedeuten:

R¹ und R² jeweils unabhängig voneinander

- (i) Wasserstoff oder
- (ii) -COOR⁴ (mit R⁴ gleich C1-3-Alkyl);

A

- (i) eine Einfachbindung,
- (ii) C1-4-Alkylen oder
- (iii) -C(R⁵)=C(R⁶)- (mit R⁵ und R⁶ unabhängig voneinander gleich Wasserstoff oder C1-4-Alkyl), und

R³

- (i) -CON (R⁷)R⁸.
- (ii) -CONR⁹-CH(R⁷)R⁸ oder
- (iii)



(worin R⁷ und R⁸ unabhängig voneinander für:

- (1) Wasserstoff,
- (2) Phenyl,
- (3) -(C1-4-Alkylen)-Phenyl,
- (4) -(C1-4-Alkylen)-Phenyl, welches 1- oder 2-fach-R¹¹-COOR¹² substituiert ist (worin R¹¹ eine Einfachbindung oder C1-8-Alkylen darstellt und R¹² Wasserstoff oder C1-4-Alkyl bedeutet),
- (5) C1-5-Alkyl,
- (6) C2-10-Alkenyl mit einer bis drei Doppelbindung(en),
- (7) -R^{11a}-COOR¹² (worin R^{11a}


(a) eine Einfachbindung,

- (b) C1-8-Alkylen,
 (c) C2-8-Alkenylen oder
 (d) C4-8-Alkenylen, wobei ein oder zwei Kohlenstoffatom(e) in der Hauptkette durch Schwefel ersetzt ist (sind) und R^{12} die zuvor angegebenen Bedeutung besitzt), oder

(8) C3-7-Cycloalkyl

stehen,
 R^9

- (1) Wasserstoff,
 (2) $-R^{11}-COOR^{12}$ (worin die verschiedenen Symbole die zuvor angegebene Bedeutung besitzen), oder
 (3) C2-6-Alkoxyalkyl
 darstellt,
 die Gruppierung:



für einen 4- bis 7-gliedrigen Monoheteroring mit einem oder zwei Stickstoff(en) steht, und

R^{10}


- (1) Wasserstoff oder
 (2) $-(C1-4-Alkylen)-Phenyl$
 darstellt, wobei gilt, daß

- (1) beide Reste R^7 und R^8 nicht gleichzeitig Wasserstoff bedeuten dürfen,
 (2) im Falle, daß mindestens eine Gruppe in R^7 , R^8 und R^9 für die $-COO-tert.-Bu$ enthaltende Gruppe steht, die anderen Gruppen nicht carboxyhaltige Gruppen darstellen dürfen);

oder eines nicht-toxischen Salzes oder nicht-toxischen Säureadditionssalzes derselben bei der Herstellung eines Medikaments zur Verhinderung und/oder Behandlung rheumatischer Arthritis, entzündlicher Darmerkrankungen, Psoriasis, durch nicht-steroidale, entzündungshemmende Mittel induzierter Magenkrankheiten, Atemnotsyndrom bei Erwachsenen, Herzinfarkt, allergischer Rhinitis, durch Hemodialyse induzierter Neutropenie oder Anaphase-Asthma.


Revendications

1. Dérivé d'amidinophénol de formule (IB) :




(IB)

dans laquelle :




est un groupe de formule :

(i)




(ii)



ou


(iii)



(où R^0 est l'hydrogène ou alkyle en C_1 à C_4 ou alcoxy en C_1 à C_4),

T est NH ou l'oxygène,


E est une liaison simple, ou un groupe de formule :







A_0 est une liaison simple, alkylène en C_1 à C_4 , -oxy(alkylène en C_1 à C_4)-, -thio(alkylène en C_1 à C_4)-, alcénylène en C_2 à C_8 , ou alcénylène en C_2 à C_8 qui est substitué par carboxy ou par alcoxycarbonyle en C_1 à C_4 , R^{100} , R^{200} , R^{300} et R^{400} , chacun indépendamment, est l'hydrogène ou alkyle en C_1 à C_4 .


R est un groupe de formule :

(i)






ou



(groupements dans lesquels :




est un hétérocycle à 4 à 10 chaîons contenant un ou deux atomes d'azote),
R⁵⁰, R⁶⁰ et R⁷⁰, chacun indépendamment, est

- (i) l'hydrogène,
- (ii) alkyle en C₁ à C₈,
- (iii) alcényle en C₂ à C₈,
- (iv) -COOR¹¹⁰ (où R¹¹⁰ est l'hydrogène ou alkyle en C₁ à C₄ non substitué ou substitué par phényle),
- (v) -(alkylène en C₁ à C₈)-COOR¹¹⁰ (où R¹¹⁰ a la même signification que celle définie ci-dessus),
- (vi) -(alcénylène en C₂ à C₈)-COOR¹¹⁰ (où R¹¹⁰ a la même signification que celle définie ci-dessus),
- (viii) cycloalkyle en C₄ à C₇,
- (viii) -(alkylène en C₁ à C₄)-(hétérocycle à 4 à 7 chaîons contenant un oxygène),
- (ix) -(alkylène en C₁ à C₄)-(hétérocycle à 4 à 7 chaîons contenant un azote),

- (x) phényle,
 (xi) alkyle en C₁ à C₈ qui est substitué par un ou deux groupes phényle,
 (xii) -(alkylène en C₁ à C₄)-O-benzoyle,
 (xiii) -(alkylène en C₁ à C₄)-CONH-(alkylène en C₁ à C₄)-NR¹²⁰R¹³⁰,
 (xiv) -(alkylène en C₁ à C₄)-COO-(alkylène en C₁ à C₄)-NR¹²⁰R¹³⁰,
 (xv) -(alkylène en C₁ à C₄)-COO-amidinophényle,
 (xvi) -(alkylène en C₁ à C₄)-CONH-(alkyle en C₁ à C₄ qui est substitué par un ou deux -COOR¹¹⁰) (où R¹¹⁰ a la même signification que celle définie ci-dessus),
 (xvii) -(alkylène en C₁ à C₄)-CONR¹²⁰R¹³⁰, ou
 (xviii) (alcoxy en C₁ à C₄)alkyle en C₁ en C₄,

R⁸⁰ et R⁹⁰, chacun indépendamment, est alkyle en C₁ à C₄ ou (alkylène en C₁ à C₄)phényle,
 R¹²⁰ et R¹³⁰, chacun indépendamment, est l'hydrogène, alkyle en C₁ à C₄ ou alcényle en C₂ à C₈,
 avec les conditions suivantes :

- (1) R⁵⁰ et R⁶⁰ dans les formules (i) est (iii), et R⁵⁰, R⁶⁰ et R⁷⁰ dans les formules (ii) et (iv), ne représentent pas en même temps l'hydrogène,
 (2) quand au moins un substituant dans R⁵⁰, R⁶⁰, R⁷⁰ et A₀ représente un substituant contenant -COO-t-Bu, les autres groupes ne représentent pas des groupes contenant carboxy,
 (3) R¹²⁰ et R¹³⁰ ne représentent pas l'hydrogène en même temps,
 (4) quand T est l'oxygène, le groupement :



est de formule (i) telle que définie ci-dessus,


E est une liaison simple,

A₀ est une liaison simple, alkylène en C₁ à C₄ ou vinylène qui est éventuellement substitué par un ou deux groupes alkyle en C₁ à C₄, et

R est de formule (i) telle que définie ci-dessus, alors au moins un groupe dans R⁵⁰ et R⁶⁰ est

- (viii) -(alkylène en C₁ à C₄)-(hétérocycle à 4 à 7 chaînons contenant un oxygène),
 (ix) -(alkylène en C₁ à C₄)-(hétérocycle à 4 à 7 chaînons contenant un azote),
 (xi-1) alkyle en C₁ à C₈ qui est substitué par deux groupes phényle,
 (xii) -(alkylène en C₁ à C₄)-O-benzoyle,
 (xiii) -(alkylène en C₁ à C₄)-CONH-(alkylène en C₁ à C₄)-NR¹²⁰R¹³⁰,
 (xiv) -(alkylène en C₁ à C₄)-COO-(alkylène en C₁ à C₄)-NR¹²⁰,R¹³⁰,
 (xv) -(alkylène en C₁ à C₄)-COO-amidinophényle,
 (xvi) -(alkylène en C₁ à C₄)-CONH-(alkyle en C₁ à C₄ qui est substitué par un ou deux -COOR¹¹⁰) (où R¹¹⁰ a la même signification que celle définie ci-dessus), ou
 (xvii) -(alkylène en C₁ à C₄)-CONR¹²⁰R¹³⁰.

(5) quand T est l'oxygène, le groupement :



est de formule (i) telle que définie ci-dessus,

E est une liaison simple,

A₀ est une liaison simple, alkylène en C₁ à C₄ ou vinylène qui est éventuellement substitué par un

ou deux groupes alkyle en C₁ à C₄, et

R est de formule (ii) telle que définie ci-dessus, alors R⁵⁰, R⁶⁰ et R⁷⁰ ne représentent pas l'hydrogène ;
et


(6) le composé de formula (IB) n'est pas le p-[(2-isopropyl-1-imidazolyl)méthyl]benzoate de p-amidinophényle ; ni le N-1-(S)-éthoxycarbonyl-2-benzoyloxyéthylamide de l'acide 4-(4-amidinophénoxy-carbonyl)- α -méthylcinnamique ;

et les sels non toxiques ou sels d'addition d'acide non toxiques d'un tel composé.

2. Composé selon la revendication 1, dans lequel R est :



ou




où les divers symboles sont tels que définis dans la revendication 1.

3. Composé selon la revendication 1, dans lequel R est :




ou




où les divers symboles sont tels que définis dans la revendication 1.

4. Composé selon la revendication 1, dans lequel R est :




où les divers symboles sont tels que définis dans la revendication 1.

5. Composé selon la revendication 1, dans lequel R est :




où les divers symboles sont tels que définis dans la revendication 1.

6. Composé selon la revendication 1, dans lequel



est un hétérocycle à 4 à 10 chaînons contenant un atome d'azote.

7. Composé selon la revendication 1, dans lequel



est un hétérocycle à 4 à 10 chaînons contenant deux atomes d'azote.

8. Composé selon la revendication 1, dans lequel R⁵⁰, R⁶⁰ et R⁷⁰, chacun indépendamment, est :

- (i) l'hydrogène,
- (ii) alkyle en C₁ à C₈,
- (iii) alcényle en C₂ à C₈,
- (iv) -COOR¹¹⁰,
- (v) -(alkylène en C₁ à C₈)-COOR¹¹⁰,
- (vi) -(alcénylène en C₂ à C₈)-COOR¹¹⁰,
- (vii) cycloalkyle en C₄ à C₇,
- (x) phényle,
- (xi) alkyle en C₁ à C₈ qui est substitué par un ou deux groupes phényle,
- (xii) -(alkylène en C₁ à C₄)-O-benzoyl,
- (xiii) -(alkylène en C₁ à C₄)-CONH-(alkylène en C₁ à C₄)NR¹²⁰R¹³⁰,
- (xiv) -(alkylène en C₁ à C₄)-COO-(alkylène en C₁ à C₄)NR¹²⁰R¹³⁰,
- (xv) -(alkylène en C₁ à C₄)-COO-amidinophényle,
- (xvi) -(alkylène en C₁ à C₄)-CONH-(alkyle en C₁ à C₄ qui est substitué par un ou deux -COOR¹¹⁰),
- (xvii) -(alkylène en C₁ à C₄)-CONR¹²⁰R¹³⁰, ou
- (xviii) (alcoxy en C₁ à C₄)alkyle en C₁ en C₄,

où les divers symboles sont tels que définis dans la revendication 1.

9. Composé selon la revendication 1, dans lequel R⁵⁰, R⁶⁰ et R⁷⁰, chacun indépendamment, est :

(viii) -(alkylène en C₁ à C₄)-(hétérocycle à 4 à 7 chaînons contenant un oxygène).

10. Composé selon la revendication 1, dans lequel R⁵⁰, R⁶⁰ et R⁷⁰, chacun indépendamment, est :

(ix) -(alkylène en C₁ à C₄)-(hétérocycle à 4 à 7 chaînons contenant un azote).

11. Composé selon la revendication 1, qui est :

le N-2-propényl-N-éthoxycarbonylméthylamide de l'acide 4-(4-amidinophénoxycarbonyl)phénoxyacétique,
le N-phényl-N-phénylméthylamide de l'acide 4-(6-amidino-7,8-dihydro-2-naphtyloxy-carbonyl)benzoïque,
le N-phényl-N-benzoyloxy-carbonylméthylamide de l'acide 4-(6-amidino-7,8-dihydro-2-naphtyloxy-carbonyl)benzoïque,
le N-phényl-N-éthoxycarbonylméthylamide de l'acide 4-(6-amidino-7,8-dihydro-2-naphtyloxy-carbonyl)benzoïque,
le N-phényl-N-éthoxycarbonylméthylamide de l'acide 4-[4-(2-amidinoéthényl)phénoxycarbonyl]benzoïque,
le N-1-(S)-éthoxycarbonyl-2-benzoyloxyéthylamide de l'acide 4-(4-amidinophénoxycarbonyl)-α-méthylcinnamique,
le N-1-(S)-3-bis(éthoxycarbonyl)propylamide de l'acide 4-[2-(4-amidinophénoxycarbonyl)éthényl]benzoïque,
le N-1-(S)-3-bis(éthoxycarbonyl)propylamide de l'acide 4-[2-(4-amidinophénoxycarbonyl)-1-propényl]benzoïque,
le N-[1,1,4-tris(éthoxycarbonyl)-3-butényl]amide de l'acide 4-(4-amidinophénoxycarbonyl)-α-méthylcinnamique,
le N-éthoxycarbonylméthyl-N-allylamide de l'acide 4-[4-(2-amidinoéthényl)phénoxycarbonyl]benzoïque,
le N-(1-(S)-3-bis(éthoxycarbonyl)propylamide de l'acide 4-(4-amidinophénoxycarbonyl)phénoxyacétique,
le N-1,1,2-tris(éthoxycarbonyl)éthylamide de l'acide 4-(4-amidinophénoxycarbonyl)-α-méthylcinnamique,
le N-éthoxycarbonylméthyl-N-allylamide de l'acide 4-[2'-(4-amidinophénoxycarbonyl)-1-propényl]benzoïque,
le N-1-(S)-3-bis(éthoxycarbonyl)propylamide de l'acide 5-[4-(4-amidinophénoxycarbonyl)phényl]-2,4-pentadiénoïque,
le N-éthoxycarbonylméthyl-N-allylamide de l'acide 5-[4-(4-amidinophénoxycarbonyl)phényl]-2,4-pentadiénoïque,
le N-[1,1-bis(éthoxycarbonyl)-3-butényl]amide de l'acide 4-(4-amidinophénoxycarbonyl)-α-méthylcinnamique,
le N-1-(S)-3-bis(éthoxycarbonyl)propylamide de l'acide 4-[2-(4-amidinophénoxycarbonyl)-1-méthyléthényl]benzoïque,
le N-éthoxycarbonylméthyl-N-allylamide de l'acide 4-[2-(4-amidinophénoxycarbonyl)-1-méthyléthényl]benzoïque,
le N-1-(S)-3-bis(éthoxycarbonyl)propylamide de l'acide 3-[4-(4-amidinophénoxycarbonyl)phényl]-2-éthoxycarbonyl-2-propénoïque,
le N-éthoxycarbonylméthyl-N-allylamide de l'acide 3-[4-(4-amidinophénoxycarbonyl)phényl]-2-éthoxycarbonyl-2-propénoïque,
le N-2-diméthylaminoéthylcarbamoyleméthyl-N-allylamide de l'acide 4-(4-amidinophénoxycarbonyl)-α-méthylcinnamique,
le N-2-diméthylaminoéthoxycarbonylméthyl-N-allylamide de l'acide 4-(4-amidinophénoxycarbonyl)-α-méthylcinnamique,
le N-4-amidinophénoxycarbonylméthyl-N-allylamide de l'acide 4-(4-amidinophénoxycarbonyl)-α-méthylcinnamique,
le N-1,1-bis(éthoxycarbonyl)méthylcarbamoyleméthyl-N-allylamide de l'acide 4-(4-amidinophénoxycarbonyl)-α-méthylcinnamique,
le N-(diallylcarbamoyle)méthyl-N-allylamide de l'acide 4-(4-amidinophénoxycarbonyl)-α-méthylcinnamique,
le N-allyl-N-carboxyméthylamide de l'acide 4-(4-amidiphénylcarbamoyle)-α-méthylcinnamique, ou
le N-1,1-bis(phénylméthyl)méthyl-N-3-phénylpropylamide de l'acide 4-(4-amidino-1-méthoxyphénoxycarbonyl)benzoïque.

12. Composé selon la revendication 1, qui est :

l'ester éthylique de l'acide 2-[4-(4-amidinophénoxy-carbonyl)phénylméthylimino]-2-(N-benzyl-N-méthylaminol) acétique, ou
l'ester éthylique de l'acide 2-[N-[4-(4-amidinophénoxy-carbonyl)phényl]méthyl-N-benzylamino]acétique.

13. Composé selon la revendication 1, qui est :

le 1-[4-(4-amidinophénoxy-carbonyl)phénylméthyl]-1-(3-phénylpropyl)phosphinate d'éthyle, ou
le 2-[4-(4-amidinophénoxy-carbonyl)phényl]éthénylphosphonate de diéthyle.

14. Composé selon la revendication 1, qui est le N-(2,3,4,5-tétrahydrofurane-2-yl)méthyl-N-carboxyméthylamide de l'acide 4-(4-amidinophénoxy-carbonyl)- α -méthylcinnamique.


15. Composé selon la revendication 1, qui est le N-(4-pipéridinylméthyl)-N-carboxyméthylamide de l'acide 4-(4-amidinophénoxy-carbonyl)- α -méthylcinnamique.

16. Composé selon la revendication 1, qui est : l'ester 4-amidinophénylique de l'acide 4-(4-phénylpipéridinylméthyl) benzoïque, ou
l'ester 4-amidinophénylique de l'acide 4-(2-éthoxycarbonylindolylméthyl)benzoïque.


17. Composé selon la revendication 1, qui est l'ester 4-aminodiphénylique de l'acide 4-(2-phénylimidazolylméthyl) benzoïque.

18. Composé selon l'une quelconque des revendications précédentes, sous la forme d'un sel d'addition d'acide non toxique.


19. Procédé pour la préparation d'un composé de formule (IB) :



qui comprend la réaction pour former une liaison ester ou amide, d'un composé de formule :



(dans laquelle les divers symboles sont tels que définis dans la revendication 1), avec un composé de formule :




(dans laquelle les divers symboles sont tels que définis dans la revendication 1) ; et, éventuellement, la conversion du dérivé d'amidinophénol ainsi obtenu en un sel ou sel d'addition d'acide de celui-ci.

20. Composition pharmaceutique qui comprend, en tant qu'ingrédient actif, un dérivé d'amidinophénol de formule (IB) décrit dans la revendication 1, ou un sel non toxique de celui-ci, ou un sel d'addition non toxique de celui-ci, avec un support ou un enrobage.

21. Utilisation d'un composé de formule (IB) selon l'une quelconque des revendications 1 à 19, ou d'un sel non toxique

de celui-ci ou d'un sel d'addition d'acide non toxique de celui-ci, dans la fabrication d'un médicament pour la prévention et/ou le traitement d'un état qui peut être amélioré par l'administration d'un antagoniste de leucotriène B₄ ou d'un inhibiteur de phospholipase A₂ ou de trypsine.

22. Utilisation d'un composé de formule (IA) :



(IA)

dans laquelle

R¹ et R², chacun indépendamment, est :

- (i) l'hydrogène, ou
- (ii) -COOR⁴ (où R⁴ est alkyle en C₁ à C₃) ;

A est

- (i) une liaison simple,
- (ii) alkylène en C₁ à C₄, ou
- (iii) -C(R⁵)=C(R⁶)- (où R⁵ et R⁶, chacune indépendamment, est l'hydrogène ou alkyle en C₁ à C₄) ;

R³ est

- (i) -CON(R⁷)R⁸,
- (ii) -CONR⁹-CH(R⁷)R⁸, ou
- (iii)



(où

R⁷ et R⁸, chacun indépendamment, est

- (1) l'hydrogène,
 - (2) phényle,
 - (3) -(alkylène en C₁ à C₄)phényle,
 - (4) -(alkylène en C₁ à C₄)phényle qui est substitué par un ou deux -R¹¹-COOR¹² (où R¹¹ est une liaison simple ou alkylène en C₁ à C₈, et R¹² est l'hydrogène ou alkyle en C₁ à C₄),
 - (5) alkyle en C₁ à C₅,
 - (6) alcényle en C₂ à C₁₀ contenant de une à trois doubles liaisons,
 - (7) -R^{11a}-COOR¹²
- où R^{11a} est


- (a) une liaison simple,
- (b) alkylène en C₁ à C₈,
- (c) alcénylène en C₂ à C₈, ou
- (d) alcénylène en C₂ à C₈ dans lequel un ou deux atomes de carbone de la chaîne principale sont remplacés par du soufre,

EP 0 703 216 B1

et R¹² a la même signification que celle définie ci-dessus), ou
(8) cycloalkyle en C₃ à C₇ ;

R⁹ est

- (1) l'hydrogène,
 - (2) -R¹¹-COOR¹² (où les divers symboles ont les mêmes significations que celles définies ci-dessus), ou
 - (3) alcoxyalkyle en C₂ à C₆ ;
- le groupement :



est un hétérocycle à 4 à 7 chaînons contenant un ou deux atomes d'azote ;

R¹⁰ est

- (1) l'hydrogène, ou
 - (2) -(alkylène en C₁ à C₄)phényle,
- avec les conditions suivantes :

- (1) R⁷ et R⁸ ne représentent pas en même temps l'hydrogène,
- (2) quand au moins un groupe dans R⁷, R⁸ et R⁹ représente le groupe contenant -COO-t-Bu, les autres groupes ne représentent pas les groupes contenant carboxy ;

et des sels non toxiques et sels d'addition d'acide non toxiques d'un tel composé, dans la fabrication d'un médicament pour la prévention et/ou le traitement de la polyarthrite rhumatoïde, des affections intestinales inflammatoires, du psoriasis, des maladies stomacales induites par un agent anti-inflammatoire non-stéroïdien, du syndrome de détresse respiratoire de l'adulte, de l'infarctus cardiaque, de la rhinite allergique, de la neutropénie induite par une hémodialyse ou de l'asthme anaphasique.