

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 705 276 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
02.09.1998 Bulletin 1998/36

(51) Int Cl.⁶: **C07K 14/705**, G01N 33/53,
A61K 38/17

(21) Application number: **94918699.3**

(86) International application number:
PCT/CA94/00319

(22) Date of filing: **09.06.1994**

(87) International publication number:
WO 95/00551 (05.01.1995 Gazette 1995/02)

(54) **PROSTAGLANDIN RECEPTOR FP AND DNA ENCODING IT**

PROSTAGLANDIN REZEPTOR FP UND KODIERENDE DNS

RECEPTEUR FP DE PROSTAGLANDINES ET ADN LE CODANT

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IE IT LI LU NL PT SE

(30) Priority: **25.06.1993 US 83741**

(43) Date of publication of application:
10.04.1996 Bulletin 1996/15

(73) Proprietor: **MERCK FROSST CANADA INC.**
Kirkland Quebec H9H 3L1 (CA)

(72) Inventors:

- **ABRAMOVITZ, Mark**
Dollard des Ormeaux, Quebec H9G 2X8 (CA)
- **GRYGORCZYK, Richard**
Dollard des Ormeaux, Quebec H9A 2P8 (CA)
- **METTERS, Kathleen**
Montreal, Quebec H2X 1W4 (CA)
- **NGUYEN, Truyen**
Dorval, Quebec H9P 1L3 (CA)
- **RUSHMORE, Thomas, H.**
Dollard des Ormeaux, Quebec H9B 2R9 (CA)
- **SLIPETZ, Deborah**
Outremont, Quebec H2V 1W5 (CA)

(74) Representative: **Thompson, John Dr. et al**
Merck & Co., Inc.
European Patent Department
Terlings Park
Eastwick Road
Harlow, Essex CM20 2QR (GB)

(56) References cited:
WO-A-94/07920

- JPN. J. PHARMACOL., SUPPL. 1 vol. 61, 1993
page 85P KITANAKA, J. ET AL. 'abstract O-75'
- J. BIOL. CHEM. vol. 269, no. 4, 28 January 1994
pages 2632 - 2636 ABRAMOVITZ, M. ET AL.
'Cloning and expression of a cDNA for the
human prostanoid FP receptor'
- J. BIOL. CHEM. vol. 269, no. 2, 14 January 1994
pages 1356 - 1360 SUGIMOTO, Y. ET AL. 'Cloning
and expression of a cDNA for mouse
prostaglandin F receptor'
- J. BIOL. CHEM. vol. 269, no. 5, 4 February 1994
pages 3881 - 3886 SAKAMOTO, K. ET AL.
'Molecular cloning and expression of a cDNA of
the bovine prostaglandin F2alpha receptor'

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 705 276 B1

Description**BACKGROUND OF THE INVENTION**

The physiological actions of prostaglandin (PG) $F_{2\alpha}$ are mediated through interaction with the prostaglandin $F_{2\alpha}$ (FP) receptor. This receptor has not previously been isolated or purified. FP-encoding DNA and the amino acid sequence of the FP receptor protein was also not known. FP receptors are normally found on a wide variety of cells and tissues including the small intestine, corpus luteum, placenta, ovary, brain, myometrium, lung, kidney, stomach, muscle, eye, uterus and trachea, in humans and other animals. Binding of prostaglandin to the FP receptor protein elicits an increase in intracellular calcium levels. This signal causes the tissues to respond, for example, by muscle contraction and in the eye indirectly causes a reduction in intraocular pressure. Studies on $PGF_{2\alpha}$ binding sites (FP receptors) have been performed using primarily corpus lutea tissue since $PGF_{2\alpha}$ is a potent luteolytic agent [Powell *et al.*, 1974 *Lancet*, **1**, pp 1120; Powell *et al.*, 1974, *Eur. J. Biochem.*, **41**, pp 103-107]. Functional activities of the FP receptor have been studied using tissue preparations such as rabbit jejunum and the cat, bullock and dog iris sphincter tissues [Dong and Jones, 1982 *Br. J. Pharmacol.*, **76**, pp 149-155; Welburn and Jones, 1978 *Prostaglandins*, **15**, pp 287]. The above methods for studying FP receptor activities have several disadvantages in that they require tissue preparations containing several different but related receptor populations with different ligand binding properties making absolute potency and selectivity impossible. In addition, tissues contain very low levels of FP receptor and since tissue samples are required, compounds cannot satisfactorily be tested as effectors of the human FP receptor.

SUMMARY OF THE INVENTION

A novel prostaglandin receptor protein termed FP has been identified from human cells. A DNA molecule encoding the full length FP protein has been isolated and purified, and the nucleotide sequence has been determined. The FP encoding DNA has been cloned into expression vectors and these expression vectors, when introduced into recombinant host cells, cause the recombinant host cells to express a functional FP receptor protein. The novel FP protein, the FP-encoding DNA, the expression vectors and recombinant host cells expressing recombinant FP are useful in the identification of modulators of FP receptor activity.

A method of identifying FP receptor modulators is also disclosed which utilizes the recombinant FP expressing host cells. Modulators of FP activity are useful for the treatment of prostaglandin-related diseases and for modulating the effects of prostaglandins on the FP receptor.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1A-B - The complete DNA sequence encoding the FP receptor protein is shown.

Figure 2 - The complete deduced amino acid sequence of the FP receptor protein is shown.

Figure 3A-B - Expression of the prostaglandin $F_{2\alpha}$ receptor in FP cDNA-injected *Xenopus* oocytes. An inward Ca^{2+} -dependent Cl^- current (shown as downward deflection) evoked by bath perfusion of 20 nM $PGF_{2\alpha}$ (upper trace) and 10 nM of fluprostenol (lower trace). The oocyte was injected with 1 ng FP cDNA and voltage-clamped at -60 mV.

Figure 4A-C - $PGF_{2\alpha}$ -induced light responses in aequorin loaded oocytes expressing recombinant FP receptor. Superimposed individual responses of 5 oocytes tested at each $PGF_{2\alpha}$ concentration. The ligand was added into the recording cuvette at 10 s; the aequorin light emission is expressed in relative units with the background emission being typically 0.5-0.7 units.

Figure 5 - Average light responses evoked by different concentrations of $PGF_{2\alpha}$, fluprostenol and PGE_2 . Each bar represents average response from 5 oocytes obtained from the same donor and the numbers above each bar indicate the number of oocytes responding. Similar results were obtained with oocytes from 5 other donors.

Figure 6 - Competition for [3H] $PGF_{2\alpha}$ specific binding to pcDNA1amp-hFP transfected COS-M6 membranes. [3H] $PGF_{2\alpha}$ binding assays were performed as described in the Methods in the presence of 0.03 nM-10 μ M $PGF_{2\alpha}$ (■), fluprostenol (●), PGD_2 (□), PGE_2 (○), U46619 (Δ) and iloprost (◇).

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to cDNA encoding a novel prostaglandin receptor termed FP. The present invention is also related to recombinant host cells which express the cloned FP-encoding DNA contained in a recombinant expression plasmid. The present invention is also related to a method for the screening of substances which modulate FP receptor activity. The DNA of the present invention is isolated from FP producing cells. FP, as used herein, refers to a G protein-coupled receptor which can specifically bind prostaglandin molecules.

Mammalian cells capable of producing FP include, but are not limited to, cells derived from small intestine, kidney,

stomach, muscle, eye, placenta, uterus and trachea. Transformed mammalian cell lines which produce FP include, but are not limited to, 3T3 fibroblasts cells. The preferred cells for the present invention include normal human kidney and placental cells and the most preferred cells are human corpora lutea cells.

Other cells and cell lines may also be suitable for use to isolate FP cDNA. Selection of suitable cells may be done by screening for FP on cell surfaces. Methods for detecting FP activity are well known in the art and measure the binding of radiolabelled ligand specific for the receptor. Cells which possess FP activity in this assay may be suitable for the isolation of FP cDNA.

Any of a variety of procedures may be used to clone FP cDNA. These methods include, but are not limited to, direct functional expression of the FP cDNA following the construction of an FP-containing cDNA library in an appropriate expression vector system. Another method is to screen an FP-containing cDNA library constructed in a bacteriophage or plasmid shuttle vector with a labelled oligonucleotide probe designed from the amino acid sequence of the FP protein. The preferred method consists of screening an FP-containing cDNA library constructed in a bacteriophage or plasmid shuttle vector with a partial cDNA encoding the FP protein. This partial cDNA is obtained by the specific PCR amplification of FP DNA fragments through the design of degenerate oligonucleotide primers from the amino acid sequence known for other G protein-coupled receptors which are related to the prostaglandin FP receptors.

It is readily apparent to those skilled in the art that other types of libraries, as well as libraries constructed from other cells or cell types, may be useful for isolating FP-encoding DNA. Other types of libraries include, but are not limited to, cDNA libraries derived from other cells or cell lines and genomic DNA libraries.

It is readily apparent to those skilled in the art that suitable cDNA libraries may be prepared from cells or cell lines which have FP activity. The selection of cells or cell lines for use in preparing a cDNA library to isolate FP cDNA may be done by first measuring cell associated FP activity using the known labelled ligand binding assay cited above and used herein.

Preparation of cDNA libraries can be performed by standard techniques well known in the art. Well known cDNA library construction techniques can be found for example, in Maniatis, T., Fritsch, E.F., Sambrook, J., *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1982).

It is also readily apparent to those skilled in the art that DNA encoding FP may also be isolated from a suitable genomic DNA library. Construction of genomic DNA libraries can be performed by standard techniques well known in the art. Well known genomic DNA library construction techniques can be found in Maniatis, T., Fritsch, E.F., Sambrook, J. in *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1982).

In order to clone the FP gene by one of the preferred methods, the amino acid sequence or DNA sequence of FP or a homologous protein is necessary. To accomplish this, FP protein or a homologous protein may be purified and partial amino acid sequence determined by automated sequenators. It is not necessary to determine the entire amino acid sequence, but the linear sequence of two regions of 6 to 8 amino acids can be determined for the PCR amplification of a partial FP DNA fragment.

Once suitable amino acid sequences have been identified, the DNA sequences capable of encoding them are synthesized. Because the genetic code is degenerate, more than one codon may be used to encode a particular amino acid, and therefore, the amino acid sequence can be encoded by any of a set of similar DNA oligonucleotides. Only one member of the set will be identical to the FP sequence but others in the set will be capable of hybridizing to FP DNA even in the presence of DNA oligonucleotides with mismatches. The mismatched DNA oligonucleotides may still sufficiently hybridize to the FP DNA to permit identification and isolation of FP encoding DNA.

Using one of the preferred methods, cDNA clones encoding FP are isolated in a two-stage approach employing polymerase chain reaction (PCR) based technology and cDNA library screening. In the first stage, NH₂-terminal and internal amino acid sequence information from the purified FP or a homologous protein is used to design degenerate oligonucleotide primers for the amplification of FP-specific DNA fragments. In the second stage, these fragments are cloned to serve as probes for the isolation of full length cDNA from a cDNA libraries.

The sequence for the near full-length cDNA encoding FP is shown in Table 1, and was designated clone FP. The deduced amino acid sequence of FP from the cloned cDNA is shown in Table 2. Inspection of the determined cDNA sequence reveals the presence of a single, large open reading frame that encodes for a 359 amino acid protein.

The cloned FP cDNA obtained through the methods described above may be recombinantly expressed by molecular cloning into an expression vector containing a suitable promoter and other appropriate transcription regulatory elements, and transferred into prokaryotic or eukaryotic host cells to produce recombinant FP. Techniques for such manipulations can be found described in Maniatis, T. *et al.*, *supra*, and are well known in the art.

Expression vectors are defined herein as DNA sequences that are required for the transcription of cloned DNA and the translation of their mRNAs in an appropriate host. Such vectors can be used to express eukaryotic DNA in a variety of hosts such as bacteria, bluegreen algae, plant cells, insect cells and animal cells.

Specifically designed vectors allow the shuttling of DNA between hosts such as bacteria-yeast or bacteria-animal cells. An appropriately constructed expression vector should contain: an origin of replication for autonomous replication in host cells, selectable markers, a limited number of useful restriction enzyme sites, a potential for high copy number,

and active promoters. A promoter is defined as a DNA sequence that directs RNA polymerase to bind to DNA and initiate RNA synthesis. A strong promoter is one which causes mRNAs to be initiated at high frequency. Expression vectors may include, but are not limited to, cloning vectors, modified cloning vectors, specifically designed plasmids or viruses.

A variety of mammalian expression vectors may be used to express recombinant FP in mammalian cells. Commercially available mammalian expression vectors which may be suitable for recombinant FP expression, include but are not limited to, pMC1neo (Stratagene), pXT1 (Stratagene), pSG5 (Stratagene), pcDNA1, pcDNA1amp (Invitrogen), EBO-pSV2-neo (ATCC 37593) pBPV-1(8-2) (ATCC 37110), pdBPV-MMTneo(342-12) (ATCC 37224), pRSVgpt (ATCC 37199), pRSVneo (ATCC 37198), pSV2-dhfr (ATCC 37146), pUCTag (ATCC 37460), and 1ZD35 (ATCC 37565)

DNA encoding FP may also be cloned into an expression vector for expression in a host cell. Host cells may be prokaryotic or eukaryotic, including but not limited to bacteria, yeast, mammalian cells including but not limited to cell lines of human, bovine, porcine, monkey and rodent origin, and insect cells including but not limited to drosophila derived cell lines. Cell lines derived from mammalian species which may be suitable and which are commercially available, include but are not limited to, CV-1 (ATCC CCL 70), COS-1 (ATCC CRL 1650), COS-7 (ATCC CRL 1651), CHO-K1 (ATCC CCL 61), 3T3 (ATCC CCL 92), NIH/3T3 (ATCC CRL 1658), HeLa (ATCC CCL 2), C1271(ATCC CRL 1616), BS-C-1 (ATCC CCL 26) and MRC-5 (ATCC CCL 171).

The expression vector may be introduced into host cells via any one of a number of techniques including but not limited to transformation, transfection, protoplast fusion, and electroporation. The expression vector-containing cells are individually analyzed to determine whether they produce FP protein. Identification of FP expressing cells may be done by several means, including but not limited to immunological reactivity with anti-FP antibodies, and the presence of host cell-associated FP activity.

Expression of FP DNA may also be performed using *in vitro* produced synthetic mRNA. Synthetic mRNA can be efficiently translated in various cell-free systems, including but not limited to wheat germ extracts and reticulocyte extracts, as well as efficiently translated in cell based systems, including but not limited to microinjection into frog oocytes, with microinjection into frog oocytes being preferred.

To determine the FP cDNA sequence(s) that yields optimal levels of receptor activity and/or FP protein, FP cDNA molecules including but not limited to the following can be constructed: the full-length open reading frame of the FP cDNA and various constructs containing portions of the cDNA encoding only specific domains of the receptor protein or rearranged domains of the protein. All constructs can be designed to contain none, all or portions of the 5' and/or 3' untranslated region of FP cDNA. FP activity and levels of protein expression can be determined following the introduction, both singly and in combination, of these constructs into appropriate host cells. Following determination of the FP cDNA cassette yielding optimal expression in transient assays, this FP cDNA construct is transferred to a variety of expression vectors (including recombinant viruses), including but not limited to those for mammalian cells, plant cells, insect cells, oocytes, *E. coli*, and yeast cells.

Mammalian cell transfectants are assayed for both the levels of FP receptor activity and levels of FP protein by the following methods. Assessing FP receptor activity involves the direct introduction of a labelled ligand to the cells and determining the amount of specific binding of the ligand to the FP-expressing cells. Binding assays for receptor activity are known in the art (Frey *et al.*, 1993, Eur. J. Pharmacol., 244, pp 239-250).

Levels of FP protein in host cells is quantitated by a variety of techniques including, but not limited to, immunoaffinity and/or ligand affinity techniques. FP-specific affinity beads or FP-specific antibodies are used to isolate ³⁵S-methionine labelled or unlabelled FP protein. Labelled FP protein is analyzed by SDS-PAGE. Unlabelled FP protein is detected by Western blotting, ELISA or RIA assays employing FP specific antibodies.

Following expression of FP in a host cell, FP protein may be recovered to provide FP in active form, capable of binding FP-specific ligands. Several FP purification procedures are available and suitable for use. Recombinant FP may be purified from cell membranes by various combinations of, or individual application of standard separation techniques including but not limited to detergent solubilization, salt fractionation, ion exchange chromatography, size exclusion chromatography, hydroxylapatite adsorption chromatography and hydrophobic interaction chromatography.

In addition, recombinant FP can be separated from other cellular proteins by use of an immuno-affinity column made with monoclonal or polyclonal antibodies specific for full length nascent FP, or polypeptide fragments of FP.

Monospecific antibodies to FP are purified from mammalian antisera containing antibodies reactive against FP or are prepared as monoclonal antibodies reactive with FP using the technique of Kohler and Milstein, Nature 256: 495-497 (1975). Monospecific antibody as used herein is defined as a single antibody species or multiple antibody species with homogenous binding characteristics for FP. Homogenous binding as used herein refers to the ability of the antibody species to bind to a specific antigen or epitope, such as those associated with the FP, as described above. FP specific antibodies are raised by immunizing animals such as mice, rats, guinea pigs, rabbits, goats, horses and the like, with an appropriate concentration of FP or a peptide derived from the sequence of the FP protein either with or without an immune adjuvant.

Preimmune serum is collected prior to the first immunization. Each animal receives between about 0.1 mg and

about 1000 mg of FP or FP-related peptide associated with an acceptable immune adjuvant. Such acceptable adjuvants include, but are not limited to, Freund's complete, Freund's incomplete, alum-precipitate, water in oil emulsion containing *Corynebacterium parvum* and tRNA. The initial immunization consisted of the enzyme in, preferably, Freund's complete adjuvant at multiple sites either subcutaneously (SC), intraperitoneally (IP) or both. Each animal is bled at regular intervals, preferably weekly, to determine antibody titer. The animals may or may not receive booster injections following the initial immunization. Those animals receiving booster injections are generally given an equal amount of FP or FP-related peptide in Freund's incomplete adjuvant by the same route. Booster injections are given at about three week intervals until maximal titers are obtained. At about 7 days after each booster immunization or about weekly after a single immunization, the animals are bled, the serum collected, and aliquots are stored at about -20°C.

Monoclonal antibodies (mAb) reactive with FP or a peptide derived from the sequence of the FP protein are prepared by immunizing inbred mice, preferably Balb/c, with FP or FP-related peptide. The mice are immunized by the IP or SC route with about 1 mg to about 100 mg, preferably about 10 mg, of FP or FP-related peptide in about 0.5 ml buffer or saline incorporated in an equal volume of an acceptable adjuvant, as discussed above. Freund's complete adjuvant is preferred. The mice receive an initial immunization on day 0 and are rested for about 3 to about 30 weeks. Immunized mice are given one or more booster immunizations of about 1 to about 100 mg of FP in a buffer solution such as phosphate buffered saline by the intravenous (IV) route. Lymphocytes, from antibody positive mice, preferably splenic lymphocytes, are obtained by removing spleens from immunized mice by standard procedures known in the art. Hybridoma cells are produced by mixing the splenic lymphocytes with an appropriate fusion partner, preferably myeloma cells, under conditions which will allow the formation of stable hybridomas. Fusion partners may include, but are not limited to: mouse myelomas P3/NS1/Ag 4-1; MPC-11; S-194 and Sp 2/0, with Sp 2/0 being preferred. The antibody producing cells and myeloma cells are fused in polyethylene glycol, about 1000 mol. wt., at concentrations from about 30% to about 50%. Fused hybridoma cells are selected by growth in hypoxanthine, thymidine and aminopterin supplemented Dulbecco's Modified Eagles Medium (DMEM) by procedures known in the art. Supernatant fluids are collected from growth positive wells on about days 14, 18, and 21 and are screened for antibody production by an immunoassay such as solid phase immunoradioassay (SPIRA) using FP or FP-related peptide as the antigen. The culture fluids are also tested in the Ouchterlony precipitation assay to determine the isotype of the mAb. Hybridoma cells from antibody positive wells are cloned by a technique such as the soft agar technique of MacPherson, Soft Agar Techniques, in Tissue Culture Methods and Applications, Kruse and Paterson, Eds., Academic Press, 1973.

Monoclonal antibodies are produced *in vivo* by injection of pristine primed Balb/c mice, approximately 0.5 ml per mouse, with about 2×10^6 to about 6×10^6 hybridoma cells about 4 days after priming. Ascites fluid is collected at approximately 8-12 days after cell transfer and the monoclonal antibodies are purified by techniques known in the art.

In vitro production of anti-FP mAb is carried out by growing the hybridoma in DMEM containing about 2% fetal calf serum to obtain sufficient quantities of the specific mAb. The mAb are purified by techniques known in the art.

Antibody titers of ascites or hybridoma culture fluids are determined by various serological or immunological assays which include, but are not limited to, precipitation, passive agglutination, enzyme-linked immunosorbent antibody (ELISA) technique and radioimmunoassay (RIA) techniques. Similar assays are used to detect the presence of FP in body fluids or tissue and cell extracts.

It is readily apparent to those skilled in the art that the above described methods for producing monospecific antibodies may be utilized to produce antibodies specific for FP polypeptide fragments, or full-length FP polypeptide.

FP antibody affinity columns are made by adding the antibodies to Affigel-10 (Biorad), a gel support which is pre-activated with N-hydroxysuccinimide esters such that the antibodies form covalent linkages with the agarose gel bead support. The antibodies are then coupled to the gel via amide bonds with the spacer arm. The remaining activated esters are then quenched with 1M ethanolamine HC1 (pH 8). The column is washed with water followed by 0.23 M glycine HC1 (pH 2.6) to remove any non-conjugated antibody or extraneous protein. The column is then equilibrated in phosphate buffered saline (pH 7.3) together with appropriate membrane solubilizing such as detergents and the cell culture supernatants or cell extracts containing FP or FP fragments are slowly passed through the column. The column is then washed with phosphate buffered saline together with appropriate membrane solubilizing such as detergents until the optical density (A_{280}) falls to background, then the protein is eluted with 0.23 M glycine-HCl (pH 2.6) together with appropriate membrane solubilizing such as detergents. The purified FP protein is then dialyzed against phosphate buffered saline together with appropriate membrane solubilizing such as detergents.

One method suitable for the isolation of DNA encoding the prostaglandin receptor of the present invention involves the utilization of amino acid and/or DNA sequence information obtained from other G-protein-linked receptors. Since other prostaglandin receptors are known to be G-protein linked, certain regions or domains such as the transmembrane and/or cytoplasmic domains, are expected to have some degree of homology sufficient to produce a probe for the isolation of novel receptors.

Prostaglandins and leukotrienes are known to transduce their signals via G-protein-linked receptors. Distinct receptors for PGH_2 /thromboxane A_2 , PGI_2 , PGE_2 , PGD_2 , $\text{PGF}2\alpha$, LTB_4 , and LTD_4 present in various tissues have been described. Some of the receptors have been solubilized and partially purified (Dutta-Roy, A.K. *et al.*, (1987) JBC, 262,

pp. 12685; Tsai, A.L. *et al.*, (1989), JBC, 264, pp 61; 168 - Watawabe, T. *et al.*, (1990), JBC, 265, pp. 21237) and the human platelet TXA₂ receptor has been purified to apparent homogeneity (Ushikubi, F. *et al.*, (1989), JBC, 264, pp. 16496). The purified thromboxane receptor exhibited a very broad band on a SDS-polyacrylamide gel centered at 3 57 kDa. Enough protein was obtained for partial sequence information.

5 An approach to the isolation of other eicosanoid receptor genes by homology screening was taken, with the assumption that these receptors are related in primary structure (Sugimoto, Y. *et al.*, (1992), JBC, 267, pp. 6463). Since these receptors are of the G-protein-coupled receptor superfamily there are areas of homology which are likely to be found in the transmembrane region and in the cytoplasmic domains. Therefore, various known G-protein linked receptors related to the prostaglandin receptors may be utilized to provide DNA probes to regions of the receptor protein-
10 encoding DNA sought, which is likely to have homology, such as the transmembrane region.

Using a 0.37-kb fragment of a putative mouse FP receptor cDNA which encodes most of the transmembrane 5-7 region of this receptor was used to screen a human kidney library from which a partial human FP cDNA was isolated. This in turn was used to obtain a 2.5-kb cDNA clone hereinafter designated FP encoding a 359-amino acid receptor was isolated from a human uterus cDNA library. This protein was designated as the FP receptor. Like many other G-
15 protein coupled receptors the FP receptor shares several common features. Firstly, there are 3 potential N-linked glycosylation sites (Asn4, Asn19) and Asn277 in the putative extracellular amino terminus. Secondly, conserved cysteine residues are found in exofacial loops 1 and 2. There are multiple serine residues, potential sites of protein kinase phosphorylation, throughout the C-terminus and third cytoplasmic loops. The FP receptor does not contain an aspartic acid residue in transmembrane three which is characteristic of the receptors binding cationic amino-containing
20 ligands, however, it possesses a conserved arginine (position 295) found in all known eicosanoid receptors within transmembrane seven. This region is the most highly conserved among the eicosanoid receptors.

The novel prostaglandin receptor of the present invention is suitable for use in an assay procedure for the identification of compounds which modulate the receptor activity. Modulating receptor activity, as described herein includes the inhibition or activation of the receptor and also includes directly or indirectly affecting the normal regulation of the
25 receptor activity. Compounds which modulate the receptor activity include agonists, antagonists and compounds which directly or indirectly affect regulation of the receptor activity.

The prostaglandin receptor of the present invention may be obtained from both native and recombinant sources for use in an assay procedure to identify receptor modulators. In general, an assay procedure to identify prostaglandin receptor modulators will contain the prostaglandin receptor of the present invention, and a test compound or sample
30 which contains a putative prostaglandin receptor modulator. The test compounds or samples may be tested directly on, for example, purified receptor protein whether native or recombinant, subcellular fractions of receptor-producing cells whether native or recombinant, and/or whole cells expressing the receptor whether native or recombinant. The test compound or sample may be added to the receptor in the presence or absence of a known labelled or unlabelled receptor ligand. The modulating activity of the test compound or sample may be determined by, for example, analyzing
35 the ability of the test compound or sample to bind to the receptor, activate the receptor, inhibit receptor activity, inhibit or enhance the binding of other compounds to the receptor, modify receptor regulation, or modify an intracellular activity.

The identification of modulators of FP receptor activity are useful in treating disease states involving the FP receptor activity. Other compounds may be useful for stimulating or inhibiting activity of the receptor. Selective agonists of the FP receptor may be of use in the treatment of glaucoma through their ability to lower intraocular pressure and may have
40 utility in the synchronization of oestrus cycles in farm animals through their ability to stimulate luteolytic function. Compounds which antagonize the FP receptor could be of use in the treatment of diseases in which activation of the FP receptor results in either cellular proliferation, induction of cellular neoplastic transformations or metastatic tumor growth or pathological states where activation of the FP receptor causes smooth muscle contraction such as the uterine contractions observed in dysmenorrhea. The isolation and purification of an FP-encoding DNA molecule would be
45 useful for establishing the tissue distribution of FP receptors, studying changes in FP receptor expression in disease states, as well as establishing a process for identifying compounds which modulate FP receptor activity.

The following examples are provided for the purpose of illustrating the present invention without, however, limiting the same thereto.

50 **EXAMPLE 1**

Cloning of the FP cDNA

An antisense 16-fold degenerate 27mer oligonucleotide [5'-ATA(A,C)ACCCAGGG(A,G)TCCA(A,G)GATCTG(G,
55 A)TT-3'] (SEQ.ID.NO.:1) based on the 9 conserved amino acids (NQILDPPWVY) (SEQ.ID.NO.:2) in transmembrane domain VII was synthesized. The ³²P-labeled oligo probe was initially used to screen a mouse kidney lambda gt10 library (Clontech, Palo Alto, CA) using standard techniques (Sambrook *et al.*, 1989. Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.). A putative mouse FP partial cDNA (430

bp) was cloned and sequenced. PCR was then used to generate a 373 bp cDNA probe (see below), based on the mouse sequence, to be used to screen a human kidney lambda gt11 library (Clontech, Palo Alto, CA).

The sequence of the 373 bp PCR generated probe from a putative mouse FP partial cDNA clone is as follows:

CTCTTAGCTCTTGGTGTTCCTTCTCGTGCAATGCCGTACGGGAGTCAC
 ACTCTTAAGAGTGAAGTTCAGAAGCCAGCAGCATAGGCAAGGCAGATCT
 CACCACCTGGAGATGATCATTGAGCTCCTGGCCATAATGTGCGTCTCCTG
 CGTCTGCTGGAGTCCCTTTCTGGTAACAATGGCCAACATTGCAATAAATG
 GAAATAATTCCCCAGTGACCTGTGAAACGACACTTTTGTCTCTCCGCATG
 GCAACGTGGAATCAGATCTTAGATCCCTGGGTCTATATTCTGCTACGGAA
 GGCTGTCCTTAGGAACCTGTATAAACTTGCCAGTCGTTGCTGTGGAGTTA
 ACATCATCAGCTTGCATATCTGGG (SEQ.ID.NO.: 3).

In this manner, two partial length human FP cDNA clones were obtained of approx. 1.7 and 1.8 kb in length. A 1 kb EcoRI 5' fragment from one of the clones was purified, ³²P-labeled and subsequently used to probe a human uterus lambda gt10 library (Clontech, Palo Alto, CA). From this screening a 2.8 kb cDNA clone was plaque-purified and DNA was prepared by the plate lysate method (Sambrook *et al.*, 1989. Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.).

Subcloning and sequencing of cDNA

The 2.5 kb cDNA clone was digested with EcoRI and was found to contain two inserts of sizes 1.8 kb and 0.7 kb. Only the 1.8 kb insert was found to hybridize with the human receptor partial cDNA probe upon Southern blot analysis. The 1.8 kb EcoRI fragment (FP) was subcloned into pKS vector (Stratagene, La Jolla, CA) for sequencing using the T7 DNA polymerase sequencing kit (Pharmacia). The DNA was sequenced entirely on both strands using the KS and SK primers (Stratagene, La Jolla, CA) or primers generated from the determined sequence. The nucleotide sequence of FP is shown in Table 1. The amino acid sequence for the encoded protein is shown in Table 2. The 1.8 kb fragment (FP; Fig. 1), when sequenced, was found to contain sequence homology to the human thromboxane receptor cDNA and the putative heptahelical arrangement characteristic of G protein-coupled receptors was evident. A long open reading frame (1077 bp) was determined which would result in a 359 amino acid polypeptide with a predicted relative molecular mass of 40,060. The ATG assigned as the initiator codon matches the Kozak consensus sequence for translation initiation (Kozak, 1989 J. Cell. Biol., 108, pp 229-241). The FP cDNA contains a long 3' untranslated region of about 1200 base pairs.

EXAMPLE 2

Constuction of pcDNAIamp-FP expression vector

The 1.8 Kb EcoRI human FP cDNA fragment was subcloned into the EcoRI site of pcDNAIamp and the correct orientation was verified by Pst I digestion.

TABLE 1

5 GTGCGCGGAGGGGACGAGCGGCTGGACCACAGCCGGCGCCCGATCAGGATCTCCGCG
 CTGGGATCGGTGGAACCTGAGGCAGCGGCGGGCGGCCATGGCACACCGAGCG
 GCTCCGTCTTCTGCTCCTCAGAGAGCCCGGCTGGCGGCCTGGGATGACAAGATGTCT
 GGACTGCAATCCTGCACAGTTTTGAGAGGGAGATGACTTGAGTGGTTGGCTTTTATC
 10 TCCACAACAATGTCCATGAACAATTCCAAACAGCTAGTGTCTCCTGCAGCTGCGCTT
 CTTTCAAACACAACCTGCCAGACGGAAAACCGGCTTTCCGTATTTTTTTCAGTAATC
 TTCATGACAGTGGGAATCTTGTCAAACAGCCTTGCCATCGCCATTCTCATGAAGGCA
 TATCAGAGATTTAGACAGAAGTCCAAGGCATCGTTTCTGCTTTTGCCAGCGGCCTG
 GTAATCACTGATTTCTTTGGCCATCTCATCAATGGAGCCATAGCAGTATTTGTATAT
 GCTTCTGATAAAGAATGGATCCGCTTTGACCAATCAAATGTCCTTTGCAGTATTTTT
 15 GGTATCTGCATGGTGTCTTCTGGTCTGTGCCCCTTCTTCTAGGCAGTGTGATGGCC
 ATTGAGCGGTGTATTGGAGTCAACAAACCAATATTTTATTCTACGAAAATTACATCC
 AAACATGTGAAAATGATGTTAAGTGGTGTGTGCTTGTGTTTTCATAGCTTTG
 CTGCCCATCCTTGGACATCGAGACTATAAAATTCAGGCGTCGAGGACCTGGTGTTC
 TACAACACAGAAGACATCAAAGACTGGGAAGATAGATTTTATCTTCTACTTTTTCT
 20 TTTCTGGGGCTCTTAGCCCTTGGTGTTCATTGTTGTGCAATGCAATCACAGGAATT
 AACTTTTAAAGAGTTAAATTTAAAGTCAGCAGCACAGACAAGGCAGATCTCATCAT
 TTGGAAATGGTAATCCAGCTCCTGGCGATAATGTGTGTCTCCTGTATTTGTTGGAGC
 CCATTTCTGGTTACAATGGCCAACATTGGAATAAATGGAATCATTCTCTGGAACC
 TGTGAAACAACACTTTTTGCTCTCCGAATGGCAACATGGAATCAAATCTTAGATCCT
 25 TGGGTATATATTCTTCTACGAAAGGCTGTCCTTAAGAATCTCTATAAGCTTGCCAGT
 CAATGCTGTGGAGTGCATGTCATCAGCTTACATATTTGGGAGCTTAGTTCCATTAAA
 AATTCTTAAAGGTTGCTGCTATTTCTGAGTCACCAGTTGCAGAGAAATCAGCAAGC
 ACCTAGCTTAATAGGACAGTAAATCTGTGTGGGGCTAGAACAAAAATTAAGACATGT
 30 TTGGCAATATTTTCAAGTTAGTTAAATACCTGTAGCCTAACTGGAAAATTCAGGCTTCA
 TCATGTAGTTTG (SEQ.ID.NO.: 4)

TABLE 2

35 MSMNNSKQLVSPAAALLSNTTCQTENRLSVFFSVIFMTVGILSNSLAIAILMKAYQR
 FRQSKASFLLLASGLVITDFFGHILINGAIAVFVYASDKEWIRFDQSNVLCISIFGIC
 40 MVFSGLCPLLLGSVMAIERCIGVTKPIFHSTKITSKHVKMMLSGVCLFAVFIALLP
 LGHRDYKIQASRTWCFYNTEDIKDWDREFYLLLFSLGLLALGVSLLCNAITGITLL
 RVKFKSQQHRQGRSHHLEMVIQLLAIMCVSCICWSPFLVTMANIGINGNHSLETCET
 45 TLFALRMATWNQILDWPVYILLRKAVLKNLYKLASQCCGVHVISLHIWELSSIKNSL
 KVA AIS ESPVAEKSAST (SEQ.ID.NO.: 5)

EXAMPLE 3

50 Cloning of the FP cDNA into E. coli Expression Vectors

Recombinant FP is produced in E. coli following the transfer of the FP expression cassette into E. coli expression
 vectors, including but not limited to, the pET series (Novagen). The pET vectors place FP expression under control of
 55 the tightly regulated bacteriophage T7 promoter. Following transfer of this construct into an E. coli host which contains
 a chromosomal copy of the T7 RNA polymerase gene driven by the inducible lac promoter, expression of FP is induced
 when an appropriate lac substrate (IPTG) is added to the culture. The levels of expressed FP are determined by the
 assays described above.

The cDNA encoding the entire open reading frame for FP is inserted into the NdeI site of pET 11a. Constructs in the positive orientation are identified by sequence analysis and used to transform the expression host strain BL21. Transformants are then used to inoculate cultures for the production of FP protein. Cultures may be grown in M9 or ZB media, whose formulation is known to those skilled in the art. After growth to an approximate $OD_{600} = 1.5$, expression of FP is induced with 1 mM IPTG for 3 hours at 37°C. FP receptor binding activity will be found in membrane fractions from these cells.

EXAMPLE 4

In Vivo Translation of Synthetic FP mRNA by *Xenopus* Oocyte Microinjection and Expression in Mammalian Cells

FP cDNA constructs are ligated into in vitro transcription vectors (the pGEM series, Promega) for the production of synthetic mRNAs.

Synthetic mRNA is produced in sufficient quantity in vitro by cloning double stranded DNA encoding FP mRNA into a plasmid vector containing a bacteriophage promoter, linearizing the plasmid vector containing the cloned FP-encoding DNA, and transcribing the cloned DNA in vitro using a DNA-dependent RNA polymerase from a bacteriophage that specifically recognizes the bacteriophage promoter on the plasmid vector.

Various plasmid vectors are available containing a bacteriophage promoter recognized by a bacteriophage DNA-dependent RNA polymerase, including but not limited to plasmids pSP64, pSP65, pSP70, pSP71, pSP72, pSP73, pGEM-3Z, pGEM-4Z, pGEM-3Zf, pGEM-5Zf, pGEM-7Zf, pGEM-9Zf, and pGEM-11Zf, the entire series of plasmids is commercially available from Promega.

The double stranded FP-encoding DNA is cloned into the bacteriophage promoter containing vector in the proper orientation using one or more of the available restriction endonuclease cloning sites on the vector which are convenient and appropriate for cloning FP DNA. The vector with the ligated FP DNA is used to transform bacteria, and clonal isolates are analyzed for the presence of the vector with the FP DNA in the proper orientation.

Once a vector containing the FP-encoding DNA in the proper orientation is identified and isolated, it is linearized by cleavage with a restriction endonuclease at a site downstream from, and without disrupting, the FP transcription unit. The linearized plasmid is isolated and purified, and used as a template for in vitro transcription of FP mRNA.

The template DNA is then mixed with bacteriophage-specific DNA-dependent RNA polymerase in a reaction mixture which allows transcription of the DNA template forming FP mRNA. Several bacteriophage-specific DNA-dependent RNA polymerases are available, including but not limited to T3, T7, and SP6 RNA polymerase. The synthetic FP mRNA is then isolated and purified.

It may be advantageous to synthesize mRNA containing a 5' terminal cap structure and a 3' poly A tail to improve mRNA stability. A cap structure, or 7-methylguanosine, may be incorporated at the 5' terminus of the mRNA by simply adding 7-methylguanosine to the reaction mixture with the DNA template. The DNA-dependent RNA polymerase incorporates the cap structure at the 5' terminus as it synthesizes the mRNA. The poly A tail is found naturally occurring in many cDNAs but can be added to the 3' terminus of the mRNA by simply inserting a poly A tail-encoding DNA sequence at the 3' end of the DNA template.

The isolated and purified FP mRNA is translated using either a cell-free system, including but not limited to rabbit reticulocyte lysate and wheat germ extracts (both commercially available from Promega and New England Nuclear) or in a cell based system, including but not limited to microinjection into *Xenopus* oocytes, with microinjection into *Xenopus* oocytes being preferred.

Xenopus oocytes are microinjected with a sufficient amount of synthetic FP mRNA to produce FP protein. The microinjected oocytes are incubated to allow translation of the FP mRNA, forming FP protein.

These synthetic mRNAs are injected into *Xenopus* oocytes (stage 5 -6) by standard procedures [Gurdon, J.B. and Wickens, M.D. *Methods in Enzymol.* 101: 370-386, (1983)]. Oocytes are harvested and analyzed for FP expression as described below.

EXAMPLE 5

pcDNAIamp-FP expression in *Xenopus* oocytes

Oocytes were taken from adult females of *Xenopus laevis* using standard surgical procedure (Colman, A., 1984 *In: Transcription and Translation - A Practical Approach*, IRL Press). To remove follicle cells, oocytes were treated for 2-3 h with freshly made collagenase (2 mg/ml, type 2, Worthington Biochemical Corp., Freehold, NJ) in Ca^{2+} -free ND96 solution (ND96 in mM: NaCl 96, KCl 2, $MgCl_2$ 1, HEPES 5, Na-pyruvate 2.5, theophylline 0.5, gentamicin 50 mg/ml, +1.8 $CaCl_2$, pH 7.6). Defolliculated stage 5-6 oocytes were selected and maintained in ND96 solution. Oocyte nuclei were injected with 1-5 ng of pcDNAIamp-FP and then incubated at 18°C for 48 h before challenge with agonist. Func-

tional activity was determined by measurement of either agonist-induced Ca^{2+} -dependent Cl^- current or light emission in oocytes injected with the Ca^{2+} -specific photoprotein aequorin (J. Blinks, Friday Harbor Photoproteins, WA), (Giladi and Spindel 1991 Biotechniques, 10, pp 744-747). For the electrophysiological assays an oocyte was placed in a 0.5 ml perfusion chamber and voltage clamped at -60 mV (with microelectrodes of 0.5-2.0 MW resistance filled with 3 M KCl) using a Turbo TEC 01C amplifier (NPI Instruments, Germany). Ligand-containing solution was perfused and the current response was recorded. For the luminometric assay, aequorin-loaded oocytes (100 ng/oocyte) were placed individually in cuvettes containing 0.4 ml ND96 and the light emission provoked by ligand addition was recorded using a Bio-Orbit 1251 luminometer (Fisher Sci. Ltd.).

Functional activity was determined in pcDNA1amp-FP-injected oocytes using electrophysiological and aequorin luminescence assays. In the electrophysiological assay, perfusion of 10 to 20nM $\text{PGF}_2\alpha$ or 10nM fluprostenol, a selective FP receptor agonist, resulted in prominent current responses in oocytes injected with pcDNA1amp-FP confirming that this clone encodes a functional FP receptor that is coupled to the phosphatidylinositol/ Ca^{2+} signalling pathway (Figure 3). Such responses were absent in control (H_2O -injected) oocytes. Ligand-induced increases in intracellular Ca^{2+} were also demonstrated directly by light emission in aequorin-loaded oocytes (Figure 4). The dose-response dependencies obtained from aequorin luminescence assay indicated that $\text{PGF}_2\alpha$ and fluprostenol were more potent agonists of the expressed receptor when compared to PGE_2 (Figure 5). This rank order of potency is consistent with that reported for the FP receptor [Coleman, et al., 1991].

EXAMPLE 6

Cloning of FP cDNA into a Mammalian Expression Vector

FP cDNA expression cassettes are ligated at appropriate restriction endonuclease sites to the following vectors containing strong, universal mammalian promoters: pBC12BI [Cullen, B.R. Methods in Enzymol. 152: 684-704 1988], and pEE12 (CellTech EP O 338,841) and its derivatives pSZ9016-1 and p9019. p9019 represents the construction of a mammalian expression vector containing the hCMVIE promoter, polylinker and SV40 polyA element with a selectable marker/amplification system comprised of a mutant gene for dihydrofolate reductase (mdHFR) (Simonsen, C.C. and Levinson, A.D. Proc. Natl. Acad. Sci USA 80: 2495-2499 [1983]) driven by the SV40 early promoter. An SV40 polyadenylation sequence is generated by a PCR reaction defined by primers 13978-120 and 139778-121 using pD5 (Berker and Sharp, Nucl. Acid Res. 13: 841-857 [1985]) as template. The resulting 0.25 Kb PCR product is digested with ClaI and SpeI and ligated into the 6.7 Kb fragment of pEE12 which had been likewise digested. The resultant plasmid is digested with BglII and SfiI to liberate the 3' portion of the SV40 early promoter and the GScDNA from the vector. A 0.73 Kb SfiI-XhoII fragment isolated from plasmid pFR400 (Simonsen, C.C. and Levinson, A.D. Proc. Natl. Acad. Sci USA 80: 2495-2499 [1983]) is ligated to the 5.6 Kb vector described above, reconstituting the SV40 early promoter, and inserting the mdHFR gene. This plasmid is designated p9019. pSZ9016-1 is identical to p9019 except for the substitution of the HIV LTR for the huCMVIE promoter. This vector is constructed by digesting p9019 with XbaI and MluI to remove the huCMVIE promoter. The HIV LTR promoter, from residue -117 to +80 (as found in the vector pCD23 containing the portion of the HIV-1 LTR (Cullen, Cell 46:973 [1986]) is PCR amplified from the plasmid pCD23 using oligonucleotide primers which appended to the ends of the product the MluI and SpeI restriction sites on the 5' side while Hind III and Xba I sites are appended on the 3' side. Following the digestion of the resulting 0.2 kb PCR product with the enzymes MluI and Xba I the fragment is agarose gel-purified and ligated into the 4.3 Kb promoterless DNA fragment to generate the vector pSZ9016-1.

Cassettes containing the FP cDNA in the positive orientation with respect to the promoter are ligated into appropriate restriction sites 3' of the promoter and identified by restriction site mapping and/or sequencing. These cDNA expression vectors are introduced into various host cells including, but not limited to: COS-7 (ATCC# CRL1651), CV-1 [Sackevitz et al., Science 238: 1575 (1987)], 293, L cells (ATCC# CRL6362) by standard methods including but not limited to electroporation, or chemical procedures (cationic liposomes, DEAE dextran, calcium phosphate). Transfected cells and cell culture extracts can be harvested and analyzed for FP expression as described below.

All of the vectors used for mammalian transient expression can be used to establish stable cell lines expressing FP. Unaltered FP cDNA constructs cloned into expression vectors will be expected to program host cells to make intracellular FP protein. The transfection host cells include, but are not limited to, CV-1 [Sackevitz et al., Science 238: 1575 (1987)], tk-L [Wigler, et al. Cell 11: 223 (1977)], NS/0, and dhFr-CHO [Kaufman and Sharp, J. Mol. Biol. 159: 601, (1982)].

Co-transfection of any vector containing FP cDNA with a drug selection plasmid including, but not limited to G418, aminoglycoside phosphotransferase, pLNCX [Miller, A.D. and Rosman G. J. Biotech News 7: 980-990 (1989)]; hygromycin, hygromycin-B phosphotransferase, pLG90 [Gritz, L. and Davies, J., GENE 25: 179 (1983)]; APRT, xanthine-guanine phosphoribosyl-transferase, pMAM (Clontech) [Murray, et al., Gene 31: 233 (1984)] will allow for the selection of stably transfected clones. Levels of FP are quantitated by the assays described above.

FP cDNA constructs are ligated into vectors containing amplifiable drug-resistance markers for the production of mammalian cell clones synthesizing the highest possible levels of FP. Following introduction of these constructs into cells, clones containing the plasmid are selected with the appropriate agent, and isolation of an over-expressing clone with a high copy number of the plasmid is accomplished by selection in increasing doses of the agent. The following systems are utilized: the 9016 or the 9019 plasmid containing the mutant DHFR gene [Simonson, C. and Levinson, A., Proc. Natl. Acad. Sci. USA 80: 2495 (1983)], transfected into DHFR- CHO cells and selected in methotrexate; the pEE12 plasmid containing the glutamine synthetase gene, transfected into NS/O cells and selected in methionine sulfoximine (CellTech International Patent Application 2089/10404); and 9016 or other CMV promoter vectors, co-transfected with pDLAT-3 containing the thymidine kinase gene [Colbere and Garopin, F., Proc. Natl. Acad. Sci. 76: 3755 (1979)] in APRT and TK deficient L cells, selected in APRT (0.05 mM azaserine, 0.1 mM adenine, 4 ug/ml adenine) and amplified with HAT (100 uM hypoxanthine, 0.4 uM aminopterin, 16 uM thymidine).

EXAMPLE 7

Expression of the FP receptor in COS-M6 cells and [3H]PGF₂α binding assays

The recently cloned human prostaglandin F_{2a} (FP) receptor was subcloned into the pcDNAamp plasmid (Invitrogen) and transfected into COS-M6 cells using the DEAE-dextran method. The cells were maintained in culture for 72 h, then harvested and membranes prepared by differential centrifugation (1000 x g for 10 min, then 100,000 x g for 30 min) following lysis of the cells by nitrogen cavitation. [3H]Prostaglandin F_{2a} ([3H]PGF₂α) binding assays were performed in 10 mM MES/KOH pH 6.0, containing 0.4 mM EDTA, 10 mM MnCl₂, 0.3 nM [3H]PGF₂α and 60 μg of protein from the 100,000 x g membrane fraction. Incubations were conducted for 1 h at room temperature prior to separation of the bound and free radioligand by rapid filtration through Whatman GF/B filters presoaked at 4°C in washing buffer (10 μM MES/KOH (pH 6.0) containing 0.01% bovine serum albumin). The filters were washed with approximately 16 ml of washing buffer and the residual [3H]PGF₂α bound to the filter was quantified by liquid scintillation counting. Specific binding was defined as the difference between total binding and non-specific binding, determined in the presence of 2 μM PGF₂α.

The cloned human FP receptor was transfected into COS-M6 cells and [3H]PGF₂α binding assays were performed with membranes prepared from the transfected cells. In competition assays PGF₂α was the most potent competing ligand for [3H]PGF₂α specific binding with an IC₅₀ value of 2.8 nM (Fig. X). The related synthetic FP agonist fluprostenol was equipotent with an IC₅₀ value of 3.5 nM. The rank order of potency for prostaglandins and related analogs was: PGF₂α = fluprostenol > PGD₂ >> PGE₂ ≈ U46619 > iloprost. U46619 and iloprost are stable analogs of thromboxane and prostacyclin and display comparable potency at the TP and IP receptors, respectively. This rank order of potency has been predicted for the FP receptor from previous pharmacological studies.

EXAMPLE 8

Cloning of FP cDNA into a Baculovirus Expression Vector for Expression in Insect Cells

Baculovirus vectors, which are derived from the genome of the AcNPV virus, are designed to provide high level expression of cDNA in the Sf9 line of insect cells (ATCC CRL# 1711). Recombinant baculoviruses expressing FP cDNA are produced by the following standard methods (InVitrogen Maxbac Manual): the FP cDNA constructs are ligated downstream of the polyhedrin promoter in a variety of baculovirus transfer vectors, including the pAC360 and the pBlueBac vector (InVitrogen). Recombinant baculoviruses are generated by homologous recombination following co-transfection of the baculovirus transfer vector and linearized AcNPV genomic DNA [Kitts, P.A., Nuc. Acid. Res. 18: 5667 (1990)] into Sf9 cells. Recombinant pAC360 viruses are identified by the absence of inclusion bodies in infected cells (Summers, M.D. and Smith, G.E., Texas Agriculture Exp. Station Bulletin No. 1555) and recombinant pBlueBac viruses are identified on the basis of b-galactosidase expression (Vialard, *et al.*, 1990, J. Virol., 64, pp 37-50). Following plaque purification and infection of sf9 cells with FP recombinant baculovirus, FP expression is measured by the assays described above.

The cDNA encoding the entire open reading frame for FP is inserted into the BamHI site of pBlueBacII. Constructs in the positive orientation with respect to the polyhedrin promoter are identified by sequence analysis and used to transfect Sf9 cells in the presence of linear AcNPV wild type DNA.

Authentic, active FP is found associated with the membranes of infected cells. Membrane preparations are prepared from infected cells by standard procedures.

EXAMPLE 9Cloning of FP cDNA into a yeast expression vector

Recombinant FP is produced in the yeast *S. cerevisiae* following the insertion of the optimal FP cDNA construct into expression vectors designed to direct the intracellular expression of heterologous proteins. For intracellular expression, vectors such as EmBlyex4 or the like are ligated to the FP cistron [Rinas, U. *et al.*, Biotechnology 8: 543-545 (1990); Horowitz B. *et al.*, J. Biol. Chem. 265: 4189-4192 (1989)]. The levels of expressed FP are determined by the assays described above.

EXAMPLE 10Purification of Recombinant FP

Recombinantly produced FP may be purified by antibody affinity chromatography.

FP antibody affinity columns are made by adding the anti-FP antibodies to Affigel-10 (Biorad), a gel support which is pre-activated with N-hydroxysuccinimide esters such that the antibodies form covalent linkages with the agarose gel bead support. The antibodies are then coupled to the gel via amide bonds with the spacer arm. The remaining activated esters are then quenched with 1M ethanolamine HCl (pH 8). The column is washed with water followed by 0.23 M glycine HCl (pH 2.6) to remove any non-conjugated antibody or extraneous protein. The column is then equilibrated in phosphate buffered saline (pH 7.3) together with appropriate membrane solubilizing agents such as detergents and the cell culture supernatants or cell extracts containing solubilized FP or FP subunits are slowly passed through the column. The column is then washed with phosphate-buffered saline together with detergents until the optical density (A280) falls to background, then the protein is eluted with 0.23 M glycine-HCl (pH 2.6) together with detergents. The purified FP protein is then dialyzed against phosphate buffered saline together with detergents.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: ABRAMOVITZ, MARK
 GRYGORCZYK, RICHARD
 METTERS, KATHLEEN
 NGUYEN, TRUYEN
 RUSHMORE, THOMAS H.
 SLIPETZ, DEBORAH

(ii) TITLE OF INVENTION: DNA Encoding Prostaglandin Receptor FP

(iii) NUMBER OF SEQUENCES: 5

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: John W. Wallen, III
 (B) STREET: P. O. Box 2000, 126 E. Lincoln Avenue
 (C) CITY: Rahway
 (D) STATE: New Jersey
 (E) COUNTRY: USA
 (F) ZIP: 07065

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
 (B) COMPUTER: IBM PC compatible
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 (D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

EP 0 705 276 B1

(A) APPLICATION NUMBER: US 08/083,741
(B) FILING DATE: 25-JUN-1993
(C) CLASSIFICATION:

5 (viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: WALLEN, JOHN W.
(B) REGISTRATION NUMBER: 35,403
(C) REFERENCE/DOCKET NUMBER: 19027P

10

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (908)594-3905
(B) TELEFAX: (908)594-4720

15

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

20

(A) LENGTH: 27 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

30

ATAMACCCAG GGRTCCARGA TCTGRTT

27

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

35

(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

45

Asn Gln Ile Leu Asp Pro Trp Val Tyr
1 5

(2) INFORMATION FOR SEQ ID NO:3:

50

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 373 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

55

(ii) MOLECULE TYPE: cDNA

EP 0 705 276 B1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

5	CTCTTAGCTC TTGGTGTTTC CTTCTCGTGC AATGCCGTCA CGGGAGTCAC ACTCTTAAGA	60
	GTGAAGTTCA GAAGCCAGCA GCATAGGCAA GGCAGATCTC ACCACCTGGA GATGATCATT	120
	CAGCTCCTGG CCATAATGTG CGTCTCCTGC GTCTGCTGGA GTCCCTTTCT GGTAACAATG	180
10	GCCAACATTG CAATAAATGG AAATAATTCC CCAGTGACCT GTGAAACGAC ACTTTTGTCT	240
	CTCCGCATGG CAACGTGGAA TCAGATCTTA GATCCCTGGG TCTATATTCT GCTACGGAAG	300
	GCTGTCCTTA GGAACCTGTA TAAACTTGCC AGTCGTTGCT GTGGAGTTAA CATCATCAGC	360
15	TTGCATATCT GGG	373

(2) INFORMATION FOR SEQ ID NO:4:

20 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1437 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
25 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

35

40

45

50

55

EP 0 705 276 B1

GTGCGCGGAG GGGACGAGCG GCTGGACCAC AGCCGGCGCC CGATCAGGAT CTCCGCGCTG 60
 GGATCGGTGG AACTTGAGGC AGCGGCGGCG CGGGGCGCCA TGGCACACCG AGCGGCTCCG 120
 5 TCTTCTGCTC CTCAGAGAGC CCGGCTGGCG GCCTGGGATG ACAAGATGTC TGGACTGCAA 180
 TCCTGCACAG TTTTGAGAGG GAGATGACTT GAGTGGTTGG CTTTATCTC CACAACAATG 240
 10 TCCATGAACA ATTCCAAACA GCTAGTGTCT CCTGCAGCTG CGCTTCTTTC AAACACAACC 300
 TGCCAGACGG AAAACCGGCT TTCCGTATTT TTTTCAGTAA TCTTCATGAC AGTGGGAATC 360
 TTGTCAAACA GCCTTGCCAT CGCCATTCTC ATGAAGGCAT ATCAGAGATT TAGACAGAAG 420
 15 TCCAAGGCAT CGTTTCTGCT TTTGGCCAGC GGCTGGTAA TCACTGATTT CTTTGGCCAT 480
 CTCATCAATG GAGCCATAGC AGTATTTGTA TATGCTTCTG ATAAAGAATG GATCCGCTTT 540
 GACCAATCAA ATGTCCTTTG CAGTATTTTT GGTATCTGCA TGGTGTTTTC TGGTCTGTGC 600
 20 CCACTTCTTC TAGGCAGTGT GATGGCCATT GAGCGGTGTA TTGGAGTCAC AAAACCAATA 660
 TTTCAATTCTA CGAAAATTAC ATCCAAACAT GTGAAAATGA TGTTAAGTGG TGTGTGCTTG 720
 TTTGCTGTTT TCATAGCTTT GCTGCCCATC CTTGGACATC GAGACTATAA AATTCAGGCG 780
 25 TCGAGGACCT GGTGTTTCTA CAACACAGAA GACATCAAAG ACTGGGAAGA TAGATTTTAT 840
 CTTCTACTTT TTTCTTTTCT GGGGCTCTTA GCCCTTGGTG TTTCAATTGTT GTGCAATGCA 900
 30 ATCACAGGAA TTACACTTTT AAGAGTTAAA TTTAAAAGTC AGCAGCACAG ACAAGGCAGA 960
 TCTCATCATT TGGAAATGGT AATCCAGCTC CTGGCGATAA TGTGTGTCTC CTGTATTTGT 1020
 TGGAGCCCAT TTCTGGTTAC AATGGCCAAC ATTGGAATAA ATGGAATCA TTCTCTGGAA 1080
 35 ACCTGTGAAA CAACACTTTT TGCTCTCCGA ATGGCAACAT GGAATCAAAT CTTAGATCCT 1140
 TGGGTATATA TTCTTCTACG AAAGGCTGTC CTTAAGAATC TCTATAAGCT TGCCAGTCAA 1200
 TGCTGTGGAG TGCATGTCAT CAGCTTACAT ATTTGGGAGC TTAGTTCCAT TAAAAATTCC 1260
 40 TTAAAGGTTG CTGCTATTTT TGAGTCACCA GTTGCGAGAG AATCAGCAAG CACCTAGCTT 1320
 AATAGGACAG TAAATCTGTG TGGGGCTAGA ACAAAAATTA AGACATGTTT GGCAATATTT 1380
 45 CAGTTAGTTA AATACCTGTA GCCTAACTGG AAAATTCAGG CTTTCATCATG TAGTTTG 1437

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 359 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

EP 0 705 276 B1

		Met	Ser	Met	Asn	Asn	Ser	Lys	Gln	Leu	Val	Ser	Pro	Ala	Ala	Ala	Leu
		1				5					10					15	
5		Leu	Ser	Asn	Thr	Thr	Cys	Gln	Thr	Glu	Asn	Arg	Leu	Ser	Val	Phe	Phe
				20						25					30		
		Ser	Val	Ile	Phe	Met	Thr	Val	Gly	Ile	Leu	Ser	Asn	Ser	Leu	Ala	Ile
				35					40					45			
10		Ala	Ile	Leu	Met	Lys	Ala	Tyr	Gln	Arg	Phe	Arg	Gln	Lys	Ser	Lys	Ala
		50						55					60				
		Ser	Phe	Leu	Leu	Leu	Ala	Ser	Gly	Leu	Val	Ile	Thr	Asp	Phe	Phe	Gly
		65					70					75					80
15		His	Leu	Ile	Asn	Gly	Ala	Ile	Ala	Val	Phe	Val	Tyr	Ala	Ser	Asp	Lys
					85						90				95		
		Glu	Trp	Ile	Arg	Phe	Asp	Gln	Ser	Asn	Val	Leu	Cys	Ser	Ile	Phe	Gly
20				100						105					110		
		Ile	Cys	Met	Val	Phe	Ser	Gly	Leu	Cys	Pro	Leu	Leu	Leu	Gly	Ser	Val
				115					120					125			
		Met	Ala	Ile	Glu	Arg	Cys	Ile	Gly	Val	Thr	Lys	Pro	Ile	Phe	His	Ser
25		130						135					140				
		Thr	Lys	Ile	Thr	Ser	Lys	His	Val	Lys	Met	Met	Leu	Ser	Gly	Val	Cys
		145					150					155					160
30		Leu	Phe	Ala	Val	Phe	Ile	Ala	Leu	Leu	Pro	Ile	Leu	Gly	His	Arg	Asp
					165						170					175	
		Tyr	Lys	Ile	Gln	Ala	Ser	Arg	Thr	Trp	Cys	Phe	Tyr	Asn	Thr	Glu	Asp
				180						185					190		
35		Ile	Lys	Asp	Trp	Glu	Asp	Arg	Phe	Tyr	Leu	Leu	Leu	Phe	Ser	Phe	Leu
				195					200					205			
		Gly	Leu	Leu	Ala	Leu	Gly	Val	Ser	Leu	Leu	Cys	Asn	Ala	Ile	Thr	Gly
40		210						215					220				

Ile Thr Leu Leu Arg Val Lys Phe Lys Ser Gln Gln His Arg Gln Gly
 225 230 235 240
 5 Arg Ser His His Leu Glu Met Val Ile Gln Leu Leu Ala Ile Met Cys
 245 250 255
 Val Ser Cys Ile Cys Trp Ser Pro Phe Leu Val Thr Met Ala Asn Ile
 260 265 270
 10 Gly Ile Asn Gly Asn His Ser Leu Glu Thr Cys Glu Thr Thr Leu Phe
 275 280 285
 Ala Leu Arg Met Ala Thr Trp Asn Gln Ile Leu Asp Pro Trp Val Tyr
 290 295 300
 15 Ile Leu Leu Arg Lys Ala Val Leu Lys Asn Leu Tyr Lys Leu Ala Ser
 305 310 315 320
 Gln Cys Cys Gly Val His Val Ile Ser Leu His Ile Trp Glu Leu Ser
 325 330 335
 20 Ser Ile Lys Asn Ser Leu Lys Val Ala Ala Ile Ser Glu Ser Pro Val
 340 345 350
 25 Ala Glu Lys Ser Ala Ser Thr
 355

Claims

- 30 1. An isolated and purified human prostaglandin receptor protein wherein said protein is characterized by the amino acid sequence

35 MSMNNSKQLVSPAALLSN'ITCQTENRLSVFFSVIFMTVGILSNSLAIAILMK
 AYQRFRQKSKASFLLLASGLVITDFFGHLINGAIAVFVYASDKEWIRFDQSNV
 LCSIFGICMVFSGLCPLLLGSVMAIERCIGVTKPIFHSTKJTSKHVKMMLSGV
 CLFAVFIALLPILGHRDYKIQASRTWCFYNTEDIKDWEDRFYLLFSFLGLLAL
 40 GVSLLCNAITGITLLRVKFSQQHRQGRSHHLEMVIQLLAIMCVSCICWSPFL
 VTMANIGINGNHSLETCTTLFALRMATWNQILDWPVYILLRKA VLKNLYKL
 ASQCCGVHVISLHIWELSSIKNSLKVAAISESPVAEKSAST (SEQ.ID.NO.: 5).

- 45 2. An isolated and purified DNA molecule encoding a prostaglandin receptor protein wherein said proten is characterized by the amino acid sequence of Claim 1.

- 50 3. An isolated and purified DNA molecule encoding a prostaglandin receptor protein wherein said DNA molecule is characterized by the nucleotide sequence

55

GTGCGCGGAGGGGACGAGCGGCTGGACCACAGCCGGCGCCCGATCAGG
 ATCTCCGCGCTGGGATCGGTGGAACCTTGAGGCAGCGGCGGCGCGGGGC
 5 GCCATGGCACACCGAGCGGCTCCGTCTTCTGCTCCTCAGAGAGCCCCGGCT
 GCGGCGCTGGGATGACAAGATGTCTGGACTGCAATCCTGCACAGTTTTGA
 GAGGGAGATGACTTGAGTGGTTGGCTTTTATCTCCACAACAATGTCCATG
 10 AACAAATTCAAACAGCTAGTGTCTCCTGCAGCTGCGCTTCTTTCAAACACA
 ACCTGCCAGACGGAAAACCGGCTTTCCGTATTTTTTTTCAGTAATCTTCATG
 ACAGTGGGAATCTTGTCAAACAGCCTTGCCATCGCCATTCTCATGAAGGC
 ATATCAGAGATTTAGACAGAAGTCCAAGGCATCGTTTCTGCTTTTGGCCAG
 15 CGGCCTGGTAATCACTGATTTCTTTGGCCATCTCATCAATGGAGCCATAGC

AGTATTTGTATATGCTTCTGATAAAGAATGGATCCGCTTTGAOCCAATCAAA
 20 TGTCCCTTGCAGTATTTTTGGTATCTGCATGGTGTTTTCTGGTCTGTGCCCA
 CTTCTTCTAGGCAGTGTGATGGCCATTGAGCGGTGTATTGGAGTCACAAA
 AOCCAATATTTCAATTCTACGAAAATTACATCCAAACATGTGAAAATGATGTTA
 25 AGTGGTGTGTGCTTGTGTTGCTGTTTTCATAGCTTTGCTGCCCATCCTTGGA
 CATCGAGACTATAAAATTCAGGCGTGGAGGACCTGGTGTGTTCTACAACAC
 AGAAGACATCAAAGACTGGGAAGATAGATTTTATCTTCTACTTTTTTCTTTT
 30 CTGGGGCTCTTAGCCCTTGGTGTGTTTCATTGTTGTGCAATGCAATCACAGGA
 ATTACACTTTTAAGAGTTAAATTTAAAAGTCAGCAGCACAGACAAGGCAGA
 TCTCATCATTTGGAAATGGTAATCCAGCTCCTGGCGATAATGTGTGTCTCC
 35 TGTATTTGTTGGAGCCCATTTCTGGTTACAATGGCCAACATTGGAATAAAT
 GGAAATCATTCTCTGGAAACCTGTGAAACAACACTTTTTGCTCTCCGAATG
 GCAACATGGAATCAAATCTTAGATCCTTGGGTATATATTCTTCTACGAAAG
 GCTGTCTTAAGAATCTCTATAAGCTTGCCAGTCAATGCTGTGGAGTGCAT
 40 GTCATCAGCTTACATATTTGGGAGCTTAGTTCCATTAAAAATTCCTTAAAG
 GTTGCTGCTATTTCTGAGTCACCAAGTTGCAGAGAAATCAGCAAGCACCTA
 GCTTAATAGGACAGTAAATCTGTGTGGGGCTAGAACAAAAATTAAGACAT
 45 GTTTGGCAATATTTCAAGTTAGTTAAATACCTGTAGCCTAACTGGAAAATTC
 AGGCTTCATCATGTAGTTTG (SEQ.ID.NO.: 4).

4. An expression vector for the expression of a prostaglandin receptor protein in a recombinant host cell wherein said expression vector contains the DNA molecule of Claim 3.
5. A host cell which expresses a recombinant prostaglandin receptor protein wherein said host cell contains the expression vector of Claim 4.
6. A process for the expression of a prostaglandin receptor protein in a recombinant host cell, comprising:
 - (a) transferring the expression vector of Claim 4 into a suitable host cell; and
 - (b) culturing the host cells of step (a) under conditions which allow expression of the prostaglandin receptor

protein from the expression vector.

7. A method of identifying modulators of a prostaglandin receptor activity, comprising:

- (a) combining a modulator of prostaglandin receptor activity, with a prostaglandin receptor wherein said receptor is characterized by the amino acid sequence of Claim 1 in whole or in part; and
(b) measuring the effect of a modulator on the prostaglandin receptor.

8. An antibody which specifically binds to a prostaglandin receptor protein wherein said protein is characterized by the amino acid sequence of Claim 1.

Patentansprüche

1. Isoliertes und gereinigtes Humanprostaglandin-Rezeptorprotein, wobei das Protein gekennzeichnet ist durch die Aminosäuresequenz

MSMNNSKQLVSPAAALLSN¹ITCQTENRLSVFFSVIFMTVGLSNSLAIAILMK
AYQRFQKSKASFLLLASGLVITDFFGHLINGAIAVFVYASDKIEWIRFDQSNV
LCSIFGICMVFSGLCPDLLGSVMAIERCIGVTKPIFHSTKJTSKHVKMMLSGV
CLFAVFLALLPILGHRDYKIQASRTWCFYNTEDIKDWEDRFYLLLFSFLGLLAL
GVSLLCNAITGITLLRVKFKSQQRHSHLEMVIQLLAIMCVSCICWSPFL
VTMANIGINGNHSLETCTTLFALRMATWNQILDWPVYILLRKA VLKNLYKL
ASQCCGVHVISLHIWELSSIKNSLKVAAISESPVAEKSAST (SEQ-ID-Nr. 5).

2. Isoliertes und gereinigtes DNA-Molekül, das für ein Prostaglandin-Rezeptorprotein kodiert, wobei das Protein durch die Aminosäuresequenz von Anspruch 1 gekennzeichnet ist.

3. Isoliertes und gereinigtes DNA-Molekül, das für ein Prostaglandin-Rezeptorprotein kodiert, wobei das DNA-Molekül gekennzeichnet ist durch die Nukleotidsequenz

GTGCGCGGAGGGGACGAGCGGCTGGACCACAGCCGGCGCCCGATCAGG
ATCTCCGCGCTGGGATCGGTGGAACCTTGAGGCAGCGGCGGCGCGGGGC
GCCATGGCACACCGAGCGGCTCCGTCTTCTGCTCCTCAGAGAGCCCGGCT
GGCGGCCTGGGATGACAAGATGTCTGGACTGCAATCCTGCACAGTTTTGA
GAGGGAGATGACTTGAGTGGTTGGCTTTTATCTCCACAACAATGTCCATG
AACAAATCCAAACAGCTAGTGTCTCCTGCAGCTGCGCTTCTTTCAAACACA
ACCTGCCAGACGGAAAACCGGCTTTCCGTATTTTTTTCAGTAATCTTCATG
ACAGTGGGAATCTTGTCAAACAGCCTTGCCATCGCCATTCTCATGAAGGC
ATATCAGAGATTTAGACAGAAGTCCAAGGCATCGTTTCTGCTTTTGGCCAG
CGGCCTGGTAATCACTGATTTCTTTGGCCATCTCATCAATGGAGCCATAGC

AGTATTTGTATATGCTTCTGATAAAGAATGGATCCGCTTTGACCAATCAAA
 TGTCCCTTTCAGTATTTTTGGTATCTGCATGGTGTCTTCTGGTCTGTGCCCA
 5 CTTCTTCTAGGCAGTGTGATGGCCATTGAGCGGTGTATTGGAGTCACAAA
 ACCAATATTTCAATTCTACGAAAATTACATCCAAACATGTGAAAATGATGTTA
 AGTGGTGTGTGCTTGTCTTGTCTGTTTTCATAGCTTTGCTGCCCATCCTTGGA
 10 CATCGAGACTATAAAATTCAGGCGTCGAGGACCTGGTGTCTTCTACAACAC
 AGAAGACATCAAAGACTGGGAAGATAGATTTTATCTTCTACTTTTTCTTTT
 CTGGGGCTCTTAGCCCTTGGTGTCTTCAATTGTTGTGCAATGCAATCACAGGA
 15 ATTACACTTTTAAGAGTTAAATTTAAAAGTCAGCAGCACAGACAAGGCAGA
 TCTCATCATTTGGAAATGGTAATCCAGCTCCTGGCGATAATGTGTGTCTCC
 TGTATTTGTTGGAGCCCATTTCTGGTTACAATGGCCAACATTGGAATAAAT
 GGAAATCATTCTCTGGAAACCTGTGAAACAACACTTTTTGCTCTCOGAATG
 20 GCAACATGGAATCAAATCTTAGATCCTTGGGTATATATTCTTCTACGAAAG
 GCTGTCCTTAAGAATCTCTATAAGCTTGCCAGTCAATGCTGTGGAGTGCAT
 GTCATCAGCTTACATATTTGGGAGCTTAGTTCATTAAAAATTCCTTAAAG
 25 GTTGCTGCTATTTCTGAGTCACCAAGTTGCAGAGAAATCAGCAAGCACCTA
 GCTTAATAGGACAGTAAATCTGTGTGGGGCTAGAACAAAAATTAAGACAT
 GTTTGGCAATATTTCAAGTTAGTTAAATACCTGTAGCCTAACTGGAAAATTC
 30 AGGCTTCATCATGTAGTTTG (SEQ-ID-Nr. 4).

4. Expressionsvektor für die Expression eines Prostaglandin-Rezeptorproteins in einer rekombinanten Wirtszelle, wobei der Expressionsvektor das DNA-Molekül nach Anspruch 3 enthält.
5. Wirtszelle, die ein rekombinantes Prostaglandin-Rezeptorprotein exprimiert, wobei die Wirtszelle den Expressionsvektor nach Anspruch 4 enthält.
6. Verfahren zur Expression eines Prostaglandin-Rezeptorproteins in einer rekombinanten Wirtszelle, welches umfaßt:
 - (a) Überführung des Expressionsvektors nach Anspruch 4 in eine geeignete Wirtszelle; und
 - (b) Kultivierung der Wirtszellen von Schritt (a) unter Bedingungen, welche die Expression des Prostaglandin-Rezeptorproteins aus dem Expressionsvektor erlauben.
7. Verfahren zur Identifizierung von Modulatoren einer Prostaglandin-Rezeptoraktivität, welches umfaßt:
 - (a) Kombination eines Modulators von Prostaglandin-Rezeptoraktivität mit einem Prostaglandin-Rezeptor, wobei der Rezeptor durch die vollständige oder partielle Aminosäuresequenz von Anspruch 1 gekennzeichnet ist; und
 - (b) Messung der Wirkung eines Modulators auf den Prostaglandin-Rezeptor.
8. Antikörper, der spezifisch an ein Prostaglandin-Rezeptorprotein bindet, wobei das Protein durch die Aminosäuresequenz von Anspruch 1 gekennzeichnet ist.

Revendications

1. Protéine de récepteur de prostaglandine humaine isolée et purifiée, ladite protéine étant caractérisée par la séquence d'acides aminés

MISMINNSKQLVSPAAALLSNITTCQTENRLSVFFSVIFMITVGI LSN SLAIAILMK
 AYQRFQRQKSKASFLLLASGLVITDFFGHLLINGAJAVFVYASDKEWIRFDQSNV
 LCSIFGICMVFSGLCPLLLGSVMAIERCIGVTKPIFHSITKJTSKHVKNMMLSGV
 CLFAVFIALLPILGHRDYKJQASRTWCFYNTEDIKDWEDRFYLLLFSLGLLAL
 GVSLLCNAITGITLLRVKFKSQQHRQGRSHHLEMVIQLLAIMCVSCICWSPFL
 VTMANIGINGNHSLETCTTLFALRMATWNQILDWPVYILLRKAVLKNLYKL
 ASQCCGVHVISLHIWELSSIKNSLKVA AIS ESPVAEKSAST (SEQ.ID.NO.: 5).

2. Molécule d'ADN isolée et purifiée codant pour une protéine de récepteur de prostaglandine, ladite protéine étant caractérisée par la séquence d'acides aminés de la revendication 1.
3. Molécule d'ADN isolée et purifiée codant pour une protéine de récepteur de prostaglandine, ladite molécule d'ADN étant caractérisée par la séquence nucléotidique

GTGCGCGGAGGGGACGAGCGGCTGGACCACAGCCGGCGCCCGATCAGG
 ATCTCCGCGCTGGGATCGGTGGAACCTTGAGGCAGCGGCGGCGCGGGGC
 GCCATGGCACACCGAGCGGCTCCGTCTTCTGCTCCTCAGAGAGCCCGGCT
 GGCGGCCTGGGATGACAAGATGTCTGGACTGCAATCCTGCACAGTTTGA
 GAGGGAGATGACITGAGTGGTTGGCTTTTATCTCCACAACAATGTCCATG
 AACAATTCCAAACAGCTAGTGTCTCCTGCAGCTGCGCTTCTTCAAACACA
 ACCTGCCAGACGGAAAACCGGCTTCCGTATTTTTTTCAGTAATCTTCATG
 ACAGTGGGAATCTTGTCAAAACAGCCTTGCCATCGCCATTCTCATGAAGGC
 ATATCAGAGATTTAGACAGAAGTCCAAGGCATCGTTTCTGCTTTTGGCCAG
 CGGCCCTGGTAATCACTGATTTCTTTGGCCATCTCATCAATGGAGCCATAGC
 AGTATTTGTATATGCTTCTGATAAAGAATGGATCCGCTTTGACCAATCAAA
 TGTCCTTTGCAGTATTTTTGGTATCTGCATGGTGTTTTCTGGTCTGTGCCCA
 CTTCTTCTAGGCAGTGTGATGGCCATTGAGCGGTGTATTGGAGTCACAAA
 ACOCAATATTTCAATTCTACGAAAATTACATCCAAACATGTGAAAATGATGTTA

AGTGGTGTGTGCTTGTGTTTCTGCTTTTCATAGCTTTGCTGCCCATCCTTGGA
 CATCGAGACTATAAAATTCAGGCGTCGAGGACCTGGTGTGTTCTACAACAC
 5 AGAAGACATCAAAGACTGGGAAGATAGATTTTATCTTCTACTTTTTCTTTT
 CTGGGGCTCTTAGCCCTTGGTGTGTTTCATTGTTGTGCAATGCAATCACAGGA
 ATTACACTTTTAAGAGTTAAATTTAAAAGTCAGCAGCACAGACAAGGCAGA
 10 TCTCATCATTTGGAAATGGTAATCCAGCTCCTGGCGATAATGTGTGTCTCC
 TGTATTTGTTGGAGCCCATTTCTGGTTACAATGGCCAAATTGGAATAAAT
 GGAAATCATTCTCTGGAAACCTGTGAAACAACACTTTTTGCTCTCCGAATG
 15 GCAACATGGAATCAAATCTTAGATCCTTGGGTATATATTCTTCTACGAAAG
 GCTGTCTTAAGAATCTCTATAAGCTTGCCAGTCAATGCTGTGGAGTGCA
 GTCATCAGCTTACATATTTGGGAGCTTAGTTCCATTAAAAATTCCTTAAAG
 GTTGCTGCTATTTCTGAGTCACCAGTTGCAGAGAAATCAGCAAGCACCTA
 20 GCTTAATAGGACAGTAAATCTGTGTGGGGCTAGAACAAAAATTAAGACAT
 GTTTGGCAATATTTCAAGTTAGTTAAATACCTGTAGCCTAACTGGAAAATTC
 AGGCTTCATCATGTAGTTTG (SEQ.ID.NO.: 4).

- 25
4. Vecteur d'expression pour l'expression d'une protéine de récepteur de prostaglandine dans une cellule hôte recombinante, caractérisé en ce que ledit vecteur d'expression contient la molécule d'ADN de la revendication 3.
- 30
5. Cellule hôte exprimant une protéine de récepteur de prostaglandine recombinante, caractérisée en ce que ladite cellule hôte contient le vecteur d'expression de la revendication 4.
6. Procédé pour l'expression d'une protéine de récepteur de prostaglandine dans une cellule hôte recombinante, comprenant:
- 35
- (a) le transfert du vecteur d'expression de la revendication 4 dans une cellule hôte appropriée; et
 (b) la culture des cellules hôtes de l'étape a) dans des conditions permettant l'expression de la protéine de récepteur de prostaglandine à partir du vecteur d'expression.
7. Procédé d'identification de modulateurs d'une activité de récepteur de prostaglandine, comprenant:
- 40
- (a) l'association d'un modulateur d'activité de récepteur de prostaglandine avec un récepteur de prostaglandine, ledit récepteur étant caractérisé par la séquence d'acides aminés de la revendication 1 en partie ou en totalité; et
 (b) la mesure de l'effet d'un modulateur sur le récepteur de prostaglandine.
- 45
8. Anticorps se fixant spécifiquement à la protéine de récepteur de prostaglandine, ladite protéine étant caractérisée par la séquence d'acides aminés de la revendication 1.
- 50
- 55

GTGCGGAGGGACGAGGGCTGGACCACAGCCGGGCGCCGATCAGGATCTCCGG
 CTGGATCGGTGGAACCTGAGGCAGCGGCGGGCGCCATGGCACACCGAGCG
 GCTCCGTCTTCTGCTCCTCAGAGAGCCCGGCTGGCGCCTGGGATGACAAGATGTCT
 GGACTGCAATCCTGCACAGTTTGTAGAGGGAGATGACTTGAGTGGTTGGCTTTTATC
 TCCACAACAATGTCCATGAACAATTCCAACAGCTAGTGCTCCTGCAGCTGCGCTT
 CTTTCAACACAACCTGCCAGACGGAAACCGGCTTTCCGTATTTTTTTCAGTAATC
 TTCAATGACAGTGGGAATCTTGTCAAACAGCCTTGCCATCGCCATTCTCATGAAGGCA
 TATCAGAGATTTAGACAGAAGTCCAAGGCATCGTTTCTGCTTTTGGCCAGCGCCTG
 GTAACTACTGATTTCTTGGCCCATCTCATCAATGGAGCCATAGCAGTATTTGTATAT
 GCTTCTGATAAAGAATGGATCCGCTTTTGACCAATCAAATGTCCCTTGCAGTATTTTT
 GGTATCTGCATGGTGTCTTCTGGTCTGTGCCCACTTCTTCTAGGCAGTGTGATGGCC
 ATTGAGCGGTGATTGGAGTCACAAACCAATATTTCATTCTACGAAATTACATCC
 AAACATGTGAAATGATGTTAAGTGGTGTGCTGTGCTTGTGCTTTTCATAGCTTTG
 CTGCCCATCCTTGGACATCGAGACTATAAAATTCAGGCGTCGAGGACCTGGTGTTTC
 TACAACACAGAAACATCAAAGACTGGGAAGATAGATTTTATCTTCTACTTTTCTCT
 TTTCTGGGGCTCTTAGCCCTTGGTGTCTTTCATTTGTTGTGCAATGCAATCACAGGAATT
 ACACCTTTTAAGAGTTAAATTTAAAGTCAGCAGCACAGACAAGGCAGATCTCATCAT
 TTGGAAATGGTAATCCAGCTCCTGGCGATAATGTGTGCTCCTGTATTTGTTGGAGC
 CCATTTCTGGTTACAATGGCCCAACATTGGAATAAATGGAAATCATTTCTCTGGAAACC

FIG.1A


TGTGAAACAACACTTTTGGCTCTCCGAATGGCAACATGGAATCAAATCTTAGATCCT
 TGGGTATATATTCTTCTACGAAAGGCTGTCCCTTAAGAAATCTCTATAAGCTTGCCAGT
 CAATGCTGTGGAGTGCATGTCAATCAGCTTACATATTGGGAGCTTAGTTCATTAAA
 AATTCCTTAAAGGTTGCTGCTATTCTGAGTCACCAGTTGCAGAGAAATCAGCAAGC
 ACCTAGCTTAATAGGACAGTAAATCTGTGTGGGCTAGAACAAAAATTAAGACATGT
 TTGGCAATATTTCAAGTTAGTTAAATACCTGTAGCCTAACTGGAAATTCAGGCTTCA
 TCATGTAGTTTG

FIG.1B

MSMNNSKQLVSPAAALLSNTTCQTENRLSVFFSVIFMTVGILSNSLAIAI
 LMKAYQRFQKSKASFLLLASGLVITDFFGHLINGAIAVFVYASDKEWIR
 FDQSNVLCISIFGICMVFSGLCPLLLGSVMAIERCIGVTKPIFHSTKITSK
 HVKMMLSGVCLFAVFIALLPILGHRDYKIQASRTWCIFYNTEDIKDWE DRF
 YLLFSFLGLLALGVSLLCNAITGITLLRVKFKSQHRQGRSHHLEMVIQ
 LLAIMCVSCICWSPFLVTMANIGINGNHSLETCTTLFALRMATWNQILD
 PWVYILLRKAVLKNLYKLASQCCGVHVISLHIWELSSIKNSLKVA AIS ES
 PVAEKSAST

FIG. 2

20 nM PGF_{2α}



50 nA
2min

FIG. 3A

10 nM Fluprostenol




FIG. 3B




FIG. 4A




FIG. 4B




FIG.4C




FIG.5




FIG.6