

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 721 337 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:

05.12.2001 Bulletin 2001/49

(51) Int Cl.7: **A61K 31/495**, A61K 9/08,
A61K 47/40, A61K 47/10

(21) Application number: **94927639.8**

(86) International application number:
PCT/EP94/03169

(22) Date of filing: **22.09.1994**

(87) International publication number:
WO 95/08993 (06.04.1995 Gazette 1995/15)

(54) **ORAL FORMULATIONS OF AN ANTIFUNGAL**

ORALE ZUBEREITUNGEN VON ANTIMYKOTIKA

FORMULATIONS ORALES POUR UN ANTIFONGIQUE

(84) Designated Contracting States:

**AT BE CH DE DK ES FR GB GR IE IT LI LU NL PT
SE**

Designated Extension States:

LT SI

(74) Representative: **Quaghebeur, Luc et al**

**Janssen Pharmaceutica N.V.,
Patent Department,
Turnhoutseweg 30
2340 Beerse (BE)**

(30) Priority: **30.09.1993 US 129504**

(56) References cited:

WO-A-93/19061 US-A- 4 916 134

(43) Date of publication of application:

17.07.1996 Bulletin 1996/29

(73) Proprietor: **JANSSEN PHARMACEUTICA N.V.**
2340 Beerse (BE)

- **ANTIMICROB. AGENTS CHEMOTHER.**, vol.36,
no.2, February 1992 pages 477 - 480 J.S.
HOSTETLER ET AL. 'Effect of cyclodextrin on
the pharmacology of antifungal oral azoles'
- **INT. J. PHARMACEUT.**, vol.80, 1992 pages 253 -
258 J. PITHA ET AL. 'Preparation of drug:
hydroxypropylcyclodextrin complexes by a
method using ethanol or aqueous ammonium
hydroxide as co-solubilizer'

(72) Inventors:

- **FRANCOIS, Marc Karel Jozef**
B-2920 Kalmthout (BE)
- **DRIES, Willy Maria Albert Carlo**
B-2330 Merksplas (BE)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 721 337 B1

Description

[0001] The present invention is concerned with novel compositions of antifungal agents which have low solubility in aqueous media, a process for preparing said compositions and pharmaceutical dosage forms for oral administration comprising said novel compositions.

[0002] The development of efficacious pharmaceutical compositions of azole antifungals such as for example, itraconazole and saperconazole, is hampered considerably by the fact that said antifungals are only very sparingly soluble in water. The solubility and bioavailability of said compounds can be increased by complexation with cyclodextrins or derivatives thereof as described in WO 85/02767 and US-4,764,604. Alternatively, strongly acidic formulations ($\text{pH} \leq 1.5$) of itraconazole and saperconazole can be formed in which the active ingredients are partially dissolved. Obviously such strongly acidic formulations are useless for oral administration. Aqueous formulations comprising a co-solvent such as PEG 400 completely dissolve itraconazole at $\text{pH} 2.3 - 2.5$. However, these acidic formulations have problems with regard to ease-of-preparation, acceptability, palatability and especially bioavailability: upon administration said formulations can precipitate irreversibly, e.g. in the stomach. Acidic formulations comprising cyclodextrin or a derivative thereof might appear an obvious alternative, but the mere combinations prove to suffer from a number of similar problems, in particular difficulty-of-preparation, lack of stability (shelf life) and palatability, and unreliable absorption.

[0003] WO-93/19061, published on 30 September 1993, discloses that the individual isomers of itraconazole (and saperconazole) in aqueous cyclodextrin formulations have greater solubilities than itraconazole or saperconazole themselves (which consist of a mixture of four isomers), but does not address the palatability of the formulations.

[0004] In the publication by J. Hostetler et al., "Effect of cyclodextrin on the pharmacology of antifungal oral azoles", Antimicrobial Agents and Chemotherapy, Feb. 1992, pp 477-480, various aqueous oral solutions of saperconazole and itraconazole are described (one version uses hydroxypropylcyclodextrin ["HPCD"] in an acidified solution that also contains propylene glycol). The solutions are administered to mice by "gavage", i. e. directly into the stomach, in order to compare the bioavailability of the active ingredient from various vehicles.

[0005] US-4,916,134 discloses a series of novel antifungal agents. Examples 12 and 13 therein disclose low dose [0.1 % (w/v)] oral formulations that are practically non-aqueous but comprise an alcoholic solvent (polyethyleneglycol or glycerol) and lack cyclodextrin complexing agent.

[0006] In short, there still exists an important demand for easily prepared formulations of antifungal agents with good bioavailability and acceptable organoleptic

properties for oral administration.

[0007] The present invention relates to formulations for oral administration which comprise an antifungal, e. g. itraconazole or saperconazole, as active ingredient, a sufficient amount of a cyclodextrin or a derivative thereof as a solubilizer, an aqueous acidic medium as bulk liquid carrier and an alcoholic co-solvent that greatly simplifies the preparation of the composition. The formulations are rendered more palatable by adding one or more pharmaceutically acceptable intense sweeteners and one or more bulk sweeteners, and one or more pharmaceutically acceptable flavours.

[0008] A low-dosage formulation according to the present invention is suitable for treating patients suffering from fungal infections, particularly for treating AIDS patients with oral candidiasis infections. The need for reliable formulations of itraconazole (and saperconazole) in this indication is especially high because of resistance to fluconazole developing in *Candida* strains. Generally, 400 mg/day represents the minimum dose required to obtain meaningful plasma levels. Suitable oral formulations typically comprise from 0.5% to 1.5% (w/v), preferably about 1% (w/v) of the active ingredient.

[0009] A high-dosage formulation according to the present invention is suitable for treating patients suffering from systemic fungal infections. Suitable oral formulations for combatting systemic fungal infections typically comprise from 3% to 5%, preferably about 4% (w/v) of the active ingredient.

[0010] The formulations of the present invention are also suitable for the treatment of fungal infections in non-human animals, in particular for the treatment of dermatophytoses.

[0011] Itraconazole or (+)-cis-4-[4-[4-[4-[2-(2,4-dichlorophenyl)-2-(1H-1,2,4-triazol-1-yl-methyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-2,4-dihydro-2-(1-methylpropyl)-3H-1,2,4-triazol-3-one, is a broadspectrum antifungal compound developed for oral, parenteral and topical use and is disclosed in US-4,267,179. Its difluoro analog, saperconazole or (+)-cis-4-[4-[4-[4-[2-(2,4-difluorophenyl)-2-(1H-1,2,4-triazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]-phenyl]-2,4-dihydro-2-(1-methoxypropyl)-3H-1,2,4-triazol-3-one, has improved activity against *Aspergillus* spp. and is disclosed in US-4,916,134.

[0012] Appropriate cyclodextrin derivatives are α -, β -, γ -cyclodextrins or ethers and mixed ethers thereof wherein one or more of the hydroxy groups of the anhydroglucose units of the cyclodextrin are substituted with C_{1-6} alkyl, particularly methyl, ethyl or isopropyl; hydroxy C_{1-6} alkyl, particularly hydroxyethyl, hydroxypropyl or hydroxybutyl; carboxy C_{1-6} alkyl, particularly carboxymethyl or carboxyethyl; C_{1-6} alkylcarbonyl, particularly acetyl; C_{1-6} alkyloxy C_{1-6} alkyl or carboxy C_{1-6} alkyloxy C_{1-6} alkyl, particularly carboxymethoxypropyl or carboxyethoxypropyl; C_{1-6} alkylcarbonyloxy C_{1-6} alkyl, particularly 2-acetyloxy-

propyl. Especially noteworthy as complexants and/or solubilizers are β -CD, 2,6-dimethyl- β -CD, 2-hydroxyethyl- β -CD, 2-hydroxyethyl- γ -CD, 2-hydroxypropyl- γ -CD and (2-carboxymethoxy)propyl- β -CD, and in particular 2-hydroxypropyl- β -CD.

[0013] The term mixed ether denotes cyclodextrin derivatives wherein at least two cyclodextrin hydroxy groups are etherified with different groups such as, for example, hydroxypropyl and hydroxyethyl.

[0014] The average molar substitution (M.S.) is used as a measure of the average number of moles of alkoxy units per mole of anhydroglucose. In the cyclodextrin derivatives for use in the compositions according to the present invention the M.S. is in the range of 0.125 to 10, in particular of 0.3 to 3, or from 0.3 to 1.5. Preferably the M.S. ranges from 0.3 to 0.8, in particular from 0.35 to 0.5 and most particularly is about 0.4. M.S. values determined by NMR or IR preferably range from 0.3 to 1, in particular from 0.55 to 0.75.

[0015] The average substitution degree (D.S.) refers to the average number of substituted hydroxyls per anhydroglucose unit. In the cyclodextrin derivatives for use in the compositions according to the present invention the D.S. is in the range of 0.125 to 3, in particular of 0.2 to 2 or from 0.2 to 1.5. Preferably the D.S. ranges from 0.2 to 0.7, in particular from 0.35 to 0.5 and most particularly is about 0.4. D.S. values determined by NMR or IR preferably range from 0.3 to 1, in particular from 0.55 to 0.75.

[0016] More particular β - and γ -cyclodextrin hydroxyalkyl derivatives for use in the compositions according to the present invention are partially substituted cyclodextrin derivatives wherein the average degree of alkylation at hydroxyl groups of different positions of the anhydroglucose units is about 0% to 20% for the 3 position, 2% to 70% for the 2 position and about 5% to 90% for the 6 position. Preferably the amount of unsubstituted β - or γ -cyclodextrin is less than 5% of the total cyclodextrin content and in particular is less than 1.5%. Another particularly interesting cyclodextrin derivative is randomly methylated β -cyclodextrin.

[0017] Most preferred cyclodextrin derivatives for use in the present invention are those partially substituted β -cyclodextrin ethers or mixed ethers having hydroxypropyl, hydroxyethyl and in particular 2-hydroxypropyl and/or 2-(1-hydroxypropyl) substituents.

[0018] The most preferred cyclodextrin derivative for use in the compositions of the present invention is hydroxypropyl- β -cyclodextrin having a M.S. in the range of from 0.35 to 0.50 and containing less than 1.5% unsubstituted β -cyclodextrin. M.S. values determined by NMR or IR preferably range from 0.55 to 0.75.

[0019] Substituted cyclodextrins can be prepared according to procedures described in US-3,459,731, EP-A-0,149,197, EP-A-0,197,571, US-4,535,152, WO-90/12035 and GB-2,189,245. Other references describing cyclodextrins for use in the compositions according to the present invention, and which provide a guide for

the preparation, purification and analysis of cyclodextrins include the following : "Cyclodextrin Technology" by József Szejtli, Kluwer Academic Publishers (1988) in the chapter Cyclodextrins in Pharmaceuticals; "Cyclodextrin Chemistry" by M.L. Bender et al., Springer-Verlag, Berlin (1978); "Advances in Carbohydrate Chemistry", Vol. 12 Ed. by M.L. Wolfrom, Academic Press, New York (1978) in the chapter The Schardinger Dextrins by Dexter French at p. 189-260; "Cyclodextrins and their Inclusions Complexes" by J. Szejtli, Akademiai Kiado, Budapest, Hungary (1982); I. Tabushi in Acc. Chem. Research, 1982, 15, p. 66-72; W. Sanger, Angewandte Chemie, 92, p. 343-361 (1981); A. P. Croft and R. A. Bartsch in Tetrahedron, 39, p. 1417-1474 (1983); Irie et al. Pharmaceutical Research, 5, p. 713-716, (1988); Pitha et al. Int. J. Pharm. 29, 73, (1986); DE 3,118,218; DE-3,317,064; EP-A-94,157; US-4,659,696; and US-4,383,992. The low-dosage oral formulations according to the present invention typically comprise from 20% to 60% (w/v), preferably about 40% (w/v) of the cyclodextrin. The high-dosage formulations typically comprise from 50% to 80% (w/v), preferably about 60% (w/v) of the cyclodextrin derivative.

[0020] In order to increase the rate of dissolution of the poorly soluble antifungal during the manufacturing process, an alcoholic co-solvent is employed in the formulations according to the present invention. For this purpose, preference is given to those alcoholic co-solvents that have good dissolving power for itraconazole and/or saperconazole, in particular ethanol, propylene glycol and glycerol, especially propylene glycol. Without the alcoholic co-solvent, the dissolution of itraconazole or saperconazole in an aqueous acidic cyclodextrin medium is very slow, requiring a viscous suspension to be stirred for a prohibitively long time until complete dissolution is obtained. Addition of the alcoholic co-solvent, in the range of 1% (v/v) to 20% (v/v), preferably about 10% (v/v), increases the dissolution rate of the antifungal agent in an aqueous acidic cyclodextrin medium by a factor of at least 5 (when used at 10% (v/v)) and thus considerably shortens and simplifies the production process.

[0021] As a bulk liquid carrier there is used an acidic aqueous medium. Preferably the acidity of said carrier derives from a strong, pharmaceutically acceptable acid such as hydrochloric acid. The bioavailability of the antifungal agent and the organoleptic properties of the oral formulations are affected contrariwise by the acidity. An optimum effect can be obtained at pH 2.0 ± 0.1 : that is, at this pH value, a sufficiently stable and bioavailable antifungal formulation is obtainable, the organoleptic properties of which can be rendered acceptable.

[0022] Not surprisingly, the ingredients thus far described yield a fairly strong-tasting potion when mixed with one another. Besides the acid taste due to the low pH, a bitter taste originating from the active ingredient, and possibly from the co-solvent (e.g. in the case of propylene glycol), is also present. Optimum taste masking

can be obtained by the use of two types of adjuvants, namely pharmaceutically acceptable sweeteners and flavours. Sweeteners are the more important additives in the low-dosage formulations, whereas the flavours are more important in the high-dosage formulations.

[0023] The pharmaceutically acceptable sweeteners comprise at least one intense sweetener such as saccharin, sodium or calcium saccharin, aspartame, acesulfame potassium, sodium cyclamate, alitame, a dihydrochalcone sweetener, monellin, stevioside or sucralose (4,1',6'-trichloro-4,1',6'-trideoxygalactosucrose), preferably saccharin, sodium or calcium saccharin, and one or more bulk sweeteners such as sorbitol, mannitol, fructose, sucrose, maltose, isomalt, glucose, hydrogenated glucose syrup, xylitol, caramel or honey.

[0024] The intense sweetener is conveniently employed in low concentrations. For example, in the case of sodium saccharin, the concentration may range from 0.04% to 0.1% (w/v) based on the total volume of the final formulation, and preferably is about 0.06% in the low-dosage formulations and about 0.08% in the high-dosage ones. The bulk sweetener can effectively be used in larger quantities ranging from about 10% to about 35%, preferably from about 10% to 15% (w/v). In the high-dosage formulations the cyclodextrin derivative behaves as a bulk sweetener and none of the aforementioned bulk sweeteners needs to be added.

[0025] The pharmaceutically acceptable flavours which can mask the bitter tasting ingredients in the low-dosage formulations are preferably fruit flavours such as cherry, raspberry, black currant or strawberry flavour. A combination of two cherry flavours was found to yield very good results in an itraconazole formulation both as regards physico-chemical stability as well as regards organoleptic acceptability. In the high-dosage formulations stronger flavours are required such as Caramel Chocolate flavour, Mint Cool flavour, Fantasy flavour and the like pharmaceutically acceptable strong flavours. Each flavour may be present in the final composition in a concentration ranging from 0.05% to 1% (w/v). Combinations of said strong flavours are advantageously used. Preferably a flavour is used that does not undergo any change or loss of taste and colour under the acidic conditions of the formulation.

[0026] A preferred high-dosage formulation according to the present invention comprises by weight or by volume based on the total volume of the formulation :

- (a) 4% (w/v) itraconazole;
- (b) 60% (w/v) hydroxypropyl- β -cyclodextrin;
- (c) 10% (v/v) propylene glycol;
- (d) acid and base to adjust the pH of the composition within the range of 2.0 ± 0.1 ;
- (e) 0.08% (w/v) sodium saccharin;
- (f) up to 1% (w/v) of one or more strong flavours; and
- (g) water.

[0027] The preparation of the formulations according

to the present invention will hereafter be described with regard to a preferred low-dosage formulation having the following composition (% are by weight or by volume based on the total volume of the formulation):

- (a) 1% (w/v) itraconazole;
- (b) 40% (w/v) hydroxypropyl- β -cyclodextrin;
- (c) 10% (v/v) propylene glycol;
- (d) acid and base to adjust the pH of the composition within the range of 2.0 ± 0.1 ;
- (e) 0.06% (w/v) sodium saccharin;
- (f) 19% (v/v) sorbitol (70%) non-crystallizing solution;
- (g) up to 1% (w/v) of one or more cherry flavours; and
- (h) water.

[0028] Optionally, the above preferred low-dosage formulation further comprises up to 0.1%, in particular 0.02% caramel sweetener.

[0029] Similar formulations can be prepared with saperconazole, though other flavours may be preferred then.

[0030] Said process of preparation comprises the steps of

- (a) dissolving the active ingredient in the alcoholic co-solvent and acid;
- (b) dissolving the cyclodextrin in water and adding thereto the solution prepared in (a) while stirring until homogenous;
- (c) adding the sweeteners and the flavour(s);
- (d) adjusting the acidity to pH 2.0 ± 0.1 and
- (e) diluting the formulation to the desired end-volume.

[0031] In particular, for preparing 1 litre of the aforementioned preferred formulation 100 ml of propylene glycol is treated with 3.76 ml concentrated HCl, stirred and slightly heated. 10 g itraconazole is added and stirring is continued until homogeneous.

In a separate vessel, 400 g hydroxypropyl- β -cyclodextrin is dissolved in 400 ml distilled water. The solution of the active ingredient is added slowly to the cyclodextrin solution while stirring. The sorbitol solution (190 ml) is added and stirred till homogeneous. The sodium saccharin (0.6 g) is dissolved in 50 ml distilled water and added to the mixture. The flavours are added and the pH of the mixture (about 1.7) is adjusted with a 10 N NaOH solution to pH 2.0 ± 0.1 . The resulting solution is diluted with distilled water to an end volume of 1 litre. A pharmaceutical dosage form is obtained by filtering the previous solution and filling it into suitable containers. e.g. in 100 ml glass bottles with a screw cap. The pharmaceutical dosage form advantageously comprises a minimal volume of air above the solution, preferably an inert gas such as nitrogen. Besides the exclusion of air (oxygen), storage at temperatures below 25°C also ben-

officially affects the maximum shelf life of the formulation for oral administration.

Claims

1. An aqueous solution for oral administration comprising :

- (a) itraconazole or saperconazole;
- (b) a sufficient amount of a cyclodextrin or a derivative thereof, to act as a solubilizer for the itraconazole or saperconazole;
- (c) an aqueous acidic medium as bulk liquid carrier;
- (d) an alcoholic co-solvent selected from the group consisting of ethanol, propylene glycol and glycerol;
- (e) sufficient amounts of one or more pharmaceutically acceptable intense sweeteners and one or more bulk sweeteners; and
- (f) a sufficient amount of one or more pharmaceutically acceptable flavors, so as to taste mask the aqueous solution.

2. A formulation according to claim 1 wherein the cyclodextrin is hydroxypropyl- β -cyclodextrin having an M.S. in the range of 0.35 to 3 and containing less than 5% unsubstituted β -cyclodextrin.

3. A formulation according to claim 1 wherein the alcoholic co-solvent is propylene glycol.

4. A formulation according to claim 1 having a pH of 2.0 ± 0.1 .

5. A formulation according to claim 1 wherein the intense sweetener is selected from the group consisting of saccharin, sodium or calcium saccharin and the bulk sweetener is selected from the group consisting of sorbitol, mannitol, fructose, sucrose, maltose, glucose, caramel or honey.

6. A formulation according to claim 1 comprising by weight or by volume based on the total volume of the formulation :

- (a) 4% (w/v) itraconazole;
- (b) 60% (w/v) hydroxypropyl- β -cyclodextrin;
- (c) 10% (v/v) propylene glycol;
- (d) acid and base to adjust the pH of the composition within the range of 2.0 ± 0.1 ;
- (e) 0.08% (w/v) sodium saccharin;
- (f) up to 1% (w/v) of one or more flavours; and
- (g) water.

7. A formulation according to claim 1 comprising by weight or by volume based on the total volume of

the formulation :

- (a) 1 % (w/v) itraconazole or saperconazole;
- (b) 40% (w/v) hydroxypropyl- β -cyclodextrin;
- (c) 10% (v/v) propylene glycol;
- (d) acid or base to adjust the pH of the composition within the range of 2.0 ± 0.1 ;
- (e) 0.06% (w/v) sodium saccharin;
- (f) 19% (v/v) sorbitol (70%) non-crystallizing solution;
- (g) up to 1% (w/v) of one or more flavours;
- (h) 0.02 % of a caramel sweetener, and
- (i) water.

8. A process of preparing a formulation as claimed in claim 1, **characterized in that** said process comprises the steps of :

- (a) dissolving the active ingredient in the alcoholic co-solvent and acid;
- (b) dissolving the cyclodextrin in water and adding thereto the solution prepared in (a) while stirring until homogenous;
- (c) adding the sweetener(s) and the flavour(s);
- (d) adjusting the acidity to pH 2.0 ± 0.1 ; and
- (e) diluting the formulation to the desired end-volume.

Patentansprüche

1. Wäßrige Lösung zur oralen Verabreichung mit:

- (a) Itraconazol oder Saperconazol;
- (b) einem Cyclodextrin oder einem seiner Derivate in einer zur Solubilisierung von Itraconazol oder Saperconazol ausreichenden Menge;
- (c) einem wäßrig-sauren Medium als hauptsächlichem flüssigen Träger;
- (d) einem alkoholischen Hilfslösungsmittel aus der Gruppe Ethanol, Propylenglykol und Glycerin;
- (e) einem oder mehreren pharmazeutisch unbedenklichen Süßmitteln mit hoher Süßkraft sowie einem oder mehreren Bulk-Süßmitteln in ausreichenden Mengen; sowie
- (f) einem oder mehreren pharmazeutisch unbedenklichen Aromastoffen in einer zur Geschmacksverdeckung der wäßrigen Lösung ausreichenden Menge.

2. Formulierung nach Anspruch 1, wobei es sich bei dem Cyclodextrin um Hydroxypropyl- β -cyclodextrin mit einem M.S. im Bereich von 0,35 bis 3, das weniger als 5% unsubstituiertes β -Cyclodextrin enthält, handelt.

3. Formulierung nach Anspruch 1, wobei es sich bei

dem alkoholischen Hilfslösungsmittel um Propylenglykol handelt.

4. Formulierung nach Anspruch 1 mit einem pH-Wert von $2,0 \pm 0,1$.

5

5. Formulierung nach Anspruch 1, wobei das Süßmittel mit hoher Süßkraft aus der Gruppe Saccharin, Saccharin-Natrium und Saccharin-Calcium und das Bulk-Süßmittel aus der Gruppe Sorbit, Mannit, Fructose, Saccharose, Maltose, Glucose, Karamel und Honig stammt.

10

6. Formulierung nach Anspruch 1 mit den folgenden Bestandteilen (Gewicht bzw. Volumen in Bezug auf das Gesamtvolumen der Formulierung):

15

- (a) 4% (w/v) Itraconazol;
- (b) 60% (w/v) Hydroxypropyl- β -cyclodextrin;
- (c) 10% (v/v) Propylenglykol;
- (d) Säure und Base zur Einstellung des pH-Werts der Zusammensetzung in dem Bereich von $2,0 \pm 0,1$;
- (e) 0,08% (w/v) Saccharin-Natrium;
- (f) bis zu 1% (w/v) eines Aromastoffs bzw. mehrerer Aromastoffe; sowie
- (g) Wasser.

20

25

7. Formulierung nach Anspruch 1 mit den folgenden Bestandteilen (Gewicht bzw. Volumen in Bezug auf das Gesamtvolumen der Formulierung):

30

- (a) 1% (w/v) Itraconazol oder Saperconazol;
- (b) 40% (w/v) Hydroxypropyl- β -cyclodextrin;
- (c) 10% (v/v) Propylenglykol;
- (d) Säure und Base zur Einstellung des pH-Werts der Zusammensetzung in dem Bereich von $2,0 \pm 0,1$;
- (e) 0,06% (w/v) Saccharin-Natrium;
- (f) 19% (v/v) Sorbit-Lösung (70%), nicht kristallisierend;
- (g) bis zu 1% (w/v) eines Aromastoffs bzw. mehrerer Aromastoffe;
- (h) 0,02% eines Karamel-Süßmittels; sowie
- (i) Wasser.

35

40

45

8. Verfahren zur Herstellung einer Formulierung nach Anspruch 1, **dadurch gekennzeichnet, daß** das Verfahren folgende Schritte umfaßt:

50

- (a) Lösen des Wirkstoffs in dem alkoholischen Hilfslösungsmittel und der Säure;
- (b) Lösen des Cyclodextrins in Wasser, Zugabe der unter (a) hergestellten Lösung unter Rühren, bis die Mischung homogen ist;
- (c) Zugabe des Süßmittels bzw. der Süßmittel sowie des Aromastoffs bzw. der Aromastoffe;
- (d) Einstellen der Azidität auf einen pH-Wert

55

von $2,0 \pm 0,1$; sowie

(e) Verdünnen der Formulierung auf das gewünschte Endvolumen.

Revendications

1. Solution aqueuse destinée à l'administration par voie orale comprenant :

- (a) l'itraconazole ou le saperconazole ;
- (b) une quantité suffisante d'une cyclodextrine ou un dérivé de celle-ci, pour agir comme solubilisant pour l'itraconazole ou le saperconazole ;
- (c) un milieu acide aqueux comme support liquide en masse ;
- (d) un co-solvant alcoolique choisi dans le groupe constitué d'éthanol, de propylèneglycol et de glycérol ;
- (e) des quantités suffisantes de un ou plusieurs édulcorants intenses pharmaceutiquement acceptables et un ou plusieurs édulcorants de masse ; et
- (f) une quantité suffisante de un ou plusieurs arômes pharmaceutiquement acceptables, de manière à masquer le goût de la solution aqueuse.

2. Formulation selon la revendication 1, **caractérisée en ce que** la cyclodextrine est l'hydroxypropyl- β -cyclodextrine ayant une S.M. dans la gamme de 0,35 à 3 et contenant moins de 5% de β -cyclodextrine non substituée.

3. Formulation selon la revendication 1, **caractérisée en ce que** le co-solvant alcoolique est le propylèneglycol.

4. Formulation selon la revendication 1, ayant un pH de $2,0 \pm 0,1$.

5. Formulation selon la revendication 1, **caractérisée en ce que** l'édulcorant intense est choisi dans le groupe constitué de saccharine, de saccharine sodique ou calcique et l'édulcorant de masse est choisi dans le groupe constitué de sorbitol, de mannitol, de fructose, de saccharose, de maltose, de glucose, de caramel ou de miel.

6. Formulation selon la revendication 1, comprenant en poids ou en volume basé sur le volume total de la formulation :

- (a) 4% (p/v) d'itraconazole ;
- (b) 60% (p/v) d'hydroxypropyl- β -cyclodextrine ;
- (c) 10% (v/v) de propylèneglycol ;
- (d) d'acide ou de base pour régler le pH de la

composition dans la gamme de $2,0 \pm 0,1$;
 (e) 0,08% (p/v) de saccharine sodique ;
 (f) jusqu'à 1% (p/v) de un ou plusieurs arômes ;
 et
 (g) de l'eau.

5

7. Formulation selon la revendication 1, comprenant en poids ou en volume basé sur le volume total de la formulation :

10

(a) 1% (p/v) d'itraconazole ou de saperconazole ;
 (b) 40% (p/v) d'hydroxypropyl- β -cyclodextrine ;
 (c) 10% (v/v) de propylèneglycol ;
 (d) d'acide ou de base pour régler le pH de la composition dans la gamme de $2,0 \pm 0,1$;
 (e) 0,06% (p/v) de saccharine sodique ;
 (f) 19% (v/v) de solution de sorbitol (70%) non cristallisable ;
 (g) jusqu'à 1% (p/v) de un ou plusieurs arômes ;
 (h) 0,02% d'un édulcorant de caramel, et
 (i) de l'eau.

15

20

8. Procédé de préparation d'une formulation selon la revendication 1, **caractérisé en ce que** ledit procédé comprend les étapes consistant à :

25

(a) dissoudre le principe actif dans le co-solvant alcoolique et l'acide ;
 (b) dissoudre la cyclodextrine dans l'eau et ajouter à celle-ci la solution préparée en (a) sous agitation jusqu'à homogénéité ;
 (c) ajouter le ou les édulcorant(s) et 1'(les) arôme(s) ;
 (d) régler l'acidité à pH $2,0 \pm 0,1$; et
 (e) diluer la formulation jusqu'au volume final désiré.

35

40

45

50

55