

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 969 080 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
05.01.2000 Bulletin 2000/01

(51) Int. Cl.⁷: **C11D 1/83**, C11D 3/33,
C11D 3/43, C11D 3/20

(21) Application number: **99112847.1**

(22) Date of filing: **02.07.1999**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **03.07.1998 JP 18872398**

(71) Applicant:
**Showa Denko Kabushiki Kaisha
Tokyo 105-8518 (JP)**

(72) Inventors:
• **Saito, Makoto,**
c/o KAWASAKI WORKS, SHOWA DENKO K.K.
Kawasaki-shi, Kanagawa 210-0865 (JP)

- **Yamamoto, Tohru,**
KAWASAKI WORKS, SHOWA DENKO K.K.
Kawasaki-shi, Kanagawa 210-0865 (JP)
- **Nambu, Nobuyoshi,**
c/o YOKKAICHI WORKS
Yokkaichi-shi, Mie 510-0886 (JP)
- **Furukawa, Masanori,**
c/o YOKKAICHI WORKS
Yokkaichi-shi, Mie 510-0886 (JP)

(74) Representative:
Strehl Schübel-Hopf & Partner
Maximilianstrasse 54
80538 München (DE)

(54) **Liquid detergent composition**

(57) A high-concentration liquid detergent composition, which does not have strong alkalinity, which has excellent compatibility even if the surfactant concentration is increased, and which is free of white turbidity, comprising (A) one or more surfactants selected from a nonionic surfactant and an anionic surfactant, (B) one or more builders selected from an amine salt and an alkali metal-amine mixed salt of aminopolycarboxylic acid chelating agent, and (C) one or more solubilizing agents selected from alcohols, with component (A) being present in a concentration of 20 wt% or more.

EP 0 969 080 A1

DescriptionFIELD OF THE INVENTION

5 **[0001]** The present invention relates to a liquid detergent composition comprising a surfactant and an aminopolycarboxylic add-based organic builder. More specifically, the present invention relates to a high concentration liquid detergent composition having excellent compatibility with a surfactant, superior resistance against white turbidity and good biodegradability.

10 BACKGROUND OF THE INVENTION

[0002] In liquid detergents, particularly in liquid detergents used as a household detergent a nonionic surfactant and/or an anionic surfactant are used as the surfactant and a builder of various types is added to increase detergency. Various types of organic or inorganic builders can be used but almost all builders have a Na salt structure.

15 **[0003]** In recent years, from the standpoint of environmental conservation, studies are aggressively being made on the use of an aminopolycarboxylic acid having excellent biodegradability, particularly an amino acid diacetic acid-type chelating agent as the builder. Aminopolycarboxylic acids including amino add diacetic acid chelating agents are used in the form of a synthetic reaction solution as it is on account of the cost. Therefore, in almost all cases, these are Na salts which cause strong alkalinity. For example, JP-A-10-25494 (the term "JP-A" as used herein means an "unexamined published Japanese patent application") discloses a liquid detergent for dishes, comprising an aspartic acid diacetic acid builder, where an alkali metal salt is used. JP-A-10-36888 discloses a liquid detergent comprising an aminopolycarboxylic acid builder, where an amine salt is disclosed on the same level with an alkali metal salt and in Examples, a Na salt is used. Moreover, the maximum concentration of the surfactant in this composition is as low as 10 wt%, thus, the problem of white turbidity on the high concentration side is not solved.

25 **[0004]** The use of a builder has a problem in that the solubility or clouding point of the surfactant as the main agent decreases and thereby the compatibility is deteriorated to cause, for example, white turbidity or separation of the white turbidity into two layers with the elapse of time. In order to prevent these, a technique of diluting the liquid detergent composition with water to reduce the concentration of the surfactant or a technique of adding a solubilizing agent has been heretofore used. However, the former case of diluting the liquid detergent composition with water inevitably results in an increase in the cost of transporting the detergent and this is commercially disadvantageous. In the latter case, it is known to newly add a surfactant as a solubilizing agent or add urea, an ether alcohol or a lower alcohol. However, the effect is not sufficiently high for obtaining a high concentration liquid detergent composition.

SUMMARY OF THE INVENTION

35 **[0005]** An object of the present invention is to obtain a liquid detergent which has excellent biodegradability, which does not assume strong alkalinity, which has excellent compatibility even if the surfactant concentration is increased, and which is free of white turbidity.

[0006] As a result of extensive investigations to solve the above-described problems, the present inventors have found that when a nonionic surfactant and/or anionic surfactant is used at a high concentration as the main surfactant of a liquid detergent and an aminopolycarboxylic acid chelating agent is used as a builder, by using an amine salt or an alkali metal-amine mixed salt as the counter ion of the builder and at the same time, adding an alcohol thereto, the compatibility is increased and the white turbidity can be prevented. The present invention has been accomplished based on this finding.

45 **[0007]** The present invention relates to the liquid detergent composition described below.

[0008] In one embodiment, the invention provides a liquid detergent composition comprising:

- (A) one or more surfactants selected from the group consisting of a nonionic surfactant and an anionic surfactant,
- (B) one or more builders selected from the group consisting of an amine salt and an alkali metal-amine mixed salt
- 50 of an aminopolycarboxylic acid chelating agent, and
- (C) one or more solubilizing agents selected from the group consisting of alcohols,

with component (A) being present in a concentration of 20 wt% or more.

[0009] In another embodiment, the invention provides a liquid detergent composition comprising:

- 55 (A) from 20 to 60 wt% of one or more surfactants selected from the group consisting of a nonionic surfactant and an anionic surfactant,

(B) from 0.1 to 15 wt% of one or more builders selected from the group consisting of an amine salt and an alkali metal-amine mixed salt of aminopolycarboxylic acid chelating agent,

(C) from 0.5 to 30 wt% of one or more solubilizing agents selected from the group consisting of alcohols, and

(D) the balance of water.

[0010] A further embodiment comprises a liquid detergent composition as described in (1) or (2), wherein the aminopolycarboxylic acid is an amino acid-N,N-diacetic acid.

[0011] An even further embodiment is a liquid detergent composition as described in (1) or (2), wherein the aminopolycarboxylic acid is glutamic acid-N,N-diacetic acid, α -alanine-N,N-diacetic acid, β -alanine-N,N-diacetic acid, aspartic acid-N,N-diacetic acid, glycine-N,N-diacetic acid or serine-N,N-diacetic acid.

[0012] Additionally, another embodiment is a liquid detergent composition as described in any one of (1) to (4), wherein the alcohol is a glycol.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0013] The surfactant used as component (A) in the present invention is one or more selected from a nonionic surfactant and an anionic surfactant. Among these, a nonionic surfactant has a higher solubility than an anionic surfactant, accordingly, it is suitable for preparing a high concentration liquid detergent composition. The surfactant is used at a concentration of 20 wt% or more in the liquid detergent composition.

[0014] Examples of suitable nonionic surfactants include polyoxyethylene alkyl ether, polyoxyethylene alkyl phenyl ether, fatty acid diethanolamide, amine oxide and amidoamine oxide. Examples of appropriate anionic surfactants include alkyl sulfate, alkylpolyoxyethylene sulfate, alkylbenzenesulfonate, α -olefinsulfonate, fatty acid salts, ether carboxylates and monoalkylphosphates. A number of these maybe commercially available.

[0015] The aminopolycarboxylic acid-based chelating agent used in the builder as component (B) of the present invention is a chelating agent containing one or more, preferably one or two amino groups and two or more, preferably three or four carboxyl groups, and having a chelating action. The amino group may be a secondary amino group or a tertiary amino group. The chelating agent is preferably an amino acid-N,N-diacetic acid.

[0016] An amino acid-N,N-diacetic acid is a compound represented by the following formula (1)





wherein A-NH₂ represents an amino acid, and -N represents an amino group in the amino acid.



[0017] Examples of amino acid-N,N-diacetic acids include glutamic acid-N,N-diacetic acid, α -alanine-N,N-diacetic acid, β -alanine-N,N-diacetic acid, aspartic acid-N,N-diacetic acid, glycine-N,N-diacetic acid and serine-N,N-diacetic acid. These amino acid-N,N-diacetic acids may be synthesized by the known Strecker reaction using, for example, an amino acid as the raw material or may be synthesized from a monochloroacetic acid.

[0018] Other than the amino acid-N,N-diacetic acid, examples of aminopolycarboxylic acid chelating agents include ethylenediaminedisuccinic acid and iminodisuccinic acid. These may also be synthesized by a known method and used. Examples of preferred aminopolycarboxylic acid chelating agents are shown below by their structural formulae.


Glutamic acid-N,N-diacetic acid

 α -Alanine-N,N-diacetic acid


Aspartic acid-N,N-diacetic acid

 β -Alanine-N,N-diacetic acid


Glycine-N,N-diacetic acid




Ethylenediaminedisuccinic acid



Iminodisuccinic acid



Serine-N,N-diacetic acid



[0019] On account of the cost, amino acid-N,N-diacetic acids and other aminopolycarboxylic acid-based chelating agents are generally obtained as an aqueous solution where all carboxyl groups have a Na salt as the counter ion. However, in the present invention, the aminopolycarboxylic acid for use in the builder as component (B) is used in the form of an amine salt or an alkali metal-amine mixed salt. The effect may be obtained using primary, secondary and tertiary amines, however, the amine salt used is preferably an alkanolamine salt, more preferably triethanolamine salt or dieth-

anolamine salt. The alkali metal salt in the alkali metal-amine mixed salt is preferably a K salt because it has a higher compatibility with the surfactant than an Na salt, however, in view of the cost, a Na salt is more preferred.

[0020] In order to convert the aminopolycarboxylic acid chelating agent obtained in the form of a Na salt into an alkali metal-amine mixed salt, the Na salt aqueous solution obtained may be treated to remove Na using an H-type cation exchange resin or by electrodialysis, or may be subjected to acid deposition to obtain a crystal having an H type carboxyl group. Thereafter, the resulting solution as it is or in the case of a crystal, a solution obtained by dissolving the crystals in water, is mixed with a mixture of an alkali metal hydroxide such as potassium hydroxide or sodium hydroxide with an amine and thereby an aqueous solution of the objective salt can be obtained. Also, after the removal of Na, the solution or crystal may be added to the formulation of a liquid detergent and then an amine or another alkali metal hydroxide may be added thereto so as to obtain the objective salt.

[0021] An aminopolycarboxylic acid-based chelating agent, particularly the aminodiacetic acid-type chelating agent is known to exert a good chelate effect namely a builder effect on the neutral to weakly alkali side at a pH of from 8 to 12. Therefore, the amount of the amine or alkali metal hydroxide added is preferably controlled so that the above-described pH region can be created. The larger the amount of the amine salt is, the higher is the compatibility with the surfactant, however, the ratio in the mixed salt of alkali metal and amine should be determined taking account of the cost for the removal of Na or the cost of amine.

[0022] The alcohol used as the solubilizing agent in the present invention is not particularly limited. However, a primary, secondary or tertiary alcohol, preferably a primary or secondary alcohol, having from 1 to 5 carbon atoms, preferably from 2 to 4 carbon atoms is preferred. The alcohol maybe a general alcohol such as ethanol and isopropyl alcohol, however, since these are volatile and highly inflammable, a commonly used and inexpensive glycol such as propylene glycol and ethylene glycol is preferably used.

[0023] The liquid detergent composition of the present invention preferably has a composition ratio such that the surfactant as component (A) is from 20 to 60 wt%, preferably from 30 to 50 wt%, the builder as component (B) is from 0.1 to 15 wt%, preferably from 1 to 10 wt%, the solubilizing agent as component (C) is from 0.5 to 30 wt%, preferably from 1 to 20 wt%, and the balance is water as component (D). This composition ratio is the proportion among four components (A) to (D). The ratio of builder as component (B) based on the surfactant as component (A) is preferably from 2 to 20 wt%, more preferably from 5 to 15 wt%, and the solubilizing agent as component (C) based on the surfactant as component (A) is preferably from 5 to 40 wt%, more preferably from 10 to 30 wt%.

[0024] If the ratio of the surfactant as component (A) is less than the above-described range, the cost for transportation or packaging of the liquid detergent increases and this is commercially disadvantageous. If it exceeds the range, the desired compatibility can be attained only within narrow limits. As a result, flexibility in the formulation may decrease or the viscosity may increase and the handleability is disadvantageously impaired.

[0025] If the ratio of the builder as component (B) is less than the above-described range, the cleaning effect of the liquid detergent composition decreases, whereas if it exceeds the range, the compatibility with the surfactant occurs only within narrow limits and flexibility in formulation disadvantageously decreases.

[0026] If the ratio of the solubilizing agent as the component (C) is less than the above-described range, compatibility between components (A) and (B) is reduced, whereas if it exceeds the range, flexibility in formulation disadvantageously decreases.

[0027] The liquid detergent composition of the present invention may contain, if desired, other additives in any optional ratio, such as a viscosity controlling agent, a moisture retaining agent, an antiseptic, an antifungal agent, a dye and a perfume.

[0028] The detergent composition of the present invention may be used for all uses as a detergent. However, this is suitably used in particular as a home detergent for cleaning dotes or household materials.

[0029] The detergent composition of the present invention may be used as it is or may be used after diluting it with water because the concentration is high.

[0030] According to the present invention, a builder comprising an amine salt or amine-alkali metal mixed salt of an aminopolycarboxylic add and a solubilizing agent comprising an alcohol are mixed each in a specific ratio with a specific surfactant, therefore, a high-concentration liquid detergent composition which does not have strong alkalinity, which has excellent compatibility even if the surfactant concentration is increased,, and which is free of white turbidity can be obtained.

EXAMPLES

[0031] The present invention is described below by referring to the Examples. In the Examples, "%" is "% by weight (wt%)".

Examples 1 to 16 and Comparative Examples 1 to 18:

[0032] The following surfactant, builder and solubilizing agent comprising an alcohol were added at the ratio shown in Tables 1 and 2, and the balance was water. The resulting solutions each was thoroughly mixed and left standing at an ordinary temperature or 40°C. Then, compatibility was visually observed. Those underwent separation into two layers after standing each was again mixed when observing and the solution state was observed. The formulations and visual results are shown together in Tables 1 and 2 below.

[0033] The following compounds were used.

Surfactant:

Nonionic surfactant:

EMULGEN 108, produced by Kao Corporation (trademark, ethylene oxide alkyl ether nonionic)

Anionic surfactant:

LAS, reagent, sodium linear alkylbenzenesulfonate (free of sodium sulfate)

Builder:

GLDA: glutamic acid-N,N-diacetic acid
 α AD: α -alanine-N,N-diacetic acid
 β AD: β -alanine-N,N-diacetic acid
SDA: serine-N,N-diacetic acid
ASDA: aspartic acid-N,N-diacetic acid
NTA: glycine-N,N-diacetic acid
EDDS: ethylenediaminedisuccinic acid
IDS: iminodisuccinic acid

Amine:

TEA: triethanolamine
DEA: diethanolamine
MEA: monoethanolamine

Solubilizing Agent:

PG: propylene glycol
IPA: isopropyl alcohol

[0034] In Tables 1 and 2, "-2Na2TEA" after "GLDA" indicates the counter ion of the carboxyl group within the molecule and means a mixed salt of 2Na and 2TEA. "-2Na2H" means 2Na salt in the low pH state. In either case, the concentration is the value calculated when the carboxyl group is the Na salt.

Table 1

Example	Surfactant	Builder	Solubilizing Agent	Compatibility
1	EMULGEN 108 40%	GLDA-2Na2TEA 5%	PG 12.5%	ordinary temperature: clear 40°C: clear
2	EMULGEN 108 40%	GLDA-2Na2TEA 5%	PG 6.9%	ordinary temperature: clear 40°C: clear
3	EMULGEN 108 40%	GLDA-1Na3TEA 5%	PG 4.8%	ordinary temperature: clear 40°C: clear

EP 0 969 080 A1

Table 1 (continued)

Example	Surfactant	Builder	Solubilizing Agent	Compatibility
4	EMULGEN 108 40%	GLDA-2Na2DEA 5%	PG 6.7%	ordinary temperature: clear 40°C: clear
5	EMULGEN 108 40%	GLDA-1Na3DEA 5%	PG 4.7%	ordinary temperature: clear 40°C: clear
6	EMULGEN 108 40%	GLDA-1Na3DEA 5%	IPA 4.7%	ordinary temperature: clear 44°C: clear
7	EMULGEN 108 40%	GLDA-1Na3MEA 5%	PG 5.0%	ordinary temperature: clear 40°C: clear
8	EMULGEN 108 40%	GLDA-4TEA 5%	PG 5.0%	ordinary temperature: clear 40°C: clear
9	EMULGEN 108 40%	α AD-1Na2TEA 3%	PG 10.0%	ordinary temperature: clear 40°C: clear
10	EMULGEN 108 40%	β AD-1Na2TEA 3%	PG 11.2%	ordinary temperature: clear 40°C: clear
11	EMULGEN 108 40%	SDA-1Na2TEA 5%	PG 5.0%	ordinary temperature: clear 40°C: clear
12	EMULGEN 108 40%	ASDA-1Na3TEA 3%	PG 12.1%	ordinary temperature: clear 40°C: clear
13	EMULGEN 108 40%	NTA-1Na2TEA 3%	PG 10.9%	ordinary temperature: clear 40°C: clear
14	EMULGEN 108 40%	EDDS-2Na2TEA 2%	PG 13.0%	ordinary temperature: clear 4°C: clear
15	EMULGEN 108 40%	IDS-2Na2TEA 3%	PG 12.0%	ordinary temperature: clear 40°C: clear
16	EMULGEN 108 40% LAS 5%	GLDA-1Na3TEA 3%	PG 12.0%	ordinary temperature: clear 40°C: clear

Table 2

Comparative Example	Surfactant	Builder	Solubilizing Agent	Compatibility
1	EMULGEN 108 40%	-	-	ordinary temperature: clear 40°C: clear
2	EMULGEN 108 40%	GLDA-4Na 5%	-	ordinary temperature: white turbidity 40°C: white turbidity
3	EMULGEN 108 40%	GLDA-4Na 5%	IPA 12.5%	ordinary temperature: white turbidity 40°C: white turbidity
4	EMULGEN 108 40%	GLDA-4Na 5%	PG 12.5%	ordinary temperature: white turbidity 40°C: white turbidity

Table 2 (continued)

	Comparative Example	Surfactant	Builder	Solubilizing Agent	Compatibility
5	5	EMULGEN 108 40%	GLDA-2Na2H 5%	-	ordinary temperature: white turbidity 40°C: white turbidity
10	6	EMULGEN 108 40%	GLDA-2Na2MEA 5%	-	ordinary temperature: white turbidity 40°C: white turbidity
	7	EMULGEN 108 40%	GLDA-2Na2H 5%	IPA 12.5%	ordinary temperature: white turbidity 40°C: white turbidity
15	8	EMULGEN 108 40%	GLDA-2Na2TEA 5%	-	ordinary temperature: white turbidity 40°C: white turbidity
20	9	EMULGEN 108 40%	GLDA-4Na 5%	PG 20.7%	ordinary temperature: white turbidity 40°C: white turbidity
	10	EMULGEN 108 40%	GLDA-2Na2TEA 5%	TEA 15.1%	ordinary temperature: white turbidity 40°C: white turbidity
25	11	EMULGEN 108 40%	GLDA-1Na3TEA 5%	-	ordinary temperature: white turbidity 40°C: white turbidity
30	12	EMULGEN 108 40%	α AD-3Na 3%	-	ordinary temperature: white turbidity 40°C: white turbidity
	13	EMULGEN 108 40%	β AD-3Na 3%	-	ordinary temperature: white turbidity 40°C: white turbidity
35	14	EMULGEN 108 40%	SDA-3Na 5%	-	ordinary temperature: white turbidity 40°C: white turbidity
40	15	EMULGEN 108 40%	ASDA-4Na 3%	-	ordinary temperature: white turbidity 40°C: white turbidity
45	16	EMULGEN 108 40%	NTA-3Na 3%	-	ordinary temperature: white turbidity 40°C: white turbidity
	17	EMULGEN 108 40%	IDS-4Na 3%	-	ordinary temperature: white turbidity 40°C: white turbidity
50	18	EMULGEN 108 40% LAS 5%	GLDA-4Na 3%	PG 12.0%	ordinary temperature: white turbidity 40°C: white turbidity

[0035] It is seen from the results in Tables 1 and 2 that, when Example 1 was compared with Comparative Examples 3, 4, 7, 8 and 10, the composition using an amine salt or an alkali metal-amine mixed salt as the builder and using an alcohol as the solubilizing agent had compatibility, whereas in the compositions using a Na salt as the builder, using no solubilizing agent or using alkanolamine as the solubilizing agent, the compatibility was poor and white turbidity occurred.

[0036] The same can be seen from a comparison between Example 3 and Comparative Example 11, Example 7 and Comparative Example 6, Example 9 and Comparative Example 12, Example 10 and Comparative Example 13, Example 11 and Comparative Example 14, Example 15 and Comparative Example 17, and Example 16 and Comparative Example 18.

[0037] As verified in the foregoing, by mixing an amine salt or alkali metal-amine mixed salt of an aminopolycarboxylic acid as the builder and an alcohol as the solubilizing agent each in a specific amount with a high-concentration nonionic or anionic surfactant, a high-concentration liquid detergent composition having excellent compatibility and free of white turbidity can be obtained.

[0038] While the invention has been described in detail and with reference to specific embodiments thereof, it will be apparent to one skilled in the art that various changes and modifications can be made therein without departing from the spirit and scope thereof.

Claims

1. A liquid detergent composition comprising:

(A) one or more surfactants selected from the group consisting of a nonionic surfactant and an anionic surfactant,

(B) one or more builders selected from the group consisting of an amine salt and an alkali metal-amine mixed salt of an aminopolycarboxylic acid-based chelating agent, and

(C) one or more solubilizing agents selected from the group consisting of alcohols,

with component (A) being present in a concentration of 20 wt% or more.

2. A liquid detergent composition comprising:

(A) from 20 to 60 wt% of one or more surfactants selected from the group consisting of a nonionic surfactant and an anionic surfactant,

(B) from 0.1 to 15 wt% of one or more builders selected from the group consisting of an amine salt and an alkali metal-amine mixed salt of an aminopolycarboxylic acid-based chelating agent,

(C) from 0.5 to 30 wt% of one or more solubilizing agents selected from the group consisting of alcohols, and
(D) the balance water.

3. The liquid detergent composition as claimed in claim 1 or 2, where the aminopolycarboxylic acid is an amino acid-N,N-diacetic acid.

4. The liquid detergent composition as claimed in claim 1 or 2, wherein the aminopolycarboxylic acid is glutamic acid-N,N-diacetic acid, α -alanine-N,N-diacetic acid, β -alanine-N,N-diacetic acid, aspartic acid-N,N-diacetic acid, glycine-N,N-diacetic acid or serine-N,N-diacetic acid.

5. The liquid detergent composition as claimed in any of the claims 1 to 4, wherein the alcohol is a glycol.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 11 2847

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.CI.7)
X	US 3 914 185 A (INAMORATO JACK THOMAS) 21 October 1975 (1975-10-21) * claim 1 *	1-3	C11D1/83 C11D3/33 C11D3/43 C11D3/20
X	GB 2 003 173 A (SANDOZ LTD) 7 March 1979 (1979-03-07) * claims 1,5,9 *	1-3	
A	US 5 536 452 A (BLACK ROBERT H) 16 July 1996 (1996-07-16) * claim 1 *	1-5	
A	GB 2 193 505 A (UNILEVER PLC) 10 February 1988 (1988-02-10) * claim 1 * * page 2, line 33 - line 48 *	1-5	
A	US 5 186 856 A (HOLLAND RICHARD J) 16 February 1993 (1993-02-16) * claims 1,2 *	1-5	
A	US 4 020 016 A (SOKOL CHARLES S) 26 April 1977 (1977-04-26) * claims 1-7 *	1-5	TECHNICAL FIELDS SEARCHED (Int.CI.7) C11D
A	EP 0 513 948 A (GRACE W R & CO) 19 November 1992 (1992-11-19) * claims 1-8 * * page 4, line 13 - line 41 *	1-5	
A	EP 0 511 091 A (COLGATE PALMOLIVE CO) 28 October 1992 (1992-10-28) * page 4, line 16 - line 18 * * page 6, line 26 - line 48 *	1-5	
		-/--	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 5 October 1999	Examiner Richards, M
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03/82 (P04/C01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 11 2847

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	DATABASE WPI Section Ch, Week 8815 Derwent Publications Ltd., London, GB; Class A97, AN 88-101776 XP002076232 & JP 63 051500 A (LION CORP) * abstract * -----	1-5	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
THE HAGUE		5 October 1999	Richards, M
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03.82 (P4/C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 11 2847

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

05-10-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 3914185 A	21-10-1975	NONE	
GB 2003173 A	07-03-1979	CH 629847 A	14-05-1982
		DE 2836567 A	08-03-1979
		FR 2401986 A	30-03-1979
		IT 1099594 B	18-09-1985
		JP 54046207 A	12-04-1979
		NL 7808788 A	02-03-1979
		US 4285840 A	25-08-1981
		BE 870010 A	28-02-1979
US 5536452 A	16-07-1996	AU 700901 B	14-01-1999
		AU 4704496 A	07-08-1996
		CA 2210760 A	25-07-1996
		EP 0804527 A	05-11-1997
		JP 10512612 T	02-12-1998
		NZ 301582 A	28-01-1999
		WO 9622346 A	25-07-1996
		US 5587022 A	24-12-1996
GB 2193505 A	10-02-1988	NONE	
US 5186856 A	16-02-1993	NONE	
US 4020016 A	26-04-1977	CA 1056251 A	12-06-1979
EP 0513948 A	19-11-1992	AT 169668 T	15-08-1998
		DE 69226557 D	17-09-1998
		DE 69226557 T	06-05-1999
		ES 2119796 T	16-10-1998
EP 0511091 A	28-10-1992	US 5254290 A	19-10-1993
		AU 647902 B	31-03-1994
		AU 1393092 A	29-10-1992
		CA 2067005 A	26-10-1992
		GR 1001376 B	29-10-1993
		JP 5117694 A	14-05-1993
		MX 9201851 A	01-10-1992
		NZ 242018 A	23-12-1993
		PT 100419 A	30-06-1993
JP 63051500 A	04-03-1988	JP 1958439 C	10-08-1995
		JP 6084514 B	26-10-1994

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82