

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 969 080 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
27.10.2004 Bulletin 2004/44

(51) Int Cl.7: **C11D 1/83**, C11D 3/33,
C11D 3/43, C11D 3/20

(21) Application number: **99112847.1**

(22) Date of filing: **02.07.1999**

(54) **Liquid detergent composition**

Flüssige Reinigungsmittel

Composition détergente liquide

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

(30) Priority: **03.07.1998 JP 18872398**

(43) Date of publication of application:
05.01.2000 Bulletin 2000/01

(73) Proprietor: **Showa Denko Kabushiki Kaisha
Tokyo 105-8518 (JP)**

(72) Inventors:
• **Saito, Makoto,**
c/o KAWASAKI WORKS, SHOWA DENKO K.K.
Kawasaki-shi, Kanagawa 210-0865 (JP)
• **Yamamoto, Tohru,**
KAWASAKI WORKS, SHOWA DENKO K.K.
Kawasaki-shi, Kanagawa 210-0865 (JP)

• **Nambu, Nobuyoshi, c/o YOKKAICHI WORKS**
Yokkaichi-shi, Mie 510-0886 (JP)
• **Furukawa, Masanori, c/o YOKKAICHI WORKS**
Yokkaichi-shi, Mie 510-0886 (JP)

(74) Representative: **Strehl Schübel-Hopf & Partner**
Maximilianstrasse 54
80538 München (DE)

(56) References cited:
EP-A- 0 511 091 EP-A- 0 513 948
GB-A- 2 003 173 GB-A- 2 193 505
US-A- 3 914 185 US-A- 4 020 016
US-A- 5 186 856 US-A- 5 536 452

• **DATABASE WPI Section Ch, Week 8815 Derwent**
Publications Ltd., London, GB; Class A97, AN
88-101776 XP002076232 & JP 63 051500 A (LION
CORP)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 969 080 B1

Description**FIELD OF THE INVENTION**

[0001] The present invention relates to a liquid detergent composition comprising a surfactant and an aminopolycarboxylic acid-based organic builder. More specifically, the present invention relates to a high concentration liquid detergent composition having excellent compatibility with a surfactant, superior resistance against white turbidity and good biodegradability.

BACKGROUND OF THE INVENTION

[0002] In liquid detergents, particularly in liquid detergents used as a household detergent, a nonionic surfactant and/or an anionic surfactant are used as the surfactant and a builder of various types is added to increase detergency. Various types of organic or inorganic builders can be used but almost all builders have a Na salt structure

[0003] For example, US-A-3 914 185 discloses clear, liquid, heavy-duty detergents based on mixed surfactant systems with nitrilotriacetic acid salt builders, which can be prepared in the absence of heat, without gel formation.

[0004] In recent years, from the standpoint of environmental conservation, studies are aggressively being made on the use of an aminopolycarboxylic acid having excellent biodegradability, particularly an amino acid diacetic acid-type chelating agent, as the builder. Aminopolycarboxylic acids including amino add diacetic acid chelating agents are used in the form of a synthetic reaction solution as it is on account of the cost. Therefore, in almost all cases, these are Na salts which cause strong alkalinity. For example, JP-A-10-25494 (the term "JP-A" as used herein means an "unexamined published Japanese patent application") discloses a liquid detergent for dishes, comprising an aspartic acid diacetic acid builder, where an alkali metal salt is used. JP-A-10-36888 discloses a liquid detergent comprising an aminopolycarboxylic acid builder, where an amine salt is disclosed on the same level with an alkali metal salt and in Examples, a Na salt is used. Moreover, the maximum concentration of the surfactant in this composition is as low as 10 wt%, thus, the problem of white turbidity on the high concentration side is not solved.

[0005] The use of a builder has a problem in that the solubility or clouding point of the surfactant as the main agent decreases and thereby the compatibility is deteriorated to cause, for example, white turbidity or separation of the white turbidity into two layers with the elapse of time. In order to prevent these, a technique of diluting the liquid detergent composition with water to reduce the concentration of the surfactant or a technique of adding a solubilizing agent has been heretofore used. However, the former case of diluting the liquid detergent composition with water inevitably results in an increase in the cost of transporting the detergent and this is commercially disadvantageous. In the latter case, it is known to newly add a surfactant as a solubilizing agent or add urea, an ether alcohol or a lower alcohol. However, the effect is not sufficiently high for obtaining a high concentration liquid detergent composition.

SUMMARY OF THE INVENTION

[0006] An object of the present invention is to obtain a liquid detergent which has excellent biodegradability, which does not assume strong alkalinity, which has excellent compatibility even if the surfactant concentration is increased, and which is free of white turbidity.

[0007] As a result of extensive investigations to solve the above-described problems, the present inventors have found that when a nonionic surfactant and/or anionic surfactant is used at a high concentration as the main surfactant of a liquid detergent and an aminopolycarboxylic acid chelating agent is used as a builder, by using an amine salt or an alkali metal-amine mixed salt as the counter ion of the builder and at the same time, adding an alcohol thereto, the compatibility is increased and the white turbidity can be prevented. The present invention has been accomplished based on this finding.

[0008] The present invention relates to the liquid detergent composition described below.

[0009] In one embodiment, the invention provides a liquid detergent composition comprising:

- (A) one or more surfactants selected from the group consisting of a nonionic surfactant and an anionic surfactant,
- (B) one or more builders selected from the group consisting of an amine salt and an alkali metal-amine mixed salt of an aminopolycarboxylic acid chelating agent, and
- (C) one or more solubilizing agents selected from the group consisting of alcohols,

with component (A) being present in a concentration of 20 wt% or more.

[0010] In another embodiment, the invention provides a liquid detergent composition comprising:

- (A) from 20 to 60 wt% of one or more surfactants selected from the group consisting of a nonionic surfactant and

an anionic surfactant,

(B) from 0.1 to 15 wt% of one or more builders selected from the group consisting of an amine salt and an alkali metal-amine mixed salt of aminopolycarboxylic acid chelating agent,

(C) from 0.5 to 30 wt% of one or more solubilizing agents selected from the group consisting of alcohols, and

(D) the balance of water.

[0011] A further embodiment comprises a liquid detergent composition as described in (1) or (2), wherein the aminopolycarboxylic acid is an amino acid-N,N-diacetic acid.

[0012] An even further embodiment is a liquid detergent composition as described in (1) or (2), wherein the aminopolycarboxylic acid is glutamic acid-N,N-diacetic acid, α -alanine-N,N-diacetic acid, β -alanine-N,N-diacetic acid, aspartic acid-N,N-diacetic acid, glycine-N,N-diacetic acid or serine-N,N-diacetic acid.

[0013] Additionally, another embodiment is a liquid detergent composition as described in any one of (1) to (4), wherein the alcohol is a glycol.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0014] The surfactant used as component (A) in the present invention is one or more selected from a nonionic surfactant and an anionic surfactant. Among these, a nonionic surfactant has a higher solubility than an anionic surfactant, accordingly, it is suitable for preparing a high concentration liquid detergent composition. The surfactant is used at a concentration of 20 wt% or more in the liquid detergent composition.

[0015] Examples of suitable nonionic surfactants include polyoxyethylene alkyl ether, polyoxyethylene alkyl phenyl ether, fatty acid diethanolamide, amine oxide and amidoamine oxide. Examples of appropriate anionic surfactants include alkyl sulfate, alkylpolyoxyethylene sulfate, alkylbenzenesulfonate, α -olefinsulfonate, fatty acid salts, ether carboxylates and monoalkylphosphates. A number of these may be commercially available.

[0016] The aminopolycarboxylic acid-based chelating agent used in the builder as component (B) of the present invention is a chelating agent containing one or more, preferably one or two amino groups and two or more, preferably three or four carboxyl groups, and having a chelating action. The amino group may be a secondary amino group or a tertiary amino group. The chelating agent is preferably an amino acid-N,N-diacetic acid.

[0017] An amino acid-N,N-diacetic acid is a compound represented by the following formula (1)




wherein A-NH₂ represents an amino acid, and -N represents an amino group in the amino acid.


[0018] Examples of amino acid-N,N-diacetic acids include glutamic acid-N,N-diacetic acid, α -alanine-N,N-diacetic acid, β -alanine-N,N-diacetic acid, aspartic acid-N,N-diacetic acid, glycine-N,N-diacetic acid and serine-N,N-diacetic acid. These amino acid-N,N-diacetic acids may be synthesized by the known Strecker reaction using, for example, an amino acid as the raw material or may be synthesized from a monochloroacetic acid.







[0019] Other than the amino acid-N,N-diacetic acid, examples of aminopolycarboxylic acid chelating agents include ethylenediaminedisuccinic acid and iminodisuccinic acid. These may also be synthesized by a known method and used. Examples of preferred aminopolycarboxylic acid chelating agents are shown below by their structural formulae.

Glutamic acid-N,N-diacetic acid



α -Alanine-N,N-diacetic acid



Aspartic acid-N,N-diacetic acid **β -Alanine-N,N-diacetic acid****Glycine-N,N-diacetic acid****Ethylenediaminedisuccinic acid****Iminodisuccinic acid****Serine-N,N-diacetic acid**

[0020] On account of the cost, amino acid-N,N-diacetic acids and other aminopolycarboxylic acid-based chelating agents are generally obtained as an aqueous solution where all carboxyl groups have a Na salt as the counter ion. However, in the present invention, the aminopolycarboxylic acid for use in the builder as component (B) is used in the form of an amine salt or an alkali metal-amine mixed salt. The effect may be obtained using primary, secondary and tertiary amines, however, the amine salt used is preferably an alkanolamine salt, more preferably triethanolamine salt or diethanolamine salt. The alkali metal salt in the alkali metal-amine mixed salt is preferably a K salt because it has a higher compatibility with the surfactant than an Na salt, however, in view of the cost, a Na salt is more preferred.

[0021] In order to convert the aminopolycarboxylic acid chelating agent obtained in the form of a Na salt into an alkali metal-amine mixed salt, the Na salt aqueous solution obtained may be treated to remove Na using an H-type cation exchange resin or by electrodialysis, or may be subjected to acid deposition to obtain a crystal having an H type carboxyl group. Thereafter, the resulting solution as it is or in the case of a crystal, a solution obtained by dissolving the crystals in water, is mixed with a mixture of an alkali metal hydroxide such as potassium hydroxide or sodium hydroxide with an amine and thereby an aqueous solution of the objective salt can be obtained. Also, after the removal of Na, the solution or crystal may be added to the formulation of a liquid detergent and then an amine or another alkali metal hydroxide may be added thereto so as to obtain the objective salt.

[0022] An aminopolycarboxylic acid-based chelating agent, particularly the aminodiacetic acid-type chelating agent is known to exert a good chelate effect namely a builder effect, on the neutral to weakly alkali side at a pH of from 8 to 12. Therefore, the amount of the amine or alkali metal hydroxide added is preferably controlled so that the above-

described pH region can be created. The larger the amount of the amine salt is, the higher is the compatibility with the surfactant, however, the ratio in the mixed salt of alkali metal and amine should be determined taking account of the cost for the removal of Na or the cost of amine.

[0023] The alcohol used as the solubilizing agent in the present invention is not particularly limited. However, a primary, secondary or tertiary alcohol, preferably a primary or secondary alcohol, having from 1 to 5 carbon atoms, preferably from 2 to 4 carbon atoms is preferred. The alcohol may be a general alcohol such as ethanol and isopropyl alcohol, however, since these are volatile and highly inflammable, a commonly used and inexpensive glycol such as propylene glycol and ethylene glycol is preferably used.

[0024] The liquid detergent composition of the present invention preferably has a composition ratio such that the surfactant as component (A) is from 20 to 60 wt%, preferably from 30 to 50 wt%, the builder as component (B) is from 0.1 to 15 wt%, preferably from 1 to 10 wt%, the solubilizing agent as component (C) is from 0.5 to 30 wt%, preferably from 1 to 20 wt%, and the balance is water as component (D). This composition ratio is the proportion among four components (A) to (D). The ratio of builder as component (B) based on the surfactant as component (A) is preferably from 2 to 20 wt%, more preferably from 5 to 15 wt%, and the solubilizing agent as component (C) based on the surfactant as component (A) is preferably from 5 to 40 wt%, more preferably from 10 to 30 wt%.

[0025] If the ratio of the surfactant as component (A) is less than the above-described range, the cost for transportation or packaging of the liquid detergent increases and this is commercially disadvantageous. If it exceeds the range, the desired compatibility can be attained only within narrow limits. As a result, flexibility in the formulation may decrease or the viscosity may increase and the handleability is disadvantageously impaired.

[0026] If the ratio of the builder as component (B) is less than the above-described range, the cleaning effect of the liquid detergent composition decreases, whereas if it exceeds the range, the compatibility with the surfactant occurs only within narrow limits and flexibility in formulation disadvantageously decreases.

[0027] If the ratio of the solubilizing agent as the component (C) is less than the above-described range, compatibility between components (A) and (B) is reduced, whereas if it exceeds the range, flexibility in formulation disadvantageously decreases.

[0028] The liquid detergent composition of the present invention may contain, if desired, other additives in any optional ratio, such as a viscosity controlling agent, a moisture retaining agent, an antiseptic, an antifungal agent, a dye and perfume.

[0029] The detergent composition of the present invention may be used for all uses as a detergent. However, this is suitably used in particular as a home detergent for cleaning clothes or household materials.

[0030] The detergent composition of the present invention may be used as it is or may be used after diluting it with water because the concentration is high.

[0031] According to the present invention, a builder comprising an amine salt or amine-alkali metal mixed salt of an aminopolycarboxylic acid and a solubilizing agent comprising an alcohol are mixed each in a specific ratio with a specific surfactant, therefore, a high-concentration liquid detergent composition which does not have strong alkalinity, which has excellent compatibility even if the surfactant concentration is increased, and which is free of white turbidity can be obtained.

EXAMPLES

[0032] The present invention is described below by referring to the Examples. In the Examples, "%" is "% by weight (wt%)".

Examples 1 to 16 and Comparative Examples 1 to 18:

[0033] The following surfactant, builder and solubilizing agent comprising an alcohol were added at the ratio shown in Tables 1 and 2, and the balance was water. The resulting solutions each was thoroughly mixed and left standing at an ordinary temperature or 40°C. Then, compatibility was visually observed. Those underwent separation into two layers after standing each was again mixed when observing and the solution state was observed. The formulations and visual results are shown together in Tables 1 and 2 below.

[0034] The following compounds were used.

Surfactant:

[0035]

Nonionic surfactant:

EP 0 969 080 B1

EMULGEN 108, produced by Kao Corporation (trademark, ethylene oxide alkyl ether nonionic)

Anionic surfactant:

LAS, reagent, sodium linear alkylbenzenesulfonate (free of sodium sulfate)

Builder:

[0036]

GLDA	glutamic acid-N,N-diacetic acid
α AD	α -alanine-N,N-diacetic acid
β AD	β -alanine-N,N-diacetic acid
SDA	serine-N,N-diacetic acid
ASDA	aspartic acid-N,N-diacetic acid
NTA	glycine-N,N-diacetic acid
EDDS	ethylenediaminedisuccinic acid
IDS	iminodisuccinic acid

Amine:

[0037]

TEA	triethanolamine
DEA	diethanolamine
MEA	monoethanolamine

Solubilizing Agent:

[0038]

PG	propylene glycol
IPA	isopropyl alcohol

[0039] In Tables 1 and 2, "-2Na2TEA" after "GLDA" indicates the counter ion of the carboxyl group within the molecule and means a mixed salt of 2Na and 2TEA. "-2Na2H" means 2Na salt in the low pH state. In either case, the concentration is the value calculated when the carboxyl group is the Na salt.

Table 1

Example	Surfactant	Builder	Solubilizing Agent	Compatibility
1	EMULGEN 108 40%	GLDA-2Na2TEA 5%	PG 12.5%	ordinary temperature: clear 40°C: clear
2	EMULGEN 108 40%	GLDA-2Na2TEA 5%	PG 6.9%	ordinary temperature: clear 40°C: clear
3	EMULGEN 108 40%	GLDA-1Na3TEA 5%	PG 4.8%	ordinary temperature: clear 40°C: clear
4	EMULGEN 108 40%	GLDA-2Na2DEA 5%	PG 6.7%	ordinary temperature: clear 40°C: clear
5	EMULGEN 108 40%	GLDA-1Na3DEA 5%	PG 4.7%	ordinary temperature: clear 40°C: clear
6	EMULGEN 108 40%	GLDA-1Na3DEA 5%	IPA 4.7%	ordinary temperature: clear 44°C: clear

EP 0 969 080 B1

Table 1 (continued)

Example	Surfactant	Builder	Solubilizing Agent	Compatibility
7	EMULGEN 108 40%	GLDA-1Na3MEA 5%	PG 5.0%	ordinary temperature: clear 40°C: clear
8	EMULGEN 108 40%	GLDA-4TEA 5%	PG 5.0%	ordinary temperature: clear 40°C: clear
9	EMULGEN 108 40%	α AD-1Na2TEA 3%	PG 10.0%	ordinary temperature: clear 40°C: clear
10	EMULGEN 108 40%	β AD-1Na2TEA 3%	PG 11.2%	ordinary temperature: clear 40°C: clear
11	EMULGEN 108 40%	SDA-1Na2TEA 5%	PG 5.0%	ordinary temperature: clear 40°C: clear
12	EMULGEN 108 40%	ASDA-1Na3TEA 3%	PG 12.1%	ordinary temperature: clear 40°C: clear
13	EMULGEN 108 40%	NTA-1Na2TEA 3%	PG 10.9%	ordinary temperature: clear 40°C: clear
14	EMULGEN 108 40%	EDDS-2Na2TEA 2%	PG 13.0%	ordinary temperature: clear 40°C: clear
15	EMULGEN 108 40%	IDS-2Na2TEA 3%	PG 12.0%	ordinary temperature: clear 40°C: clear
16	EMULGEN 108 40%	GLDA-1Na3TEA 3%	PG 12.0%	ordinary temperature: clear 40°C: clear
	LAS 5%			

Table 2

Comparative Example	Surfactant	Builder	Solubilizing Agent	Compatibility
1	EMULGEN 108 40%	-	-	ordinary temperature: clear 40°C: clear
2	EMULGEN 108 40%	GLDA-4Na 5%	-	ordinary temperature: white turbidity 40°C: white turbidity
3	EMULGEN 108 40%	GLDA-4Na 5%	IPA 12.5%	ordinary temperature: white turbidity 40°C: white turbidity
4	EMULGEN 108 40%	GLDA-4Na 5%	PG 12.5%	ordinary temperature: white turbidity 40°C: white turbidity
5	EMULGEN 108 40%	GLDA-2Na2H 5%	-	ordinary temperature: white turbidity 40°C: white turbidity
6	EMULGEN 108 40%	GLDA-2Na2MEA 5%	-	ordinary temperature: white turbidity 40°C: white turbidity

Table 2 (continued)

Comparative Example	Surfactant	Builder	Solubilizing Agent	Compatibility
7	EMULGEN 108 40%	GLDA-2Na2H 5%	IPA 12.5%	ordinary temperature: white turbidity 40°C: white turbidity
8	EMULGEN 108 40%	GLDA-2Na2TEA 5%	-	ordinary temperature: white turbidity 40°C: white turbidity
9	EMULGEN 108 40%	GLDA-4Na 5%	PG 20.7%	ordinary temperature: white turbidity 40°C: white turbidity
10	EMULGEN 108 40%	GLDA-2Na2TEA 5%	TEA 15.1%	ordinary temperature: white turbidity 40°C: white turbidity
11	EMULGEN 108 40%	GLDA-1Na3TEA 5%	-	ordinary temperature: white turbidity 40°C: white turbidity
12	EMULGEN 108 40%	α AD-3Na 3%	-	ordinary temperature: white turbidity 40°C: white turbidity
13	EMULGEN 108 40%	β AD-3Na 3%	-	ordinary temperature: white turbidity 40°C: white turbidity
14	EMULGEN 108 40%	SDA-3Na 5%	-	ordinary temperature: white turbidity 40°C: white turbidity
15	EMULGEN 108 40%	ASDA-4Na 3%	-	ordinary temperature: white turbidity 40°C: white turbidity
16	EMULGEN 108 40%	NTA-3Na 3%	-	ordinary temperature: white turbidity 40°C: white turbidity
17	EMULGEN 108 40%	IDS-4Na 3%	-	ordinary temperature: white turbidity 40°C: white turbidity
18	EMULGEN 108 40%	GLDA-4Na 3%	PG 12.0%	ordinary temperature: white turbidity 40°C: white turbidity
	LAS 5%			

[0040] It is seen from the results in Tables 1 and 2 that, when Example 1 was compared with Comparative Examples 3, 4, 7, 8 and 10, the composition using an amine salt or an alkali metal-amine mixed salt as the builder and using an alcohol as the solubilizing agent had compatibility, whereas in the compositions using a Na salt as the builder, using no solubilizing agent or using alkanolamine as the solubilizing agent, the compatibility was poor and white turbidity occurred.

[0041] The same can be seen from a comparison between Example 3 and Comparative Example 11, Example 7 and Comparative Example 6, Example 9 and Comparative Example 12, Example 10 and Comparative Example 13, Example 11 and Comparative Example 14, Example 15 and Comparative Example 17, and Example 16 and Comparative Example 18.

[0042] As verified in the foregoing, by mixing an amine salt or alkali metal-amine mixed salt of an aminopolycarboxylic acid as the builder and an alcohol as the solubilizing agent each in a specific amount with a high-concentration nonionic

or anionic surfactant, a high-concentration liquid detergent composition having excellent compatibility and free of white turbidity can be obtained.

Claims

1. A liquid detergent composition comprising:

- (A) one or more surfactants selected from the group consisting of a nonionic surfactant and an anionic surfactant,
- (B) one or more builders selected from the group consisting of an amine salt and an alkali metal-amine mixed salt of an aminopolycarboxylic acid-based chelating agent, and
- (C) one or more solubilizing agents selected from the group consisting of alcohols,

with component (A) being present in a concentration of 20 wt% or more.

2. A liquid detergent composition according to claim 1, comprising:

- (A) from 20 to 60 wt% of one or more surfactants selected from the group consisting of a nonionic surfactant and an anionic surfactant,
- (B) from 0.1 to 15 wt% of one or more builders selected from the group consisting of an amine salt and an alkali metal-amine mixed salt of an aminopolycarboxylic acid-based chelating agent,
- (C) from 0.5 to 30 wt% of one or more solubilizing agents selected from the group consisting of alcohols, and
- (D) the balance water.

3. The liquid detergent composition as claimed in claim 1 or 2, wherein the aminopolycarboxylic acid is an amino acid-N,N-diacetic acid.

4. The liquid detergent composition as claimed in claim 1 or 2, wherein the aminopolycarboxylic acid is glutamic acid-N,N-diacetic acid, α -alanine-N,N-diacetic acid, β -alanine-N,N-diacetic acid, aspartic acid-N,N-diacetic acid, glycine-N,N-diacetic acid or serine-N,N-diacetic acid.

5. The liquid detergent composition as claimed in any of the claims 1 to 4, wherein the alcohol is a glycol.

Patentansprüche

1. Flüssige Detergens-Zusammensetzung, umfassend:

- (A) ein oder mehrere oberflächenaktive Mittel, die aus der Gruppe ausgewählt sind, die aus einem nicht-ionischen oberflächenaktiven Mittel und einem anionischen oberflächenaktiven Mittel besteht,
- (B) ein oder mehrere Gerüststoffe, die aus der Gruppe ausgewählt sind, bestehend aus einem Aminsatz und einem Alkalimetall-Amin-Mischsalz eines Chelatbildners auf Aminopolycarbonsäure-Basis, und
- (C) ein oder mehrere Solubilisierungsmittel, die aus der Gruppe ausgewählt sind, die aus Alkoholen besteht,

wobei Bestandteil (A) in einer Konzentration von 20 Gew.-% oder mehr vorliegt.

2. Flüssige Detergens-Zusammensetzung, umfassend:

- (A) 20 bis 60 Gew.-% eines oder mehrerer oberflächenaktiver Mittel, die aus der Gruppe ausgewählt sind, die aus einem nicht-ionischen oberflächenaktiven Mittel und einem anionischen oberflächenaktiven Mittel besteht,
- (B) 0,1 bis 15 Gew.-% eines oder mehrerer Gerüststoffe, die aus der Gruppe ausgewählt sind, bestehend aus einem Aminsatz und einem Alkalimetall-Amin-Mischsalz eines Chelatbildners auf Aminopolycarbonsäure-Basis,
- (C) 0,5 bis 30 Gew.-% eines oder mehrerer Solubilisierungsmittel, die aus der Gruppe ausgewählt sind, die aus Alkoholen besteht, und
- (D) als Restbestandteil Wasser.

3. Flüssige Detergens-Zusammensetzung nach Anspruch 1 oder 2, wobei die Aminopolycarbonsäure eine Aminosäure-N,N-diessigsäure ist.
4. Flüssige Detergens-Zusammensetzung nach Anspruch 1 oder 2, wobei die Aminopolycarbonsäure Glutaminsäure-N,N-diessigsäure, α -Alanin-N,N-diessigsäure, β -Alanin-N,N-diessigsäure, Asparaginsäure-N,N-diessigsäure, Glycin-N,N-diessigsäure oder Serin-N,N-diessigsäure ist.
5. Flüssige Detergens-Zusammensetzung nach einem der Ansprüche 1 bis 4, wobei der Alkohol Glycol ist.

Revendications

1. Composition liquide détergente comprenant :

- (A) un ou plusieurs tensioactifs choisis dans le groupe constitué par un tensioactif non-ionique et un tensioactif anionique,
- (B) un ou plusieurs adjuvants choisis dans le groupe constitué par un sel d' amine et un sel mixte de métal alcalin et d'amine d'un agent chélatant à base d'acide aminopolycarboxylique, et
- (C) un ou plusieurs agents solubilisants choisis dans le groupe constitué par les alcools,

le composant (A) étant présent à une concentration de 20 % en masse ou plus.

2. Composition liquide détergente selon la revendication 1, comprenant :

- (A) 20 à 60 % en masse d'un ou plusieurs tensioactifs choisis dans le groupe constitué par un tensioactif non-ionique et un tensioactif anionique,
- (B) 0,1 à 15 % en masse d'un ou plusieurs adjuvants choisis dans le groupe constitué par un sel d'amine et un sel mixte de métal alcalin et d'amine d'un agent chélatant à base d'acide aminopolycarboxylique,
- (C) 0,5 à 30 % en masse d'un ou plusieurs agents solubilisants choisis dans le groupe constitué par les alcools, et
- (D) de l'eau pour la partie restante.

3. Composition liquide détergente selon la revendication 1 ou 2, dans laquelle l'acide aminopolycarboxylique est un acide aminoacide-N,N-diacétique.

4. Composition liquide détergente selon la revendication 1 ou 2, dans laquelle l'acide aminopolycarboxylique est l'acide (acide glutamique)-N,N-diacétique, l'acide α -alanine-N,N-diacétique, l'acide β -alanine-N,N-diacétique, l'acide (acide aspartique)-N,N-diacétique, l'acide glycine-N,N-diacétique ou l'acide sérine-N,N-diacétique.

5. Composition détergente liquide selon l'une quelconque des revendications 1 à 4, dans laquelle l'alcool est un glycol.