



(11) **EP 0 981 549 B9**

(12) **CORRECTED NEW EUROPEAN PATENT SPECIFICATION**

Note: Bibliography reflects the latest situation

(15) Correction information:
Corrected version no 1 (W1 B2)
Corrections, see
Claims EN

(51) Int Cl.:
C07K 16/18 (2006.01) **A61K 48/00** (2006.01)

(86) International application number:
PCT/EP1998/002808

(48) Corrigendum issued on:
27.02.2008 Bulletin 2008/09

(87) International publication number:
WO 1998/051708 (19.11.1998 Gazette 1998/46)

(45) Date of publication and mention
of the opposition decision:
27.06.2007 Bulletin 2007/26

(45) Mention of the grant of the patent:
25.09.2002 Bulletin 2002/39

(21) Application number: **98924307.6**

(22) Date of filing: **13.05.1998**

(54) **USE OF AN INHIBITOR SUBSTANCE FOR THE IMPROVEMENT OF NEURONAL REGENERATION**

VERWENDUNG EINER INHIBITORSUBSTANZ ZUR FÖRDERUNG DER NEURONALEN
REGENERATION

UTILISATION D'UN INHIBITEUR PERMETTANT D'AMÉLIORER LA REGENERATION DES
NEURONES

(84) Designated Contracting States:
AT BE CH DE DK ES FI FR GB IE IT LI NL SE

(30) Priority: **14.05.1997 EP 97107846**

(43) Date of publication of application:
01.03.2000 Bulletin 2000/09

(73) Proprietor: **Neuraxo Biopharmaceuticals GmbH**
40699 Erkrath (DE)

(72) Inventors:
• **Müller, H.W., Prof.Dr.**
40225 Düsseldorf (DE)
• **Stichel-Gunkel, Christine, Dr.**
44780 Bochum (DE)

(74) Representative: **Meyers, Hans-Wilhelm et al**
Patentanwälte
von Kreisler-Selting-Werner
Postfach 10 22 41
50462 Köln (DE)

(56) References cited:
WO-A-83/04104 **WO-A-93/19783**
WO-A-94/17831 **WO-A-95/10305**
WO-A-95/13291 **WO-A-96/00582**
JP-A- 6 103 773 **US-A- 4 214 000**
US-A- 5 021 404

• **K.Abe et al, Brain Res. vol. 723, 1996, p. 206 - 209**
• **P.E. Hughes et al, Prog. Neurobiol. vol. 57, 1999,**
p. 421 - 450
• **L.D.F. Moon et al, Eur. J. Neurosci. vol. 14, 2001,**
p. 1667 - 1677
• **Feringa E R et al. (1980), Ann Neurol 8, 148-154**

EP 0 981 549 B9

Description

[0001] The present invention refers to the use of an inhibitor substance for the improvement of neuronal regeneration, and a medicament for the improvement of neuronal regeneration.

[0002] Injury to adult mammalian CNS fiber tracts leads to the formation of a lesion scar consisting of a convoluted fringe of astroglial processes lined by a basal membrane (BM). This lesion scar is implicated as a major extrinsic constraint to effective axon regeneration in brain and spinal cord (1 - 4). While the dense astrocytic network is a permissive substrate for axon growth (5, 6), the presence of BM has been hypothesized as a crucial impediment for regeneration (7). However, experimental evidence was not shown. To the contrary, when the BM formed after a lesion of neuronal tissue was removed (24), no improved regeneration could be reproducibly monitored (25). Therefore, it is still of great importance to have a method for improving regeneration of injured neurons.

[0003] WO 93/19783 discloses a method for preventing, suppressing or treating a CNS pathology characterized by a deleterious accumulation of extracellular matrix in a tissue by contacting the tissue with an agent that inhibits the extracellular matrix producing activity of TGF- β . The disclosed methods can be used to prevent, suppress or treat scar formation in the CNS. As useful agents are addressed neutralizing anti-TGF- β antibodies, Arg-Gly-Asp-containing peptides, decorin and its functional equivalence such as biglycan and TGF- β antagonists. TGF- β has a wide spectrum of physiological functions such as activation of cell of the immune system, inhibition of cell proliferation, neurotrophic effects on sensory neurons, inhibition of Schwann cell myelination, anti-proliferative effects on glial cells, immunosuppressive effects, stimulation of extracellular matrix deposition and chemotraction of microglia cells. The anti-TGF- β treatment would induce the opposite effects. Inhibition of TGF- β activity leads to numerous non-specific cellular responses, which may even lead to unwanted side effects. One object of the invention is to avoid such potential unwanted side effects.

[0004] US-A-5,082,926 discloses the capability of competing peptide to promote nerve regeneration and indicates the clinical use in promoting regeneration of damaged nerves. It is speculated that the peptide disclosed is important in many diverse and clinically relevant processes such as cell attachment and migration in wound healing, tumor cell invasion and metastasis, diabetic microangiopathy, vascular hypertrophy due to hypertension and several kidney diseases such as diabetic nephropathy and nephrotic syndromes of variable etiology.

[0005] US-A-5,493,008 discloses high affinity binding of nidogen to laminin to be mediated by an EGF-like repeat γ 1III4 of the mouse laminin γ 1 chain.

[0006] WO-A-96/00582 discloses the use of isolated

domains of type IV collagen to modify cell and tissue interactions. The 7S domain of type IV collagen disrupts cell aggregation and tissue development. Structural changes in mesoglea, inhibition of cell proliferation, and changes in cell differentiation patterns accompanies the blockage of cell aggregates which indicate that blockage may be due to alterations in mesoglea (extracellular matrix) structure with accompanying effects on cell behavior. Type IV collagen has a critical role in the initial formation of mesoglea and the perturbation of mesoglea formation affects cell division, cell differentiation, and morphogenesis.

[0007] Surprisingly, improvement of regeneration of neuronal tissue after lesion is achieved by the use of an inhibitor substance of the present invention for the manufacturing of a medicament for the improvement of neuronal regeneration.

[0008] According to the method of the invention improved regeneration of injured neuronal tissue is achieved by prevention or specific inhibition of basal membrane formation induced by a lesion of neuronal tissue by applying an inhibitor substance of the synthesis of basal membrane building elements, or the assembly of basal membrane building elements, or both the synthesis of basal membrane building elements and the assembly of basal membrane building elements to a body in need thereof wherein the inhibitor substance is selected from the group consisting of antibodies against collagen IV, laminin, entactin, accessory substances for proper function, or the assembly of the basal membrane; Fe-chelating agents; inhibitors of amino acids hydroxylases, such as prolyl-4-hydroxylase, lysine-hydroxylase; 2-oxoglutarate competitors.

[0009] The basal membrane is a structure which is composed of different elements. Elements of the basal membrane are collagen IV, laminin, entactin (Nidogen) accessory substances. The assembly of the elements to the basal membrane is performed by enzymes which may be assisted by cofactors.

[0010] Inhibitors of TGF- β are not involved with a specific prevention or specific inhibition of basal membrane formation induced by lesion of neuronal tissue. According to the present invention it is achieved in an advantageous manner that a specific interaction is provided by the use according to the invention.

[0011] Preferably, the formation of the basal membrane is prevented or inhibited by applying a specific inhibitor substance of the synthesis of basal membrane building elements, or the assembly of basal membrane building elements, or both the synthesis of basal membrane building elements and the assembly of basal membrane building elements to a body in need thereof. The building elements of the basal membrane are in particular those which are involved with the formation of the basal membrane, for instance molecular structures building up the basal membrane, such as monomeric compounds, accessory substances, substances for the assembly of the components of the basal membrane and the like.

[0012] In particular, the basal membrane building elements are selected from the group consisting of collagen IV, laminin, entactin, accessory substances for proper function, or the assembly of the basal membrane, or both the proper function and the assembly of the basal membrane.

[0013] A specific inhibitor substance of the invention is capable of preventing or inhibiting the formation of the basal membrane and/or is specifically interfering with the assembly process of the basal membrane. The specific inhibitor substance of the invention is selected from the group consisting of antibodies against collagen IV, laminin, entactin, accessory substances for proper function, or the assembly of the basal membrane; Fe-chelating agents; inhibitors of amino acids hydroxylases, such as prolyl-4-hydroxylase, lysine-hydroxylase; 2-oxoglutarate competitors; which inhibitor substance is able to prevent or inhibit the expression of basal membrane building elements, and the like.

[0014] According to the invention are preferred those inhibitor substances which are selected from the group consisting of N-oxaloglycine; Zn salts; pyridine derivatives, such as 5-arylcarbonylamino- or 5-arylcarbamoylester derivatives, 2-carboxylate, 2,5 dicarboxylate, their ethyl esters or ethyl amides or -5-acyl sulfonamides, 2,4 dicarboxylate, their ethyl esters or ethylamides, or dimethoxyethylamides; 3,4 bipyridine, such as 5 amino-6-(1H)-one, 1,6-dihydro-2-methyl-6-oxo-5-carbonitril; 2,2'-bipyridine, such as 5,5'-dicarboxylic acid or its pharmaceutically acceptable salts, 4,4'-dicarboxylic acid ethyl ester or ethyl amide; 3,4'-dihydroxybenzoate, such as the diethyl ester; proline and its structural and functional analogues; β -aminopropionitrile; desferrioxamine; anthracyclines; 2,7,8-trihydroxy anthraquinones, fibrostatin-C; coumalic acid or its pharmaceutically acceptable salts; 5-oxaproline, β -lactam antibiotics.

[0015] In a preferred embodiment of the present invention the specific inhibitor substance(s) are applied in combination with one or more substances being capable of stimulating neuronal growth or inducing the expression of growth promoting proteins. Such neuronal growth stimulating substances are neurotrophic growth factors of the neurotrophin family and other growth factor families such as fibroblast growth factors, insulin and insulin-like growth factors, as well as epidermal growth factor, ciliary neurotrophic growth factor (CNTF), glial cell-derived growth factor (GDNF), cytokines, neurotrophic proteoglycans and glycosamino-glycans, neural cell adhesion molecules like L1 (NILE), growth-associated proteins like GAP43 and anti-apoptotic proteins like bcl-2.

[0016] According to the invention it is preferred to locally apply the specific inhibitor substances in the neuronal tissue, intraventricularly, or systemically, in particular orally or intravenously.

[0017] The concentration of the specific inhibitor substance varies in view of the chemical nature. For example, antisense inhibitor substances may have more specific effects so that lesser amounts can be applied.

[0018] Typically, the specific inhibitor substance is applied in therapeutically effective amounts, such as 1 ng/kg to 1 mg/kg body weight, when low molecular compounds such as bipyridyl-derivatives are applied.

[0019] The invention also provides a medicament for the improvement of neuronal regeneration comprising a therapeutically effective amount of a specific inhibitor substance which is capable of prevention or inhibition of basal membrane formation induced by a lesion of neuronal tissue. According to the invention the inhibitor substances are those as claimed in claim 5. The medicament may further comprise carrier substances or adjuvants in order to facilitate an appropriate application. The medicament further comprises substances which are capable of stimulating neuronal growth as claimed in claim 5.

[0020] Fig. 1:

Expression of collagen IV and axonal sprouting after transection of the postcommissural fornix in untreated animals (b-d) and after injection of anti-Coll IV (e) or DPY (f) at two weeks postsurgery, a, Sagittal view of the adult rat brain showing the course of the fornix and the location of the transection site. Marked deposition of collagen IV in the lesion site (arrow) and proximal stump (P) of an untreated animal at low (b) and high magnification (c). Note, the fine structure and the spatial orientation of collagen IV deposits perpendicular to the trajectory of the tract, d, In untreated animals regrowing fornix axons stop sharply at the lesion site (arrow). Collagen IV deposition is markedly reduced in the lesion site after anti-Coll IV (e) or DPY injection (f). Scale bars, 100 μ m.

[0021] Fig. 2: Regeneration of transected fornix fibers across the lesion site in rats treated with anti-Coll IV (a, c, e) or DPY (b, d, f) at 6 weeks postsurgery. Sagittal serial sections reacted for NF-immunohistochemistry show that in both experimental groups fibers traverse the former lesion site (arrows) (c, d) and elongate within the distal stump (e, f) up to the mammillary body (MB). Scale bars, 100 μ m.

[0022] Fig. 3: Recovery of structural features of the regenerating fornix tract. a, b Anterograde tracing with biocytin of an anti-Coll IV treated animal at 6 weeks postsurgery reveals the large number of regenerating axons (a), their elongation within the former pathway (a) and their fine varicose morphology (b). c, Large WGA-HRP-filled axon (arrowhead) in the mammillary body surrounded by compact myelin (arrows), d, e Electron micrographs of anterogradely WGA-HRP-labeled presynaptic terminals (arrowheads) in the mammillary body at 6 weeks after anti-Coll IV treatment. Scale bars, 100 μ m (a), 50 μ m (b), 0.1 μ m (c), 0.5 μ m (d), 1 μ m (e).

[0023] Fig. 4: Electrophysiological properties of fornix fibers in unlesioned rats and lesioned/injected animals with regeneration. a, Schematic illustration showing the location of the stimulating (S) and recording (R) electrode at various conditions, b, Characteristic recordings of extracellular action potentials in a sagittal slice prepared from an animal with regeneration. Recordings were obtained under conditions as illustrated in a. Application of

Tetrodotoxin (TTX) blocks the stimulus-evoked response. The net action potential is shown in trace 5. c and d, Distribution of conduction velocity and action potential response amplitude in unlesioned and lesioned/injected animals with regeneration.

[0024] The mechanically transected postcommissural fornix of the adult rat, a unidirectional and well-characterized fiber tract (8,9), was used to determine whether specific biochemical or immunochemical modulation of BM formation would provide a means to stimulate axon regeneration. Here we report that lesion-induced BM deposition can be significantly reduced by local injection of anti-collagen IV antibodies or α,α -dipyridyl, an inhibitor of collagen triple helix formation and synthesis. Reducing the collagen network allowed massive axon elongation across the lesion site. The regenerating fornix fibers followed the original pathway, reinnervated their appropriate target, the mammillary body, were remyelinated and attained nearly normal conduction properties, on failure of adult mammalian CNS axons we examined its spatio-temporal distribution pattern after penetrant CNS lesion and determined whether its remodelling allows structural and functional regeneration of a transected CNS fiber tract.

[0025] The left postcommissural fornix was stereotactically transected in adult Wistar rats Fig. 1a) and the postlesion deposition of BM was analyzed using antibodies against collagen IV (Coll IV) and laminin (LN), the major and unique components of BM (10,11). By the end of the second week after lesion the center of the wound was filled by Coll IV- and LN-rich BM (Fig. 1b, c). These newly formed BM were either arranged in long continuous layers or associated with numerous blood vessels. Within the center of the wound the BM layers formed a parallel array aligned perpendicular to the course of the fiber tract (Fig. 1b, c). In the vicinity of the transected stumps, however, BM layers were deposited as hook-like turns extending along the longitudinal tract axis for about 200 μ m into the fornix stumps (Fig. 1c). In parallel with the deposition of the BM, sprouting axons in the proximal stump reached the lesion site. They failed to cross or bypass it but stopped growing at the wound border at about 2 weeks after lesion (Fig. 1d). The spatio-temporal coincidence of BM formation with the abrupt axonal growth arrest at the tract-lesion border strongly suggests that the newly formed perpendicular layers of BM could be a physical impediment for regenerating axons.

[0026] In an effort to modulate postlesion BM deposition, either polyclonal antibodies against collagen IV (anti-Coll IV; n=14) or the iron chelator α,α -dipyridyl (DPY; n=9) were injected locally into the lesion center immediately after transection. DPY is a competitive inhibitor of prolyl 4-hydroxylase (12) and has been shown to prevent collagen triple helix formation (12), which results in feedback inhibition of procollagen synthesis (13) and enhanced procollagen degradation (14). Control animals received a PBS injection (n=9) or were sham operated (n=3). Basal membrane formation was studied in re-

sponse to antibody and drug treatment using immunohistochemical methods. Animals receiving a single injection of anti-Coll IV (80-160 ng) or DPY (1.6-16 μ mol) showed a massive and specific reduction in Collimmunopositive laminae and blood vessels in the lesion center and the fornix stumps at all examined survival time points. At 2 weeks after lesion+injection only a very small number of Collimmunoreactive structures perpendicular to the tract course had developed (Fig. 1e, f). Control animals, however, exhibited dense BM deposition as previously described for lesion only animals. The applied substances reduced the deposition of BM at the lesion site but did not affect the number or the distribution of vascular BM in the surrounding neuropil. Therefore, we conclude that the lesion-induced BM formation can be specifically reduced by immediate application of either anti-Coll IV antibodies or DPY.

[0027] To determine whether reduction of BM deposition would permit regeneration of transected axons across the lesion site, we studied the elongation of fornix axons after anti-Coll IV or DPY treatment using immunocytochemical staining. While sprouting fornix fibers in control animals ceased growing at the proximal stump-lesion interface (Fig. 1d) large numbers of axons entered and traversed the lesion center between 2 and 4 weeks after lesion+injection in those animals receiving anti-Coll IV (n= 11) (Fig. 2 a, c, e) or DPY treatment (n= 6) (Fig. 2 b, d, f). Most regenerating axons formed a loop over the lesion site, entered the distal stump and continued in a parallel bundle of fine and beaded axons within their previous pathway (Fig. 3a, b). They reached their appropriate target, the mammillary body, at about 4-6 weeks postsurgery. Anterograde tracing with WGA-HRP into the subiculum, the origin of the fornix (not shown), or biocytin application into the proximal fornix stump (Fig. 3a) provided proof, that the vast majority of fibers emerge from the formerly transected fornix tract. All regenerating fornix axons remained within their original pathway and did not invade the surrounding neuropil. The present results demonstrate that the failure of postcommissural fornix regeneration in rat brain, in fact, depends upon the formation of an axon growth-inhibiting BM at the lesion site that is oriented perpendicular to the tract course. Reduction of BM deposition seems to be a prerequisite but also a sufficient condition for the transected axons to regenerate across the lesion site.

[0028] Further preferred embodiments for restitution of functional circuitry after traumatic CNS lesion are the remyelination of regenerated fibers, the re-establishment of synaptic connections with the appropriate target and the restoration of normal conduction properties. Structural and functional properties of the regenerating axons were investigated using immunohistochemical, morphological and electrophysiological methods. Immunohistochemistry with an antibody against myelin basic protein demonstrated the remyelination of regenerated fornix axons along their entire length as early as 4 weeks after surgery (data not shown). This observation was con-

firmed by ultrastructural analysis of anterogradely WGA-HRP labeled axons in the distal stump which showed clear evidence of compact myelin sheath formation (Fig. 3c). In addition, ultrastructural studies provided evidence for the re-establishment of synaptic connections of regenerating axons within the mammillary body. Tracer re-action product was identified in presynaptic profiles with round vesicles that formed asymmetric synaptic junctions at unlabeled dendrites (Fig. 3d, e). The ultrastructural features of the labelled presynaptic profiles correspond to those described for the RA-type (round, asymmetric) of synaptic terminal, which is considered to be of subicular origin (8). The electrophysiological properties of regenerated fibers were studied using extracellular in vitro recording techniques applied to sagittal brain slices (400 μ m) of 8 unlesioned rats and 4 treated animals showing regenerated fiber tracts. In unlesioned animals electrical stimulation of the fornix fibers elicited an extracellular action potential with an amplitude of 1.02 ± 0.14 mV and a conduction velocity of 0.48 ± 0.05 m/s (mean $\pm$ SEM, n=16, Fig. 4b-d). This axonal conduction velocity corresponds well to previously reported measurements (about 0.5 m/s for hippocampal Schaffer collaterals (15). Similar values for action potential amplitude and conduction velocity (1.12 ± 0.21 mV, 0.46 ± 0.1 m/s, n=5) were obtained in regenerating animals when the stimulating (S) and the recording (R) electrodes were positioned proximally to the lesion site (see S1 and R1 in Fig. 4a). In the latter animals, functionally intact fibers showing normal extracellular action potential amplitude and conduction velocity could also be demonstrated across (S3 and R3 in Fig. 4a; 0.8 ± 0.29 mV, 0.54 ± 0.14 m/s, n=3) and distal to the lesion site (S2 and R2 in Fig. 4a; 0.91 ± 0.24 mV, 0.43 ± 0.06 m/s, n=4) (Fig. 4c, d). In all animals, the stimulus-evoked extracellular responses were blocked by Tetrodotoxin, confirming their nature as Na⁺-dependent action potentials (Fig. 4b). From these data we conclude that the reorganization of the fornix tract is accompanied by structural and functional recovery of the regenerated axons.

[0029] Our results demonstrate that structural and functional restoration of lesioned mature fornix pathway can be achieved by reduction of BM formation in the lesion site. Data described here underscore the importance of extrinsic determinants in axonal regeneration but also demonstrates that once the axons have crossed the lesion scar other potential extrinsic regeneration constraints, like CNS myelin and oligodendrocytes (9,16-18), dense astrogliosis (6) and sulfated proteoglycans (19,20), do not impede their progress. The results further indicate that similar to other CNS circuits (21,22), fornix axons have an innate potential for regeneration and self-organization. These results give rise to new and promising concepts for therapeutic strategies that might contribute to the reduction of neurological deficits after CNS lesions.

[0030] The following examples are intended for further illustration of the invention but are not limiting.

[0031] Surgery. The left postcommissural fornix of 42 Wistar rats (180-210g) was transected stereotactically at a distance of about 1 mm proximal to the target, the mammillary body, using a Scouten wire knife as described previously (9). The completeness of transection was confirmed by serial reconstruction of the lesion site for each of the animals. Immediately after transection animals received a topical application (1.6 μ l) of either polyclonal antibodies against collagen IV (anti-Coll IV, Biogenex, 50-100 μ g/ml, n=14) or the iron chelator a, a'-dipyridyl (DPY, 1-10 mM, n=9). Substances were pressure injected (injection time 10 min) directly into the lesion site via a micropipette coupled to a microsyringe. Controls received equal amounts of phosphate-buffered saline (n=9) or sham operation (n=3).

[0032] Anterograde tracing was performed for analysis of fiber course, ultrastructural morphology and target reinnervation. After a survival time of 6 weeks, anti-Coll IV-treated animals (n=4) received two injections of a 2% (w/v) solution of wheat-germagglutinin-HRP (WGA-HRP) into the left subicular complex (dorsal and caudal pole). Rats were perfused 3 days later with 2% paraformaldehyde and 2% glutaraldehyde in 0.1M phosphate buffer. Vibratome sections were reacted for WGA-HRP using tetramethylbenzidine as substrate (23).

[0033] Electron microscopy. For ultrastructural analysis vibratome sections of anti-Coll IV-treated animals were reacted for WGA-HRP, immersed for 12h in 1% osmium tetroxide and embedded in epon. Ultrathin sections were examined using a Hitachi H600 electron microscope.

[0034] Immunohistochemical staining. After a survival time of 4 days (d), 6d, 2 weeks (w), 4w and 6w after surgery brains were removed, frozen in isopentane (-50/-60°C) and cut into serial sagittal 10 μ m thick sections. Sections were fixed with acetone (-20°C), preincubated in 3% H₂O₂ (v/v) in methanol to block endogeneous peroxidase, followed by PBS containing 3% (v/v) normal horse or normal goat serum to reduce unspecific staining and then incubated with one of the following primary antibodies: polyclonal anti-collagen IV (anti-Coll IV, Biogenex, 1:3), polyclonal anti-laminin (anti-LN, Biogenex, 1:5) or monoclonal cocktail against phosphorylated neurofilaments (anti-NF, Affinity, 1:800). Following, avidin-biotin-peroxidase complex staining (Vector Labs) was done using standard procedures. For evaluation of remyelination brains were fixed with 4% paraformaldehyde, paraffinized, cut into 3- μ m thick serial sagittal sections, deparaffinized and incubated as described above with a polyclonal anti-myelin basic protein (anti-MBP, Biogenex, 1:2) or anti-NF as primary antibodies. Specificity of the stainings was confirmed by omission of the primary antibody.

[0035] Electrophysiology and biocytin injections. Sagittal slices of 400 μ m thickness were cut on a vibratome and maintained at 34-35°C in an interface-type recording chamber. Artificial cerebrospinal fluid (ACSF) consisted of (in mM) 124 NaCl, 3 KCl, 1.25 NaH₂PO₄, 1.8 MgSO₄,

1.6 CaCl₂, 26 NaHCO₃ and 10 glucose with a pH of 7.4 when saturated with 95% O₂ - 5% CO₂. Stimuli: 100 μ s, 5-20 V were delivered via a bipolar tungsten electrode. Extracellular action potentials were registered with a recording electrode (3-5 MW) located in the middle of the postcommissural fornix. Tetrodotoxin (TTX, Sigma) was applied locally in a concentration of 10 μ M (dissolved in ACSF) with a broken micropipette placed on the slice surface near the recording site. Injections of a small biocytin (Sigma) crystal into the fornix were performed with a miniature needle. After an incubation period of 8-10 h in the interface chamber, slices were fixed in 4 % paraformaldehyde, resectioned and reacted with ABC peroxidase reagent (Vector Labs)

1. Reier, P. J., Stensaas, L. J. & Guth, L. Spinal cord reconstruction (eds Kao, C.C., Bunge, R.P. & Reier, P.J.) 163-196 (Raven Press, New York, 1983).

2. Clemente, C. D. Regeneration in the nervous system (ed Windle, W.F.) 147-161 (Charles C. Thomas, Springfield, IL, 1955).

3. Schnell, L. & Schwab, M. E. Eur. J. Neurosci 5, 1156-1171 (1993).

4. Krüger, S., Sievers, J., Hansen, C., Sadler, M. & Berry, M. J. Comp. Neural 249, 103-116 (1986).

5. Berry, M., Carlile, J. & Hunter, A. J. Neurocytol 25, 147-170 (1996).

6. Stichel, C. C. & Müller, H. W. J. Neurocytol 23, 615-630 (1994).

7. Feringa, E. R., Kowalski, T. F. & Vahlsing, H. L. Ann. Neurol 8, 148-154 (1980).

8. Allen, G. V. & Hopkins, D. A. J. Comp. Neurol 275, 39-64 (1988).

9. Stichel, C. C., Wunderlich, G., Schwab, M. E. & Müller, H. W. Europ. J. Neurosci 7, 401-411 (1995).

10. Yurchenco, P. D. & Schittny, J. C. FASEB. J 4, 1577-1590 (1996).

11. Timpl, R. & Brown, J. Curr. Opin. Cell. Biol 8, 618-624 (1996).

12. Kivirikko, K. L., Myllylä, R. & Pihlajaniemi, T. FASEB. J 3, 1609-1617 (1989).

13. Ikeda, H., Wu, G. Y. & Wu, C. H. Hepatology 15, 282-287 (1992).

14. Bienkowski, R. S. J. Cell. Physiol 121, 152-158 (1984).

15. Andersen, P., Silfvenius, H., Sundberg, S., Sveen, O. & Wigström, H. Brain. Res 144, 11-18 (1978).

16. McKerracher, L. et al. Neuron 13, 805-811 (1994).

17. Mukhopadhyay, G., Doherty, P., Walsh, F. S., Crocker, P. R. & Filbin, M. T. Neuron 13, 757-767 (1994).

18. Schwab, M. E. & Caroni, P. J. Neurosci 8, 2381-2393 (1988).

19. Snow, A. D., Lemmon, V., Carrino, D. A., Caplan, A. I. & Silver, J. Exp. Neurol 109, 111-130 (1990).

20. Lips, K., Stichel, C. C. & Müller, H. W. J. Neurocytol 24, 449-464 (1995).

21. Iwashita, Y., Kawaguchi, S. & Murata, M. Nature 367, 167-170 (1994).

22. Kalderon, N. & Fuks, Z. Proc. Natl. Acad. Sci 93, 11179-11184 (1996).

23. Olucha, F., Martinez-Garcia, F. & Lopez-Garcia, C. J. Neurosci. Meth 13, 131-138 (1985).

24. Puchala, E. & Windle, W. F., Exp. Neurol. 55, 1 - 42 (1977).

25. Guth, L., Albuquerque, E. X., Deshpande, S. S., Barret, C. P., Donati, E. J. & Warnick, J. E., J. Neurosurg. 52, 73 - 86 (1980).

Claims

1. Use of an inhibitor substance selected from the group consisting of antibodies against collagen IV, laminin, entactin, accessory substances for proper function, or the assembly of the basal membrane; Fe-chelating agents; inhibitors of amino acids hydroxylases, such as prolyl-4-hydroxylase, lysine-hydroxylase; 2-oxoglutarate competitors; which inhibitor substance is capable of prevention or specific inhibition of basal membrane formation induced by a lesion of neuronal tissue for the manufacturing of a medicament for the improvement of neuronal regeneration.
2. Use of claim 1, wherein the inhibitor substance is a substance inhibiting the synthesis of basal membrane building elements, or the assembly of basal membrane building elements, or both the synthesis of basal membrane building elements and the assembly of basal membrane building elements.

3. Use of claim 2, wherein the basal membrane building elements are selected from the group consisting of collagen IV, laminin, entactin, accessory substances for proper function, or the assembly of the basal membrane, or both the proper function and the assembly of the basal membrane. 5
4. Use of claim 1, wherein the inhibitor substance is selected from the group consisting of N-oxaloglycine; Zn salts; pyridine derivatives, such as 5-aryl-carbonylamino- or 5-arylcarbamoyle-derivatives, 2-carboxylate, 2,5-dicarboxylate, their ethyl esters or ethyl amides or -5-acyl sulfonamides, 2,4 dicarboxylate, their ethyl esters or ethylamides, or dimethoxyethylamides; 3,4'-bipyridine, such as 5 amino-6-(1H)-one, 1,6-dihydro-2-methyl-6-oxo-5-carbonitril; 2,2'-bipyridine, such as 5,5'-dicarboxylic acid or its pharmaceutically acceptable salts, 4,4'-dicarboxylic acid ethyl ester or ethyl amide; 3,4'-dihydroxybenzoate, such as the diethyl ester; proline and its structural and functional analogues; β -aminopropionitrile; desferrioxamine; anthracyclines; 2,7,8-trihydroxy anthraquinones, fibrostatin-C; coumalic acid or its pharmaceutically acceptable salts; 5-oxaproline, β -lactam antibiotics. 25
5. A medicament for the improvement of neuronal regeneration comprising a therapeutically effective amount of an inhibitor substance which is capable of prevention or specific inhibition of basal membrane formation induced by a lesion of neuronal tissue comprising the inhibitor substance(s) selected from the group consisting of antibodies against collagen IV, laminin, entactin, accessory substances for proper function, or the assembly of the basal membrane; Fe-chelating agents, selected from the group consisting of pyridine derivatives, such as 5-aryl-carbonylamino- or 5-arylcarbamoyle-derivatives, 2-carboxylate, 2,5-dicarboxylate, their ethyl esters or ethyl amides or -5-acyl sulfonamides, 2,4 dicarboxylate, their ethyl esters or ethylamides, or dimethoxyethylamides; 3,4'-bipyridine, such as 5 amino-6-(1H)-one, 1,6-dihydro-2-methyl-6-oxo-5-carbonitril; 2,2'-bipyridine, such as 5,5'-dicarboxylic acid or its pharmaceutically acceptable salts, 4,4'-dicarboxylic acid ethyl ester or ethyl amide; 3,4'-dihydroxybenzoate, such as the diethyl ester; β -aminopropionitrile; desferrioxamine; anthracyclines; 2,7,8-trihydroxy anthraquinones, coumalic acid or its pharmaceutically acceptable salts; β -lactam antibiotics; 2-oxoglutarate competitors; in combination with a substance being capable of stimulating neuronal growth or inducing the expression of growth promoting proteins such as fibroblast growth factors, neural cell adhesion molecules like L1 (NILE), growth-associated proteins like GAP43 and anti-apoptotic proteins like bcl-2. 30 35 40 45 50 55

6. The medicament according to claim 5, wherein the inhibitor substance is applied in therapeutically effective amounts, such as 1 ng/kg to 1 mg/kg body weight.

Patentansprüche

1. Verwendung einer Inhibitorsubstanz, die aus der Gruppe ausgewählt ist, die aus folgenden besteht: Antikörpern gegen Collagen IV, Laminin, Entactin, Hilfssubstanzen für die richtige Funktion oder den Zusammenbau der Basalmembran; Fe-Chelatisierungsmitteln; Inhibitoren von Aminosäure-Hydroxylasen, wie Prolyl-4-Hydroxylase, Lysin-Hydroxylase; 2-Oxoglutarat-Konkurrenten; wobei die Inhibitorsubstanz die durch eine Läsion von neuronalem Gewebe induzierte Bildung der Basalmembran zu verhindern oder spezifisch zu hemmen vermag, zur Herstellung eines Medikaments zur Verbesserung der neuronalen Regeneration. 10 15
2. Verwendung gemäß Anspruch 1, wobei die Inhibitorsubstanz eine Substanz ist, die die Synthese von die Basalmembran aufbauenden Elementen oder den Zusammenbau von die Basalmembran aufbauenden Elementen oder sowohl die Synthese von die Basalmembran aufbauenden Elementen als auch den Zusammenbau von die Basalmembran aufbauenden Elementen hemmt. 20 25
3. Verwendung gemäß Anspruch 2, wobei die die Basalmembran aufbauenden Elemente aus der Gruppe ausgewählt sind, die aus Collagen IV, Laminin, Entactin, Hilfssubstanzen für die richtige Funktion oder den Zusammenbau der Basalmembran oder sowohl für die richtige Funktion als auch für den Zusammenbau der Basalmembran besteht. 30 35
4. Verwendung gemäß Anspruch 1, wobei die Inhibitorsubstanz aus der Gruppe ausgewählt ist, die aus folgenden besteht: N-Oxaloglycin; Zn-Salzen; Pyridinderivaten, wie 5-Arylcarbonylamino- oder 5-Arylcarbamoyle-Derivaten, 2-Carboxylat, 2,5-Dicarboxylat, ihren Ethylestern oder Ethylamiden oder 5-Acylsulfonamiden, 2,4-Dicarboxylat, ihren Ethylestern oder Ethylamiden oder Dimethoxyethylamiden; 3,4'-Bipyridin, wie 5-Amino-6-(1H)-on, 1,6-Dihydro-2-methyl-6-oxo-5-carbonitril; 2,2'-Bipyridin, wie 5,5'-Dicarbonsäure oder ihren pharmazeutisch annehmbaren Salzen, 4,4'-Dicarbonsäureethylester oder -ethylamid; 3,4'-Dihydroxybenzoat, wie dem Diethylester; Prolin und seinen strukturellen und funktionellen Analoga; β -Aminopropionitril; Desferrioxamin; Anthracyclinen; 2,7,8-Trihydroxyanthrachinonen, Fibrostatin-C; Cumarinsäure oder seinen pharmazeutisch annehmbaren Salzen; 5-Oxaprolin, β -Lactam-Antibiotika. 40 45 50 55

5. Medikament zur Verbesserung der neuronalen Regeneration, das folgendes umfasst: eine therapeutisch wirksame Menge einer Inhibitorsubstanz, welche die durch eine Läsion von neuronalem Gewebe induzierte Bildung der Basalmembran zu verhindern oder spezifisch zu hemmen vermag und die Inhibitorsubstanz(en) umfasst, die aus der Gruppe ausgewählt sind, die aus Folgenden besteht: Antikörpern gegen Collagen IV, Laminin, Entactin, Hilfssubstanzen für die richtige Funktion oder den Zusammenbau der Basalmembran; Fe-Chelatisierungsmitteln, die aus der Gruppe ausgewählt sind, die aus Pyridinderivaten, wie 5-Arylcarbonylamino- oder 5-Arylcarbamoyle-Derivaten, 2-Carboxylat, 2,5-Dicarboxylat, ihren Ethylestern oder Ethylamiden oder 5-Acylsulfonamiden, 2,4-dicarboxylat, ihren Ethylestern oder Ethylamiden oder Dimethoxyethylamiden; 3,4'-Bipyridin, wie 5-Amino-6-(1H)-on, 1,6-Dihydro-2-methyl-6-oxo-5-carbonitril; 2,2'-Bipyridin, wie 5,5'-Dicarbonsäure oder ihren pharmazeutisch annehmbaren Salzen, 4,4'-Dicarbonsäureethylester oder -ethylamid; 3,4'-Dihydroxybenzoat, wie dem Diethylester; β -Aminopropionitril; Desferrioxamin; Anthracyclinen; 2,7,8-Trihydroxyanthraquinonen, Cumarinsäure oder seinen pharmazeutisch annehmbaren Salzen; β -Lactam-Antibiotika besteht; 2-Oxoglutarat-Konkurrenten; in Kombination mit einer Substanz, die das neuronale Wachstum zu stimulieren oder die Expression von wachstumsfördernden Proteinen, wie Fibroblasten-Wachstumsfaktoren, Nervenzellenadhäsionsmolekülen, wie L1 (NILE), wachstumsassoziierten Proteinen, wie GAP43, und antiapoptotischen Proteinen, wie bcl-2, zu induzieren vermag.
6. Medikament gemäß Anspruch 5, wobei die Inhibitorsubstanz in therapeutisch wirksamen Mengen, wie 1 ng/kg bis 1 mg/kg Körpergewicht, angewendet wird.

Revendications

1. Utilisation d'une substance inhibitrice choisie dans la classe formée par les anticorps dirigés contre le collagène IV, la laminine, l'entactine, des substances auxiliaires pour la fonction correcte ou l'assemblage de la membrane basale ; des agents chélateurs de Fe ; des inhibiteurs d'acide-amino-hydroxylases telles que la prol-4-hydroxylase, la lysine-hydroxylase ; des compétiteurs de 2-oxoglutarate ; laquelle substance inhibitrice est capable de prévention ou d'inhibition spécifique de la formation de membrane basale induite par une lésion de tissu neuronal, pour la fabrication d'un médicament destiné à améliorer la régénération neuronale.
2. Utilisation selon la revendication 1, dans laquelle la

substance inhibitrice est une substance inhibant la synthèse d'éléments constitutifs de membrane basale ou l'assemblage d'éléments constitutifs de membrane basale, ou à la fois la synthèse d'éléments constitutifs de membrane basale et l'assemblage d'éléments constitutifs de membrane basale.

3. Utilisation selon la revendication 2, dans laquelle les éléments constitutifs de membrane basale sont choisis dans la classe formée par le collagène IV, la laminine, l'entactine, les substances auxiliaires pour la fonction correcte ou l'assemblage de la membrane basale ou à la fois la fonction correcte et l'assemblage de la membrane basale.
4. Utilisation selon la revendication 1, dans laquelle la substance inhibitrice est choisie dans le groupe formé par la N-oxaloglycine ; les sels de Zn ; des dérivés de pyridine tels que les dérivés à substitution 5-arylcarbonylamino ou 5-arylcarbamoyle, le 2-carboxylate, le 2,5-dicarboxylate, leurs esters d'éthyle ou éthylamides ou -5-acylsulfonamides, le 2,4-dicarboxylate, ses esters d'éthyle ou éthylamides ou diméthoxyéthylamides ; une 3,4'-bipyridine telle que la 5-amino-6-(1H)-one, le 1,6-dihydro-2-méthyl-6-oxo-5-carbonitrile ; une 2,2'-bipyridine telle que l'acide 5,5'-dicarboxylique ou ses sels pharmaceutiquement acceptables, l'ester éthylique ou l'éthylamide d'acide 4,4'-dicarboxylique ; un 3,4'-dihydroxybenzoate tel que l'ester diéthylique ; la proline et ses analogues structuraux et fonctionnels ; le β -aminopropionitrile ; la desferrioxamine ; les anthracyclines ; les 2,7,8-trihydroxyanthraquinones, la fibrostatine-C ; l'acide coumalique ou ses sels pharmaceutiquement acceptables ; la 5-oxaproline, les antibiotiques de la famille des β -lactames.
5. Un médicament destiné à l'amélioration de la régénération neuronale, comprenant une quantité à effet thérapeutique d'une substance inhibitrice qui est capable de prévention ou d'inhibition spécifique de la formation de membrane basale induite par une lésion de tissu neuronal, comprenant une ou plusieurs substances inhibitrices choisies dans la classe formée par des anticorps dirigés contre le collagène IV, la laminine, l'entactine, des substances auxiliaires pour la fonction correcte ou l'assemblage de la membrane basale ; des agents chélateurs de Fe sélectionnés dans le groupe formé par des dérivés de pyridine tels que les dérivés 5-arylcarbonylamino ou 5-arylcarbamoyle, le 2-carboxylate, le 2,5-dicarboxylate, leurs esters d'éthyle ou éthylamides ou -5-acylsulfonamides, le 2,4-dicarboxylate, ses esters d'éthyle ou éthylamides ou diméthoxyéthylamides ; une 3,4'-bipyridine telle que la 5-amino-6-(1H)-one, le 1,6-dihydro-2-méthyl-6-oxo-5-carbonitrile ; une 2,2'-bipyridine telle que l'acide 5,5'-dicarboxylique ou ses sels pharmaceutiquement acceptables, l'es-

ter éthylique ou l'éthylamide d'acide 4,4'-dicarboxylique ; un 3,4'-dihydroxybenzoate tel que l'ester diéthylique ; le β -aminopropionitrile ; la desferrioxamine ; les anthracyclines ; les 2,7,8-trihydroxyanthraquinones, l'acide coumalique ou ses sels pharmaceutiquement acceptables ; les antibiotiques de la famille des β -lactames ; des compétiteurs de 2-oxoglutarate ; en association avec une substance capable de stimuler la croissance neuronale ou d'induire l'expression de protéines activatrices de croissance telle que des facteurs de croissance des fibroblastes, des molécules d'adhérence des cellules nerveuses comme L1 (NILE), des protéines associées à la croissance comme GAP43 et des protéines anti-apoptotiques comme bcl-2.

6. Le médicament selon la revendication 5, dans lequel la substance inhibitrice est appliquée en des quantités à effet thérapeutique, par exemple de 1 ng/kg à 1 mg/kg de poids corporel.




Fig. 1




Fig. 2




Fig. 3




Fig. 4

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 9319783 A [0003]
- US 5082926 A [0004]
- US 5493008 A [0005]
- WO 9600582 A [0006]

Non-patent literature cited in the description

- REIER, P. J ; STENSAAS, L. J. ; GUTH, L. Spinal cord reconstruction. Raven Press, 1983, 163-196 [0035]
- CLEMENTE, C. D. Regeneration in the nervous system. Springfield, IL, 1955, 147-161 [0035]
- SCHNELL, L ; SCHWAB, M. E. *Eur. J. Neurosci*, 1993, vol. 5, 1156-1171 [0035]
- KRÜGER, S ; SIEVERS, J ; HANSEN, C ; SADLER, M ; BERRY, M. *J. Comp. Neurol*, 1986, vol. 249, 103-116 [0035]
- BERRY, M ; CARLILE, J ; HUNTER, A. *J. Neurocytol*, 1996, vol. 25, 147-170 [0035]
- STICHEL, C. C ; MÜLLER, H. W. *J. Neurocytol*, 1994, vol. 23, 615-630 [0035]
- FERINGA, E. R ; KOWALSKI, T. F ; VAHLSING, H. L. *Ann. Neurol*, 1980, vol. 8, 148-154 [0035]
- ALLEN, G. V ; HOPKINS, D. A. *J. Comp. Neurol*, 1988, vol. 275, 39-64 [0035]
- STICHEL, C. C ; WUNDERLICH, G ; SCHWAB, M. E ; MÜLLER, H. W. *Europ. J. Neurosci*, 1995, vol. 7, 401-411 [0035]
- YURCHENCO, P. D ; SCHITTNY, J. C. *FASEB. J*, 1996, vol. 4, 1577-1590 [0035]
- TIMPL, R ; BROWN, J. *Curr. Opin. Cell. Biol*, 1996, vol. 8, 618-624 [0035]
- KIVIRIKKO, K. L ; MYLLYLÄ, R ; PIHLAJANIEMI, T. *FASEB. J*, 1989, vol. 3, 1609-1617 [0035]
- IKEDA, H ; WU, G. Y ; WU, C. H. *Hepatology*, 1992, vol. 15, 282-287 [0035]
- BIENKOWSKI, R. S. *J. Cell. Physiol*, 1984, vol. 121, 152-158 [0035]
- ANDERSEN, P ; SILFVENIUS, H ; SUNDBERG, S ; SVEEN, O ; WIGSTRÖM, H. *Brain. Res*, 1978, vol. 144, 11-18 [0035]
- MCKERRACHER, L et al. *Neuron*, 1994, vol. 13, 805-811 [0035]
- MUKHOPADHYAY, G ; DOHERTY, P ; WALSH, F. S ; CROCKER, P. R ; FILBIN, M. T. *Neuron*, 1994, vol. 13, 757-767 [0035]
- SCHWAB, M. E ; CARONI, P. *J. Neurosci*, 1988, vol. 8, 2381-2393 [0035]
- SNOW, A. D ; LEMMON, V. ; CARRINO, D. A ; CAPLAN, A. I ; SILVER, J. *Exp. Neurol*, 1990, vol. 109, 111-130 [0035]
- LIPS, K ; STICHEL, C. C ; MÜLLER, H. W. *J. Neurocytol*, 1995, vol. 24, 449-464 [0035]
- IWASHITA, Y ; KAWAGUCHI, S ; MURATA, M. *Nature*, 1994, vol. 367, 167-170 [0035]
- KALDERON, N ; FUKS, Z. *Proc. Natl. Acad. Sci*, 1996, vol. 93, 11179-11184 [0035]
- OLUCHA, F ; MARTINEZ-GARCIA, F ; LOPEZ-GARCIA, C. *J. Neurosci. Meth*, 1985, vol. 13, 131-138 [0035]
- PUCHALA, E ; WINDLE, W. F. *Exp. Neurol.*, 1977, vol. 55, 1-42 [0035]
- GUTH, L ; ALBUQUERQUE, E. X ; DESHPANDE, S. S ; BARRET, C. P ; DONATI, E. J ; WARNICK, J. E. *J. Neurosurg.*, 1980, vol. 52, 73-86 [0035]