

(19)



(11)

**EP 1 002 040 B2**

(12)

**NEW EUROPEAN PATENT SPECIFICATION**

After opposition procedure

(45) Date of publication and mention  
of the opposition decision:

**21.12.2011 Bulletin 2011/51**

(45) Mention of the grant of the patent:

**18.02.2004 Bulletin 2004/08**

(21) Application number: **97933510.6**

(22) Date of filing: **09.07.1997**

(51) Int Cl.:

**C11D 3/386** (2006.01)

(86) International application number:

**PCT/US1997/012439**

(87) International publication number:

**WO 1999/002639 (21.01.1999 Gazette 1999/03)**

(54) **CLEANING COMPOSITIONS COMPRISING A SPECIFIC OXYGENASE**

REINIGUNGSZUSAMMENSETZUNGEN ENTHALTEND EINE SPEZIFISCHE OXYGENASE

COMPOSITIONS DE LAVAGE/NETTOYAGE CONTENANT UNE OXYGENASE SPECIFIQUE

(84) Designated Contracting States:

**AT BE CH DE DK ES FI FR GB GR IE IT LI LU NL  
PT SE**

(43) Date of publication of application:

**24.05.2000 Bulletin 2000/21**

(73) Proprietor: **THE PROCTER & GAMBLE COMPANY**

**Cincinnati, OH 45202 (US)**

(72) Inventors:

- **HERBOTS, Ivan, Maurice, Alfons, Jan**  
**B-9230 Wetteren (BE)**
- **BUSCH, Alfred**  
**B-1840 Londerzeel (BE)**

- **BARNABAS, Mary, Vijayarani**  
**West Chester, OH 45069 (US)**
- **BETTIOL, Jean-Luc, Philippe**  
**B-1200 Brussels (BE)**

(74) Representative: **Peet, Jillian Wendy et al**

**Procter & Gamble Technical Centres Limited**  
**Whitley Road**  
**Longbenton**  
**Newcastle upon Tyne**  
**NE12 9TS (GB)**

(56) References cited:

**EP-A- 0 086 139 WO-A-92/18683**  
**WO-A-95/01426 US-A- 5 558 812**

**EP 1 002 040 B2**

**Description**Field of the Invention

- 5 **[0001]** The present invention relates to cleaning compositions, including laundry, dishwashing, hard surface cleaner, oral/dental cleaning compositions, comprising a polyphenol / heterocyclic substrate based oxygenase.

Background of the invention

- 10 **[0002]** Performance of a cleaning product, for use in washing or cleaning method, is judged by a number of factors, including the ability to remove soils, and the ability to prevent the redeposition of the soils, or the breakdown products of the soils on the articles in the wash.

- [0003]** Coloured stains / soils are often difficult to remove effectively from a soiled item. Highly coloured stains and soils i.e. derived from fruit and/or vegetables are particularly challenging soils to remove. This stains and soils contain colour-bodies based on carotenoids compounds such as  $\alpha$ -,  $\beta$ - and  $\gamma$ -carotene and lycopene, on porphyrins such as chlorophyll and on flavonoid pigments and dye components. This latter group of natural flavonoid based dye components comprises the highly coloured anthocyanins dyes and pigments based on pelargonidin, cyanidin, delphinidin and their methyl esters and the antoxanthins. These compounds are the origin of most of the orange, red, violet and blue colours occurring in fruits and are abundant in all berries, cherry, red and black currents, grapefruits, passion fruit, oranges, lemons, apples, pears, pomegranate, red cabbage, red beets and also flowers. Derivatives of cyanidin are present in up to 80% of the pigmented leaves, in up to 70% of fruits and in up to 50% of flowers. Specific examples of such soils would include tea, coffee, spices such as curry and paprika, orange, tomato, banana, tea, mango, broccoli, carrot, beetroot, spinach soils and grass. Ball pens' ink are also known to be highly difficult coloured stains to be removed. These highly coloured flavonoid and carotenoid dyes are often polycyclic and heterocyclic compounds having a conjugated double bond systems. This chemical structure is often responsible for the pronounced colour of the compounds.

- [0004]** In addition, the complex nature of everyday "body" soils typically found on pillow cases, T-shirts, collars and socks, provides a continuous thorough cleaning challenge for detergents. These soils are difficult to remove completely and often residues build up on fabric leading to dinginess and yellowing. Everyday body soils are also found on sanitary and kitchen surfaces such as bathtubs, toilet bowl and dishware.

- 30 **[0005]** The items can be fabrics, hard surfaces, dishware such as plasticware, glassware or chinaware, teeth and mouth.

- [0006]** Traditionally, high levels of bleaching compounds, optionally with bleach precursors and/or bleach enhancers, are incorporated in cleaning compositions. Bleaching agents are compounds which are precursors of hydrogen peroxide which is formed in the course of the washing procedure. Perborates and percarbonates are the most important examples of such hydrogen peroxide precursors.

- 35 **[0007]** In view of the above, there exists clearly a continuous need to provide cleaning compositions which have an excellent detergency performance. Accordingly it is an object of the present invention to provide a cleaning composition which provides effective and efficient cleaning of coloured and everyday body soils and/or stains. It is a further object to provide a cleaning composition which provides fabric realistic items cleaning and whitening while providing excellent safety for the fabrics' colours.

- 40 **[0008]** The above objective has been met by formulating cleaning compositions comprising a polyphenol / heterocyclic substrate based oxygenase, in accordance with claim 1.

- [0009]** It has been surprisingly found an enzymatic bleach system based on a polyphenol / heterocyclic substrate based oxygenase in accordance with claim 1 delivers in a cleaning composition, bleach-like benefits in an unexpected broad range of performance areas such as dingy cleaning, whiteness maintenance and stain removal, especially on polyphenols and heterocyclic compounds-containing and everyday body stains/soils. It has also been found that the cleaning compositions of the present invention provide sanitisation of the treated surfaces.

It has been further found that the performance of the cleaning compositions of the present invention is improved by the addition of another enzymatic bleach system, a conventional activated bleach system, a metallo catalyst based bleach system and/or another detergent enzyme.

- 50 **[0010]** In a preferred embodiment, the present invention relates to a laundry composition comprising a polyphenol / heterocyclic substrate based oxygenase in accordance with claim 1, further providing fabric realistic items cleaning and whitening while providing excellent safety for the fabrics' colours. In a second embodiment, the present invention relates to dishwashing or household cleaning compositions comprising a polyphenol / heterocyclic substrate based oxygenase in accordance with claim 1. In a third embodiment, the present invention relates to oral/dental care compositions comprising a polyphenol / heterocyclic substrate based oxygenase in accordance with claim 1.

- 55 **[0011]** WO92/18683 describes a process for bleaching of dyed textiles such as coloured denim, comprising the treatment with a bleach liquor containing hydrogen peroxide or a source thereof and an enzyme exhibiting peroxidase activity or oxygen and an enzyme exhibiting a suitable oxidase activity. EP 086 139 discloses the cloning and expression of the

catechol 2,3 oxygenase for the food, pharmaceutical and cosmetic industries. This enzyme is also used for the decontamination of textiles from urushiol and its derivatives and can be added in laundry detergents or body cleaning compositions therefor.

**[0012]** WO95/01426 relates to agents capable of enhancing the activity of laccases and related enzymes. WO95/01426 is also directed to methods of oxidising a substrate in the presence of a laccase enzyme and an enhancing agent. Those enzymes are laccase EC 1.10.3.2, catechol oxidase EC.1.10.3.1, bilirubin oxidase EC 1.3.3.5 and monophenol monooxygenase EC 1.14.99.1.

**[0013]** However, the use of a polyphenol / heterocyclic substrate based oxygenase in accordance with claim 1 in a cleaning composition for polyphenols and heterocyclic compounds-containing stains/soils and everyday body soils substantive removal and for sanitisation, has never been previously recognised. Nor have been recognised the substantive fabric realistic items cleaning and whitening performance and excellent fabric colour safety benefit when the polyphenol / heterocyclic substrate based oxygenase in accordance with claim 1 is used within a laundry detergent composition.

#### Summary of the invention

**[0014]** The present invention relates to cleaning compositions, including laundry, dishwashing, hard surface cleaner, oral/dental cleaning compositions, comprising a specific polyphenol / heterocyclic substrate based oxygenase, which provide effective and efficient cleaning of coloured stains and/or soils and everyday body soils, and provide sanitisation of the treated surfaces.

**[0015]** Furthermore, the cleaning compositions of the present invention provide fabric realistic items cleaning and whitening performance while providing excellent fabric colour safety when formulated as a laundry detergent composition.

#### Detailed description of the invention

**[0016]** The essential component of the cleaning compositions of the present invention is a polyphenol and/or heterocyclic substrate based oxygenase in accordance with claim 1. Preferably said enzyme is further characterised by being an iron sulphur or iron heme oxygenase and/or a heavy metal dependant oxygenase.

**[0017]** All oxygenases of the present invention will pass the following stain removal performance test achieved on standard test fabrics and being compared to a nil-oxygenase reference in same conditions.

The small scale test is done in a launderometer Washtec ROACHES equipped with stainless steel jars of 500 ml. The small scale test is done at 30-40 °C with total wash time of 30 minutes. Tests is done in a 1% detergent solution using the detergent as described in example 10, composition 5. The detergent is dissolved in 400 ml of water with a total hardness between 2.0 - 3.0 mmol Ca<sup>2+</sup>/L. The pH of the 1% detergent solution is adjusted with an acid (citric acid) or an alkali (NaOH) to pH 8-9. Each jar also contains 15 steel ball for better agitation. Standard stain test fabrics are supplied by wfk-Testgewebe GmbH (Christenfeld, 10-Brueggen, Germany). At least two replicates are run per test. The size of the stains is 3 cmx4 cm. The standard test fabric having the polyphenol/heterocyclic substrate are the wfk10 J (tea-stained cotton); wfk10 K (coffee-stained cotton) and wfk10L (red-wine stained cotton). The enzyme level in this small scale test is 1 mg enzyme protein/L. Levels of cofactor(s) when required are calculated according to the enzyme to cofactor(s) ratio known from literature.

For the reference treatment, no enzyme or co-factor is added to the solution. In the test treatment the pre-weighted enzyme and cofactor is added to the solution and in a second test treatment only the co-factor is added to the solution. After that the test tracers are put in the jars. The launderometer is set on the required temperature. The launderometer is preheated to the right temperature before starting the test so the test is run at a constant temperature.

The jars are closed and put in the preheated launderometer. The cycle time is set and the test is started. After 30 minutes the test is stopped, the jars are taken out of the equipment and are opened. The standard test fabrics are taken out and rinsed 3 times in cold city water (hardness 2-3 mmole Ca<sup>2+</sup>/L). After rinsing the tracers are tumble dried in a normal household drier to completely dry.

The stain removal performance of the enzyme is analyzed by visual grading by an expert panel or preferentially by an instrumental stain removal measurement e.g. with the Spectraflash 500 apparatus from Datacolor. The Spectraflash 500 settings used for this test are : specular excluded, aperture small and U.V. filter FL40 (= UV cut off filter at 400nm) and calibration is done versus a standard white and black.

The result expressed in color difference (Cielab) dE, is calculated between the test and reference treatment. Test 1 comprises the oxygenase of the present invention, test 2 comprises a monophenol monooxygenase of EC 1.14.18.1. The enzymes of the invention (Test 1) have a dE of 1 or more versus the monophenol monooxygenase (Test 2) tested in same conditions.

**[0018]** It has been found that the cleaning compositions of the present invention provide effective and efficient cleaning of everyday body soils and coloured stains and/or soils and in particular provide fabric realistic items cleaning and whitening while providing colour safety.

**[0019]** Without wishing to be bound by theory, it is believed that the polyphenol / heterocyclic substrate based oxygenase in accordance with claim 1, decyclizes and fragments aromatic compounds, often substituted benzenediol based and benzoate based compounds and the heterocyclic compounds with or without multiple hydroxyl substitution. The latter compounds are often based on pyridine, pyrimidine or indole basic structures. It has also been found that hydroxylation or epoxidation of the coloured polyphenolic and heterocyclic compounds can occur, rendering the reaction products more soluble in the wash solution and hence easier to remove. Said enzymatic reactions are achieved by catalytic direct insertion of molecular oxygen in to the coloured compound structure. When the cleaning composition of the present invention is formulated as a laundry detergent, it has been found that the polyphenol/ heterocyclic substrate based oxygenase in accordance with claim 1 therein included do not darken the polyphenolic stains by radical induced polymerisation, nor fade the sensitive dyes thereby do provide excellent safety to the fabrics' colours.

**[0020]** In addition, the cleaning compositions of the present invention provide sanitisation of the treated surfaces. Sanitisation includes all positive effects obtained by the inhibition or reduction of microbial activity on fabrics and other surfaces, such as the prevention of malodour development and bacterial/fungal growth. For example, it provides prevention of malodour development on stored and wearied fabrics, on stored dishware, especially plastic kitchen gear and in toilets. In particular, the composition of the invention will inhibit or at least reduce the bacterial and/or fungal development on moist fabric waiting for further laundry processing and thereby preventing the formation of malodour. In addition, bacterial and/or fungal growth on hard surfaces such as tiles and their silicone joints, sanitary installations, will be prevented.

The sanitisation potential of the cleaning compositions of the present invention can be enhanced by the addition of chemical sanitisers such as Triclosan and/or hexemidine. *Parfums Cosmetiques Actualités* No 125, Nov, 1995, 51-4 describes suitable chemical sanitisers.

The sanitisation benefits of the cleaning compositions of the present invention can be evaluated by the Minimum Inhibitory Concentration (MIC) as described in *Tuber. Lung. Dis.* 1994 Aug; 75(4):286-90; *J. Clin. Microbiol.* 1994 May; 32(5): 1261-7 and *J. Clin. Microbiol.* 1992 Oct; 30(10):2692-7.

**[0021]** Without wishing to be bound by theory, it is believed that "everyday body soils" contain sebum excreted by the human body. Sebum is believed to contain large quantities of saturated and unsaturated fatty acids, sterols and sterols esters (The physiology and Pathology of the skin, Vol. 9, A. Jarret, (1986)). The fragmentation of the substrate and the formation of ionizable groups or hydrophilic substituents by a of a polyphenol / heterocyclic substrate based oxygenase enzyme in accordance with claim 1 renders the enzymatic reaction products more soluble and hence easier to be removed from the soiled items.

**[0022]** The polyphenol / heterocyclic substrate based oxygenases for the purpose of the present invention are listed below :

EC Number	Recommended Name
1.13.11.1	CATECHOL 1,2-DIOXYGENASE
1.13.11.3	PROTocatechuate 3,4-DIOXYGENASE
1.13.11.4	GENTISATE 1,2-DIOXYGENASE
1.13.11.5	HOMOGENTISATE 1,2-DIOXYGENASE
1.13.11.6	3-HYDROXYANTHRANILATE 3,4-DIOXYGENASE
1.13.11.8	PROTocatechuate 4,5-DIOXYGENASE
1.13.11.9	2,5-DIHYDROXYPYRIDINE 5,6-DIOXYGENASE
1.13.11.10	7,8-DIHYDROXYKYNURENATE 8,8A-DIOXYGENASE
1.13.11.14	2,3-DIHYDROXYBENZOATE 3,4-DIOXYGENASE
1.13.11.15	3,4-DIHYDROXYPHENYLACETATE 2,3-DIOXYGENASE
1.13.11.16	3-CARBOXYETHYLCATECHOL 2,3-DIOXYGENASE
1.13.11.17	INDOLE 2,3-DIOXYGENASE
1.13.11.22	CAFFEATE 3,4-DIOXYGENASE
1.13.11.23	2,3-DIHYDROXYINDOLE 2,3-DIOXYGENASE
1.13.11.28	2,3-DIHYDROXYBENZOATE 2,3-DIOXYGENASE
1.13.11.35	PYROGALLOL 1,2-OXYGENASE
1.13.11.36	CHLORIDAZON-CATECHOL DIOXYGENASE
1.13.11.37	HYDROXYQUINOL 1,2 DIOXYGENASE
1.13.11.38	1-HYDROXY-2 NAPHTHOATE 1,2 DIOXYGENASE
1.13.11.39	BIPHENYL-2,3-DIOL 1,2 DIOXYGENASE
1.13.11.41	2,4'-DIHYDROXYACETOPHENONE DIOXYGENASE

(continued)

EC Number	Recommended Name
5	1.13.11.42 INDOLEAMINE-PYRROLE 2,3 DIOXYGENASE
	1.13.11. 43 LIGNOSTILBENE $\alpha\beta$ -DIOXYGENASE
	1.13.12.11 METHYLPHENYL TETRAHYDROPYRIDINE N-MONOOXYGENASE
	1.14.11.3 PYRIMIDINE-DEOXYNUCLEOSIDE 2'-DIOXYGENASE
	1.14.11.6 THYMINE DIOXYGENASE
10	1.14.11.9 NARINGENIN 3-DIOXYGENASE
	1.14.11.10 PYRIMIDINE-DEOXYNUCLEOSIDE 1'-DIOXYGENASE
	1.14.11.11 HYOSCIAMINE (6S)-DIOXYGENASE
	1.14.11.12 GIBERELLIN-44 DIOXYGENASE
	1.14.11.13 GIBERELLIN 2 $\beta$ -DIOXYGENASE
15	1.14.11.14 6 $\beta$ -HYDROXYHYOSCIAMINE EPOXIDASE
	1.14.11.15 GIBERELLIN 3 $\beta$ -DIOXYGENASE
	1.14.12.1 ANTHRANILATE 1,2-DIOXYGENASE (DEAMINATING, DECARBOXYLATING)
	1.14.12.2 ANTHRANILATE 2,3-DIOXYGENASE
20	1.14.12.4 3-HYDROXY-2-METHYLPYRIDINECARBOXYLATE DIOXYGENASE
	1.14.12.5 5-PYRIDOXATE DIOXYGENASE
	1.14.12.6 2-HYDROXYCYCLOHEXANONE 2-MONOOXYGENASE
	1.14.12.7 PHTALATE 4,5 DIOXYGENASE
	1.14.12.10 BENZOATE 1,2-DIOXYGENASE
25	1.1.4.12.11 TOLUENE DIOXYGENASE
	1.14.12.12 NAPHTALENE 1,2 DIOXYGENASE
	1.14.13.1 SALICYLATE 1-MONOOXYGENASE
	1.14.13.2 4-HYDROXYBENZOATE 3-MONOOXYGENASE/HYDROLASE
30	1.14.13.3 4-HYDROXYPHENYLACETATE 3-MONOOXYGENASE
	1.14.13.4 MELILOTATE 3-MONOOXYGENASE
	1.14.13.5 IMIDAZOLACETATE 4-MONOOXYGENASE
	1.14.13.6 ORCINOL 2-MONOOXYGENASE
	1.14.13.8 DIMETHYLANILINE MONOOXYGENASE (N-OXIDE-FORMING)
35	1.14.13.9 KYNURENINE 3-MONOOXYGENASE
	1.14.13.10 2,6-DIHYDROXYPYRIDINE 3-MONOOXYGENASE
	1.14.13.11 TRANS-CINNAMATE 4-MONOOXYGENASE
	1.14.13.12 BENZOATE 4-MONOOXYGENASE
40	1.14.13.14 TRANS-CINNAMATE 2-MONOOXYGENASE
	1.14.13.16 CYCLOPENTANONE MONOOXYGENASE
	1.14.13.18 4-PHENYLHYDROXYACETATE 1-MONOOXYGENASE
	1.14.13.21 FLAVONOID 3'-MONOOXYGENASE
	1.14.13.22 CYCLOHEXANONE MONOOXYGENASE
45	1.14.13.23 3-HYDROXYBENZOATE 4-MONOOXYGENASE
	1.14.13.24 3-HYDROXYBENZOATE 6-MONOOXYGENASE
	1.14.13.27 4-AMINOBENZOATE 1-MONOOXYGENASE
	1.14.13. 29 4-NITROPHENOL 2-MONOOXYGENASE
50	1.14.13. 31 2-NITROPHENOL 2-MONOOXYGENASE
	1.14.13. 33 4-HYDROXYBENZOATE 3-MONOOXYGENASE
	1.14.13.35 ANTHRANILATE 3-MONOOXYGENASE
	1.14.13. 36 5-O-(4-COUMAROYL)-D-QUINATE 3'-MONOOXYGENASE
	1.14.13. 38 ANHYDROTETRACYCLINE MONOOXYFENASE
55	1.14.13. 40 ANTHRANILOYL-CoA MONOOXYGENASE
	1.14.13.43 QUESTIN MONOOXYGENASE
	1.14.13. 44 2-HYDROXYBIPHENYL 3-MONOOXYGENASE

(continued)

EC Number	Recommended Name
1.14.13.46	(-)-MENTHOL MONOOXYGENASE
1.14.14.3	ALKANAL MONOOXYGENASE (FMN-LINKED) /BACTERIAL LUCIFERASE
1.14.15.2	CAMPOR 1,2-MONOOXYGENASE
1.14.16.3	ANTHRANILATE 3-MONOOXYGENASE
1.14.16.6	MANDELATE 4-MONOOXYGENASE
1.14.17.1	DOPAMINE B-MONOOXYGENASE /DOPAMINE B-HYDROXYLASE
1.14.99.2	KYNURENINE 7,8-HYDROXYLASE
1.14.99.15	4-METHOXYBENZOATE MONOOXYGENASE (O-DEMETHYLATING)
1.14.99.20	PHYLLLOQUINONE MONOOXYGENASE (2,3-EPOXIDIZING)
1.14.99.23	3-HYDROXYBENZOATE 2-MONOOXYGENASE

**[0023]** Preferred polyphenol / heterocyclic substrate based oxygenase enzymes for the present invention are the decyclising and hydroxylating mono- and di-oxygenases, more preferred are the following enzymes :

EC Number	recommended Name
1.13.11.3	PROTocatechuate 3,4-dioxygenase
1.13.11.14	2,3-dihydroxybenzoate 3,4-dioxygenase
1.13.11.17	Indole 2,3-dioxygenase
1.13.11.22	CAFFEATE 3,4-dioxygenase
1.13.11.35	PYROGALLOL 1,2-OXYGENASE
1.14.11.9	NARINGENIN 3-dioxygenase
1.14.12.7	PHTALATE 4,5 DIOXYGENASE
1.14.12.10	BENZOATE 1,2-DIOXYGENASE
1.14.12.11	TOLUENE DIOXYGENASE
1.14.13.2	4-HYDROXYBENZOATE 3-MONOOXYGENASE/HYDROXYLASE
1.14.13.12	BENZOATE 4-MONOOXYGENASE
1.14.13.21	FLAVONOID 3'-MONOOXYGENASE

**[0024]** Some of the polyphenol / heterocyclic substrate based oxygenases of the present invention require the presence of a cofactor. In this instance, the cleaning compositions of the present invention will further comprise a cofactor such as Ascorbate, oxoglutarate, Flavine Mononucleotide (FMN), Flavine Adenine Dinucleotide (FAD), Nicotine amide Adenine Dinucleotide (Phosphate) (NAD(P)H), tetrahydropterine. When therein included, the cofactor will be comprised at a weight ratio of pure oxygenase to cofactor comprised generally between 10:1 to 1:10, preferably between 5:1 to 1:8, more preferably between 1:2 to 1:5.

**[0025]** The polyphenol / heterocyclic substrate based oxygenase enzyme in accordance with claim 1 is incorporated into the cleaning compositions in accordance with the invention preferably at a level of from 0.0001 % to 2%, more preferably from 0.001% to 0.5%, most preferably from 0.002% to 0.1% pure enzyme by weight of the composition.

**[0026]** Preferred polyphenol / heterocyclic substrate based oxygenases in accordance with claim 1 for specific applications are alkaline polyphenol / heterocyclic substrate based oxygenases, i.e. enzymes having an enzymatic activity of at least 10%, preferably at least 25%, more preferably at least 40% of its maximum activity at a pH ranging from 7 to 12. More preferred polyphenol / heterocyclic substrate based oxygenases are enzymes having their maximum activity at a pH ranging from 7 to 12.

**[0027]** Enzymes homologue to the polyphenol / heterocyclic substrate based oxygenase of the present invention are also contemplated. The term "homologue" is intended to indicate a polypeptide encoded by DNA which hybridises to the same probe as the DNA coding for the polyphenol / heterocyclic substrate based oxygenase enzyme with this amino acid sequence under certain specific conditions (such as presoaking in 5xSSC and prehybridising for 1 h at ~40°C in a solution of 20% formamide, 5xDenhardt's solution, 50 mM sodium phosphate, pH 6.8, and 50 µg of denaturated sonicated calf thymus DNA, followed by hybridisation in the same solution supplemented with 100 µM ATP for 18 h at ~40°C). The term is intended to include derivatives of the polyphenol / heterocyclic substrate based oxygenase enzyme sequence obtained by addition of one or more amino acid residues to either or both the C- and N-terminal of the native sequence,

substitution of one or more amino acid residues at one or more sites in the native sequence, deletion of one or more amino acid residues at either or both ends of the native amino acid sequence or at one or more sites within the native sequence, or insertion of one or more amino acid residues at one or more sites of the native sequence.

**[0028]** The above-mentioned enzymes may be of any suitable origin, such as vegetable, animal, bacterial, fungal and yeast origin. Origin can further be mesophilic or extremophilic (psychrophilic, psychrotrophic, thermophilic, barophilic, alkalophilic, acidophilic, halophilic, etc.). Purified or non-purified forms of these enzymes may be used. Nowadays, it is common practice to modify wild-type enzymes via protein / genetic engineering techniques in order to optimise their performance efficiency in the cleaning compositions of the invention. For example, the variants may be designed such that the compatibility of the enzyme to commonly encountered ingredients of such compositions is increased. Alternatively, the variant may be designed such that the optimal pH, bleach or chelant stability, catalytic activity and the like, of the enzyme variant is tailored to suit the particular cleaning application.

**[0029]** In particular, attention should be focused on amino acids sensitive to oxidation in the case of bleach stability and on surface charges for the surfactant compatibility. The isoelectric point of such enzymes may be modified by the substitution of some charged amino acids, e.g. an increase in isoelectric point may help to improve compatibility with anionic surfactants. The stability of the enzymes may be further enhanced by the creation of e.g. additional salt bridges and enforcing calcium binding sites to increase chelant stability.

### **Detergent components**

**[0030]** The cleaning compositions of the present invention may also contain additional detergent components. The precise nature of these additional components, and levels of incorporation thereof will depend on the physical form of the composition, and the nature of the cleaning operation for which it is to be used.

**[0031]** The cleaning compositions preferably further comprise another enzymatic bleach system, a conventional activated bleach system, a metallo catalyst based bleach system and/or another detergent enzyme.

**[0032]** In a preferred embodiment, the present invention relates to a laundry and/or fabric care composition comprising a polyphenol / heterocyclic substrate based oxygenase in accordance with claim 1 (Examples 1-18). In a second embodiment, the present invention relates to dishwashing or household cleaning compositions including sanitisation compositions (Examples 19-28), and in a third embodiment, the present invention relates to oral/dental care compositions (Examples 29-31).

**[0033]** The cleaning compositions according to the invention can be liquid, paste, gels, bars, tablets, spray, foam, powder or granular forms. Granular compositions can also be in "compact" form, the liquid compositions can also be in a "concentrated" form.

**[0034]** The compositions of the invention may for example, be formulated as hand and machine dishwashing compositions, hand and machine laundry detergent compositions including laundry additive compositions and compositions suitable for use in the soaking and/or pretreatment of stained fabrics, rinse added fabric softener compositions, and compositions for use in general household hard surface cleaning operations. Compositions containing such polyphenol / heterocyclic substrate based oxygenase can also be formulated as oral/dental care compositions.

**[0035]** When formulated as compositions for use in manual dishwashing methods the compositions of the invention preferably contain a surfactant and preferably other detergent compounds selected from organic polymeric compounds, suds enhancing agents, group II metal ions, solvents, hydrotropes and additional enzymes.

**[0036]** When formulated as compositions suitable for use in a laundry machine washing method, the compositions of the invention preferably contain both a surfactant and a builder compound and additionally one or more detergent components preferably selected from organic polymeric compounds, bleaching agents, additional enzymes, suds suppressors, dispersants, lime-soap dispersants, soil suspension and anti-redeposition agents and corrosion inhibitors. Laundry compositions can also contain softening agents, as additional detergent components.

Such compositions containing polyphenol / heterocyclic substrate based oxygenase in accordance with claim 1 can provide fabric cleaning, stain removal, whiteness maintenance, softening, colour appearance, dye transfer inhibition and sanitisation when formulated as laundry detergent compositions.

**[0037]** The compositions of the invention can also be used as detergent additive products. Such additive products are intended to supplement or boost the performance of conventional detergent compositions.

**[0038]** If needed the density of the laundry detergent compositions herein ranges from 400 to 1200 g/litre, preferably 600 to 950 g/litre of composition measured at 20°C.

The "compact" form of the compositions herein is best reflected by density and, in terms of composition, by the amount of inorganic filler salt; inorganic filler salts are conventional ingredients of detergent compositions in powder form; in conventional detergent compositions, the filler salts are present in substantial amounts, typically 17-35% by weight of the total composition.

In the compact compositions, the filler salt is present in amounts not exceeding 15% of the total composition, preferably not exceeding 10%, most preferably not exceeding 5% by weight of the composition.

The inorganic filler salts, such as meant in the present compositions are selected from the alkali and alkaline-earth-metal salts of sulphates and chlorides.

A preferred filler salt is sodium sulphate.

[0039] Liquid detergent compositions according to the present invention can also be in a "concentrated form", in such case, the liquid detergent compositions according to the present invention will contain a lower amount of water, compared to conventional liquid detergents.

Typically the water content of the concentrated liquid detergent is preferably less than 40%, more preferably less than 30%, most preferably less than 20% by weight of the detergent composition.

### Conventional detergent enzymes

[0040] The cleaning compositions of the present invention can in addition to the polyphenol / heterocyclic substrate based oxygenase enzyme in accordance with claim 1 further comprise one or more enzymes which provide cleaning performance, fabric care and/or sanitisation benefits. It has been found that the combination of said specific oxygenase with detergent enzyme provides improved cleaning of polyphenol / heterocyclic coloured and everyday body stains and/or soils and when formulated as laundry composition, its achieves improved fabric realistic items cleaning and whitening while providing colour safety.

[0041] Said enzymes include enzymes selected from cellulases, hemicellulases, peroxidases, proteases, gluco-amylases, amylases, xylanases, lipases, phospholipases, esterases, cutinases, pectinases, keratanases, reductases, oxidases, phenoloxidases, lipoxygenases, ligninases, pullulanases, tannases, pentosanases, malanases,  $\beta$ -glucanases, arabinosidases, hyaluronidase, chondroitinase, laccase or mixtures thereof.

[0042] A preferred combination is a cleaning composition having cocktail of conventional applicable enzymes like protease, amylase, lipase, cutinase and/or cellulase in conjunction with one or more plant cell wall degrading enzymes.

[0043] The cellulases usable in the present invention include both bacterial or fungal cellulases. Preferably, they will have a pH optimum of between 5 and 12 and an activity above 50 CEVU (Cellulose Viscosity Unit). Suitable cellulases are disclosed in U.S. Patent 4,435,307, Barbesgoard et al, J6107834 and WO96/02653 which discloses fungal cellulase produced respectively from *Humicola insolens*, *Trichoderma*, *Thielavia* and *Sporotrichum*. EP 739 982 describes cellulases isolated from novel *Bacillus* species. Suitable cellulases are also disclosed in GB-A-2.075.028; GB-A-2.095.275; DE-OS-2.247.832 and WO95/26398.

Examples of such cellulases are cellulases produced by a strain of *Humicola insolens* (*Humicola grisea* var. *thermoidea*), particularly the *Humicola* strain DSM 1800.

Other suitable cellulases are cellulases originated from *Humicola insolens* having a molecular weight of about 50KDa, an isoelectric point of 5.5 and containing 415 amino acids; and a 43kD endoglucanase derived from *Humicola insolens*, DSM 1800, exhibiting cellulase activity; a preferred endoglucanase component has the amino acid sequence disclosed in PCT Patent Application No. WO 91/17243. Also suitable cellulases are the EGIII cellulases from *Trichoderma longibrachiatum* described in WO94/21801, Genencor, published September 29, 1994. Especially suitable cellulases are the cellulases having color care benefits. Examples of such cellulases are cellulases described in European patent application No. 91202879.2, filed November 6, 1991 (Novo). Carezyme and Celluzyme (Novo Nordisk A/S) are especially useful. See also WO91/17244 and WO91/21801. Other suitable cellulases for fabric care and/or cleaning properties are described in WO96/34092, WO96/17994 and WO95/24471.

Said cellulases are normally incorporated in the detergent composition at levels from 0.0001% to 2% of active enzyme by weight of the detergent composition.

[0044] Other preferred enzymes that can be included in the detergent compositions of the present invention include lipases. Suitable lipase enzymes for detergent usage include those produced by microorganisms of the *Pseudomonas* group, such as *Pseudomonas stutzeri* ATCC 19.154, as disclosed in British Patent 1,372,034. Suitable lipases include those which show a positive immunological cross-reaction with the antibody of the lipase, produced by the microorganism *Pseudomonas fluorescent* IAM 1057. This lipase is available from Amano Pharmaceutical Co. Ltd., Nagoya, Japan, under the trade name Lipase P "Amano," hereinafter referred to as "Amano-P". Other suitable commercial lipases include Amano-CES, lipases ex *Chromobacter viscosum*, e.g. *Chromobacter viscosum* var. *lipolyticum* NRRLB 3673 from Toyo Jozo Co., Tagata, Japan; *Chromobacter viscosum* lipases from U.S. Biochemical Corp., U.S.A. and Disoynt Co., The Netherlands, and lipases ex *Pseudomonas gladioli*. Especially suitable lipases are lipases such as M1 Lipase<sup>R</sup> and Lipomax<sup>R</sup> (Gist-Brocades) and Lipolase<sup>R</sup> and Lipolase Ultra<sup>R</sup> (Novo) which have found to be very effective when used in combination with the compositions of the present invention. Also suitable are the lipolytic enzymes described in EP 258 068, WO 92/05249 and WO 95/22615 by Novo Nordisk and in WO 94/03578, WO 95/35381 and WO 96/00292 by Unilever.

Also suitable are cutinases [EC 3.1.1.50] which can be considered as a special kind of lipase, namely lipases which do not require interfacial activation. Addition of cutinases to detergent compositions have been described in e.g. WO-A-88/09367 (Genencor); WO 90/09446 (Plant Genetic System) and WO 94/14963 and WO 94/14964 (Unilever).



The lipases and/or cutinases are normally incorporated in the detergent composition at levels from 0.0001% to 2% of active enzyme by weight of the detergent composition.

[0045] Suitable proteases are the subtilisins which are obtained from particular strains of *B. subtilis* and *B. licheniformis* (subtilisin BPN and BPN'). One suitable protease is obtained from a strain of *Bacillus*, having maximum activity throughout the pH range of 8-12, developed and sold as ESPERASE® by Novo Industries A/S of Denmark, hereinafter "Novo". The preparation of this enzyme and analogous enzymes is described in GB 1,243,784 to Novo. Other suitable proteases include ALCALASE®, DURAZYM® and SAVINASE® from Novo and MAXATASE®, MAXACAL®, PROPERASE® and MAXAPEM® (protein engineered Maxacal) from Gist-Brocades. Proteolytic enzymes also encompass modified bacterial serine proteases, such as those described in European Patent Application EP 251 446 filed April 28, 1987 (particularly pages 17, 24 and 98), and which is called herein "Protease B", and in European Patent Application 199,404, Venegas, published October 29, 1986, which refers to a modified bacterial serine proteolytic enzyme which is called "Protease A" herein. Suitable is what is called herein "Protease C", which is a variant of an alkaline serine protease from *Bacillus* in which lysine replaced arginine at position 27, tyrosine replaced valine at position 104, serine replaced asparagine at position 123, and alanine replaced threonine at position 274. Protease C is described in EP 90915958:4, corresponding to WO 91/06637, Published May 16, 1991. Genetically modified variants, particularly of Protease C, are also included herein.

A preferred protease referred to as "Protease D" is a carbonyl hydrolase variant having an amino acid sequence not found in nature, which is derived from a precursor carbonyl hydrolase by substituting a different amino acid for a plurality of amino acid residues at a position in said carbonyl hydrolase equivalent to those selected from the group consisting of +99, +101, +103, +104, +107, +123, +27, +105, +109, +126, +128, +135, +156, +166, +195, +197, +204, +206, +210, +216, +217, +218, +222, +260, +265, and/or +274 according to the numbering of *Bacillus amyloliquefaciens* subtilisin, as described in WO95/10591 and in the patent application of C. Ghosh, et al, "Bleaching Compositions Comprising Protease Enzymes" WO95/10592, filed October 13, 1994.

Also suitable for the present invention are proteases described in patent applications EP 251 446 and WO 91/06637, protease BLAP® described in WO91/02792 and their variants described in WO 95/23221. See also a high pH protease from *Bacillus* sp. NCIMB 40338 described in WO 93/18140 A to Novo. Enzymatic detergents comprising protease, one or more other enzymes, and a reversible protease inhibitor are described in WO 92/03529 A to Novo. When desired, a protease having decreased adsorption and increased hydrolysis is available as described in WO 95/07791 to Procter & Gamble. A recombinant trypsin-like protease for detergents suitable herein is described in WO 94/25583 to Novo. Other suitable proteases are described in EP 516 200 by Unilever.

The proteolytic enzymes are incorporated in the detergent compositions of the present invention a level of from 0.0001% to 2%, preferably from 0.001% to 0.2%, more preferably from 0.005% to 0.1% pure enzyme by weight of the composition.

[0046] Amylases ( $\alpha$  and/or  $\beta$ ) can be included for removal of carbohydrate-based stains. WO94102597, Novo Nordisk A/S published February 03, 1994, describes cleaning compositions which incorporate mutant amylases. See also WO95/10603, Novo Nordisk A/S, published April 20, 1995. Other amylases known for use in cleaning compositions include both  $\alpha$ - and  $\beta$ -amylases.  $\alpha$ -Amylases are known in the art and include those disclosed in US Pat. no. 5,003,257; EP 252,666; WO/91/00353; FR 2,676,456; EP 285,123; EP 525,610; EP 368,341; and British Patent specification no. 1,296,839 (Novo). Other suitable amylases are stability-enhanced amylases described in WO94/18314, published August 18, 1994 and WO96/05295, Genencor, published February 22, 1996 and amylase variants having additional modification in the immediate parent available from Novo Nordisk A/S, disclosed in WO 95/10603, published April 95. Also suitable are amylases described in EP 277 216, WO95/26397 and WO96/23873 (all by Novo Nordisk).

Examples of commercial  $\alpha$ -amylases products are Purafect Ox Am® from Genencor and Termamyl®, Ban®, Fungamyl® and Duramyl®, all available from Novo Nordisk A/S Denmark. WO95/26397 describes other suitable amylases:  $\alpha$ -amylases characterised by having a specific activity at least 25% higher than the specific activity of Termamyl® at a temperature range of 25°C to 55°C and at a pH value in the range of 8 to 10, measured by the Phadebas®  $\alpha$ -amylase activity assay. Suitable are variants of the above enzymes, described in WO96/23873 (Novo Nordisk). Other amylolytic enzymes with improved properties with respect to the activity level and the combination of thermostability and a higher activity level are described in WO95/35382.

The amylolytic enzymes are incorporated in the detergent compositions of the present invention a level of from 0.0001% to 2%, preferably from 0.00018% to 0.06%, more preferably from 0.00024% to 0.048% pure enzyme by weight of the composition.

[0047] The above-mentioned enzymes may be of any suitable origin, such as vegetable, animal, bacterial, fungal and yeast origin. Origin can further be mesophilic or extremophilic (psychrophilic, psychrotrophic, thermophilic, barophilic, alkalophilic, acidophilic, halophilic, etc.). Purified or non-purified forms of these enzymes may be used. Also included by definition, are mutants of native enzymes. Mutants can be obtained e.g. by protein and/or genetic engineering, chemical and/or physical modifications of native enzymes. Common practice as well is the expression of the enzyme via host organisms in which the genetic material responsible for the production of the enzyme has been cloned.

**[0048]** Said enzymes are normally incorporated in the detergent composition at levels from 0.0001% to 2% of active enzyme by weight of the detergent composition. The enzymes can be added as separate single ingredients (prills, granulates, stabilized liquids, etc... containing one enzyme ) or as mixtures of two or more enzymes (e.g. cogranulates).

**[0049]** Other suitable detergent ingredients that can be added are enzyme oxidation scavengers which are described in Copending European Patent application 92870018.6 filed on January 31, 1992. Examples of such enzyme oxidation scavengers are ethoxylated tetraethylene polyamines.

**[0050]** A range of enzyme materials and means for their incorporation into synthetic detergent compositions is also disclosed in WO 9307263 A and WO 9307260 A to Genencor International, WO 8908694 A to Novo, and U.S. 3,553,139, January 5, 1971 to McCarty et al. Enzymes are further disclosed in U.S. 4,101,457, Place et al, July 18, 1978, and in U.S. 4,507,219, Hughes, March 26, 1985. Enzyme materials useful for liquid detergent formulations, and their incorporation into such formulations, are disclosed in U.S. 4,261,868, Hora et al, April 14, 1981. Enzymes for use in detergents can be stabilised by various techniques. Enzyme stabilisation techniques are disclosed and exemplified in U.S. 3,600,319, August 17, 1971, Gedge et al, EP 199,405 and EP 200,586, October 29, 1986, Venegas. Enzyme stabilisation systems are also described, for example, in U.S. 3,519,570. A useful *Bacillus*, sp. AC13 giving proteases, xylanases and cellulases, is described in WO 9401532 A to Novo.


### ***Bleaching agent***

**[0051]** Preferred additional optional detergent ingredients that can be included in the cleaning compositions of the present invention include conventional activated-, other enzymatic- and/or metallo catalyst- based bleach systems. It has been found that the combination of said specific oxygenase with another bleach system provides improved cleaning of polyphenol / heterocyclic coloured and everyday body stains and/or soils and when formulated as laundry composition, it achieves improved fabric realistic items cleaning and whitening while providing colour safety.

**[0052]** The bleaching agent component for use herein can be any of the bleaching agents useful for cleaning compositions including oxygen bleaches as well as others known in the art. The bleaching agent suitable for the present invention can be an activated or non-activated bleaching agent.

**[0053]** Bleaching agents are such as hydrogen peroxide, PB1, PB4 and percarbonate with a particle size of 400-800 microns. These bleaching agent components can include one or more oxygen bleaching agents and, depending upon the bleaching agent chosen, one or more bleach activators. When present oxygen bleaching compounds will typically be present at levels of from about 1% to about 25%.

**[0054]** The hydrogen peroxide releasing agents can be used in combination with bleach activators such as tetraacetythylenediamine (TAED), nonanoyloxybenzene-sulfonate (NOBS, described in US 4,412,934), 3,5,-trimethylhexanoyloxybenzenesulfonate (ISONOBS, described in EP 120,591) or pentaacetylglucose (PAG) or Phenolsulfonate ester of N-nonanoyl-6-aminocaproic acid (NACA-OBS, described in WO94/28106), which are perhydrolyzed to form a peracid as the active bleaching species, leading to improved bleaching effect. Also suitable activators are acylated citrate esters such as disclosed in Copending European Patent Application EP 624 154 and unsymmetrical acyclic imide bleach activator of the following formula as disclosed in the Procter & Gamble co-pending patent application:



WO 98/04664

wherein  $\text{R}_1$  is a  $\text{C}_7$ - $\text{C}_{13}$  linear or branched chain saturated or unsaturated alkyl group,  $\text{R}_2$  is a  $\text{C}_1$ - $\text{C}_8$ , linear or branched chain saturated or unsaturated alkyl group and  $\text{R}_3$  is a  $\text{C}_1$ - $\text{C}_4$  linear or branched chain saturated or unsaturated alkyl group.

**[0055]** One category of oxygen bleaching agent that can be used encompasses percarboxylic acid bleaching agents and salts thereof. Suitable examples of this class of agents include magnesium monoperoxyphthalate hexahydrate, the magnesium salt of meta-chloro perbenzoic acid, 4-nonylamino-4-oxoperoxybutyric acid and diperoxydodecanedioic acid. Such bleaching agents are disclosed in U.S. Patent 4,483,781, U.S. Patent Application 740,446, European Patent Application 0,133,354 and U.S. Patent 4,412,934. Highly preferred bleaching agents also include 6-nonytamino-6-oxoperoxyacaproic acid as described in U.S. Patent 4,634,551.

Another category of bleaching agents that can be used encompasses the halogen bleaching agents. Examples of hypohalite bleaching agents, for example, include trichloro isocyanuric acid and the sodium and potassium dichloroisocyanurates and N-chloro and N-bromo alkane sulphonamides. Such materials are normally added at 0.5-10% by weight

of the finished product, preferably 1-5% by weight.

**[0056]** Useful bleaching agents, including peroxyacids and bleaching systems comprising bleach activators and per-oxygen bleaching compounds for use in detergent compositions according to the invention are described in our copending applications WO95/10592, WO97/00937, WO95/27772, WO95/27773, WO95/27774 and WO95/27775.

**[0057]** The hydrogen peroxide may also be present by adding an enzymatic system (i.e. an enzyme and a substrate therefore) which is capable of generating hydrogen peroxide at the beginning or during the washing and/or rinsing process. Such enzymatic systems are disclosed in EP Patent Application EP 537 381 filed October 9, 1991.

Peroxidase enzymes are used in combination with oxygen, hydrogen peroxide sources, e.g. percarbonate, perborate, persulfate, hydrogen peroxide, etc and a bleach enhancer. They are used for "solution bleaching", i.e. to prevent transfer of dyes or pigments removed from substrates during wash operations to other substrates in the wash solution. Peroxidase enzymes are known in the art, and include, for example, horseradish peroxidase, ligninase and hatoperoxidase such as chloro- and bromo-peroxidase. Peroxidase-containing detergent compositions are disclosed, for example, in PCT International Application WO 89/099813, WO89/09813 and in European Patent application EP 540 784, filed on November 6, 1991. Also suitable is the laccase enzyme.

Enhancers are generally comprised at a level of from 0.1% to 5% by weight of total composition. Preferred enhancers are substituted phenothiazine and phenoxazine 10-Phenothiazinepropionic acid (PPT), 10-ethylphenothiazine-4-carboxylic acid (EPC), 10-phenoxazinepropionic acid (POP) and 10-methylphenoxazine (described in WO 94/12621) and substituted syringates (C3-C5 substituted alkyl syringates) and phenols. Sodium percarbonate or perborate are preferred sources of hydrogen peroxide.

Said peroxidases are normally incorporated in the detergent composition at levels from 0.0001 % to 2% of active enzyme by weight of the detergent composition.

**[0058]** Metal-containing catalysts for use in bleach compositions, include cobalt-containing catalysts such as Pentaamine acetate cobalt(III) salts and manganese-containing catalysts such as those described in EPA 549 271; EPA 549 272; EPA 458 397; US 5,246,621; EPA 458 398; EPA 458 397; US 5,194,416 and US 5,114,611. Bleaching composition comprising a peroxy compound, a manganese-containing bleach catalyst and a chelating agent is described in the patent application EP 718 398.

Preferred metal-containing catalyst for the purpose of the present invention is a transition metal complex of a macropolycyclic rigid ligand. The phrase "macropolycyclic rigid ligand" is sometimes abbreviated as "MRL" in discussion below. The amount used is a catalytically effective amount, suitably about 1 ppb or more, for example up to about 99.9%, more typically about 0.001 ppm or more, preferably from about 0.05 ppm to about 500 ppm (wherein "ppb" denotes parts per billion by weight and "ppm" denotes parts per million by weight).

Suitable transition metals e.g., Mn are illustrated hereinafter. "Macropolycyclic" means a MRL is both a macrocycle and is polycyclic. "Polycyclic" means at least bicyclic. The term "rigid" as used herein includes "having a superstructure" and "cross-bridged". "Rigid" has been defined as the constrained converse of flexibility: see D.H. Busch., Chemical Reviews, (1993), 93, 847-860, incorporated by reference. More particularly, "rigid" as used herein means that the MRL must be determinably more rigid than a macrocycle ("parent macrocycle") which is otherwise identical (having the same ring size and type and number of atoms in the main ring) but lacking a superstructure (especially linking moieties or, preferably cross-bridging moieties) found in the MRL's. In determining the comparative rigidity of macrocycles with and without superstructures, the practitioner will use the free form (not the metal-bound form) of the macrocycles. Rigidity is well-known to be useful in comparing macrocycles; suitable tools for determining, measuring or comparing rigidity include computational methods (see, for example, Zimmer, Chemical Reviews, (1995), 95(38), 2629-2648 or Hancock et al., Inorganica Chimica Acta, (1989), 164, 73-84. A determination of whether one macrocycle is more rigid than another can be often made by simply making a molecular model, thus it is not in general essential to know configurational energies in absolute terms or to precisely compute them. Excellent comparative determinations of rigidity of one macrocycle vs. another can be made using inexpensive personal computer-based computational tools, such as ALCHEMY III, commercially available from Tripos Associates. Tripos also has available more expensive software permitting not only comparative, but absolute determinations; alternately, SHAPES can be used (see Zimmer cited supra). One observation which is significant in the context of the present invention is that there is an optimum for the present purposes when the parent macrocycle is distinctly flexible as compared to the cross-bridged form. Thus, unexpectedly, it is preferred to use parent macrocycles containing at least four donor atoms, such as cyclam derivatives, and to cross-bridge them, rather than to start with a more rigid parent macrocycle. Another observation is that cross-bridged macrocycles are significantly preferred over macrocycles which are bridged in other manners.

Preferred MRL's herein are a special type of ultra-rigid ligand which is cross-bridged. A "cross-bridge" is nonlimitingly illustrated in 1.11 hereinbelow. In 1.11, the cross-bridge is a  $-CH_2CH_2-$  moiety. It bridges N<sup>1</sup> and N<sup>8</sup> in the illustrative structure. By comparison, a "same-side" bridge, for example if one were to be introduced across N<sup>1</sup> and N<sup>12</sup> in 1.11, would not be sufficient to constitute a "cross-bridge" and accordingly would not be preferred.

Suitable metals in the rigid ligand complexes include Mn(II), Mn(III), Mn(IV), Mn(V), Fe(II), Fe(III), Fe(IV), Co(I), Co(II), Co(III), Ni(I), Ni(II), Ni(III), Cu(I), Cu(II), Cu(III), Cr(II), Cr(III), Cr(IV), Cr(V), Cr(VI), V(III), V(IV), V(V), Mo(IV), Mo(V), Mo

(VI), W(IV), W(V), W(VI), Pd(II), Ru(II), Ru(III), and Ru(IV). Preferred transition-metals in the instant transition-metal bleach catalyst include manganese, iron and chromium. Preferred oxidation states include the (II) and (III) oxidation states. Manganese(II) in both the low-spin configuration and high spin complexes are included. It is to be noted that complexes such as low-spin Mn(II) complexes are rather rare in all of coordination chemistry. The designation (II) or (III) denotes a coordinated transition metal having the requisite oxidation state; the coordinated metal atom is not a free ion or one having only water as a ligand.

In general, as used herein, a "ligand" is any moiety capable of direct covalent bonding to a metal ion. Ligands can be charged or neutral and may range widely, including simple monovalent donors, such as chloride, or simple amines which form a single coordinate bond and a single point of attachment to a metal; to oxygen or ethylene, which can form a three-membered ring with a metal and thus can be said to have two potential points of attachment, to larger moieties such as ethylenediamine or aza macrocycles, which form up to the maximum number of single bonds to one or more metals that are allowed by the available sites on the metal and the number of lone pairs or alternate bonding sites of the free ligand. Numerous ligands can form bonds other than simple donor bonds, and can have multiple points of attachment.

Ligands useful herein can fall into several groups: the MRL, preferably a cross-bridged macropolycycle (preferably there will be one MRL in a useful transition-metal complex, but more, for example two, can be present, but not in preferred mononuclear transition-metal complexes); other, optional ligands, which in general are different from the MRL (generally there will be from 0 to 4, preferably from 1 to 3 such ligands); and ligands associated transiently with the metal as part of the catalytic cycle, these latter typically being related to water, hydroxide, oxygen or peroxides. Ligands of the third group are not essential for defining the metal bleach catalyst, which is a stable, isolable chemical compound that can be fully characterized. Ligands which bind to metals through donor atoms each having at least a single lone pair of electrons available for donation to a metal have a donor capability, or potential denticity, at least equal to the number of donor atoms. In general, that donor capability may be fully or only partially exercised.

Generally, the MRL's herein can be viewed as the result of imposing additional structural rigidity on specifically selected "parent macrocycles".


More generally, the MRL's (and the corresponding transition-metal catalysts) herein suitably comprise:

- (a) at least one macrocycle main ring comprising four or more heteroatoms; and
- (b) a covalently connected non-metal superstructure capable of increasing the rigidity of the macrocycle, preferably selected from


- (i) a bridging superstructure, such as a linking moiety;
- (ii) a cross-bridging superstructure, such as a cross-bridging linking moiety; and
- (iii) combinations thereof.

The term "superstructure" is used herein as defined in the literature by Busch et al., see, for example, articles by Busch in "Chemical Reviews".

Preferred superstructures herein not only enhance the rigidity of the parent macrocycle, but also favor folding of the macrocycle so that it co-ordinates to a metal in a cleft. Suitable superstructures can be remarkably simple, for example a linking moiety such as any of those illustrated in 1.9 and 1.10 below, can be used.



wherein n is an integer, for example from 2 to 8, preferably less than 6, typically 2 to 4, or



1.10


wherein m and n are integers from about 1 to 8, more preferably from 1 to 3; Z is N or CH; and T is a compatible substituent, for example H, alkyl, trialkylammonium, halogen, nitro, sulfonate, or the like. The aromatic ring in 1.10 can be replaced by a saturated ring, in which the atom in Z connecting into the ring can contain N, O, S or C.

Without intending to be limited by theory, it is believed that the preorganization built into the MRL's herein that leads to extra kinetic and/or thermodynamic stability of their metal complexes arises from either or both of topological constraints and enhanced rigidity (loss of flexibility) compared to the free parent macrocycle which has no superstructure. The MRL's as defined herein and their preferred cross-bridged sub-family, which can be said to be "ultra-rigid", combine two sources of fixed preorganization. In preferred MRL's herein, the linking moieties and parent macrocycle rings are combined to form ligands which have a significant extent of "fold", typically greater than in many known superstured ligands in which a superstructure is attached to a largely planar, often unsaturated macrocycle. See, for example: D.H. Busch, *Chemical Reviews*, (1993), 93, 847 - 880. Further, the preferred MRL's herein have a number of particular properties, including (1) they are characterized by very high proton affinities, as in so-called "proton sponges"; (2) they tend to react slowly with multivalent transition metals, which when combined with (1) above, renders synthesis of their complexes with certain hydrolyzable metal ions difficult in hydroxylic solvents; (3) when they are coordinated to transition metal atoms as identified herein, the MRL's result in complexes that have exceptional kinetic stability such that the metal ions only dissociate extremely slowly under conditions that would destroy complexes with ordinary ligands; and (4) these complexes have exceptional thermodynamic stability; however, the unusual kinetics of MRL dissociation from the transition metal may defeat conventional equilibrium measurements that might quantitate this property.

In one aspect of the present invention, the MRL's include those comprising:

- (i) an organic macrocycle ring containing four or more donor atoms (preferably at least 3, more preferably at least 4, of these donor atoms are N) separated from each other by covalent linkages of at least one, preferably 2 or 3, non-donor atoms, two to five (preferably three to four, more preferably four) of these donor atoms being coordinated to the same transition metal in the complex; and
- (ii) a linking moiety, preferably a cross-bridging chain, which covalently connects at least 2 (preferably non-adjacent) donor atoms of the organic macrocycle ring, said covalently connected (preferably non-adjacent) donor atoms being bridgehead donor atoms which are coordinated to the same transition metal in the complex, and wherein said linking moiety (preferably a cross-bridged chain) comprises from 2 to about 10 atoms (preferably the cross-bridged chain is selected from 2, 3 or 4 non-donor atoms, and 4-6 non-donor atoms with a further donor atom).

Suitable MRL's are further nonlimitingly illustrated by the following compound:



1.11

**[0059]** This is a MRL in accordance with the invention which is a highly preferred, cross-bridged, methyl-substituted (all nitrogen atoms tertiary) derivative of cyclam. Formally, this ligand is named 5,12-dimethyl-1,5,8,12-tetraazabicyclo

[6.6.2]hexadecane using the extended von Baeyer system. See "A Guide to IUPAC Nomenclature of Organic Compounds: Recommendations 1993", R. Panico, W.H. Powell and J-C Richer (Eds.), Blackwell Scientific Publications, Boston, 1993; see especially section R-2.4.2.1. According to conventional terminology, N1 and N8 are "bridgehead atoms"; as defined herein, more particularly "bridgehead donor atoms" since they have lone pairs capable of donation to a metal. N1 is connected to two non-bridgehead donor atoms, N5 and N12, by distinct saturated carbon chains 2,3,4 and 14,13 and to bridgehead donor atom N8 by a "linking moiety" a,b which here is a saturated carbon chain of two carbon atoms. N8 is connected to two non-bridgehead donor atoms, N5 and N12, by distinct chains 6,7 and 9,10,11. Chain a,b is a "linking moiety" as defined herein, and is of the special, preferred type referred to as a "cross-bridging" moiety. The "macrocyclic ring" of the ligand supra, or "main ring" (IUPAC), includes all four donor atoms and chains 2,3,4; 6,7; 9,10,11 and 13,14 but not a,b. This ligand is conventionally bicyclic. The short bridge or "linking moiety" a,b is a "cross-bridge" as defined herein, with a,b bisecting the macrocyclic ring.

The MRL's herein are of course not limited to being synthesized from any preformed macrocycle plus preformed "rigidizing" or "conformation-modifying" element: rather, a wide variety of synthetic means, such as template syntheses, are useful. See for example Busch et al., reviewed in "Heterocyclic compounds: Aza-crown macrocycles", J.S. Bradshaw et. al.

Transition-metal bleach catalysts useful in the invention compositions can in general include known compounds where they conform with the definition herein, as well as, more preferably, any of a large number of novel compounds expressly designed for the present laundry or cleaning uses, and non-limitingly illustrated by any of the following:

- Dichloro-5,12-dimethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-4,10-dimethyl-1,4,7,10-tetraazabicyclo[5.5.2]tetradecane Manganese(II)
- Diaquo-5,12-dimethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II) Hexafluorophosphate
- Aquo-hydroxy-5,12-dimethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(III) Hexafluorophosphate
- Diaquo-4,10-dimethyl-1,4,7,10-tetraazabicyclo[5.5.2]tetradecane Manganese(II) Hexafluorophosphate
- Diaquo-5,12-dimethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II) Tetrafluoroborate
- Diaquo-4,10-dimethyl-1,4,7,10-tetraazabicyclo[5.5.2]tetradecane Manganese(II) Tetrafluoroborate
- Dichloro-5,12-dimethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(III) Hexafluorophosphate
- Dichloro-5,12-di-n-butyl-1,5,8,12-tetraaza- bicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-5,12-dibenzyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-5-n-butyl-12-methyl-1, 5,8,12-tetraaza- bicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-5-n-octyl-12-methyl-1,5,8,12-tetraaza- bicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-5-n-butyl-12-methyl-1,5,8,12-tetraaza- bicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-5,12-dimethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Iron(II)
- Dichloro-4,10-dimethyl-1,4,7,10-tetraazabicyclo[5.5.2]tetradecane Iron(II)
- Dichloro-5,12-dimethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Copper(II)
- Dichloro-4,10-dimethyl-1,4,7,10-tetraazabicyclo[5.5.2]tetradecane Copper(II)
- Dichloro-5,12-dimethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Cobalt(II)
- Dichloro-4,10-dimethyl-1,4,7,10-tetraazabicyclo[5.5.2]tetradecane Cobalt(II)
- Dichloro 5,12-dimethyl--4-phenyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-4,10-dimethyl-3-phenyl-1,4,7,10-tetraazabicyclo[5.5.2]tetradecane Manganese(II)
- Dichloro-5,12-dimethyl-4,9-diphenyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-4,10-dimethyl-3,8-diphenyl-1,4,7,10-tetraazabicyclo[5.5.2]tetradecane Manganese(II)
- Dichloro-5,12-dimethyl-2,11 -diphenyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-4,10-dimethyl-4,9-diphenyl-1,4,7,10-tetraazabicyclo[5.5.2]tetradecane Manganese(II)
- Dichloro-2,4,5,9,11,12-hexamethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-2,3,5,9,10,12-hexamethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-2,2,4,5,9,11,12-octamethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-2,2,4,5,9,11,11,12-octamethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-3,3,5,10,10,12-hexamethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-3,5,10,12-tetramethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-3-butyl-5,10,12-trimethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)
- Dichloro-1,4,7,10-tetraazabicyclo[5.5.2]tetradecane Manganese(II)
- Dichloro-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Iron(II)
- Dichloro-1,4,7,10-tetraazabicyclo[5.5.2]tetradecane Iron(II)
- Aquo-chloro-2-(2-hydroxyphenyl)-5,12-dimethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)
- Aquo-chloro-10-(2-hydroxybenzyl)-4,10-dimethyl-1,4,7,10-tetraazabicyclo[5.5.2]tetradecane Manganese(II)
- Chloro-2-(2-hydroxybenzyl)-5-methyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)

- Chloro-10-(2-hydroxybenzyl)-4-methyl-1,4,7,10-tetraazabicyclo[5.5.2]tetradecane Manganese(II)  
 Chloro-5-methyl-12-(2-picoyl)-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II) Chloride  
 Chloro-4-methyl-10-(2-picoyl)-1,4,7,10-tetraazabicyclo[5.5.2]tetradecane Manganese(II) Chloride  
 Dichloro-5-(2-sulfato)dodecyl-12-methyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(III)  
 5 Aquo-Chloro-5-(2-sulfato)dodecyl-12-methyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)  
 Aquo-Chloro-5-(3-sulfonopropyl)-12-methyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)  
 Dichloro-5-(Trimethylammoniopropyl)dodecyl-12-methyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese  
 (III) Chloride  
 Dichloro-5,12-dimethyl-1,4,7,10,13-pentaazabicyclo[8.5.2]heptadecane Manganese(II)  
 10 Dichloro-14,20-dimethyl-1,10,14,20-tetraazabicyclo[8.6.6]docosa-3(8),4,6-triene Manganese(II)  
 Dichloro-4,11-dimethyl-1,4,7,11-tetraazabicyclo[6.5.2]pentadecane Manganese(II)  
 Dichloro-5,12-dimethyl-1,5,8,12-tetraazabicyclo[7.6.2]heptadecane Manganese(II)  
 Dichloro-5,13-dimethyl-1,5,9,13-tetraazabicyclo[7.7.2]heptadecane Manganese(II)  
 Dichloro-3,10-bis(butylcarboxy)-5,12-dimethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)  
 15 Diaquo-3,10-dicarboxy-5,12-dimethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane Manganese(II)  
 Chloro-20-methyl-1,9,20,24,25-pentaaza-tetracyclo[7.7.7.1<sup>3,7</sup>.1<sup>11,15</sup>]pentacosa-3,5,7(24),11,13,15(25)-hexaene  
 Manganese(II) Hexafluorophosphate  
 Trifluoromethanesulfono-20-methyl-1,9,20,24,25-pentaazatetracyclo[7.7.7.1<sup>3,7</sup>.1<sup>11,15</sup>]pentacosa-3,5,7(24),  
 11,13,15(25)-hexaene Manganese(II) Trifluoromethanesulfonate  
 20 Trifluoromethanesulfono-20-methyl-1,9,20,24,25-pentaazatetracyclo[7.7.7.1<sup>3,7</sup>.1<sup>11,15</sup>]pentacosa-3,5,7(24),  
 11,13,15(25)-hexaene Iron(II) Trifluoromethanesulfonate  
 Chloro-5,12,17-trimethyl-1,5,8,12,17-pentaazabicyclo[6.6.5]nonadecane Manganese(II) Hexafluorophosphate  
 Chloro-4,10,15-trimethyl-1,4,7,10,15-pentaazabicyclo[5.5.5]heptadecane Manganese(II) Hexafluorophosphate  
 Chloro-5,12,17-trimethyl-1,5,8,12,17-pentaazabicyclo[6.6.5]nonadecane Manganese(II) Chloride  
 25 Chloro-4,10,15-trimethyl-1,4,7,10,15-pentaazabicyclo[5.5.5]heptadecane Manganese(II) Chloride


**[0060]** The practitioner may further benefit if certain terms receive additional definition and illustration. As used herein, "macrocyclic rings" are covalently connected rings formed from four or more donor atoms (i.e., heteroatoms such as nitrogen or oxygen) with carbon chains connecting them, and any macrocycle ring as defined herein must contain a total  
 30 of at least ten, preferably at least twelve, atoms in the macrocycle ring. A MRL herein may contain more than one ring of any sort per ligand, but at least one macrocycle ring must be identifiable. Moreover, in the preferred embodiments, no two hetero-atoms are directly connected. Preferred transition-metal bleach catalysts are those wherein the MRL comprises an organic macrocycle ring (main ring) containing at least 10-20 atoms, preferably 12-18 atoms, more preferably from about 12 to about 20 atoms, most preferably 12 to 16 atoms.

**[0061]** "Donor atoms" herein are heteroatoms such as nitrogen, oxygen, phosphorus or sulfur, which when incorporated into a ligand still have at least one lone pair of electrons available for forming a donor-acceptor bond with a metal. Preferred transition-metal bleach catalysts are those wherein the donor atoms in the organic macrocycle ring of the cross-bridged MRL are selected from the group consisting of N, O, S, and P, preferably N and O, and most preferably all N. Also preferred are cross-bridged MRL's comprising 4 or 5 donor atoms, all of which are coordinated to the same  
 40 transition metal. Most preferred transition-metal bleach catalysts are those wherein the cross-bridged MRL comprises 4 nitrogen donor atoms all coordinated to the same transition metal, and those wherein the cross-bridged MRL comprises 5 nitrogen atoms all coordinated to the same transition metal.

**[0062]** "Non-donor atoms" of the MRL herein are most commonly carbon, though a number of atom types can be included, especially in optional exocyclic substituents (such as "pendant" moieties, illustrated hereinafter) of the macrocycles, which are neither donor atoms for purposes essential to form the metal catalysts, nor are they carbon. Thus, in the broadest sense, the term "non-donor atoms" can refer to any atom not essential to forming donor bonds with the metal of the catalyst. Examples of such atoms could include heteroatoms such as sulfur as incorporated in a non-coordinatable sulfonate group, phosphorus as incorporated into a phosphonium salt moiety, phosphorus as incorporated into a P(V) oxide, a non-transition metal, or the like. In certain preferred embodiments, all non-donor atoms are carbon.  
 50 Transition metal complexes of MRL's can be prepared in any convenient manner. Two such preparations are illustrated as follows:

#### Synthesis of [Mn(Bcyclam)Cl<sub>2</sub>]

**[0063]**



#### (a) Method I.

**[0064]** "Bcyclam" (5,12-dimethyl-1,5,8,12-tetraaza-bicyclo[6.6.2]hexadecane) is prepared by a synthesis method described by G.R. Weisman, et al., *J.Amer.Chem.Soc.*, (1990), 112, 8604. Bcyclam (1.00 g., 3.93 mmol) is dissolved in dry  $\text{CH}_3\text{CN}$  (35 mL, distilled from  $\text{CaH}_2$ ). The solution is then evacuated at 15 mm until the  $\text{CH}_3\text{CN}$  begins to boil. The flask is then brought to atmospheric pressure with Ar. This degassing procedure is repeated 4 times.  $\text{Mn}(\text{pyridine})_2\text{Cl}_2$  (1.12 g., 3.93 mmol), synthesized according to the literature procedure of H. T. Witteveen et al., *J. Inorg. Nucl. Chem.*, (1974), 36, 1535, is added under Ar. The cloudy reaction solution slowly begins to darken. After stirring overnight at room temperature, the reaction solution becomes dark brown with suspended fine particulates. The reaction solution is filtered with a  $0.2\mu$  filter. The filtrate is a light tan color. This filtrate is evaporated to dryness using a rotoevaporator. After drying overnight at 0.05 mm at room temperature, 1.35 g. off-white solid product is collected, 90% yield. Elemental Analysis: %Mn, 14.45; %C, 44.22; %H, 7.95; theoretical for  $[\text{Mn}(\text{Bcyclam})\text{Cl}_2]$ ,  $\text{MnC}_{14}\text{H}_{30}\text{N}_4\text{Cl}_2$ , MW = 380.26. Found: %Mn, 14.98; %C, 44.48; %H, 7.86; Ion Spray Mass Spectroscopy shows one major peak at 354 mu corresponding to  $[\text{Mn}(\text{Bcyclam})(\text{formate})]^+$ .

#### (b) Method II.

**[0065]** Freshly distilled Bcyclam (25.00 g., 0.0984 mol), which is prepared by the same method as above, is dissolved in dry  $\text{CH}_3\text{CN}$  (900 mL, distilled from  $\text{CaH}_2$ ). The solution is then evacuated at 15 mm until the  $\text{CH}_3\text{CN}$  begins to boil. The flask is then brought to atmospheric pressure with Ar. This degassing procedure is repeated 4 times.  $\text{MnCl}_2$  (11.25 g., 0.0894 mol) is added under Ar. The cloudy reaction solution immediately darkens. After stirring 4 hrs. under reflux, the reaction solution becomes dark brown with suspended fine particulates. The reaction solution is filtered through a  $0.2\mu$  filter under dry conditions. The filtrate is a light tan color. This filtrate is evaporated to dryness using a rotoevaporator. The resulting tan solid is dried overnight at 0.05 mm at room temperature. The solid is suspended in toluene (100 mL) and heated to reflux. The toluene is decanted off and the procedure is repeated with another 100 mL of toluene. The balance of the toluene is removed using a rotoevaporator. After drying overnight at 0.05 mm at room temperature, 31.75 g. of a light blue solid product is collected, 93.5% yield. Elemental Analysis: %Mn, 14.45; %C, 44.22; %H, 7.95; %N, 14.73; %Cl, 18.65; theoretical for  $[\text{Mn}(\text{Bcyclam})\text{Cl}_2]$ ,  $\text{MnC}_{14}\text{H}_{30}\text{N}_4\text{Cl}_2$ , MW = 380.26. Found: %Mn, 14.69; %C, 44.69; %H, 7.99; %N, 14.78; %Cl, 18.90 (Karl Fischer Water, 0.68%). Ion Spray Mass Spectroscopy shows one major peak at 354 mu corresponding to  $[\text{Mn}(\text{Bcyclam})(\text{formate})]^+$ .

**[0066]** Bleaching agents other than oxygen bleaching agents are also known in the art and can be utilized herein. One type of non-oxygen bleaching agent of particular interest includes photoactivated bleaching agents such as the sulfonated zinc and/or aluminum phthalocyanines. These materials can be deposited upon the substrate during the washing process. Upon irradiation with light, in the presence of oxygen, such as by hanging clothes out to dry in the daylight, the sulfonated zinc phthalocyanine is activated and, consequently, the substrate is bleached. Preferred zinc phthalocyanine and a photoactivated bleaching process are described in U.S. Patent 4,033,718. Typically, detergent compositions will contain about 0.025% to about 1.25%, by weight, of sulfonated zinc phthalocyanine.

#### Surfactant system

**[0067]** The cleaning compositions according to the present invention generally comprise a surfactant system wherein the surfactant can be selected from nonionic and/or anionic and/or cationic and/or ampholytic and/or zwitterionic and/or semi-polar surfactants.

**[0068]** The surfactant is typically present at a level of from 0.1 % to 60% by weight. More preferred levels of incorporation are 1% to 35% by weight, most preferably from 1% to 30% by weight of cleaning compositions in accord with the invention.

**[0069]** The surfactant is preferably formulated to be compatible with enzyme components present in the composition. In liquid or gel compositions the surfactant is most preferably formulated such that it promotes, or at least does not degrade, the stability of any enzyme in these compositions.

**[0070]** Preferred surfactant systems to be used according to the present invention comprise as a surfactant one or




more of the nonionic and/or anionic surfactants described herein.

**[0071]** Polyethylene, polypropylene, and polybutylene oxide condensates of alkyl phenols are suitable for use as the nonionic surfactant of the surfactant systems of the present invention, with the polyethylene oxide condensates being preferred. These compounds include the condensation products of alkyl phenols having an alkyl group containing from about 6 to about 14 carbon atoms, preferably from about 8 to about 14 carbon atoms, in either a straight-chain or branched-chain configuration with the alkylene oxide. In a preferred embodiment, the ethylene oxide is present in an amount equal to from about 2 to about 25 moles, more preferably from about 3 to about 15 moles, of ethylene oxide per mole of alkyl phenol. Commercially available nonionic surfactants of this type include Igepal™ CO-630, marketed by the GAF Corporation; and Triton™ X-45, X-114, X-100 and X-102, all marketed by the Rohm & Haas Company. These surfactants are commonly referred to as alkylphenol alkoxylates (e.g., alkyl phenol ethoxylates).

**[0072]** The condensation products of primary and secondary aliphatic alcohols with from about 1 to about 25 moles of ethylene oxide are suitable for use as the nonionic surfactant of the nonionic surfactant systems of the present invention. The alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from about 8 to about 22 carbon atoms. Preferred are the condensation products of alcohols having an alkyl group containing from about 8 to about 20 carbon atoms, more preferably from about 10 to about 18 carbon atoms, with from about 2 to about 10 moles of ethylene oxide per mole of alcohol. About 2 to about 7 moles of ethylene oxide and most preferably from 2 to 5 moles of ethylene oxide per mole of alcohol are present in said condensation products. Examples of commercially available nonionic surfactants of this type include Tergitol™ 15-S-9 (the condensation product of C<sub>11</sub>-C<sub>15</sub> linear alcohol with 9 moles ethylene oxide), Tergitol™ 24-L-6 NMW (the condensation product of C<sub>12</sub>-C<sub>14</sub> primary alcohol with 6 moles ethylene oxide with a narrow molecular weight distribution), both marketed by Union Carbide Corporation; Neodol™ 45-9 (the condensation product of C<sub>14</sub>-C<sub>15</sub> linear alcohol with 9 moles of ethylene oxide), Neodol™ 23-3 (the condensation product of C<sub>12</sub>-C<sub>13</sub> linear alcohol with 3.0 moles of ethylene oxide), Neodol™ 45-7 (the condensation product of C<sub>14</sub>-C<sub>15</sub> linear alcohol with 7 moles of ethylene oxide), Neodol™ 45-5 (the condensation product of C<sub>14</sub>-C<sub>15</sub> linear alcohol with 5 moles of ethylene oxide) marketed by Shell Chemical Company, Kyro™ EOB (the condensation product of C<sub>13</sub>-C<sub>15</sub> alcohol with 9 moles ethylene oxide), marketed by The Procter & Gamble Company, and Genapol LA O3O or O5O (the condensation product of C<sub>12</sub>-C<sub>14</sub> alcohol with 3 or 5 moles of ethylene oxide) marketed by Hoechst. Preferred range of HLB in these products is from 8-11 and most preferred from 8-10.

**[0073]** Also useful as the nonionic surfactant of the surfactant systems of the present invention are the alkylpolysaccharides disclosed in U.S. Patent 4,565,647, Llenado, issued January 21, 1986, having a hydrophobic group containing from about 6 to about 30 carbon atoms, preferably from about 10 to about 16 carbon atoms and a polysaccharide, e.g. a polyglycoside, hydrophilic group containing from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7 saccharide units. Any reducing saccharide containing 5 or 6 carbon atoms can be used, e.g., glucose, galactose and galactosyl moieties can be substituted for the glucosyl moieties (optionally the hydrophobic group is attached at the 2-, 3-, 4-, etc. positions thus giving a glucose or galactose as opposed to a glucoside or galactoside). The intersaccharide bonds can be, e.g., between the one position of the additional saccharide units and the 2-, 3-, 4-, and/or 6- positions on the preceding saccharide units. The preferred alkylpolyglycosides have the formula



wherein R<sup>2</sup> is selected from the group consisting of alkyl, alkylphenyl, hydroxyalkyl, hydroxyalkylphenyl, and mixtures thereof in which the alkyl groups contain from about 10 to about 18, preferably from about 12 to about 14, carbon atoms; n is 2 or 3, preferably 2; t is from 0 to about 10, preferably 0; and x is from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7. The glycosyl is preferably derived from glucose. To prepare these compounds, the alcohol or alkylpolyethoxy alcohol is formed first and then reacted with glucose, or a source of glucose, to form the glucoside (attachment at the 1-position). The additional glycosyl units can then be attached between their 1-position and the preceding glycosyl units 2-, 3-, 4- and/or 6-position, preferably predominately the 2-position.


**[0074]** The condensation products of ethylene oxide with a hydrophobic base formed by the condensation of propylene oxide with propylene glycol are also suitable for use as the additional nonionic surfactant systems of the present invention. The hydrophobic portion of these compounds will preferably have a molecular weight of from about 1500 to about 1800 and will exhibit water insolubility. The addition of polyoxyethylene moieties to this hydrophobic portion tends to increase the water solubility of the molecule as a whole, and the liquid character of the product is retained up to the point where the polyoxyethylene content is about 50% of the total weight of the condensation product, which corresponds to condensation with up to about 40 moles of ethylene oxide. Examples of compounds of this type include certain of the commercially-available Plurafac™ LF404 and Pluronic™ surfactants, marketed by BASF.

**[0075]** Also suitable for use as the nonionic surfactant of the nonionic surfactant system of the present invention, are the condensation products of ethylene oxide with the product resulting from the reaction of propylene oxide and ethylenediamine. The hydrophobic moiety of these products consists of the reaction product of ethylenediamine and excess propylene oxide, and generally has a molecular weight of from about 2500 to about 3000. This hydrophobic moiety is

condensed with ethylene oxide to the extent that the condensation product contains from about 40% to about 80% by weight of polyoxyethylene and has a molecular weight of from about 5,000 to about 11,000. Examples of this type of nonionic surfactant include certain of the commercially available Tetronic™ compounds, marketed by BASF.


**[0076]** Preferred for use as the nonionic surfactant of the surfactant systems of the present invention are polyethylene oxide condensates of alkyl phenols, condensation products of primary and secondary aliphatic alcohols with from about 1 to about 25 moles of ethylene oxide, alkylpolysaccharides, and mixtures thereof. Most preferred are C<sub>8</sub>-C<sub>14</sub> alkyl phenol ethoxylates having from 3 to 15 ethoxy groups and C<sub>8</sub>-C<sub>18</sub> alcohol ethoxylates (preferably C<sub>10</sub> avg.) having from 2 to 10 ethoxy groups, and mixtures thereof.

**[0077]** Highly preferred nonionic surfactants are polyhydroxy fatty acid amide surfactants of the formula.



wherein R<sup>1</sup> is H, or R<sup>1</sup> is C<sub>1-4</sub> hydrocarbyl, 2-hydroxy ethyl, 2-hydroxy propyl or a mixture thereof, R<sup>2</sup> is C<sub>5-31</sub> hydrocarbyl, and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxylated derivative thereof. Preferably, R<sup>1</sup> is methyl, R<sup>2</sup> is a straight C<sub>11-15</sub> alkyl or C<sub>16-18</sub> alkyl or alkenyl chain such as coconut alkyl or mixtures thereof, and Z is derived from a reducing sugar such as glucose, fructose, maltose, lactose, in a reductive amination reaction.

**[0078]** Suitable anionic surfactants to be used are linear alkyl benzene sulfonate, alkyl ester sulfonate surfactants including linear esters of C<sub>8</sub>-C<sub>20</sub> carboxylic acids (i.e., fatty acids) which are sulfonated with gaseous SO<sub>3</sub> according to "The Journal of the American Oil Chemists Society", 52 (1975), pp. 323-329. Suitable starting materials would include natural fatty substances as derived from tallow, palm oil, etc. The preferred alkyl ester sulfonate surfactant, especially for laundry applications, comprise alkyl ester sulfonate surfactants of the structural formula :



wherein R<sup>3</sup> is a C<sub>8</sub>-C<sub>20</sub> hydrocarbyl, preferably an alkyl, or combination thereof, R<sup>4</sup> is a C<sub>1</sub>-C<sub>6</sub> hydrocarbyl, preferably an alkyl, or combination thereof, and M is a cation which forms a water soluble salt with the alkyl ester sulfonate. Suitable salt-forming cations include metals such as sodium, potassium, and lithium, and substituted or unsubstituted ammonium cations, such as monoethanolamine, diethanolamine, and triethanolamine. Preferably, R<sup>3</sup> is C<sub>10</sub>-C<sub>16</sub> alkyl, and R<sup>4</sup> is methyl, ethyl or isopropyl. Especially preferred are the methyl ester sulfonates wherein R<sup>3</sup> is C<sub>10</sub>-C<sub>16</sub> alkyl.

**[0079]** Other suitable anionic surfactants include the alkyl sulfate surfactants which are water soluble salts or acids of the formula ROSO<sub>3</sub>M wherein R preferably is a C<sub>10</sub>-C<sub>24</sub> hydrocarbyl, preferably an alkyl or hydroxyalkyl having a C<sub>10</sub>-C<sub>20</sub> alkyl component, more preferably a C<sub>12</sub>-C<sub>18</sub> alkyl or hydroxyalkyl, and M is H or a cation, e.g., an alkali metal cation (e.g. sodium, potassium, lithium), or ammonium or substituted ammonium (e.g. methyl-, dimethyl-, and trimethyl ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperdinium cations and quaternary ammonium cations derived from alkylamines such as ethylamine, diethylamine, triethylamine, and mixtures thereof, and the like). Typically, alkyl chains of C<sub>12</sub>-C<sub>16</sub> are preferred for lower wash temperatures (e.g. below about 50°C) and C<sub>16-18</sub> alkyl chains are preferred for higher wash temperatures (e.g. above about 50°C).

**[0080]** Other anionic surfactants useful for deterative purposes can also be included in the cleaning compositions of the present invention. These can include salts (including, for example, sodium, potassium, ammonium, and substituted ammonium salts such as mono-, di- and triethanolamine salts) of soap, C<sub>8</sub>-C<sub>22</sub> primary or secondary alkanesulfonates, C<sub>8</sub>-C<sub>24</sub> olefinsulfonates, sulfonated polycarboxylic acids prepared by sulfonation of the pyrolyzed product of alkaline earth metal citrates, e.g., as described in British patent specification No. 1,082,179, C<sub>8</sub>-C<sub>24</sub> alkylpolyglycol ethersulfates (containing up to 10 moles of ethylene oxide); alkyl glycerol sulfonates, fatty acyl glycerol sulfonates, fatty oleyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, paraffin sulfonates, alkyl phosphates, isethionates such as the acyl

isethionates, N-acyl taurates, alkyl succinamates and sulfosuccinates, monoesters of sulfosuccinates (especially saturated and unsaturated C<sub>12</sub>-C<sub>18</sub> monoesters) and diesters of sulfosuccinates (especially saturated and unsaturated C<sub>6</sub>-C<sub>12</sub> diesters), acyl sarcosinates, sulfates of alkylpolysaccharides such as the sulfates of alkylpolyglucoside (the nonionic nonsulfated compounds being described below), branched primary alkyl sulfates, and alkyl polyethoxy carboxylates such as those of the formula RO(CH<sub>2</sub>CH<sub>2</sub>O)<sub>k</sub>-CH<sub>2</sub>COO-M<sup>+</sup> wherein R is a C<sub>8</sub>-C<sub>22</sub> alkyl, k is an integer from 1 to 10, and M is a soluble salt-forming cation. Resin acids and hydrogenated resin acids are also suitable, such as rosin, hydrogenated rosin, and resin acids and hydrogenated resin acids present in or derived from tall oil.


**[0081]** Further examples are described in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch). A variety of such surfactants are also generally disclosed in U.S. Patent 3,929,678, issued December 30, 1975 to Laughlin, et al. at Column 23, line 58 through Column 29, line 23 (herein incorporated by reference).

When included therein, the laundry detergent compositions of the present invention typically comprise from about 1% to about 40%, preferably from about 3% to about 20% by weight of such anionic surfactants.

**[0082]** Highly preferred anionic surfactants include alkyl alkoxyated sulfate surfactants hereof are water soluble salts or acids of the formula RO(A)<sub>m</sub>SO<sub>3</sub>M wherein R is an unsubstituted C<sub>10</sub>-C<sub>24</sub> alkyl or hydroxyalkyl group having a C<sub>10</sub>-C<sub>24</sub> alkyl component, preferably a C<sub>12</sub>-C<sub>20</sub> alkyl or hydroxyalkyl, more preferably C<sub>12</sub>-C<sub>18</sub> alkyl or hydroxyalkyl, A is an ethoxy or propoxy unit, m is greater than zero, typically between about 0.5 and about 6, more preferably between about 0.5 and about 3, and M is H or a cation which can be, for example, a metal cation (e.g., sodium, potassium, lithium, calcium, magnesium, etc.), ammonium or substituted-ammonium cation. Alkyl ethoxylated sulfates as well as alkyl propoxylated sulfates are contemplated herein. Specific examples of substituted ammonium cations include methyl-, dimethyl, trimethyl-ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperidinium cations and those derived from alkylamines such as ethylamine, diethylamine, triethylamine, mixtures thereof, and the like. Exemplary surfactants are C<sub>12</sub>-C<sub>18</sub> alkyl polyethoxylate (1.0) sulfate (C<sub>12</sub>-C<sub>18</sub>E(1.0)M), C<sub>12</sub>-C<sub>18</sub> alkyl polyethoxylate (2.25) sulfate (C<sub>12</sub>-C<sub>18</sub>E(2.25)M), C<sub>12</sub>-C<sub>18</sub> alkyl polyethoxylate (3.0) sulfate (C<sub>12</sub>-C<sub>18</sub>E(3.0)M), and C<sub>12</sub>-C<sub>18</sub> alkyl polyethoxylate (4.0) sulfate (C<sub>12</sub>-C<sub>18</sub>E(4.0)M), wherein M is conveniently selected from sodium and potassium.


**[0083]** The cleaning compositions of the present invention may also contain cationic, ampholytic, zwitterionic, and semi-polar surfactants, as well as the nonionic and/or anionic surfactants other than those already described herein.

**[0084]** Cationic deterative surfactants suitable for use in the cleaning compositions of the present invention are those having one long-chain hydrocarbyl group. Examples of such cationic surfactants include the ammonium surfactants such as alkyltrimethylammonium halogenides, and those surfactants having the formula:




wherein R<sup>2</sup> is an alkyl or alkyl benzyl group having from about 8 to about 18 carbon atoms in the alkyl chain, each R<sup>3</sup> is selected from the group consisting of -CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH(CH<sub>3</sub>)-, -CH<sub>2</sub>CH(CH<sub>2</sub>OH)-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, and mixtures thereof; each R<sup>4</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl, benzyl ring structures formed by joining the two R<sup>4</sup> groups, -CH<sub>2</sub>CHOH-CHOHCOR<sup>6</sup>CHOHCH<sub>2</sub>OH wherein R<sup>6</sup> is any hexose or hexose polymer having a molecular weight less than about 1000, and hydrogen when y is not 0; R<sup>5</sup> is the same as R<sup>4</sup> or is an alkyl chain wherein the total number of carbon atoms of R<sup>2</sup> plus R<sup>5</sup> is not more than about 18; each y is from 0 to about 10 and the sum of the y values is from 0 to about 15; and X is any compatible anion.

**[0085]** Quaternary ammonium surfactant suitable for the present invention has the formula (I):



Formula I

whereby R1 is a short chainlength alkyl (C<sub>6</sub>-C<sub>10</sub>) or alkylamidoalkyl of the formula (II):



### Formula II


y is 2-4, preferably 3.

whereby R2 is H or a C1-C3 alkyl,

whereby x is 0-4, preferably 0-2, most preferably 0,

whereby R3, R4 and R5 are either the same or different and can be either a short chain alkyl (C1-C3) or alkoxyalkyl of the formula III.

whereby X<sup>-</sup> is a counterion, preferably a halide, e.g. chloride or methylsulfate.



### Formula III


R6 is C<sub>1</sub>-C<sub>4</sub> and z is 1 or 2.

[0086] Preferred quat ammonium surfactants are those as defined in formula I whereby

$R_1$  is  $C_8$ ,  $C_{10}$  or mixtures thereof,  $x=0$ .

$R_3, R_4 = \text{CH}_3$  and  $R_5 = \text{CH}_2\text{CH}_2\text{OH}$ .

**[0087]** Highly preferred cationic surfactants are the water-soluble quaternary ammonium compounds useful in the present composition having the formula :



wherein R<sub>1</sub> is C<sub>8</sub>-C<sub>16</sub> alkyl, each of R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub> is independently C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> hydroxy alkyl, benzyl, and -(C<sub>2</sub>H<sub>4</sub>)<sub>x</sub>-H where x has a value from 2 to 5, and X is an anion. Not more than one of R<sub>2</sub>, R<sub>3</sub> or R<sub>4</sub> should be benzyl.

The preferred alkyl chain length for  $R_1$  is  $C_{12}$ - $C_{15}$  particularly where the alkyl group is a mixture of chain lengths derived from coconut or palm kernel fat or is derived synthetically by olefin build up or OXO alcohols synthesis. Preferred groups for  $R_2R_3$  and  $R_4$  are methyl and hydroxyethyl groups and the anion X may be selected from halide, methosulphate, acetate and phosphate ions.

Examples of suitable quaternary ammonium compounds of formulae (i) for use herein are:

coconut trimethyl ammonium chloride or bromide;

coconut methyl dihydroxyethyl ammonium chloride or bromide:

decyl triethyl ammonium chloride;

decyl dimethyl hydroxyethyl ammonium chloride or bromide:

C<sub>12-15</sub> dimethyl hydroxyethyl ammonium chloride or bromide:


coconut dimethyl hydroxyethyl ammonium chloride or bromide;

myristyl trimethyl ammonium methyl sulphate:

lauryl dimethyl benzyl ammonium chloride or bromide:

lauryl dimethyl (ethenoxy)<sub>4</sub> ammonium chloride or bromide;

choline esters (compounds of formula (i) wherein R<sub>1</sub> is



alkyl and  $R_2R_3R_4$  are methyl).

di-alkyl imidazolines [compounds of formula (i)].

**[0088]** Other cationic surfactants useful herein are also described in U.S. Patent 4,228,044, Cambre, issued October 14, 1980 and in European Patent Application EP 000,224.


**[0089]** Typical cationic fabric softening components include the water-insoluble quaternary-ammonium fabric softening actives or their corresponding amine precursor, the most commonly used having been di-long alkyl chain ammonium chloride or methyl sulfate.

Preferred cationic softeners among these include the following:

- 1) ditallow dimethylammonium chloride (DTDMAC);
- 2) dihydrogenated tallow dimethylammonium chloride;
- 3) dihydrogenated tallow dimethylammonium methylsulfate;
- 4) distearyl dimethylammonium chloride;
- 5) dioleyl dimethylammonium chloride;
- 6) dipalmityl hydroxyethyl methylammonium chloride;
- 7) stearyl benzyl dimethylammonium chloride;
- 8) tallow trimethylammonium chloride;
- 9) hydrogenated tallow trimethylammonium chloride;
- 10)  $C_{12-14}$  alkyl hydroxyethyl dimethylammonium chloride;
- 11)  $C_{12-18}$  alkyl dihydroxyethyl methylammonium chloride;
- 12) di(stearoyloxyethyl) dimethylammonium chloride (DSOEDMAC);
- 13) di(tallow-oxy-ethyl) dimethylammonium chloride;
- 14) ditallow imidazolinium methylsulfate;
- 15) 1-(2-tallowylamidoethyl)-2-tallowyl imidazolinium methylsulfate.

**[0090]** Biodegradable quaternary ammonium compounds have been presented as alternatives to the traditionally used di-long alkyl chain ammonium chlorides and methyl sulfates. Such quaternary ammonium compounds contain long chain alk(en)yl groups interrupted by functional groups such as carboxy groups. Said materials and fabric softening compositions containing them are disclosed in numerous publications such as EP-A-0,040,562, and EP-A-0,239,910.

**[0091]** The quaternary ammonium compounds and amine precursors herein have the formula (I) or (II). below :



wherein Q is selected from -O-C(O)-, -C(O)-O-, -O-C(O)-O-, -NR<sup>4</sup>-C(O)-, -C(O)-NR<sup>4</sup>-;

R<sup>1</sup> is (CH<sub>2</sub>)<sub>n</sub>-Q-T<sup>2</sup> or T<sup>3</sup>;

R<sup>2</sup> is (CH<sub>2</sub>)<sub>m</sub>-Q-T<sup>4</sup> or T<sup>5</sup> or R<sup>3</sup>;

R<sup>3</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl or C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl or H;

R<sup>4</sup> is H or C<sub>1</sub>-C<sub>4</sub> alkyl or C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl;

T<sup>1</sup>, T<sup>2</sup>, T<sup>3</sup>, T<sup>4</sup>, T<sup>5</sup> are independently C<sub>11</sub>-C<sub>22</sub> alkyl or alkenyl;

n and m are integers from 1 to 4; and

X<sup>-</sup> is a softener-compatible anion.

Non-limiting examples of softener-compatible anions include chloride or methyl sulfate.

**[0092]** The alkyl, or alkenyl, chain T<sup>1</sup>, T<sup>2</sup>, T<sup>3</sup>, T<sup>4</sup>, T<sup>5</sup> must contain at least 11 carbon atoms, preferably at least 16 carbon atoms. The chain may be straight or branched.

**[0093]** Tallow is a convenient and inexpensive source of long chain alkyl and alkenyl material. The compounds wherein T<sup>1</sup>, T<sup>2</sup>, T<sup>3</sup>, T<sup>4</sup>, T<sup>5</sup> represents the mixture of long chain materials typical for tallow are particularly preferred. Specific examples of quaternary ammonium compounds suitable for use in the aqueous fabric softening compositions herein include:

- 1) N,N-di(tallowyl-oxy-ethyl)-N,N-dimethyl ammonium chloride;
- 2) N,N-di(tallowyl-oxy-ethyl)-N-methyl, N-(2-hydroxyethyl) ammonium methyl sulfate;
- 3) N,N-di(2-tallowyl-oxy-2-oxo-ethyl)-N,N-dimethyl ammonium chloride;
- 4) N,N-di(2-tallowyl-oxy-ethylcarbonyl-oxy-ethyl)-N,N-dimethyl ammonium chloride;
- 5) N-(2-tallowyl-oxy-2-ethyl)-N-(2-tallowyl-oxy-2-oxo-ethyl)-N,N-dimethyl ammonium chloride;
- 6) N,N,N-tri(tallowyl-oxy-ethyl)-N-methyl ammonium chloride;
- 7) N-(2-tallowyl-oxy-2-oxo-ethyl)-N-(tallowyl-N,N-dimethyl-ammonium chloride; and
- 8) 1,2-ditallowyl-oxy-3-trimethylammoniopropane chloride;

and mixtures of any of the above materials.

**[0094]** When included therein, the cleaning compositions of the present invention typically comprise from 0.2% to about 25%, preferably from about 1% to about 8% by weight of such cationic surfactants.


**[0095]** Ampholytic surfactants are also suitable for use in the cleaning compositions of the present invention. These surfactants can be broadly described as aliphatic derivatives of secondary or tertiary amines, or aliphatic derivatives of heterocyclic secondary and tertiary amines in which the aliphatic radical can be straight- or branched-chain. One of the aliphatic substituents contains at least about 8 carbon atoms, typically from about 8 to about 18 carbon atoms, and at least one contains an anionic water-solubilizing group, e.g. carboxy, sulfonate, sulfate. See U.S. Patent No. 3,929,678 to Laughlin et al., issued December 30, 1975 at column 19, lines 18-35, for examples of ampholytic surfactants.

**[0096]** When included therein, the cleaning compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such ampholytic surfactants.

**[0097]** Zwitterionic surfactants are also suitable for use in cleaning compositions. These surfactants can be broadly described as derivatives of secondary and tertiary amines, derivatives of heterocyclic secondary and tertiary amines, or derivatives of quaternary ammonium, quaternary phosphonium or tertiary sulfonium compounds. See U.S. Patent No. 3,929,678 to Laughlin et al., issued December 30, 1975 at column 19, line 38 through column 22, line 48, for examples of zwitterionic surfactants.

**[0098]** When included therein, the cleaning compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such zwitterionic surfactants.

**[0099]** Semi-polar nonionic surfactants are a special category of nonionic surfactants which include water-soluble amine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; water-soluble phosphine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; and water-soluble sulfoxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and a moiety selected from the group consisting of alkyl and hydroxyalkyl moieties of from about 1 to about 3 carbon atoms. Semi-polar nonionic detergent surfactants include the amine oxide surfactants having the formula



wherein R<sup>3</sup> is an alkyl, hydroxyalkyl, or alkyl phenyl group or mixtures thereof containing from about 8 to about 22 carbon atoms; R<sup>4</sup> is an alkylene or hydroxyalkylene group containing from about 2 to about 3 carbon atoms or mixtures thereof; x is from 0 to about 3; and each R<sup>5</sup> is an alkyl or hydroxyalkyl group containing from about 1 to about 3 carbon atoms or a polyethylene oxide group containing from about 1 to about 3 ethylene oxide groups. The R<sup>5</sup> groups can be attached to each other, e.g., through an oxygen or nitrogen atom, to form a ring structure.

**[0100]** These amine oxide surfactants in particular include C<sub>10</sub>-C<sub>18</sub> alkyl dimethyl amine oxides and C<sub>8</sub>-C<sub>12</sub> alkoxy ethyl dihydroxy ethyl amine oxides.


**[0101]** When included therein, the cleaning compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such semi-polar nonionic surfactants.

**[0102]** The cleaning composition of the present invention may further comprise a cosurfactant selected from the group

of primary or tertiary amines.

Suitable primary amines for use herein include amines according to the formula  $R_1NH_2$  wherein  $R_1$  is a  $C_6$ - $C_{12}$ , preferably  $C_6$ - $C_{10}$  alkyl chain or  $R_4X(CH_2)_n$ ,  $X$  is -O-, -C(O)NH- or -NH-,  $R_4$  is a  $C_6$ - $C_{12}$  alkyl chain  $n$  is between 1 to 5, preferably 3.  $R_1$  alkyl chains may be straight or branched and may be interrupted with up to 12, preferably less than 5 ethylene oxide moieties. Preferred amines according to the formula herein above are  $n$ -alkyl amines. Suitable amines for use herein may be selected from 1-hexylamine, 1-octylamine, 1-decylamine and laurylamine. Other preferred primary amines include C8-C10 oxypropylamine, octyloxypropylamine, 2-ethylhexyloxypropylamine, lauryl amido propylamine and amido propylamine.


**[0103]** Suitable tertiary amines for use herein include tertiary amines having the formula  $R_1R_2R_3N$  wherein  $R_1$  and  $R_2$  are  $C_1$ - $C_8$  alkylchains or



$R_3$  is either a  $C_6$ - $C_{12}$ , preferably  $C_6$ - $C_{10}$  alkyl chain, or  $R_3$  is  $R_4X(CH_2)_n$ , whereby  $X$  is -O-, -C(O)NH- or -NH-,  $R_4$  is a  $C_4$ - $C_{12}$ ,  $n$  is between 1 to 5, preferably 2-3.  $R_5$  is H or  $C_1$ - $C_2$  alkyl and  $x$  is between 1 to 6.


$R_3$  and  $R_4$  may be linear or branched;  $R_3$  alkyl chains may be interrupted with up to 12, preferably less than 5, ethylene oxide moieties.

**[0104]** Preferred tertiary amines are  $R_1R_2R_3N$  where  $R_1$  is a  $C_6$ - $C_{12}$  alkyl chain,  $R_2$  and  $R_3$  are  $C_1$ - $C_3$  alkyl or



where  $R_5$  is H or  $CH_3$  and  $x = 1-2$ .

**[0105]** Also preferred are the amidoamines of the formula:



wherein  $R_1$  is  $C_6$ - $C_{12}$  alkyl;  $n$  is 2-4, preferably  $n$  is 3;  $R_2$  and  $R_3$  is  $C_1$ - $C_4$

**[0106]** Most preferred amines of the present invention include 1-octylamine, 1-hexylamine, 1-decylamine, 1-dodecylamine, C8-10oxypropylamine, N coco 1-3diaminopropane, coconutalkyldimethylamine, lauryldimethylamine, lauryl bis(hydroxyethyl)amine, coco bis(hydroxyethyl)amine, lauryl amine 2 moles propoxylated, octyl amine 2 moles propoxylated, lauryl amidopropyldimethylamine, C8-10 amidopropyldimethylamine and C10 amidopropyldimethylamine.

The most preferred amines for use in the compositions herein are 1-hexylamine, 1-octylamine, 1-decylamine, 1-dodecylamine. Especially desirable are  $n$ -dodecyldimethylamine and bishydroxyethylcoconutalkylamine and oleylamine 7 times ethoxylated, lauryl amido propylamine and cocoamido propylamine.

### Color care and fabric care benefits

**[0107]** Technologies which provide a type of color care benefit can also be included. Examples of these technologies are metallo catalysts for color maintenance. Such metallo catalysts are described in copending European Patent Application No. 92870181.2. Dye fixing agents, polyolefin dispersion for anti-wrinkles and improved water absorbancy, perfume and amino-functional polymer for color care treatment and perfume substantivity are further examples of color care / fabric care technologies and are described in the co-pending Patent Application No. 96870140.9, filed November 07, 1996.

**[0108]** Fabric softening agents can also be incorporated into cleaning compositions in accordance with the present invention. These agents may be inorganic or organic in type. Inorganic softening agents are exemplified by the smectite clays disclosed in GB-A-1 400 898 and in USP 5,019,292. Organic fabric softening agents include the water insoluble tertiary amines as disclosed in GB-A1 514 276 and EP-B0 011 340 and their combination with mono C12-C14 Quaternary ammonium salts are disclosed in EP-B-0 026 527 and EP-B-0 026 528 and di-long-chain amides as disclosed in EP-B-

0 242 919. Other useful organic ingredients of fabric softening systems include high molecular weight polyethylene oxide materials as disclosed in EP-A-0 299 575 and 0 313 146.

**[0109]** Levels of smectite clay are normally in the range from 2% to 20%, more preferably from 5% to 15% by weight, with the material being added as a dry mixed component to the remainder of the formulation. Organic fabric softening agents such as the water-insoluble tertiary amines or dilong chain amide materials are incorporated at levels of from 0.5% to 5% by weight, normally from 1% to 3% by weight whilst the high molecular weight polyethylene oxide materials and the water soluble cationic materials are added at levels of from 0.1% to 2%, normally from 0.15% to 1.5% by weight. These materials are normally added to the spray dried portion of the composition, although in some instances it may be more convenient to add them as a dry mixed particulate, or spray them as molten liquid on to other solid components of the composition.

### **Builder system**

**[0110]** The compositions according to the present invention may further comprise a builder system. Any conventional builder system is suitable for use herein including aluminosilicate materials, silicates, polycarboxylates, alkyl- or alkenylsuccinic acid and fatty acids, materials such as ethylenediamine tetraacetate, diethylene triamine pentamethyleneacetate, metal ion sequestrants such as aminopolyphosphonates, particularly ethylenediamine tetramethylene phosphonic acid and diethylene triamine pentamethylenephosphonic add. Phosphate builders can also be used herein.

**[0111]** Suitable builders can be an inorganic ion exchange material, commonly an inorganic hydrated aluminosilicate material, more particularly a hydrated synthetic zeolite such as hydrated zeolite A, X, B, HS or MAP.

Another suitable inorganic builder material is layered silicate, e.g. SKS-6 (Hoechst). SKS-6 is a crystalline layered silicate consisting of sodium silicate ( $\text{Na}_2\text{Si}_2\text{O}_5$ ).

Suitable polycarboxylates containing one carboxy group include lactic acid, glycolic acid and ether derivatives thereof as disclosed in Belgian Patent Nos. 831,368, 821,369 and 821,370. Polycarboxylates containing two carboxy groups include the water-soluble salts of succinic acid, malonic acid, (ethylenedioxy) diacetic acid, maleic acid, diglycolic acid, tartaric acid, tartronic acid and fumaric acid, as well as the ether carboxylates described in German Offentegenschrift 2,446,686, and 2,446,687 and U.S. Patent No. 3,935,257 and the sulfinyl carboxylates described in Belgian Patent No. 840,623. Polycarboxylates containing three carboxy groups include, in particular, water-soluble citrates, aconitrates and citraconates as well as succinate derivatives such as the carboxymethyloxysuccinates described in British Patent No. 1,379,241, lactoxysuccinates described in Netherlands Application 7205873, and the oxypolycarboxylate materials such as 2-oxa-1,1,3-propane tricarboxylates described in British Patent No. 1,387,447.

**[0112]** Polycarboxylates containing four carboxy groups include oxydisuccinates disclosed in British Patent No. 1,261,829, 1,1,2,2-ethane tetracarboxylates, 1,1,3,3-propane tetracarboxylates and 1,1,2,3-propane tetracarboxylates. Polycarboxylates containing sulfo substituents include the sulfosuccinate derivatives disclosed in British Patent Nos. 1,398,421 and 1,398,422 and in U.S. Patent No. 3,936,448, and the sulfonated pyrolysed citrates described in British Patent No. 1,082,179, while polycarboxylates containing phosphone substituents are disclosed in British Patent No. 1,439,000.

**[0113]** Alicyclic and heterocyclic polycarboxylates include cyclopentane-cis,cis,cis-tetracarboxylates, cyclopentadienide pentacarboxylates, 2,3,4,5-tetrahydro-furan - cis, cis, cis-tetracarboxylates, 2,5-tetrahydro-furan -cis -dicarboxylates, 2,2,5,5-tetrahydrofuran - tetracarboxylates, 1,2,3,4,5,6-hexane -hexacarboxylates and and carboxymethyl derivatives of polyhydric alcohols such as sorbitol, mannitol and xylitol. Aromatic poly-carboxylates include mellitic acid, pyromellitic acid and the phthalic acid derivatives disclosed in British Patent No. 1,425,343.

Of the above, the preferred polycarboxylates are hydroxycarboxylates containing up to three carboxy groups per molecule, more particularly citrates.

**[0114]** Preferred builder systems for use in the present compositions include a mixture of a water-insoluble aluminosilicate builder such as zeolite A or of a layered silicate (SKS-6), and a water-soluble carboxylate chelating agent such as citric acid.

**[0115]** Preferred builder systems include a mixture of a water-insoluble aluminosilicate builder such as zeolite A, and a watersoluble carboxylate chelating agent such as citric acid. Preferred builder systems for use in liquid detergent compositions of the present invention are soaps and polycarboxylates.

**[0116]** Other builder materials that can form part of the builder system for use in granular compositions include inorganic materials such as alkali metal carbonates, bicarbonates, silicates, and organic materials such as the organic phosphonates, amino polyalkylene phosphonates and amino polycarboxylates.

**[0117]** Other suitable water-soluble organic salts are the homo- or co-polymeric acids or their salts, in which the polycarboxylic acid comprises at least two carboxyl radicals separated from each other by not more than two carbon atoms. Polymers of this type are disclosed in GB-A-1,596,756. Examples of such salts are polyacrylates of MW 2000-5000 and their copolymers with maleic anhydride, such copolymers having a molecular weight of from 20,000 to 70,000, especially about 40,000.



**[0118]** Detergency builder salts are normally included in amounts of from 5% to 80% by weight of the composition preferably from 10% to 70% and most usually from 30% to 60% by weight.

### **Chelating Agents**

**[0119]** The cleaning compositions herein may also optionally contain one or more iron and/or manganese chelating agents. Such chelating agents can be selected from the group consisting of amino carboxylates, amino phosphonates, polyfunctionally-substituted aromatic chelating agents and mixtures therein, all as hereinafter defined. Without intending to be bound by theory, it is believed that the benefit of these materials is due in part to their exceptional ability to remove iron and manganese ions from washing solutions by formation of soluble chelates.

**[0120]** Amino carboxylates useful as optional chelating agents include ethylenediaminetetracetates, N-hydroxyethyl-ethylenediaminetriacetates, nitrilotriacetates, ethylenediamine tetrapropionates, triethylenetetraaminehexacetates, diethylenetriaminepentaacetates, and ethanoldiglycines, alkali metal, ammonium, and substituted ammonium salts therein and mixtures therein.

**[0121]** Amino phosphonates are also suitable for use as chelating agents in the compositions of the invention when at least low levels of total phosphorus are permitted in detergent compositions, and include ethylenediaminetetrakis (methylenephosphonates) as DEQUEST. Preferred, these amino phosphonates to not contain alkyl or alkenyl groups with more than about 6 carbon atoms.

**[0122]** Polyfunctionally-substituted aromatic chelating agents are also useful in the compositions herein. See U.S. Patent 3,812,044, issued May 21, 1974, to Connor et al. Preferred compounds of this type in acid form are dihydroxy-disulfobenzenes such as 1,2-dihydroxy-3,5-disulfobenzene.

**[0123]** A preferred biodegradable chelator for use herein is ethylenediamine disuccinate ("EDDS"), especially the [S, S] isomer as described in U.S. Patent 4,704,233, November 3, 1987, to Hartman and Perkins.

**[0124]** The compositions herein may also contain water-soluble methyl glycine diacetic acid (MGDA) salts (or acid form) as a chelant or co-builder useful with, for example, insoluble builders such as zeolites, layered silicates and the like.

**[0125]** If utilized, these chelating agents will generally comprise from about 0.1 % to about 15% by weight of the detergent compositions herein. More preferably, if utilized, the chelating agents will comprise from about 0.1% to about 3.0% by weight of such compositions.

### **Suds suppressor**

**[0126]** Another optional ingredient is a suds suppressor, exemplified by silicones, and silica-silicone mixtures. Silicones can be generally represented by alkylated polysiloxane materials while silica is normally used in finely divided forms exemplified by silica aerogels and xerogels and hydrophobic silicas of various types. These materials can be incorporated as particulates in which the suds suppressor is advantageously releasably incorporated in a water-soluble or water-dispersible, substantially non-surface-active detergent impermeable carrier. Alternatively the suds suppressor can be dissolved or dispersed in a liquid carrier and applied by spraying on to one or more of the other components.

A preferred silicone suds controlling agent is disclosed in Bartollota et al. U.S. Patent 3 933 672. Other particularly useful suds suppressors are the self-emulsifying silicone suds suppressors, described in German Patent Application DTOS 2 646 126 published April 28, 1977. An example of such a compound is DC-544, commercially available from Dow Corning, which is a siloxane-glycol copolymer. Especially preferred suds controlling agents are the suds suppressor system comprising a mixture of silicone oils and 2-alkyl-alkanols. Suitable 2-alkyl-alkanols are 2-butyl-octanol which are commercially available under the trade name Isofol 12 R.

Such suds suppressor system are described in Copending European Patent application N 92870174.7 filed 10 November, 1992.

Especially preferred silicone suds controlling agents are described in Copending European Patent application N°92201649.8. Said compositions can comprise a silicone/silica mixture in combination with fumed nonporous silica such as Aerosil®.

**[0127]** The suds suppressors described above are normally employed at levels of from 0.001 % to 2% by weight of the composition, preferably from 0.01 % to 1% by weight.

### **Others**

**[0128]** Other components used in cleaning compositions may be employed, such as soil-suspending agents, soil-release agents, optical brighteners, abrasives, bactericides, tarnish inhibitors, coloring agents, and/or encapsulated or non-encapsulated perfumes.

**[0129]** Especially suitable encapsulating materials are water soluble capsules which consist of a matrix of polysaccharide and polyhydroxy compounds such as described in GB 1,464,616.


**[0130]** Other suitable water soluble encapsulating materials comprise dextrans derived from ungelatinized starch acid-esters of substituted dicarboxylic acids such as described in US 3,455,838. These acid-ester dextrans are, preferably, prepared from such starches as waxy maize, waxy sorghum, sago, tapioca and potato. Suitable examples of said encapsulating materials include N-Lok manufactured by National Starch. The N-Lok encapsulating material consists of a modified maize starch and glucose. The starch is modified by adding monofunctional substituted groups such as octenyl succinic acid anhydride.

**[0131]** Antiredeposition and soil suspension agents suitable herein include cellulose derivatives such as methylcellulose, carboxymethylcellulose and hydroxyethylcellulose, and homo- or co-polymeric polycarboxylic acids or their salts. Polymers of this type include the polyacrylates and maleic anhydride-acrylic acid copolymers previously mentioned as builders, as well as copolymers of maleic anhydride with ethylene, methylvinyl ether or methacrylic acid, the maleic anhydride constituting at least 20 mole percent of the copolymer. These materials are normally used at levels of from 0.5% to 10% by weight, more preferably from 0.75% to 8%, most preferably from 1% to 6% by weight of the composition.

**[0132]** Preferred optical brighteners are anionic in character, examples of which are disodium 4,4'-bis-(2-diethanolamino-4-anilino-s-triazin-6-ylamino)stilbene-2:2' disulphonate, disodium 4,4'-bis-(2-morpholino-4-anilino-s-triazin-6-ylamino)stilbene-2:2' - disulphonate, disodium 4,4' - bis-(2,4-dianilino-s-triazin-6-ylamino)stilbene-2:2' - disulphonate, monosodium 4',4'' -bis-(2,4-dianilino-s-triazin-6-ylamino)stilbene-2-sulphonate, disodium 4,4' -bis-(2-anilino-4-(N-methyl-N-2-hydroxyethylamino)-s-triazin-6-ylamino)stilbene-2,2' - disulphonate, di-sodium 4,4' -bis-(4-phenyl-2,1,3-triazol-2-yl)-stilbene-2,2' disulphonate, di-sodium 4,4'bis(2-anilino-4-(1-methyl-2-hydroxyethylamino)-s-triazin-6-ylamino)stilbene-2,2' disulphonate, sodium 2(stilbyl-4'')-(naphtho-1',2',4,5)-1,2,3 - triazole-2''-sulphonate and 4,4'-bis(2-sulphostyryl) biphenyl. Highly preferred brighteners are the specific brighteners of copending European Patent application No. 95201943.8.

**[0133]** Other useful polymeric materials are the polyethylene glycols, particularly those of molecular weight 1000-10000, more particularly 2000 to 8000 and most preferably about 4000. These are used at levels of from 0.20% to 5% more preferably from 0.25% to 2.5% by weight. These polymers and the previously mentioned homo- or co-polymeric polycarboxylate salts are valuable for improving whiteness maintenance, fabric ash deposition, and cleaning performance on clay, proteinaceous and oxidizable soils in the presence of transition metal impurities.

**[0134]** Soil release agents useful in compositions of the present invention are conventionally copolymers or terpolymers of terephthalic acid with ethylene glycol and/or propylene glycol units in various arrangements. Examples of such polymers are disclosed in the commonly assigned US Patent Nos. 4116885 and 4711730 and European Published Patent Application No. 0 272 033. A particular preferred polymer in accordance with EP-A-0 272 033 has the formula



where PEG is  $-(\text{OC}_2\text{H}_4)_n\text{O}-$ , PO is  $(\text{OC}_3\text{H}_6\text{O})$  and T is  $(\text{pCOC}_6\text{H}_4\text{CO})$ .

**[0135]** Also very useful are modified polyesters as random copolymers of dimethyl terephthalate, dimethyl sulfoisophthalate, ethylene glycol and 1-2 propane diol, the end groups consisting primarily of sulphobenzoate and secondarily of mono esters of ethylene glycol and/or propane-diol. The target is to obtain a polymer capped at both end by sulphobenzoate groups, "primarily", in the present context most of said copolymers herein will be end-capped by sulphobenzoate groups. However, some copolymers will be less than fully capped, and therefore their end groups may consist of monoester of ethylene glycol and/or propane 1-2 diol, thereof consist "secondarily" of such species.

**[0136]** The selected polyesters herein contain about 46% by weight of dimethyl terephthalic acid, about 16% by weight of propane-1,2 diol, about 10% by weight ethylene glycol about 13% by weight of dimethyl sulfobenzoic acid and about 15% by weight of sulfoisophthalic acid, and have a molecular weight of about 3,000. The polyesters and their method of preparation are described in detail in EPA 311 342.

**[0137]** It is well known in the art that free chlorine in tap water rapidly deactivates the enzymes comprised in detergent compositions. Therefore, using chlorine scavenger such as perborate, ammonium sulfate, sodium sulphite or polyethyleneimine at a level above 0.1 % by weight of total composition, in the formulas will provide improved through the wash stability of the detergent enzymes. Compositions comprising chlorine scavenger are described in the European patent application 92870018.6 filed January 31, 1992.

**[0138]** Alkoxylated polycarboxylates such as those prepared from polyacrylates are useful herein to provide additional grease removal performance. Such materials are described in WO 91/08281 and PCT 90/01815 at p. 4 et seq., incorporated herein by reference. Chemically, these materials comprise polyacrylates having one ethoxy side-chain per every 7-8 acrylate units. The side-chains are of the formula  $-(\text{CH}_2\text{CH}_2\text{O})_m(\text{CH}_2)_n\text{CH}_3$  wherein m is 2-3 and n is 6-12. The side-chains are ester-linked to the polyacrylate "backbone" to provide a "comb" polymer type structure. The molecular weight can vary, but is typically in the range of about 2000 to about 50,000. Such alkoxylated polycarboxylates can comprise from about 0.05% to about 10%, by weight, of the compositions herein.

**Dispersants**

[0139] The cleaning compositions of the present invention can also contain dispersants : Suitable water-soluble organic salts are the homo- or co-polymeric acids or their salts, in which the polycarboxylic acid comprises at least two carboxyl radicals separated from each other by not more than two carbon atoms. Polymers of this type are disclosed in GB-A-1,596,756. Examples of such salts are polyacrylates of MW 2000-5000 and their copolymers with maleic anhydride, such copolymers having a molecular weight of from 1,000 to 100,000.

[0140] Especially, copolymer of acrylate and methylacrylate such as the 480N having a molecular weight of 4000, at a level from 0.5-20% by weight of composition can be added in the cleaning compositions of the present invention.

[0141] The compositions of the invention may contain a lime soap peptiser compound, which has preferably a lime soap dispersing power (LSDP), as defined hereinafter of no more than 8, preferably no more than 7, most preferably no more than 6. The lime soap peptiser compound is preferably present at a level from 0% to 20% by weight.

[0142] A numerical measure of the effectiveness of a lime soap peptiser is given by the lime soap dispersant power (LSDP) which is determined using the lime soap dispersant test as described in an article by H.C. Borghetty and C.A. Bergman, J. Am. Oil. Chem. Soc., volume 27, pages 88-90, (1950). This lime soap dispersion test method is widely used by practitioners in this art field being referred to, for example, in the following review articles; W.N. Linfield, Surfactant science Series, Volume 7, page 3; W.N. Linfield, Tenside surf. det., volume 27, pages 159-163, (1990); and M.K. Nagarajan, W.F. Masler, Cosmetics and Toiletries, volume 104, pages 71-73, (1989). The LSDP is the % weight ratio of dispersing agent to sodium oleate required to disperse the lime soap deposits formed by 0.025g of sodium oleate in 30ml of water of 333ppm  $\text{CaCO}_3$  (Ca:Mg=3:2) equivalent hardness.

[0143] Surfactants having good lime soap peptiser capability will include certain amine oxides, betaines, sulfobetaines, alkyl ethoxysulfates and ethoxylated alcohols.

[0144] Exemplary surfactants having a LSDP of no more than 8 for use in accord with the present invention include  $\text{C}_{16}$ - $\text{C}_{18}$  dimethyl amine oxide,  $\text{C}_{12}$ - $\text{C}_{18}$  alkyl ethoxysulfates with an average degree of ethoxylation of from 1-5, particularly  $\text{C}_{12}$ - $\text{C}_{15}$  alkyl ethoxysulfate surfactant with a degree of ethoxylation of amount 3 (LSDP=4), and the  $\text{C}_{14}$ - $\text{C}_{15}$  ethoxylated alcohols with an average degree of ethoxylation of either 12 (LSDP=6) or 30, sold under the tradenames Lutensol A012 and Lutensol A030 respectively, by BASF GmbH.

[0145] Polymeric lime soap peptisers suitable for use herein are described in the article by M.K. Nagarajan, W.F. Master, to be found in Cosmetics and Toiletries, volume 104, pages 71-73, (1989).

[0146] Hydrophobic bleaches such as 4-[N-octanoyl-6-aminohexanoyl]benzene sulfonate, 4-[N-nonanoyl-6-amino-hexanoyl]benzene sulfonate, 4-[N-decanoyl-6-aminohexanoyl]benzene sulfonate and mixtures thereof; and nonanoyloxy benzene sulfonate together with hydrophilic / hydrophobic bleach formulations can also be used as lime soap peptisers compounds.

**Dye transfer inhibition**

[0147] The cleaning compositions of the present invention can also include compounds for inhibiting dye transfer from one fabric to another of solubilized and suspended dyes encountered during fabric laundering operations involving colored fabrics.


**Polymeric dye transfer inhibiting agents**

[0148] The cleaning compositions according to the present invention also comprise from 0.001% to 10 %, preferably from 0.01% to 2%, more preferably from 0.05% to 1% by weight of polymeric dye transfer inhibiting agents. Said polymeric dye transfer inhibiting agents are normally incorporated into cleaning compositions in order to inhibit the transfer of dyes from colored fabrics onto fabrics washed therewith. These polymers have the ability to complex or adsorb the fugitive dyes washed out of dyed fabrics before the dyes have the opportunity to become attached to other articles in the wash. Especially suitable polymeric dye transfer inhibiting agents are polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylpyrrolidone polymers, polyvinylloxazolidones and polyvinylimidazoles or mixtures thereof.

Addition of such polymers also enhances the performance of the enzymes according the invention.


**a) Polyamine N-oxide polymers**

[0149] The polyamine N-oxide polymers suitable for use contain units having the following structure formula :




wherein P is a polymerisable unit, whereto the R-N-O group can be attached to or wherein the R-N-O group forms part of the polymerisable unit or a combination of both.

A is



-O-, -S-, -N-; x is 0 or 1; R are aliphatic, ethoxylated aliphatics, aromatic, heterocyclic or alicyclic groups or any combination thereof whereto the nitrogen of the N-O group can be attached or wherein the nitrogen of the N-O group is part of these groups.

**[0150]** The N-O group can be represented by the following general structures :



wherein R1, R2, and R3 are aliphatic groups, aromatic, heterocyclic or alicyclic groups or combinations thereof, x or/and y or/and z is 0 or 1 and wherein the nitrogen of the N-O group can be attached to or wherein the nitrogen of the N-O group forms part of these groups.

**[0151]** The N-O group can be part of the polymerisable unit (P) or can be attached to the polymeric backbone or a combination of both.

Suitable polyamine N-oxides wherein the N-O group forms part of the polymerisable unit comprise polyamine N-oxides wherein R is selected from aliphatic, aromatic, alicyclic or heterocyclic groups.

One class of said polyamine N-oxides comprises the group of polyamine N-oxides wherein the nitrogen of the N-O group forms part of the R-group. Preferred polyamine N-oxides are those wherein R is a heterocyclic group such as pyridine, pyrrole, imidazole, pyrrolidine, piperidine, quinoline, acridine and derivatives thereof.

Another class of said polyamine N-oxides comprises the group of polyamine N-oxides wherein the nitrogen of the N-O group is attached to the R-group.

**[0152]** Other suitable polyamine N-oxides are the polyamine oxides whereto the N-O group is attached to the polymerisable unit.

Preferred class of these polyamine N-oxides are the polyamine N-oxides having the general formula (I) wherein R is an aromatic, heterocyclic or alicyclic groups wherein the nitrogen of the N-O functional group is part of said R group.

Examples of these classes are polyamine oxides wherein R is a heterocyclic compound such as pyridine, pyrrole, imidazole and derivatives thereof.

Another preferred class of polyamine N-oxides are the polyamine oxides having the general formula (I) wherein R are aromatic, heterocyclic or alicyclic groups wherein the nitrogen of the N-O functional group is attached to said R groups.

Examples of these classes are polyamine oxides wherein R groups can be aromatic such as phenyl.

**[0153]** Any polymer backbone can be used as long as the amine oxide polymer formed is water-soluble and has dye transfer inhibiting properties. Examples of suitable polymeric backbones are polyvinyls, polyalkylenes, polyesters, polyethers, polyamide, polyimides, polyacrylates and mixtures thereof.

**[0154]** The amine N-oxide polymers of the present invention typically have a ratio of amine to the amine N-oxide of 10:1 to 1:1000000. However the amount of amine oxide groups present in the polyamine oxide polymer can be varied by appropriate copolymerization or by appropriate degree of N-oxidation. Preferably, the ratio of amine to amine N-oxide is from 2:3 to 1:1000000. More preferably from 1:4 to 1:1000000, most preferably from 1:7 to 1:1000000. The polymers of the present invention actually encompass random or block copolymers where one monomer type is an amine N-oxide and the other monomer type is either an amine N-oxide or not. The amine oxide unit of the polyamine N-oxides has a  $PK_a < 10$ , preferably  $PK_a < 7$ , more preferred  $PK_a < 6$ .

The polyamine oxides can be obtained in almost any degree of polymerisation. The degree of polymerisation is not critical provided the material has the desired water-solubility and dye-suspending power.

Typically, the average molecular weight is within the range of 500 to 1000,000; preferably from 1,000 to 50,000, more preferably from 2,000 to 30,000, most preferably from 3,000 to 20,000.

#### b) Copolymers of N-vinylpyrrolidone and N-vinylimidazole

**[0155]** The N-vinylimidazole N-vinylpyrrolidone polymers used in the present invention have an average molecular weight range from 5,000-1,000,000, preferably from 5,000-200,000.

**[0156]** Highly preferred polymers for use in detergent compositions according to the present invention comprise a polymer selected from N-vinylimidazole N-vinylpyrrolidone copolymers wherein said polymer has an average molecular weight range from 5,000 to 50,000 more preferably from 8,000 to 30,000, most preferably from 10,000 to 20,000.

The average molecular weight range was determined by light scattering as described in Barth H.G. and Mays J.W. Chemical Analysis Vol 113, "Modern Methods of Polymer Characterization".

Highly preferred N-vinylimidazole N-vinylpyrrolidone copolymers have an average molecular weight range from 5,000 to 50,000; more preferably from 8,000 to 30,000; most preferably from 10,000 to 20,000.

**[0157]** The N-vinylimidazole N-vinylpyrrolidone copolymers characterized by having said average molecular weight range provide excellent dye transfer inhibiting properties while not adversely affecting the cleaning performance of detergent compositions formulated therewith.

The N-vinylimidazole N-vinylpyrrolidone copolymer of the present invention has a molar ratio of N-vinylimidazole to N-vinylpyrrolidone from 1 to 0.2, more preferably from 0.8 to 0.3, most preferably from 0.6 to 0.4.

#### c) Polyvinylpyrrolidone

**[0158]** The detergent compositions of the present invention may also utilize polyvinylpyrrolidone ("PVP") having an average molecular weight of from about 2,500 to about 400,000, preferably from about 5,000 to about 200,000, more preferably from about 5,000 to about 50,000, and most preferably from about 5,000 to about 15,000. Suitable polyvinylpyrrolidones are commercially available from ISP Corporation, New York, NY and Montreal, Canada under the product names PVP K-15 (viscosity molecular weight of 10,000), PVP K-30 (average molecular weight of 40,000), PVP K-60 (average molecular weight of 160,000), and PVP K-90 (average molecular weight of 360,000). Other suitable polyvinylpyrrolidones which are commercially available from BASF Cooperation include Sokalan HP 165 and Sokalan HP 12; polyvinylpyrrolidones known to persons skilled in the detergent field (see for example EP-A-262,897 and EP-A-256,696).

#### d) Polyvinylloxazolidone :

**[0159]** The detergent compositions of the present invention may also utilize polyvinylloxazolidone as a polymeric dye transfer inhibiting agent. Said polyvinylloxazolidones have an average molecular weight of from about 2,500 to about 400,000, preferably from about 5,000 to about 200,000, more preferably from about 5,000 to about 50,000, and most preferably from about 5,000 to about 15,000.

#### e) Polyvinylimidazole :

**[0160]** The detergent compositions of the present invention may also utilize polyvinylimidazole as polymeric dye transfer inhibiting agent. Said polyvinylimidazoles have an average about 2,500 to about 400,000, preferably from about 5,000 to about 200,000, more preferably from about 5,000 to about 50,000, and most preferably from about 5,000 to about 15,000.

Cross-linked polymers :

**[0161]** Cross-linked polymers are polymers whose backbone are interconnected to a certain degree; these links can be of chemical or physical nature, possibly with active groups in the backbone or on branches; cross-linked polymers have been described in the Journal of Polymer Science, volume 22, pages 1035-1039.

In one embodiment, the cross-linked polymers are made in such a way that they form a three-dimensional rigid structure, which can entrap dyes in the pores formed by the three-dimensional structure. In another embodiment, the cross-linked polymers entrap the dyes by swelling.

Such cross-linked polymers are described in the co-pending patent application 94870213.9

#### **Method of washing**

**[0162]** The compositions of the invention may be used in essentially any washing or cleaning methods, including soaking methods, pretreatment methods and methods with rinsing steps for which a separate rinse aid composition may be added.

**[0163]** The process described herein comprises contacting fabrics with a laundering solution in the usual manner and exemplified hereunder.

**[0164]** The process of the invention is conveniently carried out in the course of the cleaning process. The method of cleaning is preferably carried out at 5°C to 95°C, especially between 10°C and 60°C. The pH of the treatment solution is preferably from 7 to 12.

**[0165]** A preferred machine dishwashing method comprises treating soiled articles with an aqueous liquid having dissolved or dispensed therein an effective amount of the machine dishwashing or rinsing composition. A conventional effective amount of the machine dishwashing composition means from 8-60 g of product dissolved or dispersed in a wash volume from 3-10 litres.

According to a manual dishwashing method, soiled dishes are contacted with an effective amount of the dishwashing composition, typically from 0.5-20g (per 25 dishes being treated). Preferred manual dishwashing methods include the application of a concentrated solution to the surfaces of the dishes or the soaking in large volume of dilute solution of the detergent composition.

**[0166]** The following examples are meant to exemplify compositions of the present invention, but are not necessarily meant to limit or otherwise define the scope of the invention.

**[0167]** In the cleaning compositions, the enzymes levels are expressed by pure enzyme by weight of the total composition and unless otherwise specified, the detergent ingredients are expressed by weight of the total compositions. The abbreviated component identifications therein have the following meanings:

LAS	Sodium linear C <sub>11-13</sub> alkyl benzene sulphonate.
TAS	Sodium tallow alkyl sulphate.
CxyAS	Sodium C <sub>1x</sub> - C <sub>1y</sub> alkyl sulfate.
CxySAS	Sodium C <sub>1x</sub> - C <sub>1y</sub> secondary (2,3) alkyl sulfate.
CxyEz	C <sub>1x</sub> - C <sub>1y</sub> predominantly linear primary alcohol condensed with an average of z moles of ethylene oxide.
CxyEzS	C <sub>1x</sub> - C <sub>1y</sub> sodium alkyl sulfate condensed with an average of z moles of ethylene oxide.
QAS	R <sub>2</sub> -N+(CH <sub>3</sub> ) <sub>2</sub> (C <sub>2</sub> H <sub>4</sub> OH) with R <sub>2</sub> = C <sub>12</sub> -C <sub>14</sub> .
QAS 1	R <sub>2</sub> -N+(CH <sub>3</sub> ) <sub>2</sub> (C <sub>2</sub> H <sub>4</sub> OH) with R <sub>2</sub> = C <sub>8</sub> -C <sub>11</sub> .
APA	C <sub>8-10</sub> amido propyl dimethyl amine.
Soap	Sodium linear alkyl carboxylate derived from a 80/20 mixture of tallow and coconut fatty acids.
Nonionic	C <sub>13</sub> -C <sub>15</sub> mixed ethoxylated/propoxylated fatty alcohol with an average degree of ethoxylation of 3.8 and an average degree of propoxylation of 4.5.
Neodol 45-13	C <sub>14</sub> -C <sub>15</sub> linear primary alcohol ethoxylate, sold by Shell Chemical CO.
STS	Sodium toluene sulphonate.
CFAA	C <sub>12</sub> -C <sub>14</sub> alkyl N-methyl glucamide.
TFAA	C <sub>16</sub> -C <sub>18</sub> alkyl N-methyl glucamide.
TPKFA	C <sub>12</sub> -C <sub>14</sub> topped whole cut fatty acids.
DEQA	Di-(tallow-oxy-ethyl) dimethyl ammonium chloride.
DEQA (2)	Di-(soft-tallowyloxyethyl) hydroxyethyl methyl ammonium methylsulfate.

(continued)

	DTDMAMS	Ditalllow dimethyl ammonium methylsulfate.
	SDASA	1:2 ratio of stearyldimethyl amine:triple-pressed stearic acid.
5	Silicate	Amorphous Sodium Silicate ( $\text{SiO}_2\text{:Na}_2\text{O}$ ratio = 1.6-3.2).
	Metasilicate	Sodium metasilicate ( $\text{SiO}_2\text{:Na}_2\text{O}$ ratio = 1.0).
	Zeolite A	Hydrated Sodium Aluminosilicate of formula $\text{Na}_{12}(\text{AlO}_2\text{SiO}_2)_{12} \cdot 27\text{H}_2\text{O}$ having a primary particle size in the range from 0.1 to 10 micrometers (Weight expressed on an anhydrous basis).
10	Na-SKS-6	Crystalline layered silicate of formula $\delta\text{-Na}_2\text{Si}_2\text{O}_5$ .
	Citrate	Tri-sodium citrate dihydrate of activity 86.4% with a particle size distribution between 425 and 850 micrometres.
	Citric	Anhydrous citric acid.
15	Borate	Sodium borate
	Carbonate	Anhydrous sodium carbonate with a particle size between 200 and 900 micrometres.
	Bicarbonate	Anhydrous sodium hydrogen carbonate with a particle size distribution between 400 and 1200 micrometres.
	Sulphate	Anhydrous sodium sulphate.
20	Mg Sulphate	Anhydrous magnesium sulfate.
	STPP	Sodium tripolyphosphate.
	TSP	Tetrasodium pyrophosphate.
	MA/AA	Random copolymer of 4:1 acrylate/maleate, average molecular weight about 70,000-80,000.
25	MA/AA 1	Random copolymer of 6:4 acrylate/maleate, average molecular weight about 10,000.
	AA	Sodium polyacrylate polymer of average molecular weight 4,500.
	PA30	Polyacrylic acid of average molecular weight of between about 4,500 - 8,000.
	480N	Random copolymer of 7:3 acrylate/methacrylate, average molecular weight about 3,500.
30	Polygel/carbopol	High molecular weight crosslinked polyacrylates.
	PB1	Anhydrous sodium perborate monohydrate of nominal formula $\text{NaBO}_2 \cdot \text{H}_2\text{O}_2$ .
	PB4	Sodium perborate tetrahydrate of nominal formula $\text{NaBO}_2 \cdot 3\text{H}_2\text{O} \cdot \text{H}_2\text{O}_2$ .
	Percarbonate	Anhydrous sodium percarbonate of nominal formula $2\text{Na}_2\text{CO}_3 \cdot 3\text{H}_2\text{O}_2$
	NaDCC	Sodium dichloroisocyanurate.
35	TAED	Tetraacetylenediamine.
	NOBS	Nonanoyloxybenzene sulfonate in the form of the sodium salt.
	NACA-OBS	(6-nonamidocaproyl) oxybenzene sulfonate.
	DTPA	Diethylene triamine pentaacetic acid.
40	HEDP	1,1-hydroxyethane diphosphonic acid.
	DETPMP	Diethyltriamine penta (methylene) phosphonate, marketed by Monsanto under the Trade name Dequest 2060.
	EDDS	Ethylenediamine-N,N'-disuccinic acid, (S,S) isomer in the form of its sodium salt
	MnTACN	Manganese 1,4,7-trimethyl-1,4,7-triazacyclononane.
45	Photoactivated Bleach	Sulfonated zinc phthalocyanine encapsulated in dextrin soluble polymer.
	Photoactivated Bleach 1	Sulfonated aluminophthalocyanine encapsulated in dextrin soluble polymer.
	PAAC	Pentaamine acetate cobalt(III) salt.
50	Paraffin	Paraffin oil sold under the tradename Winog 70 by Wintershall.
	NaBz	Sodium benzoate.
	BzP	Benzoyl Peroxide.
	Oxygenase	Protocatechuate 3,4 dioxygenase sold by Sigma under the number P8279.
55	Protease	Proteolytic enzyme sold under the tradename Savinase, Alcalase, Durazym by Novo Nordisk A/S, Maxacal, Maxapem sold by Gist-Brocades and proteases described in patents WO91/06637 and/or WO95/1059 and/or EP 251 446.

(continued)

5	Amylase	Amylolytic enzyme sold under the tradename Purafact Ox Am <sup>R</sup> described in WO 94/18314, WO96/05295 sold by Genencor; Termamyl®, Fungamyl® and Duramyl®, all available from Novo Nordisk A/S and those described in WO95/26397.
	Lipase	Lipolytic enzyme sold under the tradename Lipolase, Lipolase Ultra by Novo Nordisk A/S and Lipomax by Gist-Brocades.
	Cellulase	Cellulytic enzyme sold under the tradename Carezyme, Celluzyme and/or Endolase by Novo Nordisk A/S.
10	CMC	Sodium carboxymethyl cellulose.
	PVP	Polyvinyl polymer, with an average molecular weight of 60,000.
	PVNO	Polyvinylpyridine-N-Oxide, with an average molecular weight of 50,000.
	PVPVI	Copolymer of vinylimidazole and vinylpyrrolidone, with an average molecular weight of 20,000.
15	Brightener 1	Disodium 4,4'-bis(2-sulphostyryl)biphenyl.
	Brightener 2	Disodium 4,4'-bis(4-anilino-6-morpholino-1.3.5-triazin-2-yl) stilbene-2:2'-disulfonate.
	Silicone antifoam	Polydimethylsiloxane foam controller with siloxane-oxyalkylene copolymer as dispersing agent with a ratio of said foam controller to said dispersing agent of 10:1 to 100:1.
20	Suds Suppressor	12% Silicone/silica, 18% stearyl alcohol, 70% starch in granular form.
	Opacifier	Water based monostyrene latex mixture, sold by BASF Aktiengesellschaft under the tradename Lytron 621.
	SRP 1	Anionically end capped poly esters.
	SRP 2	Diethoxylated poly (1,2 propylene terephthalate) short block polymer.
25	QEA	bis((C <sub>2</sub> H <sub>5</sub> O)(C <sub>2</sub> H <sub>4</sub> O) <sub>n</sub> )(CH <sub>3</sub> )-N <sup>+</sup> -C <sub>6</sub> H <sub>12</sub> -N <sup>+</sup> -(CH <sub>3</sub> ) bis((C <sub>2</sub> H <sub>5</sub> O)-(C <sub>2</sub> H <sub>4</sub> O)) <sub>n</sub> , wherein n = from 20 to 30.
	PEI	Polyethyleneimine with an average molecular weight of 1800 and an average ethoxylation degree of 7 ethyleneoxy residues per nitrogen.
30	SCS	Sodium cumene sulphonate.
	HMWPEO	High molecular weight polyethylene oxide.
	PEGx	Polyethylene glycol, of a molecular weight of x.
	PEO	Polyethylene oxide, with an average molecular weight of 5,000.
	TEPAE	Tetraethylenepentaamine ethoxylate.
35	BTA	Benzotriazole.
	Silica dental abrasive	Precipitated silica identified as Zeodent 119 offered by J.M. Huber.
	Carboxyvinyl polymer	Carbopol offered by B.F. Goodrich Chemical Company.
40	Carrageenan	Iota Carrageenan offered by Hercules Chemical Company.
	pH	Measured as a 1% solution in distilled water at 20°C.

Example 1

**[0168]** The following high density laundry detergent compositions were prepared according to the present invention :

	I	II	III	IV	V	VI
LAS	8.0	8.0	8.0	2.0	6.0	6.0
TAS	-	0.5	-	0.5	1.0	0.1
C46(S)AS	2.0	2.5	-	-	-	-
C25AS	-	-	-	7.0	4.5	5.5
C68AS	2.0	5.0	7.0	-	-	-
C25E5	-	-	3.4	10.0	4.6	4.6
C25E7	3.4	3.4	1.0	-	-	-
C25E3S	-	-	-	2.0	5.0	4.5
QAS	-	0.8	-	-	-	-



# EP 1 002 040 B2

(continued)

		I	II	III	IV	V	VI
5	QAS 1	-	-	-	0.8	0.5	1.0
	Zeolite A	18.1	18.0	14.1	18.1	20.0	18.1
	Citric	-	-	-	2.5	-	2.5
	Carbonate	13.0	13.0	27.0	10.0	10.0	13.0
	Na-SKS-6	-	-	-	10.0	-	10.0
10	Silicate	1.4	1.4	3.0	0.3	0.5	0.3
	Citrate	-	1.0	-	3.0	-	-
	Sulfate	26.1	26.1	26.1	6.0	-	-
	Mg sulfate	0.3	-	-	0.2	-	0.2
	MA/AA	0.3	0.3	0.3	4.0	1.0	1.0
15	CMC	0.2	0.2	0.2	0.2	0.4	0.4
	PB4	9.0	9.0	5.0	-	-	-
	Percarbonate	-	-	-	-	18.0	18.0
	TAED	1.5	0.4	1.5	-	3.9	4.2
	NACA-OBS	-	2.0	1.0	-	-	-
20	DETPMP	0.25	0.25	0.25	0.25	-	-
	SRP 1	-	-	-	0.2	-	0.2
	EDDS	-	0.25	0.4	-	0.5	0.5
	CFAA	-	1.0	-	2.0	-	-
	HEDP	0.3	0.3	0.3	0.3	0.4	0.4
25	QEA	-	-	-	0.2	-	0.5
	Oxygenase	0.05	0.05	0.02	0.002	0.005	0.005
	Protease	0.009	0.009	0.01	0.04	0.05	0.03
	Amylase	0.002	0.002	0.002	0.006	0.008	0.008
	Cellulase	0.0007	-	-	0.0007	0.0007	0.0007
30	Lipase	0.006	-	-	0.01	0.01	0.01
	Photoactivated	15	15	15	-	20	20
	bleach (ppm)						
	PVNO/PVPVI	-	-	-	0.1	-	-
	Brightener 1	0.09	0.09	0.09	-	0.09	0.09
35	Perfume	0.3	0.3	0.3	0.4	0.4	0.4
	Silicone antifoam	0.5	0.5	0.5	-	0.3	0.3
	Density in g/litre	850	850	850	850	850	850
	Miscellaneous and minors				Up to 100%		
	40						

## Example 2

45 **[0169]** The following granular laundry detergent compositions of particular utility under European machine wash conditions were prepared according to the present invention :

		I	II	III	IV	V	VI
50	LAS	5.5	7.5	5.0	5.0	6.0	7.0
	TAS	1.25	1.9	-	0.8	0.4	0.3
	C24AS/C25AS	-	2.2	5.0	5.0	5.0	2.2
	C25E3S	-	0.8	1.0	1.5	3.0	1.0
	C45E7	3.25	-	-	-	-	3.0
55	TFAA	-	-	2.0	-	-	-
	C25E5	-	5.5	-	-	-	-
	QAS	0.8	-	-	-	-	-

(continued)

### Example 3

	I	II	III	IV
Blown Powder				
LAS	6.0	5.0	11.0	6.0
TAS	2.0	-	-	2.0

# EP 1 002 040 B2

(continued)

	I	II	III	IV
Blown Powder				
Zeolite A	24.0	-	-	20.0
STPP	-	27.0	24.0	-
Sulfate	4.0	6.0	13.0	-
MA/AA	1.0	4.0	6.0	2.0
Silicate	1.0	7.0	3.0	3.0
CMC	1.0	1.0	0.5	0.6
Brightener 1	0.2	0.2	0.2	0.2
Silicone antifoam	1.0	1.0	1.0	0.3
DETPMP	0.4	0.4	0.2	0.4
Spray On				
Brightener	0.02	-	-	0.02
C45E7	-	-	-	5.0
C45E2	2.5	2.5	2.0	-
C45E3	2.6	2.5	2.0	-
Perfume	0.5	0.3	0.5	0.2
Silicone antifoam	0.3	0.3	0.3	-
Dry additives				
QEA	-	-	-	1.0
EDDS	0.3	-	-	-
Sulfate	2.0	3.0	5.0	10.0
Carbonate	6.0	13.0	15.0	14.0
Citric	2.5	-	-	2.0
QAS 1	0.5	-	-	0.5
Na-SKS-6	10.0	-	-	-
Percarbonate	18.5	-	-	-
PB4	-	18.0	10.0	21.5
TAED	2.0	2.0	-	2.0
NACA-OBS	3.0	2.0	4.0	-
Oxygenase	0.05	0.07	0.1	0.01
Protease	0.03	0.03	0.03	0.03
Lipase	0.008	0.008	0.008	0.004
Amylase	0.003	0.003	0.003	0.006
Brightener 1	0.05	-	-	0.05
Miscellaneous and minors	Up to 100%			

## Example 4

[0171] The following granular detergent compositions were prepared according to the present invention :

	I	II	III	IV	V	VI
Blown Powder						
LAS	23.0	8.0	7.0	9.0	7.0	7.0
TAS	-	-	-	-	1.0	-
C45AS	6.0	6.0	5.0	8.0	-	-
C45AES	-	1.0	1.0	1.0	-	-
C45E35	-	-	-	-	2.0	4.0

**EP 1 002 040 B2**

(continued)

		<b>I</b>	<b>II</b>	<b>III</b>	<b>IV</b>	<b>V</b>	<b>VI</b>
5	Blown Powder						
	Zeolite A	10.0	18.0	14.0	12.0	10.0	10.0
	MA/AA	-	0.5	-	-	-	2.0
	MA/AA 1	7.0	-	-	-	-	-
	AA	-	3.0	3.0	2.0	3.0	3.0
10	Sulfate	5.0	6.3	14.3	11.0	15.0	19.3
	Silicate	10.0	1.0	1.0	1.0	1.0	1.0
	Carbonate	15.0	20.0	10.0	20.7	8.0	6.0
	PEG 4000	0.4	1.5	1.5	1.0	1.0	1.0
15	DTPA	-	0.9	0.5	-	-	0.5
	Brightener 2	0.3	0.2	0.3	-	0.1	0.3
	Spray On						
	C45E7	-	2.0	-	-	2.0	2.0
20	C25E9	3.0	-	-	-	-	-
	C23E9	-	-	1.5	2.0	-	2.0
	Perfume	0.3	0.3	0.3	2.0	0.3	0.3
	Agglomerates						
25	C45AS	-	5.0	5.0	2.0	-	5.0
	LAS	-	2.0	2.0	-	-	2.0
	Zeolite A	-	7.5	7.5	8.0	-	7.5
	Carbonate	-	4.0	4.0	5.0	-	4.0
	PEG 4000	-	0.5	0.5	-	-	0.5
30	Misc (Water etc.)	-	2.0	2.0	2.0	-	2.0
	Dry additives						
	QAS	-	-	-	-	1.0	-
	Citric	-	-	-	-	2.0	-
35	PB4	-	-	-	-	12.0	1.0
	PB1	4.0	1.0	3.0	2.0	-	-
	Percarbonate	-	-	-	-	2.0	10.0
	Carbonate	-	5.3	1.8	-	4.0	4.0
	NOBS	4.0	-	6.0	-	-	0.6
40	Methyl cellulose	0.2	-	-	-	-	-
	Na-SKS-6	8.0	-	-	-	-	-
	STS	-	-	2.0	-	1.0	-
	Culmene sulfonic acid	-	1.0	-	-	-	2.0
45	Oxygenase	0.2	0.02	0.05	0.09	0.1	0.05
	Protease	0.02	0.02	0.02	0.01	0.02	0.02
	Lipase	0.004	-	0.004	-	0.004	0.008
	Amylase	0.003	-	0.002	-	0.003	-
50	Cellulase	0.0005	0.0005	0.0005	0.0007	0.0005	0.0005
	PVPVI	-	-	-	-	0.5	0.1
	PVP	-	-	-	-	0.5	-
	PVNO	-	-	0.5	0.3	-	-
55	QEA	-	-	-	-	1.0	-

# EP 1 002 040 B2

(continued)

Dry additives						
SRP 1	0.2	0.5	0.3	-	0.2	-
Silicone antifoam	0.2	0.4	0.2	0.4	0.1	-
Mg sulfate	-	-	0.2	-	0.2	-
Miscellaneous and minors	Up to 100%					

## Example 5

**[0172]** The following nil bleach-containing detergent compositions of particular use in the washing of coloured clothing were prepared according to the present invention :

	I	II	III
Blown Powder			
Zeolite A	15.0	15.0	-
Sulfate	-	5.0	-
LAS	3.0	3.0	-
DETPMP	0.4	0.5	-
CMC	0.4	0.4	-
MA/AA	4.0	4.0	-
Agglomerates			
C45AS	-	-	11.0
LAS	6.0	5.0	-
TAS	3.0	2.0	-
Silicate	4.0	4.0	-
Zeolite A	10.0	15.0	13.0
CMC	-	-	0.5
MA/AA	-	-	2.0
Carbonate	9.0	7.0	7.0
Spray-on			
Perfume	0.3	0.3	0.5
C45E7	4.0	4.0	4.0
C25E3	2.0	2.0	2.0
Dry additives			
MA/AA	-	-	3.0
Na-SKS-6	-	-	12.0
Citrate	10.0	-	8.0
Bicarbonate	7.0	3.0	5.0
Carbonate	8.0	5.0	7.0
PVPVI/PVNO	0.5	0.5	0.5
Oxygenase	0.0009	0.002	0.005
Protease	0.03	0.02	0.05
Lipase	0.008	0.008	0.008
Amylase	0.01	0.01	0.01
Cellulase	0.001	0.001	0.001

## EP 1 002 040 B2

(continued)

Dry additives			
Silicone antifoam	5.0	5.0	5.0
Sulfate	-	9.0	-
Density (g/litre)	700	700	700
Miscellaneous and minors	Up to 100%		

### Example 6

**[0173]** The following detergent compositions were prepared according to the present invention :

	I	II	III	IV
Base granule				
Zeolite A	30.0	22.0	24.0	10.0
Sulfate	10.0	5.0	10.0	7.0
MA/AA	3.0	-	-	-
AA	-	1.6	2.0	-
MA/AA 1	-	12.0	-	6.0
LAS	14.0	10.0	9.0	20.0
C45AS	8.0	7.0	9.0	7.0
C45AES	-	1.0	1.0	-
Silicate	-	1.0	0.5	10.0
Soap	-	2.0	-	-
Brightener 1	0.2	0.2	0.2	0.2
Carbonate	6.0	9.0	10.0	10.0
PEG 4000	-	1.0	1.5	-
DTPA	-	0.4	-	-
Spray On				
C25E9	-	-	-	5.0
C45E7	1.0	1.0	-	-
C23E9	-	1.0	2.5	-
Perfume	0.2	0.3	0.3	-
Dry additives				
Carbonate	5.0	10.0	18.0	8.0
PVPVI/PVNO	0.5	-	0.3	-
Oxygenase	0.002	0.05	0.0015	0.1
Protease	0.03	0.03	0.03	0.02
Lipase	0.008	-	-	0.008
Amylase	0.002	-	-	0.002
Cellulase	0.0002	0.0005	0.0005	0.0002
NOBS	-	4.0	-	4.5
PB1	1.0	5.0	1.5	6.0
Sulfate	4.0	5.0	-	5.0
SRP 1	-	0.4	-	-
Suds suppressor	-	0.5	0.5	-
Miscellaneous and minors	Up to 100%			

### Example 7

**[0174]** The following granular detergent compositions were prepared according to the present invention :

**EP 1 002 040 B2**

	I	II	III
Blown Powder			
Zeolite A	20.0	-	15.0
STPP	-	20.0	-
Sulfate	-	-	5.0
Carbonate	-	-	5.0
TAS	-	-	1.0
LAS	6.0	6.0	6.0
C68AS	2.0	2.0	-
Silicate	3.0	8.0	-
MA/AA	4.0	2.0	2.0
CMC	0.6	0.6	0.2
Brightener 1	0.2	0.2	0.1
DETPMP	0.4	0.4	0.1
STS	-	-	1.0
Spray On			
C45E7	5.0	5.0	4.0
Silicone antifoam	0.3	0.3	0.1
Perfume	0.2	0.2	0.3
Dry additives			
QEA	-	-	1.0
Carbonate	14.0	9.0	10.0
PB1	1.5	2.0	-
PB4	18.5	13.0	13.0
TAED	2.0	2.0	2.0
QAS	-	-	1.0
Photoactivated bleach	15 ppm	15 ppm	15 ppm
Na-SKS-6	-	-	3.0
Oxygenase	0.01	0.02	0.007
Protease	0.03	0.03	0.007
Lipase	0.004	0.004	0.004
Amylase	0.006	0.006	0.003
Cellulase	0.0002	0.0002	0.0005
Sulfate	10.0	20.0	5.0
Density (g/litre)	700	700	700
Miscellaneous and minors	Up to 100%		

**Example 8**

**[0175]** The following detergent compositions were prepared according to the present invention :

	I	II	III
Blown Powder			
Zeolite A	15.0	15.0	15.0
Sulfate	-	5.0	-
LAS	3.0	3.0	3.0
QAS	-	1.5	1.5
DETPMP	0.4	0.2	0.4

# EP 1 002 040 B2

(continued)

		I	II	III
5	Blown Powder			
	EDDS	-	0.4	0.2
	CMC	0.4	0.4	0.4
	MA/AA	4.0	2.0	2.0
10	Agglomerate			
	LAS	5.0	5.0	5.0
	TAS	2.0	2.0	1.0
	Silicate	3.0	3.0	4.0
15	Zeolite A	8.0	8.0	8.0
	Carbonate	8.0	8.0	4.0
	Spray On			
	Perfume	0.3	0.3	0.3
20	C45E7	2.0	2.0	2.0
	C25E3	2.0	-	-
	Dry Additives			
	Citrate	5.0	-	2.0
25	Bicarbonate	-	3.0	-
	Carbonate	8.0	15.0	10.0
	TAED	6.0	2.0	5.0
	PB1	14.0	7.0	10.0
	PEO	-	-	0.2
30	Bentonite clay	-	-	10.0
	Oxygenase	0.008	0.01	0.009
	Protease	0.03	0.03	0.03
	Lipase	0.008	0.008	0.008
35	Cellulase	0.001	0.001	0.001
	Amylase	0.01	0.01	0.01
	Silicone antifoam	5.0	5.0	5.0
	Sulfate	-	3.0	-
	Density (g/litre)	850	850	850
40	Miscellaneous and minors	Up to 100%		

## Example 9

**[0176]** The following detergent compositions were prepared according to the present invention :

	I	II	III	IV
45	LAS	18.0	14.0	24.0
	QAS	0.7	1.0	0.7
50	TFAA	-	1.0	-
	C23E56.5	-	-	1.0
	C45E7	-	1.0	-
	C45E3S	1.0	2.5	1.0
55	STPP	32.0	18.0	30.0
	Silicate	9.0	5.0	9.0
	Carbonate	11.0	7.5	10.0
	Bicarbonate	-	7.5	-



## EP 1 002 040 B2

(continued)

5

10

15

20

25

	I	II	III	IV
PB1	3.0	1.0	-	-
PB4	-	1.0	-	-
NOBS	2.0	1.0	-	-
DETPMP	-	1.0	-	-
DTPA	0.5	-	0.2	0.3
SRP 1	0.3	0.2	-	0.1
MA/AA	1.0	1.5	2.0	0.5
CMC	0.8	0.4	0.4	0.2
PEI	-	-	0.4	-
Sulfate	20.0	10.0	20.0	30.0
Mg sulfate	0.2	-	0.4	0.9
Oxygenase	0.01	0.01	0.02	0.02
Protease	0.03	0.03	0.02	0.02
Amylase	0.008	0.007	-	0.004
Lipase	0.004	-	0.002	-
Cellulase	0.0003	-	-	0.0001
Photoactivated bleach	30 ppm	20 ppm	-	10 ppm
Perfume	0.3	0.3	0.1	0.2
Brightener 1/2	0.05	0.02	0.08	0.1
Miscellaneous and minors	up to 100%			

### Example 10

30

**[0177]** The following liquid detergent formulations were prepared according to the present invention (Levels are given in parts per weight, enzymes are expressed in pure enzyme) :

35

40

45

50

55

	I	II	III	IV	V
LAS	11.5	8.8	-	3.9	-
C25E2.5S	-	3.0	18.0	-	16.0
C45E2.25S	11.5	3.0	-	15.7	-
C23E9	-	2.7	1.8	2.0	1.0
C23E7	3.2	-	-	-	-
CFAA	-	-	5.2	-	3.1
TPKFA	1.6	-	2.0	0.5	2.0
Citric (50%)	6.5	1.2	2.5	4.4	2.5
Ca formate	0.1 1	0.06	0.1	-	-
Na formate	0.5	0.06	0.1	0.05	0.05
SCS	4.0	1.0	3.0	1.2	-
Borate	0.6	-	3.0	2.0	2.9
Na hydroxide	5.8	2.0	3.5	3.7	2.7
Ethanol	1.75	1.0	3.6	4.2	2.9
1,2 Propanediol	3.3	2.0	8.0	7.9	5.3
Monoethanolamine	3.0	1.5	1.3	2.5	0.8
TEPAE	1.6	-	1.3	1.2	1.2
Oxygenase	0.007	0.005	0.002	0.01	0.01
Protease	0.03	0.01	0.03	0.02	0.02
Lipase	-	-	0.002	-	-
Amylase	-	-	-	0.002	-
Cellulase	-	-	0.0002	0.0005	0.0001

## EP 1 002 040 B2

(continued)

	I	II	III	IV	V
SRP 1	0.2	-	0.1	-	-
DTPA	-	-	0.3	-	-
PVNO	-	-	0.3	-	0.2
Brightener 1	0.2	0.07	0.1	-	-
Silicone antifoam	0.04	0.02	0.1	0.1	0.1
Miscellaneous and water					

### Example 11

**[0178]** The following liquid detergent formulations were prepared according to the present invention (Levels are given in parts per weight, enzymes are expressed in pure enzyme) :

	I	II	III	IV
LAS	10.0	13.0	9.0	-
C25AS	4.0	1.0	2.0	10.0
C25E3S	1.0	-	-	3.0
C25E7	6.0	8.0	13.0	2.5
TFAA	-	-	-	4.5
APA	-	1.4	-	-
TPKFA	2.0	-	13.0	7.0
Citric	2.0	3.0	1.0	1.5
Dodecenyl / tetradecenyl succinic acid	12.0	10.0	-	-
Rapeseed fatty acid	4.0	2.0	1.0	-
Ethanol	4.0	4.0	7.0	2.0
1,2 Propanediol	4.0	4.0	2.0	7.0
Monoethanolamine	-	-	-	5.0
Triethanolamine	-	-	8.0	-
TEPAE	0.5	-	0.5	0.2
DETPMP	1.0	1.0	0.5	1.0
Oxygenase	0.02	0.02	0.009	0.009
Protease	0.02	0.02	0.01	0.008
Lipase	-	0.002	-	0.002
Amylase	0.004	0.004	0.01	0.008
Cellulase	-	-	-	0.002
SRP 2	0.3	-	0.3	0.1
Boric acid	0.1	0.2	1.0	2.0
Ca chloride	-	0.02	-	0.01
Brightener 1	-	0.4	-	-
Suds suppressor	0.1	0.3	-	0.1
Opacifier	0.5	0.4	-	0.3
NaOH up to pH	8.0	8.0	7.6	7.7
Miscellaneous and water				

### Example 12

**[0179]** The following liquid detergent compositions were prepared according to the present invention (Levels are given in parts per weight, enzymes are expressed in pure enzyme) :

	I	II	III	IV
LAS	25.0	-	-	-
C25AS	-	13.0	18.0	15.0
C25E3S	-	2.0	2.0	4.0
C25E7	-	-	4.0	4.0
TFAA	-	6.0	8.0	8.0
APA	3.0	1.0	2.0	-
TPKFA	-	15.0	11.0	11.0
Citric	1.0	1.0	1.0	1.0
Dodecetyl / tetradecyl succinic acid	15.0	-	-	-
Rapeseed fatty acid	1.0	-	3.5	-
Ethanol	7.0	2.0	3.0	2.0
1,2 Propanediol	6.0	8.0	10.0	13.0
Monoethanolamine	-	-	9.0	9.0
TEPAE	-	-	0.4	0.3
DETPMP	2.0	1.2	1.0	-
Oxygenase	0.02	0.04	0.08	0.1
Protease	0.08	0.02	0.01	0.02
Lipase	-	-	0.003	0.003
Amylase	0.004	0.01	0.01	0.01
Cellulase	-	-	0.004	0.003
SRP 2	-	-	0.2	0.1
Boric acid	1.0	1.5	2.5	2.5
Bentonite clay	4.0	4.0	-	-
Brightener 1	0.1	0.2	0.3	-
Suds suppressor	0.4	-	-	-
Opacifier	0.8	0.7	-	-
NaOH up to pH	8.0	7.5	8.0	8.2
Miscellaneous and water				

### Example 13

**[0180]** The following liquid detergent compositions were prepared according to the present invention (Levels are given in parts by weight, enzymes are expressed in pure enzyme) :

	I	II
LAS	27.6	18.9
C45AS	13.8	5.9
C13E8	3.0	3.1
Oleic acid	3.4	2.5
Citric	5.4	5.4
Na hydroxide	0.4	3.6
Ca Formate	0.2	0.1
Na Formate	-	0.5
Ethanol	7.0	-
Monoethanolamine	16.5	8.0
1,2 propanediol	5.9	5.5
Xylene sulfonic acid	-	2.4
TEPAE	1.5	0.8

## EP 1 002 040 B2

(continued)

	I	II
Protease	0.05	0.02
Oxygenase	0.05	0.08
PEG	-	0.7
Brightener 2	0.4	0.1
Perfume	0.5	0.3
Miscellaneous and water		

### Example 14

**[0181]** The following granular fabric detergent compositions which provide "softening through the wash" capability were prepared according to the present invention :

	I	II
C45AS	-	10.0
LAS	7.6	-
C68AS	1.3	
C45E7	4.0	-
C25E3	-	5.0
Coco-alkyl-dimethyl hydroxy-ethyl ammonium chloride	1.4	1.0
Citrate	5.0	3.0
Na-SKS-6	-	11.0
Zeolite A	15.0	15.0
MA/AA	4.0	4.0
DETPMP	0.4	0.4
PB1	15.0	-
Percarbonate	-	15.0
TAED	5.0	5.0
Smectite clay	10.0	10.0
HMWPEO	-	0.1
Oxygenase	0.05	0.05
Protease	0.02	0.01
Lipase	0.02	0.01
Amylase	0.03	0.005
Cellulase	0.001	-
Silicate	3.0	5.0
Carbonate	10.0	10.0
Suds suppressor	1.0	4.0
CMC	0.2	0.1
Water and minors	Up to 100%	

### Example 15

**[0182]** The following rinse added fabric softener composition was prepared according to the present invention :

DEQA (2)	20.0
Oxygenase	0.01
Cellulase	0.001
HCL	0.03

## EP 1 002 040 B2

(continued)

Antifoam agent	0.01
Blue dye	25ppm
CaCl <sub>2</sub>	0.20
Perfume	0.90
Miscellaneous and water	Up to 100%

### Example 16

**[0183]** The following fabric softener and dryer added fabric conditioner compositions were prepared according to the present invention :

	I	II	III	IV	V
DEQA	2.6	19.0	-	-	-
DEQA(2)	-	-	-	-	51.8
DTMAMS	-	-	-	26.0	-
SDASA	-	-	70.0	42.0	40.2
Stearic acid of IV=0	0.3	-	-	-	-
Neodol 45-13	-	-	13.0	-	-
Hydrochloride acid	0.02	0.02	-	-	-
Ethanol	-	-	1.0	-	-
Oxygenase	0.02	0.05	0.07	0.07	0.1
Perfume	1.0	1.0	0.75	1.0	1.5
Glycoperse S-20	-	-	-	-	15.4
Glycerol monostearate	-	-	-	26.0	-
Digeranyl Succinate	-	-	0.38	-	-
Silicone antifoam	0.01	0.01	-	-	-
Electrolyte	-	0.1	-	-	-
Clay	-	-	-	3.0	-
Dye	10ppm	25ppm	0.01	-	-
Water and minors	100%	100%	-	-	-

### Example 17

**[0184]** The following laundry bar detergent compositions were prepared according to the present invention (Levels are given in parts per weight, enzymes are expressed in pure enzyme):

	I	II	III	VI	V	III	VI	V
LAS	-	-	19.0	15.0	21.0	6.75	8.8	-
C28AS	30.0	13.5	-	-	-	15.75	11.2	22.5
Na Laurate	2.5	9.0	-	-	-	-	-	-
Zeolite A	2.0	1.25	-	-	-	1.25	1.25	1.25
Carbonate	20.0	3.0	13.0	8.0	10.0	15.0	15.0	10.0
Ca Carbonate	27.5	39.0	35.0	-	-	40.0	-	40.0
Sulfate	5.0	5.0	3.0	5.0	3.0	-	-	5.0
TSP	5.0	-	-	-	-	5.0	2.5	-
STPP	5.0	15.0	10.0	-	-	7.0	8.0	10.0
Bentonite clay	-	10.0	-	-	5.0	-	-	-
DETPMP	-	0.7	0.6	-	0.6	0.7	0.7	0.7
CMC	-	1.0	1.0	1.0	1.0	-	-	1.0
Talc	-	-	10.0	15.0	10.0	-	-	-

# EP 1 002 040 B2

(continued)

	I	II	III	VI	V	III	VI	V
Silicate	-	-	4.0	5.0	3.0	-	-	-
PVNO	0.02	0.03	-	0.01	-	0.02	-	-
MA/AA	0.4	1.0	-	-	0.2	0.4	0.5	0.4
SRP 1	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Oxygenase	0.001	0.002	0.005	0.005	0.008	0.01	0.01	0.5
Amylase	-	-	0.01	-	-	-	0.002	-
Protease	-	0.004	-	0.003	0.003	-	-	0.003
Lipase	-	0.002	-	0.002	-	-	-	-
Cellulase	-	.0003	-	-	.0003	.0002	-	-
PEO	-	0.2	-	0.2	0.3	-	-	0.3
Perfume	1.0	0.5	0.3	0.2	0.4	-	-	0.4
Mg sulfate	-	-	3.0	3.0	3.0	-	-	-
Brightener	0.15	0.1	0.15	-	-	-	-	0.1
Photoactivated bleach (ppm)	-	15.0	15.0	15.0	15.0	-	-	15.0

## Example 18

**[0185]** The following detergent additive compositions were prepared according to the present invention :

	I	II	III
LAS	-	5.0	5.0
STPP	30.0	-	20.0
Zeolite A	-	35.0	20.0
PB1	20.0	15.0	-
TAED	10.0	8.0	-
Protease	-	0.3	0.3
Amylase	-	0.06	0.06
Oxygenase	0.1	0.1	0.1
Minors, water and miscellaneous	Up to 100%		

## Example 19

**[0186]** The following compact high density (0.96Kg/l) dishwashing detergent compositions were prepared according to the present invention :

	I	II	III	IV	V	VI	VII	VIII
STPP	-	-	54.3	51.4	51.4	-	-	50.9
Citrate	35.0	17.0	-	-	-	46.1	40.2	-
Carbonate	-	17.5	14.0	14.0	14.0	-	8.0	32.1
Bicarbonate	-	-	-	-	-	25.4	-	-
Silicate	32.0	14.8	14.8	10.0	10.0	1.0	25.0	3.1
Metasilicate	-	2.5	-	9.0	9.0	-	-	-
PB1	1.9	9.7	7.8	7.8	7.8	-	-	-
PB4	8.6	-	-	-	-	-	-	-
Percarbonate	-	-	-	-	-	6.7	11.8	4.8
Nonionic	1.5	2.0	1.5	1.7	1.5	2.6	1.9	5.3
TAED	5.2	2.4	-	-	-	2.2	-	1.4
HEDP	-	1.0	-	-	-	-	-	-
DETPMP	-	0.6	-	-	-	-	-	-

## EP 1 002 040 B2

(continued)

	I	II	III	IV	V	VI	VII	VIII
5 MnTACN	-	-	-	-	-	-	0.008	-
PAAC	-	-	0.008	0.01	0.007	-	-	-
BzP	-	-	-	-	1.4	-	-	-
Paraffin	0.5	0.5	0.5	0.5	0.5	0.6	-	-
Oxygenase	0.002	0.004	0.008	0.01	0.02	0.05	0.1	0.2
10 Protease	0.072	0.072	0.029	0.053	0.046	0.026	0.059	0.06
Amylase	0.012	0.012	0.006	0.012	0.013	0.009	0.017	0.03
Lipase	-	0.001	-	0.005	-	-	-	-
BTA	0.3	0.3	0.3	0.3	0.3	-	0.3	0.3
MA/AA	-	-	-	-	-	-	4.2	-
15 480N	3.3	6.0	-	-	-	-	-	0.9
Perfume	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1
Sulphate	7.0	20.0	5.0	2.2	0.8	12.0	4.6	-
Ph	10.8	11.0	10.8	11.3	11.3	9.6	10.8	10.9
20 Miscellaneous and water	Up to 100%							

### Example 20

25 **[0187]** The following granular dishwashing detergent compositions of bulk density 1.02Kg/L were prepared according to the present invention :

	I	II	III	IV	V	VI	VII	VIII
30 STPP	30.0	30.0	33.0	34.2	29.6	31.1	26.6	17.6
Carbonate	30.5	30.5	31.0	30.0	23.0	39.4	4.2	45.0
Silicate	7.4	7.4	7.5	7.2	13.3	3.4	43.7	12.4
Metasilicate	-	-	4.5	5.1	-	-	-	-
Percarbonate	-	-	-	-	-	4.0	-	-
PB1	4.4	4.2	4.5	4.5	-	-	-	-
35 NADCC	-	-	-	-	2.0	-	1.6	1.0
Nonionic	1.2	1.0	0.7	0.8	1.9	0.7	0.6	0.3
TAED	1.0	-	-	-	-	0.8	-	-
PAAC	-	0.004	0.004	0.004	-	-	-	-
40 BzP	-	-	-	1.4	-	-	-	-
Paraffin	0.25	0.25	0.25	0.25	-	-	-	-
Oxygenase	0.007	0.005	0.01	0.02	0.04	0.06	0.08	0.1
Protease	0.036	0.015	0.026	0.028	-	0.01	-	-
Amylase	0.003	0.003	0.006	0.006	-	0.006	-	-
45 Lipase	0.005	-	0.001	-	-	-	-	-
BTA	0.15	0.15	0.15	0.15	-	-	-	-
Perfume	0.2	0.2	0.2	0.2	0.1	0.2	0.2	-
Sulphate	23.4	25.0	22.0	18.5	30.1	19.3	23.1	23.6
50 pH	10.8	10.8	11.3	11.3	10.7	11.5	12.7	10.9
Miscellaneous and water	Up to 100%							

### Example 21

55 **[0188]** The following tablet detergent compositions were prepared according to the present invention by compression of a granular dishwashing detergent composition at a pressure of 13KN/cm<sup>2</sup> using a standard 12 head rotary press:

# EP 1 002 040 B2

	I	II	III	IV	V	VI
STPP	-	48.8	49.2	38.0	-	46.8
Citrate	26.4	-	-	-	31.1	-
Carbonate	-	5.0	14.0	15.4	14.4	23.0
Silicate	26.4	14.8	15.0	12.6	17.7	2.4
Oxygenase	0.05	0.07	0.02	0.01	0.04	0.1
Protease	0.058	0.072	0.041	0.033	0.052	0.013
Amylase	0.01	0.012	0.012	0.007	0.016	0.002
Lipase	0.005	-	-	-	-	-
PB1	1.6	7.7	12.2	10.6	15.7	-
PB4	6.9	-	-	-	-	14.4
Nonionic	1.5	2.0	1.5	1.65	0.8	6.3
PAAC	-	-	0.02	0.009	-	-
MnTACN	-	-	-	-	0.007	-
TAED	4.3	2.5	-	-	1.3	1.8
HEDP	0.7	-	-	0.7	-	0.4
DETPMP	0.65	-	-	-	-	-
Paraffin	0.4	0.5	0.5	0.55	-	-
BTA	0.2	0.3	0.3	0.3	-	-
PA30	3.2	-	-	-	-	-
MA/AA	-	-	-	-	4.5	0.55
Perfume	-	-	0.05	0.05	0.2	0.2
Sulphate	24.0	13.0	2.3	-	10.7	3.4
Weight of tablet	25g	25g	20g	30g	18g	20g
pH	10.6	10.6	10.7	10.7	10.9	11.2
Miscellaneous and water	Up to 100%					

## Example 22

**[0189]** The following liquid dishwashing detergent compositions of density 1.40Kg/L were prepared according to the present invention :

	I	II	III	IV
STPP	17.5	17.5	17.2	16.0
Carbonate	2.0	-	2.4	-
Silicate	5.3	6.1	14.6	15.7
NaOCl	1.15	1.15	1.15	1.25
Polygen/carbopol	1.1	1.0	1.1	1.25
Nonionic	-	-	0.1	-
NaBz	0.75	0.75	-	-
Oxygenase	0.008	0.005	0.01	0.02
NaOH	-	1.9	-	3.5
KOH	2.8	3.5	3.0	-
pH	11.0	11.7	10.9	11.0
Sulphate, miscellaneous and water	up to 100%			

## Example 23

**[0190]** The following liquid rinse aid compositions were prepared according to the present invention :



# EP 1 002 040 B2

	I	II	III
Nonionic	12.0	-	14.5
Nonionic blend	-	64.0	-
Citric	3.2	-	6.5
HEDP	0.5	-	-
PEG	-	5.0	-
Oxygenase	0.01	0.03	0.02
SCS	4.8	-	7.0
Ethanol	6.0	8.0	-
pH of the liquid	2.0	7.5	/

## Example 24

**[0191]** The following liquid dishwashing compositions were prepared according to the present invention :

	I	II	III	IV	V
C17ES	28.5	27.4	19.2	34.1	34.1
Amine oxide	2.6	5.0	2.0	3.0	3.0
C12 glucose amide	-	-	6.0	-	-
Betaine	0.9	-	-	2.0	2.0
Xylene sulfonate	2.0	4.0	-	2.0	-
Neodol C11E9	-	-	5.0	-	-
Polyhydroxy fatty acid amide	-	-	-	6.5	6.5
Sodium diethylene penta acetate (40%)	-	-	0.03	-	-
TAED	-	-	-	0.06	0.06
Sucrose	-	-	-	1.5	1.5
Ethanol	4.0	5.5	5.5	9.1	9.1
Alkyl diphenyl oxide disulfonate	-	-	-	-	2.3
Ca formate	-	-	-	0.5	1.1
Ammonium citrate	0.06	0.1	-	-	-
Na-chloride	-	1.0	-	-	-
Mg chloride	3.3	-	0.7	-	-
Ca chloride	-	-	0.4	-	-
Na sulfate	-	-	0.06	-	-
Mg sulfate	0.08	-	-	-	-
Mg hydroxide	-	-	-	2.2	2.2
Na hydroxide	-	-	-	1.1	1.1
Hydrogen peroxide	200ppm	0.16	0.006	-	-
Oxygenase	0.01	0.01	0.02	0.02	0.2
Protease	0.017	0.005	.0035	0.003	0.002
Perfume	0.18	0.09	0.09	0.2	0.2
Water and minors	Up to 100%				

## Example 25

**[0192]** The following liquid hard surface cleaning compositions were prepared according to the present invention :

	I	II	III
Oxygenase	0.1	0.002	0.01

(continued)

	I	II	III
Amylase	0.01	0.002	0.005
Protease	0.05	0.01	0.02
EDTA*	0.05	0.05	0.05
Citrate	2.9	2.9	2.9
LAS	0.5	0.5	0.5
C12 AS	0.5	0.5	0.5
C12(E)S	0.5	0.5	0.5
C12, 13 E6.5 nonionic	7.0	7.0	7.0
Perfume	1.0	1.0	1.0
Hexyl carbitol**	1.0	1.0	1.0
SCS	1.3	1.3	1.3
Water	Balance to 100%		
*Na4 ethylenediamine diacetic acid			
**Diethylene glycol monohexyl ether			
***All formulas adjusted to pH 7-12			

Example 26

**[0193]** The following spray composition for cleaning of hard surfaces and removing household mildew was prepared according to the present invention :

Oxygenase	0.05
Amylase	0.01
Protease	0.01
Na octyl sulfate	2.0
Na dodecyl sulfate	4.0
Na hydroxide	0.8
Silicate	0.04
Butyl carbitol*	4.0
Perfume	0.35
Water/minors	up to 100%
*Diethylene glycol monobutyl ether	

Example 27

**[0194]** The following lavatory cleansing block compositions were prepared according to the present invention.

	I	II	III
C16-18 fatty alcohol/50EO	80.0	-	-
LAS	-	-	80.0
Nonionic	-	1.0	-
Oleoamide surfactant	-	26.0	-
Partially esterified copolymer of vinylmethyl ether and maleic anhydride, viscosity 0.1-0.5	5.0	-	-
Polyethylene glycol MW 8000	-	39.0	-
Water-soluble K-polyacrylate MW 4000-8000	-	12.0	-
Water-soluble Na-copolymer of acrylamide (70%) and acrylic acid (30%) low MW	-	19.0	-

# EP 1 002 040 B2

(continued)

	I	II	III
Na triphosphate	10.0	-	-
Oxygenase	0.05	0.09	0.15
Dye	2.5	1.0	1.0
Perfume	3.0	-	7.0
KOH / HCL solution	pH 6-11		

## Example 28

**[0195]** The following toilet bowl cleaning composition was prepared according to the present invention.

	I	II
C14-15 linear alcohol 7EO	2.0	10.0
Citric acid	10.0	5.0
Oxygenase	0.05	0.1
DETPMP	-	1.0
Dye	2.0	1.0
Perfume	3.0	3.0
NaOH	pH 6-11	
Water and minors	Up to 100%	

## Example 29

**[0196]** The following single layer effervescent denture cleansing tablets were prepared according to the present invention :

	I	II
Oxygenase	0.1	0.08
Protease	0.05	2.0
Sodium bicarbonate	39.0	39.0
Malic acid	14.0	14.0
Sulphamic acid	3.0	3.0
TAED	2.0	2.0
Dye / Flavor	2.0	2.0
PB1	16.0	16.0
EDTA	3.0	3.0
PEG 10,000	6.0	6.0
K monopersulfate	13.0	13.0
Na carbonate	1.0	1.0
LAS	1.0	1.0
Pyrogenic silica	2.0	2.0

## Example 30

**[0197]** The following dentifrice compositions were prepared according to the present invention :

	I	II	III	IV
Sorbitol (70% aqueous solution)	35.0	35.0	35.0	35.0
PEG-6	1.0	1.0	1.0	1.0

(continued)

	I	II	III	IV
Silica dental abrasive	20.0	20.0	20.0	20.0
Sodium fluoride	0.2	0.2	0.2	0.2
Titanium dioxide	0.5	0.5	0.5	0.5
Sodium saccharin	0.3	0.3	0.3	0.3
Oxygenase	0.05	0.08	0.1	0.2
Protease	0.05	0.1	0.9	2.0
Sodium alkyl sulfate (27.9% aqueous solution)	4.0	4.0	4.0	4.0
Flavor	1.0	1.0	1.0	1.0
Carboxyvinyl polymer	0.3	0.3	0.3	0.3
Carrageenan	0.8	0.8	0.8	0.8
Miscellaneous and water	Up to 100%			

**Example 31**

**[0198]** The following mouthwash compositions were prepared according to the present invention :

	I	II	III	IV
SDA 40 Alcohol	8.0	8.0	8.0	8.0
Flavor	0.08	0.08	0.08	0.08
Emulsifier	0.08	0.08	0.08	0.08
Sodium fluoride	0.05	0.05	0.05	0.05
Glycerin	10.0	10.0	10.0	10.0
Sweetener	0.02	0.02	0.02	0.02
Oxygenase	0.05	0.08	0.1	0.2
Protease	0.01	0.09	0.2	2.0
Benzoic acid	0.05	0.05	0.05	0.05
Sodium hydroxide	0.2	0.2	0.2	0.2
Dye	0.04	0.04	0.04	0.04
Miscellaneous and water	Up to 100%			

**Claims**

1. A cleaning composition comprising a surfactant and a polyphenol and/or heterocyclic substrate based oxygenase selected from the group consisting of

1.13.11.1	CATECHOL 1,2-DIOXYGENASE
1.13.11.3	PROTocatechuate 3,4-DIOXYGENASE
1.13.11.4	GENTISATE 1,2-DIOXYGENASE
1.13.11.5	HOMOGENTISATE 1,2-DIOXYGENASE
1.13.11.6	3-HYDROXYANTHRANILATE 3,4-DIOXYGENASE
1.13.11.8	PROTocatechuate 4,5-DIOXYGENASE
1.13.11.9	2,5-DIHYDROXYPYRIDINE 5,6-DIOXYGENASE
1.13.11.10	7,8-DIHYDROXYKYNURENATE 8,8A-DIOXYGENASE
1.13.11.14	2,3-DIHYDROXYBENZOATE 3,4-DIOXYGENASE
1.13.11.15	3,4-DIHYDROXYPHENYLACETATE 2,3-DIOXYGENASE
1.13.11.16	3-CARBOXYETHYLCATECHOL 2,3-DIOXYGENASE

(continued)

5

10

15

20

25

30

35

40

45

50

55

1.13.11.17	INDOLE 2,3-DIOXYGENASE
1.13.11.22	CAFFEATE 3,4-DIOXYGENASE
1.13.11.23	2,3-DIHYDROXYINDOLE 2,3-DIOXYGENASE
1.13.11.28	2,3-DIHYDROXYBENZOATE 2,3-DIOXYGENASE
1.13.11.35	PYROGALLOL 1,2-OXYGENASE
1.13.11.36	CHLORIDAZON-CATECHOL DIOXYGENASE
1.13.11.37	HYDROXYQUINOL 1,2 DIOXYGENASE
1.13.11.38	1 -HYDROXY-2 NAPHTHOATE 1,2 DIOXYGENASE
1.13.11.39	BIPHENYL-2,3-DIOL 1,2 DIOXYGENASE
1.13.11.41	2,4'-DIHYDROXYACETOPHENONE DIOXYGENASE
1.13.11.42	INDOLEAMINE-PYRROLE 2,3 DIOXYGENASE
1.13.11. 43	LIGNOSTILBENE $\alpha\beta$ -DIOXYGENASE
1.13.12.11	METHYLPHENYL TETRAHYDROPYRIDINE N-MONOOXYGENASE
1.14.11.3	PYRIMIDINE-DEOXYNUCLEOSIDE 2'-DIOXYGENASE
1.14.11.6	THYMINE DIOXYGENASE
1.14.11.9	NARINGENIN 3-DIOXYGENASE
1.14.11.10	PYRIMIDINE-DEOXYNUCLEOSIDE 1'-DIOXYGENASE
1.14.11.11	HYOSCIAMINE (6S)-DIOXYGENASE
1.14.11.12	GIBERELLIN-44 DIOXYGENASE
1.14.11.13	GIBERELLIN 2 $\beta$ -DIOXYGENASE
1.14.11.14	6 $\beta$ -HYDROXYHYOSCIAMINE EPOXIDASE
1.14.11.15	GIBERELLIN 3 $\beta$ -DIOXYGENASE
1.14.12.1	ANTHRANILATE 1,2-DIOXYGENASE (DEAMINATING, DECARBOXYLATING)
1.14.12.2	ANTHRANILATE 2,3-DIOXYGENASE
1.14.12.4	3-HYDROXY-2-METHYLPYRIDINECARBOXYLATE DIOXYGENASE
1.14.12.5	5-PYRIDOXATE DIOXYGENASE
1.14.12.6	2-HYDROXYCYCLOHEXANONE 2-MONOOXYGENASE
1.14.12.7	PHTALATE 4,5 DIOXYGENASE
1.14.12.10	BENZOATE 1,2-DIOXYGENASE
1.14.12.11	TOLUENE DIOXYGENASE
1.14.12.12	NAPHTALENE 1,2 DIOXYGENASE
1.14.13.1	SALICYLATE 1-MONOOXYGENASE
1.14.13.2	4-HYDROXYBENZOATE 3-MONOOXYGENASE /HYDROLASE
1.14.13.3	4-HYDROXYPHENYLACETATE 3-MONOOXYGENASE
1.14.13.4	MELILOTATE 3-MONOOXYGENASE
1.14.13.5	IMIDAZOLACETATE 4-MONOOXYGENASE
1.14.13.6	ORCINOL 2-MONOOXYGENASE
1.14.13.8	DIMETHYLANILINE MONOOXYGENASE (N-OXIDE-FORMING)
1.14.13.9	KYNURENINE 3-MONOOXYGENASE
1.14.13.10	2,6-DIHYDROXYPYRIDINE 3-MONOOXYGENASE
1.14.13.11	TRANS-CINNAMATE 4-MONOOXYGENASE
1.14.13.12	BENZOATE 4-MONOOXYGENASE
1.14.13.14	TRANS-CINNAMATE 2-MONOOXYGENASE
1.14.13.16	CYCLOPENTANONE MONOOXYGENASE
1.14.13.18	4-PHENYLHYDROXYACETATE 1-MONOOXYGENASE
1.14.13.21	FLAVONOID 3'-MONOOXYGENASE
1.14.13.22	CYCLOHEXANONE MONOOXYGENASE
1.14.13.23	3-HYDROXYBENZOATE 4-MONOOXYGENASE
1.14.13.24	3-HYDROXYBENZOATE 6-MONOOXYGENASE
1.14.13.27	4-AMINOBENZOATE 1-MONOOXYGENASE
1.14.13. 29	4-NITROPHENOL 2-MONOOXYGENASE

(continued)

1.14.13.31	2-NITROPHENOL 2-MONOOXYGENASE
1.14.13.33	4-HYDROXYBENZOATE 3-MONOOXYGENASE
1.14.13.35	ANTHRANILATE 3-MONOOXYGENASE
1.14.13.36	5-O-(4-COUMAROYL)-D-QUINATE 3'-MONOOXYGENASE
1.14.13.38	ANHYDROTETRACYCLINE MONOOXYFENASE
1.14.13.40	ANTHRANILOYL-CoA MONOOXYGENASE
1.14.13.43	QUESTIN MONOOXYGENASE
1.14.13.44	2-HYDROXYBIPHENYL 3-MONOOXYGENASE
1.14.13.46	(-)-MENTHOL MONOOXYGENASE
1.14.14.3	ALKANAL MONOOXYGENASE (FMN-LINKED) /BACTERIAL LUCIFERASE
1.14.15.2	CAMPBOR 1,2-MONOOXYGENASE
1.14.16.3	ANTHRANILATE 3-MONOOXYGENASE
1.14.16.6	MANDELATE 4-MONOOXYGENASE
1.14.17.1	DOPAMINE B-MONOOXYGENASE /DOPAMINE B-HYDROXYLASE
1.14.99.2	KYNURENINE 7,8-HYDROXYLASE
1.14.99.15	4-METHOXYBENZOATE MONOOXYGENASE (O-DEMETHYLATING)
1.14.99.20	PHYLLLOQUINONE MONOOXYGENASE (2,3-EPOXIDIZING)
1.14.99.23	3-HYDROXYBENZOATE 2-MONOOXYGENASE, and/or mixtures thereof.

2. A cleaning composition according to claim 1 wherein said polyphenol and/or heterocyclic substrate based oxygenase is further **characterised by** being an iron sulphur or iron heme oxygenase and/or a heavy metal dependant oxygenase.

3. A cleaning composition according to claims 1-2 wherein said polyphenol and/or heterocyclic substrate based oxygenase is present at a level of from 0.0001% to 2%, preferably from 0.001% to 0.5%, more preferably from 0.002% to 0.2% pure enzyme by weight of total composition.

4. A cleaning composition according to claims 1 to 3 wherein said polyphenol and/or heterocyclic substrate based oxygenase is a decyclising and/or hydroxylating oxygenase.

5. A cleaning composition according claims 1 to 4 wherein said said polyphenol and/or heterocyclic substrate based oxygenase is selected from

- (a) PROTOCATECHUATE 3,4-DIOXYGENASE
- (b) 2,3-DIHYDROXYBENZOATE 3,4-DIOXYGENASE
- (c) INDOLE 2,3-DIOXYGENASE
- (d) CAFFEATE 3,4-DIOXYGENASE
- (f) PYROGALLOL 1,2-OXYGENASE
- (g) NARINGENIN 3-DIOXYGENASE
- (h) PHTALATE 4,5 DIOXYGENASE
- (i) BENZOATE 1,2-DIOXYGENASE
- (j) TOLUENE DIOXYGENASE
- (k) 4-HYDROXYBENZOATE 3-MONOOXYGENASE /- HYDROXYLASE
- (l) BENZOATE 4-MONOOXYGENASE
- (m) FLAVONOID 3'-MONOOXYGENASE

and/or mixtures thereof.

6. A cleaning composition according to any of the preceding claims wherein said polyphenol and/or heterocyclic substrate based oxygenase is alkaline.

7. A cleaning composition according to any of the preceding claims further comprising a cofactor.

8. A cleaning composition according to claim 7 wherein the cofactor is comprised at a weight ratio of pure polyphenol

and/or heterocyclic substrate based oxygenase to cofactor between 10:1 to 1:10, preferably 5:1 to 1:8, more preferably between 1:2 to 1:5.

9. A cleaning composition according to any of the preceding claims further comprising a detergent enzyme, preferably selected from cellulase, lipase, protease, amylase and/or mixtures thereof.

10. A cleaning composition according to any of the preceding claims further comprising another bleach system.

11. A cleaning composition according to claim 10 wherein said bleach system is a conventional activated bleach system.

12. A cleaning composition according to claim 11 wherein the bleaching agent is selected from perborate and/or percarbonate and the activator selected from tetraacetythylenediamine, nonanoyloxybenzenesulfonate and/or 3,5,-trimethylhexanolxybenzenesulfonate.

13. A cleaning composition according to claim 10 wherein said bleach system is another enzymatic bleach system.

14. A cleaning composition according to claim 10 wherein said bleach system is a metallo catalyst based bleach system.

15. A cleaning composition according to claim 14 wherein said metallo catalyst is a transition metal complex of a macropolycyclic rigid ligand.

16. A cleaning composition according to claims 14-15 wherein said metallo catalyst is manganese.

17. A cleaning composition according to any of the preceding claims which is in the form of an additive.

18. A fabric softening composition comprising a polyphenol and/or heterocyclic substrate based oxygenase as defined in claim 1 and a cationic surfactant comprising two long chain lengths of  $C_{11}$ - $C_{22}$ .

19. Use of a polyphenol and/or heterocyclic substrate based oxygenase as defined in claim 1 in a cleaning and/or softening composition for fabric cleaning and/or fabric stain removal and/or fabric whiteness maintenance and/or fabric softening and/or fabric colour appearance and/or fabric dye transfer inhibition.

20. Use of a polyphenol and/or heterocyclic substrate based oxygenase as defined in claim 1 in a cleaning composition for cleaning hard surfaces such as floors, walls, bathroom tiles and the like.

21. Use of a polyphenol and/or heterocyclic substrate based oxygenase as defined in claim 1 in a cleaning composition for hand and machine dishwashing.

22. Use of a polyphenol and/or heterocyclic substrate based oxygenase as defined in claim 1 in a cleaning composition for oral and/or dental applications.

23. Use of a polyphenol and/or heterocyclic substrate based oxygenase as defined in claim 1 in a cleaning and/or softening composition for the sanitisation of the treated surfaces.

## Patentansprüche

1. Reinigungszusammensetzung, umfassend ein Tensid und eine auf Polyphenol und/oder heterocyclischem Substrat basierende Oxygenase, die ausgewählt ist aus der Gruppe bestehend aus:

1.13.11.1 CATECHIN-1,2-DIOXYGENASE

1.13.11.3 PROTOCATECHUAT-3,4-DIOXYGENASE

1.13.11.4 GENTISAT-1,2-DIOXYGENASE

## EP 1 002 040 B2

(fortgesetzt)

	1.13.11.5	HOMOGENISAT-1,2-DIOXYGENASE
5	1.13.11.6	3-HYDROXYANTHRANILAT-3,4-DIOXYGENASE
	1.13.11.8	PROTOKATECHUAT-4,5-DIOXYGENASE
10	1.13.11.9	2,5-DIHYDROXYPYRIDIN-5,6-DIOXYGENASE
	1.13.11.10	7,8-DIHYDROXYKYNURENAT-8,8a-DIOXYGENASE
	1.13.11.14	2,3-DIHYDROXYBENZOAT-3,4-DIOXYGENASE
15	1.13.11.15	3,4-DIHYDROXYPHENYLACETAT-2,3-DIOXYGENASE
	1.13.11.16	3-CARBOXYETHYLCATECHIN-2,3-DIOXYGENASE
20	1.13.11.17	INDOL-2,3-DIOXYGENASE
	1.13.11.22	KAFFEAT-3,4-DIOXYGENASE
25	1.13.11.23	2,3-DIHYDROXYINDOL-2,3-DIOXYGENASE
	1.13.11.28	2,3-DIHYDROXYBENZOAT-2,3-DIOXYGENASE
	1.13.11.35	PYROGALLOL-1,2-OXYGENASE
30	1.13.11.36	CHLORIDAZON-CATECHIN-DIOXYGENASE
	1.13.11.37	HYDROXYQUINOL-1,2-DIOXYGENASE
35	1.13.11.38	1-HYDROXY-2-NAPHTHOAT-1,2-DIOXYGENASE
	1.13.11.39	BIPHENYL-2,3-DIOL-1,2-DIOXYGENASE
40	1.13.11.41	2,4'-DIHYDROXYACETOPHENON-DIOXYGENASE
	1.13.11.42	INDOLEAMIN-PYRROL-2,3-DIOXYGENASE
45	1.13.11.43	LIGNOSTILBEN $\alpha\beta$ -DIOXYGENASE
	1.13.12.11	METHYLPHENYL-TETRAHYDROPYRIDIN-N-MONOOXYGENASE
50	1.14.11.3	PYRIMIDIN-DEOXYNUCLEOSID-2'-DIOXYGENASE
	1.14.11.6	THYMIN-DIOXYGENASE
	1.14.11.9	NARINGENIN-3-DIOXYGENASE
55	1.14.11.10	PYRIMIDIN-DEOXYNUCLEOSID-1'-DIOXYGENASE
	1.14.11.11	HYOSCYAMIN-(6S)-DIOXYGENASE



(fortgesetzt)

5	1.14.11.12	GIBERELLIN-44 DIOXYGENASE
	1.14.11.13	GIBERELLIN-2 $\beta$ -DIOXYGENASE
	1.14.11.14	6 $\beta$ -HYDROXYHYOSCYAMIN-EPOXIDASE
10	1.14.11.15	GIBERELLIN-3 $\beta$ -DIOXYGENASE
	1.14.12.1	ANTHRANILAT-1,2-DIOXYGENASE (DESAMINIEREND, DECARBOXYLIEREND)
15	1.14.12.2	ANTHRANILAT-2,3-DIOXYGENASE
	1.14.12.4	3-HYDROXY-2-METHYLPYRIDINCARBOXYLAT-DIOXYGENASE
20	1.14.12.5	5-PYRIDOXAT-DIOXYGENASE
	1.14.12.6	2-HYDROXYCYCLOHEXANON-2-MONOOXYGENASE
	1.14.12.7	PHTALAT-4,5-DIOXYGENASE
25	1.14.12.10	BENZOAT-1,2-DIOXYGENASE
	1.1.4.12.11	TOLUOL-DIOXYGENASE
30	1.14.12.12	NAPHTALEN-1,2 DIOXYGENASE
	1.14.13.1	SALICYLAT-1-MONOOXYGENASE
35	1.14.13.2	4-HYDROXYBENZOAT-3-MONOOXYGENASE / -HYDROLASE
	1.14.13.3	4-HYDROXYPHENYLACETAT-3-MONOOXYGENASE
40	1.14.13.4	MELILOTAT-3-MONOOXYGENASE
	1.14.13.5	IMIDAZOLACETAT-4-MONOOXYGENASE
	1.14.13.6	ORCINOL-2-MONOOXYGENASE
45	1.14.13.8	DIMETHYLANILIN-MONOOXYGENASE (N-OXID-BILDEND)
	1.14.13.9	KYNURENIN-3-MONOOXYGENASE
50	1.14.13.10	2,6-DIHYDROXYPYRIDIN-3-MONOOXYGENASE
	1.14.13.11	TRANS-CINNAMAT-4-MONOOXYGENASE
55	1.14.13.12	BENZOAT-4-MONOOXYGENASE
	1.14.13.14	TRANS-CINNAMAT-2-MONOOXYGENASE

## EP 1 002 040 B2

(fortgesetzt)

	1.14.13.16	CYCLOPENTANON-MONOOXYGENASE
5	1.14.13.18	4-PHENYLHYDROXYACETAT-1-MONOOXYGENASE
	1.14.13.21	FLAVONOID-3'-MONOOXYGENASE
10	1.14.13.22	CYCLOHEXANON-MONOOXYGENASE
	1.14.13.23	3-HYDROXYBENZOAT-4-MONOOXYGENASE
	1.14.13.24	3-HYDROXYBENZOAT-6-MONOOXYGENASE
15	1.14.13.27	4-AMINOBENZOAT-1-MONOOXYGENASE
	1.14.13.29	4-NITROPHENOL-2-MONOOXYGENASE
20	1.14.13.31	2-NITROPHENOL-2-MONOOXYGENASE
	1.14.13.33	4-HYDROXYBENZOAT-3-MONOOXYGENASE
25	1.14.13.35	ANTHRANILAT-3-MONOOXYGENASE
	1.14.13.36	5-O-(4-COUMAROYL)-D-QUINAT-3'-MONOOXYGENASE
30	1.14.13.38	ANHYPDROTETRACYCLIN-MONOOXYFENASE
	1.14.13.40	ANTHRANILOYL-CoA-MONOOXYGENASE
	1.14.13.43	QUESTIN-MONOOXYGENASE
35	1.14.13.44	2-HYDROXYBIPHENYL-3-MONOOXYGENASE
	1.14.13.46	(-)-MENTHOL-MONOOXYGENASE
40	1.14.14.3	ALKANAL-MONOOXYGENASE (FMN-VERBUNDEN) / BAKTERIELLE LUCIFERASE
	1.14.15.2	KAMPFER-1,2-MONOOXYGENASE
45	1.14.16.3	ANTHRANILAT-3-MONOOXYGENASE
	1.14.16.6	MANDELAT-4-MONOOXYGENASE
50	1.14.17.1	DOPAMIN-B-MONOOXYGENASE /DOPAMIN-B-HYDROXYLASE
	1.14.99.2	KYNURENIN-7,8-HYDROXYLASE
	1.14.99.15	4-METHOXYBENZOAT-MONOOXYGENASE (O-DEMETHYLIEREND)
55	1.14.99.20	PHYLLOQUINON-MONOOXYGENASE (2,3-EPOXIDIEREND)
	1.14.99.23	3-HYDROXYBENZOAT-2-MONOOXYGENASE und/oder Mischungen davon.

2. Reinigungszusammensetzung nach Anspruch 1, wobei die auf Polyphenol und/oder heterocyclischem Substrat basierende Oxygenase ferner **dadurch gekennzeichnet ist, dass** sie eine Eisen-Schwefel- oder Eisen-Häm-Oxygenase und/oder eine schwermetallabhängige Oxygenase ist.

3. Reinigungszusammensetzung nach Anspruch 1 bis 2, wobei die auf Polyphenol und/oder heterocyclischem Substrat basierende Oxygenase bezogen auf das Gesamtgewicht der Zusammensetzung in einer Menge von 0,0001 Gew.-% bis 2 Gew.-%, vorzugsweise von 0,001 Gew.-% bis 0,5 Gew.-%, mehr bevorzugt von 0,002 Gew.-% bis 0,2 Gew.-% reinem Enzym vorhanden ist.

4. Reinigungszusammensetzung nach Anspruch 1 bis 3, wobei die auf Polyphenol und/oder heterocyclischem Substrat basierende Oxygenase eine decyclisierende und/oder dehydroxylierende Oxygenase ist.

5. Reinigungszusammensetzung nach Anspruch 1 bis 4, wobei die auf Polyphenol und/oder heterocyclischem Substrat basierende Oxygenase ausgewählt ist aus:

- (a) PROTOCATECHUAT-3,4-DIOXYGENASE
- (b) 2,3-DIHYDROXYBENZOAT-3,4-DIOXYGENASE
- (c) INDOL-2,3-DIOXYGENASE
- (d) KAFFEAT-3,4-DIOXYGENASE
- (f) PYROGALLOL-1,2-OXYGENASE
- (g) NARINGENIN-3-DIOXYGENASE
- (h) PHTALAT-4,5-DIOXYGENASE
- (i) BENZOAT-1,2-DIOXYGENASE
- (j) TOLUOL-DIOXYGENASE
- (k) 4-HYDROXYBENZOAT-3-MONOOXYGENASE /-HYDROXYLASE
- (l) BENZOAT-4-MONOOXYGENASE
- (m) FLAVONOID-3'-MONOOXYGENASE

und/oder Mischungen davon.

6. Reinigungszusammensetzung nach einem der vorstehenden Ansprüche, wobei die auf Polyphenol und/oder heterocyclischem Substrat basierende Oxygenase alkalisch ist.

7. Reinigungszusammensetzung nach einem der vorstehenden Ansprüche, die ferner einen Cofaktor umfasst.

8. Reinigungszusammensetzung nach Anspruch 7, wobei der Cofaktor in einem Gewichtsverhältnis von reiner, auf Polyphenol und/oder heterocyclischem Substrat basierender Oxygenase zu Cofaktor zwischen 10:1 bis 1:10, vorzugsweise 5:1 bis 1:8, mehr bevorzugt zwischen 1:2 bis 1:5 enthalten ist.

9. Reinigungszusammensetzung nach einem der vorstehenden Ansprüche, die ferner ein Reinigungsenzym umfasst, das vorzugsweise aus Cellulase, Lipase, Protease, Amylase und/oder Mischungen davon ausgewählt ist.

10. Reinigungszusammensetzung nach einem der vorstehenden Ansprüche, die ferner ein weiteres Bleichsystem umfasst.

11. Reinigungszusammensetzung nach Anspruch 10, wobei das Bleichsystem ein herkömmliches aktiviertes Bleichsystem ist.

12. Reinigungszusammensetzung nach Anspruch 11, wobei das Bleichmittel aus Perborat und/oder Percarbonat ausgewählt ist und der Aktivator aus Tetraacetylenylendiamin, Nonanoyloxybenzolsulfonat und/oder 3,5,-Trimethylhexanoyloxybenzolsulfonat ausgewählt ist.

13. Reinigungszusammensetzung nach Anspruch 10, wobei das Bleichsystem ein weiteres enzymatisches Bleichsystem ist.

14. Reinigungszusammensetzung nach Anspruch 10, wobei das Bleichsystem ein Bleichsystem auf Metallkatalysatorbasis ist.

15. Reinigungszusammensetzung nach Anspruch 14, wobei der Metallkatalysator ein Übergangsmetallkomplex eines makropolycyclischen unnachgiebigen Liganden ist.
16. Reinigungszusammensetzung nach Ansprüchen 14-15, wobei der Metallkatalysator Mangan ist.
17. Reinigungszusammensetzung nach einem der vorstehenden Ansprüche, die in der Form eines Zusatzes ist.
18. Stoffweichmacherzusammensetzung, umfassend eine auf Polyphenol und/oder heterocyclischem Substrat basierende Oxygenase nach Anspruch 1 und ein kationisches Tensid, das zwei langkettige Längen von C<sub>11</sub>-C<sub>22</sub> umfasst.
19. Verwendung einer auf Polyphenol und/oder heterocyclischem Substrat basierenden Oxygenase nach Anspruch 1 in einer Reinigungs- und/oder Weichmacherzusammensetzung zur Stoffreinigung und/oder Stofffleckenentfernung und/oder Stoffweißbeständigkeit und/oder Weichmachen von Stoff und/oder Stofffarberscheinung und/oder Hemmung der Farbstoffübertragung von Stoff.
20. Verwendung einer auf Polyphenol und/oder heterocyclischem Substrat basierenden Oxygenase nach Anspruch 1 in einer Reinigungszusammensetzung zum Reinigen von harten Oberflächen wie Fußböden, Wänden, Badezimmerfliesen und dergleichen.
21. Verwendung einer auf Polyphenol und/oder heterocyclischem Substrat basierenden Oxygenase nach Anspruch 1 in einer Reinigungszusammensetzung zum Geschirrspülen per Hand und zum maschinellen Geschirrspülen.
22. Verwendung einer auf Polyphenol und/oder heterocyclischem Substrat basierenden Oxygenase nach Anspruch 1 in einer Reinigungszusammensetzung für orale und/oder dentale Anwendungen.
23. Verwendung einer auf Polyphenol und/oder heterocyclischem Substrat basierenden Oxygenase nach Anspruch 1 in einer Reinigungs- und/oder Weichmacherzusammensetzung zur Desinfektion der behandelten Oberflächen.

## Revendications

1. Composition de nettoyage comprenant un agent tensioactif et une oxygénase basée sur un substrat polyphénol et/ou hétérocyclique choisi dans le groupe constitué de

- |            |  |
|------------|--|
| 1.13.11.1  | CATÉCHOL 1,2-DIOXYGÉNASE                   |
| 1.13.11.3  | PROCATÉCHUATE 3,4-DIOXYGÉNASE              |
| 1.13.11.4  | GENTISATE 1,2-DIOXYGÉNASE                  |
| 1.13.11.5  | HOMOGENTISATE 1,2-DIOXYGÉNASE              |
| 1.13.11.6  | 3-HYDROXYANTHRANILATE 3,4-DIOXYGÉNASE      |
| 1.13.11.8  | PROCATÉCHUATE 4,5-DIOXYGÉNASE              |
| 1.13.11.9  | 2,5-DIHYDROXYPYRIDINE 5,6-DIOXYGÉNASE      |
| 1.13.11.10 | 7,8-DIHYDROXYKYNURÉNATE 8,8A-DIOXYGÉNASE   |
| 1.13.11.14 | 2,3-DIHYDROXYBENZOATE 3,4-DIOXYGÉNASE      |
| 1.13.11.15 | 3,4-DIHYDROXYPHÉNYLACÉTATE 2,3-DIOXYGÉNASE |
| 1.13.11.16 | 3-CARBOXYÉTHYLCATÉCHOL 2,3-DIOXYGÉNASE     |
| 1.13.11.17 | INDOLE 2,3-DIOXYGÉNASE                     |
| 1.13.11.22 | CAFFÉATE 3,4-DIOXYGÉNASE                   |
| 1.13.11.23 | 2,3-DIHYDROXYINDOLE 2,3-DIOXYGÉNASE        |
| 1.13.11.28 | 2,3-DIHYDROXYBENZOATE 2,3-DIOXYGÉNASE      |
| 1.13.11.35 | PYROGALLOL 1,2-OXYGÉNASE                   |
| 1.13.11.36 | CHLORIDAZON-CATÉCHOL DIOXYGÉNASE           |
| 1.13.11.37 | HYDROXYQUINOL 1,2 DIOXYGÉNASE              |
| 1.13.11.38 | 1-HYDROXY-2 NAPHTOATE 1,2 DIOXYGÉNASE      |
| 1.13.11.39 | BIPHÉNYL-2,3-DIOL 1,2 DIOXYGÉNASE          |

(suite)

	1.13.11.41	2,4'-DIHYDROXYACÉTOPHÉNONE DIOXYGÉNASE
	1.13.11.42	INDOLEAMINE-PYRROLE 2,3 DIOXYGÉNASE
5	1.13.11. 43	LIGNOSTILBÈNE $\alpha\beta$ -DIOXYGÉNASE
	1.13.12.11	MÉTHYLPHÉNYLTÉTRAHYDROPYRIDINE N-MONO-OXYGÉNASE
	1.14.11.3	PYRIMIDINE-DÉOXYNUCLÉOSIDE 2'-DIOXYGÉNASE
	1.14.11.6	THYMINÉ DIOXYGÉNASE
10	1.14.11.9	NARINGÉNINE 3-DIOXYGÉNASE
	1.14.11.10	PYRIMIDINE-DÉOXYNUCLÉOSIDE 1'-DIOXYGÉNASE
	1.14.11.11	HYOSCIAMINE (6S)-DIOXYGÉNASE
	1.14.11.12	GIBÉRELLIN-44 DIOXYGÉNASE
	1.14.11.13	GIBÉRELLIN 2 $\beta$ -DIOXYGÉNASE
15	1.14.11.14	6 $\beta$ -HYDROXYHYOSCIAMINE ÉPOXIDASE
	1.14.11.15	GIBÉRELLIN 3 $\beta$ -DIOXYGÉNASE
	1.14.12.1	ANTHRANILATE 1,2-DIOXYGÉNASE (À DÉSAMINATION, DÉCARBOXYLATION)
	1.14.12.2	ANTHRANILATE 2,3-DIOXYGÉNASE
20	1.14.12.4	3-HYDROXY-2-MÉTHYLPYRIDINE CARBOXYLATE DIOXYGÉNASE
	1.14.12.5	5-PYRIDOXATE DIOXYGÉNASE
	1.14.12.6	2-HYDROXYCYCLOHEXANONE 2-MONO-OXYGÉNASE
	1.14.12.7	PHTALATE 4,5 DIOXYGÉNASE
	1.14.12.10	BENZOATE 1,2-DIOXYGÉNASE
25	1.14.12.11	TOLUÈNE DIOXYGÉNASE
	1.14.12.12	NAPHTALÈNE 1,2 DIOXYGÉNASE
	1.14.13.1	SALICYLATE 1-MONO-OXYGÉNASE
	1.14.13.2	4-HYDROXYBENZOATE 3-MONO-OXYGÉNASE /HYDROLASE
	1.14.13.3	4-HYDROXYPHENYLACÉTATE 3-MONO-OXYGÉNASE
30	1.14.13.4	MÉLILOTATE 3-MONO-OXYGÉNASE
	1.14.13.5	IMIDAZOLACÉTATE 4-MONO-OXYGÉNASE
	1.14.13.6	ORCINOL 2-MONO-OXYGÉNASE
	1.14.13.8	DIMÉTHYLANILINE MONO-OXYGÉNASE (À FORMATION DE N-OXYDE)
35	1.14.13.9	KYNURÉNINE 3-MONO-OXYGÉNASE
	1.14.13.10	2,6-DIHYDROXYPYRIDINE 3-MONO-OXYGÉNASE
	1.14.13.11	TRANS-CINNAMATE 4-MONO-OXYGÉNASE
	1.14.13.12	BENZOATE 4-MONO-OXYGÉNASE
	1.14.13.14	TRANS-CINNAMATE 2-MONO-OXYGÉNASE
40	1.14.13.16	CYCLOPENTANONE MONO-OXYGÉNASE
	1.14.13.18	4-PHÉNYLHYDROXYACÉTATE 1-MONO-OXYGÉNASE
	1.14.13.21	FLAVONOÏDE 3'-MONO-OXYGÉNASE
	1.14.13.22	CYCLOHEXANONE MONO-OXYGÉNASE
45	1.14.13.23	3-HYDROXYBENZOATE 4-MONO-OXYGÉNASE
	1.14.13.24	3-HYDROXYBENZOATE 6-MONO-OXYGÉNASE
	1.14.13.27	4-AMINOBENZOATE 1-MONO-OXYGÉNASE
	1.14.13.29	4-NITROPHÉNOL 2-MONO-OXYGÉNASE
	1.14.13.31	2-NITROPHÉNOL 2-MONO-OXYGÉNASE
50	1.14.13.33	4-HYDROXYBENZOATE 3-MONO-OXYGÉNASE
	1.14.13.35	ANTHRANILATE 3-MONO-OXYGÉNASE
	1.14.13.36	5-O-(4-COUMAROYL)-D-QUINATE 3'-MONO-OXYGÉNASE
	1.14.13.38	ANHYDROTÉTRACYCLINE MONO-OXYGÉNASE
55	1.14.13. 40	ANTHRANILOYL-CoA MONO-OXYGÉNASE
	1.14.13.43	QUESTIN MONO-OXYGÉNASE
	1.14.13.44	2-HYDROXYBIPHÉNYL 3-MONO-OXYGÉNASE
	1.14.13.46	(-)-MENTHOL MONO-OXYGÉNASE

(suite)

- 5 1.14.14.3 ALCANAL MONO-OXYGÉNASE (LIÉ À FMN) /LUCIFÉrase BACTÉRIENNE  
 1.14.15.2 CAMPHRE 1,2-MONO-OXYGÉNASE  
 1.14.16.3 ANTHRANILATE 3-MONO-OXYGÉNASE  
 1.14.16.6 MANDÉLATE 4-MONO-OXYGÉNASE  
 1.14.17.1 DOPAMINE B-MONO-OXYGÉNASE /DOPAMINE B-HYDROXYLASE  
 1.14.99.2 KYNURÉNINE 7,8-HYDROXYLASE  
 10 1.14.99.15 4-MÉTHOXYBENZOATE MONO-OXYGÉNASE (O-DÉMÉTHYLATION)  
 1.14.99.20 PHYLLOQUINONE MONO-OXYGÉNASE (2,3-EPOXIDIZING)  
 1.14.99.23 3-HYDROXYBENZOATE 2-MONO-OXYGÉNASE, et/ou leurs mélanges.
- 15 2. Composition de nettoyage selon la revendication 1, dans lequel ladite oxygénase à base de substrat polyphénol et/ou hétérocyclique est en outre **caractérisée en ce qu'**elle est une oxygénase de soufre ferrique ou d'hème ferrique et/ou une oxygénase dépendant d'un métal lourd.
- 20 3. Composition de nettoyage selon les revendications 1 à 2, dans laquelle ladite oxygénase à base de substrat polyphénol et/ou hétérocyclique est présente à un taux allant de 0,0001 % à 2 %, de préférence de 0,001 % à 0,5 %, plus préférentiellement de 0,002 % à 0,2 % d'enzyme pure en poids de composition totale.
4. Composition de nettoyage selon les revendications 1 à 3, dans lequel ladite oxygénase à base de substrat polyphénol et/ou hétérocyclique est une oxygénase à décyclisation et/ou déshydroxylation.
- 25 5. Composition de nettoyage selon la revendication 1 à 4, dans lequel ladite oxygénase à base de substrat polyphénol et/ou hétérocyclique est choisie parmi :
- 30 (a) PROTOCATÉCHUATE 3,4-DIOXYGÉNASE  
 (b) 2,3-DIHYDROXYBENZOATE 3,4-DIOXYGÉNASE  
 (c) INDOLE 2,3-DIOXYGÉNASE  
 (d) CAFFÉATE 3,4-DIOXYGÉNASE  
 (f) PYROGALLOL 1,2-OXYGÉNASE  
 (g) NARINGÉNIN 3-DIOXYGÉNASE  
 (h) PHTALATE 4,5 DIOXYGÉNASE  
 35 (i) BENZOATE 1,2-DIOXYGÉNASE  
 (j) TOLUÈNE DIOXYGÉNASE  
 (k) 4-HYDROXYBENZOATE 3-MONO-OXYGÉNASE /-HYDROXYLASE  
 (l) BENZOATE 4-MONO-OXYGÉNASE  
 (m) FLAVONOÏDE 3'-MONO-OXYGÉNASE  
 40 et/ou leurs mélanges.
- 45 6. Composition de nettoyage selon l'une quelconque des revendications précédentes, dans laquelle ladite oxygénase à base de substrat polyphénol et/ou hétérocyclique est alcaline.
7. Composition de nettoyage selon l'une quelconque des revendications précédentes, comprenant en outre un cofacteur.
- 50 8. Composition de nettoyage selon la revendication 7, dans laquelle le cofacteur est compris à un rapport pondéral d'oxygénase à base de substrat polyphénol et/ou hétérocyclique pure sur cofacteur compris entre 10:1 et 1:10, de préférence 5:1 à 1:8, plus préférentiellement entre 1:2 à 1:5.
9. Composition de nettoyage selon l'une quelconque des revendications précédentes, comprenant en outre une enzyme détergente, de préférence choisie parmi la cellulase, la lipase, la protéase, l'amylase et/ou leurs mélanges.
- 55 10. Composition de nettoyage selon l'une quelconque des revendications précédentes, comprenant en outre un autre système de blanchiment.

11. Composition de nettoyage selon la revendication 10, dans laquelle ledit système de blanchiment est un système de blanchiment activé classique.
- 5 12. Composition de nettoyage selon la revendication 11, dans laquelle l'agent de blanchiment est choisi parmi le perborate et/ou le percarbonate et l'activateur choisi parmi la tétraacétyléthylènediamine, le nonanoyloxybenzènesulfonate et/ou le 3,5,-triméthylhexanoloxybenzènesulfonate.
- 10 13. Composition de nettoyage selon la revendication 10, dans laquelle ledit système de blanchiment est un autre système de blanchiment enzymatique.
14. Composition de nettoyage selon la revendication 10, dans laquelle ledit système de blanchiment est un système de blanchiment à base de catalyseur métallocène.
- 15 15. Composition de nettoyage selon la revendication 14, dans laquelle ledit catalyseur métallocène est un complexe de métaux de transition d'un ligand rigide macropolycyclique.
16. Composition de nettoyage selon les revendications 14 à 15, dans laquelle ledit catalyseur métallocène est du manganèse.
- 20 17. Composition de nettoyage selon l'une quelconque des revendications précédentes, laquelle est sous la forme d'un additif.
- 25 18. Composition d'adoucissement des tissus comprenant une oxygénase à base de substrat polyphénol et/ou hétérocyclique selon la revendication 1 et un agent tensioactif cationique comprenant deux longues chaînes en C<sub>11</sub> à C<sub>22</sub>.
- 30 19. Utilisation d'une oxygénase à base de substrat polyphénol et/ou hétérocyclique selon la revendication 1 dans une composition de nettoyage et/ou adoucissante pour le nettoyage de tissu et/ou l'élimination de salissures d'un tissu et/ou le maintien de la blancheur d'un tissu et/ou l'adoucissement des tissus et/ou l'apparence des couleurs d'un tissu et/ou l'inhibition de décoloration d'un tissu.
- 35 20. Utilisation d'une oxygénase à base de substrat polyphénol et/ou hétérocyclique selon la revendication 1 dans une composition de nettoyage pour nettoyer des surfaces dures telles que des sols, des murs, des carreaux de salle de bain et similaires.
- 40 21. Utilisation d'une oxygénase à base de substrat polyphénol et/ou hétérocyclique selon la revendication 1 dans une composition de nettoyage pour le lavage de la vaisselle à la main et en machine.
- 45 22. Utilisation d'une oxygénase à base de substrat polyphénol et/ou hétérocyclique selon la revendication 1 dans une composition pour des applications orales et/ou dentaires.
- 50 23. Utilisation d'une oxygénase à base de substrat polyphénol et/ou hétérocyclique selon la revendication 1, dans une composition de nettoyage et/ou adoucissante pour l'assainissement des surfaces traitées.
- 55

## REFERENCES CITED IN THE DESCRIPTION

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

## Patent documents cited in the description

- WO 9218683 A [0011]
- EP 086139 A [0011]
- WO 9501426 A [0012]
- US 4435307 A [0043]
- WO 9602653 A [0043]
- EP 739982 A [0043]
- GB 2075028 A [0043]
- GB 2095275 A [0043]
- DE 2247832 [0043]
- WO 9526398 A [0043]
- WO 9117243 A [0043]
- WO 9421801 A [0043]
- EP 91202879 A [0043]
- WO 9117244 A [0043]
- WO 9121801 A [0043]
- WO 9634092 A [0043]
- WO 9617994 A [0043]
- WO 9524471 A [0043]
- GB 1372034 A [0044]
- EP 258068 A [0044]
- WO 9205249 A [0044]
- WO 9522615 A [0044]
- WO 9403578 A [0044]
- WO 9535381 A [0044]
- WO 9600292 A [0044]
- WO 8809367 A [0044]
- WO 9009446 A [0044]
- WO 9414963 A [0044]
- WO 9414964 A [0044]
- GB 1243784 A [0045]
- EP 251446 A [0045] [0167]
- EP 199404 A [0045]
- EP 90915958 A [0045]
- WO 9106637 A [0045] [0167]
- WO 9510591 A [0045]
- WO 9510592 A [0045] [0056]
- WO 9102792 A [0045]
- WO 9523221 A [0045]
- WO 9318140 A [0045]
- WO 9203529 A [0045]
- WO 9507791 A [0045]
- WO 9425583 A [0045]
- EP 516200 A [0045]
- WO 94102597 A [0046]
- WO 9510603 A [0046]
- US 5003257 A [0046]
- EP 252666 A [0046]
- WO 9100353 A [0046]
- FR 2676456 [0046]
- EP 285123 A [0046]
- EP 525610 A [0046]
- EP 368341 A [0046]
- GB 1296839 A [0046]
- WO 9418314 A [0046] [0167]
- WO 9605295 A [0046] [0167]
- EP 277216 A [0046]
- WO 9526397 A [0046] [0167]
- WO 9623873 A [0046]
- WO 9535382 A [0046]
- EP 92870018 A [0049]
- WO 9307263 A [0050]
- WO 9307260 A [0050]
- WO 8908694 A [0050]
- US 3553139 A [0050]
- US 4101457 A [0050]
- US 4507219 A [0050]
- US 4261868 A [0050]
- US 3600319 A [0050]
- EP 199405 A [0050]
- EP 200586 A [0050]
- US 3519570 A [0050]
- WO 9401532 A [0050]
- US 4412934 A [0054] [0055]
- EP 120591 A [0054]
- WO 9428106 A [0054]
- EP 624154 A [0054]
- WO 9804664 A [0054]
- US 4483781 A [0055]
- US 740446 A [0055]
- EP 0133354 A [0055]
- US 4634551 A [0055]
- WO 9700937 A [0056]
- WO 9527772 A [0056]
- WO 9527773 A [0056]
- WO 9527774 A [0056]
- WO 9527775 A [0056]
- EP 537381 A [0057]
- WO 89099813 A [0057]
- WO 8909813 A [0057]
- EP 540784 A [0057]
- WO 9412621 A [0057]
- US 5246621 A [0058]
- US 5194416 A [0058]
- US 5114611 A [0058]
- EP 718398 A [0058]
- US 4033718 A [0066]
- US 4565647 A [0073]
- GB 1082179 A [0080] [0112]



## EP 1 002 040 B2

- US 3929678 A [0081] [0095] [0097]
- US 4228044 A [0088]
- EP 000224 A [0088]
- EP 0040562 A [0090]
- EP 0239910 A [0090]
- EP 92870181 A [0107]
- WO 96870140 A [0107]
- GB 1400898 A [0108]
- US 5019292 A [0108]
- GB 514276 A1 [0108]
- EP 0011340 B [0108]
- EP 0026527 B [0108]
- EP 0026528 B [0108]
- EP 0242919 B [0108]
- EP 0299575 A [0108]
- EP 0313146 A [0108]
- US 3935257 A [0111]
- GB 1379241 A [0111]
- NL 7205873 [0111]
- GB 1398421 A [0112]
- GB 1398422 A [0112]
- US 3936448 A [0112]
- GB 1425343 A [0113]
- GB 1596756 A [0117] [0139]
- US 3812044 A [0122]
- US 4704233 A [0123]
- US 3933672 A [0126]
- US 3455838 A [0130]
- US 4116885 A [0134]
- US 4711730 A [0134]
- EP 0272033 A [0134]
- WO 9108281 A [0138]
- EP 262897 A [0158]
- EP 256696 A [0158]
- WO 951059 A [0167]

### Non-patent literature cited in the description

- *J. Clin. Microbiol.*, 1994, vol. 32 (5), 1261-7 [0020]
- *J. Clin. Microbiol.*, 1992, vol. 30 (10), 2692-7 [0020]
- *The Journal of the American Oil Chemists Society*, 1975, vol. 52, 323-329 [0078]
- **H.C. Borghetty ; C.A. Bergman.** *J. Am. Oil. Chem. Soc.*, 1950, vol. 27, 88-90 [0142]
- **W.N. Linfield.** *Surfactant science Series*, vol. 7, 3 [0142]
- **W.N. Linfield.** *Tenside surf. det.*, 1990, vol. 27, 159-163 [0142]
- **M.K. Nagarajan ; W.F. Masler.** *Cosmetics and Toiletries*, 1989, vol. 104, 71-73 [0142]
- **M.K. Nagarajan ; W.F. Master.** *Cosmetics and Toiletries*, 1989, vol. 104, 71-73 [0145]
- *Journal of Polymer Science*, vol. 22, 1035-1039 [0161]