



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 013 665 A1**

(12) **EUROPEAN PATENT APPLICATION**
published in accordance with Art. 158(3) EPC

(43) Date of publication:
28.06.2000 Bulletin 2000/26

(51) Int. Cl.⁷: **C07K 14/47, C12N 15/12**

(21) Application number: **98911039.0**

(86) International application number:
PCT/JP98/01371

(22) Date of filing: **27.03.1998**

(87) International publication number:
WO 98/43997 (08.10.1998 Gazette 1998/40)

(84) Designated Contracting States:
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE**

(30) Priority: **28.03.1997 JP 9484597**

(71) Applicant:
SANTEN PHARMACEUTICAL CO., LTD.
Higashiyodogawa-ku, Osaka-shi, Osaka 533-
8651 (JP)

(72) Inventors:
• **OKUBO, Kousaku**
Mino-shi, Osaka 562-0045 (JP)

- **KURIYAMA, Hiroshi**
Toyonaka-shi, Osaka 560-0004 (JP)
- **MITA, Shiro**
Ashiya-shi, Hyogo 659-0014 (JP)
- **ISHIDA, Naruhiro**
Ikoma-shi, Nara 630-0243 (JP)

(74) Representative:
Nevant, Marc et al
Cabinet Beau de Loménie,
158, rue de l'Université
75340 Paris Cedex 07 (FR)

(54) **NOVEL POLYPEPTIDE HAVING WATER CHANNEL ACTIVITY AND DNA SEQUENCE**

(57) The present invention has its objects to provide a novel polypeptide having water channel activity and to a DNA sequence encoding for the polypeptide.

This invention is related to a novel polypeptide having water channel activity which has the amino acid sequence, within the molecule thereof, shown in the sequence listing under SEQ ID NO:1.

EP 1 013 665 A1

Description

TECHNICAL FIELD

- 5 [0001] The present invention relates to a novel, human adipose tissue-derived polypeptide having water channel activity and to a DNA sequence encoding for the polypeptide.

BACKGROUND ART

- 10 [0002] The permeation of water through a cell membrane generally occurs slowly by way of diffusion into the lipid bilayer which is the main structure of the cell membrane. Recently, however, it was discovered that, in certain kinds of cells, water is transferred rapidly through the cell membrane, suggesting the involvement in the above phenomenon of some membrane protein selectively permeable to water. Thereafter, such membrane proteins of various kinds have actually been isolated. Such membrane proteins are designated as water channels. In this specification, the function of the above water channels which has selective permeation of water through the cell membrane is referred to as "water channel activity". The water channels may be permeable to water alone or permeable to not only water but also low-molecular-weight substances such as glycerol and urea.

- [0003] As a membrane protein having such water channel activity, there have been isolated a group of membrane proteins known as aquaporins (AQPs). Furthermore, some aquaporin genes have so far been cloned, and aquaporins such as AQP1 through AQP5, FA-CHIP and AQP-γTIP have been discovered in mammals, amphibians, plants, etc. [cf. e.g. Akira Sasaki, Igaku no Ayumi (Advances in Medicine), vol. 173, No. 9, 1995].

- 20 [0004] P. Agre et al. reported, in Science (vol. 256, pp. 385 to 387, 1992) that *Xenopus laevis* oocytes in which the *in vitro* transcript RNA for CHIP28, the current designation of which is AQP1, had been introduced showed increased water permeability. In Science (vol. 264, pp. 92 to 95, 1994), B. A. van Oost et al. disclosed the amino acid sequence of human AQP2 and suggested that this should be involved in vasopressin-dependent urine concentration.

- 25 [0005] In Proc. Natl. Acad. Sci. USA (vol. 91, pp. 6269 to 6273, 1994), Ishibashi et al. disclosed the nucleotide sequence of the gene for renal collecting tubule-derived AQP3 and the amino acid sequence encoded thereof. Ishibashi et al. confirmed its water channel activity by injecting the AQP3 cRNA into *Xenopus laevis* oocytes and measuring the water permeability thereof. Ishibashi et al. reported that this AQP3 transported not only water but also nonionic small molecules such as urea and glycerol.

- 30 [0006] In Proc. Natl. Acad. Sci. USA (vol. 91, pp. 13052 to 13056, 1994), J. S. Jung et al. reported about the isolation of AQP4. This AQP4 is known to occur most abundantly in mammalian brains and have mercury resistance. In J. Biol. Chem. (vol. 270, pp. 1908 to 1912, 1995), S. Raina et al. who prepared rat salivary gland-derived AQP5 cDNA describe the nucleotide sequence of the cDNA and the amino acid sequence encoded thereby. S. Raina et al. cloned the cDNA by utilizing the occurrence of an NPA sequence and confirmed its function by observing that the cRNA enhances the water permeability of *Xenopus laevis* oocytes.

- [0007] The aquaporin family mentioned above is considered to be involved in water metabolism in mammals and, for example, it has been confirmed that AQP2 is found only in the renal collecting tubule luminal membrane, which is indicative of its close association with the vasopressin-urea concentration system, and its involvement in renal diseases has become acknowledged. Therefore, such membrane proteins having water channel activity are of importance in any attempt to develop novel therapies for water-associated diseases.

- 40 [0008] Meanwhile, the expression of the aquaporin family mentioned above has been confirmed in such organs as kidney, brain, gall bladder, eye, intestine, salivary gland and bronchus but there is no report as yet about the occurrence of membrane proteins having water channel activity in other organs or tissues, particularly in adipose tissue.

SUMMARY OF THE INVENTION

- [0009] In view of the above-mentioned state of the art, the present invention has for its object to provide a novel membrane protein having water channel activity and a DNA sequence encoding for the polypeptide.

- 50 [0010] The present invention is related to a novel polypeptide having water channel activity which has the amino acid sequence, within the molecule thereof, shown in the sequence listing under SEQ ID NO:1.

- [0011] The present invention is also related to a nucleotide sequence itself which codes for a polypeptide having, within the molecule thereof, the amino acid sequence shown in the sequence listing under SEQ ID NO:1 and having water channel activity.

- 55 [0012] The present invention is further related to the DNA sequence shown in the sequence listing under SEQ ID NO:2.

- [0013] The present invention is still further related to a polypeptide having water channel activity which has the amino acid sequence, within the molecule thereof, encoded by the nucleotide No. 173 to No. 1198 of the nucleotide

sequence shown in the sequence listing under SEQ ID NO:2.

DETAILED DESCRIPTION OF THE INVENTION

[0014] In the following, the present invention is described in detail.

[0015] The polypeptide of the present invention has the amino acid sequence shown in the sequence listing under SEQ ID NO:1. This polypeptide has a sequence composed of three amino acids, namely asparagine-proline-alanine, as the amino acid Nos. 195 to 197. However, the characteristic feature common to the so-far known AQP, that said asparagine-proline-alanine sequence occurs twice, is not found in the polypeptide of the present invention. That this polypeptide has water channel activity can be confirmed from the fact that it enhances the water permeability of *Xenopus laevis* oocytes.

[0016] The above polypeptide may be generated by translation by a protein synthesis system constituted, *in vivo* or *in vitro*, based on the nucleotide sequence coding for the amino acid sequence of said polypeptide. The nucleotide sequence of the present invention substantially has a region coding for the amino acid sequence of said polypeptide and, where necessary, may contain one or more other regions such as a promoter region. In the protein synthesis based on genetic information, the information carried by the gene DNA is transcribed into mRNA as the result of DNA-dependent RNA synthesis aided by RNA polymerase. And, this mRNA is translated into the amino acid sequence in a tRNA-containing protein synthesis system. Therefore, the nucleotide sequence of the present invention includes not only the DNA sequence but also the RNA sequence. Furthermore, since it is generally known that, for an amino acid, there is one or a plurality of codons corresponding thereto, it is a matter of course that the above-mentioned nucleotide sequence is not limited to only one sequence but may include nucleotide sequences resulting from substitution of another synonymous codon coding for the same amino acid.

[0017] The above polypeptide can be formed based on the genetic information carried by the DNA sequence shown in the sequence listing under SEQ ID NO:2. This polypeptide is encoded by that portion of the nucleotide sequence shown in the sequence listing under SEQ ID NO:2 which ranges from the nucleotide No. 173 to No. 1198. Of the DNA sequence shown in the sequence listing under SEQ ID NO:2, the nucleotide sequences other than the portion of said nucleotide numbers are noncoding regions, among which the polyadenylation consensus sequence AATAAA occurs at the nucleotide No.1234 to No.1239. Other possible reading frames of said DNA sequence shown under SEQ ID NO:2 can be excluded from consideration, since the polypeptides encoded are very small-sized, hence considered to be incapable of performing any water channel function.

[0018] It has been confirmed by the inventors that the full-length sequence of the above nucleic acid bases has no counterpart sequence either in GenBank or in dbEST.

[0019] The polypeptide of the present invention has water channel activity in adipose tissue. While adipose tissue is distributed in various parts of the living organism, the polypeptide of the present invention has an action to control the transfer of water in such adipose tissue and is expected to be effective in upholding normal functions of adipose tissue at various sites.

BEST MODES FOR CARRYING OUT THE INVENTION

[0020] The above-mentioned DNA sequence given under SEQ ID NO:2 corresponds to the nucleotide sequence of cDNA obtained from human adipose tissue by cloning. Human adipose tissue is a tissue which stores fat as energy reserves. It is known that various proteins are formed in this adipose tissue. A 3'-directed DNA library is known as a cDNA library from which the genes actually expressed in this adipose tissue or, in other words, the mRNA composition in this adipose tissue can be copied faithfully. This 3'-directed DNA library contains only those specified 3'-terminal regions of mRNAs which range from poly(A) to the *Mbo*I site which is a restriction enzyme recognition site upstream of said poly(A) and, therefore, said library is suited for template preparation by the PCR technique. Therefore, by extracting a clone from this library and using it to determine a longer nucleotide sequence including the amino acid coding region from this complete adipose tissue cDNA library, it becomes possible to obtain the genetic information concerning the protein which is actually formed in adipose tissue. The DNA sequence of the present invention as shown under the above-mentioned SEQ ID NO:2 is found by such cloning. A method of obtaining the cDNA by cloning from human adipose tissue is now described in detail.

[0021] Known as said method is, for example, the method described in Biochem. Biophys. Res. Commun., 221, 286 to 289 (1996). According to this method, the total RNA is first separated from adipose tissue and, when necessary, purified to give poly(A) RNA. For this purification, commercially available purification kits can be used. For example, Pharmacia's Quick prep mRNA purification kit or the like in which oligo(dT)-cellulose and various buffers are used in combination can judiciously be employed. Then, a double-stranded cDNA is synthesized using a pUC19 system vector primer and the double-stranded cDNA so synthesized is selectively cleaved with the restriction enzyme *Mbo*I (which recognizes the nucleotide sequence GATC). On that occasion, the GATC sequence on the vector molecule side, which

can be methylated to give G^mATC when replication is effected in dam⁺ bacterial cells, is not cleaved with MboI. As the cleaved cDNA is subjected to self-cyclization using *E. coli* ligase, a plasmid containing a cDNA fragment extending from poly(A) to the nearest MboI site is completed. This plasmid is introduced into *Escherichia coli*, followed by cultivation and selection of a transformant *E. coli* colony. Then, the cDNA in said colony is amplified by the PCR technique using appropriate PCR primers.

[0022] On the other hand, the full-length double-stranded cDNA synthesized using the pUC19 system vector primer is cleaved at the 5' end using T4 polymerase and subjected to cyclization using T4 ligase and introduction into *Escherichia coli* for transformation. From among the thus-obtained transformant colonies, the desired colony is obtained by screening using, as a probe, a labeled form of the adipose tissue-specific cDNA obtained from said 3'-directed DNA library by the method mentioned above. The insert cDNA in this colony is amplified by the PCR technique using appropriate PCR primers. The amplification product is purified and, after sonication, subcloned into the M13 phage.

[0023] The nucleotide sequence of the thus-cloned cDNA can be determined, for example, by reaction with a primer dye, purification and analysis using an automated sequencer or the like. In this manner, the DNA sequence of the present invention can be obtained.

[0024] The polypeptide of the present invention has water channel activity. This water channel activity can be confirmed by observing an enhancement of water permeability in *Xenopus laevis* oocytes. It is known that no AQP family gene has been expressed in *Xenopus laevis* oocytes and, therefore, any increase in water permeability as caused by the injected mRNA can be easily confirmed. For this reason, said oocytes are widely used in confirming water channel activity. For example, in Proc. Natl. Acad. Sci. USA, vol. 91, pp. 6269 to 6273 (1994), Ishibashi et al. confirmed the water channel activity by inserting the AQP3 cDNA into the pSP64T-derived BlueScript vector, synthesizing the cRNA using T7 RNA polymerase, injecting this cRNA into *Xenopus laevis* oocytes and, after 48 to 62 hours of incubation following injection, observing an increase in water permeability and in the volume of the oocyte.

[0025] The polypeptide encoded by the DNA sequence of the present invention can be identified by analyzing the amino acid sequence of the polypeptide synthesized in an *Escherichia coli* protein synthesis system constituted *in vitro*. In this case, the methods of identifying N- and C-terminal sequences of expression products as described in Shin Seikagaku Jikken Koza (Experiments in Biochemistry, A New Course) 1 (published by Tokyo Kagaku Dojin), pages 22 to 24 can be employed.

[0026] The following examples illustrate the present invention in further detail. These examples are, however, by no means limitative of the scope of the present invention.

Example 1

Determination of DNA nucleotide sequence

[0027] According to a recent report [Biochem. Biophys. Res. Commun., 221, 286 to 289 (1996)], a DNA for an adipose tissue-specific collagen-like factor has been cloned using a 3'-directed DNA library containing only those specified 3'-terminal regions of RNA molecules which range from poly(A) to the restriction enzyme MboI site upstream thereof [Nature Genet., 2, 173 to 179 (1992)], which library makes it possible not only to identify the gene under expression but also to examine the frequency of expression or amount of expression. In accordance with the method described in the above report, the DNA nucleotide sequence of the present invention was determined using an adipose tissue-specific 3'-directed DNA library.

Method

[0028] Reverse transcriptase was added to poly(A)⁺ RNA separated and purified from human adipose tissue using a Quick prep mRNA purification kit, followed by insertion into λ ZAP II containing the pUC19 system vector pBluescript and introduction into *Escherichia coli*. Screening carried out using an adipose tissue-specific partial DNA as a probe gave a colony of transformant *Escherichia coli*. Then, together with two primers (SK: 5'CGCTCTAGAACTAGTGGATC3'; T7: 5'GTAATACGACTCACTATAGGGC3'), PCR [polymerase chain reaction; (30 seconds at 95°C + 30 seconds at 50°C + 60 seconds at 70°C) x 15 cycles, followed by (30 seconds at 95°C + 60 seconds at 70°C) x 15 cycles], was carried out and, after sonication, the product was subcloned in M13. A primer dye was added thereto and, after purification, the nucleotide sequence of the DNA was analyzed using an automated sequencer. The nucleotide sequence obtained is shown in the sequence listing under SEQ ID NO:2.

Example 2

Examination as to water permeability

[0029] As for the studies on the water permeability of the AQP family, there are reports about AQP1 [Science, 256, 385-387 (1992)] and AQP3 [Proc. Natl. Acad. Sci. USA, 91, 6269-6273 (1994)] in which RNA was introduced into *Xenopus laevis* oocytes, in which AQP family genes have not been expressed, and then the water permeability was calculated from changes in the surface area and volume of said oocytes in a hypotonic culture medium. Therefore, the membrane protein encoded by the DNA of the present invention was checked for water permeability according to the method described in the references cited above.

Test Method

[0030] The RNA (10 ng) obtained in Example 1 from human adipose tissue was introduced into *Xenopus laevis* oocytes by microinjection and the oocytes were incubated at 20°C for 3 days in an isotonic culture medium (about 200 mOsm). The cultured oocytes were transferred to a hypotonic culture medium (about 40 mOsm). Photograph was taken 20 seconds and 40 seconds after transfer, and the sectional area and volume of each oocyte were determined using an image analyzer. The water permeability of the membrane protein was calculated as follows:

$$\text{Permeability (cm/sec)} = [V_{40} - V_{20}] / 20 / [A_{20} \times 10^{-2} \times 4] \times 1.384$$

where V_{20} denotes the oocyte volume (cm^3) after 20 seconds, V_{40} denotes the oocyte volume (cm^3) after 40 seconds, and A_{20} denotes the oocyte sectional area (mm^2) after 20 seconds.

[0031] The result is shown in Table 1. The water permeability found when purified water was used for microinjection in lieu of the RNA is also shown.

[0032] Further, the expression of the polypeptide of the present invention in said oocytes was confirmed by a C-terminal region immunoassay using the rabbit antiserum obtained by using the polypeptide of the present invention as synthesized *in vitro*.

Table 1

	Water permeability (cm/sec)
Group in which RNA introduction was not made	30.0×10^{-4}
Group in which RNA introduction was made	292.5×10^{-4}

[0033] As is evident from Table 1, the polypeptide encoded by the DNA sequence of the present invention caused an increase in water permeability after introduction into the oocytes. This result indicates that the polypeptide of the present invention has water channel activity.

INDUSTRIAL UTILIZABILITY

[0034] The present invention provides a novel protein having water channel activity and a novel DNA sequence encoding the protein. Said protein is one found in human adipose tissue for which the occurrence of water channels has not been reported as yet. The present invention makes it possible to develop novel therapies for water or fat metabolism-associated diseases in which said tissue is involved.

[Sequence listings]

5 SEQ ID No.: 1

Length: 342

Type: amino acid

10 Topology: linear

Species: peptide

15 Sequence

Met Val Gln Ala Ser Gly His Arg Arg Ser Thr Arg Gly Ser Lys Met
 5 10 15
 Val Ser Trp Ser Val Ile Ala Lys Ile Gln Glu Ile Leu Gln Arg Lys
 20 25 30
 Met Val Arg Glu Phe Leu Ala Glu Phe Met Ser Thr Tyr Val Met Met
 35 40 45
 Val Phe Gly Leu Gly Ser Val Ala His Met Val Leu Asn Lys Lys Tyr
 50 55 60
 Gly Ser Tyr Leu Gly Val Asn Leu Gly Phe Gly Phe Gly Val Thr Met
 65 70 75 80
 Gly Val His Val Ala Gly Arg Ile Ser Gly Ala His Met Asn Ala Ala
 85 90 95
 Val Thr Phe Ala Asn Cys Ala Leu Gly Arg Val Pro Trp Arg Lys Phe
 100 105 110
 Pro Val Tyr Val Leu Gly Gln Phe Leu Gly Ser Phe Leu Ala Ala Ala
 115 120 125
 Thr Ile Tyr Ser Leu Phe Tyr Thr Ala Ile Leu His Phe Ser Gly Gly
 130 135 140
 Gln Leu Met Val Thr Gly Pro Val Ala Thr Ala Gly Ile Phe Ala Thr
 145 150 155 160
 Tyr Leu Pro Asp His Met Thr Leu Trp Arg Gly Phe Leu Asn Glu Ala
 165 170 175
 Trp Leu Thr Gly Met Leu Gln Leu Cys Leu Phe Ala Thr Thr Asp Gln
 180 185 190
 Glu Asn Asn Pro Ala Leu Pro Gly Thr Glu Ala Leu Val Ile Gly Ile
 195 200 205

	Leu	Val	Val	Ile	Ile	Gly	Val	Ser	Leu	Gly	Met	Asn	Thr	Gly	Tyr	Ala		
5	210						215						220					
	Ile	Asn	Pro	Ser	Arg	Asp	Leu	Pro	Pro	Arg	Ile	Phe	Thr	Phe	Ile	Ala		
	225						230						235			240		
10	Gly	Trp	Gly	Lys	Gln	Val	Phe	Ser	Asn	Gly	Glu	Asn	Trp	Trp	Trp	Val		
				245						250						255		
	Pro	Val	Val	Ala	Pro	Leu	Leu	Gly	Ala	Tyr	Leu	Gly	Gly	Ile	Ile	Tyr		
				260						265						270		
15	Leu	Val	Phe	Ile	Gly	Ser	Thr	Ile	Pro	Arg	Glu	Pro	Leu	Lys	Leu	Glu		
	275						280						285					
	Asp	Ser	Val	Ala	Tyr	Glu	Asp	His	Gly	Ile	Thr	Val	Leu	Pro	Lys	Met		
20	290						295						300					
	Gly	Ser	His	Glu	Pro	Thr	Ile	Ser	Pro	Leu	Thr	Pro	Val	Ser	Val	Ser		
	305						310						315			320		
25	Pro	Ala	Asn	Arg	Ser	Ser	Val	His	Pro	Ala	Pro	Pro	Leu	His	Glu	Ser		
				325						330						335		
	Met	Ala	Leu	Glu	His	Phe												
	340																	

SEQ ID No: 2

Length: 1258

Type: nucleotide

Strandedness: double strand

Topology: linear

Species: cDNA to mRNA

Original source

Organism: human

Tissue type: adipose tissue

Feature Information

Name/Key: peptide

Location: 173..1198

Identification method: E

Sequence

```

25  GGCTCTGGAC TGGGGACACA GGGATAGCTG AGCCCCAGCT GGGGGTGGAA GCTGAGCCAG 60
    GGACAGTCAC GGAGGAACAA GATCAAGATG CGCTGTAAC TGAAGAGCCCC CAAGGCGGAG 120
    GCTGAGAATC AGAGACATTT CAGCAGACAT CTACAAATCT GAAAGACAAA AC ATG GTT 178
                                     Met Val
30                                     1

    CAA GCA TCC GGG CAC AGG CGG TCC ACC CGT GGC TCC AAA ATG GTC TCC 226
    Gln Ala Ser Gly His Arg Arg Ser Thr Arg Gly Ser Lys Met Val Ser
35          5              10              15

    TGG TCC GTG ATA GCA AAG ATC CAG GAA ATA CTG CAG AGG AAG ATG GTG 274
    Trp Ser Val Ile Ala Lys Ile Gln Glu Ile Leu Gln Arg Lys Met Val
40          20              25              30

    CGA GAG TTC CTG GCC GAG TTC ATG AGC ACA TAT GTC ATG ATG GTA TTC 322
    Arg Glu Phe Leu Ala Glu Phe Met Ser Thr Tyr Val Met Met Val Phe
45          35              40              45              50

    GGC CTT GGT TCC GTG GCC CAT ATG GTT CTA AAT AAA AAA TAT GGG AGC 370
    Gly Leu Gly Ser Val Ala His Met Val Leu Asn Lys Lys Tyr Gly Ser
50          55              60              65

    TAC CTT GGT GTC AAC TTG GGT TTT GGC TTC GGA GTC ACC ATG GGA GTG 418
    Tyr Leu Gly Val Asn Leu Gly Phe Gly Phe Gly Val Thr Met Gly Val
          70              75              80

```


	CAC GTG GCA GGC CGC ATC TCT GGA GCC CAC ATG AAC GCA GCT GTG ACC	466
	His Val Ala Gly Arg Ile Ser Gly Ala His Met Asn Ala Ala Val Thr	
5	85 90 95	
	TTT GCT AAC TGT GCG CTG GGC CGC GTG CCC TGG AGG AAG TTT CCG GTC	514
	Phe Ala Asn Cys Ala Leu Gly Arg Val Pro Trp Arg Lys Phe Pro Val	
10	100 105 110	
	TAT GTG CTG GGG CAG TTC CTG GGC TCC TTC CTG GCG GCT GCC ACC ATC	562
	Tyr Val Leu Gly Gln Phe Leu Gly Ser Phe Leu Ala Ala Ala Thr Ile	
	115 120 125 130	
15	TAC AGT CTC TTC TAC ACG GCC ATT CTC CAC TTT TCG GGT GGA CAG CTG	610
	Tyr Ser Leu Phe Tyr Thr Ala Ile Leu His Phe Ser Gly Gly Gln Leu	
	135 140 145	
20	ATG GTG ACC GGT CCC GTC GCT ACA GCT GGC ATT TTT GCC ACC TAC CTT	658
	Met Val Thr Gly Pro Val Ala Thr Ala Gly Ile Phe Ala Thr Tyr Leu	
	150 155 160	
	CCT GAT CAC ATG ACA TTG TGG CGG GGC TTC CTG AAT GAG GCG TGG CTG	706
25	Pro Asp His Met Thr Leu Trp Arg Gly Phe Leu Asn Glu Ala Trp Leu	
	165 170 175	
	ACC GGG ATG CTC CAG CTG TGT CTC TTC GCC ATC ACG GAC CAG GAG AAC	754
	Thr Gly Met Leu Gln Leu Cys Leu Phe Ala Thr Thr Asp Gln Glu Asn	
30	180 185 190	
	AAC CCA GCA CTG CCA GGA ACA GAG GCG CTG GTG ATA GGC ATC CTC GTG	802
	Asn Pro Ala Leu Pro Gly Thr Glu Ala Leu Val Ile Gly Ile Leu Val	
35	195 200 205 210	
	GTC ATC ATC GGG GTG TCC CTT GGC ATG AAC ACA GGA TAT GCC ATC AAC	850
	Val Ile Ile Gly Val Ser Leu Gly Met Asn Thr Gly Tyr Ala Ile Asn	
	215 220 225	
40	CCG TCC CGG GAC CTG CCC CCC CGC ATC TTC ACC TTC ATT GCT GGT TGG	898
	Pro Ser Arg Asp Leu Pro Pro Arg Ile Phe Thr Phe Ile Ala Gly Trp	
	230 235 240	
45	GGC AAA CAG GTC TTC AGC AAT GGG GAG AAC TGG TGG TGG GTG CCA GTG	946
	Gly Lys Gln Val Phe Ser Asn Gly Glu Asn Trp Trp Trp Val Pro Val	
	245 250 255	
50	GTG GCA CCA CTT CTG GGT GCC TAT CTA GGT GGC ATC ATC TAC CTG GTC	994
	Val Ala Pro Leu Leu Gly Ala Tyr Leu Gly Gly Ile Ile Tyr Leu Val	
	260 265 270	

55

[illegible]

30 Claims

1. A novel polypeptide having water channel activity which has the amino acid sequence, within the molecule thereof, shown in the sequence listing under SEQ ID NO:1.
2. A nucleotide sequence which codes for a polypeptide having, within the molecule thereof, the amino acid sequence shown in the sequence listing under SEQ ID NO:1 and having water channel activity.
3. The DNA sequence shown in the sequence listing under SEQ DI NO:2.
4. A polypeptide having water channel activity which has the amino acid sequence, within the molecule thereof, encoded by the nucleotide No.173 to No.1198 of the nucleotide sequence shown in the sequence listing under SEQ ID NO:2.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP98/01371

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl ⁶ C07K14/47, C12N15/12		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) Int.Cl ⁶ C07K14/47, C12N15/12		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) BIOSIS (DERWENT), WPI (DERWENT), GenBank/EMBL (geneseq)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	Biochem. Biophys. Res. Commun., Vol. 241, No. 1 (1997, December) H. Kuriyama et al., "Molecular Cloning and Expression of a Novel Human Aquaporin from Adipose Tissue with Glycerol Permeability", p.53-58	1-4
P, Y	J. Biol. Chem., Vol. 272, No. 33 (1997, August) K. Ishibashi et al., "Cloning and Functional Expression of a New Water Channel Abundantly Expressed in the Testis Permeable to Water, Glycerol, and Urea", p.20782-20786	1-4
☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search July 13, 1998 (13. 07. 98)		Date of mailing of the international search report July 21, 1998 (21. 07. 98)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)