



(11) **EP 1 044 014 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

Note: Bibliography reflects the latest situation

- (15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see page(s) 12,13
- (48) Corrigendum issued on:
03.01.2007 Bulletin 2007/01
- (45) Date of publication and mention of the grant of the patent:
09.08.2006 Bulletin 2006/32
- (21) Application number: **99900567.1**
- (22) Date of filing: **08.01.1999**
- (51) Int Cl.:
A61K 38/16^(2006.01) A61P 37/08^(2006.01)
- (86) International application number:
PCT/GB1999/000070
- (87) International publication number:
WO 1999/034817 (15.07.1999 Gazette 1999/28)

(54) **AGENT FOR TREATING ALLERGIC OR HYPERSENSITIVITY CONDITION**

MITTEL ZUR BEHANDLUNG VON ALLERGISCHEN ODER
ÜBEREMPFINDLICHKEITZUSTÄNDEN

AGENT POUR TRAITER UN ÉTAT ALLERGIQUE OU D'HYPERSENSIBILITÉ

- | | |
|---|---|
| <p>(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE</p> <p>(30) Priority: 09.01.1998 GB 9800487</p> <p>(43) Date of publication of application:
18.10.2000 Bulletin 2000/42</p> <p>(60) Divisional application:
06016549.5 / 1 736 170
06076571.6</p> <p>(73) Proprietor: Trident Pharmaceuticals, Inc.
Boston, MA 02109 (US)</p> <p>(72) Inventors:
• WILLIAMS, Neil Andrew
Somerset BS26 2EF (GB)
• HIRST, Timothy Raymond
North Somerset BS21 7RR (GB)
• BIENENSTOCK, John
Toronto, Ontario M5V 1E3 (CA)</p> | <p>(74) Representative: Crump, Julian Richard John
Mintz Levin Cohn Ferris Glovsky and Popeo
Intellectual Property LLP
The Rectory
9, Ironmonger Lane
London EC2V 8EY (GB)</p> <p>(56) References cited:
WO-A-95/10301 WO-A-97/02045</p> <ul style="list-style-type: none">• NASHAR T O ET AL: "Modulation of B-cell activation by the B subunit of Escherichia coli enterotoxin: receptor interaction up-regulates MHC class II, B7, CD40, CD25 and ICAM-1." IMMUNOLOGY, (1997 AUG) 91 (4) 572-8, XP002102696• NASHAR T O ET AL: "POTENT IMMUNOGENICITY OF THE B SUBUNITS OF ESCHERICHIA COLI HEAT -LABILE ENTEROTOXIN: RECEPTOR BINDING IS ESSENTIAL AND INDUCES DIFFERENTIAL MODULATION OF LYMPHOCYTE SUBSETS" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 93, no. 1, 1 January 1996, pages 226-230, XP002019422 |
|---|---|

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 044 014 B9

- **YANKELEVICH B ET AL: "PREVENTION OF ACUTE GRAFT-VERSUS-HOST DISEASE BY TREATMENT WITH A NOVEL IMMUNOSUPPRESSANT. CHOLERA TOXIN B SUBUNIT" JOURNAL OF IMMUNOLOGY, vol. 154, no. 7, 1 January 1995, pages 3611-3617, XP002019424**

Description

[0001] The present invention relates to a medicament.

[0002] In particular, the present invention relates to a medicament useful to affect an allergic condition and/or a hypersensitivity condition.

[0003] More in particular, in one aspect the present invention relates to an immunological tolerance inducing agent.

[0004] More in particular, the present invention relates to such an agent optionally co-administered with a specific antigen for use in the treatment of mammalian particularly human, allergic and other hypersensitivity diseases.

[0005] When an adaptive immune response occurs in an exaggerated or inappropriate form, the term allergy or hypersensitivity is applied. Allergic or hypersensitivity reactions are the result of normally beneficial immune responses acting inappropriately to foreign antigens (usually environmental macromolecules) and sometimes cause inflammatory reactions and tissue damage. In these situations, a normally harmless environmental stimulus, called an "allergen", triggers an immune response which upon re-exposure, is re-activated to generate pathological damage. Allergies or hypersensitivities are distinguished into four types of reactions. The first three are antibody-mediated, the fourth is mediated mainly by T cells and macrophages.

[0006] In Type I Immediate Hypersensitivity/Atopic Allergy, the principal immune response to the allergen involves the production of IgE antibodies. Such disorders are by far the most prevalent in humans and are seen as principal targets for new therapeutic approaches. Although these diseases are not exclusively IgE mediated, IgE binds to cells within the tissues such as mast cells and basophils and the cross-linking of IgE on the cells surfaces by allergen invokes the release of many inflammatory mediators.

[0007] Typical examples of such diseases include asthma, allergic cough, allergic rhinitis and conjunctivitis, atopic eczema and dermatitis, urticaria, hives, insect bite allergy, dietary and certain drug allergies. In many cases, the particular allergens are known. By way of example, the principal allergen in asthma is DerP1 from house dust mite but other triggers of asthma such as pet dander antigens also exist.

[0008] Type II or antibody dependent cytotoxic hypersensitivity occurs when antibodies of a different type, usually IgG and IgM, binds to either self antigen or foreign antigen on cells and leads to phagocytosis, killer cell activity or complement mediated lysis. These types of allergies are relatively unusual but can include some allergies to drugs.

[0009] Type III hypersensitivity develops when immune complexes are formed in large quantities or cannot be cleared adequately by the reticuloendothelial system. The immune complexes usually result from the deposition of antibody, usually IgM or IgG, allergen complexes at these sites. In normal circumstances, antibody binds to allergen and is cleared by a variety of tissue cells. However, a number of factors may influence the persistence of the immune complexes and where they remain in the blood for prolonged periods, they can lodge and establish inflammation in the kidneys, skin (where they cause rashes) and joints (where they can cause a type of arthritis other than rheumatoid arthritis).

[0010] Type IV or delayed type hypersensitivity (DTH) does not involve antibody but instead the prolonged activation of T lymphocytes. These T cells are capable of secreting soluble factors causing tissue damage and enhancing the recruitment and activation of other cell types to the tissues. Incoming cells themselves contribute to the inflammation and tissue damage. DTH is most seriously manifested when antigens (for example those associated with mycobacteria tuberculosis) are trapped in a macrophage and cannot be cleared. T cells are then stimulated to elaborate cytokines which mediate a range of inflammatory responses. DTH reactions are less common than Type I reactions but are seen in graft rejection and allergic contact dermatitis which is generally manifested as a contact sensitivity (allergy usually involving skin rash) to environmental "contact allergens" such as heavy metals.

[0011] Oral administration of antigens - such as allergens and autoantigens - has long been recognised as a method to prevent peripheral T cell responses and, in the case of autoantigens, has also been shown to prevent or delay the onset of several experimental autoimmune diseases including experimental allergic encephalomyelitis (EAE). Major problems recognised with such strategies are that it usually requires feeding of large, if not massive, doses of autoantigens and it is generally less efficient in an immune as opposed to a naive host. The latter problem has limited the therapeutic potential of this strategy. However, it has now been shown by Sun *et al* (1994 Proc Natl Acad Sci 91: 10795-10799) that oral administration of minute amounts of prototype particulate and soluble protein antigens conjugated to cholera toxin B subunit (CtxB), the nontoxic receptor-binding moiety of cholera toxin, can readily induce tolerance in the peripheral T-cell compartment and is effective not only in naive but also in systemically sensitised animals. In addition, oral administration of minute amounts of an autoantigen, myelin basic protein (MBP), coupled to CtxB can prevent EAE in Lewis rats (Sun *et al* 1996 Proc Natl Acad Sci 93: 7196-7201). Other researchers have also shown that feeding even a single dose of minute amounts (microgram) of antigens conjugated to the receptor binding nontoxic B subunit moiety of cholera toxin (CtxB) can markedly suppress systemic T cell mediated inflammatory reactions in naive as well as in experimental animals (Bergerot *et al* 1997 Proc Natl Acad Sci 94: 4610-4614).

[0012] *Escherichia coli* (*E. coli*) heat labile enterotoxin (Etx) and its closely related homologue, cholera toxin (Ctx), are examples of protein toxins which bind to glycolipid receptors on host cell surfaces. Each toxin consists of six non-covalently linked polypeptide chains, including a single A subunit (27 kDa) and five identical B subunits (11.6 kDa) which

principally bind to GM 1 ganglioside receptors found on the surfaces of mammalian cells (Nashar *et al* 1996 Proc Natl Acad Sci 93: 226-230). The A subunit is responsible for toxicity possessing adenosine diphosphate (ADP) ADP-ribosyl-transferase activity, whereas the B subunits (EtxB and CtxB) are non-toxic oligomers which bind and cross-link a ubiquitous cell surface glycolipid ganglioside, called GM1, thus facilitating A subunit entry into the cell.

[0013] The GM1 ganglioside receptor is a member of family of gangliosides comprising sialic acid containing glycolipids (also called glycosphingolipids) which are formed by a hydrophobic portion, the ceramide, and a hydrophilic part, that is the oligosaccharide chain. Gangliosides are defined as any ceramide oligosaccharide carrying, in addition to other sugar residues, one or more sialic residues (Oxford Dictionary of biochemistry and molecular biology. Oxford University Press. 1997. Eds Smith AD, Datta SP, Howard Smith G, Campbell PN, Bentley R and McKenzie HA). Although first described in neural tissue, several studies have shown that gangliosides are almost ubiquitous molecules expressed in all vertebrate tissues. Within cells, gangliosides are usually associated with plasma membranes, where they may act as receptors for a variety of molecules and take part in cell-to-cell interaction and in signal transduction. In addition, gangliosides are expressed in cytosol membranes like those of secretory granules of some endocrine cells such as the pancreatic islets and adrenal medulla.

[0014] Gangliosides contain in their oligosaccharide head groups one or more residues of a sialic acid which gives the polar head of the gangliosides a net negative charge at pH 7.0. The sialic acid usually found in human gangliosides is N-acetylneuraminic acid. Over 20 different types of gangliosides have been identified, differing in the number and relative positions of the hexose and sialic residues which form the basis of their classification. Nearly all of the known gangliosides have a glucose residue in glycosidic linkage with ceramide, residues of D-galactose and N-acetyl-D-galactosamine are also present.

[0015] In the ganglioside nomenclature of gangliosides, devised by Svennerholm (Biochemistry Lehninger 2nd Ed 1975 Worth Publishers Inc p 294-295), the subscript letters indicate the number of sialic groups. M is monosialo, D is disialo and T is trisialo.

[0016] One of the best studied members of the ganglioside family is the monosialosylganglioside, GM1, which has been shown to be the natural receptor for the cholera toxin. Soluble ganglioside GM1 binds to the toxin with high affinity and inactivates it (Svennerholm 1976 Adv Exp Med Biol 71: 191-204).

[0017] The chemical formula for GM1 can be represented as:

Gal β 3GalNAc β 4(NeuAc α 3)Gal β 4Glc β 1 Cer

where Glc is D-glucose, Gal is D-galactose, GalNAc is N-acetyl-D-galactosamine; NeuAc is N-acetylneuraminic acid, Cer is ceramide.

[0018] The chemical formula for GM1 can also be represented as
galactosyl-N-acetylgalactosaminyl {sialosyl}lactosyl ceramide
or

galactosyl-N-acetyl-galactosaminyl-(sialyl)-galactosylglucosylceramide

[0019] The x-ray crystal structures of Etx bound to lactose (Sixma *et al* 1992 Nature (London) 355: 561-564) and CtxB bound to the pentasaccharide of GM1 (Merritt *et al* 1994 Protein Sci 3: 166-175) have revealed that CtxB and EtxB bind to the terminal galactose and sialic acid moieties of GM 1 which can be represented as

Gal β -1-3-3GalNAc

and that such binding does not induce any striking changes in B subunit conformation.

[0020] Furthermore the cholera toxin has been shown to demonstrate an absolute requirement for terminal galactose and internal sialic acid residues (as in GM1) with tolerance for substitution with a second internal sialic acid (as in GD1b).

[0021] Etx, like Ctx also probably binds to the terminal sugar sequence

Gal β 1-3GalNAc β 1-4(NeuAc α 2-3)Gal

where GalNAc is the N-acetylgalactosamine and NeuAc is N-acetylneuraminic acid.

[0022] In addition to binding to GM1, EtxB binds weakly to other gangliosides, including non-galactose containing GM2 and asialo-GM1 as well as galactoproteins (Nashar *et al* Immunology 1997 91: 572-578). Other researchers have shown that EtxB is capable of binding to GM 1 and tolerated removal or extension of the internal sialic acid residue (as in asialo-GM1 and GD1b respectively) but not substitution of the terminal galactose of GM1 (Umesaki and Setoyama 1992 Immunology 75: 386-388).

[0023] In contrast to the poor immunogenicity of the A subunit alone, both EtxB and CtxB are exceptionally potent immunogens and their respective holotoxins, Etx and Ctx, are known to be exceptionally potent adjuvants when given orally in combination with unrelated antigens (Ruedl *et al* 1996 Vaccine 14: 792-798; Nashar *et al* 1993 Vaccine 11: 235; Nashar and Hirst 1995 Vaccine 13: 803; Elson and Ealding 1984 J Immunol 133: 2892; Lycke and Holmgren 1986 Immunology 59: 301). Because of their remarkable immunogenicity, both EtxB and CtxB have been used as carriers for other epitopes and antigens (Nashar *et al* 1993 *ibid*) and have been used as components of vaccines against cholera and E.coli diarrhoea (Jetborn *et al* 1992 Vaccine 10: 130).

[0024] The ability of the B subunit of Ctx and Etx to interact with receptors present on mammalian cells has been shown to exert modulatory effects on the function of those cells. It is known that cells of the immune system are differentially

affected following such interaction. In particular, WO 95/020045 discloses that EtxB binds to GM1 ganglioside receptors which are found on the surfaces of mammalian cells and that this binding induces differential effects on lymphocyte populations including a specific depletion of CD8+ T cells and an associated activation of B cells. These effects are absent when a mutant EtxB protein lacking GM1 binding activity is employed. These observations have led to the use of agents capable of binding to GM1 in the prevention and treatment of autoimmune disease, transplant rejection and graft versus host disease (GVHD). These studies suggest that agents that bind to GM1 or mimic binding to GM1 promote the induction of immunological tolerance.

[0025] Researchers have shown that a state of immunological unresponsiveness, also known as "immunological or oral tolerance", can be induced by the oral administration of dietary protein antigens. (Sun *et al* 1994 *ibid*; Sun *et al* 1996 *ibid*; Bergerot *et al* 1997 *ibid*). The inhalation of antigens can also induce a state of specific immunological unresponsiveness or "nasal tolerance". Thus, systemic immunological tolerance can be induced when antigen is administered orally or nasally by a mucosal route. WO 95/01301 discloses an immunological tolerance-inducing agent comprising a mucosa-binding agent linked to a specific tolerogen. WO95/10301 also includes mention of the treatment of allergy using a mucosa binding agent coupled to an allergen. Other researchers such as Tamura *et al* (1997 *Vaccine* 15: 225-229) have taken directly the protocol of WO 95/10301 and tested its efficacy in preventing allergy in a murine model of Type I allergy. They reported a significant lowering of IgE levels which are a strong predictor of efficacy and they cite data, following administration of EtxB coupled to ovalbumin (the results were not included), which shows that EtxB was not effective once IgE levels are established. It has also been shown that orally administered Ctx and Etx can act on several humoral and cellular immune responses not only at the gastrointestinal tract, but also in distant mucosal effector sites such as the respiratory tract. These data suggest that these mucosal adjuvants have a potential use in oral immunisation strategies to improve the local immune responses in remote mucosal tissues, in accordance with the concept of a common mucosal immune system (Bienenstock J 1974 *The physiology of the local immune system and the gastrointestinal tract*. In: *Progress in Immunology II*, vol 4: clinical aspects, I.L. Brent, J.Holborrow, Eds. Amsterdam, North Holland, pp197-207; Ruedl *et al* 1996 *ibid*; Umesaki 1992 *ibid*; Czerkinsky and Holmgren (1994 *Cell Mol Biol* 40: 37-44).

[0026] The induction of immunological tolerance may include a number of different mechanisms which may be summarised as follows:

- (i) a process whereby antigen reactive cells are removed through triggering them to commit suicide (apoptosis);
- (ii) an induction of anergy or the long term inactivation of the antigen reactive cells;
- (iii) immune deviation of the antigen reactive cells away from the production of pathological responses;
- (iv) suppression of the antigen reactive cells or their regulation by specific factors or regulatory cells

[0027] In the treatment of allergy, it is possible that the induction of any of these mechanisms may be useful. However, while the deletion of antigen reactive cells and/or the induction of anergy are useful strategies once the precise allergens are known, invoking these mechanisms will usually silence only those cells which respond to the allergen which was given in the treatment regime. On the other hand, the implementation of immune deviation or suppression strategies has the advantage of potential regulation of responses to antigens which are involved in the condition but were not part of the treatment. This phenomenon, known as "bystander suppression" allows the "spread" of tolerance to other antigens (such as allergens) in the target tissues through either the possible secretion of non-antigen specific suppressor molecules or through suppressive cellular interactions in that tissue as a result of the interaction between the antigen specific cells and the specific immunising antigen. In this way, as long as at least one of the antigens involved in the disorder is known, the condition may be treated even if there are other antigens implicated as well. Thus, the goal of a good treatment is the induction of a specific immune deviation or suppression.

[0028] Nashar and co-workers (Proc Natl Acad Sci 1996 93: 223-226; Int Immunol 1996 8: 731-736; Immunol 1997 91: 572-578) have demonstrated that the administration of EtxB and other homologues can modulate the immune response away from the production of Th1 cytokines such as IFN γ and interleukin 2 (IL-2) and towards the secretion of Th2 cytokines such as IL-4, IL-10 and IL-13. IFN γ is the classical Th1 cytokine, IL-4 is the classical Th2 cytokine. This "immune deviation" is the basis of the disclosure in WO 97/02045 and has been shown to be effective in the treatment of autoimmune diseases. The experimental results in WO 97/02045 would suggest that GM1 binding agents would not find use in the treatment of allergic conditions and/or hypersensitivity conditions since such conditions involve IgE, the production of which is generally accepted to be promoted by IL-4 and down regulated by IFN γ .

[0029] The present invention now seeks to provide new ways of treating allergic conditions and/or hypersensitivity conditions through the induction of a specific immune deviation or suppression.

[0030] According to a first aspect of the present invention, there is provided the use of an agent in the manufacture of a medicament for use in the treatment and/or prevention of an allergic condition and/or a hypersensitivity condition;

wherein the agent is selected from Etx, Ctx and EtxB; and wherein the agent is not coupled to an antigen.

[0031] There is also provided the use of an agent in the manufacture of a medicament for use in the treatment and/or prevention of an allergic condition; wherein the agent is CtxB; and wherein the agent is not linked to an antigen.

[0032] Preferably the subject is a human - e.g. a human patient.

[0033] Preferably the agent is EtxB.

[0034] In a particularly preferred embodiment the agent is the wild type EtxB.

[0035] Alternatively, preferably the agent is either a mutant of EtxB which is capable of modulating a ganglioside associated activity or other equivalent proteins thereof

[0036] Preferably the agent(s) is/are non-toxic.

[0037] Preferably the agent is CtxB and mutants thereof which are capable of modulating a ganglioside associated activity.

[0038] Preferably the medicament is used for the treatment or prophylaxis of a Type I and/or a Type IV allergic condition as defined herein and/or other hypersensitivity conditions such as contact hypersensitivity.

[0039] Preferably the medicament includes one or more antigens which are optionally co-administered with antigen.

[0040] Preferably the agent may be administered to a mammal with or without coadministration of an antigen.

[0041] Preferably the mammal is a human - e.g. a human patient.

[0042] We have surprisingly found that the use of agents capable of modulating a ganglioside associated activity, when given alone or when co-administered with suitable antigens, can be used as an effective treatment for allergic and/or hypersensitivity conditions. Previous workers have either not attempted to find a mechanism (Sun *et al* 1996 *ibid*) or have argued that agents capable of modulating a ganglioside associated activity, such as EtxB and CtxB, cause a Th1 to Th2 switch in the immune response to antigen (WO97/02045). Since allergic conditions are known in the art to be promoted by Th2 responses, then the previous findings suggest that such agents would either be ineffective in treating allergies or may even worsen them.

[0043] We have surprisingly found that while EtxB and CtxB promote some aspects of Th2-associated responses, in some cases, they may not stimulate the production of the key factor in triggering allergy, IgE. Thus allergic conditions and/or hypersensitivity conditions can be treated with an agent capable of modulating a ganglioside associated activity, for instance, which is not coupled with an antigen.

[0044] Significantly, the linkage of the components was not found to be necessary. Furthermore, our findings indicate that the mechanisms of protection against allergic conditions and/or hypersensitivity conditions may include, though not be limited to either the suppression of antigen specific IgE secretion and/or the upregulated production of non-inflammatory antigen specific antibody isotypes (particularly IgG and IgA).

[0045] The term "ganglioside" as used with respect to the present invention include its normal definition in the art (such as that defined above) as well as active fragments thereof.

[0046] The ganglioside can be made synthetically or isolated from natural sources. Alternatively, it can be obtained from commercial sources.

[0047] The term "ganglioside associated activity" includes any one or more of modulating or immunomodulating a ganglioside receptor, modulating any signalling event prior to, during or subsequent to ganglioside receptor binding.

[0048] The term "Ctx" refers to the cholera toxin and CtxB refers to the B subunit of the cholera toxin. In other texts, these may sometimes be identified as CT or Ct or CTB or CtB respectively.

[0049] The term "Etx" herein means the *E. coli* heat labile enterotoxin and EtxB is the B subunit of Etx. In other texts, these may sometimes be identified as LT or Lt and LTB or LtB respectively.

[0050] The term "adjuvant" includes a substance that enhances an immune response to an antigen.

[0051] The term "mucosal adjuvant" includes an agent which is delivered mucosally with an unrelated antigen, such that the agent is capable of facilitating a mucosal immune response to the unrelated antigen. In this way, the agent acts as a so-called mucosal adjuvant.

[0052] The term "mucosal surfaces" includes but is not limited to oral, sublingual, intranasal, vaginal, rectal, salivary, intestinal and conjunctival surfaces.

[0053] The term "mucosal membrane" and/or "mucosal tissue" includes but is not limited to the intestine, the lung, the mouth, the genital tract, the nose and the eye.

[0054] A "vaccine carrier" includes a carrier of relevant antigens (Szostak *et al* 1996 J Biotechnol 44: 161-170)

[0055] The term "mucosal immunogen" includes an agent administerable by a mucosal route that has the capability to evoke local and/or systemic antibody production and/or cell mediated immune reactions and/or delayed type hypersensitivity reactions.

[0056] A "hapten" means a small molecule which can act as an epitope but is incapable by itself of eliciting an antibody response.

[0057] The term "immunological or oral tolerance" means a reduction in immunological reactivity of a host towards a specific tolerated antigen(s). Immunological or oral tolerance may not mean a complete suppression of the immune response to a particular antigen but it is a form of tolerance also known as "immune deviation" or "split tolerance".

EP 1 044 014 B9

[0058] The term "immune deviation" or "split tolerance" can be used to describe the likely preservation of local IgA responses with the retention of some IgG responses but with the down regulation of delayed hypersensitivity and/or IgE responses.

[0059] The term "tolerance" means a state of specific immunological unresponsiveness.

5 **[0060]** A "tolerogen" means a tolerated antigen.

[0061] The term "autoimmunity" is used to describe the process by which the body generates an immune response to self-antigens.

[0062] The term "agent capable of modulating a ganglioside associated activity" can be used to describe any agent which acts as an immunomodulator through interacting with a ganglioside.

10 **[0063]** The term "GM 1 binding agent" includes any agent which acts as an immunomodulator through interacting with a GM1 ganglioside receptor.

[0064] The term "immunomodulator" includes any agent that alters the extent of the immune response to an antigen, by altering the antigenicity of the antigen or by altering in a nonspecific manner the specific reactivity or the nonspecific effector associated mechanisms of the host.

15 **[0065]** The term "administered" includes delivery by viral or non-viral techniques. Viral delivery mechanisms include but are not limited to adenoviral vectors, adeno-associated viral (AAV) vectors, herpes viral vectors, retroviral vectors, lentiviral vectors, and baculoviral vectors. Non-viral delivery mechanisms include lipid mediated transfection, liposomes, immunoliposomes, lipofectin, cationic facial amphiphiles (CFAs) and combinations thereof. The routes for such delivery mechanisms include but are not limited to mucosal, nasal, oral, parenteral, gastrointestinal, topical, or sublingual routes.

20 **[0066]** The term "co-administered" means that the site and time of administration of each of the agent and the antigen are such that the necessary modulation of the immune system is achieved. Thus, whilst the agent and the antigen may be administered at the same moment in time and at the same site, there may be advantages in administering the agent at a different time and to a different site from the antigen. The agent and antigen may even be delivered in the same delivery vehicle (such as Macrosol™ - see WO95/13795 and WO96/14871) - but with the proviso that the agent and the antigen are uncoupled.

25 **[0067]** The term "administered" includes but is not limited to delivery by a mucosal route, for example, as a nasal spray or aerosol for inhalation or as an ingestible solution; a parenteral route where delivery is by an injectable form, such as, for example, an intravenous, intramuscular or subcutaneous route.

30 **[0068]** The term "systemic immunisation" means the introduction of an antigen into a non-mucosal tissue such as the skin or the blood.

[0069] The term "self antigens" means components derived from host tissues.

[0070] The term "target interaction components" includes but is not limited to an agent capable of modulating a ganglioside associated activity, a ganglioside and/or an antigen.

35 **[0071]** The term "coupled" - which is synonymous with the term "linked" - means the linkage of the agent with the antigen - which includes but is not limited to direct linkage (such as by an ionic or covalent bond) or indirect linkage by the provision of suitable spacer groups.

[0072] The term "uncoupled" - which is synonymous with the term "unlinked" - means that the agent is not coupled to the antigen.

[0073] However, in accordance with the present invention, the agent and/or antigen can be coupled to another entity.

40 **[0074]** The term "affect" includes modulation, such as treatment, prevention, suppression, alleviation, restoration or other alteration of pre-existing condition and/or to potentially affect a future condition, as well as any combination thereof.

[0075] An "antigen" means an agent which, when introduced into an immunocompetent animal, stimulates the production of a specific antibody or antibodies that can combine with the agent. The antigen may be a pure substance, a mixture of substance or soluble or particulate material (including cells or cell fragments). In this sense, the term includes any suitable antigenic determinant, auto-antigen, self-antigen, tolerogen, allergen, hapten, and immunogen, or parts thereof, as well as any combination thereof, and these terms are used interchangeably throughout the text.

[0076] An "allergen" includes any antigen that stimulates an allergic reaction, inducing a Type I hypersensitivity reaction.

[0077] Examples of common allergen sources are outlined in the Table below.

50

Group	Examples of Allergens
Airborne	
grass pollens	ragweed, rye, couch, wild oat, timothy, Bermuda, Kentucky blue, mugwort
tree pollens	alder, birch, hazel, beech, Cupressae, oak, olive
55 moulds	<i>Aspergillus</i> spp., <i>Cladosporium</i> spp., <i>Alternaria</i> spp., Basidiospores, Ascomycetes
cereal grains	wheat, rye,
animal dander and urine	oat cat, dog, horse, rabbit, guinea pig, hamster

(continued)

Group	Examples of Allergens
Airborne bird feathers house dust mite insects	budgerigar, parrot, pigeon, duck, chicken <i>Dermatophagoides pteronyssinus</i> , <i>D.farinae</i> , <i>Euroglyphus maynei</i> cockroach, fly, locust, midge
Oral foods drugs	seafood, legumes, peanuts, nuts, cereals, dairy products, eggs, fruits, tomatoes, mushrooms, alcoholic beverages, coffee, chocolate penicillins, sulphonamides and other antibiotics, sulphasalazine, carbamazepine
Injected insects drugs	bee and wasp stings, ant and mosquito bites blood products, sera, vaccines, contrast media, drugs (including anti-asthma drugs and antibiotics)

[0078] The term "allergic condition" includes but is not limited to asthma, allergic cough, allergic rhinitis and conjunctivitis, atopic eczema and dermatitis, urticaria, hives, insect bite allergy, dietary allergy (peanut, fish milk, wheat etc) and drug allergies

[0079] The term "hypersensitivity condition" includes but is not limited to conditions such as contact hypersensitivity induced by plant poison ivy.

[0080] The term "agent" includes entities capable of modulating a ganglioside associated activity. The agent can be one or more of an inorganic or organic chemical, as well as combinations thereof. By way of example the agent can be a polypeptide as well as a variant/homologue/derivative/fragment thereof so long as they retain the required immunomodulatory activity. It also includes mimics and equivalents and mutants thereof. Other agents for the treatment of allergic conditions or hypersensitivity conditions include antibodies to the target interaction components. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, Fab fragments, fragments produced by a Fab expression library and specifically designed humanised monoclonal antibodies.

[0081] Agents capable of modulating a ganglioside associated activity may be designed and produced as outlined above, by methods which are known in the art. By way of example, when the agent of the invention is a protein such as the EtxB subunit or the CtxB subunit, it may be produced, for use in all aspects of this invention by a method in which the gene or genes coding for the specific polypeptide chain (or chains) from which the protein is formed, is inserted into a suitable vector and then used to transfect a suitable host. For example, the gene coding for the polypeptide chain from which the EtxB assemble may be inserted into, for example, plasmid pMM68, which is then used to transfect host cells, such as *Vibrio sp.60*. The protein is purified and isolated in a manner known *per se*. Mutant genes expressing active mutant EtxB protein may then be produced by known methods from the wild type gene.

[0082] Where a target interaction component is a protein, procedures well known in the art may be used for the production of antibodies to that component.

[0083] For the production of antibodies, various hosts including goats, rabbits, rats, mice, etc. may be immunized by injection with the target interaction component or any derivative or homologue thereof or oligopeptide which retains immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminium hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, and dinitrophenol. BCG (*Bacilli Calmette-Guerin*) and *Corynebacterium parvum* are potentially useful human adjuvants.

[0084] Where a target interaction component is a protein, monoclonal antibodies to that component may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique originally described by Koehler and Milstein (1975 Nature 256: 495-497), the human B-cell hybridoma technique (Kosbor *et al* (1983) Immunol Today 4:72; Cote *et al* (1983) Proc Natl Acad Sci 80:2026-2030) and the EBV-hybridoma technique (Cole *et al* (1985) Monoclonal Antibodies and Cancer Therapy, Alan R Liss Inc, pp 77-96). In addition, techniques developed for the production of "chimeric antibodies", the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity can be used (Morrison *et al* (1984) Proc Natl Acad Sci 81:6851-6855; Neuberger *et al* (1984) Nature 312:604-608; Takeda *et al* (1985) Nature 314:452-454). Alternatively, techniques described for the production of single chain antibodies (US Patent No. 4,946,779) can be adapted to produce target interaction component specific single chain antibodies.

[0085] Antibodies may also be produced by inducing *in vivo* production in the lymphocyte population or by screening

recombinant immunoglobulin libraries or panels of highly specific binding reagents as disclosed in Orlandi *et al* (1989, Proc Natl Acad Sci 86: 3833-3837), and Winter G and Milstein C (1991; Nature 349:293-299).

5 [0086] Antibody fragments which contain specific binding sites for a target interaction components may also be generated. For example, such fragments include, but are not limited to, the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by inducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity (Huse WD *et al* (1989) Science 256:1275-1281).

10 [0087] The target interaction components or a derivative or homologue thereof and/or a cell line that expresses the target interaction components or a derivative or homologue thereof may be used to screen for antibodies, peptides, or other agent, such as organic or inorganic molecules, that act as modulators of the target interaction, thereby identifying a therapeutic agent capable of modulating the target interaction. For example, antibodies capable of modulating the target interaction may be identified.

15 [0088] Alternatively, screening of peptide libraries or organic libraries made by combinatorial chemistry with recombinantly expressed target interaction components or a derivative or homologue thereof or cell lines expressing the target interaction components or a derivative or homologue thereof may be useful for identification of therapeutic agents that function by modulating the target interaction. Synthetic compounds, natural products, and other sources of potentially biologically active materials can be screened in a number of ways deemed to be routine to those of skill in the art.

20 [0089] A target interaction component polypeptide, its immunogenic fragments or oligopeptides thereof can be used for screening therapeutic compounds in any of a variety of drug screening techniques. The polypeptide employed in such a test may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The abolition of activity or the formation of binding complexes between the target interaction component and the agent being tested may be measured.

[0090] Alternatively, phage display can be employed in the identification of candidate agents which affect the target interaction components.

25 [0091] Phage display is a protocol of molecular screening which utilises recombinant bacteriophage. The technology involves transforming bacteriophage with a gene that encodes an appropriate ligand (in this case a candidate agent) capable of reacting with a target interaction component (or a derivative or homologue thereof) or the nucleotide sequence (or a derivative or homologue thereof) encoding same. The transformed bacteriophage (which preferably is tethered to a solid support) expresses the appropriate ligand (such as the candidate agent) and displays it on their phage coat. The entity or entities (such as cells) bearing the target molecules which recognises the candidate agent are isolated and amplified. The successful candidate agents are then characterised. Phage display has advantages over standard affinity ligand screening technologies. The phage surface displays the candidate agent in a three dimensional configuration, more closely resembling its naturally occurring conformation. This allows for more specific and higher affinity binding for screening purposes.

30 [0092] A method for screening a plurality of agents for specific binding affinity with the target interaction component or a derivative or homologue thereof comprising may comprise the following steps: providing a plurality of agents; combining the target interaction components or a derivative or homologue thereof with each of a plurality of agents for a time sufficient to allow binding under suitable conditions; and detecting binding of the target interaction components, or a derivative or homologue thereof to each of the plurality of agents, thereby identifying the agent or agents which specifically bind the target interaction components, In such an assay, the plurality of agents may be produced by combinatorial chemistry techniques known to those of skill in the art.

35 [0093] Another technique for screening provides for high throughput screening of agents having suitable binding affinity to the target interaction components polypeptides and is based upon the method described in detail in WO 84/03564. In summary, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The peptide test agents are reacted with the target interaction component fragments and washed. A bound target interaction component is then detected - such as by appropriately adapting methods well known in the art. A purified target interaction component can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

40 [0094] The agent used in the present invention may be part of a pharmaceutical composition also comprising a pharmaceutically acceptable carrier, diluent, excipient or adjuvant for treating a subject in need of same by administration of a therapeutically effective amount.

45 [0095] The pharmaceutical compositions may be for human or animal usage and will typically comprise any one or more of a pharmaceutically acceptable diluent, carrier, excipient or adjuvant. The choice of pharmaceutical carrier, excipient or diluent can be selected with regard to the intended route of administration and standard pharmaceutical practice. The pharmaceutical compositions may comprise as - or in addition to - the carrier, excipient or diluent any suitable binder(s), lubricant(s), suspending agent(s), coating agent(s), solubilising agent(s).

50 [0096] The pharmaceutical composition may be formulated together with an appropriate antigen.

[0097] Alternatively, a kit may be provided comprising separate compositions for each of the therapeutic agent and the antigen.

[0098] In use, the medicament may be formulated to be delivered by a mucosal route, for example, as a nasal spray or aerosol for inhalation or ingestible solution, or parenterally in which the composition is formulated by an injectable form, for delivery, by, for example, an intravenous, intramuscular or subcutaneous route. Alternatively, the medicament may be designed to be delivered by both routes.

[0099] Where the agent is delivered mucosally through the gastrointestinal mucosa, it is preferably stable during transit through the gastrointestinal tract; for example, it is preferably resistant to proteolytic degradation, stable at acid pH and resistant to the detergent effects of bile.

[0100] Typically, a physician will determine the actual dosage which will be most suitable for a subject and it will vary with the age, weight and response of the particular subject. While a single dose of the agent and optionally the antigenic determinant may be satisfactory, multiple doses are contemplated within the scope of the invention.

[0101] Where appropriate, the medicament can be administered by inhalation, in the form of a suppository or pessary, topically in the form of a lotion, solution, cream, ointment or dusting powder, by use of a skin patch, orally in the form of tablets containing excipients such as starch or lactose, or in capsules or ovules either alone or in admixture with excipients, or in the form of elixirs, solutions or suspensions containing flavouring or colouring agents, or they can be injected parenterally, for example intracavernosally, intravenously, intramuscularly or subcutaneously. For parenteral administration, the medicament may be best used in the form of a sterile aqueous solution which may contain other substances, for example enough salts or monosaccharides to make the solution isotonic with blood. For buccal or sublingual administration the medicament may be administered in the form of tablets or lozenges which can be formulated in a conventional manner.

[0102] There may be different delivery requirements dependent on the different composition/formulation systems.

[0103] Expression vectors derived from retroviruses, adenovirus, herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of the agent to the targeted tissue and/or cell population. Methods which are well known to those skilled in the art can be used to construct recombinant vectors containing the agent. Alternatively, the agent can be delivered to target cells in liposomes.

[0104] By way of example, the controlled release of antigens on mucosal surfaces using biodegradable polymer microspheres may help to target antigens and reduce the numbers of doses required for primary immunisation (Gupta and Siber 1995 Vaccine 13: 1263-1276).

[0105] Encapsulation of vaccines in biodegradable microspheres provides excellent mucosal immunogens. Recombinant Norwalk Virus-like (rNV) particles may also be used for mucosal antigen delivery (Ball *et al* 1996 Arch Virol Suppl 12:243-249).

[0106] Viral Like Particles (VLPs) have been utilised as vaccine delivery system for multiple immunogens including B and T cell epitopes (Roy 1996 Intervirology 39: 62-71).

[0107] One preferred method of oral delivery uses formulations as described in WO95/13795, WO96/17593 and WO96/17594. These formulations allow macromolecules such as proteins to be solubilised in "oils" for oral delivery. Such formulations therefore allow delivery of the macromolecules to mucosal surfaces in the gut.

[0108] In a further approach, it is possible to deliver the agent by means of a bacterial delivery system such as that described in WO 93/17117. This system utilises the bacterium *Lactococcus lactis* to deliver proteins, for instance orally or indeed by other mucosal routes such as nasally.

[0109] In another broad aspect, the present invention provides an immunological tolerance inducing agent comprising an agent capable of modulating a ganglioside associated activity which is not coupled to an antigen.

[0110] Other aspects of the present invention are now presented below by way of numbered paragraphs, which include:

1. The use of an agent having GM1 binding activity, or an agent having an effect on GM1 mediated intracellular signalling events, but no GM1 binding activity, in the preparation of a medicament to treat an allergic or other hypersensitive condition, with the proviso that said agent is not coupled with an allergen and/or an antigen.

2. The use as defined in paragraph 1 wherein the agent is a GM1 binding agent such as Ctx, Etx, CtxB or EtxB or a mutant form or derivative thereof.

3. The use as defined in paragraph 1 and paragraph 2 wherein the medicament is for the prophylaxis or treatment of asthma, allergic cough, allergic rhinitis, conjunctivitis, atopic eczema, dermatitis, urticaria, hives, insect bite allergy, dietary allergy (peanut, fish, milk, wheat etc), drug allergies or contact and other hypersensitivities.

4. A method for the treatment or prophylaxis of an allergic or other hypersensitive condition which comprises administering to a subject an effective amount of an agent having GM1 binding activity, or an agent having an effect on GM1 mediated intracellular signalling events, but no GM1 binding activity, with the proviso that said agent is not

coupled with an allergen and/or an antigen.

5. A method as defined in paragraph 4 wherein the agent is a GM1 binding agent such as Ctx, Etx, CtxB or EtxB or a mutant form or derivative thereof.

6. A method as defined in paragraph 4 or paragraph 5 wherein the method is for the prophylaxis or treatment of asthma, allergic cough, allergic rhinitis, conjunctivitis, atopic eczema, dermatitis, urticaria, hives, insect bite allergy, dietary allergy (peanut, fish, milk, wheat etc), drug allergies or contact hypersensitivity.

7. A pharmaceutical composition for the treatment of a human allergic and/or hypersensitivity disease comprising

(i) an agent having GM1 binding activity; or

(ii) an agent having an effect on GM1 mediated intracellular signalling events, but no GM1 binding activity;

with the proviso that the agent is not coupled with an allergen/antigen; and a pharmaceutically acceptable carrier or diluent therefor.

8. A product comprising an agent having GM1 binding activity, or an agent having an effect on GM1 mediated intracellular signalling events, but no GM1 binding activity, in the preparation of a medicament to treat an allergic or other hypersensitive condition, with the proviso that said agent is not coupled with an allergen and/or an antigen, and at least one antigen/allergen as a combined preparation for simultaneous, separate or sequential use.

[0111] The present invention will now be described only by way of example.

EXAMPLES

Screens for Agents capable of modulating ganglioside associated activity

[0112] Agents capable of modulating ganglioside associated activity are tested by any one of a variety of methods.

[0113] Examples of such methods include, but are not limited to the following methods:

1. Binding to a ganglioside receptor, such as GM1, is determined by using purified GM1 to coat microtiter plates. Following blocking of further non-specific protein binding to the plate, the agent under investigation is applied to the plate and allowed to interact prior to washing and detection with specific antibodies to said agent. Conjugation of the antibodies either directly or indirectly to an enzyme or radiolabel allows subsequent quantification of binding either using colorimetric or radioactivity based methods (ELISA or RIA respectively).

2. The pentasaccharide moiety of a ganglioside, such as GM1, is bound to a suitable column matrix in order to allow standard affinity chromatography to be performed. Removal of known compounds applied to the column from the diluent are used as evidence for binding activity. Alternatively, where mixtures of compounds are applied to the column, elution and subsequent analysis allows the properties of the agent capable of modulating ganglioside associated activity to be determined.

Protein analysis includes peptide sequencing and tryptic digest mapping followed by comparisons with available databases. If eluted proteins cannot be identified in this way, then standard biochemical analysis, such as, for example, mass determination by laser desorption mass spectrometry is used to further characterise the compound. Non-proteins eluted from GMI-affinity columns are analysed by HPLC and mass spectrometry of single homogenous peaks.

3. The ability to bind to gangliosides, such as GM1, and the precise affinity of the interaction may be determined using plasmon surface resonance as previously reported [Kuziemko *et al* (1996) *Biochem* 35:6375-6384].

Evaluation of identified agents

[0114] The identification of agents capable of modulating ganglioside associated activity such that the modulation of the ganglioside associated activity affects an allergic condition and/or a hypersensitivity condition is determined as follows:

[0115] Laboratory animals are stimulated to produce antigen-specific IgE by methods well known in the art. By way of example, mice are challenged with alum precipitated soluble protein antigen (e.g. ovalbumin or allergens known to

be involved in human allergic diseases such as ragweed or house dust mite antigens) either subcutaneously or intraperitoneally.

[0116] In the unmanipulated animal, this procedure routinely leads to the production of antigen-specific IgE which is easily detected in the serum, by standard ELISAs, using the antigen to coat suitable microtiter plates. Serum from the immunised mice is applied to the plates after non-specific protein binding has been blocked and the presence of IgE is determined using widely available labelled antibodies specific for murine IgE.

[0117] In order to screen agents for their capability to prevent or treat allergy, agents capable of modulating ganglioside associated activity are administered to mice either in the presence or absence of the challenge antigen at a range of doses, and by a variety of routes. Although the oral route is the preferred method of administration, delivery can be by other mucosal surfaces or parenterally. The frequency of such administration as well as the timing of repetitive dosing is also investigated. Such intervention strategies are utilised either prior to the IgE inducing antigen challenge (prophylaxis) or after the IgE inducing antigen challenge (treatment). Antigen challenge can be either with (i) the antigen used as part of the prophylactic or treatment protocol; (ii) an unrelated antigen or (iii) a mixture of the challenge and unrelated antigen in order to test the specificity of the response and the induction of bystander suppression respectively.

[0118] Efficacy is determined in a variety of ways and is manifested as a number of different outcomes.

1. Antigen-specific IgE levels. Measurement of serum IgE by specific ELISA (as described) is used to determine whether prophylactic or treatment protocols are capable of reducing levels of serum antigen-specific IgE. Other methods known in the art for the determination of IgE response are used either as alternatives to ELISA or in order to provide complementary data. Such methods include the so-called "Ussing Chamber test" or "passive cutaneous anaphylaxis" assay. A reduction in specific IgE, as determined by any of these assays, is a strong marker of potential clinical efficacy.

2. Antigen specific T-cell reactivity. The responses of T-cells, derived from secondary lymphoid organs of the treated animals to the challenge antigen, is investigated using established methodology. Cell suspensions are prepared and cultured, in the presence or absence of the challenge antigen. At appropriate time intervals after the initiation of the cultures, samples are assessed for cell proliferation and cytokine production.

[0119] Cytokines are measured by specific capture ELISA, by intracellular staining followed by cytometric analysis, by RT-PCR or by other established procedures. Comparison of cell proliferation and cytokine production, in the presence of antigen as opposed to its absence, provides in each case a measure of that part of the response which is specific to the challenge antigen. Evidence of efficacy of prophylactic or treatment protocols is demonstrated by a reduction in the production of Th2 associated cytokines (in particular IL-4) or by an increased expression of cytokines which are involved in down-regulating the allergic response (for example, IL-10 or TGF β).

3. IgG and IgA levels. Protocols which do not reduce the levels of antigen specific IgE can still be considered as potentially effective in the event that they are also able to enhance the production of other non-allergy associated antibody isotopes. Thus investigation of serum and mucosal secretions from animals which have been either untreated or given agents under investigation as part of prophylactic or treatment protocols for the presence of IgG and IgA are also carried out. Standard antigen specific ELISA assays (as described) utilising detecting antibodies specific for IgG and specific subclass thereof, and IgA are used for this purpose. Enhanced production of secreted or serum IgG or IgA antibodies indicate efficacy since such antibodies can be expected to prevent an allergen from cross-linking IgE bound to mast cells, basophils and eosinophils or limit the uptake of antigen across the mucosal epithelium and hence prevent the subsequent allergic inflammatory response.

Enzyme Linked Immunosorbent Assays (ELISAs)

[0120] Binding of EtxB or EtxB (G33D) to GM1 is examined by a GM1-ELISA (Amin, T., & Hirst, T.R. (1994) Prot. Express. and Purif. 5, 198-204).

[0121] Sera and gut secretions are examined for the presence of anti-B subunit IgG and IgA antibodies by ELISAs in which samples are applied to microtitre plates (Immulon 1, Dynateck, USA) coated with 5 μ g/ml of either EtxB or EtxB (G33D) in PBS. Anti-B subunits IgA antibodies in gut secretion supernatants are extrapolated from a standard curve made by coating 2 rows of wells on each plate with 1 μ g/ml rabbit anti-mouse IgA (α chain specific; Zymed Lab, USA) in PBS followed by addition of 1 μ g/ml of mouse myeloma IgA (MOPC 315, Sigma, USA). To measure total IgA, wells are coated with rabbit anti-mouse IgA followed by addition of gut secretion supernatants. All samples are serially diluted. Goat anti-mouse IgG (Fc fragment specific; Jackson Lab., USA) or goat anti-mouse IgA (a chain specific; Sigma) peroxidase conjugate are diluted and added to all wells. The anti-B subunit IgG titer, giving an $A_{450nm} \geq 0.2$, is determined. The IgA anti-B subunit response for each of EtxB and EtxB (G33D) in gut secretions is calculated as "IgA specific activity"

[mean IgA anti-B subunit ($\mu\text{g/ml}$)/total IgA ($\mu\text{g/ml}$)].

[0122] A known ELISA method for measuring cytokine levels of IL-2, IL-4, IL-5, IL-10 and IFN- γ is used. Briefly, microtiter plates are coated with rat antibodies to mouse IL-2, IL-4, IL-5, IL-10 and IFN- γ . Plates are blocked with 2% (w/v) bovine serum albumin. Supernatants from culture medium are added to wells and diluted down. One row on each plate for each cytokine contains a standard amount of recombinant cytokines. Plates are then incubated with 0.5 $\mu\text{g/ml}$ of biotinylated anti-cytokine monoclonal antibodies followed by addition of avidine-peroxidase and 3,3', 5,5'-Tetramethylbenzidine (TMB) substrate and read at $A_{450\text{nm}}$.

Claims

1. The use of an agent in the manufacture of a medicament for use in the treatment and/or prevention of an allergic condition and/or a hypersensitivity condition; wherein the agent is selected from Enterotoxin (Etx), Cholera toxin (Ctx) and Enterotoxin Subunit B (CtxB); and wherein the agent is not linked to an antigen.
2. The use according to claim 1, wherein the agent is CtxB.
3. The use of an agent in the manufacture of a medicament for use in the treatment and/or prevention of an allergic condition; wherein the agent is cholera toxin Subunit B (CtxB) and wherein the agent is not linked to an antigen.
4. The use according to any preceding claim, wherein the agent is coadministered with an antigen.
5. The use according to claim 4, wherein the antigen is an allergen.
6. The use according to any preceding claim, wherein the condition is selected from asthma, allergic cough, allergic rhinitis, conjunctivitis, atopic eczema, dermatitis, urticaria, hives, insect bite allergy, dietary allergy, drug allergies or contact hypersensitivity.
7. The use according to claim 6, wherein the condition is asthma.

Patentansprüche

1. Verwendung eines Stoffs bei der Herstellung eines Medikaments zur Behandlung und/oder Prävention einer Allergieerkrankung und/oder einer Hypersensitivitätserkrankung; wobei der Stoff gewählt ist aus Enterotoxin (Etx), Cholera toxin (Ctx) und Enterotoxin Untereinheit B (CtxB); und wobei der Stoff nicht an ein Antigen gekoppelt ist.
2. Verwendung nach Anspruch 1, wobei der Stoff CtxB ist.
3. Verwendung eines Stoffs bei der Herstellung eines Medikaments zur Behandlung und/oder Prävention einer Allergieerkrankung; wobei der Stoff Cholera toxin Untereinheit B (CtxB) ist; und wobei der Stoff nicht an ein Antigen gekoppelt ist.
4. Verwendung nach einem der vorangehenden Ansprüche, wobei der Stoff gemeinsam mit einem Antigen verabreicht wird.
5. Verwendung nach Anspruch 4, wobei das Antigen ein Allergen ist.
6. Verwendung nach einem der vorangehenden Ansprüche, wobei die Erkrankung gewählt ist aus Asthma, allergischer Husten, allergische Rhinitis, Conjunctivitis, atopisches Ekzem, Dermatitis, Urticaria, Nesselausschlag, Insektenbissallergie, Nahrungsmittelallergie, Medikamentenallergien oder Kontakthypersensitivität.
7. Verwendung nach Anspruch 6, wobei die Erkrankung Asthma ist.

Revendications

- 5
1. Utilisation d'un agent dans la fabrication d'un médicament pour utilisation dans le traitement et/ou la prévention d'un état allergique et/ou d'un état d'hypersensibilité ;
dans laquelle l'agent est choisi entre l'Entérotoxine (Etx), la toxine du choléra (Ctx) et la sous-unité B de l'Entérotoxine (EtxB) ; et
dans laquelle l'agent n'est pas lié à un antigène.
- 10
2. Utilisation selon la revendication 1, dans laquelle l'agent est l'EtxB.
- 15
3. Utilisation d'un agent dans la fabrication d'un médicament pour utilisation dans le traitement et/ou la prévention d'un état allergique ;
dans laquelle l'agent est la sous-unité B de la toxine du choléra (CtxB) ; et
dans laquelle l'agent n'est pas lié à un antigène.
- 20
4. Utilisation selon l'une quelconque des revendications précédentes, dans laquelle l'agent est co-administré avec un antigène.
- 25
5. Utilisation selon la revendication 4, dans laquelle l'antigène est un allergène.
- 30
6. Utilisation selon l'une quelconque des revendications précédentes, dans laquelle l'état est choisi parmi l'asthme, la toux allergique, la rhinite allergique, la conjonctivite, l'eczéma atopique, la dermatite, les urticaires, les éruptions, l'allergie due à une morsure d'insecte, l'allergie alimentaire, l'allergie médicamenteuse et l'hypersensibilité aux contacts.
- 35
- 40
- 45
- 50
- 55
7. Utilisation selon la revendication 6, dans laquelle l'état est l'asthme.