



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 133 489 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

Note: Bibliography reflects the latest situation

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see page(s) 41, 42, 45

(48) Corrigendum issued on:
03.05.2006 Bulletin 2006/18

(45) Date of publication and mention
of the grant of the patent:
27.07.2005 Bulletin 2005/30

(21) Application number: **00966823.7**

(22) Date of filing: **22.09.2000**

(51) Int Cl.:
C07D 401/14 ^(1990.01) **C07D 401/12** ^(1990.01)
C07D 401/06 ^(1990.01) **C07D 211/62** ^(1985.01)
C07D 405/14 ^(1990.01) **C07D 409/14** ^(1990.01)
C07D 417/12 ^(1990.01) **A61K 31/454** ^(2000.01)
A61P 7/02 ^(2000.01)

(86) International application number:
PCT/US2000/026163

(87) International publication number:
WO 2001/023376 (05.04.2001 Gazette 2001/14)

(54) **ISONIPECOTAMIDES FOR THE TREATMENT OF INTEGRIN-MEDIATED DISORDERS**

ISONIPECOTAMIDE ZUR BEHANDLUNG VON INTEGRIN VERMITTELTEN STÖRUNGEN

ISONIPECOTAMIDES DESTINEES AU TRAITEMENT DE MALADIES PROVOQUEES PAR
L'INTEGRINE

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

(30) Priority: **29.09.1999 US 156617 P**
21.09.2000 US 667134

(43) Date of publication of application:
19.09.2001 Bulletin 2001/38

(60) Divisional application:
05075325.0 / 1 533 310

(73) Proprietor: **Ortho-McNeil Pharmaceutical, Inc.**
Raritan, NJ 08869-0602 (US)

(72) Inventors:
• **COSTANZO, Michael, J.**
Ivyland, PA 18974 (US)
• **HOEKSTRA, William, J.**
Chapel Hill, NC 27514 (US)
• **MARYANOFF, Bruce, E.**
New Hope, PA 18938 (US)

(74) Representative: **Mercer, Christopher Paul et al**
Carpmaels & Ransford
43, Bloomsbury Square
London WC1A 2RA (GB)

(56) References cited:
WO-A-99/67230 **US-A- 5 700 801**
US-A- 5 922 717

- **VARON, DAVID ET AL: "Inhibition of integrin-mediated platelet aggregation, fibrinogen-binding, and interactions with extracellular matrix by nonpeptidic mimetics of Arg-Gly-Asp" THROMB. HAEMOSTASIS (1993), 70(6), 1030-6 , XP000979521**
- **SU, TING ET AL: "Fibrinogen Receptor (GPIIb-IIIa) Antagonists Derived from 5,6-Bicyclic Templates. Amidinoindoles, Amidinoindazoles, and Amidinobenzofurans Containing the N-.alpha.-Sulfonamide Carboxylic Acid Function as Potent Platelet Aggregation Inhibitors" J. MED. CHEM. (1997), 40(26), 4308-4318 , XP000915289**
- **CHEMICAL ABSTRACTS, vol. 118, no. 25, 21 June 1993 (1993-06-21) Columbus, Ohio, US; abstract no. 255342, HOSODA, AKIHIKO ET AL: "Preparation of N-(heterocyclylcarbonyl)amino acids and analogs as prolyl endopeptidase inhibitors" XP002159418 & JP 04 334357 A (FUJIREBIO, INC., JAPAN) 20 November 1992 (1992-11-20)**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 133 489 B9

Description


[0001] This invention relates to certain novel compounds, their synthesis and methods for their use in treating integrin-mediated disorders. More particularly, this invention relates to isonipecotamide compounds that are $\alpha v\beta 3$, $\alpha v\beta 5$, GPIIb/IIIa, dual $\alpha v\beta 3$ /GPIIb/IIIa or dual $\alpha v\beta 3/\alpha v\beta 5$ integrin antagonists and methods for their use in treating a variety of integrin-mediated disorders.

Background of the Invention

[0002] Integrins are a widely expressed family of calcium- or magnesium-dependent $\alpha\beta$ heterodimeric cell surface receptors which bind to extracellular matrix adhesive proteins such as fibrinogen, fibronectin, vitronectin, and osteopontin. These transmembrane glycoproteins (GP's), known for their large extracellular domains, are classified by at least 8 known β subunits and 14 α subunits (S. A. Mousa, et al., *Emerging Therapeutic Targets*, **2000**, 4, (2), 143-153). For example, the $\beta 1$ subfamily, also known as the very late antigen (VLA) subfamily, has the largest number of integrins (S. A. Mousa, et al., *Emerging Therapeutic Targets*, **2000**, 4, (2), 144). The $\alpha v\beta 1$ subfamily further associates with various β subunits: $\beta 3$, $\beta 5$, $\beta 6$, $\beta 8$ and $\alpha IIb\beta 3$ (also referred to as GPIIb/IIIa) (S. A. Mousa, et al., *Emerging Therapeutic Targets*, **2000**, 4, (2), 144, 147). Some of the disease states that have a strong $\beta 3$, $\beta 5$ and GPIIb/IIIa integrin component in their etiologies are unstable angina, thromboembolic disorders or atherosclerosis (GPIIb/IIIa); thrombosis or restenosis (GPIIb/IIIa or $\alpha v\beta 3$); restenosis (dual $\alpha v\beta 3$ /GPIIb/IIIa); rheumatoid arthritis, vascular disorders or osteoporosis ($\alpha v\beta 3$); tumor angiogenesis, multiple sclerosis, neurological disorders, asthma, vascular injury or diabetic retinopathy ($\alpha v\beta 3$ or $\alpha v\beta 5$); and, angiogenesis (dual $\alpha v\beta 3/\alpha v\beta 5$) (S. A. Mousa, et al., *Emerging Therapeutic Targets*, **2000**, 4, (2), 148-149; W. H. Miller, et al., *Drug Discovery Today* **2000**, 5 (9), 397-407; and, S. A. Mousa, et al., *Exp. Opin. Ther. Patents*, **1999**, 9 (9), 1237-1248). The $\beta 3$ subunit has received significant attention in recent drug discovery efforts. (W. J. Hoekstra, *Current Medicinal Chemistry* **1998**, 5, 195) and antibodies and/or low-molecular weight compound antagonists of $\alpha v\beta 3$ have shown efficacy in animal models (J. Samanen, *Current Pharmaceutical Design* **1997**, 3, 545).

[0003] Furthermore, GPIIb/IIIa and $\alpha v\beta 3$ antagonists have typically been designed after the bioactive arginine-glycine-aspartate (RGD) conformations of peptides derived from their primary ligands, fibrinogen and vitronectin, respectively. The RGD motif is the general cell attachment sequence of many extracellular matrix, blood, and cell surface proteins, as half of the ca. 20 known integrins bind the RGD-containing adhesion ligands. To discover RGD peptides with integrin selectivity, peptides with both restricted conformations and alterations of flanking residues have been studied. In particular, the structural requirements for interaction of the RGD sequence with GPIIb/IIIa and the inhibitory potential of a series of nonpeptidic mimetics on platelet aggregation and interactions with the extracellular matrix have been described (D. Varon, et al., *Thromb. Haemostasis*, **1993**, 70(6), 1030-1036). Iterative synthesis of cyclic and alicyclic peptides and computer modelling have provided potent, selective agents as a platform for nonpeptide αv integrin antagonist design.

[0004] For example, PCT Application WO98/25892 to Fisher, et al., describes a series of α -sulfonamido and α -sulfinamido carboxylic acid compounds of the formula:



wherein


Y is selected from the group consisting of $-\text{COOH}$, $-\text{PO}_3\text{H}_2$, $-\text{SO}_3\text{H}$ and $-\text{COOR}^4$; where R^4 is selected from the group consisting of $\text{C}_{1-10}\text{alkyl}$, $\text{C}_{1-8}\text{alkylaryl}$, $\text{arylC}_{1-8}\text{alkyl}$, $\text{C}_{1-8}\text{alkyloxycarbonyloxyC}_{1-8}\text{alkyl}$, $\text{aryloxycarbonyloxyC}_{1-8}\text{alkyl}$, $\text{C}_{1-8}\text{alkyloxycarbonyloxyaryl}$, $\text{C}_{1-10}\text{alkylcarbonyloxyC}_{1-8}\text{alkyl}$ and $\text{C}_{1-8}\text{alkylcarbonyloxyaryl}$; A is selected from the group consisting of $\text{C}_{6-12}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-NR}^5\text{-CO-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-CO-NR}^5\text{-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-O-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-NR}^5\text{-CO-C}_{1-8}\text{alkyl-NR}^5\text{-CO-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-NR}^5\text{-CO-C}_{1-8}\text{alkyl-CO-NR}^5\text{-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-CO-NR}^5\text{-C}_{1-8}\text{alkyl-NR}^5\text{-CO-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-CO-NR}^5\text{-C}_{0-8}\text{alkyl-CO-NR}^5\text{-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-CO-NR}^5\text{-C}_{0-8}\text{alkyl-CO-NR}^5\text{-C}_{1-8}\text{alkyl-CO-NR}^5\text{-CO-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-O-C}_{2-8}\text{alkyl-NR}^5\text{-CO-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-O-C}_{0-8}\text{alkyl-CO-NR}^5\text{-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-O-C}_{2-8}\text{alkyl-O-C}_{0-8}\text{alkyl-CO-NR}^5\text{-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-S-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-S(O}_n\text{)-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-S-C}_{2-8}\text{alkyl-NR}^5\text{-CO-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-S(O}_n\text{)-C}_{2-8}\text{alkyl-NR}^5\text{-CO-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-S-C}_{0-8}\text{alkyl-CO-NR}^5\text{-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-S(O}_n\text{)-C}_{1-8}\text{alkyl-CO-NR}^5\text{-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-NR}^5\text{-CO-C}_{0-8}\text{alkyl-S-C}_{0-8}\text{alkyl}$, $\text{C}_{0-8}\text{alkyl-NR}^5\text{-CO-C}_{1-8}\text{alkyl-S}$

(O_n)-C₀₋₈alkyl, C₀₋₈alkyl-CO-NR⁵-C₂₋₈alkyl-S-C₀₋₈alkyl, C₀₋₈alkyl-CO-NR⁵-C₂₋₈alkyl-S(O_n)-C₀₋₈alkyl, C₀₋₈alkyl-NR⁵-C₀₋₈alkyl-CO₂-C₀₋₈alkyl, C₀₋₈alkyl-NR⁵-C₀₋₈alkyl-CS-O-C₀₋₈alkyl, C₀₋₈alkyl-NR⁵-C₀₋₈alkyl-CO-NR⁵-C₀₋₈alkyl, C₀₋₈alkyl-NR⁵-C₀₋₈alkyl-CS-NR⁵-C₀₋₈alkyl, C₀₋₈alkyl-O-C₀₋₈alkyl-CO₂-C₀₋₈alkyl, C₀₋₈alkyl-O-C₀₋₈alkyl-CS-O-C₀₋₈alkyl, C₀₋₈alkyl-SiR⁷R⁸-C₀₋₈alkyl, C₀₋₈alkyl-SiR⁷R⁸-C₀₋₈alkyl-NR⁶-CO-C₀₋₈alkyl and C₀₋₈alkyl-SiR⁷R⁸-C₀₋₈alkyl-CO-NR⁶-C₀₋₈alkyl; where R⁵, R⁶, R⁷ and R⁸ are independently selected from the group consisting of H and C₁₋₆alkyl; and where n=1 or 2; Z is selected from the group consisting of -NH-C(NR⁹R¹⁰)=R¹¹, -NH-C(R⁹)=R¹¹, -C(NR⁹R¹⁰)=R¹¹ and piperidinyl; where R⁹, R¹⁰ and R¹¹ are independently selected from the group consisting of H, C₁₋₆alkyl, arylC₁₋₃alkyl and aryl; or where two of the R⁹, R¹⁰ or R¹¹ substituents form a cyclic ring containing (CH₂)_p, where p=2-5; R₁ is H and R₂ is selected from the group consisting of -SO_m-aryl-, -SO_m-C₁₋₁₀alkyl-, -SO_m-heteroaryl-, where m=1-2; R³ is selected from the group consisting of H, C₁₋₈alkyl, aryl, C₁₋₈alkylaryl and heteroaryl as inhibitors of RGD-dependent integrins for the treatment of thrombotic or restenotic disorders.

[0005] Accordingly, it is an object of the present invention to provide compounds that are antagonists of integrins. It is another object to provide isonipicotamide compounds that are αvβ3, αvβ5, GPIIb/IIIa, dual αvβ3/GPIIb/IIIa or dual αvβ3/αvβ5 integrin antagonists. It is a further object to provide methods for treating a variety of integrin-mediated disorders including, but not limited to, unstable angina, thromboembolic disorders, atherosclerosis, arterial and/or venous thrombosis, restenosis, rheumatoid arthritis, vaso-occlusive disorders, osteoporosis, tumor angiogenesis, multiple sclerosis, neurological disorders, asthma, vascular injury, macular degeneration or diabetic complications including diabetic retinopathy.

Summary of the Invention

[0006] The present invention is directed to compounds represented by the following general Formula (I):



wherein

M is selected from the group consisting of ethylene (optionally substituted within the carbon chain with methyl and substituted on the terminal carbon with one substituent selected from A), propylene (optionally substituted within the carbon chain with a substituent selected from the group consisting of methyl, ethenyl, cyclohexylidene [wherein a ring carbon atom forms the point of attachment to the carbon chain] and 4-Cl-phenyl and substituted on the terminal carbon with one substituent selected from A), allylene (substituted with one substituent selected from A); piperidin-4-yl, piperidin-4-ylene (substituted with one substituent selected from A), 1,4,5-dihydro-2-cyclopenten-1-ylene (substituted with one substituent selected from A) and 4-methylphenylene (substituted on methylene with one substituent selected from A); A is optionally present and is selected from the group consisting of 1H-imidazol-1-yl, 1H-imidazol-2-yl, 4,5-dihydro-1H-imidazol-2-yl (optionally substituted with a substituent selected from the group consisting of C₁₋₄ alkoxycarbonyl and aryl(C₁₋₄)alkoxycarbonyl), pyridin-2-yl (optionally substituted with a substituent selected from the group consisting of C₁₋₄ alkyl, heteroaryl [optionally substituted with C₁₋₄ alkyl], halogen, hydroxy, nitro, cyano, amino, amino(C₁₋₄) alkyl, C₁₋₄ alkylamino(C₁₋₄)alkyl and di(C₁₋₄ alkyl)amino(C₁₋₄)alkyl), pyrimidin-2-yl, 1,4,5,6-tetrahydro-pyrimidin-2-yl (optionally substituted with one to two substituents independently selected from the group consisting of C₁₋₄ alkyl, hydroxy and amino), piperidin-4-yl, benzimidazol-2-yl, 1,2,3,4-tetrahydro-1,8-naphthyridin-7-yl, 3H-imidazo[4,5-b]pyridin-2-yl, amino, (C₁₋₆ alkyl)amino, (1H-imidazol-1-yl)amino, (1H-imidazol-2-yl)amino, (4,5-dihydro-1H-imidazol-2-yl) amino (optionally substituted on 4,5-dihydro-1H-imidazol-2-yl with a substituent selected from the group consisting of (C₁₋₆)alkoxycarbonyl and aryl(C₁₋₆)alkoxycarbonyl), (pyridin-2-yl)amino (optionally substituted on pyridin-2-yl with a substituent selected from the group consisting of C₁₋₄ alkyl, heteroaryl [optionally substituted with C₁₋₄ alkyl], halogen, hydroxy, nitro, cyano, amino, amino(C₁₋₄)alkyl, C₁₋₄ alkylamino(C₁₋₄)alkyl and di(C₁₋₄ alkyl)amino(C₁₋₄)alkyl), (pyrimidin-2-yl)amino, (1,4,5,6-tetrahydropyrimidin-2-yl)amino (optionally substituted on 1,4,5,6-tetrahydropyrimidin-2-yl with one to two substituents independently selected from the group consisting of C₁₋₄ alkyl, hydroxy and amino), (4,5,6,7-tetrahydro-1H-1,3-diazepin-2-yl)amino, (thiazol-2-yl)amino, (benzimidazol-2-yl)amino, (3H-imidazo[4,5-b]pyridin-2-yl)amino, R₃HNC(=NH)NH-, R₃HNC(=O)NH-, (4,5,6,7-tetrahydro-1H-1,3-diazepin-2-yl)aminoxy, (4,5-dihydro-

1H-imidazol-2-yl)aminoxy and $R_3\text{HNC(=NH)NHO-}$;

with the proviso that if A is $\text{H}_2\text{NC(=NH)NH-}$, then, dependently, W is not hydrogen when Q is $-\text{CH}_2-$;

L is selected from the group consisting of $-\text{C(=O)-}$, $-\text{OC(=O)-}$ and $-\text{HNC(=O)-}$;

R_1 is selected from the group consisting of hydrogen and $\text{C}_1\text{-C}_4$ alkyl;

5 R_2 is hydrogen;

R_3 is selected from the group consisting of hydrogen, $\text{C}_1\text{-C}_8$ alkyl, aryl($\text{C}_1\text{-C}_8$)alkyl and hydroxy;

Q is selected from the group consisting of $-\text{CH}_2-$, $-\text{CH}(\text{C}_1\text{-C}_8\text{alkyl})-$, $-\text{CH}(\text{aryl})-$ (wherein aryl is optionally substituted with one to five substituents independently selected from the group consisting of $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy, $-\text{O}(\text{C}_1\text{-C}_3\text{alkyl})-\text{O-}$, halogen, hydroxy and trihalo($\text{C}_1\text{-C}_3$)alkyl), $-\text{CH}(\text{heteroaryl})-$ (wherein heteroaryl is optionally substituted with a substituent selected from halogen) and $-\text{CH}(\text{aryl}(\text{C}_1\text{-C}_8)\text{alkyl})-$;

10 W is selected from the group consisting of hydrogen and $\text{N}(\text{R}_4)\text{T}$;

r is 1;

R_4 is selected from the group consisting of hydrogen and $\text{C}_1\text{-C}_4$ alkyl;

T is selected from the group consisting of $\text{R}_5\text{C(=O)-}$ and $\text{R}_5\text{OC(=O)-}$;

15 R_5 is selected from the group consisting of aryl and aryl($\text{C}_1\text{-C}_4$)alkyl;

R_6 is selected from the group consisting of hydrogen, methyl and $(\text{R}_8)(\text{R}_7)\text{NC(=O)CH}_2$; and,

R_7 and R_8 are independently selected from the group consisting of hydrogen, methyl and ethyl; and pharmaceutically acceptable salts thereof.

20 3. The compound of Claim 1, wherein

[0007] A is optionally present and is selected from the group consisting of 1H-imidazol-1-yl, 4,5-dihydro-1H-imidazol-2-yl, pyridin-2-yl (optionally substituted with a substituent selected from the group consisting of $\text{C}_1\text{-C}_4$ alkyl and heteroaryl [optionally substituted with $\text{C}_1\text{-C}_4$ alkyl]), piperidin-4-yl, benzimidazol-2-yl, 1,2,3,4-tetrahydro-1,8-naphthyridin-7-yl, 3H-imidazo[4,5-b]pyridin-2-yl, amino, (4,5-dihydro-1H-imidazol-2-yl)amino (optionally substituted on 4,5-dihydro-1H-imidazol-2-yl with $\text{C}_1\text{-C}_6$ alkoxy carbonyl), (pyridin-2-yl)amino (optionally substituted on pyridinyl with a substituent selected from the group consisting of $\text{C}_1\text{-C}_4$ alkyl and heteroaryl [optionally substituted with $\text{C}_1\text{-C}_4$ alkyl]), (1,4,5,6-tetrahydro-5-hydroxypyrimidin-2-yl)amino, (1,4,5,6-tetrahydro-5-methylpyrimidin-2-yl)amino, (1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-yl)amino, (thiazol-2-yl)amino, (3H-imidazo[4,5-b]pyridin-2-yl)amino, $\text{R}_3\text{HNC(=NH)NH-}$, $\text{R}_3\text{HNC(=O)NH-}$, (4,5-dihydro-1H-imidazol-2-yl)aminoxy and $\text{R}_3\text{HNC(=NH)NHO-}$;

with the proviso that if A is $\text{H}_2\text{NC(=NH)NH-}$, then, dependently, W is not hydrogen when Q is $-\text{CH}_2-$;

R_1 is hydrogen;

R_3 is selected from the group consisting of hydrogen, butyl, benzyl and hydroxy;

35 Q is selected from the group consisting of $-\text{CH}_2-$, $-\text{CH}(\text{methyl})-$, $-\text{CH}(\text{ethyl})-$, $-\text{CH}(\text{phenyl})-$ (wherein phenyl is optionally substituted with one to five substituents independently selected from the group consisting of methyl, ethyl, propyl, methoxy, ethoxy, propoxy, bromine, chlorine, fluorine, iodine, hydroxy and trifluoromethyl), $-\text{CH}(\text{naphthalen-1-yl})-$, $-\text{CH}(\text{naphthalen-2-yl})-$, $-\text{CH}[(3,4\text{-dioxymethylene})\text{phenyl}]-$, $-\text{CH}[(3,4\text{-dioxylethylene})\text{phenyl}]-$, $-\text{CH}[(3\text{-bromo-5-chloro-2-hydroxy})\text{phenyl}]-$, $-\text{CH}(\text{thien-3-yl})-$, $-\text{CH}(\text{quinolin-3-yl})-$, $-\text{CH}(\text{pyridin-3-yl})-$ (wherein pyridinyl is optionally substituted with chlorine) and $-\text{CH}(\text{benzyl})-$;

40 R_4 is hydrogen;

T is selected from $\text{R}_5\text{OC(=O)-}$;

R_5 is selected from aryl($\text{C}_1\text{-C}_4$)alkyl;

R_6 is hydrogen; and,

R_7 and R_8 are independently ethyl;

45 and pharmaceutically acceptable salts thereof.

[0008] Illustrative of the invention is a pharmaceutical composition comprising a pharmaceutically acceptable carrier and any of the compounds described above. Illustrating the invention is a pharmaceutical composition made by mixing any of the compounds described above and a pharmaceutically acceptable carrier. An illustration of the invention is a process for making a pharmaceutical composition comprising mixing any of the compounds described above and a pharmaceutically acceptable carrier.

50 **[0009]** An example of the invention is a method of treating integrin-mediated disorders in a subject in need thereof comprising administering to the subject a therapeutically effective amount of any of the compounds or pharmaceutical compositions described above. Examples of integrin-mediated disorders include, but are not limited to, unstable angina, thromboembolic disorders, atherosclerosis, arterial and/or venous thrombosis, restenosis, rheumatoid arthritis, vaso-occlusive disorders, osteoporosis, tumor angiogenesis, multiple sclerosis, neurological disorders, asthma, vascular injury, macular degeneration or diabetic complications, including diabetic retinopathy.

[0010] Further exemplifying the invention is the method of treating integrin-mediated disorders, wherein the therapeutically effective amount of the compound is from about 0.01 mg/kg/day to about 300 mg/kg/day.

[0011] Also included in the invention is the use of a compound of Formula (I) for the preparation of a medicament for treating an integrin-mediated disorder in a subject in need thereof.

[0012] The isonipecotamide compounds of the present invention are integrin antagonists; particularly, $\alpha_v\beta_3$, $\alpha_v\beta_5$, GPIIb/IIIa, dual $\alpha_v\beta_3$ /GPIIb/IIIa and dual $\alpha_v\beta_3/\alpha_v\beta_5$ integrin antagonists. The instant compounds are useful in treating thrombotic disorders such as restenosis, arterial and venous thrombosis, acute myocardial infarction, re-occlusion following thrombolytic therapy and angioplasty, inflammation, unstable angina, atherosclerosis, angiogenesis and a variety of vaso-occlusive disorders. These compounds are also useful as antithrombotics used in conjunction with fibrinolytic therapy (e.g., t-PA or streptokinase). Additionally, the compounds are useful in treating and preventing osteoporosis, rheumatoid arthritis, bone resorption, cancer, macular degeneration, and diabetic complications including diabetic retinopathy.

Detailed Description of the Invention

[0013] Relative to the above generic description, certain compounds of Formula (I) are preferred.

[0014] The preferred embodiments of the present invention are those compounds wherein, independently, M is selected from ethylene (optionally substituted within the carbon chain with methyl and substituted on the terminal carbon with one substituent selected from A), propylene (optionally substituted within the carbon chain with methyl, ethenyl, cyclohexylidene (wherein a ring carbon atom forms the point of attachment to the carbon chain) or 4-Cl-phenyl and substituted on the terminal carbon with one substituent selected from A), allylene (substituted with one substituent selected from A); piperidin-4-ylene (optionally substituted with one substituent selected from A), 1,4,5-dihydro-2-cyclopenten-1-ylene (substituted with one substituent selected from A) or 4-methylenepheryl (substituted on methylene with one substituent selected from A);

A is optionally present and is selected from 1*H*-imidazol-1-yl, 1*H*-imidazol-2-yl, 4,5-dihydro-1*H*-imidazol-2-yl (optionally substituted with a substituent selected from C₁-C₄ alkoxy carbonyl or aryl(C₁-C₄)alkoxy carbonyl), pyridin-2-yl (optionally substituted with a substituent selected from C₁-C₄ alkyl, heteroaryl (optionally substituted with C₁-C₄ alkyl), halogen, hydroxy, nitro, cyano, amino, amino(C₁-C₄)alkyl, C₁-C₄ alkylamino(C₁-C₄)alkyl or di(C₁-C₄ alkyl)amino(C₁-C₄)alkyl), pyrimidin-2-yl, 1,4,5,6-tetrahydro-pyrimidin-2-yl (optionally substituted with one to two substituents independently selected from C₁-C₄ alkyl, hydroxy or amino), piperidin-4-yl, benzimidazol-2-yl, 1,2,3,4-tetrahydro-1,8-naphthyridin-7-yl, 3*H*-imidazo[4,5-*b*]pyridin-2-yl, amino, (C₁-C₆ alkyl)amino, (1*H*-imidazol-1-yl)amino, (1*H*-imidazol-2-yl)amino, (4,5-dihydro-1*H*-imidazol-2-yl)amino (optionally substituted on 4,5-dihydro-1*H*-imidazol-2-yl with a substituent selected from C₁-C₆ alkoxy carbonyl or aryl(C₁-C₆)alkoxy carbonyl), (pyridin-2-yl)amino (optionally substituted on pyridin-2-yl with a substituent selected from C₁-C₄ alkyl, heteroaryl [optionally substituted with C₁-C₄ alkyl], halogen, hydroxy, nitro, cyano, amino, amino(C₁-C₄)alkyl, C₁-C₄ alkylamino(C₁-C₄)alkyl or di(C₁-C₄ alkyl)amino(C₁-C₄)alkyl), (pyrimidin-2-yl)amino, (1,4,5,6-tetrahydro-pyrimidin-2-yl)amino (optionally substituted on 1,4,5,6-tetrahydro-pyrimidin-2-yl with one to two substituents independently selected from C₁-C₄ alkyl, hydroxy or amino), (4,5,6,7-tetrahydro-1*H*-1,3-diazepin-2-yl)amino, (thiazol-2-yl)amino, (benzimidazol-2-yl)amino, (3*H*-imidazo[4,5-*b*]pyridin-2-yl)amino, R₃HNC(=NH)NH-, R₃HNC(=O)NH-, (4,5,6,7-tetrahydro-1*H*-1,3-diazepin-2-yl)aminooxy, (4,5-dihydro-1*H*-imidazol-2-yl)aminooxy or R₃HNC(=NH)NHO-;

more preferably, A is optionally present and is selected from 1*H*-imidazol-1-yl, 4,5-dihydro-1*H*-imidazol-2-yl, pyridin-2-yl (optionally substituted with a substituent selected from C₁-C₄ alkyl or heteroaryl [optionally substituted with C₁-C₄ alkyl]), piperidin-4-yl, benzimidazol-2-yl, 1,2,3,4-tetrahydro-1,8-naphthyridin-7-yl, 3*H*-imidazo[4,5-*b*]pyridin-2-yl, amino, (4,5-dihydro-1*H*-imidazol-2-yl)amino (optionally substituted on 4,5-dihydro-1*H*-imidazol-2-yl with C₁-C₆ alkoxy carbonyl), (pyridin-2-yl)amino (optionally substituted on pyridinyl with a substituent selected from C₁-C₄ alkyl or heteroaryl [optionally substituted with C₁-C₄ alkyl]), (1,4,5,6-tetrahydro-5-hydroxypyrimidin-2-yl)amino, (1,4,5,6-tetrahydro-5-methylpyrimidin-2-yl)amino, (1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-yl)amino, (thiazol-2-yl)amino, (3*H*-imidazo[4,5-*b*]pyridin-2-yl)amino, R₃HNC(=NH)NH-, R₃HNC(=O)NH-, (4,5-dihydro-1*H*-imidazol-2-yl)aminooxy or R₃HNC(=NH)NHO-;

most preferably, A is optionally present and is selected from 4,5-dihydro-1*H*-imidazol-2-yl, 1,2,3,4-tetrahydro-1,8-naphthyridin-7-yl, 3*H*-imidazo[4,5-*b*]pyridin-2-yl, amino, (4,5-dihydro-1*H*-imidazol-2-yl)amino, (pyridin-2-yl)amino, (3-methylpyridin-2-yl)amino, (thiazol-2-yl)amino, (3*H*-imidazo[4,5-*b*]pyridin-2-yl)amino, NH₂C(=NH)NH-, R₃HNC(=O)NH-, (4,5-dihydro-1*H*-imidazol-2-yl)aminooxy or NH₂C(=NH)NHO-;

with the proviso that if A is H₂NC(=NH)NH-, then, dependently, W is not hydrogen when Q is -CH₂-;

L is selected from -C(=O)-, -OC(=O)- or -HNC(=O)-;

R₁ is selected from hydrogen or C₁-C₄ alkyl;

more preferably, R₁ is hydrogen;

R₂ is hydrogen;

R₃ is selected from hydrogen, C₁-C₈ alkyl, aryl(C₁-C₈)alkyl or hydroxy; more preferably, R₃ is selected from hydrogen, butyl, benzyl or hydroxy;

Q is selected from -CH₂-, -CH(C₁-C₈alkyl)-, -CH(aryl)- (wherein aryl is optionally substituted with one to five substituents

independently selected from C₁-C₄ alkyl, C₁-C₄ alkoxy, -O-(C₁-C₃ alkyl)-O-, halogen, hydroxy or trihalo(C₁-C₃)alkyl), -CH(heteroaryl)- (wherein heteroaryl is optionally substituted with a substituent selected from halogen) or -CH(aryl(C₁-C₈)alkyl)-;

more preferably, Q is selected from -CH₂-, -CH(methyl)-, -CH(ethyl)-, -CH(phenyl)- (wherein phenyl is optionally substituted with one to five substituents independently selected from methyl, ethyl, propyl, methoxy, ethoxy, propoxy, bromine, chlorine, fluorine, iodine, hydroxy or trifluoromethyl), -CH(naphthalen-1-yl)-, -CH(naphthalen-2-yl)-, -CH[(3,4-dioxymethylene)phenyl]-, -CH[(3,4-dioxyethylene)phenyl]-, -CH[(3-bromo-5-chloro-2-hydroxy)phenyl]-, -CH(thien-3-yl)-, -CH(quinolin-3-yl)-, -CH(pyridin-3-yl)- (wherein pyridinyl is optionally substituted with chlorine) or -CH(benzyl)-;

most preferably, Q is selected from -CH₂-, -CH(methyl)-, -CH(phenyl)- (wherein phenyl is optionally substituted with one to five substituents independently selected from methyl, methoxy, bromine, chlorine, fluorine, hydroxy or trifluoromethyl), -CH(naphthalen-1-yl)-, -CH(naphthalen-2-yl)-, -CH[(3,4-dioxymethylene)phenyl]-, -CH[(3,4-dioxyethylene)phenyl]-, -CH[(3-bromo-5-chloro-2-hydroxy)phenyl]-, -CH(thien-3-yl)-, -CH(quinolin-3-yl)-, -CH(pyridin-3-yl)- (wherein pyridinyl is optionally substituted with chlorine) or -CH(benzyl)-;

r is 1;

R₄ is selected from hydrogen or C₁-C₄ alkyl;

more preferably, R₄ is hydrogen;

T is selected from R₅C(=O)- or R₅OC(=O)-;

more preferably, T is selected from R₅OC(=O)-;

R₅ is selected from aryl or aryl(C₁-C₄)alkyl;

more preferably, R₅ is selected from aryl(C₁-C₄)alkyl;

most preferably, R₅ is benzyl;


R₆ is selected from hydrogen, methyl or (R₇)(R₈)NC(=O)CH₂;

more preferably, R₆ is hydrogen;


R₇ and R₈ are independently selected from hydrogen, methyl or ethyl; and, more preferably, R₇ and R₈ are independently ethyl;

and pharmaceutically acceptable salts thereof.


[0015] Exemplifying the invention is a compound of Formula (I) selected from:




1




2




3




4




5




6




7




8




9




10




11




12




13




14




15




16




17




18




19




20




21




22




23




24




25




26




27




28




29




30




31




32




33




34




35




36




37




38




39




40




41







42




43




44


**45****46****47****48**




49




50




51




52




53




54




55




56




57





58



59



60

**61, or****62**

and pharmaceutically acceptable salts thereof.

[0016] Particularly preferred compounds of Formula (I) are selected from:

Cpd	Name
1	β -[[[1-[4-[(4,5-dihydro-1 <i>H</i> -imidazol-2-yl)amino]-1-oxobutyl]-4 piperidinyl]carbonyl]amino]-, (β^3S)- 3-quinolinepropanoic acid;
2	3-[[[1-[4-[(4,5-dihydro-1 <i>H</i> -imidazol-2-yl)amino]-1-oxobutyl]-4 piperidinyl]carbonyl]amino]- <i>N</i> -[(phenylmethoxy)carbonyl]-L-alanine;
6	β -[[[1-[1-oxo-4-(2-pyridinylamino)butyl]-4-piperidinyl]carbonyl]amino]-, (β^3S)-3-quinolinepropanoic acid;
11	3-[[[1-[1-[(4,5-dihydro-1 <i>H</i> -imidazol-2-yl)-4-piperidinyl]carbonyl]-4-piperidinyl]carbonyl]amino]- <i>N</i> -[(phenylmethoxy)carbonyl]-L-alanine;
14	3-[[[1-[[(1 <i>S</i> ,4 <i>R</i>)-4-[(4,5-dihydro-1 <i>H</i> -imidazol-2-yl)amino]-2-cyclopenten-1-yl]carbonyl]-4-piperidinyl]carbonyl]amino]- <i>N</i> -[(phenylmethoxy)carbonyl]-L-alanine;
16	3-[[[1-[3-[(4,5-dihydro-1 <i>H</i> -imidazol-2-yl)amino]-2-methyl-1-oxopropyl]-4-piperidinyl]carbonyl]amino]- <i>N</i> -[(phenylmethoxy)carbonyl]-L-alanine;
17	3-[[[1-[[(1 <i>R</i> ,4 <i>S</i>)-4-[(4,5-dihydro-1 <i>H</i> -imidazol-2-yl)amino]-2-cyclopenten-1-yl]carbonyl]-4-piperidinyl]carbonyl]amino]- <i>N</i> -[(phenylmethoxy)carbonyl]-L-alanine;
28	4-[[[[(2 <i>S</i>)-2-carboxy-2-[(phenylmethoxy)carbonyl]amino]ethyl]amino]carbonyl]-, 2-[(4,5-dihydro-1 <i>H</i> -imidazol-2-yl)amino]ethyl ester 1-piperidinecarboxylic acid;
39	β -[[[1-[2-[(4,5-dihydro-1 <i>H</i> -imidazol-2-yl)amino]ethoxy]carbonyl]-4-piperidinyl]carbonyl]amino]-, (β^3S)- 3-quinolinepropanoic acid;
47	3-[[[1-[4-[(4,5-dihydro-1 <i>H</i> -imidazol-2-yl)amino]-2-methyl-1-oxobutyl]-4 piperidinyl]carbonyl]amino]- <i>N</i> -[(phenylmethoxy)carbonyl]-L-alanine;
50	3-[[[1-(4-amino-1-oxo-5-hexenyl)-4-piperidinyl]carbonyl]amino]- <i>N</i> -[(phenylmethoxy)carbonyl]-L-alanine;
53	3-[[[1-[4-[(4,5-dihydro-1 <i>H</i> -imidazol-2-yl)amino]methyl]benzoyl]-4-piperidinyl]carbonyl]amino]- <i>N</i> -[(phenylmethoxy)carbonyl]-L-alanine;
61	β -[[[1-[3-[(4,5-dihydro-1 <i>H</i> -imidazol-2-yl)amino]oxy]-1-oxopropyl]-4-piperidinyl]carbonyl]amino]- 1,3-benzodioxole-5-propanoic acid; or,
62	β -[[[1-[1-oxo-4-(2-pyridinylamino)butyl]-4-piperidinyl]carbonyl]amino]-1,3-benzodioxole-5-propanoic acid;

and pharmaceutically acceptable salts thereof.

[0017] Representative Chemical Abstracts Service (CAS) Index-like names for the compounds of the present invention were derived using the ACD/LABS SOFTWARE™ Index Name Pro Version 4.0 nomenclature software program provided by Advanced Chemistry Development, Inc., Toronto, Ontario, Canada.

[0018] The compounds of the present invention may also be present in the form of pharmaceutically acceptable salts. For use in medicine, the salts of the compounds of this invention refer to non-toxic "pharmaceutically acceptable salts." Other salts may, however, be useful in the preparation of compounds according to this invention or of their pharmaceutically acceptable salts. The pharmaceutically acceptable salts generally take a form in which a basic nitrogen or nitrogens on the compound of the Formula (I) are protonated with an inorganic or organic acid. Representative inorganic or organic acids include, but are not limited to, hydrochloric, hydrobromic, hydriodic, perchloric, sulfuric, nitric, phosphoric, acetic, propionic, glycolic, lactic, succinic, maleic, fumaric, malic, tartaric, citric, benzoic, mandelic, methanesulfonic, hydroxyethanesulfonic, benzenesulfonic, oxalic, pamoic, 2-naphthalenesulfonic, p-toluenesulfonic, cyclohexanesulfamic, salicylic, saccharinic or trifluoroacetic acid. Similarly, pharmaceutically acceptable salts also include compounds wherein a carboxylic acid group of the compound of the Formula (I) combines with a organic or inorganic base. Representative inorganic or organic bases include, but are not limited to lithium, sodium, potassium, magnesium, calcium, aluminum, zinc, benzathine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (*N*-methyl glucamine), tromethamine (2-amino-2-(hydroxymethyl)-1,3-propanediol) or procaine (4-amino-[2-(diethylamino)ethyl ester] benzoic acid).

[0019] As used herein, unless otherwise noted alkyl and alkoxy when used alone include straight and branched chains having 1 to 8 carbon atoms, or any number within this range. Similarly, alkenyl and alkynyl groups include straight and branched chain alkenes and alkynes having 2 to 8 carbon atoms, or any number within this range. The term alkylene, alkenylene and alkynylene refers to alkyl, alkenyl and alkynyl chains, respectively, that are further substituted and act as linking groups. Alkoxy radicals are oxygen ethers formed from the previously described straight or branched chain alkyl groups. Cycloalkyl groups contain 3 to 8 ring carbons; preferably, 5 to 8 ring carbons; and, more preferably, 5 to 7 ring carbons.

[0020] The term "heterocyclyl" as used herein refers to an optionally substituted, stable, saturated 3 to 10 (preferably 4 to 8) membered monocyclic or bicyclic ring system which consists of carbon atoms and from one to three heteroatoms selected from N, O or S. The term "heterocyclenyl" as used herein refers to an optionally substituted, stable, unsaturated 3 to 10 (preferably 4 to 8) membered monocyclic or bicyclic ring system which consists of carbon atoms and from one to three heteroatoms selected from N, O or S. The heterocyclyl or heterocyclenyl group may be attached at any heteroatom or carbon atom which results in the creation of a stable structure and, accordingly, may be further attached to, for example, alkyl or alkoxy chains. The term heterocyclylene and heterocyclenylene refers to heterocyclyl and heterocyclenyl groups, respectively, that are further substituted and act as linking groups; wherein, one or both rings may be optionally substituted with one to five substituents attached at any heteroatom or carbon atom which results in the creation of a stable structure.

[0021] The term "aryl", as used herein, refers to optionally substituted aromatic groups such as phenyl and naphthyl. The term arylene refers to aryl groups that are further substituted and act as linking groups; wherein, one or both rings may be optionally substituted with one to five substituents attached at any carbon atom which results in the creation of a stable structure.

[0022] The term "heteroaryl" as used herein represents a stable five or six membered monocyclic aromatic ring system or a nine or ten membered benzo-fused heteroaromatic ring system which consists of carbon atoms and from one to three heteroatoms selected from N, O or S. The heteroaryl group may be attached or optionally substituted at any heteroatom or carbon atom which results in the creation of a stable structure.

[0023] The term "arylalkyl" means an alkyl group substituted with an aryl group (e.g., benzyl, phenethyl). Similarly, the term "arylalkoxy" indicates an alkoxy group substituted with an aryl group (e.g., benzyloxy).


[0024] The term "acyl" as used herein means an organic radical having 2 to 6 carbon atoms (branched or straight chain) derived from an organic acid by removal of the hydroxy group.

[0025] Whenever the term "alkyl" or "aryl" or either of their prefix roots appear in a name of a substituent (e.g., aralkyl, alkylamino) it shall be interpreted as including those limitations given above for "alkyl" and "aryl." When present, designated numbers of carbon atoms (e.g., C₁-C₈) shall refer independently to the number of carbon atoms in a moiety or to the alkyl portion of a larger substituent in which alkyl appears as its prefix root.

[0026] Any of the foregoing cycloalkyl, heterocyclyl, heterocyclenyl, aryl and heteroaryl substituent groups and their corresponding heterocyclylene, heterocyclenylene, arylene and heteroarylene linking groups, as a monocyclic or bicyclic ring system may be further bridged at any heteroatom or carbon atom which results in the creation of a stable ring structure; wherein the monocyclic bridged ring system consists of 6 to 8 members and the bicyclic bridged ring system consists of 10 to 12 members.

[0027] It is intended that the definition of any substituent or variable at a particular location in a molecule be independent of its definitions elsewhere in that molecule. It is understood that substituents and substitution patterns on the compounds of this invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art as well as those methods set forth herein.

[0028] Under standard nomenclature used throughout this disclosure, the terminal portion of the designated side chain is described first, followed by the adjacent functionality toward the point of attachment. Thus, for example, a "phenylC₁-C₆ alkylamidoC₁-C₆alkyl" substituent refers to a group of the formula:



[0029] The present invention includes within its scope prodrugs of the compounds of this invention. In general, such prodrugs will be functional derivatives of the compounds which are readily convertible *in vivo* into the required compound. Examples of prodrugs include but are not limited to various esters, hydroxyamidines and hydroxyguanidines of compounds of the present invention. Thus, in the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various disorders described with the compound specifically disclosed or with a compound which may not be specifically disclosed, but which converts to the specified compound *in vivo* after administration to the patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

[0030] Where the compounds according to this invention have at least one chiral center, they may accordingly exist as enantiomers. Where the compounds possess two or more chiral centers, they may additionally exist as diastereomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention. Furthermore, some of the crystalline forms for the compounds may exist as polymorphs and as such are intended to be included in the present invention. In addition, some of the compounds may form solvates with water (i.e., hydrates) or common organic solvents, and such solvates are also intended to be encompassed within the scope of this invention.

[0031] The term "subject" as used herein, refers to an animal, preferably a mammal, most preferably a human, who has been the object of treatment, observation or experiment.

[0032] The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human, that is being sought by a researcher, veterinarian, medical doctor, or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

[0033] The term "bone resorption," as used herein, refers to the process by which osteoclasts degrade bone.

[0034] The present invention provides isonipecotamide compounds that are useful as $\alpha\nu\beta 3$, $\alpha\nu\beta 5$ and GPIIb/IIIa integrin antagonists. Compounds of Formula (I) that are particularly preferred as $\alpha\nu\beta 3$ integrin antagonists are selected from Compound 1, 2, 3, 5, 6, 12, 14, 16, 18, 21, 24, 25, 28, 29, 31, 33, 34, 39, 40, 44, 47, 49, 54, 55, 58 or 60. Compounds of Formula (I) that are particularly preferred as GPIIb/IIIa integrin antagonists are selected from Compound 1, 2, 5, 11, 12, 14, 16, 17, 18, 19, 39, 47, 49, 50, 53, 54, 55 or 56. Compounds of Formula (I) that are particularly preferred as $\alpha\nu\beta 5$ integrin antagonists are selected from Compound 1, 2, 16, 45, 58 or 60. Compounds of Formula (I) that are particularly preferred as dual $\alpha\nu\beta 3$ /GPIIb/IIIa integrin antagonists are selected from Compound 2, 5, 12, 14, 16, 18, 39, 47, 49, 54 or 55. Compounds of Formula (I) that are particularly preferred as dual $\alpha\nu\beta 3$ / $\alpha\nu\beta 5$ integrin antagonists are selected from Compound 1, 2, 16, 58 or 60.

[0035] Compounds of Formula (I) that are more particularly preferred as $\alpha\nu\beta 3$ integrin antagonists are selected from Compound 1, 2, 6, 28, 39 or 47. Compounds of Formula (I) that are more particularly preferred as GPIIb/IIIa integrin antagonists are selected from Compound 2, 11, 14, 17, 50 or 53. Compounds of Formula (I) that are more particularly preferred as $\alpha\nu\beta 5$ integrin antagonists are selected from Compound 1, 2 or 16. Compounds of Formula (I) that are more particularly preferred as dual $\alpha\nu\beta 3$ /GPIIb/IIIa integrin antagonists are Compound 2. Compounds of Formula (I) that are more particularly preferred as dual $\alpha\nu\beta 3$ / $\alpha\nu\beta 5$ integrin antagonists are selected from Compound 1 or 2.

[0036] The compounds of Formula (I) inhibit the binding of adhesive proteins such as fibrinogen, vitronectin, and osteopontin to the integrin class of receptors. As demonstrated by the results of the biological studies described hereinafter, the compounds block vitronectin binding to isolated $\alpha\nu\beta 3$ (IC_{50} 's of ca. 1-300 nM), and inhibit fibrinogen binding to isolated GPIIb/IIIa as well. Because the compounds of this invention inhibit integrin-mediated cell-cell or cell-matrix adhesion, they may be useful against restenosis, thrombosis, inflammation, atherosclerosis, arthritis, angiogenesis, osteoporosis, bone resorption, tumor cell metastasis, tumor growth, macular degeneration, diabetic retinopathy, diseases of lung/airway resistance, etc. (D. Cox, *Drug News & Perspectives* 1995, 8, 197).

[0037] The compounds of Formula (I) are also useful as antithrombotics in combination with fibrinolytic therapy (e.g., t-PA or streptokinase). Additionally, the compounds of Formula (I) are useful in combination with one or more agents useful in the prevention or treatment of osteoporosis and arthritis. For example, the compounds of the instant invention may be effectively administered in combination with other agents used in the treatment of osteoporosis such as bisphosphonate bone resorption inhibitors; preferably, the bisphosphonate bone resorption inhibitor is alendronate, sold as FOSAMAX®. Preferred combinations are simultaneous or alternating treatments of an integrin antagonist of the present invention and alendronate.

[0038] In accordance with the methods of the present invention, the individual components of the combination can be administered separately at different times during the course of therapy or concurrently in divided or single combination forms. The instant invention is therefore to be understood as embracing all such regimes of simultaneous or alternating treatment and the term "administering" is to be interpreted accordingly.

[0039] As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

[0040] The utility of the compounds to treat integrin-mediated disorders can be determined according to the procedures herein. The present invention therefore provides a method of treating integrin-mediated disorders in a subject in need thereof which comprises administering any of the compounds as defined herein in a quantity effective to treat thrombotic disorders, osteoporosis, cancer, and diabetic complications. The compound may be administered to a patient by any conventional route of administration, including, but not limited to, intravenous, oral, subcutaneous, intramuscular, intradermal and parenteral.

[0041] The present invention also provides pharmaceutical compositions comprising one or more compounds of this invention in association with a pharmaceutically acceptable carrier.

[0042] To prepare the pharmaceutical compositions of this invention, one or more compounds of Formula (I) or salt thereof of the invention as the active ingredient, is intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending of the form of preparation desired for administration, e.g., oral or parenteral such as intramuscular. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed. Thus, for liquid oral preparations, such as for example, suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like; for solid oral preparations such as, for example, powders, capsules, caplets, gelcaps and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated or enteric coated by standard techniques. For parenterals, the carrier will usually comprise sterile water, through other ingredients, for example, for purposes such as aiding solubility or for preservation, may be included. Injectable suspensions may also be prepared, in which case appropriate liquid carriers, suspending agents and the like may be employed. The pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder, injection, teaspoonful and the like, an amount of the active ingredient necessary to deliver an effective dose as described above. The pharmaceutical compositions herein will contain, per unit dosage unit, e.g., tablet, capsule, powder, injection, suppository, teaspoonful and the like, of from about 0.01 mg/kg to 100 mg/kg (preferably, from about 0.1 mg/kg to about 30 mg/kg) and may be given at a dosage of from about 0.01 mg/kg/day to about 300 mg/kg/day (preferably, from about 1 mg/kg/day to about 50 mg/kg/day). The dosages, however, may be varied depending upon the requirement of the patients, the severity of the condition being treated and the compound being employed. The use of either daily administration or post-periodic dosing may be employed.

[0043] Preferably these compositions are in unit dosage forms from such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, autoinjector devices or suppositories; for oral parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. Alternatively, the composition may be presented in a form suitable for once-weekly or once-monthly administration; for example, an insoluble salt of the active compound, such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of material can be used for such enteric layers or coatings, such materials including a number of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

[0044] The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include, aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured

emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions, include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinylpyrrolidone or gelatin.

[0045] Where the processes for the preparation of the compounds according to the invention give rise to mixture of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The compounds may, for example, be resolved into their components enantiomers by standard techniques, such as the formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-d-tartaric acid and/or (+)-di-p-toluoyl-l-tartaric acid followed by fractional crystallization and regeneration of the free base. The compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary. Alternatively, the compounds may be resolved using a chiral HPLC column.

[0046] During any of the processes for preparation of the compounds of the present invention, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis; Third Edition, John Wiley & Sons, 1999. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

[0047] The method of treating integrin-mediated disorders described in the present invention may also be carried out using a pharmaceutical composition comprising any of the compounds as defined herein and a pharmaceutically acceptable carrier. The pharmaceutical composition may contain between about 0.01 mg and 100 mg, preferably about 5 to 50 mg, of the compound, and may be constituted into any form suitable for the mode of administration selected. Carriers include necessary and inert pharmaceutical excipients, including, but not limited to, binders, suspending agents, lubricants, flavorings, sweeteners, preservatives, dyes, and coatings. Compositions suitable for oral administration include solid forms, such as pills, tablets, caplets, capsules (each including immediate release, timed release and sustained release formulations), granules, and powders, and liquid forms, such as solutions, syrups, elixirs, emulsions, and suspensions. Forms useful for parenteral administration include sterile solutions, emulsions and suspensions.

[0048] Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

[0049] For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders; lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

[0050] The liquid forms in suitably flavored suspending or dispersing agents such as the synthetic and natural gums, for example, tragacanth, acacia, methylcellulose and the like. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable preservatives are employed when intravenous administration is desired.

[0051] The compound of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

[0052] Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targeted drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamidephenol, polyhydroxy-ethylaspartamidephenol, or polyethyl enoxidepolylysine substituted with palmitoyl residue. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

[0053] Compounds of this invention may be administered in any of the foregoing compositions and according to dosage regimens established in the art whenever treatment of thrombotic disorders is required.

[0054] The daily dosage of the products may be varied over a wide range from about 0.01 mg to about 21,000 mg

per adult human per day. For oral administration, the compositions are preferably provided in the form of tablets containing, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 150, 200, 250 and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. A therapeutically effective amount of the drug is ordinarily supplied at a dosage level of from about 0.01 mg/kg to about 300 mg/kg of body weight per day. Preferably, the range is from about 0.01 mg/kg to about 100 mg/kg of body weight per day; and more preferably, the range is from about 0.01 mg/kg to about 50 mg/kg of body weight per day. The compounds may be administered on a regimen of 1 to 4 times per day.

[0055] Optimal dosages to be administered may be readily determined by those skilled in the art, and will vary with the particular compound used, the mode of administration, the strength of the preparation, the mode of administration, and the advancement of the disease condition. In addition, factors associated with the particular patient being treated, including patient age, weight, diet and time of administration, will result in the need to adjust dosages.

[0056] Abbreviations used in the instant specification, particularly the Schemes and Examples, are as follows:

	Aq. =	aqueous
15	Bn or Bzl =	benzyl
	Boc =	<i>tert</i> -butoxycarbonyl
	Boc ₂ O	di- <i>tert</i> -butyl dicarbonate
	BSA =	bovine serum albumin
	<i>n</i> -Bu =	<i>n</i> -butyl
20	Cbz =	benzyloxycarbonyl
	CP or Cpd =	Compound
	DCM =	dichloromethane
	DIC =	diisopropylcarbodiimide
	DIPEA =	diisopropylethylamine
25	DMAP =	4-dimethylaminopyridine
	DMF =	N, N-dimethylformamide
	DMSO =	dimethylsulfoxide
	EDC =	ethyl dimethylaminopropyl-carbodiimide
	EDTA =	ethylenediaminetetraacetic acid
30	ES =	electrospray
	Et =	ethyl
	Et ₂ O =	diethyl ether
	EtOH =	ethanol
	¹ H NMR =	proton nuclear magnetic resonance spectrum
35	HBTU =	2-(1 <i>H</i> -benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
	HEPES =	4-(2-hydroxyethyl)-1- piperazine-ethanesulfonic acid
	HOBT =	hydroxybenzotriazole
	HBTU =	2-(1 <i>H</i> -benzotriazole-1-yl)-1,1,3,3-tetramethyl-uronium hexafluorophosphate
	IPA =	2-propanol
40	Me =	methyl
	MeOH =	methanol
	MH ⁺ =	observed molecular ion
	MPK =	milligrams per kilogram
	MS =	mass spectrum
45	NMM =	N-methylmorpholine
	NT =	not tested
	Ph =	phenyl
	PPT =	precipitate
	Pyr. =	pyridine
50	RP-HPLC =	preparative reversed-phase high pressure liquid chromatography
	RT =	room temperature
	Sch =	Scheme
	TEA =	triethylamine
	THF =	tetrahydrofuran
55	TFA =	trifluoroacetic acid
	TMS =	trimethylsilane
	Z =	benzyloxycarbonyl

General Synthetic Methods

[0057] Although the following Schemes depict synthetic routes to specific compounds of Formula (I), they are intended to also represent general synthetic routes to other compounds of the present invention to those skilled in the art of synthetic organic chemistry. By way of example, the synthetic route shown in Scheme AA used to prepare Compound **1** is also illustrative of a general method for the preparation of other instant compounds whereby other starting materials and appropriate reagents and solvents are used. Most of the reaction unless otherwise indicated were performed at room temperature.

SCHEME AA

[0058] Compounds of the invention wherein **A** is (4,5-dihydro-1*H*-imidazol-2-yl)amino and its homologs may be prepared as shown below. The β -Aminoacid derivative **AA5** was prepared as detailed in PCT International Application WO 97/41102 and as published (J. Rico, J. *Org. Chem.* **1993**, *58*, 7948). Such β -aminoacid derivatives can also be prepared by the methods described in WO 97/08145 and in U.S. Patent 6,100,423. The reagent **AA3** was purchased from Maybridge Chemical Company.


[0059] For the preparation of compounds exemplified by final target Compound **1**, secondary amine **AA1** was coupled with 4-*N*-Boc-aminobutyric acid in the presence, and the Boc group subsequently removed with trifluoroacetic acid to give primary amine **AA2** as the TFA salt. This salt was acylated with reagent **AA3**, the 2-aminoimidazoline product was protected with di-*tert*-butyl dicarbonate, and the methyl ester was saponified with lithium hydroxide to afford **AA4**. Intermediate **AA4** was coupled with amino ester **AA5** in the presence of HBTU, and the product was then de-protected with aqueous hydrochloric acid to give the target Compound **1**. Similarly, the Boc group of coupling product from **AA4** and **AA5** can be removed with a mixture of trifluoroacetic acid in dichloromethane (1:1) and the ester moiety subsequently saponified with aqueous LiOH in tetrahydrofuran to afford the lithium salt of Compound **1**. If the ester derivative is desired, then the saponification step is eliminated.

[0060] For the preparation of α -diaminoalkanoic acid derivative compounds of Formula (I), such as those exemplified by the Compound **11**, the secondary amine **AA1** was reacted with (N-*t*-butoxycarbonyl)isonipecotic acid (purchased from BaChem) instead of 4-*N*-Boc-aminobutyric acid and the product was carried forward to replace **AA2** in Scheme AA, wherein **AB2** was used to replace **AA5** to produce the target Compound **11**.

[0061] Similarly, for the preparation of compounds of Formula (I), such as those exemplified by Compound **16**, the secondary amine **AA1** was reacted with commercially available *N*-Boc-(3-amino-2-methyl)propionic acid instead of 4-*N*-Boc-aminobutyric acid and carried forward to replace **AA2** in Scheme AA, wherein **AB2** was used to replace **AA5** to produce the target Compound **16**.


[0062] For the preparation of compounds of Formula (I) exemplified by Compound **53**, the secondary amine **AA1** was reacted with the commercially available *N*-Boc-(4-aminomethyl)benzoic acid instead of 4-*N*-Boc-aminobutyric acid and carried forward to replace **AA2** in Scheme AA to produce the target Compound **53**.

[0063] For the preparation of compounds of Formula (I), such as those exemplified by Compound **14** and Compound **17**, the secondary amine **AA1** was reacted with commercially available (-)-(1*R*,4*S*)-*N*-Boc-1-aminocyclopent-2-ene-4-carboxylic acid (for Compound **14**) or (+)-(1*S*,4*R*)-*N*-Boc-1-aminocyclopent-2-ene-4-carboxylic acid (for Compound **17**) instead of 4-*N*-Boc-aminobutyric acid. The resulting product replaced **AA2** in Scheme AA and was carried forward, with **AB2** replacing **AA5**, to produce Compound **14** and Compound **17**, respectively.



SCHEME AB


[0064] The α -diaminoalkanoic acid derivatives of Formula (I), such as Compound **2**, can be prepared from ester intermediates, such as the free base of **AB2** (methyl *N*- α -(benzyloxycarbonyl)-L-2,3-diaminopropionate), to replace **AA5** in Scheme AA. In this example, Cbz-protected acid **AB1** (purchased from BaChem) was dissolved in methanol and treated with 4N HCl in 1,4-dioxane in the presence of 1,2-dimethoxypropane to afford the intermediate amino ester **AB2**. Using the method described in Scheme AA, the free base of **AB2** was used to replace **AA5** to produce Compound **2**.




SCHEME AC

[0065] Compounds of Formula (I) wherein A is heteroaryl, heterocyclyl, (heteroaryl)amino or (heterocyclyl)amino, such as Compound **6**, can be prepared by reacting the heteroaryl, heterocyclyl, (heteroaryl)amino or (heterocyclyl)amino starting material with the appropriate alkyl halide. For example, Compound **AC1** was acylated with 4-bromobutyl chloride. The resulting alkyl bromide product was reacted with (pyridin-2-yl)amino at reflux to provide **AC2**. The ester **AC2** was saponified with lithium hydroxide and carried forward to replace **AA4** in Scheme AA to give Compound **6**. Other compounds of Formula (I), such as Compounds **30**, **41** and **62**, can be prepared using Scheme AC by varying

the starting material used.




[0066] Another method used to prepare heterocyclyl compounds, such as benzimidazol-2-yl, was the N-acylation of starting materials using methodology analogous to that shown in Scheme AA. For example, the N-acylation of AC3 with AD2 followed by saponification and deprotection by removal of the Boc group afforded Compounds 8, 13 and 15.



[0067] Certain other (heterocyclyl)amino compounds may be prepared by reacting a 2-fluoroheterocyclyl compound with the appropriate aminoalkanoate (e.g., AA2) as reported by C. Senanayake in *Tetrahedron Lett.* **1999**, 40, 6875. For compounds where A is (tetrahydro-pyrimidin-2-yl)amino or (pyrimidin-2-yl)amino, 2-bromopyrimidine may be reacted with the appropriate aminoalkanoate (e.g., AA2) as reported by G. Hartman in WO 95/32710. For compounds where A is 1H-imidazo[4,5b]pyridin-2-yl, an appropriate alkyl dicarboxylic acid (e.g., succinic acid) may be reacted with 2,3-diaminopyridine as reported by R. Keenan in *Bioorg. Med. Chem. Lett.* **1998**, 8, 3171.


SCHEME AD

[0068] Scheme AD provides an alternate method of synthesis for the α -diaminoalkanoic acid derivatives of Formula (I), such as Compound 2, by changing the order of the coupling steps depicted in Scheme AA. For example, the isonipecotamide intermediate AD2 used to further synthesize other α -diaminoalkanoic acid derivative compounds of the invention was prepared by the HBTU-mediated coupling of AD1 (purchased from Advanced Chemtech) and AB2.



[0069] The acid AD3 was prepared (as described by J. V. Greenhill, *Chem. Soc., Perkin Trans. 2* **1985**, (8), 1255-1264)

and was reacted with di-*tert*-butyl dicarbonate in the presence of NaOH. The intermediate sodium salt was coupled with **AD2** using HBTU to afford **AD4**. Deprotection of **AD4** with 4 M HCl (aq.) yielded Compound **2**.




SCHEME AE

25 **[0070]** Compounds of Formula (I) wherein A includes an aminoxy moiety can be prepared by the methods represented below. Boc-protected aminoxy intermediate **AE5** (used to prepare Compound **58**) and aminoxy intermediate **AE8** (used to prepare Compounds **60** and **61**) were synthesized according to the methods described by C. Gilon, et. al. in *Tetrahedron* **1967**, 23, 441-4447 and B. J. Ludwig in *J. Med. Chem.* **1970**, 13, 60-63. Commercially available β -propiolactone **AE1** and N-hydroxysuccinimide **AE2** were reacted to afford the corresponding oxyamino ester, which was hydrolyzed in a refluxing mixture of 6 M HCl (aq.) and glacial acetic acid. The resulting HCl salt was converted to the free base **AE3** by treatment with NaOMe in 2-propanol. Guanylation of **AE3** with commercially available **AE4** followed by reaction with di-*tert*-butyl dicarbonate furnished the protected oxyguanidine intermediate **AE5**. To produce the target Compound **53**, the secondary amine **AA1** in Scheme AA was reacted with the protected oxyguanidine intermediate **AE5** instead of 4-*N*-Boc-aminobutyric acid and carried forward as **AA2**.


30

35



55 **[0071]** Similarly, Boc-protection of **AE6** (purchased from Aldrich) produced **AE7**, followed by reaction of **AE7** with **AE3** to afford the protected (4,5-dihydro-1*H*-imidazol-2-yl)aminoxy intermediate **AE8**. To produce the target Compounds **60** and **61**, the secondary amine **AA1** in Scheme AA was reacted with the protected (4,5-dihydro-1*H*-imidazol-2-yl) aminoxy intermediate **AE8** instead of 4-*N*-Boc-aminobutyric acid and carried forward as **AA2**.


5



10 **[0074]** Similarly, thiourea **AG2**, prepared as described in WO/0034255, is activated with methyl iodide followed by reaction with **AG3** as described in WO/9708145 to provide **AG4**.

15

20



25

SCHEME AH


30

35

[0075] Compounds of Formula (I) wherein L is -OC(=O)- or -HNC(=O)- can be prepared as outlined below. The protected aminoalcohol **AH1** can be converted to a corresponding *p*-nitrophenylcarbonate intermediate by reaction with **AH2**. Reaction of the *p*-nitrophenylcarbonate intermediate product with **AF1**, followed by removal of the Boc group furnished an amine intermediate. The amine intermediate was subsequently reacted with reagent **AA3** to give a corresponding (imidazolin-2-yl)amino intermediate. Boc protection of the (imidazolin-2-yl)amino intermediate followed by saponification yielded **AH3**, which was used to replace intermediate **AA4** in Scheme AA to produce Compounds **28, 29, 31, 35, 39** and **40**.

40

45



50

[0076] Similarly, **AD2** was converted to **AH4** by reaction with **AH2** to produce **AH4**. Reaction of **AH4** with **AH5** and saponification provided Compound **10**.

55



SCHEME AI

25

30



Specific Synthetic Methods

[0078] The following Examples are set forth to aid in the understanding of the invention, and are not intended and should not be construed to limit in any way the invention set forth in the claims which follow thereafter.

[0079] Protected amino acids were purchased from Bachem Bioscience Inc. *N*- α -Cbz-L-2,3-diaminopropionic acid was purchased from Fluka. Aromatic aldehydes were purchased from Aldrich Chemical Company, as were all other chemicals. High field ^1H NMR spectra were recorded on a Bruker AC-360 spectrometer at 360 MHz, and coupling constants are given in Hertz. Melting points were determined on a Mel-Temp II melting point apparatus and are uncorrected. Microanalyses were performed at Robertson Microlit Laboratories, Inc., Madison, New Jersey and are expressed in percentage by weight of each element per total molecular weight. In those cases where the product is obtained as a salt, the free base is obtained by methods known to those skilled in the art, *e.g.* by basic ion exchange purification. Nuclear magnetic resonance (NMR) spectra for hydrogen atoms were measured in the indicated solvent with tetramethylsilane (TMS) as the internal standard on a Bruker AM-360 (360 MHz) spectrometer. The values are expressed in parts per million down field from TMS. The mass spectra (MS) were determined on a Micromass Platform LC single quadrupole mass analyser fitted with an electrospray ionisation source coupled to a Hewlett Packard 1050 HPLC and a Jones Chromatography diverter valve in the loop injection mode. Positive mode electrospray was used eluting with acetonitrile/water/acetic acid (50:50:1) flowing at 0.3 mL/min with a sprayer temperature of 120°C and an inlet desolvation cone set to 40 kV.

[0080] Unless otherwise noted, the materials used in the examples were obtained from readily available commercial suppliers or synthesized by standard methods known to anyone skilled in the art of chemical synthesis. The substituent groups, which vary between examples, are hydrogen unless otherwise noted.

EXAMPLE 1

β -[[[1-[4-[(4,5-dihydro-1*H*-imidazol-2-yl)amino]-1-oxobutyl]-4-piperidiny]carbonyl]amino]-, (β^3S)-3-quinolinepropanoic acid (Cpd 1)

[0081] A mixture of **AA1** (2.0 g, HCl salt, 0.011 mol), CH_2Cl_2 (100 mL), 4-*N*-Boc-aminobutyric acid (2.3 g, 1 eq), NMM (2.4 mL, 2 eq), HOBT (10 mg), and EDC•HCl (3.2 g, 1.5 eq) was stirred at 5 °C for 1 h, the ice bath was removed, and stirred for 2 h. The reaction was diluted with saturated aqueous NH_4Cl (50 mL) and the layers were separated. The organic layer was washed with NaOH (0.5 *N*, 15 mL), dried (Na_2SO_4), and evaporated to give a glass (3.6 g, 0.011 mol). This glass was dissolved in CH_2Cl_2 (15 mL) and anisole (1 mL), treated with TFA (15 mL), and stirred for 2 h. The solution was evaporated to give an oil (**AA2**, 3.8 g). **AA2** was dissolved in dioxane (40 mL), treated with NMM (2.4 mL, 2 eq) and 2-(3,5-dimethylpyrazolyl)-4,5-dihydroimidazole-HBr **AA3** (2.7 g, 1 eq) and heated at reflux for 2 h. This mixture was cooled to RT and filtered. The filtrate was treated with MeOH (80 mL), NMM (2.4 mL, 2 eq), di-*tert*-butyl dicarbonate (4.8 g, 2 eq), and sodium carbonate (2.4 g, 2 eq), and stirred for 3 days. The solvents were evaporated, and the residue was partitioned between CH_2Cl_2 (100 mL) and water (10 mL). The organic layer was dried (Na_2SO_4), evaporated, and the oil purified by silica gel chromatography (1% NH_4OH /10% MeOH/ CH_2Cl_2) to give a tan glass (1.8 g). The tan glass (0.09 g, 0.0023 mol) was dissolved in THF (5 mL), cooled to 5°C, treated with aqueous LiOH (0.11 g in 15 mL water), and stirred for 2 h. The solvents were evaporated to give **AA4** as a pale yellow solid (1.0 g). This solid was slurried with MeCN (100 mL), **AA5** (0.82 g, 1 eq, prepared starting from quinoline-3-carboxaldehyde as reported in M. Costanzo, *et al.*, WO 97/41102), HOBT (0.19 g, 0.5 eq), NMM (0.87 mL, 3 eq), and HBTU (1.2 g, 1.2 eq) at 5°C, stirred for 6 h, diluted with saturated aqueous NH_4Cl (12 mL), and the MeCN evaporated. The residue was partitioned between CHCl_3 (100 mL) and water (15 mL) and filtered. The filtrate layers were separated, and the organic layer was dried (Na_2SO_4), evaporated, and purified by silica gel chromatography (1% NH_4OH /4% MeOH/7% EtOH/ CH_2Cl_2) to give a clear glass (0.60 g). The glass was treated with water (12 mL) and HCl (conc., 10 mL), and stirred for 18 h. The aqueous HCl was evaporated, the resultant oil treated with MeCN (20 mL), and the precipitate collected and dried to afford Compound 1 as a white foam: ^1H NMR ($\text{DMSO}-d_6$) δ 1.3 (m, 2 H), 1.6 (m, 4 H), 2.3 (m, 2 H), 2.6 (m, 1 H), 2.8 (m, 1 H), 3.1 (m, 6 H), 3.4 (m, 1 H), 3.7 (m, 3 H), 3.8 (m, 1 H), 4.0 (m, 2 H), 4.2 (m, 1 H), 5.3 (m, 1 H), 7.7 (t, $J = 4$ Hz, 1 H), 7.9 (t, $J = 4$ Hz, 1 H), 8.2 (m, 2 H), 8.3 (m, 1 H), 8.6 (m, 2 H), 9.2 (s, 1 H), 10.9 (m, 1 H); MS m/e 481.4 (MH^+); $[\alpha]^{25}_{\text{D}} -42.1^\circ$ (c 0.114, MeOH).

EXAMPLE 2

3-[[[1-[4-[(4,5-dihydro-1*H*-imidazol-2-yl)amino]-1-oxobutyl]-4-piperidiny]carbonyl]amino]-*N*-[(phenylmethoxy)carbonyl]-L-alanine (Cpd 2)

[0082] Compound 2 was prepared as described for Compound 1 in Scheme AA using **AA4** (0.10 g) and methyl *N*- α -Cbz-L-2,3-diaminopropionate•HCl **AB2** to replace **AA5** (0.09 g), and isolated as a tan glass: ^1H NMR ($\text{DMSO}-d_6$) δ 1.3

(m, 1 H), 1.4 (m, 1 H), 1.7 (m, 4 H), 2.4 (m, 3 H), 2.6 (m, 2 H), 2.8 (t, J = 4 Hz, 1 H), 3.0 (m, 1 H), 3.1 (m, 1 H), 3.3 (m, 2 H), 3.4 (m, 1 H), 3.8 (m, 2 H), 4.1 (t, J = 4 Hz, 1 H), 4.3 (m, 1 H), 5.0 (t, J = 7 Hz, 2 H), 7.3 (m, 5 H), 7.6 (m, 3 H), 8.0 (m, 3 H), 8.2 (m, 1 H); MS m/e 503.4 (MH⁺).

EXAMPLE 3

6-chloro-β-[[[1-[4-[(4,5-dihydro-1*H*-imidazol-2-yl)amino]-1-oxobutyl]-4-piperidiny]carbonyl]amino]-, (β³S)-3-pyridine-propanoic acid (Cpd 3)

[0083] Compound **3** was prepared as described for Compound **1** in Scheme AA using **AA4** (0.15 g) and methyl (3*S*)-amino-3-(6-chloro-3-pyridyl)propionate•2HCl (0.11 g; prepared starting from 6-chloropyridine-3-carboxaldehyde as reported by M. Costanzo, et al. in WO 97/41102) for **AA5**, and isolated as a clear glass: ¹H NMR (DMSO-d₆) δ 1.2 (m, 4 H), 1.5 (m, 2 H), 2.2 (m, 3 H), 2.5 (m, 2 H), 2.7 (m, 2 H), 2.8 (m, 1 H), 3.1 (m, 1 H), 3.5 (m, 3 H), 3.8 (m, 1 H), 4.1 (m, 3 H), 4.3 (m, 1 H), 5.1 (m, 1 H), 7.4 (d, J = 5 Hz, 1 H), 7.7 (d, J = 5 Hz, 1 H), 8.1 (m, 3 H), 8.3 (s, 1 H), 8.5 (m, 1 H); MS m/e 464.9 (MH⁺).

EXAMPLE 4

3-[[[1-[4-[(4,5-dihydro-1*H*-imidazol-2-yl)amino]-1-oxobutyl]-4-piperidiny]carbonyl]amino]-, (3*R*)-butanoic acid (Cpd 4)

[0084] Compound **4** was prepared as described for Compound **1** in Scheme AA using **AA4** (0.90 g) and *tert*-butyl (3*R*)-aminobutyrate (0.37 g, purchased from Oxford Asymmetry) for **AA5**, and isolated as a clear glass: ¹H NMR (DMSO-d₆) δ 1.0(d,J=5Hz,3H), 1.3(m,2H), 1.7 (m, 4 H), 2.3 (m, 6 H), 2.6 (m, 1 H), 3.0 (m, 1 H), 3.2 (m, 2 H), 3.6 (s, 4 H), 3.8 (m, 1 H), 4.0 (m, 1 H), 4.4 (m, 1 H), 7.8 (d, J = 7 Hz, 2 H), 8.2 (m, 2 H); MS m/e 368.4 (MH⁺).

EXAMPLE 5

β-[[[1-[3-[(4,5-dihydro-1*H*-imidazol-2-yl)amino]-2-methyl-1-oxopropyl]-4-piperidiny]carbonyl]amino]-, (β³S)-3-quinoline-propanoic acid (Cpd 5)

[0085] Compound **5** was prepared as described for Compound **1** in Scheme AA using *N*-Boc-imidazolin-2-yl-(3-aminoisobutyl)-isonipecotic acid (0.10 g) for **AA4** and **AA5** (0.12 g), and isolated as glass: ¹H NMR (DMSO-d₆) δ 1.1 (d, J = 6 Hz, 3 H), 1.3 (m, 1 H), 1.4 (m, 1 H), 1.7 (m, 4 H), 2.3 (m, 2 H), 2.6 (m, 2 H), 2.8 (m, 1 H), 3.0 (m, 2 H), 3.1 (m, 1 H), 3.3 (m, 2 H), 3.4 (m, 1 H), 3.8 (m, 1 H), 4.1 (m, 1H), 4.3 (m, 1 H), 5.0 (t, J = 7 Hz, 2 H), 7.3 (m, 5 H), 7.6 (m, 2 H), 8.0 (m, 3 H); MS m/e 503.4 (MH⁺).

EXAMPLE 6

β-[[[1-[1-oxo-4-(2-pyridinylamino)butyl]-4-piperidiny]carbonyl]amino]-, (β³S)-3-quinolinepropanoic acid (Cpd 6)

[0086] A mixture of **AC1** (2.0 g, 0.011 mol), CH₂Cl₂ (70 mL), and NMM (2.4 mL, 2 eq) at 5 °C was treated with 4-bromobutryl chloride (1.3 mL, 1 eq), stirred for 6 h, and diluted with saturated aqueous NH₄Cl (15 mL). The organic layer was dried (Na₂SO₄), evaporated, and the resultant oil purified by silica gel chromatography (4% MeOH/ CH₂Cl₂) to give an oil (1.8 g). The oil was dissolved in MeOH (10 mL) and isopropanol (15 mL) and then treated with 2-aminopyridine (0.70 g, 1.2 eq) and sodium carbonate (2 mg). This mixture was heated at reflux in a sealed flask for 2 h, cooled to RT, and evaporated to an oil. The oil was purified by silica gel chromatography (2% MeOH/3% EtOH/CH₂Cl₂) to give **AC2** as a glass (0.3 g) intermediate. The intermediate **AC2** was then saponified with lithium hydroxide and carried forward to replace **AA4** as exemplified in Scheme AA to produce Compound **6** as clear flakes: ¹H NMR (DMSO-d₆) δ 1.0-1.2 (m, 2 H), 1.7 (m, 4 H), 2.2 (m, 2 H), 3.0 (m, 3 H), 3.3 (m, 3 H), 3.5 (t, J = 4 Hz, 1 H), 3.9 (m, 2 H), 5.5 (t, J = 4 Hz, 1 H), 6.9 (m, 1 H), 7.1 (m, 1 H), 7.9 (m, 4 H), 8.0 (t, J = 4 Hz, 1), 8.2 (m, 2 H), 8.8 (m, 3 H), 9.2 (s, 1 H), 10.9 (m, 1 H), 13.5 (m, 1 H); mp 87-90°C; MS m/e 490 (MH⁺).

EXAMPLES 7-62

[0087] Using the procedures described in Examples 1-6 or outlined in Schemes AA-AH and the appropriate reagents and starting materials known to those skilled in the art, other compounds of the present invention may be prepared including, but not limited to:

Cpd	Sch	Salt	MH ⁺	Cpd	Sch	Salt	MH ⁺	Cpd	Sch	Salt	MH ⁺
7	AC	TFA	500	27	AA	HCl	498	47	AA	TFA	517
8	AC	TFA	522	28	AH	HCl	505	48	AA	TFA	458
9	AA	Li	439	29	AH	HCl	476	49	AA	TFA	529
10	AH	Li	501	30	AC	HCl	440	50	AA	TFA	461
11	AA	TFA	529	31	AH	HCl	433	51	AA	HCl	517
12	AA	TFA	489	32	AA	HCl	448	52	AA	HCl	431
13	AC	Li	507	33	AA	HCl	431	53	AA	TFA	551
14	AA	TFA	527	34	AA	HCl	490	54	AA	TFA	517
15	AC	TFA	499	35	AH	HCl	468	55	AA	HCl	501
16	AA	TFA	503	36	AA	HCl	480	56	AA	HCl	435
17	AA	TFA	527	37	AA	HCl	480	57	AG	HCl	520
18	AA	TFA	614	38	AA	HCl	466	58	AE	HCl	479
19	AA	TFA	534	39	AH	HCl	483	59	AF	HCl	568
20	AA	HCl	436	40	AH	HCl	433	60	AE	HCl	505
21	AA	HCl	474	41	AC	HCl	526	61	AE	HCl	476
22	AA	HCl	444	42	AA	HCl	517	62	AC	HCl	483
23	AA	HCl	472	43	AA	HCl	466				
24	AA	HCl	490	44	AA	HCl	495				
25	AA	HCl	431	45	AA	---	694				
	26	AA	HCl	499	46	AA	HCl	594			

EXAMPLE 63

[0088] As a specific embodiment of an oral composition, 100 mg of the Compound **1** of Example 1 is formulated with sufficient finely divided lactose to provide a total amount of 580 to 590 mg to fill a size O hard gel capsule.

EXAMPLE 64*IN VITRO SOLID PHASE PURIFIED $\alpha\text{v}\beta 3$ BINDING ASSAY*

[0089] The vitronectin/ $\alpha\text{v}\beta 3$ binding assay methods were derived from Mehta et al. (*Biochem J.* **1998**, 330, 861). Human $\alpha\text{v}\beta 3$ (Chemicon International Inc., Temecula, CA), at a concentration of 1 $\mu\text{g}/\text{ml}$ dissolved in Tris buffer (20 mM Tris, 1 mM CaCl_2 , 1 mM MgCl_2 , 10 μM MnCl_2 , 150 mM NaCl), was immobilized on Immulon 96 well plates (Dynex Technologies, Chantilly, VA) overnight at 4 °C. Plates were washed and treated with blocking buffer (3 % **BSA** in Tris buffer) for 2 h at 37 °C. Plates were then rinsed 2 times with Tris buffer containing 0.3% BSA and 0.2% Tween20 (polyoxyethylenesorbitan monolaurate). Five minutes prior to the addition of 5 nM vitronectin (Sigma, St. Louis, MO), synthesized compounds were added to wells in duplicate. Each plate included c-RGDfV as an internal control. Following a 3 hour incubation at 37°C, plates were washed 5 times in assay buffer. An anti-human vitronectin IgG rabbit polyclonal antibody (Calbiochem, San Diego, CA) was added (1:2000) and plates were incubated for 1 hour at room temperature. VectaStain ABC peroxidase kit reagents (Vector Laboratories, Burlingame, CA) employing a biotin labeled anti-rabbit IgG, were utilized for detection of bound antibody. Plates were read at 490 nm on a Molecular Devices (Sunnyvale, CA) microplate reader. Table 1 shows the results of the in vitro solid phase purified $\alpha\text{v}\beta 3$ binding assay for representative compounds of the present invention.

EXAMPLE 65*IN VITRO SOLID PHASE PURIFIED GP IIB/IIIA BINDING ASSAY*

[0090] A 96 well Immulon-2 microtiter plate (Dynatech-Immulon) was coated with 50 $\mu\text{l}/\text{well}$ of RGD-affinity purified GPIIb/IIIA (effective range 0.5-10 $\mu\text{g}/\text{mL}$) in 10 mM HEPES, 150 mM NaCl, 1 mM MgCl_2 at pH 7.4. The plate was covered and incubated overnight at 4 °C. The GPIIb/IIIA solution was discarded and 150 μl of 5% BSA was added and incubated at RT for 1-3 h. The plate was washed extensively with modified Tyrodes buffer. Biotinylated fibrinogen (25 $\mu\text{l}/\text{well}$) at

2 x final concentration was added to the wells that contain the test compounds (25 μ l/well). The plate was covered and incubated at RT for 2-4 h. Twenty minutes prior to incubation completion, one drop of Reagent A (VectaStain ABC Horse Radish Peroxidase kit, Vector Laboratories, Inc.) and one drop Reagent B were added with mixing to 5 mL modified Tyrodes buffer mix and let stand. The ligand solution was discarded and the plate washed (5 x 200 μ l/well) with modified Tyrodes buffer. Vecta Stain HRP-Biotin-Avidin reagent (50 μ l/well, as prepared above) was added and incubated at RT for 15 min. The Vecta Stain solution was discarded and the wells washed (5 x 200 μ l/well) with modified Tyrodes buffer. Developing buffer (10 mL of 50 mM citrate/phosphate buffer @ pH 5.3, 6 mg o-phenylenediamine, 6 μ l 30% H₂O₂; 50 μ l/well) was added and incubated at RT for 3-5 min, and then 2 N H₂SO₄ (50 μ l/well) was added. The absorbance was read at 490 nM. Table 1 shows the results of the in vitro solid phase purified GPIIb/IIIa binding assay for representative compounds of the present invention.

EXAMPLE 66

IN VITRO SOLID PHASE PURIFIED $\alpha_v\beta_5$ BINDING ASSAY

[0091] The vitronectin/ $\alpha_v\beta_5$ binding assay method was performed in the same manner as the vitronectin/ $\alpha_v\beta_3$ binding assay (EXAMPLE 64) immobilizing with 1 μ g/ml of human purified $\alpha_v\beta_5$ (Chemicon International, Inc.) on to Immulon 96 well plates (Dynex Technologies) instead of $\alpha_v\beta_3$. All other aspects of the assay including buffers, reagents, and incubation times remain unchanged.

TABLE I

No.	$\alpha_v\beta_3$ IC ₅₀ (μ M)	IIb/IIIa IC ₅₀ (μ M)	$\alpha_v\beta_5$ IC ₅₀ (μ M)
1	0.0037	0.030	0.025
2	0.0058	0.0089	0.021
3	0.028	0.13	NT
4	0.15	1.70	NT
5	0.015	0.023	NT
6	0.0040	0.46	NT
7	0.49	7.9	NT
8	0.55	1.1	NT
9	30	0.30	NT
10	>50	0.95	NT
11	1.3	0.0026	NT
12	0.020	0.014	NT
13	10	29	NT
14	0.050	0.0092	NT
15	>5	17	NT
16	0.019	0.018	0.038
17	0.38	0.0054	NT
18	0.050	0.015	NT
19	0.32	0.071	NT
20	0.53	4.9	NT
21	0.023	0.41	NT
22	>0.5	38	NT
23	>0.5	7.3	NT
24	0.064	2.2	NT

EP 1 133 489 B9

Table continued


No.	$\alpha_v\beta_3$ IC ₅₀ (uM)	IIb/IIIa IC ₅₀ (uM)	$\alpha_v\beta_5$ IC ₅₀ (uM)
25	0.032	2.9	NT
26	0.12	1.8	NT
27	>0.5	13	NT
28	0.0082	1.5	NT
29	0.034	0.21	NT
30	0.14	56	NT
31	0.029	4.2	NT
32	0.12	3.6	NT
33	0.019	1.8	NT
34	0.029	1.5	>5.0
35	0.14	1.6	NT
36	>0.3	12	NT
37	0.29	1.7	NT
38	>0.3	3.3	NT
39	0.0023	0.036	NT
40	0.012	1.1	NT
41	>0.3	0.11	NT
42	0.17	0.14	NT
43	0.23	1.3	NT
44	0.028	0.36	2.9
45	3.7	14	40%*
46	0.15	1.6	>5
47	0.0040	0.028	NT
48	0.51	1.1	NT
49	0.039	0.012	0.21
50	3.9	0.0074	38%*
51	0.27	0.29	2
52	0.25	2.7	0.76
53	0.57	0.0015	NT
54	0.072	0.013	NT
55	0.012	0.026	0
56	>5	0.049	NT
57	0.35	0.13	NT
58	0.040	0.26	0.071
59	0.16	0.51	NT
60	0.027	0.19	0.086
61	0.18	2.7	NT

Table continued

No.	$\alpha_v\beta_3$ IC ₅₀ (uM)	IIb/IIIa IC ₅₀ (uM)	$\alpha_v\beta_5$ IC ₅₀ (uM)
62	>0.50	0.38	NT
*Percent inhibition at 5 μ M.			

Claims

1. A compound of the Formula (I):



wherein

M is selected from the group consisting of ethylene (optionally substituted within the carbon chain with methyl and substituted on the terminal carbon with one substituent selected from A), propylene (optionally substituted within the carbon chain with a substituent selected from the group consisting of methyl, ethenyl, cyclohexylidene [wherein a ring carbon atom forms the point of attachment to the carbon chain] and 4-Cl-phenyl and substituted on the terminal carbon with one substituent selected from A), allylene (substituted with one substituent selected from A); piperidin-4-yl, piperidin-4-ylene (substituted with one substituent selected from A), 1,4,5-dihydro-2-cyclopenten-1-ylene (substituted with one substituent selected from A) and 4-methylphenylene (substituted on methylene with one substituent selected from A);

A is optionally present and is selected from the group consisting of 1*H*-imidazol-1-yl, 1*H*-imidazol-2-yl, 4,5-dihydro-1*H*-imidazol-2-yl (optionally substituted with a substituent selected from the group consisting of C₁-C₄ alkoxycarbonyl and aryl(C₁-C₄)alkoxycarbonyl), pyridin-2-yl (optionally substituted with a substituent selected from the group consisting of C₁-C₄ alkyl, heteroaryl [optionally substituted with C₁-C₄ alkyl], halogen, hydroxy, nitro, cyano, amino, amino(C₁-C₄)alkyl, C₁-C₄ alkylamino(C₁-C₄)alkyl and di(C₁-C₄ alkyl)amino(C₁-C₄)alkyl), pyrimidin-2-yl, 1,4,5,6-tetrahydro-pyrimidin-2-yl (optionally substituted with one to two substituents independently selected from the group consisting of C₁-C₄ alkyl, hydroxy and amino), piperidin-4-yl, benzimidazol-2-yl, 1,2,3,4-tetrahydro-1,8-naphthyridin-7-yl, 3*H*-imidazo[4,5-*b*]pyridin-2-yl, amino, (C₁-C₆ alkyl)amino, (1*H*-imidazol-1-yl)amino, (1*H*-imidazol-2-yl)amino, (4,5-dihydro-1*H*-imidazol-2-yl)amino (optionally substituted on 4,5-dihydro-1*H*-imidazol-2-yl with a substituent selected from the group consisting of (C₁-C₆)alkoxycarbonyl and aryl(C₁-C₆)alkoxycarbonyl), (pyridin-2-yl)amino (optionally substituted on pyridin-2-yl with a substituent selected from the group consisting of C₁-C₄ alkyl, heteroaryl [optionally substituted with C₁-C₄ alkyl], halogen, hydroxy, nitro, cyano, amino, amino(C₁-C₄)alkyl, C₁-C₄ alkylamino(C₁-C₄)alkyl and di(C₁-C₄ alkyl)amino(C₁-C₄)alkyl), (pyrimidin-2-yl)amino, (1,4,5,6-tetrahydropyrimidin-2-yl)amino (optionally substituted on 1,4,5,6-tetrahydropyrimidin-2-yl with one to two substituents independently selected from the group consisting of C₁-C₄ alkyl, hydroxy and amino), (4,5,6,7-tetrahydro-1*H*-1,3-diazepin-2-yl)amino, (thiazol-2-yl)amino, (benzimidazol-2-yl)amino, (3*H*-imidazo[4,5-*b*]pyridin-2-yl)amino, R₃HNC(=NH)NH-, R₃HNC(=O)NH-, (4,5,6,7-tetrahydro-1*H*-1,3-diazepin-2-yl)aminooxy, (4,5-dihydro-1*H*-imidazol-2-yl)aminooxy and R₃HNC(=NH)NHO-;

with the proviso that if A is H₂NC(=NH)NH-, then, dependently, W is not hydrogen when Q is -CH₂-;

L is selected from the group consisting of -C(=O)-, -OC(=O)- and -HNC(=O)-;

R₁ is selected from the group consisting of hydrogen and C₁-C₄ alkyl;

R₂ is hydrogen;

R₃ is selected from the group consisting of hydrogen, C₁-C₈ alkyl, aryl(C₁-C₈)alkyl and hydroxy;

Q is selected from the group consisting of -CH₂-, -CH(C₁-C₈alkyl)-, -CH(aryl)- (wherein aryl is optionally substituted with one to five substituents independently selected from the group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, -O-(C₁-C₃ alkyl)-O-, halogen, hydroxy and trihalo(C₁-C₃)alkyl), -CH(heteroaryl)- (wherein heteroaryl is optionally substituted

with a substituent selected from halogen) and -CH(aryl(C₁-C₈)alkyl)-;

W is selected from the group consisting of hydrogen and N(R₄)T;

r is 1;

R₄ is selected from the group consisting of hydrogen and C₁-C₄ alkyl;

T is selected from the group consisting of R₅C(=O)- and R₅OC(=O)-;

R₅ is selected from the group consisting of aryl and aryl(C₁-C₄)alkyl;

R₆ is selected from the group consisting of hydrogen, methyl and (R₈)(R₇)NC(=O)CH₂; and,

R₇ and R₈ are independently selected from the group consisting of hydrogen, methyl and ethyl; and pharmaceutically acceptable salts thereof.

2. The compound of Claim 1, wherein

A is optionally present and is selected from the group consisting of 1H-imidazol-1-yl, 4,5-dihydro-1H-imidazol-2-yl, pyridin-2-yl (optionally substituted with a substituent selected from the group consisting of C₁-C₄ alkyl and heteroaryl [optionally substituted with C₁-C₄ alkyl]), piperidin-4-yl, benzimidazol-2-yl, 1,2,3,4-tetrahydro-1,8-naphthyridin-7-yl, 3H-imidazo[4,5-b]pyridin-2-yl, amino, (4,5-dihydro-1H-imidazol-2-yl)amino (optionally substituted on 4,5-dihydro-1H-imidazol-2-yl with C₁-C₆ alkoxycarbonyl), (pyridin-2-yl)amino (optionally substituted on pyridinyl with a substituent selected from the group consisting of C₁-C₄ alkyl and heteroaryl [optionally substituted with C₁-C₄ alkyl]), (1,4,5,6-tetrahydro-5-hydroxypyrimidin-2-yl)amino, (1,4,5,6-tetrahydro-5-methylpyrimidin-2-yl)amino, (1,4,5,6-tetrahydro-5,5-dimethylpyrimidin-2-yl)amino, (thiazol-2-yl)amino, (3H-imidazo[4,5-b]pyridin-2-yl)amino, R₃HNC(=NH)NH-, R₃HNC(=O)NH-, (4,5-dihydro-1H-imidazol-2-yl)aminooxy and R₃HNC(=NH)NHO-;

with the proviso that if A is H₂NC(=NH)NH-, then, dependently, W is not hydrogen when Q is -CH₂-;

R₁ is hydrogen;

R₃ is selected from the group consisting of hydrogen, butyl, benzyl and hydroxy;

Q is selected from the group consisting of -CH₂-, -CH(methyl)-, -CH(ethyl)-, -CH(phenyl)- (wherein phenyl is optionally substituted with one to five substituents independently selected from the group consisting of methyl, ethyl, propyl, methoxy, ethoxy, propoxy, bromine, chlorine, fluorine, iodine, hydroxy and trifluoromethyl), -CH(naphthalen-1-yl)-, -CH(naphthalen-2-yl)-, -CH[(3,4-dioxymethylene)phenyl]-, -CH[(3,4-dioxyethylene)phenyl]-, -CH[(3-bromo-5-chloro-2-hydroxy)phenyl]-, -CH(thien-3-yl)-, -CH(quinolin-3-yl)-, -CH(pyridin-3-yl)- (wherein pyridinyl is optionally substituted with chlorine) and -CH(benzyl)-;

R₄ is hydrogen;

T is selected from R₅OC(=O)-;

R₅ is selected from aryl(C₁-C₄)alkyl;

R₆ is hydrogen; and,

R₇ and R₈ are independently ethyl;

and pharmaceutically acceptable salts thereof.

3. The compound of Claim 2, wherein

A is optionally present and is selected from the group consisting of 4,5-dihydro-1H-imidazol-2-yl, 1,2,3,4-tetrahydro-1,8-naphthyridin-7-yl, 3H-imidazo[4,5-b]pyridin-2-yl, amino, (4,5-dihydro-1H-imidazol-2-yl)amino, (pyridin-2-yl)amino, (3-methylpyridin-2-yl)amino, (thiazol-2-yl)amino, (3H-imidazo[4,5-b]pyridin-2-yl)amino, NH₂C(=NH)NH-, R₃HNC(=O)NH-, (4,5-dihydro-1H-imidazol-2-yl)aminooxy and NH₂C(=NH)NHO-,

with the proviso that if A is H₂NC(=NH)NH-, then, dependently, W is not hydrogen when Q is -CH₂-;

Q is selected from the group consisting of -CH₂-, -CH(methyl)-, -CH(phenyl)- (wherein phenyl is optionally substituted with one to five substituents independently selected from the group consisting of methyl, methoxy, bromine, chlorine, fluorine, hydroxy and trifluoromethyl), -CH(naphthalen-1-yl)-, -CH(naphthalen-2-yl)-, -CH[(3,4-dioxymethylene)phenyl]-, -CH[(3,4-dioxyethylene)phenyl]-, -CH[(3-bromo-5-chloro-2-hydroxy)phenyl]-, -CH(thien-3-yl)-, -CH(quinolin-3-yl)-, -CH(pyridin-3-yl)- (wherein pyridinyl is optionally substituted with chlorine) and -CH(benzyl)-; and,

R₅ is benzyl;

and pharmaceutically acceptable salts thereof.

4. The compound of Claim 1 selected from the group consisting of:

β-[[[1-[4-[(4,5-dihydro-1H-imidazol-2-yl)amino]-1-oxobutyl]-4-piperidinyl]carbonyl]amino]-, (β³S)- 3-quinolinepropanoic acid;

3-[[[1-[4-[(4,5-dihydro-1H-imidazol-2-yl)amino]-1-oxobutyl]-4-piperidinyl]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanine;

β-[[[1-[1-oxo-4-(2-pyridinylamino)butyl]-4-piperidinyl]carbonyl]amino]-, (β³S)- 3-quinolinepropanoic acid;

3-[[[1-[1-[(4,5-dihydro-1H-imidazol-2-yl)-4-piperidinyl]carbonyl]-4-piperidinyl]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanine;


3-[[[1-[[[(1S,4R)-4-[(4,5-dihydro-1H-imidazol-2-yl)amino]-2-cyclopenten-1-yl]carbonyl]-4-piperidinyl]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanine;
 3-[[[1-[3-[(4,5-dihydro-1H-imidazol-2-yl)amino]-2-methyl-1-oxopropyl]-4-piperidinyl]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanine;
 3-[[[1-[[[(1R,4S)-4-[(4,5-dihydro-1H-imidazol-2-yl)amino]-2-cyclopenten-1-yl]carbonyl]-4-piperidinyl]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanine;
 4-[[[[(2S)-2-carboxy-2-[(phenylmethoxy)carbonyl]amino]ethyl]amino]carbonyl]-, 2-[(4,5-dihydro-1H-imidazol-2-yl)amino]ethylester-1-piperidinecarboxylic acid;
 β -[[[1-[2-[(4,5-dihydro-1H-imidazol-2-yl)amino]ethoxy]carbonyl]-4-piperidinyl]carbonyl]amino]-, (β^3 S)-3-quinoline-propanoic acid;
 3-[[[1-[4-[(4,5-dihydro-1H-imidazol-2-yl)amino]-2-methyl-1-oxobutyl]-4-piperidinyl]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanine;
 3-[[[1-(4-amino-1-oxo-5-hexenyl)-4-piperidinyl]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanine;
 3-[[[1-[4-[(4,5-dihydro-1H-imidazol-2-yl)amino]methyl]benzoyl]-4-piperidinyl]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanine;
 β -[[[1-[3-[(4,5-dihydro-1H-imidazol-2-yl)amino]oxy]-1-oxopropyl]-4-piperidinyl]carbonyl]amino]-1,3-benzodioxole-5-propanoic acid;
 and,
 β -[[[1-[1-oxo-4-(2-pyridinylamino)butyl]-4-piperidinyl]carbonyl]amino]-1,3-benzodioxole-5-propanoic acid;
 and pharmaceutically acceptable salts thereof.

5. The compound of any one of Claims 1 to 7 which is an effective integrin antagonist.
6. The compound of Claim 5 which is an effective $\alpha\text{v}\beta 3$, $\alpha\text{v}\beta 5$, GP11b/111a, dual $\alpha\text{v}\beta 3$ /GP11b/111a or dual $\alpha\text{v}\beta 3/\alpha\text{v}\beta 5$ integrin antagonist.
7. The compound of Claim 6 which is an effective $\alpha\text{v}\beta 3$ integrin antagonist selected from the group consisting of Compound 1, 2, 3, 5, 6, 12, 14, 16, 18, 21, 24, 25, 28, 29, 31, 33, 34, 39, 40, 44, 47, 49, 54, 55, 58 and 60.
8. The compound of Claim 7 wherein the compound is selected from the group consisting of Compound 1, 2, 6, 28, 39 and 47.
9. The compound of Claim 6 which is an effective GP11b/111a integrin antagonist selected from the group consisting of Compound 1, 2, 5, 11, 12, 14, 16, 17, 18, 19, 39, 47, 49, 50, 53, 54, 55 and 56.
10. The compound of Claim 9 wherein the compound is selected from the group consisting of Compound 2, 11, 14, 17, 50 and 53.
11. The compound of Claim 6 which is an effective $\alpha\text{v}\beta 5$ integrin antagonist selected from the group consisting of Compound 1, 2, 16, 45, 58 and 60.
12. The compound of Claim 11 wherein the compound is selected from the group consisting of Compound 1, 2 and 16.
13. The compound of Claim 6 which is an effective dual $\alpha\text{v}\beta 3$ /GP11b/111a integrin antagonist selected from the group consisting of Compound 2, 5, 12, 14, 16, 18, 39, 47, 49, 54 and 55.
14. The compound of Claim 13 wherein the compound is Compound 2.
15. The compound of Claim 6 which is an effective dual $\alpha\text{v}\beta 3/\alpha\text{v}\beta 5$ integrin antagonist selected from the group consisting of Compound 1, 2, 16, 58 and 60.
16. The compound of claim 15 wherein the compound is selected from the group consisting of Compound 1 and 2.
17. The compound of any one of Claims 1 to 16, which are effective prodrugs selected from the group consisting of esters, hydroxyamidines and hydroxyguanidines of a compound of any one of Claims 1 to 16.
18. The compound of any one of Claims 1 to 17 which is an effective agent for the treatment of an integrin-mediated disorder ameliorated by administration of an integrin antagonist.

19. The compound of Claim 18 wherein the integrin-mediated disorder is restenosis, unstable angina, a thromboembolic disorder, vascular injury, atherosclerosis, arterial thrombosis, venous thrombosis, a vaso-occlusive disorder, acute myocardial infarction, re-occlusion following thrombolytic therapy, re-occlusion following angioplasty, inflammation, rheumatoid arthritis, osteoporosis, a bone resorption disorder, cancer, tumor growth, angiogenesis, multiple sclerosis, a neurological disorder, asthma, macular degeneration, a diabetic complication or diabetic retinopathy.
20. A pharmaceutical composition comprising a compound of any one of Claims 1 to 19 and a pharmaceutically acceptable carrier.
21. The composition of Claim 20 adapted to provide a therapeutically effective amount of the compound of any one of Claims 1 to 19 which is between 0.01 mg/kg/day and 300 mg/kg/day.
22. A process for making a pharmaceutical composition comprising mixing a compound of any one of Claims 1 to 19 and a pharmaceutically acceptable carrier.
23. A compound of any one of Claims 1 to 19 or a pharmaceutical composition of Claim 20 or Claim 21 for use in treating an integrin-mediated disorder, such as restenosis, unstable angina, a thromboembolic disorder, vascular injury, atherosclerosis, arterial thrombosis, venous thrombosis, a vaso-occlusive disorder, acute myocardial infarction, re-occlusion following thrombolytic therapy, re-occlusion following angioplasty, inflammation, rheumatoid arthritis, osteoporosis, a bone resorption disorder, cancer, tumor growth, angiogenesis, multiple sclerosis, a neurological disorder, asthma, macular degeneration, a diabetic complication or diabetic retinopathy.

Patentansprüche

1. Verbindung der Formel (I):



wobei

M ausgewählt ist aus der Gruppe, bestehend aus Ethylen (fakultativ substituiert innerhalb der Kohlenstoffkette mit Methyl und am terminalen Kohlenstoff mit einem Substituenten substituiert, ausgewählt aus A), Propylen (fakultativ substituiert innerhalb der Kohlenstoffkette mit einem Substituenten, ausgewählt aus der Gruppe, bestehend aus Methyl, Ethenyl, Cyclohexyliden [wobei ein Ringkohlenstoffatom den Anhängungspunkt an die Kohlenstoffkette bildet] und 4-Cl-Phenyl, und am terminalen Kohlenstoff mit einem Substituenten substituiert, ausgewählt aus A), Allylen (substituiert mit einem Substituenten, ausgewählt aus A); Piperidin-4-yl, Piperidin-4-ylen (substituiert mit einem Substituenten, ausgewählt aus A), 1,4,5-Dihydro-2-cyclopenten-1-ylen (substituiert mit einem Substituenten, ausgewählt aus A) und 4-Methylphenylen (substituiert an Methylen mit einem Substituenten, ausgewählt aus A); A fakultativ vorhanden ist und ausgewählt ist aus der Gruppe, bestehend aus 1H-Imidazol-1-yl, 1H-Imidazol-2-yl, 4,5-Dihydro-1H-imidazol-2-yl (fakultativ mit einem Substituenten substituiert, ausgewählt aus der Gruppe, bestehend aus C₁-C₄-Alkoxycarbonyl und Aryl(C₁-C₄)-Alkoxycarbonyl), Pyridin-2-yl (fakultativ mit einem Substituenten substituiert, ausgewählt aus der Gruppe, bestehend aus C₁-C₄-Alkyl, Heteroaryl [fakultativ mit C₁-C₄-Alkyl substituiert], Halogen, Hydroxy, Nitro, Cyano, Amino, Amino(C₁-C₄)-Alkyl, C₁-C₄-Alkylamino(C₁-C₄)-Alkyl und Di(C₁-C₄-Alkyl)-amino-(C₁-C₄)-alkyl), Pyrimidin-2-yl, 1,4,5,6-Tetrahydro-pyrimidin-2-yl (fakultativ mit einem bis zwei Substituenten substituiert, unabhängig ausgewählt aus der Gruppe, bestehend aus C₁-C₄-Alkyl, Hydroxy und Amino), Piperidin-4-yl, Benzimidazol-2-yl, 1,2,3,4-Tetrahydro-1,8-naphthyridin-7-yl, 3H-Imidazo[4,5-b]-pyridin-2-yl, Amino, (C₁-C₆-Alkyl)-amino, (1H-Imidazol-1-yl)-amino, (1H-Imidazol-2-yl)-amino, (4,5-Dihydro-1H-imidazol-2-yl)-amino (fakultativ an 4,5-Dihydro-1H-imidazol-2-yl mit einem Substituenten substituiert, ausgewählt aus der Gruppe, bestehend aus (C₁-C₆)-Alkoxycarbonyl und Aryl(C₁-C₆)-alkoxycarbonyl), (Pyridin-2-yl)-amino (fakultativ am Py-

ridin-2-yl mit einem Substituenten substituiert, ausgewählt aus der Gruppe, bestehend aus C₁-C₄-Alkyl, Heteroaryl [fakultativ substituiert mit C₁-C₄-Alkyl], Halogen, Hydroxy, Nitro, Cyano, Amino, Amino(C₁-C₄)-alkyl, C₁-C₄-Aylamino (C₁-C₄)-Alkyl und Di(C₁-C₄-Alkyl)-amino(C₁-C₄)-alkyl), (Pyrimidin-2-yl)-amino, (1,4,5,6-Tetrahydropyrimidin-2-yl)-amino (fakultativ an 1,4,5,6-Tetrahydropyrimidin-2-yl mit einem bis zwei Substituenten substituiert, unabhängig ausgewählt aus der Gruppe, bestehend aus C₁-C₄-Alkyl, Hydroxy und Amino), (4,5,6,7-Tetrahydro-1H-1,3-diazepin-2-yl)-amino, (Thiazol-2-yl)-amino, (Benzimidazol-2-yl)-amino, (3H-Imidazo[4,5-b]-pyridin-2-yl)-amino, R₃HNC(=NH)NH-, R₃HNC(=O)NH-, (4,5,6,7-Tetrahydro-1H-1,3-diazepin-2-yl)aminoxy, (4,5-Dihydro-1H-imidazol-2-yl)aminoxy und R₃HNC(=NH)NHO-;

unter der Voraussetzung, daß, wenn A H₂NC(=NH)NH- ist, dann ist, abhängig, W nicht Wasserstoff, wenn Q -CH₂- ist; L ausgewählt ist aus der Gruppe, bestehend aus -C(=O)-, -OC(=O)- und -HNC(=O)-;

R₁ ausgewählt ist aus der Gruppe, bestehend aus Wasserstoff und C₁-C₄-Alkyl;

R₂ Wasserstoff ist;

R₃ ausgewählt ist aus der Gruppe, bestehend aus Wasserstoff, C₁-C₈-Alkyl, Aryl(C₁-C₈)-Alkyl und Hydroxy;

Q ausgewählt ist aus der Gruppe, bestehend aus -CH₂-, -CH(C₁-C₈-Alkyl)-, -CH(Aryl)-, (wobei Aryl fakultativ mit einem bis fünf Substituenten substituiert ist, unabhängig ausgewählt aus der Gruppe, bestehend aus C₁-C₄-Alkyl, C₁-C₄-Alkoxy, -O-(C₁-C₃-Alkyl)-O-, Halogen, Hydroxy und Trihalogen(C₁-C₃)-alkyl), -CH(Heteroaryl)-, (wobei Heteroaryl fakultativ mit einem Substituenten substituiert ist, ausgewählt aus Halogen) und -CH(Aryl(C₁-C₈)-alkyl)-; W ausgewählt ist aus der Gruppe, bestehend aus Wasserstoff und N(R₄)T;

r 1 ist;

R₄ ausgewählt ist aus der Gruppe, bestehend aus Wasserstoff und C₁-C₄-Alkyl;

T ausgewählt ist aus der Gruppe, bestehend aus R₅C(=O)- und R₅OC(=O)-;

R₅ ausgewählt ist aus der Gruppe, bestehend aus Aryl und Aryl(C₁-C₄)-alkyl;

R₆ ausgewählt ist aus der Gruppe, bestehend aus Wasserstoff, Methyl und (R₈)(R₇)NC(=O)CH₂; und

R₇ und R₈ unabhängig ausgewählt sind aus der Gruppe, bestehend aus Wasserstoff, Methyl und Ethyl;

und pharmazeutisch akzeptable Salze davon.

2. Verbindung nach Anspruch 1, wobei

A fakultativ vorhanden ist und ausgewählt ist aus der Gruppe, bestehend aus 1H-Imidazol-1-yl, 4,5-Dihydro-1H-imidazol-2-yl, Pyridin-2-yl (fakultativ mit einem Substituenten substituiert, ausgewählt aus der Gruppe, bestehend aus C₁-C₄-Alkyl und Heteroaryl [fakultativ mit C₁-C₄-Alkyl substituiert]), Piperidin-4-yl, Benzimidazol-2-yl, 1,2,3,4-Tetrahydro-1,8-naphthyridin-7-yl, 3H-Imidazo[4,5-b]pyridin-2-yl, Amino, (4,5-Dihydro-1H-imidazol-2-yl)-amino (fakultativ substituiert an 4,5-Dihydro-1H-imidazol-2-yl mit C₁-C₆-alkoxycarbonyl), (Pyridin-2-yl)-amino (fakultativ substituiert am Pyridinyl mit einem Substituenten, ausgewählt aus der Gruppe, bestehend aus C₁-C₄-Alkyl und Heteroaryl [fakultativ mit C₁-C₄-Alkyl substituiert]), (1,4,5,6-Tetrahydro-5-hydroxypyrimidin-2-yl)-amino, (1,4,5,6-Tetrahydro-5-methylpyrimidin-2-yl)-amino, (1,4,5,6-Tetrahydro-5,5-dimethylpyrimidin-2-yl)-amino, (Thiazol-2-yl)-amino, (3H-Imidazo[4,5-b]-pyridin-2-yl)-amino, R₃HNC(=NH)NH-, R₃HNC(=O)NH-, (4,5-Dihydro-1H-imidazol-2-yl)-aminoxy und R₃HNC(=NH)NHO-;

unter der Voraussetzung, daß, wenn A H₂NC(=NH)NH- ist, dann, abhängig, W nicht Wasserstoff ist, wenn Q -CH₂- ist;

R₁ Wasserstoff ist;

R₃ ausgewählt ist aus der Gruppe, bestehend aus Wasserstoff, Butyl, Benzyl und Hydroxy;

Q ausgewählt ist aus der Gruppe, bestehend aus -CH₂-, -CH(Methyl)-, CH(Ethyl)-, -CH(Phenyl)-, wobei Phenyl fakultativ mit einem bis fünf Substituenten substituiert ist, unabhängig ausgewählt aus der Gruppe, bestehend aus Methyl, Ethyl, Propyl, Methoxy, Ethoxy, Propoxy, Brom, Chlor, Fluor, Iod, Hydroxy und Trifluormethyl), -CH(Naphthalen-1-yl)-, -CH(Naphthalen-2-yl)-, -CH[(3,4-Dioxymethylen)phenyl]-, -CH[(3,4-Dioxyethylen)phenyl]-, -CH[(3-Brom-5-chlor-2-hydroxy)phenyl]-, -CH(Thien-3-yl)-, -CH(Chinolin-3-yl)-, -CH(Pyridin-3-yl)- (wobei Pyridinyl fakultativ mit Chlor substituiert ist) und -CH(Benzyl)-;

R₄ Wasserstoff ist;

T ausgewählt ist aus R₅OC(=O)-;

R₅ ausgewählt ist aus Aryl(C₁-C₄)-alkyl;

R₆ Wasserstoff ist; und

R₇ und R₈ unabhängig Ethyl sind;

und pharmazeutisch akzeptable Salze davon.

3. Verbindung nach Anspruch 2, wobei

A fakultativ vorhanden ist und ausgewählt ist aus der Gruppe, bestehend aus 4,5-Dihydro-1H-imidazol-2-yl, 1,2,3,4-Tetrahydro-1,8-naphthyridin-7-yl, 3H-Imidazo[4,5-b]pyridin-2-yl, Amino, (4,5-Dihydro-1H-imidazol-2-yl)-amino, (Pyridin-2-yl)-amino, (3-Methylpyridin-2-yl)-amino, (Thiazol-2-yl)-amino, (3H-Imidazo[4,5-b]-pyridin-2-yl)-amino, NH₂C(=NH)NH-, R₃HNC(=O)NH-, (4,5-Dihydro-1H-imidazol-2-yl)aminoxy und NH₂C(=NH)NHO-,

unter der Voraussetzung, daß, wenn $AH_2NC(=NH)NH-$ ist, dann, abhängig, W nicht Wasserstoff ist, wenn $Q-CH_2-$ ist; Q ausgewählt ist aus der Gruppe, bestehend aus $-CH_2-$, $-CH(\text{Methyl})-$, $-CH(\text{Phenyl})-$ (wobei Phenyl fakultativ mit einem bis fünf Substituenten substituiert ist, unabhängig ausgewählt aus der Gruppe, bestehend aus Methyl, Methoxy, Brom, Chlor, Fluor, Hydroxy und Trifluormethyl), $-CH(\text{Naphthalen-1-yl})-$, $-CH(\text{Naphthalen-2-yl})-$, $-CH[(3,4\text{-Dioxymethylen})\text{phenyl}]-$, $-CH[(3,4\text{-Dioxyethylen})\text{phenyl}]-$, $-CH[(3\text{-Brom-5-chlor-2-hydroxy})\text{phenyl}]-$, $-CH(\text{Thien-3-yl})-$, $-CH(\text{Chinolin-3-yl})-$, $-CH(\text{Pyridin-3-yl})-$, (wobei Pyridinyl fakultativ mit Chlor substituiert ist), und $-CH(\text{Benzyl})-$; und R_5 Benzyl ist; und pharmazeutisch akzeptable Salze davon.

4. Verbindung nach Anspruch 1, ausgewählt aus der Gruppe, bestehend aus:

β -[[[1-[4-[(4,5-Dihydro-1H-imidazol-2-yl)amino]-1-oxobutyl]-4-piperidiny]carbonyl]amino]-, (β^3S)-3-chinolinpropansäure;
 3-[[[1-[4-[(4,5-Dihydro-1H-imidazol-2-yl)amino]-1-oxobutyl]-4-piperidiny]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanin;
 β -[[[1-[1-Oxo-4-(2-pyridinylamino)butyl]-4-piperidiny]carbonyl]amino]-, (β^3S)-3-chinolinpropansäure;
 3-[[[1-[1-(4,5-Dihydro-1H-imidazol-2-yl)-4-piperidiny]carbonyl]-4-piperidiny]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanin;
 3-[[[1-[[1S, 4R)-4-[(4,5-Dihydro-1H-imidazol-2-yl)amino]-2-cyclopenten-1-yl]carbonyl]-4-piperidiny]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanin;
 3-[[[1-[3-[(4,5-Dihydro-1H-imidazol-2-yl)amino]-2-methyl-1-oxopropyl]-4-piperidiny]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanin;
 3-[[[1-[[1R, 4S)-4-[(4,5-Dihydro-1H-imidazol-2-yl)amino]-2-cyclopenten-1-yl]carbonyl]-4-piperidiny]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanin;
 4-[[[2S)-2-Carboxy-2-[(phenylmethoxy)carbonyl]amino]ethyl]amino]carbonyl]-, 2-[(4,5-dihydro-1H-imidazol-2-yl)amino]ethylester-1-piperidincarbonsäure;
 β -[[[1-[2-[(4,5-Dihydro-1H-imidazol-2-yl)amino]ethoxy]carbonyl]-4-piperidiny]carbonyl]amino]-, (β^3S)-3-chinolinpropansäure;
 3-[[[1-[4-[(4,5-Dihydro-1H-imidazol-2-yl)amino]-2-methyl-1-oxobutyl]-4-piperidiny]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanin;
 3-[[[1-(4-amino-1-oxo-5-hexenyl)-4-piperidiny]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanin;
 3-[[[1-[4-[(4,5-Dihydro-1H-imidazol-2-yl)amino]methyl]-benzoyl]-4-piperidiny]carbonyl]amino]-N-[(phenylmethoxy)carbonyl]-L-alanin;
 β -[[[1-[3-[(4,5-Dihydro-1H-imidazol-2-yl)amino]oxy]-1-oxopropyl]-4-piperidiny]carbonyl]amino]-1,3-benzodioxol-5-propansäure; und
 β -[[[1-[1-oxo-4-(2-pyridinylamino)butyl]-4-piperidiny]carbonyl]amino]-1,3-benzodioxol-5-propansäure;

und pharmazeutisch akzeptable Salze davon.

5. Verbindung nach einem der Ansprüche 1 bis 7, die ein effektiver Integrin-Antagonist ist.

6. Verbindung nach Anspruch 5, die ein effektiver $\alpha v\beta 3$ -, $\alpha v\beta 5$ -, GP11b/111a, dualer $\alpha v\beta 3$ /GP11b/111a - oder dualer $\alpha v\beta 3$ / $\alpha v\beta 5$ -Integrin-Antagonist ist.

7. Verbindung nach Anspruch 6, die ein effektiver $\alpha v\beta 3$ -Integrin-Antagonist ist, ausgewählt aus der Gruppe, bestehend aus Verbindung 1, 2, 3, 5, 6, 12, 14, 16, 18, 21, 24, 25, 28, 29, 31, 33, 34, 39, 40, 44, 47, 49, 54, 55, 58 und 60.

8. Verbindung nach Anspruch 7, wobei die Verbindung ausgewählt ist aus der Gruppe, bestehend aus Verbindung 1, 2, 6, 28, 39 und 47.

9. Verbindung nach Anspruch 6, die ein effektiver GP11b/111a-Integrin-Antagonist ist, ausgewählt aus der Gruppe, bestehend aus Verbindung 1, 2, 5, 11, 12, 14, 16, 17, 18, 19, 39, 47, 49, 50, 53, 54, 55 und 56.

10. Verbindung nach Anspruch 9, wobei die Verbindung ausgewählt ist aus der Gruppe, bestehend aus Verbindung 2, 11, 14, 17, 50 und 53.

11. Verbindung nach Anspruch 6, die ein effektiver $\alpha v\beta 5$ -Integrin-Antagonist ist, ausgewählt aus der Gruppe, bestehend

aus Verbindung 1, 2, 16, 45, 58 und 60.

12. Verbindung nach Anspruch 11, wobei die Verbindung ausgewählt ist aus der Gruppe, bestehend aus Verbindung 1, 2 und 16.

13. Verbindung nach Anspruch 6, die ein effektiver dualer $\alpha v \beta 3 / GP11b / 111a$ -Integrin-Antagonist ist, ausgewählt aus der Gruppe, bestehend aus Verbindung 2, 5, 12, 14, 16, 18, 39, 47, 49, 54 und 55.

14. Verbindung nach Anspruch 13, wobei die Verbindung Verbindung 2 ist.

15. Verbindung nach Anspruch 6, die ein effektiver dualer $\alpha v \beta 3 / \alpha v \beta 5$ -Integrin-Antagonist ist, ausgewählt aus der Gruppe, bestehend aus Verbindung 1, 2, 16, 58 und 60.

16. Verbindung nach Anspruch 15, wobei die Verbindung ausgewählt ist aus der Gruppe, bestehend aus Verbindung 1 und 2.

17. Verbindung nach einem der Ansprüche 1 bis 16, die effektive Prodrugs sind, ausgewählt aus der Gruppe, bestehend aus Estern, Hydroxyamidinen und Hydroxyguanidinen einer Verbindung nach einem der Ansprüche 1 bis 16.

18. Verbindung nach einem der Ansprüche 1 bis 17, die ein effektives Mittel für die Behandlung einer Integrin-vermittelten Erkrankung ist, die durch die Verabreichung eines Integrin-Antagonisten gelindert wird.

19. Verbindung nach Anspruch 18, wobei die Integrin-vermittelte Erkrankung Restenose, instabile Angina, eine thromboembolische Erkrankung, Gefäßerkrankung, Arteriosklerose, arterielle Thrombose, venöse Thrombose, eine Gefäßverschlusserkrankung, akuter Myokardinfarkt, Wiederverschluß nach thrombolytischer Therapie, Wiederverschluß nach Angioplastie, Entzündung, rheumatoide Arthritis, Osteoporose, eine Knochenresorptionserkrankung, Krebs, Tumorwachstum, Angiogenese, multiple Sklerose, eine neurologische Erkrankung, Asthma, Macula-Degeneration, eine Diabeteskomplikation oder Diabetes-Retinopathie ist.

20. Pharmazeutische Zusammensetzung, umfassend eine Verbindung nach einem der Ansprüche 1 bis 19 und einen pharmazeutisch akzeptablen Träger.


21. Verbindung nach Anspruch 20, die angepaßt ist, eine therapeutisch wirksame Menge der Verbindung nach einem der Ansprüche 1 bis 19 bereitzustellen, die zwischen 0,01 mg/kg/Tag und 300 mg/kg/Tag ist.

22. Verfahren zum Herstellen einer pharmazeutischen Zusammensetzung, umfassend das Mischen einer Verbindung nach einem der Ansprüche 1 bis 19 und einen pharmazeutisch akzeptablen Träger.

23. Verbindung nach einem der Ansprüche 1 bis 19 oder eine pharmazeutische Zusammensetzung nach Anspruch 20 oder Anspruch 21 zur Verwendung bei der Behandlung einer Integrin-vermittelten Erkrankung, wie etwa Restenose, instabile Angina, eine thromboembolische Erkrankung, Gefäßerkrankung, Arteriosklerose, arterielle Thrombose, venöse Thrombose, eine Gefäßverschlusserkrankung, akuter Myokardinfarkt, Wiederverschluß nach thrombolytischer Therapie, Wiederverschluß nach Angioplastie, Entzündung, rheumatoide Arthritis, Osteoporose, eine Knochenresorptionserkrankung, Krebs, Tumorwachstum, Angiogenese, multiple Sklerose, eine neurologische Erkrankung, Asthma, Macula-Degeneration, eine Diabetes-Komplikation oder Diabetes-Retinopathie.

Revendications

1. Composé de formule (I) :



dans laquelle

M est choisi dans le groupe constitué par l'éthylène (facultativement substitué à l'intérieur de la chaîne de carbone par un groupe méthyle et substitué sur l'atome de carbone terminal par un substituant choisi parmi A), le propylène (facultativement substitué à l'intérieur de la chaîne de carbone par un substituant choisi dans le groupe constitué par un groupe méthyle, un groupe éthényle, un groupe cyclohexylidène [dans lequel un atome de carbone cyclique forme le point de liaison à la chaîne de carbone] et un groupe 4-Cl-phényle et substitué sur l'atome de carbone terminal par un substituant choisi parmi A), l'allylène (substitué par un substituant choisi parmi A) ; le pipéridin-4-yl, le pipéridin-4-ylène (substitué par un substituant choisi parmi A), le 1,4,5-dihydro-2-cyclopenten-1-ylène (substitué par un substituant choisi parmi A) et le 4-méthylphénylène (substitué sur le méthylène par un substituant choisi parmi A) ;

A est facultativement présent et est choisi dans le groupe constitué par le 1H-imidazol-1-yl, le 1H-imidazol-2-yl, le 4,5-dihydro-1H-imidazol-2-yl (facultativement substitué par un substituant choisi dans le groupe constitué par un groupe alcoxy en C₁ à C₄ carbonyle et un groupe aryl(alcoxycarbonyle en C₁ à C₄)), le pyridin-2-yl (facultativement substitué par un substituant choisi dans le groupe constitué par un groupe alkyle en C₁ à C₄, un groupe hétéroaryle [facultativement substitué par un groupe alkyle en C₁ à C₄], un atome d'halogène, un groupe hydroxy, un groupe nitro, un groupe cyano, un groupe amino, un groupe amino-(alkyle en C₁ à C₄), un groupe (alkylamino en C₁ à C₄)-(alkyle en C₁ à C₄) et un groupe di(alkyle en C₁ à C₄) amino (alkyle en C₁ à C₄)), le pyrimidin-2-yl, le 1,4,5,6-tétrahydropyrimidin-2-yl (facultativement substitué par un ou deux substituants indépendamment choisis dans le groupe constitué par un groupe alkyle en C₁ à C₄, un groupe hydroxy et un groupe amino), le pipéridin-4-yl, le benzimidazol-2-yl, le 1,2,3,4-tétrahydro-1,8-naphtyridin-7-yl, le 3H-imidazo[4,5-b]pyridin-2-yl, un groupe amino, un groupe (alkyl en C₁ à C₆)amino, un groupe (1H-imidazol-1-yl)amino, un groupe (1H-imidazol-2-yl)amino, un groupe (4,5-dihydro-1H-imidazol-2-yl)-amino (facultativement substitué sur le 4,5-dihydro-1H-imidazol-2-yl par un substituant choisi dans le groupe constitué par un groupe alcoxy en C₁ à C₆ carbonyle et un groupe aryl(alcoxy en C₁ à C₆ carbonyle), un groupe (pyridin-2-yl)amino (facultativement substitué sur le pyridin-2-yl par un substituant choisi dans le groupe constitué par un groupe alkyle en C₁ à C₄, un groupe hétéroaryle [facultativement substitué par un groupe alkyle en C₁ à C₄], un atome d'halogène, un groupe hydroxy, un groupe nitro, un groupe cyano, un groupe amino, un groupe amino(alkyle en C₁ à C₄), un groupe (alkylamino en C₁ à C₄) (alkyle en C₁ à C₄) et un groupe di(alkyl en C₁ à C₄) amino (alkyle en C₁ à C₄), un groupe (pyrimidin-2-yl)amino, un groupe (1,4,5,6-tétrahydropyrimidin-2-yl)amino (facultativement substitué sur le 1,4,5,6-tétrahydropyrimidin-2-yl par un ou deux substituants indépendamment choisis dans le groupe constitué par un groupe alkyle en C₁ à C₄, un groupe hydroxy et un groupe amino), un groupe (4,5,6,7-tétrahydro-1H-1,3-diazépin-2-yl)amino, un groupe (thiazol-2-yl)amino, un groupe (benzimidazol-2-yl)amino, un groupe (3H-imidazo-[4, 5-b] pyridin-2-yl) amino, R₃HNC(=NH)NH-, R₃HNC(=O)NH-, un groupe (4,5,6,7-tétrahydro-1H-1,3-diazépin-2-yl)aminooxy, un groupe (4,5-dihydro-1H-imidazol-2-yl)aminooxy et R₃HNC(=NH)NHO- ;

à condition que si A est H₂NC(=NH)NH-, alors, de manière dépendante, W n'est pas un atome d'hydrogène lorsque Q est -CH₂- ;

L est choisi dans le groupe constitué par -C(=O)-, -OC(=O)- et -HNC(=O)- ;

R₁ est choisi dans le groupe constitué par un atome d'hydrogène et un groupe alkyle en C₁ à C₄ ;

R₂ est un atome d'hydrogène ;

R₃ est choisi dans le groupe constitué par un atome d'hydrogène, un groupe alkyle en C₁ à C₈, un groupe aryl(alkyle en C₁ à C₈) et un groupe hydroxy ;

Q est choisi dans le groupe constitué par un groupe -CH₂-, un groupe -CH(alkyle en C₁ à C₈)-, un groupe -CH(aryle)- (dans lequel le groupe aryle est facultativement substitué par un à cinq substituants indépendamment choisis dans le groupe constitué par un groupe alkyle en C₁ à C₄, un groupe alcoxy en C₁ à C₄, un groupe -O-(alkyl en C₁ à C₃)-O-, un atome d'halogène, un groupe hydroxy et un groupe trihalo(alkyle en C₁ à C₃)), un groupe -CH(hétéroaryle)- (dans lequel le groupe hétéroaryle est facultativement substitué par un substituant choisi parmi un atome d'halogène) et un groupe -CH(aryl(alkyle en C₁ à C₈))- ;

W est choisi dans le groupe constitué par un atome d'hydrogène et N(R₄)T ;

r est 1 ;

R₄ est choisi dans le groupe constitué par un atome d'hydrogène et un groupe alkyle en C₁ à C₄ ;

T est choisi dans le groupe constitué par R₅C(=O)- et R₅OC(=O)- ;

R₅ est choisi dans le groupe constitué par un groupe aryle et un groupe aryl(alkyle en C₁ à C₄) ;

R₆ est choisi dans le groupe constitué par un atome d'hydrogène, un groupe méthyle et (R₈)(R₇)NC(=O)CH₂ ; et, R₇ et R₈ sont indépendamment choisis dans le groupe constitué par un atome d'hydrogène, un groupe méthyle et un groupe éthyle ;

et les sels pharmaceutiquement acceptables de celui-ci.

2. Composé selon la revendication 1, dans lequel

A est facultativement présent et est choisi dans le groupe constitué par le 1H-imidazol-1-yl, le 4,5-dihydro-1H-imidazol-2-yl, le pyridin-2-yl (facultativement substitué par un substituant choisi dans le groupe constitué par un groupe alkyle en C₁ à C₄ et un groupe hétéroaryle [facultativement substitué par un groupe alkyle en C₁ à C₄]), le pipéridin-4-yl, le benzimidazol-2-yl, le 1,2,3,4-tétrahydro-1,8-naphtyridin-7-yl, le 3H-imidazo[4,5-b]pyridin-2-yl, un groupe amino, un groupe (4,5-dihydro-1H-imidazol-2-yl)amino (facultativement substitué sur le 4,5-dihydro-1H-imidazol-2-yl par un groupe alcoxy en C₁ à C₆ carbonyle), un groupe (pyridin-2-yl)amino (facultativement substitué sur le groupe pyridinyle par un substituant choisi dans le groupe constitué par un groupe alkyle en C₁ à C₄ et un groupe hétéroaryle [facultativement substitué par un groupe alkyle en C₁ à C₄]), un groupe (1,4,5,6-tétrahydro-5-hydroxypyrimidin-2-yl)amino, un groupe (1,4,5,6-tétrahydro-5-méthylpyrimidin-2-yl)amino, un groupe (1,4,5,6-tétrahydro-5,5-diméthylpyrimidin-2-yl)amino, un groupe (thiazol-2-yl)amino, un groupe (3H-imidazo[4,5-b]pyridin-2-yl) amino, R₃HNC(=NH)NH-, R₃HNC(=O)NH-, un groupe (4,5-dihydro-1H-imidazol-2-yl)aminoxy et R₃HNC(=NH)NHO- ;

à condition que si A est H₂NC(=NH)NH-, alors, de manière dépendante, W n'est pas un atome d'hydrogène lorsque Q est -CH₂- ;

R₁ est un atome d'hydrogène ;

R₃ est choisi dans le groupe constitué par un atome d'hydrogène, un groupe butyle, un groupe benzyle et un groupe hydroxy ;

Q est choisi dans le groupe constitué par un groupe -CH₂-, un groupe -CH(méthyle)-, un groupe -CH(éthyle)-, un groupe -CH(phényle)- (dans lequel le groupe phényle est facultativement substitué par un à cinq substituants indépendamment choisis dans le groupe constitué par un groupe méthyle, un groupe éthyle, un groupe propyle, un groupe méthoxy, un groupe éthoxy, un groupe propoxy, un atome de brome, un atome de chlore, un atome de fluor, un atome d'iode, un groupe hydroxy et un groupe trifluorométhyle), un groupe -CH(naphtalène-1-yl)-, un groupe -CH(naphtalène-2-yl)-, un groupe -CH[(3,4-dioxyéthylène)phényle]-, un groupe -CH[(3,4-dioxyéthylène)phényle]-, un groupe -CH[(3-bromo-5-chloro-2-hydroxy)phényle]-, un groupe -CH(thien-3-yl)-, un groupe -CH(quinolin-3-yl)-, un groupe -CH(pyridin-3-yl)- (dans lequel le groupe pyridinyle est facultativement substitué par un atome de chlore) et un groupe -CH(benzyle)- ;

R₄ est un atome d'hydrogène ;

T est choisi parmi R₅OC(=O)- ;

R₅ est choisi parmi un groupe aryl(alkyle en C₁ à C₄) ;

R₆ est un atome d'hydrogène ; et,

R₇ et R₈ sont indépendamment un groupe éthyle ;

et les sels pharmaceutiquement acceptables de celui-ci.

3. Composé selon la revendication 2, dans lequel

A est facultativement présent et est choisi dans le groupe constitué par le 4,5-dihydro-1H-imidazol-2-yl, le 1,2,3,4-tétrahydro-1,8-naphtyridin-7-yl, le 3H-imidazo[4,5b]pyridin-2-yl, un groupe amino, un groupe (4,5-dihydro-1H-imidazol-2-yl)amino, un groupe (pyridin-2-yl)amino, un groupe (3-méthylpyridin-2-yl)amino, un groupe (thiazol-2-yl) amino, un groupe (3H-imidazo[4,5-b]pyridin-2-yl)amino, NH₂C(=NH)NH-, R₃HNC(=O)NH-, un groupe (4,5-dihydro-1H-imidazol-2-yl)aminoxy et NH₂C(=NH)NHO-,

à condition que si A est H₂NC(=NH)NH-, alors, de manière dépendante, W n'est pas un atome d'hydrogène lorsque Q est -CH₂- ;

Q est choisi dans le groupe constitué par un groupe -CH₂-, un groupe -CH(méthyle)-, un groupe -CH(phényle)- (dans lequel le groupe phényle est facultativement substitué par un à cinq substituants indépendamment choisis dans le groupe constitué par un groupe méthyle, un groupe méthoxy, un atome de brome, un atome de chlore, un atome de fluor, un groupe hydroxy et un groupe trifluorométhyle), un groupe -CH(naphtalène-1-yl)-, un groupe -CH(naphtalène-2-yl)-, un groupe -CH[(3,4-dioxyéthylène)phényle]-, un groupe -CH[(3,4-dioxyéthylène)phényle]-, un groupe -CH[(3-bromo-5-chloro-2-hydroxy)phényle]-, un groupe -CH(thien-3-yl)-, un groupe -CH(quinolin-3-yl)-, un groupe -CH(pyridin-3-yl)- (dans lequel le groupe pyridinyle est facultativement substitué par un atome de chlore) et un

groupe -CH(benzyle)- ; et,
 R_5 est un groupe benzyle ;
 et les sels pharmaceutiquement acceptables de celui-ci.

5 4. Composé selon la revendication 1, choisi dans le groupe constitué par :

β -[[[1-[4-[(4,5-dihydro-1H-imidazol-2-yl)amino]-1-oxobutyl]-4-pipéridinyl]carbonyl]amino]-, acide (β^3S)-3-quinolinepropanoïque ;
 3-[[[1-(4-[(4,5-dihydro-1H-imidazol-2-yl)amino]-1-oxobutyl]-4-pipéridinyl]carbonyl]amino]-N-[(phénylméthoxy)carbonyl]-L-alanine ;
 10 β -[[[1-[1-oxo-4-(2-pyridinylamino)butyl]-4-pipéridinyl]carbonyl]amino]-, acide (β^3S)-3-quinoline-propanoïque ;
 3-[[[1-[[1-(4,5-dihydro-1H-imidazol-2-yl)-4-pipéridinyl]carbonyl]4-pipéridinyl]carbonyl]amino]-N-[(phénylméthoxy)carbonyl]-L-alanine ;
 15 3-[[[1-[[1S,4R)-4-[(4,5-dihydro-1H-imidazol-2-yl)-amino]-2-cyclopenten-1-yl]carbonyl]-4-pipéridinyl]carbonyl]amino]-N-[(phénylméthoxy)carbonyl]-L-alanine ;
 3-[[[1-[3-[(4,5-dihydro-1H-imidazol-2-yl)amino]-2-méthyl-1-oxopropyl]-4-pipéridinyl]carbonyl]amino]-N-[(phénylméthoxy)carbonyl]-L-alanine ;
 3-[[[1-[[1R,4S)-4-[(4,5-dihydro-1H-imidazol-2-yl)amino]-2-cyclopenten-1-yl]carbonyl]-4-pipéridinyl]carbonyl]amino]-N-[(phénylméthoxy)carbonyl]-L-alanine ;
 20 4-[[[1-(2S)-2-carboxy-2-[(phénylméthoxy)carbonyl]-amino]éthyl]amino]carbonyl]-, ester 2-[(4,5-dihydro-1H-imidazol-2-yl)amino]éthylrique de l'acide 1-pipéridine-carboxylique ;
 β -[[[1-[[2-[(4,5-dihydro-1H-imidazol-2-yl)amino]éthoxy]-carbonyl]-4-pipéridinyl]carbonyl]amino]-, acide (β^3S)-3-quinoline-propanoïque ;
 25 3-[[[1-[4-[(4,5-dihydro-1H-imidazol-2-yl)amino]-2-méthyl-1-oxobutyl]-4-pipéridinyl]carbonyl]amino]-N-[(phénylméthoxy)carbonyl]-L-alanine ;
 3-[[[1-(4-amino-1-oxo-5-hexenyl)-4-pipéridinyl]carbonyl]amino]-N-[(phénylméthoxy)carbonyl]-L-alanine ;
 3-[[[1-[4-[(4,5-dihydro-1H-imidazol-2-yl)amino]méthyl]benzoyl]-4-pipéridinyl]carbonyl]amino]-N-[(phénylméthoxy)carbonyl]-L-alanine ;
 30 acide β -[[[1-[3-[(4,5-dihydro-1H-imidazol-2-yl)amino]-oxy]-1-oxopropyl]-4-pipéridinyl]carbonyl]amino]-1,3-benzodioxole-5-propanoïque ; et,
 acide β -[[[1-[1-oxo-4-(2-pyridinylamino)butyl]-4-pipéridinyl]carbonyl]amino]-1,3-benzodioxole-5-propanoïque ;

et les sels pharmaceutiquement acceptables de celui-ci.

35 5. Composé selon l'une quelconque des revendications 1 à 4, qui est un antagoniste efficace de l'intégrine.

6. Composé selon la revendication 5, qui est un antagoniste efficace des intégrines $\alpha v \beta 3$, $\alpha v \beta 5$, GP11b/111a, double de $\alpha v \beta 3$ /GP11b/111a ou double de $\alpha v \beta 3/\alpha v \beta 5$.

40 7. Composé selon la revendication 6, qui est un antagoniste efficace de l'intégrine $\alpha v \beta 3$ choisi dans le groupe constitué par les composés 1, 2, 3, 5, 6, 12, 14, 16, 18, 21, 24, 25, 28, 29, 31, 33, 34, 39, 40, 44, 47, 49, 54, 55, 58 et 60.

8. Composé selon la revendication 7, dans lequel le composé est choisi dans le groupe constitué par les composés 1, 2, 6, 28, 39 et 47.

45 9. Composé selon la revendication 6, qui est un antagoniste efficace de l'intégrine GP11b/111a choisi dans le groupe constitué par les composés 1, 2, 5, 11, 12, 14, 16, 17, 18, 19, 39, 47, 49, 50, 53, 54, 55 et 56.

50 10. Composé selon la revendication 9, dans lequel le composé est choisi dans le groupe constitué par les composés 2, 11, 14, 17, 50 et 53.

11. Composé selon la revendication 6, qui est un antagoniste efficace de l'intégrine $\alpha v \beta 5$ choisi dans le groupe constitué par les composés 1, 2, 16, 45, 58 et 60.

55 12. Composé selon la revendication 11, dans lequel le composé est choisi dans le groupe constitué par les composés 1, 2 et 16.

13. Composé selon la revendication 6, qui est un antagoniste efficace et double des intégrines $\alpha v \beta 3$ /GP11b/111a choisi

dans le groupe constitué par les composés 2, 5, 12, 14, 16, 18, 39, 47, 49, 54 et 55.

14. Composé selon la revendication 13, dans lequel le composé est le composé 2.

15. Composé selon la revendication 6, qui est un antagoniste efficace et double des intégrines $\alpha v\beta 3/\alpha v\beta 5$ choisi dans le groupe constitué par les composés 1, 2, 16, 58 et 60.

16. Composé selon la revendication 15, dans lequel le composé est choisi dans le groupe constitué par les composés 1 et 2.

17. Composé selon l'une quelconque des revendications 1 à 16, qui est un promédicament efficace choisi dans le groupe constitué par les esters, les hydroxyamidines et les hydroxyguanidines d'un composé selon l'une quelconque des revendications 1 à 16.

18. Composé selon l'une quelconque des revendications 1 à 17, qui est un agent efficace pour le traitement d'un trouble provoqué par l'intégrine et amélioré par l'administration d'un antagoniste de l'intégrine.

19. Composé selon la revendication 18, dans lequel le trouble provoqué par l'intégrine est la resténose, un angor instable, un trouble thromboembolique, une lésion vasculaire, l'athérosclérose, la thrombose artérielle, la thrombose veineuse, un trouble vaso-occlusif, un infarctus du myocarde aigu, une ré-occlusion après une thérapie thrombolytique, une ré-occlusion après une angioplastie, une inflammation, la polyarthrite rhumatoïde, l'ostéoporose, un trouble de la résorption osseuse, un cancer, une croissance tumorale, l'angiogenèse, la sclérose en plaques, un trouble neurologique, l'asthme, la dégénérescence maculaire, une complication du diabète et la rétinopathie diabétique.

20. Composition pharmaceutique comprenant un composé selon l'une quelconque des revendications 1 à 19 et un vecteur pharmaceutiquement acceptable.

21. Composition selon la revendication 20, adaptée pour fournir une quantité thérapeutiquement efficace du composé selon l'une quelconque des revendications 1 à 19 qui est comprise entre 0,01 mg/kg/jour et 300 mg/kg/jour.

22. Procédé de fabrication d'une composition pharmaceutique comprenant le mélange d'un composé selon l'une quelconque des revendications 1 à 19 et un vecteur pharmaceutiquement acceptable.

23. Composé selon l'une quelconque des revendications 1 à 19 ou une composition pharmaceutique selon la revendication 20 ou la revendication 21, destiné(e) à être utilisé(e) dans le traitement d'un trouble provoqué par l'intégrine, tel que la resténose, un angor instable, un trouble thromboembolique, une lésion vasculaire, l'athérosclérose, la thrombose artérielle, la thrombose veineuse, un trouble vaso-occlusif, un infarctus du myocarde aigu, une ré-occlusion après une thérapie thrombolytique, une ré-occlusion après une angioplastie, une inflammation, la polyarthrite rhumatoïde, l'ostéoporose, un trouble de la résorption osseuse, un cancer, une croissance tumorale, l'angiogenèse, la sclérose en plaques, un trouble neurologique, l'asthme, la dégénérescence maculaire, une complication du diabète et la rétinopathie diabétique.