



(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

Note: Bibliography reflects the latest situation

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see page(s) 2

(51) Int Cl.7: **A61K 35/74**, A61K 35/72,
A23L 1/03, A23L 1/308

(48) Corrigendum issued on:
27.10.2004 Bulletin 2004/44

(86) International application number:
PCT/NL1999/000755

(45) Date of publication and mention
of the grant of the patent:
07.04.2004 Bulletin 2004/15

(87) International publication number:
WO 2000/033854 (15.06.2000 Gazette 2000/24)

(21) Application number: **99962557.7**

(22) Date of filing: **09.12.1999**

(54) **PREPARATION WHICH CONTAINS OLIGOSACCHARIDES AND PROBIOTICS**

OLIGOSACCHARIDE UND PROBIOTICA ENTHALTENDES PRÄPARAT

PREPARATION CONTENANT DES OLIGOSACCHARIDES ET DES PROBIOTIQUES

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

(30) Priority: **09.12.1998 NL 1010770**

(43) Date of publication of application:
04.10.2001 Bulletin 2001/40

(60) Divisional application:
04076034.0

(73) Proprietor: **N.V. Nutricia**
2700 MA Zoetermeer (NL)

(72) Inventors:
• **DE JONG, Patricia**
NL-6026 AM Maarheeze (NL)
• **VAN LAERE, Katrien**
NL-6666 WS Heteren (NL)

(74) Representative: **van Westenbrugge, Andries et al**
Nederlandsch Octrooibureau
P.O. Box 29720
2502 LS The Hague (NL)

(56) References cited:
EP-A- 0 856 259 EP-A- 0 904 784
WO-A-96/26732 WO-A-97/35596
WO-A-98/26787 US-A- 5 021 245

Description

[0001] The present invention relates to a preparation which has a health-promoting action, in particular for the prevention and/or treatment of disorders of the digestive tract, more particularly of the intestines.

[0002] The application relates in particular to a preparation of this type which contains probiotics and non-digestible oligosaccharides.

[0003] It is known that certain microorganisms have both a prophylactic and a therapeutic effect on intestinal diseases, such as intestinal infections. When these microorganisms are administered to humans or animals they will compete with pathogenic bacteria for nutrients and/or adhesion sites on the intestinal wall, as a result of which the number of pathogenic bacteria will decrease and infections are prevented or reduced. Such microorganisms are generally designated by the term "probiotics".

[0004] If these microorganisms are to have an optimum action they must reach the intestines alive. A further beneficial effect of the administration of live microorganisms to the intestines is, for example, that they are able to ferment the oligosaccharides present in the intestines, whereby, for example, fatty acids with short chains are formed.

[0005] In addition it is advantageous on economic grounds if as many microorganisms as possible reach the intestines alive. With the customary preparations which contain probiotics the percentage of microorganisms that reaches the intestines alive is frequently low.

[0006] It is therefore an object of the present invention to provide a preparation that contains such probiotics with which a high percentage of the microorganisms administered reaches the intestines alive.

[0007] It is also an object of the present invention to provide a preparation that can be used for the treatment of disorders of the intestines and/or that can be used for prophylactic treatment of the intestines.

[0008] The present invention now relates to a preparation having a health-promoting action, in particular for the prevention and/or treatment of disorders of the digestive tract, more particularly of the intestines, which contains probiotics and non-digestible oligosaccharides, wherein the probiotics comprise at least one bacterial strain and at least one yeast strain and wherein the oligosaccharides are trans-galacto-oligo-saccharides or have been obtained by the hydrolysis of one or more polysaccharides chosen from β -(arabino)-galactans, β -(arabino)xylans, β -glucans, β -glucomannans, β -galactomannans, α -arabans and combinations thereof.

[0009] Without wishing to be tied to any theory, it is assumed that the oligosaccharides form a substrate for the probiotics, as a result of which the likelihood that said ~~deletion(s)~~ microorganisms reach the intestines alive increases and as a result of which the likelihood that they are able, in combination with the oligosaccharides present in the intestines, to exert their beneficial action increases. The oligosaccharides could therefore also be designated as "prebiotics".

[0010] The probiotics and non-digestible oligosaccharides are present in the preparation in a ratio of 1 to 5 g oligosaccharides per 10^8 to 10^{11} cells of the probiotic.

[0011] The oligosaccharides used are chosen from the so-called "non-digestible oligosaccharides", that is to say oligosaccharides which are not absorbed by the human or animal body. These oligosaccharides as a rule have a degree of polymerisation of 2 to 20. This implies that the oligosaccharides consist of 2 to 20 monosaccharide units. In the case of a degree of polymerisation of less than 2, that is to say in the case of the monosaccharide, the preparation is not effective since such monosaccharides are absorbed by the human or animal body. At a degree of polymerisation of greater than 20 the oligosaccharides lose their beneficial action.

[0012] Preferably the oligosaccharides have a degree of polymerisation of 2 to 10, more preferentially of 2 to 6, more particularly 3 to 10 and more preferentially 3 to 6.

[0013] Where reference is made in the present application to oligosaccharides this term is used to refer both to oligosaccharides having one specific chain length and to mixtures of oligosaccharides having different chain lengths. However, a mixture of oligosaccharides having different chain lengths is preferred.

[0014] Furthermore, the oligosaccharides usually do not consist entirely of disaccharides. The disaccharide content is usually less than 90 % and sometimes less than 60 %.

[0015] The average degree of polymerisation is as a rule more than 2.1, usually more than 2.5.

[0016] The oligosaccharides used in the invention are, as a rule, furthermore so chosen that they are at least 20 % usable as substrate for the probiotic microorganisms present in the preparation, as determined by high performance anion exchange chromatography.

[0017] Examples of suitable oligosaccharides are transgalacto-oligosaccharides (TOS), fructo-oligosaccharides (FOS) and combinations thereof.

[0018] It is particularly advantageous if the oligosaccharides are added to the preparation in the form of a hydrolysis product of one or more polysaccharides, for example chosen from β -(arabino)galactans, β -(arabino)xylans, β -glucans, β -glucomannans, β -galactomannans, α -arabans, inulin and combinations thereof. In addition to the oligosaccharides, such a hydrolysis product can also contain yet further components, such as monosaccharides and saccharides having a degree of polymerisation higher than 20. However, the hydrolysis product must contain at least 50 % non-digestible

oligosaccharides, preferably at least 70%.

[0019] The polysaccharides which are preferably hydrolysed are β -(arabino)galactans, β -mannans, and xylans.

[0020] It is also possible to use a hydrolysis product of one or more fibres which are mainly made up of the above-mentioned oligosaccharides, such as fibres from oats, wheat, potatoes, sugar beet, soya polysaccharides and the like.

[0021] The hydrolysis of the polysaccharide(s) and/or the fibre(s) can be carried out in a manner known per se, for example by the use of suitable enzymes.

[0022] The concentration of oligosaccharides in the preparation is such that 0.5 to 20 gram per day can be administered. If desired, this administration can be spread over the day, as long as the total quantity of the oligosaccharides administered remains in the range described above. In general the preparation will be administered 2 to 4 times per day.

[0023] In general the oligosaccharides will make up 5 wt.% to 50 wt.% of the total preparation.

[0024] The probiotics suitable for the present invention are generally known. They comprise, in general, one or more bacterial strains suitable for use in food preparations, such as lactic acid bacteria suitable for use in food preparations, or one or more yeast strains suitable for use in food preparations, or a combination thereof. These probiotics will usually have GRAS ("Generally Recognised As Safe") status.

[0025] Suitable bacterial strains are, for example, chosen from those which are described in European Application 97202900.3 (EP 0,904,784) in the name of the Applicant. Other possible probiotics are the *Pediococci*, *Propionibacteria* or *Leuconostoc* species. The *Lactobacillus*, and *Bifidobacterium* genera and combinations thereof are to be preferred.

[0026] The *Bifidobacterium* strain used can be any strain which is suitable for, and preferably is also approved for, administration to humans and animals, such as *Bifidobacterium*, *bifidum*, *Bifidobacterium breve*, *Bifidobacterium lactis* or *Bifidobacterium longum*, or a combination thereof.

[0027] The *Lactobacillus* strain is preferably so chosen that this produces mainly, preferably exclusively, dextrorotatory (L+) lactate. What is meant by this is that the lactate produced is less than 5 %, preferably less than 2 %, laevorotatory lactate. It is, of course, possible that the microorganism produces other metabolites in addition to lactate and the beneficial action of the microorganism can (also) be based on the formation of these further metabolites.

[0028] Examples of these are *Lactobacillus acidophilus*, *Lactobacillus brevis*, *Lactobacillus (para)casei*, *Lactobacillus fermentum*, *Lactobacillus plantarum* and *Lactobacillus rhamnosus*, and combinations thereof.

[0029] The yeast strain used in the invention can be any strain which is suitable for, and preferably is also approved for, administration to humans and animals, such as of the genus *Saccharomyces*. Examples of suitable *Saccharomyces* species are *Saccharomyces cerevisiae* and *Saccharomyces boulardii*.

[0030] The yeast in the preparation can be alive or dead. Both dead and live yeast contains a high content of manoproteins, which are able to prevent the adhesion of bacteria to the intestinal wall to a large extent. Administration of dead yeast offers advantages in the case of people who are suffering from inflammatory intestinal diseases.

[0031] If the yeast is used in dead form, at least one other live microorganism must be used. Said live organism can once again be chosen from both the *Lactobacillus*, *Bifidobacterium*, *Pediococci*, *Propionibacteria* and *Leuconostoc* strains, but can also be live *Saccharomyces*.

[0032] The total concentration of the probiotics is 10^6 to 10^{12} , preferably 10^9 to 10^{10} , live cells per gram of total product. If a combination of microorganisms is used, the minimum concentration of each of the microorganisms must be such that there can still be said to be live organisms, that is to say at least approximately 10 per gram of product. The total concentration of microorganisms must always be within the above specified range of 10^6 to 10^{12} live cells per gram of total product.

[0033] If dead *Saccharomyces cerevisiae* is also used, this is administered in a quantity of 0.5 to 5 g per day.

[0034] The combination of *Lactobacillus rhamnosus* with transgalacto-oligosaccharide or hydrolysis products of (potato) galactan is found to be a particularly suitable combination of oligosaccharide and probiotic.

[0035] The suitability of a specific oligosaccharide for a specific probiotic can be determined by determining the capacity of the microorganism concerned to ferment said oligosaccharide or said oligosaccharide fraction. A specific method for this is given below with the examples.

[0036] The administration forms of the preparations according to the invention are as a rule analogous to those which have been described in European Application 97202900.3 in the name of the Applicant, the contents of which are incorporated herein by reference. In this context it must be pointed out that according to the present invention also only one probiotic microorganism (including a yeast) can be present and that in addition to the one or more probiotic microorganisms one or more oligosaccharides can also be incorporated, in the quantities specified above.

[0037] The preparations according to the present invention can, among others, be administered in the form of a nutritional supplement, total nutrition and clinical nutrition. Reference is likewise made to European Application 97202900:3 (EP 0,904,784) for the specific additives which are added to such foods and the preparation and applications of such foods.

[0038] The probiotics are preferably added to the preparation in (freeze-)dried form. It is also possible to produce liquid preparations, but these must be stored cool. Furthermore, one or more of the microorganisms can be used in encapsulated form, for example in order to improve the shelf life.

[0039] If the preparation according to the invention is used as a nutritional supplement, it can be administered as such, can be mixed with a suitable drinkable liquid, such as water, yoghurt, milk or fruit juice, or can be mixed with solid or liquid food. In this context the nutritional supplement can be in the form of tablets, capsules, powders, sachets, pastilles, sweets, bars and corresponding administration forms, usually in the form of a unit dose.

[0040] A supplement according to the invention can, for example, have the following composition:

- probiotics: 10 - 40 wt. %
- oligosaccharides: 40 - 80 wt. %
- further additives: 0 - 40 wt. %,

to a total of 100 wt. %.

[0041] The preparation can also be in the form of a food preparation that is suitable for direct consumption, such as total or clinical nutrition. This can be either oral nutrition or nutrition for administration via a tube or catheter.

[0042] Such foods can be in solid form or liquid/drinkable form and can contain all customary additives for (total and/or clinical) nutrition, including proteins, vitamins, minerals, trace elements and the like.

[0043] A total nutrition according to the invention can, for example, have the following composition:

- probiotics: 0.1 - 10 wt. %
- oligosaccharides: 1 - 20 wt. %
- further additives: 75 - 95 wt. %

to a total of 100 wt. %.

[0044] According to a particular embodiment, the preparation according to the invention is in the form of a (supplement for a) baby food or a nutritional supplement for babies.

[0045] The preparations described above can be used for the same applications as those described in European Application 97202900.3 (EP 0,904,784) in the name of the Applicant, in particular in the treatment of disorders of the intestines, such as diarrhoea, such as can arise when travelling or after treatment with antibiotics, or which results from food poisoning. Another application is in the treatment of inflammatory bowel diseases (IBD), such as colitis ulcerosa and Crohn's disease. The preparations according to the invention are also suitable for patients who have a food allergy, such as an allergy to cow's milk or to gluten.

[0046] The probiotics and non-digestible oligosaccharides can also be used in baby food to prevent or treat intestinal problems.

[0047] The invention will now be explained with the aid of the following non-limiting examples.

Examples

[0048] To determine suitable combinations of oligosaccharides and probiotics the microorganisms listed below were tested to determine their capacity to ferment structurally different oligosaccharide fractions. Strains pre-cultured in liquid medium based on thioglycolate (Oxoid, Unipath Ltd, Basingstoke, Hampshire, UK) were subjected to subculture in thioglycolate to which 0.5 % (m/V) oligosaccharides were added. The sugar-free thioglycolate medium and the oligosaccharide solutions were sterilised separately for 15 minutes at 121 °C.

[0049] Following anaerobic incubation for 48 hours at 37 °C the pH was measured with the aid of a micro-pH meter (Sentron, Roden, The Netherlands). The changes in the residual oligosaccharide content and the formation of reaction products were determined using HPAEC (high performance anionic exchange chromatography). For HPAEC analysis the cultures were centrifuged and the supernatant liquor was diluted 10-fold with H₂O and boiled for 5 minutes to stop the enzymatic activity. The purity of the strains was checked before and after fermentation.

[0050] The HPAEC system consisted of a Dionex Bio-LC GPM-II quaternary gradient module (Dionex Corporation, Sunnyval, CA, USA) equipped with a Dionex Carbopac PA-100 column (4 * 250 mm) in combination with a Carbopac PA-100 guard column (3 * 25 mm). The samples (20 µl) were injected using a Spectra Physics SP8880 autosampler. The oligomers were analysed using a gradient of sodium acetate in 100 mmol.l⁻¹ NaOH.

[0051] The results of these experiments are given below in the table for a number of combinations of probiotics and substrates, where:

- ++ indicates complete fermentation
- + indicates partial fermentation
- indicates no or very limited fermentation

Bacteria	Substrate			
	Hydrolysis product of:			
	Arabino-galactans	Arabans	Arabinoxylans	Fructans
Bi. Breve	++	+	-	++
Bi. Longum	++	++	+	++
Bi. adolescentis	++	+	++	++
L. acidophilus	++	-	-	+
L. fermentum	++	-	-	+
K. pneumoniae	++	-	-	+
C. perfringens	-	-	-	++

[0052] A few examples of preparations according to the present invention are given below.

Example I: Supplement (Not part of the invention)

[0053] A suspension of *Lactobacillus rhamnosus* ATCC 7469 (Lb) was freeze-dried, a powder being obtained which contained at least 10^9 viable cells per gram powder. Transgalacto-oligosaccharides (TOS), obtained from lactose (Borculo Whey Products), were dissolved in water at 40 °C to a solids content of 25 % and this solution was spraydried. The two powders were mixed in a TOS/Lb ratio of 4/1 until a homogeneous product was obtained. Sachets were filled with 2 - 5 g of this mixture, depending on the dosage regime (5 g for one sachet per day; 3 g for two sachets per day). The contents of one sachet can, for example, be taken mixed in a glass of orange juice or milk.

Example II: Synbiotic bar

[0054] A 23 g bar was prepared from 4.0 g oat flakes, 4.0 g wheat flakes, 3.0 g puffed rice, 1.0 g crushed hazelnuts, 0.25 g honey, 3.0 g raisins, 1.5 g maltodextrin, 1.0 g freeze-dried *Lb rhamnosus*, 0.5 g baker's yeast (*Saccharomyces cerevisiae*; Gist Brocades) and 5.0 g transgalacto-oligosaccharides.

Example III: Method for hydrolysing vegetable fibres

[0055] A 20 % suspension of fibres, for example from wheat, potatoes, oats, soya polysaccharides, carob gum or sugar beet, in water was prepared. These sources of fibre are commercially available. The suspension was heated to a temperature of between 20 and 50 °C (preferably 35 - 45 °C), after which enzymes were added in a quantity of one part enzyme per 5 - 500 parts (m/m) substrate. The choice of the type of enzyme is dependent on the type of polysaccharide. Examples of suitable enzymes are Novoferm Pectinex Ultra s.p.-L, Pentopan and Ultra.s.p. (NOVO).

[0056] After 0.5 - 4 hours the reaction was terminated by heating, after which the solution thus obtained can be used as the oligosaccharide fraction in the preparations according to the invention, optionally after further filtration/purification or after concentrating.

Example IV: Synbiotic mixture for mixing with a complete enteral clinical nutrition (Not part of the invention)

[0057] A mother batch of a powder mixture was prepared in accordance with the method of Example I. The powder consisted of 20 % hydrolysed wheat arabinoxylans, 20 % hydrolysed potato arabinogalactans, 20 % hydrolysed carob gum, 20 % hydrolysed sugar beet fibre (arabans), 15 % hydrolysed oat fibres (glucans) and 5 % *Bifidobacterium longum*. 5 g of the powder mixture is placed in a sachet. The contents of this sachet can be added to a standard enteral clinical nutrition a maximum of 30 min before use.

Example V: Synbiotic powder mixture for fortifying baby food (Not part of the invention)

[0058] A synbiotic mixture was prepared in accordance with the method of Example I. The composition contains 10 % baker's yeast (Gist Brocades), 40 % mannoproteins, obtained from yeast, 25 % inulin and 25 % raffinose.

Example VI: Sweet that contains a synbiotic mixture (Not part of the invention)

[0059] A 2 g sweet was prepared starting from 58 % digestible carbohydrates (glucose syrup), 35 % TOS, 4 % *Lactobacillus rhamnosus* ATCC 7469, and 2 % flavourings and colourants.

Claims

1. Preparation having a health promoting action, in particular for the prevention and/or treatment of disorders of the digestive tract, which contains probiotics comprising at least one bacterial strain and at least one yeast strain and one or more non-digestible oligosaccharides; wherein the oligosaccharides are trans-galacto-oligosaccharides or have been obtained by the hydrolysis of one or more polysaccharides, chosen from β -(arabino)galactans, β -(arabino)xylans, β -glucans, β -glucomannans, β -galactomannans, α -arabans and combinations thereof.
2. Preparation according to Claim 1, wherein the oligosaccharides have a degree of polymerisation of 2 to 20, preferably of 2 to 10, more preferentially of 2 to 6.
3. Preparation according to Claim 1 or 2, wherein the polysaccharides are chosen from β -(arabino)galactans and combinations thereof.
4. Preparation according to Claim 1 or 2, wherein the oligosaccharides originate from the hydrolysis of one or more fibres, such as fibres originating from oats, wheat, potatoes, sugar beet, soya polysaccharides and the like.
5. Preparation according to one of the preceding claims, wherein at least one of the bacterial strains is chosen from one or more strains of a *Lactobacillus* or a *Bifidobacterium* or a *Pediococci* or a *Propionibacteria* or a *Leuconostoc* species and the yeast strain is a strain of a *Saccharomyces* species.
6. Preparation according to one of the preceding claims which also contains dead yeast cells.
7. Preparation according to one of the preceding claims, wherein the ratio between the one or more probiotics and the one or more non-digestible oligosaccharides is in the range of 1 to 5 g oligosaccharides per 10^8 to 10^{11} cells of the probiotic.
8. Preparation according to one of the preceding claims which contains the probiotics in a concentration of 10^7 to 10^{11} live cells per gram of total product.

Patentansprüche

1. Präparat mit einer gesundheitsfördernden Wirkung, insbesondere zur Verhinderung und/oder Behandlung von Störungen des Verdauungstraktes, welches Probiotika enthält, umfassend wenigstens einen Bakterienstamm und wenigstens einen Hefestamm und ein oder mehrere nicht-verdaubare Oligosaccharide, worin die Oligosaccharide Transgalactooligosaccharide sind oder durch die Hydrolyse eines oder mehrerer Polysaccharide erhalten worden sind, ausgewählt aus β -(Arabino)galactanen, β -(Arabino)xylanen, β -Glucanen, β -Gtucomannanen, β -Galactomannanen, α -Arabanen und Kombinationen davon.
2. Präparat gemäß Anspruch 1, worin die Oligosaccharide einen Polymerisationsgrad von 2 bis 20, bevorzugt von 2 bis 10 und bevorzugter von 2 bis 6 haben.
3. Präparat gemäß Anspruch 1 oder 2, worin die Polysaccharide ausgewählt sind aus β -(Arabino)galactanen und Kombinationen davon.
4. Präparat gemäß Anspruch 1 oder 2, worin die Oligosaccharide aus der Hydrolyse von einer oder mehreren Fasern herrühren, wie von Hafer, Weizen, Kartoffeln, Zuckerrübe, Sojapolysacchariden und Ähnlichem, herrührenden Fasern.
5. Präparat gemäß einem der vorhergehenden Ansprüche, worin wenigstens einer der Bakterienstämme ausgewählt ist aus einem oder mehreren Stämmen einer *Lactobacillus*- oder einer *Bifidobacterium*- oder einer *Pediococcus*-

oder einer *Propionibacterium*- oder einer *Leuconostoc*-Art, und der Hefestamm ein Stamm einer *Saccharomyces*-Art ist.

- 5
6. Präparat gemäß einem der vorhergehenden Ansprüche, welches auch tote Hefezellen enthält.
7. Präparat gemäß einem der vorhergehenden Ansprüche, worin das Verhältnis zwischen dem einen Probiotikum oder mehreren Probiotika und dem einen nicht-verdaubaren Oligosaccharid oder mehreren nicht-verdaubaren Oligosacchariden in dem Bereich von 1 bis 5 g Oligosacchariden pro 10^8 bis 10^{11} Zellen des Probiotikums liegt.
- 10
8. Präparat gemäß einem der vorhergehenden Ansprüche, welches die Probiotika in einer Konzentration von 10^7 bis 10^{11} lebenden Zellen pro Gramm des Gesamtproduktes enthält.

Revendications

- 15
1. Préparation ayant une action favorisant la santé, en particulier pour la prévention et/ou le traitement de troubles des voies digestives, laquelle contient des probiotiques comprenant au moins une souche bactérienne et au moins une souche de levure et un ou plusieurs oligosaccharides non digestibles, dans laquelle les oligosaccharides sont des trans-galacto-oligosaccharides ou ont été obtenus par l'hydrolyse d'un ou de plusieurs polysaccharides, choisis
- 20
- parmi les β -(arabino)galactanes, les β -(arabino)xylanes, les β -glucanes, les β -glucomannanes, les β -galactomannanes, les α -arabanes et des combinaisons de ceux-ci.
2. Préparation selon la revendication 1, dans laquelle les oligosaccharides ont un degré de polymérisation de 2 à 20, de préférence de 2 à 10, de manière encore préférable de 2 à 6.
- 25
3. Préparation selon la revendication 1 ou 2, dans laquelle les polysaccharides sont choisis parmi les β -(arabino)galactanes et des combinaisons de celles-ci.
4. Préparation selon la revendication 1 ou 2, dans laquelle les oligosaccharides ont pour origine l'hydrolyse d'une ou de plusieurs fibres, telles que des fibres provenant de l'avoine, du blé, de la pomme de terre, de la betterave à sucre, des polysaccharides du soja et des fibres similaires.
- 30
5. Préparation selon l'une des revendications précédentes, dans laquelle au moins une des souches bactériennes est choisie parmi une ou plusieurs souches d'une espèce de *Lactobacillus*, de *Bifidobacterium*, de *Pediococci*, de *Propionibacteria* ou de *Leuconostoc* et la souche de levure est une souche d'une espèce de *Saccharomyces*.
- 35
6. Préparation selon l'une des revendications précédentes qui contient également des cellules de levure mortes.
7. Préparation selon l'une des revendications précédentes, dans laquelle le rapport entre le ou les probiotique(s) et le ou les oligosaccharide(s) non digestible(s) est de 1 à 5 g d'oligosaccharides par 10^8 à 10^{11} cellules du probiotique.
- 40
8. Préparation selon l'une des revendications précédentes qui contient les probiotiques à une concentration de 10^7 à 10^{11} cellules vivantes par gramme de produit total.
- 45
- 50
- 55