

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 195 168 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

10.04.2002 Bulletin 2002/15

(51) Int Cl.7: **A61K 51/08**

// A61K123:00, A61K103:10

(21) Application number: **01123928.2**

(22) Date of filing: **03.04.1996**

(84) Designated Contracting States:

**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE**

(30) Priority: **03.04.1995 US 415908**

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:

96914548.1 / 0 820 312

(71) Applicant: **DuPont Pharmaceuticals Company
Wilmington, DE 19805 (US)**

(72) Inventors:

- **Edwards, David S.
Burlington, MA 01803 (US)**

• **Liu, Shuang**

Chelmsford, MA 01824 (US)

(74) Representative:

**Helbing, Jörg, Dr. Dipl.-Chem. et al
Patentanwälte
von Kreisler-Selting-Werner,
Postfach 10 22 41
50462 Köln (DE)**

Remarks:

This application was filed on 06 - 10 - 2001 as a
divisional application to the application mentioned
under INID code 62.

(54) **Ternary radiopharmaceutical complexes**

(57) This invention provides novel radiopharmaceuticals which are useful as imaging agents for the diagnosis of cardiovascular disorders, infectious diseases and cancer. The radiopharmaceuticals are comprised of phosphine or arsine ligated technetium-99m labelled hydrazine or diazino modified biologically active molecules that selectively localize at sites of disease and thus allow an image to be obtained of the loci using gamma

scintigraphy. This invention also provides methods for using the radiopharmaceuticals and kits comprising radiopharmaceutical precursors. The radiopharmaceuticals of this invention have the structure: $[(Q)_d(L_n-C_h)]_x-M_t(A_{L1})_y(A_{L2})_z$ wherein the variables are as defined therein.

EP 1 195 168 A2

DescriptionCROSS-REFERENCE TO RELATED APPLICATIONS

5 **[0001]** The present application is a continuation-in-part of our copending application U.S.S.N. 08/218,861 which is a continuation-in-part of U.S.S.N. 08/040,336 filed March 30, 1993, the disclosures of which are incorporated herein by reference.

FIELD OF THE INVENTION

10 **[0002]** This invention relates to novel radiopharmaceuticals which are useful as imaging agents for the diagnosis of cardiovascular disorders, infectious disease and cancer, and to kits useful for their preparation. The radiopharmaceuticals are comprised of phosphine or arsine ligated technetium-99m labeled hydrazino or diazino modified biologically active molecules that selectively localize at sites of disease and thus allow an image to be obtained of the loci using
15 gamma scintigraphy.

BACKGROUND OF THE INVENTION

20 **[0003]** There is a current need for new methods for the non-invasive diagnosis of a variety of diseases such as thromboembolic disease, atherosclerosis, infection and cancer. Radiopharmaceuticals comprised of gamma-ray emitting radionuclide labeled biologically active molecules can fulfill the need. The biologically active molecules serve to localize the radionuclides at the sites of disease and thus allow the sites to be visualized by gamma scintigraphy. The molecules can be either proteins, antibodies, antibody fragments, peptides or polypeptides, or peptidomimetics. The molecules interact with a receptor or binding site expressed at the sites of the disease or with a receptor or binding
25 site on an endogenous blood component, such as platelets and leukocytes, that accumulate at the sites. This interaction results in selective localization of a percentage of the injected radiopharmaceutical while the remainder is cleared either through the renal or hepatobiliary systems. The localized radiopharmaceutical is then imaged externally using gamma scintigraphy. The relative rates of sequestration, clearance and radionuclidic decay determine the ease of visualization, often expressed as the target-to-background ratio. Frequently, only certain portions of the biologically active molecules
30 bind to the receptors; these portions are termed the recognition sequences or units.

[0004] A number of radiopharmaceuticals comprised of radionuclide labeled proteins, antibodies or antibody fragments are under development, however, to date only one has been approved by the Food and Drug Administration. This sparse record results from a combination of factors that make developing these radiopharmaceuticals difficult, including problems with manufacturing and quality control, non-optimal sequestration and clearance rates, and the
35 occurrence of antigenic or allergic responses to the radiopharmaceuticals. These problems are mainly due to the macromolecular nature of the proteins, antibodies and antibody fragments. Their high molecular weight makes direct chemical synthesis impractical, therefore they must be synthesized by recombinant or cloning techniques that typically give low yields and require extensive isolation and purification procedures. Their molecular weight can slow their rates of localization and preclude their clearance by an active elimination mechanism via the kidneys or liver, resulting in prolonged retention in the circulation which causes a high background level during imaging. Also, the body's immune
40 system tends to recognize more efficiently larger exogenous species.

[0005] The use of lower molecular weight peptides, polypeptides or peptidomimetics as the biologically active molecules obviates a number of these problems. These molecules can be synthesized directly using classical solution chemistry or by an automated peptide synthesizer. They can be formed in higher yields and require less complicated
45 purification procedures. They tend to clear more rapidly from the circulation by an active elimination pathway resulting in a lower background in the images. They are also usually not immunogenic. The first radionuclide labeled polypeptide radiopharmaceutical has been recently approved by the Food and Drug Administration.

[0006] There are two general methods for labeling biologically active molecules with radionuclides for use as radiopharmaceuticals termed direct and indirect labeling. Direct labeling involves attaching the radionuclide to atoms on
50 the biologically active molecule; while the indirect method involves attaching the radionuclide via a chelator. The chelator can either be attached to the biologically active molecule prior to reaction with the radionuclide or the radionuclide labeled chelator moiety can be attached to the biologically active molecule. Several recent reviews describe these labeling methods and are incorporated herein by reference: S. Jurisson et. al., Chem. Rev., 1993, 93, 1137; A. Verbruggen, Eur. J. Nuc. Med., 1990, 17, 346; and M. Derwanjee, Semin. Nuc. Med., 1990, 20, 5.

55 **[0007]** The use of hydrazines and hydrazides as chelators to modify proteins for labeling with radionuclides has been recently disclosed in Schwartz et. al., U.S. Patent 5,206,370. For labeling with technetium-99m, the hydrazino-modified protein is reacted with a reduced technetium species, formed by reacting pertechnetate with a reducing agent in the presence of a chelating dioxygen ligand. The technetium becomes bound to the protein through what are believed to

be hydrazido or diazenido linkages with the coordination sphere completed by the ancillary dioxygen ligands. Examples of ancillary dioxygen ligands include glucoheptonate, gluconate, 2-hydroxyisobutyrate, and lactate.

[0008] Certain dioxygen ligands have been recently reported to be particularly advantageous for labeling hydrazino-modified proteins with technetium-99m. Bridger et. al., European Patent Application 93302712.0, disclose a series of functionalized aminocarboxylates the use of which are reported to improve the labeling process of hydrazino-modified macromolecules such as monoclonal antibodies. The improvements are manifest by shorter reaction times and higher specific activities. Examples of these improved dioxygen ligands include hydroxyalkyl substituted glycine derivatives such as tricine.

[0009] In co-pending U.S. Ser. No. 08/218,861 (equivalent to WO 94/22494), filed March 28, 1993, the synthesis of novel radiolabeled platelet IIb/IIIa receptor antagonists as imaging agents for thromboembolic disorders is disclosed. These reagents comprise radionuclide labeled chelator modified cyclic compounds. A preferred chelator for modifying the cyclic compounds is the hydrazino or diazenido moiety.

[0010] The present invention provides novel technetium-99m labeled hydrazino or diazino modified biologically active molecules that are formed as a minimal number of isomers, the relative ratios of which do not change with time. These compounds are more straightforward to develop, requiring less complicated manufacturing and labeling process controls.

SUMMARY OF THE INVENTION

[0011] This invention provides novel radiopharmaceuticals which are useful as imaging agents for the diagnosis of cardiovascular disorders, such as thromboembolic disease or atherosclerosis, infectious disease and cancer. The radiopharmaceuticals are comprised of phosphine or arsine ligated technetium-99m labeled hydrazino or diazenido modified biologically active molecules that selectively localize at sites of disease and thus allow an image to be obtained of the loci using gamma scintigraphy. The invention also provides methods of using said radiopharmaceuticals as imaging agents for the diagnosis of cardiovascular disorders, such as thromboembolic disease or atherosclerosis, infectious disease and cancer. It further provides kits for the preparation of said radiopharmaceuticals.

Brief Description of the Figures

[0012] Figure 1. HPLC chromatograms, using both Methods 1 and 2, of the final product obtained in Example 1 of the present invention.

[0013] Figure 2. Data from the Canine Deep Vein Thrombosis model for the radiopharmaceuticals of Examples 1 and 2 of the present invention and Tc-albumin (negative control); thrombus to blood and thrombus to muscle ratios.

[0014] Figure 3. Blood clearance curves from the Arteriovenous Shunt model for radiopharmaceuticals of Examples 1 and 2 of the present invention and Tc-albumin (negative control).

DETAILED DESCRIPTION OF THE INVENTION

[0015] The present invention is directed to novel radiopharmaceuticals for the diagnosis of cardiovascular disorders, such as thromboembolic disease and atherosclerosis, infectious disease or cancer of the formula, methods of using said radiopharmaceuticals in the diagnosis of diseases and kits useful for the preparation of said radiopharmaceutical.

[1] One embodiment of the present prevention is a radiopharmaceutical comprising a transition metal radionuclide, a transition metal chelator, a biologically active group connected to said chelator, a first ancillary ligand, a second ancillary ligand capable of stabilizing the radiopharmaceutical, optionally having a linking group between said chelator and said biologically active group.

[2] Another embodiment of the present invention is a radiopharmaceutical of embodiment [1] having a linking group between said chelator and said biologically active group.

[3] Another embodiment of the present invention is a radiopharmaceutical of embodiment [2] of formula:




wherein:

EP 1 195 168 A2

Q is a biologically active molecule;

d' is 1 to 20;

5 L_n is a linking group of formula:



10 wherein:

M¹ is -[(CH₂)_gZ¹]_{g'}-(CR⁵⁵R⁵⁶)_{g''};

15 M² is -(CR⁵⁵R⁵⁶)_{g'}-[Z¹(CH₂)_g]_{g''};

g is independently 0-10;

g' is independently 0-1;

20 g'' is independently 0-10;

f is independently 0-10;

25 f' is independently 0-10;

f'' is independently 0-1;

30 Y¹ and Y², at each occurrence, are independently selected from:
a bond, O, NR⁵⁶, C=O, C(=O)O, OC(=O)O, C(=O)NH-, C=NR⁵⁶, S, SO,
SO₂, SO₃, NHC(=O), (NH)₂C(=O), (NH)₂C=S;

35 Z¹ is independently selected at each occurrence from a C₆-C₁₄ saturated, partially saturated, or aromatic carbocyclic ring system, substituted with 0-4 R⁵⁷; and a heterocyclic ring system, optionally substituted with 0-4 R⁵⁷;

R⁵⁵ and R⁵⁶ are independently selected at each occurrence from:
hydrogen; C₁-C₁₀ alkyl substituted with 0-5 R⁵⁷; alkaryl wherein the aryl is substituted with 0-5 R⁵⁷;

40 R⁵⁷ is independently selected at each occurrence from the group: hydrogen, OH, NHR⁵⁸, C(=O)R⁵⁸, OC(=O)R⁵⁸, OC(=O)OR⁵⁸, C(=O)OR⁵⁸, C(=O)NR⁵⁸-, C=N, SR⁵⁸, SOR⁵⁸, SO₂R⁵⁸, NHC(=O)R⁵⁸, NHC(=O)NHR⁵⁸, NHC(=S)NHR⁵⁸; or, alternatively, when attached to an additional molecule Q, R⁵⁷ is independently selected at each occurrence from the group: O, NR⁵⁸, C=O, C(=O)O, OC(=O)O, C(=O)N-, C=NR⁵⁸, S, SO, SO₂, SO₃, NHC(=O), (NH)₂C(=O), (NH)₂C=S; and,

45 R⁵⁸ is independently selected at each occurrence from the group: hydrogen; C₁-C₆ alkyl; benzyl, and phenyl;

50 x and y are independently 1 or 2;

z is independently 1-4;

55 M_t is a transition metal radionuclide selected from the group: ^{99m}Tc, ¹⁸⁶Re and ¹⁸⁸Re;

C_h is a radionuclide metal chelator coordinated to transition metal radionuclide M_t, and is independently selected at each occurrence, from the group: R⁴⁰N=N⁺=, R⁴⁰R⁴¹N-N=, R⁴⁰N=,

EP 1 195 168 A2

and $R^{40}N=N(H)-$, wherein

R^{40} is independently selected at each occurrence from the group: a bond to L_n , C_1 - C_{10} alkyl substituted with 0-3 R^{52} , aryl substituted with 0-3 R^{52} , cycloalkyl substituted with 0-3 R^{52} , heterocycle substituted with 0-3 R^{52} , heterocycloalkyl substituted with 0-3 R^{52} , aralkyl substituted with 0-3 R^{52} and alkaryl substituted with 0-3 R^{52} ;

R^{41} is independently selected from the group: hydrogen, aryl substituted with 0-3 R^{52} , C_1 - C_{10} alkyl substituted with 0-3 R^{52} , and a heterocycle substituted with 0-3 R^{52} ;

R^{52} is independently selected at each occurrence from the group: a bond to L_n , $=O$, F , Cl , Br , I , $-CF_3$, $-CN$, $-CO_2R^{53}$, $-C(=O)R^{53}$, $-C(=O)N(R^{53})_2$, $-CHO$, $-CH_2OR^{53}$, $-OC(=O)R^{53}$, $-OC(=O)OR^{53a}$, $-OR^{53}$, $-OC(=O)N(R^{53})_2$, $-NR^{53}C(=O)R^{53}$, $-NR^{54}C(=O)OR^{53a}$, $-NR^{53}C(=O)N(R^{53})_2$, $-NR^{54}SO_2N(R^{53})_2$, $-NR^{54}SO_2R^{53a}$, $-SO_3H$, $-SO_2R^{53a}$, $-SR^{53}$, $-S(=O)R^{53a}$, $-SO_2N(R^{53})_2$, $-N(R^{53})_2$, $-NHC(=NH)NHR^{53}$, $-C(=NH)NHR^{53}$, $=NOR^{53}$, NO_2 , $-C(=O)NHR^{53}$, $-C(=O)NHN(R^{53})_2$, $-OCH_2CO_2H$, 2-(1-morpholino)ethoxy;

R^{53} , R^{53a} , and R^{54} are each independently selected at each occurrence from the group: hydrogen, C_1 - C_6 alkyl, and a bond to L_n ;

A_{L1} is a first ancillary ligand selected from the group:
dioxxygen ligand,
functionalized aminocarboxylate, and
halide;

A_{L2} is an ancillary ligand capable of stabilizing the radiopharmaceutical selected from the group:
 A^9 and A^{10} - W - A^{11} ,

wherein:

A^9 is independently selected at each occurrence from the group: $PR^{61}R^{62}R^{63}$ and $AsR^{61}R^{62}R^{63}$;

A^{10} and A^{11} are independently selected at each occurrence from the group: $PR^{61}R^{62}$ and $AsR^{61}R^{62}$;

W is a spacer group selected from the group: C_1 - C_{10} alkyl substituted with 0-3 R^{70} , aryl substituted with 0-3 R^{70} , cycloalkyl substituted with 0-3 R^{70} , heterocycle substituted with 0-3 R^{70} , heterocycloalkyl substituted with 0-3 R^{70} , aralkyl substituted with 0-3 R^{70} and alkaryl substituted with 0-3 R^{70} ;

R^{61} , R^{62} , and R^{63} are independently selected at each occurrence from the group: C_1 - C_{10} alkyl substituted with 0-3 R^{70} , aryl substituted with 0-3 R^{70} , cycloalkyl substituted with 0-3 R^{70} , heterocycle substituted with 0-3 R^{70} , aralkyl substituted with 0-3 R^{70} , alkaryl substituted with 0-3 R^{70} , and arylalkaryl substituted with 0-3 R^{70} ;

R^{70} is independently selected at each occurrence from the group: F , Cl , Br , I , $-CF_3$, $-CN$, $-CO_2R^{71}$, $-C(=O)R^{71}$, $-C(=O)N(R^{71})_2$, $-CH_2OR^{71}$, $-OC(=O)R^{71}$, $-OC(=O)OR^{71a}$, $-OR^{71}$, $-OC(=O)N(R^{71})_2$, $-NR^{71}C(=O)R^{71}$, $-NR^{71}C(=O)OR^{71}$, $-NR^{71}C(=O)N(R^{71})_2$, SO_3^- , $-NR^{71}SO_2N(R^{71})_2$,

EP 1 195 168 A2

-NR⁷¹SO₂R^{71a}, -SO₃H, -SO₂R⁷¹, -S(=O)R⁷¹, -SO₂N(R⁷¹)₂, -N(R⁷¹)₂, -N(R⁷¹)₃⁺, -NHC(=NH)NHR⁷¹, -C(=NH)NHR⁷¹, =NOR⁷¹, NO₂, -C(=O)NHOR⁷¹, -C(=O)NHN(R⁷¹)₂, -OCH₂CO₂H; and

5 R⁷¹ and R^{71a} are independently selected at each occurrence from the group: hydrogen and C₁-C₆ alkyl; and

pharmaceutically acceptable salts thereof.

10 [4] Another embodiment of the present invention is a radiopharmaceutical of embodiment [3] wherein:

Q is a biologically active molecule selected from the group: IIb/IIIa receptor antagonists, IIb/IIIa receptor ligands, fibrin binding peptides, leukocyte binding peptides, chemotactic peptides, somatostatin analogs, and selectin binding peptides;

15 d' is 1 to 3;

L_n is:

20
$$-(CR^{55}R^{56})_{g''}-[Y^1(CR^{55}R^{56})_fY^2]_{f'}-(CR^{55}R^{56})_{g''},$$

wherein:

25 g'' is 0-5;

f is 0-5;

f' is 1-5;

Y¹ and Y², at each occurrence, are independently selected from:

30 O, NR⁵⁶, C=O, C(=O)O, OC(=O)O, C(=O)NH-, C=NR⁵⁶, S, SO, SO₂, SO₃, NHC(=O), (NH)₂C(=O), (NH)₂C=S;

R⁵⁵ and R⁵⁶ are independently selected at each occurrence from: hydrogen, C₁-C₁₀ alkyl, and alkaryl;

35 x and y are independently 1 or 2;

z is independently 1-2;

M_t is ^{99m}Tc;

40 C_h is a radionuclide metal chelator coordinated to transition metal radionuclide M_t, and is independently selected at each occurrence, from the group: R⁴⁰N=N⁺=, R⁴⁰R⁴¹N=N-, R⁴⁰N=, and R⁴⁰N=N(H)-;

45 R⁴⁰ is independently selected at each occurrence from the group: aryl substituted with 0-3 R⁵², and heterocycle substituted with 0-3 R⁵²;

R⁴¹ is independently selected from the group: hydrogen, aryl substituted with 0-1 R⁵², C₁-C₃ alkyl substituted with 0-1 R⁵², and a heterocycle substituted with 0-1 R⁵²;

50 R⁵² is independently selected at each occurrence from the group: a bond to L_n, -CO₂R⁵³, -CH₂OR⁵³, -SO₃H, -SO₂R^{53a}, -N(R⁵³)₂, -N(R⁵³)₃⁺, -NHC(=NH)NHR⁵³, and -OCH₂CO₂H;

55 R⁵³, R^{53a} are each independently selected at each occurrence from the group: hydrogen and C₁-C₃ alkyl;

A_{L1} is selected from the group:

pyrones, pyridinones, and
functionalized aminocarboxylates;

A_{L2} is selected from the group:

A^9 and A^{10} -W- A^{11} ,

wherein:

A^9 is $PR^{61}R^{62}R^{63}$;

A^{10} and A^{11} are $PR^{61}R^{62}$;

W is a spacer group selected from the group: C_1 - C_3 alkyl substituted with 0-3 R^{70} , aryl substituted with 0-3 R^{70} , and heterocycle substituted with 0-3 R^{70} ; R^{61} , R^{62} , and R^{63} are independently selected at each occurrence from the group: C_1 - C_3 alkyl substituted with 0-3 R^{70} , aryl substituted with 0-3 R^{70} , and heterocycle substituted with 0-3 R^{70} ;

R^{70} is independently selected at each occurrence from the group: $-CO_2R^{71}$, $-OR^{71}$, $-SO_3^-$ and $-SO_3H$; and


R^{71} is hydrogen.

[5] Another embodiment of the present invention is a radiopharmaceutical of embodiment [4] wherein:

Q represents a biologically active molecule selected from the group: IIb/IIIa receptor antagonists and chemotactic peptides;

d' is 1;

L_n is:



wherein:

g'' is 0-5;

f is 0-5;

f' is 1-5;

Y^1 and Y^2 , at each occurrence, are independently selected from:
O, NR^{56} , $C=O$, $C(=O)O$, $OC(=O)O$, $C(=O)NH-$, $C=NR^{56}$, S, $NHC(=O)$, $(NH)_2C(=O)$, $(NH)_2C=S$;

R^{55} and R^{56} are hydrogen;

x and y are 1;

z is 1;

$C_{h'}$ is a radionuclide metal chelator coordinated to transition metal radionuclide Mt, and is independently selected at each occurrence, from the group: $R^{40}N=N^+=$, and $R^{40}R^{41}N-N=$;

R^{40} is independently selected at each occurrence from the group: heterocycle substituted with R^{52} ;

R^{41} is hydrogen;

R^{52} is a bond to L_n ;

A_{L1} is tricline;


A_{L2} is $PR^{61}R^{62}R^{63}$, wherein

R⁶¹, R⁶², and R⁶³ are independently selected at each occurrence from the group: C₁-C₃ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰;

R⁷⁰ is independently selected at each occurrence from the group: -CO₂H, -OH, -SO₃H, -SO₃⁻.



[5] Another embodiment of the present invention is the radiopharmaceutical of embodiment [3] wherein:

Q is



d' is 1;

L_n is attached to Q at the carbon atom designated with a * and has the formula:

 C_h is

and is attached to Ln at the carbon atom designated with a *;

M_t is ^{99m}Tc ;


A_{L1} is tricine;

A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹, R⁶² and R⁶³ are each phenyl bearing an SO₃H or SO₃⁻ group in the meta position; and

x, y and z are 1.


[7] Another embodiment of the present invention is the radiopharmaceutical of embodiment [3] wherein:

Q is




d' is 1;

L_n is attached to Q at the carbon atom designated with a * and has the formula:



$C_{h'}$ is



and is attached to L_n at the carbon atom designated with a *;

M_t is ^{99m}Tc ;


A_{L1} is tricine;

A_{L2} is $\text{PR}^{61}\text{R}^{62}\text{R}^{63}$, wherein R^{61} is phenyl, R^{62} and R^{63} are each phenyl bearing an SO_3H or SO_3^- group in the meta position; and

x, y and z are 1.


[8] Another embodiment of the present invention is the radiopharmaceutical of embodiment [3] wherein:

Q is




d' is 1;

L_n is attached to Q at the carbon atom designated with a * and has the formula:



C_{n'} is



and is attached to L_n at the carbon atom designated with a *;

M_t is ^{99m}Tc;


A_{L1} is tricine;

A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹ and R⁶² are phenyl, and R⁶³ is phenyl bearing an SO₃H or SO₃⁻ group in the meta position; and

x, y and z are 1.


[9] Another embodiment of the present invention is the radiopharmaceutical of embodiment [3] wherein:

Q is




d' is 1;

L_n is attached to Q at the carbon atom designated with a * and has the formula:



C_{h'} is



and is attached to L_n at the carbon atom designated with a *;

M_t is ^{99m}Tc;


A_{L1} is tricine;

A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹, R⁶² and R⁶³ are each p-(2-phenylethyl)phenyl wherein the phenylethyl bears an SO₃H or SO₃⁻ group in the para position; and

x, y and z are 1.


embodiment [10] Another embodiment of the present invention is the radiopharmaceutical of embodiment [3] wherein:

Q is




d' is 1;

L_n is attached to Q at the carbon atom designated with a * and has the formula:



$C_{h'}$ is



and is attached to L_n at the carbon atom designated with a *;

M_t is ^{99m}Tc ;


A_{L1} is tricine;

A_{L2} is $\text{PR}^{61}\text{R}^{62}\text{R}^{63}$, wherein R^{61} , R^{62} and R^{63} are each p-(2-phenylpropyl)phenyl wherein the phenylpropyl bears an SO_3H or SO_3^- group in the para position; and

x, y and z are 1.


[11] Another embodiment of the present invention is the radiopharmaceutical of embodiment [3] wherein:

Q is




d' is 1;

L_n is attached to Q at the carbon atom designated with a * and has the formula:



C_{h'} is



and is attached to L_n at the carbon atom designated with a *;

M_t is ^{99m}Tc;


A_{L1} is tricine;

A_{L2} is R⁶¹R⁶²PCH₂CH₂PR⁶¹R⁶², wherein R⁶¹, R⁶² are each phenyl substituted with an SO₃H or SO₃⁻ group in the meta position; and

x, y and z are 1.


[12] Another embodiment of the present invention is the radiopharmaceutical of embodiment [3] wherein:

Q is




d' is 1;

L_n is attached to Q at the carbon atom designated with a * and has the formula:



$C_{h'}$ is



and is attached to L_n at the carbon atom designated with a *;

M_t is ^{99m}Tc ;


A_{L1} is tricine;

A_{L2} is $\text{PR}^{61}\text{R}^{62}\text{R}^{63}$, wherein R^{61} , R^{62} and R^{63} are C_3 -alkyl substituted with 1 OH group; and

x, y and z are 1.


[13] Another embodiment of the present invention is the radiopharmaceutical of embodiment [3] wherein:

Q is




d' is 1;

L_n is attached to Q at the carbon atom designated with a * and has the formula:



C_{h'} is



and is attached to L_n at the carbon atom designated with a *;

M_t is ^{99m}Tc;


A_{L1} is tricine;

A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹, R⁶² and R⁶³ are CH₂CH₂COOH; and

x, y and z are 1.


[14] Another embodiment of the present invention is the radiopharmaceutical of embodiment [3] wherein:

Q is




d' is 1;

L_n is attached to Q at the carbon atom designated with a * and has the formula:



C_n is



and is attached to L_n at the carbon atom designated with a *;

M_t is ^{99m}Tc ;

A_{L1} is kojic acid;

A_{L2} is $\text{PR}^{61}\text{R}^{62}\text{R}^{63}$, wherein R^{61} , R^{62} and R^{63} are each phenyl bearing an SO_3H or SO_3^- group in the meta position;

x and z are 1; and

y is 2.

[15] Another embodiment of the present invention is a method for radioimaging a mammal comprising (i) administering to said mammal an effective amount of a radiopharmaceutical of any of embodiments [1]-[14], and (ii) scanning the mammal using a radioimaging device.

[16] Another embodiment of the present invention is a method for visualizing sites of platelet deposition in a mammal by radioimaging, comprising (i) administering to said mammal an effective amount of a radiopharmaceutical of any of embodiments [6]-[14], and (ii) scanning the mammal using a radioimaging device.


[17] Another embodiment of the present invention is a method of determining platelet deposition in a mammal comprising administering to said mammal a radiopharmaceutical composition of any of embodiments [6]-[14], and

imaging said mammal.

[18] Another embodiment of the present invention is a method of diagnosing a disorder associated with platelet deposition in a mammal comprising administering to said mammal a radiopharmaceutical composition of any of
5
embodiments [6]-[14], and imaging said mammal.

[19] Another embodiment of the present invention is a kit for preparing a radiopharmaceutical comprising:

(a) a predetermined quantity of a sterile, pharmaceutically acceptable reagent of formulae:



(b) a predetermined quantity of a sterile, pharmaceutically acceptable first ancillary ligand, A_{L1} , selected from
15
the group:

dioxygen ligand,
functionalized aminocarboxylate, and
halide; and

(c) a predetermined quantity of a sterile, pharmaceutically acceptable second ancillary ligand, A_{L2} , selected
20
from the group:


A^9 and A^{10} -W- A^{11} ;

wherein:

Q is a biologically active molecule;

d' is 1 to 20;

L_n is a linking group of formula:



wherein:

M^1 is $-(CH_2)_gZ^1]_g-(CR^{55}R^{56})_{g''}$;

M^2 is $-(CR^{55}R^{56})_{g''}-[Z^1(CH_2)_g]_g$;

g is independently 0-10;

g' is independently 0-1;

g'' is independently 0-10;

f is independently 0-10;

f' is independently 0-10;

f'' is independently 0-1;

Y^1 and Y^2 , at each occurrence, are independently selected from:
a bond, O, NR^{56} , C=O, C(=O)O, OC(=O)O, C(=O)NH-, C=NR⁵⁶, S, SO,
SO₂, SO₃, NHC(=O), (NH)₂C(=O), (NH)₂C=S;

EP 1 195 168 A2

	Z ¹	is independently selected at each occurrence from a C ₆ -C ₁₄ saturated, partially saturated, or aromatic carbocyclic ring system, substituted with 0-4 R ⁵⁷ ; and a heterocyclic ring system, optionally substituted with 0-4 R ⁵⁷ ;
5	R ⁵⁵ and R ⁵⁶	are independently selected at each occurrence from: hydrogen; C ₁ -C ₁₀ alkyl substituted with 0-5 R ⁵⁷ ; alkaryl wherein the aryl is substituted with 0-5 R ⁵⁷ ;
10	R ⁵⁷	is independently selected at each occurrence from the group: hydrogen, OH, NHR ⁵⁸ , C(=O)R ⁵⁸ , OC(=O)R ⁵⁸ , OC(=O)OR ⁵⁸ , C(=O)OR ⁵⁸ , C(=O)NR ⁵⁸ -, C≡N, SR ⁵⁸ , SOR ⁵⁸ , SO ₂ R ⁵⁸ , NHC(=O)R ⁵⁸ , NHC(=O)NHR ⁵⁸ , NHC(=S)NHR ⁵⁸ ; or, alternatively, when attached to an additional molecule Q, R ⁵⁷ is independently selected at each occurrence from the group: O, NR ⁵⁸ , C=O, C(=O)O, OC(=O)O, C(=O)N-, C=NR ⁵⁸ , S, SO, SO ₂ , SO ₃ , NHC(=O), (NH) ₂ C(=O), (NH) ₂ C=S; and,
15	R ⁵⁸	is independently selected at each occurrence from the group: hydrogen; C ₁ -C ₆ alkyl; benzyl, and phenyl;
20	C _h	is a radionuclide metal chelator independently selected at each occurrence from the group: R ⁴⁰ R ⁴¹ N-N=C(C ₁ -C ₃ alkyl) ₂ and R ⁴⁰ NNH ₂ -, wherein::
25	R ⁴⁰	is independently selected at each occurrence from the group: a bond to L _n , C ₁ -C ₁₀ alkyl substituted with 0-3 R ⁵² , aryl substituted with 0-3 R ⁵² , cycloalkyl substituted with 0-3 R ⁵² , heterocycle substituted with 0-3 R ⁵² , heterocycloalkyl substituted with 0-3 R ⁵² , aralkyl substituted with 0-3 R ⁵² and alkaryl substituted with 0-3 R ⁵² ;
30	R ⁴¹	is independently selected from the group: hydrogen, aryl substituted with 0-3 R ⁵² , C ₁ -C ₁₀ alkyl substituted with 0-3 R ⁵² , and a heterocycle substituted with 0-3 R ⁵² ;
35	R ⁵²	is independently selected at each occurrence from the group: a bond to L _n , =O, F, Cl, Br, I, -CF ₃ , -CN, -CO ₂ R ⁵³ , -C(=O)R ⁵³ , -C(=O)N(R ⁵³) ₂ , -CHO, -CH ₂ OR ⁵³ , -OC(=O)R ⁵³ , -OC(=O)OR ^{53a} , -OR ⁵³ , -OC(=O)N(R ⁵³) ₂ , -NR ⁵³ C(=O)R ⁵³ , -NR ⁵⁴ C(=O)OR ^{53a} , -NR ⁵³ C(=O)N(R ⁵³) ₂ , -NR ⁵⁴ SO ₂ N(R ⁵³) ₂ , -NR ⁵⁴ SO ₂ R ^{53a} , -SO ₃ H, -SO ₂ R ^{53a} , -SR ⁵³ , -S(=O)R ^{53a} , -SO ₂ N(R ⁵³) ₂ , -N(R ⁵³) ₂ , -NHC(=NH)NHR ⁵³ , -C(=NH)NHR ⁵³ , =NOR ⁵³ , NO ₂ , -C(=O)NHOR ⁵³ , -C(=O)NHN(R ⁵³)R ^{53a} , -OCH ₂ CO ₂ H, 2-(1-morpholino)ethoxy;
40	R ⁵³ , R ^{53a} , and R ⁵⁴	are each independently selected at each occurrence from the group: hydrogen, C ₁ -C ₆ alkyl, and a bond to L _n ;
45	A ⁹	is independently selected at each occurrence from the group: PR ⁶¹ R ⁶² R ⁶³ and AsR ⁶¹ R ⁶² R ⁶³ ;
	A ¹⁰ and A ¹¹	are independently selected at each occurrence from the group: PR ⁶¹ R ⁶² and AsR ⁶¹ R ⁶² ;
50	W	is a spacer group selected from the group: C ₁ -C ₁₀ alkyl substituted with 0-3 R ⁷⁰ , aryl substituted with 0-3 R ⁷⁰ , cycloalkyl substituted with 0-3 R ⁷⁰ , heterocycle substituted with 0-3 R ⁷⁰ , heterocycloalkyl substituted with 0-3 R ⁷⁰ , aralkyl substituted with 0-3 R ⁷⁰ and alkaryl substituted with 0-3 R ⁷⁰ ;
55	R ⁶¹ , R ⁶² , and R ⁶³	are independently selected at each occurrence from the group: C ₁ -C ₁₀ alkyl substituted with 0-3 R ⁷⁰ , aryl substituted with 0-3 R ⁷⁰ , cycloalkyl substituted with 0-3 R ⁷⁰ , heterocycle substituted with 0-3 R ⁷⁰ , aralkyl substituted with 0-3 R ⁷⁰ , alkaryl substituted with 0-3 R ⁷⁰ , and aryla-

lkaryl substituted with 0-3 R⁷⁰;

R⁷⁰ is independently selected at each occurrence from the group: F, Cl, Br, I, -CF₃, -CN, -CO₂R⁷¹, -C(=O)R⁷¹, -C(=O)N(R⁷¹)₂, -CH₂OR⁷¹, -OC(=O)R⁷¹, -OC(=O)OR^{71a}, -OR⁷¹, -OC(=O)N(R⁷¹)₂, -NR⁷¹C(=O)R⁷¹, -NR⁷¹C(=O)OR⁷¹, -NR⁷¹C(=O)N(R⁷¹)₂, SO₃⁻, -NR⁷¹SO₂N(R⁷¹)₂, -NR⁷¹SO₂R^{71a}, -SO₃H, -SO₂R⁷¹, -S(=O)R⁷¹, -SO₂N(R⁷¹)₂, -N(R⁷¹)₂, -N(R⁷¹)₃⁺, -NHC(=NH)NHR⁷¹, -C(=NH)NHR⁷¹, =NOR⁷¹, NO₂, -C(=O)NHOR⁷¹, -C(=O)NHN(R⁷¹)R^{71a}, -OCH₂CO₂H; and


R⁷¹ and R^{71a} are independently selected at each occurrence from the group: hydrogen and C₁-C₆ alkyl.

[20] Another embodiment of the present invention is the kit of embodiment [19] wherein:

Q is a biologically active molecule selected from the group: IIb/IIIa receptor antagonists, IIb/IIIa receptor ligands, fibrin binding peptides, leukocyte binding peptides, chemotactic peptides, somatostatin analogs, and selectin binding peptides;

d' is 1 to 3;

L_n is:



wherein:

g'' is 0-5;

f is 0-5;

f' is 1-5;

Y¹ and Y², at each occurrence, are independently selected from: O, NR⁵⁶, C=O, C(=O)O, OC(=O)O, C(=O)NH-, C=NR⁵⁶, S, SO, SO₂, SO₃, NHC(=O), (NH)₂C(=O), (NH)₂C=S;

R⁵⁵ and R⁵⁶ are independently selected at each occurrence from: hydrogen, C₁-C₁₀ alkyl, and (C₁-C₁₀ alkyl)aryl;

A_{L1} is selected from the group:

pyrones, pyridinones, and functionalized aminocarboxylates;

A_{L2} is selected from the group:

A⁹ and A¹⁰-W-A¹¹,

wherein:

A⁹ is PR⁶¹R⁶²R⁶³;

A¹⁰ and A¹¹ are PR⁶¹R⁶²;

W is a spacer group selected from the group: C₁-C₃ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰, and heterocycle substituted with 0-3 R⁷⁰; R⁶¹, R⁶², and R⁶³ are independently selected at each occurrence from the group: C₁-C₃ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰, and heterocycle substituted with 0-3 R⁷⁰;

R^{70} is independently selected at each occurrence from the group: $-\text{CO}_2R^{71}$, $-\text{OR}^{71}$, $-\text{SO}_3^-$ and $-\text{SO}_3\text{H}$; and


R^{71} is hydrogen.

[21] Another embodiment of the present invention is the kit of embodiment [20] wherein:

Q is a biologically active molecule selected from the group: IIb/IIIa receptor antagonists, and chemotactic peptides;

d' is 1;

L_n is:



wherein:

g'' is 0-5;

f is 0-5;

f' is 1-5;

Y^1 and Y^2 , at each occurrence, are independently selected from:
O, NR^{56} , $\text{C}=\text{O}$, $\text{C}(=\text{O})\text{O}$, $\text{OC}(=\text{O})\text{O}$, $\text{C}(=\text{O})\text{NH}-$, $\text{C}=\text{NR}^{56}$, S, $\text{NHC}(=\text{O})$, $(\text{NH})_2\text{C}(=\text{O})$, $(\text{NH})_2\text{C}=\text{S}$;

R^{55} and R^{56} are hydrogen;

A_{L1} is tricine;


A_{L2} is $\text{PR}^{61}\text{R}^{62}\text{R}^{63}$, wherein

R^{61} , R^{62} , and R^{63} are independently selected at each occurrence from the group: C_1 - C_3 alkyl substituted with 0-3 R^{70} , aryl substituted with 0-3 R^{70} ; and

R^{70} is independently selected at each occurrence from the group: $-\text{CO}_2\text{H}$, $-\text{OH}$, $-\text{SO}_3\text{H}$, $-\text{SO}_3^-$.


[22] Another embodiment of the present invention is the kit of embodiment [21] wherein:

Q is



d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is $\text{PR}^{61}\text{R}^{62}\text{R}^{63}$, wherein R^{61} , R^{62} and R^{63} are each phenyl bearing an SO_3H or SO_3^- group in the meta position.


[23] Another embodiment of the present invention is the kit of embodiment [21] wherein:

Q is



d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is $\text{PR}^{61}\text{R}^{62}\text{R}^{63}$, wherein R^{61} is phenyl, R^{62} and R^{63} are each phenyl bearing an SO_3H or SO_3^- group in the meta position.


[24] Another embodiment of the present invention is the kit of embodiment [21] wherein:

Q is



d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹ and R⁶² are phenyl, and R⁶³ is phenyl bearing an SO₃H or SO₃⁻ group in the meta position.


[25] Another embodiment of the present invention is the kit of embodiment [21] wherein:

Q is



d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹, R⁶² and R⁶³ are each p-(2-phenylethyl)phenyl wherein the phenylethyl bears an SO₃H or SO₃⁻ group in the para position.

[26] Another embodiment of the present invention is the kit of embodiment [21] wherein:


Q is



;

d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is $PR^{61}R^{62}R^{63}$, wherein R^{61} , R^{62} and R^{63} are each p-(2-phenylpropyl)phenyl wherein the phenylpropyl bears an SO_3H or SO_3^- group in the para position.

[27] Another embodiment of the present invention is the kit of embodiment [21] wherein:


Q is



;

d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is $R^{61}R^{62}PCH_2CH_2PR^{61}R^{62}$, wherein R^{61} , R^{62} are each phenyl substituted with an SO_3H or SO_3^- group in the meta position.


[28] Another embodiment of the present invention is the kit of embodiment [21] wherein:

Q is



d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹, R⁶² and R⁶³ are C3-alkyl substituted with 1 OH group.


[29] Another embodiment of the present invention is the kit of embodiment [21] wherein:

Q is



d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is $PR^{61}R^{62}R^{63}$, wherein R^{61} , R^{62} and R^{63} are CH_2CH_2COOH .


[30] Another embodiment of the present invention is the kit of embodiment [20] wherein:

Q is



d' is 1;

L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L1} is kojic acid;

A_{L2} is $PR^{61}R^{62}R^{63}$, wherein R^{61} , R^{62} and R^{63} are each phenyl bearing an SO_3H or SO_3^- group in the meta position.

[31] Another embodiment of the invention is the kits of any of embodiments [19]-[30] wherein a reducing agent is also present.

[32] A preferred embodiment of the invention is the kits of embodiment [31] wherein the reducing agent is stannous chloride.

[0016] When any variable occurs more than one time in any constituent or in any formula, its definition on each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-2 R^{52} , then said group may optionally be substituted with up to two R^{52} and R^{52} at each occurrence is selected independently from the defined list of possible R^{52} . Also, by way of example, for the group $-N(R^{53})_2$, each of the two R^{53} substituents on N is independently selected from the defined list of possible R^{53} . Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

[0017] By "stable compound" or "stable structure" is meant herein a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious diagnostic agent.

[0018] The term "capable of stabilizing", as used herein to describe the second ancillary ligand A_{L2} , means that the ligand is capable of coordinating to the transition metal radionuclide in the presence of the first ancillary ligand and the transition metal chelator, under the conditions specified herein, resulting in a radiopharmaceutical of Formula 1 having a minimal number of isomeric forms, the relative ratios of which do not change significantly with time, and that remains substantially intact upon dilution.

[0019] The term "substituted", as used herein, means that one or more hydrogens on the designated atom or group is replaced with a selection from the indicated group, provided that the designated atom's or group's normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (i.e., $=O$), then 2 hydrogens on the atom are replaced.

[0020] The term "bond", as used herein, means either a single or double bond.

[0021] The term "salt", as used herein, is used as defined in the CRC Handbook of Chemistry and Physics, 65th Edition, CRC Press, Boca Raton, Fla, 1984, as any substance which yields ions, other than hydrogen or hydroxyl ions.

[0022] As used herein, "alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms; "cycloalkyl" is intended to include saturated ring groups, including mono-, bi- or poly-cyclic ring systems, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl and adamantyl; and "bicycloalkyl" is intended to include saturated bicyclic ring groups such as [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), [2.2.2]bicyclooctane, and so forth.

[0023] As used herein, "aryl" or "aromatic residue" is intended to mean phenyl or naphthyl, which when substituted, the substitution can be at any position.

[0024] As used herein, the term "heterocycle" or "heterocyclic ring system" is intended to mean a stable 5- to 7-membered monocyclic or bicyclic or 7- to 10- membered bicyclic heterocyclic ring which may be saturated, partially unsaturated, or aromatic, and which consists of carbon atoms and from 1 to 4 heteroatoms selected independently from the group consisting of N, O and S and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen may optionally be quaternized, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom which results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. Examples of such heterocycles include, but are not limited to, benzopyranyl, thiadiazine, tetrazolyl, benzofuranyl, benzothiophenyl, indolene, quinoline, isoquinolinyl or benzimidazolyl, piperidinyl, 4-piperidone, 2-pyrrolidone, tetrahydrofuran, tetrahydroquinoline, tetrahydroisoquinoline, decahydroquinoline, octahydroisoquinoline, azocine, triazine (including 1,2,3-, 1,2,4-, and 1,3,5-triazine), 6H-1,2,5-thiadiazine, 2H,6H-1,5,2-dithiazine, thiophene, tetrahydrothiophene, thianthrene, furan, pyran, isobenzofuran, chromene, xanthene, phenoxathiin, 2H-pyrrole, pyrrole, imidazole, pyrazole, thiazole, isothiazole, oxazole (including 1,2,4- and 1,3,4-oxazole), isoxazole, triazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, 3H-indole, indole, 1H-indazole, purine, 4H-quinolizine, isoquinoline, quinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, 4aH-carbazole, carbazole, β -carboline, phenanthridine, acridine, perimidine, phenanthroline, phenazine, phenarsazine, phenothiazine, furazan, phenoxazine, isochroman, chroman, pyrrolidine, pyrroline, imidazolidine, imidazoline, pyrazolidine, pyrazoline, piperazine, indoline, isoindoline, quinuclidine, or morpholine. Also included are fused ring and spiro compounds containing, for example, the above heterocycles.

[0025] As used herein, the term "alkaryl" means an aryl group bearing an alkyl group of 1-10 carbon atoms; the term "aralkyl" means an alkyl group of 1-10 carbon atoms bearing an aryl group; the term "arylalkaryl" means an aryl group bearing an alkyl group of 1-10 carbon atoms bearing an aryl group; and the term "heterocycloalkyl" means an alkyl group of 1-10 carbon atoms bearing a heterocycle.

[0026] The biologically active molecule Q can be a protein, antibody, antibody fragment, peptide or polypeptide, or peptidomimetic that is comprised of a recognition sequence or unit for a receptor or binding site expressed at the site of the disease, or for a receptor or binding site expressed on platelets or leukocytes. The exact chemical composition of Q is selected based on the disease state to be diagnosed, the mechanism of localization to be utilized, and to provide an optimum combination of rates of localization, clearance and radionuclidic decay.

[0027] For the purposes of this invention, the term thromboembolic disease is taken to include both venous and arterial disorders and pulmonary embolism, resulting from the formation of blood clots.

[0028] For the diagnosis of thromboembolic disorders or atherosclerosis, Q is selected from the group including the cyclic IIb/IIIa receptor antagonist compounds described in co-pending U.S. Ser. No.08/218,861 (equivalent to WO 94/22494); the RGD containing peptides described in U.S. Patents 4,578,079, 4,792,525, the applications PCT US88/04403, PCT US89/01742, PCT US90/03788, PCT US91/02356 and by Ojima et. al., 204th Meeting of the Amer. Chem. Soc., 1992, Abstract 44; the peptides that are fibrinogen receptor antagonists described in European Patent Applications 90202015.5, 90202030.4, 90202032.2, 90202032.0, 90311148.2, 90311151.6, 90311537.6, the specific binding peptides and polypeptides described as IIb/IIIa receptor ligands, ligands for the polymerization site of fibrin, laminin derivatives, ligands for fibrinogen, or thrombin ligands in PCT WO 93/23085 (excluding the technetium binding groups); the oligopeptides that correspond to the IIIa protein described in PCT WO90/00178; the hirudin-based peptides described in PCT WO90/03391; the IIb/IIIa receptor ligands described in PCT WO90/15818; the thrombus, platelet binding or atherosclerotic plaque binding peptides described in PCT WO92/13572 (excluding the technetium binding group) or GB 9313965.7; the fibrin binding peptides described in U.S. Patents 4,427,646 and 5,270,030; the hirudin-based peptides described in U.S. Patent 5,279,812; or the fibrin binding proteins described in U.S. Patent 5,217,705; the guanine derivatives that bind to the IIb/IIIa receptor described in U.S. Patent 5,086,069; or the tyrosine derivatives described in European Patent Application 0478328A1, and by Hartman et. al., J. Med. Chem., 1992, 35, 4640; or oxidized low density lipoprotein (LDL).

[0029] For the diagnosis of infection, inflammation or transplant rejection, Q is selected from the group including the leukocyte binding peptides described in PCT WO93/17719 (excluding the technetium binding group), PCT

WO92/13572 (excluding the technetium binding group) or U.S. Ser. No. 08-140000; the chemotactic peptides described in Eur. Pat. Appl. 90108734.6 or A. Fischman et. al., Semin. Nuc. Med., 1994, 24, 154; or the leukostimulatory agents described in U.S. Patent 5,277,892.

[0030] For the diagnosis of cancer, Q is selected from the group of somatostatin analogs described in UK Application 8927255.3 or PCT WO94/00489, the selectin binding peptides described in PCT WO94/05269, the biological-function domains described in PCT WO93/12819, Platelet Factor 4 or the growth factors (PDGF, EGF, FGF, TNF MCSF or I11-8).

[0031] Q may also represent proteins, antibodies, antibody fragments, peptides, polypeptides, or peptidomimetics that bind to receptors or binding sites on other tissues, organs, enzymes or fluids. Examples include the β -amyloid proteins that have been demonstrated to accumulate in patients with Alzheimer's disease, atrial natriuretic factor derived peptides that bind to myocardial and renal receptors, antimyosin antibodies that bind to areas of infarcted tissues, or nitroimidazole derivatives that localize in hypoxic areas in vivo.

[0032] Ancillary dioxygen ligands include ligands that coordinate to the metal ion through at least two oxygen donor atoms. Examples include but are not limited to: glucoheptonate, gluconate, 2-hydroxyisobutyrate, lactate, tartrate, mannitol, glucarate, maltol, Kojic acid, 2,2-bis(hydroxymethyl)propionic acid, 4,5-dihydroxy-1,3-benzene disulfonate, or substituted or unsubstituted 1,2 or 3,4 hydroxypyridinones. (The names for the ligands in these examples refer to either the protonated or non-protonated forms of the ligands.)

[0033] Functionalized aminocarboxylates include ligands that have a combination of nitrogen and oxygen donor atoms. Examples include but are not limited to: iminodiacetic acid, 2,3 diaminopropionic acid, nitrilotriacetic acid, N, N'-ethylenediamine diacetic acid, N,N,N'-ethylenediamine triacetic acid, hydroxyethylethylenediamine triacetic acid, N,N'-ethylenediamine bis-hydroxyphenylglycine, or the ligands described in Eur. Pat. Appl. 93302712.0. (The names for the ligands in these examples refer to either the protonated or non-protonated forms of the ligands.)

[0034] The radiopharmaceuticals of the present invention for the diagnosis of thromboembolic disease can be easily prepared by admixing a salt of a radionuclide, a reagent of Formula 2, an ancillary ligand A_{L1} , an ancillary ligand A_{L2} , and optionally a reducing agent, in an aqueous solution at temperatures from room temperature to 100 °C.



and pharmaceutically acceptable salts thereof, wherein: Q, d', L_n are as defined above and C_h is a radionuclide metal chelator independently selected at each occurrence from the group: $R^{40}R^{41}N-N=C(C_1-C_3 \text{ alkyl})_2$ and $R^{40}NNH_2^-$, wherein R^{40} , R^{41} are as described above, and pharmaceutically acceptable salts thereof.

[0035] Alternatively, the radiopharmaceuticals of the present invention can be prepared by first admixing a salt of a radionuclide, an ancillary ligand A_{L1} , and a reducing agent in an aqueous solution at temperatures from room temperature to 100 °C to form an intermediate radionuclide complex with the ancillary ligand A_{L1} then adding a reagent of Formula 2 and an ancillary ligand A_{L2} and reacting further at temperatures from room temperature to 100 °C.

[0036] Alternatively, the radiopharmaceuticals of the present invention can be prepared by first admixing a salt of a radionuclide, an ancillary ligand A_{L1} , a reagent of Formula 2, and a reducing agent in an aqueous solution at temperatures from room temperature to 100 °C to form an intermediate radionuclide complex, as described in co-pending U. S. Ser. No.08/218,861 (equivalent to WO 94/22494), and then adding an ancillary ligand A_{L2} and reacting further at temperatures from room temperature to 100 °C.

[0037] The total time of preparation will vary depending on the identity of the radionuclide, the identities and amounts of the reactants and the procedure used for the preparation. The preparations may be complete, resulting in > 80% yield of the radiopharmaceutical, in 1 minute or may require more time. If higher purity radiopharmaceuticals are needed or desired, the products can be purified by any of a number of techniques well known to those skilled in the art such as liquid chromatography, solid phase extraction, solvent extraction, dialysis or ultrafiltration.

[0038] The radionuclides for the present invention are selected from the group ^{99m}Tc , ^{186}Re , or ^{188}Re . For diagnostic purposes ^{99m}Tc is the preferred isotope. Its 6 hour half-life and 140 keV gamma ray emission energy are almost ideal for gamma scintigraphy using equipment and procedures well established for those skilled in the art. The rhenium isotopes also have gamma ray emission energies that are compatible with gamma scintigraphy, however, they also emit high energy beta particles that are more damaging to living tissues. These beta particle emissions can be utilized for therapeutic purposes, for example, cancer radiotherapy.

[0039] The salt of ^{99m}Tc is preferably in the chemical form of pertechnetate and a pharmaceutically acceptable cation. The pertechnetate salt form is preferably sodium pertechnetate such as obtained from commercial Tc-99m generators. The amount of pertechnetate used to prepare the radiopharmaceuticals of the present invention can range from 0.1 mCi to 1 Ci, or more preferably from 1 to 200 mCi.

[0040] The reagents of Formula 2 can be synthesized as described in co-pending U.S. Ser. No.08/218,861 (equivalent to WO 94/22494). The amount of the reagents used to prepare the radiopharmaceuticals of the present invention

can range from 0.1 μg to 10 mg, or more preferably from 0.5 μg to 100 μg . The amount used will be dictated by the amounts of the other reactants and the identity of the radiopharmaceuticals of Formula 1 to be prepared.

[0041] The ancillary ligands A_{L1} used to synthesize the radiopharmaceuticals of the present invention can either be synthesized or obtained from commercial sources and include, halides, dioxygen ligands and functionalized aminocarboxylates. Dioxygen ligands are ligands that coordinate to the radionuclide through at least two oxygen donor atoms. Examples include but are not limited to: glucoheptonate, gluconate, 2-hydroxyisobutyrate, lactate, tartrate, mannitol, glucarate, maltol, Kojic acid, 2,2-bis(hydroxymethyl)propionic acid, 4,5-dihydroxy-1,3-benzene disulfonate, or substituted or unsubstituted 1,2- or 3,4-hydroxypyridinones, or pharmaceutically acceptable salts thereof.

[0042] Functionalized aminocarboxylates include ligands that coordinate to the radionuclide through a combination of nitrogen and oxygen donor atoms. Examples include but are not limited to: iminodiacetic acid, 2,3-diaminopropionic acid, nitrilotriacetic acid, N,N'-ethylenediamine diacetic acid, N,N,N'-ethylenediamine triacetic acid, hydroxyethylethylenediamine triacetic acid, N,N'-ethylenediamine bis-hydroxyphenylglycine, or the ligands described in Eur. Pat. Appl. 93302712.0, or pharmaceutically acceptable salts thereof.

[0043] Halides can be fluoride, chloride, bromide or iodide.

[0044] The selection of an ancillary ligand A_{L1} is determined by several factors including the chemical and physical properties of the ancillary ligand, the rate of formation, the yield, and the number of isomeric forms of the resulting radiopharmaceuticals, and the compatibility of the ligand in a lyophilized kit formulation. The charge and lipophilicity of the ancillary ligand will effect the charge and lipophilicity of the radiopharmaceuticals. For example, the use of 4,5-dihydroxy-1,3-benzene disulfonate results in radiopharmaceuticals with an additional two anionic groups because the sulfonate groups will be anionic under physiological conditions. The use of N-alkyl substituted 3,4-hydroxypyridinones results in radiopharmaceuticals with varying degrees of lipophilicity depending on the size of the alkyl substituents.








[0045] A series of functionalized aminocarboxylates are disclosed by Bridger et. al. that result in improved rates of formation of technetium labeled hydrazino modified proteins. We have determined that certain of these aminocarboxylates result in improved yields and a minimal number of isomeric forms of the radiopharmaceuticals of the present invention. The preferred ancillary ligands A_{L1} are the dioxygen ligands pyrones or pyridinones and functionalized aminocarboxylates that are derivatives of glycine; the most preferred is tricine (tris(hydroxymethyl)methylglycine).

[0046] The amounts of the ancillary ligands A_{L1} used can range from 0.1 mg to 1 g, or more preferably from 1 mg to 100 mg. The exact amount for a particular radiopharmaceutical is a function of the the procedure used and the amounts and identities of the other reactants. Too large an amount of A_{L1} will result in the formation of by-products comprised of technetium labeled A_{L1} without a biologically active molecule or by-products comprised of technetium labeled biologically active molecules with the ancillary ligand A_{L1} but without the ancillary ligand A_{L2} . Too small an amount of A_{L1} will result in other by-products such as reduced hydrolyzed technetium, or technetium colloid.

[0047] The preferred ancillary ligands A_{L2} are trisubstituted phosphines or trisubstituted arsines. The substituents can be alkyl, aryl, alkoxy, heterocycle, aralkyl, alkaryl and arylalkaryl and may or may not bear functional groups comprised of heteroatoms such as oxygen, nitrogen, phosphorus or sulfur. Examples of such functional groups include but are not limited to: hydroxyl, carboxyl, carboxamide, ether, ketone, amino, ammonium, sulfonate, sulfonamide, phosphonate, and phosphonamide. These phosphine and arsine ligands can be obtained either from commercial sources or can be synthesized by a variety of methods known to those skilled in the art. A number of methods can be found in Kosolapoff and Maier, Organic Phosphorus Compounds: Wiley-Interscience: New York, 1972; Vol. 1.

[0048] The selection of an ancillary ligand A_{L2} is determined by several factors including the chemical and physical properties of the ancillary ligand, the rate of formation, the yield, and the number of isomeric forms of the resulting radiopharmaceuticals, and the suitability of the ligand for a lyophilized kit formulation. Preferred ancillary ligands for the present invention are those that bear at least one functionality. The presence of the functionality effects the chemical and physical properties of the ancillary ligands such as basicity, charge, lipophilicity, size, stability to oxidation, solubility in water, and physical state at room temperature. The preferred ancillary ligands have a solubility in water of at least 0.001 mg/mL. This solubility allows the ligands to be used to synthesize the radiopharmaceuticals of the present invention without an added solubilizing agent or co-solvent.

[0049] The more preferred ancillary ligands A_{L2} include trisubstituted phosphines and trisubstituted arsines that have at least one functionality comprised of the heteroatoms oxygen, sulfur or nitrogen. These ligands can either be obtained commercially or synthesized. References for the synthesis of specific more preferred ligands can be obtained as follows: Tris(3-sulfonatophenyl)phosphine, sodium salt (TPPTS) was synthesized as described in Bartik et. al., Inorg. Chem., 1992, 31, 2667. Bis(3-sulfonatophenyl)phenylphosphine, sodium salt (TPPDS) and (3-sulfonatophenyl)diphenylphosphine, sodium salt (TPPMS) were synthesized as described in Kuntz, E., U.S. Patent 4,248,802. Tris(2-(p-sulfonatophenyl)ethyl) phosphine, sodium salt (TPEPTS) and Tris(3-(p-sulfonatophenyl)propyl)phosphine, sodium salt (TPPPTS) were prepared as described in Bartik et. al., Organometallics, 1993, 12, 164. 1,2-Bis[bis(3-sulfonatophenyl)phosphino]ethane, sodium salt (DPPETS) was synthesized as described in Bartik et. al., Inorg. Chem., 1994, 33, 164. References for the



synthesis of other more preferred ancillary ligands A_{L2} include Kuntz, E., Br. Pat. 1,540,242, Sinou, D., et. al., J. Chem. Soc. Chem Commun., 1986, 202, and Ahrlund, S., et. al., J. Chem. Soc., 1950, 264, 276.

[0050] The more preferred ligands A_{L2} have at least one functionality comprised of heteroatoms which do not bind to the technetium in competition with the donor atoms of the ancillary ligand A_{L1} or the hydrazino or diazino moiety of the reagents of Formula 2. The ligands bind only through the phosphorus or arsenic donors. This insures that the resulting radiopharmaceuticals of Formula 1 are formed as a mixture of a minimal number of isomeric forms. The ligands are also hydrophilic as evidenced by a solubility in water of at least 0.01 mg/mL. This insures that a sufficient concentration can be used to synthesize the radiopharmaceuticals in high yield. There is no maximum solubility limit for use in this invention. Therefore, the hydrophilicity of the more preferred ancillary ligands A_{L2} can still cover a wide range.

[0051] The charge and hydrophilicity of the ancillary ligand will effect the charge and hydrophilicity of the radiopharmaceuticals. As can be seen in Table 1, the hydrophilicity of a series of radiopharmaceuticals of Formula 1 that differ only in the identity of the ancillary ligand A_{L2} varies systematically as determined by the retention times on reverse-phase HPLC.

[0052] The amounts of the ancillary ligands A_{L2} used can range from 0.001 mg to 1 g, or more preferably from 0.01 mg to 10 mg. The exact amount for a particular radiopharmaceutical is a function of the procedure used and the amounts

and identities of the other reactants. Too large an amount of A_{L2} will result in the formation of by-products comprised of technetium labeled A_{L2} without a biologically active molecule or by-products comprised of technetium labeled biologically active molecules with the ancillary ligand A_{L2} but without the ancillary ligand A_{L1} .

[0053] A reducing agent can optionally be used for the synthesis of the radiopharmaceuticals of Formula 1. Suitable reducing agents include stannous salts, dithionite or bisulfite salts, borohydride salts, and formamidinesulfinic acid, wherein the salts are of any pharmaceutically acceptable form. The preferred reducing agent is a stannous salt. The use of a reducing agent is optional because the ancillary ligand A_{L2} can also serve to reduce the Tc-99m-pertechnetate. The amount of a reducing agent used can range from 0.001 mg to 10 mg, or more preferably from 0.005 mg to 1 mg.

[0054] Kits in accord with the present invention comprise a sterile, non-pyrogenic, mixture of a reagent of Formula 2, an ancillary ligand A_{L1} , an ancillary ligand A_{L2} , and optionally a reducing agent. Preferably, such kits are comprised of a lyophilized mixture of a predetermined amount of a reagent of Formula 2, a predetermined amount of an ancillary ligand A_{L1} , a predetermined amount of an ancillary ligand A_{L2} , and optionally a predetermined amount of a reducing agent. The kits may also optionally include a bulking agent or lyophilization aid or a buffer. A list of acceptable bulking agents or lyophilization aids and a list of acceptable buffers can be found in the United States Pharmacopeia.

[0055] The specific structure of a radiopharmaceutical of the present invention will depend on the identity of the biologically active molecule Q, the number d', the identity of the linker L_n , the identity of the chelator moiety C_h , the identity of the ancillary ligand A_{L1} , the identity of the ancillary ligand A_{L2} , and the identity of the radionuclide M_t . The identities of Q, L_n , and C_h , and the number d' are determined by the choice of the reagent of Formula 2. For a given reagent of Formula 2, the amount of the reagent, the amount and identity of the ancillary ligands A_{L1} and A_{L2} , the identity of the radionuclide M_t and the synthesis conditions employed will determine the structure of the radiopharmaceutical of Formula 1.

[0056] Radiopharmaceuticals synthesized using concentrations of reagents of Formula 2 of <100 $\mu\text{g/mL}$, will be comprised of one hydrazido or diazenido group C_h ; the value of x will be 1. Those synthesized using >1 mg/mL concentrations will be comprised of two hydrazido or diazenido groups; the value of x will be 2. The two C_h groups may be the same or different. For most applications, only a limited amount of the biologically active molecule can be injected and not result in undesired side-effects, such as chemical toxicity, interference with a biological process or an altered biodistribution of the radiopharmaceutical. Therefore, the radiopharmaceuticals with x equal to 2, which require higher concentrations of the reagents of Formula 2 comprised in part of the biologically active molecule, will have to be diluted or purified after synthesis to avoid such side-effects.

[0057] The identities and amounts used of the ancillary ligands A_{L1} and A_{L2} will determine the values of the variables y and z. The values of y can be an integer from 0 to 3, while the values of z can be an integer from 1 to 4. In combination, the values of y and z will result in a technetium coordination sphere that is made up of at least five and no more than seven donor atoms, preferably six donor atoms. For monodentate phosphines or arsines of the formula A^9 , z can be an integer from 1 to 4; for bidentate phosphines or arsines of the formula A^{10} - A^{11} , z can be either 1 or 2. The preferred combination for monodentate phosphines or arsines is y equal to 1 or 2 and z equal to 1. The preferred combination for bidentate phosphines or arsines is y equal to 0 or 1 and z equal to 1 or 2.

[0058] The radiopharmaceuticals are injected intravenously, usually in saline solution, at a dose of 1 to 100 mCi per 70 kg body weight, or preferably at a dose of 5 to 50 mCi. Imaging is performed using known procedures.

EXAMPLE SECTION

[0059] The materials used to synthesize the radiopharmaceuticals of the present invention described in the following examples were obtained as follows. The reagents of Formula 2 were synthesized as described in co-pending U.S. Ser. No. 08/218,861 (equivalent to WO 94/22494). The ancillary ligands tricine and Kojic Acid were obtained from Research Organics Inc. and Aldrich Chemical Co., respectively. The phosphines were synthesized as described above, except for tris(hydroxypropyl)phosphine which was obtained from Cytac Canada Limited and tris(carboxyethyl)phosphine which was obtained from Aldrich Chemical Co. Deionized water was obtained from a Milli-Q Water System and was of > 18 M Ω quality. Technetium-99m-pertechnetate ($^{99m}\text{TcO}_4^-$) was obtained from a DuPont Pharma $^{99}\text{Mo}/^{99m}\text{Tc}$ generator. Stannous chloride dihydrate was obtained from Aldrich Chemical Co.. D-Phe(OMe) was obtained from Bachem Bioscience Inc..

[0060] The following abbreviations are used herein:

TPPTS	Tris(3-sulfonatophenyl)phosphine, sodium salt
TPPDS	Bis (3-sulfonatophenyl)phenylphosphine, sodium salt
TPPMS	(3-sulfonatophenyl)diphenylphosphine, sodium salt
TPEPTS	Tris(2-(p-sulfonatophenyl)ethyl)phosphine, sodium salt
TPPPTS	Tris (3-(p-sulfonatophenyl)propyl)phosphine, sodium salt
THPP	Tris(3-hydroxypropyl)phosphine

EP 1 195 168 A2

TCEP Tris(2-carboxyethyl)phosphine
DPPETS 1,2-Bis[bis(3-sulfonatophenyl)phosphino] ethane, sodium salt

Example 1

Synthesis of $^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})\text{-Cyclo(D-Val-NMeArg-Gly-Asp-Mamb(hydrazino-nicotinyl-5-Aca))}$

[0061] To a clean 10 cc vial was added 40 mg tricine dissolved in 0.7 mL deionized H_2O , 5 μg Cyclo(D-Val-NMeArg-Gly-Asp-Mamb(hydrazino-nicotinyl-5-Aca)) dissolved in H_2O , 20 mCi $^{99m}\text{TcO}_4^-$ in saline, 1 mg TPPTS dissolved in H_2O , and 20 μg $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ dissolved in 0.1 N HCl. The total reaction volume was 1 - 1.5 mL. The pH of the solution was adjusted to 4 with 1 N HCl. The solution was heated at 50 °C for 30 minutes and then was analyzed by HPLC Method 1 and ITLC Method 1. Analytical and yield data are shown in Table 1.

Example 2

Synthesis of $^{99m}\text{Tc}(\text{tricine})(\text{TPPDS})\text{-Cyclo(D-Val-NMeArg-Gly-Asp-Mamb(hydrazino-nicotinyl-5-Aca))}$

[0062] The synthesis was performed as described in Example 1 substituting TPPDS as the phosphine co-ligand and heating at 80 °C for 30 minutes. Analytical and yield data are shown in Table 1.

Example 3

Synthesis of $^{99m}\text{Tc}(\text{tricine})(\text{TPPMS})\text{-Cyclo(D-Val-NMeArg-Gly-Asp-Mamb(hydrazino-nicotinyl-5-Aca))}$

[0063] The synthesis was performed as described in Example 2 substituting TPPMS as the phosphine co-ligand. Analytical and yield data are shown in Table 1.

Example 4

Synthesis of $^{99m}\text{Tc}(\text{tricine})(\text{TPEPTS})\text{-Cyclo(D-Val-NMeArg-Gly-Asp-Mamb(hydrazino-nicotinyl-5-Aca))}$

[0064] To a 10 cc vial was added 40 mg Tricine in 0.5 mL H_2O , 5 μg XV-120 in 100 μl H_2O , 50 mCi $^{99m}\text{TcO}_4^-$ in 0.5 mL 0.9% saline, 1.0 mg of TPEPTS in 0.2 mL H_2O , and 20 μg of $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ dissolved in 0.1 N HCl. Total Volume 1.4 mL. The pH of the solution was adjusted to 7 using 1 N NaOH. The solution was heated at 80 °C for 30 minutes and then was analyzed by HPLC Method 1 and ITLC Method 1. Analytical and yield data are shown in Table 1.

Example 5

Synthesis of $^{99m}\text{Tc}(\text{tricine})(\text{TPPPTS})\text{-Cyclo(D-Val-NMeArg-Gly-Asp-Mamb(hydrazino-nicotinyl-5-Aca))}$

[0065] The synthesis was performed as described in Example 4 substituting TPPPTS as the phosphine co-ligand. Analytical and yield data are shown in Table 1.

Example 6

Synthesis of $^{99m}\text{Tc}(\text{tricine})(\text{DPPETS})\text{-Cyclo(D-Val-NMeArg-Gly-Asp-Mamb(hydrazino-nicotinyl-5-Aca))}$

[0066] To a clean 10 cc vial was added 40 mg tricine dissolved in 0.7 mL deionized H_2O , 5 μg Cyclo(D-Val-NMeArg-Gly-Asp-Mamb(hydrazino-nicotinyl-5-Aca)) dissolved in H_2O , 20 mCi $^{99m}\text{TcO}_4^-$ in saline, and 20 μg $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ dissolved in 0.1 N HCl. The total reaction volume was 1 - 1.5 mL. The solution was maintained at room temperature for 5 minutes and then 1 mg DPPETS dissolved in H_2O was added. The pH of the solution was adjusted to 4 and then the solution was heated at 80 °C for 20 minutes. The resulting solution was analyzed by HPLC Method 1 and ITLC Method 1. Analytical and yield data are shown in Table 1.

Example 7

Synthesis of $^{99m}\text{Tc}(\text{tricine})(\text{THPP})\text{-Cyclo}(\text{D-Val-NMeArg-Gly-Asp-Mamb}(\text{hydrazino-nicotinyl-5-Aca}))$

- 5 **[0067]** The reagent is synthesized in two steps by first forming the reagent $^{99m}\text{Tc}(\text{tricine})\text{-Cyclo}(\text{D-Val-NMeArg-Gly-Asp-Mamb}(\text{hydrazino-nicotinyl-5-Aca}))$ and then reacting it with THPP.

Step 1. Synthesis of $^{99m}\text{Tc}(\text{tricine})\text{-Cyclo}(\text{D-Val-NMeArg-Gly-Asp-Mamb}(\text{hydrazino-nicotinyl-5-Aca}))$

- 10 **[0068]** To a 10 mL vial was added 0.3 mL of $^{99m}\text{TcO}_4^-$ (~ 100 mCi/mL) in saline, followed by 10 μg of Cyclo(D-Val-NMeArg-Gly-Asp-Mamb(hydrazino-nicotinyl-5-Aca)) dissolved in saline, 20 mg tricine dissolved in water at pH 7, and 20 μg of $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ dissolved in 1 N HCl. The reaction mixture was allowed to stand at room temperature for 15-20 min. and then analyzed by HPLC Method 1 and ITLC Method 1. The complex was formed in 90 - 95% yield.

- 15 Step 2. Reaction with THPP

[0069] To the reaction solution above was added 5 mg of THPP dissolved in saline. The mixture was heated at 50 °C for 15 - 20 min. The resulting solution was analyzed by HPLC Method 1 and ITLC Method 1. Analytical and yield data are shown in Table 1.

20

Example 8

Synthesis of $^{99m}\text{Tc}(\text{tricine})(\text{TCEP})\text{-Cyclo}(\text{D-Val-NMeArg-Gly-Asp-Mamb}(\text{hydrazino-nicotinyl-5-Aca}))$

- 25 **[0070]** The reagent is synthesized in two steps by first forming the reagent $^{99m}\text{Tc}(\text{tricine})\text{-Cyclo}(\text{D-Val-NMeArg-Gly-Asp-Mamb}(\text{hydrazino-nicotinyl-5-Aca}))$ and then reacting it with TCEP.

Step 1. Synthesis of $^{99m}\text{Tc}(\text{tricine})\text{-Cyclo}(\text{D-Val-NMeArg-Gly-Asp-Mamb}(\text{hydrazino-nicotinyl-5-Aca}))$

- 30 **[0071]** To a 10 mL vial was added 40 mg tricine dissolved in 0.5 mL H_2O , 5 μg of Cyclo(D-Val-NMeArg-Gly-Asp-Mamb(hydrazino-nicotinyl-5-Aca)) dissolved in 100 μL water, 0.5 mL of $^{99m}\text{TcO}_4^-$ (~ 100 mCi/mL) in saline, and 20 μg of $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ dissolved in 1 N HCl. The total reaction volume was 1 - 1.5 mL. The reaction mixture was allowed to stand at room temperature for 15-20 min. and then analyzed by HPLC Method 1 and ITLC Method 1. The complex was formed in 90 - 95% yield.

35

Step 2. Reaction with TCEP

- [0072]** To the reaction solution above was added 1.0 mg of TCEP dissolved in 0.2 mL water. The pH was adjusted to 4 using 1 N HCl. The mixture was heated at 50 °C for 15-20 min. The resulting solution was analyzed by HPLC Method 1 and ITLC Method 1. Analytical and yield data are shown in Table 1. (The product exists as two resolvable isomeric forms.)

40

Example 9

- 45 Synthesis of $^{99m}\text{Tc}(\text{Kojic Acid})(\text{TPPTS})\text{-Cyclo}(\text{D-Val-NMeArg-Gly-Asp-Mamb}(\text{hydrazino-nicotinyl-5-Aca}))$

[0073] The synthesis was performed as described in Example 1, substituting Kojic Acid (30 mg) for the tricine. Analytical and yield data are shown in Table 1.

- 50 Example 10

Synthesis of $^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})(\text{Hydrazino-nicotinyl-D-Phe}(\text{OMe}))$

Step 1. Synthesis of 2-Hydrazino-nicotinyl-D-Phe(OMe)

55

[0074] The synthesis was performed as described in co-pending U.S. Ser. No. , Example 3, substituting D-Phe(OMe) for the Cyclo(D-Val-NMeArg-Gly-Asp-Mamb(5-Aca)).

EP 1 195 168 A2

Step 2. Synthesis of $^{99m}\text{Tc}(\text{tricine})(\text{TPPTS})(\text{Hydrazino-nicotinyl-D-Phe(OMe)})$

[0075] The synthesis was performed as described in Example 1, substituting 2-hydrazino-nicotinyl-D-Phe(OMe) for the Cyclo(D-Val-NMeArg-Gly-Asp-Mamb(hydrazino-nicotinyl-5-Aca). The product is characterized by retention times of 17.6 and 18.0 minutes (HPLC Method 1) and is formed in 85% yield.

Purification

[0076] As a general rule, compounds provided by the methods described herein are pure, as shown by the analytical techniques described directly below. However, if greater purity is desired, compounds provided herein may be further purified on HPLC, by collecting the compound as it elutes from the HPLC column using Method 1, shown below. The volatiles are then evaporated and the residue redissolved in a 2% tricine in saline solution.

Analytical Methods:

HPLC Method 1

[0077]

Column: Vydac, C_{18} , 250 mm x 4.6 mm, 300 Å pore size

Flow: 1.0 mL/min

Solvent A: 10 mM sodium monophosphate, pH = 6.0

Solvent B: 100% acetonitrile

Gradient:

0% B	30% B	75% B	0% B
0 min	15 min	25 min	30 min

Detection by NaI probe

HPLC Method 2

[0078]

Column: Zorbax-Rx, C_{18} , 250 mm x 4.6 mm

Flow: 1.0 mL/min

Solvent A: 95% 5 mM tetrabutylammonium ion, 30 mM phosphate, pH = 3.7; 5% acetonitrile

Solvent B: 20% solvent A in acetonitrile

Gradient:

0% B	10% B	40% B	60% B	100% B
0 min	20 min	30 min	35 min	40 min

Detection by NaI probe

ITLC Method 1

[0079] Gelman ITLC-SG strips, 1 cm x 7.5 cm, developed in 1:1 acetone:saline (0.9%).

Table 1

Analytical and Yield Data for ^{99m}Tc Reagents		
	HPLC Retention time Method 1 (min)	% Yield
Example 1	10.4	95
Example 2	12.8	93

Table 1 (continued)

Analytical and Yield Data for ^{99m} Tc Reagents		
	HPLC Retention time Method 1 (min)	% Yield
Example 3	15.9	93
Example 4	10.0	70
Example 5	12.6	83
Example 6	9.6	88
Example 7	12.3	92
Example 8	8.7, 9.2	70
Example 9	9.3	80

[0080] The values reported in Table 1 were obtained using HPLC Method 1. One retention time is shown for most of these examples. The two species that comprise these radiopharmaceuticals are usually not completely resolved by this HPLC method. Typically there is a shoulder on the main peak reported.

Utility

[0081] The radiopharmaceuticals provided herein are useful as imaging agents for the diagnosis of cardiovascular disorders, such as thromboembolic disease or atherosclerosis, infectious disease and cancer. The radiopharmaceuticals are comprised of phosphine or arsine ligated technetium-99m labeled hydrazino or diazenido modified biologically active molecules that selectively localize at sites of disease and thus allow an image to be obtained of the loci using gamma scintigraphy. The complexes described in Examples 1-3 were evaluated for potential clinical utility as radiopharmaceuticals for the diagnosis of thromboembolic disease by performing imaging studies in a canine model of deep vein thrombosis. The blood clearance rates for the complexes were determined in the arteriovenous shunt model. Said imaging studies showed that the radiopharmaceuticals provided herein are useful in imaging thrombosis.

[0082] Canine Deep Vein Thrombosis Model: This model incorporates the triad of events (hypercoagulable state, period of stasis, low shear environment) essential for the formation of a venous fibrin-rich actively growing thrombus. The procedure was as follows: Adult mongrel dogs of either sex (9-13 kg) were anesthetized with pentobarbital sodium (35 mg/kg, i.v.) and ventilated with room air via an endotracheal tube (12 strokes/min, 25 ml/kg). For arterial pressure determination, the right femoral artery was cannulated with a saline-filled polyethylene catheter (PE-240) and connected to a Statham pressure transducer (P23ID; Oxnard, CA). Mean arterial blood pressure was determined via damping the pulsatile pressure signal. Heart rate was monitored using a cardiometer (Biotach, Grass Quincy, MA) triggered from a lead II electrocardiogram generated by limb leads. The right femoral vein was cannulated (PE-240) for drug administration. A 5 cm segment of both jugular veins was isolated, freed from fascia and circumscribed with silk suture. A microthermister probe was placed on the vessel which serves as an indirect measure of venous flow. A balloon embolectomy catheter was utilized to induce the 15 min period of stasis during which time a hypercoagulable state was then induced using 5 U thrombin (American Diagnostica, Greenwich CT) administered into the occluded segment. Fifteen minutes later, flow was reestablished by deflating the balloon. The radiopharmaceutical was infused during the first 5 minutes of reflow and the rate of incorporation monitored using gamma scintigraphy.

[0083] Arteriovenous Shunt Model: Adult mongrel dogs of either sex (9-13kg) were anesthetized with pentobarbital sodium (35 mg/kg, i.v.) and ventilated with room air via an endotracheal tube (12 strokes/min, 25 ml/kg). For arterial pressure determination, the left carotid artery was cannulated with a saline-filled polyethylene catheter (PE-240) and connected to a Statham pressure transducer (P23ID; Oxnard, CA). Mean arterial blood pressure was determined via damping the pulsatile pressure signal. Heart rate was monitored using a cardiometer (Biotach, Grass Quincy, MA) triggered from a lead II electrocardiogram generated by limb leads. A jugular vein was cannulated (PE-240) for drug administration. The both femoral arteries and femoral veins were cannulated with silicon treated (Sigmacote, Sigma Chemical Co. St Louis, MO), saline filled polyethylene tubing (PE-200) and connected with a 5 cm section of silicon treated tubing (PE-240) to form an extracorporeal arterio-venous shunts (A-V). Shunt patency was monitored using a doppler flow system (model VF-1, Crystal Biotech Inc, Hopkinton, MA) and flow probe (2-2.3 mm, Titronics Med. Inst., Iowa City, IA) placed proximal to the locus of the shunt. All parameters were monitored continuously on a polygraph recorder (model 7D Grass) at a paper speed of 10 mm/min or 25 mm/sec.

[0084] On completion of a 15 min post surgical stabilization period, an occlusive thrombus was formed by the introduction of a thrombogenic surface (4-0 braided silk thread, 5 cm in length, Ethicon Inc., Somerville, NJ) into the shunt

one shunt with the other serving as a control. Two consecutive 1hr shunt periods were employed with the test agent administered as an infusion over 5 min beginning 5 min before insertion of the thrombogenic surface. At the end of each 1 hr shunt period the silk was carefully removed and weighed and the % incorporation determined via well counting. Thrombus weight was calculated by subtracting the weight of the silk prior to placement from the total weight of the silk on removal from the shunt. Arterial blood was withdrawn prior to the first shunt and every 30 min thereafter for determination of blood clearance, whole blood collagen-induced platelet aggregation, thrombin-induced platelet degranulation (platelet ATP release), prothrombin time and platelet count. Template bleeding time was also performed at 30 min intervals.

Results

[0085] The results of the imaging studies performed on the radiopharmaceuticals of Examples 1 and 2 are shown in Figure 2 and Tc-99m-albumin, a negative control. The top graph shows the thrombus-to-blood ratios, the bottom graph shows the thrombus-to muscle ratios obtained from the images by drawing appropriate regions of interest and comparing the number of counts in each region. The values reported are for the images obtained at 15, 60 and 120 minutes after the end of the infusion of the compounds. Even as early as 15 minutes, the three radiopharmaceuticals have higher ratios than the negative control; the differences are pronounced by 60 - 120 minutes.

[0086] Complexes in which the biologically active molecules, Q, are chemotactic peptides can be evaluated for potential clinical utility as radiopharmaceuticals for the diagnosis of infection by performing imaging studies in a rabbit model of focal infection.

Rabbit Focal Infection Model

[0087] Using aseptic technique, adult rabbits of either sex (2-3 kg) were anesthetized with Ketamine/xylazine (15/1.5 mg/kg,i.v.) via the marginal ear vein. Each animal was administered a 1 ml suspension of 2×10^9 of e Coli in the posterior thigh muscle. At the appropriate time point, 18-48 hrs later, each animal was anesthetized with pentobarbital sodium (35 mg/kg,i.v.). A tracheotomy was then performed and the animal ventilated with room air using a rodent respirator. For arterial pressure determination, the left carotid artery was cannulated with a saline-filled polyethylene catheter and connected to a pressure transducer. Mean arterial blood pressure was determined via damping the pulsatile pressure signal. Heart rate was monitored using a cardiometer triggered from a lead II electrocardiogram generated by limb leads. A jugular vein was cannulated for drug administration. All parameters were monitored continuously on a polygraph recorder.

[0088] On completion of a 15 min post surgical stabilization period, the agent was infused over 1-5 min (1-20 mCi). On line assessment of the rate of incorporation into the inflammatory site was accomplished using serial scintigrams acquired at 0-3 and 18-24 hrs posttreatment. Images were acquired for a preset time of 5 min/view. To characterize the location of the peptide, region of interest analysis was performed comparing the infected thigh to the contralateral normal muscle at the corresponding time. Arterial blood was withdrawn prior to administration and every 30 min thereafter for determination of blood clearance, hematological profile and white blood cell function. On completion of the protocol, the animal was euthanized and the biodistribution of the compound determined via gamma well counting.

[0089] A radiopharmaceutical comprising a transition metal radionuclide, a transition metal chelator, a biologically active group connected to said chelator, a first ancillary ligand, a second ancillary ligand capable of stabilizing the radiopharmaceutical, optionally having a linking group between said chelator and said biologically active group.

[0090] A radiopharmaceutical as defined above having a linking group between said chelator and said biologically active group.

[0091] A radiopharmaceutical as defined above of formula:




wherein:

Q is a biologically active molecule;

d' is 1 to 20;

L_n is a linking group of formula:



wherein:

5		
	M ¹	is $-(CH_2)_gZ^1)_{g'}-(CR^{55}R^{56})_{g''}$;
	M ²	is $-(CR^{55}R^{56})_{g''}[Z^1(CH_2)_g]_{g'}$;
10	g	is independently 0-10;
	g'	is independently 0-1;
	g''	is independently 0-10;
15	f	is independently 0-10;
	f'	is independently 0-10;
20	f''	is independently 0-1;
	Y ¹ and Y ² ,	at each occurrence, are independently selected from: a bond, O, NR ⁵⁶ , C=O, C(=O)O, OC(=O)O, C(=O)NH-, C=NR ⁵⁶ , S, SO, SO ₂ , SO ₃ , NHC(=O), (NH) ₂ C(=O), (NH) ₂ C=S;
25	Z ¹	is independently selected at each occurrence from a C ₆ -C ₁₄ saturated, partially saturated, or aromatic carbocyclic ring system, substituted with 0-4 R ⁵⁷ ; and a heterocyclic ring system, optionally substituted with 0-4 R ⁵⁷ ;
30	R ⁵⁵ and R ⁵⁶	are independently selected at each occurrence from: hydrogen; C ₁ -C ₁₀ alkyl substituted with 0-5 R ⁵⁷ ; alkaryl wherein the aryl is substituted with 0-5 R ⁵⁷ ;
35	R ⁵⁷	is independently selected at each occurrence from the group: hydrogen, OH, NHR ⁵⁸ , C(=O)R ⁵⁸ , OC(=O)R ⁵⁸ , OC(=O)OR ⁵⁸ , C(=O)OR ⁵⁸ , C(=O)NR ⁵⁸ -, C≡N, SR ⁵⁸ , SOR ⁵⁸ , SO ₂ R ⁵⁸ , NHC(=O)R ⁵⁸ , NHC(=O)NHR ⁵⁸ , NHC(=S)NHR ⁵⁸ ; or, alternatively, when attached to an additional molecule Q, R ⁵⁷ is independently selected at each occurrence from the group: O, NR ⁵⁸ , C=O, C(=O)O, OC(=O)O, C(=O)N-, C=NR ⁵⁸ , S, SO, SO ₂ , SO ₃ , NHC(=O), (NH) ₂ C(=O), (NH) ₂ C=S; and,
40	R ⁵⁸	is independently selected at each occurrence from the group: hydrogen; C ₁ -C ₆ alkyl; benzyl, and phenyl;
	x and y	are independently 1 or 2;
45	z	is independently 1-4;
	M _t	is a transition metal radionuclide selected from the group: ^{99m} Tc, ¹⁸⁶ Re and ¹⁸⁸ Re;
50	C _h	is a radionuclide metal chelator coordinated to transition metal radionuclide M _t , and is independently selected at each occurrence, from the group: R ⁴⁰ N=N+=, R ⁴⁰ R ⁴¹ N=N=, R ⁴⁰ N=, and R ⁴⁰ N=N(H)-, wherein
55	R ⁴⁰	is independently selected at each occurrence from the group: a bond to L _n , C ₁ -C ₁₀ alkyl substituted with 0-3 R ⁵² , aryl substituted with 0-3 R ⁵² , cycloalkyl substituted with 0-3 R ⁵² , heterocycle substituted with 0-3 R ⁵² , heterocycloalkyl substituted with 0-3 R ⁵² , aralkyl substituted with 0-3 R ⁵² and alkaryl substituted with 0-3 R ⁵² ;

EP 1 195 168 A2

	R ⁴¹	is independently selected from the group: hydrogen, aryl substituted with 0-3 R ⁵² , C ₁ -C ₁₀ alkyl substituted with 0-3 R ⁵² , and a heterocycle substituted with 0-3 R ⁵² ;
5	R ⁵²	is independently selected at each occurrence from the group: a bond to L _n , =O, F, Cl, Br, I, -CF ₃ , -CN, -CO ₂ R ⁵³ , -C(=O)R ⁵³ , -C(=O)N(R ⁵³) ₂ , -CHO, -CH ₂ OR ⁵³ , -OC(=O)R ⁵³ , -OC(=O)OR ^{53a} , -OR ⁵³ , -OC(=O)N(R ⁵³) ₂ , -NR ⁵³ C(=O)R ⁵³ , -NR ⁵⁴ C(=O)OR ^{53a} , -NR ⁵³ C(=O)N(R ⁵³) ₂ , -NR ⁵⁴ SO ₂ N(R ⁵³) ₂ , -NR ⁵⁴ SO ₂ R ^{53a} , -SO ₃ H, -SO ₂ R ^{53a} , -SR ⁵³ , -S(=O)R ^{53a} , -SO ₂ N(R ⁵³) ₂ , -N(R ⁵³) ₂ , -NHC(=NH)NHR ⁵³ , -C(=NH)NHR ⁵³ , =NOR ⁵³ , NO ₂ , -C(=O)NHOR ⁵³ , -C(=O)NHN(R ⁵³) ₂ , -OCH ₂ CO ₂ H, 2-(1-morpholino)ethoxy;
10		
15	R ⁵³ , R ^{53a} , and R ⁵⁴	are each independently selected at each occurrence from the group: hydrogen, C ₁ -C ₆ alkyl, and a bond to L _n ;
	A _{L1}	is a first ancillary ligand selected from the group:
20		dioxygen ligand, functionalized aminocarboxylate, and halide;
	A _{L2}	is an ancillary ligand capable of stabilizing the radiopharmaceutical selected from the group:
25	A ⁹ and A ¹⁰ -W-A ¹¹ ,	
	wherein:	
30	A ⁹	is independently selected at each occurrence from the group: PR ⁶¹ R ⁶² R ⁶³ and AsR ⁶¹ R ⁶² R ⁶³ ;
	A ¹⁰ and A ¹¹	are independently selected at each occurrence from the group: PR ⁶¹ R ⁶² and AsR ⁶¹ R ⁶² ;
35	W	is a spacer group selected from the group: C ₁ -C ₁₀ alkyl substituted with 0-3 R ⁷⁰ , aryl substituted with 0-3 R ⁷⁰ , cycloalkyl substituted with 0-3 R ⁷⁰ , heterocycle substituted with 0-3 R ⁷⁰ , heterocycloalkyl substituted with 0-3 R ⁷⁰ , aralkyl substituted with 0-3 R ⁷⁰ and alkaryl substituted with 0-3 R ⁷⁰ ;
40	R ⁶¹ , R ⁶² , and R ⁶³	are independently selected at each occurrence from the group: C ₁ -C ₁₀ alkyl substituted with 0-3 R ⁷⁰ , aryl substituted with 0-3 R ⁷⁰ , cycloalkyl substituted with 0-3 R ⁷⁰ , heterocycle substituted with 0-3 R ⁷⁰ , aralkyl substituted with 0-3 R ⁷⁰ , alkaryl substituted with 0-3 R ⁷⁰ , and arylalkaryl substituted with 0-3 R ⁷⁰ ;
45		
50	R ⁷⁰	is independently selected at each occurrence from the group: F, Cl, Br, I, -CF ₃ , -CN, -CO ₂ R ⁷¹ , -C(=O)R ⁷¹ , -C(=O)N(R ⁷¹) ₂ , -CH ₂ OR ⁷¹ , -OC(=O)R ⁷¹ , -OC(=O)OR ^{71a} , -OR ⁷¹ , -OC(=O)N(R ⁷¹) ₂ , -NR ⁷¹ C(=O)R ⁷¹ , -NR ⁷¹ C(=O)OR ⁷¹ , -NR ⁷¹ C(=O)N(R ⁷¹) ₂ , SO ₃ ⁻ , -NR ⁷¹ SO ₂ N(R ⁷¹) ₂ , -NR ⁷¹ SO ₂ R ^{71a} , -SO ₃ H, -SO ₂ R ⁷¹ , -S(=O)R ⁷¹ , -SO ₂ N(R ⁷¹) ₂ , -N(R ⁷¹) ₂ , -N(R ⁷¹) ₃ ⁺ , -NHC(=NH)NHR ⁷¹ , -C(=NH)NHR ⁷¹ , =NOR ⁷¹ , NO ₂ , -C(=O)NHOR ⁷¹ , -C(=O)NHN(R ⁷¹) ₂ , -OCH ₂ CO ₂ H; and
55	R ⁷¹ and R ^{71a}	are independently selected at each occurrence from the group: hydrogen and C ₁ -C ₆ alkyl; and

pharmaceutically acceptable salts thereof.

[0092] A radiopharmaceutical as defined above wherein:

	Q	is a biologically active molecule selected from the group: IIb/IIIa receptor antagonists, IIb/IIIa receptor ligands, fibrin binding peptides, leukocyte binding peptides, chemotactic peptides, somatostatin analogs, and selectin binding peptides;
5	d'	is 1 to 3;
	L _n	is:
10		$-(\text{CR}^{55}\text{R}^{56})_{\text{g}''}-[\text{Y}^1(\text{CR}^{55}\text{R}^{56})_{\text{f}}\text{Y}^2]_{\text{f}'}-(\text{CR}^{55}\text{R}^{56})_{\text{g}''}-,$
		wherein:
15	g''	is 0-5;
	f	is 0-5;
	f'	is 1-5;
	Y ¹ and Y ² ,	at each occurrence, are independently selected from: O, NR ⁵⁶ , C=O, C(=O)O, OC(=O)O, C(=O)NH-, C=NR ⁵⁶ , S, SO, SO ₂ , SO ₃ , NHC(=O), (NH) ₂ C(=O), (NH) ₂ C=S;
20	R ⁵⁵ and R ⁵⁶	are independently selected at each occurrence from: hydrogen, C ₁ -C ₁₀ alkyl, and alkaryl;
	x and y	are independently 1 or 2;
25	z	is independently 1-2;
	M _t	is ^{99m} Tc;
30	C _h	is a radionuclide metal chelator coordinated to transition metal radionuclide M _t , and is independently selected at each occurrence, from the group: R ⁴⁰ N=N ⁺ =, R ⁴⁰ R ⁴¹ N-N=, R ⁴⁰ N=, and R ⁴⁰ N=N(H)-;
	R ⁴⁰	is independently selected at each occurrence from the group: aryl substituted with 0-3 R ⁵² , and heterocycle substituted with 0-3 R ⁵² ;
35	R ⁴¹	is independently selected from the group: hydrogen, aryl substituted with 0-1 R ⁵² , C ₁ -C ₃ alkyl substituted with 0-1 R ⁵² , and a heterocycle substituted with 0-1 R ⁵² ;
	R ⁵²	is independently selected at each occurrence from the group: a bond to L _n , -CO ₂ R ⁵³ , -CH ₂ OR ⁵³ , -SO ₃ H, -SO ₂ R ^{53a} , -N(R ⁵³) ₂ , -N(R ⁵³) ₃ ⁺ , -NHC(=NH)NHR ⁵³ , and -OCH ₂ CO ₂ H;
40	R ⁵³ , R ^{53a}	are each independently selected at each occurrence from the group: hydrogen and C ₁ -C ₃ alkyl;
45	A _{L1}	is selected from the group: pyrones, pyridinones, and functionalized aminocarboxylates;
50	A _{L2}	is selected from the group: A ⁹ and A ¹⁰ -W-A ¹¹ ,
		wherein:
55	A ⁹	is PR ⁶¹ R ⁶² R ⁶³ ;
	A ¹⁰ and A ¹¹	are PR ⁶¹ R ⁶² ;

EP 1 195 168 A2

W is a spacer group selected from the group: C₁-C₃ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰, and heterocycle substituted with 0-3 R⁷⁰; R⁶¹, R⁶², and R⁶³ are independently selected at each occurrence from the group: C₁-C₃ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰, and heterocycle substituted with 0-3 R⁷⁰;

R⁷⁰ is independently selected at each occurrence from the group: -CO₂R⁷¹, -OR⁷¹, -SO₃⁻ and -SO₃H; and


R⁷¹ is hydrogen.

[0093] A radiopharmaceutical as defined above wherein:

Q represents a biologically active molecule selected from the group: IIb/IIIa receptor antagonists and chemotactic peptides;

d' is 1;

L_n is:



wherein:

g'' is 0-5;

f is 0-5;

f' is 1-5;

Y¹ and Y², at each occurrence, are independently selected from:

O, NR⁵⁶, C=O, C(=O)O, OC(=O)O, C(=O)NH-, C=NR⁵⁶, S, NHC(=O), (NH)₂C(=O), (NH)₂C=S;

R⁵⁵ and R⁵⁶ are hydrogen;

x and y are 1;

z is 1;

C_{h'} is a radionuclide metal chelator coordinated to transition metal radionuclide Mt, and is independently selected at each occurrence, from the group: R⁴⁰N=N⁺=, and R⁴⁰R⁴¹N=N=;

R⁴⁰ is independently selected at each occurrence from the group: heterocycle substituted with R⁵²

R⁴¹ is hydrogen;

R⁵² is a bond to L_n;

A_{L1} is tricine;


A_{L2} is PR⁶¹R⁶²R⁶³, wherein

R⁶¹, R⁶², and R⁶³ are independently selected at each occurrence from the group: C₁-C₃ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰;

R⁷⁰ is independently selected at each occurrence from the group: -CO₂H, -OH, -SO₃H, -SO₃⁻.


[0094] The radiopharmaceutical as defined above wherein:

Q is




d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



C_h is



or



and is attached to L_n at the carbon atom designated with a *;

M_t is ^{99m}Tc;


A_{L1} is tricine;

A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹, R⁶² and R⁶³ are each phenyl bearing an SO₃H or SO₃⁻ group in the meta position; and

x, y and z are 1.


[0095] The radiopharmaceutical as defined above wherein:

Q is




d' is 1;

L_n is attached to Q at the carbon atom designated with a * and has the formula:



C_{h'} is



and is attached to L_n at the carbon atom designated with a *;

M_t is ^{99m}Tc;

A_{L1} is tricine;

A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹ is phenyl, R⁶² and R⁶³ are each phenyl bearing an SO₃H or SO₃⁻ group in the meta position; and

x, y and z are 1.

[0096] The radiopharmaceutical as defined above wherein:

Q is



15

20

20

N#Nc1ccncc1


•

35

40


45

50




d' is 1;

L_n is attached to Q at the carbon atom designated with a * and has the formula:



C_h' is



and is attached to L_n at the carbon atom designated with a *;

M_t is ^{99m}Tc;

A_{L1} is tricine;

A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹, R⁶² and R⁶³ are each p- (2-phenylethyl)phenyl wherein the phenylethyl bears an SO₃H or SO₃⁻ group in the para position; and

x, y and z are 1.

[0098] The radiopharmaceutical as defined above wherein:

Q is



15

20

25

30




40

45


50

55




d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



C_{h'} is



or



and is attached to L_n at the carbon atom designated with a *;

M_t is ^{99m}Tc;


A_{L1} is tricine;

A_{L2} is R⁶¹R⁶²PCH₂CH₂PR⁶¹R⁶², wherein R⁶¹, R⁶² are each phenyl substituted with an SO₃H or SO₃⁻ group in the meta position; and

x, y and z are 1.


[0100] The radiopharmaceutical as defined above wherein:

Q is




d' is 1;

L_n is attached to Q at the carbon atom designated with a * and has the formula:



C_{h'} is



and is attached to L_n at the carbon atom designated with a *;

M_t is ^{99m}Tc;


A_{L1} is tricine;

A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹, R⁶² and R⁶³ are C3-alkyl substituted with 1 OH group; and

x, y and z are 1.


[0101] The radiopharmaceutical as defined above wherein:

Q is




d' is 1;

L_n is attached to Q at the carbon atom designated with a * and has the formula:



C_{h'} is



and is attached to L_n at the carbon atom designated with a *;

M_t is ^{99m}Tc;

A_{L1} is tricine;

A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹, R⁶² and R⁶³ are CH₂CH₂COOH; and

x, y and z are 1.

[0102] The radiopharmaceutical as defined above wherein:

Q is



15

20

 $C_{h'}$ 

35

 M_t

40

A₁ 2

45

y

50


[01

55

[01

[01

(a) a predetermined quantity of a sterile, pharmaceutically acceptable reagent of formulae:



(b) a predetermined quantity of a sterile, pharmaceutically acceptable first ancillary ligand, A_{L1} , selected from the group:

dioxygen ligand,
functionalized aminocarboxylate, and
halide; and

(c) a predetermined quantity of a sterile, pharmaceutically acceptable second ancillary ligand, A_{L2} , selected from the group:


A^9 and $A^{10}-W-A^{11}$;

wherein:

Q is a biologically active molecule;

d' is 1 to 20;

L_n is a linking group of formula:



wherein:

M^1 is $[(CH_2)_gZ^1]_g-(CR^{55}R^{56})_{g''}$;

M^2 is $-(CR^{55}R^{56})_{g''}-[Z^1(CH_2)_g]_g$;

g is independently 0-10;

g' is independently 0-1;

g'' is independently 0-10;

f is independently 0-10;

f' is independently 0-10;

f'' is independently 0-1;

Y^1 and Y^2 , at each occurrence, are independently selected from:
a bond, O, NR^{56} , $C=O$, $C(=O)O$, $OC(=O)O$, $C(=O)NH-$, $C=NR^{56}$, S, SO,
 SO_2 , SO_3 , $NHC(=O)$, $(NH)_2C(=O)$, $(NH)_2C=S$;

Z^1 is independently selected at each occurrence from a C_6-C_{14} saturated, partially saturated, or aromatic carbocyclic ring system, substituted with 0-4 R^{57} ; and a heterocyclic ring system, optionally substituted with 0-4 R^{57} ;

R^{55} and R^{56} are independently selected at each occurrence from:
hydrogen; C_1-C_{10} alkyl substituted with 0-5 R^{57} ; alkaryl wherein the aryl is substituted with 0-5 R^{57} ;

EP 1 195 168 A2

5	R ⁵⁷	is independently selected at each occurrence from the group: hydrogen, OH, NHR ⁵⁸ , C(=O)R ⁵⁸ , OC(=O)R ⁵⁸ , OC(=O)OR ⁵⁸ , C(=O)OR ⁵⁸ , C(=O)NR ⁵⁸ -, C≡N, SR ⁵⁸ , SOR ⁵⁸ , SO ₂ R ⁵⁸ , NHC(=O)R ⁵⁸ , NHC(=O)NHR ⁵⁸ , NHC(=S)NHR ⁵⁸ ; or, alternatively, when attached to an additional molecule Q, R ⁵⁷ is independently selected at each occurrence from the group: O, NR ⁵⁸ , C=O, C(=O)O, OC(=O)O, C(=O)N-, C=NR ⁵⁸ , S, SO, SO ₂ , SO ₃ , NHC(=O), (NH) ₂ C(=O), (NH) ₂ C=S; and,
10	R ⁵⁸	is independently selected at each occurrence from the group: hydrogen; C ₁ -C ₆ alkyl; benzyl, and phenyl;
15	C _h	is a radionuclide metal chelator independently selected at each occurrence from the group: R ⁴⁰ R ⁴¹ N-N=C(C ₁ -C ₃ alkyl) ₂ and R ⁴⁰ NNH ₂ -, wherein::
20	R ⁴⁰	is independently selected at each occurrence from the group: a bond to L _n , C ₁ -C ₁₀ alkyl substituted with 0-3 R ⁵² , aryl substituted with 0-3 R ⁵² , cycloalkyl substituted with 0-3 R ⁵² , heterocycle substituted with 0-3 R ⁵² , heterocycloalkyl substituted with 0-3 R ⁵² , aralkyl substituted with 0-3 R ⁵² and alkaryl substituted with 0-3 R ⁵² ;
25	R ⁴¹	is independently selected from the group: hydrogen, aryl substituted with 0-3 R ⁵² , C ₁ -C ₁₀ alkyl substituted with 0-3 R ⁵² , and a heterocycle substituted with 0-3 R ⁵² ;
30	R ⁵²	is independently selected at each occurrence from the group: a bond to L _n , =O, F, Cl, Br, I, -CF ₃ , -CN, -CO ₂ R ⁵³ , -C(=O)R ⁵³ , -C(=O)N(R ⁵³) ₂ , -CHO, -CH ₂ OR ⁵³ , -OC(=O)R ⁵³ , -OC(=O)OR ^{53a} , -OR ⁵³ , -OC(=O)N(R ⁵³) ₂ , -NR ⁵³ C(=O)R ⁵³ , -NR ⁵⁴ C(=O)OR ^{53a} , -NR ⁵³ C(=O)N(R ⁵³) ₂ , -NR ⁵⁴ SO ₂ N(R ⁵³) ₂ , -NR ⁵⁴ SO ₂ R ^{53a} , -SO ₃ H, -SO ₂ R ^{53a} , -SR ⁵³ , -S(=O)R ^{53a} , -SO ₂ N(R ⁵³) ₂ , -N(R ⁵³) ₂ , -NHC(=NH)NHR ⁵³ , -C(=NH)NHR ⁵³ , =NOR ⁵³ , NO ₂ , -C(=O)NHOR ⁵³ , -C(=O)NHN(R ⁵³) ₂ , -OCH ₂ CO ₂ H, 2-(1-morpholino)ethoxy;
35	R ⁵³ , R ^{53a} , and R ⁵⁴	are each independently selected at each occurrence from the group: hydrogen, C ₁ -C ₆ alkyl, and a bond to L _n ;
40	A ⁹	is independently selected at each occurrence from the group: PR ⁶¹ R ⁶² R ⁶³ and AsR ⁶¹ R ⁶² R ⁶³ ;
45	A ¹⁰ and A ¹¹	are independently selected at each occurrence from the group: PR ⁶¹ R ⁶² and AsR ⁶¹ R ⁶² ;
50	W	is a spacer group selected from the group: C ₁ -C ₁₀ alkyl substituted with 0-3 R ⁷⁰ , aryl substituted with 0-3 R ⁷⁰ , cycloalkyl substituted with 0-3 R ⁷⁰ , heterocycle substituted with 0-3 R ⁷⁰ , heterocycloalkyl substituted with 0-3 R ⁷⁰ , aralkyl substituted with 0-3 R ⁷⁰ and alkaryl substituted with 0-3 R ⁷⁰ ;
55	R ⁶¹ , R ⁶² , and R ⁶³	are independently selected at each occurrence from the group: C ₁ -C ₁₀ alkyl substituted with 0-3 R ⁷⁰ , aryl substituted with 0-3 R ⁷⁰ , cycloalkyl substituted with 0-3 R ⁷⁰ , heterocycle substituted with 0-3 R ⁷⁰ , aralkyl substituted with 0-3 R ⁷⁰ , alkaryl substituted with 0-3 R ⁷⁰ , and arylalkaryl substituted with 0-3 R ⁷⁰ ;
60	R ⁷⁰	is independently selected at each occurrence from the group: F, Cl, Br, I, -CF ₃ , -CN, -CO ₂ R ⁷¹ , -C(=O)R ⁷¹ , -C(=O)N(R ⁷¹) ₂ , -CH ₂ OR ⁷¹ , -OC(=O)R ⁷¹ , -OC(=O)OR ^{71a} , -OR ⁷¹ , -OC(=O)N(R ⁷¹) ₂ , -NR ⁷¹ C(=O)R ⁷¹ , -NR ⁷¹ C(=O)OR ⁷¹ , -NR ⁷¹ C(=O)N(R ⁷¹) ₂ , SO ₃ -, -NR ⁷¹ SO ₂ N(R ⁷¹) ₂ , -NR ⁷¹ SO ₂ R ^{71a} , -SO ₃ H, -SO ₂ R ⁷¹ , -S(=O)R ⁷¹ , -SO ₂ N(R ⁷¹) ₂ , -N(R ⁷¹) ₂ , -N(R ⁷¹) ₃ +, -NHC(=NH)NHR ⁷¹ , -C(=NH)NHR ⁷¹ , =NOR ⁷¹ , NO ₂ , -C(=O)

EP 1 195 168 A2

NHOR⁷¹, -C(=O)NHN⁷¹R^{71a}, -OCH₂CO₂H; and


R⁷¹ and R^{71a} are independently selected at each occurrence from the group: hydrogen and C₁-C₆ alkyl.

[0108] The kit as defined above wherein:

Q is a biologically active molecule selected from the group: IIb/IIIa receptor antagonists, IIb/IIIa receptor ligands, fibrin binding peptides, leukocyte binding peptides, chemotactic peptides, somatostatin analogs, and selectin binding peptides;

d' is 1 to 3;

L_n is:



wherein:

g'' is 0-5;

f is 0-5;

f' is 1-5;

Y¹ and Y², at each occurrence, are independently selected from:
O, NR⁵⁶, C=O, C(=O)O, OC(=O)O, C(=O)NH-, C=NR⁵⁶, S, SO, SO₂, SO₃, NHC(=O), (NH)₂C(=O), (NH)₂C=S;

R⁵⁵ and R⁵⁶ are independently selected at each occurrence from: hydrogen, C₁-C₁₀ alkyl, and (C₁-C₁₀ alkyl)aryl;

A_{L1} is selected from the group:

pyrones, pyridinones, and functionalized aminocarboxylates;

A_{L2} is selected from the group:

A⁹ and A¹⁰-W-A¹¹,

wherein:

A⁹ is PR⁶¹R⁶²R⁶³;

A¹⁰ and A¹¹ are PR⁶¹R⁶²;

W is a spacer group selected from the group: C₁-C₃ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰, and heterocycle substituted with 0-3 R⁷⁰;
R⁶¹, R⁶², and R⁶³ are independently selected at each occurrence from the group: C₁-C₃ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰, and heterocycle substituted with 0-3 R⁷⁰;

R⁷⁰ is independently selected at each occurrence from the group: -CO₂R⁷¹, -OR⁷¹, -SO₃⁻ and -SO₃H; and

R⁷¹ is hydrogen.


[0109] The kit as defined above wherein:

Q is a biologically active molecule selected from the group: IIb/IIIa receptor antagonists, and chemotactic

peptides;

d' is 1;

5 L_n is:



10 wherein:

g'' is 0-5;

f is 0-5;

f' is 1-5;

15 Y¹ and Y², at each occurrence, are independently selected from:

O, NR⁵⁶, C=O, C(=O)O, OC(=O)O, C(=O)NH-, C=NR⁵⁶, S, NHC(=O), (NH)₂C(=O), (NH)₂C=S;

R⁵⁵ and R⁵⁶ are hydrogen;

20 A_{L1} is tricine;


A_{L2} is PR⁶¹R⁶²R⁶³, wherein

25 R⁶¹, R⁶², and R⁶³ are independently selected at each occurrence from the group: C₁-C₃ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰; and

R⁷⁰ is independently selected at each occurrence from the group: -CO₂H, -OH, -SO₃H, -SO₃⁻.

30 **[0110]** The kit as defined above wherein:

Q is



d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹, R⁶² and R⁶³ are each phenyl bearing an SO₃H or SO₃⁻ group in the meta position.


[0111] The kit as defined above wherein:

Q is



d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹ is phenyl, R⁶² and R⁶³ are each phenyl bearing an SO₃H or SO₃⁻ group in the meta position.


[0112] The kit as defined above wherein:

Q is



d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is $PR^{61}R^{62}R^{63}$, wherein R^{61} and R^{62} are phenyl, and R^{63} is phenyl bearing an SO_3H or SO_3^- group in the meta position.


[0113] The kit as defined above wherein:

Q is



d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is $PR^{61}R^{62}R^{63}$, wherein R^{61} , R^{62} and R^{63} are each p -(2-phenylethyl)phenyl wherein the phenylethyl bears an SO_3H or SO_3^- group in the para position.


[0114] The kit as defined above wherein:

Q is



d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is $\text{PR}^{61}\text{R}^{62}\text{R}^{63}$, wherein R^{61} , R^{62} and R^{63} are each p -(2-phenylpropyl)phenyl wherein the phenylpropyl bears an SO_3H or SO_3^- group in the para position.


[0115] The kit as defined above wherein:

Q is



d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is $\text{R}^{61}\text{R}^{62}\text{PCH}_2\text{CH}_2\text{PR}^{61}\text{R}^{62}$, wherein R^{61} , R^{62} are each phenyl substituted with an SO_3H or SO_3^- group in the meta position.


[0116] The kit as defined above wherein:

Q is



d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹, R⁶² and R⁶³ are C3-alkyl substituted with 1 OH group.


[0117] The kit as defined above wherein:

Q is



d' is 1;


L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹, R⁶² and R⁶³ are CH₂CH₂COOH.


[0118] The kit as defined above wherein:

Q is



d' is 1;

L_n is attached to Q at the carbon atom designated with a * and has the formula:



A_{L1} is kojic acid;

A_{L2} is PR⁶¹R⁶²R⁶³, wherein R⁶¹, R⁶² and R⁶³ are each phenyl bearing an SO₃H or SO₃⁻ group in the meta position.

[0119] The kits as defined above wherein a reducing agent is also present.

[0120] The kits as defined above wherein the reducing agent is stannous chloride.

Claims

1. A radiopharmaceutical comprising a transition metal radionuclide, a transition metal chelator, a biologically active group connected to said chelator, a first ancillary ligand, a second ancillary ligand capable of stabilizing the radiopharmaceutical, said pharmaceutical having the formula:



wherein:

Q is a biologically active molecule;

d' is 1 to 20;

x and y are independently 1 or 2;

z is independently 1-4;

M_t is a transition metal radionuclide selected from: ^{99m}Tc, ¹⁸⁶Re and ¹⁸⁸Re;

C_h is a radionuclide metal chelator coordinated to transition metal radionuclide M_t, and is independently selected at each occurrence, from: R⁴⁰N=N⁺=, R⁴⁰R⁴¹N=N=, R⁴⁰N=, and R⁴⁰N=N(H)-, wherein

R⁴⁰ is independently selected at each occurrence from : a bond to Q C₁-C₁₀ alkyl substituted with 0-3 R⁵², aryl substituted with 0-3 R⁵², cycloalkyl substituted with 0-3 R⁵², heterocycle substituted with 0-3 R⁵², heterocycloalkyl substituted with 0-3 R⁵², aralkyl substituted with 0-3 R⁵² and alkaryl substituted with 0-3 R⁵²;

R⁴¹ is independently selected from: hydrogen, aryl substituted with 0-3 R⁵², C₁-C₁₀ alkyl substituted with 0-3 R⁵², and a heterocycle substituted with 0-3 R⁵²;

R⁵² is independently selected at each occurrence from : a bond to Q =O, F, Cl, Br, I, -CF₃, -CN, -CO₂R⁵³, -C(=O)R⁵³, -C(=O)N(R⁵³)₂, -CHO, -CH₂OR⁵³, -OC(=O)R⁵³, -OC(=O)OR^{53a}, -OR⁵³, -OC(=O)N(R⁵³)₂, -NR⁵³C(=O)R⁵³, -NR⁵⁴C(=O)OR^{53a}, -NR⁵³C(=O)N(R⁵³)₂, -NR⁵⁴SO₂N(R⁵³)₂, -NR⁵⁴SO₂R^{53a}, -SO₃H, -SO₂R^{53a}, -SR⁵³, -S(=O)R^{53a}, -SO₂N(R⁵³)₂, -N(R⁵³)₂, -NHC(=NH)NHR⁵³, -C(=NH)NHR⁵³, =NOR⁵³, NO₂, -C(=O)NHOR⁵³, -C(=O)NHNHR⁵³R^{53a}, -OCH₂CO₂H,

EP 1 195 168 A2

2-(1-morpholino)ethoxy;

R⁵³, R^{53a}, and R⁵⁴ are each independently selected at each occurrence from:

hydrogen, C₁-C₆ alkyl, and a bond to Q;

A_{L1} is a first ancillary ligand selected from:

dioxygen ligand,
functionalized aminocarboxylate, and
halide;

A_{L2} is an ancillary ligand capable of stabilizing the radiopharmaceutical selected from:

A⁹ and A¹⁰-W-A¹¹,

wherein:

A⁹ is independently selected at each occurrence from:

PR⁶¹R⁶²R⁶³ and AsR⁶¹R⁶²R⁶³;

A¹⁰ and A¹¹ are independently selected at each occurrence from:

PR⁶¹R⁶² and AsR⁶¹R⁶²;

W is a spacer group selected from:

C₁-C₁₀ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰,
cycloalkyl substituted with 0-3 R⁷⁰, heterocycle substituted with
0-3 R⁷⁰, heterocycloalkyl substituted with 0-3 R⁷⁰, aralkyl substi-
tuted with 0-3 R⁷⁰ and alkaryl substituted with 0-3 R⁷⁰;

R⁶¹, R⁶², and R⁶³ are independently selected at each occurrence from:

C₁-C₁₀ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰,
cycloalkyl substituted with 0-3 R⁷⁰, heterocycle substituted with
0-3 R⁷⁰, aralkyl substituted with 0-3 R⁷⁰, alkaryl substituted with
0-3 R⁷⁰, and arylalkaryl substituted with 0-3 R⁷⁰;

R⁷⁰ is independently selected at each occurrence from: F, Cl, Br, I, -CF₃,
-CN, -CO₂R⁷¹, -C(=O)R⁷¹, -C(=O)N(R⁷¹)₂, -CH₂OR⁷¹, -OC(=O)R⁷¹,
-OC(=O)OR^{71a}, -OR⁷¹, -OC(=O)N(R⁷¹)₂, -NR⁷¹C(=O)R⁷¹, -NR⁷¹C
(=O)OR⁷¹, -NR⁷¹C(=O)N(R⁷¹)₂, SO₃⁻, -NR⁷¹SO₂N(R⁷¹)₂,
-NR⁷¹SO₂R^{71a}, -SO₃H, -SO₂R⁷¹, -S(=O)R⁷¹, -SO₂N(R⁷¹)₂, -N(R⁷¹)₂,
-N(R⁷¹)₃⁺, -NHC(=NH)NHR⁷¹, -C(=NH)NHR⁷¹, =NOR⁷¹, NO₂, -C(=O)
NHOR⁷¹, -C(=O)NHN(R⁷¹)₂, -OCH₂CO₂H; and

R⁷¹ and R^{71a} are independently selected at each occurrence from:

hydrogen and C₁-C₆ alkyl; and

pharmaceutically acceptable salts thereof.

2. A radiopharmaceutical of Claim 1 wherein:

Q is a biologically active molecule selected from:

IIb/IIIa receptor antagonists, IIb/IIIa receptor ligands, fibrin binding peptides, leukocyte binding peptides, chemotactic peptides, somatostatin analogs, and selectin binding peptides;

5	d'	is 1 to 3;
	x and y	are independently 1 or 2;
	z	is independently 1-2;
10	M _t	is ^{99m} Tc;
	C _h	is a radionuclide metal chelator coordinated to transition metal radionuclide M _t , and is independently selected at each occurrence, from: R ⁴⁰ N=N ⁺ =, R ⁴⁰ R ⁴¹ N-N=, R ⁴⁰ N=, and R ⁴⁰ N=N(H)-;
15	R ⁴⁰	is independently selected at each occurrence from: aryl substituted with 0-3 R ⁵² , and heterocycle substituted with 0-3 R ⁵² ;
	R ⁴¹	is independently selected from:
20		hydrogen, aryl substituted with 0-1 R ⁵² , C ₁ -C ₃ alkyl substituted with 0-1 R ⁵² , and a heterocycle substituted with 0-1 R ⁵² ;
	R ⁵²	is independently selected at each occurrence from: a bond to Q, -CO ₂ R ⁵³ , -CH ₂ OR ⁵³ , -SO ₃ H, -SO ₂ R ^{53a} , -N(R ⁵³) ₂ , -N(R ⁵³) ₃ ⁺ , -NHC(=NH)NHR ⁵³ , and -OCH ₂ CO ₂ H;
25	R ⁵³ , R ^{53a}	are each independently selected at each occurrence from:
		hydrogen and C ₁ -C ₃ alkyl;
30	A _{L1}	is selected from:
		pyrones, pyridinones, and functionalized aminocarboxylates;
35	A _{L2}	is selected from;
		A ⁹ and A ¹⁰ -W-A ¹¹ ,
40		wherein:
	A ⁹	is PR ⁶¹ R ⁶² R ⁶³ ;
	A ¹⁰ and A ¹¹	are PR ⁶¹ R ⁶² ;
45	W	is a spacer group selected from:
		C ₁ -C ₃ alkyl substituted with 0-3 R ⁷⁰ , aryl substituted with 0-3 R ⁷⁰ , and heterocycle substituted with 0-3 R ⁷⁰ ;
50		R ⁶¹ , R ⁶² , and R ⁶³ are independently selected at each occurrence from:
		C ₁ -C ₃ alkyl substituted with 0-3 R ⁷⁰ , aryl substituted with 0-3 R ⁷⁰ , and heterocycle substituted with 0-3 R ⁷⁰ ;
55	R ⁷⁰	is independently selected at each occurrence from: -CO ₂ R ⁷¹ , -OR ⁷¹ , -SO ₃ ⁻ and -SO ₃ H; and
	R ⁷¹	is hydrogen.

3. A radiopharmaceutical of Claim 2 wherein:

Q represents a biologically active molecule selected from the group: IIb/IIIa receptor antagonists and chemotactic peptides;

d' is 1;

x and y are 1;

z is 1;

C_h is a radionuclide metal chelator coordinated to transition metal radionuclide M_t, and is independently selected at each occurrence, from: R⁴⁰N=N⁺=, and R⁴⁰R⁴¹N-N=;

R⁴⁰ is independently selected at each occurrence from; heterocycle substituted with R⁵²;

R⁴¹ is hydrogen;

R⁵² is a bond to Q;

A_{L1} is tricine;

A_{L2} is PR⁶¹R⁶²R⁶³, wherein

R⁶¹, R⁶², and R⁶³ are independently selected at each occurrence from:


C₁-C₃ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰;

R⁷⁰ is independently selected at each occurrence from: -CO₂H, -OH, -SO₃H, -SO₃⁻.

4. Use of a radiopharmaceutical as defined in any of claims 1-3 for preparing an agent for radioimaging a mammal.

5. A kit for preparing a radiopharmaceutical comprising:

(a) a predetermined quantity of a sterile, pharmaceutically acceptable reagent of formulae:



wherein Q and d' are as defined in claim 1, and

C_h is a radionuclide metal chelator independently selected at each occurrence from: R⁴⁰R⁴¹N-N=C(C₁-C₃ alkyl)₂ and R⁴⁰NNH₂⁻, wherein R⁴⁰ and R⁴¹ are as defined in claim 1

(b) a predetermined quantity of a sterile, pharmaceutically acceptable first ancillary ligand, A_{L1}, selected from;

dioxygen ligand,
functionalized aminocarboxylate, and
halide; and

(c) a predetermined quantity of a sterile, pharmaceutically acceptable second ancillary ligand, A_{L2}, selected from:

A⁹ and A¹⁰-W-A¹¹,

wherein A⁹, A¹⁰, A¹¹ and W are as defined in claim 1.

6. The kit of Claim 5 wherein

Q is a biologically active molecule selected from:

IIb/IIIa receptor antagonists, IIb/IIIa receptor ligands, fibrin binding peptides, leukocyte binding peptides, chemotactic peptides, somatostatin analogs, and selectin binding peptides;

d' is 1 to 3;

A_{L1} is selected from:

pyrones, pyridinones, and functionalized aminocarboxylates;

A_{L2} is selected from:

A⁹ and A¹⁰-W-A¹¹,

wherein:

A⁹ is PR⁶¹R⁶²R⁶³;

A¹⁰ and A¹¹ are PR⁶¹R⁶²;

W is a spacer group selected from:

C₁-C₃ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰, and heterocycle substituted with 0-3 R⁷⁰;

R⁶¹, R⁶², and R⁶³ are independently selected at each occurrence from:

C₁-C₃ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰, and heterocycle substituted with 0-3 R⁷⁰;

R⁷⁰ is independently selected at each occurrence from: -CO₂R⁷¹, -OR⁷¹, -SO₃⁻ and -SO₃H; and

R⁷¹ is hydrogen.

7. The kit of Claim 6 wherein:

Q is a biologically active molecule selected from:

IIb/IIIa receptor antagonists, and chemotactic peptides;

A_{L1} is tricine;

A_{L2} is PR⁶¹R⁶²R⁶³, wherein

R⁶¹, R⁶², and R⁶³ are independently selected at each occurrence from:

C₁-C₃ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰; and

R⁷⁰ is independently selected at each occurrence from: -CO₂H, -OH, -SO₃H, -SO₃⁻.

8. The kits of any of Claims 5-7 wherein a reducing agent is also present.

9. The kits of Claim 8 wherein the reducing agent is stannous chloride.

10. A method for radioimaging a mammal for the diagnosis of cardiovascular disorders, comprising:


- (i) administering to said mammal an effective amount of a radiopharmaceutical; and
- (ii) scanning the mammal using a radioimaging device,

wherein the radiopharmaceutical is of the formula:



wherein:

Q is a biologically active molecule;
d' is 1 to 20;
L_n is a linking group of formula:



wherein:

M¹ is $-(CH_2)_gZ^1]_{g''}-(CR^{55}R^{56})_{g''}$;
M² is $-(CR^{55}R^{56})_{g''}-[Z^1(CH_2)_g]_{g''}$;
g is independently 0-10;
g' is independently 0-1;
g'' is independently 0-10;
f is independently 0-10;
f' is independently 0-10;
f'' is independently 0-1;
Y¹ and Y², at each occurrence, are independently selected from:

a bond, O, NR⁵⁶, C=O, C(=O)O, OC(=O)O, C(=O)NH-, C=NR⁵⁶, S, SO, SO₂, SO₃, NHC(=O), (NH)₂C(=O), (NH)₂C=S;

Z¹ is independently selected at each occurrence from a C₆-C₁₄ saturated, partially saturated, or aromatic carbocyclic ring system, substituted with 0-4

R⁵⁷; and a heterocyclic ring system, optionally substituted with 0-4 R⁵⁷;
R⁵⁵ and R⁵⁶ are independently selected at each occurrence from:

hydrogen; C₁-C₁₀ alkyl substituted with 0-5 R⁵⁷; alkaryl wherein the aryl is substituted with 0-5 R⁵⁷;

R⁵⁷ is independently selected at each occurrence from the group: hydrogen, OH, NHR⁵⁸, C(=O)R⁵⁸, OC(=O)R⁵⁸, OC(=O)OR⁵⁸, C(=O)OR⁵⁸, C(=O)NR⁵⁸-, C≡N, SR⁵⁸, SOR⁵⁸, SO₂R⁵⁸, NHC(=O)R⁵⁸, NHC(=O)NHR⁵⁸, NHC(=S)NHR⁵⁸; or, alternatively, when attached to an additional molecule Q, R⁵⁷ is independently selected at each occurrence from the group: O, NR⁵⁸, C=O, C(=O)O, OC(=O)O, C(=O)N-, C=NR⁵⁸, S, SO, SO₂, SO₃, NHC(=O), (NH)₂C(=O), (NH)₂C=S; and,

R⁵⁸ is independently selected at each occurrence from the group: hydrogen; C₁-C₆ alkyl; benzyl, and phenyl;
x and y are independently 1 or 2;

z is independently 1-4;

M_t is a transition metal radionuclide selected from the group: ^{99m}Tc, ¹⁸⁶Re and ¹⁸⁸Re;

C_{h'} is a radionuclide metal chelator coordinated to transition metal radionuclide M_t, and is independently selected at each occurrence, from the group: R⁴⁰N=N⁺=, R⁴⁰R⁴¹N-N=, R⁴⁰N=, and R⁴⁰N=N(H)-, wherein

R⁴⁰ is independently selected at each occurrence from the group: a bond to L_n, C₁-C₁₀ alkyl substituted with 0-3 R⁵², aryl substituted with 0-3 R⁵², cycloalkyl substituted with 0-3 R⁵², heterocycle substituted with 0-3 R⁵², heterocycloalkyl substituted with 0-3 R⁵², alkaryl substituted with 0-3 R⁵² and alkaryl substituted with 0-3 R⁵²;
R⁴¹ is independently selected from the group: hydrogen, aryl substituted with 0-3 R⁵², C₁-C₁₀ alkyl substituted with 0-3 R⁵², and a heterocycle substituted with 0-3 R⁵²;

R⁵² is independently selected at each occurrence from the group: a bond to L_n, =O, F, Cl, Br, I, -CF₃, -CN, -CO₂R⁵³, -C(=O)R⁵³, -C(=O)N(R⁵³)₂, -CHO, -CH₂OR⁵³, -OC(=O)R⁵³, -OC(=O)OR^{53a}, -OR⁵³, -OC(=O)N

$(R^{53})_2$, $-NR^{53}C(=O)R^{53}$, $-NR^{54}C(=O)OR^{53a}$, $-NR^{53}C(=O)N(R^{53})_2$, $-NR^{54}SO_2N(R^{53})_2$, $-NR^{54}SO_2R^{53a}$, $-SO_3H$, $-SO_2R^{53a}$, $-SR^{53}$, $-S(=O)R^{53a}$, $-SO_2N(R^{53})_2$, $-N(R^{53})_2$, $-NHC(=NH)NHR^{53}$, $-C(=NH)NHR^{53}$, $=NOR^{53}$, NO_2 , $-C(=O)NHOR^{53}$, $-C(=O)NHN(R^{53})R^{53a}$, $-OCH_2CO_2H$, 2-(1-morpholino)ethoxy;

R^{53} , R^{53a} , and R^{54} are each independently selected at each occurrence from the group: hydrogen, C_1 - C_6 alkyl, and a bond to L_n ;

A_{L1} is a first ancillary ligand selected from the group:

dioxygen ligand,
functionalized aminocarboxylate, and
halide;

A_{L2} is an ancillary ligand capable of stabilizing the radiopharmaceutical selected from the group:

A^9 and A^{10} - W - A^{11} ,

wherein:

A^9 is independently selected at each occurrence from the group:

$PR^{61}R^{62}R^{63}$ and $AsR^{61}R^{62}R^{63}$;

A^{10} and A^{11} are independently selected at each occurrence from the group:

$PR^{61}R^{62}$ and $AsR^{61}R^{62}$;

W is a spacer group selected from the group: C_1 - C_{10} alkyl substituted with 0-3 R^{70} , aryl substituted with 0-3 R^{70} , cycloalkyl substituted with 0-3 R^{70} , heterocycle substituted with 0-3 R^{70} , heterocycloalkyl substituted with 0-3 R^{70} , aralkyl substituted with 0-3 R^{70} and alkaryl substituted with 0-3 R^{70} ;

R^{61} , R^{62} , and R^{63} are independently selected at each occurrence from the group: C_1 - C_{10} alkyl substituted with 0-3 R^{70} , aryl substituted with 0-3 R^{70} , cycloalkyl substituted with 0-3 R^{70} , heterocycle substituted with 0-3 R^{70} , aralkyl substituted with 0-3 R^{70} , alkaryl substituted with 0-3 R^{70} , and arylalkaryl substituted with 0-3 R^{70} ;

R^{70} is independently selected at each occurrence from the group: F, Cl, Br, I, $-CF_3$, $-CN$, $-CO_2R^{71}$, $-C(=O)R^{71}$, $-C(=O)N(R^{71})_2$, $-CH_2OR^{71}$, $-OC(=O)R^{71}$, $-OC(=O)OR^{71a}$, $-OR^{71}$, $-OC(=O)N(R^{71})_2$, $-NR^{71}C(=O)R^{71}$, $-NR^{71}C(=O)OR^{71}$, $-NR^{71}C(=O)N(R^{71})_2$, SO_3^- , $-NR^{71}SO_2N(R^{71})_2$, $-NR^{71}SO_2R^{71a}$, $-SO_3H$, $-SO_2R^{71}$, $-S(=O)R^{71}$, $-SO_2N(R^{71})_2$, $-N(R^{71})_2$, $-N(R^{71})_3^+$, $-NHC(=NH)NHR^{71}$, $-C(=NH)NHR^{71}$, $=NOR^{71}$, NO_2 , $-C(=O)NHOR^{71}$, $-C(=O)NHN(R^{71})R^{71a}$, $-OCH_2CO_2H$; and

R^{71} and R^{71a} are independently selected at each occurrence from the group:

hydrogen and C_1 - C_6 alkyl; and

pharmaceutically acceptable salts thereof.

11. A method according to claim 10, wherein the cardiovascular disorder is selected from the group consisting of thromboembolic disease or atherosclerosis.

12. A method according to claim 11, wherein the thromboembolic disease results from the formation of blood clots and is selected from the group consisting of venous disorders, arterial disorders and pulmonary embolism.

13. A method according to claim 11, wherein the thromboembolic disease or atherosclerosis results from IIb/IIIa receptor activity, fibrinogen receptor activity, thrombin activity, fibrin binding activity, and lipoprotein receptor activity.

14. A method according to claim 12, wherein Q is selected from the group consisting of IIb/IIIa receptor ligands, laminin derivatives, hirudin-based peptides, atherosclerotic plaque binding peptides and fibrin binding peptides.

15. A method for radioimaging a mammal for the diagnosis of infection, inflammation or transplant rejection, comprising:

- (i) administering to said mammal an effective amount of a radiopharmaceutical; and
- (ii) scanning the mammal using a radioimaging device,

wherein the radiopharmaceutical is of the formula:




wherein:

Q is a biologically active molecule;

d' is 1 to 20;

L_n is a linking group of formula:



wherein:

M^1 is $-(CH_2)_gZ^1]_{g'}-(CR^{55}R^{56})_{g''}-$;

M^2 is $-(CR^{55}R^{56})_{g'''}-[Z^1(CH_2)_g]_{g''}-$;

g is independently 0-10;

g' is independently 0-1;

g'' is independently 0-10;

f is independently 0-10;

f' is independently 0-10;

f'' is independently 0-1;

Y^1 and Y^2 , at each occurrence, are independently selected from:

a bond, O, NR^{56} , C=O, C(=O)O, OC(=O)O, C(=O)NH-, C=NR⁵⁶, S, SO, SO₂, SO₃, NHC(=O), (NH)₂C(=O), (NH)₂C=S;

Z^1 is independently selected at each occurrence from a C₆-C₁₄ saturated, partially saturated, or aromatic carbocyclic ring system, substituted with 0-4

R⁵⁷; and a heterocyclic ring system, optionally substituted with 0-4 R⁵⁷;

R⁵⁵ and R⁵⁶ are independently selected at each occurrence from:

hydrogen; C₁-C₁₀ alkyl substituted with 0-5 R⁵⁷; alkaryl wherein the aryl is substituted with 0-5 R⁵⁷;

R⁵⁷ is independently selected at each occurrence from the group: hydrogen, OH, NHR⁵⁸, C(=O)R⁵⁸, OC(=O)R⁵⁸, OC(=O)OR⁵⁸, C(=O)OR⁵⁸, C(=O)NR⁵⁸-, C≡N, SR⁵⁸, SOR⁵⁸, SO₂R⁵⁸, NHC(=O)R⁵⁸, NHC(=O)NHR⁵⁸, NHC(=S)NHR⁵⁸; or, alternatively, when attached to an additional molecule Q, R⁵⁷ is independently selected at each occurrence from the group: O, NR⁵⁸, C=O, C(=O)O, OC(=O)O, C(=O)N-, C=NR⁵⁸, S, SO, SO₂, SO₃, NHC(=O), (NH)₂C(=O), (NH)₂C=S; and,

R⁵⁸ is independently selected at each occurrence from the group: hydrogen; C₁-C₆ alkyl; benzyl, and phenyl;

x and y are independently 1 or 2;

z is independently 1-4;

M_t is a transition metal radionuclide selected from the group: ^{99m}Tc, ¹⁸⁶Re and ¹⁸⁸Re;

C_h is a radionuclide metal chelator coordinated to transition metal radionuclide M_t , and is independently selected at each occurrence, from the group: R⁴⁰N=N⁺-, R⁴⁰R⁴¹N-N=, R⁴⁰N=, and R⁴⁰N=N(H)-, wherein

R⁴⁰ is independently selected at each occurrence from the group: a bond to L_n , C₁-C₁₀ alkyl substituted with 0-3 R⁵², aryl substituted with 0-3 R⁵², cycloalkyl substituted with 0-3 R⁵², heterocycle substituted with 0-3 R⁵², heterocycloalkyl substituted with 0-3 R⁵², alkaryl substituted with 0-3 R⁵² and alkaryl substituted with 0-3 R⁵²; R⁴¹ is independently selected from the group: hydrogen, aryl substituted with 0-3 R⁵², C₁-C₁₀ alkyl substituted with 0-3 R⁵², and a heterocycle substituted with 0-3 R⁵²;

R⁵² is independently selected at each occurrence from the group: a bond to L_n , =O, F, Cl, Br, I, -CF₃-, -CN, -CO₂R⁵³, -C(=O)R⁵³, -C(=O)N(R⁵³)₂, -CHO, -CH₂OR⁵³, -OC(=O)R⁵³, -OC(=O)OR^{53a}-, -OR⁵³, -OC(=O)N(R⁵³)₂-, -NR⁵³C(=O)R⁵³, -NR⁵⁴C(=O)OR^{53a}-, -NR⁵³C(=O)N(R⁵³)₂-, -NR⁵⁴SO₂N(R⁵³)₂-, -NR⁵⁴SO₂R^{53a}-, -SO₃H, -SO₂R^{53a}-, -SR⁵³-, -S(=O)R^{53a}-, -SO₂N(R⁵³)₂-, -N(R⁵³)₂-, -NHC(=NH)NHR⁵³-, -C(=NH)NHR⁵³-, =NOR⁵³, NO₂-, -C

(=O)NHOR⁵³, -C(=O)NHN⁵³R^{53a}, -OCH₂CO₂H, 2-(1-morpholino)ethoxy;

R⁵³, R^{53a}, and R⁵⁴ are each independently selected at each occurrence from the group: hydrogen, C₁-C₆ alkyl, and a bond to L_n;

A_{L1} is a first ancillary ligand selected from the group:

dioxygen ligand,
functionalized aminocarboxylate, and
halide;

A_{L2} is an ancillary ligand capable of stabilizing the radiopharmaceutical selected from the group:

A⁹ and A¹⁰-W-A¹¹,

wherein:

A⁹ is independently selected at each occurrence from the group:

PR⁶¹R⁶²R⁶³ and AsR⁶¹R⁶²R⁶³;

A¹⁰ and A¹¹ are independently selected at each occurrence from the group:

PR⁶¹R⁶² and AsR⁶¹R⁶²;

W is a spacer group selected from the group: C₁-C₁₀ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰, cycloalkyl substituted with 0-3 R⁷⁰, heterocycle substituted with 0-3 R⁷⁰, heterocycloalkyl substituted with 0-3 R⁷⁰, aralkyl substituted with 0-3 R⁷⁰ and alkaryl substituted with 0-3 R⁷⁰;

R⁶¹, R⁶², and R⁶³ are independently selected at each occurrence from the group: C₁-C₁₀ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰, cycloalkyl substituted with 0-3 R⁷⁰, heterocycle substituted with 0-3 R⁷⁰, aralkyl substituted with 0-3 R⁷⁰, alkaryl substituted with 0-3 R⁷⁰, and arylalkaryl substituted with 0-3 R⁷⁰;

R⁷⁰ is independently selected at each occurrence from the group: F, Cl, Br, I, -CF₃, -CN, -CO₂R⁷¹, -C(=O)R⁷¹, -C(=O)N(R⁷¹)₂, -CH₂OR⁷¹, -OC(=O)R⁷¹, -OC(=O)OR^{71a}, -OR⁷¹, -OC(=O)N(R⁷¹)₂, -NR⁷¹C(=O)R⁷¹, -NR⁷¹C(=O)OR⁷¹, -NR⁷¹C(=O)N(R⁷¹)₂, SO₃⁻, -NR⁷¹SO₂N(R⁷¹)₂, -NR⁷¹SO₂R^{71a}, -SO₃H, -SO₂R⁷¹, -S(=O)R⁷¹, -SO₂N(R⁷¹)₂, -N(R⁷¹)₂, -N(R⁷¹)₃⁺, -NHC(=NH)NHR⁷¹, -C(=NH)NHR⁷¹, =NOR⁷¹, NO₂, -C(=O)NHOR⁷¹, -C(=O)NHN⁷¹R^{71a}, -OCH₂CO₂H; and

R⁷¹ and R^{71a} are independently selected at each occurrence from the group:

hydrogen and C₁-C₆ alkyl; and

pharmaceutically acceptable salts thereof.

16. A method according to claim 15, wherein:

Q is selected from the group consisting of:

leukocyte binding peptides and chemotactic peptides.

17. A method for radioimaging a mammal for the diagnosis of cancer, comprising:

- (i) administering to said mammal an effective amount of a radiopharmaceutical; and
- (ii) scanning the mammal using a radioimaging device,

wherein the radiopharmaceutical is of the formula:




wherein:

Q is a biologically active molecule;

d' is 1 to 20;

L_n is a linking group of formula:

5



wherein:

10

M¹ is $-(CH_2)_gZ^1]_{g'}-(CR^{55}R^{56})_{g''}-$;

M² is $-(CR^{55}R^{56})_{g'''}-[Z^1(CH_2)_g]_{g'}$;

g is independently 0-10;

g' is independently 0-1;

g'' is independently 0-10;

15

f is independently 0-10;

f' is independently 0-10;

f'' is independently 0-1;

Y¹ and Y², at each occurrence, are independently selected from:

20

a bond, O, NR⁵⁶, C=O, C(=O)O, OC(=O)O, C(=O)NH-, C=NR⁵⁶, S, SO, SO₂, SO₃, NHC(=O), (NH)₂C(=O), (NH)₂C=S;

Z¹ is independently selected at each occurrence from a C₆-C₁₄ saturated, partially saturated, or aromatic carbocyclic ring system, substituted with 0-4

25

R⁵⁷; and a heterocyclic ring system, optionally substituted with 0-4 R⁵⁷;

R⁵⁵ and R⁵⁶ are independently selected at each occurrence from:

hydrogen; C₁-C₁₀ alkyl substituted with 0-5 R⁵⁷; alkaryl wherein the aryl is substituted with 0-5 R⁵⁷;

30

R⁵⁷ is independently selected at each occurrence from the group: hydrogen, OH, NHR⁵⁸, C(=O)R⁵⁸, OC(=O)R⁵⁸, OC(=O)OR⁵⁸, C(=O)OR⁵⁸, C(=O)NR⁵⁸-, C≡N, SR⁵⁸, SOR⁵⁸, SO₂R⁵⁸, NHC(=O)R⁵⁸, NHC(=O)NHR⁵⁸, NHC(=S)NHR⁵⁸; or, alternatively, when attached to an additional molecule Q, R⁵⁷ is independently selected at each occurrence from the group: O, NR⁵⁸, C=O, C(=O)O, OC(=O)O, C(=O)N-, C=NR⁵⁸, S, SO, SO₂, SO₃, NHC(=O), (NH)₂C(=O), (NH)₂C=S; and,

35

R⁵⁸ is independently selected at each occurrence from the group: hydrogen; C₁-C₆ alkyl; benzyl, and phenyl; x and y are independently 1 or 2;

z is independently 1-4;

M_i is a transition metal radionuclide selected from the group: ^{99m}Tc, ¹⁸⁶Re and ¹⁸⁸Re;

40

C_h is a radionuclide metal chelator coordinated to transition metal radionuclide M_i, and is independently selected at each occurrence, from the group: R⁴⁰N=N⁺=, R⁴⁰R⁴¹N-N=, R⁴⁰N=, and R⁴⁰N=N(H)-, wherein

R⁴⁰ is independently selected at each occurrence from the group: a bond to L_n, C₁-C₁₀ alkyl substituted with 0-3 R⁵², aryl substituted with 0-3 R⁵², cycloalkyl substituted with 0-3 R⁵², heterocycle substituted with 0-3 R⁵², heterocycloalkyl substituted with 0-3 R⁵², aralkyl substituted with 0-3 R⁵² and alkaryl substituted with 0-3 R⁵²;

45

R⁴¹ is independently selected from the group: hydrogen, aryl substituted with 0-3 R⁵², C₁-C₁₀ alkyl substituted with 0-3 R⁵², and a heterocycle substituted with 0-3 R⁵²;

R⁵² is independently selected at each occurrence from the group: a bond to L_n, =O, F, Cl, Br, I, -CF₃, -CN, -CO₂R⁵³, -C(=O)R⁵³, -C(=O)N(R⁵³)₂, -CHO, -CH₂OR⁵³, -OC(=O)R⁵³, -OC(=O)OR^{53a}, -OR⁵³, -OC(=O)N(R⁵³)₂, -NR⁵³C(=O)R⁵³, -NR⁵⁴C(=O)OR^{53a}, -NR⁵³C(=O)N(R⁵³)₂, -NR⁵⁴SO₂N(R⁵³)₂, -NR⁵⁴SO₂R^{53a}, -SO₃H, -SO₂R^{53a}, -SR⁵³, -S(=O)R^{53a}, -SO₂N(R⁵³)₂, -N(R⁵³)₂, -NHC(=NH)NHR⁵³, -C(=NH)NHR⁵³, =NOR⁵³, NO₂, -C(=O)NHR⁵³, -C(=O)NHNHR⁵³R^{53a}, -OCH₂CO₂H, 2-(1-morpholino)ethoxy;

50

R⁵³, R^{53a}, and R⁵⁴ are each independently selected at each occurrence from the group: hydrogen, C₁-C₆ alkyl, and a bond to L_n;

A_{L1} is a first ancillary ligand selected from the group:

55

dioxygen ligand,
functionalized aminocarboxylate, and
halide;

A_{L2} is an ancillary ligand capable of stabilizing the radiopharmaceutical selected from the group:

A⁹ and A¹⁰-W-A¹¹,

wherein:

A⁹ is independently selected at each occurrence from the group:

PR⁶¹R⁶²R⁶³ and AsR⁶¹R⁶²R⁶³;

A¹⁰ and A¹¹ are independently selected at each occurrence from the group:

PR⁶¹R⁶² and AsR⁶¹R⁶²;

W is a spacer group selected from the group: C₁-C₁₀ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰, cycloalkyl substituted with 0-3 R⁷⁰, heterocycle substituted with 0-3 R⁷⁰, heterocycloalkyl substituted with 0-3 R⁷⁰, aralkyl substituted with 0-3 R⁷⁰ and alkaryl substituted with 0-3 R⁷⁰;

R⁶¹, R⁶², and R⁶³ are independently selected at each occurrence from the group: C₁-C₁₀ alkyl substituted with 0-3 R⁷⁰, aryl substituted with 0-3 R⁷⁰, cycloalkyl substituted with 0-3 R⁷⁰, heterocycle substituted with 0-3 R⁷⁰, aralkyl substituted with 0-3 R⁷⁰, alkaryl substituted with 0-3 R⁷⁰, and arylalkaryl substituted with 0-3 R⁷⁰;

R⁷⁰ is independently selected at each occurrence from the group: F, Cl, Br, I, -CF₃, -CN, -CO₂R⁷¹, -C(=O)R⁷¹, -C(=O)N(R⁷¹)₂, -CH₂OR⁷¹, -OC(=O)R⁷¹, -OC(=O)OR^{71a}, -OR⁷¹, -OC(=O)N(R⁷¹)₂, -NR⁷¹C(=O)R⁷¹, -NR⁷¹C(=O)OR⁷¹, -NR⁷¹C(=O)N(R⁷¹)₂, SO₃⁻, -NR⁷¹SO₂N(R⁷¹)₂, -NR⁷¹SO₂R^{71a}, -SO₃H, -SO₂R⁷¹, -S(=O)R⁷¹, -SO₂N(R⁷¹)₂, -N(R⁷¹)₂, -N(R⁷¹)₃⁺, -NHC(=NH)NHR⁷¹, -C(=NH)NHR⁷¹, =NOR⁷¹, NO₂, -C(=O)NHOR⁷¹, -C(=O)NHNR⁷¹R^{71a}, -OCH₂CO₂H; and

R⁷¹ and R^{71a} are independently selected at each occurrence from the group:

hydrogen and C₁-C₆ alkyl; and
pharmaceutically acceptable salts thereof.


18. A method according to claim 17, wherein the cancer is associated with Platelet Factor 4 activity.

19. A method according to claim 17, wherein the cancer is associated with a growth factor selected from the group consisting of PDGF, EGF, FGF, TNF MCSF and II1-8.


20. A method according to claim 17, wherein:

Q is selected from the group consisting of somatostatin analogs and selectin binding peptides

FIGURE 1



HPLC Chromatogram of Example 1 of this application
using Method 1.



HPLC Chromatogram of Example 1 of this application
using Method 2.

Figure 2




FIGURE 3

