

(19)



(11)

EP 1 197 227 B9

(12)

CORRECTED EUROPEAN PATENT SPECIFICATION

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Description Paragraph(s) 22

(48) Corrigendum issued on:
11.06.2008 Bulletin 2008/24

(45) Date of publication and mention
 of the grant of the patent:
12.09.2007 Bulletin 2007/37

(21) Application number: **00940773.5**

(22) Date of filing: **21.06.2000**

(51) Int Cl.:
A61K 45/06 ^(2006.01) **A61K 51/00** ^(2006.01)
A61K 47/06 ^(2006.01)

(86) International application number:
PCT/JP2000/004039

(87) International publication number:
WO 2000/078352 (28.12.2000 Gazette 2000/52)

(54) **METHOD OF THE ADMINISTRATION OF DRUGS HAVING BINDING AFFINITY WITH PLASMA PROTEIN AND PREPARATION TO BE USED IN THE METHOD**

VERFAHREN ZUR VERABREICHUNG VON ARZNEIMITTELN MIT BINDUNGS-AFFINITÄT ZUM PLASMAPROTEIN UND VERWENDUNG DER ZUSAMMENSETZUNG IN DEM VERFAHREN

PROCEDE D'ADMINISTRATION DE MEDICAMENTS POSSEDANT UNE AFFINITE DE LIAISON AUX PROTEINES PLASMATQUES ET PREPARATION UTILISEE POUR LA MISE EN OEUVRE DUDIT PROCEDE

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

(30) Priority: **21.06.1999 JP 17351499**

(43) Date of publication of application:
17.04.2002 Bulletin 2002/16

(73) Proprietor: **NIHON MEDI-PHYSICS CO., LTD.**
Nishinomiya-shi,
Hyogo 662-0918 (JP)

(72) Inventors:
 • **KAWAI, Keiichi**
Miyazaki-gun, Miyazaki 889-1601 (JP)
 • **TAKAMURA, Norito**
Miyazaki-shi, Miyazaki 880-0925 (JP)
 • **NISHII, Ryuichi**
Miyazaki-shi, Miyazaki 880-0916 (JP)

(74) Representative: **Keen, Celia Mary**
J.A. Kemp & Co.
14 South Square
Gray's Inn
London WC1R 5JJ (GB)

(56) References cited:
WO-A-8/39037 DE-A- 19 648 629
US-A- 4 642 284 US-A- 4 976 950
US-A- 5 792 444

- PRITCHARD J F ET AL: "PLASMA PROTEIN BINDING OF BEPRIDIL" JOURNAL OF CLINICAL PHARMACOLOGY, vol. 25, no. 5, 1985, pages 347-353, XP001083994 ISSN: 0091-2700
- BERTUCCI C ET AL: "The binding of 5-fluorouracil to native and modified human serum albumin: UV, CD, and 1H and 19F NMR investigation" JOURNAL OF PHARMACEUTICAL AND BIOMEDICAL ANALYSIS, NEW YORK, NY, US, vol. 13, no. 9, August 1995 (1995-08), pages 1087-1093, XP002101869 ISSN: 0731-7085

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 197 227 B9

- MASON R W ET AL: "IN-VITRO DISPLACEMENT OF INDOMETHACIN FROM PLASMA PROTEIN BINDING BY IBUPROFEN PHENYL BUTAZONE AND SALICYLATE" PROCEEDINGS OF THE UNIVERSITY OF OTAGO MEDICAL SCHOOL, vol. 52, no. 3, 1974, pages 49-50, XP009000051 ISSN: 0301-6331
- KAWAI K. ET AL.: 'Competitive displacement of 99mTc-MAG3 serum protein binding in in-vitro and in-vivo' JOURNAL OF LABELLED COMPOUNDS AND RADIOPHARMACEUTICALS vol. 42, no. SUPPL. 1, June 1999, pages S584 - S586, XP002932501
- BUBECK BERND ET AL.: 'Pharmacokinetics of technetium-99m-MAG3 in humans' THE JOURNAL OF NUCLEAR MEDICINE vol. 31, no. 8, 1990, pages 2285 - 1293, XP002932508
- MATSUI,M. ET AL.: 'Application of fractional uptake method to 99mTc-ECD SPECT for quantification of brain perfusion using cardiac output index' KOBE DAIGAKU IGAKUBU KIYO (DEPARTMENT OF RADIOLOGY, KOBE UNIVERSITY SCHOOL OF MEDICINE) vol. 58, no. 4, 1998, pages 191 - 196, XP002932502
- IVARSEN,PER RAMLOEV ET AL.: 'Displacement of bilirubin from adult and newborn serum albumin by a drug and fatty acid' DEV. PHARMACOL. THER. vol. 12, no. 1, 1989, pages 19 - 29, XP002932503
- BRIAND C. ET AL.: 'Study of the interaction between human serum albumin and some cephalosporins' MOL. PHARMACOL. vol. 21, no. 1, 1982, pages 92 - 99, XP002932504
- SEMMES,ROBIN L.O. ET AL.: 'Nonlinear binding of valproic acid (VPA) and E-delta2-valproic acid to rat plasma proteins' PHARM. RES. vol. 7, no. 5, 1990, pages 461 - 467, XP002932505
- LOCKWOOD,GRAHAM F. ET AL.: 'Pharmacokinetics of ibuprofen in man-III: Plasma protein binding' J. PHARMACOKINET. BIOPHARM. vol. 11, no. 5, 1983, pages 469 - 482, XP002932506
- ABDEL-RAHMAN M. ET AL.: 'Interaction between verapamil and vincristine binding to plasma proteins' INTERNATIONAL JOURNAL OF CLINICAL PHARMACOLOGY, THERAPY AND TOXICOLOGY vol. 30, no. 11, 1992, pages 536 - 537, XP002932507

Description

[0001] The present invention is based on the administration of drugs with binding affinity for plasma protein and drugs regulating the effective ingredient dose of drugs with binding affinity for plasma protein; and a pharmaceutical preparation whereby the effective ingredient dose of drugs with binding affinity for plasma protein is regulated.

[0002] Generally, drugs administered for the purpose of medical treatment or diagnosis once go through the systemic blood circulation, and then take the process of absorption, distribution, metabolism, excretion and the like. In the process of absorption and distribution, the drug moves along on flow of the blood, while it transfers to each spaces of intravascular, interstice and intracellular by diffusion and transportation of a free drug being in the state of unbound form with proteins, and finally the drug arrives at the active region of target. When movement of the drug reaches a steady state, then the free drug concentration in each space become uniform, thus the whole pattern of the concentration of the drug is determined by the binding level with proteins. Hence, in accordance with the property, a drug *in vivo*, may partially exists in the form of reversible binding state with biopolymers such as plasma proteins. Generally, drugs permeable through capillary wall or cell membrane are free drugs, therefore, the transfer of such free drugs being unbound with plasma proteins to the active region of target may be greatly influenced by the binding level with plasma proteins.

[0003] For example, mercaptoacetylglycylglycylglycine labeled with 99m-technetium ($^{99m}\text{Tc-MAG}_3$) is widely-used in renal scintigraphy, especially the renal plasma flow can be effectively exhibited by its efficient renal extraction and renal tubular secretion. It is known that about 90% of $^{99m}\text{Tc-MAG}_3$ binds to plasma protein in an ordinary clinical dose (Bubeck B. et al., J. Nucl. Med., 31, 1285-1295, 1990). If the binding of $^{99m}\text{Tc-MAG}_3$ with plasma protein is inhibited by drugs having high binding affinity to the same binding site on protein with $^{99m}\text{Tc-MAG}_3$, then more clear renal imaging can be obtained in the earlier stage after the administration, thus it may be thought that the dose of radioactivity to the patient can be reduced at the same time.

[0004] On the contrary, if the binding of drugs with plasma protein is increased, then the concentration of the free drugs in the blood can be kept in lower level for long period, therefore, it may be possible to achieve continuous appearance of pharmacological effects.

[0005] However, at the present stage, little is known as the research work for improving therapeutic effect or diagnostic effect of the drugs by regulating the concentrations of the free drugs, using the binding affinity of the second drug with plasma proteins.

[0006] Set against the above-mentioned problems, the present invention provides the use of a second drug in the manufacture of a medicament to regulate the binding of a first drug to a plasma protein, wherein the first drug is a radiodiagnostic drug for *in vivo* use or a radiotherapeutic drug for *in vivo* use and has one group selected from bisaminothiol or its derivatives, monoaminomonoamidobisthiol or its derivatives, bisamidobisthiol or its derivatives, mercaptoacetylglycylglycylglycine or its derivatives, hexamethylpropyleneamineoxime or its derivatives, ethylenebis[bis(2-ethoxyethyl)phosphine] (tetrafosmin) or its derivatives, 2,3-dimercaptosuccinic acid or its derivatives, ethylenecysteine dimer derivatives, methoxyisobutylisonitrile derivatives, polyamine derivatives, pyridoxylideneamine derivatives, methylene diphosphonate, hydroxymethylene diphosphonate derivatives, β -methyl- ω -phenylpentadecanoic acid or its derivatives, N-isopropyl-amphetamine, hippuric acid, benzylguanidine, and tropane derivatives, the group being radiolabeled with one nuclide selected from 11-carbon (^{11}C), 15-oxygen (^{15}O), 18-fluorine (^{18}F), 32-phosphorus (^{32}P), 59-iron (^{59}Fe), 67-copper (^{67}Cu), 67-gallium (^{67}Ga), 81m-krypton (^{81m}Kr), 81-rubidium (^{81}Rb), 89-strontium (^{89}Sr), 90-yttrium (^{90}Y), 99m-technetium (^{99m}Tc), 111-indium (^{111}In), 123-iodine (^{123}I), 125-iodine (^{125}I), 131-iodine (^{131}I), 133-xenon (^{133}Xe), 117m-tin (^{117m}Sn), 153-samarium (^{153}Sm), 186-rhenium (^{186}Re), 188-rhenium (^{188}Re), 201-thallium (^{201}Tl), 212-bismuth (^{212}Bi), 213-bismuth (^{213}Bi) and 211-astatine (^{211}At);

the second drug is selected from bucolome, cefazolin, etoposide, phenylbutazone, aspirin, salicylic acid, ceftriaxone, sulfamethizole, valproic acid, nabumetone, 6-methoxy-2-naphthyl acetic acid, ibuprofen, probenecid, dansyl-L-asparagine (DNSA), verapamil and disopyramide;

and the first drug and the second drug have binding affinity for said plasma protein.

[0007] The second drug is administered simultaneously with the first drug or before or after the administration of the first drug.

[0008] In a preferred embodiment, the second drug has binding affinity to the same binding sites on the plasma protein to which the first drug has binding affinity. The medicament comprising the second drug may be administered before, after or simultaneously with the administration of the first drug, and such administration timing of the medicament comprising the second drug can be suitably selected in connection with the timing when the free drug concentration of the first drug reaches to the level so as to obtain an adequate effect. Additionally, a single drug may be used as the second drug, or more than one of the said second drugs may be used, in which case synergistic effect can be expected.

[0009] Each of the first drug and the second drug may be separately filled in a container, and may be prepared as kit form for supply. Accordingly, the present invention further provides a product which comprises

(a) a second drug selected from bucolome, cefazolin, etoposide, phenylbutazone, aspirin, salicylic acid, ceftriaxone,

sulfamethizole, valproic acid, nabumetone, 6-methoxy-2-naphthyl acetic acid, ibuprofen, probenecid, dansyl-L-asparagine, verapamil and disopyramide;

(b) a first drug which is a radiodiagnostic drug for *in vivo* use or a radiotherapeutic drug for *in vivo* use and has one group selected from bisaminothiol or its derivatives, monoaminomonoamidobisthiol or its derivatives, bisamido-bisthiol or its derivatives, mercaptoacetyl-glycyl-glycyl- glycine or its derivatives, hexamethylpropylene-amineoxime or its derivatives, ethylenebis[bis(2-ethoxyethyl)phosphine](tetrofosmin) or its derivatives, 2,3-dimercaptosuccinic acid or its derivatives, ethylenecysteine dimer derivatives, methoxyisobutylisonitrile derivatives, polyamine derivatives, pyridoxylideneamine derivatives, methylene diphosphonate, hydroxymethylene diphosphonate derivatives, β -methyl- ω -phenylpentadecanoic acid or its derivatives, N-isopropyl-amphetamine, hippuric acid, benzylguanidine, and tropane derivatives, the group being radiolabelled with one nuclide selected from 11-carbon (^{11}C), 15-oxygen (^{15}O), 18-fluorine (^{18}F), 32-phosphorus (^{32}P), 59-iron (^{59}Fe), 67-copper (^{67}Cu), 67-gallium (^{67}Ga), 81m-krypton ($^{81\text{m}}\text{Kr}$), 81-rubidium (^{81}Rb), 89-strontium (^{89}Sr), 90-yttrium (^{90}Y), 99m-technetium ($^{99\text{m}}\text{Tc}$), 111-indium (^{111}In), 123-iodine (^{123}I), 125-iodine (^{125}I), 131-iodine (^{131}I), 133-xenon (^{133}Xe), 117m-tin ($^{117\text{m}}\text{Sn}$), 153-samarium (^{153}Sm), 186-rhenium (^{186}Re), 188-rhenium (^{188}Re), 201-thallium (^{201}Tl), 212-bismuth (^{212}Bi), 213-bismuth (^{213}Bi) and 211-astatine (^{211}At); which first drug has binding affinity for a plasma protein for which said second drug has binding affinity; for simultaneous, separate or sequential use in regulating the binding affinity of the first drug to the plasma protein.

[0010] In case of such a kit form with separate containers, they may be administered simultaneously by mixing together when used, or each one of the first drug and the second drug can be administered at different times separately or by different routes. The first drug may be a commercially available pharmaceutical.

[0011] In the accompanying drawings:

Fig. 1 shows the free fraction of $^{99\text{m}}\text{Tc}$ -MAG₃ in human plasma in the presence of site specific agent.

Fig. 2 shows the free fraction of $^{99\text{m}}\text{Tc}$ -MAG₃ in rat plasma in the presence of site specific agent.

Fig. 3 shows the effect of bucolome on blood clearance of $^{99\text{m}}\text{Tc}$ -MAG₃ in rat.

Fig. 4 shows the effect of bucolome on the free fraction of $^{99\text{m}}\text{Tc}$ -MAG₃ in rat blood after administration of bucolome.

Fig. 5 shows the effect of bucolome on the accumulation of $^{99\text{m}}\text{Tc}$ -MAG₃ in rat kidney after administration of bucolome.

Fig. 6 shows the effect of bucolome loading on the biodistribution of $^{99\text{m}}\text{Tc}$ -MAG₃ in rat.

Fig. 7 shows the renogram of $^{99\text{m}}\text{Tc}$ -MAG₃ in rat.

[0012] When the second drug as defined above is administered simultaneously with the first drug, or before or after the administration of the first drug, then competitive displacement will take place at the binding site, thus it can be thought that the first drug may be released in a higher concentration (displacement effect). Therefore, it can be expected that the higher pharmacological activity of the first drug can be obtained as compared with the case that the first drug is administered singly. On the contrary, when the binding fraction of the first drug to plasma protein increases by the effect of the second drug (reducing effect of the free drug concentration), continuous appearance of pharmacological effect of the first drug can be expected to achieve by keeping the free fraction of the first drug in blood at a lower level for long period.

[0013] In the present invention, the first drug with binding affinity for plasma protein may be either one of therapeutic agent or diagnostic agent, as long as it meets the purpose of administration.

[0014] Regardless of therapeutic or diagnostic purpose, in case of obtaining the above-mentioned displacement effect, the second drug may be preferably selected from those having competitive binding affinity for the same plasma protein as the first drug has; increasing the free fraction of the first drug by the binding inhibition of the first drug with plasma protein; having the affinity for the same binding site of the first drug on plasma protein; and having the higher binding affinity for plasma protein.

[0015] On the contrary, in case of obtaining the above-mentioned reducing effect of the free drug concentration, the object is achieved by selecting the second drug from those having effect to increase the binding affinity of the first drug for plasma protein by the second drug bound to the same plasma proteins.

[0016] At the present, a report relating to research for clarifying the entity of the reducing effect of the free drug concentration has not been found yet. While, it can be considered that said reducing effect may appear, for example by a mechanism similar to the allosteric effect of an enzyme, and surprisingly, it was found that the binding affinity for the plasma protein could be increased by using the combination of the drugs shown in Example 8 of the present invention.

[0017] Regarding dose forms of the drug, in the case that the first drug and second drug are administered simultaneously without necessarily considering any chemical change such as decomposition thereof by mixing together, it is possible to supply a pharmaceutical product prepared by mixing the first drug with the second drug. In such a mixed-type of pharmaceutical preparation, medicinally acceptable ingredients, such as pH-adjusting agents, inorganic salts for adjusting the osmotic pressure, stabilizing agents for stabilizing each one of these ingredients may be added thereto. The mixed-type of pharmaceutical preparations can be processed into the suitable dose form, for example a liquid form preparation, a lyophilized form preparation and the like, in consideration of the constitutional ingredients, preservation

stability thereof, etc. Further, the first drug and the second drug may be separately filled in a container and prepared in kit form for supply. Similar to the mixed-type preparation, medicinally acceptable ingredients, such as stabilizing agents or the like may be added to each one of these separate type of drugs, and in consideration of administration method, stabilization and the like, these separate type of the drugs can be processed into the suitable form of preparations, such as liquid form preparation, a lyophilized form preparation and the like.

[0018] In case of the kit form mentioned above, the first drug and the second drug can be administered separately, or can be administered simultaneously by mixing together at the time of use. Especially, in the case of predicting changes of quality of the product, such as decomposition of the ingredients during the storage after mixing the first drug and the second drug, and in the case that these drugs are administered by different route, or in the case that these drugs are administered necessarily in different timings, the above-mentioned kit form in which the first drug and the second drugs are filled in separate containers are useful.

[0019] Generally, as the plasma proteins bound to drug, human serum albumin (HSA), α_1 -acidic glycoprotein (AGP), γ -globulin, lipoprotein and the like are exemplified, and many drugs may bind to HSA or AGP. In selecting the second drug, for example when the first drug has the property of mainly binding to HSA, it may be preferably selected from an acidic drug having the binding affinity for HSA. When the first drug the property of binding to AGP, it may be preferably selected from a basic drug having the binding affinity for AGP. Further, in the case that the first drug has the affinity for plural plasma proteins or has the affinity for different binding sites on the single protein, the use of plural drugs as second drugs may be effective. Furthermore, in case of selecting the second drug, other properties than the binding affinity with the above-mentioned plasma protein should be considered, such as clinically acceptable appearance of the original pharmacological activity, a broad range of usual dose, and maintenance of high blood concentration after administration, etc.

[0020] Administration timing of the second drug may be either simultaneously with the first drug or before or after the administration of the first drug, thus the timing is selected suitably so as to obtain the effect to meet the administration purpose of the first drug. Administration route of the drugs may be suitably selected from either one of intravenous injection, intraarterial injection, subcutaneous injection, lymphaginal injection or oral administration.

[0021] Specifically, HSA has three specific binding sites such as site I, site II and site III on its molecule. Examples of the second drug with binding specificity at the site I, are bucolome (5-n-butyl-1-cyclohexyl-2,4,6-trioxoperhydropyrimidine), cefazolin (7-[1-(H)-tetrazolylacetamido]-3-[2-(5-methyl-1,3,4-thiazolyl)thiomethyl]-3-cephem-4-carboxylate), phenylbutazone (1,2-diphenyl-3,5-dioxo-4-n-butylpyrazolidine), valproic acid (sodium 2-propylpentanoate), aspirin (2-acetoxybenzoic acid), salicylic acid (O-hydroxybenzoic acid), ceftriaxone (disodium (6R,7R)-7-[2-amino-4-thiazoyl]-2-methoxyiminoacetamide)-3-(2,5-dihydro-2-methyl-6-oxide-5-oxo-1,2,4-triazin-3-ylthiomethyl)-8-oxo-5-thia-1-azobicyclo [4.2.0]octo-2-ene-2-carboxylate), sulfamethizol (N-(5-methyl-1,3,4-thiadiazol-2-yl)sulfanylamide), canrenoic acid (17-hydroxy-3-oxo-17 α -pregna-4,6-dien-21-carboxylate), and dansyl-L-asparagine. Examples of the second drug with binding specificity at the site II, are ibuprofen (2-(4-isobutylphenyl)propionic acid), nabumetone (4-(6-methoxy-2-naphthyl)-2-butanone (6-methoxy-2-naphthylacetic acid, which is a metabolite of nabumetone, shows binding specificity at the site II) and probenecid (4-(N,N-dipropylsulfamoyl)benzoic acid), etc. Further, etoposide ((5S, 5aR, 8aR, 9S)-9-[(4,6,0-(R)-ethylidene- β -D-glucopyranosyl)oxy]-5,8,8a,9-tetrahydro-5-(4-hydroxy-3,5-dimethoxyphenylisobenzofuro [5,6-f][1,3]benzodioxol-6(5aH)-one) also has binding specificity for HSA, though the binding site on the HSA has not been assigned. As the second drug with binding specificity for AGP, the following drugs can be exemplified, disopyramide (α -(2-diisopropylaminoethyl)- α -phenyl-2-pyridineacetamide), verapamil (α -[3-[[2-(3,4-dimethoxyphenyl)ethyl]-methylamino]propyl]-3,4-dimethoxy- α -(1-methylethyl)benzeneacetonitrile) and propranolol(1-isopropylamino-3-(1-naphthyl-oxy)-2-propanol), etc.

[0022] As compounds, such as chelating group or receptor ligand, of radio-therapeutic drug for *in vivo* use or radiodiagnostic drug for *in vivo* use, both having binding affinity for plasma protein and being labeled with radioactive nuclides, following compounds can be exemplified; mercaptoacetylglycylglycylglycine (MAG₃) or its derivatives, hexamethylpropyleneaminoxime(HMPAO) or its derivatives, ethylenebis[bis(2-ethoxyethyl)phosphine] (tetrofosmin) or its derivatives, 2,3-dimercaptosuccinic acid (DMSA) or its derivatives, ethylene cysteine dimer (ECD) derivatives such as N,N'-ethylene-L-cystein diethylester and the like, methoxyisobutylisonitrile (MIBI) derivatives, polyamine derivatives such as diethylenetriaminepentaacetic acid (DTPA) and the like, pyridoxylideneamine derivatives such as pyridoxylideneisoleucine and the like; other chelating groups which can form complex with radioactive metals such as methylene diphosphonate (MDP), hydroxymethylene diphosphonate (HMDP) and the like; and compounds labeled with radioactive iodine such as β -methyl-p-iodophenylpentadecanoic acid (BMIPP), N-isopropyl-p-iodoamphetamine (IMP), iodinated hippuric acid (OIH), 3-iodobenzylguanidine (MIBG), tropane derivatives such as N-(3-fluoropropyl)-2 β -carbomethoxy-3 β -(4-iodophenyl)nortropane (FP-CIT), N-methyl-2 β -carbomethoxy-3 β -(4-iodophenyl)nortropane (CIT) and the like. As to the diagnostic purposes, 18-fluorine (¹⁸F), 99m-technetium (^{99m}Tc), 67-gallium (⁶⁷Ga), 111-indium (¹¹¹In), 123-iodine (¹²³I), 131-iodine (¹³¹I) and the like are frequently used.

[0023] 99m-Technetium complex of MAG₃(^{99m}Tc-MAG₃) is a radiopharmaceutical for *in vivo* use and is widely used for the purpose of diagnosis of renopathy and uropathy, because it possesses accumulation property to the kidney. It

is known that about 90% of $^{99m}\text{Tc-MAG}_3$ bind to plasma protein. For this reason, *in vitro* study was conducted by using $^{99m}\text{Tc-MAG}_3$ as the first drug, the serum as the plasma protein, wherein the blood cells and blood coagulation factors are removed, and several pharmaceuticals with the binding affinity for serum proteins as the second drug. As the result, when bucolome, valproic acid, warfarin or the like was added, then displacement of $^{99m}\text{Tc-MAG}_3$ occurred either in human serum albumin or in rat serum albumin, thus the free fraction of $^{99m}\text{Tc-MAG}_3$ in serum albumin increased. In case of bucolome, the free fraction of $^{99m}\text{Tc-MAG}_3$ particularly increased (Table 1). Fig. 5 shows time course of accumulation of $^{99m}\text{Tc-MAG}_3$ in the rat kidney after administering 20 mg/kg of Bucolome. Fig. 6 shows the biodistributions in rats at 10 minutes after the administration of $^{99m}\text{Tc-MAG}_3$. In this case, 10 minutes before the administration of $^{99m}\text{Tc-MAG}_3$, 100 mg/kg of bucolome was administered. These results show that the amount of free $^{99m}\text{Tc-MAG}_3$ was increased by bucolome loading, and rapid clearance from the blood and accumulation of $^{99m}\text{Tc-MAG}_3$ into the kidney took place.

[0024] Regarding ^{99m}Tc -technetium complex of diethyl ester of N,N'-ethylene-L-cystein ($^{99m}\text{Tc-ECD}$), which is a radiopharmaceutical used for scintigraphy of regional cerebral blood flow, in the *in vitro* experiment by using a human serum, the displacement effect was observed by adding Etoposide (cf. Example 4 and Table 8).

[0025] For the purpose to prove the displacement effect on organic compounds, *in vitro* and *in vivo* experiments were conducted by using N-isopropyl-p-iodoamphetamine ($^{123}\text{I-IMP}$) as one example of organic compounds. In *in vitro* experiments, the displacement effects were observed by adding warfarin or 6-methoxy-2-naphthylacetic acid (6-MNA), both of which have the specificity to HSA, or by adding verapamil which has the binding specificity to AGP (cf. Example 5 and Table 9), thus the displacement effect on organic compounds was observed and proved. Further, in the experiments by using 6-MNA and Verapamil in which those were added separately or added simultaneously, the synergistic effect of the displacement effect was observed, thus it is indicated that the displacement effect can be enhanced by using plural second drugs together (cf. Example 6 and Table 10).

[0026] In *in vivo* experiments in rats, as compared with the control group (unloaded with verapamil), the higher concentration of free $^{123}\text{I-IMP}$ in blood was observed in the test group (loaded with verapamil). Reflecting the fact, 10 minutes after the administration, the brain uptake of $^{123}\text{I-IMP}$ in the test group (loaded with verapamil) was about 2 times that in the control group (Example 7). In this *in vivo* experiments, the test solution containing both of $^{123}\text{I-IMP}$ and verapamil was prepared in advance (Example 7 (1)) and used it in the experiment. Results of Example 7 indicates that it is possible to regulate the free drug concentration by simultaneous administration of the first and second drugs using their mixture as well as by the separate administration of the first drug and the second drugs and the biodistribution of the first drug could reflect it.

[0027] As for an example of the reducing effect of the free drug concentration, decrease in the free fraction (i.e., increase in binding fraction to protein) was observed in *in vitro* experiment using N-(3-fluoropropyl)-2 β -carbomethoxy-3 β -(4-iodophenyl)-nortropane labeled with radioactive iodine (I-125) ($^{125}\text{I-FP-CIT}$) together with human serum by adding dansyl-L-asparagine (DNSA) which is specific to the site I on albumin (cf. Example 8 and Table 15).

EXAMPLES

[0028] The present invention will be explained in more detail by illustrating the following examples, but the invention will not be restricted only to these examples.

[0029] Methods for testing the compounds obtained and the reagents used are as follows.

(1) Ultrafiltration: Filtration was conducted by using an equipment of ULTRACENT-10 which treats for up to 1.5 ml (manufactured by Tosoh Corp.).

(2) $^{99m}\text{TcO}_4^-$: Prepared by use of a $^{99}\text{Mo}/^{99m}\text{Tc}$ generator of MEDITECH (manufactured by NIHON MEDI-PHYSICS CO., LTD.) and used its eluent as in the form of a physiological saline solution.

(3) Reagents: The all reagents used were "Extra-pure reagent grade".

(4) Test animals: The all test animals used were Wister-strain male rats (body weight: 200-250g). Prior to the test, the animals were bred under light-dark cyclic condition in every 12 hours for 1 week, and had free access to food and water.

Example 1

Examination of displacement effects of the second drugs on $^{99m}\text{Tc-MAG}_3$ binding to plasma protein

[0030] Displacement test of $^{99m}\text{Tc-MAG}_3$ binding to serum albumin was conducted as follows by use of human serum or rat serum and site-specific drugs (second drugs) with binding affinity for the binding site I or site II on albumin. Bucolome, valproic acid, warfarin and cefazolin were used as site-specific drugs with binding affinity for the site I, and ibuprofen, sodium octanoate and sodium oleate were used as site-specific drugs with binding affinity for the site II.

[0031] First, albumin content in a normal human serum was measured previously, and the concentration of the human

serum albumin (HSA) was adjusted to 500 μM by phosphate buffer (pH=7.4).

[0032] Further, a site-specific drug with the binding affinity for the site I or the site II on HSA was added to the above-mentioned serum solution, as in the form of a methanol solution or an aqueous solution. As the sample solution for control group, only methanol or water was added to the above-mentioned serum solution.

[0033] Next, a certain amount of $^{99\text{m}}\text{Tc-MAG}_3$ (about 740 kBq/20 μl) was added to each of the sample and a certain amount (20-50 μl) of the sample was taken out as the specimen before ultrafiltration. Each 0.9 mL of the samples was put in a ultrafilter and conducted ultrafiltration under the condition of 1500 X g, for 10 minutes. Then, 20-50 μl each of the filtrates was taken out as the specimen after ultrafiltration. Radioactivities (cpm) of the specimens before and after ultrafiltration were measured, and the free fraction (%) of $^{99\text{m}}\text{Tc-MAG}_3$ was calculated by the following equation:

$$\text{free fraction of } ^{99\text{m}}\text{Tc-MAG}_3 (\%) = [A]/[B]$$

[A]: Radioactivity (cpm) after ultrafiltration,

[B]: Radioactivity (cpm) before ultrafiltration

[0034] Similarly, albumin content in normal rat serum was previously measured, and the concentration of the rat serum albumin (RSA) was adjusted to 375 μM by phosphate buffer (pH=7.4) so as to conduct the test similar to that of the case of human serum. The results are shown in Table 1, Fig.1 and Fig.2.

[0035] In the case of human serum, the free fraction (%) of $^{99\text{m}}\text{Tc-MAG}_3$ in the test samples to which a site-specific drug to site I was added, such as bucolome, valproic acid, warfarin or cefazolin, was significantly increased as compared with the free fraction (10.2%) of $^{99\text{m}}\text{Tc-MAG}_3$ of the control sample.

[0036] On the other hand, in other test samples to which a site-specific drug to site II was added, such as ibuprofen, sodium octoate or sodium oleate, increase in the free fraction was not observed.

[0037] Similarly, in test samples of the rat serum to which a site-specific drug to site I was added, increase in the free fraction (%) of $^{99\text{m}}\text{Tc-MAG}_3$ was observed.

[0038] As can be seen from the above results, it is clearly indicated that the free fraction of $^{99\text{m}}\text{Tc-MAG}_3$ in the blood can be increased by adding a site-specific drug to site I. Although, warfarin, octanoic acid and oleic acid might be considered clinically unsuitable for the purpose of this invention, they were used for the confirmation of the effects of site-specific drugs to the binding site.

Table 1. Displacement effects of the second drugs on $^{99\text{m}}\text{Tc-MAG}_3$ binding to plasma protein

Site-specific drug (second drug)	Human serum albumin (HAS) $^{99\text{m}}\text{Tc-MAG}_3$ free fraction (%)		Rat serum albumin (RSA) $^{99\text{m}}\text{Tc-MAG}_3$ free fraction (%)	
concentration	200 μM	400 μM	200 μM	400 μM
Control	10.20%		24.75%	
bucolome	12.23%	13.74%	32.76%	43.85%
valproic acid	11.98%	13.02%	28.48%	29.30%
warfarin	11.50%	13.57%	33.57%	43.28%
cefazolin	11.13%	14.76%	28.58%	33.52%
ibuprofen	10.18%	10.53%	28.48%	33.04%
octanoic acid	9.60%	9.86%	-	-
oleic acid	8.74%	9.44%	-	-

Example 2

Biodistribution of $^{99\text{m}}\text{Tc-MAG}_3$ in rat loaded with bucolome

[0039] Effect of second drug on biodistribution of $^{99\text{m}}\text{Tc-MAG}_3$ in rat was examined using the control group and the test group with bucolome loading. $^{99\text{m}}\text{Tc-MAG}_3$ (740 kBq/100 μl) was administered to the tail vein of Wister-strain rat. The rats were decapitated at 2, 5, 10 and 15 minutes after administration of $^{99\text{m}}\text{Tc-MAG}_3$, then the blood and the organs of interest were excised. After measured the weight of these organs, the radioactivities were determined. After decay correction of the radioactivity, the accumulation ratios (% dose/organ and % dose/g of the tissue) were determined.

[0040] As for the rat of test group loaded with bucolome, 5 minutes before the administration of $^{99\text{m}}\text{Tc-MAG}_3$, 20

mg/kg of body weight or 100 mg/kg of body weight of bucolome was administered to the tail vein.

[0041] The results are shown in Table 2 and Table 3 (control group), Table 4 and Table 5 (test group, loaded with 20 mg/kg of bucolome) and Table 6 (test group, loaded with 100 mg/kg of bucolome).

[0042] In the control group and test group with bucolome loading of 20 mg/kg of body weight, wherein the dose and other conditions were the same as mentioned above, except the decapitation time was prescribed at 2, 5 and 10 minutes, administrations and decapitations of rats were conducted so that 3-5 ml of blood per one rat were collected. Serum was separated using a sample tube, after that the free fraction was determined by the procedures as described in Example 1. The time course of the free fraction of $^{99m}\text{Tc-MAG}_3$ *in vivo* are shown in Fig. 4.

[0043] From the results shown above, it has become apparent that blood clearance was accelerated in the test group with bucolome loading (Fig. 3), and the free fraction of $^{99m}\text{Tc-MAG}_3$ *in vivo* in test group was remarkably increased (Fig. 4).

[0044] In the control group, accumulation of $^{99m}\text{Tc-MAG}_3$ in the kidney (% dose/organ) increased from 2 minutes to 5 minutes after the administration, then gradually decreased and disappeared. While, in test group with bucolome loading, accumulation of $^{99m}\text{Tc-MAG}_3$ in the kidney rapidly increased to the maximum value just after the administration (in 2 minutes), then decreased and disappeared quickly as compared with that of the control group (Fig. 5).

[0045] Biodistribution of $^{99m}\text{Tc-MAG}_3$ (% dose/g tissue) in rat 10 minutes after the administration is shown in Fig. 6. As can be seen from Fig. 6, in the test group with bucolome loading, $^{99m}\text{Tc-MAG}_3$ was rapidly cleared from the kidney which is the target organ of $^{99m}\text{Tc-MAG}_3$, thus radioactivity was quickly cleared as compared with that of the control group. Clearance from the blood and other organs were also rapid.

Table 2. Biodistribution of $^{99m}\text{Tc-MAG}_3$ in rats (Control group: % dose/organ)

Organs	2 minutes	5 minutes	10 minutes	15 minutes
Spleen	0.110 ± 0.024	0.064 ± 0.001	0.025 ± 0.006	0.01,9 ± 0.001
Pancreas	0.137 ± 0.026	0.079 ± 0.013	0.084 ± 0.050	0.030 ± 0.001
Stomach	0.276 ± 0.011	0.169 ± 0.003	0.119 ± 0.043	0.178 ± 0.009
Liver	5.196 ± 0.387	5.187 ± 2.759	1.671 ± 0.099	0.973 ± 0.266
Kidney	23.882 ± 4.669	31.329 ± 4.979	29.198 ± 3.729	15.864 ± 3.960
Heart	0.262 ± 0.039	0.184 ± 0.046	0.079 ± 0.028	0.034 ± 0.004
Lung	0.635 ± 0.116	0.594 ± 0.106	0.275 ± 0.042	0.129 ± 0.084
Urine	0.236 ± 0.119	1.309 ± 0.941	16.872 ± 4.042	38.419 ± 2.150

Table 3. Biodistribution of $^{99m}\text{Tc-MAG}_3$ in rats (Control group: % dose/g tissue)

Organs	2 minutes	5 minutes	10 minutes	15 minutes
Blood	1.482 ± 0.137	0.968 ± 0.163	0.387 ± 0.018	0.160 ± 0.022
Spleen	0.171 ± 0.031	0.104 ± 0.014	0.094 ± 0.008	0.026 ± 0.006
Pancreas	0.233 ± 0.029	0.150 ± 0.001	0.096 ± 0.027	0.098 ± 0.007
Stomach	0.176 ± 0.011	0.031 ± 0.012	0.015 ± 0.010	0.110 ± 0.050
Liver	0.579 ± 0.081	0.523 ± 0.268	0.145 ± 0.009	0.100 ± 0.024
Kidney	13.039 ± 3.199	16.721 ± 0.992	15.526 ± 2.763	8.282 ± 1.222
Heart	0.448 ± 0.056	0.290 ± 0.099	0.128 ± 0.031	0.057 ± 0.010
Lung	0.621 ± 0.100	0.970 ± 0.069	0.213 ± 0.008	0.112 ± 0.057

Table 4. Biodistribution of $^{99m}\text{Tc-MAG}_3$ in rats (Test group with 20 mg/kg bucolome loading: % dose/organ)

Organs	2 minutes	5 minutes	10 minutes	15 minutes
Spleen	0.103 ± 0.001	0.048 ± 0.006	0.018 ± 0.009	0.011 ± 0.003
Pancreas	0.239 ± 0.072	0.139 ± 0.030	0.060 ± 0.023	0.075 ± 0.053
Stomach	0.289 ± 0.057	0.153 ± 0.023	0.111 ± 0.032	0.104 ± 0.040
Liver	7.289 ± 0.333	3.140 ± 0.745	1.217 ± 0.471	0.806 ± 0.187
Kidney	26.404 ± 2.243	22.952 ± 9.437	17.118 ± 8.295	9.544 ± 3.655
Heart	0.210 ± 0.034	0.114 ± 0.019	0.037 ± 0.012	0.029 ± 0.014
Lung	0.792 ± 0.044	0.456 ± 0.137	0.148 ± 0.079	0.085 ± 0.025

(continued)

Organs	2 minutes	5 minutes	10 minutes	15 minutes
Urine	0.802±0.709	2.692±2.721	14.792±9.307	23.969±18.025

Table 5. Biodistribution of $^{99m}\text{Tc-MAG}_3$ in rats (Test group with 20 mg/kg bucolome loading: % dose/g tissue)

Organs	2 minutes	5 minutes	10 minutes	15 minutes
Blood	1.050±0.057	0.544±0.043	0.186±0.076	0.152±0.088
Spleen	0.153±0.018	0.083±0.005	0.026±0.011	0.018±0.006
Pancreas	0.314±0.013	0.145±0.017	0.062±0.021	0.088±0.052
Stomach	0.145±0.121	0.033±0.017	0.033±0.020	0.032±0.012
Liver	0.853±0.135	0.280±0.017	0.117±0.035	0.088±0.028
Kidney	13.069±0.379	11.050±4.260	8.558±3.867	9.809±1.823
Heart	0.329±0.034	0.172±0.021	0.057±0.017	0.095±0.021
Lung	0.613±0.013	0.373±0.073	0.120±0.099	0.081±0.020

Table 6. Biodistribution of $^{99m}\text{Tc-MAG}_3$ in rats 10 minutes after the administration (Test group with 100 mg/kg bucolome loading: % dose/g tissue)

Organs	Control group	Test group with bucolome loading
Blood	0.317±0.073	0.047±0.044
Brain	0.010±0.001	0.001±0.001
Spleen	0.052±0.008	0.009±0.008
Pancreas	0.046±0.000	0.006±0.007
Stomach	0.024±0.024	0.040±0.036
Liver	0.151±0.001	0.033±0.026
Kidney	6.191±0.187	0.651±0.324
Heart	0.101±0.016	0.014±0.010
Lung	0.195±0.030	0.043±0.037

Example 3

Examination of displacement effect on $^{99m}\text{Tc-MAG}_3$ by means of renography in rats

[0046] By using Wistar-strain rats (body weight: 400g), displacement effect of bucolome on $^{99m}\text{Tc-MAG}_3$ was examined by means of renography in rats. Prism 3000 (picker) was used as an apparatus.

[0047] A catheter was inserted into the femoral vein of the rat, then $^{99m}\text{Tc-MAG}_3$ (11.1 MBq) was injected through the catheter and the control renogram was obtained. The dynamic imaging was conducted 10 seconds/scan for 20 minutes. About 2 hours after, confirming the urination and decrease in background radioactivity, then bucolome was loaded to the same rat. Bucolome was dissolved in ethanol and adjusted as in the dose of 20 mg/kg, then intravenously injected by using a microinjector taken in 10 minutes. About 5 minutes after finish of the intravenous injection of bucolome, $^{99m}\text{Tc-MAG}_3$ was intravenously injected through the catheter and the renogram was taken similarly by 10 seconds/scan for 20 minutes. Fig. 7 shows the renogram (time-radioactivity curves in the kidney) used for functional analysis of the kidney. As can be seen from the Fig. 7, in the control group, the radioactivity curve went up gradually at the initial stage after the administration, and the peak time was 240 seconds. On the other hand, in the test group with bucolome loading, the radioactivity curve rose quickly, and the peak time was 120 seconds which was a half length of that of the control group. Renal function is usually analyzed by determination of the peak time in this renogram and a slope of the straight line in a linear regression. By inhibiting the binding ability of $^{99m}\text{Tc-MAG}_3$ to plasma protein, the renogram was obtained as close to an ideal and was approximated to a simple curve. Thus, the functional analysis of the kidney can be easily carried out, and the time for functional analysis can be shortened by shortening the peak time.

Table 7. Analytical results of renogram of ^{99m}Tc -MAG₃ in rats

		Peak time (Second)	Slope (Count/second)
5	Rat 1		
	Control group	240	1.166
	Test group (with bucolome loading)	110	2.208
10	Rat 2		
	Control group	170	0.941
	Test group (with bucolome loading)	120	2.000

Example 4

Examination of displacement effects of the second drugs on ^{99m}Tc -EDC binding to plasma protein

[0048] Displacement experiment of ^{99m}Tc -ECD binding to a serum albumin was conducted by procedures similar to those shown in Example 1 using human serum; and bucolome, valproic acid, warfarin and cefazolin with binding specificity to the site I on albumin; ibuprofen and sodium octanoate with binding specificity to the site II on albumin; and etoposide with binding specificity to HAS on which binding site is not identified. The results are shown in Table 8.

[0049] As compared with the free fraction (26.03%) of ^{99m}Tc -ECD in human serum shown in control group, the free fraction of ^{99m}Tc -ECD in human serum were remarkably increased in test group by etoposide, both at the concentration of 200 μM and 400 μM . Similarly, the free fraction of ^{99m}Tc -ECD in human serum were also increased by bucolome, valproic acid and warfarin, but were not remarkably shown as compared with that of etoposide. On the contrary, the free fraction of ^{99m}Tc -ECD in human serum were not clearly increased by ibuprofen and sodium octanoate which have the specificity to the site II on albumine.

Table 8 Displacement of ^{99m}Tc -ECD binding to plasma protein

Site-specific drug (second drug)	free fraction of ^{99m}Tc -ECD (%)	
Concentration	200 μM	400 μM
Control	26.03%	
Bucolome	28.62%	30.25%
Valproic acid	28.36%	30.25%
Warfarin	31.00%	31.37%
Cefazolin	25.92%	27.40%
Etoposide	33.26%	37.38%
Ibuprofen	23.09%	24.09%
Octanoic acid	28.22%	29.64%

Example 5

Examination of displacement effects of the second drugs on ^{123}I -IMP binding to plasma protein

[0050] Displacement experiment of ^{123}I -IMP binding to a serum albumin was conducted by procedures similar to those shown in Example 1 using human serum; and as the second drug, bucolome and warfarin with binding specificity to the binding site I on albumin; ibuprofen, sodium octanoate, 6-methoxy-2-naphthylacetic acid (6-MNA) with binding specificity to the binding site II on albumin; and verapamil which has the specificity to α_1 -acid glycoprotein (AGP). Concentration of the second drug (e.g., bucolome) was 400 μM and added amount of ^{123}I -IMP was about 220 kBq/20 μL . The results are shown in Table 9.

[0051] As compared with free fraction (29.29%) of ^{123}I -IMP in human serum shown in control group, the free fraction of ^{123}I -IMP in human serum in test group was remarkably increased by adding verapamil with binding specificity to AGP. Further, the free fraction of ^{123}I -IMP binding in human serum in test group was also increased by warfarin and 6-MNA mainly bound to albumin. In view of these facts, it is suggested that ^{123}I -IMP binds to the binding site on both albumin and AGP, and it is clearly understood that the free fraction of ^{123}I -IMP can be increased by a drug having the specificity

to each binding sites of these proteins.

Table 9. Displacement of ^{123}I -IMP binding to human plasma protein (Concentration of the site-specific drug was 400 μM)

Site-specific drug (second drug)	free of fraction of ^{123}I -IMP (%)
Control	29.29%
bucolome	30.26%
warfarin	34.69%
ibuprofen	28.43%
octanoic acid	28.74%
6-MNA	32.70%
verapamil	38.34%

Example 6

Examination of displacement effects of the second drugs on ^{123}I -IMP binding to plasma protein; synergistic effect

[0052] Displacement experiment of ^{123}I -IMP binding to serum albumin was conducted by procedures similar to those shown in Example 5 using human serum and as the second drugs, 6-MNA having the specificity to the binding site II on albumin and verapamil having the specificity to the binding site on AGP. Concentration of the second drugs were 400 μM and the added amount of ^{123}I -IMP was about 220 kBq/20 μL .

[0053] The tests were conducted in one group by using 6-MNA or verapamil independently, and in another group by using 6-MNA and verapamil simultaneously to study synergistic effect. In both groups, concentrations of the second drugs were 400 μM . The results are shown in Table 10.

[0054] In case of using 6-MNA and verapamil simultaneously, the free fraction of ^{123}I -IMP was over than the sum of the corresponding values obtained by single use of 6-MNA or verapamil, respectively. In view of the above facts, synergistic effect can be expected by using the plural second drugs.

Table 10. Displacement of ^{123}I -IMP binding to human plasma protein: synergistic effect

Site-specific drug (second drug)	free fraction of ^{123}I -IMP
Control	26.52%
6-MNA	30.00%
verapamil	33.87%
6-MNA + verapamil	39.26%

Example 7

Biodistribution of ^{123}I -IMP in rat with verapamil loading (1) Preparation of ^{123}I -IMP-verapamil mixed solution

[0055] 35 Milligrams of verapamil bulk drug powder was dissolved in 2 ml of Vasolan injection (verapamil 5 mg/2 ml, manufactured by Eisai Co., Ltd.), then 34 μL of ^{123}I -IMP injection (111 MBq/ml, manufactured by NIHON MEDI-PHYSICS CO., LTD.) was added thereto and mixed throughly.

(2) Biodistribution of ^{123}I -IMP in rats

[0056] Control group: ^{123}I -IMP Injection solution (185 kBq/300 μL) being diluted with physiological saline was administered via the caudal vein of rats of control group. The rats were decapitated at 2, 5, 10, 30 and 60 minutes after administration. Then the blood was sampled and the organs of interest were excised. After measured the weight of these specimens, the radioactivities of the blood and organs were measured. After half-life of the radioactivities were corrected, the accumulation rate (% dose/g of the tissue) was obtained.

[0057] Test group: 100 μL of ^{123}I -IMP-verapamil mixed solution was administered via the caudal vein of rats of the

test group (about 10 mg/kg loaded as verapamil), then the rats were treated similarly to those of control group. The results of biodistribution of ^{123}I -IMP are shown in Table 11 (control group), Table 12 (test group with verapamil loading), and Table 13 (comparison of both control and test groups of 10 minutes after the administration).

(3) Examination of displacement effect on ^{123}I -IMP binding to plasma protein in rat

[0058] Under the same conditions as mentioned above concerning the constitution of control and testing group, timings of decapitation, and dose of drugs, administrations of the drugs and decapitations of rats were conducted, and 3-5 ml of blood per one rat was sampled. Serum was separated using a sample tube, after that the free fraction of ^{123}I -IMP was determined by the procedures as described in Example 1. The free fraction of ^{123}I -IMP in the blood sample of rat obtained at each decapitation timing are shown in Table 14.

[0059] As shown in Table 14, it is clearly indicated that the free fraction of ^{123}I -IMP in the blood sample of rat were increased by the loading with the verapamil. As shown in Table 11 through Table 13, corresponding to increasing in the free fraction of ^{123}I -IMP in the blood due to the loading with verapamil, the uptake of ^{123}I -IMP into the brain which is the target organ of ^{123}I -IMP was rapidly increased after administration of ^{123}I -IMP-verapamil mixed solution in the test group, thus the brain uptake of ^{123}I -IMP in the test group after the administration was increased about 2 times higher than that of shown in control group. These facts indicate that, even if a mixed drug of the first drug and the second drug is administered (simultaneous administration of the first drug and the second drug), the free fraction of ^{123}I -IMP can be regulated by the second drug, and the biodistribution of the first drug could reflect it.

Table 11. Biodistribution of ^{123}I -IMP in rats (Control group: % Dose/g Tissue)

Tissues	2 Minutes	5 Minutes	10 Minutes	30 Minutes	60 Minutes
Blood	0.198±0.052	0.133±0.005	0.116±0.011	0.136±0.028	0.181±0.006
Brain	1.800±0.418	1.476±0.225	1.006±0.379	1.396±0.395	1.511±0.011
Pancreas	1.503±0.353	1.923±0.445	1.721±0.217	2.032±0.505	1.957±0.345
Spleen	0.880±0.216	0.999±0.355	1.008±0.074	1.356±0.277	1.290±0.138
Stomach	0.302±0.065	0.500±0.078	0.407±0.230	0.885±0.366	1.295±0.393
Liver	0.506±0.109	0.699±0.061	0.711±0.143	1.192±0.536	1.442±0.164
Kidney	3.406±0.905	2.285±0.256	1.303±0.190	1.359±0.222	1.585±0.132
Heart	1.949±0.293	1.014±0.070	0.631±0.111	0.529±0.037	0.540±0.026
Lung	11.236±0.780	9.000±0.600	6.279±1.026	5.209±1.446	5.186±0.616

Table 12. Biodistribution of ^{123}I -IMP in rats (Test group (Loaded with verapamil): % Dose/g Tissue)

Tissues	2 Minutes	5 Minutes	10 Minutes	30 Minutes	60 Minutes
Blood	0.238±0.083	0.228±0.012	0.139±0.003	0.098±0.044	0.110±0.002
Brain	1.584±0.425	1.916±0.131	2.145±0.410	1.529±0.811	1.449±0.281
Pancreas	1.268±0.375	1.659±0.496	1.911±0.685	1.877±0.886	1.478±0.161
Spleen	0.052±0.025	0.063±0.250	0.213±0.118	0.886±0.319	1.193±0.129
Stomach	0.234±0.111	0.164±0.078	0.377±0.013	0.782±0.621	1.058±0.126
Liver	0.287±0.156	0.350±0.130	0.688±0.237	1.185±0.751	1.639±0.051
Kidney	1.424±0.313	1.278±0.381	1.766±0.678	1.231±0.632	1.242±0.146
Heart	3.769±0.911	2.260±0.680	1.247±0.209	0.471±0.209	0.456±0.039
Lung	9.234±1.748	8.377±0.563	6.947±1.486	3.890±2.223	9.133±0.079

Table 13. Biodistribution of ^{123}I -IMP in rats 10 minutes after the administration (% Dose/g Tissue)

Tissue	Control group	Test group with verapamil loading
Blood	0.116±0.011	0.139±0.003
Brain	1.006±0.379	2.145±0.410
Pancreas	1.721±0.217	1.911±0.685
Spleen	1.008±0.074	0.213±0.118

(continued)

Tissue	Control group	Test group with verapamil loading
Stomach	0.407±0.230	0.377±0.013
Liver	0.711±0.143	0.688±0.237
Kidney	1.303±0.190	1.766±0.678
Heart	0.631±0.111	1.247±0.209
Lung	6.279±1.026	6.947±1.486

Table 14. Free fraction (%) of ^{123}I -IMP in the blood of rats

	2 Minutes	5 Minutes	10 Minutes	30 Minutes	60 Minutes
Control group	56.75±9.21	50.70±10.37	45.91±3.12	27.29±4.85	16.77±4.11
Test group with verapamil loading	52.40±6.00	56.52±4.38	66.86±6.34	38.03±6.69	31.86±8.23

Example 8

Examination of regulating the free fraction of ^{125}I -FP-CIT

[0060] Experiment was conducted by procedures similar to those shown in Example 5 using human serum and as the second drug, bucolome, phenylbutazone, warfarin and dansyl-L-asparagine (DNSA) with binding specificity to the binding site I on albumin, and ibuprofen, 6-methoxy-2-naphthylacetic acid (6-MNA) with binding specificity to the binding site II on albumin. Concentration of the second drug (e.g., bucolome and the like) were 400 μM and added amount of ^{123}I -FP-CIT was about 74 kBq/20 μL . The result are shown in Table 15.

[0061] As compared with the free fraction (17.26%) of ^{123}I -FP-CIT in human serum shown in control group, the free fraction of ^{123}I -FP-CIT in test group was remarkably decreased by DNSA. Furthermore, the free fraction of ^{123}I -FP-CIT in test group was also decreased by phenylbutazone and ibuprofen. In view of these facts, it is clearly understood that the free fraction of the first drug can be decreased by the second drug having the binding affinity for the plasma proteins.

Table 15. Free fraction of ^{123}I -FP-CIT in human serum (concentration of the site-specific drug was 400 μM)

Site-specific drug (second drug)	free fraction (%) of ^{123}I -FP-CIT
Control	17.26%
bucolome	18.40%
phenylbutazone	14.92%
warfarin	17.88%
DNSA	12.80%
ibuprofe	15.92%
6-MNA	18.10%

Claims

1. Use of a second drug in the manufacture of a medicament to regulate the binding of a first drug to a plasma protein, wherein:

the first drug is a radiodiagnostic drug for *in vivo* use or a radiotherapeutic drug for *in vivo* use and has one group selected from bisaminothiol or its derivatives, monoaminomonoamidobisthiol or its derivatives, bisamidobisthiol or its derivatives, mercaptoacetylglcylglycylglycine or its derivatives, hexamethylpropyleneamine-oxime or its derivatives, ethylenebis[bis(2-ethoxyethyl) phosphine] (tetrofosmin) or its derivatives, 2,3-dimercaptosuccinic acid or its derivatives, ethylenecysteine dimer derivatives, methoxyisobutylisonitrile derivatives, polyamine derivatives, pyridoxylideneamine derivatives, methylene diphosphonate, hydroxymethylene diphosphonate derivatives, β -methyl- ω -phenylpentadecanoic acid or its derivatives, N-isopropylamphetamine,

hippuric acid, benzylguanidine and tropane derivatives, the group being radiolabeled with one nuclide selected from 11-carbon (^{11}C), 15-oxygen (^{15}O), 18-fluorine, (^{18}F), 32-phosphorus (^{32}P), 59-iron (^{59}Fe), 67-copper (^{67}Cu), 67-gallium (^{67}Ga), 81m-krypton ($^{81\text{m}}\text{Kr}$), 81-rubidium (^{81}Rb), 89-strontium (^{89}Sr), 90-yttrium (^{90}Y), 99m-technetium ($^{99\text{m}}\text{Tc}$), 111-indium (^{111}In), 123-iodine (^{123}I), 125-iodine (^{125}I), 131-iodine (^{131}I), 133-xenon (^{133}Xe), 117m-tin ($^{117\text{m}}\text{Sn}$), 153-samarium (^{153}Sm), 186-rhenium (^{186}Re), 188-rhenium (^{188}Re), 201-thallium (^{201}Tl), 212-bismuth (^{212}Bi), 213-bismuth (^{213}Bi) and 211-astatine (^{211}At);

the second drug is selected from bucolome, cefazolin, etoposide, phenylbutazone, aspirin, salicylic acid, ceftriaxone, sulfamethizole, valproic acid, nabumetone, 6-methoxy-2-naphthyl acetic acid, ibuprofen, probenecid, dansyl-L-asparagine, verapamil and disopyramide; and

the first drug and the second drug have binding affinity for said plasma protein.

2. Use according to claim 1, wherein the second drug has binding affinity to the same binding sites on the plasma protein to which the first drug has binding affinity.

3. A product which comprises:

(a) a second drug selected from bucolome, cefazolin, etoposide, phenylbutazone, aspirin, salicylic acid, ceftriaxone, sulfamethizole, valproic acid, nabumetone, 6-methoxy-2-naphthyl acetic acid, ibuprofen, probenecid, dansyl-L-asparagine, verapamil and disopyramide; and

(b) a first drug which is a radiodiagnostic drug for *in vivo* use or a radiotherapeutic drug for *in vivo* use and has one group selected from bisaminothiol or its derivatives, monoaminomonoamidobisthiol or its derivatives, bisamidobisthiol or its derivatives, mercaptoacetylglucylglycylglycine or its derivatives, hexamethylpropyleneaminioxime or its derivatives, ethylenebis[bis(2-ethoxyethyl) phosphine] (tetrofosmin) or its derivatives, 2,3-dimercaptosuccinic acid or its derivatives, ethylenecysteine dimer derivatives, methoxyisobutylisonitrile derivatives, polyamine derivatives, pyridoxylideneaminat derivatives, methylene diphosphonate, hydroxymethylene diphosphonate derivatives, β -methyl- ω -phenylpentadecanoic acid or its derivatives, N-isopropylamphetamine, hippuric acid, benzylguanidine and tropane derivatives, the group being radiolabeled with one nuclide selected from 11-carbon (^{11}C), 15-oxygen (^{15}O), 18-fluorine, (^{18}F), 32-phosphorus (^{32}P), 59-iron (^{59}Fe), 67-copper (^{67}Cu), 67-gallium (^{67}Ga), 81m-krypton ($^{81\text{m}}\text{Kr}$), 81-rubidium (^{81}Rb), 89-strontium (^{89}Sr), 90-yttrium (^{90}Y), 99m-technetium ($^{99\text{m}}\text{Tc}$), 111-indium (^{111}In), 123-iodine (^{123}I), 125-iodine (^{125}I), 131-iodine (^{131}I), 133-xenon (^{133}Xe), 117m-tin ($^{117\text{m}}\text{Sn}$), 153-samarium (^{153}Sm), 186-rhenium (^{186}Re), 188-rhenium (^{188}Re), 201-thallium (^{201}Tl), 212-bismuth (^{212}Bi), 213-bismuth (^{213}Bi) and 211-astatine (^{211}At); which first drug has binding affinity for a plasma protein for which said second drug has binding affinity;

for simultaneous, separate or sequential use in regulating the binding affinity of the first drug to said plasma protein.

4. A product according to claim 3, wherein each of the first drug and the second drug is separately filled in a container, and prepared as kit form for supply.

5. A product according to claim 3 or 4, wherein the second drug has binding affinity to the same binding sites on the plasma protein to which the first drug has binding affinity.

Patentansprüche

1. Verwendung eines zweiten Arzneimittels zur Herstellung eines Medikaments, um die Bindung eines ersten Arzneimittels an ein Plasmaprotein zu regulieren, wobei:

das erste Arzneimittel ein radiodiagnostisches Arzneimittel zur Verwendung *in vivo* oder ein radiotherapeutisches Arzneimittel zur Verwendung *in vivo* ist und einen Rest aufweist, ausgewählt aus Bisaminothiol oder Derivaten davon, Monoaminomonoamidobisthiol oder Derivaten davon, Bisamidobisthiol oder Derivaten davon, Mercaptoacetylglucylglycylglycine oder Derivaten davon, Hexamethylpropylenaminioxim oder Derivaten davon, Ethylenbis[bis(2-ethoxyethyl)phosphin] (Tetrofosmin) oder Derivaten davon, 2,3-Dimercaptobernsteinsäure oder Derivaten davon, Ethylencysteindimerderivaten, Methoxyisobutylisonitrilderivaten, Polyaminderivaten, Pyridoxylidenaminatderivaten, Methylendiphosphonat, Hydroxymethylendiphosphonatderivaten, β -Methyl- ω -phenylpentadecansäure oder Derivaten davon, N-Isopropylamphetamin, Hippursäure, Benzylguanidin und Tropanderivaten, wobei der Rest mit einem Nuklid radiomarkiert ist, ausgewählt aus 11-Kohlenstoff (^{11}C), 15-Sauerstoff (^{15}O), 18-Fluor (^{18}F), 32-Phosphor (^{32}P), 59-Eisen (^{59}Fe), 67-Kupfer (^{67}Cu), 67-Gallium (^{67}Ga), 81m-

Krypton ($^{81\text{m}}\text{Kr}$), 81-Rubidium (^{81}Rb), 89-Strontium (^{89}Sr), 90-Yttrium (^{90}Y), 99m-Technetium ($^{99\text{m}}\text{Tc}$), 111-Indium (^{111}In), 123-Iod (^{123}I), 125-Iod (^{125}I), 131-Iod (^{131}I), 133-Xenon (^{133}Xe), 117m-Zinn ($^{117\text{m}}\text{Sn}$), 153-Samarium (^{153}Sm), 186-Rhenium (^{186}Re), 188-Rhenium (^{188}Re), 201-Thallium (^{201}Tl), 212-Bismut (^{212}Bi), 213-Bismut (^{213}Bi) und 211-Astatin (^{211}At);

das zweite Arzneimittel aus Bucolom, Cefazolin, Etoposid, Phenylbutazon, Aspirin, Salicylsäure, Ceftriaxon, Sulfamethizol, Valprionsäure, Nabumeton, 6-Methoxy-2-naphthylethylsäure, Ibuprofen, Probenecid, Dansyl-L-asparagin, Verapamil und Disopyramid ausgewählt ist; und

das erste Arzneimittel und das zweite Arzneimittel Bindungsaffinität zu dem Plasmaprotein aufweisen.

2. Verwendung gemäß Anspruch 1, wobei das zweite Arzneimittel Bindungsaffinität zu den selben Bindungsstellen auf dem Plasmaprotein aufweist, zu denen das erste Arzneimittel Bindungsaffinität aufweist.

3. Produkt, umfassend:

(a) ein zweites Arzneimittel, ausgewählt aus Bucolom, Cefazolin, Etoposid, Phenylbutazon, Aspirin, Salicylsäure, Ceftriaxon, Sulfamethizol, Valprionsäure, Nabumeton, 6-Methoxy-2-naphthylethylsäure, Ibuprofen, Probenecid, Dansyl-L-asparagin, Verapamil und Disopyramid; und

(b) ein erstes Arzneimittel, welches ein radiodiagnostisches Arzneimittel zur Verwendung in vivo oder ein radiotherapeutisches Arzneimittel zur Verwendung in vivo ist und einen Rest aufweist, ausgewählt aus Bisaminthiol oder Derivaten davon, Monoaminomonoamidobisthiol oder Derivaten davon, Bisamidobisthiol oder Derivaten davon, Mercaptoacetylglucylglycylglycine oder Derivaten davon, Hexamethylpropylenaminnoxim oder Derivaten davon, Ethylenbis[bis(2-ethoxyethyl)phosphin] (Tetrofosmin) oder Derivaten davon, 2,3-Dimercaptobernsteinsäure oder Derivaten davon, Ethylencysteinidimerderivaten, Methoxyisobutylisonitrilderivaten, Polyaminderivaten, Pyridoxylidenaminatderivaten, Methylendiphosphonat, Hydroxymethylendiphosphonatderivaten, β -Methyl- ω -phenylpentadecansäure oder Derivaten davon, N-Isopropylamphetamin, Hippursäure, Benzylguanidin und Tropanderivaten, wobei der Rest mit einem Nuklid radiomarkiert ist, ausgewählt aus 11-Kohlenstoff (^{11}C), 15-Sauerstoff (^{15}O), 18-Fluor (^{18}F), 32-Phosphor (^{32}P), 59-Eisen (^{59}Fe), 67-Kupfer (^{67}Cu), 67-Gallium (^{67}Ga), 81m-Krypton ($^{81\text{m}}\text{Kr}$), 81-Rubidium (^{81}Rb), 89-Strontium (^{89}Sr), 90-Yttrium (^{90}Y), 99m-Technetium ($^{99\text{m}}\text{Tc}$), 111-Indium (^{111}In), 123-Iod (^{123}I), 125-Iod (^{125}I), 131-Iod (^{131}I), 133-Xenon (^{133}Xe), 117m-Zinn ($^{117\text{m}}\text{Sn}$), 153-Samarium (^{153}Sm), 186-Rhenium (^{186}Re), 188-Rhenium (^{188}Re), 201-Thallium (^{201}Tl), 212-Bismut (^{212}Bi), 213-Bismut (^{213}Bi) und 211-Astatin (^{211}At); wobei das erste Arzneimittel Bindungsaffinität zu einem Plasmaprotein aufweist, zu dem das zweite Arzneimittel Bindungsaffinität aufweist; zur gleichzeitigen, getrennten oder aufeinanderfolgenden Verwendung bei der Regulierung der Bindungsaffinität des ersten Arzneimittels zu dem Plasmaprotein.

4. Produkt gemäß Anspruch 3, wobei das erste und zweite Arzneimittel jeweils getrennt in einen Behälter gefüllt sind und zur Bereitstellung als Kit hergestellt sind.

5. Produkt gemäß Anspruch 3 oder 4, wobei das zweite Arzneimittel Bindungsaffinität zu den selben Bindungsstellen auf dem Plasmaprotein aufweist, zu denen das erste Arzneimittel Bindungsaffinität aufweist.

Revendications

1. Utilisation d'une seconde drogue dans la fabrication d'un médicament pour réguler la liaison d'une première drogue à une protéine du plasma, dans laquelle :

la première drogue est une drogue de radiodiagnostic pour une utilisation in vivo ou une drogue radiothérapeutique pour une utilisation in vivo et a un groupe choisi parmi le bisaminthiol ou ses dérivés, le monoaminomonoamidobisthiol ou ses dérivés, le bisamidobisthiol ou ses dérivés, le mercaptoacétylglycylglycylglycine ou ses dérivés, l'hexaméthylpropylèneamineoxime ou ses dérivés, l'éthylènebis[bis(2-éthoxyéthyl) phosphine] (tétrofosmine) ou ses dérivés, l'acide 2,3-dimercaptosuccinique ou ses dérivés, des dérivés dimères d'éthylène-cystéine, des dérivés de méthoxyisobutylisocyanure, des dérivés de polyamine, des dérivés de pyridoxylidèneaminat, le diphosphonate de méthylène, des dérivés de disphosphonate d'hydroxyméthylène, l'acide β -phényl- ω -phénylpentadécanoïque ou ses dérivés, la N-isopropylamphétamine, l'acide hippurique, des dérivés benzylguanidine et tropane, le groupe étant radiomarké avec un nucléide choisi parmi le carbone-11 (^{11}C), l'oxygène 15 (^{15}O), le fluor 18 (^{18}F), le phosphore 32 (^{32}P), le fer 59 (^{59}Fe), le cuivre 67 (^{67}Cu), le gallium 67 (^{67}Ga), le krypton 81m ($^{81\text{m}}\text{Kr}$), le rubidium 81 (^{81}Rb), le strontium 89 (^{89}Sr), l'yttrium 90 (^{90}Y), le Technetium

99m (^{99m}Tc), l'indium 111 (¹¹¹In), l'iode 123 (¹²³I), l'iode 125 (¹²⁵I), l'iode 131 (¹³¹I), le Xénon 133 (¹³³Xe), l'étain 117m (^{117m}Sn), le samarium 153 (¹⁵³Sm), le rhénium 186 (¹⁸⁶Re), le rhénium 188 (¹⁸⁸Re), le thallium 201 (²⁰¹Tl), le bismuth 212 (²¹²Bi), le bismuth 213 (²¹³Bi) et l'Astatine 211 (²¹¹At);

la seconde drogue est choisie parmi le bucolome, la céfazoline, l'étoposide, la phénylbutazone, l'aspirine, l'acide salicylique, la céfatriaxone, la sulfaméthizole, l'acide valproïque, la nabumétone, l'acide 6-méthoxy-2-naphtyl acétique, l'ibuprofène, le probénécide, la dansyl-L-asparagine, le vérapamil et le disopyramide ; et la première drogue et la seconde drogue ont une affinité de liaison pour ladite protéine du plasma.

2. Utilisation selon la revendication 1, dans laquelle la seconde drogue a une affinité de liaison aux mêmes sites de liaison sur la protéine de plasma à laquelle la première drogue a une affinité de liaison.

3. Produit qui comprend :

(a) une seconde drogue choisie parmi le bucolome, la céfazoline, l'étoposide, la phénylbutazone, l'aspirine, l'acide salicylique, la céfatriaxone, la sulfaméthizole, l'acide valproïque, la nabumétone, l'acide 6-méthoxy-2-naphtyl acétique, l'ibuprofène, le probénécide, la dansyl-L-asparagine, le vérapamil et le disopyramide ; et

(b) une première drogue qui est une drogue de radiodiagnostic pour une utilisation in vivo ou une drogue radiothérapeutique pour une utilisation in vivo et a un groupe choisi parmi le bisaminothiols ou ses dérivés, le monoaminomonoamidobisthiol ou ses dérivés, le bisamidobisthiol ou ses dérivés, le mercaptoacétylglycylglycylglycine ou ses dérivés, l'hexaméthylpropylèneamineoxime ou ses dérivés, l'éthylènebis[bis(2-éthoxyéthyl)phosphine](tétrofosmine) ou ses dérivés, l'acide 2,3-dimercaptosuccinique ou ses dérivés, des dérivés dimère d'éthylène cystéine, des dérivés de méthoxyisobutylisonitrile, des dérivés de polyamine, des dérivés de pyridoxylidèneamine, le diphosphonate de méthylène, des dérivés de disphosphonate d'hydroxyméthylène, l'acide β-phényl-ω-phénylpentadécanoïque ou ses dérivés, la N-isopropylamphétamine, l'acide hippurique, des dérivés benzylguanidine et tropane le groupe étant radiomarké avec un nucléide choisi parmi le carbone-11 (¹¹C), l'oxygène 15 (¹⁵O), le fluor 18 (¹⁸F), le phosphore 32 (³²P), le fer 59 (⁵⁹Fe), le cuivre 67 (⁶⁷Cu), le gallium 67 (⁶⁷Ga), le krypton 81m (^{81m}Kr), le rubidium 81 (⁸¹Rb), le strontium 89 (⁸⁹Sr), l'yttrium 90 (⁹⁰Y), le Technetium 99m (^{99m}Tc), l'indium 111 (¹¹¹In), l'iode 123 (¹²³I), l'iode 125 (¹²⁵I), l'iode 131 (¹³¹I), le Xénon 133 (¹³³Xe), l'étain 117m (^{117m}Sn), le samarium 153 (¹⁵³Sm), le rhénium 186 (¹⁸⁶Re), le rhénium 188 (¹⁸⁸Re), le thallium 201 (²⁰¹Tl), le bismuth 212 (²¹²Bi), le bismuth 213 (²¹³Bi) et l'Astatine 211 (²¹¹At) ; laquelle première drogue a une affinité de liaison pour une protéine du plasma pour laquelle ladite seconde drogue a une affinité de liaison ;

pour une utilisation simultanée, séparée ou séquentielle dans la régulation de l'affinité de liaison de la première drogue à ladite protéine du plasma.

4. Produit selon la revendication 3, dans lequel chacune de la première drogue et la seconde drogue est séparément remplie dans un conteneur, et préparée sous forme de nécessaire pour approvisionnement.

5. Produit selon la revendication 3 ou 4, dans lequel la seconde drogue a une affinité de liaison aux mêmes sites de liaison sur la protéine de plasma pour laquelle la première drogue a une affinité de liaison.

FIG.1




FIG.2




FIG.3




FIG.4




FIG.5




FIG.6





FIG.7



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Non-patent literature cited in the description

- **BUBECK B. et al.** *J. Nucl. Med.*, 1990, vol. 31, 1285-1295 [0003]