

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 294 720 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
05.04.2006 Bulletin 2006/14

(21) Application number: **01949124.0**

(22) Date of filing: **14.06.2001**

(51) Int Cl.:
C07D 471/14 ^(2006.01) **A61K 31/55** ^(2006.01)
C07D 471/14 ^(2006.01) **C07D 243/00** ^(2006.01)
C07D 221/00 ^(2006.01) **C07D 221/00** ^(2006.01)

(86) International application number:
PCT/CA2001/000890

(87) International publication number:
WO 2001/096338 (20.12.2001 Gazette 2001/51)

(54) NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS

NICHT-NUKLEOSID REVERSE TRANSKRIPTASE INHIBITOREN

INHIBITEURS NON NUCLEOSIDIQUES DE TRANSCRIPTASE INVERSE

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
LT LV RO SI

(30) Priority: **16.06.2000 US 212329 P**
18.12.2000 US 256638 P

(43) Date of publication of application:
26.03.2003 Bulletin 2003/13

(60) Divisional application:
06100695.3

(73) Proprietor: **BOEHRINGER INGELHEIM (CANADA)
LTD.**
Laval, Quebec, H7S 2G5 (CA)

(72) Inventor: **SIMONEAU, Bruno**
Laval, Québec H7L 3V6 (CA)

(74) Representative: **Mahlbacher, Volker et al**
Boehringer Ingelheim GmbH,
CD Patents
55216 Ingelheim (DE)

(56) References cited:
EP-A- 0 767 172

- **KLUNDER, JANICE M. ET AL: "Novel Nonnucleoside Inhibitors of HIV-1 Reverse Transcriptase. 7. 8-Arylethylidipyridodiazepinones as Potent Broad-Spectrum Inhibitors of Wild-Type and Mutant Enzymes" J. MED. CHEM. (1998), 41(16), 2960-2971, XP002181339 cited in the application**
- **CYWIN, CHARLES L. ET AL: "Novel Nonnucleoside Inhibitors of HIV-1 Reverse Transcriptase. 8. 8-Aryloxymethyl- and 8-Arylthiomethylidipyridodiazepinones" J. MED. CHEM. (1998), 41(16), 2972 -2984, XP002181340**

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

EP 1 294 720 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**TECHNICAL FIELD OF THE INVENTION**

[0001] The invention relates to novel compounds and pharmaceutically acceptable salts thereof, their use, either alone or in combination with other therapeutic agents, in the treatment or prophylaxis of HIV infection, and to pharmaceutical compositions comprising the compounds.

BACKGROUND OF THE INVENTION

[0002] The disease known as acquired immune deficiency syndrome (AIDS) is caused by the human immunodeficiency virus (HIV), particularly the strain known as HIV-1. In order for HIV to be replicated by a host cell, the information of the viral genome must be integrated into the host cell's DNA. However, HIV is a retrovirus, meaning that its genetic information is in the form of RNA. The HIV replication cycle therefore requires a step of transcription of the viral genome (RNA) into DNA, which is the reverse of the normal chain of events. The transcription of the viral RNA into DNA is accomplished by an enzyme that has been aptly dubbed reverse transcriptase (RT). The HIV virion includes a copy of RT along with the viral RNA.

[0003] Reverse transcriptase has three known enzymatic functions; it acts as an RNA-dependent DNA polymerase, as a ribonuclease, and as a DNA-dependent DNA polymerase. Acting as an RNA-dependent DNA polymerase, RT transcribes a single-stranded DNA copy of the viral RNA. Acting as a ribonuclease, RT destroys the original viral RNA, and frees the DNA just produced from the original RNA. Finally, acting as a DNA-dependent DNA polymerase, RT makes a second, complementary DNA strand, using the first DNA strand as a template. The two strands form double-stranded DNA, which is integrated into the host cell's genome by another enzyme called integrase.

[0004] Compounds that inhibit the enzymatic functions of HIV-1 reverse transcriptase will inhibit replication of HIV-1 in infected cells. Such compounds are useful in the prevention or treatment of HIV-1 infection in human subjects, as demonstrated by known RT inhibitors such as 3'-azido-3'-deoxythymidine (AZT), 2',3'-dideoxyinosine (ddI), 2',3'-dideoxycytidine (ddC), d4T, 3TC, Nevirapine, Delavirdine, Efavirenz and Abacavir, the main drugs thus far approved for use in the treatment of AIDS.

[0005] As with any antiviral therapy, use of RT inhibitors in the treatment of AIDS eventually leads to a virus that is less sensitive to the given drug. Resistance (reduced sensitivity) to these drugs is the result of mutations that occur in the reverse transcriptase segment of the *pol* gene. Several mutant strains of HIV have been characterised, and resistance to known therapeutic agents is due to mutations in the RT gene. Some of the most commonly observed mutants clinically are: the Y181C mutant, in which a tyrosine (Y), at codon 181, has been mutated to a cysteine (C) residue, and K103N where the lysine (K) at position 103 has been replaced by asparagine (N). Other mutants which emerge with increasing frequency during treatment with known antivirals include the single mutants V106A, G190A, Y188C, and P236L; and the double mutants K103N/Y181C, K103N/P225H, K103N/V108I, and K103N/L100I.

[0006] As therapy and prevention of HIV infection using antivirals continues, the emergence of new resistant strains is expected to increase. There is therefore an ongoing need for new inhibitors of RT, with different patterns of effectiveness against the various mutants.

[0007] Compounds having tricyclic structures which are inhibitors of HIV-1 are described in U.S. Pat. No. 5,366,972. Other inhibitors of HIV-1 reverse transcriptase are described in Hargrave et al., J. Med. Chem., 34, 2231 (1991).

[0008] U.S. Pat. No. 5,705,499 proposes 8-arylalkyl- and 8-arylheteroalkyl-5,11-dihydro-6H-dipyrido[3,2-B:2',3'-E][1,4]diazepines as inhibitors of RT. The exemplified compounds are shown to have some activity against wild type and mutant HIV-1 RT, particularly Y181C and other single mutants such as K103N albeit less effectively.


Specifically, the compounds of the present invention are effective at inhibiting the Y181C and K103N mutants as well as a broad range of other single mutants and certain commonly found double mutants such as K103N/Y181C and K103N/P225H.

[0009] J. Klunder et al. (J. Med. Chem. 1998,41, 2960 - 2971) and Ch. L. Cywin et al. (J. Med. Chem. 1988, 41, 2972 - 2984) describe novel non-nucleoside inhibitors of HIV-1 reverse transcriptase, which differ from the tricyclic dipyrido-diazepinone skeleton of nevirapine by the introduction of a substituent at the 8-position.

SUMMARY OF THE INVENTION

[0010] The invention reduces the difficulties and disadvantages of the prior art by providing novel compounds that are potent inhibitors of single and double mutant strains of HIV-1 RT.

In a first aspect the invention provides a compound of the general formula I:



wherein


R² is selected from the group consisting of H, F, Cl, C₁₋₄ alkyl, C₃₋₄ cycloalkyl and CF₃;

R⁴ is H or Me;

R⁵ is H, Me or Et, with the proviso that R⁴ and R⁵ are not both Me, and if R⁴ is Me then R⁵ cannot be Et;

R¹¹ is Me, Et, cyclopropyl, propyl, isopropyl, or cyclobutyl;

Q is selected from the group consisting of:



or a pharmaceutically acceptable salt thereof.

[0011] In a second aspect, the invention provides an inhibitor of HIV replication, of the general formula I, or a pharmaceutically acceptable salt thereof.

[0012] In a third aspect, the invention provides an inhibitor of a reverse transcriptase enzyme of HIV, of the general formula I, or a pharmaceutically acceptable salt thereof.

[0013] In a fourth aspect, the invention provides the use of a compound of formula I, or a pharmaceutically acceptable salt thereof in the manufacture of a medicament useful for the treatment or prevention of HIV infection.

[0014] In a fifth aspect, the invention provides a pharmaceutical composition for the treatment or prevention of HIV infection, comprising a compound of formula I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

[0015] In a sixth aspect, the invention provides a method for preparation of a compound of formula I, or a pharmaceutically acceptable salt thereof.

DETAILED DESCRIPTION OF THE INVENTION**Definitions**

[0016] As used herein, the term "C₁₋₄ alkyl" is intended to mean linear or branched alkyl radicals containing from one to four carbon atoms and includes methyl, ethyl, propyl, isopropyl, butyl, sec-butyl and tert-butyl.

[0017] As used herein, the term "C₃₋₄ cycloalkyl" is intended to mean saturated cyclic hydrocarbon radicals containing three to four carbon atoms and includes cyclopropyl and cyclobutyl.

Detailed description of preferred embodiments


[0018] According to a preferred embodiment, compounds of the invention are defined according to formula I wherein R² is preferably Cl, F, or H. More preferably, R² is Cl or H. Most preferably, R² is H.

[0019] According to a preferred embodiment, compounds of the invention are defined according to formula I wherein R⁴ is preferably H.


[0020] According to an alternative embodiment, compounds of the invention are defined according to formula I wherein preferably R⁵ is Me.

[0021] Preferably, compounds of the invention are defined according to formula I wherein R¹¹ is Et or cyclopropyl. More preferably, R¹¹ is Et.


[0022] According to a preferred embodiment, compounds of the invention are defined according to formula I wherein Q is preferably selected from the group consisting of:




[0023] More preferably, Q is:




[0024] Alternatively, preferred embodiments of the invention include compounds selected from the group consisting of:





and



[0025] The compounds of the invention are effective inhibitors of wild type reverse transcriptase as well as inhibiting, for example, the single mutation enzymes Y181C, K103N, V106A, G190A, Y188C, and P236L. The compounds also inhibit the double mutation enzymes K103N/Y181C, K103N/P225H, K103N/V108I and K103N/L100I.

[0026] The compounds of formula I possess inhibitory activity against HIV-1 reverse transcriptase. When administered in suitable dosage forms, they are useful in the treatment of AIDS, ARC and related disorders associated with HIV-1 infection. Another aspect of the invention, therefore, is a method for treating HIV-1 infection which comprises administering to a human being, infected by HIV-1, a therapeutically effective amount of a novel compound of formula I, as described above. Whether it be termed treatment or prophylaxis, the compounds may also be used to prevent perinatal transmission of HIV-1 from mother to baby, by administration to the mother prior to birth.

[0027] The compounds of formula I may be administered in single or divided doses by the oral or parenteral routes. A suitable oral dosage for a compound of formula I would be in the range of about 0.5 mg to 1 g per day. A preferred oral dosage for a compound of formula I would be in the range of about 100 mg to 800 mg per day for a patient weighing 70 kg. In parenteral formulations, a suitable dosage unit may contain from 0.1 to 250 mg of said compounds, preferably 1 mg to 200 mg. It should be understood, however, that the dosage administration from patient to patient will vary and the dosage for any particular patient will depend upon the clinician's judgement, who will use as criteria for fixing a proper dosage the size and condition of the patient as well as the patient's response to the drug.

[0028] When the compounds of the present invention are to be administered by the oral route, they may be administered as medicaments in the form of pharmaceutical preparations, which contain them in association with a compatible pharmaceutical carrier material. Such carrier material can be an inert organic or inorganic carrier material suitable for oral administration. Examples of such carrier materials are water, gelatin, talc, starch, magnesium stearate, gum arabic, vegetable oils, polyalkylene-glycols, petroleum jelly and the like.

[0029] The pharmaceutical preparations can be prepared in a conventional manner and finished dosage forms can be solid dosage forms, for example, tablets, dragees, capsules, and the like, or liquid dosage forms, for example solutions, suspensions, emulsions and the like. The pharmaceutical preparations may be subjected to conventional pharmaceutical operations such as sterilisation. Further, the pharmaceutical preparations may contain conventional adjuvants such as preservatives, stabilisers, emulsifiers, flavour-improvers, wetting agents, buffers, salts for varying the osmotic pressure and the like. Solid carrier material which can be used include, for example, starch, lactose, mannitol, methyl cellulose, microcrystalline cellulose, talc, silica, dibasic calcium phosphate, and high molecular weight polymers (such as polyethylene glycol).

[0030] For parenteral use, a compound of formula I can be administered in an aqueous or non-aqueous solution, suspension or emulsion in a pharmaceutically acceptable oil or a mixture of liquids, which may contain bacteriostatic agents, antioxidants, preservatives, buffers or other solutes to render the solution isotonic with the blood, thickening agents, suspending agents or other pharmaceutically acceptable additives. Additives of this type include, for example,


tartrate, citrate and acetate buffers, ethanol, propylene glycol, polyethylene glycol, complex formers (such as EDTA), antioxidants (such as sodium bisulfite, sodium metabisulfite, and ascorbic acid), high molecular weight polymers (such as liquid polyethylene oxides) for viscosity regulation and polyethylene derivatives of sorbitol anhydrides. Preservatives may also be added if necessary, such as benzoic acid, methyl or propyl paraben, benzalkonium chloride and other quaternary ammonium compounds.


[0031] The compounds of this invention may also be administered as solutions for nasal application and may contain in addition to the compounds of this invention suitable buffers, tonicity adjusters, microbial preservatives, antioxidants and viscosity-increasing agents in an aqueous vehicle. Examples of agents used to increase viscosity are polyvinyl alcohol, cellulose derivatives, polyvinylpyrrolidone, polysorbates or glycerin. Microbial preservatives added may include benzalkonium chloride, thimerosal, chlorobutanol or phenylethyl alcohol.

[0032] Additionally, the compounds provided by the invention can be administered by suppository.


[0033] The compounds of the invention may be made using the skills of a synthetic organic chemist. An exemplary reaction scheme is shown in schemes 1 to 6.


Scheme 1: Preparation of intermediates in which R⁴ is Me




Scheme 2: Preparation of intermediates in which R⁵ is Me


[0034] The sequence of scheme 2 is analogous to one described by J.M. Klunder *et al.*; *J. Med. Chem.* **1998**, 41, 2960-71, and C.L. Cywin *et al.*; *J. Med. Chem.* **1998**, 41, 2972-84.

Scheme 3: Preparation of intermediates in which R² is C₁₋₃ alkyl

Scheme 4: Preparation of intermediates in which R² is CF₃

Scheme 5: Introduction of the quinoline nucleus



Scheme 6: Alternate route to compounds in which R⁴ is Me

[0035] As stated before, the compounds provided by the invention inhibit the enzymatic activity of HIV-1 RT. Based upon testing of these compounds, as described below, it is known that they inhibit the RNA-dependent DNA polymerase activity of HIV-1 RT. It is known (data not shown) that they also inhibit the DNA-dependent DNA polymerase activity of HIV-1 RT. Utilising the Reverse Transcriptase (RT) Assay described below, compounds can be tested for their ability to inhibit the RNA-dependent DNA polymerase activity of HIV-1 RT. Certain specific compounds described in the Examples which appear below, were so tested. The results of this testing appear in Table 2 as IC₅₀ (nM) and Table 3 as EC₅₀ (nM).

EXAMPLES

[0036] The present invention is illustrated in further detail by the following nonlimiting examples. All reactions were performed in a nitrogen or argon atmosphere. Temperatures are given in degrees Celsius. Solution percentages or ratios express a volume to volume relationship, unless stated otherwise. Abbreviations or symbols used herein include:

DEAD:	diethyl azodicarboxylate;
DIAD:	diisopropyl azodicarboxylate;
DIEA:	diisopropylethylamine;
DMAP:	4-(dimethylamino)pyridine;
DMSO:	dimethylsulfoxide;
DMF:	dimethylformamide;
ES MS:	electron spray mass spectrometry;
Et:	ethyl;
EtOAc:	ethyl acetate;
Et ₂ O:	diethyl ether;
HPLC:	high performance liquid chromatography;
iPr:	isopropyl;
Me:	methyl;
MeOH:	methanol;
MeCN:	acetonitrile;
NBS:	N-bromosuccinimide;
Ph:	phenyl;
TBE:	tris-borate-EDTA;
TBTU:	2-(1 <i>H</i> -benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate;
TFA:	trifluoroacetic acid;
THF:	tetrahydrofuran;
MS (ES):	electrospray mass spectrometry;
MS (FAB) or FAB/ MS:	fast atom bombardment mass spectrometry;
HRMS:	high resolution mass spectrometry;
PFU:	plaque forming units;

DEPC:	diethyl pyrocarbonate;
DTT:	dithiothreitol;
EDTA:	ethylenediaminetetraacetate;
UMP:	uridine 5'-monophosphate;
5 UTP:	uridine 5'-triphosphate;
MES:	2-(n-morpholino)ethanesulfonic acid;
SD5-PAGE:	sodium dodecyl sulfate-polyacrylamide gel electrophoresis;
MWCO:	molecular weight cut-off;
Bis-Tris	Propane: 1,3-Bis[tris(hydroxymethyl)-methylamino]propane;
10 GSH:	reduced glutathione;
OBG:	n-Octyl-β-D-glucoside.

SYNTHESES

[0037] The following examples illustrate methods for preparing compounds of the invention.

Example I (entry 24, Table 1)

5,11-Dihydro-11-ethyl-5-methyl-8-{2-(4-quinolinyloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

a) 2-(Ethylamino)-3-nitropyridine

[0038] To a solution of 2-chloro-3-nitropyridine (51 g, 325 mmol) in THF (650 mL) was added a 2 M solution of ethylamine in THF (365 mL, 731 mmol). The reaction was stirred at room temperature overnight. The reaction mixture was poured into water (~1.5 L) and the resulting solid was filtered and dried under reduced pressure to give the title compound (52 g).

b) 3-Amino-2-(ethylamino)pyridine

[0039] A solution of 2-(ethylamino)-3-nitropyridine (52 g) in MeOH (600 mL) was stirred overnight at room temperature under hydrogen (1 atm.) in the presence of 20% Pd(OH)₂/C (10.4 g). The catalyst was removed by filtration through diatomaceous earth. The filtrate was concentrated under reduced pressure to give the title compound as a black solid (39 g, 88% yield over steps a) and b).

c) 2-Chloro-N-(2-(ethylamino)-3-pyridinyl)-5-bromo-3-pyridinecarboxamide

[0040] To a cooled solution of 3-amino-2-(ethylamino)pyridine (30.6 g, 223 mmol) in MeCN (740 mL) was added solid NaHCO₃ (56.3 g, 669 mmol). After 5 min, crude 5-bromo-2-chloro-3-pyridinecarbonyl chloride (prepared from 5-bromo-2-hydroxy-3-pyridinecarboxylic acid and SOCl₂ [as described by T. W. Gero *et al.* in *Synth. Commun.* 1989, 19, 553-559 (incorporated herein by reference) but with omission of the aqueous work-up] was added (1 equiv., 223 mmol). After 2 h, the reaction mixture was poured over ice/H₂O (1.5 L) and the resulting solid was filtered, rinsed with H₂O and then hexane. After drying under reduced pressure overnight, the title compound was obtained as a black solid (54.9 g, 69 % yield).

d) 8-Bromo-5,11-dihydro-11-ethyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one

[0041] To a solution of 2-chloro-N-(2-(ethylamino)-3-pyridinyl)-5-bromo-3-pyridinecarboxamide (54.9 g, 154.4 mmol) in pyridine (308 mL) at 50 °C was added drop-wise a 1 M solution of NaHMDS (sodium hexamethyldisilazide) in THF (355 mL, 355 mmol). After 10 min, the reaction was allowed to cool to room temperature, and then was poured over ice water (2 L). The resulting solid was filtered, rinsed with water and then hexane. The solid was dried under reduced pressure to give the title compound (36 g, 75% yield) as a dark green solid.

e) 8-Bromo-5,11-dihydro-11-ethyl-5-methyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one

[0042] To a solution of the 8-bromo-5,11-dihydro-11-ethyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (36.7 g, 115 mmol) in DMF (380 mL) was added NaH (3.5 g, 138 mmol), and the mixture was heated to 50 °C for 30 min. The reaction mixture was cooled to room temperature and treated with MeI (14.3 mL, 230 mmol). After 1.5 h, the reaction mixture was poured over ice water. The solid was filtered, washed with water and then hexane to give after drying the title

compound (37.9 g 99% yield) as a dark grey solid.

f) 5,11-Dihydro-11-ethyl-5-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one

[0043] Allyltributyltin (30.7 mL, 99.0 mmol) and Pd(Ph₃P)₄ (5.20 g, 4.50 mmol) were added to a degassed (N₂ through solution for 30 min) solution of 8-bromo-5,11-dihydro-11-ethyl-5-methyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (30.0 g, 90.0 mmol) in DMF (450 mL) at room temperature. The mixture was stirred at 90 °C for 1.5 h then was cooled to room temperature and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane: EtOAc, 8:2 to 7:3) to give the title compound (22.19 g, 84% yield).

g) 5,11-Dihydro-11-ethyl-8-(2-hydroxyethyl)-5-methyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one

[0044] A stream of ozonised oxygen was bubbled through a cold (-78 °C) solution of 5,11-dihydro-11-ethyl-5-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (22.19 g, 75.4 mmol) in CH₂Cl₂ (150 mL) and MeOH (150 mL) for 2.5 h. A stream of N₂ was next bubbled through the solution for 15 min and then solid NaBH₄ (4.99 g, 132 mmol) was added to the solution. The reaction mixture was allowed to warm to room temperature. After 1 h, aqueous saturated NH₄Cl (200 mL) was added and the mixture was stirred at room temperature for 2 h. The organic solvents were removed under reduced pressure. Water (300 mL) and CHCl₃ (300 mL) were added to the residue. The phases were separated and the aqueous layer was extracted with CHCl₃ (3 × 300 mL). The combined organic layers were dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:CHCl₃, 4:1) to give the title compound (16.1 g, 72% yield) as a white solid.

h) 5,11-Dihydro-11-ethyl-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2b:2',3'-e] [1,4]diazepin-6-one

[0045] Diethyl azodicarboxylate (DEAD) (12.8 mL, 81.0 mmol) was added dropwise to a solution of 5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-5-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (16.1 g, 54.0 mmol), 4-hydroxyquinoline (11.6 g, 81.0 mmol) and Ph₃P (21.3 g, 81.0 mmol) in THF (270 mL) at room temperature. The mixture was stirred at room temperature for 1h then was concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:MeOH; 95:5) to give the title compound (17.7 g, 77% yield) as a white solid: MS (ESI) m/z 426 (MH)⁺.

Example II (entry 2, Table 1)

2-Chloro-5,11-dihydro-11-ethyl-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

a) 6-Chloro-2-(ethylamino)-3-nitropyridine

[0046] An ice-cold solution of EtNH₂ (49.8 g, 1.10 mol) in toluene (200 mL) was added over 15 min to an ice-cold solution of 2,6-dichloro-3-nitropyridine (100.0 g, 0.52 mol) in toluene (225 mL). The mixture was stirred at 0 °C for 45 min. Water (500 mL) and EtOAc (500 mL) were added and the phases were separated. The organic layer was successively washed with water (200 mL) and brine (200 mL), dried (MgSO₄), filtered and concentrated under reduced pressure. The residual solid was recrystallised from MeOH to give the title compound (83.7 g, 80% yield) as yellow needles.

b) 3-Amino-6-chloro-2-(ethylamino)pyridine

[0047] A solution of SnCl₂·H₂O (616.3 g, 2.73 mol) in aqueous 12 N HCl (500 mL) was rapidly added to a solution of 6-chloro-2-(ethylamino)-3-nitropyridine (169.5 g, 0.84 mol) in AcOH (1.7 L) at room temperature. After 20 min, the mixture was cooled to 0 °C and water (250 mL) was added. Solid NaOH (240 g) was then added in small portions. The resulting suspension was filtered to remove tin salts. The filtrate was diluted with water (3.5 L), the solution rendered basic by addition of aqueous 10 N NaOH and then extracted with EtOAc (3 × 1.7 L). The combined organic layers were washed with brine (1 L), dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 3:2) to give the title compound (89.9 g, 62% yield) as a brown solid: MS (ESI) m/z 172/174 (MH)⁺.

c) 5-Bromo-2-chloro-N-{2-(ethylamino)-6-chloro-3-pyridinyl}-3 pyridinecarboxamide

[0048] A solution of 5-bromo-2-chloro-3-pyridinecarbonyl chloride (30.0 g, 97.0 mmol) in MeCN (100 mL) was added via cannula to a solution of 3-amino-6-chloro-2-(ethylamino)pyridine (16.6 g, 97.0 mmol) in MeCN (180 mL) containing

solid NaHCO₃ (14.2 g, 169 mmol) at room temperature. The mixture was stirred at room temperature for 1 h. Water (200 mL) was added and the mixture was stirred for 10 min. The resulting suspension was filtered. The solid was washed with Et₂O (50 mL) and concentrated from pyridine solution (3 × 50 mL) to give the title compound (28.4 g, 75% yield).

d) 8-Bromo-2-chloro-5,11-dihydro-11-ethyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0049] A 1 M solution of NaHMDS in THF (167.5 mL, 167.5 mmol) was slowly added to a solution of 5-bromo-2-chloro-N-{2-(ethylamino)-6-chloro-3-pyridinyl}-3-pyridinecarboxamide (28.4 g, 72.8 mmol) in pyridine (146 mL) heated to 50 °C. The reaction mixture was stirred at 50 °C for 1.5 h. The mixture was then poured into a mixture of water and ice (1 L) and, after 1 h, the resulting suspension was filtered. The solid was washed with water and dried under reduced pressure to give the title compound (23.4 g, 91 % yield).

e) 8-Bromo-2-chloro-5,11-dihydro-11-ethyl-5-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0050] Solid NaH (60% oil dispersion, 3.46 g, 86.1 mmol) was added over 30 min to a solution of 8-bromo-2-chloro-5,11-dihydro-11-ethyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (23.4 g, 66.3 mmol) in DMF (220 mL) at 50 °C. The mixture was stirred at 50 °C for 1 h then was allowed to cool to room temperature. The mixture was poured into water (1 L) and the resulting suspension was filtered. The solid was successively washed with water and hexane then dried under reduced pressure to give the title compound (23.0 g, 94% yield).

f) 2-Chloro-5,11-dihydro-11-ethyl-5-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0051] Allyltributyltin (21.3 mL, 68.7 mmol) and Pd(Ph₃P)₄ (3.61 g, 3.12 mmol) were added to a degassed (N₂ through solution for 30 min) solution of 8-bromo-2-chloro-5,11-dihydro-11-ethyl-5-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (23.0 g, 62.5 mmol) in DMF (312 mL). The mixture was heated to 90 °C for 2 h. The mixture was concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 7:3) to give the title compound (13.4 g, 65% yield).

g) 2-Chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-5-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0052] Ozonised oxygen was introduced into a cold (-78 °C) solution of 2-chloro-5,11-dihydro-11-ethyl-5-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (13.4 g, 40.7 mmol) in MeOH (102 mL) and CH₂Cl₂ (102 mL) until complete disappearance of the alkene. Nitrogen was bubbled through the solution to remove excess O₃. Solid NaBH₄ (2.69 g, 71.1 mmol) was next added in small portions and the mixture was allowed to warm slowly to room temperature. After 1 h, aqueous saturated NH₄Cl (150 mL) was added and the mixture stirred for 20 min. The organic solvents were removed under reduced pressure. Water (100 mL) was added to the aqueous solution. The solution was extracted with CHCl₃ (3 × 200 mL). The combined organic layers were dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:CHCl₃, 4:1) to give the title compound (10.4 g, 77% yield).

h) 2-Chloro-5,11-dihydro-11-ethyl-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0053] Diethyl azodicarboxylate (DEAD) (4.26 mL, 27.0 mmol) was added dropwise to a solution of 2-chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-5-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (6.00 g, 18.0 mmol), 4-hydroxyquinoline (3.92 g, 27.0 mmol) and Ph₃P (7.09 g, 27.0 mmol) in THF (90 mL) at room temperature. The mixture was stirred at room temperature for 1 h then was concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:MeOH, 95:5) to give the title compound (6.22 g, 75% yield) as a white solid: MS (ESI) m/z 460/462.

Example III (entry 6, Table 1)

2-Chloro-11-cyclopropyl-5,11-dihydro-5-methyl-8-{2-(4-quinolinylloxy)-ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

a) 6-Chloro-2-(cyclopropylamino)-3-nitropyridine

[0054] A solution of cyclopropylamine (1.25 g, 22.0 mmol) in toluene (11 mL) was added over 10 min to an ice-cold

solution of 2,6-dichloro-3-nitropyridine (2.00 g, 10.4 mmol) in toluene (10 mL). The mixture was stirred at 0 °C for 1 h and at room temperature for 2 h. Water (50 mL) was added to the mixture and the phases were separated. The organic layer was washed with brine (25 mL), dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 3:2) to give the title compound (1.97 g, 89% yield) as a yellow solid.

b) 2-Chloro-11-cyclopropyl-5,11-dihydro-8-(2-hydroxyethyl)-5-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0055] The title compound was prepared from 6-chloro-2-(cyclopropylamino)-3-nitropyridine using a method similar to that described for the 11-ethyl analog in example II.

c) 2-Chloro-11-cyclopropyl-5,11-dihydro-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0056] Diethyl azodicarboxylate (DEAD) (85.6 µL, 0.54 mmol) was added dropwise to a solution of 2-chloro-11-cyclopropyl-5,11-dihydro-8-(2-hydroxyethyl)-5-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (125 mg, 0.36 mmol), 4-hydroxyquinoline (78.9 mg, 0.54 mmol) and Pb₃P (143 mg, 0.54 mmol) in THF (1.8 mL) at room temperature. The mixture was stirred at room temperature for 3 h then was concentrated under reduced pressure. The residue was partially purified by flash chromatography (EtOAc:MeOH; 95:5). The solid was further purified by reverse phase HPLC (CombiPrep ADS-AQ 50 × 20 mm, 5µ, 120 Å, 5-100% MeCN + 0.10% TFA / water + 0.10% TFA in 25 min) to give the trifluoroacetic acid salt of the title compound (17.7 g, 77% yield) as a white solid: MS (ESI) m/z 472/474 (MH)⁺.

Example IV (entry 4, Table 1)

5,11-Dihydro-11-ethyl-2-fluoro-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

a) 2,6-Difluoro-3-nitropyridine

[0057] To a mixture of concentrated sulphuric acid (600 mL) and fuming nitric acid (90%, 400mL) in a ice-bath (internal temperature maintained at 5-10 °C) was added drop-wise 2,6-difluoropyridine (200 g, 1.74 mol). The resulting mixture was stirred overnight at room temperature. The mixture was poured slowly into 3 kg of ice and extracted with Et₂O (2 × 2 L). The combined organic layers were washed with aqueous 1.5 N NaOH (2 × 1 L), then with aqueous saturated NaHCO₃ (400 mL) or until pH is around 8-9. The organic layers were dried over MgSO₄, filtrated and concentrated under reduced pressure until constant weight (to remove unreacted 2,6-difluoropyridine: 10-12%). The title compound was obtained as a yellow liquid (207.3 g, 74% yield).

b) 2-(Ethylamino)-6-fluoro-3-nitropyridine

[0058] To a solution of 2,6-difluoro-3-nitropyridine (45.7 g, 285 mmol) in THF (500 mL) at -40 °C was added drop-wise a solution of ethylamine (25.7 g, 570 mmol) in THF (250 mL). After 30 min, the reaction mixture was concentrated under reduced pressure and the residue was dissolved in EtOAc. The organic phase was washed with brine, dried (MgSO₄), filtered and concentrated. The resulting yellow solid was purified by flash chromatography (15% EtOAc in hexane) to give the title compound (43.2 g, 82% yield) as a yellow solid.

c) 3-Amino-2-(ethylamino)-6-fluoropyridine

[0059] A solution of 2-(ethylamino)-6-fluoro-3-nitropyridine (43.2 g, 230 mmol) in THF (1 L) was stirred overnight at room temperature under hydrogen (1 atm.) in the presence of 20% Pd(OH)₂/C (4.35 g). The catalyst was removed by filtration through diatomaceous earth. The filtrate was concentrated under reduced pressure to give the title compound (36.3 g, 95% yield) as a black solid.

d) 2-Chloro-N-{2-(ethylamino)-6-fluoro-3-pyridinyl}-5-bromo-3-pyridinecarboxamide

[0060] To a cooled solution (4 °C) of 3-amino-2-(ethylamino)-6-fluoropyridine (31.0 g, 200 mmol) in MeCN (160 mL) was added solid NaHCO₃ (50.4 g, 600 mmol). After 15 min, a solution of 5-bromo-2-chloro-3-pyridinecarbonyl chloride (1 equiv., 200 mmol) in MeCN (155 mL) was added. After 60 min at room temperature, the reaction mixture was poured into water (1.2 L) and stirred for 30 min. The resulting solid was filtered, dried under reduced pressure at 50°C overnight.

The title compound (73.7 g, 99% yield) was obtained as a black solid.

e) 8-Bromo-5,11-dihydro-11-ethyl-2-fluoro-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one

[0061] To a solution of the 2-chloro-N{2-(ethylamino)-6-fluoro-3-pyridinyl}-5-bromo-3-pyridinecarboxamide (73.5 g, 216 mmol) in pyridine (435 mL) at 50 °C was added drop-wise a 1 M solution of NaHMDS in THF (520 mL, 520 mmol). After 10 min, the reaction was allowed to cool to room temperature, then poured over ice water (2 L). The resulting solid was filtered, rinse with water and then hexane. The solid was dried under reduced pressure to give the title compound (50.6 g, 69% yield) as a dark green solid.

f) 8-Bromo-5,11-dihydro-11-ethyl-2-fluoro-5-methyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one

[0062] To a solution of the 8-bromo-5,11-dihydro-12-ethyl-2-fluoro-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (44 g, 130.5 mmol) in DMF (520 mL) was added NaH (4.28 g, 178 mmol), and the mixture was heated to 50 °C for 30 min. The reaction mixture was cooled to room temperature and treated with MeI (24.4 mL, 522 mmol). After 2.5 h, the reaction mixture was poured over ice water. The solid was filtered, washed with water and then hexane, dried under reduced pressure to give the title compound (43.2 g, 94% yield) as dark grey solid.

g) 5,11-Dihydro-11-ethyl-2-fluoro-5-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one

[0063] Allyltributyltin (32.0 mL, 103.4 mmol) and Pd(Pb₃P)₄ (5.43 g, 4.70 mmol) were added to a degassed (N₂ through solution for 45 min) solution of 8-bromo-5,11-dihydro-11-ethyl-2-fluoro-5-methyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (33.0 g, 94.0 mmol) in DMF (470 mL). Additional amounts of Pd(Ph₃)₄ (1.09 g, 0.94 mmol added after 1, 2, 3, 4, 5 h of reaction) were added to complete the reaction. The mixture was heated to 90 °C for 6 h. The mixture was concentrated under reduced pressure. The residue was purified by flash chromatography (hexanes:EtOAc, 8:2 to 7:3) to give the title compound (22.4 g, 76% yield).

h) 5,11-Dihydro-11-ethyl-2-fluoro-8-(2-hydroxyethyl)-5-methyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one

[0064] A stream of ozonised oxygen was bubbled through a cold (-78 °C) solution of 5,11-dihydro-11-ethyl-2-fluoro-5-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (22.38 g, 71.6 mmol) in CH₂Cl₂ (100 mL) and MeOH (100 mL) for 3 h. A stream of N₂ was next bubbled through the solution for 15 min and then solid NaBH₄ (5.05 g, 133 mmol) was added to the solution. The reaction mixture was allowed to warm to room temperature. After 1 h, an additional portion of NaBH₄ (1.62 g, 43.0 mmol) was added to the reaction mixture. After an additional hour, aqueous saturated NH₄Cl (150 mL) was added and the mixture was stirred at room temperature for 30 min. The organic solvents were removed under reduced pressure. Water (200 mL) was added and the mixture was extracted with CHCl₃ (3 × 300 mL). The combined organic layers were dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:CHCl₃, 4:1) to give the title compound (19.7 g, 72% yield) as a white solid.

i) 5,11-Dihydro-11-ethyl-2-fluoro-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one

[0065] Diethyl azodicarboxylate (DEAD) (14.3 mL, 91.0 mmol) was added dropwise to a solution of 5,11-dihydro-11-ethyl-2-fluoro-8-(2-hydroxyethyl)-5-methyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (19.2 g, 60.7 mmol), 4-hydroxyquinoline (13.2 g; 91.0 mmol) and Ph₃P (23.9 g, 91.0 mmol) in THF (300 mL) at room temperature. The mixture was stirred at room temperature for 1h then was concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:MeOH; 95:5) to give the title compound (17.9 g, 67% yield) as a white solid: MS (ESI) m/z 444 (MH)⁺.

Example V (entry 1, Table 1)

2-Chloro-5,11-dihydro-11-ethyl-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one

a) 2-Chloro-N-(2,6-dichloro-4-methyl-3-pyridinyl)-3-pyridinecarboxamide

[0066] NaHCO₃ (21.72 g, 258.6 mmol) was added to a solution of 3-amino-2,6-dichloro-4-methylpyridine (41.61 g, 235.1 mmol; prepared as described by K. G. Grozinger *et al.* J. Heterocyclic Chem.1995, 32, 259-263) in MeCN (350

mL) and the resulting suspension was stirred for 15 min. A solution of 2-chloronicotiny chloride (43.44 g, 246.8 mmol) in MeCN (500 mL) was then introduced over 30 min. The resulting suspension was stirred at room temperature. After 24 h, the solution was found to be acidic and so an additional amount of NaHCO₃ (3.00 g, 35.7 mmol) was added. The suspension was then stirred at room temperature for an additional 2 days. The mixture was then poured into a mixture of water (2 L) and ice (200 g) and stirred for 20 min. The solid was filtered and washed with water (500 mL) and the resulting solid was then dried over P₂O₅ under reduced pressure to give the title compound (62.55 g, 84% yield) as a white powder.

b) *N*-(2,6-dichloro-4-methyl-3-pyridinyl)-2-(ethylamino)-3-pyridinecarboxamide

[0067] A solution of 2-chloro-*N*-(2,6-dichloro-4-methyl-3-pyridinyl)-3-pyridinecarboxamide (63.17 g, 194.0 mmol) and ethylamine (28.0 g, 583 mmol) in xylenes (250 mL) was stirred at 120 to 125 °C in a steel autoclave for 7 h. The resulting suspension was poured into water (1 L), stirred for 15 min and filtered. The residue was dissolved in EtOAc and washed with water (3 ×), brine and was dried over MgSO₄. The filtrate was extracted with EtOAc and the combined organic extracts were washed with water and brine and dried over MgSO₄. The two fractions were then combined and excess xylenes removed by co-distillation with benzene to give the title compound (61.98 g, pink solid) as the major component of a mixture of compounds. The material, not purified at this point, was used directly in the subsequent reaction.

c) 5-Bromo-*N*-(2,6-dichloro-4-methyl-3-pyridinyl)-2-(ethylamino)-3-pyridinecarboxamide

[0068] Solid KOAc (22.2 g, 229 mmol) was added to a solution of crude *N*-(2,6-dichloro-4-methyl-3-pyridinyl)-2-(ethylamino)-3-pyridinecarboxamide (62.98 g, < 190 mmol) in AcOH (600 mL). A solution of Br₂ (9.7 mL, 191 mmol) was then added over 30 min. After stirring 30 min, additional amounts of KOAc and Br₂ were introduced until TLC indicated that no starting material remained (a total of 11.2 g of KOAc and 5.91 mL of Br₂). The solution was then poured into water (2 L) and stirred for 20 min. The solid was removed by filtration and then dissolved in Et₂O (1 L). The ethereal solution was washed with water (3 ×), aqueous saturated NaHCO₃, brine then dried (MgSO₄) to give a mixture containing, as the major component, the title compound (77.7 g, yellow foam): The material was used directly in the subsequent reaction.

d) 8-Bromo-2-chloro-5,11-dihydro-11-ethyl-4-methyl-6*H*-dipyrido[3,2-*b*:2',3'-*e*] [1,4]diazepin-6-one

[0069] A solution of crude 5-bromo-*N*-(2,6-dichloro-4-methyl-3-pyridinyl)-2-(ethylamino)-3-pyridinecarboxamide (77.7 g, mmol) in anhydrous pyridine (1.0 L) was heated to 50 °C. A 1 M solution of NaHMDS in THF (418 mL, 418 mmol) was then added dropwise and stirring was continued for an additional 15 min. After cooling to room temperature, water (50 mL) was added and the mixture concentrated to two-thirds volume on a rotary evaporator. The residue was then poured into water (3 L). Filtration gave the title compound (23.44 g) as a tan solid. The filtrate was extracted with EtOAc (3 ×) and the combined extracts were washed with water and brine then dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 9:1 or CHCl₃) to give an additional amount of the title compound (14.06 g, for a total of 37.50 g, 50% yield for 3 steps).

e) 2-Chloro-5,11-dihydro-11-ethyl-4-methyl-8-(2-propenyl)-6*H*-dipyrido[3,2-*b*:2',3'-*e*] [1,4]diazepin-6-one

[0070] A solution of 8-bromo-2-chloro-5,11-dihydro-11-ethyl-4-methyl-6*H*-dipyrido[3,2-*b*:2',3'-*e*] [1,4]diazepin-6-one (12.6 g, 34.3 mmol) in DMF (250 mL) was degassed under reduced pressure for 20 min. Pd(PPh₃)₄ (775 mg, 0.7 mmol) was then added followed by allyltributyltin (12.5 mL, 41.1 mmol). After degassing under reduced pressure for 10 min, the mixture was heated to 100 °C for 7 h. The mixture was then concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 19:1) to give the title compound (8.04 g, 71% yield) as a yellow solid.

f) 2-Chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-4-methyl-6*H*-dipyrido[3,2-*b*:2',3'-*e*] [1,4]diazepin-6-one

[0071] A stream of O₃ was introduced into a cold (-78 °C) solution of 2-chloro-5,11-dihydro-11-ethyl-4-methyl-8-(2-propenyl)-6*H*-dipyrido[3,2-*b*:2',3'-*e*] [1,4]diazepin-6-one (8.04 g, 24.4 mmol) and Sudan III in CH₂Cl₂ (120 mL) and MeOH (120 mL). When the pink solution turned brown, O₂ was bubbled through the solution for 10 min. NaBH₄ (1.12 g, 29.4 mmol) was then added and the solution was allowed to warm to room temperature. After 30 min, an additional amount of NaBH₄ (500 mg) was added. After 1 h, aqueous saturated NH₄Cl was then added and the mixture stirred for 20 min. The solution was concentrated under reduced pressure and extracted with CH₂Cl₂ (3 ×). The combined organic extracts were washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (CHCl₃:EtOH, 97:3) to give the title compound (6.01 g, 74% yield) as a white solid.

g) 2-Chloro-5,11-dihydro-11-ethyl-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0072] Diethyl azodicarboxylate (DEAD) (83 μ L, 0.53 mmol) was added drop-wise to a solution of 2-chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (125 mg, 0.38 mmol), 4-hydroxyquinoline (76.3 mg, 0.53 mmol) and Ph_3P (138 mg, 0.53 mmol) in THF (2.5 mL) at room temperature. The mixture was stirred at room temperature for 3 h then was concentrated under reduced pressure. The residue was dissolved in EtOAc (60 mL) and the solution was successively washed with 1 N aqueous NaOH (3×10 mL) and brine (15 mL) then was dried (MgSO_4), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:MeOH, 9:1) to give the title compound (52 mg, 30% yield) as a white solid: MS (ESI) m/z 460/462 (MH)⁺.

Example VI (entry 7, Table 1)**5,11-Dihydro-11-ethyl-2-fluoro-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one****a) 8-Bromo-5,11-dihydro-11-ethyl-2-{{(4-methoxyphenyl)methyl}amino}-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

[0073] 8-Bromo-2-chloro-5,11-dihydro-11-ethyl-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (2.50 g, 6.80 mmol) was dissolved in dioxane (10 mL) in a glass pressure bottle and 4-methoxybenzylamine (4.0 mL, 30.6 mmol) was added to the solution. The mixture was heated to 170 °C for 5 days. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The residue was taken up in EtOAc and washed with 10% aqueous NH_4Cl , dried over MgSO_4 , filtered and concentrated under reduced pressure to give a brown oil which was chromatographed over silica gel (hexane:EtOAc, 4:1). The title compound was obtained as a yellow solid (1.35 g, 42% yield).

b) 2-Amino-8-bromo-5,11-dihydro-11-ethyl-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0074] 8-Bromo-5,11-dihydro-11-ethyl-2-{{(4-methoxyphenyl)methyl}amino}-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (6.52 g, 18.7 mmol) was dissolved in trifluoroacetic acid (100 mL) and the solution was stirred magnetically at room temperature for 18 h. Trifluoroacetic acid was evaporated under reduced pressure and the residue was poured into a mixture containing water (450 mL), concentrated NH_4OH (50 mL) and EtOAc (400 mL). The mixture was stirred magnetically for 2 h at room temperature. The organic phase was concentrated under reduced pressure.

[0075] The yellow precipitate was filtered off and washed with water, dried under reduced pressure to give 7.51 g of a yellow solid which was triturated in Et_2O to give the title compound (5.12 g, 105% yield) as a yellow solid.

c) 8-Bromo-5,11-dihydro-11-ethyl-2-fluoro-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0076] A plastic bottle was charged with 2-amino-8-bromo-5,11-dihydro-11-ethyl-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (5.10 g, 14.6 mmol). HF·pyridine (100 g) was added and the resulting suspension was cooled to 0° C. Sodium nitrite (1.12 g, 16.1 mmol) was added in several portions over 30 min to produce a purple solution. The mixture was then stirred for 16 h at room temperature. The reaction mixture was poured onto ice and 6 N NaOH. The beige suspension was extracted with EtOAc. The organic extracts were, combined, dried over MgSO_4 , filtered and concentrated. The residue was triturated with Et_2O /hexane to give the title compound (4.30 g, 84% yield).

d) 5,11-Dihydro-11-ethyl-2-fluoro-4-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4] diazepin-6-one

[0077] Allyltributyltin (7.77 mL, 25.1 mmol) and $\text{Pd}(\text{Ph}_3\text{P})_4$ (1.32 g, 1.14 mmol) were added to a degassed (N_2 through solution for 30 min) solution of 8-bromo-5,11-dihydro-11-ethyl-2-fluoro-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (8.0 g, 22.8 mmol) in DMF (114 mL). The mixture was heated to 90 °C for 3h. The mixture was poured into water (250 mL) and extracted with EtOAc (4×250 mL). The combined organic layers were washed with 20% aqueous NH_4OH (250 mL) and brine (250 mL), dried (MgSO_4), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (hexanes:EtOAc, 4:1 to 1:1) to give the title compound (3.16 g, 51% yield).

e) 5,11-Dihydro-11-ethyl-2-fluoro-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0078] A stream of ozonised oxygen was bubbled through a cold (-78 °C) solution of 5,11-dihydro-11-ethyl-2-fluoro-4-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (3.16 g, 10.1 mmol) in CH_2Cl_2 (25 mL) and MeOH

(25 mL) for 45 min. A stream of N₂ was next bubbled through the solution for 15 min and then solid NaBH₄ (669 mg, 17.1 mmol) was added to the solution. The reaction mixture was allowed to warm to room temperature. After 2 h, aqueous saturated NH₄Cl (50 mL) was added and the mixture was stirred at room temperature for 15 min. The organic solvents were removed under reduced pressure. The residual aqueous solution was extracted with CHCl₃ (3 × 100 mL). The combined organic layers were dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:CHCl₃, 1:1) to give the title compound (2.40 g, 75% yield) as a white solid.

f) 5,11-Dihydro-11-ethyl-2-fluoro-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0079] Diisopropyl azodicarboxylate (DIAD) (117 µL, 0.59 mmol) was added dropwise to a solution of 5,11-dihydro-11-ethyl-2-fluoro-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (125 mg, 0.40 mmol), 4-hydroxyquinoline (86.0 mg, 0.59 mmol) and Ph₃P (156 mg, 0.59 mmol) in THF (1.9 mL) at room temperature. The mixture was stirred at room temperature for 3 h then was concentrated under reduced pressure. The residue was dissolved in EtOAc (40 mL) and the solution was successively washed with 3 N aqueous NaOH (3 × 20 mL), aqueous saturated NH₄Cl (15 mL), brine (15 mL) then was dried (MgSO₄) filtered and concentrated under reduced pressure. The residue was partially purified by flash chromatography (EtOAc:MeOH; 95:5). The solid was further purified by reverse phase HPLC (CombiPrep ADS-AQ 50 × 20 mm, 5 µ, 120 Å, 5-100% MeCN + 0.10% TFA / water + 0.10% TFA in 25 min) to give the trifluoroacetic acid salt of the title compound (106 mg, 48% yield) as a white solid: MS (ESI) m/z 444 (MH)⁺.

Example VII (entry 5, Table 1)

11-(Cyclopropyl)-5,11-dihydro-2-fluoro-4-methyl-8-{2-(4-quinolinylloxy)-ethyl}-6H-dipyrido [3,2-b: 2', 3'- e] [1,4] diazepin- 6-one

a) 11-Cyclopropyl-5,11-dihydro-2-fluoro-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0080] The title compound was prepared from 8-bromo-2-chloro-11-cyclopropyl-5,11-dihydro-4-methyl-6H-dipyrido [3,2-b:2',3'-e] [1,4]diazepin-6-one using a procedure similar to that of the corresponding 11-ethyl analog described in example VI.

b) 11-Cyclopropyl-5,11-dihydro-2-fluoro-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4] diazepin-6-one

[0081] Diisopropyl azodicarboxylate (DIAD) (131 µL, 0.66 mmol) was added dropwise to a solution of 11-(cyclopropyl)-5,11-dihydro-2-fluoro-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (140 mg, 0.44 mmol), 4-hydroxyquinoline (96.4 mg, 0.66 mmol) and Ph₃P (124 mg, 0.66 mmol) in THF (2.2 mL) at room temperature. The mixture was stirred at room temperature for 3 h then was concentrated under reduced pressure. The residue was dissolved in EtOAc (60 mL) and the solution was successively washed with 3 N aqueous NaOH (3 × 20 mL), aqueous saturated NH₄Cl (20 mL) and brine (20 mL) then was dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was partially purified by flash chromatography (EtOAc:MeOH; 95:5). The solid was further purified by reverse phase HPLC (CombiPrep ADS-AQ 50 × 20 mm, 5 µ, 120 Å, 5-100% MeCN + 0.10% TFA / water + 0.10% TFA in 25 min) to give the trifluoroacetic acid salt of the title compound (104 mg, 43% yield) as a white solid: MS (ESI) m/z 456 (MH)⁺.

Example VIII (entry 8, Table 1)

5,11-Dihydro-11-ethyl-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

a) 5,11-Dihydro-11-ethyl-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0082] A mixture of 2-chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (268 mg, 0.51 mmol), ammonium formate (318 mg, 5.1 mmol) and 10% Pd/C (50 mg) in EtOH (6 mL) was stirred at room temperature. After 24 h, additional amounts of ammonium formate (200 mg, mmol) and 10% Pd/C (50 mg) were added. After stirring an additional 24 h, the reaction was concentrated under reduced pressure. The residue was dissolved in CHCl₃ and the mixture was washed with aqueous saturated NaHCO₃, dried (MgSO₄), filtered and concentrated under reduced pressure to give the title compound (122 mg, 81%) as a yellow oil.

b) 5,11-Dihydro-11-ethyl-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0083] Diisopropyl azodicarboxylate (DIAD) (42 μ L, 0.21 mmol) was added dropwise to a solution of 5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (42.0 mg, 0.14 mmol), 4-hydroxyquinoline (31 mg, 0.21 mmol) and Ph_3P (55.4 mg, 0.21 mmol) in THF (0.7 mL) at room temperature. The mixture was stirred at room temperature for 2 h then was concentrated under reduced pressure. The residue was dissolved in EtOAc (60 mL) and the solution was successively washed with 3 N aqueous NaOH (3×20 mL), aqueous saturated NH_4Cl (20 mL) and brine (20 mL) then was dried (MgSO_4), filtered and concentrated under reduced pressure. The residue was partially purified by flash chromatography (EtOAc:MeOH; 95:5). The solid was further purified by reverse phase HPLC (CombiPrep ADS-AQ 50×20 mm, 5 μ , 120 A, 5-100% MeCN + 0.10% TFA /water + 0.10% TFA in 25 min) to give the trifluoroacetic acid salt of the title compound (14.2 mg, 19% yield) as a white solid: MS (ESI) m/z 426 (MH)⁺.

Example IX (entry 3, Table 1)**11-(Cyclopropyl)-5,11-dihydro-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one****a) 5-Bromo-2-chloro-N-(2,6-dichloro-4-methyl-3-pyridinyl)-3-pyridinecarboxamide**

[0084] 3-Amino-2,6-dichloro-4-methylpyridine (0.51 g, 2.85 mmol) was dissolved in toluene (35 mL) and pyridine (0.27 mL, 3.28 mmol) was added. 5-Bromo-2-chloro-3-pyridinecarbonyl chloride (0.80 g, 3.14 mmol) was then added dropwise over 30 min. The resulting mixture was stirred at room temperature for 1 h, diluted with water and extracted with toluene (2 $\times$). The combined organic extracts were dried (MgSO_4), filtered and concentrated under reduced pressure. The resulting thick oil was triturated with CH_2Cl_2 , and the white solid collected via suction filtration to give the title compound (0.41 g, 36% yield).

b) 5-Bromo-2-(cyclopropylamino)-N-(2,6-dichloro-4-methyl-3-pyridinyl)-3-pyridinecarboxamide

[0085] A solution of 5-bromo-2-chloro-N-(2,6-dichloro-4-methyl-3-pyridinyl)-3-pyridinecarboxamide (3.00 g, 7.61 mmol) and cyclopropylamine (2.11 mL, 30.5 mmol) in xylenes was heated in a sealed tube at 120 $^\circ\text{C}$ for 24 h. The reaction was cooled to 0 $^\circ\text{C}$, diluted with water and extracted with EtOAc (2 $\times$). The combined organic extracts were dried (MgSO_4), filtered and concentrated under reduced pressure. The resulting residue was purified by trituration (Et_2O : hexane, 3:1) to give the title compound (2.35 g, 75% yield) as a pale yellow solid.

c) 8-Bromo-2-chloro-11-cyclopropyl-5,11-dihydro-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0086] A solution of 5-bromo-2-(cyclopropylamino)-N-(2,6-dichloro-4-methyl-3-pyridinyl)-3-pyridinecarboxamide (105 mg, 0.254 mmol) in pyridine (3.0 mL) was heated to 50 $^\circ\text{C}$ while a 1 M solution of NaHMDS in THF (0.53 mL, 0.53 mmol) was added. The resulting solution was stirred at 50 $^\circ\text{C}$ for 3 h. The reaction was quenched with aqueous saturated NH_4Cl , diluted with water and extracted with EtOAc (3 $\times$). The combined organic extracts were washed with brine, dried (MgSO_4), filtered and concentrated under reduced pressure. The resulting residue was purified via flash chromatography (hexane:EtOAc, 7:3) to give the title compound (32 mg, 33% yield).

d) 2-Chloro-11-cyclopropyl-5,11-dihydro-4-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0087] A solution of 8-bromo-2-chloro-11-cyclopropyl-5,11-dihydro-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (0.046 g, 0.122 mmol) in DMF (2.0 mL) was degassed by sparging with N_2 for 10 min. $\text{Pd}(\text{PPh}_3)_4$ (2.8 mg, 0.003 mmol) and allyltributyltin (0.45 mL, 0.146 mmol) were added and the resulting solution was heated to 90 $^\circ\text{C}$ for 2.5 h. The reaction mixture was diluted with water and extracted with CH_2Cl_2 (3 $\times$) and EtOAc (2 $\times$). The combined organic extracts were dried (MgSO_4), filtered and concentrated under reduced pressure. The resulting residue was purified via flash chromatography (hexane to hexane:EtOAc, 7:3) to give the title compound (0.033 g, 80% yield).

e) 2-Chloro-11-cyclopropyl-5,11-dihydro-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0088] A cold (-78 $^\circ\text{C}$) solution of 2-chloro-11-cyclopropyl-5,11-dihydro-4-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (0.82 g, 2.43 mmol) in CH_2Cl_2 (7.5 mL) and MeOH (7.5 mL) was saturated with O_3 until the

solution turned pale blue. The reaction mixture was left at -78 °C for 10 min, then NaBH₄ (0.23 g, 9.71 mmol) was added and the mixture was allowed to warm slowly to room temperature over 2 h. The reaction was quenched with aqueous 10% citric acid solution and extracted with EtOAc (3 ×). The organic extracts were combined, washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure. The resulting residue was purified via flash chromatography (MeOH:CH₂Cl₂, 2 to 10%) to give the title compound (0.41 g, 60% yield) as a white solid.

f) 11-Cyclopropyl-5,11-dihydro-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0089] The title compound was prepared using a method similar to that described for the 11-ethyl analog in example VIIIa.

g) 11-Cyclopropyl-5,11-dihydro-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0090] Diisopropyl azodicarboxylate (DIAD) (167 µL, 0.85 mmol) was added dropwise to a solution of 11-(cyclopropyl)-5,11-dihydro-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (175 mg, 0.57 mmol), 4-hydroxyquinoline (123 mg, 0.85 mmol) and Ph₃P (223 mg, 0.85 mmol) in THF (2.8 mL) at room temperature. The mixture was stirred at room temperature for 3 h then was concentrated under reduced pressure. The residue was dissolved in EtOAc (60 mL) and the solution was successively washed with 3 N aqueous NaOH (3 × 20 mL), aqueous saturated NH₄Cl (20 mL) and brine (20 mL) then was dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:MeOH; 95:5) to give the title compound (74 mg, 30% yield) as a white solid: MS (ESI) m/z 438 (MH)⁺.

Example X (entry 21, Table 1)

5,11-Dihydro-11-ethyl-5-methyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido[3,2-b:2'3'-e] [1,4]diazepin-6-one

[0091] Solid mCPBA (80-85%, *m*-chloroperbenzoic acid) (2.21 g, 10.2 mmol) was added to a solution of 5,11-dihydro-11-ethyl-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (2.56 g, 6.02 mmol) in CH₂Cl₂ (30 mL) and THF (30 mL) at room temperature (note that the reaction can be performed in CH₂Cl₂ alone). The mixture was stirred at room temperature for 1 h then was concentrated under reduced pressure. The residue was dissolved in CHCl₃ (400 mL), the solution was successively washed with aqueous 10% Na₂S₂O₃ (3 × 75 mL), aqueous saturated NaHCO₃ (3 × 75 mL) and brine (75 mL), dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (CHCl₃:MeOH, 9:1) to give the title compound (2.01 g, 76% yield) as a white solid: MS (ESI) m/z 442 (MH)⁺.

Example XI (entry 17, Table 1)

2-Chloro-5,11-dihydro-11-ethyl-5-methyl-8-{2-[(2-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido [3,2-b: 2', 3'- e] [1,4] diazepin- 6-one

[0092] Using a procedure identical to that of example X, 2-chloro-5,11-dihydro-11-ethyl-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (2.70 g, 5.87 mmol) gave the title compound (2.05 g, 73% yield) as a white solid: MS (ESI) m/z 476/478 (MH)⁺.

Example XII (entry 11, Table 1)

5,11-Dihydro-11-ethyl-2-fluoro-5-methyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido [3,2-b: 2', 3'- e] [1,4] diazepin- 6-one

[0093] Solid mCPBA (457 mg, 2.12 mmol) was added to a solution of 5,11-dihydro-2-fluoro-11-ethyl-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (470 mg, 1.06 mmol) in CH₂Cl₂ (10 mL). The mixture was stirred at room temperature for 13 h then was concentrated under reduced pressure. The residue was dissolved in CHCl₃ (75 mL), the solution was successively washed with aqueous 10% Na₂S₂O₃ (3 × 25 mL), aqueous saturated NaHCO₃ (3 × 25 mL) and brine (25 mL), dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (CHCl₃:MeOH, 19:1 to 9:1) to give the title compound (250 mg, 51% yield) as a white solid: MS (ESI) m/z 460 (MH)⁺.

Example XIII (entry 10, Table 1)**2-Chloro-11-cyclopropyl-5,11-dihydro-5-methyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e][1,4] diazepin- 6-one**

[0094] Solid mCPBA (177 mg, 0.82 mmol) was added to a solution of 2-chloro-11-cyclopropyl-5,11-dihydro-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (143 mg, 0.30 mmol) in CH₂Cl₂ (3 mL) and THF (3 mL). The mixture was stirred at room temperature for 14 h then was concentrated under reduced pressure. The residue was dissolved in EtOAc (50 mL), the solution was successively washed with aqueous 10% Na₂S₂O₃ (3 × 10 mL), aqueous saturated NaHCO₃ (3 × 10 mL) and brine (10 mL), dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was partially purified by flash chromatography (EtOAc:MeOH, 95:5 to CH₂Cl₂:MeOH, 9:1). The impure compound was further purified by reverse phase HPLC (CombiPrep ADS-AQ 50 × 20 mm, 5 μ, 120 A, 5-100% MeCN + 0.10% TFA / water + 0.10% TFA in 25 min). The pure fractions were treated with aqueous saturated NaHCO₃, extracted with EtOAc and the solution was dried (MgSO₄) filtered and concentrated under reduced pressure to give the title compound (28.3 mg, 19%) as a white solid: MS (ESI) m/z 488/490 (MH)⁺.

Example XIV (entry 14, Table 1)**2-Chloro-5,11-dihydro-11-ethyl-4-methyl-8-{2-[(1-oxido-4-quinolinyl)-oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e][1,4] diazepin- 6-one**

[0095] Solid mCPBA (46.9 mg, 0.22 mmol) was added to a solution of 2-chloro-5,11-dihydro-11-ethyl-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (50.0 mg, 0.11 mmol) in CH₂Cl₂ (2.2 mL). The mixture was stirred at room temperature for 1 h then was concentrated under reduced pressure. The residue was dissolved in EtOAc (50 mL), the solution was successively washed with aqueous 10% Na₂S₂O₃ (3 × 10 mL), aqueous saturated NaHCO₃ (3 × 10 mL) and brine (10 mL), dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by reverse phase HPLC (CombiPrep ADS-AQ 50 × 20 mm, 5 μ, 120 A, 5-100% MeCN + 0.10% TFA / water + 0.10% TFA in 25 min). The pure fractions were treated with aqueous saturated NaHCO₃, extracted with EtOAc and the solution was dried (MgSO₄), filtered and concentrated under reduced pressure to give the title compound (32.5 mg, 63%) as a white solid: MS (ESI) m/z 488/490 (MH)⁺.

Example XV (entry 9, Table 1)**5,11-Dihydro-11-ethyl-2-fluoro-4-methyl-8-{2-[(2-oxido-4-quinolinyl)-oxy]ethyl}-6H-dipyrido [3,2-b: 2', 3'- e] [1,4] diazepin- 6-one**

[0096] Using a procedure similar to that of example XIV, 5,11-dihydro-11-ethyl-2-fluoro-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (125 mg, 0.28 mmol) gave the title compound (40 mg, 31 %) as a white solid: MS (ESI) m/z 460 (MH)⁺.

Example XVI (entry 12, Table 1)**5,11-Dihydro-11-ethyl-4-methyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin- 6-one**

[0097] Solid mCPBA (163 mg, 0.75 mmol) was added to a solution of 5,11-dihydro-11-ethyl-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (144 mg, 0.34 mmol) in CH₂Cl₂ (7.5 mL). The solution was stirred at room temperature for 2 h. The mixture was diluted with CH₂Cl₂ (125 mL), washed successively with aqueous 10% Na₂S₂O₃ (2 × 25 mL), aqueous saturated NaHCO₃ (2 × 25 mL) and brine (25 mL), then was dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (CH₂Cl₂:MeOH, 10:1) to give the title compound (102 mg, 68% yield) as a white solid: MS (ESI) m/z 442 (MH).

Example XVII (entry 15, Table 1)**11-Cyclopropyl-5,11-dihydro-4-methyl-8-{2-[(1-oxido-4-quinonlinyl)oxy]ethyl}-6H-dipyrido [3,2-b: 2', 3'- e] [1,4] diazepin- 6-one**

[0098] Using a procedure similar to that of example XIV, 11-cyclopropyl-5,11-dihydro-4-methyl-8-{2-(4-quinolinylloxy)

ethyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (37.1 mg, 0.085 mmol) gave the title compound (35 mg, 91%) as a white solid: MS (ESI) m/z 454 (MH)⁺.

Example XVIII (entry 25, Table 1)

2-Chloro-5,11-dihydro-11-ethyl-4-methyl-8-{2-[(5-quinolinylloxy)ethyl]}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0099] Diethyl azodicarboxylate (DEAD) (77 μ L, 0.49 mmol) was added drop wise to a solution of 2-chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (125 mg, 0.38 mmol), 5-hydroxyquinoline (70.9 mg, 0.49 mmol) and Ph₃P (128 mg, 0.49 mmol) in THF (2.5 mL) at room temperature. After 1.5 h, additional amounts of DEAD (35 μ L, 0.22 mmol) and PPh₃ (59 mg, 0.22 mmol) were added to the mixture. The mixture was stirred at room temperature for 1 h then was diluted in EtOAc (50 mL) and the solution was successively washed with aqueous 1 N NaOH (4 $\times$ 10 mL) and brine (15 mL) then was dried (MgSO₄) filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (CH₂Cl₂:acetone, 85:15). The resulting solid was triturated with Et₂O to give the title compound (38 mg, 22% yield) as a white solid: MS (ESI) m/z 460/462 (MH)⁺.

Example XIX (entry 20, Table 1)

2-Chloro-5,11-dihydro-11-ethyl-5-methyl-8-{2-[(1-oxido-5-quinolinyl)-oxy]ethyl}-6H-dipyrido [3,2-b: 2', 3'- e] [1,4] diazepin- 6-one

[0100] Diethyl azodicarboxylate (DEAD) (160 μ L, 1.01 mmol) was added drop-wise to a solution of 2-chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-5-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (225 mg, 0.68 mmol), 5-hydroxyquinoline (147 mg, 1.01 mmol) and Ph₃P (266 mg, 1.01 mmol) in THF (3.4 mL) at room temperature. The mixture was stirred at room temperature for 3 h then was concentrated under reduced pressure. The residue was partially purified by flash chromatography (EtOAc:MeOH; 95:5). The fractions containing the 5-quinolinylloxy derivative were concentrated, dissolved in CH₂Cl₂ (2 mL) and THF (5 mL) and treated with mCPBA (80%, 248 mg, 1.15 mmol) at room temperature. The solution was stirred at room temperature for 2.5 h, then was concentrated under reduced pressure. The residue was dissolved in EtOAc (50 mL), washed successively with aqueous 10% Na₂S₂O₃ (3 $\times$ 10 mL), aqueous saturated NaHCO₃ (3 $\times$ 10 mL) and brine (10 mL), then was dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:MeOH, 95:5 to CH₂Cl₂:MeOH, 9:1) to give the title compound (113 mg, 35% yield for 2 steps) as a white solid: MS (ESI) m/z 476/478 (NH)⁺.

Example XX (entry 18, Table 1)

5,11-Dihydro-11-ethyl-2-fluoro-5-methyl-8-{2-[(1-oxido-5-quinolinyl)-oxy]ethyl}-6H-dipyrido [3,2-b: 2,3'- e] [1,4] diazepin- 6-one

[0101] Using a procedure identical to that of example XIX, 5,11-dihydro-11-ethyl-2-fluoro-8-(2-hydroxyethyl)-5-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (267 mg, 0.84 mmol) gave the title compound (204 mg, 52% yield for 2 steps) as a white solid: MS (ESI) m/z 460 (MH)⁺.

Example XXI (entry 19, Table 1)

2-Chloro-5,11-dihydro-11-ethyl-4-methyl-8-{2-[(1-oxido-5-quinolinyl)-oxy]ethyl}-6H-dipyrido [3,2-b: 2', 3'- e] [1,4] diazepin- 6-one

[0102] Using a procedure similar to that of example XIX but using diisopropyl azodicarboxylate (DIAD) instead of diethyl azodicarboxylate, 2-chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido [3,2-b:2',3'-e] [1,4]diazepin-6-one (223 mg, 0.67 mmol) gave the title compound (43.4 mg, 14% yield for 2 steps) as a white solid: MS (ESI) m/z 476/478 (MH)⁺.

Example XXII (entry 23, Table 1)

5,11-Dihydro-11-ethyl-2-fluoro-4-methyl-8-{2-[(1-oxido-5-quinolinyl)-oxy]ethyl}-6H-dipyrido [3,2-b: 2,3'-e] [1,4] diazepin- 6-one

[0103] Using a procedure similar to that of example XXI, 5,11-dihydro-11-ethyl-2-fluoro-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (250 mg, 0.79 mmol) gave the title compound (64.7 mg, 18% yield for 2 steps) as a white solid: MS (ESI) m/z 460 (MH)⁺.

Example XXIII, Entry 22, Table 1**2-Chloro-5,11-dihydro-11-ethyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

[0104] MS (ESI) m/z 462/464 (MH)⁺.

Example XXIV, Entry 26, Table 1**5,11-Dihydro-5,11-diethyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

[0105] MS (ESI) m/z 440 (MH)⁺.

Example XXV, Entry 27, Table 1**5,21-Dihydro-5,11-diethyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

[0106] MS (ESI) m/z 456 (MH)⁺.

Example XXVI (Entry 28, Table 1)**2,5-Dimethyl-5,11-dihydro-11-ethyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one****a) 5-Bromo-2-chloro-N-(6-chloro-2-methyl-3-pyridinyl)-3-pyridinecarboxamide**

[0107] NaHCO₃ (3.9 g, 46.4 mmol) was added to a solution of 3-amino-2-chloro-6-methylpyridine (2.2 g, 15.4 mmol; prepared as described by K G. Grozinger et al. J. Heterocyclic Chem. 1995, 32, 259-263) in MeCN (50 mL). The resulting suspension was stirred for 15 min. and a solution of crude 5-bromo-2-chloro-3-pyridinecarbonyl chloride (prepared from 5-bromo-2-hydroxy-3-pyridinecarboxylic acid and SOCl₂ [as described by T. W. Gero et al. in Synth. Commun. 1989, 19, 553-559 (incorporated herein by reference) but with omission of the aqueous work-up] (4 g.) in MeCN (10 mL) was introduced over 30 min. The resulting suspension was stirred at room temperature. After 24 h, the mixture was poured into a mixture of water (100 mL) and ice (10 g) and stirred for 20 min. The suspension was filtered and the resulting solid was washed with water (50 mL) and hexane (25 mL), then dried over P₂O₅ under reduced pressure to give the title compound (1.8 g, 31% yield) as a white powder.

b) 5-Bromo-N-(2-chloro-6-methyl-3-pyridinyl)-2-(ethylamino)-3-pyridinecarboxamide

[0108] A solution of 5-bromo-2-chloro-N-(2-chloro-6-methyl-3-pyridinyl)-3-pyridinecarboxamide (1.7 g, 4.8 mmol) and ethylamine (2M in THF, 10.5 mmol, 5.2 mL) in THF (5 mL) was stirred at 90 to 95 °C in a steel autoclave for 16 h. The resulting mixture was poured into water (100 mL), stirred for 15 min and filtered. The solid was washed with water followed by hexane to give the title compound as a yellow solid (1.58 g, 89% yield).

c) 8-Bromo-5,11-dihydro-2,5-dimethyl-11-ethyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0109] A solution of crude 5-bromo-N-(2-dichloro-6-methyl-3-pyridinyl)-2-(ethylamino)-3-pyridinecarboxamide (0.7 g, 1.8 mmol) in anhydrous pyridine (18 mL) was heated to 50 °C. A 1 M solution of NaHMDS in THF (7.2 mL, 7.2 mmol) was then added dropwise and stirring was continued for an additional 3h. After cooling to room temperature, methyl iodide (0.6 mL, 9 mmol) was added and the reaction was stirred overnight. Water (25 mL) was added and the mixture was extracted with EtOAc (3 ×) and the combined extracts were washed with water and brine then dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc,

8:2) to give the title compound (259 mg, 41% yield).

d) 5,11-Dihydro-2,5-dimethyl-11-ethyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0110] A solution of 8-bromo-5,11-dihydro-2,5-dimethyl-11-ethyl-6H-dipyrido[3,2-b:2',3'-e] [1,4] diazepin-6-one (258 mg, 0.7 mmol) in DMF (7.4 mL) was degassed under reduced pressure for 20 min. Pd(PPh₃)₄ (43 mg, 0.04 mmol) was then added followed by allyltributyltin (0.3 mL, 0.85 mmol). After degassing under reduced pressure for 10 min, the mixture was heated to 100 °C for 1.5 h. The mixture was then concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 8:2) to give the title compound (184 mg, 98% yield) as a yellow solid.

e) 5,11-Dihydro-2,5-dimethyl-11-ethyl-8-(2-hydroxyethyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0111] A stream of O₃ was introduced into a cold (-78 °C) solution of 5,11-dihydro-2,5-dimethyl-11-ethyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (187 mg, 0.6 mmol) and Sudan III in CH₂Cl₂ (3 mL) and MeOH (3 mL). When the pink solution turned brown, O₂ was bubbled through the solution for 10 min. NaBH₄ (57 mg, 1.52 mmol) was then added and the solution was allowed to warm to room temperature. After 30 min, an additional amount of NaBH₄ (20 mg) was added. After 2 h, aqueous saturated NH₄Cl was then added and the mixture stirred for 20 min. The solution was concentrated under reduced pressure and extracted with CH₂Cl₂ (3x). The combined organic extracts were washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:Hexane 6:4) to give the title compound (111 mg, 58% yield) as a white solid.

f) 2,5-Dimethyl-5,11-dihydro-11-ethyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0112] Diethyl azodicarboxylate (DEAD) (139 µL, 0.88 mmol) was added dropwise to a solution of 5,11-dihydro-2,5-dimethyl-11-ethyl-8-(2-hydroxyethyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (110 mg, 0.35 mmol), 4-hydroxyquinoline (128 mg, 0.88 mmol) and Pb₃P (231 mg, 0.88 mmol) in THF (3.5 mL) at room temperature. The mixture was stirred at room temperature for 16h. The reaction mixture was diluted in EtOAc (60 mL) and the solution was successively washed with 1 N aqueous NaOH (3 × 10 mL) and brine (15 mL) then was dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc then EtOAc:MeOH, 9.5:0.5) to give the title compound (57 mg, 37% yield) as a white solid: MS (ESI) m/z 440 (MH)⁺.

g) 5,11-Dihydro-2,5-dimethyl-11-ethyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

[0113] Solid mCPBA (80-85%, m-chloroperbenzoic acid) (60 mg, 0.28 mmol) was added to a solution of 5,11-dihydro-2,5-dimethyl-11-ethyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (55 mg, 0.13 mmol) in CH₂Cl₂ (1.3 mL) at room temperature. The mixture was stirred at room temperature for 2.5 h then diluted with CH₂Cl₂. The resulting solution was successively washed with aqueous 10% Na₂S₂O₃, aqueous saturated NaHCO₃ and brine, dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (MeOH/EtOAc, 2% to 5%) to give the title compound (59 mg, 99% yield) as a white solid: MS (ESI) m/z 456 (MH)⁺.

Example XXVII (Entry 29, Table 1)

5,11-Dihydro-11-ethyl-5-methyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one

a) 2-(Ethylamino)-3-nitro-6-(trifluoromethyl)pyridine

[0114] To a solution of 2-nitroacetamide ammonium salt (4 g, 33 mmol) in water (165 mL) was added piperidinium acetate (33 mmol) in water followed by slow addition of 4-ethoxy-1,1,1-trifluoro-3-buten-2-one (6.1 mL, 43 mmol) in MeOH (11 mL). The reaction was stirred for 2 h at room temperature and at reflux for 3 h. The reaction mixture was allowed to cool to 45 °C and aqueous HCl (1N) was added until the pH was acidic. After 1 h at room temperature, the reaction mixture was extracted with CH₂Cl₂ (3x). The combined organic layer was successively washed with water and brine, dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc, then EtOAc:MeOH, 9:1) to give 3-nitro-6-(trifluoromethyl)-2(1H)-pyridinone (2.7 g, 40% yield) as a yellow solid.

[0115] To a solution of 3-nitro-6-(trifluoromethyl)-2(1H)-pyridinone (2.4 g, 11.5 mmol) in DMF (69 mL) was added

sodium hydride (350 mg, 13.8 mmol). The reaction was stirred for 30 min at 45 °C, cooled to room temperature and N-phenyltrifluoromethanesulfonimide (4.9 g, 13.8 mmol) was added. After 1 h, a 2 M solution of ethylamine in THF (13 mL, 25 mmol) was added. The reaction was stirred at room temperature overnight. The reaction mixture was poured into water (100 mL) and extracted with EtOAc (3x). The combined organic layer was successively washed with water and brine, dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 9:1) to give the desired compound (1.1 g, 41 % yield) as a yellow oil.

b) 3-Amino-2-(ethylamino)-6-(trifluoromethyl)pyridine

[0116] A solution of 2-(ethylamino)-3-nitro-6-(trifluoromethyl)pyridine (1.1 g, 4.9 mmol) in MeOH (40 mL) was stirred overnight at room temperature under hydrogen (1 atm.) in the presence of 20% Pd(OH)₂/C (100 mg). The catalyst was removed by filtration through diatomaceous earth. The filtrate was concentrated under reduced pressure to give the title compound as an orange oil (1 g).

c) 2-Chloro-N-{2-(ethylamino)-6-(trifluoromethyl)-3-pyridinyl}-5-bromo-3-pyridinecarboxamide

[0117] To a cooled solution of 3-amino-2-(ethylamino)-6-trifluoromethyl pyridine (1 g, 4.9 mmol) in MeCN (24 mL) was added solid NaHCO₃ (906 mg, 11 mmol). After 5 min, crude 5-bromo-2-chloro-3-pyridinecarbonyl chloride (prepared from 5-bromo-2-hydroxy-3-pyridinecarboxylic acid and SOCl₂ [as described by T. W. Gero et al. in Synth. Commun. 1989, 19, 553-559 (incorporated herein by reference) but with omission of the aqueous work-up] was added (1 equiv., 4.9 mmol). After 2 h, the reaction mixture was poured over ice/H₂O (1.5 L) and the resulting solid was filtered, rinsed with H₂O and then hexane. After drying under reduced pressure overnight, the title compound was obtained as a light brown solid (1.6 g, 77% yield).

d) 8-Bromo-5,11-dihydro-11-ethyl-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one

[0118] To a solution of 2-chloro-N-{2-(ethylamino)-6-(trifluoromethyl)-3-pyridinyl}-5-bromo-3-pyridinecarboxamide (766 mg, 1.8 mmol) in DMF (18 mL) was added NaH (137 mg, 5.4 mmol). The reaction was stirred at 80 °C. After 1 h the reaction was allowed to cool to room temperature, and then was poured in water (20 mL). The resulting mixture was extracted with EtOAc (3x). The combined organic layer was successively washed with water and brine, dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 7:3) to give the desired compound (200 mg, 28% yield) as a yellow solid.

e) 8-Bromo-5,11-dihydro-11-ethyl-5-methyl-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one

[0119] To a solution of the 8-bromo-5,11-dihydro-11-ethyl-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (200 mg, 0.52 mmol) in DMF (2.6 mL) was added NaH (20 mg, 0.78 mmol), and the mixture was heated to 50 °C for 30 min. The reaction mixture was cooled to room temperature and treated with MeI (97 µL, 1.56 mmol). After 16 h, the reaction mixture was diluted with water. The resulting mixture was extracted with EtOAc (3x). The combined organic layer was successively washed with water and brine, dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 8.5:1.5) to give the title compound (110 mg 53% yield) as a white foam.

f) 5,11-Dihydro-11-ethyl-5-methyl-8-(2-propenyl)-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one

[0120] Allyltributyltin (98 µL, 0.32 mmol) and Pd(Ph₃P)₄ (16 mg, 0.02 mmol) were added to a degassed (N₂ through solution for 30 min) solution of 8-bromo-5,11-dihydro-11-ethyl-5-methyl-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (11 mg, 0.27 mmol) in DMF (1.4 mL) at room temperature. The mixture was stirred at 90 °C for 1.5 h then was cooled to room temperature and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 8:2 to 7:3) to give the title compound (101 mg, 99% yield).

g) 5,11-Dihydro-11-ethyl-8-(2-hydroxyethyl)-5-methyl-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one

[0121] A stream of ozonised oxygen was bubbled through a cold (-78 °C) solution of 5,11-dihydro-11-ethyl-5-methyl-8-(2-propenyl)-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (100 mg, 0.27 mmol) in CH₂Cl₂ (2.7 mL) and MeOH (2.7 mL) for 2.5 h. A stream of N₂ was next bubbled through the solution for 15 min and then solid NaBH₄

(52 mg, 1.4 mmol) was added to the solution. The reaction mixture was allowed to warm to room temperature. After 1 h, aqueous saturated NH_4Cl (5 mL) was added and the mixture was stirred at room temperature for 2 h. The reaction mixture was diluted with CH_2Cl_2 and the aqueous layer was re-extracted with CH_2Cl_2 . The combined organic layer was successively washed with water and brine, dried (MgSO_4), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:hexane: 75:25 to EtOAc 100%) to give the title compound (60 mg 59% yield) as a white solid.


h) 5,11-Dihydro-11-ethyl-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-2-(trifluoromethyl)-6H-dipyrido[3,2b:2',3'-e][1,4]diazepin-6-one

[0122] Diethyl azodicarboxylate (DEAD) (38 μL , 0.24 mmol) was added dropwise to a solution of 5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-5-methyl-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (58 mg, 0.16 mmol), 4-hydroxyquinoline (35 mg, 0.24 mmol) and Ph_3P (62 g, 0.24 mmol) in THF (1.6 mL) at room temperature. The mixture was stirred at room temperature for 1h then was concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:MeOH; 95:5) to give the title compound (24 mg, 31 % yield) as a white solid: MS (ESI) m/z 494, 495 (MH)⁺.

i) 5,11-Dihydro-11-ethyl-5-methyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one

[0123] Solid mCPBA (80-85%, m-chloroperbenzoic acid) (16 mg, 0.08 mmol) was added to a solution of 5,11-dihydro-11-ethyl-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (22 mg, 0.04 mmol) in CH_2Cl_2 (0.5 mL) at room temperature. The mixture was stirred at room temperature for 2.5 h then diluted with CH_2Cl_2 . The resulting solution was successively washed with aqueous 10% $\text{Na}_2\text{S}_2\text{O}_3$, aqueous saturated NaHCO_3 and brine, dried (MgSO_4), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (MeOH: CHCl_3 , 5% to 10%) to give the title compound (11 mg, 52% yield) as a white solid: MS (ESI) m/z 511 (MH)⁺.

TABLE 1
Compounds of formula I



I

Entry #	R ²	R ⁴	R ⁵	R ¹¹	Q
1	Cl	Me	H	Et	
2	Cl	H	Me	Et	
3	H	Me	H	CycPr	
4	F	H	Me	Et	
5	F	Me	H	CycPr	
6	Cl	H	Me	CycPr	
7	F	Me	H	Et	






Table 1 Entry #	R ²	R ⁴	R ⁵	R ¹¹	Q
8	H	Me	H	Et	
9	F	Me	H	Et	
10	Cl	H	Me	CycPr	
11	F	H	Me	Et	
12	H	Me	H	Et	
14	Cl	Me	H	Et	
15	H	Me	H	CycPr	



I

Table 1 Entry #	R ²	R ⁴	R ⁵	R ¹¹	Q
17	Cl	H	Me	Et	
18	F	H	Me	Et	
19	Cl	Me	H	Et	
20	Cl	H	Me	Et	
21	H	H	Me	Et	
22	Cl	H	H	Et	
23	F	Me	H	Et	



I

Entry #	R ²	R ⁴	R ⁵	R ¹¹	Q
24	H	H	Me	Et	
25	Cl	Me	H	Et	
26	H	H	Et	Et	
27	H	H	Et	Et	
28	Me	H	Me	Et	
29	CF ₃	H	H CH ₃	Et	

REVERSE TRANSCRIPTASE (RT) ASSAYS

ASSAY THEORY:

[0124] Among the enzymes for which Human Immunodeficiency Virus (HIV-1) encodes is a reverse transcriptase (1), so-named because it transcribes a DNA copy from an RNA template. This activity can be quantitatively measured in a cell-free enzyme assay and is based upon the observation that reverse transcriptase is able to use a synthetic template poly r(C) primed with a biotinylated oligo d(G) to transcribe a radio-labelled DNA strand utilising 3H-dGTP as a substrate. The assay described below utilises the wild-type enzyme (which is the predominant form of the enzyme observed in

patients infected with HIV-1) and can also be used with mutant RT enzymes (for example, Y181C, prepared by site-directed mutagenesis in which the tyrosine residue at codon 181 has been replaced by a cysteine residue) in analogous assay conditions. This assay allows compounds to be evaluated for their effectiveness at inhibiting the mutant enzymes.

MATERIALS:

a) Preparation of the enzyme

[0125] Some HIV-1 IIB clone BH10 RT mutants were provided by Dr. C.-K. Shih (Boehringer Ingelheim Pharmaceuticals Inc., U.S.A.) in the vector pKK233-2 (Pharmacia). Briefly an HIV RT clone pKRT2 containing only the RT p66 gene regulated by the lac operon/trc promoter was obtained from Dr. W. Summers (Yale University) (2). A variety of specific amino acid substitutions were introduced into the wild-type RT gene by site-directed mutagenesis. RT clones were subcloned into the pKK233-2 bacterial expression vector. Clones provided included wild-type, Val106Ala, Tyr181Cys, Tyr188Cys, Tyr188Leu, Gly190Ala and Pro236Leu. Others were made in-house by site-directed mutagenesis of the pKK233-2 RT clones including Lys103Asn, Lys103Asn/Tyr181Cys, Lys103Asn/Leu100Ile, Lys103Asn/Pro225His, and Lys103Asn/Val108Ile.

b) Purification of Enzyme

[0126] Purification of recombinant reverse transcriptase was performed using a combination of methods previously described (3). A single colony from a fresh plate of transformed JM109 cells was used to initiate growth of a pre-culture grown o/n at 37°C. Two liters of growth medium were inoculated with this pre-culture. At OD₆₀₀ ~ 1.5 (5-6 h at 37°C), RT gene expression was induced with IPTG (1 mM final), and the fermentation was continued for a few more hours at 37 °C. After centrifugation, supernatants were discarded while cell pellets were pooled and stored at -80 °C until purification. Cells were thawed at 4 °C overnight and suspended in lysis buffer (MES 50mM pH 6, EDTA 1mM, 10% v/v glycerol, 0.02% w/v OBG, 0.02% w/v sodium azide). Lysozyme was added and the mixture was incubated on ice for 40 minutes. After homogenization using a Dounce in presence of lysozyme and sonication, the cells were centrifuged for 30 minutes. Supernatant (S1) was saved and stored at 4°C. The centrifuged pellet was resuspended in extraction buffer (MES 50mM pH 6, KPO₄ 50 mM pH 6, KCl 100mM, 10 % v/v glycerol, 0.02% w/v OBG, 0.02% w/v sodium azide) and stirred for 30 minutes at 4°C. This second mixture was centrifuged again and the supernatant (S2) was saved. The above procedure was repeated 2 more times saving supernatants S3 and S4 and one last extraction was performed overnight (S5). Polymyxin P (0.005% final) was added to the combined supernatants to remove nucleic acids. This solution was stirred for 75 minutes at 4 °C and centrifuged for 1 h. The supernatant (SS1) was precipitated on ice with 20% w/v ammonium sulfate and stirred for 1 h at 4 °C. The mixture was then centrifuged and the resulting supernatant (SS2) was precipitated with additional 40% w/v ammonium sulfate (60% total), stirred for 1 h and centrifuged again. The final pellet (P1) was stored overnight at 4 °C before undergoing purification the following day. All steps of the purification were performed at 4 °C unless otherwise stated.

[0127] Pellet (P1) was resuspended into MES 50mM pH 6, KPO₄ 10 mM pH 6, KCl 100mM, 10% v/v glycerol, 0.02% w/v OBG, 0.02% w/v sodium azide. The suspension was dialysed against the same buffer overnight using 12-14kD MWCO dialysis tubing. The dialysate was centrifuged and the supernatant was filtered through Millex-PF 0.8 µm filter units. The filtered sample was loaded on a Hydroxy Apatite column (30-mL bed volume) and washed with the same buffer. The bound enzyme was eluted with ~220 mL of a linear gradient of 10 to 300mM KPO₄ in the above buffer. The fractions containing p66/p51 heterodimer (as determined by SDS-PAGE 8% and Western blotting) were pooled for the next column. The RT containing fractions were diluted two-fold with Bis-Tris propane 50 mM pH 7.0, 0.02% w/v OBG, 10% v/v glycerol, 0.02% w/v sodium azide and loaded on a Hi-Trap Heparin Sepharose column (5-mL bed volume) and washed with the same buffer. The bound RT was then eluted with 75 mL of a linear gradient of 0 to 1 M ammonium sulfate in the same buffer. RT-containing fractions were pooled according to SDS-PAGE and Western blotting analyses. Protein concentration of this pool was determined by the Bradford method using BSA as standard. The final enzyme preparation was dialyzed in MES 50mM pH 6, KPO₄ 300 mM pH 6, KCl 175mM, 10% v/v glycerol, 0.02% w/v sodium azide and aliquoted and frozen at -80° C.

ASSAY PROCEDURE:

[0128] The radiometric enzyme assay has been adapted to a 96-well microtiter plate format and uses streptavidin scintillation proximity beads. The assay is briefly described below. The HIV-1 RT enzyme was thawed and appropriately diluted into Tris/HCl 50 mM pH 7.8 containing NaCl 60mM, MgCl₂ hexahydrate 2mM, DTT 6mM, GSH 2mM and 0.02% w/v Chaps to give ~3 nM enzyme. To 30 µL of this enzyme solution was added 10 µL of inhibitor solution (50 µM to 2.5 nM inhibitor in same assay buffer as above containing 15% v/v DMSO). The plate was pre-incubated for 15 minutes at

room temperature before proceeding to the next step. In this pre-incubation step, the highest and lowest inhibitor concentrations were 12.5 μ M and 0.62 nM respectively and the concentration of DMSO was 3.75% v/v. Then the enzymatic reaction was initiated by addition of 10 μ L of substrates solution. The final reaction mixture contained Tris/HCl 50 mM pH 7.8, NaCl 60 mM, $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ 2 mM, DTT 6 mM, GSH 2 mM, Chaps 0.02% w/v DMSO 3% v/v, Poly rC 179 nM, Biotin dG₁₅ 18 nM, dGTP 288 nM, ³H-dGTP 71 nM, and 1-2 nM enzyme.

[0129] In this incubation step, the highest and lowest inhibitor concentrations were 10 μ M and 0.5 nM respectively. After addition of substrates, the plate was covered with a plastic seal and incubated for 1 hour at 37°C in a dry incubator. Then the reaction was quenched by addition of 75 μ L of EDTA 0.5M containing 5mg/mL of streptavidin scintillation proximity beads.

[0130] The plate was shaken for 2 minutes at medium speed and incubated 1 hour at room temperature. Then 75 μ L of cesium chloride 7 M solution was added, the plate was shaken for 2 minutes at medium speed and incubated again for 1 hour at room temperature. The plate was then covered with a plastic seal and counted using the TopCount-NXT™ Microplate Scintillation & Luminescence Counter, (Packard). Each well was counted for 60 seconds.

[0131] Each row contained at its extremities a blank and a control well.

[0132] The calculation for percent inhibition is as follows:

$$\% \cdot \text{inhibition} = \left(1 - \left[\frac{\text{cpm} \cdot \text{well} - \text{cpm} \cdot \text{blank}}{\text{cpm} \cdot \text{control} - \text{cpm} \cdot \text{blank}} \right] \right) * 100$$

[0133] Using the above assay, compounds of the invention were tested for inhibition of RT wild-type (WT) and mutant enzymes. The results are listed in Table 2, as IC₅₀ (nM).

[0134] To confirm the ability of these compounds to inhibit HIV replication, they were also tested in the human T-Cell Culture (Syncytia) Assay described below.

ELISA assay for assessment of activity in cell culture

[0135] Compounds of the invention were tested for their ability to inhibit HIV replication in cell culture in a 96-well plate assay. Complete RPMI 1640, consisting of RPMI 1640 +10% fetal bovine serum, 10 μ g/ml gentamycin and 10 μ M β -mercaptoethanol was used for dilution of the compound as well as cell growth medium. The T lymphocyte cell line C8166 was infected at a multiplicity of infection of 0.001 with viruses coding for wild-type and mutant reverse transcriptase. Cells were then incubated for three days in the presence of serial dilutions of the compounds of the invention. The supernatant was pooled from eight replica wells and the concentration of extracellular p24 was determined using a commercially available HIV-1 p24 antigen assay kit (Beckman-Coulter®). The level of inhibition (% inhibition) was calculated with the following equation:

$$\% \cdot \text{inhibition} = \left(1 - \left[\frac{\text{p24 pg / mL} \cdot \text{inhibitor}}{\text{p24 pg / mL} \cdot \text{control}} \right] \right) * 100$$

[0136] The results are listed in Table 3, as EC₅₀ (nM).

Specificity Assays

[0137] In order to assess the specificity of the enzyme inhibitory activity of the compounds provided by the invention, a few were tested, for their ability to inhibit other viral polymerases (such as Hepatitis C virus and Respiratory Syncytial virus RNA-dependent RNA polymerases) as well as mammalian polymerases (such as Calf Thymus RNA-dependent DNA polymerase and Human telomerase) using known assay methods. None of the compounds so tested was observed to possess any significant inhibitory activity against these enzymes. These results indicate that the enzyme inhibitory activity of the compounds provided by the invention is directed rather specifically against HIV-1 RT.

REFERENCES (INCORPORATED HEREIN BY REFERENCE)

[0138]

1. Benn, S., et al. Science 230:949,1985.

2. D'Aquila, R. T. and Summers, W. C. J. Acq. Imm. Def. Syn. 2:579,1989.

3. a) Warren, T. C. et al. Protein Expression & Purification 3:479,1992; b) Kohlstaedt, L. A. Science 256(5065): 1783,1992.

TABLE 2
Inhibition of Wild-type and mutant strains of RT for compounds of formula I

compound no. (see Table 1)	IC ₅₀ W/T RT (nM)	IC ₅₀ K103N (nM)	IC ₅₀ Y181C (nM)	IC ₅₀ V106A (nM)	IC ₅₀ P236L (nM)	IC ₅₀ Y188L (nM)	IC ₅₀ G190A (nM)	IC ₅₀ K103N/ Y181C (nM)	IC ₅₀ K103N/ P225H (nM)	IC ₅₀ Y188C (nM)	IC ₅₀ K103N/ V108I (nM)	IC ₅₀ K103N/ L100I (nM)
1	10.5	19	20	132		2740		52	31.2		29	28.5
2	5.5	28.5	25	148	50.5	258	14.5	24.5	27	7.1	31	46
3	16.5	45	73	79		3236		612	56		47	124
4	11	22.5	26	206		520		52	27		32	108
5	19	37.5	76	61		2541		536	38		38	98
6	25											
7	9.8	26	38	74		4318		199	46		45	98
8	10	37	43	190		7511		301	67		63	135
9	14	23	43	98		3395		175	25		58	88
10	41	67	206	1372		1422		710			255	229
11	9.5	18	27	336	56	661	19	55	15	6.6	55	102
12	12	31	38	121		>10000		212	38		75	124
14	15	19	28	77		2298		61	30		59	24
15	61	66	226	206		>10000		1522	93		215	270
17	14	25	29	319	60	198	25	48	23	8.1	83	40
18	11	37	41	737		1167		146	20		96	158
19	7	25	20	89		3938		91	16		56	27

compound no. (see Table 1)	IC ₅₀ WTRT (nM)	IC ₅₀ K103N (nM)	IC ₅₀ Y181C (nM)	IC ₅₀ V106A (nM)	IC ₅₀ P236L (nM)	IC ₅₀ Y188L (nM)	IC ₅₀ G190A (nM)	IC ₅₀ K103N/ Y181C (nM)	IC ₅₀ K103N/ P225H (nM)	IC ₅₀ Y188C (nM)	IC ₅₀ K103N/ V108I (nM)	IC ₅₀ K103N/ L100I (nM)
20	10	36	22	661		439		51	17		64	44
21	17	44	51	1106	129	1357	26	95	40	9.9	120	168
22	45			2493		2990		336				
23	6.1			105		5614		247				
24	5.6			344		627	25	70				169
25	8	19	17	84		2537		64	23		38	28
26	8.9	40	51	1745		5204		200	85		147	443
27	38			2271		5335		299				
28	19			605		1190	25	141	25	9.0	35	28
29	35			893		578		84				


TABLE 3
Inhibition of wild type and mutant strains of HIV in cell cultures by compounds of formula I

compound no. (see Table 1)	EC ₅₀ WTRT (nM)	EC ₅₀ K103N (nM)	EC ₅₀ Y181C (nM)	EC ₅₀ V106A (nM)	EC ₅₀ Y188L (nM)	EC ₅₀ K103N/ Y181C (nM)	EC ₅₀ K103N/ P225H (nM)	EC ₅₀ K103N/ V108I (nM)	EC ₅₀ K103N/ L100I (nM)
1	2.6								
2	2.2								
3	1.3								
4	0.98								
5	0.83								
6									
7	1.1								
8	0.93								
9	0.92			5	118	7.6			
10									
11	0.34	2	4.1	28	52	7.5	2.0	3.9	3.2
12	0.26	2.8	2.7	12.4			1.5	5.0	2.9
14	0.67								
15									
17	0.34	1.5	2.8	24	58	2.3	1.2	2.7	1.2
18	0.34					9.5			
19	0.37					22			

compound no. (see Table 1)	EC ₅₀ WTRT (nM)	EC ₅₀ K103N (nM)	EC ₅₀ Y181C (nM)	EC ₅₀ V106A (nM)	EC ₅₀ Y188L (nM)	EC ₅₀ K103N/ Y181C (nM)	EC ₅₀ K103N/ P225H (nM)	EC ₅₀ K103N/ V108I (nM)	EC ₅₀ K103N/ L100I (nM)
20	0.40					11			
21	0.53	3.5	4.8	65	67	4.3	2.9	5.0	5.9
22									
23									
24									
25	2.2								
26	2.2								
28	1.3	2.3	2.6	42	70	7.5			
29	1.6			83	65	3.5			

Claims

1. A compound of the general formula I:



wherein


R² is selected from the group consisting of H, F, Cl C₁₋₄ alkyl, C₃₋₄ cycloalkyl and CF₃;

R⁴ is H or Me;

R⁵ is H, Me or Et, with the proviso that R⁴ and R⁵ are not both Me, and if R⁴ is Me then R⁵ cannot be Et;

R¹¹ is Me, Et, cyclopropyl, propyl, isopropyl, or cyclobutyl; and .


Q is selected from the group consisting of:



or a pharmaceutically acceptable salt thereof.


2. A compound according to claim 1, wherein R¹¹ is Et or cyclopropyl or a pharmaceutically acceptable salt thereof.

3. A compound according to claim 1, wherein Q is selected from the group consisting of:




or a pharmaceutically acceptable salt thereof.

4. A compound according to claim 1, wherein Q is:




or a pharmaceutically acceptable salt thereof.

5. A compound according to claim 1, wherein R⁵ is Me, or a pharmaceutically acceptable salt thereof.
6. A compound according to claim 1, wherein R¹¹ is Et or a pharmaceutically acceptable salt thereof.
7. A compound according to claim 1 of the formula:




or a pharmaceutically acceptable salt thereof.

8. A compound according to claim 1 of the formula:



or a pharmaceutically acceptable salt thereof.

9. A compound according to claim 1 of the formula:




or a pharmaceutically acceptable salt thereof.

10. An inhibitor of HIV replication of the general formula I according to one of the claims 1 to 9, or a pharmaceutically acceptable salt thereof.
11. Use of a compound of the general formula I according to one of the claims 1 to 9, or a pharmaceutically acceptable salt thereof, for the manufacture of an inhibitor of HIV replication.
12. Use of a compound of formula I according to claim 1, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment or prevention of HIV infection.
13. A pharmaceutical composition for the treatment or prevention of HIV infection, comprising a compound of formula I according to claim 1, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

Patentansprüche

1. Verbindung der allgemeinen Formel I



worin:


R^2 ausgewählt ist aus der Gruppe bestehend aus H, F, Cl, C_{1-4} -Alkyl, C_{3-4} -Cycloalkyl und CF_3 ;

R^4 H oder Me ist;

R^5 H, Me oder Et ist, mit der Maßgabe, dass R^4 und R^5 nicht beide Me sind und, wenn R^4 Me ist, R^5 dann nicht Et sein kann;

R^{11} Me, Et, Cyclopropyl, Propyl, Isopropyl oder Cyclobutyl ist; und


Q ausgewählt ist aus der Gruppe bestehend aus



oder ein pharmazeutisch verträgliches Salz davon.


2. Verbindung nach Anspruch 1, worin R¹¹ Et oder Cyclopropyl ist, oder ein pharmazeutisch verträgliches Salz davon.

3. Verbindung nach Anspruch 1, worin Q ausgewählt ist aus der Gruppe bestehend aus



oder ein pharmazeutisch verträgliches Salz davon.

4. Verbindung nach Anspruch 1, worin Q




ist, oder ein pharmazeutisch verträgliches Salz davon.

5. Verbindung nach Anspruch 1, worin R⁵ Me ist, oder ein pharmazeutisch verträgliches Salz davon.


6. Verbindung nach Anspruch 1, worin R¹¹ Et ist, oder ein pharmazeutisch verträgliches Salz davon.

7. Verbindung nach Anspruch 1 der Formel




oder ein pharmazeutisch verträgliches Salz davon.

8. Verbindung nach Anspruch 1 der Formel



oder ein pharmazeutisch verträgliches Salz davon.

9. Verbindung nach Anspruch 1 der Formel



oder ein pharmazeutisch verträgliches Salz davon.

10. Inhibitor der HIV-Replikation der allgemeinen Formel I nach einem der Ansprüche 1 bis 9 oder ein pharmazeutisch verträgliches Salz davon.


11. Verwendung einer Verbindung der allgemeinen Formel I nach einem der Ansprüche 1 bis 9 oder eines pharmazeutisch verträglichen Salzes davon für die Herstellung eines Inhibitors der HIV-Replikation.

12. Verwendung einer Verbindung der Formel I nach Anspruch 1 oder eines pharmazeutisch verträglichen Salzes davon zur Herstellung eines Medikaments zur Behandlung oder Vermeidung von HIV-Infektion.

13. Pharmazeutische Zusammensetzung zur Behandlung oder Vermeidung von HIV-Infektion, umfassend eine Verbindung der Formel I nach Anspruch 1 oder ein pharmazeutisch verträgliches Salz davon und einen pharmazeutisch verträglichen Träger.

Revendications

1. Composé de formule générale I :



où


R² est choisi dans le groupe consistant en H, F, Cl, C₁₋₄ alkyle, C₃₋₄ cycloalkyle et CF₃ ;

R⁴ est H ou Me ;

R⁵ est H, Me ou Et, avec la condition que R⁴ et R⁵ ne soient pas l'un et l'autre Me, et si R⁴ est Me, alors R⁵ ne peut pas être Et ;

R¹¹ est Me, Et, cyclopropyle, propyle, isopropyle ou cyclobutyle ; et


Q est choisi dans le groupe consistant en :



ou sel pharmaceutiquement acceptable de celui-ci.


2. Composé selon la revendication 1 où R¹¹ est Et ou cyclopropyle ou sel pharmaceutiquement acceptable de celui-ci.

3. Composé selon la revendication 1 où Q est choisi dans le groupe consistant en :



ou sel pharmaceutiquement acceptable de celui-ci.

4. Composé selon la revendication 1 où Q est :




ou sel pharmaceutiquement acceptable de celui-ci.

5. Composé selon la revendication 1 où R⁵ est Me, ou sel pharmaceutiquement acceptable de celui-ci.


6. Composé selon la revendication 1 où R¹¹ est Et, ou sel pharmaceutiquement acceptable de celui-ci.

7. Composé selon la revendication 1 de formule :




ou sel pharmaceutiquement acceptable de celui-ci.

8. Composé selon la revendication 1 de formule :



ou sel pharmaceutiquement acceptable de celui-ci.

9. Composé selon la revendication 1 de formule :



ou sel pharmaceutiquement acceptable de celui-ci.

10. Inhibiteur de réplication de VIH de formule générale I selon l'une des revendications 1 à 9 ou sel pharmaceutiquement acceptable de celui-ci.

11. Utilisation d'un composé de formule générale 1 selon l'une des revendications 1 à 9, ou d'un sel pharmaceutiquement acceptable de celui-ci, pour la fabrication d'un inhibiteur de réplication de VIH.

12. Utilisation d'un composé de formule I selon la revendication 1, ou d'un sel pharmaceutiquement acceptable de celui-ci, pour la fabrication d'un médicament pour le traitement ou la prévention d'une infection à VIH.

13. Composition pharmaceutique pour le traitement ou la prévention d'une infection à VIH comprenant un composé de formule I selon la revendication 1, ou un sel pharmaceutiquement acceptable de celui-ci, et un support pharmaceutiquement acceptable.