

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 308 436 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

07.05.2003 Bulletin 2003/19(21) Application number: **01955658.8**(22) Date of filing: **10.08.2001**

(51) Int Cl.7: **C07C 217/10**, C07D 213/30,
C07D 333/16, C07D 307/42,
C07D 277/24, A61K 31/135,
A61K 31/18, A61K 31/275,
A61K 31/4402, A61K 31/381,
A61K 31/341, A61K 31/426

(86) International application number:

PCT/JP01/06903

(87) International publication number:

WO 02/014259 (21.02.2002 Gazette 2002/08)

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**

Designated Extension States:

AL LT LV MK RO SI(30) Priority: **11.08.2000 JP 2000244536****27.04.2001 JP 2001132879**(71) Applicant: **Japan Tobacco Inc.****Tokyo 105-8422 (JP)**

(72) Inventors:

- **SHINAGAWA, Yuko**
c/o Central Pharm. Research Inst.
Takatsuki-shi, Osaka 569-1125 (JP)

- **KATSUSHIMA, Takeo**

c/o Central Pharm. Res. Inst.
Takatsuki-shi, Osaka 569-1125 (JP)

- **NAKAGAWA, Takashi**


c/o Central Pharm. Res. Inst.
Takatsuki-shi, Osaka 569-1125 (JP)

(74) Representative:

von Kreisler, Alek, Dipl.-Chem. et al
Patentanwälte
von Kreisler-Selting-Werner
Postfach 10 22 41
50462 Köln (DE)

(54) **CALCIUM RECEPTOR ANTAGONISTS**

(57) A compound of the formula [I]



wherein R¹ is optionally substituted aryl group or optionally substituted heteroaryl group; R² is optionally substituted C₁₋₆ alkyl group, C₃₋₇ cycloalkyl group and the like; R³ is hydrogen atom, C₁₋₆ alkyl group, hydroxyl group and the like; R⁴ is hydrogen atom, C₁₋₆ alkyl group and the like; R⁵ and R⁶ are each C₁₋₆ alkyl group and the like; R⁷ is optionally substituted aryl group or optionally substituted heteroaryl group; X¹, X² and X³ are each C₁₋₆ alkylene group and the like; and X⁴ and X⁵ are each a single bond, methylene group and the like, a salt thereof, a solvate thereof or a prodrug thereof, and a pharmaceutical composition containing the compound, particularly a calcium receptor antagonist and a therapeutic agent for osteoporosis, are provided. The compound of the present invention is useful as a therapeutic drug of diseases accompanied by abnormal calcium homeostasis, or osteoporosis, hypoparathyroidism, osteosarcoma,

EP 1 308 436 A1

periodontal disease, bone fracture, steoarthrosis, chronic rheumatoid arthritis, Paget's disease, humoral hypercalcemia, autosomal dominant hypocalcemia and the like.

In addition, an intermediate for the compound is provided.

Description**Technical Field**

[0001] The present invention relates to a compound having a calcium-sensing receptor (CaSR, hereinafter to be simply referred to as a calcium receptor) antagonistic action, a pharmaceutical composition containing the compound, particularly a calcium receptor antagonist and a therapeutic agent of osteoporosis, and to an intermediate compound useful for synthesizing the compound.

Background Art

[0002] Calcium receptors sense extracellular Ca^{2+} concentration and increase intracellular Ca^{2+} , thereby suppressing the production of parathyroid hormone (PTH) involved in the control of Ca^{2+} metabolism and bone metabolism.

[0003] The serum calcium concentration of healthy mammal is strictly maintained at about 9-10 mg/100 ml (ca. 2.5 mM), which is referred to as calcium homeostasis of living organisms. When this value falls to not more than 50%, tetania occurs, and when it increases by 50%, consciousness is clouded, both cases threatening the lives. For maintaining calcium homeostasis, duodenum acts as a Ca^{2+} uptake organ, bone acts as a Ca^{2+} storage organ, and kidney acts as a Ca^{2+} excretory organ. These Ca^{2+} kinetics are controlled by various hormones generally referred to as "calcium controlling hormone". Representative hormone includes active vitamin D [$1,25(\text{OH})_2\text{D}_3$], PTH, calcitonin, Parathyroid Hormone-Related Protein (PTH-related Protein (PTHrP)) and the like.

[0004] Bone plays an important role not only as a supporting framework and motor organ of the body, but also as a storage organ of Ca^{2+} , which is its constituent component. To fulfill such functions, bone tissues repeat formation thereof (osteogenesis) and absorption thereof (bone resorption) throughout the entire life. For osteogenesis, osteoblast derived from mesenchymal cell plays a major role, and for bone resorption, osteoclast derived from hematopoietic cell plays a major role. Osteogenesis is a mechanism including osteoid formation by bone organic matrix (bone matrix proteins such as type I collagen and the like) produced by osteoblast present on the osteogenesis surface, and subsequent calcification. Bone resorption is a mechanism including adhesion of osteoclast to the bone surface, intracellular absorption of Ca^{2+} via acid secretion and ion transport, and excretion of absorbed Ca^{2+} to the bone marrow side, thereby releasing Ca^{2+} into blood. The deficient part of the bone absorbed by osteoclast is repaired by osteogenesis by osteoblast. This series of phenomena are called remodeling of bone, and by the remodeling, old bones are replaced by new bones, thus maintaining the strength of the entire bone while maintaining calcium homeostasis.

[0005] PTH is a hormone that plays a key role in maintaining the calcium homeostasis. When blood Ca^{2+} concentration decreases, secretion of PTH from the parathyroid gland is promoted immediately, which, in the bone, acts on osteoblast (activation of osteoclast by osteoblast, production of bone organic matrix decomposition enzyme and the like) to promote osteoclastic bone resorption, whereby Ca^{2+} is transferred from the bone into the blood. In kidney, PTH promotes resorption of Ca^{2+} in the distal convoluted tubule, and activates $25(\text{OH})$ vitamin D_3 in the proximal tubule, thereby promoting the production of active vitamin D_3 [$1,25(\text{OH})_2\text{D}_3$] having a function of promoting resorption of Ca^{2+} from the intestine. It also suppresses resorption of phosphorus. As mentioned above, PTH directly or indirectly increases blood Ca^{2+} concentration.

[0006] When blood Ca^{2+} concentration increases, calcium receptor senses it, immediately suppresses secretion of PTH from the parathyroid gland to decrease the amount of Ca^{2+} to be supplied into the blood [Brown, E.M., Homeostatic mechanisms regulating extracellular and intracellular calcium metabolism, in the parathyroids, p.19, (1994), Raven press, New York]. Secretion of PTH is also suppressed by active vitamin D [$1,25(\text{OH})_2\text{D}_3$].

[0007] Because PTH is a hormone assuming an important role in controlling Ca^{2+} metabolism and bone metabolism, attempts have been made to apply PTH to the treatment of osteoporosis. In 1982, Tam et al. found that sustained administration of bovine PTH (1-84) to thyroid/parathyroid gland enucleated rat results in promotion of both osteogenesis and bone resorption of femoral cancellous bone, leading to a decrease in net bone mass, but subcutaneous intermittent administration thereof does not result in promotion of bone resorption but in promotion of osteogenesis alone, leading to an increase in the bone mass [Endocrinology, 110, 506-512 (1982)]. Furthermore, Uzawa et al. compared the actions of sustained administration and intermittent administration of PTH with regard to epiphyseal long bone and metaphyseal cancellous bone of young rat. As a result, they clarified that sustained administration of PTH results in remarkable increase in bone mass in metaphyseal cancellous bone highly susceptible to the effect of enchondral ossification, though associated with abnormal findings such as hyperplasia of epiphyseal plate cartilage, fibrous osteitis and the like, and in marked promotion of bone resorption and decrease in bone mass accompanied by rarefaction of cortical bone, in epiphyseal cancellous bone where the effect is small [Bone, 16, 477-484 (1995)]. In addition, it has been reported that intermittent administration of PTH results in significant increases in bone mass and bone trabecula in both epiphyseal and metaphyseal cancellous bones without increase in osteoclast or decrease of cortical bone.

[0008] Moreover, Scutt et al. have reported that, in chicken calvaria derived osteoblast, a short time (10-20 min)

treatment with PTH promotes cell growth as compared to a long time (18 hr) treatment [Calcif. Tissue Int., 55, 208-215 (1994)]. This suggests that some of the actions of PTH on osteoblast are temporary and that expression of the action by the treatment for an extremely short time may be related to the fact that sustained administration and intermittent administration of PTH in vivo show different actions on bone tissues.

[0009] Ishizuya et al. further clarified through investigation of the action of PTH on differentiation of osteoblast using an in vitro experiment system that the action of PTH varies depending on the treatment time. They have reported that sustained action of PTH on osteoblast derived from rat calvaria resulted in strong inhibition of differentiation of osteoblast and nearly complete inhibition of osteogenesis in vitro, but repeated PTH action for the first 6 hr of 48 hr as one cycle resulted in significant promotion of differentiation of osteoblast and promotion of osteogenesis in vitro.

[0010] PTH is considered to not only prevent decrease in bone mass of osteoporosis model, but also has a bone mass recovery effect even on an animal already suffering from marked decrease in bone mass. Wronski et al. intermittently administered human PTH (1-34) to 90-day-old SD rat at 4 weeks post-ovariectomy and showing an obvious decrease in cancellous bone, for 15 weeks from 4 weeks post-ovariectomy. As a result, promotion of osteogenesis and inhibition of bone resorption were observed during the period of from week 5 to week 10 after the start of the administration, showing increased bone mass of about twice the bone mass of sham operation group [Endocrinology, 132, 823-831 (1993)]. They have also reported that, in this experiment, estrogen and bisphosphonate prevented decrease in bone mass caused by ovariectomy but did not show increase in bone mass, unlike PTH. They detailedly analyzed the cortical bone of this experiment system and found images showing promoted osteogenesis and bone mass increase on the periosteal side and endosteum side by intermittent administration of human PTH (1-34), based on which they have clarified that the increase in cancellous bone due to PTH did not accompany decrease in cortical bone [Bone, 15, 51-58 (1994)].

[0011] Furthermore, Mosekilde et al. have reported that intermittent administration of human PTH (1-34) or human PTH (1-84) causes not only an increase in bone mass but also a dose-dependent increase in compression strength and bending strength, which are indices of bone substance, of cancellous bone [Endocrinology, 129, 421-428 (1991)] and cortical bone [J. Bone Miner. Res., 8, 1097-1101 (1993)] of rat vertebral bone. As discussed above, since PTH shows an obvious bone mass increasing action in experimental animals, various investigations are ongoing as regards the restrictive conditions expected in actual clinical applications. Mizoguchi studied whether or not a pharmacological effect is observed by intermittent administration of PTH, even when PTH in blood, which is considered to be one of the factors responsible for osteoporosis, has significantly increased, and concluded that the bone mass increased as usual [Journal of Japanese Society of Bone Morphometry, vol. 5, pp. 33 - 39 (1995)]. Takao et al. have studied the frequency of PTH administration and reported that administration of once a week for 12 weeks to healthy rat scarcely promoted bone absorption but dose-dependently increased the bone mass [Japanese Journal of Bone Metabolism, vol. 12 (Suppl.), p. S343 (1994)], suggesting possible effectiveness of clinically useful low frequency administration. The foregoing achievements suggest the possibility of PTH for making a potent and promising therapeutic drug for the treatment of postmenopausal osteoporosis or postovariectomy osteoporosis, which increases bone mass and decreases bone fracture rate.

[0012] These results clearly indicate that intermittent administration of PTH would enable treatment of osteoporosis. On the other hand, PTH problematically requires injection as an administration route, which is painful for many patients. However, an orally administrable pharmaceutical agent that can intermittently increase PTH concentration in blood is greatly expected to become a therapeutic drug of osteoporosis, which is based on a new action mechanism different from that of the above-mentioned PTH and conventional calcitonin.


[0013] Calcium receptor is a G protein coupled receptor, which is cloned as a molecule essential for controlling PTH secretion, and which penetrates cell membrane 7 times. Human calcium receptor consists of 1078 amino acids, and shows 93% amino acid homology with bovine calcium receptor. Human calcium receptor consists of a large N terminal extracellular region consisting of 612 amino acids, a cell membrane penetration region consisting of 250 amino acids and a C terminal intracellular region consisting of 216 amino acids.

[0014] Expression of calcium receptor has been found in parathyroid gland, kidney, thyroid C cell, brain and the like, as well as in bone (bone marrow cells).

[0015] When calcium receptor is bound with a ligand such as Ca^{2+} and the like, it activates phospholipase C in conjugation with G protein, causes production of inositol triphosphate and increase in intracellular Ca^{2+} concentration, and as a result, suppresses secretion of PTH [Nature, 366, 575-580 (1993)].

[0016] As mentioned above, a pharmaceutical agent that inhibits activation of calcium receptor, or a pharmaceutical agent that antagonizes calcium receptor, removes suppression of PTH secretion in parathyroid gland cells, and promotes secretion of PTH. If the antagonistic action can increase blood PTH concentration discontinuously and intermittently, its antagonist is expected to show the same effect as provided by intermittent administration of PTH, and a pharmaceutical agent extremely effective for the treatment of osteoporosis is considered to be provided.

[0017] As a CaSR antagonist, international publication WO99/51569 describes a compound of the following formula



wherein

Y_1 is a covalent bond, alkylene or alkenylene of up to 4 carbon atoms, unsubstituted or substituted by C_{1-4} alkyl, or O;
 Y_2 is methylene, unsubstituted or substituted by C_{1-4} alkyl or haloalkyl;

Y_3 is a covalent bond or O, S, N- R^{IV} or C_{1-4} alkylene-O, C_{1-4} alkylene-S, C_{1-4} alkylene-N- R^{IV} ;

R_3 and R_4 are, independently, methyl or ethyl, or, together, form cyclopropyl;

R_5 is aryl or fused aryl, dihydro or tetrahydro fused aryl, unsubstituted or substituted with any substituents being selected from the group consisting of OH, halogen, C_{1-4} alkyl, C_{1-4} alkoxy, C_{3-6} cycloalkyl, OSO_2R^{IV} , CN, NO_2 , OCF_3 , CF_3 , CH_2CF_3 , $(CH_2)_nCO_2R^{IV}$, and $O-(CH_2)_nCO_2R^{IV}$, wherein n is an integer from 0 to 3 and R^{IV} is selected from the group consisting of H, C_{1-4} alkyl, and C_{3-6} cycloalkyl;

or R_5 is heteroaryl or fused heteroaryl; wherein the hetero-ring contains N, O or S, and is aromatic, dihydro or tetrahydro, unsubstituted or substituted with any substituents being selected from the group consisting of OH, OCH_3 , $CH(CH_3)_2$, halogen, C_{1-4} alkyl, C_{1-4} alkoxy, C_{3-6} cycloalkyl, OSO_2R^{IV} , CN, NO_2 , OCF_3 , CF_3 , CH_2CF_3 , $(CH_2)_nCO_2H$, $(CH_2)_nCO_2R^{IV}$, and $O-(CH_2)_nCO_2R^{IV}$; G is a covalent bond, CHR_6 or $C-R_6$, wherein R_6 is H, OH or O (forming a ketone) ;

R_7 is H, OH, or O- C_{1-4} alkyl;

R_8 is H or C_{1-4} alkyl; or R_7 and R_8 together form a ketone;

A and B are, independently, selected from the group consisting of a bond, CH_2 , NH, O, S and C=O, provided that A or B is selected from CH_2 and NH; or A and B together form a bond; or the A-B moiety is represented by $CH=CH$ or $C\equiv C$;

wherein


X_1 and X_5 are independently selected from the group consisting of H, halogen, CN, NO_2 , C_{1-4} alkyl, cycloalkyl, CH_2 -aryl, and CH_2 -heteroaryl; provided that either X_1 or X_5 is H;

X_2 , X_3 and X_4 are selected from the group consisting of H, halogen, O- C_{1-4} alkyl, O-aryl, O-heteroaryl, CH_2 -aryl, CH_2 -heteroaryl, alkyl, C(O)aryl, C(O)heteroaryl, $CH(OH)$ aryl, $CH(OH)$ heteroaryl and J-K;

J is a covalent bond, alkylene, O-alkylene or alkenylene of up to 5 carbon atoms, unsubstituted or substituted by a substituent selected from the group consisting of C_{1-4} alkyl, OH, O(forming a ketone), aryl, heteroaryl, and $NR'R''$, wherein R' and R'' are independently selected from the group consisting of H, alkyl, aryl, heteroaryl, C(O)alkyl, C(O)aryl, and C(O)heteroaryl;


K is selected from the group consisting of CO_2R^{IV} , OH, and CN; and pharmaceutically acceptable salts and complexes thereof.

[0018] Particularly, a compound of the formula wherein Y_3 is C_{1-4} alkylene-O



is suggested, though not directly described. In this case, however, "C₁₋₄ alkylene" means a straight chain, and the branched chain such as in the present invention is not described or suggested.

[0019] In addition, international publication WO99/51241 describes, as a CaSR antagonist, a compound of the following formula



wherein

Y₁ is a covalent bond, alkylene or alkenylene of up to 4 carbon atoms, unsubstituted or substituted by C₁₋₄ alkyl or O;

Y₂ is methylene, unsubstituted or substituted by C₁₋₄ alkyl or haloalkyl;

Y₃ is covalent bond or selected from the group consisting of O, S, N-R^{IV}, C₁₋₄ alkylene-O, C₁₋₄ alkylene-S, and C₁₋₄ alkylene-N-R^{IV};

R^{IV} is selected from the group consisting of H, C₁₋₄ alkyl, and C₃₋₆ cycloalkyl;

R₃ and R₄ are, independently, methyl or ethyl, or, together, form cyclopropyl;

R₅ is heteroaryl or fused heteroaryl; wherein the hetero-ring contains N, O or S, and is aromatic, dihydro or tetrahydro, unsubstituted or substituted with any substituents being selected from the group consisting of OH, OCH₃, CH(CH₃)₂, halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, OSO₂R^{IV}, CN, NO₂, OCF₃, CF₃, CH₂CF₃, (CH₂)_nCO₂H, (CH₂)_nCO₂R^{IV}, and O-(CH₂)_nCO₂R^{IV}; n is an integer of from 0 to 3;


G is a covalent bond, CHR₆ or C-R₆, wherein R₆ is H, OH or O (forming a ketone) ;


R₇ is H, OH, or O-C₁₋₄ alkyl;

R₈ is H or C₁₋₄ alkyl; or R₇ and R₈ together form a ketone;

A and B are, independently, selected from the group consisting of a bond, CH₂, NH, O, S and C=O, provided that either A or B is selected from CH₂ and NH; or A and B together form a bond; or A-B moiety is represented by CH=CH or C≡C;

X is selected from sub formulas (Ia) to (Ie) hereinbelow:





wherein

W is selected from the group consisting of R_1 , SO_2R_1 , $C(O)R_1$, $SO_2NR_1R_1'$, $C(O)NR_1R_1'$, $C(O)OR_1$, and SO_3R_1' , wherein R_1 and R_1' are independently selected from the group consisting of hydrogen, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{2-5} alkenyl, C_{2-5} alkynyl, heterocycloalkyl, aryl and aryl C_{1-4} alkyl; or R_1 and R_1' together form a 3 to 7 membered optionally substituted heterocyclic ring; wherein any substituents are selected from the group consisting of CN, aryl, CO_2R , CO_2NHR , OH, OR, NH_2 , halo, CF_3 , OCF_3 and NO_2 ; wherein R represents C_{1-4} alkyl, or C_{3-6} cycloalkyl; X_1 is selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R', OR', CF_3 , OCF_3 and OSO_2R' , wherein R' represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;

X_2 , X_3 and X_4 are, independently, selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R'', OR'', CF_3 , OCF_3 and OSO_2R'' , provided that either X_1 or X_3 is H, wherein R'' is C_{1-4} alkyl or haloalkyl; or X_1 and X_2 together form an aryl or heteroaryl ring, substituted or unsubstituted; wherein the heteroatom is selected from N, S, and O; and any substituents are selected from the group consisting of halo, C_{1-4} alkyl, OCF_3 , CF_3 , OMe, CN, OSO_2R' and NO_2 ; or X_3 and X_4 independently represent $C(O)R_1$; and

R_2 is selected from the group consisting of hydrogen, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{2-5} alkenyl, C_{2-5} alkynyl, heterocycloalkyl, aryl and aryl- C_{1-4} alkyl;

X_1'' is selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R, OR, CF_3 , OCF_3 and OSO_2R , wherein R represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;

X_2'' , X_3'' and X_4'' are, independently, selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R', OR', CF_3 , OCF_3 and OSO_2R' , provided that either X_1'' or X_3'' is H, wherein R' is C_{1-4} alkyl or haloalkyl; or X_1'' and X_2'' together form an aryl or heteroaryl ring, substituted or unsubstituted; wherein the heteroatom is selected from N, S and O and any substituents are selected from the group consisting of halo, C_{1-4} alkyl, OCF_3 , CF_3 , OMe, CN, OSO_2-C_{1-4} alkyl, OSO_2-C_{3-6} cycloalkyl and NO_2 ;

or X_3'' and X_4'' independently represent $C(O)R_1$; and

R_1'' and R_2'' are, independently, selected from the group consisting of hydrogen, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{2-5} alkenyl, C_{2-5} alkynyl, heterocycloalkyl and aryl; or R_1'' and R_2'' together form a 3 to 7 membered optionally substituted heterocyclic ring; wherein any substituents are selected from the group consisting of CN, aryl, CO_2R'' , CO_2NHR'' , OH, OR'', NH_2 , halo, CF_3 , OCF_3 and NO_2 ;

wherein R'' represents C_{1-4} alkyl, or C_{3-6} cycloalkyl; X_1''' is selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R, OR, CF_3 , OCF_3 and OSO_2R , wherein R represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;

X_2''' , X_3''' and X_4''' are, independently, selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R', OR', CF_3 , OCF_3 and OSO_2R' , provided that either X_1''' or X_3''' is H, wherein R' is C_{1-4} alkyl or haloalkyl;

or X_1''' and X_2''' together form an aryl or heteroaryl ring, substituted or unsubstituted; wherein the heteroatom is selected from N, S and O and the substituents are selected from the group consisting of halo, C_{1-4} alkyl, OCF_3 , CF_3 , OMe, CN, OSO_2-C_{1-4} alkyl, OSO_2-C_{3-6} cycloalkyl and NO_2 ;

or X_3''' and X_4''' independently represent $C(O)R_1$;

R_1''' and R_2''' are, independently, selected from the group consisting of hydrogen, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{2-5} alkenyl, C_{2-5} alkynyl, heterocycloalkyl and aryl; or R_1''' and R_2''' together form a 3 to 7 membered optionally substituted heterocyclic ring; wherein any substituents are selected from the group consisting of CN, aryl, CO_2R''' , CO_2NHR''' , OH, OR'', NH_2 , halo, CF_3 , OCF_3 and NO_2 ; wherein R'' represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;

D is selected from the group consisting of H, CN, NO_2 , Cl, F, Br, I, R, OR, SR, CF_3 , OCF_3 and OSO_2R , wherein R represents C_{1-4} alkyl, C_{3-6} cycloalkyl, or C_{1-10} aryl or heteroaryl wherein the heteroatom is selected from N, S and O and substituents are selected from the group consisting of halo, C_{1-4} alkyl, OCF_3 , CF_3 , OMe, CN, OSO_2-C_{1-4} alkyl, OSO_2-C_{3-6} cycloalkyl and NO_2 ; n is the integer of 1 or 2;

each E is independently C or N, provided that less than two E moieties is N; further provided that when n is 2, each E is C; a and b are optionally present bonds;

R_1^{IV} is selected from the group consisting of $(CR_2)_nCO_2R'$, $(CH_2)_nCO_2H$, $(CH_2)_nCONR_2'$, $(CH_2)_nCH_2OR'$, OR',

SR', CN, NO₂, Cl, F, BR, I, H, CF₃, OCF₃, OSO₂R', R' and H; wherein R' is C₁₋₄ alkyl, or C₃₋₆ cycloalkyl; or R₁^{IV} is O, forming a ketone such that YR₁^{IV} represents -C=O; R₂^{IV} is selected from the group consisting of hydrogen, CN, NO₂, Cl, F, Br, I, H, R'', OR'', CF₃, OCF₃, and OSO₂R''; wherein R'' represents C₁₋₄ alkyl, or C₃₋₆ cycloalkyl.

Y is selected from the group consisting of C, CH, O, N and S; provided that when Y is S, R₁^{IV} is O or not present; further provided that when Y is O, R₁^{IV} is not present;

X' is CH₂, NH, O and S.

R₉ is O-alkyl, O-CH₂-aryl, and O-aryl;


X₁^{'''} is selected from the group consisting of CN, NO₂, Cl, F, Br, I, H, R, OR, CF₃, OCF₃ and OSO₂R, wherein R represents C₁₋₄ alkyl, or C₃₋₆ cycloalkyl;

X₂^{'''}, X₃^{'''}, and X₄^{'''} are, independently, selected from the group consisting of CN, NO₂, Cl, F, Br, I, H, R', OR', CF₃, OCF₃ and OSO₂R', provided that either X^{'''}₁ or X^{'''}₃ is H, wherein R' is C₁₋₄ alkyl or haloalkyl;

or X₁^{'''} and X₂^{'''} together form an aryl or heteroaryl ring, substituted or unsubstituted; wherein the heteroatom is selected from N, S and O and the substituents are selected from the group consisting of halo, C₁₋₄ alkyl, OCF₃, CF₃, OMe, CN, OSO₂-C₁₋₄ alkyl, OSO₂-C₃₋₆ cycloalkyl and NO₂;


or X₃^{'''} and X₄^{'''} independently represent C(O)R₁; and pharmaceutically acceptable salts and complex thereof.

[0020] Again, a compound of the formula wherein Y₃ is C₁₋₄ alkylene-O



is suggested, though not directly described. In this case, however, "C₁₋₄ alkylene" means a straight chain, and the branched chain such as in the present invention is not described or suggested.

[0021] International publication WO98/45255 (EP-A-973730) also describes a compound of the following formula as a CaSR antagonist



wherein

Y₁ is a covalent bond, alkylene or alkenylene of up to 4 carbon atoms, unsubstituted or substituted by C₁₋₄ alkyl;

Y₂ is methylene, unsubstituted or substituted by C₁₋₄ alkyl or CF₃;

Z is selected from the group consisting of a covalent bond, O, S, NH, N-C₁₋₄ alkyl, O(CH₂)_n, (CH₂)_nO, NR^{'''}C=O and C=ONR^{'''}, where R^{'''} is C₁₋₄ alkyl and n is an integer from 1 to 3,

R₃ and R₄ are, independently, methyl or ethyl, or, together, form cyclopropyl;

R₅ is phenyl or naphthyl, unsubstituted or substituted with one or more substituents selected from the group consisting of OH, C₁₋₄ alkyl CH(CH₃)₂, halo, halo C₁₋₄ alkyl, C₁₋₄ alkoxy, C₃₋₆ cycloalkyl, OSO₂R^{IV}, CN, NO₂, OCF₃, CF₃ and CH₂CF₃, wherein R^{IV} represents C₁₋₄ alkyl or C₃₋₆ cycloalkyl;


G is a covalent bond or C-R₆ wherein R₆ is H, OH or O (forming a carbonyl moiety);

R₇ is H, OH, or O-C₁₋₄ alkyl;

R₈ is H or C₁₋₄ alkyl; or R₇ and R₈ together form a carbonyl moiety;

the A-B moiety is represented by CH₂CH₂, a covalent bond, -CH=CH- or -C≡C-; and

X is selected from the group consisting of sub formulae (Ia), (Ib), (Ic), (Id) and (Ie) hereinbelow:



wherein

in sub formula (Ia):

W is selected from the group consisting of R_1 , SO_2R_1 , $C(O)R_1$, $SO_2NR_1R_1'$, $C(O)NR_1R_1'$ and $C(O)OR_1SO_3R_1'$, wherein R_1 and R_1' are independently selected from the group consisting of hydrogen,
 C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{2-5} alkenyl, C_{2-5} alkynyl, heterocycloalkyl aryl and aryl C_{1-4} alkyl; or R_1 and R_1' together form a 3- to 7-membered optionally substituted heterocyclic ring; wherein any substituents are selected from the group consisting of CN, aryl, CO_2R , CO_2NHR , OH, OR, NH_2 , halo, CF_3 , OCF_3 and NO_2 ; wherein R represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;
 X_1 is selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R' , OR', CF_3 , OCF_3 and OSO_2R' , wherein R' represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;
 X_2 , X_3 and X_4 are, independently, selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R'' , OR'', CF_3 , OCF_3 and OSO_2R'' , wherein R'' is C_{1-4} alkyl or haloalkyl; or X_1 and X_2 together form an aryl or heteroaryl ring, substituted or unsubstituted; wherein the heteroatom is selected from N, S and O; and any substituents are selected from the group consisting of halo, C_{1-4} alkyl, OCF_3 , CF_3 , OMe, CN, OSO_2R' and NO_2 ; or X_3 and X_4 independently represent $C(O)R_1$; provided that when there are multiple halo substitutions in the haloalkyl, halo represents F; also provide that either X_1 or X_3 is hydrogen; and
 R_2 is selected from the group consisting of hydrogen, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{2-5} alkenyl, C_{2-5} alkynyl, heterocycloalkyl aryl and aryl- C_{1-4} alkyl;

in sub formula (Ib):

X_1'' is selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R, OR, CF_3 , OCF_3 and OSO_2R , wherein R represents C_{1-4} alkyl, or C_{3-6} cycloalkyl;
 X_2'' , X_3'' and X_4'' are, independently, selected from the group consisting of CN, NO_2 , Cl, F, Br, I, H, R' , OR', CF_3 , OCF_3 and OSO_2R' , wherein R' is C_{1-4} alkyl or haloalkyl; provided that when there are multiple halo substitutions in the haloalkyl, halo represents F, or X_1'' and X_2'' together form an aryl or heteroaryl ring, substituted or unsubstituted; wherein the heteroatom is selected from N, S and O and any substituents are selected from the group con-

sisting of halo, C₁₋₄ alkyl, OCF₃, CF₃, OMe, CN, OSO₂-C₁₋₄ alkyl, OSO₂-C₃₋₆ cycloalkyl and NO₂; or X₃" and X₄" independently represent C(O)R₁;

provided that either X₁" or X₃" is hydrogen; and

R₁" and R₂" are, independently, selected from the group consisting of hydrogen, C₁₋₄ alkyl, C₃₋₆ cycloalkyl, C₂₋₅ alkenyl, C₂₋₅ alkynyl, heterocycloalkyl and aryl; or R₁" and R₂" together form a 3 to 7 membered optionally substituted heterocyclic ring; optionally containing an additional heteroatom selected from O, S and N; wherein any substituents are selected from the group consisting of CN, aryl, CO₂R", CO₂NHR", OH, OR", NH₂, halo, CF₃, OCF₃ and NO₂; wherein R" represents C₁₋₄ alkyl, or C₃₋₆ cycloalkyl;

in sub formula (Ic):

X₁" is selected from the group consisting of CN, NO₂, Cl, F, Br, I, H, R, OR, CF₃, OCF₃ and OSO₂R, wherein R represents C₁₋₄ alkyl, or C₃₋₆ cycloalkyl;

X₂", X₃" and X₄" are, independently, selected from the group consisting of CN, NO₂, Cl, F, Br, I, H, R', OR', CF₃, OCF₃ and

OSO₂R', wherein R' is C₁₋₄ alkyl or haloalkyl; provided that when there are multiple halo substitutions in the haloalkyl, halo represents F; or X₁" and X₂" together form an aryl or heteroaryl ring, substituted or unsubstituted; wherein the heteroatom is selected from N, S and O and the substituents are selected from the group consisting of halo, C₁₋₄ alkyl, OCF₃, CF₃, OMe, CN, OSO₂-C₁₋₄ alkyl, OSO₂-C₃₋₆ cycloalkyl and NO₂; or X₃" and X₄" independently represent C(O)R₁;

provided that either X₁" or X₃" represents H; and

R₁" and R₂" are, independently, selected from the group consisting of hydrogen, C₁₋₄ alkyl, C₃₋₆ cycloalkyl, C₂₋₅ alkenyl,

C₂₋₅ alkynyl, heterocycloalkyl and aryl; or R₁" and R₂" together form a 3 to 7 membered optionally substituted heterocyclic ring optionally containing an additional heteroatom selected from O, S and N; wherein the substituents are selected from the group consisting of CN, aryl, CO₂R", CO₂NHR", OH, OR", NH₂, halo, CF₃, OCF₃ and NO₂; wherein R" represents C₁₋₄ alkyl, or C₃₋₆ cycloalkyl;

in sub formula (Id):

D is selected from the group consisting of CN, NO₂, Cl, F, Br,

I, R, OR, SR, CF₃, OCF₃ and OSO₂R, wherein R represents C₁₋₄ alkyl, C₃₋₆ cycloalkyl, or C₁₋₁₀ aryl or heteroaryl wherein the heteroatom is selected from N, S and O and substituents are selected from the group consisting of halo, C₁₋₄ alkyl, OCF₃, CF₃, OMe, CN, OSO₂-C₁₋₄ alkyl, OSO₂-C₃₋₆ cycloalkyl and NO₂;

n is the integer of 1 or 2;

each E is independently C or N, provided that no more than two E moieties are N;

further provided that when n is 2, each E is C;

a and b are optionally present bond;

R₁^{IV} is selected from the group consisting of (CH₂)_nCO₂R', (CH₂)_nCO₂H, (CH₂)_nCONR'₂, (CH₂)_nCH₂OR', OR', SR', CN, NO₂, Cl, F, Br, I, CF₃, OCF₃, OSO₂R', R' and H; wherein R' is C₁₋₄ alkyl, or C₃₋₆ cycloalkyl; or R₁^{IV} is O, forming a ketone such that YR₁^{IV} represents -C=O;

R₂^{IV} is selected from the group consisting of hydrogen, CN, NO₂,

Cl, F, Br, I, H, R", OR", CF₃, OCF₃, and OSO₂R"; wherein R" represents C₁₋₄ alkyl, or C₃₋₆ cycloalkyl.

Y is selected from C, CH, O, N and S; provided that when Y is S, R₁^{IV} is O; further provided that when Y is O, R₁^{IV} is not present;

X' is CH₂, NH, O and S; and

attachment is at the carbon atom marked 3;

in sub formula (Ie) :

X₁" is selected from the group consisting of CN, NO₂, Cl, F, Br, I, H, R', OR', CF₃, OCF₃ and OSO₂R', wherein R' represents


C₁₋₄ alkyl, or C₃₋₆ cycloalkyl;

X₂", X₃" and X₄" are, independently, selected from the group consisting of CN, NO₂, Cl, F, Br, I, H, R", OR", CF₃, OCF₃ and

OSO₂R", wherein R" is C₁₋₄ alkyl or haloalkyl; or X₁" and X₂" together form an aryl or heteroaryl ring, substituted or unsubstituted; wherein the heteroatom is selected from N, S and

O; and any substituents are selected from the group consisting of halo, C₁₋₄ alkyl, OCF₃, CF₃, OMe, CN, OSO₂R' and NO₂; or X₃^{'''} and X₄^{'''} independently represent C(O)R₁;
 provided that when there are multiple halo substitutions in the haloalkyl, halo represents F; also provided that
 either X₁^{'''} or X₃^{'''} is hydrogen;
 and R₉ is O-CH₂-alkyl, O-CH₂-aryl and O-aryl.

[0022] In addition, Japanese Patent Application under PCT laid-open under kohyo No. 2001-501584 (WO97/37967, EP-A-901459, US Patent No. 6022894) also describes a compound of the following formula as a CaSR antagonist.



wherein

R₁ is selected from the group consisting of aryl, longer-length alk, and cycloalk;

R₂ is selected from the group consisting of lower alk, cycloalk, alkoxy, H, OH, =O, C(O)OH, C(O)O-lower alk, C(O)NH-lower alk, C(O)N(lower alk)₂, SH, S-lower alk, NH₂, NH-lower alk, and N(lower alk)₂;

R₃ and R₄ is each independently lower alk or together cyclopropyl;

R₅ is either an optionally substituted naphthyl having 1 - 4 substituents independently selected from the group consisting of methyl, ethyl, isopropyl, methoxy, Cl, F, Br, and lower haloalkoxy, or a substituted phenyl having 1 - 4 substituents with at least one substituent in a meta or para position selected from the group consisting of lower alkyl, methoxy, Cl, F, Br, and lower haloalkoxy, provided that said substituted phenyl may also have 2 or 3 additional substituents;

R₆ if present is either hydrogen, lower alkyl or lower alkenyl, wherein R₆ is not present if R₂ is =O;


Y₁ is either a covalent bond, alkylene or alkenylene;

Y₂ is alkylene;

Y₃ is alkylene; and

Z is selected from the group consisting of a covalent bond, O, S, NH, N-lower alk, alkylene, alkenylene, and alkenylene, provided that if Z is O, S, NH, or N-lower alk, then Y₁ is not a covalent bond; further provided that Y₁ and Z may together form a covalent bond; provided that R₁ is not 6-CN-2-pyridyl; further provided that if R₅ is 3,4-dimethoxy-phenyl, then R₁ is not CH₃(CH₂)₅O-phenyl; 2-cyclopentyl-phenyl; 2-Cl-phenyl; 2-CN-phenyl; 2-(3-furanyl)phenyl; or 4-(1,2-benzisothiazole); further provided that if R₅ is 4-methoxy-phenyl, then R₁ is not 2-cyclopentyl-phenyl; 2-CH₃-phenyl; 2-benzyl-phenyl; 3-CH₃, 4-CH₃SO₂-phenyl; 4-(1,2-benzisothiazole); further provided that if R₅ is 4-Cl-phenyl, then R₁ is not 2-CH₃-phenyl; 5-iso-propyl-phenyl; 2-CH₃-phenyl; 4-CH₃-phenyl; phenyl; 2-Cl-phenyl; 4-Cl-phenyl; 2-methoxy; 4-CH₃CHCH-phenyl; 3,4-CH₃-phenyl; 2,4-CH₃-phenyl; 2,3-CH₃-phenyl; 2-iso-propyl; 5-CH₃-phenyl; pyridyl; 1-imidazole; or 4-(1,2-benzisothiazole); and further provided that if R₅ is 3,5-dimethyl, or 4-methoxy-phenyl, then R₁ is not 4-CH₃, 6-CN-2-pyridyl; or thiophenecarboxamide; and pharmaceutical acceptable salts and complexes thereof; wherein said compound has an IC₅₀ ≤ 10 μM using the Calcium Receptor Inhibitor Assay.

[0023] Maxine Gowen et al. examined the effect of a compound having a CaSR antagonistic action and called NPS-2143



NPS-2143

on osteogenesis by orally administering NPS-2143 to OVX rat and measuring the concentration in blood and bone density, and reported the results [J. Clin. Invest., 105, 1595-1604 (2000)].

[0024] According to this publication, NPS-2143 significantly promotes release of PTH, but has no direct effect on osteoblast and osteoclast in vitro, as a result of which no bone increase or decrease was found. One of the reasons was considered to be the too long half-life of NPS-2143 in blood. When rat PTH (1-34) was administered to OVX rat at a dose of 5 µg/kg, the PTH concentration in blood shows a peak at about 175 pg/ml in 30 min, and restores its original level in 2 hr. When NPS-2143 was administered at a dose of 100 µmol/kg, PTH concentration kept increasing even after the PTH concentration in blood reached about 115 pg/ml in 30 min, and the concentration was about 140 pg/ml even after 4 hr [see J. Clin. Invest., 105, 1595-1604 (2000), p. 1598, Fig. 3].

[0025] At this time, the concentration in blood of NPS-2143 itself remained above 100 ng/ml even at 8 hr after administration and it was only after 24 hr when the concentration became not more than 10 ng/ml and could not be detected.


[0026] The reference of the above-mentioned Maxine Gowen et al. suggests that a calcium receptor antagonist having a too long half-life in blood brings about the same results as in sustained administration of PTH and teaches that an increase in the bone mass cannot be expected.

[0027] The present invention aims at providing a compound having a calcium receptor antagonistic action. The present invention also aims at providing a pharmaceutical composition comprising said compound, which is effective as an agent for the treatment of a disease showing abnormal calcium homeostasis, namely, osteoporosis, hypoparathyreosis, osteosarcoma, periodontal disease, bone fracture, steoarthritis, chronic rheumatoid arthritis, Paget's disease, humoral hypercalcemia, autosomal dominant hypocalcemia and the like, particularly as a therapeutic agent for osteoporosis, which is capable of oral administration and intermittent administration. Moreover, the present invention aims at providing a synthetic intermediate for a compound having a calcium receptor antagonistic action.


Disclosure of the Invention

[0028] The present inventors have conducted intensive studies in an attempt to solve the above-mentioned problems and as a result, found that a compound of the following formula [I] has a superior calcium receptor antagonistic action and can be administered orally and intermittently, which resulted in the completion of the present invention. All the conventionally known calcium receptor antagonists increase PTH concentration in blood in a sustained manner and a sufficient osteogenesis promoting action cannot be expected. In contrast, the compound of the following formula is surprisingly capable of increasing the PTH concentration in blood intermittently in a non-sustained manner, and is expected to be put into practical use as a superior therapeutic agent for osteoporosis.


[0029] The compound represented by the following formula of the present invention is characterized by its structure wherein carbon atom adjacent to oxygen atom has R¹ and R² as substituents, i.e.,



[0030] As is clear from the Experimental Example below, the compound of the present invention having the structure




has not only a superior calcium receptor antagonistic action, but a temporary non-sustained PTH secretion promoting action, as compared to conventional compounds having the structure



Accordingly, it is considered that, by administration of the compound of the present invention, an effect similar to that obtained by intermittent administration of PTH is obtained, which is extremely effective for the treatment of osteoporosis.


[0031] Accordingly, the present invention relates to a compound represented by the following formula, a calcium receptor antagonist and a therapeutic agent for osteoporosis, which contains the compound as an active ingredient, and an intermediate compound useful for the synthesis of the compound. More particularly, the present invention provides the following (1) to (21).

(1) A compound of the formula [I]



wherein

R¹ is aryl group or heteroaryl group wherein said aryl group and heteroaryl group are optionally substituted by 1 to 3 substituents selected from halogen atom, C₁₋₆ alkyl group, halo(C₁₋₆)alkyl group, hydroxy(C₁₋₆)alkyl group, C₁₋₆ alkoxy(C₁₋₆)alkyl group, hydroxyl group, C₁₋₆ alkoxy group, halo(C₁₋₆)alkoxy group, mercapto group, C₁₋₆ alkylthio group, C₁₋₆ alkylsulfanyl group, C₁₋₆ alkylsulfonyl group, aminosulfonyl group, C₁₋₆ alkyl-sulfamoyl group, di(C₁₋₆)alkylsulfamoyl group, carboxy group, (C₁₋₆ alkoxy)carbonyl group, C₁₋₇ acyl group, carbamoyl group, (C₁₋₆ alkyl)carbamoyl group, di(C₁₋₆ alkyl)carbamoyl group, cyano group, nitro group, amino group, C₁₋₆ alkylamino group, di(C₁₋₆)alkylamino group, C₁₋₇ acylamino group, C₁₋₃ alkylenedioxy group,



wherein R^A is (C₁₋₆ alkoxy)carbonyl group or carboxy group, and R^B is hydrogen atom or C₁₋₆ alkyl group;
R² is C₁₋₆ alkyl group

wherein said C₁₋₆ alkyl group is optionally substituted by 1 to 3 substituents selected from halogen atom, hydroxyl group, C₁₋₆ alkoxy group, carboxy group, amino group, C₁₋₆ alkylamino group, di(C₁₋₆)alkylamino group and oxo group, C₃₋₇ cycloalkyl group, C₂₋₆ alkenyl group, C₂₋₆ alkynyl group, aralkyl group, carboxy group, (C₁₋₆ alkoxy)carbonyl group or cyano group;

R³ is hydrogen atom, C₁₋₆ alkyl group, hydroxyl group, C₁₋₆ alkoxy group, mercapto group, C₁₋₆ alkylthio group, carboxy group, (C₁₋₆ alkoxy)carbonyl group, (C₁₋₆ alkyl)carbamoyl group, di(C₁₋₆ alkyl)carbamoyl group, amino group, C₁₋₆ alkylamino group or di(C₁₋₆)alkylamino group;

R⁴ is hydrogen atom, C₁₋₆ alkyl group or C₂₋₆ alkenyl group, or R³ and R⁴ in combination show oxo group;

R⁵ and R⁶ are the same or different and each is C₁₋₆ alkyl group, or R⁵ and R⁶ in combination show cyclopropyl group together with the carbon atom they bind to;

R⁷ is aryl group or heteroaryl group wherein said aryl group and heteroaryl group are optionally substituted by 1 to 3 substituents selected from halogen atom, C₁₋₆ alkyl group, halo(C₁₋₆)alkyl group, hydroxy(C₁₋₆)alkyl group, C₃₋₇ cycloalkyl group, hydroxyl group, C₁₋₆ alkoxy group, halo(C₁₋₆)alkoxy group, carboxy group, (C₁₋₆ alkoxy)carbonyl group, nitro group, cyano group, C₁₋₆ alkylsulfonyloxy group, carbamoyl group and C₁₋₃ alkylenedioxy group;

X¹ is a single bond, C₁₋₆ alkylene group or C₂₋₆ alkynylene group wherein said C₁₋₆ alkylene group and C₂₋₆ alkynylene group are optionally substituted by C₁₋₆ alkyl group or oxo group;


X² is C₁₋₆ alkylene group wherein said C₁₋₆ alkylene group is optionally substituted by C₁₋₆ alkyl group or halo(C₁₋₆)alkyl group;

X³ is a single bond or C₁₋₆ alkylene group wherein said C₁₋₆ alkylene group is optionally substituted by hydroxyl group or oxo group;

X⁴ and X⁵ in combination show a single bond, methylene group, -NH-, oxygen atom, sulfur atom, -C(=O)-, -CH₂NH-, -CH₂O-, -CH₂S-, -CH₂CO-, -NHCH₂-, -OCH₂-, -SCH₂-, -COCH₂-, -CH=CH- or -C≡C- (hereinafter sometimes to be referred to as compound [I]),

a salt thereof, a solvate thereof or a prodrug thereof.

(2) A compound of the formula [I']



wherein


R¹' is aryl group or heteroaryl group wherein said aryl group and heteroaryl group are optionally substituted by 1 to 3 substituents selected from halogen atom, C₁₋₆ alkyl group, halo(C₁₋₆)alkyl group, hydroxy (C₁₋₆)alkyl group, C₁₋₆ alkoxy(C₁₋₆)alkyl group, hydroxyl group, C₁₋₆ alkoxy group, halo(C₁₋₆)alkoxy group, mercapto group, C₁₋₆ alkylthio group, C₁₋₆ alkylsulfanyl group, C₁₋₆ alkylsulfonyl group, aminosulfonyl group, C₁₋₆ alkylsulfamoyl group, di(C₁₋₆)alkylsulfamoyl group, carboxy group, (C₁₋₆ alkoxy)carbonyl group, C₁₋₇ acyl group, carbamoyl group, (C₁₋₆ alkyl)carbamoyl group, di(C₁₋₆ alkyl)carbamoyl group, cyano group, nitro group, amino group, C₁₋₆ alkylamino group, di(C₁₋₆)alkylamino group and C₁₋₇ acylamino group;

R⁷' is aryl group or heteroaryl group wherein said aryl group and heteroaryl group are optionally substituted by 1 to 3 substituents selected from halogen atom, C₁₋₆ alkyl group, halo(C₁₋₆)alkyl group, C₃₋₆ cycloalkyl group, hydroxyl group, C₁₋₆ alkoxy group, halo(C₁₋₆)alkoxy group, carboxy group, (C₁₋₆ alkoxy)carbonyl group, nitro group, cyano group and C₁₋₆ alkylsulfonyloxy group; and

R², R³, R⁴, R⁵, R⁶, X¹, X², X³, X⁴ and X⁵ are same as defined above in (1) (hereinafter sometimes to be referred to as compound [I']) respectively,


a salt thereof, a solvate thereof or a prodrug thereof.

(3) A compound of the formula [I'']



wherein

R¹'' is aryl group or heteroaryl group wherein said aryl group and heteroaryl group are optionally substituted by 1 to 3 substituents selected from C₁₋₃ alkylendioxy group,



wherein R^A is (C₁₋₆ alkoxy)carbonyl group or carboxy group and


R^B is hydrogen atom or C₁₋₆ alkyl group;

$R^{7''}$ is aryl group or heteroaryl group wherein said aryl group and heteroaryl group are optionally substituted by 1 to 3 substituents selected from hydroxy(C₁₋₆)alkyl group, carbamoyl group and C₁₋₃ alkylenedioxy group; and

R^2 , R^3 , R^4 , R^5 , R^6 , X^1 , X^2 , X^3 , X^4 and X^5 are same as defined above (1) (hereinafter sometimes to be referred to as compound [I'']) respectively,

a salt thereof, a solvate thereof or a prodrug thereof.

(4) A compound of the formula [I-2]



wherein R^{11} and R^{12} are the same or different and each is hydrogen atom, halogen atom, C₁₋₄ alkyl group, hydroxy (C₁₋₆)alkyl group, C₁₋₆ alkoxy(C₁₋₆)alkyl group, hydroxyl group, C₁₋₄ alkoxy group, cyano group or nitro group, or R^{11} and R^{12} in combination show C₁₋₃ alkylenedioxy group,

R^{21} is C₁₋₄ alkyl group


wherein said alkyl group is optionally substituted by

C₁₋₄ alkoxy group,

C₃₋₅ cycloalkyl group, C₂₋₄ alkenyl group, or aralkyl group, and R^3 , R^4 , R^5 , R^6 , R^7 , X^1 , X^2 , X^3 , X^4 and X^5 are same as defined above (1) (hereinafter sometimes to be referred to as compound [I-2]) respectively,

a salt thereof, a solvate thereof or a prodrug thereof.

(5) The compound of the above-mentioned (4), which is represented by the formula [I-3]



wherein R^{71} and R^{72} are the same or different and each is hydrogen atom, C₁₋₄ alkyl group or C₁₋₄ alkoxy group, or in combination show -CH=CH-CH=CH- or C₁₋₃ alkylenedioxy group, and

R^{11} , R^{12} and R^{21} are same as defined in the above (4) (hereinafter sometimes to be referred to as compound [I-3]) respectively, a salt thereof, a solvate thereof or a prodrug thereof.

(6) The compound of the above-mentioned (5), wherein R^{11} and R^{12} are the same or different and each is hydrogen atom, chlorine atom, methyl group, hydroxymethyl group, hydroxyl group, methoxy group, cyano group or nitro group, or R^{11} and R^{12} in combination show methylenedioxy group, and R^{21} is optionally branched C₁₋₄ alkyl group or C₃₋₅ cycloalkyl group, a salt thereof, a solvate thereof or a prodrug thereof.

(7) The compound of the above-mentioned (6), wherein R^{21} is methyl group, ethyl group, cyclopropyl group or cyclobutyl group,

a salt thereof, a solvate thereof or a prodrug thereof.

(8) The compound of the above-mentioned (7), wherein R^{11} is hydrogen atom, R^{12} is methyl group, methoxy group or hydroxymethyl group, or R^{11} and R^{12} in combination show methylenedioxy group, and R^{21} is methyl group or

cyclopropyl group,

a salt thereof, a solvate thereof or a prodrug thereof.

(9) The compound of the above-mentioned (8), wherein R^{21} is cyclopropyl group,

a salt thereof, a solvate thereof or a prodrug thereof.

(10) The compound of the above-mentioned (9), wherein R^{71} and R^{72} in combination show $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$,
a salt thereof, a solvate thereof or a prodrug thereof.

(11) The compound of the above-mentioned (9), wherein R^{71} and R^{72} are groups selected from C_{1-4} alkyl group
and C_{1-4} alkoxy group,

a salt thereof, a solvate thereof or a prodrug thereof.

(12) The compound of the above-mentioned (1), which is selected from

(2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(2-methoxyphenyl)methoxy]propan-
2-ol,

(2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(2-methylphenyl)methoxy]propan-2-ol,

(2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(phenyl)methoxy]propan-2-ol,

(2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methoxyphenyl)ethoxy]propan-2-ol,

(2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methylphenyl)ethoxy]propan-2-ol,

(2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methoxyphenyl)propoxy]propan-2-ol,

(2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-cyanophenyl)ethoxy]propan-2-ol,

(2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(3-methoxyphenyl)ethoxy]propan-2-ol and

(2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(3-methylphenyl)ethoxy]propan-2-ol,

a salt thereof, a solvate thereof or a prodrug thereof.


(13) (2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(2-hydroxymethylphenyl)methoxy]pro-
pan-2-ol, a salt thereof, a solvate thereof or a prodrug thereof.

(14) A pharmaceutical composition comprising the compound of any of the above-mentioned (1) to (13), a salt
thereof, a solvate thereof or a prodrug thereof as an active ingredient.

(15) A calcium receptor antagonist comprising the compound of any of the above-mentioned (1) to (13), a salt
thereof, a solvate thereof or a prodrug thereof as an active ingredient.

(16) A therapeutic agent for osteoporosis comprising the compound of any of the above-mentioned (1) to (13), a
salt thereof, a solvate thereof or a prodrug thereof as an active ingredient.

(17) A compound of the formula [II]




wherein R^{11} is halogen atom, C_{1-4} alkyl group, hydroxy(C_{1-6})alkyl group, C_{1-6} alkoxy(C_{1-6})alkyl group, hydroxyl
group, C_{1-4} alkoxy group, tert-butyldimethylsilyloxymethyl group, cyano group or nitro group (hereinafter some-
times to be referred to as compound [II]),

a salt thereof or a solvate thereof.

(18) The compound of the above-mentioned (17), wherein, in the formula [II], R^{11} is C_{1-4} alkyl group, hydroxy(C_{1-6})
alkyl group or C_{1-4} alkoxy group,

a salt thereof or a solvate thereof.


(19) A compound of the formula [III]



wherein R¹¹ is halogen atom, hydroxy(C₁₋₆)alkyl group, C₁₋₆ alkoxy(C₁₋₆)alkyl group, hydroxyl group, C₁₋₄ alkoxy group, tert-butyldimethylsilyloxymethyl group, cyano group or nitro group (hereinafter sometimes to be referred to as compound [III]), a salt thereof or a solvate thereof.

(20) The compound of the above-mentioned (19), wherein, in the formula [III], R¹¹ is C₁₋₄ alkoxy group, a salt thereof or a solvate thereof.

(21) A compound of the formula [IV]



wherein R⁸ is carboxy group, nitro group, tert-butoxycarbonylamino group or benzyloxycarbonylamino group (hereinafter sometimes to be referred to as compound [IV]), a salt thereof or a solvate thereof.

Brief Description of the Drawings

[0032]

Fig. 1 shows the time-course changes of serum PTH concentration when 30 mg/kg of a compound of Example 22 was administered to rat.

Fig. 2 shows the time-course changes of serum PTH concentration when 30 mg/kg of a compound of Example 23 was administered to rat.

Fig. 3 shows the time-course changes of serum PTH concentration when 30 mg/kg of a compound of Example 24 was administered to rat.

Fig. 4 shows the time-course changes of serum PTH concentration when 100 mg/kg of a compound of Comparative Example 1 was administered to rat.

Fig. 5 shows the time-course changes of serum PTH concentration when 30 mg/kg of NPS-2143 was administered to rat.

Embodiment of the Invention

[0033] The terms used in the present specification are defined in the following.

[0034] The "aryl group" is aromatic hydrocarbon group having 6 to 12 carbon atoms, which may be partly saturated. Examples thereof include phenyl group, biphenyl group, indenyl group, naphthyl and the like. Preferred are phenyl group and naphthyl group, and particularly preferred is phenyl group. These aryl groups are optionally substituted by the substituents to be mentioned later. The position of the bond of these aryl groups and the position of the substituent when substituted are not particularly limited as long as they are chemically acceptable.

[0035] The "heteroaryl group" shows a 5-membered to 6-membered unsaturated ring containing 1 to 3 heteroatoms in the ring, which is selected from nitrogen atom, oxygen atom and sulfur atom, including a fused ring with a benzene ring or other hetero ring. Examples of these heteroaryl groups include pyrrolyl group, furyl group, thienyl group, imidazolyl group, oxazolyl group, thiazolyl group, pyrazolyl group, isoxazolyl group, isothiazolyl group, oxadiazolyl group, triazolyl group, indolyl group, benzofuryl group, benzothienyl group, benzimidazolyl group, benzoxazolyl group, benzothiazolyl group, pyridyl group, pyrimidinyl group, quinolyl group, isoquinolyl group and the like. Preferred are benzofuryl group, benzothienyl group, benzimidazolyl group, benzoxazolyl group, benzothiazolyl group, pyridyl group and quinolyl group. These heteroaryl groups may be substituted by the substituents to be mentioned later. The position of

the bond of these heteroaryl groups and the position of the substituent when substituted are not particularly limited as long as they are chemically acceptable.

[0036] The "halogen atom" is fluorine atom, chlorine atom, bromine atom or iodine atom, preferably fluorine atom and chlorine atom, and particularly preferably chlorine atom.

[0037] The " C_{1-6} alkyl group" is linear or branched chain alkyl group having 1 to 6, preferably 1 to 4, carbon atoms, and is exemplified by methyl group, ethyl group, propyl group, isopropyl group, butyl group, isobutyl group, tert-butyl group, pentyl group, isopentyl group, tert-pentyl group, hexyl group and the like, with preference given to C_{1-4} alkyl group selected from methyl group, ethyl group, propyl group, isopropyl group, butyl group, isobutyl group and tert-butyl group.

[0038] The "halo(C_{1-6})alkyl group" is haloalkyl group wherein the aforementioned " C_{1-6} alkyl group" is substituted by one or more halogen atoms. The position of the substituent is not particularly limited as long as it is chemically acceptable. Examples of "halo(C_{1-6})alkyl group" include fluoromethyl group, difluoromethyl group, trifluoromethyl group, chloromethyl group, dichloromethyl group, trichloromethyl group, bromomethyl group, dibromomethyl group, tribromomethyl group, iodomethyl group, diiodomethyl group, triiodomethyl group, 2-fluoroethyl group, 2,2-difluoroethyl group, 2,2,2-trifluoroethyl group, 2-chloroethyl group, 2,2-dichloroethyl group, 2,2,2-trichloroethyl group, 2-bromoethyl group, 2,2-dibromoethyl group, 2,2,2-tribromoethyl group, 3-chloropropyl group, 4-chlorobutyl group and the like, with preference given to halo(C_{1-2})alkyl group such as trifluoromethyl group and 2,2,2-trichloroethyl group.

[0039] The "hydroxy(C_{1-6})alkyl group" is hydroxyalkyl group wherein the aforementioned " C_{1-6} alkyl group" is substituted by hydroxyl group. The position of the substituent is not particularly limited as long as it is chemically acceptable. Examples of "hydroxy(C_{1-6})alkyl group" include hydroxymethyl group, 1-hydroxyethyl group, 2-hydroxyethyl group, 1-hydroxypropyl group, 2-hydroxypropyl group, 3-hydroxypropyl group, 2-hydroxy-1-methylethyl group, 1-hydroxybutyl group, 2-hydroxybutyl group, 3-hydroxybutyl group, 4-hydroxybutyl group, 3-hydroxy-2-methylpropyl group, 2-hydroxy-1,1-dimethylethyl group, 5-hydroxypentyl group, 6-hydroxyhexyl group and the like, with preference given to hydroxy C_{1-4} alkyl group selected from hydroxymethyl group, 2-hydroxyethyl group, 3-hydroxypropyl group and 4-hydroxybutyl group.

[0040] The " C_{1-6} alkoxy group" is linear or branched chain alkoxy group having 1 to 6, preferably 1 to 4, carbon atoms. Examples thereof include methoxy group, ethoxy group, propoxy group, isopropoxy group, butoxy group, tert-butoxy group, pentyloxy group, tert-pentyloxy group, hexyloxy group and the like, with preference given to C_{1-4} alkoxy group selected from methoxy group, ethoxy group, propoxy group, isopropoxy group, butoxy group and tert-butoxy group.

[0041] The " C_{1-6} alkoxy(C_{1-6})alkyl group" is alkoxyalkyl group wherein the aforementioned " C_{1-6} alkyl group" is substituted by the aforementioned " C_{1-6} alkoxy group". The position of the substituent is not particularly limited as long as it is chemically acceptable. Examples of " C_{1-6} alkoxy(C_{1-6})alkyl group" include methoxymethyl group, ethoxymethyl group, propoxymethyl group, butoxymethyl group, pentyloxymethyl group, hexyloxymethyl group, 1-methoxyethyl group, 1-ethoxyethyl group, 2-methoxyethyl group, 2-ethoxyethyl group, 1-methoxypropyl group, 1-ethoxypropyl group, 2-methoxypropyl group, 2-ethoxypropyl group, 3-methoxypropyl group, 3-ethoxypropyl group, 2-methoxy-1-methylethyl group, 1-methoxybutyl group, 1-ethoxybutyl group, 2-methoxybutyl group, 2-ethoxybutyl group, 3-methoxybutyl group, 3-ethoxybutyl group, 4-methoxybutyl group, 4-ethoxybutyl group, 3-methoxy-2-methylpropyl group, 2-methoxy-1,1-dimethylethyl group, 2-ethoxy-1,1-dimethylethyl group, 5-methoxypentyl group, 6-methoxyhexyl group and the like, with preference given to C_{1-4} alkoxy(C_{1-4})alkyl group selected from methoxymethyl group, ethoxymethyl group, propoxymethyl group, butoxymethyl group, 2-methoxyethyl group, 3-methoxypropyl group and 4-methoxybutyl group.

[0042] The "halo(C_{1-6})alkoxy group" is haloalkoxy group wherein the aforementioned " C_{1-6} alkoxy group" is substituted by one or more halogen atoms. The position of the substituent is not particularly limited as long as it is chemically acceptable. Examples of "halo(C_{1-6})alkoxy group" include fluoromethoxy group, difluoromethoxy group, trifluoromethoxy group, chloromethoxy group, dichloromethoxy group, trichloromethoxy group, bromomethoxy group, dibromomethoxy group, tribromomethoxy group, iodomethoxy group, diiodomethoxy group, triiodomethoxy group, 2-fluoroethoxy group, 2,2-difluoroethoxy group, 2,2,2-trifluoroethoxy group, 2-chloroethoxy group, 2,2-dichloroethoxy group, 2,2,2-trichloroethoxy group, 2-bromoethoxy group, 2,2-dibromoethoxy group, 2,2,2-tribromoethoxy group, 3-chloropropoxy group, 4-chlorobutoxy group and the like, with preference given to halo(C_{1-2})alkoxy group such as trifluoromethoxy group and 2,2,2-trichloroethoxy group.

[0043] The " C_{1-6} alkylthio group" is linear or branched chain alkylthio group having 1 to 6, preferably 1 to 4, carbon atoms. Examples thereof include methylthio group, ethylthio group, propylthio group, isopropylthio group, butylthio group, tert-butylthio group, pentylthio group, tert-pentylthio group, hexylthio group and the like, with preference given to C_{1-4} alkylthio group selected from methylthio group, ethylthio group, propylthio group, isopropylthio group, butylthio group and tert-butylthio group.

[0044] The " C_{1-6} alkylsulfanyl group" is linear or branched chain alkylsulfanyl group having 1 to 6 carbon atoms. Examples thereof include methylsulfanyl group, ethylsulfanyl group, propylsulfanyl group, isopropylsulfanyl group, butylsulfanyl group, tert-butylsulfanyl group, pentylsulfanyl group, tert-pentylsulfanyl group, hexylsulfanyl group and the like, with preference given to C_{1-4} alkylsulfanyl group selected from methylsulfanyl group, ethylsulfanyl group, propylsulfanyl group, isopropylsulfanyl group, butylsulfanyl group and tert-butylsulfanyl group.

tylsulfanyl group, tent-butylsulfanyl group, pentylsulfanyl group, tert-pentylsulfanyl group, hexylsulfanyl group and the like, with preference given to methylsulfanyl group, ethylsulfanyl group, propylsulfanyl group, isopropylsulfanyl group, butylsulfanyl group and tert-butylsulfanyl group, all of which having 1 to 4 carbon atoms.

[0045] The " C_{1-6} alkylsulfonyl group" is linear or branched chain alkylsulfonyl group having 1 to 6 carbon atoms. Examples thereof include methylsulfonyl group, ethylsulfonyl group, propylsulfonyl group, isopropylsulfonyl group, butylsulfonyl group, tert-butylsulfonyl group, pentylsulfonyl group, tert-pentylsulfonyl group, hexylsulfonyl group and the like, with preference given to methylsulfonyl group, ethylsulfonyl group, propylsulfonyl group, isopropylsulfonyl group, butylsulfonyl group and tert-butylsulfonyl group, all of which having 1 to 4 carbon atoms.

[0046] The " C_{1-6} alkylsulfamoyl group" is sulfamoyl group monosubstituted by the aforementioned " C_{1-6} alkyl group". Examples thereof include methylsulfamoyl group, ethylsulfamoyl group, propylsulfamoyl group, isopropylsulfamoyl group, butylsulfamoyl group, tert-butylsulfamoyl group, pentylsulfamoyl group, tert-pentylsulfamoyl group; hexylsulfamoyl group and the like, with preference given to methylsulfamoyl group, ethylsulfamoyl group, propylsulfamoyl group, isopropylsulfamoyl group, butylsulfamoyl group and tert-butylsulfamoyl group, all of which having 1 to 4 carbon atoms.

[0047] The " $di(C_{1-6})$ alkylsulfamoyl group" is sulfamoyl group disubstituted by the aforementioned " C_{1-6} alkyl group". Examples thereof include dimethylsulfamoyl group, diethylsulfamoyl group, dipropylsulfamoyl group, diisopropylsulfamoyl group, dibutylsulfamoyl group, diisobutylsulfamoyl group, di-tert-butylsulfamoyl group, dipentylsulfamoyl group, ethylmethylsulfamoyl group, methylpropylsulfamoyl group, butylmethylsulfamoyl group, ethylpropylsulfamoyl group, ethylbutylsulfamoyl group and the like, with preference given to dimethylsulfamoyl group, diethylsulfamoyl group and dipropylsulfamoyl group.

[0048] The " $(C_{1-6}$ alkoxy)carbonyl group" is alkoxycarbonyl group wherein the C_{1-6} alkoxy moiety is exemplified by the aforementioned " C_{1-6} alkoxy group". Examples thereof include methoxycarbonyl group, ethoxycarbonyl group, propoxycarbonyl group, isopropoxycarbonyl group, butoxycarbonyl group, isobutoxycarbonyl group, tert-butoxycarbonyl group, pentyloxycarbonyl group, hexyloxycarbonyl group and the like, with preference given to (C_{1-4}) alkoxy)carbonyl group selected from methoxycarbonyl group, ethoxycarbonyl group, propoxycarbonyl group, isopropoxycarbonyl group, butoxycarbonyl group and tert-butoxycarbonyl group.

[0049] The " C_{1-7} acyl group" is alkanoyl group or aroyl group having 1 to 7 carbon atoms. Examples thereof include formyl group, acetyl group, propionyl group, butyryl group, pivaloyl group, benzoyl group and the like, with preference given to formyl group, acetyl group, pivaloyl group and benzoyl group.

[0050] The " $(C_{1-6}$ alkyl)carbamoyl group" is alkylcarbamoyl group wherein the carbamoyl group is substituted by C_{1-6} alkyl group exemplified for the aforementioned " C_{1-6} alkyl group". Examples thereof include methylcarbamoyl group, ethylcarbamoyl group, propylcarbamoyl group, isopropylcarbamoyl group, butylcarbamoyl group, tert-butylcarbamoyl group, pentylcarbamoyl group, tert-pentylcarbamoyl group, hexylcarbamoyl group and the like, with preference given to (C_{1-4}) alkyl)carbamoyl group selected from methylcarbamoyl group, ethylcarbamoyl group, propylcarbamoyl group, isopropylcarbamoyl group, butylcarbamoyl group and tert-butylcarbamoyl group.

[0051] The " $di(C_{1-6})$ alkyl)carbamoyl group" is dialkylcarbamoyl group wherein the carbamoyl group is disubstituted by the aforementioned " C_{1-6} alkyl group", and the kind of the alkyl groups may be different. Examples thereof include dimethylcarbamoyl group, diethylcarbamoyl group, dipropylcarbamoyl group, diisopropylcarbamoyl group, dibutylcarbamoyl group, di-tert-butylcarbamoyl group, dipentylcarbamoyl group, di-tert-pentylcarbamoyl group, dihexylcarbamoyl group, methylethylcarbamoyl group, methylpropylcarbamoyl group, methylbutylcarbamoyl group, ethylpropylcarbamoyl group, ethylbutylcarbamoyl group and the like, with preference given to $di(C_{1-4})$ alkyl)carbamoyl group selected from dimethylcarbamoyl group, diethylcarbamoyl group, dipropylcarbamoyl group, dibutylcarbamoyl group and di-tert-butylcarbamoyl group.

[0052] The " C_{1-6} alkylamino group" is alkylamino group wherein the amino group is substituted by the aforementioned " C_{1-6} alkyl group". Examples thereof include methylamino group, ethylamino group, propylamino group, isopropylamino group, butylamino group, isobutylamino group, tert-butylamino group, pentylamino group, isopentylamino group, tert-pentylamino group, hexylamino group and the like, with preference given to C_{1-4} alkylamino group selected from methylamino group, ethylamino group, propylamino group, isopropylamino group, butylamino group, isobutylamino group and tert-butylamino group.

[0053] The " $di C_{1-6}$ alkylamino group" is dialkylamino group wherein the amino group is disubstituted by the aforementioned " C_{1-6} alkyl group", and the kind of the alkyl groups may be different. Examples thereof include dimethylamino group, ethylmethylamino group, diethylamino group, methylpropylamino group, ethylpropylamino group, dipropylamino group, diisopropylamino group, dibutylamino group, diisobutylamino group, di-tert-butylamino group, dipentylamino group, diisopentylamino group, di-tert-pentylamino group, dihexylamino group and the like, with preference given to $di(C_{1-4})$ alkylamino group selected from dimethylamino group, diethylamino group, dipropylamino group, diisopropylamino group, dibutylamino group, diisobutylamino group and di-tert-butylamino group.

[0054] The " C_{1-7} acylamino group" is amino group substituted by the aforementioned " C_{1-7} acyl group". Examples thereof include alkanoylamino group such as formylamino group, acetylamino group, propionylamino group, butyrylamino group, pivaloylamino group and the like; and aroylamino group wherein aryl group optionally has 1 to 3 sub-

stituents, such as benzoylamino group and the like, with preference given to formylamino group, acetylamino group, pivaloylamino group and benzoylamino group.

[0055] The " C_{3-7} cycloalkyl group" is cyclic alkyl group having 3 to 7, preferably 3 to 6, carbon atoms. Examples thereof include cyclopropyl group, cyclobutyl group, cyclopentyl group, cyclohexyl group, cycloheptyl group and the like. Preferred are C_{3-5} cycloalkyl group such as cyclopropyl group, cyclobutyl group, cyclopentyl group and the like, more preferred are cyclopropyl group and cyclobutyl group, and particularly preferred is cyclopropyl group.

[0056] The " C_{2-6} alkenyl group" is alkenyl group having 2 to 6 carbon atoms. Examples thereof include vinyl group, 1-propenyl group, allyl group, 1-butenyl group, 2-butenyl group, 3-butenyl group, 1-pentenyl group, 2-pentenyl group, 3-pentenyl group, 4-pentenyl group, 5-hexenyl group and the like, with preference given to C_{2-4} alkenyl group such as vinyl group, allyl group and the like.

[0057] The " C_{2-6} alkynyl group" is alkynyl group having 2 to 6, preferably 2 to 4, carbon atoms. Examples thereof include ethynyl group, 1-propynyl group, 2-propynyl group, 1-butylnyl group, 2-butylnyl group, 3-butylnyl group, 1-pentynyl group, 2-pentynyl group, 3-pentynyl group, 4-pentynyl group, 5-hexynyl group and the like, with preference given to C_{2-4} alkynyl group selected from ethynyl group, 1-propynyl group and 2-propynyl group.

[0058] The "aralkyl group" is arylalkyl group wherein the aforementioned " C_{1-6} alkyl group" is substituted by the aforementioned "aryl group". The position of the substituent is not particularly limited as long as it is chemically acceptable. Examples of "aralkyl group" include benzyl group, phenethyl group, 3-phenylpropyl group, 4-phenylbutyl group, 5-phenylpentyl group, 6-phenylhexyl group, 1-naphthylmethyl group, 2-naphthylmethyl group and the like, with preference given to phenyl(C_{1-2})alkyl group wherein the "aryl" moiety is phenyl and the "alkyl" moiety is methyl or ethyl, such as benzyl group and phenethyl group.


[0059] The " C_{1-6} alkylsulfonyloxy group" is alkylsulfonyloxy group wherein the C_{1-6} alkyl moiety is the aforementioned " C_{1-6} alkyl group". Examples thereof include methylsulfonyloxy group, ethylsulfonyloxy group, propylsulfonyloxy group, isopropylsulfonyloxy group, butylsulfonyloxy group, isobutylsulfonyloxy group, tert-butylsulfonyloxy group, pentylsulfonyloxy group, isopentylsulfonyloxy group, tert-pentylsulfonyloxy group and hexylsulfonyloxy group, with preference given to C_{1-4} alkylsulfonyloxy group selected from methylsulfonyloxy group, ethylsulfonyloxy group, propylsulfonyloxy group, isopropylsulfonyloxy group, butylsulfonyloxy group, isobutylsulfonyloxy group and tert-butylsulfonyloxy group.

[0060] The " C_{1-3} alkylenedioxy group" is methylenedioxy group, ethylenedioxy group or propylenedioxy group, preferably methylenedioxy group or ethylenedioxy group, and particularly preferably methylenedioxy group.

[0061] The " C_{1-6} alkylene group" is alkylene group having 1 to 6 carbon atoms. Examples thereof include methylene group, ethylene group, propylene group, butylene group, pentylene group, hexylene group and the like, with preference given to C_{1-4} alkylene group selected from methylene group, ethylene group and propylene group.

[0062] The " C_{2-6} alkynylene group" is alkynylene group having 2 to 6, preferably 2 or 3, carbon atoms. Examples thereof include vinylene group, 1-propenylenylene group, 2-propenylenylene group, 1-butenylene group, 2-butenylene group, 3-butenylene group, 1-pentenylene group, 2-pentenylene group, 3-pentenylene group, 4-pentenylene group, 1-hexenylenylene group, 2-hexenylenylene group, 3-hexenylenylene group, 4-hexenylenylene group, 5-hexenylenylene group and the like, with preference given to C_{2-3} alkynylene group selected from vinylene group, 1-propenylenylene group and 2-propenylenylene group.

[0063] The aryl group and heteroaryl group of the present invention are optionally substituted by 1 to 3 substituents selected from halogen atom, C_{1-6} alkyl group, halo(C_{1-6})alkyl group, hydroxy(C_{1-6})alkyl group, C_{1-6} alkoxy(C_{1-6})alkyl group, hydroxyl group, C_{1-6} alkoxy group, halo(C_{1-6})alkoxy group, mercapto group, C_{1-6} alkylthio group, C_{1-6} alkylsulfanyl group, C_{1-6} alkylsulfonyl group, aminosulfonyl group, C_{1-6} alkylsulfamoyl group, di(C_{1-6})alkylsulfamoyl group, carboxy group, (C_{1-6} alkoxy)carbonyl group, C_{1-7} acyl group, carbamoyl group, (C_{1-6} alkyl)carbamoyl group, di(C_{1-6} alkyl)carbamoyl group, cyano group, nitro group, amino group, C_{1-6} alkylamino group, di(C_{1-6})alkylamino group, C_{1-7} acylamino group, C_{1-3} alkylenedioxy group,



wherein R^A is (C_{1-6} alkoxy)carbonyl group or carboxy group, and R^B is hydrogen atom or C_{1-6} alkyl group.

[0064] The "salt" of the compound of the present invention is exemplified by, but not limited to, inorganic acid addition salts such as hydrochloride, hydrobromide, sulfate, phosphate, nitrate and the like; organic acid addition salts such as acetate, propionate, succinate, glycolate, lactate, malate, oxalate, tartrate, citrate, maleate, fumarate, methanesulfonate, benzenesulfonate, p-toluenesulfonate, ascorbate and the like; amino acid addition salts such as aspartate, glutamate and the like. Preferable salts are hydrochloride and fumarate, particularly preferably fumarate.

[0065] The present invention encompasses solvates, wherein a "solvate" of the compound means that the compound of the present invention is bonded to a solvent molecule such as water, alcohol and the like via a comparatively weak bond due to van der Waals force, electrostatic interaction, hydrogen bond, charge transfer bond, coordinate bond and the like in a solid such as crystal, amorphous and the like or a solution. In some cases, a solvent may be taken in a solid as exemplified by hydrate, alcoholate and the like. Preferable solvate is hydrate.

[0066] A "prodrug" of the compound means a derivative of the compound of the present invention, which has a chemically or metabolically decomposable group, and shows pharmaceutical activity upon hydrolysis, solvolysis or decomposition under physiological conditions.

[0067] The compounds of the formula [I], [I-2] and [I-3] of the present invention may be present as various isomers such as optical isomer, stereoisomer, geometric isomer, tautomer and the like. The present invention encompasses all these isomers and mixtures thereof.

[0068] In the compound of the formula [I] of the present invention, R¹ is preferably aryl group, more preferably phenyl group or naphthyl group and particularly preferably phenyl group. When R¹ is aryl group, the substituent is preferably halogen atom, C₁₋₄ alkyl group, hydroxy(C₁₋₆)alkyl group, C₁₋₆ alkoxy(C₁₋₆)alkyl group, hydroxyl group, C₁₋₄ alkoxy group, cyano group, nitro group or C₁₋₃ alkylenedioxy group, more preferably chlorine atom, methyl group, hydroxymethyl group, methoxy group, cyano group, nitro group, hydroxyl group or methylenedioxy group, particularly preferably methyl group, hydroxymethyl group, methoxy group or methylenedioxy group.

[0069] When R¹ is heteroaryl group, R¹ is preferably selected from thienyl group, furyl group, pyridyl group and thiazolyl group, and the substituent when R¹ is heteroaryl group is preferably C₁₋₆ alkyl group or halogen atom, particularly preferably methyl group or bromine atom.


[0070] R² is preferably optionally substituted and optionally branched C₁₋₄ alkyl group, C₃₋₅ cycloalkyl group, C₂₋₄ alkenyl group or aralkyl group, and the substituent of C₁₋₄ alkyl group is preferably C₁₋₄ alkoxy group. R² is more preferably methyl group, ethyl group, cyclopropyl group or cyclobutyl group, most preferably methyl group or cyclopropyl group, and particularly preferably cyclopropyl group.

[0071] R³ is preferably hydroxyl group, and R⁴ is preferably hydrogen atom. R⁵ and R⁶ are each preferably methyl group. R⁷ is preferably aryl group, more preferably phenyl group or naphthyl group. The substituent of R⁷ is preferably C₁₋₆ alkyl group, C₁₋₆ alkoxy group, hydroxyl group or C₁₋₃ alkylenedioxy group, more preferably C₁₋₄ alkyl group, C₁₋₄ alkoxy group or C₁₋₃ alkylenedioxy group, most preferably C₁₋₄ alkyl group or C₁₋₄ alkoxy group, particularly preferably methyl group or methoxy group. R⁷ is preferably naphthyl group, 4-methoxy-3-methylphenyl group or 3,4-methylenedioxyphenyl group, and particularly preferably naphthyl group or 4-methoxy-3-methylphenyl group. X¹ is preferably C₁₋₆ alkylene group, particularly preferably methylene group. X² is preferably methylene group. X³ is preferably methylene group. X⁴ and X⁵ are each preferably a single bond.

[0072] R¹¹ is preferably hydrogen atom, halogen atom, C₁₋₄ alkyl group, hydroxy(C₁₋₆)alkyl group, C₁₋₆ alkoxy(C₁₋₆)alkyl group, hydroxyl group, C₁₋₄ alkoxy group, cyano group or nitro group, more preferably hydrogen atom, chlorine atom, methyl group, hydroxymethyl group, methoxy group, cyano group, hydroxyl group or nitro group, particularly preferably hydrogen atom.

[0073] R¹² is preferably hydrogen atom, halogen atom, C₁₋₄ alkyl group, hydroxy(C₁₋₆)alkyl group, C₁₋₆ alkoxy(C₁₋₆)alkyl group, hydroxyl group, C₁₋₄ alkoxy group, cyano group or nitro group, more preferably chlorine atom, methyl group, hydroxymethyl group, methoxy group, cyano group, hydroxyl group or nitro group, particularly preferably methyl group, hydroxymethyl group or methoxy group.

[0074] R²¹ is preferably methyl group, ethyl group, cyclopropyl group or cyclobutyl group, more preferably methyl group or cyclopropyl group, particularly preferably cyclopropyl group. The configuration of



[0075] A preferable embodiment of the compound [I] of the present invention is compound [I-2], and more preferably compound [I-3].

[0076] Specific examples of the preferable compounds of the present invention are given in the following:


(2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(2-methoxyphenyl)methoxy]propan-2-ol,
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(2-methylphenyl)methoxy]propan-2-ol,
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(phenyl)methoxy]propan-2-ol,
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-methoxyphenyl]ethoxy]propan-2-ol,
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methylphenyl)ethoxy]propan-2-ol,

(2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methoxyphenyl)propoxy]propan-2-ol,
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-cyanophenyl)ethoxy]propan-2-ol,
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(3-methoxyphenyl)ethoxy]propan-2-ol, and
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(3-methylphenyl)ethoxy]propan-2-ol; and
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(2-hydroxymethylphenyl)methoxy]propan-
 2-ol.

[0077] When the compound of the present invention is used as a pharmaceutical product, its form is the compound itself (free form), a salt of the compound, a solvate of the compound or a prodrug of the compound, which is preferably a free form, a salt of the compound or a solvate of the compound, particularly preferably a salt of the compound.

[0078] Now the production method of the compound of the formula [I] of the present invention is explained in detail. It is needless to say that the present invention is not limited to these production methods. When constructing the compound of the present invention, the construction can be started from the moiety easily produced. When a reactive functional group is involved in each step, it may be protected or deprotected as appropriate. For easy progress of the reaction, a reagent other than the exemplified reagent can be used as appropriate.

[0079] Any compound obtained in each step can be isolated and purified by a conventional method, and in some cases, the next step is performed without isolation or purification.



wherein X is halogen atom, L^1 and L^2 are each a leaving group, such as halogen atom and sulfonyloxy group (e.g., 3-nitrobenzenesulfonyloxy group, p-toluenesulfonyloxy group, benzenesulfonyloxy group, p-bromobenzenesulfonyloxy group, methanesulfonyloxy group, trifluoromethanesulfonyloxy group and the like), R^3 is oxygen atom or sulfur atom, and other symbols are as defined above.

Step 1

[0080] By reacting aldehyde compound (4) with Grignard reagent (5) in diethyl ether, tetrahydrofuran, 1,4-dioxane and the like or a mixed solvent thereof at a temperature of from $-80^\circ C$ to room temperature, compound (6) can be

obtained. The Grignard reagent can be prepared by a known method.

[0081] Alternatively, ketone compound (4') is reduced with a reducing agent such as lithium aluminum hydride, sodium borohydride, lithium borohydride and the like in diethyl ether, tetrahydrofuran, 1,4-dioxane, isopropanol and the like or a mixed solvent thereof at a temperature of from -10°C to room temperature to give compound (6). In this case, asymmetric reduction using an asymmetric reducing agent such as B-chlorodiisopinocampheylborane and the like, or asymmetric hydrogenation using a ruthenium complex such as dichloro[(S)-(-)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl]ruthenium (II) and the like can be also conducted.

Step 2

[0082] By reacting the compound (6) to be obtained in Step 1 with compound (7) in N,N-dimethylformamide, dimethyl sulfoxide, tetrahydrofuran and the like or a mixed solvent thereof in the presence of a base such as sodium hydride, sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium hydrogencarbonate, potassium hydrogencarbonate, triethylamine, N,N-diisopropylethylamine, N-methylmorpholine, pyridine, 4-dimethylaminopyridine and the like at a temperature of from 0°C to room temperature, compound (8) can be obtained. In this case, L¹ is a leaving group having greater reactivity than does L².

[0083] When a compound wherein R³ is hydroxyl group or mercapto group, R⁴ is hydrogen atom and X² is methylene group is desired, compound (7') is used instead of compound (7). In this case, alkylammonium hydrogensulfate such as tetrabutylammonium hydrogensulfate and the like can be added.

[0084] Upon determination of the reagent and leaving group to be used, a stereoselective reaction can be carried out.

Step 3


[0085] By reacting compound (8) to be obtained in Step 2 with compound (9) in N,N-dimethylformamide, dimethyl sulfoxide, tetrahydrofuran and the like or a mixed solvent thereof in the presence of a base such as sodium hydride, sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate and the like at a temperature of from 0°C to room temperature, a compound of the formula [I] can be obtained.

[0086] In addition, by reacting compound (8') with compound (9) in methanol, ethanol, n-propanol, isopropanol, tetrahydrofuran, 1,4-dioxane, acetonitrile, toluene and the like or a mixed solvent thereof at a temperature of from room temperature to refluxing temperature, a compound of the formula [I] wherein R³ is hydroxyl group or mercapto group, R⁴ is hydrogen atom and X² is methylene group can be obtained. In this case, alkali perchlorate such as lithium perchlorate and the like is preferably added.

[0087] When an acid addition salt of compound of the formula [I] is desired, a known method can be used. For example, compound of the formula [I] is dissolved in water, methanol, ethanol, n-propanol, isopropanol, diethyl ether, tetrahydrofuran, 1,4-dioxane, ethyl acetate, dichloromethane, 1,2-dichloroethane, chloroform and the like or a mixed solvent of these, the above-mentioned solvent, in which a desired acid has been dissolved, is added and the precipitated crystals are collected by filtration, or the solution is concentrated under reduced pressure.

[0088] When the acid addition salt of the compound of the formula [I] is converted to a free form, the acid addition salt of the compound of the formula [I] is subjected to partitioning between two phases of an aqueous solution of a base such as sodium hydrogencarbonate, potassium hydrogencarbonate, sodium carbonate, potassium carbonate, sodium hydroxide, potassium hydroxide, lithium hydroxide and the like and a solvent such as ethyl acetate, dichloromethane, 1,2-dichloroethane, chloroform, methyl ethyl ketone, toluene and the like to give a free form of the compound of the formula [I].

[0089] The compound (4') can be prepared by the following method.



wherein each symbol is as defined above.


Step 4

[0090] By reacting compound (10) with N,O-dimethylhydroxylamine or a salt thereof in a solvent such as dichloromethane, chloroform, 1,2-dichloroethane, tetrahydrofuran, 1,4-dioxane and the like in the presence of a base such as triethylamine, N,N-diisopropylethylamine and the like, compound (11) can be obtained.

Step 5

[0091] By reacting compound (11) to be obtained in Step 4 with Grignard reagent (5) in diethyl ether, tetrahydrofuran, 1,4-dioxane and the like or a mixed solvent thereof at a temperature of from -80°C to room temperature, compound (4') can be obtained.

[0092] The compound (9) can be prepared by the following method.



wherein Rh is carboxy-protecting group such as benzyl group, tert-butyl group and the like and each of other symbols is as defined above.

Step 6

[0093] By reacting compound (12) with compound (13) in a solvent such as tetrahydrofuran, n-hexane and the like in the presence of a base such as n-butyllithium and the like and hexamethylphosphoramide, compound (14) can be obtained.


Step 7

[0094] In this step, compound (14) to be obtained in Step 6 is subjected to Curtius rearrangement to give compound (15). The compound (14) is reacted with alkyl halocarbonate such as ethyl chlorocarbonate and the like in water, acetone, methyl ethyl ketone and the like or a mixed solvent thereof in the presence of a base such as triethylamine, N,N-diisopropylethylamine and the like. Then, sodium azide is reacted, and the obtained compound is subjected to rearrangement under heating and reacted with alcohol represented by Rh-OH to give compound (15).

Step 8

[0095] In this step, -CO₂Rh of compound (15) to be obtained in Step 7 is deprotected by a method generally employed for removing this protecting group. For example, when Rh is benzyl group, compound (15) is subjected to hydrogenation using a catalyst such as palladium carbon, palladium black, palladium hydroxide on carbon, Raney-nickel and the like in a solvent such as methanol, ethanol, n-propanol, isopropanol, tetrahydrofuran, 1,4-dioxane and the like to give compound (9). When, for example, Rh is tert-butyl group, a reaction using an acid such as hydrogen chloride, sulfuric acid, hydrogen bromide and the like in water, methanol, ethanol, n-propanol, isopropanol, tetrahydrofuran, 1,4-dioxane, acetic acid and the like or a mixed solvent thereof gives compound (9).

[0096] When, of the compounds (9), compound (9') wherein -X³-X⁴-X⁵- is methylene is desired, the following method can be employed.



wherein each symbol is as defined above.

Step 9

[0097] By reacting compound (17) with compound (16) in a solvent such as tetrahydrofuran, n-hexane and the like in the presence of a base such as n-butyllithium and the like, compound (18) can be obtained.

Step 10


[0098] By reacting compound (18) to be obtained in Step 9 in acetonitrile and acetic acid by adding sulfuric acid, compound (19) is obtained.

Step 11

[0099] By reacting compound (19) to be obtained in Step 10 in water, methanol, ethanol, n-propanol, isopropanol, tetrahydrofuran, 1,4-dioxane, acetic acid and the like or a mixed solvent thereof using an acid such as hydrogen chloride, sulfuric acid, hydrogen bromide and the like under heating, compound (9') is obtained.

[0100] The compound (9') can be also obtained by reacting compound (19) in a solvent such as water, methanol, ethanol, n-propanol, isopropanol, tetrahydrofuran, 1,4-dioxane, ethylene glycol and the like using a base such as sodium hydroxide, potassium hydroxide, lithium hydroxide and the like under heating.

[0101] The compound (9') can be also obtained by the following method.



wherein each symbol is as defined above.

Step 12

[0102] By reacting compound (21) with compound (20) in a solvent such as tetrahydrofuran, N,N-dimethylformamide, dimethyl sulfoxide and the like, in the presence of tetraalkylammonium halide such as tetrabutylammonium fluoride and the like and trialkylhalosilane such as tert-butyldimethylchlorosilane and the like, compound (22) can be obtained.

Step 13

[0103] By subjecting compound (22) to be obtained in Step 12 to halogenation using a halogenating agent such as thionyl chloride, oxalyl chloride and the like, compound (23) can be obtained. In this reaction, the halogenating agent itself to be used may be used as a solvent, or a solvent such as dichloromethane, chloroform, 1,2-dichloroethane, tetrahydrofuran, 1,4-dioxane and the like may be used.

Step 14

[0104] By subjecting compound (23) to be obtained in Step 13 to hydrogenation in a solvent such as methanol, ethanol, n-propanol, isopropanol, tetrahydrofuran, 1,4-dioxane, ethyl acetate and the like in the presence of a catalyst such as palladium carbon, palladium black, palladium hydroxide on carbon and the like, compound (24) can be obtained. In this reaction, pressurization to some extent is preferable.

Step 15

[0105] By subjecting compound (24) to be obtained in Step 14 to hydrogenation using a catalyst such as Raney-nickel and the like in a solvent such as methanol, ethanol, n-propanol, isopropanol, tetrahydrofuran, 1,4-dioxane and the like, compound (9') can be obtained. In this reaction, pressurization to some extent is preferable.

[0106] The intermediate compounds shown in the above-mentioned production method are useful as an intermediate for compound [I]. Of these intermediate compounds, the following compounds are novel compounds:

(A) compound [II] consisting of part of compound (6) and compound to be obtained by Step 2 of Example 57 below, a salt thereof and a solvate thereof (R^{11} is preferably C_{1-4} alkyl group, hydroxy(C_{1-6})alkyl group or C_{1-4} alkoxy group),

(B) compound [III] consisting of part of compound (4') and compound to be obtained by Step 2 of Example 58 below, a salt thereof and a solvate thereof (R^{11} is preferably C_{1-4} alkoxy group),

(C) compound [IV] consisting of compound (14), compound (15) and compound (24) (in the formulas (14), (15) and (24), R^5 and R^6 are methyl group, and $-X^3-X^4-X^5-R^7$ group is 2-naphthylmethyl group), a salt thereof and a solvate thereof.

[0107] The thus-obtained compound of the formula [I] of the present invention has a superior calcium receptor antagonistic action. When the compound of the present invention is to be used as a therapeutic agent of osteoporosis, hypoparathyroidism, osteosarcoma, periodontal disease, bone fracture, osteoarthritis, chronic rheumatoid arthritis, Paget's disease, humoral hypercalcemia, autosomal dominant hypocalcemia and the like, it is generally administered systemically or topically, and orally or parenterally.

[0108] While the dose varies depending on age, body weight, condition, treatment effect, administration method, treatment period and the like, it is generally 0.01 mg to 10 g for an adult per day, which is given once or in several portions a day by oral or parenteral administration.

[0109] When the compound of the present invention is prepared into a solid composition for oral administration, a dosage form of tablet, pill, powder, granule and the like can be employed. In such a solid composition, one or more active ingredient is admixed with at least one inert diluent, dispersing agent, absorbent and the like, such as lactose, mannitol, glucose, hydroxypropyl cellulose, crystalline cellulose, starch, polyvinyl hydrin, magnesium aluminometasilicate, anhydrous silicic acid powder and the like. The composition may contain an additive other than diluent according to a conventional method.

[0110] For preparation of tablets or pills, gastric or enteric film of sucrose, gelatin, hydroxypropyl cellulose, hydroxymethylcellulose phthalate and the like may be applied or two or more layers may be formed. In addition, they may be prepared into capsules of gelatin or ethylcellulose.

[0111] For preparation of liquid composition for oral administration, a dosage form such as pharmaceutically acceptable emulsion, solution, suspension, syrup, elixir and the like can be employed. The diluent to be used is, for example, purified water, ethanol, vegetable oil, emulsifier and the like. This composition may contain diluent and an adjuvant other than the diluent, such as wetting agent, suspending agent, sweetener, flavor, perfume, preservative and the like.

[0112] For preparation of parenteral injection, sterile aqueous or nonaqueous solvent, solubilizer, suspending agent or emulsifier is used. Examples of the aqueous solvent, solubilizer and suspending agent include distilled water for injection, physiological saline, cyclodextrin and derivatives thereof, organic amines such as triethanolamine, diethanolamine, monoethanolamine, triethylamine and the like, inorganic alkali solution and the like.

[0113] When a water-soluble solvent is to be prepared, for example, propylene glycol, polyethylene glycol, vegetable oil such as olive oil, alcohol such as ethanol, and the like may be used. As the solubilizer, for example, surfactant (forming a mixed micelle) such as polyoxyethylene hydrogenated castor oil, sucrose esters of fatty acid and the like, lecithin or hydrogenated lecithin (forming a liposome) and the like can be used. In addition, an emulsion preparation consisting of a water-insoluble solvent such as vegetable oil and the like, and lecithin, polyoxyethylene hydrogenated castor oil, polyoxyethylene polyoxypropylene glycol and the like may be formed.

[0114] As other compositions for parenteral administration, an external liquid, liniment such as ointment, suppository, pessary and the like, containing one or more active ingredients and prepared by a method known per se may be formulated.

Examples

[0115] The compound of the formula [I] of the present invention and its production methods are explained in detail by referring to the following Examples, which are not to be construed as limitative.


Example 1

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(2-methoxyphenyl)methoxy]propan-2-ol.

Step 1

(Cyclopropyl) (2-methoxyphenyl)methanol.

[0116]




[0117] Magnesium (10.7 g) was suspended in tetrahydrofuran (80 ml), and to the suspension was added iodine (5 mg). Bromocyclopropane (32.0 ml) was added dropwise thereto over 1.5 hr and the mixture was heated under reflux for 1.5 hr. Thereto was added tetrahydrofuran to give a 1M cyclopropylmagnesium bromide-tetrahydrofuran solution. Subsequently, o-anisaldehyde (8.17 g) was dissolved in tetrahydrofuran (150 ml), the 1M cyclopropylmagnesium bromide-tetrahydrofuran solution (90 ml) was added dropwise thereto over 50 min under ice-cooling, and the mixture was stirred at room temperature for 12 hr. The reaction mixture was ice-cooled, a saturated aqueous ammonium chloride solution (9 ml) was added, and the reaction mixture was stirred at room temperature for 30 min. The organic layer was dried over magnesium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (n-hexane:ethyl acetate = 9:1) to give the title compound (6.13 g).

¹H-NMR(300MHz, δppm, DMSO-d₆) 7.41(1H,d,J=7.8Hz), 7.19(1H,dt,J=1.6, 7.7Hz), 6.92(2H,t,J=7.2Hz), 4.88(1H,d,J=4.8Hz), 4.55(1H,t,J=5.5Hz), 3.76(3H,s), 1.09-1.00(1H,m), 0.33-0.25(4H,m). MS(ESI,m/z) 161(M+H-H₂O)⁺.

Step 2

(R)-2-[(Cyclopropyl)(2-methoxyphenyl)methoxymethyl]oxirane

[0118]




[0119] (Cyclopropyl)(2-methoxyphenyl)methanol (3.57 g) obtained in Step 1 was dissolved in N,N-dimethylformamide (50 ml), sodium hydride (960 mg, 60% oil) was added and the mixture was stirred for 3 min. To the resulting mixture was added (R)-glycidyl 3-nitrobenzenesulfonate (6.22 g) and the mixture was stirred at room temperature for 12 hr. The reaction mixture was poured into water and extracted with diethyl ether. The organic layer was washed successively with water and saturated aqueous sodium chloride solution, dried over sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (n-hexane:ethyl acetate=88:12) to give the title compound (1.10 g).

Step 3

2-Methyl-1-(naphthalen-2-yl)-2-nitropropanol


[0120]



[0121] To tetrabutylammonium fluoride 3 hydrate (2.41 g) was added tetrahydrofuran (20 ml) and the mixture was ice-cooled. To the mixture were added 2-nitropropane (2.7 ml), 2-naphthaldehyde (3.12 g) and triethylamine (2.8 ml) under an argon atmosphere. To the mixture was added a solution of tert-butyldimethylchlorosilane (4.51 g) in tetrahydrofuran (20 ml). The resulting mixture was stirred for 40 min while allowing to warm to room temperature from ice-cooling. After removing insoluble matter by filtration, the filtrate was poured into a solution (500 ml) of diethyl ether-n-hexane=1:3 and the mixture was washed twice with water (40 ml). The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained residue was subjected to crystallization from n-hexane to give the title compound (3.78 g).

Step 4


2-(1-Chloro-2-methyl-2-nitropropyl)naphthalene

[0122]

[0123] To 2-methyl-1-(naphthalen-2-yl)-2-nitropropanol (1.23 g) obtained in Step 3 was added thionyl chloride (3.1 ml), and the resulting mixture was heated under reflux for 1 hr, and stirred at room temperature for 12 hr. The reaction mixture was concentrated under reduced pressure and the residue thus obtained was poured into water and extracted with ethyl acetate. The organic layer was washed successively with water and saturated aqueous sodium hydrogen-carbonate solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (n-hexane:ethyl acetate=93:7). Recrystallization from a solution of n-hexane:ethyl acetate=4:1, yielded title compound (776 mg).

Step 5

2-(2-Methyl-2-nitropropyl)naphthalene


[0124]

[0125] 2-(1-Chloro-2-methyl-2-nitropropyl)naphthalene (200 mg) obtained in Step 4 was dissolved in methanol (5 ml) and ethyl acetate (5 ml) and to the solution was added 10% palladium carbon (20 mg). A hydrogenation was performed at pressure of 3 atm for 2 hr. After filtration of the reaction mixture using Celite, the title compound (139 mg) was obtained by purification using silica gel column chromatography (n-hexane:ethyl acetate=95:5).

¹H-NMR(300MHz, δ ppm, CDCl₃) 7.85-7.76(3H,m), 7.58(1H,s), 7.50-7.44(2H,m), 7.22(1H,dd,J=1.7, 8.4Hz), 3.37(2H,s), 1.69(6H,s). MS(APCI,m/z) 183(M+H-NO₂)⁺.

Step 6

[2-Methyl-1-(naphthalen-2-yl)propan-2-yl]amine

[0126]


[0127] Raney nickel W2 (200 mg) was suspended in ethanol (10 ml), to the suspension was added 2-(2-methyl-2-nitropropyl)naphthalene (134 mg) obtained in Step 5, and hydrogenation was performed at a pressure of 3.5 atm for 12 hr. The reaction mixture was filtered through Celite and concentrated under reduced pressure. The obtained

residue was subjected to crystallization from ethyl acetate to give the title compound (70 mg).

Step 7

2-Methyl-1-(naphthalen-2-yl)-2-propanol

[0128]




[0129] 2-Methylnaphthalene (7.11 g) was dissolved in tetrahydrofuran (100 ml) and the solution was cooled to -69°C. To the solution was added dropwise a 1.6M n-butyllithium-tetrahydrofuran solution (34 ml), and then a solution of acetone (4.41 ml) in tetrahydrofuran (4.41 ml). The reaction mixture was stirred for 12 hr while allowing to warm to room temperature. To the reaction mixture was added dropwise saturated ammonium chloride solution (6 ml), and the resulting mixture was poured into water (200 ml) and extracted with diethyl ether. The organic layer was washed with water, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (n-hexane:ethyl acetate=85:15) to give the title compound (3.80 g).

Step 8

N-[2-Methyl-1-(naphthalen-2-yl)propan-2-yl]acetamide

[0130]




[0131] To 2-methyl-1-(naphthalen-2-yl)-2-propanol (500 mg) obtained in Step 7 were added successively acetonitrile (0.5 ml) and acetic acid (0.5 ml). After ice-cooling the mixture, sulfuric acid (0.5 ml) was added dropwise. The mixture was stirred under ice-cooling for 20 min and poured into a 1N aqueous sodium hydroxide solution, and the mixture was extracted with diethyl ether. The organic layer was washed with water, dried over sodium sulfate, and concentrated under reduced pressure. The obtained residue was subjected to recrystallization from ethyl acetate-n-hexane to give the title compound (303 mg).

Step 9

[2-Methyl-1-(naphthalen-2-yl)propan-2-yl]amine

[0132]




[0133] To N-[2-methyl-1-(naphthalen-2-yl)propan-2-yl]acetamide (26 mg) obtained in Step 8 was added 6N hydrochloric acid (2 ml) and the mixture was heated under reflux for 5 hr. The reaction mixture was poured into water, made

basic with a 4N aqueous sodium hydroxide solution, and extracted with diethyl ether. The organic layer was washed with water, dried over potassium carbonate and concentrated under reduced pressure to give the title compound (19 mg).

Step 10

2,2-Dimethyl-3-(naphthalen-2-yl)propionic acid

[0134]




[0135] A solution of diisopropylamine (576 ml) in tetrahydrofuran (3.5 L) was cooled to -68°C, and a 2.6M n-butyllithium-n-hexane solution (1.5 L) was slowly added under an argon atmosphere. After addition of a solution of isobutyric acid (181 ml) in hexamethylphosphoramide (340 ml), the reaction mixture was stirred at room temperature for 30 min. After ice-cooling the mixture, a solution of 2-bromomethylnaphthalene (392 g) in tetrahydrofuran (1 L) was added dropwise, and the obtained solution was stirred at room temperature for 1 day. The reaction mixture was ice-cooled, 6N hydrochloric acid (665 ml) was added thereto, and the organic layer was separated from the aqueous layer. The separated organic layer was concentrated under reduced pressure to yield a residue. On the other hand, ethyl acetate and water were added to an aqueous layer and the organic layer was separated from the aqueous layer. The organic layer and the aforementioned residue were combined and washed three times with water. The organic layer was extracted with a solution of sodium hydroxide (80 g) in water (600 ml), and then extracted three times with a 4N aqueous sodium hydroxide solution (200 ml). The aqueous layer was washed with ethyl acetate, acidified with conc. hydrochloric acid, and extracted with ethyl acetate. The ethyl acetate layer was washed with aqueous sodium chloride solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (177.5 g). ¹H-NMR (300MHz, δppm, DMSO-d₆) 7.88-7.80(3H,m), 7.66(1H,s), 7.51-7.44(2H,m), 7.33(1H,dd,J=8.4, 1.6Hz), 2.96(2H,s), 1.13(6H,s). MS(FAB,m/z) 228(M)⁺.

Step 11

N-[2-Methyl-1-(naphthalen-2-yl)propan-2-yl]-benzyloxycarboxamide

[0136]



[0137] 2,2-Dimethyl-3-(naphthalen-2-yl)propionic acid (205.4 g) obtained in Step 10 was dissolved in water (173 ml), triethylamine (131 ml) and acetone (800 ml), and a solution of ethyl chlorocarbonate (101.5 ml) in acetone (400 ml) was added under ice-cooling. A solution of sodium azide (73.1 g) in water (400 ml) was added dropwise, and the reaction mixture was stirred at room temperature for 2 hr. To the reaction mixture were added water (1.5 L) and toluene (1.2 L), and the organic layer was separated from the aqueous layer. The organic layer was washed twice with water and three times with a saturated aqueous sodium chloride solution, and dried over anhydrous sodium sulfate. After the drying agent was filtered off, the filtrate was heated to 100°C over 4 hr under an argon atmosphere, and stirred at 100°C for 1 hr. The reaction mixture was concentrated under reduced pressure and benzyl alcohol (500 ml) was added. The mixture was stirred at 105°C for 1 day. The reaction mixture was concentrated under reduced pressure, and ethyl acetate and n-hexane were added to the obtained residue. The solution was treated with activated carbon and concentrated under reduced pressure, and the obtained residue was purified by silica gel column chromatography (n-

hexane:ethyl acetate=20:1) to give the title compound (240.5 g).


$^1\text{H-NMR}$ (300MHz, δppm , CDCl_3) 8.83-7.78(1H,m), 7.71-7.66(2H,m), 7.45-7.36(7H,m), 7.22(1H,dd,J=1.6, 8.4Hz), 5.13(2H,s), 4.54(1H,br s), 3.15(2H,s), 1.34(6H,s).

MS(FAB,m/z) 334(M+H) $^+$.

Step 12

[2-Methyl-1-(naphthalen-2-yl)propan-2-yl]amine

[0138]




[0139] Palladium hydroxide on carbon (19.7 g) was suspended in methanol (500 ml), to the suspension was added a solution of N-[2-methyl-1-(naphthalen-2-yl)propan-2-yl]-benzyloxycarboxamide (237.6 g) obtained in Step 11 in methanol (2 L), and a hydrogenation reaction was performed at room temperature overnight. After filtration of the reaction mixture, the filtrate was concentrated under reduced pressure. To the obtained residue was added ethyl acetate, and to the solution was added conc. hydrochloric acid (70 ml) to give crystals. The obtained crystals were suspended in water, and after an addition of 4N aqueous sodium hydroxide solution (350 ml), the mixture was extracted with ethyl acetate. The ethyl acetate layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give crystals of the title compound (138 g).

Step 13

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(2-methoxyphenyl)methoxy]propan-2-ol

[0140]



[0141] (R)-2-[(Cyclopropyl)(2-methoxyphenyl)methoxymethyl]-oxirane (703 mg) obtained in Step 2 was dissolved in ethanol (12 ml). To the solution was added [2-methyl-1-(naphthalen-2-yl)propan-2-yl]amine (120 ml) obtained either in Step 6, 9 or 12, and the reaction mixture was stirred at 60°C for 20 hr. The reaction mixture was concentrated under reduced pressure, and the obtained residue was purified by silica gel column chromatography (chloroform:methanol=97:3) to give the title compound (954 mg).

$^1\text{H-NMR}$ (300MHz, δppm , DMSO-d_6) 7.90-7.70(3H,m), 7.67(1H,s), 7.50-7.20(5H,m) 7.00-6.90(2H,m), 4.62(1H, br s), 4.29(1H,d,J=7.0Hz), 3.77(3H,s), 3.70-3.50(1H,m), 3.30-3.10(2H,m), 2.80-2.50(4H,m), 1.50-0.85(8H,m), 0.50-0.20(4H,m).


MS(APCI,m/z) 433(M+H) $^+$.

Example 2


(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(2-methylphenyl)methoxy]propan-2-ol

Step 1


(Cyclopropyl)(2-methylphenyl)methanol

[0142]**[0143]** Employing the same procedure described in Step 1 of Example 1, the title compound (5.36 g) was obtained using o-tolualdehyde (4.81 g) instead of o-anisaldehyde.¹H-NMR(300MHz, δ ppm, DMSO- d_6) 7.43(1H,d,J=7.2Hz), 7.18-7.10(3H,m), 4.95(1H,d,J=4.8Hz), 4.36(1H,dd,J=4.4, 6.5Hz), 2.31(3H,s), 1.20-1.07(1H,m), 0.47-0.20(4H,m).MS(APCI,m/z) 145(M+H-H₂O)⁺.**Step 2**

(R)-2-[(Cyclopropyl)(2-methylphenyl)methoxymethyl]oxirane

[0144]**[0145]** Employing the same procedure described in Step 2 of Example 1, the title compound (1.53 g) was obtained from (cyclopropyl)(2-methylphenyl)methanol (3.25 g) obtained in Step 1.**Step 3**

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(2-methylphenyl)methoxy]propan-2-ol

[0146]

[0147] Employing the same procedure described in Step 13 of Example 1, the title compound (849 mg) was obtained from (R)-2-[(cyclopropyl)(2-methylphenyl)methoxymethyl]oxirane (655 mg) obtained in Step 2.

¹H-NMR(300MHz, δ ppm, DMSO-d₆) 7.90-7.70(3H,m), 7.67(1H,s), 7.50-7.25(4H,m) 7.20-7.10(3H,m), 4.65(1H,br s), 4.06(1H,d,J=7.4Hz), 3.70-3.50(1H,m), 3.30-3.10(2H,m), 2.80-2.50(4H,m), 2.31(3H,s), 1.50-0.80(8H,m), 0.60-0.10(4H,m).

MS(APCI,m/z) 417(M+H)⁺.


Example 3

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(phenyl)methoxy]propan-2-ol

Step 1

(R)-2-[(Cyclopropyl)(phenyl)methoxymethyl]oxirane

[0148]




[0149] In the same manner as in Step 2 of Example 1, the title compound (296 mg) was obtained from α -cyclopropylbenzyl alcohol (740 mg).

Step 2

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(phenyl)methoxy]propan-2-ol

[0150]



[0151] In the same manner as in Step 13 of Example 1, the title compound (334 mg) was obtained from (R)-2-[(cyclopropyl)(phenyl)methoxymethyl]oxirane (204 mg) obtained in Step 1.

¹H-NMR(300MHz, δ ppm, DMSO-d₆) 7.90-7.70(3H,m), 7.67(1H,s), 7.50-7.20(8H,m) 4.65(1H,br s), 3.80-3.50(2H,m), 3.30-3.10(2H,m), 2.80-2.50(4H,m), 1.50-0.80(8H,m), 0.60-0.10(4H,m).


MS(APCI,m/z) 403(M+H)⁺.

Example 4

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methoxyphenyl)ethoxy]propan-2-ol

Step 1


1-(2-Methoxyphenyl)ethanol

[0152]

[0153] Lithium aluminum hydride (1.52 g) was suspended in tetrahydrofuran (100 ml), 2'-methoxyacetophenone (2.76 ml) was added under ice-cooling, and the mixture was stirred for 30 min. To the reaction mixture were successively added water (1.5 ml), 15% aqueous sodium hydroxide solution (1.5 ml) and water (4.5 ml). The organic layer was dried over magnesium sulfate and concentrated under reduced pressure to give the title compound (3.05 g).

Step 2


(R)-2-[1-(2-Methoxyphenyl)ethoxymethyl]oxirane

[0154]

[0155] In the same manner as in Step 2 of Example 1, the title compound (123 mg) was obtained from 1-(2-methoxyphenyl)ethanol (837 mg) obtained in Step 1.

Step 3

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methoxyphenyl)ethoxy]propan-2-ol

[0156]


[0157] In the same manner as in Step 13 of Example 1, the title compound (148 mg) was obtained from (R)-2-[1-(2-methoxyphenyl)ethoxymethyl]oxirane (115 mg) obtained in Step 2.

¹H-NMR(300MHz, δ ppm, DMSO-d₆) 7.85-7.70(3H,m), 7.61(1H,s), 7.50-7.20(5H,m), 7.00-6.80(2H,m), 4.88(1H,q,

J=6.4Hz), 3.85-3.75(4H,m), 3.45-3.35(2H,m), 2.85-2.60(4H,m), 1.40-1.37(3H,m), 1.10-1.08(6H,m).
MS(APCI,m/z) 408 (M+H)⁺.

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methoxyphenyl)ethoxy]propan-2-ol hydrochloride

[0158]



[0159] (2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methoxyphenyl)ethoxy]propan-2-ol (141 mg) obtained in Example 4 was dissolved in diethyl ether (5 ml), and 4N hydrogen chloride-ethyl acetate solution was added. The reaction mixture was concentrated under reduced pressure and diethyl ether (5 ml) was added. The mixture was concentrated under reduced pressure to give the title compound (154 mg).

¹H-NMR(300MHz, δppm, DMSO-d₆) 8.91(1H,br s), 8.54(1H,br s), 7.95-7.86(3H,m), 7.67(1H,s), 7.55-7.45(2H,m), 7.40-7.20(3H,m), 7.02-6.93(2H,m), 5.66-5.61(1H,m), 4.90-4.80(1H,m), 4.05-3.95(1H,m), 3.80(3H,s), 3.40-3.10(5H,m), 3.05-2.85(1H,m), 1.32(3H,d,J=6.4Hz), 1.09(6H,s).

MS(APCI,m/z) 408(M+H-HCl)⁺.


Example 6

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methylphenyl)ethoxy]propan-2-ol

Step 1

1-(2-Methylphenyl)ethanol

[0160]




[0161] In the same manner as in Step 1 of Example 4, the title compound (755 mg) was obtained from 2'-methylacetophenone (2.68 g).

Step 2

(R)-2-[1-(2-Methylphenyl)ethoxymethyl]oxirane

[0162]




[0163] In the same manner as in Step 2 of Example 1, the title compound (123 mg) was obtained from 1-(2-methylphenyl)ethanol (749 mg) obtained in Step 1.

Step 3

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methylphenyl)ethoxy]propan-2-ol

[0164]




[0165] In the same manner as in Step 13 of Example 1, the title compound (345 mg) was obtained from (R)-2-[1-(2-methylphenyl)ethoxymethyl]oxirane (192 mg) obtained in Step 2. ¹H-NMR(300MHz, δppm, CDCl₃) 7.85-7.70 (3H,m), 7.60(1H,s), 7.50-7.10(7H,m), 4.75-4.65(1H,m), 3.80-3.70(1H,m), 3.40-3.30(2H,m), 2.90-2.60(4H,m), 2.31(3H,s), 1.39(3H,d,J=6.4Hz), 1.10-1.07(6H,m). MS(APCI,m/z) 392(M+H)⁺.

Example 7

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methylphenyl)ethoxy]propan-2-ol hydrochloride

[0166]



[0167] In the same manner as in Example 5, the title compound (337 mg) was obtained from (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methylphenyl)ethoxy]propan-2-ol (339 mg) obtained in Example 6.

¹H-NMR(300MHz, δppm, DMSO-d₆) 8.97(1H,br s), 8.56(1H,br s), 7.95-7.85(3H,m), 7.77(1H,s), 7.75-7.65(2H,m), 7.40-7.32(2H,m), 7.25-7.12(3H,m), 5.66-5.61(1H,m), 4.73(1H,q,J=6.3Hz), 4.10-3.95(1H,br s), 3.40-3.10(5H,m), 3.05-2.90(1H,m), 2.31(3H,s), 1.34(3H,d,J=6.6Hz), 1.27(3H,s), 1.26(3H,s).

MS(APCI,m/z) 392(M+H-HCl)⁺.


Example 8

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methoxyphenyl)propoxy]propan-2-ol

Step 1

1-(2-Methoxyphenyl)propanol

[0168]




[0169] o-Anisaldehyde (2.72 g) was dissolved in tetrahydrofuran (50 ml), and 0.93M magnesium ethylbromide - tetrahydrofuran solution (31.3 ml) was added dropwise under ice-cooling over 10 min. The mixture was stirred at room temperature for 2 hr. The reaction mixture was ice-cooled, and saturated aqueous ammonium chloride solution (40 ml) and water (40 ml) were added and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium chloride solution, dried over magnesium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (n-hexane:ethyl acetate=4:1) to give the title compound (3.15 g).

Step 2

(R)-2-[1-(2-Methoxyphenyl)propoxymethyl]oxirane

[0170]




[0171] In the same manner as in Step 2 of Example 1, the title compound (264 mg) was obtained from 1-(2-methoxyphenyl)propanol (831 mg) obtained in Step 1.

Step 3

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methoxyphenyl)propoxy]propan-2-ol

[0172]



[0173] In the same manner as in Step 13 of Example 1, the title compound (403 mg) was obtained from (R)-2-[1-(2-methoxyphenyl)propoxymethyl]oxirane (264 mg) obtained in Step 2.

¹H-NMR (300MHz, δ ppm, CDCl₃) 7.85-7.70(3H,m), 7.61(1H,s), 7.50-7.20(5H,m), 7.00-6.90(1H,m), 6.85-6.75(1H,m), 4.67(1H,t,J=6.3Hz), 3.85-3.70(4H,m), 3.42-3.27(2H,m), 2.90-2.65(6H,m), 1.15-1.05(6H,m), 0.91(3H,t,J=7.4Hz).

MS(APCI,m/z) 422(M+H)⁺.

Example 9 - Example 20

[0174] In the same manner as in Examples 1-8, the compounds of Example 9 - Example 20 were obtained. They are shown in Table 1 and Table 2.

Table 1








Ex. 9		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 7.90-7.30(11H,m), 4.80-4.60(2H,m), 3.75-3.55(1H,m) 3.45-3.25(2H,m), 2.80-2.40(4H,m), 1.45-1.30(3H,m), 1.05-0.90(6H,m). MS(APCI,m/z) 403(M+H) ⁺ .
Ex. 10		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 9.03(1H,br s), 8.61(1H,br s), 7.95-7.70(6H,m), 7.65(1H,t,J=6.4Hz), 7.60-7.40(3H,m), 7.38(1H,d,J=8.4Hz) 5.75-5.65(1H,m), 4.90-4.70(1H,m), 4.15-3.90(1H,m), 3.50-2.80(6H,m), 1.47(3H,d,J=6.4Hz), 1.07(6H,s). MS(APCI,m/z) 403(M+H-HCl) ⁺ .
Ex. 11		¹ H-NMR(300MHz, δ ppm,CDCl ₃) 7.85-7.70(3H,m), 7.61(1H,s), 7.50-7.20(5H,m), 7.00-6.80(2H,m), 4.88(1H,q,J=6.4Hz), 3.85-3.75(4H,m), 3.45-3.35(2H,m), 2.85-2.60(4H,m), 1.40-1.37(3H,m), 1.10-1.08(6H,m). MS(APCI,m/z) 408(M+H) ⁺ .
Ex. 12		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 8.99(1H,br s), 8.56(1H,br s), 7.95-7.85(3H,m), 7.76(1H,s), 7.57-7.47(2H,m), 7.42-7.35(1H,m), 7.33-7.23(1H,m), 6.95-6.80(3H,m), 5.61-5.65(1H,m), 4.55-4.42(1H,m), 4.10-3.95(1H,m), 3.76(3H,s), 3.40-3.10(5H,m), 3.00-2.80(1H,m), 1.37(3H,d,J=6.6Hz), 1.09(6H,s). MS(APCI,m/z) 408(M+H-HCl) ⁺ .
Ex. 13		¹ H-NMR(300MHz, δ ppm,CDCl ₃) 7.85-7.70(3H,m), 7.60(1H,s), 7.50-7.40(2H,m), 7.35-7.15(2H,m), 7.10-7.00(3H,m), 4.45-4.35(1H,m), 3.80-3.70(1H,m), 3.35-3.25(2H,m), 2.85-2.60(4H,m), 2.34(3H,s), 1.43-1.40(3H,m), 1.10-1.07(6H,m). MS(APCI,m/z) 392(M+H) ⁺ .
Ex. 14		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 9.03(1H,br s), 8.57(1H,br s), 7.95-7.85(3H,m), 7.76(1H,s), 7.57-7.45(2H,m), 7.41-7.33(1H,m), 7.30-7.20(1H,m), 7.18-7.07(3H,m), 5.65-5.62(1H,m), 4.50-4.40(1H,m), 4.08-3.92(1H,m), 3.40-3.08(5H,m), 3.00-2.80(1H,m), 2.32(3H,s), 1.36(3H,d,J=6.4Hz), 1.26(6H,s). MS(APCI,m/z) 392(M+H-HCl)
Ex. 15		¹ H-NMR(300MHz, δ ppm,CDCl ₃) 7.85-7.70(3H,m), 7.60(1H,s), 7.50-7.40(2H,m), 7.35-7.20(6H,m), 4.45-4.35(1H,m), 3.80-3.70(1H,m), 3.40-3.25(2H,m), 2.90-2.60(4H,m), 1.43(3H,d,J=6.5Hz), 1.10-1.06(6H,m). MS(APCI,m/z) 378(M+H) ⁺ .

Table 2

Ex. 16		¹ H-NMR(300MHz, δ ppm,DMSO- d_6) 8.99 (1H,br s), 8.56(1H,br s), 7.90-7.87(m,3H), 7.77(1H,s), 7.55-7.48(2H,m), 7.40-7.27(6H,m), 5.63(1H,t,J=5.1Hz), 4.52-4.48(1H,m), 4.08-3.93(1H,m), 3.40-2.80(6H,m), 1.37(3H,d,J=6.3Hz), 1.27(3H,s), 1.26(3H,s). MS(APCI,m/z) 378(M+H-HCl) ⁺ .
Ex. 17		¹ H-NMR(300MHz, δ ppm,DMSO- d_6) 8.97 (1H,br s), 8.59(1H,br s), 7.95-7.85(3H,m), 7.77(1H,s), 7.57-7.47(2H,m), 7.45-7.22(3H,m), 7.05-6.95(2H,m), 5.64-5.59(1H,m), 4.66 (1H,t,J=6.0Hz), 4.10-3.95(1H,m), 3.80(3H,s), 3.35-3.10(5H,m), 3.05-2.85(1H,m), 1.70-1.55(2H,m), 1.27(6H,s), 0.92-0.82(3H,m). MS(APCI,m/z) 422(M+H-HCl) ⁺ .
Ex. 18		¹ H-NMR(300MHz, δ ppm,DMSO- d_6) 9.09 (1H,br s), 8.61(1H,br s), 7.92-7.88(3H,m), 7.78(1H,s), 7.60-7.45(2H,m), 7.40-7.20(6H,m), 5.65-5.55(1H,m), 4.30-4.20(1H,m), 4.15-3.90(1H,m), 3.40-2.80(6H,m), 1.85-1.52(2H,m), 1.27(6H,s), 0.90-0.75(3H,m). MS(APCI,m/z) 392(M+H-HCl) ⁺ .
Ex. 19		¹ H-NMR(300MHz, δ ppm,CDCl ₃) 7.85-7.70(3H,m), 7.61(1H,s), 7.50-7.40(2H,m), 7.35-7.20(6H,m), 4.15-4.05(1H,m), 3.80-3.70(1H,m), 3.40-3.20(2H,m), 2.90-2.45(5H,m), 2.10-1.60(6H,m), 1.10-1.00(6H,m). MS(APCI,m/z) 418(M+H) ⁺ .
Ex. 20		¹ H-NMR(300MHz, δ ppm,DMSO- d_6) 8.90 (1H,m), 8.56(1H,m), 7.92-7.80(3H,m), 7.77 (1H,s), 7.55-7.45(2H,m), 7.40-7.22(6H,m), 5.61-5.54(1H,m), 4.26(1H,d,J=8.4Hz), 4.02(1H,br s), 3.40-2.80(6H,m), 2.05-1.92(2H,m), 1.83-1.60(5H,m), 1.26(6H,s). MS(APCI,m/z) 418(M+H-HCl) ⁺ .


Example 21

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1R)-1-(2-methoxyphenyl)ethoxy]propan-2-ol

Step 1

(R) -1- (2-Methoxyphenyl)ethanol

[0175]




[0176] To a solution of (+)-B-chlorodiisopinocampheylborane (10.5 g) in tetrahydrofuran (50 ml) was added dropwise 2'-methoxyacetophenone (10.5 g) at -25°C. The mixture was stirred at -25°C for 1 hr and concentrated under reduced pressure. To the residue were added diethyl ether (100 ml) and diethanolamine (18.1 g) and the reaction mixture was stirred at room temperature for 2 hr. The precipitated solid was filtered off. The filtrate was concentrated under reduced pressure and purified by distillation under reduced pressure (bp 74-77°C/1 mmHg) to give the title compound (8.00 g).

Step 2

(R)-2-[(1R)-1-(2-Methoxyphenyl)ethoxymethyl]oxirane

[0177]




[0178] In the same manner as in Step 2 of Example 1, the title compound (930 mg) was obtained from (R)-1-(2-methoxyphenyl)ethanol (1.52 g) obtained in Step 1.

Step 3

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1R)-1-(2-methoxyphenyl)ethoxy]propan-2-ol

[0179]




[0180] In the same manner as in Step 13 of Example 1, the title compound (770 mg) was obtained from (R)-2-[(1R)-1-(2-methoxyphenyl)ethoxymethyl]oxirane (417 mg) obtained in Step 2.

¹H-NMR(300MHz, δ ppm, CDCl₃) 7.85-7.70(3H,m), 7.62(1H,s), 7.50-7.20(5H,m), 7.00-6.80(2H,m), 4.88(1H,q, J=6.5Hz), 3.85-3.75(4H,m), 3.38(2H,d,J=5.1Hz), 2.90-2.60(4H,m), 1.38(3H,d,J=6.6Hz), 1.10(3H,s), 1.07(3H,s). MS(APCI,m/z) 408(M+H)⁺.

Example 22

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1R)-1-(2-methoxyphenyl)ethoxy]propan-2-ol hydrochloride

[0181]

[0182] In the same manner as in Example 5, the title compound (810 mg) was obtained from (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1R)-1-(2-methoxyphenyl)ethoxy]propan-2-ol obtained in Example 21.

¹H-NMR(300MHz, δ ppm, DMSO- d_6) 8.94(1H,br s), 8.56(1H,br s), 7.95-7.85(3H, m), 7.77(1H,s), 7.57-7.45(2H,m), 7.42-7.20(3H,m), 7.05-6.95(2H,m), 5.61(1H,d,J=4.8Hz), 4.84(1H,q,J=6.3Hz), 4.10-3.95(1H,m), 3.80(3H,s), 3.40-3.10(5H,m), 3.00-2.85(1H,m), 1.32(3H,d,J=6.3Hz), 1.26(6H,s).


MS(APCI,m/z) 408(M+H-HCl)⁺.

Example 23

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1R)-(cyclopropyl)(2-methylphenyl)methoxy]propan-2-ol

Step 1

N-Methoxy-2, N-dimethylbenzamide


[0183]

[0184] To a solution of o-toluy chloride (3.09 g) and triethylamine (5.58 ml) in dichloromethane (150 ml) was added N,O-dimethylhydroxylamine hydrochloride (3.90 g) under ice-cooling and the obtained mixture was stirred at room temperature for 12 hr. The reaction mixture was concentrated under reduced pressure, and the residue was diluted with ethyl acetate, washed successively with 1N hydrochloric acid, water, a saturated aqueous sodium hydrogencarbonate solution and saturated brine. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give the title compound (3.58 g).

Step 2

Cyclopropyl 2-methylphenyl ketone

[0185]




[0186] To a solution of N-methoxy-2,N-dimethylbenzamide (3.55 g) obtained in Step 1 in tetrahydrofuran (40 ml) was added dropwise a solution (29.7 ml) of 1M cyclopropylmagnesium bromide-tetrahydrofuran under ice-cooling, and the mixture was stirred at room temperature for 12 hr. To the reaction mixture was added a 4N hydrogen chloride-ethyl acetate solution (10 ml), and the resulting mixture was concentrated under reduced pressure. The residue was diluted with ethyl acetate, washed successively with 1N hydrochloric acid, water, saturated sodium hydrogencarbonate and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate:hexane=1:5) to give the title compound (1.05 g).

Step 3

(R) - (Cyclopropyl) (2-methylphenyl)methanol

[0187]




[0188] To a suspension of dichloro[(S)-(-)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl]ruthenium(II) (851 mg), (1S, 2S)-(-)-1,2-diphenylethylenediamine (191 mg) and potassium-tert-butoxide (270 mg) in isopropanol (90 ml) was added cyclopropyl 2-methylphenyl ketone (9.61 g) obtained in Step 2, and the mixture was hydrogenated at room temperature and at medium pressure (3.0 kgf/cm²) for 60 hr. The reaction mixture was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (ethyl acetate:hexane=1:9) to give the title compound (6.92 g).

¹H-NMR(300MHz, δ ppm, DMSO-d₆) 7.43(1H,d,J=6.8Hz), 7.18-7.09(3H,m), 4.97(1H,d,J=4.6Hz), 4.36(1H,dd,J=4.6, 6.4Hz), 2.30(3H,s), 1.17-1.07(1H,m), 0.41-0.24(4H,m).

MS(APCI,m/z) 145(M+H-H₂O)⁺.

Step 4


(R)-2-[(1R)-(Cyclopropyl)(2-methylphenyl)methoxymethyl]oxirane

[0189]

[0190] Employing the same procedure described in Step 2 of Example 1, the title compound (1.80 g) was obtained from (R)-(cyclopropyl)(2-methylphenyl)methanol (3.24 g) obtained in Step 3.

Step 5

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1R)-(cyclopropyl)(2-methylphenyl)methoxy]propan-2-ol

[0191]

[0192] Employing the same procedure described in Step 13 of Example 1, the title compound (1.17g) was obtained from (R)-2-[(1R)-(cyclopropyl)(2-methylphenyl)methoxymethyl]oxirane (655 mg) obtained in Step 4.

¹H-NMR(300MHz, δ ppm, DMSO- d_6) 7.90-7.70(3H,m), 7.66(1H,s), 7.50-7.25(4H,m) 7.20-7.10(3H,m), 4.64(1H,br s), 4.06(1H,d,J=7.4Hz), 3.70-3.50(1H,m), 3.30-3.10(2H,m), 2.80-2.50(4H,m), 2.31(3H,s), 1.50-0.85(2H,m), 0.60-0.10(4H,m)


MS(APCI,m/z) 417(M+H)⁺.

Example 24

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1R)-(cyclopropyl)(2-methoxyphenyl)methoxy]propan-2-ol

Step 1


2,N-Dimethoxy-N-methylbenzamide

[0193]

Step 2

Cyclopropyl 2-methoxyphenyl ketone

[0194]



[0195] In the same manner as in Step 1 of Example 23, the title compound (3.56 g) was obtained from 2-methoxybenzoyl chloride (3.41 g).

[0196] In the same manner as in Step 2 of Example 23, the title compound (2.27 g) was obtained from 2,N-dimethoxy-N-methylbenzamide (3.55 g) obtained in Step 1.


¹H-NMR(300MHz, δppm, CDCl₃) 7.59(1H,dd,J=7.5, 1.7Hz), 7.48-7.42(1H,m), 7.02-6.97(2H,m), 3.91(3H,s), 2.77-2.68(1H,m), 1.25-1.20(2H,m), 1.01-0.95(1H,m).

MS(ESI,m/z) 177(M+H)⁺.

Step 3

(R)-(Cyclopropyl)(2-methoxyphenyl)methanol

[0197]




[0198] In the same manner as in Step 3 of Example 23, the title compound (3.55 g) was obtained from cyclopropyl 2-methoxyphenyl ketone (3.52 g) obtained in Step 2. ¹H-NMR(300MHz, δppm, CDCl₃) 7.40(1H,dd,J=1.6, 7.5Hz), 7.26(1H,dt,J=1.7, 7.8Hz), 7.00-6.89(2H,m), 4.20(1H,br d,J=8.4Hz), 3.87(3H,s) 2.83(1H,br s), 1.42-1.30(1H,m), 0.70-0.61(1H,m), 0.57-0.44(2H,m), 0.37-0.27(1H,m).

MS(ESI,m/z) 161(M+H-H₂O)⁺.

Step 4


(R)-2-[(1R)-(Cyclopropyl)(2-methoxyphenyl)methoxymethyl]oxirane

[0199]

[0200] In the same manner as in Step 2 of Example 1, the title compound (840 mg) was obtained from (R)-(cyclopropyl)(2-methoxyphenyl)methanol (1.78 g) obtained in Step 3.

Step 5

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1R)-(cyclopropyl)(2-methoxyphenyl)methoxy]propan-2-ol

[0201]

[0202] In the same manner as in Step 13 of Example 1, the title compound (970 mg) was obtained from (R)-2-[(1R)-(cyclopropyl)(2-methoxyphenyl)methoxymethyl]oxirane (586 mg) obtained in Step 4.

¹H-NMR (300MHz, δ ppm, CDCl₃) 7.83-7.70(3H,m), 7.61(1H,s), 7.50-7.37(3H,m), 7.35-7.20(2H,m), 7.00-6.82(2H,m), 4.28(1H,d,J=7.8Hz), 3.83-3.70(4H,m), 3.42-3.30(2H,m), 2.90-2.60(4H,m), 1.25-1.10(7H,m), 0.60-0.50(1H,m), 0.45-0.30(3H,m).

MS(APCI,m/z) 434(M+H)⁺.

Example 25 - Example 56

[0203] In the same manner as in Examples 1 - 24, the compounds of Example 25 - Example 56 were obtained. They are shown in Table 3 - Table 7.

Table 3

Ex. 25		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 9.02 (1H,brs), 8.57(1H,brs), 7.90-7.85(3H,m), 7.76 (1H,s), 7.55-7.45(2H,m), 7.40-7.15(6H,m), 5.6 2(1H,d,J=6.3Hz), 4.51(1H,q,J=6.3Hz), 4.03(1 H,brs), 3.40-3.10(5H,m), 2.95-2.80(1H,m), 1.3 7(1H,d,J=6.3Hz), 1.26(6H,s). MS(APCI,m/z) 378(M+H-HCl)*.
Ex. 26		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 9.02 (1H,brs), 8.59(1H,brs), 7.91-7.88(3H,m), 7.77 (1H,s), 7.55-7.48(2H,m), 7.40-7.25(6H,m), 5.6 2-5.57(1H,m), 4.34-4.31(1H,m), 4.01(1H,brs), 3.35-3.16(5H,m), 3.05-2.80(1H,m), 1.80-1.65(1 H,m), 1.60-1.45(1H,m), 1.40-1.15(8H,m), 0.87 (3H,t,J=7.2Hz). MS(APCI,m/z) 406(M+H-HCl)*.
Ex. 27		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 9.04 (1H,brs), 8.60(1H,brs), 7.90-7.85(3H,m), 7.77 (1H,s), 7.55-7.48(2H,m), 7.40-7.25(6H,m), 5.6 2-5.57(1H,m), 4.35-4.25(1H,m), 4.02(1H,brs), 3.35-3.10(5H,m), 3.05-2.80(1H,m), 1.80-1.65(1 H,m), 1.60-1.50(1H,m), 1.40-1.10(10H,m), 0.8 3(3H,t,J=7.2Hz). MS(APCI,m/z) 420(M+H-HCl)*.
Ex. 28		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 8.98 (1H,brs), 8.58(1H,brs), 7.95-7.85(3H,m), 7.77 (1H,s), 7.55-7.45(2H,m), 7.40-7.25(6H,m), 5.8 5-5.70(1H,m), 5.63-5.58(1H,m), 4.41(1H,t,J=6. 6Hz), 4.01(1H,brs), 3.40-3.10(8H,m), 3.00-2.8 5(1H,m), 2.45-2.30(1H,m), 1.27(3H,s), 1.26(3 H,s). MS(APCI,m/z) 404(M+H-HCl)*.
Ex. 29		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 8.80 (1H,brs), 8.50(1H,brs), 7.95-7.85(3H,m), 7.76 (1H,s), 7.55-7.45(2H,m), 7.40-7.15(11H,m), 5. 65-5.59(1H,m), 4.40-4.30(1H,m), 4.00(1H,brs), 3.35-3.10(7H,m), 2.75-2.60(1H,m), 2.10-2.00 (1H,m), 2.00-1.80(1H,m), 1.27(3H,s), 1.26(3 H,s). MS(APCI,m/z) 468(M+H-HCl)*.
Ex. 30		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 9.00 (1H,brs), 8.58(1H,brs), 7.95-7.85(3H,m), 7.76 (1H,s), 7.55-7.45(2H,m), 7.40-7.25(6H,m), 6.0 0-5.85(1H,m), 5.67(1H,d,J=3.9Hz), 5.35-5.18(2 H,m), 4.89(1H,d,J=6.6Hz), 4.06(1H,brs), 3.55- 3.25(5H,m), 3.05-2.90(1H,m), 1.27(6H,s). MS(APCI,m/z) 390(M+H-HCl)*.

Table 4







Ex. 31		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 8.84 (1H,brs), 8.54(1H,brs), 7.95-7.85(3H,m), 7.77 (1H, s), 7.55-7.45(2H,m), 7.40-7.25(6H,m), 5.64-5.59(1H,m), 4.60-4.50(1H,m), 4.02(1H,brs), 3.60-3.10(11H,m), 1.26(6H,s). MS(APCLm/z) 408(M+H-HCl)*.
Ex. 32		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 8.90 (1H,brs), 8.56(1H,brs), 7.95-7.88(3H,m), 7.77 (1H,s), 7.55-7.45(2H,m), 7.40-7.20(6H,m), 5.57-5.51(1H,m), 4.07-3.90(2H,m), 3.30-3.10(6H,m), 3.05-2.80(1H,m), 2.20-2.05(1H,m), 1.85-1.70(1H,m), 1.65-1.35(4H,m), 1.30-1.05(8H,m). MS(APCLm/z) 432(M+H-HCl)*.
Ex. 33		¹ H-NMR(300MHz, δ ppm,CDCl ₃) 7.83-7.73 (3H,m), 7.60(1H,s), 7.50-7.40(2H,m), 7.35-7.20(5H,m), 4.41-4.35(1H,m), 3.80-3.65(1H,m), 3.40-3.25(2H,m), 2.95-2.55(4H,m), 1.39(3H,d, J=6.5Hz), 1.25-1.10(6H,m). MS(APCLm/z) 412(M+H)*.
Ex. 34		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 8.93 (1H,brs), 8.55(1H,brs), 7.95-7.85(3H,m), 7.77 (1H,s), 7.60-7.25(5H,m), 5.70-5.65(1H,m), 4.87(1H,q,J=6.6Hz), 4.03(1H,brs), 3.45-3.10(7H,m), 3.05-2.90(1H,m), 1.38(3H,d,J=6.3Hz), 1.27(6H,s). MS(APCLm/z) 412(M+H-HCl)*.
Ex. 35		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 9.00 (1H,brs), 8.57(1H,brs), 7.95-7.85(3H,m), 7.77 (1H,s), 7.55-7.25(5H,m), 5.68-5.63(1H,m), 4.54(1H,q,J=6.6Hz), 4.02(1H,brs), 3.45-3.10(7H,m), 3.05-2.90(1H,m), 1.37(3H,d,J=6.3Hz), 1.27(6H,s). MS(APCLm/z) 412(M+H-HCl)*.
Ex. 36		¹ H-NMR(300MHz, δ ppm,CDCl ₃) 7.85-7.70 (3H,m), 7.61(1H,s), 7.50-7.40(2H,m), 7.35-7.15(5H,m), 3.85(1H,d,J=7.3Hz), 3.80-3.70(1H,m), 3.35-3.20(2H,m), 2.90-2.65(4H,m), 2.00-1.85(2H,m), 1.20-1.05(6H,m), 0.99-0.97(3H,m), 0.72-0.69(3H,m). MS(APCLm/z) 406(M+H)*.

Table 5







Ex. 37		¹ H-NMR(300MHz, δ ppm,DMSO- d_6) 8.90 (1H,brs), 8.50(1H,brs), 7.90-7.80(3H,m), 7.76 (1H,s), 7.55-7.45(2H,m), 7.40-7.35(1H,m), 7.2 5-7.10(4H,m), 5.61(1H,t,J=4.8Hz), 4.45(1H,qJ =6.2Hz), 3.97(1H,brs), 3.40-3.10(5H,m), 3.00-2.80(1H,m), 2.29(3H,s), 1.35(3H,d,J=6.4Hz), 1.26-1.25(6H,m). MS(APCI,m/z) 392(M+H-HCl)*.
Ex. 38		¹ H-NMR(300MHz, δ ppm,DMSO- d_6) 8.90 (1H,brs), 8.50(1H,brs), 7.95-7.85(3H,m), 7.76 (1H,s), 7.55-7.45(2H,m), 7.40-7.25(6H,m), 5.6 2-5.56(1H,m), 4.40-4.30(1H,m), 4.00(1H,brs), 3.40-3.10(5H,m), 3.05-2.80(1H,m), 1.80-1.60(2 H,m), 1.45-1.20(7H,m), 0.90(6H,d,J=6.4Hz). MS(APCI,m/z) 420(M+H-HCl)*.
Ex. 39		¹ H-NMR(300MHz, δ ppm,CDCl ₃) 7.85-7.70 (3H,m), 7.76(1H,s), 7.50-7.40(2H,m), 7.30-7.1 0(3H,m), 6.90-6.80(2H,m), 4.40-4.30(1H,m), 3.80-3.65(4H,m), 3.40-3.25(2H,m), 2.90-2.60(4 H,m), 1.41(3H,d,J=6.5Hz), 1.10-1.05(6H,m). MS(APCI,m/z) 408(M+H)*.
Ex. 40		¹ H-NMR(300MHz, δ ppm,DMSO- d_6) 8.99 (1H,brs), 8.57(1H,brs), 7.95-7.85(3H,m), 7.76 (1H,s), 7.65-7.45(5H,m), 7.40-7.35(1H,m), 5.7 5-5.65(1H,m), 4.84(1H,qJ=6.2Hz), 4.03(1H,br s), 3.40-3.10(5H,m), 3.05-2.85(1H,m), 1.37(3 H,d,J=6.4Hz), 1.27(6H,s). MS(APCI,m/z) 446(M+H-HCl)*.
Ex. 41		¹ H-NMR(300MHz, δ ppm,DMSO- d_6) 8.90 (1H,brs), 8.55(1H,brs), 7.95-7.85(3H,m), 7.76 (1H,s), 7.55-7.45(4H,m), 7.40-7.30(2H,m), 5.7 0-5.60(1H,m), 5.29(1H,qJ=6.7Hz), 4.05-3.90(1 H,m), 3.40-2.80(6H,m), 1.25(3H,d,J=5.5Hz), 1.26(3H,s), 1.24(3H,s). MS(APCI,m/z) 446(M+H-HCl)*.
Ex. 42		¹ H-NMR(300MHz, δ ppm,DMSO- d_6) 8.96 (1H,brs), 8.55(1H,brs), 7.95-7.85(3H,m), 7.76 (1H,s), 7.65-7.60(2H,m), 7.55-7.45(2H,m), 7.4 0-7.30(2H,m), 5.69-5.65(1H,m), 4.55(1H,qJ= 6.4Hz), 4.00(1H,brs), 3.45-3.10(5H,m), 3.05-2. 85(1H,m), 1.37(3H,d,J=6.4Hz), 1.26(6H,s). MS(APCI,m/z) 446(M+H-HCl)*.

Table 6








Ex. 43		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 8.93 (1H,brs), 8.59(1H,brs), 7.95-7.85(3H,m), 7.77-7.75(1H,m), 7.60-7.30(6H,m), 5.64-5.61(1H,m), 5.07(1H,tJ=8.1Hz), 4.10-3.95(1H,m), 3.40-3.00(7H,m), 2.90-2.80(1H,m), 2.15-1.80(2H,m), 1.27-1.25(6H,m), 0.95-0.80(3H,m). MS(APCI,m/z) 460(M+H-HCl)*.
Ex. 44		¹ H-NMR(300MHz, δ ppm,DMSO-d ₆) 8.95 (1H,brs), 8.60(1H,brs), 7.95-7.85(3H,m), 7.75 (1H,s), 7.55-7.45(4H,m), 7.40-7.30(2H,m), 5.6 2-5.58(1H,m), 4.74-4.70(1H,m), 4.10-3.90(1H,m), 3.40-3.00(6H,m), 2.90-2.80(1H,m), 1.30-1.10(9H,m), 0.67-0.60(3H,m). MS(APCI,m/z) 474(M+H-HCl)*.
Ex. 45		¹ H-NMR(300MHz, δ ppm,CDCl ₃) 8.20-8.15 (2H,m), 7.85-7.70(3H,m), 7.60(1H,s), 7.50-7.4 0(4H,m), 7.35-7.25(1H,m), 4.60-4.50(1H,m), 3.80-3.70(1H,m), 3.45-3.30(2H,m), 2.90-2.75(3 H,m), 2.70-2.60(1H,m), 1.43-1.41(3H,m), 1.12 -1.09(6H,m). MS(APCI,m/z) 423(M+H)*.
Ex. 46		¹ H-NMR(300MHz, δ ppm,CDCl ₃) 8.20-8.05 (2H,m), 7.80-7.70(3H,m), 7.65-7.55(2H,m), 7. 50-7.35(3H,m), 7.35-7.25(1H,m), 4.60-4.45(1 H,m), 3.80-3.70(1H,m), 3.50-3.30(2H,m), 2.90 -2.60(4H,m), 1.45-1.42(3H,m), 1.12-1.09(6H, m). MS(APCI,m/z) 423(M+H)*.
Ex. 47		7.90-7.70(3H,m), 7.66(1H,dJ=6.8Hz), 7.50-7.2 0(6H,m), 4.65(1H,brs), 4.35(1H,dJ=9.4Hz), 3. 70-3.50(1H,m), 3.40-3.00(2H,m), 2.80-2.40(4 H,m), 1.80-0.80(8H,m), 0.80-0.10(4H,m). MS(APCI,m/z) 472(M+H)*.
Ex. 48		¹ H-NMR(300MHz, δ ppm, DMSO-d ₆) 7.90 -7.70(3H,m), 7.67(1H,s), 7.50-7.30(3H,m), 7.2 5-7.00(4H,m), 4.64(1H,brs), 3.70-3.50(2H,m), 3.40-3.10(2H,m), 2.80-2.50(4H,m), 2.30(3H,s), 1.50-0.80(8H,m), 0.60-0.10(4H,m). MS(APCI,m/z) 418(M+H)*.
Ex. 49		¹ H-NMR(300MHz, δ ppm, DMSO-d ₆) 7.90 -7.30(11H,m), 4.72(1H,brs), 4.10-4.00(1H,m), 3.70-3.55(1H,m), 3.45-3.10(2H,m), 2.80-2.50(4 H,m), 1.50-0.80(8H,m), 0.70-0.20(4H,m). MS(APCI,m/z) 429(M+H)*.

Table 7

Ex. 50		¹ H-NMR(300MHz, δ ppm, CDCl ₃) 7.85-7.70 (3H,m), 7.61(1H,s), 7.50-7.40(2H,m), 7.35-7.10(4H,m), 4.40-4.30(1H,m), 3.80-3.70(1H,m), 3.40-3.25(2H,m), 2.90-2.60(4H,m), 1.41-1.37(3H,m), 1.12-1.08(6H,m). MS(APCL,m/z) 446(M+H) ⁺ .
Ex. 51		¹ H-NMR(300MHz, δ ppm, CDCl ₃) 7.85-7.70 (3H,m), 7.60(1H,s), 7.50-7.40(2H,m), 7.35-7.15(2H,m), 7.10-7.00(3H,m), 4.15-4.05(1H,m), 3.80-3.70(1H,m), 3.45-3.25(2H,m), 2.90-2.60(4H,m), 2.34(3H,s), 1.90-1.60(2H,m), 1.15-1.05(6H,m), 0.85(3H,t, J=7.4Hz). MS(APCL,m/z) 406(M+H) ⁺ .
Ex. 52		¹ H-NMR(300MHz, δ ppm, CDCl ₃) 7.85-7.70 (3H,m), 7.60(1H,s), 7.50-7.40(2H,m), 7.35-7.05(5H,m), 4.50-4.40(1H,m), 3.80-3.70(1H,m), 3.40-3.20(2H,m), 2.90-2.60(4H,m), 2.31(3H,s), 1.80-1.60(2H,m), 1.15-1.05(6H,m), 0.92(3H,t, J=7.2Hz). MS(APCL,m/z) 406(M+H) ⁺ .
Ex. 53		¹ H-NMR(300MHz, δ ppm, DMSO-d ₆) 7.90-7.64(4H,m), 7.50-7.30(3H,m), 7.25-7.05(4H,m), 4.61(1H,brs), 4.50-4.40(1H,m), 3.75-3.60(1H,m), 3.17(2H,d, J=5.7Hz), 2.85-2.45(4H,m), 2.31(3H,s), 1.95-1.50(6H,m), 1.45-0.90(7H,m). MS(APCL,m/z) 432(M+H) ⁺ .
Ex. 54		¹ H-NMR(300MHz, δ ppm, DMSO-d ₆) 7.90-7.70(3H,m), 7.68(1H,s), 7.50-7.10(5H,m), 7.00-6.80(2H,m), 4.70-4.50(2H,m), 3.78(3H,s), 3.70-3.50(1H,m), 2.85-2.50(4H,m), 1.95-1.50(6H,m), 1.45-0.90(7H,m). MS(APCL,m/z) 448(M+H) ⁺ .
Ex. 55		¹ H-NMR(300MHz, δ ppm, DMSO-d ₆) 7.90-7.70(3H,m), 7.67(1H,s), 7.50-7.25(4H,m), 7.20-7.10(3H,m), 4.65(1H,brs), 4.06(1H,d, J=7.4Hz), 3.70-3.50(1H,m), 3.30-3.10(2H,m), 2.80-2.50(4H,m), 2.31(3H,s), 1.50-0.80(8H,m), 0.60-0.10(4H,m). MS(APCL,m/z) 418(M+H) ⁺ .
Ex. 56		¹ H-NMR(300MHz, δ ppm, CDCl ₃) 7.85-7.70 (3H,m), 7.61(1H,s), 7.50-7.40(3H,m), 7.35-7.20(2H,m), 7.00-6.80(2H,m), 4.27(1H,d, J=7.5Hz), 3.85-3.75(4H,m), 3.50-3.30(2H,m), 2.85-2.65(4H,m), 1.25-1.00(7H,m), 0.60-0.50(1H,m), 0.45-0.30(3H,m). MS(APCL,m/z) 434(M+H) ⁺ .


Example 57

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(2-hydroxymethylphenyl)methoxy]propan-2-ol

Step 1

2-Bromo-1-(tert-butyldimethylsilyloxymethyl)benzene

[0204]




[0205] 2-Bromobenzyl alcohol (25.0 g) was dissolved in N,N-dimethylformamide (150 ml) and imidazole (20.0 g) and tert-butyldimethylchlorosilane (22.2 g) were added. The reaction mixture was stirred at room temperature for 2 hr. The reaction mixture was poured into 5% aqueous sodium hydrogencarbonate solution and extracted with ethyl acetate. The organic layer was washed successively with 5% aqueous sodium hydrogencarbonate solution, water and saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (40.9 g).

Step 2

(Cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)-phenyl]methanol

[0206]




[0207] Magnesium (2.20 g) was suspended in tetrahydrofuran (4 ml) and iodine (2 mg) was added. Thereto was added dropwise a solution of 2-bromo-1-(tert-butyldimethylsilyloxymethyl)benzene (24.0 g) obtained in Step 1 in tetrahydrofuran (10 ml) over 30 min and the reaction mixture was heated under reflux for 1 hr. Thereto was added tetrahydrofuran to give 1M 2-(tert-butyldimethylsilyloxymethyl)phenylmagnesium bromide-tetrahydrofuran solution. Then cyclopropanecarboxaldehyde (2.80 g) was dissolved in tetrahydrofuran (120 ml), and 1M 2-(tert-butyldimethylsilyloxymethyl)phenylmagnesium bromide-tetrahydrofuran solution (80 ml) was added dropwise under ice-cooling over 50 min. The reaction mixture was stirred at room temperature for 12 hr. To the reaction mixture was added saturated aqueous ammonium chloride solution (8 ml) under ice-cooling and the reaction mixture was stirred at room temperature for 30 min. The reaction mixture was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (n-hexane:ethyl acetate=95:5) to give the title compound (8.20 g).

¹H-NMR(300MHz, δppm, DMSO-d₆) 7.53-7.34(2H,m), 7.25-7.20(2H,m), 4.91(1H,d,J=4.6Hz), 4.81(2H,s), 4.31(1H,dd, J=4.7, 6.5Hz), 1.20-1.10(1H,m), 0.91(9H,s), 0.47-0.23(4H,m), 0.10(3H,s), 0.08(3H,s).
MS(APCI,m/z) 275(M+H-H₂O)⁺.

Step 3

(R)-2-[(Cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)-phenyl]methoxymethyl]oxirane

[0208]




[0209] In the same manner as in Step 2 of Example 1, the title compound (1.04 g) was obtained from (cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)phenyl]methanol (2.11 g) obtained in Step 2.

Step 4

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)-phenyl]methoxy]propan-2-ol

[0210]




[0211] In the same manner as in Step 13 of Example 1, the title compound (345 mg) was obtained from (R)-2-[(cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)phenyl]methoxymethyl]oxirane (546 mg) obtained in Step 3.

Step 5

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)[2-(hydroxymethyl)phenyl]methoxy]propan-2-ol

[0212]



[0213] (2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)-phenyl]methoxypropan-2-ol (365 mg) obtained in Step 4 was dissolved in tetrahydrofuran (4 ml) and tetrabutylammonium fluoride-1M tetrahydrofuran solution (0.73 ml) was added under ice-cooling. The mixture was stirred at room

temperature for 4 hr. To the reaction mixture was added saturated aqueous ammonium chloride solution (1 ml). The reaction mixture was poured into water and extracted with ethyl acetate. The organic layer was washed successively with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (ethyl acetate:methanol=98:2) to give the title compound (221 mg).

$^1\text{H-NMR}$ (300MHz, δ ppm, DMSO- d_6) 7.90-7.70(3H,m), 7.67(1H,s), 7.50-7.20(7H,m), 5.13(1H,brs), 4.90-4.40(3H,m), 4.20-4.05(1H,m), 3.70-3.50(1H,m), 3.30-3.10(2H,m), 2.80-2.50(4H,m), 1.40-0.85(7H,m), 0.60-0.10(4H,m).

MS(APCI,m/z) 434(N+H) $^+$.


Example 58

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1R)-(cyclopropyl)[2-(hydroxymethyl)phenyl]methoxy]propan-2-ol hemifumarate

Step 1

2-tert-Butyldimethylsilyloxymethyl-N-methoxy-N-methylbenzamide

[0214]




[0215] To a solution of phthalide' (26.8 g) in methylene chloride (600 ml) were successively added N,O-dimethylhydroxylamine hydrochloride (58.5 g) and aluminum chloride (40.0 g) under ice-cooling, and triethylamine (139 ml) was added dropwise over 40 min. The mixture was stirred at room temperature for 12 hr. The reaction mixture was poured into diluted hydrochloric acid, and the organic layer was washed successively with water and saturated brine, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure to give 2-hydroxymethyl-N-methoxy-N-methylbenzamide. The obtained 2-hydroxymethyl-N-methoxy-N-methylbenzamide was dissolved in N,N-dimethylformamide (300 ml), and imidazole (9.53 g) and tert-butyldimethylchlorosilane (21.1 g) were added. The mixture was stirred at room temperature for 3.5 hr and the reaction mixture was poured into water and extracted with ethyl acetate. The organic layer was washed successively with water and saturated brine, dried over anhydrous sodium sulfate, and concentration under reduced pressure. The obtained residue was purified by silica gel column chromatography (ethyl acetate:hexane=1:3) to give the title compound (42.0 g).

Step 2

(Cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)-phenyl]methanone

[0216]



[0217] In the same manner as in Step 2 of Example 23, the title compound (18.4 g) was obtained from 2-tert-butylid-


dimethylsilyloxymethyl-N-methoxy-N-methylbenzamide (21.7 g) obtained in Step 1.

$^1\text{H-NMR}$ (300 MHz, δ ppm, CDCl_3) 7.88 (1H, dd, $J=1.1$, 7.7 Hz), 7.78 (1H, d, $J=7.3$ Hz), 7.53 (1H, dt, $J=1.2$, 7.6 Hz), 7.35 (1H, t, $J=7.2$ Hz), 4.97 (2H, s), 2.57-2.48 (1H, m), 1.24-1.20 (2H, m), 1.06-1.01 (2H, m), 0.95 (9H, s), 0.11 (6H, s).

Step 3

(R)-(Cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)-phenyl]methanol

[0218]



[0219] To a suspension of dichloro[(S)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl][(S)-1,1'-bis(p-methoxyphenyl)-2-isopropylethane-1,2-diamine]ruthenium (II) (111 mg) and potassium-tert-butoxide (44.9 mg) in isopropanol (100 ml) was added (cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)phenyl]methanone (5.81 g) obtained in Step 2, and the mixture was subjected to medium pressure hydrogenation (5.0 kgf/cm²) at room temperature for 36 hr. The reaction mixture was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (ethyl acetate:hexane=1:5) to give the title compound (5.82 g).


$^1\text{H-NMR}$ (300 MHz, δ ppm, DMSO-d_6) 7.49-7.46 (1H, m), 7.39-7.37 (1H, m), 7.25-7.22 (2H, m), 4.99 (1H, br s), 4.81 (2H, s), 4.32 (1H, d, $J=6.3$ Hz), 1.32-1.09 (1H, m), 0.91 (9H, s), 0.49-0.22 (4H, m), 0.10 (3H, s), 0.08 (3H, s).

MS (ESI, m/z) 275 ($\text{M}+\text{H}-\text{H}_2\text{O}$)⁺.

Step 4

(R)-2-[(1R)-(Cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)-phenyl]methoxymethyl]oxirane

[0220]




[0221] In the same manner as in Step 2 of Example 1, the title compound (2.52 g) was obtained from (R)-(cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)phenyl]methanol (2.92 g) obtained in Step 3.

Step 5

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1R)-(cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)-phenyl]methoxypropan-2-ol

[0222]




[0223] In the same manner as in Step 13 of Example 1, the title compound (1.79 g) was obtained from (R)-2-[(1R)-(cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)-phenyl]methoxymethyl]oxirane (1.22 g) obtained in Step 4.

Step 6

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1R)-(cyclopropyl)[2-(hydroxymethyl)phenyl]methoxy]propan-2-ol

[0224]




[0225] In the same manner as in Step 5 of Example 57, the title compound (710 mg) was obtained from (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1R)-(cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)phenyl]methoxy]propan-2-ol (1.70 g) obtained in Step 5.

Step 7

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1R)-(cyclopropyl)[2-(hydroxymethyl)phenyl]methoxy]propan-2-ol hemifumarate

[0226]



[0227] (2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1R)-(cyclopropyl)[2-(hydroxymethyl)phenyl]methoxy]propan-2-ol (700 mg) obtained in Step 6 was dissolved in methanol (15 ml), and fumaric acid (94 mg) was added. The mixture was stirred for 30 min. The reaction mixture was concentrated under reduced pressure, and ether was added to the residue. The precipitated solid was collected by filtration to give the title compound (720 mg). ¹H-NMR (300MHz, δ ppm, DMSO- d_6) 7.95-7.85(3H,m), 7.73(1H,s), 7.55-7.20(7H,m), 6.51(1H,s), 4.63(1H,d,J=13.2Hz), 4.56(1H,d,J=13.2Hz), 4.13(1H,d,J=7.5Hz), 3.90-3.80(1H,brs), 3.40-3.15(2H,m), 3.05-2.95(3H,m), 2.80-2.65(1H,m), 1.20-1.00(7H,m), 0.60-0.20(4H,m). MS(ESI,m/z) 434(M+H-1/2C₄R₄O₄)⁺.


Example 59

(2R)-2-[1,1-Dimethyl-2-[naphthalen-2-yl)ethylamino]-3-[(1S)-(cyclopropyl)[2-(hydroxymethyl)phenyl]methoxy]propan-2-ol hemifumarate

Step 1

(S)-(Cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)-phenyl]methanol

[0228]




[0229] To a suspension of dichloro[(R)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl][(R)-1,1'-bis(pmethoxyphenyl)-2-isopropylethane-1,2-diamine]ruthenium (II) (111 mg) and potassium-tert-butoxide (44.9 mg) in isopropanol (100 ml) was added (cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)phenyl]methanone (5.81 g) obtained in Step 2 of Example 58, and the mixture was subjected to medium pressure hydrogenation (5.0 kgf/cm²) at room temperature for 36 hr. The reaction mixture was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (ethyl acetate:hexane=1:5) to give the title compound (5.50 g). ¹H-NMR (300MHz, δ ppm, DMSO- d_6) 7.49-7.46(1H,m), 7.40-7.37(1H,m), 7.22-7.20(2H,m), 4.99(1H,d,J=4.2Hz), 4.81(2H,s), 4.31(1H,dd,J=4.5, 6.6Hz), 1.20-1.09(1H,m), 0.91(9H,s), 0.47-0.21(4H,m), 0.10(3H,s), 0.08(3H,s). MS(ESI,m/z) 275(M+H-H₂O)⁺.

Step 2

(R)-2-[(1S)-(Cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)-phenyl]methoxymethyl]oxirane

[0230]




[0231] In the same manner as in Step 2 of Example 1, the title compound (2.62 g) was obtained from (S)-(cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)phenyl]methanol (2.92 g) obtained in Step 1.

Step 3

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1S)-(cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)-phenyl]methoxy]propan-2-ol

[0232]




[0233] In the same manner as in Step 13 of Example 1, the title compound (1.80 g) was obtained from (R)-2-[(1S)-(cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)-phenyl]methoxymethyl]oxirane (1.22 g) obtained in Step 2.

Step 4

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1S)-(cyclopropyl)[2-(hydroxymethyl)phenyl]methoxy]propan-2-ol

[0234]




[0235] In the same manner as in Step 5 of Example 57, the title compound (1.26 g) was obtained from (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1S)-(cyclopropyl)[2-(tert-butyldimethylsilyloxymethyl)phenyl]methoxy]propan-2-ol (1.70 g) obtained in Step 3.

Step 5

(2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1S)-(cyclopropyl)[2-(hydroxymethyl)phenyl]methoxy]propan-2-ol hemifumarate

[0236]



EP 1 308 436 A1

[0237] In the same manner as in Step 7 of Example 58, the title compound (1.33 g) was obtained from (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(1S)-(cyclopropyl)[2-(hydroxymethyl)phenyl]methoxy]propan-2-ol (1.25 g) obtained in Step 4.

¹H-NMR(300MHz, δppm, DMSO-d₆) 7.90-7.80(3H,m), 7.71(1H,s), 7.50-7.20(7H,m), 6.48(1H,s), 4.63(1H,d,J=13.2Hz), 4.56(1H,d,J=13.2Hz), 4.12(1H,d,J=7.8Hz), 3.80-3.70(1H,brs), 3.35-3.15(2H,m), 2.95-2.85(3H,m), 2.80-2.65(1H,m), 1.20-1.00(7H,m), 0.60-0.20(4H,m).






MS(ESI,m/z) 434(M+H-1/2C₄H₄O₄)⁺.

Example 60 - Example 160

[0238] In the same manner as in Examples 1 - 59, the compounds of Example 60 - Example 160 were obtained. They are shown in

Table 8 - Table 23.

Table 8

Ex. 60	 1/2 HO₂C-CH=CH-CO₂H	¹ H-NMR (300MHz, δppm, DMSO-d ₆) 7.95-7.75 (3H, m), 7.71 (1H, s), 7.55-7.45 (2H, m), 7.40- 7.20 (3H, m), 7.05-6.90 (2H, m), 6.48 (1H, s), 4.80 (1H, q, J=6.4Hz), 3.90-3.70 (4H, m), 3.40- 3.20 (2H, m), 3.00-2.80 (3H, m), 2.75-2.60 (1H, m), 1.28 (3H, d, J=6.4Hz), 1.15- 1.85 (6H, m). MS (APCI, m/z) 408 (M+H- 1/2 C ₄ H ₄ O ₄) ⁺ .
Ex. 61	 1/2 HO₂C-CH=CH-CO₂H	¹ H-NMR (300MHz, δppm, CDCl ₃) 7.95-7.75 (3H, m), 7.71 (1H, s), 7.60-7.15 (5H, m), 7.05- 6.85 (2H, m), 6.48 (1H, s), 4.29 (1H, d, J=7.2Hz), 3.85- 3.70 (4H, m), 3.40-3.10 (2H, m), 3.00-2.80 (4H, m), 2.75- 2.60 (1H, m), 1.20-1.00 (7H, m), 0.60-0.20 (4H, m). MS (APCI, m/z) 434 (M+H- 1/2 C ₄ H ₄ O ₄) ⁺ .
Ex. 62	 1/2 HO₂C-CH=CH-CO₂H	¹ H-NMR (300MHz, δppm, DMSO-d ₆) 7.90-7.80 (3H, m), 7.71 (1H, s), 7.60-7.25 (4H, m), 7.20- 7.10 (3H, m), 6.49 (1H, s), 4.09 (1H, d, J=7.5Hz), 3.80 (1H, brs), 3.40-3.10 (2H, m), 3.00-2.80 (3H, m), 2.75- 2.60 (1H, m), 2.32 (3H, s), 1.25- 1.00 (7H, m), 0.60-0.10 (4H, m). MS (APCI, m/z) 418 (M+H- 1/2 C ₄ H ₄ O ₄) ⁺ .
Ex. 63	 HCl	¹ H-NMR (300MHz, δppm, DMSO-d ₆) 9.00 (1H, bs), 8.57 (1H, bs), 7.95- 7.85 (3H, m), 7.62 (1H, s), 7.65- 7.35 (6H, m), 5.75-5.65 (1H, m), 4.95-4.85 (1H, m), 4.10- 4.00 (1H, m), 3.50-3.30 (2H, m), 3.30-3.10 (3H, m), 3.05- 2.90 (1H, m), 1.39 (3H, d, J=6.4Hz), 1.27 (6H, s). MS (APCI, m/z) 446 (M+H-HCl) ⁺ .
Ex. 64	 HCl	¹ H-NMR (300MHz, δppm, DMSO-d ₆) 8.98 (1H, bs), 8.57 (1H, bs), 7.95- 7.85 (3H, m), 7.59 (1H, s), 7.60- 7.45 (4H, m), 7.45-7.35 (2H, m), 5.80-5.65 (1H, m), 4.85- 4.75 (1H, m), 4.10-4.00 (1H, m), 3.50-3.30 (2H, m), 3.25- 3.10 (3H, m), 3.05-2.90 (1H, m), 1.40-1.35 (3H, m), 1.27 (6H, s). MS (APCI, m/z) 446 (M+H-HCl) ⁺ .


Ex. 65	 · 2HCl	¹ H-NMR (300MHz, δ ppm, DMSO-d ₆) 9.33 (1H, bs), 8.85-8.65 (2H, m), 8.50-8.40 (1H, m), 8.05- 7.70 (6H, m), 7.60-7.35 (3H, m), 5.05-4.90 (1H, m), 4.20- 4.10 (1H, m), 3.70-3.55 (2H, m), 3.50-3.40 (1H, m), 3.30- 3.15 (3H, m), 3.10-2.95 (1H, m), 1.53 (3H, d, J=6.6Hz), 1.29 (6H, s). MS (APCI, m/z) 379 (M+H-2HCl) ⁺ .
-----------	---	--

Table 9

Ex. 66		¹ H-NMR (300MHz, δ ppm, DMSO-d ₆) 8.94 (1H, bs), 8.56 (1H, bs), 7.95-7.85 (3H, m), 7.77 (1H, s), 7.55-7.45 (2H, m), 7.40-7.20 (3H, m), 7.05-6.90 (2H, m), 5.61 (1H, d, J=4.8Hz), 4.84 (1H, dd, J=9.5, 6.3Hz), 4.10-4.00 (1H, m), 3.80 (3H, s), 3.45-3.10 (5H, m), 3.00-2.85 (1H, m), 1.32 (3H, d, J=6.3Hz), 1.26 (6H, s). MS (APCI, m/z) 408 (M+H-HCl) ⁺ .
Ex. 67		¹ H-NMR (300MHz, δ ppm, DMSO-d ₆) 8.87 (1H, bs), 8.53 (1H, bs), 7.95-7.85 (3H, m), 7.77 (1H, s), 7.55-7.45 (2H, m), 7.45-7.35 (2H, m), 7.30-7.20 (1H, m), 7.05-6.90 (2H, m), 5.64 (1H, d, J=4.8Hz), 4.84 (1H, dd, J=9.6, 6.3Hz), 4.05-3.95 (1H, m), 3.80 (3H, s), 3.40-3.30 (2H, m), 3.25-3.10 (3H, m), 3.10-2.95 (1H, m), 1.32 (3H, d, J=6.6Hz), 1.27 (6H, s). MS (APCI, m/z) 408 (M+H-HCl) ⁺ .
Ex. 68		¹ H-NMR (300MHz, δ ppm, DMSO-d ₆) 7.90-7.70 (3H, m), 7.67 (1H, s), 7.50-7.25 (4H, m), 7.20-7.10 (3H, m), 4.65 (1H, brs), 4.06 (1H, d, J=7.4Hz), 3.70-3.50 (1H, m), 3.30-3.10 (2H, m), 2.80-2.50 (4H, m), 2.31 (3H, s), 1.50-0.80 (8H, m), 0.60-0.10 (4H, m). MS (APCI, m/z) 418 (M+H) ⁺ .
Ex. 69		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.90-7.20 (11H, m), 4.82 (1H, d, J=7.3Hz), 3.88 (3H, s), 3.95-3.70 (1H, m), 3.55-3.20 (2H, m), 2.90-2.60 (4H, m), 1.30-1.00 (7H, m), 0.70-0.30 (4H, m). MS (ESI, m/z) 462 (M+H) ⁺ .
Ex. 70		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 8.10-8.00 (2H, m), 7.85-7.70 (3H, m), 7.60 (1H, s), 7.50-7.20 (5H, m), 4.55-4.45 (1H, m), 4.37 (2H, dd, J=10.7, 5.9Hz), 3.85-3.70 (1H, m), 3.40-3.25 (2H, m), 2.90-2.55 (4H, m), 1.45-1.35 (6H, m), 1.10-1.00 (6H, m). MS (APCI, m/z) 450 (M+H) ⁺ .
Ex. 71		¹ H-NMR (300MHz, δ ppm, DMSO-d ₆) 7.40-7.00 (6H, m), 6.83 (2H, d, J=8.5Hz), 4.08 (1H, d, J=7.5Hz), 3.80-3.60 (4H, m), 3.40-3.10 (2H, m), 2.90-2.50 (4H, m), 2.32 (3H, s), 1.30-0.80 (7H, m), 0.60-


		0.15 (4H, m) . MS (ESI, m/z) 398 (M+H) ⁺ .
Ex. 72		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 8.05-8.00 (2H, m), 7.85-7.70 (3H, m), 7.61 (1H, s), 7.50-7.25 (5H, m), 3.91 (3H, s), 3.90-3.70 (2H, m), 3.45-3.30 (2H, m), 2.90-2.60 (4H, m), 1.15-1.00 (7H, m), 0.65-0.55 (1H, m), 0.50-0.35 (2H, m), 0.30-0.20 (1H, m) . MS (ESI, m/z) 462 (M+H) ⁺ .

Table 10

Ex. 73		¹ H-NMR (400MHz, δ ppm, CDCl ₃) 7.85-7.70 (3H, m), 7.61 (1H, s), 7.50-7.20 (7H, m), 4.67 (2H, d, J=4.4Hz), 3.80-3.65 (2H, m), 3.45-3.30 (2H, m), 2.90-2.60 (4H, m), 1.20-1.00 (7H, m), 0.65-0.60 (1H, m), 0.50-0.40 (2H, m), 0.30-0.20 (1H, m). MS (ESI, m/z) 434 (M+H) ⁺ .
Ex. 74		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 8.00-7.90 (2H, m), 7.85-7.70 (3H, m), 7.61 (1H, s), 7.55-7.25 (5H, m), 3.91 (3H, s), 3.85-3.70 (2H, m), 3.50-3.30 (2H, m), 2.90-2.60 (4H, m), 1.20-1.00 (7H, m), 0.65-0.55 (1H, m), 0.50-0.40 (2H, m), 0.30-0.20 (1H, m). MS (ESI, m/z) 462 (M+H) ⁺ .
Ex. 75		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.85-7.70 (3H, m), 7.61 (1H, s), 7.50-7.15 (7H, m), 4.69 (2H, d, J=3.5Hz), 3.80-3.60 (2H, m), 3.45-3.35 (2H, m), 2.90-2.70 (4H, m), 1.20-1.00 (7H, m), 0.70-0.55 (1H, m), 0.50-0.40 (2H, m), 0.30-0.20 (1H, m). MS (ESI, m/z) 434 (M+H) ⁺ .
Ex. 76		¹ H-NMR (300MHz, δ ppm, DMSO-d ₆) 8.00-7.80 (5H, m), 7.79 (1H, s), 7.60-7.30 (5H, m), 3.90-2.60 (8H, m), 1.20-1.00 (7H, m), 0.60-0.20 (4H, m). MS (ESI, m/z) 448 (M+H) ⁺ .
Ex. 77		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.80-7.70 (3H, m), 7.61 (1H, s), 7.50-7.20 (5H, m), 6.80-6.70 (1H, m), 4.22 (1H, d, J=7.8Hz), 3.85-3.70 (4H, m), 3.50-3.30 (2H, m), 3.00-2.60 (4H, m), 1.20-1.00 (7H, m), 0.60-0.50 (1H, m), 0.50-0.30 (3H, m). MS (ESI, m/z) 513 (M+H) ⁺ .
Ex. 78		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.83-7.74 (3H, m), 7.61 (1H, brs), 7.48-7.41 (4H, m), 7.32-7.19 (3H, m), 7.02-6.99 (1H, m), 4.41-4.39 (1H, m), 3.80-3.70 (1H, m), 3.38-3.34 (2H, m), 2.87-2.81 (3H, m), 2.70-2.60 (1H, m), 2.15 (3H, s), 1.43-1.37 (3H, m), 1.12-1.09 (6H, m). MS (ESI, m/z) 435 (M+H) ⁺ .







Ex. 79		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.80-7.74 (3H, m), 7.65-7.60 (1H, m), 7.46-7.43 (2H, m), 7.30-7.19 (2H, m), 6.66-6.63 (3H, m), 4.37-4.35 (1H, m), 3.85-3.75 (1H, m), 3.40-3.34 (2H, m), 2.95 (6H, s), 2.98-2.60 (4H, m), 1.44-1.40 (3H, m), 1.12-1.08 (6H, m). MS (ESI, m/z) 421. (M+H) $^+$.
-----------	---	--

Table 11

Ex. 80		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.90-7.70 (5H, m), 7.60 (1H, s), 7.50-7.20 (5H, m), 6.30 (1H, bs), 5.70 (1H, bs), 3.85-3.70 (2H, m), 3.50-3.30 (2H, m), 2.90-2.60 (4H, m), 1.20-1.00 (7H, m), 0.70-0.60 (1H, m), 0.55-0.40 (2H, m), 0.35-0.25 (1H, m). MS (ESI, m/z) 447 (M+H) $^+$.
Ex. 81		$^1\text{H-NMR}$ (300MHz, δ ppm, $\text{DMSO}-d_6$) 8.05-7.30 (10H, m), 4.71 (1H, brs), 3.90-3.80 (1H, m), 3.75-3.50 (1H, m), 3.45-3.10 (2H, m), 2.90-2.50 (4H, m), 1.10-0.80 (7H, m), 0.70-0.20 (4H, m). MS (ESI, m/z) 4830 (M+H) $^+$.
Ex. 82		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.79-7.74 (3H, m), 7.62 (1H, brs), 7.46-7.43 (3H, m), 7.33-7.11 (4H, m), 4.49-4.43 (1H, m), 3.80-3.70 (1H, m), 3.48-3.33 (2H, m), 2.86-2.69 (4H, m), 2.63 (3H, s), 2.62 (3H, s), 1.20-1.09 (7H, m), 0.60-0.20 (4H, m). MS (ESI, m/z) 447 (M+H) $^+$.
Ex. 83	 · 1/2 $\text{HO}_2\text{C}-\text{CH}=\text{CH}-\text{CO}_2\text{H}$	$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.80-7.76 (3H, m), 7.64 (1H, s), 7.46-7.44 (2H, m), 7.30-7.17 (4H, m), 6.99-6.92 (2H, m), 6.85 (1H, s), 4.10-3.95 (2H, m), 3.50-3.30 (2H, m), 3.18-2.80 (4H, m), 2.27 (3H, s), 2.26 (3H, s), 1.29-1.22 (6H, m), 1.20-1.00 (1H, m), 0.60-0.10 (4H, m). MS (ESI, m/z) 432 (M+H-1/2 $\text{C}_4\text{H}_4\text{O}_4$) $^+$.
Ex. 84		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.82-7.73 (3H, m), 7.60 (1H, s), 7.45-7.43 (2H, m), 7.32-7.28 (1H, m), 7.16 (1H, s), 7.03-6.96 (2H, m), 4.04 (1H, d, $J=7.5\text{Hz}$), 3.82-3.74 (1H, m), 3.40-3.32 (2H, m), 2.86-2.69 (4H, m), 2.31-2.28 (6H, m), 1.22-1.09 (7H, m), 0.59-0.23 (4H, m). MS (ESI, m/z) 432 (M+H) $^+$.



Ex. 85	 • 1/2 HO₂C-CH=CH-CO₂H	¹ H-NMR (300MHz, δppm, CDCl ₃) 7.80-7.74 (3H, m), 7.63 (1H, s), 7.45-7.43 (2H, m), 7.32-7.28 (1H, m), 7.12 (1H, s), 7.03-6.97 (2H, m), 6.85 (1H, s), 4.26-4.12 (1H, m), 4.01 (1H, d, J=7.8Hz), 3.62-3.28 (2H, m), 3.20-2.76 (4H, m), 2.28-2.24 (6H, m), 1.29-1.05 (7H, m), 0.60-0.48 (1H, m), 0.40-0.26 (2H, m), 0.22-0.14 (1H, m). MS (ESI, m/z) 432 (M+H-1/2C ₄ H ₄ O ₄) ⁺ .
Ex. 86		¹ H-NMR (300MHz, δppm, DMSO-d ₆) 8.19 (1H, s), 7.90-7.30 (12H, m), 7.20-7.10 (1H, m), 4.40-4.30 (1H, m), 3.90-3.80 (4H, m), 3.45-3.25 (2H, m), 3.00-2.85 (3H, m), 2.80-2.65 (1H, m), 1.25-1.00 (7H, m), 0.60-0.30 (4H, m). MS (ESI, m/z) 554 (M+H) ⁺ .

Table 12

Ex. 87		¹ H-NMR (300MHz, δ ppm, DMSO-d ₆) 8.10-8.00 (2H, m), 7.90-7.70 (3H, m), 7.50-7.14 (9H, m), 6.80-6.70 (1H, m), 4.36-4.34 (1H, m), 3.90-3.80 (1H, m), 3.20-2.60 (9H, m), 1.26-1.20 (9H, m). MS (ESI, m/z) 591 (M+H) ⁺ .
Ex. 88		¹ H-NMR (400MHz, δ ppm, CDCl ₃) 7.90-7.70 (4H, m), 7.68-7.60 (1H, m), 7.60-7.40 (3H, m), 7.30-7.15 (3H, m), 4.18-4.09 (2H, m), 3.91-3.90 (3H, m), 3.81-3.79 (3H, m), 3.60-3.40 (2H, m), 3.30-2.90 (4H, m), 1.35-1.28 (6H, m), 1.20-0.80 (1H, m), 0.60-0.20 (4H, m). MS (ESI, m/z) 492 (M+H) ⁺ .
Ex. 89		¹ H-NMR (300MHz, δ ppm, DMSO-d ₆) 7.68 (1H, d, J=8.4Hz), 7.59-7.55 (2H, m), 7.33-7.32 (1H, m), 7.25-7.18 (4H, m), 7.08-7.04 (2H, m), 6.48 (1H, s), 4.08 (1H, d, J=7.8Hz), 3.78 (1H, brs), 3.36-3.17 (4H, m), 2.90-2.85 (2H, m), 2.31 (3H, s), 1.10-1.07 (7H, m), 0.52-0.50 (1H, m), 0.37-0.33 (2H, m), 0.14-0.10 (1H, m). MS (ESI, m/z) 434 (M+H-1/2C ₄ H ₄ O ₄) ⁺ .
Ex. 90		¹ H-NMR (300MHz, δ ppm, DMSO-d ₆) 7.70 (1H, d, J=8.7Hz), 7.61-7.57 (2H, m), 7.35-7.32 (1H, m), 7.26-7.22 (2H, m), 7.08-6.94 (4H, m), 6.50 (1H, s), 4.27 (1H, d, J=7.1Hz), 3.84-3.72 (4H, m), 3.48-3.18 (4H, m), 3.06-2.92 (2H, m), 1.17-1.08 (7H, m), 0.50-0.24 (4H, m). MS (ESI, m/z) 450 (M+H-1/2C ₄ H ₄ O ₄) ⁺ .
Ex. 91		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.85-7.70 (3H, m), 7.61 (1H, s), 7.50-7.40 (2H, m), 7.35-7.15 (4H, m), 4.10-4.00 (1H, m), 3.85-3.70 (1H, m), 3.40-3.25 (2H, m), 2.90-2.60 (4H, m), 2.29 (3H, m), 1.20-1.00 (7H, m), 0.65-0.55 (1H, m), 0.50-0.15 (3H, m). MS (ESI, m/z) 497 (M+H) ⁺ .



Ex. 92		¹H-NMR (300MHz, δppm, DMSO-d_6) 8.00-7.94 (1H, m), 7.80-7.70 (3H, m), 7.61-7.55 (2H, m), 7.46-7.43 (3H, m), 7.28-7.16 (2H, m), 5.80-5.70 (1H, brs), 4.22-4.17 (1H, m), 3.81-3.70 (4H, m), 3.50-3.30 (2H, m), 2.85-2.60 (4H, m), 1.15-1.07 (7H, m), 0.70-0.30 (4H, m). MS (ESI, m/z) 477 (M+H)⁺.
Ex. 93		¹H-NMR (300MHz, δppm, DMSO-d_6) 7.40-7.10 (4H, m), 6.86 (1H, d, J=8.2Hz), 6.77 (1H, s), 6.70 (1H, d, J=8.1Hz), 6.47 (1H, s), 3.90-3.60 (7H, m), 3.40-3.10 (2H, m), 2.90-2.80 (1H, m), 2.75-2.50 (3H, m), 1.30-1.00 (7H, m), 0.60-0.10 (4H, m). MS (ESI, m/z) 428 (M+H-1/2C₄H₄O₄)⁺.

Table 13

Ex. 94	• 1/2 HO₂C-CH=CH-CO₂H	¹ H-NMR (300MHz, δppm, DMSO-d ₆) 7.40-7.10 (4H, m), 7.10-6.90 (2H, m), 6.88 (1H, d, J=8.2Hz), 6.77 (1H, s), 6.71 (1H, d, J=8.2Hz), 6.50 (1H, s), 4.27 (1H, d, J=7.4Hz), 3.95-3.65 (10H, m), 3.40-3.15 (2H, m), 3.10-2.90 (1H, m), 2.85-2.60 (3H, m), 1.20-1.00 (7H, m), 0.60-0.10 (4H, m). MS (ESI, m/z) 444 (M+H-1/2C ₄ H ₄ O ₄) ⁺ .
Ex. 95	• 1/2 HO₂C-CH=CH-CO₂H	¹ H-NMR (300MHz, δppm, DMSO-d ₆) 7.40-6.90 (6H, m), 6.87 (1H, d, J=8.1Hz), 6.53 (1H, m), 4.09 (1H, d, J=7.5Hz), 4.00-3.80 (1H, m), 3.76 (3H, s), 3.45-2.95 (4H, m), 2.90-2.70 (2H, m), 2.33 (3H, s), 2.13 (3H, s), 1.20-1.00 (7H, m), 0.60-0.10 (4H, m). MS (ESI, m/z) 412 (M+H-1/2C ₄ H ₄ O ₄) ⁺ .
Ex. 96	• 1/2 HO₂C-CH=CH-CO₂H	¹ H-NMR (300MHz, δppm, DMSO-d ₆) 7.40-7.15 (2H, m), 7.05-6.90 (4H, m), 6.89 (1H, d, J=8.1Hz), 6.53 (1H, m), 4.27 (1H, d, J=7.5Hz), 3.95-3.70 (7H, m), 3.45-3.00 (4H, m), 3.90-3.65 (2H, m), 2.13 (3H, s), 1.20-1.00 (7H, m), 0.60-0.10 (4H, m). MS (ESI, m/z) 428 (M+H-1/2C ₄ H ₄ O ₄) ⁺ .
Ex. 97	• 1/2 HO₂C-CH=CH-CO₂H	¹ H-NMR (300MHz, δppm, CDCl ₃) 7.82-7.76 (3H, m), 7.48 (1H, s), 7.48-7.46 (2H, m), 7.33-7.29 (1H, m), 7.11-6.99 (3H, m), 6.88 (1H, s), 4.24-4.12 (1H, m), 3.61-3.38 (3H, m), 3.23-2.82 (4H, m), 2.25-2.23 (6H, m), 1.32-1.26 (6H, m), 1.11-0.94 (1H, m), 0.62-0.52 (1H, m), 0.46-0.35 (2H, m), 0.20-0.14 (1H, m). MS (ESI, m/z) 432 (M+H-1/2C ₄ H ₄ O ₄) ⁺ .
Ex. 98	• 1/2 HO₂C-CH=CH-CO₂H	¹ H-NMR (300MHz, δppm, CDCl ₃) 8.23-8.21 (2H, m), 7.79-7.74 (4H, m), 7.65-7.62 (2H, m), 7.50-7.43 (3H, m), 7.29-7.17 (2H, m), 7.10-6.90 (2H, m), 4.38-4.34 (1H, m), 3.91 (3H, s), 3.80-3.70 (1H, m), 3.31-3.28 (2H, m), 3.19 (3H, s), 2.89-2.69 (4H, m), 1.33-1.30 (3H, m), 1.14-1.12 (6H, m). MS (ESI, m/z) 605 (M+H) ⁺ .







Ex. 99		¹ H-NMR (300MHz, δ ppm, DMSO-d ₆) 8.20-8.17 (1H, m), 7.89- 7.64 (7H, m), 7.52-7.17 (6H, m), 6.90-6.77 (1H, m), 4.40- 4.30 (1H, m), 3.95-3.80 (1H, m), 3.65-3.55 (1H, m), 3.24- 3.13 (3H, m), 3.09 (3H, s), 3.08- 2.90 (2H, m), 1.30-1.20 (9H, m). MS (ESI, m/z) 591 (M+H) ⁺ .
-----------	---	--

Table 14

Ex. 100		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.82-7.80 (3H, m), 7.68-7.66 (1H, m), 7.50-7.47 (2H, m), 7.32-7.28 (1H, m), 7.17-7.14 (1H, m), 6.79-6.76 (1H, m), 4.20-4.10 (1H, m), 4.09-4.02 (1H, m), 3.62-3.48 (2H, m), 3.27-3.23 (1H, m), 3.12-3.00 (3H, m), 2.18-2.17 (3H, m), 1.38-1.32 (6H, m), 1.10-1.00 (1H, m), 0.60-0.20 (4H, m). MS (ESI, m/z) 424 (M+H) ⁺ .
Ex. 101		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.76-7.74 (3H, m), 7.65 (1H, s), 7.45-7.42 (2H, m), 7.31-7.19 (2H, m), 6.93-6.90 (2H, m), 6.85 (1H, s), 4.20-4.10 (1H, m), 3.90 (1H, dd, J=11.7, 8.1 Hz), 3.64-3.40 (2H, m), 3.20-2.80 (4H, m), 1.30-1.24 (6H, m), 1.10-1.00 (1H, m), 0.60-0.45 (3H, m), 0.30-0.20 (1H, m). MS (APCI, m/z) 410 (M+H-1/2C ₄ H ₄ O ₄) ⁺ .
Ex. 102		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.78-7.75 (3H, m), 7.67 (1H, s), 7.46-7.43 (2H, m), 7.32-7.26 (1H, m), 6.85 (1H, s), 6.11-6.09 (1H, m), 5.87 (1H, s), 4.33 (1H, brs), 4.20-4.10 (1H, m), 3.60-3.45 (3H, m), 3.20-2.80 (3H, m), 2.24 (3H, s), 1.28-1.22 (6H, m), 1.20-1.10 (1H, m), 0.60-0.30 (3H, m), 0.20-0.10 (1H, m). MS (ESI, m/z) 408 (M+H-1/2C ₄ H ₄ O ₄) ⁺ .
Ex. 103		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.83-7.30 (3H, m), 7.70 (1H, s), 7.50-7.47 (2H, m), 7.37-7.30 (2H, m), 6.33-6.31 (1H, m), 6.26-6.25 (1H, m), 4.30-4.20 (1H, m), 3.63-3.52 (3H, m), 3.40-3.30 (1H, m), 3.15-3.10 (3H, m), 1.42-1.37 (6H, m), 1.15-1.00 (1H, m), 0.60-0.30 (3H, m), 0.20-0.10 (1H, m). MS (ESI, m/z) 394 (M+H) ⁺ .
Ex. 104		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.83-7.78 (3H, m), 7.68 (1H, brs), 7.49-7.46 (2H, m), 7.38-7.29 (3H, m), 6.36 (1H, brs), 4.30-4.10 (1H, m), 3.63-3.49 (3H, m), 3.29-3.25 (1H, m), 3.18-3.08 (3H, m), 1.37-1.34 (6H, m), 1.05-0.80 (1H, m), 0.56-0.33 (3H, m), 0.13-0.08 (1H, m). MS (ESI, m/z) 394 (M+H) ⁺ .







Ex. 105	 <chem>N#Cc1ccc(cc1)C2(CCC2)OC[C@H](O)CN(C)Cc3ccc4ccccc4c3</chem>	$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.85-7.75 (3H, m), 7.70-7.40 (7H, m), 7.35-7.25 (1H, m), 3.90-3.80 (1H, m), 3.75-3.65 (1H, m), 3.55-3.35 (2H, m), 3.00-2.85 (3H, m), 2.85-2.70 (1H, m), 1.29-0.95 (7H, m), 0.70-0.60 (1H, m), 0.55-0.45 (2H, m), 0.30-0.20 (1H, m). $\text{MS (ESI, } m/z \text{) } 429 \text{ (M+H)}^+$.
------------	---	---

Table 15

Ex. 106		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.85-7.70 (3H, m), 7.61 (1H, s), 7.50-7.40 (2H, m), 7.35-7.20 (2H, m), 7.15-7.05 (2H, m), 4.20-4.10 (1H, m), 3.90-3.75 (1H, m), 3.45-3.30 (2H, m), 2.95-2.65 (4H, m), 2.35-2.20 (6H, m), 1.35-1.00 (7H, m), 0.65-0.55 (1H, m), 0.50-0.20 (3H, m). MS (ESI, m/z) 432 (M+H) $^+$.
Ex. 107		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 8.15-8.05 (1H, m), 8.00-7.90 (1H, m), 7.85-7.70 (3H, m), 7.65-7.60 (1H, m), 7.50-7.40 (2H, m), 7.35-7.25 (1H, m), 6.95-6.85 (1H, m), 4.25-4.20 (1H, m), 3.95-3.80 (7H, m), 3.55-3.35 (2H, m), 3.00-2.70 (4H, m), 1.25-1.00 (7H, m), 0.65-0.55 (1H, m), 0.50-0.25 (3H, m). MS (ESI, m/z) 492 (M+H) $^+$.
Ex. 108		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.85-7.70 (3H, m), 7.61 (1H, s), 7.50-7.35 (2H, m), 7.30-7.25 (1H, m), 7.10-6.95 (2H, m), 6.90-6.80 (1H, m), 4.20-4.10 (1H, m), 3.90-3.70 (7H, m), 3.50-3.30 (2H, m), 2.90-2.60 (4H, m), 1.20-1.00 (7H, m), 0.70-0.25 (4H, m). MS (ESI, m/z) 464 (M+H) $^+$.
Ex. 109		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.82-7.70 (3H, m), 7.63 (1H, s), 7.47-7.43 (2H, m), 7.31-7.26 (3H, m), 4.15 (1H, dd, $J=7.7, 5.7\text{Hz}$), 3.90-3.80 (1H, m), 3.70-3.54 (2H, m), 2.90-2.78 (4H, m), 1.28-1.15 (7H, m), 0.70-0.45 (4H, m). MS (APCI, m/z) 411 (M+H) $^+$.
Ex. 110		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.82-7.77 (3H, m), 7.65 (1H, s), 7.48-7.45 (2H, m), 7.31-7.25 (2H, m), 7.14 (1H, s), 7.04-7.01 (1H, m), 4.07-4.00 (1H, m), 3.72-3.67 (1H, m), 3.60-3.46 (2H, m), 3.16-2.90 (4H, m), 1.28-1.26 (6H, m), 1.10-0.95 (1H, m), 0.60-0.35 (3H, m), 0.20-0.15 (1H, m). MS (APCI, m/z) 409 (M+H) $^+$.









Ex. 111	 1/2HO₂C-CH=CH-CO₂H	¹H-NMR (300MHz, δppm, DMSO-d₆) 7.40-7.10 (4H, m), 7.07 (1H, d, J=7.8Hz), 6.96 (1H, s) 6.91 (1H, d, J=7.8Hz), 6.53 (1H, s), 4.10 (1H, d, J=7.5Hz), 4.00- 3.80 (1H, m), 3.40-3.00 (3H, m), 2.90-2.70 (3H, m), 2.33 (3H, s), 2.20 (3H, s), 2.19 (3H, s), 1.20- 1.00 (7H, m), 0.60-0.10 (4H, m). MS (ESI, m/z) 396 (M+H-1/2C₄H₄O₄)⁺.
Ex. 112	 1/2HO₂C-CH=CH-CO₂H	¹H-NMR (300MHz, δppm, DMSO-d₆) 7.40-7.20 (4H, m), 7.15- 6.85 (5H, m), 6.46 (1H, s), 4.35- 4.20 (1H, m), 3.90-3.65 (4H, m), 3.40-3.10 (2H, m), 2.95- 2.65 (4H, m), 2.18 (6H, s) 1.20- 1.0 (7H, m), 0.60-0.10 (4H, m). MS (ESI, m/z) 412 (M+H-1/2C₄H₄O₄)⁺.

Table 16

Ex. 113		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.40-7.20 (4H, m), 7.15-6.85 (5H, m), 6.53 (1H, s), 4.27 (1H, d, $J=7.5\text{Hz}$), 3.95-3.70 (4H, m), 3.50-3.00 (8H, m), 2.95-2.70 (3H, m), 2.20 (3H, s), 2.19 (3H, s), 1.20-1.00 (7H, m), 0.60-0.10 (4H, m). MS (ESI, m/z) 412 ($M+H-1/2C_4H_4O_4$) ⁺ .
Ex. 114		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.85-7.70 (3H, m), 7.61 (1H, s), 7.50-7.40 (3H, m), 7.35-7.15 (4H, m), 4.40-4.30 (1H, m), 3.90-3.75 (1H, m), 3.50-3.30 (2H, m), 2.95-2.65 (4H, m), 1.20-1.00 (7H, m), 0.65-0.50 (1H, m), 0.50-0.35 (3H, m). MS (ESI, m/z) 438 ($M+H$) ⁺ .
Ex. 115		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.85-7.70 (3H, m), 7.62 (1H, s), 7.50-7.40 (2H, m), 7.35-7.10 (5H, m), 3.85-3.75 (1H, m), 3.70-3.55 (1H, m), 3.50-3.30 (2H, m), 2.95-2.65 (4H, m), 1.20-1.00 (7H, m), 0.70-0.55 (1H, m), 0.50-0.40 (2H, m), 0.30-0.20 (1H, m). MS (ESI, m/z) 438 ($M+H$) ⁺ .
Ex. 116		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.85-7.70 (3H, m), 7.61 (1H, s), 7.50-7.40 (2H, m), 7.35-7.15 (5H, m), 3.85-3.75 (1H, m), 3.70-3.60 (1H, m), 3.50-3.30 (2H, m), 2.90-2.80 (3H, m), 2.75-2.65 (1H, m), 1.20-1.00 (7H, m), 0.70-0.60 (1H, m), 0.50-0.40 (2H, m), 0.30-0.20 (1H, m). MS (ESI, m/z) 438 ($M+H$) ⁺ .
Ex. 117		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.85-7.25 (11H, m), 4.35-4.20 (1H, m), 3.80-3.70 (1H, m), 3.45-3.35 (1H, m), 3.30-3.15 (1H, m), 2.90-2.75 (3H, m), 2.75-2.60 (1H, m), 1.20-1.00 (7H, m), 0.65-0.30 (4H, m). MS (ESI, m/z) 472 ($M+H$) ⁺ .
Ex. 118		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.85-7.70 (3H, m), 7.65-7.40 (7H, m), 7.35-7.25 (1H, m), 3.90-3.70 (2H, m), 3.50-3.30 (2H, m), 2.95-2.60 (4H, m), 1.20-1.00 (7H, m), 0.70-0.60 (1H, m), 0.55-0.40 (2H, m), 0.30-0.20 (1H, m). MS (ESI, m/z) 472 ($M+H$) ⁺ .







Ex. 119		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.85-7.70 (3H, m), 7.65-7.55 (3H, m), 7.50-7.35 (4H, m), 7.35-7.25 (1H, m), 3.85-3.70 (2H, m), 3.50-3.30 (2H, m), 2.95-2.80 (3H, m), 2.80-2.65 (1H, m), 1.20-1.00 (7H, m), 0.70-0.55 (1H, m), 0.50-0.40 (2H, m), 0.30-0.20 (1H, m). MS (ESI, m/z) 472 ($M+H$) $^+$.
------------	---	--

Table 17

Ex. 120		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.90-7.77 (3H, m), 7.64 (1H, brs), 7.63-7.46 (2H, m), 7.32-7.18 (5H, m), 4.17-4.08 (2H, m), 3.59-3.53 (1H, m), 3.45-3.43 (1H, m), 3.26-3.22 (1H, m), 3.10-3.03 (3H, m), 2.66-2.59 (2H, m), 1.32-1.28 (6H, m), 1.22-1.17 (3H, m), 1.11-1.00 (1H, m), 0.60-0.50 (1H, m), 0.40-0.30 (2H, m), 0.25-0.20 (1H, m). MS (ESI, m/z) 432 (M+H) ⁺ .
Ex. 121		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.90-7.78 (3H, m), 7.65 (1H, brs), 7.50-7.46 (2H, m), 7.29-7.17 (5H, m), 4.21-4.11 (2H, m), 3.60-3.52 (1H, m), 3.45-3.42 (1H, m), 3.30-3.05 (4H, m), 1.34-1.18 (12H, m), 1.20-1.00 (1H, m), 0.60-0.50 (1H, m), 0.40-0.30 (2H, m), 0.20-0.10 (1H, m). MS (ESI, m/z) 446 (M+H) ⁺ .
Ex. 122		¹ H-NMR (300MHz, δ ppm, DMSO-d ₆) 7.96-7.82 (3H, m), 7.73 (1H, s), 7.50-7.47 (3H, m), 7.41-7.38 (2H, m), 7.13 (1H, d, J=5.1Hz), 3.90-3.81 (2H, m), 3.44-3.28 (3H, m), 3.10-2.70 (4H, m), 1.20-1.05 (7H, m), 0.60-0.50 (1H, m), 0.45-0.35 (2H, m), 0.30-0.20 (1H, m). MS (ESI, m/z) 410 (M+H-1/2C ₄ H ₄ O ₄) ⁺ .
Ex. 123		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.84-7.78 (3H, m), 7.66 (1H, s), 7.48-7.46 (2H, m), 7.32-7.29 (1H, m), 6.72 (1H, t, J=3.0Hz), 6.58 (1H, s), 4.16-4.06 (1H, m), 3.81-3.75 (1H, m), 3.61-3.54 (2H, m), 3.22-3.16 (1H, m), 3.06-2.95 (3H, m), 2.44 (1H, d, J=3.0Hz), 1.32-1.28 (6H, m), 1.16-1.06 (1H, m), 0.62-0.42 (3H, m), 0.30-0.20 (1H, m). MS (ESI, m/z) 424 (M+H) ⁺ .
Ex. 124		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.83-7.78 (3H, m), 7.66 (1H, s), 7.49-7.45 (2H, m), 7.31-7.28 (1H, m), 6.74 (1H, t, J=3.4Hz), 6.62-6.60 (1H, m), 4.14-4.09 (1H, m), 3.79 (1H, t, J=8.5Hz), 3.62-3.50 (2H, m), 3.24-3.02 (4H, m), 2.84-2.75 (2H, m), 1.32-1.23 (9H, m), 1.12-1.04 (1H, m), 0.66-0.40 (3H, m), 0.28-0.20 (1H, m). MS (ESI, m/z) 437 (M+H) ⁺ .







Ex. 125		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.81-7.79 (3H, m); 7.68 (1H, s), 7.49-7.45 (2H, m), 7.32-7.29 (1H, m), 6.88 (1H, t, $J=3.7\text{Hz}$), 6.70 (1H, d, $J=6.7\text{Hz}$), 4.14-4.08 (1H, m), 3.79 (1H, t, $J=8.7\text{Hz}$), 3.62-3.50 (2H, m), 3.22-2.90 (4H, m), 1.34-1.30 (6H, m), 1.10-1.00 (1H, m), 0.60-0.40 (3H, m), 0.30-0.20 (1H, m). MS (ESI, m/z) 490 (M+H) $^+$.
------------	---	---

Table 18

Ex. 126		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 8.20-8.05 (2H, m), 7.85-7.70 (3H, m), 7.70-7.60 (2H, m), 7.55-7.40 (3H, m), 7.35-7.25 (1H, m), 3.85-3.70 (2H, m), 3.55-3.35 (2H, m), 2.95-2.60 (4H, m), 1.20-1.00 (7H, m), 0.70-0.60 (1H, m), 0.55-0.40 (2H, m), 0.35-0.20 (1H, m). MS (ESI, m/z) 449 (M+H) $^+$.
Ex. 127		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.40-7.10 (4H, m), 7.00-6.85 (3H, m), 6.75 (1H, d, $J=8.2\text{Hz}$), 4.20-4.00 (3H, m), 3.80 (3H, s), 3.60-3.30 (2H, m), 3.20-2.80 (2H, m), 2.73 (2H, s), 2.32 (3H, s), 2.18 (3H, s), 1.30-1.00 (7H, m), 0.70-0.10 (4H, m). MS (ESI, m/z) 412 (M+H) $^+$.
Ex. 128		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.98-7.95 (2H, d, $J=8.3\text{Hz}$), 7.36-7.34 (1H, m), 7.24-7.16 (5H, m), 4.11-4.08 (1H, d, $J=7.7\text{Hz}$), 4.05-3.95 (1H, m), 3.91 (3H, s), 3.52-3.37 (2H, m), 3.05-3.01 (1H, m), 2.88-2.85 (1H, m), 2.34 (3H, s), 1.23-1.20 (1H, m), 1.14 (6H, s), 0.70-0.55 (1H, m), 0.50-0.20 (3H, m). MS (ESI, m/z) 426 (M+H) $^+$.
Ex. 129		$^1\text{H-NMR}$ (300MHz, δ ppm, DMSO-d_6) 7.86-7.83 (2H, d, $J=8.1\text{Hz}$), 7.34-7.14 (6H, m), 4.10-4.07 (1H, d, $J=7.5\text{Hz}$), 3.80-3.70 (1H, m), 3.35-3.30 (1H, m), 3.22-3.17 (1H, m), 2.87-2.81 (3H, m), 2.66-2.59 (1H, m), 2.32 (3H, s), 1.25-1.10 (1H, m), 1.05-1.04 (6H, m), 0.60-0.15 (4H, m). MS (ESI, m/z) 412 (M+H) $^+$.
Ex. 130		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.81-7.76 (3H, m), 7.65-7.63 (2H, m), 7.48-7.45 (2H, m), 7.32-7.29 (1H, m), 6.94 (1H, d, $J=3.7\text{Hz}$), 4.10-4.06 (1H, m), 3.90-3.84 (4H, m), 3.64-3.60 (1H, m), 3.56-3.50 (1H, m), 3.18-2.90 (4H, m), 1.30-1.28 (6H, m), 1.12-1.02 (1H, m), 0.68-0.46 (3H, m), 0.32-0.24 (1H, m). MS (ESI, m/z) 468 (M+H) $^+$.








Ex. 131		¹ H-NMR (300MHz, δ ppm, , DMSO-d ₆) 7.91-7.85 (3H, m) , 7.73 (1H, s) , 7.49-7.46 (3H, m) , 7.36 (1H, d, J=8.1Hz) , 7.03 (1H, s) , 4.06 (1H, d, J=8.1Hz) , 3.89- 2.80 (7H, m) , 1.24-1.19 (7H, m) , 0.60-0.20 (4H, m) . MS (ESI, m/z) 454 (M+H) ⁺ .
Ex. 132		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.85- 7.40 (10H, m) , 7.35-7.30 (1H, m) , 3.85-3.70 (2H, m) , 3.50- 3.35 (2H, m) , 3.00-2.85 (3H, m) , 2.80-2.65 (7H, m) , 1.20- 1.00 (7H, m) , 0.70-0.60 (1H, m) , 0.55-0.40 (2H, m) , 0.30- 0.20 (1H, m) . MS (ESI, m/z) 511 (M+H) ⁺ .

Table 19

Ex. 133		¹ H-NMR (300MHz, δ ppm, DMSO-d ₆) 7.87-7.77 (3H, m), 7.67 (1H, s), 7.60 (1H, d, J=3.7Hz), 7.46- 7.44 (2H, m), 7.37-7.35 (1H, m), 7.01 (1H, d, J=3.7Hz), 3.99 (1H, dd, J=7.8, 3.2Hz), 3.65- 3.60 (1H, m), 3.45-3.33 (2H, m), 2.78-2.70 (4H, m), 1.10- 0.97 (7H, m), 0.60-0.40 (3H, m), 0.32-0.26 (1H, m). MS (ESI, m/z) 453 (M+H) ⁺ .
Ex. 134		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.83- 7.80 (3H, m), 7.68 (1H, s), 7.49- 7.46 (2H, m), 7.31 (1H, d, J=8.4Hz), 6.75-6.70 (2H, m), 4.18- 4.12 (1H, m), 3.75 (1H, t, J=9.3Hz), 3.64-3.51 (2H, m), 3.23- 2.99 (4H, m), 1.36-0.95 (7H, m), 0.70-0.40 (3H, m), 0.30- 0.20 (1H, m). MS (ESI, m/z) 444 (M+H) ⁺ .
Ex. 135		¹ H-NMR (300MHz, δ ppm, DMSO-d ₆) 7.88-7.78 (3H, m), 7.68 (1H, s), 7.49-7.45 (2H, m), 7.37 (1H, d, J=8.4Hz), 6.82 (2H, dd, J=14.7, 3.3Hz), 5.37 (1H, t, J=5.4Hz), 3.94 (1H, dd, J=8.1, 3.3Hz), 3.64 (1H, brs), 3.38-3.17 (2H, m), 2.81-2.75 (2H, m), 1.14- 1.06 (7H, m), 0.60-0.50 (1H, m), 0.48-0.40 (2H, m), 0.30- 0.20 (1H, m). MS (ESI, m/z) 440 (M+H) ⁺ .
Ex. 136		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.80- 7.74 (5H, m), 7.62 (1H, s), 7.47- 7.39 (4H, m), 7.27-7.25 (1H, m), 4.37 (1H, dd, J=14.4, 7.8Hz), 4.10- 4.04 (1H, m), 3.63-3.48 (2H, m), 3.16-2.97 (4H, m), 1.28- 1.10 (7H, m), 0.63-0.42 (4H, m). MS (ESI, m/z) 494 (M+H) ⁺ .
Ex. 137		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.82- 7.77 (3H, m), 7.67 (1H, s), 7.48- 7.45 (2H, m), 7.30 (1H, d, J=8.4Hz), 7.12 (1H, dd, J=4.7, 1.4Hz), 6.86 (1H, s), 4.16-4.06 (1H, m), 3.80 (1H, t, J=8.8Hz), 3.62- 3.50 (2H, m), 3.24-2.97 (4H, m), 1.34-1.25 (6H, m), 1.06- 1.02 (1H, m), 0.60-0.40 (3H, m), 0.30-0.20 (1H, m). MS (ESI, m/z) 490 (M+H) ⁺ .







Ex. 138		$^1\text{H-NMR}$ (300 MHz, δ ppm, CDCl_3) 7.81-7.78 (3H, m), 7.68 (1H, s), 7.49-7.46 (2H, m), 7.30 (1H, d, $J=8.4\text{ Hz}$), 6.90 (1H, s), 6.81 (1H, dd, $J=3.2, 1.7\text{ Hz}$), 4.20-4.14 (1H, m), 3.62-3.06 (7H, m), 1.38-1.35 (6H, m), 1.00-0.70 (1H, m), 0.60-0.30 (3H, m), 0.20-0.10 (1H, m). MS (ESI, m/z) 444 ($M+H$) $^+$.
------------	---	---

Table 20

Ex. 139		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.72-7.70 (2H, d, $J=8.2\text{Hz}$), 7.38-7.26 (1H, m), 7.24-7.16 (5H, m), 6.30-5.50 (2H, br), 4.13-4.10 (1H, d, $J=7.4\text{Hz}$), 3.85-3.70 (1H, m), 3.35-3.29 (2H, m), 2.78-2.59 (4H, m), 2.35 (3H, s), 1.30-1.20 (1H, m), 1.04 (3H, s), 1.02 (3H, s), 0.65-0.20 (4H, m). MS (ESI, m/z) 411 (M+H) $^+$.
Ex. 140		$^1\text{H-NMR}$ (300MHz, δ ppm, DMSO-d_6) 7.89 (2H, d, $J=8.1\text{Hz}$), 7.36-7.33 (3H, m), 7.28-7.22 (1H, m), 7.01-6.94 (2H, m), 4.30-4.26 (1H, m), 4.08-3.76 (7H, m), 3.24-3.16 (2H, m), 2.86 (3H, brs), 2.76-2.64 (1H, m), 1.12-1.08 (7H, m), 0.50-0.40 (1H, m), 0.45-0.20 (3H, m). MS (ESI, m/z) 442 (M+H) $^+$.
Ex. 141		$^1\text{H-NMR}$ (300MHz, δ ppm, DMSO-d_6) 7.36-7.32 (1H, m), 7.27-7.18 (3H, m), 7.18-7.11 (2H, m), 7.00-6.93 (2H, m), 5.12-5.04 (1H, m), 4.45 (2H, d, $J=4.5\text{Hz}$), 4.28 (1H, d, $J=6.9\text{Hz}$), 4.07 (1H, brs), 3.77 (3H, s), 3.61 (1H, brs), 3.24-3.17 (3H, m), 2.70-2.59 (2H, m), 1.10-1.00 (1H, m), 0.94-0.92 (6H, m), 0.50-0.40 (1H, m), 0.30-0.20 (3H, m). MS (ESI, m/z) 414 (M+H) $^+$.
Ex. 142		$^1\text{H-NMR}$ (300MHz, δ ppm, DMSO-d_6) 7.33-7.31 (1H, m), 7.20-7.08 (7H, m), 5.09 (1H, t, $J=5.7\text{Hz}$), 4.45 (2H, d, $J=5.5\text{Hz}$), 4.11-4.06 (4H, m), 3.60 (1H, brs), 3.30-3.24 (1H, m), 2.70-2.57 (2H, m), 2.31 (3H, s), 1.20-1.10 (1H, m), 0.93-0.91 (6H, m), 0.55-0.45 (1H, m), 0.40-0.30 (2H, m), 0.25-0.15 (1H, m). MS (ESI, m/z) 398 (M+H) $^+$.
Ex. 143		$^1\text{H-NMR}$ (300MHz, δ ppm, DMSO-d_6) 7.91 (2H, d, $J=8.1\text{Hz}$), 7.36-7.24 (4H, m), 7.02-6.96 (2H, m), 4.31-4.26 (1H, m), 3.88 (1H, brs), 3.78 (3H, s), 3.20-2.85 (4H, m), 2.60-2.50 (2H, m), 1.18 (7H, brs), 0.60-0.40 (1H, m), 0.40-0.20 (3H, m). MS (ESI, m/z) 428 (M+H) $^+$.


Ex. 144	 <chem>COc1ccc(cc1)C(C)(C)NC(O)COC(C2CC2)c3cc(Br)sc3</chem>	$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.00-6.89 (3H, m), 6.78-6.73 (2H, m), 4.22-4.18 (1H, m), 3.87-3.83 (1H, m), 3.80 (3H, s), 3.65-3.53 (2H, m), 3.30-3.20 (1H, m), 3.12-3.01 (1H, m), 2.88-2.85 (2H, m), 2.18 (3H, s), 1.33 (6H, d, $J=11.1\text{Hz}$), 1.15-1.00 (1H, m), 0.70-0.45 (3H, m), 0.30-0.20 (1H, m). $\text{MS (ESI, } m/z \text{) } 484 \text{ (M+H)}^+$.
------------	---	--

Table 21

Ex. 145		¹ H-NMR (300MHz, δ ppm, DMSO-d ₆) 9.18 (1H, s), 7.40-7.10 (4H, m), 6.90-6.60 (3H, m), 4.20- 4.00 ((2H, m), 3.95-3.75 (1H, m), 3.40-2.90 (3H, m), 2.85- 2.60 (3H, m), 2.33 (3H, s), 2.09 (3H, s), 1.30-1.00 (7H, m), 0.60-0.10 (4H, m). MS (ESI, m/z) 398 (M+H) ⁺ .
Ex. 146		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.40- 7.30 (1H, m), 7.20-7.00 (3H, m), 6.80-6.50 (3H, m), 5.93 (2H, s), 4.10 (1H, d, J=7.6Hz), 3.95- 3.75 (1H, m), 3.50-3.20 (2H, m), 3.00-2.55 (4H, m), 2.34 (3H, s), 1.30-1.00 (7H, m), 0.70- 0.10 (4H, m). MS (ESI, m/z) 412 (M+H) ⁺ .
Ex. 147		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.85- 7.70 (3H, m), 7.61 (1H, s), 7.50- 7.35 (3H, m), 7.35-6.95 (4H, m), 4.15-4.05 (1H, m), 3.80- 3.70 (1H, m), 3.50-3.30 (2H, m), 2.90-2.60 (4H, m), 1.20- 1.00 (7H, m), 0.70-0.55 (1H, m), 0.50-0.25 (3H, m). MS (ESI, m/z) 422 (M+H) ⁺ .
Ex. 148		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.85- 7.70 (3H, m), 7.62 (1H, s), 7.50- 7.40 (2H, m), 7.35-7.20 (2H, m), 7.10-6.90 (3H, m), 3.90- 3.75 (1H, m), 3.66 (1H, t, J=8.1Hz), 3.50-3.30 (2H, m), 2.95- 2.60 (4H, m), 1.20-1.00 (7H, m), 0.70-0.55 (1H, m), 0.50- 0.40 (2H, m), 0.30-0.20 (1H, m). MS (ESI, m/z) 422 (M+H) ⁺ .
Ex. 149		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.85- 7.70 (3H, m), 7.61 (1H, s), 7.50- 7.40 (2H, m), 7.35-7.20 (3H, m), 7.10-6.95 (2H, m), 3.85- 3.75 (1H, m), 3.66 (1H, t, J=7.8Hz), 3.45-3.30 (2H, m), 2.90- 2.65 (4H, m), 1.20-1.00 (7H, m), 0.70-0.55 (1H, m), 0.50- 0.35 (2H, m), 0.30-0.15 (1H, m). MS (ESI, m/z) 422 (M+H) ⁺ .
Ex. 150		¹ H-NMR (300MHz, δ ppm, CDCl ₃) 7.85- 7.70 (3H, m), 7.61 (1H, s), 7.55- 7.40 (4H, m), 7.40-7.20 (2H, m), 7.20-7.05 (1H, m), 4.40- 4.35 (1H, m), 3.80-3.70 (1H, m), 3.45-3.25 (2H, m), 2.90- 2.60 (4H, m), 1.20-1.00 (7H, m), 0.60-0.35 (4H, m). MS (ESI, m/z) 483 (M+H) ⁺ .







Ex. 151		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3): 7.85-7.70 (3H, m), 7.61 (1H, s), 7.50-7.10 (7H, m), 3.85-3.70 (1H, m), 3.63 (1H, t, $J=7.2\text{Hz}$), 3.50-3.30 (2H, m), 2.95-2.60 (4H, m), 1.20-1.00 (7H, m), 0.70-0.55 (1H, m), 0.50-0.40 (2H, m), 0.30-0.20 (1H, m). MS (ESI, m/z) 483 (M+H) $^+$.
------------	---	---

Table 22

Ex. 152		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.85-7.70 (3H, m), 7.61 (1H, s), 7.50-7.40 (4H, m), 7.35-7.20 (1H, m), 7.20-7.10 (2H, m), 3.85-3.75 (1H, m), 3.63 (1H, t, $J=7.8\text{Hz}$), 3.45-3.30 (2H, m), 2.95-2.80 (3H, m), 2.80-2.65 (1H, m), 1.20-1.00 (7H, m), 0.70-0.55 (1H, m), 0.50-0.35 (2H, m), 0.30-0.15 (1H, m). MS (ESI, m/z) 483 (M+H) $^+$.
Ex. 153		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.85-7.70 (3H, m), 7.61 (1H, s), 7.50-7.40 (2H, m), 7.35-7.25 (1H, m), 6.90-6.75 (3H, m), 3.95-3.75 (7H, m), 3.61 (1H, t, $J=8.5\text{Hz}$), 3.50-3.35 (2H, m), 2.95-2.80 (3H, m), 2.80-2.65 (1H, m), 1.20-1.00 (7H, m), 0.70-0.55 (1H, m), 0.50-0.40 (2H, m), 0.30-0.15 (1H, m). MS (APCI, m/z) 464 (M+H) $^+$.
Ex. 154		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.85-7.70 (3H, m), 7.61 (1H, s), 7.50-7.40 (2H, m), 7.35-7.25 (1H, m), 6.83 (1H, s), 6.80-6.65 (2H, m), 5.94 (2H, d, $J=1.1\text{Hz}$), 3.85-3.70 (1H, m), 3.57 (1H, t, $J=7.7\text{Hz}$), 3.40-3.30 (2H, m), 2.90-2.80 (3H, m), 2.80-2.65 (1H, m), 1.20-1.00 (7H, m), 0.65-0.55 (1H, m), 0.50-0.35 (2H, m), 0.30-0.15 (1H, m). MS (APCI, m/z) 448 (M+H) $^+$.
Ex. 155		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.85-7.70 (3H, m), 7.63 (1H, s), 7.50-7.40 (2H, m), 7.35-7.30 (1H, m), 6.24-6.23 (2H, m), 3.90-3.75 (1H, m), 3.70-3.60 (1H, m), 3.60-3.45 (2H, m), 2.95-2.85 (3H, m), 2.85-2.65 (1H, m), 1.30-1.05 (7H, m), 0.70-0.60 (1H, m), 0.60-0.40 (2H, m), 0.30-0.20 (1H, m). MS (APCI, m/z) 473 (M+H) $^+$.
Ex. 156		$^1\text{H-NMR}$ (300MHz, δ ppm, DMSO-d_6) 7.95-7.80 (2H, m), 7.36-7.23 (4H, m), 7.01-6.95 (2H, m), 4.31-4.27 (1H, n), 4.08 (1H, brs), 3.84 (1H, brs), 3.78 (3H, s), 3.31-2.73 (5H, m), 1.11 (7H, brs), 0.50-0.20 (4H, m). MS (ESI, m/z) 427 (M+H) $^+$.





Ex. 157		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.82-7.71 (3H, m), 7.61 (1H, s), 7.45-7.42 (2H, m), 7.32-7.20 (3H, m), 6.97 (1H, t, $J=7.4\text{Hz}$), 6.88 (1H, d, $J=8.1\text{Hz}$), 5.08-5.03 (1H, m), 4.10-3.90 (3H, m), 3.00-2.80 (4H, m), 1.52 (3H, dd, $J=6.6, 1.4\text{Hz}$), 1.14 (6H, d, $J=4.9\text{Hz}$). MS (ESI, m/z) 394 (M+H) $^+$.
------------	---	--

Table 23

Ex. 158		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.82-7.76 (3H, m), 7.65 (1H, s), 7.47-7.45 (2H, m), 7.27 (1H, d, $J=10.6\text{Hz}$), 6.56 (1H, dd, $J=3.6, 1.8\text{Hz}$), 6.01-5.99 (1H, m), 4.05-3.95 (1H, m), 3.86-3.83 (3H, m), 3.72 (1H, t, $J=7.7\text{Hz}$), 3.60-3.45 (2H, m), 3.12-2.89 (4H, m), 1.27-1.24 (6H, m), 1.09-1.00 (1H, m), 0.60-0.40 (3H, m), 0.30-0.20 (1H, m). MS (ESI, m/z) 440 (M+H) $^+$.
Ex. 159		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.82-7.73 (3H, m), 7.61 (1H, s), 7.47-7.40 (2H, m), 7.31-7.27 (2H, m), 6.51-6.43 (2H, m), 4.15 (1H, d, $J=7.8\text{Hz}$), 3.80-3.77 (7H, m), 3.48-3.33 (2H, m), 2.84-2.68 (4H, m), 1.19-1.68 (7H, m), 0.60-0.50 (1H, m), 0.38-0.25 (3H, m). MS (ESI, m/z) 464 (M+H) $^+$.
Ex. 160		$^1\text{H-NMR}$ (300MHz, δ ppm, CDCl_3) 7.40-7.10 (4H, m), 7.00-6.85 (3H, m), 6.72 (1H, d, $J=8.7\text{Hz}$), 4.09 (1H, d, $J=7.5\text{Hz}$), 4.05-3.85 (3H, m), 3.50-3.30 (2H, m), 3.20-2.75 (2H, m), 2.67 (2H, s), 2.33 (3H, s), 2.19 (3H, s), 1.41-3H, t, $J=6.9\text{Hz}$), 1.25-1.00 (7H, m), 0.65-0.10 (4H, m). MS (ESI, m/z) 426 (M+H) $^+$.

Experimental Examples

[0239] The biological activity of the compound of the present invention was examined.

Experimental Example 1


[0240] Evaluation of antagonistic action on calcium receptor using reporter gene

[0241] Luciferase cDNA and human calcium receptor cDNA were transfected into a cell strain derived from rat adrenal and the transformed cells were cultured overnight in a medium (F12 medium containing 0.5% dialyzed horse serum and 0.25% dialyzed bovine fetal serum). The next day, a dimethyl sulfoxide solution containing a test compound at 0.01 - 100 mM was diluted 100-fold with the medium and added to the test compound group at 10 μ l per well. A medium containing 50 mM calcium chloride was added to the control group at 10 μ l per well, such that the final calcium concentration of the medium became 5 mM. A medium alone was added to the blank group. After culture for 4 hr, luciferase substrate (PicaGene LT-2.0, TOYO INK) was added and the luciferase activity was measured with a photoluminometer. The inhibitory rate (%) was calculated from the obtained measured values according to the following formula.

$$\text{Inhibitory rate (\%)} = 100 - \frac{\text{measured value of compound group} - \text{measured value of blank group}}{\text{measured value of control group} - \text{measured value of blank group}} \times 100$$

[0242] Based on the results, the concentration (IC₅₀) showing 50% inhibitory rate was determined.

[0243] For reference, a compound of the above-mentioned formula [1-3] wherein R¹¹, R¹² and R²¹ are hydrogen atoms and R⁷¹ and R⁷² in combination show -CH=CH-CH=CH-



(Comparative Example 1) was also subjected to the testing. The results are shown in Table 24 and Table 25.

Experimental Example 2

PTH secretion promoting action

[0244] The test compound was orally administered to 6 to 9-week-old male SD rats (Charles River Japan, Inc.) fasted for 20 hr, using a solvent (5% ethanol, 0.5% aqueous methyl cellulose solution) at a dose of 30 mg/5 ml/kg and 100 mg/5 ml/kg. A solvent alone was orally administered to the control group at a dose of 5 ml/kg. The blood was drawn from the tail vein immediately before and 0.5, 1, 2, 4, 6 hr after the administration of the test compound, and sera were obtained. The serum PTH concentration was measured using rat PTH (1-84) ELISA kit (Nohon Medi-Physics). The results of the serum PTH concentration before and 30 min and 4 hr after the administration of the test compound are shown in Table 24, and the results of the serum PTH concentration before and 30 min and 2 hr after the administration of the test compound of the 30 mg/5 ml/kg administration group are shown in Table 25.

[0245] In Fig. 1, Fig. 2 and Fig. 3, the time-course changes of the serum PTH concentration at the dose (30 mg/kg) of Example 22, Example 23 and Example 24 are shown. For reference, moreover, Fig. 4 shows the time-course changes of the serum PTH concentration at the dose of Comparative Example 1 (100 mg/kg), and Fig. 5 shows the time-course changes of the serum PTH concentration at the dose of NPS-2143 of 30 mg/kg.

Table 24

test compound	IC ₅₀ (μM)	serum PTH concentration (pg/ml)		
		Before administration	30 min later later	240 min later
		30 mg/kg administration group (upper line)		
		100 mg/kg administration group (lower line)		
Example 1	0.041	56.9±9.4	141.5±15.3	54.1±7.7
		55.8±3.9	160.0±13.8	58.3±7.3

EP 1 308 436 A1

Table 24 (continued)

test compound	IC ₅₀ (μM)	serum PTH concentration (pg/ml)		
		Before administration	30 min later later	240 min later
		30 mg/kg administration group (upper line)		
		100 mg/kg administration group (lower line)		
Example 2	0.027	52.0±3.4	148.4±10.1	60.0±5.8
		68.9±7.6	154.8±17.1	85.8±17.3
Example 5	0.070	78.9±11.8	155.6±14.9	84.8±12.3
		-	-	-
Example 22	0.059	62.2±5.2	185.7±10.0	73.4±11.0
		-	-	-
Example 23	0.027	85.3±3.6	228.3±15.0	73.9±6.9
		-	-	-
Example 24	0.022	66.5±4.3	197.7±18.9	77.5±9.0
		-	-	-
Example 62	0.045	99.3±43.6	259.2±29.4	82.2±9.9
		-	-	-
Comparative Example 1	0.56	-	-	-
		107.1±17.5	139.0±18.8	78.9±12.3
(- means unmeasured, mean±S.E.)				

Table 25

test compound	IC ₅₀ (μM)	serum PTH concentration (pg/ml)		
		Before administration	30 min later	120 min later
Example 80	0.064	44.8±6.3	156.6±15.3	76.1±12.6
Example 82	0.051	66.7±5.0	121.9±9.0	68.5±11.4
Example 95	0.029	105.8±44.2	199.1±25.8	80.4±9.6
Example 100	0.067	52.2±4.4	130.1±53.7	49.3±4.4
Example 105	0.049	38.3±8.6	115.4±20.0	74.3±12.1
Example 111	0.024	51.2±12.8	236.3±60.8	65.3±6.4
Example 127	0.133	65.0±6.9	265.3±22.3	75.2±7.3
(- means unmeasured, mean±S.E.)				

Experimental Example 3

PTH secretion promoting action

[0246] The test compound was orally administered to 4 to 6-week-old female Fisher rats (Charles River Japan, Inc.) fasted for 20 hr, using a solvent (5% ethanol, 0.5% aqueous methyl cellulose solution) at a dose of 30 mg/5 ml/kg. A solvent alone was orally administered to the control group at a dose of 5 ml/kg. The blood was drawn from the tail vein immediately before and 0.5, 1, 2, 4 hr after the administration of the test compound, and sera were obtained. The serum PTH concentration was measured using rat PTH (1-84) ELISA kit (Nohon Medi-Physics). The results of the serum PTH concentration before and 30 min and 4 hr after the administration of the test compound are shown in Table 26.

Table 26

test compound	serum PTH concentration (pg/ml)		
	Before administration	30 min later	240 min later
Example 58	179.2±45.6	644.4±65.2	394.1±100.9
Example 59	189.8±90.9	313.4±103.3	62.2±10.3
Example 62	82.9±24.5	620.6±34.2	100.1±11.7
(mean±S.E.)			

[0247] When osteoporosis is to be treated by increasing the blood PTH concentration by inhibition of the action of calcium receptor, the compound to be used for this end should have at least the following properties.

(i) The compound has a sufficient antagonistic action on calcium receptors. In other words, the compound has a sufficiently low IC_{50} value. In the specification of WO99/51241, it is described, "In general, a compound showing a low IC_{50} value in the assay of calcium receptor inhibitor is a more superior compound. A compound showing an IC_{50} value of not lower than 50 μM is considered to be inactive. A preferable compound shows an IC_{50} value of not more than 10 μM , more preferably 1 μM , and most preferably not more than 0.1 μM ."

(ii) Administration of the compound results in a sufficient increase in the blood PTH concentration.

(iii) The time-course concentrations in blood after administration of the compound are not sustainable. Desirably, the PTH concentration before administration is restored at 3, 4 hr after administration of the compound.

[0248] From the above-mentioned test results, the compound of the present invention clearly shows the above-mentioned characteristics.

[0249] As regards (i); As shown in Table 24 and Table 25, the IC_{50} value of the compound of the present invention is not more than 1 μM , and the compound has a sufficient antagonistic action on calcium receptors. The compound of the present invention is considered to be preferable in view of the IC_{50} value.

[0250] As regards (ii); As shown in Table 24 - Table 26 and Figs. 1-3, at 30 min later, the 30 mg/kg administration group showed a 2.0 - 3.0 times higher blood PTH concentration than that before administration, and the 100 mg/kg administration group showed a 2.2 - 3.6 times higher blood PTH concentration than that before administration, and the compound of the present invention has been confirmed to have a superior PTH secretion promoting action. In contrast, as shown in Table 24 and Fig. 4, at 30 min after administration, the compound of Comparative Example 1 only showed a 1.3 times higher blood PTH concentration even at the dose of 100 mg/kg. The compound was not found to have a superior PTH secretion promoting action, and cannot be expected to be a pharmaceutical product.

[0251] As regards (iii); As shown in Table 24 - Table 26 and Figs. 1-3, PTH secretion by the compound of the present invention reached a peak at 30 min after administration, sharply decreased thereafter and returned to the blood PTH concentration before administration in about 2 - 4 hr. It is clear that the compound of the present invention is superior from this aspect. In contrast, as a result of the reproductive test of NPS-2143 of the reference, the sustained secretion promoting action of NPS-2143 was confirmed (from Fig. 5).


Industrial Applicability

[0252] As is clear from the above-mentioned Experimental Example 1, the compound of the formula [I] of the present invention has a superior calcium receptor antagonistic action. Accordingly, the compound is expected to be useful as a therapeutic drug of diseases accompanied by abnormal calcium homeostasis, such as osteoporosis, hypoparathyroidism, osteosarcoma, periodontal disease, bone fracture, steoarthritis, chronic rheumatoid arthritis, Paget's disease, humoral hypercalcemia, autosomal dominant hypocalcemia and the like. As is clear from Experimental Examples 2 and 3, the compound of the present invention has a temporary PTH secretion promoting action. Accordingly, the compound is particularly useful as a therapeutic agent for osteoporosis.

[0253] This application is based on patent application Nos. 244536/2000 and 132879/2001 filed in Japan, the contents of which are hereby incorporated by reference.


Claims

1. A compound of the formula [I]



wherein

R^1 is aryl group or heteroaryl group wherein said aryl group and heteroaryl group are optionally substituted by 1 to 3 substituents selected from halogen atom, C_{1-6} alkyl group, halo(C_{1-6})alkyl group, hydroxy(C_{1-6})alkyl group, C_{1-6} alkoxy(C_{1-6})alkyl group, hydroxyl group, C_{1-6} alkoxy group, halo(C_{1-6})alkoxy group, mercapto group, C_{1-6} alkylthio group, C_{1-6} alkylsulfanyl group, C_{1-6} alkylsulfonyl group, aminosulfonyl group, C_{1-6} alkylsulfamoyl group, di(C_{1-6})alkylsulfamoyl group, carboxy group, (C_{1-6} alkoxy)carbonyl group, C_{1-7} acyl group, carbamoyl group, (C_{1-6} alkyl)carbamoyl group, di(C_{1-6} alkyl)carbamoyl group, cyano group, nitro group, amino group, C_{1-6} alkylamino group, di(C_{1-6})alkylamino group, C_{1-7} acylamino group, C_{1-3} alkylenedioxy group,



wherein R^A is (C_{1-6} alkoxy)carbonyl group or carboxy group, and R^B is hydrogen atom or C_{1-6} alkyl group;

R^2 is C_{1-6} alkyl group wherein said C_{1-6} alkyl group is optionally substituted by 1 to 3 substituents selected from halogen atom, hydroxyl group, C_{1-6} alkoxy group, carboxy group, amino group, C_{1-6} alkylamino group, di(C_{1-6})alkylamino group and oxo group, C_{3-7} cycloalkyl group, C_{2-6} alkenyl group, C_{2-6} alkynyl group, aralkyl group, carboxy group, (C_{1-6} alkoxy)carbonyl group or cyano group;

R^3 is hydrogen atom, C_{1-6} alkyl group, hydroxyl group, C_{1-6} alkoxy group, mercapto group, C_{1-6} alkylthio group, carboxy group, (C_{1-6} alkoxy)carbonyl group, (C_{1-6} alkyl)carbamoyl group, di(C_{1-6} alkyl)carbamoyl group, amino group, C_{1-6} alkylamino group or di(C_{1-6})alkylamino group;

R^4 is hydrogen atom, C_{1-6} alkyl group or C_{2-6} alkenyl group, or R^3 and R^4 in combination show oxo group; R^5 and R^6 are the same or different and each is C_{1-6} alkyl group, or R^5 and R^6 in combination show cyclopropyl group together with the carbon atom they bind to;

R^7 is aryl group or heteroaryl group wherein said aryl group and heteroaryl group are optionally substituted by 1 to 3 substituents selected from halogen atom, C_{1-6} alkyl group, halo(C_{1-6})alkyl group, hydroxy(C_{1-6})alkyl group, C_{3-7} cycloalkyl group, hydroxyl group, C_{1-6} alkoxy group, halo(C_{1-6})alkoxy group, carboxy group, (C_{1-6} alkoxy)carbonyl group, nitro group, cyano group, C_{1-6} alkylsulfonyloxy group, carbamoyl group and C_{1-3} alkylenedioxy group;

X^1 is a single bond, C_{1-6} alkylene group or C_{2-6} alkynylene group wherein said C_{1-6} alkylene group and C_{2-6} alkynylene group are optionally substituted by C_{1-6} alkyl group or oxo group;

X^2 is C_{1-6} alkylene group wherein said C_{1-6} alkylene group is optionally substituted by C_{1-6} alkyl group or halo(C_{1-6})alkyl group;


X^3 is a single bond or C_{1-6} alkylene group wherein said C_{1-6} alkylene group is optionally substituted by hydroxyl group or oxo group;

X^4 and X^5 in combination show a single bond, methylene group,

-NH-, oxygen atom, sulfur atom, $-C(=O)-$, $-CH_2NH-$, $-CH_2O-$, $-CH_2S-$, $-CH_2CO-$, $-NHCH_2-$, $-OCH_2-$, $-SCH_2-$, $-COCH_2-$, $-CH=CH-$ or $-C\equiv C-$,

a salt thereof, a solvate thereof or a prodrug thereof.

2. A compound of the formula [I']




wherein

- R^{1'} is aryl group or heteroaryl group wherein said aryl group and heteroaryl group are optionally substituted by 1 to 3 substituents selected from halogen atom, C₁₋₆ alkyl group, halo(C₁₋₆)alkyl group, hydroxy(C₁₋₆)alkyl group, C₁₋₆ alkoxy(C₁₋₆)alkyl group, hydroxyl group, C₁₋₆ alkoxy group, halo(C₁₋₆)alkoxy group, mercapto group, C₁₋₆ alkylthio group, C₁₋₆ alkylsulfanyl group, C₁₋₆ alkylsulfonyl group, aminosulfonyl group, C₁₋₆ alkylsulfamoyl group, di(C₁₋₆)alkylsulfamoyl group, carboxy group, (C₁₋₆ alkoxy)carbonyl group, C₁₋₇ acyl group, carbamoyl group, (C₁₋₆ alkyl)carbamoyl group, di(C₁₋₆ alkyl)carbamoyl group, cyano group, nitro group, amino group, C₁₋₆ alkylamino group, di(C₁₋₆)alkylamino group and C₁₋₇ acylamino group;
- R^{7'} is aryl group or heteroaryl group wherein said aryl group and heteroaryl group are optionally substituted by 1 to 3 substituents selected from halogen atom, C₁₋₆ alkyl group, halo(C₁₋₆)alkyl group, C₃₋₆ cycloalkyl group, hydroxyl group, C₁₋₆ alkoxy group, halo(C₁₋₆)alkoxy group, carboxy group, (C₁₋₆ alkoxy)carbonyl group, nitro group, cyano group and C₁₋₆ alkylsulfonyloxy group; and
- R², R³, R⁴, R⁵, R⁶, X¹, X², X³, X⁴ and X⁵ are same as defined in

claim 1 respectively,


a salt thereof, a solvate thereof or a prodrug thereof.

3. A compound of the formula [I'']



wherein

R^{1''} is aryl group or heteroaryl group wherein said aryl group and heteroaryl group are optionally substituted by 1 to 3 substituents selected from C₁₋₃ alkylenedioxy group,




wherein R^A is (C₁₋₆ alkoxy)carbonyl group or carboxy group and R^B is hydrogen atom or C₁₋₆ alkyl group;

R^{7''} is aryl group or heteroaryl group wherein said aryl group and heteroaryl group are optionally substituted by 1 to 3 substituents selected from hydroxy(C₁₋₆)alkyl group, carbamoyl group and C₁₋₃ alkylenedioxy group; and

R², R³, R⁴, R⁵, R⁶, X¹, X², X⁴ and X⁵ are same as defined in claim 1 respectively,

a salt thereof, a solvate thereof or a prodrug thereof.

4. A compound of the formula [I-2]



wherein R^{11} and R^{12} are the same or different and each is hydrogen atom, halogen atom, C_{1-4} alkyl group, hydroxy (C_{1-6})alkyl group, C_{1-6} alkoxy(C_{1-6})alkyl group, hydroxyl group, C_{1-4} alkoxy group, cyano group or nitro group, or R^{11} and R^{12} in combination show C_{1-3} alkylenedioxy group,

R^{21} is C_{1-4} alkyl group

wherein said alkyl group is optionally substituted by


C_{1-4} alkoxy group,

C_{3-5} cycloalkyl group, C_{2-4} alkenyl group, or aralkyl group, and R^3 , R^4 , R^5 , R^6 , R^7 , X^1 , X^2 , X^3 , X^4 and X^5 are same as defined in

claim 1 respectively,

a salt thereof, a solvate thereof or a prodrug thereof.

5. The compound of claim 4, which is represented by the formula [I-3]



wherein R^{71} and R^{72} are the same or different and each is hydrogen atom, C_{1-4} alkyl group or C_{1-4} alkoxy group, or in combination show $-CH=CH-CH=CH-$ or C_{1-3} alkylenedioxy group, and R^{11} , R^{12} and R^{21} are same as defined in claim 4 respectively, a salt thereof, a solvate thereof or a prodrug thereof.

6. The compound of claim 5, wherein R^{11} and R^{12} are the same or different and each is hydrogen atom, chlorine atom, methyl group, hydroxymethyl group, hydroxyl group, methoxy group, cyano group or nitro group, or R^{11} and R^{12} in combination show methylenedioxy group, and R^{21} is optionally branched C_{1-4} alkyl group or C_{3-5} cycloalkyl group, a salt thereof, a solvate thereof or a prodrug thereof.

7. The compound of claim 6, wherein R^{21} is methyl group, ethyl group, cyclopropyl group or cyclobutyl group, a salt thereof, a solvate thereof or a prodrug thereof.

8. The compound of claim 7, wherein R^{11} is hydrogen atom, R^{12} is methyl group, methoxy group or hydroxymethyl group, or R^{11} and R^{12} in combination show methylenedioxy group, and R^{21} is methyl group or cyclopropyl group, a salt thereof, a solvate thereof or a prodrug thereof.

9. The compound of claim 8, wherein R^{21} is cyclopropyl group, a salt thereof, a solvate thereof or a prodrug thereof.

10. The compound of claim 9, wherein R^{71} and R^{72} in combination show $-CH=CH-CH=CH-$, a salt thereof, a solvate thereof or a prodrug thereof.

11. The compound of claim 9, wherein R^{71} and R^{72} are groups selected from C_{1-4} alkyl group and C_{1-4} alkoxy group, a salt thereof, a solvate thereof or a prodrug thereof.

12. The compound of claim 1, which is selected from

(2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(2-methoxyphenyl)methoxy]propan-2-ol,
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(2-methylphenyl)methoxy]propan-2-ol,
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(phenyl)methoxy]propan-2-ol,
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methoxyphenyl)ethoxy]propan-2-ol,
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methylphenyl)ethoxy]propan-2-ol,
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-methoxyphenyl)propoxy]propan-2-ol,
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(2-cyanophenyl)ethoxy]propan-2-ol,
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(3-methoxyphenyl)ethoxy]propan-2-ol and
 (2R)-1-[1,1-dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[1-(3-methylphenyl)ethoxy]propan-2-ol,

a salt thereof, a solvate thereof or a prodrug thereof.


13. (2R)-1-[1,1-Dimethyl-2-(naphthalen-2-yl)ethylamino]-3-[(cyclopropyl)(2-hydroxymethylphenyl)methoxy]propan-2-ol, a salt thereof, a solvate thereof or a prodrug thereof.

14. A pharmaceutical composition comprising the compound of any of claims 1 to 13, a salt thereof, a solvate thereof or a prodrug thereof as an active ingredient.

15. A calcium receptor antagonist comprising the compound of any of claims 1 to 13, a salt thereof, a solvate thereof or a prodrug thereof as an active ingredient.

16. A therapeutic agent for osteoporosis comprising the compound of any of claims 1 to 13, a salt thereof, a solvate thereof or a prodrug thereof as an active ingredient.


17. A compound of the formula [II]



wherein R^{11'} is halogen atom, C₁₋₄ alkyl group, hydroxy(C₁₋₆)alkyl group, C₁₋₆ alkoxy(C₁₋₆)alkyl group, hydroxyl group, C₁₋₄ alkoxy group, tert-butyldimethylsilyloxymethyl group, cyano group or nitro group, a salt thereof or a solvate thereof.

18. The compound of claim 17, wherein, in the formula [II], R^{11'} is C₁₋₄ alkyl group, hydroxy(C₁₋₆)alkyl group or C₁₋₄ alkoxy group, a salt thereof or a solvate thereof.


19. A compound of the formula [III]



wherein R^{11a} is halogen atom, hydroxy(C_{1-6})alkyl group, C_{1-6} alkoxy(C_{1-6})alkyl group, hydroxyl group, C_{1-4} alkoxy group, tert-butyldimethylsilyloxymethyl group, cyano group or nitro group, a salt thereof or a solvate thereof.

20. The compound of claim 19, wherein, in the formula [III], R^{11a} is C_{1-4} alkoxy group, a salt thereof or a solvate thereof.

21. A compound of the formula [IV]



wherein R^8 is carboxy group, nitro group, tert-butoxycarbonylamino group or benzyloxycarbonylamino group, a salt thereof or a solvate thereof.

FIG. 1




FIG. 2




FIG. 3




FIG. 4





FIG. 5



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP01/06903

A. CLASSIFICATION OF SUBJECT MATTER Int. Cl. ⁷ C07C217/10, C07D213/30, C07D333/16, C07D307/42, C07D277/24, A61K31/135, A61K31/18, A61K31/275, A61K31/4402, A61K31/381, A61K31/341, A61K31/426, A61P43/00, A61P3/14, A61P19/10, A61P1/02, A61P29/00, A61P19/02, C07D317/54, A61K31/36, C07D333/28, C07D333/40, C07D333/56, C07D333/32, C07C205/16, C07C205/09, C07C205/04, C07C43/23, C07C49/84 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int. Cl. ⁷ C07C217/10, C07D213/30, C07D333/16, C07D307/42, C07D277/24, A61K31/135, A61K31/18, A61K31/275, A61K31/4402, A61K31/381, A61K31/341, A61K31/426, A61P43/00, A61P3/14, A61P19/10, A61P1/02, A61P29/00, A61P19/02, C07D317/54, A61K31/36, C07D333/28, C07D333/40, C07D333/56, C07D333/32, C07C205/16, C07C205/09, C07C205/04, C07C43/23, C07C49/84 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAPLUS (STN), REGISTRY (STN)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	ADAMS, Mortimer H. et al., "The Synthesis of New Bronchodilator Prodrugs", Synth. Commun., (1999), Vol.29, No.14, pages 2419 to 2430	1-14 15-16
A	Chem. abstr., Vol.88, (1978), (Columbus, Ohio, USA), page 358, column 2, abstract No.62084(f), IBANEZ, Antonio et al., "New derivatives of 2-diphenylmethoxyethylamine with vasodilator and spasmolytic activity", Eur. J. Med. Chem. Chim. Ther., (1977), 12(5), pages 459 to 462, (Eng.)	1-16
A	GOWEN, Maxine et al., "Antagonizing the parathyroid calcium receptor stimulates parathyroid hormone secretion and bone formation in osteopenic rats", J. Clin. Invest., (2000), Vol.105, No.11, pages 1595 to 1604	1-16
X A	US 5856576 A (American Cyanamid Company), 05 January, 1999 (05.01.99), especially, example 8 & CN 1187485 A	19-20 17-18
☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 16 October, 2001 (16.10.01)		Date of mailing of the international search report 30 October, 2001 (30.10.01)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP01/06903

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	EP 852224 A1 (American Cyanamid Company), 08 July, 1998 (08.07.98), especially, example 8 & JP 10-245367 A	19-20 17-18
X A	EP 682007 A1 (American Cyanamid Company), 15 November, 1995 (15.11.95), especially, example 4 & JP 7-300445 A & US 5414136 A	19-20 17-18
X A	EP 542685 A1 (Ciba-Geigy AG), 19 May, 1993 (19.05.93), especially, page 38, Beispiel 92 & JP 5-294920 A & US 5336663 A	19-20 17-18
X A	US 5684049 A (Janssen Pharmaceutica), 04 November, 1997 (04.11.97), especially, column 15 & EP 538301 A1 & JP 5-507931 A	19-20 17-18
X A	BECK, Anthony L. et al., "Preparation of Tetrahydrobenz [cd]indoles from 1-Tetralones", J. Chem. Soc., Perkin Trans.1, (1990), No.3, pages 689 to 693	19-20 17-18
A	EP 489670 A1 (Rhone-Poulenc Agrochimie), 10 June, 1992 (10.06.92), & FR 2670205 A1 & JP 4-342541 A	17-20
A	Chem. abstr., Vol.95, (1981), (Columbus, Ohio, USA), page 963, column 2, the abstract No.969(j), XU, Xiu-Rong et al., "Relation between radioprotection and estrogenic effect of nonsteroidal estrogens", Yao Hsueh Hsueh Pao, (1980), 15(11), pages 648 to 655, (Ch)	21

Form PCT/ISA/210 (continuation of second sheet) (July 1992)