



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 324 659 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

Note: Bibliography reflects the latest situation

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see page(s) 7,8

(48) Corrigendum issued on:
28.06.2006 Bulletin 2006/26

(45) Date of publication and mention
of the grant of the patent:
14.12.2005 Bulletin 2005/50

(21) Application number: **01977677.2**

(22) Date of filing: **09.10.2001**

(51) Int Cl.:
A01N 59/00 (1985.01) A01N 25/00 (1985.01)

(86) International application number:
PCT/US2001/031669

(87) International publication number:
WO 2002/030195 (18.04.2002 Gazette 2002/16)

(54) **AN AZIDE METHOD AND COMPOSITION FOR CONTROLLING DELETERIOUS ORGANISMS**
AZID-VERFAHREN UND -ZUSAMMENSETZUNG ZUR BEKÄMPFUNG VON SCHADORGANISMEN
METHODE ET COMPOSITION AZIDE POUR LUTTER CONTRE LES ORGANISMES NUISIBLES

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**

(30) Priority: **10.10.2000 US 238943 P**

(43) Date of publication of application:
09.07.2003 Bulletin 2003/28

(73) Proprietor: **Auburn University**
Auburn, AL 36849-5176 (US)

(72) Inventor: **RODRIGUEZ-KABANA, Rodrigo**
Auburn, AL 36830 (US)

(74) Representative: **Becker Kurig Straus**
Patentanwälte
Bavariastrasse 7
80336 München (DE)

(56) References cited:
EP-A- 0 102 924 US-A- 3 583 987
US-A- 3 771 994 US-A- 3 812 254
US-A- 4 132 780

• **DATABASE WPI Derwent Publications Ltd.,**
London, GB; AN 1998-002870, XP002906433
ANONYMOUS: 'Stable liquid immuno-
agglutination reagent' & JP 9 269 326 A 14
October 1997

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 324 659 B9

Description

[0001] The present invention relates generally to the fields of biology and agriculture and more particularly to compositions and methods for use as pesticides and herbicides.

Background

[0002] The control of insects, plant pathogens, nematodes and weeds is of central importance to the agriculture industry. In particular, the substantial reduction or elimination of nematode populations in soils is critical to initial plant growth, productivity and life-span. Pathogenic fungi and nematodes develop on the extensive root systems of both annual and perennial crops, damaging them severely. Moreover, they persist in the soil after crop removal and need to be eliminated before replanting of new crops.

[0003] Approaches which have been used successfully to combat plant pathogens and nematodes have been crop rotation, fallowing for at least four years, use of pathogen and nematode-resistant crops and soil fumigation. Resistance to plant pathogens and nematodes is available only in a few crops, and resistant cultivars may not be developed in the foreseeable future for many crops of significant commercial interest. Therefore, soil fumigation remains the best alternative for control of plant pathogens and nematodes.

[0004] Methyl bromide (CH_3Br) is the most widely used and most universal fumigant in the world. It is used extensively for soil fumigation, as a commodity quarantine treatment (export and imports) to control a variety of pests on numerous crops, and as a structural fumigant for wood destroying pests.

[0005] Methyl bromide (hereinafter referred to as "MBr") is categorized as an ozone depleting chemical with an ozone depleting potential (ODP) of greater than 0.2 compared to trichlorofluoromethane (cfc 11), a refrigerant used as a reference gas having an ODP of 1.

[0006] Evidence on the loss of MBr to the atmosphere after soil fumigation indicates that of the total amount applied to the soil for fumigation, approximately 87% is lost to the atmosphere within seven days. On reaching the stratosphere MBr undergoes photo-oxidation, releasing bromine atoms which enter the ozone depletion cycle. MBr loss from fumigated soils is further supported by studies which indicated a loss of as much as 70% of the applied MBr to the atmosphere through the tarp and after the tarp is removed.

[0007] As currently available alternatives to MBr are less effective and/or more expensive, the removal of MBr will be very costly. Annual losses to U.S. producers and consumers is estimated to be in the region of 1.5 billion dollars. This figure does not account for the losses due to post harvest and quarantine losses as well as structural fumigation losses. MBr removal would most adversely affect such commodities as tomatoes, strawberries, pep-

pers, melons and ornamentals. The loss of MBr would thus be extremely costly to both agricultural producers and consumers as well as having a substantial impact on the U.S. economy. Nonetheless, it is the general consensus of those working in the field that no approach is currently available that will achieve the same level of broad-spectrum pest management as MBr; chemical and non-chemical approaches that are available can provide some level of agricultural pest management, but generally with narrower activity and lower crop yields and quality. Therefore, there is clearly a need for alternatives to MBr.

[0008] Solid compositions for controlling plant pests comprising an ionic azide, a metal salt and a basic azide stabilizing material are described in US 4 132780.

Summary

[0009] The present invention provides both compositions and methods for controlling a population of deleterious organisms in soil without causing significant harm to environment. The deleterious organisms that are the target of this invention include any pests, such as, for example, insects, fungi, nematodes, weeds and any other organism that may adversely affect agricultural endeavors. Such deleterious organisms may be controlled by applying the composition of the present invention : a composition comprising an ionic azide; an azide stabilizer comprising an amine selected from the group consisting of ethanolamine, dimethylamine, ethylamine, butylamine, diethylamine, diethanolamine, and phenylethylamine in an amount effective to stabilize the azide and a liquid dispersal medium. Said composition can be applied to an environmental system to control a population of deleterious organisms. For example, it may be applied to soil, whereby a population of target pest organisms within the soil is controlled. Application of the composition may include applying to a soil an amount of azide effective in controlling the population of one or more targeted pests. The azide contained within the liquid is in the form of ionic azide (N_3^-) and may be derived from any appropriate azide compound. For example, the azide may be sodium azide, potassium azide or a combination of both sodium and potassium azide.

[0010] Azide can be stabilized by the presence of one or more stabilizers including, for example, a detergent, an amine, a protein or a combination thereof. A detergent may also function as a wetting agent thereby improving the delivery of the liquid composition of the present invention to the soil

[0011] The azide may further be stabilized by maintaining the pH of the liquid medium above 7.0. The alkalinity may be achieved by an organic or inorganic pH buffering agent, the addition of hydroxyl ions or a combination thereof.

[0012] The composition of the present invention may serve generally as a pesticide. The composition may be used more specifically as a herbicide, an insecticide, a

fungicide and/or a nematocide or a combination thereof, depending upon the needs of the user.

[0013] The present invention also encompasses a kit for preparing a pesticidal composition, comprising the components of the pesticides composition described above, and instructions for preparing the pesticidal composition and applying the pesticidal composition to an environmental system, such as, for example, soil, to control a population of a targeted deleterious organism therein. The kit may further comprise one or more stabilizers such as, but not limited to, a detergent, an amine, a protein, a pH buffering agent or any combination thereof.

[0014] The invention further provides agricultural compositions described above for controlling a population of deleterious organisms and one or more other components such as, for example, a fertilizer to promote crop growth.

[0015] Additional objects and aspects of the present invention will become more apparent upon review of the detailed description set forth below when taken in conjunction with the accompanying figures, which are briefly described as follows.

Description of the Drawings

[0016] In the drawings:

Figure 1 is a graph illustrating the fruit yield of eggplants grown in soil treated with increasing doses of an aqueous solution of sodium azide;

Figure 2 is a graph illustrating the weed count or infestation in soil treated with increasing doses of an aqueous solution of sodium azide;

Figure 3 is a graph illustrating the count of the nematode *Helicotylanchus dihystra* in soil treated with increasing doses of an aqueous solution of sodium azide;

Figure 4 is a graph illustrating the count of the nematode *Dorylaimida* in soil treated with increasing doses of an aqueous solution of sodium azide;

Figure 5 illustrates a graph comparing both the fresh shoot weight and fresh root weight of cotton grown in soil treated with increasing doses of an aqueous solution of sodium azide;

Figure 6 is a graph illustrating the root condition index of cotton count of cotton grown in soil treated with increasing doses of an aqueous solution of sodium azide;

Figure 7 depicts a graph illustrating the count of the nematode *Meloidogyne incognita* in soil treated with increasing doses of an aqueous solution of sodium azide stabilized in a mixture of amines and proteins; Figure 8 is a graph illustrating the weed count or infestation in soil treated with increasing doses of an aqueous solution of sodium azide stabilized in a mixture of amines and proteins; and

Figure 9 depicts a graph comparing the number of fruits and yield of tomato plants grown in soil treated

with increasing doses of an aqueous solution of sodium azide stabilized in a mixture of amines and proteins.

Detailed Description

[0017] The present invention provides methods and compositions for controlling a population of a deleterious organism such as, but not limited to, an insect, a nematode or a weed or a combination thereof, by applying a composition comprised of a liquid dispersal medium and an azide and an azide stabilizer comprising an amine selected from the group consisting of ethandamine, ethylamine, dimethylamine, butylamine, diethylamine, diethandamine, and phenylethylamine. The liquid medium is both safe and stable for use in irrigation systems, such as in a drip irrigation line and may be used in formulations with plant nutrients and other pest control agents. The azide is an ionic azide. Exemplary ionic azides include, but are not limited to, sodium and potassium salts or any combination thereof. The azide may be produced through the introduction of an azide salt to an aqueous medium. While sodium and potassium azides may be used most commonly, any azide salt, such as, for example, ammonium, calcium, zinc azides, are also appropriate. The compositions of the present invention provide effective nematocides having herbicidal properties, and do not cause significant or permanent harm to the environment.

[0018] The term "controlling" as used herein refers to regulating a population of a deleterious organism that may be harmful to an agricultural product. The population may be regulated by the compositions and methods of the present invention so that the organism is killed, thereby reducing the viable populations such as by nematocidal, fungicidal, herbicidal or insecticidal activity or the like. The methods and compositions of the present invention may maintain and not allow a population of a deleterious organism to increase or may prevent an invasion of a soil by a deleterious organism.

[0019] The term "azide" as used herein refers to any compound having the N_3^- moiety therein. The azide may be a metal azide wherein the metal is an alkali metal such as potassium, sodium, lithium, rubidium or cesium. The metal may be a transition metal such as, but not limited to, iron, cobalt, nickel, copper or zinc. It is understood that certain metallic azides may be formed in solution by mixing sodium azide or the like with a metallic salt such as, for example, copper sulfate. The azide of the present invention may also be an organic azide or ammonium azide.

[0020] The term "amine" as used herein refers to the organic aliphatic amines including, but not limited to, ethanamine, ethylamine, butylamine, diethylamine, dimethylamine and the like. The organic amine may be an aromatic amine such as, for example, phenylethylamine.

[0021] The term "liquid medium" as used herein refers to an aqueous or organic fluid at least partially in the liquid

phase under ambient conditions. Suitable fluids include, but are not limited to, water, an oil, an emulsion, a liquid organic compound such as, for example, ethanolamine, ethanol and the like.

[0022] The term "chelating agent" as used herein refers to any organic or inorganic compound that will bind to a metal ion having a valence greater than one, and includes, but is not limited to, organic chelating agents such as ethylenediaminetetracetic acid (EDTA), triethylene tetramine dihydrochloride (TRIEN), ethylene glycol-bis (beta-aminoethyl ether)-N, N, N', N'-tetracetic acid (EGTA), diethylenetriamin-pentaacetic acid (DPTA), and triethylenetetramine hexaacetic acid (TTG), deferroxamine, Dimercaprol, edetate calcium disodium, zinc citrate, penicillamine succimer and Editronate or any other chelating agent, salt or combination thereof, known to one of ordinary skill in the art, and which will chelate divalent metal ions such as, but not only Ca^{2+} , Mg^{2+} , Mn^{2+} , Fe^{2+} , and Zn^{2+} .

[0023] The term "pH buffering agent" as used herein refers to any organic or inorganic compound or combination of compounds that will maintain the pH of a solution to within about 0.5 pH units of a selected pH value. A "pH buffering agent" may be selected from, but is not limited to, Tris (hydroxymethyl) aminomethane (tromethaprim; TRIZMA base), or salts thereof, phosphates, amino acids, polypeptides or any other pH buffering agent or combination thereof.

[0024] The term "detergent" as used herein refers to an amphipathic compound, either neutral or ionic in nature, which is soluble in water, such as sodium lauryl sulfate.

[0025] As used herein the terms "oligopeptide", "polypeptide" and "protein" refer to a polymer of amino acids of three or more amino acids in a serial array, linked through peptide bonds. The term "polypeptide" includes proteins, protein fragments, protein analogues, oligopeptides and the like. The term "polypeptides" further contemplates polypeptides as defined above that include chemically modified amino acids or amino acids covalently or noncovalently linked to labeling ligands.

[0026] Sodium and potassium azides are typically first formulated as granules (attapulgitic clay, diatomaceous earth) or they can be formed in a variety of liquid formulations. The stability of azide formulations is enhanced at pH levels of greater than about 8.7. For the control of pests such as nematodes and fungi such as *Armillaria*, *Verticillium*, and the protection of deep-rooted crops such as grapes, fruit, and nut trees, liquid formulations are optionally suitable.

[0027] The liquid medium may further include a stabilizer. The delivery of an azide to a desired fumigation zone may be difficult if the reactivity of the azide in the soil-air space and atmosphere is too rapid and results in an effective concentration of the active compound that is too low for pest control. A stabilizer aids in increasing the effectivity of the liquid medium in such conditions. Thus, the liquid medium can be modified depending upon

the soil conditions in which the composition of the present invention is used such as, for example, by increasing the pH of the liquid medium for application to an acidic soil.

[0028] Aqueous azide solutions can be stabilized in an alkaline solution by the addition of pH buffering agents such as, hydroxyl ions such as in the form of sodium or potassium hydroxide or the like, or by adding carbonates and phosphates. Some pH buffering agents may not be desired, however, when used with irrigation waters having, for example, a high calcium ion content. Calcium may combine with solubilized carbonates and/or phosphates of the compositions of the present invention to form calcium carbonate or phosphate precipitates that may corrode or clog irrigation systems. If precipitation is not a concern, any appropriate pH buffering agent may be used, such as, for example, ammonium, sodium or potassium phosphates, ammonium, sodium or potassium carbonates, ammonium, sodium or potassium citrates, acid-salt systems with pH buffering capacity, organic buffers such as Tris and the like, or hydroxyl ions that will maintain, either singly or in combination, a pH in the liquid medium of greater than about 8.7.

[0029] Other stabilizers include a detergent such as, but not limited to, sodium lauryl sulfate; Tween 20 [polyoxyethylene (20) sorbitan monolaurate]; Tween 40 [polyoxyethylene (20) sorbitan monopalmitate]; Tween 40 [polyoxyethylene (20) sorbitan monopalmitate]; Tween 60 [polyoxyethylene (20) sorbitan monostearate]; Tween 80 [polyoxyethylene (20) sorbitan monostearatepalmitate]; Tween 85 [polyoxyethylene (20) sorbitan trioleate]; soaps, (i.e. salts of Na, K or ammonium and various fatty acids [ammonium oleate, potassium stearate, etc.]); and tergitol. By using a detergent, the liquid medium can be stabilized without the problem of forming calcium precipitates. The detergent may further cleanse an irrigation system that is used to apply a composition of the present invention to a soil. It is further contemplated to be within the scope of the present invention for the detergent to have wetting properties that may increase the dispersal of the composition of the present invention in the soil.

[0030] A further stabilizer contemplated for use in the compositions of the present invention and which may be used in the liquid medium formulation is an amine. An amine group of a stabilizer may stabilize the azide-containing formulations. The amine may also provide an additional source of nitrogen for a crop planted in the soil. All types of amine may be used such as, but not limited to, alkyl or aromatic amines. An exemplary amine is ethanolamine.

[0031] The amine stabilizer of the present invention can be an amino acid, an oligopeptide or a polypeptide. The amine groups of the stabilizer can be in the form of a protein that can itself provide a pH buffering effect. For example, the protein such as casein may be used. Casein is soluble in ethanolamine and, when added to soils, stimulates the activities of proteolytic microorganisms. The increased microbial proteolytic activity may further be antagonistic to plant parasitic nematodes and other soil

borne pests. Suitable proteins for use in the compositions of the present invention include, but are not limited to, proteins derived from a cereal meal such as soybean meal, cotton seed meal, maize meal or wheat meal. Other suitable proteins may include zein, gluten, gelatin and the like or protein obtained from animal sources such as whey.

[0032] The present invention is further illustrated by the following examples, which are provided by way of illustration and should not be construed as limiting.

Reference Example 1: Control of nematodes and weeds

[0033] In Example 1, the efficacy of pre-plant applications of NaN_3 for control of plant parasitic nematodes and weeds was studied in a microplot experiment using the aqueous formulation of the chemical. The formulation contained 5% sodium azide dissolved in demineralized water containing 0.5% sodium lauryl sulfate; final pH of the solution was 9.8. The square microplots of 1 ft² [929 cm²] and 2 feet [61 cms] deep having an open-bottom were used. Each microplot was delimited by a terra-cotta chimney flue. The soil filling the plot was a sandy loam [pH= 6.2; organic matter content < 1.0%; C.E.C < 10 meq/100 g soil] and was infested with the nematodes: *Meloidogyne incognita* [root-knot], *Paratrichodorus minor* [stubby root], *Tylenchorhynchus claytoni* [stunt], and *Helicotylenchus dihystra* [spiral]. Weeds introduced into the plots were principally crabgrass [*Digitaria sanguinalis*], yellow nutsedge [*Cyperus esculentum*], pigweed [*Amaranthus* spp.], and morning glories [*Ipomea* spp., *Jacquemontia tamnifolia*].

[0034] Applied to the plots was the aqueous formulation of NaN_3 diluted in irrigation water and delivered by drenching [2L/plot] at amounts of: 0, 0.5, 1.0, 1.5, 2.0, 3.0, 4.0, and 5.0 g sodium azide/plot. The plots were covered with standard polyethylene [1 mil] and after 10 days the cover was removed, the number of weeds was determined, soil samples for nematological analyses were taken and each plot was planted with two 3-week old 'Black Beauty' eggplant [*Solanum melongena*] seedlings. The plants were grown for 2 months and data were collected on yield, weed infestation and nematode populations. There were no nematodes in the planting time samples from plots treated with the solution of NaN_3 . In contrast, soil from control plots were infested with all nematode species. When planted, there were no weeds in plots treated with all but the two lowest rates of NaN_3 . Yield and number of fruits increased directly in response to NaN_3 rates from 0-1.5 g/plot and leveled out with no additional increases obtained in response to higher rates. At the end of the experiment there were no plant parasitic nematodes and no significant weed infestation in plots with the three highest rates of the chemical but there were significant populations of the parasites and weeds in plots treated with ≤ 2.0 g NaN_3 rates. The results of Example 1 are illustrated in Figures 1-4.

Reference Example 2: Control of root knot

[0035] In Example 2, the value of NaN_3 solution for the suppression of the cotton wilt complex [*Fusarium oxysporum f.sp. vasinfectum* x *Meloidogyne incognita*] was investigated in a greenhouse experiment with soil from a field having a severe wilt problem. The soil was apportioned in one kg amounts in 4L plastic bags and was treated with an aqueous formulation of NaN_3 to have rates of : 10, 20, 30, 40, 50, 60, 70, 80, 90, and 100 mgs a.i./kg soil. The aqueous formulation was a solution containing 1% [w/v] NaN_3 in water containing 0.5% sodium lauryl sulfate with pH = 9.8. After thorough mixing, the contents of a bag were poured into 1 L capacity, 10-cm-diam cylindrical plastic pots which were then covered with standard polyethylene [1ml]. Each rate and the no treatment control was represented by 7 replications [pots] arranged in a randomized complete block design. Ten days after application of the chemical solution, the pots were uncovered and 5 seed of "Rowden" cotton [*Gossypium hirsutum*] were planted in each pot. The resulting plants were grown for 8 weeks. When the plants were removed after 8 weeks, soil samples were collected for nematological analyses. The shoots and roots of the plants were weighed and the root systems were examined and indexed for disease severity symptoms. Following examination, the roots were incubated to determine nematode populations.

[0036] The results of the analysis indicated that at all NaN_3 dosages the plant parasitic nematodes were eliminated from the soil and roots. Sharp increases in weights of shoots and roots were recorded in response to NaN_3 application rates in the range 10-60 mgs/kg soil with no additional weight increments being obtained with rates ≥ 70 mgs/kg soil. Cotton root health [root condition index] was markedly improved in response to all NaN_3 dosages, root systems of plants from pots with the 10 mg-rate were as healthy looking as those from pots treated with all other dosages of the compound. Results indicate that NaN_3 may be useful for suppression of Fusarium wilt complex and that rates required for wilt control are below those needed for broad-spectrum herbicidal activity. The results of Example 2 are further illustrated in Figures 5 and 6.

Example 3: Amine-protein stabilizers in the control of nematodes and weeds

[0037] Example 3 illustrates the efficacy of pre-plant applications of an aqueous formulation of NaN_3 for control of plant parasitic nematodes and weeds in a microplot experiment with tomato [*Lycopersicon esculentum*]. NaN_3 was stabilized in the formulation by means of a mixture of amines and the commercially available protein, casein. The formulation contained 5% [w/v] NaN_3 dissolved in demineralized water containing 2% [v/v] of a 10% [w/v] casein solution in ethanolamine. Soil in the microplots was a sandy loam [pH= 6.2; organic matter

content < 1.0%; C.E.C < 10 meq/100 g soil] infested with the nematodes: *Meloidogyne incognita*, *Paratrichodorus minor*, *Tylenchorhynchus claytoni*, and *Helicotylenchus dihystra* 1]. Weeds in the plots were principally crabgrass [*Digitaria sanguinalis*], yellow nutsedge [*Cyperus esculentum*], pigweed [*Amaranthus* spp.], and morning glories [*Ipomea* spp., *Jacquemonlia tamnifolia*].

[0038] NaN_3 was applied to the 1-ft² microplots by drenching [2L/plot] at amounts of: 0, 0.5, 1.0, 1.5, 2.0, 3.0, 4.0, and 5.0 g/plot. The plots were covered with standard polyethylene [1 mil] tarp immediately after application of the chemical. After 10 days the cover was removed, the number of weeds was determined, soil samples for nematological analyses were taken and each plot was planted with two 4-week old 'Huskie' tomato seedlings. The plants were grown for 3 months and data were collected on yield, number of fruits, weed infestation and nematode populations.

[0039] Soil samples from plots treated with NaN_3 did not contain nematodes, but the soil samples from the control plots had significant numbers of all nematode species. At planting time, there were no weeds in plots treated with all but the three lowest rates of NaN_3 . Yield and number of fruits increased directly in response to NaN_3 rates from 0- 4 g/plot. At the end of the experiment there were no plant parasitic nematodes in NaN_3 -treated plots for all rates of application.

[0040] Final weed counts were inversely related to NaN_3 rate in a pattern described [$R^2=0.95^{**}$] by $W=-0.32X+5.82$, where W represents total weed population density/microplot and X, azide rates in gms NaN_3 /plot. Results indicate that the amine-protein formulation is superior for nematode control and equal in herbicidal activity to formulations stabilized with inorganic buffers. The data also suggest that it is possible to deliver NaN_3 into soil together with organic compounds that can serve as plant nutrients and stimulate beneficial soil microbial activities. The results of Example 3 are illustrated in Figures 7-9.

Reference Example 4: Control of root-knot nematode and weeds in green peppers and tomatoes

[0041] Example 4 illustrates the efficacy of pre-plant applications of NaN_3 for control of root-knot nematode [*Meloidogyne incognita*], coastal bermudagrass [*Cynodon dactylon*], yellow nutsedge [*Cyperus esculentum*] and other weeds in pepper [*Capsicum annum*] and tomato [*Lycopersicon esculentum*]. The soil was a calcareous silty clay loam with pH 7.8 and < 1% organic matter. The NaN_3 was applied at drenching rates of 100 and 200 kg/ha using a solution of sodium azide in sodium lauryl sulfate. Each dosage rate was delivered into pre-acidified soil as well as in non-acidified and administered using 3 different water levels: 3, 10, and 15 L/m². Acidification was with H_2SO_4 to lower soil pH to less than 7.00. The soil was covered with standard polyethylene tarp immediately after NaN_3 application. After 3 weeks the soil was

uncovered and a soil sample was removed for nematological analyses and the weeds were counted. Tomato and pepper seedlings were then transplanted 6 weeks after NaN_3 application.

[0042] Azide applications effectively controlled greater than 90% of the nematodes and all weeds at dosages of 100 and 200 kgs. The compound was particularly effective against nutsedge. NaN_3 performed well when drenched with any of the 3 water levels. There was no evidence of phytotoxicity to pepper or tomato plants in any of the plots treated with NaN_3 .

Reference Example 5: Nematicidal and Herbicidal Properties of Potassium Azide

[0043] In this Example, the nematotoxic properties of liquid potassium azide KN_3 are illustrated using soil from a cotton field infested with the reniform nematode (*Rotylenchulus reniformis*). The liquid potassium azide compound was added to the soil in an aqueous solution of amounts of: 1, 2, 3, 4 and 5 mgs KN_3 per kg of soil. The formulation used for this experiment 1% KN_3 and 0.5% sodium lauryl sulfate. Soil samples were collected one week after the application of the KN_3 solution for nematological analysis using such techniques as the salad bowl incubation technique. From the analysis it was shown that the numbers of reniform nematode declined exponentially in response to the increasing KN_3 . Rates of 4-5 mg of KN_3 per kg of soil showed an almost 100% control of the reniform nematodes. Numbers of microbivorous nematodes declined in an almost linear fashion in response to the increasing dosages of KN_3 .

[0044] In the application of KN_3 rates of 20-200 mg/kg of soil applied to a soil infested with crab grass (*Digitaria sanguinalis*), purple nutsedge (*Cyperus rotundus*), Jimson weed (*Datura stramonium*) and a variety of other weeds resulted in the number of weeds declining in proportion to rates used. Rates of greater than 140 mg/kg of soil resulted in over 80% control rates for weeds.

Claims

1. A composition for controlling a population of a deleterious organism comprising:
 - an ionic azide;
 - an azide stabilizer comprising an amine selected from the group consisting of ethanolamine, dimethylamine, ethylamine, butylamine, diethylamine, diethanolamine, and phenylethylamine in an amount sufficient to stabilize the azide; and
 - a liquid dispersal medium.
2. 4. The composition of claim 1, wherein the liquid dispersal medium contains water.
3. The composition of any of claims 1 or 2, wherein the

azide is selected from the group consisting of a metal azide, an organic azide and combinations thereof.

4. The composition of claim 3, wherein the metal azide is an alkali metal azide selected from the group consisting of potassium azide, sodium azide, lithium azide, and combinations thereof.

5. The composition of claim 1, wherein the azide is ammonium azide.

6. The composition of any of claims 1-5, wherein the azide stabilizer further comprises a compound is selected from the group consisting of detergents, pH buffering agents, alkalis, amines, amino acids, oligopeptides, polypeptides and combinations thereof.

7. The composition of claim 6, wherein the detergent is selected from the group consisting of polyoxyethylene (20) sorbitan monolaurate, polyoxyethylene (20) sorbitan monopalmitate, polyoxyethylene (20) sorbitan monostearate, polyoxyethylene (20) sorbitan monostearatepalmitate, polyoxyethylene (20) sorbitan trioleate, sodium lauryl sulfate and combinations thereof.

8. The composition of claim 6, wherein the pH buffering agent is selected from the group consisting of tris (hydroxymethyl)aminomethane, ammonium phosphate, sodium phosphate, potassium phosphate, ammonium carbonate, sodium carbonate, potassium carbonate, ammonium citrate, sodium citrate, potassium citrate, and combinations thereof.

9. The composition of claim 6, wherein the polypeptide is a protein selected from the group consisting of proteins derived from a cereal meal, zein, gluten, gelatin, casein, and proteins obtained from whey.

10. A method of controlling a population of a deleterious organism, in soil, comprising the step of applying to a soil a composition of any of claims 1-9, wherein the amount of azide in the soil is effective for controlling a population of a deleterious organism therein.

11. The method of claim 10, wherein the amount of azide in the soil effective for controlling the population of a deleterious organism therein is in the range of about 1 to about 200 mg per kg of soil.

12. A kit for preparing a pesticidal composition comprising packaging and having therein

the components of a pesticidal composition of any of claims 1-9, and instructions for preparing the pesticidal composition and applying the pesticidal composition to

a soil, thereby reducing a population of deleterious organisms therein.

5 Patentansprüche

1. Zusammensetzung zum Steuern einer Population eines schädlichen Organismus, umfassend:

ein ionisches Azid;
einen Azid-Stabilisator, umfassend, ein Amin ausgewählt aus der Gruppe bestehend aus Ethanolamin, Dimethylamin, Ethylamin, Butylamin, Diethylamin, Diethanolamin und Phenylethylamin in einer zur Stabilisierung des Azids ausreichenden Menge; und
ein flüssiges Dispersionsmedium.

2. Zusammensetzung nach Anspruch 1, worin das flüssige Dispersionsmedium Wasser umfasst.

3. Zusammensetzung nach einem der Ansprüche 1 oder 2, worin das Azid ausgewählt ist der Gruppe bestehend aus einem Metallazid, einem organischen Azid und Kombinationen davon.

4. Zusammensetzung nach Anspruch 3, worin das Metallazid ein Alkalimetallazid ist, ausgewählt aus der Gruppe bestehend aus Kaliumazid, Natriumazid, Lithiumazid und Kombinationen davon.

5. Zusammensetzung nach Anspruch 1, worin das Azid Ammoniumazid ist.

6. Zusammensetzung nach einem der Ansprüche 1-5, worin der Azid-Stabilisator weiterhin eine Verbindung umfasst, ausgewählt aus der Gruppe bestehend aus Detergentien, pH puffernden Agenzien, Alkali, Aminen, Aminosäuren, Oligopeptiden, Polypeptiden und Kombinationen davon.

7. Zusammensetzung nach Anspruch 6, worin das Detergens ausgewählt ist aus der Gruppe bestehend aus Polyoxyethylen (20) Sorbitanmonolaurat, Polyoxyethylen (20) Sorbitanmonopalmitat, Polyoxyethylen (20) Sorbitanmonostearat, Polyoxyethylen (20) Sorbitanmonostearatpalmitat, Polyoxyethylen (20) Sorbitantrioleat, Natriumlaurylsulfat und Kombinationen davon.

8. Zusammensetzung nach Anspruch 6, worin das pH puffernde Agens ausgewählt ist aus der Gruppe bestehend aus Tris(hydroxymethyl)aminomethan, Ammoniumphosphat, Natriumphosphat, Kaliumphosphat, Ammoniumcarbonat, Natriumcarbonat, Kaliumcarbonat, Ammoniumcitrat, Natriumcitrat, Kaliumcitrat und Kombinationen davon.

9. Zusammensetzung nach Anspruch 6, worin das Polypeptid ein Protein ist, ausgewählt aus der Gruppe bestehend aus Proteinen erhalten aus einem Getreidemehl, Zein, Gluten, Gelatine, Kasein und aus Molke erhältlichen Proteinen.

10. Verfahren zum Steuern einer Population eines schädlichen Organismus in einem Boden, umfassend den Schritt eines Anwendens einer Zusammensetzung gemäß den Ansprüchen 1-9 auf einen Boden, worin die Menge an Azid in dem Boden wirksam zum Steuern einer Population eines darin befindlichen schädlichen Organismus ist.

11. Verfahren nach Anspruch 10, worin die Menge an Azid in dem Boden, die wirksam zum Steuern der Population eines darin befindlichen schädlichen Organismus ist, in dem Bereich von ungefähr 1 bis ungefähr 200 mg pro kg Boden liegt.

12. Kit zur Herstellung einer schädlingsbekämpfenden Zusammensetzung, umfassend Verpackung und darin aufweisend

die Bestandteile einer schädlingsbekämpfenden Zusammensetzung nach einem der Ansprüche 1-9, und Anweisungen zum Herstellen der schädlingsbekämpfenden Zusammensetzung und Anwenden der schädlingsbekämpfenden Zusammensetzung auf einen Boden, wodurch eine Population eines darin befindlichen schädlichen Organismus verringert wird.

Revendications

1. Une composition pour contrôler une population d'un organisme nuisible comprenant :

un azide ionique ;
un stabilisateur d'azide comprenant une amine choisie parmi le groupe consistant en éthanolamine, diméthylamine, éthylamine, butylamine, diéthylamine, diéthanolamine et phényléthylamine en une quantité suffisante pour stabiliser l'azide ; et
un milieu de dispersion liquide.

2. La composition de la revendication 1, dans laquelle le milieu de dispersion liquide contient de l'eau.

3. La composition de l'une quelconque des revendications 1 ou 2, dans laquelle l'azide est choisi parmi le groupe consistant en un azide métallique, un azide organique et des combinaisons de ceux-ci.

4. La composition de la revendication 3, dans laquelle

l'azide métallique est un azide de métal alcalin choisi parmi le groupe consistant en azide de potassium, azide de sodium, azide de lithium et des combinaisons de ceux-ci.

5. La composition de la revendication 1, dans laquelle l'azide est l'azide d'ammonium.

6. La composition de l'une quelconque des revendications 1 à 5, dans laquelle le stabilisateur d'azide comprend en outre un composé choisi parmi le groupe consistant en détergents, agents tampons de pH, alcalis, amines, acides aminés, oligopeptides, polypeptides et des combinaisons de ceux-ci.

7. La composition de la revendication 6, dans laquelle le détergent est choisi parmi le groupe consistant en sorbitane-monolaurate de polyoxyéthylène (20), sorbitane-monopalmitate de polyoxyéthylène (20), sorbitane-monostéarate de polyoxyéthylène (20), sorbitane-monostéarate-palmitate de polyoxyéthylène (20), sorbitane-trioléate de polyoxyéthylène (20), laurylsulfate de sodium et des combinaisons de ceux-ci.

8. La composition de la revendication 6, dans laquelle l'agent tampon de pH est choisi parmi le groupe consistant en tri(hydroxyméthyl)aminoéthane, phosphate d'ammonium, phosphate de sodium, phosphate de potassium, carbonate d'ammonium, carbonate de sodium, carbonate de potassium, citrate d'ammonium, citrate de sodium, citrate de potassium et des combinaisons de ceux-ci.

9. La composition de la revendication 6, dans laquelle le polypeptide est une protéine choisie parmi le groupe consistant en des protéines dérivées d'une farine de céréale, zéine, gluten, gélatine, caséine et des protéines obtenues à partir du petit lait.

10. Une méthode pour contrôler une population d'un organisme nuisible dans le sol, comprenant l'étape d'appliquer à un sol une composition de l'une quelconque des revendications 1 à 9, dans laquelle la quantité d'azide dans le sol est efficace pour contrôler une population d'un organisme nuisible qui s'y trouve.

11. La méthode de la revendication 10, dans laquelle la quantité d'azide dans le sol efficace pour contrôler la population d'un organisme nuisible qui s'y trouve est dans le domaine d'environ 1 à environ 200 mg par kg de sol.

12. Un kit pour préparer une composition pesticide comprenant un emballage et ayant à l'intérieur

les composants d'une composition pesticide de

l'une quelconque des revendications 1 à 9, et des instructions pour préparer la composition pesticide et appliquer la composition pesticide au sol, réduisant de ce fait une population d'organismes nuisibles qui s'y trouve.

5

10

15

20

25

30

35

40

45

50

55

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5





Figure 6



INDEX SCALE: 1= BEST TO 5= WORST LOOKING ROOT SYSTEM

Figure 7




figure 8




Figure 9

