



(11) **EP 1 338 604 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

- (15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Description Paragraph(s) 10, 11, 13, 15
Claims EN
- (48) Corrigendum issued on:
21.05.2008 Bulletin 2008/21
- (45) Date of publication and mention of the grant of the patent:
12.07.2006 Bulletin 2006/28
- (21) Application number: **01972113.3**
- (22) Date of filing: **05.10.2001**
- (51) Int Cl.:
C07K 5/08^(2006.01) A61K 38/06^(2006.01)
A61P 25/04^(2006.01)
- (86) International application number:
PCT/ES2001/000369
- (87) International publication number:
WO 2002/028885 (11.04.2002 Gazette 2002/15)

(54) **N-ALKYLGLYCINE TRIMERES CAPABLE OF PROTECTING NEURONS AGAINST EXCITOTOXIC AGGRESSIONS AND COMPOSITIONS CONTAINING SAID TRIMERES**

N-ALKYLGLYCINTRIMERES, DAS DAZU IN DER LAGE IST, NEURONEN GEGEN EXCITOTOXISCHE AGGRESSIONEN ZU SCHÜTZEN, UND ZUSAMMENSETZUNGEN, DIE DIESES TRIMERE ENTHALTEN

TRIMERES DE N-ALKYLGLYCINE POUVANT PROTEGER LES NEURONES CONTRE LES AGRESSIONS EXCITOTOXIQUES, ET COMPOSITIONS LES CONTENANT

- (84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE TR
- (30) Priority: **06.10.2000 ES 200002414**
- (43) Date of publication of application:
27.08.2003 Bulletin 2003/35
- (73) Proprietor: **Diverdrugs, S.L.**
08850 Gava, Barcelona (ES)
- (72) Inventors:
 - **FELIPO ORTS, Vicente,**
Edif. Vanguard
E-08902 Barcelona (ES)
 - **MONTOLIU FELIX, Carmen,**
Edif. Vanguard
E-08902 Barcelona (ES)
 - **FERRER MONTIEL, Antonio,**
Edif. Vanguard
E-08902 Barcelona (ES)
 - **PLANELLS CASES, Rosa,**
Edif. Vanguard
E-08902 Barcelona (ES)
 - **MERINO FERNANDEZ, Jaime M.,**
Edif. Vanguard
E-08902 Barcelona (ES)
- (74) Representative: **Maldonado Jordan, Julia**
Linares, 7
46018 Valencia (ES)
- (56) References cited:
WO-A-96/40747
 - **GIBBONS J.A. ET AL.: 'Pharmacologic characterization of CHIR 2279, an N-sustituted glycine peptoid with high-affinity binding for alpha1 adrenoceptors' J. PHARMACOL. EXP. THER. vol. 277, no. 2, 1996, pages 885 - 889, XP002953307**

EP 1 338 604 B9

- GOODSON B. ET AL.: 'Characterization of novel antimicrobial peptoids' *ANTIMICROB. AGENTS CHEMOTHER.* vol. 43, no. 6, 1999, pages 1429 - 1434, XP002906265
- KIRSHENBAUM K. ET AL.: 'Sequence specific polypeptoids: A diverse family of heteropolymers with stable secondary structure' *PROC. NATL. ACAD. SCI. USA* vol. 95, no. 8, pages 4303 - 4308, XP002944445
- ZUCKERMANN R.N. ET AL.: 'Discovery of nanomolar ligands for 7-transmembrane G-protein-coupled receptors from a diverse N (substituted) glycine peptoid library' *J. MED. CHEM.* vol. 37, 1994, pages 2678 - 2685, XP000608720

Description**SCOPE OF THE INVENTION**

5 **[0001]** This invention refers to N-alkylglycine trimers capable of protecting neurones against excitotoxic aggressions, useful as neuroprotectors, to compositions containing them and to their employment in the treatment of illnesses or disorders mediated by the excitotoxicity.

BACKGROUND OF THE INVENTION

10 **[0002]** Neurodegenerative illnesses and neurological disorders constitute a serious social and economic problem. Clear examples are represented by senile dementia, Alzheimer, Huntington, and the one associated to the AIDS virus, as well as the neurodegeneration caused by ischemia associated to a cerebrovascular accident. In spite of the seriousness of the problem, the pharmacological arsenal to fight, prevent, and/or decrease its symptoms and progress, is surprisingly limited.

15 **[0003]** Though the biological mechanisms that lead to the neurodegeneration are not clearly established, in many neurodegenerative illnesses, such as amiotrophic lateral sclerosis, dementia associated to AIDS and to Alzheimer, the presence of high and chronic levels of the L-glutamate excitotoxic neurotransmitter (1-3), has been observed in the cerebral parenchyma. This neurotransmitter has also been involved in the etiology of neurological disorders such as cerebral ischemia (4). The glutamate activates membrane receptors that have an ionic channel activity (ionotropic receptors) or that transduce the signal through G proteins (metabotropic receptors) (5). The ionotropic receptors, especially those of the NMDA type [N-methyl-D-aspartate activated glutamate receptors (NMDA)], have been involved in the glutamatergic neurodegeneration due to their high Ca^{2+} ion permeability (1-5). The proposed molecular mechanism indicates that high and chronic levels of glutamate cause a prolonged activation (hyperactivation) of the NMDA receptor that "overloads" the neurones with Ca^{2+} ions, triggering off the massive activation and excessive intracellular cascades that, inevitably lead to neuronal death (1-8). In fact, it has been described that antagonists of this receptor are capable of preventing the glutamate neurotoxicity (8,9). From what is expounded it can be deduced, that a strategy for preventing or decreasing the neurodegeneration is to control the functional activity of this ionotropic receptor, especially, under conditions in which a high pathology of the glutamate levels exist.

25 **[0004]** In spite of the advance made in the past few years, potent, selective and toxicity-free neuroprotectors have not yet been developed. Up to the moment, a large part of the effort has been focused, towards the development of competitive inhibitors that recognise the glutamatergical receptors of the central nervous system (1,2). For example, an important effort has been made to develop competitive and non competitive antagonists of glutamate and/or glycine [a coagonist that participates in the activation of the NMDA type glutamate]. These molecules, though powerful neuroprotectors, present important secondary effects, such as cognitive anomalies, that limit their clinical use (10-12). The main disadvantage of using competitive and non competitive antagonist is that they interact with their receptors, non specifically inhibiting the neurotransmission, and affecting both the pathological activity of the glutamate and its physiological activity (13). A strategy to overcome this therapeutic obstacle would be to use non competitive and/or acompetitive antagonists that preferably join the agonists-receptor complex. The most important advantage of using this type of antagonists is that these agents mainly act on hyperactivated receptors (pathological receptors), showing a marginal interaction over receptors that perform on rapid excitatory neurotransmission processes (physiological receptors) (13). This preferred activity over the "pathological" receptors makes these types of antagonists valued as promising therapeutic agents to prevent the neurodegeneration (13-18). Molecules such as phencyclidine and dizolcipine are powerful acompetitive antagonists of the NMDA receptor that act as efficient in vitro neuroprotectors (12-18). However, their clinical use is questioned due to the psycotomimetic effects (13).

SUMMARY OF THE INVENTION

50 **[0005]** The invention faces the problem of searching for new neuroprotector compounds capable of preventing, decreasing or treating neurodegeneration, preferably, the excitotoxic, that overcome totally or partly the previously indicated disadvantages.

[0006] The solution provided by this invention is based on the development of N-alkylglycine trimers that are capable of blocking the ionotropic glutamate receptors and that may be used to prevent, decrease or treat neurodegeneration, as well as prevent or treat illnesses or disorders mediated by the neurodegeneration. The capacity of said N-alkylglycine trimers for blocking the NMDA type glutamate ionotropic receptors as well as their capacity to prevent excitotoxic neuronal death caused by a prolonged exposure to L-glutamic in the absence or presence of glycine, has been shown by means of the tests described in Example 1.2.

[0007] Consequently, an object of this invention is constituted by N-alkylglycine trimers capable of blocking the


glutamate ionotropic receptors, useful for blocking the excitotoxic aggressions response.

[0008] An additional object of this invention is constituted by a composition that comprises at least one of said N-alkylglycine trimers, such as a pharmaceutical composition.

[0009] The employment of said N-alkylglycine trimers in the elaboration of a medicine for the prevention or treatment of illnesses or disorders mediated by the neurodegeneration constitutes another additional object of this invention.

DETAILED DESCRIPTION OF THE INVENTION

[0010] The invention provides general formula (I) N-alkylglycine trimers



where,

R_1 , R_2 , R_3 , equal or different, with independence from each other, are selected from among cyclopropyl, *sec*-butyl, 2-methoxyethyl, 3-methylbutyl, cyclohexyl, 2-(*N*-pyrrolidinyl) ethyl, 2-(methylcarbonylamine)ethyl, 3-(2-oxo-*N*-pyrrolydiny) propyl, 2-(2-pyridyl)ethyl, 2-phenylethyl, 1-(2-tetrahydrofuryl)methyl, 2-(*N*-imidazolyl)ethyl, 2-(9-metoxyphenyl)ethyl, 2-(3,4-dimetoxyphenyl)ethyl, 2-(2,4-dichlorophenyl)ethyl, 2-[2-(*N*-methyl)pyrrolidinyl]ethyl, 2-(4-aminosulfonylphenyl) ethyl, 2-(morpholine)ethyl, 3-(*N,N*-diethylamine)propyl, 3,3-diphenylpropyl, 3-(*N,N*-dimethylamine)propyl and 2-(*N,N*-diethylamine) ethyl,

[0011] Their stereoisometric forms and mixtures, racemic or non racemic, of the same, and their pharmaceutically acceptable salts.

[0012] Some formula (I) N-alkylglycine trimers may have one or more chiral centres. Consequently, said formula (I) compounds may exist under any of their stereoisomeric forms (enantiomeric or diastereoisomeric) or in mixtures, racemic or non racemic, of the same, all of which fall within the scope of the present invention. Illustrative examples of formula (I) N-alkylglycine trimers that contain chiral centres include those that contain *sec*-butyl groups [(*R,S*)-*sec*-butyl, (*R*)-*sec*-butyl or (*S*)-*sec*-butyl]; 1-(2-tetrahydrofuryl) methyl [(*R,S*)-1-(2-tetrahydrofuryl)methyl, (*R*)-1-(2-tetrahydrofuryl) methyl or (*S*)-1-(2-tetrahydrofuryl) methyl]; or 2-[2-(*N*-methyl)pyrrolidinyl] ethyl, [(*R,S*)-2-[2-(*N*-methyl)pyrrolidinyl]ethyl, (*R*)-2-[2-(*N*-methyl) pyrrolidinyl] ethyl or (*S*)-2-[2-(*N*-methyl)pyrrolidinyl] ethyl].

[0013] In a particular embodiment, the invention provides formula (I) N-alkylglycine trimers in which the R_1 is *sec*-butyl, 2-phenylethyl, *N*-acetamidoethyl, *N,N*-dimethylaminopropyl, or 2-(*N*-imidazolyl)ethyl, and their stereoisomeric forms and mixtures, racemic or non racemic; of the same.

[0014] In another particular embodiment, the invention provides formula (I) N-alkylglycine trimers in which R_2 is cyclopropyl, *N,N*-diethylaminopropyl, 2-(morpholine)ethyl or 3,3-diphenylpropyl and its stereoisomeric forms and mixtures, racemic or non racemic, of the same.

[0015] In another particular embodiment, the invention provides formula (I) N-alkylglycine trimers in which R_3 is 2-(*N*-pyrrolidyl)ethyl, cyclopropyl, 3,3-diphenylpropyl or *N*-methylpyrrolidyl-2-ethyl, and its stereoisomeric forms and mixtures, racemic or non racemic, of the same.

[0016] In another particular embodiment, the invention provides an N-alkylglycine trimere that additionally contains a reversible modification with the purpose of increasing its bioavailability and ease of passage of the hematoencephalic barrier and epithelial tissue.

[0017] Illustrative examples of formula (I) N-alkylglycine trimeres provided by this invention are gathered in Table 1 [Example 1.1].

[0018] Within the scope of this invention, are to be found, the pharmaceutically acceptable salts of the formula (I) N-alkylglycine trimeres provided by this invention. The term "pharmaceutically acceptable salts" includes the salts normally

used to form metallic salts or salts with acid additives. The nature of the salt is not critical, provided it is pharmaceutically acceptable. The pharmaceutically acceptable salts of formula (I) N-alkylglycine trimers may be obtained as from acids, organic or inorganic. Said salts may be obtained by conventional techniques, well known to experts in the art, reacting the appropriate acid with the formula (I) N-alkylglycine trimers.

5 [0019] Formula (I) N-alkylglycine trimers can be obtained by conventional methods, for example, by means of copolymerization and amidation reactions, among the glycine derivatives that constitute the structural components of said trimers. The glycine derivatives can in turn be obtained by means of conventional methods of amino acid modification. The stereoisomeric forms of the formula (I) N-alkylglycine trimers can be synthesised as from the corresponding enantiomers or from racemic or non racemic mixtures of the raw products. When parting from enantiomeric mixtures, the
10 obtained stereoisomers can be separated by conventional methods of stereoisomeric resolution (enantiomers and diastereoisomers), for example, fractionated crystallisation, chromatography or salts formation.

[0020] Formula (I) N-alkylglycine trimers are capable of blocking the glutamate ionotropic receptors and can be used to prevent, decrease or treat neurodegeneration, preferably, the neurodegeneration by excitotoxicity, as well as to prevent or treat illnesses or disorders mediated by neurodegeneration. These formula (I) N-alkylglycine trimers are also capable
15 of preventing excitatory neuronal death caused by prolonged exposure of neuronal cultures to L-glutamine.

[0021] The capacity of formula (I) N-alkylglycine trimers to prevent excitotoxic neurodegeneration can be demonstrated by means of a test that evaluates the efficiency and power of said N-alkylglycine trimers abating neuronal death caused by prolonged exposure of primary cultures of rat cerebellum neurones to L-glutamine and glycine [see Example 1.2]. The neuroprotector mechanism implies, at least in part, the blocking of the glutamate ionotropic receptors, as is
20 made manifest by the inhibition of the ionic current activated by the NMDA type of glutamatergic receptor agonist expressed in *X. laevis oocytes* (19,20). An important advantage of these biological tests is that they permit the search for neuroprotectors in functionally active systems, increasing the potentiality of its *in vivo* use.

[0022] Formula (I) N-alkylglycine trimers can form part of diverse types of compositions for application in the body of mammals, preferably human beings. In this sense, the invention provides a composition that comprises at least one
25 formula (I) N-alkylglycine trimer. In a particular embodiment, said composition is a pharmaceutical composition.

[0023] The pharmaceutical composition provided by this invention comprises a therapeutically effective amount of at least one formula (I) N-alkylglycine trimer together with at least, one pharmaceutically acceptable excipient.

[0024] Formula (I) N-alkylglycine trimers can be administered in order to treat the neurodegeneration, by any means that produces contact of the formula (I) N-alkylglycine trimers with the site of action thereof in the body of a mammal,
30 preferably a human being.

[0025] The amount of therapeutically efficient formula (I) N-alkylglycine trimers that must be administered as well as its dose for treating a pathological condition mediated by neurodegeneration shall depend on numerous factors, including the age, condition of the patient, severity of the alteration or disorder, the route and frequency of the administration and the particular formula (I) N-alkylglycine trimer to be used.

[0026] The pharmaceutical composition provided by this invention may be presented under any administrative form, for example, solid or liquid, and may be administered in any appropriate way, for example, orally, parenterally, rectal or
35 topic, for which it shall include the necessary pharmaceutically acceptable excipients for the formulation of the desired administrative form. A revision of the different pharmaceutical forms of administration of the medicines and of the necessary excipients for their obtention may, for example, be found in the "Treaty of Galenic Pharmacy", C.Fauli i Trillo, 1993, Luzán 5, S.A. Ediciones, Madrid.

[0027] Consequently, an additional object of this invention is constituted by the employment of a formula (I) N-alkylglycine trimer in the elaboration of a medicine for the attenuation of the nervous activity of neurones involved in the neurodegeneration mediated by the application of exogenous chemical substances or by the endogenous liberation of
40 chemical substances that cause excitotoxicity in the nervous system (excitotoxic or excitotoxic substances) or in the elaboration of a medicine that inhibits the ionic channels activated by exogenous chemical substances or by excitotoxins that lead to neurodegeneration.

[0028] More specifically, the invention refers to the use of a formula (I) N-alkylglycine trimer in the elaboration of a medicine for the treatment of illnesses and pathological alterations mediated by the activity of the ionic channels of the L-glutamate ionotropic receptors, such as type NMDA receptors, for example, the neurodegeneration in response to a
50 noxious stimulus.

[0029] More specifically, the invention refers to the use of a formula (I) N-alkylglycine trimer in the elaboration of a medicine for the treatment, slowing down, reduction, decrease and/or prevention of neurodegeneration, as well as the use of a formula (I) N-alkylglycine trimer in the elaboration of a medicine for the treatment of cerebral ischemia, cerebrovascular accident, migraine, depression, Huntington, Parkinson, Alzheimer, senile dementia, epilepsy and multiple
55 and amiotropic sclerosis.

[0030] The invention additionally provides a method for the treatment in a patient of illnesses and pathological disorders mediated by the activity of the ionic channels and of the glutamate ionotropic receptors, for example, the neurodegeneration mediated by glutamate ionotropic receptors in response to diverse noxious stimuli, for example, mechanical,

chemical and thermal, that comprise the administration to said patient, suffering from said illness or pathological disorder, of a therapeutically effective amount of a formula (I) N-alkylglycine trimere, preferably, in the form of a pharmaceutical composition that contains it.

[0031] The amount of formula (I) N-alkylglycine trimeres to be administered shall depend on numerous factors, among which is to be found the degree of neurodegeneration produced by the excitotoxic aggressions and on the N-alkylglycine trimere to be used.

[0032] On the other hand, a fundamental requirement for the identification of bioactive molecules is to perform a test that permits the determination of its biological activity on the therapeutic targets. The inventors have developed a biological test that permits the evaluation of the power of the molecules that block the ionic current activated by agonist in *X. laevis* oocytes that express neuronal receptors, such as glutamatergical receptors (19, 20). An important advantage of this biological test is that it permits the search for antagonists in functionally active receptors, increasing the power of its use *in vivo*.

[0033] The receptor heterologous expression methods in *X. laevis* oocytes have been described in detail by Ferrer-Montiel and Montal (19).

[0034] The following examples serve to illustrate the nature of the present invention and shall not be considered in its limitative sense.

EXAMPLE 1

N-alkylglycine trimeres capable of blocking the NMDA receptor

1.1 N-alkylglycine trimeres synthesis

[0035] The N-alkylglycine trimeres identified in Table 1 were synthesised by means of conventional methods of solid phase peptide synthesis (21). The trimeres were purified by means of high resolution liquid chromatography.

Table 1 Formula (I) N-alkylglycine trimeres

[1]:	[N-[3,3-diphenylpropyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[Z-(methylcarbonylamine)ethyl]glycinamide;
[2]:	[N-[3,3-diphenylpropyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(2-pyridyl)ethyl]glycinamide;
[3]:	[N-[3,3-diphenylpropyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(N-imidazolyl)ethyl]glycinamide;
[4]:	[N-[3,3-diphenylpropyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[3-(N,N-dimethylamine)propyl]glycinamide;
[5a]:	[N-[(R,S)-2-[2-(N-methyl)pyrrolidinyl]ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(methylcarbonylamine)ethyl]glycinamide;
[5b]:	[N-[(R)-2-[2-(N-methyl)pyrrolidinyl]ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(methylcarbonylamine)ethyl]glycinamide;
[5c]:	[N-[(S)-2-[2-(N-methyl)pyrrolidinyl]ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(methylcarbonylamine)ethyl]glycinamide;
[6a]:	[N-[(R,S)-2-[2-(N-methyl)pyrrolidinyl]ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(2-pyridyl)ethyl]glycinamide;
[6b]:	[N-[(R)-2-[2-(N-methyl)pyrrolidinyl]ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(2-pyridyl)ethyl]glycinamide;
[6c]:	[N-[(S)-2-[2-N-[2-(N-methyl)pyrrolidinyl]ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(2-pyridyl)ethyl]glycinamide;
[7a]:	[N-[(R,S)-2-[2-[N-methyl]pyrrolidinyl]ethyl]glycyl]-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(N-imidazolyl)ethyl]glycinamide;
[7b]:	[N-[(R)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(H-imidazolyl)ethyl]glycinamide;
[7c]:	[N-[(S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(N-imidazolyl)ethyl]glycinamide;
[8a]:	[N-[(R,3)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[3-(N,N-dimethylamine)propyl]glycinamide;
[8b]:	[N-[(R)-2-(2-(N-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl)-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[3-(N,N-dimethylamine)propyl]glycinamide;

(continued)

- [8c]: [N-[(S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[3-N,N-dimethylamine)propyl] glycineamide;
- 5 [9a]: [N-[(R,S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[2-(methylcarbonylamine)ethyl]glycineamide;
- [9b]: [N-[(R)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[2-(methylcarbonylamine)ethyl]glycineamide;
- [9c]: [N-[(S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[2-(methylcarbonylamine)ethyl]glycineamide;
- 10 [10a]: [N-[(R,S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-N-[3,3-diphenylpropyl]glycyl]-N-[2-(pyridyl)ethyl]glycineamide;
- [10b]: [N-[(R)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[2-(2-pyridyl)ethyl]glycineamide;
- 15 [10c]: [N-[(S)-2-(2-N-methyl)pyrrolidinyl]ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-(2-(2-pyridyl)ethyl)glycineamide;
- [11a]: [N-[(R,S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[2-(N-diphenylpropyl)glycyl]-N-[2-(N-imidazolyl)ethyl]glycineamide;
- [11b]: [N-[(R)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[2-(N-imidazolyl)ethyl]glycineamide;
- 20 [11c]: [N-[(S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[2-(N-imidazolyl)ethyl]glycineamide;
- [12a]: [N-[(R,S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[3-(N,N-dimethylamine)propyl]glycineamide;
- 25 [12b]: [N-[(R)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[3-(N,N-dimethylamine)propyl]glycineamide;
- [12c]: [N-[(S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[3-(N,N-dimethylamine)propyl]glycineamide;
- 30 [13a]: [N-(cyclopropyl)glycyl]-[N-(cyclopropyl)glycyl]-N-((R,S)-*sec*-butyl)glycineamide;
- [13b]: [N-(cyclopropyl)glycyl]-[N-(cyclopropyl)glycyl]-N-((R)-*sec*-butyl)glycineamide;
- [13c]: [N-(cyclopropyl)glycyl]-[N-(cyclopropyl)glycyl]-N-((S)-*sec*-butyl)glycineamide.
- [14]: [N-(cyclopropyl)glycyl]-[N-(cyclopropyl)glycyl]-N-(phenethyl)glycineamide;
- [15]: [N-(cyclopropyl)glycyl]-N-(cyclopropyl)glycyl]-N-[2-(4-aminosulfonylphenyl)ethyl]glycineamide;
- 35 [16a]: [N-(cyclopropyl)glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-((R,S)-*sec*-butyl)glycineamide;
- [16b]: [N-(cyclopropyl)glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-((R)-*sec*-butyl)glycineamide;
- [16c]: [N-(cyclopropyl)glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-((S)-*sec*-butyl)glycineamide;
- [17]: [N-(cyclopropyl)glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-(phenethyl)glycineamide;
- 40 [18]: [N-(cyclopropyl)glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-[2-(4-aminosulfonylphenyl)ethyl]glycineamide;
- [19a]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-(cyclopropyl)glycyl]-N-((R,S)-*sec*-butyl)glycineamide;
- [19b]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-(cyclopropyl)glycyl]-N-((R)-*sec*-butyl)glycineamide;
- [19c]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-(cyclopropyl)glycyl]-N-((5)-*sec*-butyl)glycineamide;
- 45 [20]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-(cyclopropyl)glycyl]-N-(phenethyl)glycineamide
- [21]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-(cyclopropyl)glycyl]-[N-[2-(4-aminosulfonylphenyl)ethyl]glycyl]-N-((R,S)-*sec*-butyl)glycineamide;
- [22a]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-((R,S)-*sec*-butyl)glycineamide;
- [22b]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-((R)-*sec*-butyl)glycineamide;
- 50 [22c]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-((S)-*sec*-butyl)glycineamide ;
- [23]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-(phenethyl)glycineamide and
- [24]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-(2-(9-aminosulfonylphenyl)ethyl)glycineamide

55 1.2. Evaluation of the biological activity

[0036] Two complementary biological tests were used in order to evaluate the biological activity of the N-alkylglycine trimers obtained in Example 1.1. The first evaluates the efficiency and power with which said compounds block the

ionic current activated by agonists in *X.laevis* oocytes that express the NMDA receptor. The second, determines the neuroprotector efficiency of the N-alkylglycine trimers in primary cultures of neurones subjected to an excitotoxic aggression, such as prolonged exposure to L-glutamate and glycine.

5 [0037] The heterologous expression of receptors in *X. laevis* oocytes can be performed according to the procedure described by Ferrer-Montiel and Montal (19). Briefly, the *X. laevis* oocytes of adult frogs are gathered, manipulated and injected with cDNA that codes NR1 and NR2A sub-units of NMDA receptor (5,20). The ionic currents activated by the L-glutamate agonist in the presence of the glycine coagonist are recorded with the maintenance of the constant voltage method with two microelectrodes (two-microelectrode voltage clamp) (19, 20). The oocytes that express the receptor are transferred to the recording chamber and are perfused using an 8 output perfusion system. The agonist and the coagonist, both in the absence and in the presence of the N-alkylglycine trimer to be tested, are dissolved in a Ringer buffer (Hepes 10 mM pH 7,4 NaCl 100 mM, BaCl₂ 2,0 mM, RCl 3,0 mM) supplemented with 100 μM fluphenamic acid (20). This buffered solution is used to minimize the contribution of the chloride endogenous ionic channel activated by calcium to the ionic current of the glutamatergic receptor (19, 20). The transmembrane voltage is maintained constant at -80 mV, and the ionic currents are activated by applying pulses of the L-glutamate/glycine solution (100 μM/20 μM) in the presence or absence of growing concentrations of the N-alkylglycine trimers to be tested. The inhibiting activity is detected by measuring the ionic current activated by the agonist in the presence and absence of the N-alkylglycine trimers.

10 [0038] The power and efficiency of the inhibiting activity of the N-alkylglycine trimers is determined by means of the obtention of dose-response curves. For this, the magnitude of the ionic channel blocking activated by L-glutamate/glycine in oocytes that express the NMDA receptor in the presence of growing concentrations of the N-alkylglycine trimer is examined. The ratio of ionic current intensities in the presence and absence of said N-alkylglycine trimer is used to obtain the dose-response curves (19, 20). These graphs meet logarithmic functions to determine the maximum blockage (power) and the antagonist concentration that is produced by half the maximum blockage (IC₅₀, efficiency)

15 [0039] The results obtained showed that all the tested N-alkylglycine trimers blocked the ionic channel activity that is characteristic of the NMDA receptor expressed in *X. laevis* of frog oocytes. The N-alkylglycine trimer concentrations that inhibited half the NMDA receptor response (IC₅₀) activated by L-glutamate/glycine at a concentration of 100 μM / 20 μM, oscillated between 0,1 μM and 100 μM for the different N-alkylglycine trimers tested (Table 1). Thus, for example, the IC₅₀ of the N-alkylglycine trimer identified as compound [2] is of approximately 2 μM and of approximately 10 μM for the N-alkylglycine trimer identified as compound [3]. The rest of the compounds showed IC₅₀ values comprised between 10 and 100 μM.

20 [0040] Additionally, it was observed that the N-alkylglycine trimers tested did not behave as competitive antagonists of the natural agonist.

25 [0041] It was also observed that the N-alkylglycine trimers tested protected primary neuronal cultures coming from the cerebellum of rats pre-exposed to 10 μM glycine versus the prolonged exposure (≤3 h) of L-glutamate or N-methyl-D-aspartate at a concentration of 1 mM. The test consists of extracting the cerebellar neurones of baby rats, 7-8 days old, cultivating them in plates in a culture medium supplemented with bovine foetal serum (22). After 13-19 days under culture, the excitotoxic aggression with L-glutamate is performed, monitoring the neuronal death in the absence and presence of the N-alkylglycine trimers 24 hours post-aggression using two fluorescent dyes: the fluorescent diacetate that indicates the number of viable cells, and the propidium iodine that informs of the number of dead cells (23). As an example, the N-alkylglycine trimer identified as compound [1] decreased a 95% of the neuronal death at a concentration of 30 μg/ml; compound [5] an 85% at 30 μg/ml; compound [23a] a 57% at a concentration of 50 μg/ml, and compound [24] a 60% at 50 μg/ml. The rest of the N-alkylglycine trimers tested, showed neuronal protection efficiencies that oscillated between 37% and 96%.

30 [0042] Additionally, it was observed that the N-alkylglycine trimers tested showed a glutamate neurotoxicity protecting effect in animal models. A neurotoxicity model was used on mice induced by the injection of high doses of ammonium, the result of which is the death of the animal due to the hyperactivation of the NMDA receptor. The test consists in the intraperitoneal injection in mice of an amount of N-alkylglycine trimer of 0,05-0,1 mg/g, followed, after 10 minutes, with an injection of 12 mmol/kg of ammonium acetate. After 24 hours post-injection, the number of dead mice was determined. The N-alkylglycine trimer identified as compound [24] protected 82% of the mice, whilst the N-alkylglycine trimer identified as compound [23a] protected 100% of the animals.

35 [0043] On the other hand, due to the fact that some formula I N-alkylglycine trimers may be presented as stereoisomers, the possibility that the neuroprotector activity of the racemic mixture preferably corresponded to one of the isomers, was evaluated. For this, the neuroprotector activity of the enantiomers identified as [23b] and [23c] was evaluated, (see Table 1) in primary cultures of cerebellar neurons exposed to excitotoxic aggression, such as the incubation with L-glutamate 1mM comparing it with the neuroprotector activity of a racemic mixture of said enantiomers. The results obtained were similar in both cases and showed that both enantiomers are neuroprotectors with equal power. Likewise, said power was similar to the one presented by the racemic mixture.

40 [0044] Given the role of the NMDA receptor in the excitatory synaptic transmission and in the etiology of the numerous

neurodegenerative illnesses, a consequence of the blocker activity of the N-alkylglycine trimers provided by this invention is their use as useful neuroprotectors to decrease or reduce the neuronal death caused by excitotoxic aggression, such as cerebral ischemia, the neurodegeneration mediated by hyperammonemia, Alzheimer dementia, epilepsy, etc.


5 BIBLIOGRAPHY

[0045]

1. Olney, J.W. (1990). Excitotoxic amino acid and neuropsychiatric disorders. *Annu. Rev. Pharmacol. Toxicol.* 30, 47-71.
2. Lipton, S.A. and Rosenburg, P.A. (1994). Excitatory amino acids as a final common pathway for neurologic disorders. *New Engl. J. Med.* 330, 613-622.
3. Choi, D.W., and Rothmann, S.M. (1990). The role of glutamate neurotoxicity in hypoxic ischemic neuronal death. *Annu. Rev. Neurosci.* 13, 171-182.
4. Collingridge, G.L. and Lester, R.A.J. (1989). Excitatory amino acid receptors in the vertebrate central nervous system. *Pharmacol. Rev.* 40, 143-210.
5. Choi, D.W. (1994). Calcium and excitotoxic neuronal injury. *Ann. N. Y. Acad. Sci.* 747, 162-171.
6. Siesjö, B.K. (1994). Calcium-mediated processes in neuronal degeneration. *Ann. N.Y. Acad. Sci.* 747, 140-161.
7. Lipton, S.A. (1994). AIDS-related dementia and calcium homeostasis, *Ann. N.Y. Acad. Sci.* 747, 205-224.
8. Montoliu, C., Llansola, M., Kosenko, E., Corbalán, R., Felipo, V. (1999). Role of cyclic GMP in glutamate neurotoxicity in primary cultures of cerebellar neurons. *Neuropharmacology* 38, 1883-1891.
9. Kornhuber, J. and Quack, G. (1995). Cerebrospinal fluid and serum concentrations of the N-methyl-D-aspartate (NMDA) receptor antagonist memantine in man. *Neurosci. Lett.* 195, 137-139.
10. Nakanishi, S. and Masu, M. (1994). Molecular diversity and functions of glutamate receptors (1994) *Annu. Rev. Biophys. Biomol. Struc.* 23, 319-348.
11. Herrling, P.L. (1994). Clinical implications of NMDA receptor. In the NMDA receptor, Eds G.L. Collingridge and J.C. Watkins, 2nd Ed. 376-394.
12. Lipton, S.A. (1993) Prospects for clinically tolerated NMDA antagonists: open channel blockers and alternative redox states of nitric oxide. *Trends neurosci.* 16, 527-532.
13. Müller, W.E., Mutschler, E. and Riederer, P. (1995). Non-competitive NMDA receptor antagonists with fast open channel blocking kinetics and strong voltage dependency as potential therapeutic agents for Alzheimer's. *Pharmacopsychiat* 28, 113-124.
14. Iversen, L.L. and Kemp, J.A. (1994). Non-competitive NMDA antagonists as drugs. In: the NMDA receptor. Eds. G.L. Collingridge and J.C. Watkins. 2nd Ed. 469-486.
15. Miñana, M.D., Hermenegildo, C., Llansola, M., Montoliu, C., Grisolia, S. and Felipo, V. (1996) Carnitine and choline derivatives containing a trimethylamine group prevent ammonia toxicity in mice and glutamate toxicity in primary cultures of neurones. *J. Pharmacol. Exp. Ther.* 279, 194-199.
16. Koek, W., Woods, J.H. and Winger, G.D. (1988). MK-801, a proposed noncompetitive antagonist of excitatory amino acid neurotransmission, produces phencyclidine-like behavioral effects in pigeons, rats and rhesus monkeys. *J. Pharmacol. Exp. Ther.* 245, 969-974.
17. Choi, D.W. (1987) Dextrorphan and dextromethorphan attenuate glutamate neurotoxicity. *Brain res.* 403, 333-336.
18. Seif el Nasr, M., Peruche, B., Rossberg, C., Mennel, H.-D. and Kriegelstein, J. (1990). Neuroprotective effect of memantine demonstrated in vivo and in vitro. *Eur. J. Pharmacol.* 185, 19-24.
19. Ferrer-Montiel, A.V. and Montal, M. (1994). Structure-function relations in ligand-gated ion channels: Reconstitution in lipid bilayers and heterologous expression in *Xenopus* oocytes. *Methods: a Companion to Methods in Enzymology* 6, 60-69.
20. Ferrer-Montiel, A.V., Sun, W. and Montal, M. (1995). Molecular design of the NMDA receptor binding site for PCP and MK-801. *Proc. Natl. Acad. Sci. USA.* 92, 8021-8025.
21. (A) Suckermann, R.N., Kerr, J.M., Kent, S.B.H., Moos, W.H. (1992). Efficient method for the preparation of peptoids [oligo(N-substituted glycines)] by submonomer solid-phase synthesis. *J. Am. Chem. Soc.* 114, 10646-10647. (B) Figliozzi, G.M., Goldsmith, R., Ng, S.C., Banville, S.C., Zuckermann, R.N. (1996) Synthesis of N-substituted glycine peptoid libraries. *Method Enzymol.* 267, 437-447.
22. Minana M.D., Montoliu, C., Llansola, M., Grisolia, S., Felipo, V. Nicotine prevents glutamate-induced proteolysis of the microtubule-associated protein MAP-2 and glutamate neurotoxicity in primary cultures of cerebellar neurons. *Neuropharmacology* 37, 847-857.
23. Maracaida, G., Miñana, M.D., Burgal, M., Grisolia, S. and Felipo, V. (1995) Ammonia prevents activation of NMDA receptors in rat cerebellar neuronal cultures. *Eur. J. Neurosci.* 7, 2389-2396.

Claims

1. A N-alkylglycine trimers of general formula (I)



where

R₁, R₂, R₃, equal or different, independent from each other, are selected from among cyclopropyl, *sec*-butyl, 2-methoxyethyl, 3-methylbutyl, cyclohexyl, 2-(N-pyrrolidiny)ethyl, 2-(methylcarbonylamine) ethyl, 3-(2-oxo-N-pyrrolidiny) propyl, 2-(2-pyridyl) ethyl, 2-phenylethyl, 1-(2-tetrahydrofuryl) methyl, 2-(N-imidazolyl) ethyl, 2-(4-methoxyphenyl) ethyl, 2-(3,4-dimethoxyphenyl) ethyl, 2-(2,4-dichlorophenyl) ethyl, 2-[2-(N methyl) pyrrolidiny] ethyl, 2-(4-aminosulphonylphenyl) ethyl, 2-(morpholine) ethyl, 3-(N,N-diethylamine) propyl, 3,3-diphenylpropyl, 3-(N,N-dimethylamine) propyl and 2-(N,N-diethylamine) ethyl.

its stereoisomeric forms and mixtures, racemic or non racemic, the same, and their pharmaceutically acceptable salts.

2. An N-alkylglycine trimers according to claim 1, in which, R₁ is *sec*-butyl, 2-phenylethyl, 2-(methylcarbonylamine) ethyl, N,N-dimethylaminopropyl, or 2-(N-imidazolyl)ethyl, its stereoisomeric forms and mixtures, racemic or non racemic, of the same.
3. An N-alkylglycine trimers according to any of the claims 1 or 2, in which R₂ is cyclopropyl, N, N-diethylaminopropyl, 2-(morpholine)ethyl or 3,3-diphenylpropyl, its stereoisomeric forms and mixtures, racemic or non racemic, of the same.
4. An N-alkylglycine trimer according to any of the claims 1 to 3, in which R₃ is 2-(N-pyrrolidiny)ethyl, cyclopropyl, 3, 3-diphenylpropyl or 2-[2-(N-methyl)pyrrolidiny] ethyl, its stereoisomeric forms and mixtures, racemic or non racemic, or the same.
5. An N-alkylglycine trimer according to claim 1, selected from among;

[1]: [N-[3,3-diphenylpropyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(methylcarbonylamine)ethyl]glycinamide;

[2]: [N-[3,3-diphenylpropyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(2-pyridyl)ethyl]glycinamide;

[3]: [N-[3,3-diphenylpropyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(N-imidazolyl)ethyl]glycinamide ;

[4]: [N-(3,3-diphenylpropyl)glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[3-(N,N-dimethylamine)propyl]glycinamide;

[5a]: [N-[(R,S)-2-[2-(N-methyl)pyrrolidiny]ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(methylcarbonylamine)ethyl]glycinamide,

[5b]: [N-[(R)-2-[2-(N-methyl)pyrrolidiny]ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(methylcarbonylamine)ethyl]glycinamide;

[5c]: [N-[(S)-2-[2-(N-methyl)pyrrolidiny]ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(methylcarbonylamine)ethyl]glycinamide;

[6a]: [N-[(R,S)-2-[2-(N-methyl)pyrrolidiny]ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(2-pyridyl)ethyl]glycinamide ;

[6b]: [N-[(R)-2-[2-(N-methyl)pyrrolidiny]ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(2-pyridyl)ethyl]glycinamide;

[6c]: [N-[(S)-2-[2-(N-methyl)pyrrolidiny]ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(2-pyridyl)ethyl]glycinamide;

[7a]: [N-[(R,S)-2-(2-(N-methyl)pyrrolidiny)ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(N-imida-

zoly)ethyl]glycinamide;
 [7b]: [N-[(R)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(N-imidazolyl)ethyl]glycinamide;
 [7c]: [N-[(S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[2-(N-imidazolyl)ethyl]glycinamide;
 5 [8a]: [N-[(R,S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[3-(N,N-dimethylamine)propyl]glycinamide;
 [8b]: [N-[(R)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[3-(N,N-dimethylamine)propyl]glycinamide;
 10 [8c]: [N-[(S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3-(N,N-diethylamine)propyl]glycyl]-N-[3-(N,N-dimethylamine)propyl]glycinamide;
 [9a]: [N-[(R,S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[2-(methylcarbonylamine)ethyl]glycinamide;
 [9b]: [N-[(R)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[2-(methylcarbonylamine)ethyl]glycinamide;
 15 [9c]: [N-[(S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[2-(methylcarbonylamine)ethyl]glycinamide;
 [10a]: [N-[(R,S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[2-(pyridyl)-ethyl]glycinamide;
 20 [10b]: [N-[(R)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[2-(2-pyridyl)ethyl]glycinamide;
 [10c]: [N-[(S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[2-(2-pyridyl)ethyl]glycinamide;
 [11a]: [N-[(R,S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[2-(N-imidazolyl)ethyl]glycinamide;
 25 [11b]: [N-[(R)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[2-(N-imidazolyl)ethyl]glycinamide;
 [11c]: [N-[(S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[2-(N-imidazolyl)ethyl]glycinamide;
 30 [12a]: [N-[(R,S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[3-(N,N-dimethylamine)propyl]glycinamide;
 [12b]: [N-[(R)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[3-(N,N-dimethylamine)propyl]glycinamide ;
 [12c]: [N-[(S)-2-(2-(N-methyl)pyrrolidinyl)ethyl]glycyl]-[N-[3,3-diphenylpropyl]glycyl]-N-[3-(N,N-dimethylamine)propyl]glycinamide ;
 35 [13a]: [N-(cyclopropyl)glycyl]-[N-(cyclopropyl)glycyl]-N-[(R,S)-*sec*-butyl]glycinamide;
 [13b]: [N-(cyclopropyl)glycyl]-[N-(cyclopropyl)glycyl]-N-[(R)-*sec*-butyl]glycinamide;
 [13c]: [N-(cyclopropyl)glycyl]-[N-(cyclopropyl)glycyl]-N-[(S)-*sec*-butyl]glycinamide;
 [14]: [N-(cyclopropyl)glycyl]-[N-(cyclopropyl)glycyl]-N-(phenethyl)glycinamide;
 40 [15]: [N-(cyclopropyl)glycyl]-[N-(cyclopropyl)glycyl]-N-[2-(4-aminosulfonylphenyl)ethyl]glycinamide;
 [16a]: [N-(cyclopropyl)glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-[(R,S)-*sec*-butyl]glycinamide;
 [16b]: [N-(cyclopropyl)glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-[(R)-*sec*-butyl]glycinamide;
 [16c]: [N-(cyclopropyl)glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-[(S)-*sec*-butyl]glycinamide;
 [17]: [N-(cyclopropyl)glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-(phenethyl)glycinamide;
 45 [18]: [N-(cyclopropyl)glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-[2-(4-aminosulfonylphenyl)ethyl]glycinamide;
 [19a]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-(cyclopropyl)glycyl]-N-[(R,S)-*sec*-butyl]glycinamide;
 [19b]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-(cyclopropyl)glycyl]-N-[(R)-*sec*-butyl]glycinamide;
 [19c]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-(cyclopropyl)glycyl]-N-[(S)-*sec*-butyl]glycinamide;
 [20]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-(cyclopropyl)glycyl]-N-(phenethyl)glycinamide;
 50 [21]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-(cyclopropyl)glycyl]-N-[2-(4-aminosulfonylphenyl)ethyl]glycinamide;
 [22a]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-[(R,S)-*sec*-butyl]glycinamide;
 [22b]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-[(R)-*sec*-butyl]glycinamide;
 [22c]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-[(S)-*sec*-butyl]glycinamide;
 [23]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-(phenethyl)glycinamide and
 55 [24]: [N-[2-(N-pyrrolidinyl)ethyl]glycyl]-[N-[2-(N-morpholine)ethyl]glycyl]-N-[2-(4-aminosulfonylphenyl)ethyl]glycinamide.

6. A pharmaceutical composition that comprises a therapeutically efficient amount of at least one formula (I) N-alkylg-

lycine trimere according to any of the claims 1 to 5 and at least one pharmaceutically acceptable excipient.

- 5 7. Use of a formula (I) N-alkylglycine trimere, according to any of the claims 1 to 5, in the elaboration of a medicine for the attenuation of the nervous activity of neurones involved in the neurodegeneration mediated by the application of exogenous chemical substances, or by the endogenous liberation of chemical substances that cause excitotoxicity in the nervous system.
- 10 8. Use of a formula (I) N-alkylglycine trimere, according to any of the claims 1 to 5, in the elaboration of a medicine that inhibits the ionic channels that are activated by exogenous chemical substances or by endogenous .chemical substances, that lead to the neurodegeneration.
- 15 9. Use of a formula (I) N-alkylglycine trimere, according to any of the claims 1 to 5, in the elaboration of a medicine for the treatment of illnesses and pathological disorders mediated by the activity of L-glutamate ionotropic receptors.
- 20 10. Use of a formula (I) N-alkylglycine trimere, according to any of the claims 1 to 5, in the elaboration of a medicine for the treatment, slowing down, reduction, decrease and/or prevention of neurodegeneration.
11. Use of a formula (I) N-alkylglycine trimere, according to any of the claims 1 to 5, in the elaboration of a medicine for the treatment of cerebral ischemia, cerebrovascular accident, migraine, depression, Huntington, Parkinson, Alzheimer, senile dementia, epilepsy and multiple and amiotropic sclerosis.

Patentansprüche

- 25 1. Ein N-Alkylglycin Trimer einer allgemeinen Formel (I) hierbei werden R1, R2, R3, gleich oder unterschiedlich, unabhängig voneinander ausgewählt aus Cyclopropyl, Sec-Butyl, 2-Methoxyethyl, 3-Methylbutyl, Cyclohexyl, 2-(N-Pyrrolidiny)Ethyl, 2-(Methylcarbonylamin) Ethyl, 3-(2-Oxo-N-Pyrrolidiny) Propyl, 2-(2-Pyridyl) Ethyl, 2-Phenylethyl, 1-(2-Tetrahydrofuryl) Methyl, 2-(N-Imidazolyl) Ethyl, 2-(4-Methoxyphenyl) Ethyl, 2-(3,4-Dimethoxyphenyl) Ethyl, 2-(2,4-Dichlorophenyl) Ethyl, 2-[2-(N Methyl) Pyrrolidiny] Ethyl, 2-(4-Aminosulphonylphenyl) Athyl, 2-(Morpholin) Ethyl, 3-(N,N-Diethylamine) Propyl, 3,3-Diphenylpropyl, 3-(N,N-Dimethylamin) Propyl und 2-(N,N-Diethylamin) Ethyl.
Seine stereoisomerischen Formen und Mischungen, razemisch oder nicht razemisch, die Gleichen, ihre pharmazeutisch akzeptable Salze.
- 30 2. Ein N-Alkylglycin Trimer gemäß Anspruch 1, hier ist R1 Sec-Butyl, 2-Phenylethyl, 2-(Methylcarbonylamine)Ethyl, N,N-Dimethylaminopropyl, oder 2-(N-Imidazolyl)Ethyl, seine stereoisomerischen Formen und Mischungen, razemisch oder nicht razemisch, des Gleichen.
- 35 3. Ein N-Alkylglycin Trimer gemäß einem beliebigen der Ansprüche 1 oder 2, hier ist R2 Cyclopropyl, N,N-Dimethylaminopropyl, 2-(Morpholin)Ethyl oder 3,3-Diphenylpropyl, seine stereoisomerischen Formen und Mischungen, razemisch oder nicht razemisch, des Gleichen.
- 40 4. Ein N-Alkylglycin Trimer gemäß einem beliebigen der Ansprüche 1 bis 3, hier ist R3 2-(N-Pyrrolidiyl)Ethyl, Cyclopropyl, 3,3-Diphenylpropyl oder 2-[2-(N-Methylpyrrolidiny)] Ethyl, seine stereoisomerischen Formen und Mischungen, razemisch oder nicht razemisch, oder die Gleichen.
- 45 5. Ein N-Alkylglycin Trimer gemäß Anspruch 1, ausgewählt aus:
- [1]: [N-[3,3-Diphenylpropyl]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[2-(Metholcarbolylamin)Ethyl]Glycinamid;
- 50 [2]: [N-[3,3-Diphenylpropyl]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[2-(2-Pyridyl)Ethyl]Glycinamid;
- [3]: [N-[3,3-Diphenylpropyl]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[2-(N-imidazolyl)Ethyl]Glycinamid;
- [4]: [N-[3,3-Diphenylpropyl]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[3-(N,N-Diethylamin)Propyl]Glycinamid;
- 55 [5a]: [N-[(R,S)-2-[2-(N-Methyl)Pyrrolidiny]Ethyl]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[2-(Methylcarbonylamin)Ethyl]Glycinamid;
- [5b]: [N-[(R)-2-[2-(N-Methyl)Pyrrolidiny]Ethy]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[2-(Methylcarbonylamin)Ethyl]Glycinamid;

- [5c]: [N-[(S)-2-[2-(N-Methyl)Pyrrolidinyl]Ethyl]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[2-(Methylcarbonylamino)Ethyl]Glycinamid;
- [6a]: [N-[(R,S)-2-[2-(N-Methyl)Pyrrolidinyl]Ethyl]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[2-[2-Pyridyl]Ethyl]Glycinamid;
- 5 [6b]: [N-[(R)-2-[2-(N-Methyl)Pyrrolidinyl]Ethyl]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[2-[2-Pyridyl]Ethyl]Glycinamid;
- [6c]: [N-[(S)-2-[2-(N-Methyl)Pyrrolidinyl]Ethyl]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[2-[2-Pyridyl]Ethyl]Glycinamid;
- 10 [7a]: [(R,S)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[2-(N-Imidazolyl)Ethyl]Glycinamid;
- [7b]: [N-[(R)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[2-(N-Imidazolyl)Ethyl]Glycinamid;
- [7c]: [N-[(S)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[2-(N-Imidazolyl)Ethyl]Glycinamid;
- 15 [8a]: [N-[(R,S)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[3-(N,N-Dimethylamin)Propyl]Glycinamid;
- [8b]: [N-[(R)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[3-(N,N-Dimethylamin)Propyl]Glycinamid;
- [8c]: [N-[(S)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3-(N,N-Diethylamin)Propyl]glycyl]-N-[3-(N,N-Dimethylamin)Propyl]Glycinamid;
- 20 [9a]: [N-[(R,S)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3,3-Diphenylpropyl]glycyl]-N-[2-Methylcarbonylamino)Ethyl]Glycinamid;
- [9b]: [N-[(R)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3,3-Diphenylpropyl]glycyl]-N-[2-Methylcarbonylamino)Ethyl]Glycinamid;
- 25 [9c]: [N-[(S)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3,3-Diphenylpropyl]glycyl]-N-[2-Methylcarbonylamino)Ethyl]Glycinamid;
- [10a]: [N-[(R,S)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3,3-Diphenylpropyl]glycyl]-N-[2-Pyridyl]Ethyl]Glycinamid;
- [10b]: [N-[(R)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3,3-Diphenylpropyl]glycyl]-N-[2-(2-Pyridyl)Ethyl]Glycinamid;
- 30 [10c]: [N-[(S)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3,3-Diphenylpropyl]glycyl]-N-[2-(2-Pyridyl)Ethyl]Glycinamid;
- [11a]: [N-[(R,S)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3,3-Diphenylpropyl]glycyl]-N-[2-N-Imidazolyl)Ethyl]Glycinamid;
- 35 [11b]: [N-[(R)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3,3-Diphenylpropyl]glycyl]-N-[2-N-Imidazolyl)Ethyl]Glycinamid;
- [11c]: [N-[(S)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3,3-Diphenylpropyl]glycyl]-N-[2-N-Imidazolyl)Ethyl]Glycinamid;
- [12a]: [N-[(R,S)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3,3-Diphenylpropyl]glycyl]-N-[3-(N,N-Dimethylamin)Propyl]Glycinamid;
- 40 [12b]: [N-[(R)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3,3-Diphenylpropyl]glycyl]-N-[3-(N,N-Dimethylamin)Propyl]Glycinamid;
- [12c]: [N-[(S)-2-(2-(N-Methyl)Pyrrolidinyl)Ethyl]Glycyl]-[N-[3,3-Diphenylpropyl]glycyl]-N-[3-(N,N-Dimethylamin)Propyl]Glycinamid;
- 45 [13a]: [N-(Cyclopropyl)Glycyl]-[N-(Cyclopropyl)Glycyl]-N-(R,S)-sec-Butyl)Glycinamid;
- [13b]: [N-(Cyclopropyl)Glycyl]-[N-(Cyclopropyl)Glycyl]-N-(R)-sec-Butyl)Glycinamid;
- [13e]: [N-(Cyclopropyl)Glycyl]-[N-(Cyclopropyl)Glycyl]-N-(S)-sec-Butyl)Glycinamid;
- [14]: [N-(Cyclopropyl)Glycyl]-[N-(Cyclopropyl)Glycyl]-N-Phenethyl)Glycinamid;
- [15]: [N-(Cyclopropyl)Glycyl]-[N-(Cyclopropyl)Glycyl]-N-[2-(4-Aminosulfonylphenyl)Ethyl]Glycinamid;
- 50 [16a]: [N-(Cyclopropyl)Glycyl]-[N-[2-(N-Morpholin)Ethyl]Glycyl]-N-((R,S)-sec-Butyl)Glycinamid;
- [16b]: [N-(Cyclopropyl)Glycyl]-[N-[2-(N-Morpholin)Ethyl]Glycyl]-N-((R)-sec-Butyl)Glycinamid;
- [16c]: [N-(Cyclopropyl)Glycyl]-[N-[2-(N-Morpholin)Ethyl]Glycyl]-N-((S)-sec-Butyl)Glycinamid;
- [17]: [N-(Cyclopropyl)Glycyl]-[N-[2-(N-Morpholin)Ethyl]Glycyl]-N-(Phenethyl)Glycinamid;
- [18]: [N-(Cyclopropyl)Glycyl]-[N-[2-(N-Morpholin)Ethyl]Glycyl]-N-[2-(4-Aminosulfonylphenyl)Ethyl]Glycinamid;
- 55 [19a]: [N-[2-(N-pyrrolidinyl)Ethyl]Glycyl]-[N-(Cyclopropyl)Glycyl]-N-((R,S)-sec-Butyl)Glycinamid;
- [19b]: [N-[2-(N-pyrrolidinyl)Ethyl]Glycyl]-[N-(Cyclopropyl)Glycyl]-N-((R)-sec-Butyl)Glycinamid;
- [19c]: [N-[2-(N-pyrrolidinyl)Ethyl]Glycyl]-[N-(Cyclopropyl)Glycyl]-N-((S)-sec-Butyl)Glycinamid;
- [20]: [N-[2-(N-pyrrolidinyl)Ethyl]Glycyl]-[N-(Cyclopropyl)Glycyl]-N-(Phenethyl)Glycinamid;

[21]: [N-[2-(N-pyrrolidinyl)Ethyl]Glycyl]-[N-(Cyclopropyl)Glycyl]-N-[2-(4-Aminosulfonylphenyl)Ethyl]Glycinamid;
 [22a]: [N-[2-(N-pyrrolidinyl)Ethyl]Glycyl]-[N-[2-(N-morpholin)Ethyl]Glycyl]-N-((R,S)-sec-Butyl)Glycinamid;
 [22b]: [N-[2-(N-pyrrolidinyl)Ethyl]Glycyl]-[N-[2-(N-morpholin)Ethyl]Glycyl]-N-((R)-sec-Butyl)Glycinamid;
 [22c]: [N-[2-(N-pyrrolidinyl)Ethyl]Glycyl]-[N-[2-(N-morpholin)Ethyl]Glycyl]-N-((S)-sec-Butyl)Glycinamid;
 [23]: [N-[2-(N-pyrrolidinyl)Ethyl]Glycyl]-[N-[2-(N-morpholin)Ethyl]Glycyl]-N-(Phenethyl)Glycinamid und
 [24]: [N-[2-(N-pyrrolidinyl)Ethyl]Glycyl]-[N-[2-(N-morpholin)Ethyl]Glycyl]-N-(2-(4-Aminosulfonylphenyl)Ethyl)Glycinamid;

- 5
- 10
- 15
- 20
- 25
- 30
- 35
6. Une pharmaceutische Zusammensetzung, bestehend aus einer therapeutisch wirksamen Menge von mindestens einer Formel (I) N-Alkylglycin Trimer laut einem beliebigen der Ansprüche 1 bis 5 und mindestens einem pharmazeutisch akzeptablen Bindemittel.
 7. Verwendung einer Formel (I) N-Alkylglycin Trimer laut einem beliebigen der Ansprüche 1 bis 5 zur Herstellung eines Arzneimittels zur Minderung der nervösen Aktivität von Neuronen, die an Neurodegeneration beteiligt ist, vermittelt durch die Anwendung exogener chemischer Substanzen; oder durch die endogene Freigabe chemischer Substanzen, die eine Erregungstoxizität im Nervensystem verursachen.
 8. Verwendung einer Formel (I) N-Alkylglycin Trimer laut einem beliebigen der Ansprüche 1 bis 5 zur Herstellung eines Arzneimittels, welches die ionischen Kanäle hemmt, die durch exogene chemischen Substanzen oder durch endogene chemische Substanzen aktiviert werden; die zu Neurodegeneration führen.
 9. Verwendung einer Formel (I) N-Alkylglycin Trimer laut einem beliebigen der Ansprüche 1 bis 5 zur Herstellung eines Arzneimittels zur Behandlung von Krankheiten und pathologischen Störungen, die durch die Aktivität von ionotropen Rezeptoren für L-Glutamat vermittelt werden.
 10. Verwendung einer Formel (I) N-Alkylglycin Trimer laut einem beliebigen der Ansprüche 1 bis 5 zur Herstellung eines Arzneimittels zur Behandlung, Verlangsamung, Reduktion, Abschwächung und/oder Vorbeugung der Neurodegeneration.
 11. Verwendung einer Formel (I) N-Alkylglycin Trimer laut einem beliebigen der Ansprüche 1 bis 5 zur Herstellung eines Arzneimittels zur Behandlung von zerebraler Ischämie, Schlaganfall, Migräne, Depression, Huntington, Parkinson, Alzheimer, seniler Demenz, Epilepsie und multipler und amyotropher Sklerose.

Revendications

1. Une file de N-alkyle-glycine dont la formule générale est (I)
Où:

40 R_1, R_2, R_3 , égaux ou différents et indépendants les uns des autres, sont sélectionnés parmi : cyclo-propyle, sec-butyle, 2- méthoxy-éthyle, 3- méthyle-butyle, cyclo-hexyle, 2- éthyle (N-pyrrolidinyle), 2- éthyle (2-méthyle carbylamine), 3- propyle (2- oxo-N-pyrrolidinyle), 2- éthyle (2- pyridyle), 2- phényléthyle, 1- méthyle (2- tétra hydrofuryle), 2- éthyle (N-imidazolyle), 2- éthyle (4-méthoxyphényle), 2- éthyle (3, 4-diméthoxyphényle), 2-éthyle (2, 4- dichlorophényle), 2- éthyle (2- (N méthyle) pyrrolidinyle), 2- éthyle (4- aminosulphonyphényle), 2- éthyle (morpholine), 3- propyle (N-N-diéthylamine), 3, 3- diphénylpropyle, 3- propyle (N, N-diméthylamine) et 2-éthyle (N,N- diéthylamine),
 45 ses formes et mélanges stéréo-isométriques, en grappes ou non, et leurs sels acceptables en pharmacie.

2. Une file de N-alkyle-glycine d'après la revendication 1, dans laquelle : R_3 est sec-butyle, 2- phényléthyle, 2- éthyle (méthylcarbylamine), N, N- diméthylaminopropyle, ou 2- éthyle (N- imidazolyle), ses formes et mélanges stéréo-isométriques, en grappes ou non.
3. Une file de N-alkyle-glycine d'après l'une ou l'autre des revendications 1 et 2, dans laquelle R_2 est cyclo-propyle, N, N- diéthylaminopropyle, 2- éthyle (morpholine) ou 3, 3-diphénylpropyle, ses formes et mélanges stéréo-isométriques, en grappes ou non.
4. Une file de N-alkyle-glycine d'après l'une ou l'autre des revendications 1 et 2, dans laquelle R_3 est 2- éthyle (N-pyrrolidinyle), cyclo-propyle, 3, 3-diphénylpropyle ou 2- (2- éthyle (N- méthylpyrrolidinyle), ses formes et mélanges

stéréo-isométriques, en grappes ou non.

5. Une file de N-alkyle-glycine d'après la revendication 1, sélectionnée parmi:

- 5 (1) N-glycyle propyle(3,3-diphényle)-(N-éthyle)(3-(N,N-diéthylamine propyle)-N-[2-glycinamide(méthylcarbylamine)éthyle];
- (2) [N-glycyle propyle](3,3-diphényle)-(N-glycyle propyle)(3-(N,N diéthylamine)-N-(1-(2-pyridyle)éthyle)glycinamide;
- 10 (3) (N-(3,3diphénylpropyle glycyle)-N-(3-(N,Ndiéthylamine propyle)glycyle) -N-(2-glycinamide éthyle(N-imidazolyle));
- (4) [N-glycyle propyle](3,3diphényle)-(N-glycyle propyle)(3-(N,N diéthylamine)-N-glycinamide propyle)(3-(N,N-diméthylamine).
- (5a) [N-(R,S)-2-[2-(N-(glycyle éthyle)méthyle pirrolidinyle)]-N-(2-(N,N-glycinamide glycyle (méthylcarbylamine)
- 15 (5b) [N-(R)-2-[2-(N-(glycyleéthyle)méthyle pirrolidinyle)]-N-(3(N,N-glycyle propyle diéthylamine)-N-12 glycina-mide éthyle(méthylcarbylamine)
- (5c) [N-(B)-2-[2-(N-(glycyle éthyle)méthyle pirrolidinyle)]N-(3-glycyle propyle(N,N-diéthylamine)-N-glycinamide éthyle(2-méthylcarbylamine);
- (6a) [N-(R,S)-2-[2-(N-(glycyleé thyle)méthyle pirrolidinyle)],-(N-(3-N,N-glycyle propyle (diéthylamine)-N-(2-gly-cinamideéthyle(pyridyle));
- 20 (6b) [N-(R)-2-[2-(N-(glycyle éthyle)méthyle(pirrolidinyle))-(N-(3-(N,N-glycyle propyle diéthylamine)-N-(2-12pyridyle)éthyl glycinamide;
- (6c) [N-(S)-2-[2-(N-méthyle)pirrolidinyle)éthyle]glycyle-(N-(3-(N,N-diéthylamine propyle) glycyle-N-(2-(2-glyci-namide éthyle pyridyle));
- (7a) [N-[R,S)-2-[2-(N-(glycyle éthyle)méthyle(pirrolidinyle)]N-(3(N,N-glycyle propyle diéthylamine)-N-(2-(2-gly-cinamide éthyle(méthylcarbylaznine);
- 25 (7b) [N-(R)-2-[2-(N-(glycyle éthyle)méthyle(pirrolidinyle))-(N-(3(N,N-glycyle propyle diéthylamine)-N-(2-(N-gly-cinamide éthyle(imidazolyle));
- (7c) [N-(S)-2-[2-(N-(glycyleéthyle)méthyle(pirrolidinyle))(N-(3(N,N-glycylepropyle diéthylamine)-N-(2-(N-glyci-namideéthyleimidazolyle);
- 30 (8a) [N-(R,S)-2-[2-(N-(glycyleéthyle)méthyle(pirrolidinyle))-(N-(3(N,N-glycylepropyle diéthylamine)N-(3-(N,N-glycinamidepropyle(diméthylamine);
- (8b) [N-(R)-2-[2-(N-(glycyleéthyle)méthyle(pirrolidinyle))-(N-(3-(N,N-glycylepropyle diéthylamine)-N-(3-(N,N-glycinamidepropyle(diméthylamine);
- (8c) [N-(S)-2-[2-(N-(glycyleéthyle)méthyle(pirrolidinyle))-(N-(3(N,N-glycylepropyle diéthylamine)-N-(3-(N,N-gly-cinamidepropyle(diméthylamine);
- 35 (9a) [N-(R,S)-2-(2-(N-(glycyleéthyle)méthyle(pirrolidinyle))(N-(3(N3,3-glycylepropyle diphényle)-N-(2-(glycina-mideéthyleméthylcarbylamine);
- (9b) [N-(R)-2-[2-(N-(glycyleéthyle)méthyle(pirrolidinyle))-(N-(3,3-glycylepropyle diphényle)-N-(2-(glycinamideé-thyleméthylcarbylamine);
- 40 (9c) [N-(S)-2-[2-(N-méthyle)pirrolidinyle)éthyle]glycyle-N-(3(N3,3-diphénylepropyle)glycyle N-2(méthylcarbyla-mine)éthyle)glycinamide;
- (10a) [N-(R,S)-2-[2-(N-(glycyleéthyle)méthyle(pirrolidinyle))N-(3,3-glycylepropyle diphényle)-N-(2-glycina-mideéthyle(pyridyle);
- (10b) [N-(R)-2-[2-(N-(glycyleéthyle)méthyle(pirrolidinyle))N-(3,3-glycylepropyle diphényle)-N-(2-(2-glycina-mideéthyle(pyridyle);
- 45 (10c) [N-(S)-2-[2-(N-(glycyleéthyle)méthyle(pirrolidinyle))-(N-(3(N3,3-glycylepropyle diphényle)-N-(2-(2-glycina-mideéthyle(pyridyle);
- (11a) [N-(R,S)-2-[2-(N-(glycyleéthyle)méthyle(pirrolidinyle))-(N-(3,3-glycylepropyle diphényle)-N-(2-(N-glycina-mideéthyleimidazolyle);
- 50 (11b) [N-(R)-2-[2-(N-(glycyle éthyle)méthyle(pirrolidinyle))-(N-(3,3-glycyle propyle diphényle)-N-(2-(N-glycina-mide éthyle imidazolyle);
- (11c) [N-(S)-2-[2-(N-(glycyle éthyle)méthyle(pirrolidinyle))-(N-(3,3-diphényle propyle)glycyle N-2(N-imidazolyle)éthyle)glycinamide;
- (12a) [N-[R,S)-2-[2-(N-méthyle)pirrolidinyle)éthyle]glycyleN-(3(N3,3glycyle propyle-diphényle)-N-(3-(N,N-glyci-namide propyle diméthylamine);
- 55 (12b) [N-(R)-2-[2-(N-(glycyle éthyle)méthyle(pirrolidinyle))N-(3,3-glycyle propyle diphényle)N-(3-(N,N-glycina-mide propyle diméthylamine);
- (12c) [N-(S)-2-[2-(N-(glycyle éthyle)méthyle(pirrolidinyle))-(N-(3(N-3,3 glycyle propyle diphényle)-N-(3-(N,N-gly-

cinamide propyle(diméthylamine);

(13a) [N-glycyle(cyclo-propyle)-(N-glycyle cyclo-propyle)-N-[(R,S)-glycinamide sec-butyle];

(13b) [N-glycyle(cyclo-propyle)-(N-glycyle cyclo-propyle)-N-[(R)-glycinamide sec-butyle];

(13c) [N-glycyle(cyclo-propyle)-(N-glycyle cyclo-propyle)-N-[(S)-glycinamide sec-butyle];

(14) [N-glycyle(cyclo-propyle)-(N-glycyle cyclo-propyle)-N-glycinamide phénétyle];

(15) [N-glycyle(cyclo-propyle)-(N-glycyle cyclo-propyle)-N-(2-(4-glycinamide éthyle aminosulphonyle phényle)];

(16a) [N-glycyle(cyclo-propyle)-(N-(2-(N-glycyle éthyle morpholine)N-[(R,S) -glycinamide sec-butyle];

(16b) [N-glycyle(cyclo-propyle)-(N-(2-(N-glycyle éthyle morpholine)N-[(R)-glycinamide sec-butyle];

(16c) [N-glycyle(cyclo-propyle)-(N-(2-(N-glycyle éthyle morpholine)N-[(S)-glycinamide sec-butyle];

(17)[N-glycyle(cyclo-propyle)-(N-(2-(N-glycyle éthyle morpholine)-N-glycinamide(phénétyle)];

(18)[N-glycyle(cyclo-propyle)-(N-(2-(N-glycyle éthyle morpholine)-N-(2-(4-glycinamide éthyle aminosulphonyle phényle)];

(19a) [N-(2(glycyle éthyle(N-pyrrolidinyle)-(N-glycyle cyclo-propyle)N-[(R, S)-glycinamide sec-butyle];

(19b) [N-(2-glycyle éthyle(N-pyrrolidinyle)-(N-glycyle cyclo-propyle)-N-[(R) -glycinamide sec-butyle];

(19c) [N-(2-glycyle)éthyle(N-pyrrolidinyle)-(N-glycyle cyclo-propyle))-N-[(S) -glycinamide sec-butyle];

(20)[N-(2-glycyle éthyle(N-pyrrolidinyle)-(N-glycyle cyclo-propyle)-N-glycinamide(phénétyle)];

(21)[N-(2-glycyle éthyle(N-pyrrolidinyle)-(N-glycyle cyclo-propyle)N-(2-(4-glycinamide a éthyle aminosulfonyle phényle)];

(22a) [N-(2-glycyle éthyle(N-pyrrolidinyle)N-(2-glycyle éthyle(N-morpholine) -N-[(R,S)glycinamide sec-butyle];

(22b) [N-(2-glycyle éthyle(N-pyrrolidinyle)N-(2-glycyle éthyle(N-morpholine) -N-[(R)glycinamide sec-butyle];

(22c) [N-(2-glycyle éthyle(N-pyrrolidinyle)-N-(2-glycyle éthyle(N-morpholine) -N-[(S)glycinamide sec-butyle];

(23)[N-(2-glycyle(éthyle N-pyrrolidinyle)-N-(2-glycyle éthyle(N-morpholine) -N-glycinamide(phénétyle),et

(24)[N-(2-glycyle éthyle(N-pyrrolidinyle)-N-(2-glycyle éthyle(N-morpholine) -N-(2-(4-glycinamide éthyle amino-sulphonyle phényle).

6. Un composé pharmaceutique contenant une quantité efficace en thérapeutique d'au moins une formule (I) de file de N-alkyle-glycine, d'après l'une ou l'autre des revendications 1 à 5 et au moins un excipient acceptable en termes de pharmacie.
7. Utilisation d'une formule (I) de file de N-alkyle-glycine, d'après l'une ou l'autre des revendications 1 à 5, dans l'élaboration d'un médicament pour atténuer l'activité nerveuse des neurones qu'implique la neurodégénérescence, provoquée par l'application de substances chimiques exogènes, ou par la libération endogène de substances chimiques qui provoquent une excito-toxicité du système nerveux.
8. Utilisation d'une formule (I) de file de N-alkyle-glycine, d'après l'une ou l'autre des revendications 1 à 5, dans l'élaboration d'un médicament qui inhibe les chaînes ioniques qui sont activées par les substances chimiques exogènes ou par des substances chimiques endogènes, qui conduisent à une neurodégénérescence.
9. Utilisation d'une formule (I) de file de N-alkyle-glycine, d'après l'une ou l'autre des revendications 1 à 5, dans l'élaboration d'un médicament pour le traitement de maladies et troubles pathologiques provoqués par l'activité des récepteurs ionotropiques du glutamate L.
10. Utilisation d'une formule (I) de file de N-alkyle-glycine, d'après l'une ou l'autre des revendications 1 à 5, dans l'élaboration d'un médicament pour le traitement, le ralentissement, la réduction, la modération et/ ou la prévention de la neurodégénérescence,
11. Utilisation d'une formule (I) de file de N-alkyle-glycine, d'après l'une ou l'autre des revendications 1 à 5, dans l'élaboration d'un médicament pour le traitement de l'ischémie cérébrale, des accidents cardio-vasculaires, de la migraine, la dépression, la maladie de Huntington, Parkinson, Alzheimer, la démence sénile, l'épilepsie et la sclérose amyotrophique multiple.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Non-patent literature cited in the description

- **OLNEY, J.W.** Excitotoxic amino acid and neuropsychiatric disorders. *Annu. Rev. Pharmacol. Toxicol.*, 1990, vol. 30, 47-71 [0045]
- **LIPTON, S.A. ; ROSENBERG, P.A.** Excitatory amino acids as a final common pathway for neurologic disorders. *New Engl. J. Med.*, 1994, vol. 330, 613-622 [0045]
- **CHOI, D.W. ; ROTHMANN, S.M.** The role of glutamate neurotoxicity in hypoxic ischemic neuronal death. *Annu. Rev. Neurosci.*, 1990, vol. 13, 171-182 [0045]
- **COLLINGRIDGE, G.L. ; LESTER, R.A.J.** Excitatory amino acid receptors in the vertebrate central nervous system. *Pharmacol. Rev.*, 1989, vol. 40, 143-210 [0045]
- **CHOI, D.W.** Calcium and excitotoxic neuronal injury. *Ann. N. Y. Acad. Sci.*, 1994, vol. 747, 162-171 [0045]
- **SIESJÖ, B.K.** Calcium-mediated processes in neuronal degeneration. *Ann. N.Y. Acad. Sci.*, 1994, vol. 747, 140-161 [0045]
- **LIPTON, S.A.** AIDS-related dementia and calcium homeostasis. *Ann. N.Y. Acad. Sci.*, 1994, vol. 747, 205-224 [0045]
- **MONTOLIU, C. ; LLANSOLA, M. ; KOSENKO, E. ; CORBALÁN, R ; FELIPO, V.** Role of cyclic GMP in glutamate neurotoxicity in primary cultures of cerebellar neurons. *Neuropharmacology*, 1999, vol. 3 (8), 1883-1891 [0045]
- **KORNHUBER, J. ; QUACK, G.** Cerebrospinal fluid and serum concentrations of the N-methyl-D-aspartate (NMDA) receptor antagonist memantine in man. *Neurosci. Lett.*, 1995, vol. 195, 137-139 [0045]
- **NAKANISHI, S. ; MASU, M.** Molecular diversity and functions of glutamate receptors (1994). *Annu. Rev. Biophys. Biomol. Struct.*, 1994, vol. 23, 319-348 [0045]
- Clinical implications of NMDA receptor. **HERRLING, PL.** NMDA receptor. 1994, 376-394 [0045]
- **LIPTON, S.A.** Prospects for clinically tolerated NMDA antagonists: open channel blockers and alternative redox states of nitric oxide. *Trends neurosci.*, 1993, vol. 16, 527-532 [0045]
- **MÜLLER, W.E. ; MUTSCHLER, E. ; RIEDERER, P.** Non-competitive NMDA receptor antagonists with fast open channel blocking kinetics and strong voltage dependency as potential therapeutic agents for Alzheimer's. *Pharmacopsychiat*, 1995, vol. 28, 113-124 [0045]
- Non-competitive NMDA antagonists as drugs. **IVERSEN, L.L ; KEMP, J.A.** NMDA receptor. 1994, 469-486 [0045]
- **MIÑANA, M.D. ; HERMENEGILDO, C. ; LLANSOLA, M. ; MONTOLIU, C. ; GRISOLIA, S. ; FELIPO, V.** Carnitine and choline derivatives containing a trimethylamine group prevent ammonia toxicity in mice and glutamate toxicity in primary cultures of neurons. *J. Pharmacol. Exp. Ther.*, 1996, vol. 279, 194-199 [0045]
- **KOEK, W. ; WOODS, J.H. ; WINGER, G.D.** MK-801, a proposed noncompetitive antagonist of excitatory amino acid neurotransmission, produces phencyclidine-like behavioral effects in pigeons, rats and rhesus monkeys. *J. Pharmacol. Exp. Ther.*, 1988, vol. 245, 969-974 [0045]
- **CHOI, D.W.** Dextrorphan and dextromethorphan attenuate glutamate neurotoxicity. *Brain res.*, 1987, vol. 403, 333-336 [0045]
- **SEIF EL NASR, M. ; PERUCHE, B. ; ROSSBERG, C. ; MENNEL, H.-D. ; KRIEGLSTEIN, J.** Neuroprotective effect of memantine demonstrated in vivo and in vitro. *Eur. J. Pharmacol.*, 1990, vol. 185, 19-24 [0045]
- **FERRER-MONTIEL, A.V. ; MONTAL, M.** Structure-function relations in ligand-gated ion channels: Reconstitution in lipid bilayers and heterologous expression in *Xenopus* oocytes. *Methods: a Companion to Methods in Enzymology*, 1994, vol. 6, 60-69 [0045]
- **FERRER-MONTIEL, A.V. ; SUN, W. ; MONTAL, M.** Molecular design of the NMDA receptor binding site for PCP and MK-801. *Proc. Natl. Acad. Sci. USA.*, 1995, vol. 92, 8021-8025 [0045]
- **SUCKERMANN, R.N. ; KERR, J.M. ; KENT, S.B.H. ; MOOS, W.H.** Efficient method for the preparation of peptoids [oligo(N-substituted glycines)] by submonomer solid-phase synthesis. *J. Am. Chem. Soc.*, 1992, vol. 114, 10646-10647 [0045]
- **FIGLIOZZI, G.M. ; GOLDSMITH, R. ; NG., S.C. ; BANVILLE, S.C. ; ZUCKERMANN, R.N.** Synthesis of N-substituted glycine peptoid libraries. *Method Enzymol.*, 1996, vol. 267, 437-447 [0045]
- **MINANA M.D. ; MONTOLIU, C. ; LLANSOLA, M. ; GRISOLIA, S. ; FELIPO, V.** Nicotine prevents glutamate-induced proteolysis of the microtubule-associated protein MAP-2 and glutamate neurotoxicity in primary cultures of cerebellar neurons. *Neuropharmacology*, vol. 37, 847-857 [0045]

- **MARACAIDA, G. ; MIÑANA, M.D. ; BURGAL, M. ; GRISOLIA, S. ; FELIPO, V.** Ammonia prevents activation of NMDA receptors in rat cerebellar neuronal cultures. *Eur. J. Neurosci.*, 1995, vol. 7, 2389-2396
[0045]