



European Patent Office

Office européen des brevets



EP 1 364 946 A1

(12)

published in accordance with Art. 158(3) EPC

26.11.2003 Bulletin 2003/48

(21) Application number: **02701539.5**

(22) Date of filing: 07.02.2002

(51) Int Cl.⁷: **C07D 261/10**, C07D 261/12,
C07D 413/12, C07D 498/04,
C07D 513/04, C07D 413/14,
C07D 417/12, C07D 261/20,
C07D 487/04, A01N 43/80

(86) International application number:
PCT/JP02/01015

(87) International publication number:
WO 02/062770 (15.08.2002 Gazette 2002/33)

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR
 Designated Extension States:
AL LT LV MK RO SI

(30) Priority: 08.02.2001 JP 2001031784

(71) Applicants:

- KUMIAI CHEMICAL INDUSTRY CO., LTD.
Tokyo 110-0008 (JP)
- IHARA CHEMICAL INDUSTRY CO., LTD.
Taitoh-ku, Tokyo 110-0008 (JP)


(72) Inventors:
• **NAKATANI, Masao**
Iwata-gun, Shizuoka 437-1213 (JP)

- KUGO, Ryotaro
Sen-nan-shi, Osaka 590-0531 (JP)
- MIYAZAKI, Masahiro
Iwata-gun, Shizuoka 437-1213 (JP)
- KAKU, Koichiro
Iwata-gun, Shizuoka 437-1207 (JP)
- FUJINAMI, Makoto
Ogasa-gun, Shizuoka 439-0031 (JP)
- UENO, Ryohei
Ogasa-gun, Shizuoka 439-0031 (JP)
- TAKAHASHI, Satoru
Shizuoka-shi, Shizuoka 420-0046 (JP)

(74) Representative: **Laufhütte, Dieter, Dr.-Ing.**
Lorenz-Seidler-Gossel
Widenmayerstrasse 23
80538 München (DE)

(54) **ISOXAZOLINE DERIVATIVE AND HERBICIDE COMPRISING THE SAME AS ACTIVE INGREDIENT**

(57) An isoxazoline derivative represented by the following general formula [I]:



wherein R¹ and R² may be the same or different and are each an alkyl group;

R³, R⁴, R⁵ and R⁶ are each a hydrogen atom;

Y is an optionally substituted 5- to 6-membered aromatic heterocyclic group or fused aromatic heterocyclic group having a hetero atom selected from a nitrogen atom, a oxygen atom and a sulfur atom; and
n is an integer of 0 to 2.

The isoxazoline derivative has an excellent herbicidal effect and an excellent selectivity between crop and weed.

Description

Background of the Invention

1. Field of the Invention

[0001] The present invention relates to a novel isoxazoline derivative and a herbicide containing the isoxazoline derivative as the active ingredient.

2. Description of the Prior Art

[0002] The herbicidal activity of isoxazoline derivatives are reported in, for example, JP-A-8-22558, JP-A-9-328477 and JP-A-9-328483. The compound of the present invention described in detail later, however, is not described in these literatures.


[0003] Herbicides applied to useful crops are desired to (a) be applicable to soil or foliage, (b) show a sufficient herbicidal effect at a low ingredient amount, and (c) show a high selectivity between crop and weed. In these respects, the compounds described in the above literatures are not fully satisfactory.

Summary of the Invention

[0004] In view of the above situation, the present inventors made a study on the herbicidal effect and selectivity between crop and weed of various compounds. As a result, the present inventors found out that a novel isoxazoline derivative has an excellent herbicidal effect and an excellent selectivity between crop and weed. The above finding has led to the completion of the present invention.

[0005] The present invention provides the followings.

(1) An isoxazoline derivative represented by the following general formula [I] or a pharmaceutically acceptable salt thereof:



wherein R^1 and R^2 may be the same or different and are each a hydrogen atom, a C1 to C10 alkyl group, a C3 to C8 cycloalkyl group or a C3 to C8 cycloalkyl C1 to C3 alkyl group, or R^1 and R^2 may be bonded to each other to form a C3 to C7 spiro ring together with the carbon atoms to which they bond;

[0006] R^3 and R^4 may be the same or different and are each a hydrogen atom, a C1 to C10 alkyl group or a C3 to C8 cycloalkyl group; or R^3 and R^4 may be bonded to each other to form a C3 to C7 spiro ring together with the carbon atoms to which they bond; or R^1 , R^2 , R^3 and R^4 may form a 5- to 8-membered ring together with the carbon atoms to which they bond;

[0007] R^5 and R^6 may be the same or different and are each a hydrogen atom or a C1 to C10 alkyl group;

[0008] Y is a 5- to 6-membered aromatic heterocyclic group or condensed aromatic heterocyclic group having one or more hetero atoms selected from a nitrogen atom, an oxygen atom and a sulfur atom; the heterocyclic group may be substituted with 0 to 6 same or different groups selected from the following substituent group α ; when the heterocyclic group is substituted at the two adjacent positions with two alkyl groups, two alkoxy groups, an alkyl group and an alkoxy group, an alkyl group and an alkylthio group, an alkyl group and an alkylsulfonyl group, an alkyl group and a monoalkylamino group, or an alkyl group and a dialkylamino group, all selected from the substituent group α , the two groups may form, together with the atoms to which they bond, a 5- to 8-membered ring which may be substituted with 1 to 4 halogen atoms; the hetero atom of the heterocyclic group, when it is a nitrogen atom, may be oxidized to become N-oxide;

n is an integer of 0 to 2.

[Substituent group α]

[0009] Hydroxyl group; thiol group; halogen atoms; C1 to C10 alkyl groups; C1 to C10 alkyl groups each mono-substituted with a group selected from the following substituent group β , C1 to C4 haloalkyl groups; C3 to C8 cycloalkyl groups; C1 to C10 alkoxy groups; C1 to C10 alkoxy groups each mono-substituted with a group selected from the following substituent group γ , C1 to C4 haloalkoxy groups; C3 to C8 cycloalkoxy groups; C3 to C8 cycloalkyl C1 to C3 alkyloxy groups; C1 to C10 alkylthio groups; C1 to C10 alkylthio groups each mono-substituted with a group selected from the substituent group γ ; C1 to C4 haloalkylthio groups; C2 to C6 alkenyl groups; C2 to C6 alkenyloxy groups; C2 to C6 alkynyl groups; C2 to C6 alkynyloxy groups; C1 to C10 alkylsulfonyl groups; C1 to C10 alkylsulfonyl groups each mono-substituted with a group selected from the substituent group γ ; C1 to C10 alkylsulfonyl groups; C1 to C10 alkylsulfonyl groups each mono-substituted with a group selected from the substituent group γ , C1 to C4 haloalkylsulfonyl groups; C1 to C10 alkylsulfonyloxy groups each mono-substituted with a group selected from the substituent group γ ; C1 to C4 haloalkylsulfonyl groups; C1 to C10 alkylsulfonyloxy groups; C1 to C4 haloalkylsulfonyloxy groups; optionally substituted phenyl group; optionally substituted phenoxy group; optionally substituted phenylthio group; optionally substituted aromatic heterocyclic groups; optionally substituted aromatic heterocyclic oxy groups; optionally substituted aromatic heterocyclic thio groups; optionally substituted phenylsulfonyl groups; optionally substituted phenylsulfonyl groups; optionally substituted aromatic heterocyclic sulfonyl groups; optionally substituted phenylsulfonyloxy groups; acyl groups; C1 to C4 haloalkylcarbonyl groups; optionally substituted benzylcarbonyl group; optionally substituted benzoyl group; carboxyl group; C1 to C10 alkoxycarbonyl groups; optionally substituted benzyloxycarbonyl group; optionally substituted phenoxycarbonyl group; cyano group; carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups and optionally substituted phenyl group); C1 to C6 acyloxy groups; C1 to C4 haloalkylcarbonyloxy groups; optionally substituted benzylcarbonyloxy group; optionally substituted benzyloxy group; nitro group; and amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, optionally substituted phenyl group, C1 to C6 acyl groups, C1 to C4 haloalkylcarbonyl groups, optionally substituted benzylcarbonyl group, optionally substituted benzoyl group, C1 to C10 alkylsulfonyl group, C1 to C4 haloalkylsulfonyl groups, optionally substituted benzylsulfonyl group, and optionally substituted phenylsulfonyl group).

[Substituent group β]

[0010] Hydroxyl group; C3 to C8 cycloalkyl groups which may be substituted with halogen atom or alkyl group); C1 to C10 alkoxy groups; C1 to C10 alkylthio groups; C1 to C10 alkylsulfonyl groups; C1 to C10 alkoxycarbonyl groups; C2 to C6 haloalkenyl groups; amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, C1 to C6 acyl groups; C1 to C4 haloalkylcarbonyl groups, C1 to C10 alkylsulfonyl groups and C1 to C4 haloalkylsulfonyl groups); carbamoyl group (its nitrogen atom may be substituted with same or different C1 to C10 alkyl groups); C1 to C6 acyl groups; C1 to C4 haloalkylcarbonyl groups; C1 to C10 alkoxyimino groups; cyano group; optionally substituted phenyl group; and optionally substituted phenoxy group.

[Substituent group γ]

[0011] C1 to C10 alkoxycarbonyl groups; optionally substituted phenyl group; optionally substituted aromatic heterocyclic groups; cyano group; and carbamoyl group (its nitrogen atom may be substituted with same or different C1 to C10 alkyl groups).

(2) An isoxazoline derivative according to (1), wherein the substituent group α on the heterocycle which may be substituted with 0 to 6 same or different groups, includes hydroxyl group; halogen atoms; C1 to C10 alkyl groups; C1 to C10 alkyl groups each mono-substituted with a group selected from the substituent group β , C1 to C4 haloalkyl groups; C3 to C8 cycloalkyl groups; C1 to C10 alkoxy groups; C1 to C10 alkoxy groups each mono-substituted with a group selected from the substituent group γ ; C1 to C4 haloalkoxy groups; C3 to C8 cycloalkoxy groups; C3 to C8 cycloalkyl C1 to C3 alkyloxy groups; C1 to C10 alkylthio groups; C1 to C10 alkylthio groups each mono-substituted with a group selected from the substituent group γ ; C1 to C4 haloalkylthio groups; C2 to C6 alkenyl groups; C2 to C6 alkenyloxy groups; C2 to C6 alkynyl groups; C2 to C6 alkynyloxy groups; C1 to C10 alkylsulfonyl groups; C1 to C4 haloalkylsulfonyl groups; optionally substituted phenyl group; optionally substituted phenoxy group; optionally substituted phenylthio group; optionally substituted aromatic heterocyclic groups; optionally substituted aromatic heterocyclic oxy groups; optionally substituted aromatic heterocyclic thio groups; optionally substituted phenylsulfonyl groups; optionally substituted aromatic heterocyclic sulfonyl groups; C1 to C6 acyl groups; C1 to C4 haloalkylcarbonyl groups; optionally substituted benzylcarbonyl group; optionally substituted benzoyl group; carboxyl group; C1 to C10 alkoxycarbonyl groups; cyano group; carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups and optionally substituted phenyl group); nitro group; and amino group (its nitrogen atom

may be substituted with same or different groups selected from C1 to C10 alkyl groups, optionally substituted phenyl group, C1 to C6 acyl groups, C1 to C4 haloalkylcarbonyl groups, optionally substituted benzylcarbonyl group, optionally substituted benzoyl group, C1 to C10 alkylsulfonyl groups, C1 to C4 haloalkylsulfonyl groups, optionally substituted benzylsulfonyl group, and optionally substituted phenylsulfonyl group); when the heterocyclic group is substituted at the two adjacent positions with two alkyl groups, two alkoxy groups, an alkyl group and an alkoxy group, an alkyl group and an alkylthio group, an alkyl group and an alkylsulfonyl group, an alkyl group and a monoalkylamino group, or an alkyl group and a dialkylamino group, all selected from the substituent group α , the two groups may form, together with the atoms to which they bond, a 5- to 8-membered ring which may be substituted with 1 to 4 halogen atoms.

(3) An isoxazoline derivative according to (2), wherein the substituent group α on the heterocycle which may be substituted with 0 to 6 same or different groups, includes halogen atoms; C1 to C10 alkyl groups; C1 to C4 haloalkyl groups; C1 to C10 alkoxy C1 to C3 alkyl groups; C3 to C8 cycloalkyl groups which may be substituted with halogen atom or alkyl group; C1 to C10 alkoxy groups; C1 to C4 haloalkoxy groups; C3 to C8 cycloalkyl C1 to C3 alkyloxy groups; optionally substituted phenoxy group; C1 to C10 alkylthio groups; C1 to C10 alkylsulfonyl groups; acyl groups; C1 to C4 haloalkylcarbonyl groups; C1 to C10 alkoxycarbonyl groups; cyano group and carbamoyl group (its nitrogen atom may be substituted with same or different C1 to C10 alkyl groups).

(4) An isoxazoline derivative according to any of (1), (2) or (3), wherein R^1 and R^2 may be the same or different and are each a methyl group or an ethyl group; and R^3 , R^4 , R^5 and R^6 are each a hydrogen atom.

(5) An isoxazoline derivative according to any of (1), (2), (3) or (4), wherein Y is a 5- or 6-membered aromatic heterocyclic group having a hetero atom selected from a nitrogen atom, an oxygen atom and a sulfur atom.

(6) An isoxazoline derivative according to (5), wherein Y is a thienyl group, a pyrazolyl group, an isoxazolyl group, an isothiazolyl group, a pyridyl group or a pyrimidinyl group.

(7) An isoxazoline derivative according to (6), wherein Y is a thiophen-3-yl group, a pyrazol-4-yl group, a pyrazol-5-yl group, an isoxazol-4-yl group, an isothiazol-4-yl group, a pyridin-3-yl group or a pyrimidin-5-yl group.

(8) An isoxazoline derivative according to (7), wherein Y is a thiophen-3-yl group and the thiophene ring is substituted with the substituent group α at the 2- and 4-positions.

(9) An isoxazoline derivative according to (7), wherein Y is a pyrazol-4-yl group and the pyrazole ring is substituted at the 3- and 5-positions with the substituent group α and at the 1-position with a hydrogen atom, a C1 to C10 alkyl group, a C1 to C10 alkyl group mono-substituted with a group selected from the substituent group β , a C1 to C4 haloalkyl group, a C3 to C8 cycloalkyl group, a C2 to C6 alkenyl group, a C2 to C6 alkynyl group, a C1 to C10 alkylsulfinyl group, a C1 to C10 alkylsulfonyl group, a C1 to C10 alkylsulfonyl group mono-substituted with a group selected from the substituent group γ , a C1 to C4 haloalkylsulfonyl group, an optionally substituted phenyl group, an optionally substituted aromatic heterocyclic group, an optionally substituted phenylsulfonyl group, an optionally substituted aromatic heterocyclic sulfonyl group, an acyl group, a C1 to C4 haloalkylcarbonyl group, an optionally substituted benzylcarbonyl group, an optionally substituted benzoyl group, a C1 to C10 alkoxycarbonyl group, an optionally substituted benzyloxycarbonyl group, an optionally substituted phenoxycarbonyl group, a carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups and optionally substituted phenyl group), or an amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, optionally substituted phenyl group, acyl groups, C1 to C4 haloalkylcarbonyl groups, optionally substituted benzylcarbonyl group, optionally substituted benzoyl group, C1 to C10 alkylsulfonyl groups, C1 to C4 haloalkylsulfonyl groups, optionally substituted benzylsulfonyl group and optionally substituted phenylsulfonyl group).

(10) An isoxazoline derivative according to (7), wherein Y is a pyrazol-5-yl group and the pyrazole ring is substituted at the 4-position with the substituent group α and at the 1-position with a hydrogen atom, a C1 to C10 alkyl group, a C1 to C10 alkyl group mono-substituted with a group selected from the substituent group β , a C1 to C4 haloalkyl group, a C3 to C8 cycloalkyl group, a C2 to C6 alkenyl group, a C2 to C6 alkynyl group, a C1 to C10 alkylsulfinyl group, a C1 to C10 alkylsulfonyl group, a C1 to C10 alkylsulfonyl group mono-substituted with a group selected from the substituent group γ , a C1 to C4 haloalkylsulfonyl group, an optionally substituted phenyl group, an optionally substituted aromatic heterocyclic group, an optionally substituted phenylsulfonyl group, an optionally substituted aromatic heterocyclic sulfonyl group, an acyl group, a C1 to C4 haloalkylcarbonyl group, an optionally substituted benzylcarbonyl group, an optionally substituted benzoyl group, a C1 to C10 alkoxycarbonyl group, an optionally substituted benzyloxycarbonyl group, an optionally substituted phenoxycarbonyl group, a carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups and optionally substituted phenyl group), or an amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, optionally substituted phenyl group, acyl groups, C1 to C4 haloalkylcarbonyl groups, optionally substituted benzylcarbonyl group, optionally substituted benzoyl group, C1 to C10 alkylsulfonyl groups, C1 to C4 haloalkylsulfonyl groups, optionally substituted benzylsulfonyl group and optionally substituted phenylsulfonyl group).

(11) An isoxazoline derivative according to (7), wherein Y is an isoxazol-4-yl group and the isoxazole ring is substituted with the substituent group α at the 3- and 5-positions.

(12) An isoxazoline derivative according to (7), wherein Y is an isothiazol-4-yl group and the isothiazole ring is substituted

tuted with the substituent group α at the 3- and 5-positions.

(13) An isoxazoline derivative according to (7), wherein Y is a pyridin-3-yl group and the pyridine ring is substituted with the substituent group α at the 2- and 4-positions.

(14) An isoxazoline derivative according to (7), wherein Y is a pyrimidin-5-yl group and the pyrimidine ring is substituted with the substituent group α at the 4- and 6-positions.

(15) An isoxazoline derivative according to any of (1) to (14), wherein n is an integer of 2.

(16) An isoxazoline derivative according to any of (1) to (14), wherein n is an integer of 1.

(17) An isoxazoline derivative according to any of (1) to (14), wherein n is an integer of 0.

(18) A herbicide containing, as the active ingredient, an isoxazoline derivative set forth in any of (1) to (17) or a pharmaceutically acceptable salt thereof.

Detailed Description of the Invention

[0012] The definitions of the terms used in the present specification are given below.

[0013] The expression of "C1 to C10", etc. indicates that the substituent appearing after the expression has 1 to 10 carbon atoms in the case of "C1 to C10".

[0014] Halogen atom refers to a fluorine atom, a chlorine atom, a bromine atom or an iodine atom.

[0015] C1 to C10 alkyl group refers to a straight or branched chain alkyl group of 1 to 10 carbon atoms unless otherwise specified; and there can be mentioned, for example, methyl group, ethyl group, n-propyl group, isopropyl group, n-butyl group, isobutyl group, sec-butyl group, tert-butyl group, n-pentyl group, isopentyl group, neopentyl group, n-hexyl group, isohexyl group, 3,3-dimethylbutyl group, heptyl group and octyl group.

[0016] C3 to C8 cycloalkyl group refers to a cycloalkyl group of 3 to 8 carbon atoms; and there can be mentioned, for example, cyclopropyl group, cyclobutyl group, cyclopentyl group and cyclohexyl group.

[0017] C3 to C8 cycloalkyl C1 to C3 alkyl group (which may be substituted with halogen atom or alkyl group) refers, unless otherwise specified, to a C1 to C3 alkyl group substituted with a C3 to C8 cycloalkyl group which may be substituted with 1 to 4 same or different halogen atoms or C1 to C3 alkyl group; and there can be mentioned, for example, cyclopropylmethyl group, 1-cyclopropylethyl group, 2-cyclopropylethyl group, 1-cyclopropylpropyl group, 2-cyclopropylpropyl group, 3-cyclopropylpropyl group, cyclobutylmethyl group, cyclopentylmethyl group, cyclohexylmethyl group, 2-chlorocyclopropylmethyl group, 2,2-dichlorocyclopropylmethyl group, 2-fluorocyclopropylmethyl group, 2,2-difluorocyclopropylmethyl group, 2-methylcyclopropylmethyl group, 2,2-dimethylcyclopropylmethyl group and 2-methylcyclopropylethyl group.

[0018] C3 to C8 cycloalkyl C1 to C3 alkyl group refers to a alkyl group of 1 to 3 carbon atoms, substituted with a cycloalkyl group of 3 to 8 carbon atoms; and there can be mentioned, for example, cyclopropylmethyl group, 1-cyclopropylethyl group, 2-cyclopropylethyl group, 1-cyclopropylpropyl group, 2-cyclopropylpropyl group, 3-cyclopropylpropyl group, cyclobutylmethyl group, cyclopentylmethyl group and cyclohexylmethyl group.

[0019] C1 to C4 haloalkyl group refers, unless otherwise specified, to a straight or branched chain alkyl group of 1 to 4 carbon atoms, substituted with 1 to 9 same or different halogen atoms; and there can be mentioned, for example, fluoromethyl group, chloromethyl group, bromomethyl group, difluoromethyl group, trifluoromethyl group, 2,2-difluoroethyl group, 2,2,2-trifluoroethyl group and pentafluoroethyl group.

[0020] C2 to C6 alkenyl group refers to a straight or branched chain alkenyl group of 2 to 6 carbon atoms; and there can be mentioned, for example, ethenyl group, 1-propenyl group, 2-propenyl group, isopropenyl group, 1-butenyl group, 2-butenyl group, 3-butenyl group and 2-pentenyl group.

[0021] C2 to C6 alkynyl group refers to a straight or branched chain alkynyl group of 2 to 6 carbon atoms; and there can be mentioned, for example, ethynyl group, 2-propynyl group, 1-methyl-2-propynyl group, 2-butylnyl group, 3-butylnyl group and 2-methyl-3-butylnyl group.

[0022] C2 to C6 haloalkenyl group refers, unless otherwise specified, to a straight or branched alkenyl group of 2 to 6 carbon atoms, substituted with 1 to 4 same or different halogen atoms; and there can be mentioned, for example, 3-chloro-2-propenyl group and 2-chloro-2-propenyl group.

[0023] C1 to C10 alkoxy group refers to an (alkyl)-O- group wherein the alkyl moiety has the above definition; and there can be mentioned, for example, methoxy group, ethoxy group, n-propoxy group, isopropoxy group, tert-butoxy group, n-butoxy group, sec-butoxy group and isobutoxy group.

[0024] C1 to C4 haloalkoxy group refers to a (haloalkyl)-O-group wherein the haloalkyl moiety has the above definition; and there can be mentioned, for example, difluoromethoxy group, trifluoromethoxy group, 2,2-difluoroethoxy group and 2,2,2-trifluoroethoxy group.

[0025] C3 to C8 cycloalkyloxy group refers to a (cycloalkyl)-O- group wherein the cycloalkyl moiety has the above definition; and there can be mentioned, for example, cyclopropyloxy group, cyclobutyloxy group, cyclopentyloxy group and cyclohexyloxy group.

[0026] C3 to C8 cycloalkyl C1 to C3 alkyloxy group refers to a (cycloalkylalkyl)-O- group wherein the cycloalkylalkyl

moiety has the above definition; and there can be mentioned, for example, cyclopropylmethoxy group, 1-cyclopropylethoxy group, 2-cyclopropylethoxy group, 1-cyclopropylpropoxy group, 2-cyclopropylpropoxy group, 3-cyclopropylpropoxy group, cyclobutylmethoxy group, cyclopentylmethoxy group and cyclohexylmethoxy group.

[0027] C2 to C6 alkenyloxy group and C2 to C6 alkynyloxy group refer, respectively, to an (alkenyl)-O- group and an (alkynyl)-O- group, in each of which the alkenyl or alkynyl moiety has the above definition; and there can be mentioned, for example, 2-propenyloxy group and 2-propynyloxy group.

[0028] C1 to C10 alkoxyimino group refers to an (alkoxy)-N= group wherein the alkoxy moiety has the above definition; and there can be mentioned, for example, methoxyimino group and ethoxyimino group.

[0029] C1 to C10 alkylthio group, C1 to C10 alkylsulfinyl group and C1 to C10 alkylsulfonyl group refer, respectively, to an (alkyl)-S- group, an (alkyl)-SO- group and an (alkyl)-SO₂- group, in each of which the alkyl moiety has the above definition; and there can be mentioned, for example, methylthio group, ethylthio group, n-propylthio group, isopropylthio group, methylsulfinyl group, methylsulfonyl group, ethylsulfonyl group, n-propylsulfonyl group and isopropylsulfonyl group.

[0030] C1 to C10 alkylsulfonyloxy group refers to an (alkylsulfonyl)-O- group wherein the alkylsulfonyl moiety has the above definition, and there can be mentioned, for example, methylsulfonyloxy group and ethylsulfonyloxy group.

[0031] C1 to C10 alkoxy carbonyl group refers to an (alkoxy)-CO- group wherein the alkoxy moiety has the above definition, and there can be mentioned, for example, methoxycarbonyl group, ethoxycarbonyl group, n-propoxycarbonyl group and isopropoxycarbonyl group.

[0032] C1 to C6 acyl group refers to a straight or branched chain aliphatic acyl group of 1 to 6 carbon atoms, and there can be mentioned, for example, formyl group, acetyl group, propionyl group, isopropionyl group, butyryl group and pivaloyl group.

[0033] C1 to C10 acyloxy group refers to an (acyl)-O- group wherein the acyl moiety has the above definition; and there can be mentioned, for example, acetoxy group, propionyloxy group, isopropionyloxy group and pivaloyloxy group.

[0034] C1 to C4 haloalkylcarbonyl group, C1 to C4 haloalkylthio group and C1 to C4 haloalkylsulfonyl group refers, respectively, to a (haloalkyl)-CO- group, a (haloalkyl)-S-group and a (haloalkyl)-SO₂- group, in each of which the haloalkyl moiety has the above definition; and there can be mentioned, for example, chloroacetyl group, trifluoroacetyl group, pentafluoropropyl group, difluoromethylthio group, trifluoromethylthio group, chloromethylsulfonyl group, difluoromethylsulfonyl group and trifluoromethylsulfonyl group.

[0035] C1 to C4 haloalkylcarbonyloxy group and C1 to C4 haloalkylsulfonyloxy group refer, respectively, to a (haloalkylcarbonyl)-O- group and a (haloalkylsulfonyl)-O-group, in each of which the haloalkylcarbonyl moiety or the haloalkylsulfonyl moiety has the above definition; and there can be mentioned, for example, chloroacetyloxy group, trifluoroacetyloxy group, chloromethylsulfonyloxy group and trifluoromethylsulfonyloxy group.

[0036] "Optionally substituted" in (optionally substituted) phenyl group, (optionally substituted) aromatic heterocyclic group, (optionally substituted) phenoxy group, (optionally substituted) aromatic heterocyclic oxy group, (optionally substituted) phenylthio group, (optionally substituted) aromatic heterocyclic thio group, (optionally substituted) phenylsulfonyl group, (optionally substituted) phenylsulfonyloxy group, (optionally substituted) aromatic heterocyclic sulfonyl group, (optionally substituted) benzylcarbonyl group, (optionally substituted) benzylcarbonyloxy group, (optionally substituted) benzylsulfonyl group, (optionally substituted) benzoyl group, (optionally substituted) benzoyloxy group, (optionally substituted) benzoyloxy carbonyl group and (optionally substituted) phenoxy carbonyl group, refers to being optionally substituted with, for example, halogen atom, C1 to C10 alkyl group, C1 to C4 haloalkyl group, C1 to C10 alkoxyalkyl group, C1 to C10 alkoxy group, C1 to C10 alkylthio group, C1 to C10 alkylsulfonyl group, acyl group, C1 to C10 alkoxy carbonyl group, cyano group, carbamoyl group (its nitrogen atom may be substituted with same or different C1 to C10 alkyl groups), nitro group or amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, C1 to C6 acyl groups, C1 to C4 haloalkylcarbonyl groups, C1 to C10 alkylsulfonyl groups and C1 to C4 haloalkylsulfonyl groups).

[0037] 5- to 6-membered aromatic heterocyclic group having a hetero atom selected from a nitrogen atom, an oxygen atom and a sulfur atom includes, for example, furyl group, thienyl group, pyrrolyl group, pyrazolyl group, isoxazolyl group, isothiazolyl group, oxazolyl group, thiazolyl group, imidazolyl group, pyridyl group, pyridazinyl group, pyrimidinyl group, pyrazinyl group, triazinyl group, triazolyl group, oxadiazolyl group and thiadiazolyl group, each having 1 to 3 hetero atoms.

[0038] Fused aromatic heterocyclic group refers to a group having 1 to 3 hetero atoms randomly selected from nitrogen atom, oxygen atom and sulfur atom; and there can be mentioned, for example, benzofuryl group, benzothienyl group, indolyl group, benzoxazolyl group, benzothiazolyl group, benzimidazolyl group, benzisoxazolyl group, benzisothiazolyl group, indazolyl group, quinolyl group, isoquinolyl group, phthalazinyl group, quinoxalinyl group, quinazolinyl group, cinnolinyl group and benzotriazolyl group.

[0039] Aromatic heterocycle in (optionally substituted) aromatic heterocyclic group, (optionally substituted) aromatic heterocyclic oxy group, (optionally substituted) aromatic heterocyclic thio group and (optionally substituted) aromatic heterocyclic sulfonyl group, refers to a 5- to 6-membered group having 1 to 3 hetero atoms randomly selected from

nitrogen atom, oxygen atom and sulfur atom; and there can be mentioned, for example, furyl group, thienyl group, pyrrolyl group, pyrazolyl group, isoxazolyl group, isothiazolyl group, oxazolyl group, thiazolyl group, imidazolyl group, pyridyl group, pyridazinyl group, pyrimidinyl group, pyrazinyl group, triazinyl group, triazolyl group, oxadiazolyl group and thiadiazolyl group.

[0040] Pharmaceutically acceptable salt is a salt of a compound of the general formula [I] having, in the structure, hydroxyl group, carboxyl group, amino group or the like, with a metal or an organic base or with a mineral acid or an organic acid. As the metal, there can be mentioned alkali metals such as sodium, potassium and the like; and alkaline earth metals such as magnesium, calcium and the like. As the organic base, there can be mentioned triethylamine, diisopropylamine, etc. As the mineral acids, there can be mentioned hydrochloric acid, sulfuric acid, etc. As the organic acid, there can be mentioned acetic acid, methanesulfonic acid, p-toluenesulfonic acid, etc.

[0041] In the above-mentioned general formula [I], it is preferred that

R¹ and R² may be the same or different and are each a methyl group or an ethyl group;

R³, R⁴, R⁵ and R⁶ are each a hydrogen atom;

n is an integer of 2; and

Y is a thiophen-3-yl group [the 2- and 4-positions of the group are substituted with same or different groups selected from halogen atoms, alkyl groups, haloalkyl groups, alkoxyalkyl groups, cycloalkyl groups, alkoxy groups, haloalkoxy groups, acyl groups, haloalkylcarbonyl groups, alkoxy carbonyl groups, cyano group and carbamoyl group (its nitrogen atom may be substituted with same or different alkyl groups)], or

a pyrazol-4-yl group [the 3- and 5-positions of the group are substituted with same or different groups selected from halogen atoms, alkyl groups, haloalkyl groups, alkoxyalkyl groups, cycloalkyl groups, alkoxy groups, haloalkoxy groups, cycloalkylalkyloxy groups, optionally substituted phenoxy group, alkylthio groups, alkylsulfonyl groups, acyl groups, haloalkylcarbonyl groups, alkoxy carbonyl groups, cyano group and carbamoyl group (its nitrogen atom may be substituted with same or different alkyl groups)]; the 1-position is substituted with hydrogen atom, alkyl group, alkyl group mono-substituted with a group selected from the substituent group β, haloalkyl group, cycloalkyl group, alkenyl group, alkynyl group, alkylsulfonyl group, alkylsulfonyl group mono-substituted with a group selected from the substituent group γ, haloalkylsulfonyl group, optionally substituted phenyl group, optionally substituted aromatic heterocyclic group, optionally substituted phenylsulfonyl group, optionally substituted aromatic heterocyclicsulfonyl group, acyl group, haloalkylcarbonyl group, optionally substituted benzylcarbonyl group, optionally substituted benzoyl group, alkoxy carbonyl group, optionally substituted benzyloxycarbonyl group, optionally substituted phenoxycarbonyl group or carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from alkyl groups and optionally substituted phenyl group)], or

a pyrazol-5-yl group [the 4-position of the group is substituted with halogen atom, alkyl group haloalkyl group, alkoxyalkyl group, haloalkoxy group, acyl group, haloalkylcarbonyl group, alkoxy carbonyl group, cyano group or carbamoyl group (its nitrogen atom maybe substituted with same or different alkyl groups)]; the 1-position is substituted with hydrogen atom, alkyl group, alkyl group mono-substituted with a group selected from the substituent group β, haloalkyl group, cycloalkyl group, or optionally substituted phenyl group], or

an isoxazol-4-yl group [the 3- and 5-positions of the group are substituted with same or different groups selected from halogen atoms, alkyl groups, haloalkyl groups, alkoxyalkyl groups, cycloalkyl groups, alkoxy groups, haloalkoxy groups, alkylthio groups, alkylsulfonyl groups, acyl groups, haloalkylcarbonyl groups, alkoxy carbonyl groups, cyano group and carbamoyl group (its nitrogen atom may be substituted with same or different alkyl groups)], or

an isothiazol-4-yl group [the 3- and 5-positions of the group are substituted with same or different groups selected from halogen atoms, alkyl groups, haloalkyl groups, alkoxyalkyl groups, cycloalkyl groups, alkoxy groups, haloalkoxy groups, optionally substituted phenoxy group, alkylthio groups, alkylsulfonyl groups, acyl groups, haloalkylcarbonyl groups, alkoxy carbonyl groups, cyano group and carbamoyl group (its nitrogen atom may be substituted with same or different alkyl groups)], or

a pyridin-3-yl group [the 2- and 4-positions of the group are substituted with same or different groups selected from halogen atoms, alkyl groups, haloalkyl groups, alkoxyalkyl groups, cycloalkyl groups, alkoxy groups, haloalkoxy groups, alkylthio groups, alkylsulfonyl groups, acyl groups, haloalkylcarbonyl groups, alkoxy carbonyl groups, cyano group and carbamoyl group (its nitrogen atom may be substituted with same or different alkyl groups)], or

a pyrimidin-5-yl group [the 4- and 6-positions of the group are substituted with same or different groups selected from halogen atoms, alkyl groups, haloalkyl groups, alkoxyalkyl groups, cycloalkyl groups, alkoxy groups, haloalkoxy groups, alkylthio groups, alkylsulfonyl groups, acyl groups, haloalkylcarbonyl groups, alkoxy carbonyl groups, cyano group and carbamoyl group (its nitrogen atom may be substituted with same or different alkyl groups)].

[0042] Next, representative examples of the present compound represented by the general formula [I] are shown in Tables 1 to 10. However, the present compound is not restricted to these examples.

[0043] The following abbreviated expressions used in the Tables refer to the following groups.


EP 1 364 946 A1

Me: methyl group	Et: ethyl group
Pr: n-propyl group	Pr-i: isopropyl group
Pr-c: cyclopropyl group	Bu: n-butyl group
Bu-i: isobutyl group	Bu-s: sec-butyl group
Bu-t: tert-butyl group	Bu-c: cyclobutyl group
Pen: n-pentyl group	Pen-c: cyclopentyl group
Hex: n-hexyl group	p Hex-c: cyclohexyl group
Ph: phenyl group	

[0044] For example, (4-Cl)Ph indicates 4-chlorophenyl group, and 3-Hex indicates 3-hexyl group.

[0045] When the present compound contains hydroxyl group as a substituent, there may exist keto-enol tautomers. Any of these tautomers and any mixture of these tautomers are included in the present compound.

Table 1

										
R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Z ¹	R ²²	R ²³	R ²⁴
Me	Me	H	H	2	H	H	S	Me	H	H
Me	Me	H	H	2	H	H	S	Cl	Me	H
Me	Me	H	H	2	H	H	S	H	H	Me
Me	Me	H	H	2	H	H	S	Cl	H	H
Me	Me	H	H	2	H	H	S	H	H	Cl
Me	Me	H	H	2	H	H	S	Cl	Cl	Cl
Me	Me	H	H	2	H	H	S	OMe	H	H
Me	Me	H	H	2	H	H	S	OEt	H	H
Me	Me	H	H	2	H	H	S	OCHF ₂	H	H
Me	Me	H	H	2	H	H	S	OCH ₂ Ph	H	H
Me	Me	H	H	2	H	H	O	H	H	H
Me	Me	H	H	2	H	H	O	H	H	C(=O)OMe
Me	Me	H	H	2	H	H	NMe	Me	H	Me
Me	Me	H	H	2	H	H	NMe	Me	C(=O)OMe	CH ₂ C(=O)OMe
Me	Me	H	H	2	H	H	NMe	Me	C(=O)OEt	CH ₂ C(=O)OEt
Me	Me	H	H	2	H	H	NMe	Me	Me	Me
Me	Me	H	H	2	H	H	NPh	OMe	H	H
Me	Me	H	H	2	H	H	NPh	OEt	H	H
Me	Me	H	H	2	H	H	NPh	OCHF ₂	H	H
H	H	H	H	2	H	H	S	OCHF ₂	H	H
Me	H	H	H	2	H	H	S	OCHF ₂	H	H
Me	H	Me	H	2	H	H	S	OCHF ₂	H	H
Me	Me	H	H	2	Me	H	S	OCHF ₂	H	H
Me	Me	H	H	2	Et	H	S	OCHF ₂	H	H

	Me	Me	H	H	2	Pr-i	H	S	OCHF ₂	H	H
	Me	Me	H	H	2	Me	Me	S	OCHF ₂	H	H
5	Me	Et	H	H	2	H	H	S	OCHF ₂	H	H
	Et	Et	H	H	2	H	H	S	OCHF ₂	H	H
	Me	Pr-i	H	H	2	H	H	S	OCHF ₂	H	H
	Me	Pr	H	H	2	H	H	S	OCHF ₂	H	H
10	Me	Pr-c	H	H	2	H	H	S	OCHF ₂	H	H
	Me	CH ₂ Pr-c	H	H	2	H	H	S	OCHF ₂	H	H
		-(CH ₂) ₂ -	H	H	2	H	H	S	Cl	Cl	Cl
		-(CH ₂) ₃ -	H	H	2	H	H	S	Cl	Cl	Cl
15		-(CH ₂) ₄ -	H	H	2	H	H	S	Cl	Cl	Cl
		-(CH ₂) ₅ -	H	H	2	H	H	S	Cl	Cl	Cl
	H	-(CH ₂) ₃ -	H	H	2	H	H	S	Cl	Cl	Cl
	H	-(CH ₂) ₄ -	H	H	2	H	H	S	Cl	Cl	Cl
20	H	-(CH ₂) ₅ -	H	H	2	H	H	S	Cl	Cl	Cl
	H	-(CH ₂) ₆ -	H	H	2	H	H	S	Cl	Cl	Cl
	Me	Me	H	H	1	H	H	S	Me	H	H
	Me	Me	H	H	1	H	H	S	Cl	Me	H
25	Me	Me	H	H	1	H	H	S	H	H	Me
	Me	Me	H	H	1	H	H	S	Cl	H	H
	Me	Me	H	H	1	H	H	S	H	H	Cl
	Me	Me	H	H	1	H	H	S	Cl	Cl	Cl
30	Me	Me	H	H	1	H	H	S	OMe	H	H
	Me	Me	H	H	1	H	H	S	OEt	H	H
	Me	Me	H	H	1	H	H	S	OCHF ₂	H	H
	Me	Me	H	H	1	H	H	S	OCH ₂ Ph	H	H
	Me	Me	H	H	1	H	H	O	H	H	H
35	Me	Me	H	H	1	H	H	O	H	H	C(=O)Ome
	Me	Me	H	H	1	H	H	NMe	Me	H	Me
	Me	Me	H	H	1	H	H	NMe	Me	C(=O)OMe	CH ₂ C(=O)OMe
	Me	Me	H	H	1	H	H	NMe	Me	C(=O)OEt	CH ₂ C(=O)OEt
40	Me	Me	H	H	1	H	H	NMe	Me	Me	Me
	Me	Me	H	H	1	H	H	NPh	OMe	H	H
	Me	Me	H	H	1	H	H	NPh	OEt	H	H
	Me	Me	H	H	1	H	H	NPh	OCHF ₂	H	H
45	H	H	H	H	1	H	H	S	OCHF ₂	H	H
	Me	H	H	H	1	H	H	S	OCHF ₂	H	H
	Me	H	Me	H	1	H	H	S	OCHF ₂	H	H
	Me	Me	H	H	1	Me	H	S	OCHF ₂	H	H
50	Me	Me	H	H	1	Et	H	S	OCHF ₂	H	H
	Me	Me	H	H	1	Pr-i	H	S	OCHF ₂	H	H
	Me	Me	H	H	1	Me	Me	S	OCHF ₂	H	H
	Me	Et	H	H	1	H	H	S	OCHF ₂	H	H
55	Et	Et	H	H	1	H	H	S	OCHF ₂	H	H

5	Me	Pr-i	H	H	1	H	H	S	OCHF ₂	H	H
	Me	Pr	H	H	1	H	H	S	OCHF ₂	H	H
	Me	Pr-c	H	H	1	H	H	S	OCHF ₂	H	H
	Me	CH ₂ Pr-c	H	H	1	H	H	S	OCHF ₂	H	H
		-(CH ₂) ₂ -	H	H	1	H	H	S	Cl	Cl	Cl
		-(CH ₂) ₃ -	H	H	1	H	H	S	Cl	Cl	Cl
10		-(CH ₂) ₄ -	H	H	1	H	H	S	Cl	Cl	Cl
		-(CH ₂) ₅ -	H	H	1	H	H	S	Cl	Cl	Cl
	H	-(CH ₂) ₃ -	H	H	1	H	H	S	Cl	Cl	Cl
	H	-(CH ₂) ₄ -	H	H	1	H	H	S	Cl	Cl	Cl
15	H	-(CH ₂) ₅ -	H	H	1	H	H	S	Cl	Cl	Cl
	H	-(CH ₂) ₆ -	H	H	1	H	H	S	Cl	Cl	Cl
	Me	Me	H	H	0	H	H	S	Me	H	H
	Me	Me	H	H	0	H	H	S	Cl	Me	H
20	Me	Me	H	H	0	H	H	S	H	H	Me
	Me	Me	H	H	0	H	H	S	Cl	H	H
	Me	Me	H	H	0	H	H	S	H	H	Cl
	Me	Me	H	H	0	H	H	S	Cl	Cl	Cl
25	Me	Me	H	H	0	H	H	S	OMe	H	H
	Me	Me	H	H	0	H	H	S	OEt	H	H
	Me	Me	H	H	0	H	H	S	OCHF ₂	H	H
	Me	Me	H	H	0	H	H	S	OCH ₂ Ph	H	H
30	Me	Me	H	H	0	H	H	O	H	H	H
	Me	Me	H	H	0	H	H	O	H	H	C(=O)Ome
	Me	Me	H	H	0	H	H	NMe	Me	H	Me
	Me	Me	H	H	0	H	H	NMe	Me	C(=O)OMe	CH ₂ C(=O)OMe
	Me	Me	H	H	0	H	H	NMe	Me	C(=O)OEt	CH ₂ C(=O)OEt
35	Me	Me	H	H	0	H	H	NMe	Me	Me	Me
	Me	Me	H	H	0	H	H	NPh	OMe	H	H
	Me	Me	H	H	0	H	H	NPh	OEt	H	H
	Me	Me	H	H	0	H	H	NPh	OCHF ₂	H	H
40	H	H	H	H	0	H	H	S	OCHF ₂	H	H
	Me	H	H	H	0	H	H	S	OCHF ₂	H	H
	Me	H	Me	H	0	H	H	S	OCHF ₂	H	H
	Me	Me	H	H	0	Me	H	S	OCHF ₂	H	H
45	Me	Me	H	H	0	Et	H	S	OCHF ₂	H	H
	Me	Me	H	H	0	Pr-i	H	S	OCHF ₂	H	H
	Me	Me	H	H	0	Me	Me	S	OCHF ₂	H	H
	Me	Et	H	H	0	H	H	S	OCHF ₂	H	H
50	Et	Et	H	H	0	H	H	S	OCHF ₂	H	H
	Me	Pr-i	H	H	0	H	H	S	OCHF ₂	H	H
	Me	Pr	H	H	0	H	H	S	OCHF ₂	H	H
	Me	Pr-c	H	H	0	H	H	S	OCHF ₂	H	H
55	Me	CH ₂ Pr-c	H	H	0	H	H	S	OCHF ₂	H	H

5	-(CH ₂) ₂ -		H	H	0	H	H	S	Cl	Cl	Cl
	-(CH ₂) ₃ -		H	H	0	H	H	S	Cl	Cl	Cl
	-(CH ₂) ₄ -		H	H	0	H	H	S	Cl	Cl	Cl
	-(CH ₂) ₅ -		H	H	0	H	H	S	Cl	Cl	Cl
10	H	-(CH ₂) ₃ -		H	0	H	H	S	Cl	Cl	Cl
	H	-(CH ₂) ₄ -		H	0	H	H	S	Cl	Cl	Cl
	H	-(CH ₂) ₅ -		H	0	H	H	S	Cl	Cl	Cl
	H	-(CH ₂) ₆ -		H	0	H	H	S	Cl	Cl	Cl
15	Me	Et	H	H	2	H	H	S	H	H	H
	Me	Et	H	H	2	H	H	O	H	H	H
	Me	Et	H	H	2	H	H	NH	H	H	H

20

25

30

35


40

45

50

55

Table 2

										
R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Z ²	R ²⁵	R ²⁶	R ²⁷
Me	Me	H	H	2	H	H	S	H	H	H
Me	Me	H	H	2	H	H	S	H	OMe	H
Me	Me	H	H	2	H	H	S	Cl	H	Cl
Me	Me	H	H	2	H	H	S	Cl	Cl	Cl
Me	Me	H	H	2	H	H	S	Cl	Me	H
Me	Me	H	H	2	H	H	S	NHMe	Me	H
Me	Me	H	H	2	H	H	S	N(Me) ₂	Me	H
Me	Me	H	H	2	H	H	S	NHC(=O)Me	Me	H
Me	Me	H	H	2	H	H	S	NHC(=O)Ph	Me	H
Me	Me	H	H	2	H	H	S	NHSO ₂ Me	Me	H
Me	Me	H	H	2	H	H	S	NHSO ₂ Ph	Me	H
Me	Me	H	H	2	H	H	S	Me	Me	Me
Me	Me	H	H	2	H	H	S	Me	C(=O)OMe	Me
Me	Me	H	H	2	H	H	S	Me	C(=O)OEt	Me
Me	Me	H	H	2	H	H	S	Me	C(=O)OPh	Me
Me	Me	H	H	2	H	H	S	Me	CN	Me
Me	Me	H	H	2	H	H	S	Me	C(=O)NHMe	Me
Me	Me	H	H	2	H	H	S	Me	C(=O)Me	Me
Me	Me	H	H	2	H	H	S	Me	C(=O)Et	Me
Me	Me	H	H	2	H	H	S	Me	C(=O)Pr-i	Me
Me	Me	H	H	2	H	H	S	Me	C(=O)Pr	Me
Me	Me	H	H	2	H	H	S	Me	C(=O)CF ₃	Me
Me	Me	H	H	2	H	H	S	Me	C(=NOMe)Me	Me
Me	Me	H	H	2	H	H	S	Ph	C(=O)Me	Me
Me	Me	H	H	2	H	H	S	Ph	C(=NOMe)Me	Me

	Me	Me	H	H	2	H	H	S	CF ₃	OMe	H
	Me	Me	H	H	2	H	H	S	CF ₃	OEt	H
5	Me	Me	H	H	2	H	H	S	CF ₃	OPr-i	H
	Me	Me	H	H	2	H	H	S	CF ₃	OPr-i	H
	Me	Me	H	H	2	H	H	S	CF ₃	OCHF ₂	H
10	Me	Me	H	H	2	H	H	S	Cl	Me	H
	Me	Me	H	H	2	H	H	S	Cl	Me	Me
	Me	Me	H	H	2	H	H	S	Cl	C(=O)OMe	Cl
	Me	Me	H	H	2	H	H	S	Cl	CN	Cl
	Me	Me	H	H	2	H	H	S	Cl	C(=O)NHMe	Cl
15	Me	Me	H	H	2	H	H	S	Cl	C(=O)N(Me) ₂	Cl
	Me	Me	H	H	2	H	H	S	Cl	C(=O)Me	Cl
	Me	Me	H	H	2	H	H	S	Cl	C(=O)Et	Cl
	Me	Me	H	H	2	H	H	S	Cl	C(=O)Pr-i	Cl
20	Me	Me	H	H	2	H	H	S	Cl	C(=O)Pr	Cl
	Me	Me	H	H	2	H	H	S	Cl	C(=O)CF ₃	Cl
	Me	Me	H	H	2	H	H	S	Cl	C(=NOMe)Me	Cl
	Me	Me	H	H	2	H	H	O	H	H	H
25	Me	Me	H	H	2	H	H	O	Me	H	Cl
	H	H	H	H	2	H	H	S	Cl	Cl	Cl
	Me	H	H	H	2	H	H	S	Cl	Cl	Cl
	Me	H	Me	H	2	H	H	S	Cl	Cl	Cl
	Me	Me	H	H	2	Me	H	S	Cl	Cl	Cl
30	Me	Me	H	H	2	Et	H	S	Cl	Cl	Cl
	Me	Me	H	H	2	Pr-i	H	S	Cl	Cl	Cl
	Me	Me	H	H	2	Me	Me	S	Cl	Cl	Cl
	Me	Et	H	H	2	H	H	S	Cl	Cl	Cl
	Et	Et	H	H	2	H	H	S	Cl	Cl	Cl
35	Me	Pr-i	H	H	2	H	H	S	Cl	Cl	Cl
	Me	Pr	H	H	2	H	H	S	Cl	Cl	Cl
	Me	Pr-c	H	H	2	H	H	S	Cl	Cl	Cl
	Me	CH ₂ Pr-c	H	H	2	H	H	S	Cl	Cl	Cl
40		-(CH ₂) ₂ -	H	H	2	H	H	S	Cl	Cl	Cl
		-(CH ₂) ₃ -	H	H	2	H	H	S	Cl	Cl	Cl
		-(CH ₂) ₄ -	H	H	2	H	H	S	Cl	Cl	Cl
		-(CH ₂) ₅ -	H	H	2	H	H	S	Cl	Cl	Cl
45	H	-(CH ₂) ₃ -	H	2	H	H	H	S	Cl	Cl	Cl
	H	-(CH ₂) ₄ -	H	2	H	H	H	S	Cl	Cl	Cl
	H	-(CH ₂) ₅ -	H	2	H	H	H	S	Cl	Cl	Cl
50	H	-(CH ₂) ₆ -	H	2	H	H	H	S	Cl	Cl	Cl
	Me	Me	H	H	1	H	H	S	H	H	H
	Me	Me	H	H	1	H	H	S	H	OMe	H
	Me	Me	H	H	1	H	H	S	Cl	H	Cl
	Me	Me	H	H	1	H	H	S	Cl	Cl	Cl
55	Me	Me	H	H	1	H	H	S	Cl	Me	H
	Me	Me	H	H	1	H	H	S	NHMe	Me	H

	Me	Me	H	H	1	H	H	S	N(Me) ₂	Me	H
	Me	Me	H	H	1	H	H	S	NHC(=O)Me	Me	H
5	Me	Me	H	H	1	H	H	S	NHC(=O)Ph	Me	H
	Me	Me	H	H	1	H	H	S	NHSO ₂ Me	Me	H
	Me	Me	H	H	1	H	H	S	NHSO ₂ Ph	Me	H
	Me	Me	H	H	1	H	H	S	Me	Me	Me
10	Me	Me	H	H	1	H	H	S	Me	C(=O)OMe	Me
	Me	Me	H	H	1	H	H	S	Me	C(=O)OEt	Me
	Me	Me	H	H	1	H	H	S	Me	C(=O)OPh	Me
	Me	Me	H	H	1	H	H	S	Me	CN	Me
	Me	Me	H	H	1	H	H	S	Me	C(=O)NHMe	Me
15	Me	Me	H	H	1	H	H	S	Me	C(=O)Me	Me
	Me	Me	H	H	1	H	H	S	Me	C(=O)Et	Me
	Me	Me	H	H	1	H	H	S	Me	C(=O)Pr-i	Me
	Me	Me	H	H	1	H	H	S	Me	C(=O)Pr	Me
20	Me	Me	H	H	1	H	H	S	Me	C(=O)CF ₃	Me
	Me	Me	H	H	1	H	H	S	Me	C(=NOMe)Me	Me
	Me	Me	H	H	1	H	H	S	Ph	C(=O)Me	Me
	Me	Me	H	H	1	H	H	S	Ph	C(=NOMe)Me	Me
	Me	Me	H	H	1	H	H	S	CF ₃	OMe	H
25	Me	Me	H	H	1	H	H	S	CF ₃	OEt	H
	Me	Me	H	H	1	H	H	S	CF ₃	OPr-i	H
	Me	Me	H	H	1	H	H	S	CF ₃	OPr-i	H
30	Me	Me	H	H	1	H	H	S	CF ₃	OCHF ₂	H
	Me	Me	H	H	1	H	H	S	Cl	Me	H
	Me	Me	H	H	1	H	H	S	Cl	Me	Me
	Me	Me	H	H	1	H	H	S	Cl	C(=O)OMe	Cl
35	Me	Me	H	H	1	H	H	S	Cl	CN	Cl
	Me	Me	H	H	1	H	H	S	Cl	C(=O)NHMe	Cl
	Me	Me	H	H	1	H	H	S	Cl	C(=O)N(Me) ₂	Cl
	Me	Me	H	H	1	H	H	S	Cl	C(=O)Me	Cl
	Me	Me	H	H	1	H	H	S	Cl	C(=O)Et	Cl
40	Me	Me	H	H	1	H	H	S	Cl	C(=O)Pr-i	Cl
	Me	Me	H	H	1	H	H	S	Cl	C(=O)Pr	Cl
	Me	Me	H	H	1	H	H	S	Cl	C(=O)CF ₃	Cl
	Me	Me	H	H	1	H	H	S	Cl	C(=NOMe)Me	Cl
45	Me	Me	H	H	1	H	H	O	H	H	H
	Me	Me	H	H	1	H	H	O	Me	H	Cl
	H	H	H	H	1	H	H	S	Cl	Cl	Cl
	Me	H	H	H	1	H	H	S	Cl	Cl	Cl
	Me	H	Me	H	1	H	H	S	Cl	Cl	Cl
50	Me	Me	H	H	1	Me	H	S	Cl	Cl	Cl
	Me	Me	H	H	1	Et	H	S	Cl	Cl	Cl
	Me	Me	H	H	1	Pr-i	H	S	Cl	Cl	Cl
	Me	Me	H	H	1	Me	Me	S	Cl	Cl	Cl
	Me	Et	H	H	1	H	H	S	Cl	Cl	Cl
55	Et	Et	H	H	1	H	H	S	Cl	Cl	Cl
	Me	Pr-i	H	H	1	H	H	S	Cl	Cl	Cl

	Me	Pr	H	H	1	H	H	S	Cl	Cl	Cl
	Me	Pr-c	H	H	1	H	H	S	Cl	Cl	Cl
5	Me	CH ₂ Pr-c	H	H	1	H	H	S	Cl	Cl	Cl
	-(CH ₂) ₂ -		H	H	1	H	H	S	Cl	Cl	Cl
	-(CH ₂) ₃ -		H	H	1	H	H	S	Cl	Cl	Cl
10	-(CH ₂) ₄ -		H	H	1	H	H	S	Cl	Cl	Cl
	-(CH ₂) ₅ -		H	H	1	H	H	S	Cl	Cl	Cl
	H	-(CH ₂) ₃ -	H	1	H	H	S	Cl	Cl	Cl	Cl
	H	-(CH ₂) ₄ -	H	1	H	H	S	Cl	Cl	Cl	Cl
15	H	-(CH ₂) ₅ -	H	1	H	H	S	Cl	Cl	Cl	Cl
	H	-(CH ₂) ₆ -	H	1	H	H	S	Cl	Cl	Cl	Cl
	Me	Me	H	H	0	H	H	S	H	H	H
	Me	Me	H	H	0	H	H	S	H	OMe	H
20	Me	Me	H	H	0	H	H	S	Cl	H	Cl
	Me	Me	H	H	0	H	H	S	Cl	Cl	Cl
	Me	Me	H	H	0	H	H	S	Cl	Me	H
	Me	Me	H	H	0	H	H	S	NHMe	Me	H
25	Me	Me	H	H	0	H	H	S	N(Me) ₂	Me	H
	Me	Me	H	H	0	H	H	S	NHC(=O)Me	Me	H
	Me	Me	H	H	0	H	H	S	NHC(=O)Ph	Me	H
	Me	Me	H	H	0	H	H	S	NHSO ₂ Me	Me	H
	Me	Me	H	H	0	H	H	S	NHSO ₂ Ph	Me	H
30	Me	Me	H	H	0	H	H	S	Me	Me	Me
	Me	Me	H	H	0	H	H	S	Me	C(=O)OMe	Me
	Me	Me	H	H	0	H	H	S	Me	C(=O)OEt	Me
	Me	Me	H	H	0	H	H	S	Me	C(=O)OPh	Me
35	Me	Me	H	H	0	H	H	S	Me	CN	Me
	Me	Me	H	H	0	H	H	S	Me	C(=O)NHMe	Me
	Me	Me	H	H	0	H	H	S	Me	C(=O)Me	Me
	Me	Me	H	H	0	H	H	S	Me	C(=O)Et	Me
	Me	Me	H	H	0	H	H	S	Me	C(=O)Pr-i	Me
40	Me	Me	H	H	0	H	H	S	Me	C(=O)Pr	Me
	Me	Me	H	H	0	H	H	S	Me	C(=O)CF ₃	Me
	Me	Me	H	H	0	H	H	S	Me	C(=NOMe)Me	Me
	Me	Me	H	H	0	H	H	S	Ph	C(=O)Me	Me
45	Me	Me	H	H	0	H	H	S	Ph	C(=NOMe)Me	Me
	Me	Me	H	H	0	H	H	S	CF ₃	OMe	H
	Me	Me	H	H	0	H	H	S	CF ₃	OEt	H
	Me	Me	H	H	0	H	H	S	CF ₃	OPr-i	H
50	Me	Me	H	H	0	H	H	S	CF ₃	OPr-i	H
	Me	Me	H	H	0	H	H	S	CF ₃	OCHF ₂	H
	Me	Me	H	H	0	H	H	S	Cl	Me	H
	Me	Me	H	H	0	H	H	S	Cl	Me	Me
55	Me	Me	H	H	0	H	H	S	Cl	C(=O)OMe	Cl
	Me	Me	H	H	0	H	H	S	Cl	CN	Cl
	Me	Me	H	H	0	H	H	S	Cl	C(=O)NHMe	Cl

5	Me	Me	H	H	0	H	H	S	Cl	C(=O)N(Me) ₂	Cl
	Me	Me	H	H	0	H	H	S	Cl	C(=O)Me	Cl
	Me	Me	H	H	0	H	H	S	Cl	C(=O)Et	Cl
	Me	Me	H	H	0	H	H	S	Cl	C(=O)Pr-i	Cl
	Me	Me	H	H	0	H	H	S	Cl	C(=O)Pr	Cl
10	Me	Me	H	H	0	H	H	S	Cl	C(=O)CF ₃	Cl
	Me	Me	H	H	0	H	H	S	Cl	C(=NOMe)Me	Cl
	Me	Me	H	H	0	H	H	O	H	H	H
	Me	Me	H	H	0	H	H	O	Me	H	Cl
15	H	H	H	H	0	H	H	S	Cl	Cl	Cl
	Me	H	H	H	0	H	H	S	Cl	Cl	Cl
	Me	H	Me	H	0	H	H	S	Cl	Cl	Cl
	Me	Me	H	H	0	Me	H	S	Cl	Cl	Cl
	Me	Me	H	H	0	Et	H	S	Cl	Cl	Cl
20	Me	Me	H	H	0	Pr-i	H	S	Cl	Cl	Cl
	Me	Me	H	H	0	Me	Me	S	Cl	Cl	Cl
	Me	Et	H	H	0	H	H	S	Cl	Cl	Cl
	Et	Et	H	H	0	H	H	S	Cl	Cl	Cl
25	Me	Pr-i	H	H	0	H	H	S	Cl	Cl	Cl
	Me	Pr	H	H	0	H	H	S	Cl	Cl	Cl
	Me	Pr-c	H	H	0	H	H	S	Cl	Cl	Cl
	Me	CH ₂ Pr-c	H	H	0	H	H	S	Cl	Cl	Cl
30	-(CH ₂) ₂ -		H	H	0	H	H	S	Cl	Cl	Cl
	-(CH ₂) ₃ -		H	H	0	H	H	S	Cl	Cl	Cl
	-(CH ₂) ₄ -		H	H	0	H	H	S	Cl	Cl	Cl
	-(CH ₂) ₅ -		H	H	0	H	H	S	Cl	Cl	Cl
35	H	-(CH ₂) ₃ -	H	0	H	H	H	S	Cl	Cl	Cl
	H	-(CH ₂) ₄ -	H	0	H	H	H	S	Cl	Cl	Cl
	H	-(CH ₂) ₅ -	H	0	H	H	H	S	Cl	Cl	Cl
40	H	-(CH ₂) ₆ -	H	0	H	H	H	S	Cl	Cl	Cl

Table 3

R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	R ²⁹	R ²⁸	R ³⁰
Me	Me	H	H	2	H	H	Cl	H	Cl
Me	Me	H	H	2	H	H	OCHF ₂	H	Cl
Me	Me	H	H	2	H	H	OCHF ₂	H	OCHF ₂
Me	Me	H	H	2	H	H	Me	H	Cl

	Me	Me	H	H	2	H	H	Me	H	OCHF ₂
	Me	Me	H	H	2	H	H	CHF ₂	H	Cl
5	Me	Me	H	H	2	H	H	CHF ₂	H	OCHF ₂
	Me	Me	H	H	2	H	H	CF ₃	H	F
	Me	Me	H	H	2	H	H	CF ₃	H	Cl
10	Me	Me	H	H	2	H	H	CF ₃	H	OMe
	Me	Me	H	H	2	H	H	CF ₃	H	OEt
	Me	Me	H	H	2	H	H	CF ₃	H	OCHF ₂
	Me	Me	H	H	2	H	H	CF ₃	H	CN
15	Me	Me	H	H	2	H	H	CF ₃	H	Me
	Me	Me	H	H	2	H	H	H	Me	Cl
	Me	Me	H	H	2	H	H	Me	Me	Me
20	Me	Me	H	H	2	H	H	Me	Me	F
	Me	Me	H	H	2	H	H	F	Me	Me
	Me	Me	H	H	2	H	H	Me	Me	Cl
	Me	Me	H	H	2	H	H	Cl	Me	Me
25	Me	Me	H	H	2	H	H	Me	Me	OMe
	Me	Me	H	H	2	H	H	OMe	Me	Me
	Me	Me	H	H	2	H	H	Me	Me	OCHF ₂
30	Me	Me	H	H	2	H	H	OCHF ₂	Me	Me
	Me	Me	H	H	2	H	H	Me	Me	CN
	Me	Me	H	H	2	H	H	CN	Me	Me
	Me	Me	H	H	2	H	H	Et	Me	F
35	Me	Me	H	H	2	H	H	F	Me	Et
	Me	Me	H	H	2	H	H	Et	Me	Cl
	Me	Me	H	H	2	H	H	Cl	Me	Et
	Me	Me	H	H	2	H	H	Et	Me	OMe
40	Me	Me	H	H	2	H	H	OMe	Me	Et
	Me	Me	H	H	2	H	H	Et	Me	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	Me	Et
45	Me	Me	H	H	2	H	H	Et	Me	CN
	Me	Me	H	H	2	H	H	CN	Me	Et
	Me	Me	H	H	2	H	H	Pr-i	Me	F
	Me	Me	H	H	2	H	H	F	Me	Pr-i
50	Me	Me	H	H	2	H	H	Pr-i	Me	Cl
	Me	Me	H	H	2	H	H	Cl	Me	Pr-i
	Me	Me	H	H	2	H	H	Pr-i	Me	OMe
	Me	Me	H	H	2	H	H	OMe	Me	Pr-i
55	Me	Me	H	H	2	H	H	Pr-i	Me	OCHF ₂

	Me	Me	H	H	2	H	H	OCHF ₂	Me	Pr-i
	Me	Me	H	H	2	H	H	Pr-i	Me	CN
5	Me	Me	H	H	2	H	H	CN	Me	Pr-i
	Me	Me	H	H	2	H	H	Bu-t	Me	F
	Me	Me	H	H	2	H	H	F	Me	Bu-t
10	Me	Me	H	H	2	H	H	Bu-t	Me	Cl
	Me	Me	H	H	2	H	H	Cl	Me	Bu-t
	Me	Me	H	H	2	H	H	Bu-t	Me	OMe
	Me	Me	H	H	2	H	H	OMe	Me	Bu-t
15	Me	Me	H	H	2	H	H	Bu-t	Me	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	Me	Bu-t
	Me	Me	H	H	2	H	H	Bu-t	Me	CN
20	Me	Me	H	H	2	H	H	CN	Me	Bu-t
	Me	Me	H	H	2	H	H	CH ₂ OMe	Me	F
	Me	Me	H	H	2	H	H	F	Me	CH ₂ OMe
	Me	Me	H	H	2	H	H	CH ₂ OMe	Me	Cl
25	Me	Me	H	H	2	H	H	Cl	Me	CH ₂ OMe
	Me	Me	H	H	2	H	H	CH ₂ OMe	Me	OMe
	Me	Me	H	H	2	H	H	OMe	Me	CH ₂ OMe
	Me	Me	H	H	2	H	H	CH ₂ OMe	Me	OCHF ₂
30	Me	Me	H	H	2	H	H	OCHF ₂	Me	CH ₂ OMe
	Me	Me	H	H	2	H	H	CH ₂ OMe	Me	CN
	Me	Me	H	H	2	H	H	CN	Me	CH ₂ OMe
35	Me	Me	H	H	2	H	H	Cl	Me	Cl
	Me	Me	H	H	2	H	H	CHF ₂	Me	Cl
	Me	Me	H	H	2	H	H	Cl	Me	CHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	Me	H
40	Me	Me	H	H	2	H	H	OCHF ₂	Me	F
	Me	Me	H	H	2	H	H	F	Me	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	Me	Cl
	Me	Me	H	H	2	H	H	Cl	Me	OCHF ₂
45	Me	Me	H	H	2	H	H	OCHF ₂	Me	OMe
	Me	Me	H	H	2	H	H	OMe	Me	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	Me	OCHF ₂
50	Me	Me	H	H	2	H	H	OCHF ₂	Me	CN
	Me	Me	H	H	2	H	H	CN	Me	OCHF ₂
	Me	Me	H	H	2	H	H	CF ₃	Me	H
	Me	Me	H	H	2	H	H	CF ₃	Me	Cl
55	Me	Me	H	H	2	H	H	Cl	Me	CF ₃

	Me	Me	H	H	2	H	H	CF ₃	Me	Br
	Me	Me	H	H	2	H	H	Br	Me	CF ₃
5	Me	Me	H	H	2	H	H	CF ₃	Me	I
	Me	Me	H	H	2	H	H	I	Me	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Me	F
	Me	Me	H	H	2	H	H	F	Me	CF ₃
10	Me	Me	H	H	2	H	H	CF ₃	Me	OH
	Me	Me	H	H	2	H	H	OH	Me	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Me	OMe
15	Me	Me	H	H	2	H	H	OMe	Me	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Me	OEt
	Me	Me	H	H	2	H	H	OEt	Me	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Me	OPr-i
20	Me	Me	H	H	2	H	H	CF ₃	Me	OPr
	Me	Me	H	H	2	H	H	CF ₃	Me	OBu-t
	Me	Me	H	H	2	H	H	CF ₃	Me	OBu-s
25	Me	Me	H	H	2	H	H	CF ₃	Me	OBu-i
	Me	Me	H	H	2	H	H	CF ₃	Me	OBu
	Me	Me	H	H	2	H	H	CF ₃	Me	O(2-Pen)
	Me	Me	H	H	2	H	H	CF ₃	Me	O(3-Pen)
30	Me	Me	H	H	2	H	H	CF ₃	Me	OPen-n
	Me	Me	H	H	2	H	H	CF ₃	Me	O(2-Hex)
	Me	Me	H	H	2	H	H	CF ₃	Me	O(3-Hex)
35	Me	Me	H	H	2	H	H	CF ₃	Me	OHex-n
	Me	Me	H	H	2	H	H	CF ₃	Me	OPen-c
	Me	Me	H	H	2	H	H	CF ₃	Me	OHex-c
	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ Pr-c
40	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ Bu-c
	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ Pen-c
	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ Hex-c
45	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ CH=CH ₂
	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ C≡CH
	Me	Me	H	H	2	H	H	CF ₃	Me	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	Me	CF ₃
50	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ CHF ₂
	Me	Me	H	H	2	H	H	OCH ₂ CHF ₂	Me	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ CF ₃
	Me	Me	H	H	2	H	H	OCH ₂ CF ₃	Me	CF ₃
55	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ CN

	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ C(=O)OEt
	Me	Me	H	H	2	H	H	CF ₃	Me	OCH(Me)C(=O)OEt
5	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ C(=O)NH ₂
	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ C(=O)NHMe
	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ C(=O)N(Me) ₂
10	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	OPh
	Me	Me	H	H	2	H	H	CF ₃	Me	O(2-Cl)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(2-Br)Ph
15	Me	Me	H	H	2	H	H	CF ₃	Me	O(2-F)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(2-Me)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(2-OMe)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(2-NO ₂)Ph
20	Me	Me	H	H	2	H	H	CF ₃	Me	O(2-CN)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(2-C(=O)OMe)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(3-Cl)Ph
25	Me	Me	H	H	2	H	H	CF ₃	Me	O(3-Br)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(3-F)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(3-Me)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(3-OMe)Ph
30	Me	Me	H	H	2	H	H	CF ₃	Me	O(3-NO ₂)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(3-CN)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(3-C(=O)OMe)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(4-Cl)Ph
35	Me	Me	H	H	2	H	H	CF ₃	Me	O(4-Br)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(4-F)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(4-Me)Ph
40	Me	Me	H	H	2	H	H	CF ₃	Me	O(4-OMe)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(4-NO ₂)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(4-CN)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	O(4-C(=O)OMe)Ph
45	Me	Me	H	H	2	H	H	CF ₃	Me	OC(=O)Me
	Me	Me	H	H	2	H	H	CF ₃	Me	OC(=O)Et
	Me	Me	H	H	2	H	H	CF ₃	Me	OC(=O)CH ₂ Ph
50	Me	Me	H	H	2	H	H	CF ₃	Me	OC(=O)CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Me	OC(=O)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	OSO ₂ Me
	Me	Me	H	H	2	H	H	CF ₃	Me	OSO ₂ Et
55	Me	Me	H	H	2	H	H	CF ₃	Me	OSO ₂ CH ₂ Ph

	Me	Me	H	H	2	H	H	CF ₃	Me	OSO ₂ CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Me	OSO ₂ Ph
5	Me	Me	H	H	2	H	H	CF ₃	Me	SMe
	Me	Me	H	H	2	H	H	CF ₃	Me	SOMe
	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ Me
10	Me	Me	H	H	2	H	H	CF ₃	Me	SEt
	Me	Me	H	H	2	H	H	CF ₃	Me	SOEt
	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ Et
	Me	Me	H	H	2	H	H	CF ₃	Me	SPr
15	Me	Me	H	H	2	H	H	CF ₃	Me	SOPr
	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ Pr
	Me	Me	H	H	2	H	H	CF ₃	Me	SPr-i
	Me	Me	H	H	2	H	H	CF ₃	Me	SOPr-i
20	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ Pr-i
	Me	Me	H	H	2	H	H	CF ₃	Me	SBu-t
	Me	Me	H	H	2	H	H	CF ₃	Me	SOBu-t
25	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ Bu-t
	Me	Me	H	H	2	H	H	CF ₃	Me	SCHF ₂
	Me	Me	H	H	2	H	H	CF ₃	Me	SOCHF ₂
	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ CHF ₂
30	Me	Me	H	H	2	H	H	CF ₃	Me	SCF ₃
	Me	Me	H	H	2	H	H	CF ₃	Me	SOCF ₃
	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ CF ₃
35	Me	Me	H	H	2	H	H	CF ₃	Me	SPh
	Me	Me	H	H	2	H	H	CF ₃	Me	SOh
	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	SCH ₂ Ph
40	Me	Me	H	H	2	H	H	CF ₃	Me	SOCH ₂ Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ CH ₂ Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	SCH ₂ C(=O)OEt
45	Me	Me	H	H	2	H	H	CF ₃	Me	SOCH ₂ C(=O)OEt
	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ CH ₂ C(=O)OEt
	Me	Me	H	H	2	H	H	CF ₃	Me	SCH(Me)C(=O)OEt
	Me	Me	H	H	2	H	H	CF ₃	Me	SOCH(Me)C(=O)OEt
50	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ CH(Me)C(=O)OEt
	Me	Me	H	H	2	H	H	CF ₃	Me	SCH ₂ C(=O)NH ₂
	Me	Me	H	H	2	H	H	CF ₃	Me	SOCH ₂ C(=O)NH ₂
55	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ CH ₂ C(=O)NH ₂
	Me	Me	H	H	2	H	H	CF ₃	Me	SCH ₂ C(=O)NHMe

	Me	Me	H	H	2	H	H	CF ₃	Me	SOCH ₂ C(=O)NHMe
	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ CH ₂ C(=O)NHMe
5	Me	Me	H	H	2	H	H	CF ₃	Me	SCH ₂ C(=O)N(Me) ₂
	Me	Me	H	H	2	H	H	CF ₃	Me	SOCH ₂ C(=O)N(Me) ₂
	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ CH ₂ C(=O)N(Me) ₂
10	Me	Me	H	H	2	H	H	CF ₃	Me	NH ₂
	Me	Me	H	H	2	H	H	CF ₃	Me	NHMe
	Me	Me	H	H	2	H	H	CF ₃	Me	N(Me) ₂
15	Me	Me	H	H	2	H	H	CF ₃	Me	NHC(=O)Me
	Me	Me	H	H	2	H	H	CF ₃	Me	N(Me)C(=O)Me
	Me	Me	H	H	2	H	H	CF ₃	Me	NHSO ₂ Me
	Me	Me	H	H	2	H	H	CF ₃	Me	N(Me)SO ₂ Me
20	Me	Me	H	H	2	H	H	CF ₃	Me	NHSO ₂ CHF ₂
	Me	Me	H	H	2	H	H	CF ₃	Me	N(Me)SO ₂ CHF ₂
	Me	Me	H	H	2	H	H	CF ₃	Me	NHSO ₂ CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Me	N(Me)SO ₂ CF ₃
25	Me	Me	H	H	2	H	H	CF ₃	Me	NHPh
	Me	Me	H	H	2	H	H	CF ₃	Me	N(Me)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	CN
30	Me	Me	H	H	2	H	H	CN	Me	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Me	C(=O)OMe
	Me	Me	H	H	2	H	H	CF ₃	Me	C(=O)OPr-i
	Me	Me	H	H	2	H	H	CF ₃	Me	C(=O)OCH ₂ Ph
35	Me	Me	H	H	2	H	H	CF ₃	Me	C(=O)OPh
	Me	Me	H	H	2	H	H	CF ₃	Me	C(=O)NH ₂
	Me	Me	H	H	2	H	H	CF ₃	Me	C(=O)NHMe
40	Me	Me	H	H	2	H	H	CF ₃	Me	C(=O)N(Me) ₂
	Me	Me	H	H	2	H	H	CF ₃	Me	C(=O)Me
	Me	Me	H	H	2	H	H	CF ₃	Me	C(=O)CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Me	C(=O)CH ₂ Ph
45	Me	Me	H	H	2	H	H	CF ₃	Me	C(=O)Ph
	Me	Me	H	H	2	H	H	CF ₃	Me	Me
	Me	Me	H	H	2	H	H	Me	Me	CF ₃
50	Me	Me	H	H	2	H	H	CF ₃	Me	Et
	Me	Me	H	H	2	H	H	CF ₃	Me	Pr-i
	Me	Me	H	H	2	H	H	CF ₃	Me	Pr
	Me	Me	H	H	2	H	H	CF ₃	Me	CH ₂ OMe
55	Me	Me	H	H	2	H	H	CF ₃	Me	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Me	CHF ₂

	Me	Me	H	H	2	H	H	CF ₃	Me	Ph
	Me	Me	H	H	2	H	H	CF ₂ CF ₃	Me	Cl
5	Me	Me	H	H	2	H	H	CN	Me	F
	Me	Me	H	H	2	H	H	F	Me	CN
	Me	Me	H	H	2	H	H	CN	Me	Cl
10	Me	Me	H	H	2	H	H	Cl	Me	CN
	Me	Me	H	H	2	H	H	CN	Me	CN
	Me	Me	H	H	2	H	H	COOMe	Me	F
15	Me	Me	H	H	2	H	H	F	Me	COOMe
	Me	Me	H	H	2	H	H	COOMe	Me	Cl
	Me	Me	H	H	2	H	H	Cl	Me	COOMe
	Me	Me	H	H	2	H	H	SO ₂ Me	Me	Cl
20	Me	Me	H	H	2	H	H	Cl	Me	SO ₂ Me
	Me	Me	H	H	2	H	H	Ph	Me	Me
	Me	Me	H	H	2	H	H	Ph	Me	Cl
	Me	Me	H	H	2	H	H	Ph	Me	OEt
25	Me	Me	H	H	2	H	H	Ph	Me	CF ₃
	Me	Me	H	H	2	H	H	Ph	Me	Ph
	Me	Me	H	H	2	H	H	Me	Et	OCHF ₂
30	Me	Me	H	H	2	H	H	OCHF ₂	Et	Me
	Me	Me	H	H	2	H	H	Me	Et	CN
	Me	Me	H	H	2	H	H	CN	Et	Me
	Me	Me	H	H	2	H	H	Pr-i	Et	OCHF ₂
35	Me	Me	H	H	2	H	H	OCHF ₂	Et	Pr-i
	Me	Me	H	H	2	H	H	Pr-i	Et	CN
	Me	Me	H	H	2	H	H	CN	Et	Pr-i
	Me	Me	H	H	2	H	H	Cl	Et	Cl
40	Me	Me	H	H	2	H	H	OCHF ₂	Et	Cl
	Me	Me	H	H	2	H	H	Cl	Et	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	Et	OCHF ₂
45	Me	Me	H	H	2	H	H	CF ₃	Et	F
	Me	Me	H	H	2	H	H	F	Et	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Et	Cl
	Me	Me	H	H	2	H	H	Cl	Et	CF ₃
50	Me	Me	H	H	2	H	H	CF ₃	Et	OMe
	Me	Me	H	H	2	H	H	OMe	Et	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Et	OEt
	Me	Me	H	H	2	H	H	OEt	Et	CF ₃
55	Me	Me	H	H	2	H	H	CF ₃	Et	OCHF ₂

	Me	Me	H	H	2	H	H	OCHF ₂	Et	CF ₃
5	Me	Me	H	H	2	H	H	CF ₃	Et	CN
	Me	Me	H	H	2	H	H	CN	Et	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Et	Me
	Me	Me	H	H	2	H	H	Me	Et	CF ₃
10	Me	Me	H	H	2	H	H	Me	Pr-i	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	Pr-i	Me
	Me	Me	H	H	2	H	H	Me	Pr-i	CN
15	Me	Me	H	H	2	H	H	CN	Pr-i	Me
	Me	Me	H	H	2	H	H	Pr-i	Pr-i	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	Pr-i	Pr-i
	Me	Me	H	H	2	H	H	Pr-i	Pr-i	CN
20	Me	Me	H	H	2	H	H	CN	Pr-i	Pr-i
	Me	Me	H	H	2	H	H	Cl	Pr-i	Cl
	Me	Me	H	H	2	H	H	OCHF ₂	Pr-i	Cl
	Me	Me	H	H	2	H	H	Cl	Pr-i	OCHF ₂
25	Me	Me	H	H	2	H	H	OCHF ₂	Pr-i	OCHF ₂
	Me	Me	H	H	2	H	H	CF ₃	Pr-i	F
	Me	Me	H	H	2	H	H	F	Pr-i	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Pr-i	Cl
30	Me	Me	H	H	2	H	H	Cl	Pr-i	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Pr-i	OMe
	Me	Me	H	H	2	H	H	OMe	Pr-i	CF ₃
35	Me	Me	H	H	2	H	H	CF ₃	Pr-i	OEt
	Me	Me	H	H	2	H	H	OEt	Pr-i	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Pr-i	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	Pr-i	CF ₃
40	Me	Me	H	H	2	H	H	CF ₃	Pr-i	CN
	Me	Me	H	H	2	H	H	CN	Pr-i	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Pr-i	Me
45	Me	Me	H	H	2	H	H	Me	Pr-i	CF ₃
	Me	Me	H	H	2	H	H	Me	Pr	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	Pr	Me
	Me	Me	H	H	2	H	H	Me	Pr	CN
50	Me	Me	H	H	2	H	H	CN	Pr	Me
	Me	Me	H	H	2	H	H	Pr-i	Pr	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	Pr	Pr-i
	Me	Me	H	H	2	H	H	Pr-i	Pr	CN
55	Me	Me	H	H	2	H	H	CN	Pr	Pr-i

	Me	Me	H	H	2	H	H	Cl	Pr	Cl
	Me	Me	H	H	2	H	H	OCHF ₂	Pr	Cl
5	Me	Me	H	H	2	H	H	Cl	Pr	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	Pr	OCHF ₂
	Me	Me	H	H	2	H	H	CF ₃	Pr	F
10	Me	Me	H	H	2	H	H	F	Pr	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Pr	Cl
	Me	Me	H	H	2	H	H	Cl	Pr	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Pr	OMe
15	Me	Me	H	H	2	H	H	OMe	Pr	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Pr	OEt
	Me	Me	H	H	2	H	H	OEt	Pr	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Pr	OCHF ₂
20	Me	Me	H	H	2	H	H	OCHF ₂	Pr	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Pr	CN
	Me	Me	H	H	2	H	H	CN	Pr	CF ₃
25	Me	Me	H	H	2	H	H	CF ₃	Pr	Me
	Me	Me	H	H	2	H	H	Me	Pr	CF ₃
	Me	Me	H	H	2	H	H	Me	Bu-t	F
	Me	Me	H	H	2	H	H	Me	Bu-t	Cl
30	Me	Me	H	H	2	H	H	Me	Bu-t	OCHF ₂
	Me	Me	H	H	2	H	H	Me	Bu-t	CN
	Me	Me	H	H	2	H	H	Cl	Bu-t	Cl
	Me	Me	H	H	2	H	H	OCHF ₂	Bu-t	Cl
35	Me	Me	H	H	2	H	H	OCHF ₂	Bu-t	OCHF ₂
	Me	Me	H	H	2	H	H	CF ₃	Bu-t	H
	Me	Me	H	H	2	H	H	CF ₃	Bu-t	F
40	Me	Me	H	H	2	H	H	CF ₃	Bu-t	Cl
	Me	Me	H	H	2	H	H	Cl	Bu-t	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Bu-t	OMe
	Me	Me	H	H	2	H	H	OMe	Bu-t	CF ₃
45	Me	Me	H	H	2	H	H	CF ₃	Bu-t	OEt
	Me	Me	H	H	2	H	H	OEt	Bu-t	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Bu-t	OCHF ₂
50	Me	Me	H	H	2	H	H	CF ₃	Bu-t	CN
	Me	Me	H	H	2	H	H	CF ₃	Bu-t	Me
	Me	Me	H	H	2	H	H	Me	Bu-t	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Bu-s	Cl
55	Me	Me	H	H	2	H	H	Cl	Bu-s	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Bu-i	Cl

	Me	Me	H	H	2	H	H	Cl	Bu-i	CF ₃
5	Me	Me	H	H	2	H	H	CF ₃	Bu	Cl
	Me	Me	H	H	2	H	H	Cl	Bu	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	1-Methylbutyl	Cl
	Me	Me	H	H	2	H	H	Cl	1-Methylbutyl	CF ₃
10	Me	Me	H	H	2	H	H	CF ₃	1-Ethylpropyl	Cl
	Me	Me	H	H	2	H	H	Cl	1-Ethylpropyl	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	1-Pentyl	Cl
15	Me	Me	H	H	2	H	H	Cl	1-Pentyl	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	1-Methylpentyl	Cl
	Me	Me	H	H	2	H	H	Cl	1-Methylpentyl	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	2-Ethylbutyl	Cl
20	Me	Me	H	H	2	H	H	Cl	2-Ethylbutyl	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	3,3-Dimethylbutyl	Cl
	Me	Me	H	H	2	H	H	Cl	3,3-Dimethylbutyl	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	1-Hexyl	Cl
25	Me	Me	H	H	2	H	H	Cl	1-Hexyl	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	1-Heptyl	Cl
	Me	Me	H	H	2	H	H	Cl	1-Heptyl	CF ₃
30	Me	Me	H	H	2	H	H	CF ₃	1-Octyl	Cl
	Me	Me	H	H	2	H	H	Cl	1-Octyl	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Ph	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ Ph	CF ₃
35	Me	Me	H	H	2	H	H	CF ₃	Pr-c	F
	Me	Me	H	H	2	H	H	CF ₃	Pr-c	Cl
	Me	Me	H	H	2	H	H	CF ₃	Pr-c	OMe
	Me	Me	H	H	2	H	H	CF ₃	Pr-c	OCHF ₂
40	Me	Me	H	H	2	H	H	CF ₃	Pr-c	CN
	Me	Me	H	H	2	H	H	CF ₃	Pen-c	Cl
	Me	Me	H	H	2	H	H	Cl	Pen-c	CF ₃
45	Me	Me	H	H	2	H	H	CF ₃	Hex-c	Cl
	Me	Me	H	H	2	H	H	Cl	Hex-c	CF ₃
	Me	Me	H	H	2	H	H	Me	CH ₂ Pr-c	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	CH ₂ Pr-c	Me
50	Me	Me	H	H	2	H	H	Cl	CH ₂ Pr-c	Cl
	Me	Me	H	H	2	H	H	OCHF ₂	CH ₂ Pr-c	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ Pr-c	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	CH ₂ Pr-c	OCHF ₂
55	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	F

5	Me	Me	H	H	2	H	H	F	CH ₂ Pr-c	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ Pr-c	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	OH
10	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	OMe
	Me	Me	H	H	2	H	H	OMe	CH ₂ Pr-c	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	OEt
	Me	Me	H	H	2	H	H	OEt	CH ₂ Pr-c	CF ₃
15	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	OPr-i
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	OPr
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	OBu-t
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	OCH ₂ Pr-c
20	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	OCH ₂ Bu-c
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	OPen-c
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	CH ₂ Pr-c	CF ₃
25	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	CN
	Me	Me	H	H	2	H	H	CN	CH ₂ Pr-c	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	Me
	Me	Me	H	H	2	H	H	Me	CH ₂ Pr-c	CF ₃
30	Me	Me	H	H	2	H	H	CF ₃	1-cyclopropylethyl	Cl
	Me	Me	H	H	2	H	H	Cl	1-cyclopropylethyl	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ (2-Methylcyclopropyl)	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ (2-Methylcyclopropyl)	CF ₃
35	Me	Me	H	H	2	H	H	CF ₃	CH ₂ (2,2-Dimethylcyclopropyl)	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ (2,2-Dimethylcyclopropyl)	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ (2-Chlorocyclopropyl)	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ (2-Chlorocyclopropyl)	CF ₃
40	Me	Me	H	H	2	H	H	CF ₃	CH ₂ (2,2-Dichlorocyclopropyl)	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ (2,2-Dichlorocyclopropyl)	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ (2-Fluorocyclopropyl)	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ (2-Fluorocyclopropyl)	CF ₃
45	Me	Me	H	H	2	H	H	CF ₃	CH ₂ (2,2-Difluorocyclopropyl)	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ (2,2-Difluorocyclopropyl)	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Bu-c	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ Bu-c	CF ₃
50	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pen-c	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ Pen-c	CF ₃
55	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pen-c	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ Pen-c	CF ₃

5	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Hex-c	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ Hex-c	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ Pr-c	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ CH ₂ Pr-c	CF ₃
10	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH=CH ₂	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ CH=CH ₂	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH=CHCl	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ CH=CHCl	CF ₃
15	Me	Me	H	H	2	H	H	Me	CH ₂ C≡CH	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	CH ₂ C≡CH	Me
	Me	Me	H	H	2	H	H	Cl	CH ₂ CCH	Cl
	Me	Me	H	H	2	H	H	OCHF ₂	CH ₂ C≡CH	Cl
20	Me	Me	H	H	2	H	H	Cl	CH ₂ C≡CH	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	CH ₂ C≡CH	OCHF ₂
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C≡CH	F
	Me	Me	H	H	2	H	H	F	CH ₂ C≡CH	CF ₃
25	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C≡CH	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ C≡CH	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C≡CH	OMe
	Me	Me	H	H	2	H	H	OMe	CH ₂ C≡CH	CF ₃
30	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C≡CH	OEt
	Me	Me	H	H	2	H	H	OEt	CH ₂ C≡CH	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C≡CH	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	CH ₂ C≡CH	CF ₃
35	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C≡CH	CN
	Me	Me	H	H	2	H	H	CN	CH ₂ C≡CH	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C≡CH	Me
	Me	Me	H	H	2	H	H	Me	CH ₂ C≡CH	CF ₃
40	Me	Me	H	H	2	H	H	CF ₃	CHMeC≡CH	Cl
	Me	Me	H	H	2	H	H	Cl	CHMeC≡CH	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C≡CMe	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ C≡CMe	CF ₃
45	Me	Me	H	H	2	H	H	Me	CHF ₂	F
	Me	Me	H	H	2	H	H	F	CHF ₂	Me
	Me	Me	H	H	2	H	H	Me	CHF ₂	Cl
	Me	Me	H	H	2	H	H	Cl	CHF ₂	Me
50	Me	Me	H	H	2	H	H	Me	CHF ₂	OMe
	Me	Me	H	H	2	H	H	OMe	CHF ₂	Me
	Me	Me	H	H	2	H	H	Me	CHF ₂	OCHF ₂
	Me	Me	H	H	2	H	H	Me	CHF ₂	OCHF ₂

	Me	Me	H	H	2	H	H	OCHF ₂	CHF ₂	Me
5	Me	Me	H	H	2	H	H	Me	CHF ₂	CN
	Me	Me	H	H	2	H	H	CN	CHF ₂	Me
	Me	Me	H	H	2	H	H	Me	CHF ₂	Me
	Me	Me	H	H	2	H	H	Et	CHF ₂	Cl
10	Me	Me	H	H	2	H	H	Cl	CHF ₂	Et
	Me	Me	H	H	2	H	H	Et	CHF ₂	Et
	Me	Me	H	H	2	H	H	Pr-i	CHF ₂	Cl
15	Me	Me	H	H	2	H	H	Cl	CHF ₂	Pr-i
	Me	Me	H	H	2	H	H	Cl	CHF ₂	Cl
	Me	Me	H	H	2	H	H	OCHF ₂	CHF ₂	Cl
	Me	Me	H	H	2	H	H	Cl	CHF ₂	OCHF ₂
20	Me	Me	H	H	2	H	H	OCHF ₂	CHF ₂	OCHF ₂
	Me	Me	H	H	2	H	H	CF ₃	CHF ₂	Cl
	Me	Me	H	H	2	H	H	Cl	CHF ₂	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CHF ₂	F
25	Me	Me	H	H	2	H	H	F	CHF ₂	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CHF ₂	OMe
	Me	Me	H	H	2	H	H	OMe	CHF ₂	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CHF ₂	OEt
30	Me	Me	H	H	2	H	H	OEt	CHF ₂	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CHF ₂	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	CHF ₂	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CHF ₂	CN
35	Me	Me	H	H	2	H	H	CN	CHF ₂	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CHF ₂	Me
	Me	Me	H	H	2	H	H	Me	CHF ₂	CF ₃
40	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CHF ₂	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ CHF ₂	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CF ₃	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ CF ₃	CF ₃
45	Me	Me	H	H	2	H	H	CF ₃	CH ₂ OH	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ OH	CF ₃
	Me	Me	H	H	2	H	H	Me	CH ₂ OMe	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	CH ₂ OMe	Me
50	Me	Me	H	H	2	H	H	Cl	CH ₂ OMe	Cl
	Me	Me	H	H	2	H	H	OCHF ₂	CH ₂ OMe	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ OMe	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	CH ₂ OMe	OCHF ₂
55	Me	Me	H	H	2	H	H	CF ₃	CH ₂ OMe	F

5	Me	Me	H	H	2	H	H	F	CH ₂ OMe	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ OMe	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ OMe	CF ₃
10	Me	Me	H	H	2	H	H	CF ₃	CH ₂ OMe	OMe
	Me	Me	H	H	2	H	H	OMe	CH ₂ OMe	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ OMe	OEt
15	Me	Me	H	H	2	H	H	OEt	CH ₂ OMe	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ OMe	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	CH ₂ OMe	CF ₃
20	Me	Me	H	H	2	H	H	CF ₃	CH ₂ OMe	CN
	Me	Me	H	H	2	H	H	CN	CH ₂ OMe	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ OMe	Me
25	Me	Me	H	H	2	H	H	Me	CH ₂ OMe	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ OEt	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ OEt	CF ₃
30	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ OH	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ CH ₂ OH	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ OMe	Cl
35	Me	Me	H	H	2	H	H	Cl	CH ₂ CH ₂ OMe	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ OEt	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ CH ₂ OEt	CF ₃
40	Me	Me	H	H	2	H	H	CF ₃	CH ₂ NHMe	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ NHMe	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ N(Me) ₂	Cl
45	Me	Me	H	H	2	H	H	Cl	CH ₂ N(Me) ₂	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ N(Me)C(=O)Me	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ N(Me)C(=O)Me	CF ₃
50	Me	Me	H	H	2	H	H	CF ₃	CH ₂ N(Me)C(=O)CF ₃	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ N(Me)C(=O)CF ₃	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ N(Me)SO ₂ Me	Cl
55	Me	Me	H	H	2	H	H	Cl	CH ₂ N(Me)SO ₂ Me	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ N(Me)SO ₂ CHF ₂	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ N(Me)SO ₂ CHF ₂	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ N(Me)SO ₂ CF ₃	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ N(Me)SO ₂ CF ₃	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ SMe	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ SMe	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ SO ₂ Me	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ SO ₂ Me	CF ₃

5	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ SMe	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ CH ₂ SMe	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ SO ₂ Me	Cl
10	Me	Me	H	H	2	H	H	Cl	CH ₂ CH ₂ SO ₂ Me	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CN	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ CN	CF ₃
15	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=O)OMe	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=O)OMe	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=O)OEt	Cl
20	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=O)OEt	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH(Me)C(=O)OMe	Cl
	Me	Me	H	H	2	H	H	Cl	CH(Me)C(=O)OMe	CF ₃
25	Me	Me	H	H	2	H	H	CF ₃	C(Me) ₂ C(=O)OMe	Cl
	Me	Me	H	H	2	H	H	Cl	C(Me) ₂ C(=O)OMe	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=O)NH ₂	Cl
30	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=O)NH ₂	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=O)NHMe	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=O)NHMe	CF ₃
35	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=O)N(Me) ₂	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=O)N(Me) ₂	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=O)Me	Cl
40	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=O)Me	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=NOMe)Me	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=NOMe)Me	CF ₃
45	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=O)CF ₃	Cl
	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=O)CF ₃	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ C(=O)Me	Cl
50	Me	Me	H	H	2	H	H	Cl	CH ₂ CH ₂ C(=O)Me	CF ₃
	Me	Me	H	H	2	H	H	Me	Ph	Me
	Me	Me	H	H	2	H	H	Me	Ph	F
55	Me	Me	H	H	2	H	H	Me	Ph	Cl
	Me	Me	H	H	2	H	H	Me	Ph	OCHF ₂
	Me	Me	H	H	2	H	H	Me	Ph	CN
55	Me	Me	H	H	2	H	H	Et	Ph	F
	Me	Me	H	H	2	H	H	Et	Ph	Cl
	Me	Me	H	H	2	H	H	Et	Ph	OCHF ₂
55	Me	Me	H	H	2	H	H	Et	Ph	CN
	Me	Me	H	H	2	H	H	Pr	Ph	F
	Me	Me	H	H	2	H	H	Pr	Ph	Cl

	Me	Me	H	H	2	H	H	Pr	Ph	OCHF ₂
5	Me	Me	H	H	2	H	H	Pr	Ph	CN
	Me	Me	H	H	2	H	H	Pr-i	Ph	F
	Me	Me	H	H	2	H	H	Pr-i	Ph	Cl
	Me	Me	H	H	2	H	H	Pr-i	Ph	OCHF ₂
10	Me	Me	H	H	2	H	H	Pr-i	Ph	CN
	Me	Me	H	H	2	H	H	Bu-t	Ph	Cl
	Me	Me	H	H	2	H	H	CH ₂ OMe	Ph	Cl
15	Me	Me	H	H	2	H	H	Cl	Ph	Cl
	Me	Me	H	H	2	H	H	OCHF ₂	Ph	Cl
	Me	Me	H	H	2	H	H	OCHF ₂	Ph	OCHF ₂
	Me	Me	H	H	2	H	H	CHF ₂	Ph	Cl
20	Me	Me	H	H	2	H	H	CF ₃	Ph	H
	Me	Me	H	H	2	H	H	CF ₃	Ph	Me
	Me	Me	H	H	2	H	H	Me	Ph	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Ph	Et
25	Me	Me	H	H	2	H	H	CF ₃	Ph	Pr-i
	Me	Me	H	H	2	H	H	CF ₃	Ph	CHF ₂
	Me	Me	H	H	2	H	H	CF ₃	Ph	CF ₃
30	Me	Me	H	H	2	H	H	CF ₃	Ph	F
	Me	Me	H	H	2	H	H	CF ₃	Ph	Cl
	Me	Me	H	H	2	H	H	Cl	Ph	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Ph	OH
35	Me	Me	H	H	2	H	H	OH	Ph	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Ph	OMe
	Me	Me	H	H	2	H	H	OMe	Ph	CF ₃
40	Me	Me	H	H	2	H	H	CF ₃	Ph	OEt
	Me	Me	H	H	2	H	H	OEt	Ph	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Ph	OPr-i
	Me	Me	H	H	2	H	H	CF ₃	Ph	OPr
45	Me	Me	H	H	2	H	H	CF ₃	Ph	OBu-t
	Me	Me	H	H	2	H	H	CF ₃	Ph	OCH ₂ Pr-c
	Me	Me	H	H	2	H	H	CF ₃	Ph	OCH ₂ CH=CH ₂
50	Me	Me	H	H	2	H	H	CF ₃	Ph	OCH ₂ C≡CH
	Me	Me	H	H	2	H	H	CF ₃	Ph	OCHF ₂
	Me	Me	H	H	2	H	H	OCHF ₂	Ph	CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Ph	OCH ₂ CHF ₂
55	Me	Me	H	H	2	H	H	CF ₃	Ph	OCH ₂ CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Ph	OCH ₂ C(=O)OMe

5	Me	Me	H	H	2	H	H	CF ₃	Ph	OCH(Me)C(=O)OMe
	Me	Me	H	H	2	H	H	CF ₃	Ph	OC(Me) ₂ C(=O)OMe
	Me	Me	H	H	2	H	H	CF ₃	Ph	OC(=O)Me
10	Me	Me	H	H	2	H	H	CF ₃	Ph	OC(=O)Et
	Me	Me	H	H	2	H	H	CF ₃	Ph	OC(=O)CH ₂ Ph
	Me	Me	H	H	2	H	H	CF ₃	Ph	OC(=O)CF ₃
15	Me	Me	H	H	2	H	H	CF ₃	Ph	OC(=O)Ph
	Me	Me	H	H	2	H	H	CF ₃	Ph	OSO ₂ Me
	Me	Me	H	H	2	H	H	CF ₃	Ph	OSO ₂ Et
20	Me	Me	H	H	2	H	H	CF ₃	Ph	OSO ₂ CH ₂ Ph
	Me	Me	H	H	2	H	H	CF ₃	Ph	OSO ₂ CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Ph	OSO ₂ Ph
25	Me	Me	H	H	2	H	H	CF ₃	Ph	SMe
	Me	Me	H	H	2	H	H	CF ₃	Ph	SOMe
	Me	Me	H	H	2	H	H	CF ₃	Ph	SO ₂ Me
30	Me	Me	H	H	2	H	H	CF ₃	Ph	SEt
	Me	Me	H	H	2	H	H	CF ₃	Ph	SOEt
	Me	Me	H	H	2	H	H	CF ₃	Ph	SO ₂ Et
35	Me	Me	H	H	2	H	H	CF ₃	Ph	SPr-i
	Me	Me	H	H	2	H	H	CF ₃	Ph	SOPr-i
	Me	Me	H	H	2	H	H	CF ₃	Ph	SO ₂ Pr-i
40	Me	Me	H	H	2	H	H	CF ₃	Ph	SPr
	Me	Me	H	H	2	H	H	CF ₃	Ph	SOPr
	Me	Me	H	H	2	H	H	CF ₃	Ph	SO ₂ Pr
45	Me	Me	H	H	2	H	H	CF ₃	Ph	SBu-t
	Me	Me	H	H	2	H	H	CF ₃	Ph	SOBu-t
	Me	Me	H	H	2	H	H	CF ₃	Ph	SO ₂ Bu-t
50	Me	Me	H	H	2	H	H	CF ₃	Ph	SCHF ₂
	Me	Me	H	H	2	H	H	CF ₃	Ph	SOCHF ₂
	Me	Me	H	H	2	H	H	CF ₃	Ph	SO ₂ CHF ₂
55	Me	Me	H	H	2	H	H	CF ₃	Ph	NH ₂
	Me	Me	H	H	2	H	H	CF ₃	Ph	NHMe
	Me	Me	H	H	2	H	H	CF ₃	Ph	N(Me) ₂
55	Me	Me	H	H	2	H	H	CF ₃	Ph	NHC(=O)Me
	Me	Me	H	H	2	H	H	CF ₃	Ph	N(Me)C(=O)Me
	Me	Me	H	H	2	H	H	CF ₃	Ph	NHSO ₂ Me
55	Me	Me	H	H	2	H	H	CF ₃	Ph	N(Me)SO ₂ Me
	Me	Me	H	H	2	H	H	CF ₃	Ph	NHSO ₂ CF ₃
	Me	Me	H	H	2	H	H	CF ₃	Ph	N(Me)SO ₂ CF ₃

	Me	Me	H	H	2	H	H	CF ₃	Ph	NHPh
5	Me	Me	H	H	2	H	H	CF ₃	Ph	N(Me)Ph
	Me	Me	H	H	2	H	H	CF ₃	Ph	CN
	Me	Me	H	H	2	H	H	CF ₃	Ph	C(=O)Me
	Me	Me	H	H	2	H	H	CF ₃	Ph	C(=O)OMe
10	Me	Me	H	H	2	H	H	CF ₃	Ph	C(=O)NH ₂
	Me	Me	H	H	2	H	H	CF ₃	Ph	C(=O)NHMe
	Me	Me	H	H	2	H	H	CF ₃	Ph	C(=O)N(Me) ₂
15	Me	Me	H	H	2	H	H	CF ₃	Ph	Imidazol-1-yl
	Me	Me	H	H	2	H	H	CF ₃	Ph	Pyrazol-1-yl
	Me	Me	H	H	2	H	H	CF ₃	Ph	1,2,4-Triazol-1-yl
	Me	Me	H	H	2	H	H	CF ₃	Ph	1,2,4-Triazol-4-yl
20	Me	Me	H	H	2	H	H	CF ₃	Ph	Tetrazol-1-yl
	Me	Me	H	H	2	H	H	CF ₃	Ph	Tetrazol-5-yl
	Me	Me	H	H	2	H	H	CF ₃	Ph	(4,6-Dimethoxypyrimidin-2-yl)oxy
25	Me	Me	H	H	2	H	H	CF ₃	Ph	(4,6-Dimethoxypyrimidin-2-yl)sulfonyl
	Me	Me	H	H	2	H	H	CF ₂ CF ₃	Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(2-Cl)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(2-F)Ph	Cl
30	Me	Me	H	H	2	H	H	CF ₃	(2-OMe)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(2-Me)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(2-NO ₂)Ph	Cl
35	Me	Me	H	H	2	H	H	CF ₃	(2-CN)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(2-C(=O)Me)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(2-C(=O)OMe)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(2-C(=O)OEt)Ph	Cl
40	Me	Me	H	H	2	H	H	CF ₃	(2-C(=O)OPr-i)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(2-C(=O)NH ₂)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(2-C(=O)NHMe)Ph	Cl
45	Me	Me	H	H	2	H	H	CF ₃	(2-C(=O)NMe ₂)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(3-Cl)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(3-F)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(3-OMe)Ph	Cl
50	Me	Me	H	H	2	H	H	CF ₃	(3-Me)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(3-NO ₂)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(3-CN)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(3-C(=O)Me)Ph	Cl
55	Me	Me	H	H	2	H	H	CF ₃	(3-C(=O)OMe)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(3-C(=O)OEt)Ph	Cl

5	Me	Me	H	H	2	H	H	CF ₃	(3-C(=O)OPr-i)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(3-C(=O)NH ₂)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(3-C(=O)NHMe)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(3-C(=O)NMe ₂)Ph	Cl
10	Me	Me	H	H	2	H	H	CF ₃	(4-Cl)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(4-F)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(4-OMe)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(4-Me)Ph	Cl
15	Me	Me	H	H	2	H	H	CF ₃	(4-NO ₂)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(4-CN)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(4-C(=O)Me)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(4-C(=O)OMe)Ph	Cl
20	Me	Me	H	H	2	H	H	CF ₃	(4-C(=O)OEt)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(4-C(=O)OPr-i)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(4-C(=O)NH ₂)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	(4-C(=O)NHMe)Ph	Cl
25	Me	Me	H	H	2	H	H	CF ₃	(4-C(=O)NMe ₂)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	Pyrimidin-2-yl	Cl
	Me	Me	H	H	2	H	H	CF ₃	4,6-Dimethoxypyrimidin-2-yl	Cl
	Me	Me	H	H	2	H	H	CF ₃	Thiophen-2-yl	Cl
30	Me	Me	H	H	2	H	H	CF ₃	Furan-2-yl	Cl
	Me	Me	H	H	2	H	H	CF ₃	SO ₂ Me	Cl
	Me	Me	H	H	2	H	H	CF ₃	SO ₂ Et	Cl
	Me	Me	H	H	2	H	H	CF ₃	SO ₂ Pr-i	Cl
35	Me	Me	H	H	2	H	H	CF ₃	SO ₂ CH ₂ Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	SO ₂ CHF ₂	Cl
	Me	Me	H	H	2	H	H	CF ₃	SO ₂ CF ₃	Cl
	Me	Me	H	H	2	H	H	CF ₃	SO ₂ Ph	Cl
40	Me	Me	H	H	2	H	H	CF ₃	C(=O)Me	Cl
	Me	Me	H	H	2	H	H	CF ₃	C(=O)Et	Cl
	Me	Me	H	H	2	H	H	CF ₃	C(=O)Pr-i	Cl
	Me	Me	H	H	2	H	H	CF ₃	C(=O)Bu-t	Cl
45	Me	Me	H	H	2	H	H	CF ₃	C(=O)Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	C(=O)CH ₂ Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	C(=O)CH ₂ Cl	Cl
	Me	Me	H	H	2	H	H	CF ₃	C(=O)CHCl ₂	Cl
50	Me	Me	H	H	2	H	H	CF ₃	C(=O)CF ₃	Cl
	Me	Me	H	H	2	H	H	CF ₃	C(=O)OMe	Cl
	Me	Me	H	H	2	H	H	CF ₃	C(=O)OPh	Cl
	Me	Me	H	H	2	H	H	CF ₃		

5	Me	Me	H	H	2	H	H	CF ₃	C(=O)OCH ₂ Ph	Cl
	Me	Me	H	H	2	H	H	CF ₃	C(=O)NHMe	Cl
	Me	Me	H	H	2	H	H	CF ₃	C(=O)N(Me) ₂	Cl
	Me	Me	H	H	2	H	H	CF ₃	C(=O)NHPh	Cl
10	Me	Me	H	H	2	H	H	CF ₃	NH ₂	Cl
	Me	Me	H	H	2	H	H	Cl	$-(CH_2)_2O-$ $-(CH_2)_3O-$ $-(CH_2)_3S-$ $-(CH_2)_3SO_2-$ $-(CH_2)_2O-$ $-(CH_2)_3O-$ $-(CH_2)_3S-$ $-(CH_2)_3SO_2-$ $-(CH_2)_4-$ $-(CH_2)_4-$	
	Me	Me	H	H	2	H	H	Cl		
	Me	Me	H	H	2	H	H	Cl		
15	Me	Me	H	H	2	H	H	Cl		
	Me	Me	H	H	2	H	H	CF ₃		
	Me	Me	H	H	2	H	H	CF ₃		
	Me	Me	H	H	2	H	H	CF ₃		
20	Me	Me	H	H	2	H	H	CF ₃		
	Me	Me	H	H	2	H	H	OMe		
	Me	Me	H	H	2	H	H	OCHF ₂		
25	H	H	H	H	2	H	H	CF ₃	Me	Cl
	Me	H	H	H	2	H	H	CF ₃	Me	Cl
	Me	H	Me	H	2	H	H	CF ₃	Me	Cl
	Me	Me	Me	H	2	H	H	CF ₃	Me	Cl
30	Me	Me	H	H	2	Me	H	CF ₃	Me	Cl
	Me	Me	H	H	2	Et	H	CF ₃	Me	Cl
	Me	Me	H	H	2	Pr-i	H	CF ₃	Me	Cl
	Me	Me	H	H	2	Me	Me	CF ₃	Me	Cl
35	Me	Et	H	H	2	H	H	CF ₃	Me	Cl
	Et	Et	H	H	2	H	H	CF ₃	Me	Cl
	Me	Pr-i	H	H	2	H	H	CF ₃	Me	Cl
40	Me	Pr	H	H	2	H	H	CF ₃	Me	Cl
	Me	Pr-c	H	H	2	H	H	CF ₃	Me	Cl
	Me	CH ₂ Pr-c	H	H	2	H	H	CF ₃	Me	Cl
45		-(CH ₂) ₂ -	H	H	2	H	H	CF ₃	Me	Cl
		-(CH ₂) ₃ -	H	H	2	H	H	CF ₃	Me	Cl
		-(CH ₂) ₄ -	H	H	2	H	H	CF ₃	Me	Cl
		-(CH ₂) ₅ -	H	H	2	H	H	CF ₃	Me	Cl
50	H	-(CH ₂) ₃ -	H	2	H	H	H	CF ₃	Me	Cl
	H	-(CH ₂) ₄ -	H	2	H	H	H	CF ₃	Me	Cl
	H	-(CH ₂) ₅ -	H	2	H	H	H	CF ₃	Me	Cl
	H	-(CH ₂) ₆ -	H	2	H	H	H	CF ₃	Me	Cl
55	Me	Me	H	H	1	H	H	Cl	H	Cl
	Me	Me	H	H	1	H	H	OCHF ₂	H	Cl

	Me	Me	H	H	1	H	H	OCHF ₂	H	OCHF ₂
5	Me	Me	H	H	1	H	H	CHF ₂	H	Cl
	Me	Me	H	H	1	H	H	CF ₃	H	F
	Me	Me	H	H	1	H	H	CF ₃	H	Cl
	Me	Me	H	H	1	H	H	CF ₃	H	OMe
10	Me	Me	H	H	1	H	H	CF ₃	H	OEt
	Me	Me	H	H	1	H	H	CF ₃	H	OCHF ₂
	Me	Me	H	H	1	H	H	CF ₃	H	CN
15	Me	Me	H	H	1	H	H	CF ₃	H	Me
	Me	Me	H	H	1	H	H	H	Me	Cl
	Me	Me	H	H	1	H	H	Me	Me	Me
	Me	Me	H	H	1	H	H	Me	Me	Cl
20	Me	Me	H	H	1	H	H	Cl	Me	Me
	Me	Me	H	H	1	H	H	Et	Me	Cl
	Me	Me	H	H	1	H	H	Cl	Me	Et
	Me	Me	H	H	1	H	H	Pr-i	Me	Cl
25	Me	Me	H	H	1	H	H	Cl	Me	Pr-i
	Me	Me	H	H	1	H	H	Bu-t	Me	Cl
	Me	Me	H	H	1	H	H	Cl	Me	Bu-t
30	Me	Me	H	H	1	H	H	Cl	Me	Cl
	Me	Me	H	H	1	H	H	CHF ₂	Me	Cl
	Me	Me	H	H	1	H	H	Cl	Me	CHF ₂
	Me	Me	H	H	1	H	H	OCHF ₂	Me	H
35	Me	Me	H	H	1	H	H	OCHF ₂	Me	Cl
	Me	Me	H	H	1	H	H	Cl	Me	OCHF ₂
	Me	Me	H	H	1	H	H	OCHF ₂	Me	OCHF ₂
40	Me	Me	H	H	1	H	H	CF ₃	Me	H
	Me	Me	H	H	1	H	H	CF ₃	Me	Cl
	Me	Me	H	H	1	H	H	Cl	Me	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Me	F
45	Me	Me	H	H	1	H	H	F	Me	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Me	OH
	Me	Me	H	H	1	H	H	OH	Me	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Me	OMe
50	Me	Me	H	H	1	H	H	OMe	Me	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Me	OEt
	Me	Me	H	H	1	H	H	OEt	Me	CF ₃
55	Me	Me	H	H	1	H	H	CF ₃	Me	OPr-i
	Me	Me	H	H	1	H	H	CF ₃	Me	OPr

5	Me	Me	H	H	1	H	H	CF ₃	Me	OBu-t
	Me	Me	H	H	1	H	H	CF ₃	Me	OBu-s
	Me	Me	H	H	1	H	H	CF ₃	Me	OBu-i
	Me	Me	H	H	1	H	H	CF ₃	Me	OBu
10	Me	Me	H	H	1	H	H	CF ₃	Me	O(2-Pen)
	Me	Me	H	H	1	H	H	CF ₃	Me	O(3-Pen)
	Me	Me	H	H	1	H	H	CF ₃	Me	OPen-n
	Me	Me	H	H	1	H	H	CF ₃	Me	O(2-Hex)
15	Me	Me	H	H	1	H	H	CF ₃	Me	O(3-Hex)
	Me	Me	H	H	1	H	H	CF ₃	Me	OHex-n
	Me	Me	H	H	1	H	H	CF ₃	Me	OPen-c
	Me	Me	H	H	1	H	H	CF ₃	Me	OHex-c
20	Me	Me	H	H	1	H	H	CF ₃	Me	OCH ₂ Pr-c
	Me	Me	H	H	1	H	H	CF ₃	Me	OCH ₂ Bu-c
	Me	Me	H	H	1	H	H	CF ₃	Me	OCH ₂ Pen-c
	Me	Me	H	H	1	H	H	CF ₃	Me	OCH ₂ Hex-c
25	Me	Me	H	H	1	H	H	CF ₃	Me	OCH ₂ CH=CH ₂
	Me	Me	H	H	1	H	H	CF ₃	Me	OCH ₂ C≡CH
	Me	Me	H	H	1	H	H	CF ₃	Me	OCHF ₂
	Me	Me	H	H	1	H	H	CF ₃	Me	CF ₃
30	Me	Me	H	H	1	H	H	OCHF ₂	Me	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Me	OCH ₂ CHF ₂
	Me	Me	H	H	1	H	H	OCH ₂ CHF ₂	Me	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Me	OCH ₂ CF ₃
35	Me	Me	H	H	1	H	H	OCH ₂ CF ₃	Me	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Me	OCH ₂ CN
	Me	Me	H	H	1	H	H	CF ₃	Me	OCH ₂ C(=O)OEt
	Me	Me	H	H	1	H	H	CF ₃	Me	OCH(Me)C(=O)OEt
40	Me	Me	H	H	1	H	H	CF ₃	Me	OCH ₂ C(=O)NH ₂
	Me	Me	H	H	1	H	H	CF ₃	Me	OCH ₂ C(=O)NHMe
	Me	Me	H	H	1	H	H	CF ₃	Me	OCH ₂ C(=O)N(Me) ₂
	Me	Me	H	H	1	H	H	CF ₃	Me	OCH ₂ Ph
45	Me	Me	H	H	1	H	H	CF ₃	Me	OPh
	Me	Me	H	H	1	H	H	CF ₃	Me	O(2-Cl)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(2-Br)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(2-F)Ph
50	Me	Me	H	H	1	H	H	CF ₃	Me	O(2-Me)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(2-OMe)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(2-NO ₂)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(2-CN)Ph

	Me	Me	H	H	1	H	H	CF ₃	Me	O(2-C(=O)OMe)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(3-Cl)Ph
5	Me	Me	H	H	1	H	H	CF ₃	Me	O(3-Br)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(3-F)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(3-Me)Ph
10	Me	Me	H	H	1	H	H	CF ₃	Me	O(3-OMe)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(3-NO ₂)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(3-CN)Ph
15	Me	Me	H	H	1	H	H	CF ₃	Me	O(3-C(=O)OMe)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(4-Cl)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(4-Br)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(4-F)Ph
20	Me	Me	H	H	1	H	H	CF ₃	Me	O(4-Me)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(4-OMe)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(4-NO ₂)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	O(4-CN)Ph
25	Me	Me	H	H	1	H	H	CF ₃	Me	O(4-C(=O)OMe)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	OC(=O)Me
	Me	Me	H	H	1	H	H	CF ₃	Me	OC(=O)Et
30	Me	Me	H	H	1	H	H	CF ₃	Me	OC(=O)CH ₂ Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	OC(=O)CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Me	OC(=O)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	OSO ₂ Me
35	Me	Me	H	H	1	H	H	CF ₃	Me	OSO ₂ Et
	Me	Me	H	H	1	H	H	CF ₃	Me	OSO ₂ CH ₂ Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	OSO ₂ CF ₃
40	Me	Me	H	H	1	H	H	CF ₃	Me	OSO ₂ Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	SMe
	Me	Me	H	H	1	H	H	CF ₃	Me	SO ₂ Me
	Me	Me	H	H	1	H	H	CF ₃	Me	SEt
45	Me	Me	H	H	1	H	H	CF ₃	Me	SO ₂ Et
	Me	Me	H	H	1	H	H	CF ₃	Me	SPr
	Me	Me	H	H	1	H	H	CF ₃	Me	SO ₂ Pr
50	Me	Me	H	H	1	H	H	CF ₃	Me	SPr-i
	Me	Me	H	H	1	H	H	CF ₃	Me	SO ₂ Pr-i
	Me	Me	H	H	1	H	H	CF ₃	Me	SBU-t
	Me	Me	H	H	1	H	H	CF ₃	Me	SO ₂ BU-t
55	Me	Me	H	H	1	H	H	CF ₃	Me	SCHF ₂
	Me	Me	H	H	1	H	H	CF ₃	Me	SO ₂ CHF ₂

	Me	Me	H	H	1	H	H	CF ₃	Me	SCF ₃
	Me	Me	H	H	1	H	H	CF ₃	Me	SO ₂ CF ₃
5	Me	Me	H	H	1	H	H	CF ₃	Me	SPh
	Me	Me	H	H	1	H	H	CF ₃	Me	SO ₂ Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	SCH ₂ Ph
10	Me	Me	H	H	1	H	H	CF ₃	Me	SO ₂ CH ₂ Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	SCH ₂ C(=O)OEt
	Me	Me	H	H	1	H	H	CF ₃	Me	SO ₂ CH ₂ C(=O)OEt
15	Me	Me	H	H	1	H	H	CF ₃	Me	SCH(Me)C(=O)OEt
	Me	Me	H	H	1	H	H	CF ₃	Me	SO ₂ CH(Me)C(=O)OEt
	Me	Me	H	H	1	H	H	CF ₃	Me	SCH ₂ C(=O)NH ₂
	Me	Me	H	H	1	H	H	CF ₃	Me	SO ₂ CH ₂ C(=O)NH ₂
20	Me	Me	H	H	1	H	H	CF ₃	Me	SCH ₂ C(=O)NHMe
	Me	Me	H	H	1	H	H	CF ₃	Me	SO ₂ CH ₂ C(=O)NHMe
	Me	Me	H	H	1	H	H	CF ₃	Me	SCH ₂ C(=O)N(Me) ₂
	Me	Me	H	H	1	H	H	CF ₃	Me	SO ₂ CH ₂ C(=O)N(Me) ₂
25	Me	Me	H	H	1	H	H	CF ₃	Me	NH ₂
	Me	Me	H	H	1	H	H	CF ₃	Me	NHMe
	Me	Me	H	H	1	H	H	CF ₃	Me	N(Me) ₂
	Me	Me	H	H	1	H	H	CF ₃	Me	NHC(=O)Me
30	Me	Me	H	H	1	H	H	CF ₃	Me	N(Me)C(=O)Me
	Me	Me	H	H	1	H	H	CF ₃	Me	NHSO ₂ Me
	Me	Me	H	H	1	H	H	CF ₃	Me	N(Me)SO ₂ Me
35	Me	Me	H	H	1	H	H	CF ₃	Me	NHSO ₂ CHF ₂
	Me	Me	H	H	1	H	H	CF ₃	Me	N(Me)SO ₂ CHF ₂
	Me	Me	H	H	1	H	H	CF ₃	Me	NHSO ₂ CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Me	N(Me)SO ₂ CF ₃
40	Me	Me	H	H	1	H	H	CF ₃	Me	NHPh
	Me	Me	H	H	1	H	H	CF ₃	Me	N(Me)Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	CN
45	Me	Me	H	H	1	H	H	CN	Me	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Me	C(=O)OMe
	Me	Me	H	H	1	H	H	CF ₃	Me	C(=O)OCH ₂ Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	C(=O)OPh
50	Me	Me	H	H	1	H	H	CF ₃	Me	C(=O)NH ₂
	Me	Me	H	H	1	H	H	CF ₃	Me	C(=O)NHMe
	Me	Me	H	H	1	H	H	CF ₃	Me	C(=O)N(Me) ₂
55	Me	Me	H	H	1	H	H	CF ₃	Me	C(=O)Me
	Me	Me	H	H	1	H	H	CF ₃	Me	C(=O)CF ₃

	Me	Me	H	H	1	H	H	CF ₃	Me	C(=O)CH ₂ Ph
	Me	Me	H	H	1	H	H	CF ₃	Me	C(=O)Ph
5	Me	Me	H	H	1	H	H	CF ₃	Me	Me
	Me	Me	H	H	1	H	H	Me	Me	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Me	Et
10	Me	Me	H	H	1	H	H	CF ₃	Me	Pr-i
	Me	Me	H	H	1	H	H	CF ₃	Me	Pr
	Me	Me	H	H	1	H	H	CF ₃	Me	CH ₂ OMe
15	Me	Me	H	H	1	H	H	CF ₃	Me	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Me	CHF ₂
	Me	Me	H	H	1	H	H	CF ₃	Me	Ph
	Me	Me	H	H	1	H	H	CF ₂ CF ₃	Me	Cl
20	Me	Me	H	H	1	H	H	Ph	Me	Me
	Me	Me	H	H	1	H	H	Ph	Me	Cl
	Me	Me	H	H	1	H	H	Ph	Me	OEt
	Me	Me	H	H	1	H	H	Ph	Me	CF ₃
25	Me	Me	H	H	1	H	H	Ph	Me	Ph
	Me	Me	H	H	1	H	H	Cl	Et	Cl
	Me	Me	H	H	1	H	H	OCHF ₂	Et	Cl
	Me	Me	H	H	1	H	H	Cl	Et	OCHF ₂
30	Me	Me	H	H	1	H	H	OCHF ₂	Et	OCHF ₂
	Me	Me	H	H	1	H	H	CF ₃	Et	F
	Me	Me	H	H	1	H	H	F	Et	CF ₃
35	Me	Me	H	H	1	H	H	CF ₃	Et	Cl
	Me	Me	H	H	1	H	H	Cl	Et	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Et	OMe
	Me	Me	H	H	1	H	H	OMe	Et	CF ₃
40	Me	Me	H	H	1	H	H	CF ₃	Et	OEt
	Me	Me	H	H	1	H	H	OEt	Et	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Et	OCHF ₂
45	Me	Me	H	H	1	H	H	OCHF ₂	Et	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Et	CN
	Me	Me	H	H	1	H	H	CN	Et	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Et	Me
50	Me	Me	H	H	1	H	H	Me	Et	CF ₃
	Me	Me	H	H	1	H	H	Cl	Pr-i	Cl
	Me	Me	H	H	1	H	H	OCHF ₂	Pr-i	Cl
	Me	Me	H	H	1	H	H	Cl	Pr-i	OCHF ₂
55	Me	Me	H	H	1	H	H	OCHF ₂	Pr-i	OCHF ₂
	Me	Me	H	H	1	H	H	CF ₃	Pr-i	F

5	Me	Me	H	H	1	H	H	F	Pr-i	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Pr-i	Cl
	Me	Me	H	H	1	H	H	Cl	Pr-i	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Pr-i	OMe
10	Me	Me	H	H	1	H	H	OMe	Pr-i	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Pr-i	OEt
	Me	Me	H	H	1	H	H	OEt	Pr-i	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Pr-i	OCHF ₂
15	Me	Me	H	H	1	H	H	OCHF ₂	Pr-i	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Pr-i	CN
	Me	Me	H	H	1	H	H	CN	Pr-i	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Pr-i	Me
20	Me	Me	H	H	1	H	H	Me	Pr-i	CF ₃
	Me	Me	H	H	1	H	H	Cl	Pr	Cl
	Me	Me	H	H	1	H	H	OCHF ₂	Pr	Cl
	Me	Me	H	H	1	H	H	Cl	Pr	OCHF ₂
25	Me	Me	H	H	1	H	H	OCHF ₂	Pr	OCHF ₂
	Me	Me	H	H	1	H	H	CF ₃	Pr	F
	Me	Me	H	H	1	H	H	F	Pr	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Pr	Cl
30	Me	Me	H	H	1	H	H	Cl	Pr	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Pr	OMe
	Me	Me	H	H	1	H	H	OMe	Pr	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Pr	OEt
35	Me	Me	H	H	1	H	H	OEt	Pr	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Pr	OCHF ₂
	Me	Me	H	H	1	H	H	OCHF ₂	Pr	CF ₃
40	Me	Me	H	H	1	H	H	CF ₃	Pr	CN
	Me	Me	H	H	1	H	H	CN	Pr	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Pr	Me
	Me	Me	H	H	1	H	H	Me	Pr	CF ₃
45	Me	Me	H	H	1	H	H	Cl	Bu-t	Cl
	Me	Me	H	H	1	H	H	OCHF ₂	Bu-t	Cl
	Me	Me	H	H	1	H	H	OCHF ₂	Bu-t	OCHF ₂
	Me	Me	H	H	1	H	H	CF ₃	Bu-t	H
50	Me	Me	H	H	1	H	H	CF ₃	Bu-t	F
	Me	Me	H	H	1	H	H	CF ₃	Bu-t	Cl
	Me	Me	H	H	1	H	H	Cl	Bu-t	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Bu-t	OMe
55	Me	Me	H	H	1	H	H	OMe	Bu-t	CF ₃

	Me	Me	H	H	1	H	H	CF ₃	Bu-t	OEt
5	Me	Me	H	H	1	H	H	OEt	Bu-t	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Bu-t	OCHF ₂
	Me	Me	H	H	1	H	H	CF ₃	Bu-t	CN
	Me	Me	H	H	1	H	H	CF ₃	Bu-t	Me
10	Me	Me	H	H	1	H	H	Me	Bu-t	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Bu-s	Cl
	Me	Me	H	H	1	H	H	Cl	Bu-s	CF ₃
15	Me	Me	H	H	1	H	H	CF ₃	Bu-i	Cl
	Me	Me	H	H	1	H	H	Cl	Bu-i	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Bu	Cl
	Me	Me	H	H	1	H	H	Cl	Bu	CF ₃
20	Me	Me	H	H	1	H	H	CF ₃	1-Methylbutyl	Cl
	Me	Me	H	H	1	H	H	Cl	1-Methylbutyl	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	1-Ethylpropyl	Cl
	Me	Me	H	H	1	H	H	Cl	1-Ethylpropyl	CF ₃
25	Me	Me	H	H	1	H	H	CF ₃	1-Pentyl	Cl
	Me	Me	H	H	1	H	H	Cl	1-Pentyl	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	1-Methylpentyl	Cl
30	Me	Me	H	H	1	H	H	Cl	1-Methylpentyl	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	2-Ethylbutyl	Cl
	Me	Me	H	H	1	H	H	Cl	2-Ethylbutyl	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	3,3-Dimethylbutyl	Cl
35	Me	Me	H	H	1	H	H	Cl	3,3-Dimethylbutyl	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	1-Hexyl	Cl
	Me	Me	H	H	1	H	H	Cl	1-Hexyl	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	1-Heptyl	Cl
40	Me	Me	H	H	1	H	H	Cl	1-Heptyl	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	1-Octyl	Cl
	Me	Me	H	H	1	H	H	Cl	1-Octyl	CF ₃
45	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Ph	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ Ph	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Pr-c	Cl
	Me	Me	H	H	1	H	H	CF ₃	Pen-c	Cl
50	Me	Me	H	H	1	H	H	Cl	Pen-c	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Hex-c	Cl
	Me	Me	H	H	1	H	H	Cl	Hex-c	CF ₃
	Me	Me	H	H	1	H	H	Cl	CH ₂ Pr-c	Cl
55	Me	Me	H	H	1	H	H	OCHF ₂	CH ₂ Pr-c	Cl

5	Me	Me	H	H	1	H	H	Cl	CH ₂ Pr-c	OCHF ₂
	Me	Me	H	H	1	H	H	OCHF ₂	CH ₂ Pr-c	OCHF ₂
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pr-c	F
10	Me	Me	H	H	1	H	H	F	CH ₂ Pr-c	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pr-c	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ Pr-c	CF ₃
15	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pr-c	CN
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pr-c	OH
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pr-c	OMe
20	Me	Me	H	H	1	H	H	OMe	CH ₂ Pr-c	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pr-c	OEt
	Me	Me	H	H	1	H	H	OEt	CH ₂ Pr-c	CF ₃
25	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pr-c	OPr-i
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pr-c	OPr
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pr-c	OBu-t
30	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pr-c	OCH ₂ Pr-c
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pr-c	OCH ₂ Bu-c
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pr-c	OPen-c
35	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pr-c	OCHF ₂
	Me	Me	H	H	1	H	H	OCHF ₂	CH ₂ Pr-c	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pr-c	CN
40	Me	Me	H	H	1	H	H	CN	CH ₂ Pr-c	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pr-c	Me
	Me	Me	H	H	1	H	H	Me	CH ₂ Pr-c	CF ₃
45	Me	Me	H	H	1	H	H	CF ₃	1-cyclopropylethyl	Cl
	Me	Me	H	H	1	H	H	Cl	1-cyclopropylethyl	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ (2-Methyl-cyclopropyl)	Cl
50	Me	Me	H	H	1	H	H	Cl	CH ₂ (2-Methyl-cyclopropyl)	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ (2,2-Dimethyl-cyclopropyl)	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ (2,2-Dimethyl-cyclopropyl)	CF ₃
55	Me	Me	H	H	1	H	H	CF ₃	CH ₂ (2-Chloro-cyclopropyl)	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ (2-Chloro-cyclopropyl)	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ (2,2-Dichloro-cyclopropyl)	Cl
50	Me	Me	H	H	1	H	H	Cl	CH ₂ (2,2-Dichloro-cyclopropyl)	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ (2-Fluoro-cyclopropyl)	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ (2-Fluoro-cyclopropyl)	CF ₃
55	Me	Me	H	H	1	H	H	CF ₃	CH ₂ (2,2-Difluoro-cyclopropyl)	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ (2,2-Difluoro-cyclopropyl)	CF ₃

	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Bu-c	Cl
5	Me	Me	H	H	1	H	H	Cl	CH ₂ Bu-c	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Pen-c	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ Pen-c	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ Hex-c	Cl
10	Me	Me	H	H	1	H	H	Cl	CH ₂ Hex-c	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ CH ₂ Pr-c	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ CH ₂ Pr-c	CF ₃
15	Me	Me	H	H	1	H	H	CF ₃	CH ₂ CH=CH ₂	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ CH=CH ₂	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ CH=CHCl	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ CH=CHCl	CF ₃
20	Me	Me	H	H	1	H	H	Cl	CH ₂ C≡CH	Cl
	Me	Me	H	H	1	H	H	OCHF ₂	CH ₂ C≡CH	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ C≡CH	OCHF ₂
	Me	Me	H	H	1	H	H	OCHF ₂	CH ₂ C≡CH	OCHF ₂
25	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C≡CH	F
	Me	Me	H	H	1	H	H	F	CH ₂ C≡CH	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C≡CH	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ C≡CH	CF ₃
30	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C≡CH	OMe
	Me	Me	H	H	1	H	H	OMe	CH ₂ C≡CH	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C≡CH	OEt
35	Me	Me	H	H	1	H	H	OEt	CH ₂ C≡CH	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C≡CH	OCHF ₂
	Me	Me	H	H	1	H	H	OCHF ₂	CH ₂ C≡CH	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C≡CH	CN
40	Me	Me	H	H	1	H	H	CN	CH ₂ C≡CH	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C≡CH	Me
	Me	Me	H	H	1	H	H	Me	CH ₂ C≡CH	CF ₃
45	Me	Me	H	H	1	H	H	CF ₃	CHMeC≡CH	Cl
	Me	Me	H	H	1	H	H	Cl	CHMeC≡CH	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C≡CMe	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ C≡CMe	CF ₃
50	Me	Me	H	H	1	H	H	Cl	CHF ₂	Cl
	Me	Me	H	H	1	H	H	OCHF ₂	CHF ₂	Cl
	Me	Me	H	H	1	H	H	Cl	CHF ₂	OCHF ₂
	Me	Me	H	H	1	H	H	OCHF ₂	CHF ₂	OCHF ₂
55	Me	Me	H	H	1	H	H	CF ₃	CHF ₂	Cl
	Me	Me	H	H	1	H	H	Cl	CHF ₂	CF ₃

	Me	Me	H	H	1	H	H	CF ₃	CHF ₂	F
5	Me	Me	H	H	1	H	H	F	CHF ₂	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CHF ₂	OMe
	Me	Me	H	H	1	H	H	OMe	CHF ₂	CF ₃
10	Me	Me	H	H	1	H	H	CF ₃	CHF ₂	OEt
	Me	Me	H	H	1	H	H	OEt	CHF ₂	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CHF ₂	OCHF ₂
	Me	Me	H	H	1	H	H	OCHF ₂	CHF ₂	CF ₃
15	Me	Me	H	H	1	H	H	CF ₃	CHF ₂	CN
	Me	Me	H	H	1	H	H	CN	CHF ₂	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CHF ₂	Me
	Me	Me	H	H	1	H	H	Me	CHF ₂	CF ₃
20	Me	Me	H	H	1	H	H	Me	CHF ₂	Cl
	Me	Me	H	H	1	H	H	Cl	CHF ₂	Me
	Me	Me	H	H	1	H	H	Et	CHF ₂	Cl
25	Me	Me	H	H	1	H	H	Cl	CHF ₂	Et
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ CHF ₂	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ CHF ₂	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ CF ₃	Cl
30	Me	Me	H	H	1	H	H	Cl	CH ₂ CF ₃	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ OH	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ OH	CF ₃
	Me	Me	H	H	1	H	H	Cl	CH ₂ OMe	Cl
35	Me	Me	H	H	1	H	H	OCHF ₂	CH ₂ OMe	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ OMe	OCHF ₂
	Me	Me	H	H	1	H	H	OCHF ₂	CH ₂ OMe	OCHF ₂
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ OMe	F
40	Me	Me	H	H	1	H	H	F	CH ₂ OMe	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ OMe	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ OMe	CF ₃
45	Me	Me	H	H	1	H	H	CF ₃	CH ₂ OMe	OMe
	Me	Me	H	H	1	H	H	OMe	CH ₂ OMe	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ OMe	OEt
	Me	Me	H	H	1	H	H	OEt	CH ₂ OMe	CF ₃
50	Me	Me	H	H	1	H	H	CF ₃	CH ₂ OMe	OCHF ₂
	Me	Me	H	H	1	H	H	OCHF ₂	CH ₂ OMe	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ OMe	CN
	Me	Me	H	H	1	H	H	CN	CH ₂ OMe	CF ₃
55	Me	Me	H	H	1	H	H	CF ₃	CH ₂ OMe	Me

5	Me	Me	H	H	1	H	H	Me	CH ₂ OMe	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ OEt	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ OEt	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ CH ₂ OH	Cl
10	Me	Me	H	H	1	H	H	Cl	CH ₂ CH ₂ OH	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ CH ₂ OMe	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ CH ₂ OMe	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ CH ₂ OEt	Cl
15	Me	Me	H	H	1	H	H	Cl	CH ₂ CH ₂ OEt	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ NHMe	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ NHMe	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ N(Me) ₂	Cl
20	Me	Me	H	H	1	H	H	Cl	CH ₂ N(Me) ₂	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ N(Me)C(=O)Me	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ N(Me)C(=O)Me	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ N(Me)C(=O)CF ₃	Cl
25	Me	Me	H	H	1	H	H	Cl	CH ₂ N(Me)C(=O)CF ₃	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ N(Me)SO ₂ Me	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ N(Me)SO ₂ Me	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ N(Me)SO ₂ CHF ₂	Cl
30	Me	Me	H	H	1	H	H	Cl	CH ₂ N(Me)SO ₂ CHF ₂	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ N(Me)SO ₂ CF ₃	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ N(Me)SO ₂ CF ₃	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ SMe	Cl
35	Me	Me	H	H	1	H	H	Cl	CH ₂ SMe	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ SO ₂ Me	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ SO ₂ Me	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ CH ₂ SMe	Cl
40	Me	Me	H	H	1	H	H	Cl	CH ₂ CH ₂ SMe	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ CH ₂ SO ₂ Me	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ CH ₂ SO ₂ Me	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ CN	Cl
45	Me	Me	H	H	1	H	H	Cl	CH ₂ CN	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C(=O)OMe	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ C(=O)OMe	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C(=O)OEt	Cl
50	Me	Me	H	H	1	H	H	Cl	CH ₂ C(=O)OEt	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH(Me)C(=O)OMe	Cl
	Me	Me	H	H	1	H	H	Cl	CH(Me)C(=O)OMe	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH(Me)C(=O)OMe	Cl
55	Me	Me	H	H	1	H	H	Cl	CH(Me)C(=O)OMe	CF ₃

5	Me	Me	H	H	1	H	H	CF ₃	C(Me) ₂ C(=O)OMe	Cl
	Me	Me	H	H	1	H	H	Cl	C(Me) ₂ C(=O)OMe	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C(=O)NH ₂	Cl
10	Me	Me	H	H	1	H	H	Cl	CH ₂ C(=O)NH ₂	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C(=O)NHMe	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ C(=O)NHMe	CF ₃
15	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C(=O)N(Me) ₂	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ C(=O)N(Me) ₂	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C(=O)Me	Cl
20	Me	Me	H	H	1	H	H	Cl	CH ₂ C(=O)Me	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C(=NOMe)Me	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ C(=NOMe)Me	CF ₃
25	Me	Me	H	H	1	H	H	CF ₃	CH ₂ C(=O)CF ₃	Cl
	Me	Me	H	H	1	H	H	Cl	CH ₂ C(=O)CF ₃	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	CH ₂ CH ₂ C(=O)Me	Cl
30	Me	Me	H	H	1	H	H	Cl	CH ₂ CH ₂ C(=O)Me	CF ₃
	Me	Me	H	H	1	H	H	Me	Ph	Me
	Me	Me	H	H	1	H	H	Me	Ph	Cl
35	Me	Me	H	H	1	H	H	Et	Ph	Cl
	Me	Me	H	H	1	H	H	Pr	Ph	Cl
	Me	Me	H	H	1	H	H	Pr-i	Ph	Cl
40	Me	Me	H	H	1	H	H	Bu-t	Ph	Cl
	Me	Me	H	H	1	H	H	CH ₂ OMe	Ph	Cl
	Me	Me	H	H	1	H	H	Cl	Ph	Cl
45	Me	Me	H	H	1	H	H	OCHF ₂	Ph	Cl
	Me	Me	H	H	1	H	H	OCHF ₂	Ph	OCHF ₂
	Me	Me	H	H	1	H	H	CHF ₂	Ph	Cl
50	Me	Me	H	H	1	H	H	CF ₃	Ph	H
	Me	Me	H	H	1	H	H	CF ₃	Ph	Me
	Me	Me	H	H	1	H	H	Me	Ph	CF ₃
55	Me	Me	H	H	1	H	H	CF ₃	Ph	Et
	Me	Me	H	H	1	H	H	CF ₃	Ph	Pr-i
	Me	Me	H	H	1	H	H	CF ₃	Ph	CHF ₂
	Me	Me	H	H	1	H	H	CF ₃	Ph	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Ph	F
	Me	Me	H	H	1	H	H	CF ₃	Ph	Cl
	Me	Me	H	H	1	H	H	Cl	Ph	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Ph	OH
	Me	Me	H	H	1	H	H	OH	Ph	CF ₃

	Me	Me	H	H	1	H	H	CF ₃	Ph	OMe
5	Me	Me	H	H	1	H	H	OMe	Ph	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Ph	OEt
	Me	Me	H	H	1	H	H	OEt	Ph	CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Ph	OPr-i
10	Me	Me	H	H	1	H	H	CF ₃	Ph	OPr
	Me	Me	H	H	1	H	H	CF ₃	Ph	OBu-t
	Me	Me	H	H	1	H	H	CF ₃	Ph	OCH ₂ Pr-c
15	Me	Me	H	H	1	H	H	CF ₃	Ph	OCH ₂ CH=CH ₂
	Me	Me	H	H	1	H	H	CF ₃	Ph	OCH ₂ C≡CH
	Me	Me	H	H	1	H	H	CF ₃	Ph	OCHF ₂
	Me	Me	H	H	1	H	H	OCHF ₂	Ph	CF ₃
20	Me	Me	H	H	1	H	H	CF ₃	Ph	OCH ₂ CHF ₂
	Me	Me	H	H	1	H	H	CF ₃	Ph	OCH ₂ CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Ph	OCH ₂ C(=O)OMe
	Me	Me	H	H	1	H	H	CF ₃	Ph	OCH(Me)C(=O)OMe
25	Me	Me	H	H	1	H	H	CF ₃	Ph	OC(Me) ₂ C(=O)OMe
	Me	Me	H	H	1	H	H	CF ₃	Ph	OC(=O)Me
	Me	Me	H	H	1	H	H	CF ₃	Ph	OC(=O)Et
	Me	Me	H	H	1	H	H	CF ₃	Ph	OC(=O)CH ₂ Ph
30	Me	Me	H	H	1	H	H	CF ₃	Ph	OC(=O)CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Ph	OC(=O)Ph
	Me	Me	H	H	1	H	H	CF ₃	Ph	OSO ₂ Me
35	Me	Me	H	H	1	H	H	CF ₃	Ph	OSO ₂ Et
	Me	Me	H	H	1	H	H	CF ₃	Ph	OSO ₂ CH ₂ Ph
	Me	Me	H	H	1	H	H	CF ₃	Ph	OSO ₂ CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Ph	OSO ₂ Ph
40	Me	Me	H	H	1	H	H	CF ₃	Ph	SMe
	Me	Me	H	H	1	H	H	CF ₃	Ph	SO ₂ Me
	Me	Me	H	H	1	H	H	CF ₃	Ph	SEt
45	Me	Me	H	H	1	H	H	CF ₃	Ph	SO ₂ Et
	Me	Me	H	H	1	H	H	CF ₃	Ph	SPr-i
	Me	Me	H	H	1	H	H	CF ₃	Ph	SO ₂ Pr-i
	Me	Me	H	H	1	H	H	CF ₃	Ph	SPr
50	Me	Me	H	H	1	H	H	CF ₃	Ph	SO ₂ Pr
	Me	Me	H	H	1	H	H	CF ₃	Ph	SBu-t
	Me	Me	H	H	1	H	H	CF ₃	Ph	SO ₂ Bu-t
	Me	Me	H	H	1	H	H	CF ₃	Ph	SCHF ₂
55	Me	Me	H	H	1	H	H	CF ₃	Ph	SO ₂ CHF ₂

	Me	Me	H	H	1	H	H	CF ₃	Ph	NH ₂
5	Me	Me	H	H	1	H	H	CF ₃	Ph	NHMe
	Me	Me	H	H	1	H	H	CF ₃	Ph	N(Me) ₂
	Me	Me	H	H	1	H	H	CF ₃	Ph	NHC(=O)Me
	Me	Me	H	H	1	H	H	CF ₃	Ph	N(Me)C(=O)Me
10	Me	Me	H	H	1	H	H	CF ₃	Ph	NHSO ₂ Me
	Me	Me	H	H	1	H	H	CF ₃	Ph	N(Me)SO ₂ Me
	Me	Me	H	H	1	H	H	CF ₃	Ph	NHSO ₂ CF ₃
15	Me	Me	H	H	1	H	H	CF ₃	Ph	N(Me)SO ₂ CF ₃
	Me	Me	H	H	1	H	H	CF ₃	Ph	NHPh
	Me	Me	H	H	1	H	H	CF ₃	Ph	N(Me)Ph
	Me	Me	H	H	1	H	H	CF ₃	Ph	CN
20	Me	Me	H	H	1	H	H	CF ₃	Ph	C(=O)Me
	Me	Me	H	H	1	H	H	CF ₃	Ph	C(=O)OMe
	Me	Me	H	H	1	H	H	CF ₃	Ph	C(=O)NH ₂
	Me	Me	H	H	1	H	H	CF ₃	Ph	C(=O)NHMe
25	Me	Me	H	H	1	H	H	CF ₃	Ph	C(=O)N(Me) ₂
	Me	Me	H	H	1	H	H	CF ₃	Ph	Imidazol-1-yl
	Me	Me	H	H	1	H	H	CF ₃	Ph	Pyrazol-1-yl
	Me	Me	H	H	1	H	H	CF ₃	Ph	1,2,4-Triazol-1-yl
30	Me	Me	H	H	1	H	H	CF ₃	Ph	1,2,4-Triazol-4-yl
	Me	Me	H	H	1	H	H	CF ₃	Ph	Tetrazol-1-yl
	Me	Me	H	H	1	H	H	CF ₃	Ph	Tetrazol-5-yl
35	Me	Me	H	H	1	H	H	CF ₃	Ph	(4,6-Dimethoxypyrimidin-2-yl)oxy
	Me	Me	H	H	1	H	H	CF ₃	Ph	(4,6-Dimethoxypyrimidin-2-yl)sulfonyl
	Me	Me	H	H	1	H	H	CF ₂ CF ₃	Ph	Cl
40	Me	Me	H	H	1	H	H	CF ₃	(2-Cl)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(2-F)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(2-OMe)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(2-Me)Ph	Cl
45	Me	Me	H	H	1	H	H	CF ₃	(2-NO ₂)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(2-CN)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(2-C(=O)Me)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(2-C(=O)OMe)Ph	Cl
50	Me	Me	H	H	1	H	H	CF ₃	(2-C(=O)OEt)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(2-C(=O)OPr-i)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(2-C(=O)NH ₂)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(2-C(=O)NHMe)Ph	Cl
55	Me	Me	H	H	1	H	H	CF ₃	(2-C(=O)NMe ₂)Ph	Cl

5	Me	Me	H	H	1	H	H	CF ₃	(3-Cl)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(3-F)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(3-OMe)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(3-Me)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(3-NO ₂)Ph	Cl
10	Me	Me	H	H	1	H	H	CF ₃	(3-CN)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(3-C(=O)Me)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(3-C(=O)OMe)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(3-C(=O)OEt)Ph	Cl
15	Me	Me	H	H	1	H	H	CF ₃	(3-C(=O)OPr-i)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(3-C(=O)NH ₂)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(3-C(=O)NHMe)Ph	Cl
20	Me	Me	H	H	1	H	H	CF ₃	(3-C(=O)NMe ₂)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(4-Cl)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(4-F)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(4-OMe)Ph	Cl
25	Me	Me	H	H	1	H	H	CF ₃	(4-Me)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(4-NO ₂)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(4-CN)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(4-C(=O)Me)Ph	Cl
30	Me	Me	H	H	1	H	H	CF ₃	(4-C(=O)OMe)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(4-C(=O)OEt)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(4-C(=O)OPr-i)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(4-C(=O)NH ₂)Ph	Cl
35	Me	Me	H	H	1	H	H	CF ₃	(4-C(=O)NHMe)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	(4-C(=O)NMe ₂)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	Pyrimidin-2-yl	Cl
40	Me	Me	H	H	1	H	H	CF ₃	4,6-Dimethoxypyrimidin-2-yl	Cl
	Me	Me	H	H	1	H	H	CF ₃	Thiophen-2-yl	Cl
	Me	Me	H	H	1	H	H	CF ₃	Furan-2-yl	Cl
	Me	Me	H	H	1	H	H	CF ₃	SO ₂ Me	Cl
45	Me	Me	H	H	1	H	H	CF ₃	SO ₂ Et	Cl
	Me	Me	H	H	1	H	H	CF ₃	SO ₂ Pr-i	Cl
	Me	Me	H	H	1	H	H	CF ₃	SO ₂ CH ₂ Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	SO ₂ CHF ₂	Cl
50	Me	Me	H	H	1	H	H	CF ₃	SO ₂ CF ₃	Cl
	Me	Me	H	H	1	H	H	CF ₃	SO ₂ Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	C(=O)Me	Cl
55	Me	Me	H	H	1	H	H	CF ₃	C(=O)Et	Cl

5	Me	Me	H	H	1	H	H	CF ₃	C(=O)Pr-i	Cl
	Me	Me	H	H	1	H	H	CF ₃	C(=O)Bu-t	Cl
	Me	Me	H	H	1	H	H	CF ₃	C(=O)Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	C(=O)CH ₂ Ph	Cl
10	Me	Me	H	H	1	H	H	CF ₃	C(=O)CH ₂ Cl	Cl
	Me	Me	H	H	1	H	H	CF ₃	C(=O)CHCl ₂	Cl
	Me	Me	H	H	1	H	H	CF ₃	C(=O)CF ₃	Cl
	Me	Me	H	H	1	H	H	CF ₃	C(=O)OMe	Cl
15	Me	Me	H	H	1	H	H	CF ₃	C(=O)OPh	Cl
	Me	Me	H	H	1	H	H	CF ₃	C(=O)OCH ₂ Ph	Cl
	Me	Me	H	H	1	H	H	CF ₃	C(=O)NHMe	Cl
	Me	Me	H	H	1	H	H	CF ₃	C(=O)N(Me) ₂	Cl
20	Me	Me	H	H	1	H	H	CF ₃	C(=O)NHPh	Cl
	Me	Me	H	H	1	H	H	CF ₃	NH ₂	Cl
	Me	Me	H	H	1	H	H	Cl	-(CH ₂) ₂ O- -(CH ₂) ₃ O- -(CH ₂) ₃ S- -(CH ₂) ₃ SO ₂ - -(CH ₂) ₂ O- -(CH ₂) ₃ O- -(CH ₂) ₃ S- -(CH ₂) ₃ SO ₂ - -(CH ₂) ₄ - -(CH ₂) ₄ -	
25	Me	Me	H	H	1	H	H	Cl		
	Me	Me	H	H	1	H	H	Cl		
	Me	Me	H	H	1	H	H	Cl		
	Me	Me	H	H	1	H	H	CF ₃		
30	Me	Me	H	H	1	H	H	CF ₃		
	Me	Me	H	H	1	H	H	CF ₃		
	Me	Me	H	H	1	H	H	CF ₃		
	Me	Me	H	H	1	H	H	OMe		
35	Me	Me	H	H	1	H	H	OCHF ₂		
	H	H	H	H	1	H	H	CF ₃	Me	Cl
	Me	H	H	H	1	H	H	CF ₃	Me	Cl
	Me	H	Me	H	1	H	H	CF ₃	Me	Cl
40	Me	Me	Me	H	1	H	H	CF ₃	Me	Cl
	Me	Me	H	H	1	Me	H	CF ₃	Me	Cl
	Me	Me	H	H	1	Et	H	CF ₃	Me	Cl
	Me	Me	H	H	1	Pr-i	H	CF ₃	Me	Cl
45	Me	Me	H	H	1	Me	Me	CF ₃	Me	Cl
	Me	Et	H	H	1	H	H	CF ₃	Me	Cl
	Et	Et	H	H	1	H	H	CF ₃	Me	Cl
50	Me	Pr-i	H	H	1	H	H	CF ₃	Me	Cl
	Me	Pr	H	H	1	H	H	CF ₃	Me	Cl
	Me	Pr-c	H	H	1	H	H	CF ₃	Me	Cl
	Me	CH ₂ Pr-c	H	H	1	H	H	CF ₃	Me	Cl
55	-(CH ₂) ₂ -		H	H	1	H	H	CF ₃	Me	Cl

5	-(CH ₂) ₃ -		H	H	1	H	H	CF ₃	Me	Cl
	-(CH ₂) ₄ -		H	H	1	H	H	CF ₃	Me	Cl
	-(CH ₂) ₅ -		H	H	1	H	H	CF ₃	Me	Cl
10	H	-(CH ₂) ₃ -		H	1	H	H	CF ₃	Me	Cl
	H	-(CH ₂) ₄ -		H	1	H	H	CF ₃	Me	Cl
	H	-(CH ₂) ₅ -		H	1	H	H	CF ₃	Me	Cl
	H	-(CH ₂) ₆ -		H	1	H	H	CF ₃	Me	Cl
15	Me	Me	H	H	0	H	H	Cl	H	Cl
	Me	Me	H	H	0	H	H	OCHF ₂	H	Cl
	Me	Me	H	H	0	H	H	OCHF ₂	H	OCHF ₂
	Me	Me	H	H	0	H	H	CHF ₂	H	Cl
	Me	Me	H	H	0	H	H	CF ₃	H	F
20	Me	Me	H	H	0	H	H	CF ₃	H	Cl
	Me	Me	H	H	0	H	H	CF ₃	H	OMe
	Me	Me	H	H	0	H	H	CF ₃	H	OEt
	Me	Me	H	H	0	H	H	CF ₃	H	OCHF ₂
	Me	Me	H	H	0	H	H	CF ₃	H	CN
25	Me	Me	H	H	0	H	H	CF ₃	H	Me
	Me	Me	H	H	0	H	H	H	Me	Cl
	Me	Me	H	H	0	H	H	Me	Me	Me
	Me	Me	H	H	0	H	H	Me	Me	Cl
	Me	Me	H	H	0	H	H	Cl	Me	Me
30	Me	Me	H	H	0	H	H	Et	Me	Cl
	Me	Me	H	H	0	H	H	Cl	Me	Et
	Me	Me	H	H	0	H	H	Pr-i	Me	Cl
	Me	Me	H	H	0	H	H	Cl	Me	Pr-i
	Me	Me	H	H	0	H	H	Bu-t	Me	Cl
35	Me	Me	H	H	0	H	H	Cl	Me	Bu-t
	Me	Me	H	H	0	H	H	Cl	Me	Cl
	Me	Me	H	H	0	H	H	CHF ₂	Me	Cl
	Me	Me	H	H	0	H	H	Cl	Me	CHF ₂
	Me	Me	H	H	0	H	H	OCHF ₂	Me	H
40	Me	Me	H	H	0	H	H	OCHF ₂	Me	Cl
	Me	Me	H	H	0	H	H	Cl	Me	OCHF ₂
	Me	Me	H	H	0	H	H	OCHF ₂	Me	OCHF ₂
	Me	Me	H	H	0	H	H	CF ₃	Me	H
	Me	Me	H	H	0	H	H	CF ₃	Me	Cl
45	Me	Me	H	H	0	H	H	Cl	Me	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Me	F
	Me	Me	H	H	0	H	H	CF ₃	Me	
	Me	Me	H	H	0	H	H	CF ₃	Me	
	Me	Me	H	H	0	H	H	CF ₃	Me	
50	Me	Me	H	H	0	H	H	CF ₃	Me	
	Me	Me	H	H	0	H	H	CF ₃	Me	
	Me	Me	H	H	0	H	H	CF ₃	Me	
	Me	Me	H	H	0	H	H	CF ₃	Me	
	Me	Me	H	H	0	H	H	CF ₃	Me	
55	Me	Me	H	H	0	H	H	CF ₃	Me	
	Me	Me	H	H	0	H	H	CF ₃	Me	
	Me	Me	H	H	0	H	H	CF ₃	Me	
	Me	Me	H	H	0	H	H	CF ₃	Me	
	Me	Me	H	H	0	H	H	CF ₃	Me	

	Me	Me	H	H	0	H	H	F	Me	CF ₃
5	Me	Me	H	H	0	H	H	CF ₃	Me	OH
	Me	Me	H	H	0	H	H	OH	Me	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Me	OMe
	Me	Me	H	H	0	H	H	OMe	Me	CF ₃
10	Me	Me	H	H	0	H	H	CF ₃	Me	OE _t
	Me	Me	H	H	0	H	H	OE _t	Me	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Me	OP _r -i
15	Me	Me	H	H	0	H	H	CF ₃	Me	OP _r
	Me	Me	H	H	0	H	H	CF ₃	Me	OBu-t
	Me	Me	H	H	0	H	H	CF ₃	Me	OBu-s
	Me	Me	H	H	0	H	H	CF ₃	Me	OBu-i
20	Me	Me	H	H	0	H	H	CF ₃	Me	OBu
	Me	Me	H	H	0	H	H	CF ₃	Me	O(2-Pen)
	Me	Me	H	H	0	H	H	CF ₃	Me	O(3-Pen)
	Me	Me	H	H	0	H	H	CF ₃	Me	OPen-n
25	Me	Me	H	H	0	H	H	CF ₃	Me	O(2-Hex)
	Me	Me	H	H	0	H	H	CF ₃	Me	O(3-Hex)
	Me	Me	H	H	0	H	H	CF ₃	Me	OHex-n
	Me	Me	H	H	0	H	H	CF ₃	Me	OPen-c
30	Me	Me	H	H	0	H	H	CF ₃	Me	OHex-c
	Me	Me	H	H	0	H	H	CF ₃	Me	OCH ₂ Pr-c
	Me	Me	H	H	0	H	H	CF ₃	Me	OCH ₂ Bu-c
35	Me	Me	H	H	0	H	H	CF ₃	Me	OCH ₂ Pen-c
	Me	Me	H	H	0	H	H	CF ₃	Me	OCH ₂ Hex-c
	Me	Me	H	H	0	H	H	CF ₃	Me	OCH ₂ CH=CH ₂
	Me	Me	H	H	0	H	H	CF ₃	Me	OCH ₂ C*CH
40	Me	Me	H	H	0	H	H	CF ₃	Me	OCHF ₂
	Me	Me	H	H	0	H	H	OCHF ₂	Me	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Me	OCH ₂ CHF ₂
	Me	Me	H	H	0	H	H	OCH ₂ CHF ₂	Me	CF ₃
45	Me	Me	H	H	0	H	H	CF ₃	Me	OCH ₂ CF ₃
	Me	Me	H	H	0	H	H	OCH ₂ CF ₃	Me	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Me	OCH ₂ CN
50	Me	Me	H	H	0	H	H	CF ₃	Me	OCH ₂ C(=O)OE _t
	Me	Me	H	H	0	H	H	CF ₃	Me	OCH(Me)C(=O)OE _t
	Me	Me	H	H	0	H	H	CF ₃	Me	OCH ₂ C(=O)NH ₂
	Me	Me	H	H	0	H	H	CF ₃	Me	OCH ₂ C(=O)NHMe
55	Me	Me	H	H	0	H	H	CF ₃	Me	OCH ₂ C(=O)N(Me) ₂

	Me	Me	H	H	0	H	H	CF ₃	Me	OCH ₂ Ph
5	Me	Me	H	H	0	H	H	CF ₃	Me	OPh
	Me	Me	H	H	0	H	H	CF ₃	Me	O(2-Cl)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(2-Br)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(2-F)Ph
10	Me	Me	H	H	0	H	H	CF ₃	Me	O(2-Me)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(2-OMe)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(2-NO ₂)Ph
15	Me	Me	H	H	0	H	H	CF ₃	Me	O(2-CN)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(2-C(=O)OMe)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(3-Cl)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(3-Br)Ph
20	Me	Me	H	H	0	H	H	CF ₃	Me	O(3-F)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(3-Me)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(3-OMe)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(3-NO ₂)Ph
25	Me	Me	H	H	0	H	H	CF ₃	Me	O(3-CN)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(3-C(=O)OMe)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(4-Cl)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(4-Br)Ph
30	Me	Me	H	H	0	H	H	CF ₃	Me	O(4-F)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(4-Me)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(4-OMe)Ph
35	Me	Me	H	H	0	H	H	CF ₃	Me	O(4-NO ₂)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(4-CN)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	O(4-C(=O)OMe)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	OC(=O)Me
40	Me	Me	H	H	0	H	H	CF ₃	Me	OC(=O)Et
	Me	Me	H	H	0	H	H	CF ₃	Me	OC(=O)CH ₂ Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	OC(=O)CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Me	OC(=O)Ph
45	Me	Me	H	H	0	H	H	CF ₃	Me	OSO ₂ Me
	Me	Me	H	H	0	H	H	CF ₃	Me	OSO ₂ Et
	Me	Me	H	H	0	H	H	CF ₃	Me	OSO ₂ CH ₂ Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	OSO ₂ CF ₃
50	Me	Me	H	H	0	H	H	CF ₃	Me	OSO ₂ Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	SMe
	Me	Me	H	H	0	H	H	CF ₃	Me	SO ₂ Me
55	Me	Me	H	H	0	H	H	CF ₃	Me	SEt

5	Me	Me	H	H	0	H	H	CF ₃	Me	SO ₂ Et
	Me	Me	H	H	0	H	H	CF ₃	Me	SPr
	Me	Me	H	H	0	H	H	CF ₃	Me	SO ₂ Pr
	Me	Me	H	H	0	H	H	CF ₃	Me	SPr-i
	Me	Me	H	H	0	H	H	CF ₃	Me	SO ₂ Pr-i
10	Me	Me	H	H	0	H	H	CF ₃	Me	SBu-t
	Me	Me	H	H	0	H	H	CF ₃	Me	SO ₂ Bu-t
	Me	Me	H	H	0	H	H	CF ₃	Me	SCHF ₂
	Me	Me	H	H	0	H	H	CF ₃	Me	SO ₂ CHF ₂
15	Me	Me	H	H	0	H	H	CF ₃	Me	SCF ₃
	Me	Me	H	H	0	H	H	CF ₃	Me	SO ₂ CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Me	SPh
20	Me	Me	H	H	0	H	H	CF ₃	Me	SO ₂ Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	SCH ₂ Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	SO ₂ CH ₂ Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	SCH ₂ C(=O)OEt
25	Me	Me	H	H	0	H	H	CF ₃	Me	SO ₂ CH ₂ C(=O)OEt
	Me	Me	H	H	0	H	H	CF ₃	Me	SCH(Me)C(=O)OEt
	Me	Me	H	H	0	H	H	CF ₃	Me	SO ₂ CH(Me)C(=O)OEt
	Me	Me	H	H	0	H	H	CF ₃	Me	SCH ₂ C(=O)NH ₂
30	Me	Me	H	H	0	H	H	CF ₃	Me	SO ₂ CH ₂ C(=O)NH ₂
	Me	Me	H	H	0	H	H	CF ₃	Me	SCH ₂ C(=O)NHMe
	Me	Me	H	H	0	H	H	CF ₃	Me	SO ₂ CH ₂ C(=O)NHMe
	Me	Me	H	H	0	H	H	CF ₃	Me	SCH ₂ C(=O)N(Me) ₂
35	Me	Me	H	H	0	H	H	CF ₃	Me	SO ₂ CH ₂ C(=O)N(Me) ₂
	Me	Me	H	H	0	H	H	CF ₃	Me	NH ₂
	Me	Me	H	H	0	H	H	CF ₃	Me	NHMe
40	Me	Me	H	H	0	H	H	CF ₃	Me	N(Me) ₂
	Me	Me	H	H	0	H	H	CF ₃	Me	NHC(=O)Me
	Me	Me	H	H	0	H	H	CF ₃	Me	N(Me)C(=O)Me
	Me	Me	H	H	0	H	H	CF ₃	Me	NHSO ₂ Me
45	Me	Me	H	H	0	H	H	CF ₃	Me	N(Me)SO ₂ Me
	Me	Me	H	H	0	H	H	CF ₃	Me	NHSO ₂ CHF ₂
	Me	Me	H	H	0	H	H	CF ₃	Me	N(Me)SO ₂ CHF ₂
	Me	Me	H	H	0	H	H	CF ₃	Me	NHSO ₂ CF ₃
50	Me	Me	H	H	0	H	H	CF ₃	Me	N(Me)SO ₂ CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Me	NHPh
	Me	Me	H	H	0	H	H	CF ₃	Me	N(Me)Ph
55	Me	Me	H	H	0	H	H	CF ₃	Me	CN

	Me	Me	H	H	0	H	H	CN	Me	CF ₃
5	Me	Me	H	H	0	H	H	CF ₃	Me	C(=O)OMe
	Me	Me	H	H	0	H	H	CF ₃	Me	C(=O)OCH ₂ Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	C(=O)OPh
	Me	Me	H	H	0	H	H	CF ₃	Me	C(=O)NH ₂
10	Me	Me	H	H	0	H	H	CF ₃	Me	C(=O)NHMe
	Me	Me	H	H	0	H	H	CF ₃	Me	C(=O)N(Me) ₂
	Me	Me	H	H	0	H	H	CF ₃	Me	C(=O)Me
15	Me	Me	H	H	0	H	H	CF ₃	Me	C(=O)CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Me	C(=O)CH ₂ Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	C(=O)Ph
	Me	Me	H	H	0	H	H	CF ₃	Me	Me
20	Me	Me	H	H	0	H	H	Me	Me	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Me	Et
	Me	Me	H	H	0	H	H	CF ₃	Me	Pr-i
	Me	Me	H	H	0	H	H	CF ₃	Me	Pr
25	Me	Me	H	H	0	H	H	CF ₃	Me	CH ₂ OMe
	Me	Me	H	H	0	H	H	CF ₃	Me	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Me	CHF ₂
	Me	Me	H	H	0	H	H	CF ₃	Me	Ph
30	Me	Me	H	H	0	H	H	CF ₂ CF ₃	Me	Cl
	Me	Me	H	H	0	H	H	Ph	Me	Me
	Me	Me	H	H	0	H	H	Ph	Me	Cl
	Me	Me	H	H	0	H	H	Ph	Me	OEt
35	Me	Me	H	H	0	H	H	Ph	Me	CF ₃
	Me	Me	H	H	0	H	H	Ph	Me	Ph
	Me	Me	H	H	0	H	H	Cl	Et	Cl
40	Me	Me	H	H	0	H	H	OCHF ₂	Et	Cl
	Me	Me	H	H	0	H	H	Cl	Et	OCHF ₂
	Me	Me	H	H	0	H	H	OCHF ₂	Et	OCHF ₂
	Me	Me	H	H	0	H	H	CF ₃	Et	F
45	Me	Me	H	H	0	H	H	F	Et	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Et	Cl
	Me	Me	H	H	0	H	H	Cl	Et	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Et	OMe
50	Me	Me	H	H	0	H	H	OMe	Et	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Et	OEt
	Me	Me	H	H	0	H	H	OEt	Et	CF ₃
55	Me	Me	H	H	0	H	H	CF ₃	Et	OCHF ₂

	Me	Me	H	H	0	H	H	OCHF ₂	Et	CF ₃
5	Me	Me	H	H	0	H	H	CF ₃	Et	CN
	Me	Me	H	H	0	H	H	CN	Et	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Et	Me
10	Me	Me	H	H	0	H	H	Me	Et	CF ₃
	Me	Me	H	H	0	H	H	Cl	Pr-i	Cl
	Me	Me	H	H	0	H	H	OCHF ₂	Pr-i	Cl
	Me	Me	H	H	0	H	H	Cl	Pr-i	OCHF ₂
15	Me	Me	H	H	0	H	H	OCHF ₂	Pr-i	OCHF ₂
	Me	Me	H	H	0	H	H	CF ₃	Pr-i	F
	Me	Me	H	H	0	H	H	F	Pr-i	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Pr-i	Cl
20	Me	Me	H	H	0	H	H	Cl	Pr-i	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Pr-i	OMe
	Me	Me	H	H	0	H	H	OMe	Pr-i	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Pr-i	OEt
25	Me	Me	H	H	0	H	H	OEt	Pr-i	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Pr-i	OCHF ₂
	Me	Me	H	H	0	H	H	OCHF ₂	Pr-i	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Pr-i	CN
30	Me	Me	H	H	0	H	H	CN	Pr-i	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Pr-i	Me
	Me	Me	H	H	0	H	H	Me	Pr-i	CF ₃
35	Me	Me	H	H	0	H	H	Cl	Pr	Cl
	Me	Me	H	H	0	H	H	OCHF ₂	Pr	Cl
	Me	Me	H	H	0	H	H	Cl	Pr	OCHF ₂
	Me	Me	H	H	0	H	H	OCHF ₂	Pr	OCHF ₂
40	Me	Me	H	H	0	H	H	CF ₃	Pr	F
	Me	Me	H	H	0	H	H	F	Pr	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Pr	Cl
	Me	Me	H	H	0	H	H	Cl	Pr	CF ₃
45	Me	Me	H	H	0	H	H	CF ₃	Pr	OMe
	Me	Me	H	H	0	H	H	OMe	Pr	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Pr	OEt
	Me	Me	H	H	0	H	H	OEt	Pr	CF ₃
50	Me	Me	H	H	0	H	H	CF ₃	Pr	OCHF ₂
	Me	Me	H	H	0	H	H	OCHF ₂	Pr	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Pr	CN
55	Me	Me	H	H	0	H	H	CN	Pr	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Pr	Me

	Me	Me	H	H	0	H	H	Me	Pr	CF ₃
5	Me	Me	H	H	0	H	H	Cl	Bu-t	Cl
	Me	Me	H	H	0	H	H	OCHF ₂	Bu-t	Cl
	Me	Me	H	H	0	H	H	OCHF ₂	Bu-t	OCHF ₂
	Me	Me	H	H	0	H	H	CF ₃	Bu-t	H
10	Me	Me	H	H	0	H	H	CF ₃	Bu-t	F
	Me	Me	H	H	0	H	H	CF ₃	Bu-t	Cl
	Me	Me	H	H	0	H	H	Cl	Bu-t	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Bu-t	OMe
15	Me	Me	H	H	0	H	H	OMe	Bu-t	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Bu-t	OEt
	Me	Me	H	H	0	H	H	OEt	Bu-t	CF ₃
20	Me	Me	H	H	0	H	H	CF ₃	Bu-t	OCHF ₂
	Me	Me	H	H	0	H	H	CF ₃	Bu-t	CN
	Me	Me	H	H	0	H	H	CF ₃	Bu-t	Me
	Me	Me	H	H	0	H	H	Me	Bu-t	CF ₃
25	Me	Me	H	H	0	H	H	CF ₃	Bu-s	Cl
	Me	Me	H	H	0	H	H	Cl	Bu-s	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Bu-i	Cl
	Me	Me	H	H	0	H	H	Cl	Bu-i	CF ₃
30	Me	Me	H	H	0	H	H	CF ₃	Bu	Cl
	Me	Me	H	H	0	H	H	Cl	Bu	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	1-Methylbutyl	Cl
35	Me	Me	H	H	0	H	H	Cl	1-Methylbutyl	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	1-Ethylpropyl	Cl
	Me	Me	H	H	0	H	H	Cl	1-Ethylpropyl	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	1-Pentyl	Cl
40	Me	Me	H	H	0	H	H	Cl	1-Pentyl	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	1-Methylpentyl	Cl
	Me	Me	H	H	0	H	H	Cl	1-Methylpentyl	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	2-Ethylbutyl	Cl
45	Me	Me	H	H	0	H	H	Cl	2-Ethylbutyl	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	3,3-Dimethylbutyl	Cl
	Me	Me	H	H	0	H	H	Cl	3,3-Dimethylbutyl	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	1-Hexyl	Cl
50	Me	Me	H	H	0	H	H	Cl	1-Hexyl	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	1-Heptyl	Cl
	Me	Me	H	H	0	H	H	Cl	1-Heptyl	CF ₃
55	Me	Me	H	H	0	H	H	CF ₃	1-Octyl	Cl

5	Me	Me	H	H	0	H	H	Cl	1-Octyl	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Ph	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ Ph	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Pr-c	Cl
10	Me	Me	H	H	0	H	H	CF ₃	Pen-c	Cl
	Me	Me	H	H	0	H	H	Cl	Pen-c	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Hex-c	Cl
	Me	Me	H	H	0	H	H	Cl	Hex-c	CF ₃
15	Me	Me	H	H	0	H	H	Cl	CH ₂ Pr-c	Cl
	Me	Me	H	H	0	H	H	OCHF ₂	CH ₂ Pr-c	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ Pr-c	OCHF ₂
	Me	Me	H	H	0	H	H	OCHF ₂	CH ₂ Pr-c	OCHF ₂
20	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pr-c	F
	Me	Me	H	H	0	H	H	F	CH ₂ Pr-c	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pr-c	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ Pr-c	CF ₃
25	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pr-c	CN
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pr-c	OH
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pr-c	OMe
	Me	Me	H	H	0	H	H	OMe	CH ₂ Pr-c	CF ₃
30	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pr-c	OEt
	Me	Me	H	H	0	H	H	OEt	CH ₂ Pr-c	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pr-c	OPr-i
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pr-c	OPr
35	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pr-c	OBu-t
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pr-c	OCH ₂ Pr-c
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pr-c	OCH ₂ Bu-c
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pr-c	OPen-c
40	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pr-c	OCHF ₂
	Me	Me	H	H	0	H	H	OCHF ₂	CH ₂ Pr-c	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pr-c	CN
	Me	Me	H	H	0	H	H	CN	CH ₂ Pr-c	CF ₃
45	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pr-c	Me
	Me	Me	H	H	0	H	H	Me	CH ₂ Pr-c	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	1-cyclopropylethyl	Cl
	Me	Me	H	H	0	H	H	Cl	1-cyclopropylethyl	CF ₃
50	Me	Me	H	H	0	H	H	CF ₃	CH ₂ (2-Methyl-cyclopropyl)	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ (2-Methyl-cyclopropyl)	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ (2,2-Dimethyl-cyclopropyl)	Cl
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ (2,2-Dimethyl-cyclopropyl)	Cl

5	Me	Me	H	H	0	H	H	Cl	CH ₂ (2,2-Dimethyl-cyclopropyl)	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ (2-Chloro-cyclopropyl)	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ (2-Chloro-cyclopropyl)	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ (2,2-Dichloro-cyclopropyl)	Cl
10	Me	Me	H	H	0	H	H	Cl	CH ₂ (2,2-Dichloro-cyclopropyl)	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ (2-Fluoro-cyclopropyl)	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ (2-Fluoro-cyclopropyl)	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ (2,2-Difluoro-cyclopropyl)	Cl
15	Me	Me	H	H	0	H	H	Cl	CH ₂ (2,2-Difluoro-cyclopropyl)	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Bu-c	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ Bu-c	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Pen-c	Cl
20	Me	Me	H	H	0	H	H	Cl	CH ₂ Pen-c	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ Hex-c	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ Hex-c	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ CH ₂ Pr-c	Cl
25	Me	Me	H	H	0	H	H	Cl	CH ₂ CH ₂ Pr-c	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ CH=CH ₂	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ CH=CH ₂	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ CH=CHCl	Cl
30	Me	Me	H	H	0	H	H	Cl	CH ₂ CH=CHCl	CF ₃
	Me	Me	H	H	0	H	H	Cl	CH ₂ C≡CH	Cl
	Me	Me	H	H	0	H	H	OCHF ₂	CH ₂ C≡CH	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ C≡CH	OCHF ₂
35	Me	Me	H	H	0	H	H	OCHF ₂	CH ₂ C≡CH	OCHF ₂
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C≡CH	F
	Me	Me	H	H	0	H	H	F	CH ₂ C≡CH	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C≡CH	Cl
40	Me	Me	H	H	0	H	H	Cl	CH ₂ C≡CH	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C≡CH	OMe
	Me	Me	H	H	0	H	H	OMe	CH ₂ C≡CH	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C≡CH	OEt
45	Me	Me	H	H	0	H	H	OEt	CH ₂ C≡CH	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C≡CH	OCHF ₂
	Me	Me	H	H	0	H	H	OCHF ₂	CH ₂ C≡CH	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C≡CH	CN
50	Me	Me	H	H	0	H	H	CN	CH ₂ C≡CH	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C≡CH	Me
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C≡CH	
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C≡CH	
55	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C≡CH	

	Me	Me	H	H	0	H	H	Me	$\text{CH}_2\text{C}\equiv\text{CH}$	CF_3
5	Me	Me	H	H	0	H	H	CF_3	$\text{CH}_2\text{C}\equiv\text{CH}$	Cl
	Me	Me	H	H	0	H	H	Cl	$\text{CH}_2\text{C}\equiv\text{CH}$	CF_3
	Me	Me	H	H	0	H	H	CF_3	$\text{CH}_2\text{C}\equiv\text{CMe}$	Cl
	Me	Me	H	H	0	H	H	Cl	$\text{CH}_2\text{C}\equiv\text{CMe}$	CF_3
10	Me	Me	H	H	0	H	H	Cl	CHF_2	Cl
	Me	Me	H	H	0	H	H	OCHF_2	CHF_2	Cl
	Me	Me	H	H	0	H	H	Cl	CHF_2	OCHF_2
	Me	Me	H	H	0	H	H	OCHF_2	CHF_2	OCHF_2
15	Me	Me	H	H	0	H	H	CF_3	CHF_2	Cl
	Me	Me	H	H	0	H	H	Cl	CHF_2	CF_3
	Me	Me	H	H	0	H	H	CF_3	CHF_2	F
20	Me	Me	H	H	0	H	H	F	CHF_2	CF_3
	Me	Me	H	H	0	H	H	CF_3	CHF_2	OMe
	Me	Me	H	H	0	H	H	OMe	CHF_2	CF_3
	Me	Me	H	H	0	H	H	CF_3	CHF_2	OEt
25	Me	Me	H	H	0	H	H	OEt	CHF_2	CF_3
	Me	Me	H	H	0	H	H	CF_3	CHF_2	OCHF_2
	Me	Me	H	H	0	H	H	OCHF_2	CHF_2	CF_3
	Me	Me	H	H	0	H	H	CF_3	CHF_2	CN
30	Me	Me	H	H	0	H	H	CN	CHF_2	CF_3
	Me	Me	H	H	0	H	H	CF_3	CHF_2	Me
	Me	Me	H	H	0	H	H	Me	CHF_2	CF_3
35	Me	Me	H	H	0	H	H	Me	CHF_2	Cl
	Me	Me	H	H	0	H	H	Cl	CHF_2	Me
	Me	Me	H	H	0	H	H	Et	CHF_2	Cl
	Me	Me	H	H	0	H	H	Cl	CHF_2	Et
40	Me	Me	H	H	0	H	H	CF_3	CH_2CHF_2	Cl
	Me	Me	H	H	0	H	H	Cl	CH_2CHF_2	CF_3
	Me	Me	H	H	0	H	H	CF_3	CH_2CF_3	Cl
	Me	Me	H	H	0	H	H	Cl	CH_2CF_3	CF_3
45	Me	Me	H	H	0	H	H	CF_3	CH_2OH	Cl
	Me	Me	H	H	0	H	H	Cl	CH_2OH	CF_3
	Me	Me	H	H	0	H	H	Cl	CH_2OMe	Cl
	Me	Me	H	H	0	H	H	OCHF_2	CH_2OMe	Cl
50	Me	Me	H	H	0	H	H	Cl	CH_2OMe	OCHF_2
	Me	Me	H	H	0	H	H	OCHF_2	CH_2OMe	OCHF_2
	Me	Me	H	H	0	H	H	CF_3	CH_2OMe	F
	Me	Me	H	H	0	H	H	F	CH_2OMe	CF_3
55	Me	Me	H	H	0	H	H	CF_3	CH_2OMe	Cl

5	Me	Me	H	H	0	H	H	Cl	CH ₂ OMe	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ OMe	OMe
	Me	Me	H	H	0	H	H	OMe	CH ₂ OMe	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ OMe	OEt
	Me	Me	H	H	0	H	H	OEt	CH ₂ OMe	CF ₃
10	Me	Me	H	H	0	H	H	CF ₃	CH ₂ OMe	OCHF ₂
	Me	Me	H	H	0	H	H	OCHF ₂	CH ₂ OMe	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ OMe	CN
15	Me	Me	H	H	0	H	H	CN	CH ₂ OMe	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ OMe	Me
	Me	Me	H	H	0	H	H	Me	CH ₂ OMe	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ OEt	Cl
20	Me	Me	H	H	0	H	H	Cl	CH ₂ OEt	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ CH ₂ OH	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ CH ₂ OH	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ CH ₂ OMe	Cl
25	Me	Me	H	H	0	H	H	Cl	CH ₂ CH ₂ OMe	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ CH ₂ OEt	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ CH ₂ OEt	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ NHMe	Cl
30	Me	Me	H	H	0	H	H	Cl	CH ₂ NHMe	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ N(Me) ₂	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ N(Me) ₂	CF ₃
35	Me	Me	H	H	0	H	H	CF ₃	CH ₂ N(Me)C(=O)Me	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ N(Me)C(=O)Me	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ N(Me)C(=O)CF ₃	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ N(Me)C(=O)CF ₃	CF ₃
40	Me	Me	H	H	0	H	H	CF ₃	CH ₂ N(Me)SO ₂ Me	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ N(Me)SO ₂ Me	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ N(Me)SO ₂ CHF ₂	Cl
45	Me	Me	H	H	0	H	H	Cl	CH ₂ N(Me)SO ₂ CHF ₂	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ N(Me)SO ₂ CF ₃	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ N(Me)SO ₂ CF ₃	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ SMe	Cl
50	Me	Me	H	H	0	H	H	Cl	CH ₂ SMe	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ SO ₂ Me	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ SO ₂ Me	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ CH ₂ SMe	Cl
55	Me	Me	H	H	0	H	H	Cl	CH ₂ CH ₂ SMe	CF ₃

	Me	Me	H	H	0	H	H	CF ₃	CH ₂ CH ₂ SO ₂ Me	Cl
5	Me	Me	H	H	0	H	H	Cl	CH ₂ CH ₂ SO ₂ Me	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ CN	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ CN	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C(=O)OMe	Cl
10	Me	Me	H	H	0	H	H	Cl	CH ₂ C(=O)OMe	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C(=O)OEt	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ C(=O)OEt	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH(Me)C(=O)OMe	Cl
15	Me	Me	H	H	0	H	H	Cl	CH(Me)C(=O)OMe	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	C(Me) ₂ C(=O)OMe	Cl
	Me	Me	H	H	0	H	H	Cl	C(Me) ₂ C(=O)OMe	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C(=O)NH ₂	Cl
20	Me	Me	H	H	0	H	H	Cl	CH ₂ C(=O)NH ₂	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C(=O)NHMe	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ C(=O)NHMe	CF ₃
25	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C(=O)N(Me) ₂	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ C(=O)N(Me) ₂	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C(=O)Me	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ C(=O)Me	CF ₃
30	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C(=NOMe)Me	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ C(=NOMe)Me	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C(=O)CF ₃	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ C(=O)CF ₃	CF ₃
35	Me	Me	H	H	0	H	H	CF ₃	CH ₂ CH ₂ C(=O)Me	Cl
	Me	Me	H	H	0	H	H	Cl	CH ₂ CH ₂ C(=O)Me	CF ₃
	Me	Me	H	H	0	H	H	Me	Ph	Me
40	Me	Me	H	H	0	H	H	Me	Ph	Cl
	Me	Me	H	H	0	H	H	Et	Ph	Cl
	Me	Me	H	H	0	H	H	Pr	Ph	Cl
	Me	Me	H	H	0	H	H	Pr-i	Ph	Cl
45	Me	Me	H	H	0	H	H	Bu-t	Ph	Cl
	Me	Me	H	H	0	H	H	CH ₂ OMe	Ph	Cl
	Me	Me	H	H	0	H	H	Cl	Ph	Cl
	Me	Me	H	H	0	H	H	OCHF ₂	Ph	Cl
50	Me	Me	H	H	0	H	H	OCHF ₂	Ph	OCHF ₂
	Me	Me	H	H	0	H	H	CHF ₂	Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	Ph	H
55	Me	Me	H	H	0	H	H	CF ₃	Ph	Me

	Me	Me	H	H	0	H	H	Me	Ph	CF ₃
5	Me	Me	H	H	0	H	H	CF ₃	Ph	Et
	Me	Me	H	H	0	H	H	CF ₃	Ph	Pr-i
	Me	Me	H	H	0	H	H	CF ₃	Ph	CHF ₂
	Me	Me	H	H	0	H	H	CF ₃	Ph	CF ₃
10	Me	Me	H	H	0	H	H	CF ₃	Ph	F
	Me	Me	H	H	0	H	H	CF ₃	Ph	Cl
	Me	Me	H	H	0	H	H	Cl	Ph	CF ₃
15	Me	Me	H	H	0	H	H	CF ₃	Ph	OH
	Me	Me	H	H	0	H	H	OH	Ph	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Ph	OMe
	Me	Me	H	H	0	H	H	OMe	Ph	CF ₃
20	Me	Me	H	H	0	H	H	CF ₃	Ph	OEt
	Me	Me	H	H	0	H	H	OEt	Ph	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Ph	OPr-i
25	Me	Me	H	H	0	H	H	CF ₃	Ph	OPr
	Me	Me	H	H	0	H	H	CF ₃	Ph	OBu-t
	Me	Me	H	H	0	H	H	CF ₃	Ph	OCH ₂ Pr-c
	Me	Me	H	H	0	H	H	CF ₃	Ph	OCH ₂ CH=CH ₂
30	Me	Me	H	H	0	H	H	CF ₃	Ph	OCH ₂ C≡CH
	Me	Me	H	H	0	H	H	CF ₃	Ph	OCHF ₂
	Me	Me	H	H	0	H	H	OCHF ₂	Ph	CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Ph	OCH ₂ CHF ₂
	Me	Me	H	H	0	H	H	CF ₃	Ph	OCH ₂ CF ₃
35	Me	Me	H	H	0	H	H	CF ₃	Ph	OCH ₂ C(=O)OMe
	Me	Me	H	H	0	H	H	CF ₃	Ph	OCH(Me)C(=O)OMe
	Me	Me	H	H	0	H	H	CF ₃	Ph	OC(Me) ₂ C(=O)OMe
40	Me	Me	H	H	0	H	H	CF ₃	Ph	OC(=O)Me
	Me	Me	H	H	0	H	H	CF ₃	Ph	OC(=O)Et
	Me	Me	H	H	0	H	H	CF ₃	Ph	OC(=O)CH ₂ Ph
	Me	Me	H	H	0	H	H	CF ₃	Ph	OC(=O)CF ₃
45	Me	Me	H	H	0	H	H	CF ₃	Ph	OC(=O)Ph
	Me	Me	H	H	0	H	H	CF ₃	Ph	OSO ₂ Me
	Me	Me	H	H	0	H	H	CF ₃	Ph	OSO ₂ Et
50	Me	Me	H	H	0	H	H	CF ₃	Ph	OSO ₂ CH ₂ Ph
	Me	Me	H	H	0	H	H	CF ₃	Ph	OSO ₂ CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Ph	OSO ₂ Ph
	Me	Me	H	H	0	H	H	CF ₃	Ph	SMe
55	Me	Me	H	H	0	H	H	CF ₃	Ph	SO ₂ Me

	Me	Me	H	H	0	H	H	CF ₃	Ph	SEt
5	Me	Me	H	H	0	H	H	CF ₃	Ph	SO ₂ Et
	Me	Me	H	H	0	H	H	CF ₃	Ph	SPr-i
	Me	Me	H	H	0	H	H	CF ₃	Ph	SO ₂ Pr-i
	Me	Me	H	H	0	H	H	CF ₃	Ph	SPr
10	Me	Me	H	H	0	H	H	CF ₃	Ph	SO ₂ Pr
	Me	Me	H	H	0	H	H	CF ₃	Ph	SBu-t
	Me	Me	H	H	0	H	H	CF ₃	Ph	SO ₂ Bu-t
15	Me	Me	H	H	0	H	H	CF ₃	Ph	SCHF ₂
	Me	Me	H	H	0	H	H	CF ₃	Ph	SO ₂ CHF ₂
	Me	Me	H	H	0	H	H	CF ₃	Ph	NH ₂
	Me	Me	H	H	0	H	H	CF ₃	Ph	NHMe
20	Me	Me	H	H	0	H	H	CF ₃	Ph	N(Me) ₂
	Me	Me	H	H	0	H	H	CF ₃	Ph	NHC(=O)Me
	Me	Me	H	H	0	H	H	CF ₃	Ph	N(Me)C(=O)Me
	Me	Me	H	H	0	H	H	CF ₃	Ph	NHSO ₂ Me
25	Me	Me	H	H	0	H	H	CF ₃	Ph	N(Me)SO ₂ Me
	Me	Me	H	H	0	H	H	CF ₃	Ph	NHSO ₂ CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Ph	N(Me)SO ₂ CF ₃
	Me	Me	H	H	0	H	H	CF ₃	Ph	NHPh
30	Me	Me	H	H	0	H	H	CF ₃	Ph	N(Me)Ph
	Me	Me	H	H	0	H	H	CF ₃	Ph	CN
	Me	Me	H	H	0	H	H	CF ₃	Ph	C(=O)Me
35	Me	Me	H	H	0	H	H	CF ₃	Ph	C(=O)OMe
	Me	Me	H	H	0	H	H	CF ₃	Ph	C(=O)NH ₂
	Me	Me	H	H	0	H	H	CF ₃	Ph	C(=O)NHMe
	Me	Me	H	H	0	H	H	CF ₃	Ph	C(=O)N(Me) ₂
40	Me	Me	H	H	0	H	H	CF ₃	Ph	Imidazol-1-yl
	Me	Me	H	H	0	H	H	CF ₃	Ph	Pyrazol-1-yl
	Me	Me	H	H	0	H	H	CF ₃	Ph	1,2,4-Triazol-1-yl
	Me	Me	H	H	0	H	H	CF ₃	Ph	1,2,4-Triazol-4-yl
45	Me	Me	H	H	0	H	H	CF ₃	Ph	Tetrazol-1-yl
	Me	Me	H	H	0	H	H	CF ₃	Ph	Tetrazol-5-yl
	Me	Me	H	H	0	H	H	CF ₃	Ph	(4,6-Dimethoxypyrimidin-2-yl)oxy
50	Me	Me	H	H	0	H	H	CF ₃	Ph	(4,6-Dimethoxypyrimidin-2-yl)sulfonyl
	Me	Me	H	H	0	H	H	CF ₂ CF ₃	Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(2-Cl)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(2-F)Ph	Cl
55	Me	Me	H	H	0	H	H	CF ₃	(2-OMe)Ph	Cl

5	Me	Me	H	H	0	H	H	CF ₃	(2-Me)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(2-NO ₂)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(2-CN)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(2-C(=O)Me)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(2-C(=O)OMe)Ph	Cl
10	Me	Me	H	H	0	H	H	CF ₃	(2-C(=O)OEt)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(2-C(=O)OPr-i)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(2-C(=O)NH ₂)Ph	Cl
15	Me	Me	H	H	0	H	H	CF ₃	(2-C(=O)NHMe)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(2-C(=O)NMe ₂)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(3-Cl)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(3-F)Ph	Cl
20	Me	Me	H	H	0	H	H	CF ₃	(3-OMe)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(3-Me)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(3-NO ₂)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(3-CN)Ph	Cl
25	Me	Me	H	H	0	H	H	CF ₃	(3-C(=O)Me)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(3-C(=O)OMe)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(3-C(=O)OEt)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(3-C(=O)OPr-i)Ph	Cl
30	Me	Me	H	H	0	H	H	CF ₃	(3-C(=O)NH ₂)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(3-C(=O)NHMe)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(3-C(=O)NMe ₂)Ph	Cl
35	Me	Me	H	H	0	H	H	CF ₃	(4-Cl)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(4-F)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(4-OMe)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(4-Me)Ph	Cl
40	Me	Me	H	H	0	H	H	CF ₃	(4-NO ₂)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(4-CN)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(4-C(=O)Me)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(4-C(=O)OMe)Ph	Cl
45	Me	Me	H	H	0	H	H	CF ₃	(4-C(=O)OEt)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(4-C(=O)OPr-i)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(4-C(=O)NH ₂)Ph	Cl
50	Me	Me	H	H	0	H	H	CF ₃	(4-C(=O)NHMe)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	(4-C(=O)NMe ₂)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	Pyrimidin-2-yl	Cl
	Me	Me	H	H	0	H	H	CF ₃	4,6-Dimethoxypyrimidin-2-yl	Cl
55	Me	Me	H	H	0	H	H	CF ₃	Thiophen-2-yl	Cl

5	Me	Me	H	H	0	H	H	CF ₃	Furan-2-yl	Cl
	Me	Me	H	H	0	H	H	CF ₃	SO ₂ Me	Cl
	Me	Me	H	H	0	H	H	CF ₃	SO ₂ Et	Cl
10	Me	Me	H	H	0	H	H	CF ₃	SO ₂ Pr-i	Cl
	Me	Me	H	H	0	H	H	CF ₃	SO ₂ CH ₂ Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	SO ₂ CHF ₂	Cl
15	Me	Me	H	H	0	H	H	CF ₃	SO ₂ CF ₃	Cl
	Me	Me	H	H	0	H	H	CF ₃	SO ₂ Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	C(=O)Me	Cl
20	Me	Me	H	H	0	H	H	CF ₃	C(=O)Et	Cl
	Me	Me	H	H	0	H	H	CF ₃	C(=O)Pr-i	Cl
	Me	Me	H	H	0	H	H	CF ₃	C(=O)Bu-t	Cl
25	Me	Me	H	H	0	H	H	CF ₃	C(=O)Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	C(=O)CH ₂ Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	C(=O)CH ₂ Cl	Cl
30	Me	Me	H	H	0	H	H	CF ₃	C(=O)CHCl ₂	Cl
	Me	Me	H	H	0	H	H	CF ₃	C(=O)CF ₃	Cl
	Me	Me	H	H	0	H	H	CF ₃	C(=O)OMe	Cl
35	Me	Me	H	H	0	H	H	CF ₃	C(=O)OPh	Cl
	Me	Me	H	H	0	H	H	CF ₃	C(=O)OCH ₂ Ph	Cl
	Me	Me	H	H	0	H	H	CF ₃	C(=O)NHMe	Cl
40	Me	Me	H	H	0	H	H	CF ₃	C(=O)N(Me) ₂	Cl
	Me	Me	H	H	0	H	H	CF ₃	C(=O)NHPh	Cl
	Me	Me	H	H	0	H	H	CF ₃	NH ₂	Cl
45	Me	Me	H	H	0	H	H	Cl	-(CH ₂) ₂ O- -(CH ₂) ₃ O- -(CH ₂) ₃ S- -(CH ₂) ₃ SO ₂ - -(CH ₂) ₂ O- -(CH ₂) ₃ O- -(CH ₂) ₃ S- -(CH ₂) ₃ SO ₂ - -(CH ₂) ₄ - -(CH ₂) ₄ -	
	Me	Me	H	H	0	H	H	Cl		
	Me	Me	H	H	0	H	H	Cl		
50	Me	Me	H	H	0	H	H	Cl		
	Me	Me	H	H	0	H	H	CF ₃		
	Me	Me	H	H	0	H	H	CF ₃		
55	Me	Me	H	H	0	H	H	CF ₃		
	Me	Me	H	H	0	H	H	CF ₃		
	Me	Me	H	H	0	H	H	OMe		
	Me	Me	H	H	0	H	H	OCHF ₂		
	H	H	H	H	0	H	H	CF ₃		
	Me	H	H	H	0	H	H	CF ₃	Me	Cl
	Me	H	Me	H	0	H	H	CF ₃	Me	Cl
	Me	Me	Me	H	0	H	H	CF ₃	Me	Cl
	Me	Me	H	H	0	Me	H	CF ₃	Me	Cl

5	Me	Me	H	H	0	Et	H	CF ₃	Me	Cl
	Me	Me	H	H	0	Pr-i	H	CF ₃	Me	Cl
	Me	Me	H	H	0	Me	Me	CF ₃	Me	Cl
	Me	Et	H	H	0	H	H	CF ₃	Me	Cl
10	Et	Et	H	H	0	H	H	CF ₃	Me	Cl
	Me	Pr-i	H	H	0	H	H	CF ₃	Me	Cl
	Me	Pr	H	H	0	H	H	CF ₃	Me	Cl
	Me	Pr-c	H	H	0	H	H	CF ₃	Me	Cl
15	Me	CH ₃ Pr-c	H	H	0	H	H	CF ₃	Me	Cl
	-(CH ₂) ₂ -		H	H	0	H	H	CF ₃	Me	Cl
	-(CH ₂) ₃ -		H	H	0	H	H	CF ₃	Me	Cl
	-(CH ₂) ₄ -		H	H	0	H	H	CF ₃	Me	Cl
20	-(CH ₂) ₅ -		H	H	0	H	H	CF ₃	Me	Cl
	H	-(CH ₂) ₃ -	H	0	H	H	CF ₃	Me	Cl	
	H	-(CH ₂) ₄ -	H	0	H	H	CF ₃	Me	Cl	
	H	-(CH ₂) ₅ -	H	0	H	H	CF ₃	Me	Cl	
25	H	-(CH ₂) ₆ -	H	0	H	H	CF ₃	Me	Cl	
	Me	Et	H	H	2	H	H	H	H	H

30

35


40

45

50

55

Table 4

									
R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Z ³	R ³¹	R ³²
Me	Me	H	H	2	H	H	O	Me	F
Me	Me	H	H	2	H	H	O	Me	Cl
Me	Me	H	H	2	H	H	O	Me	OMe
Me	Me	H	H	2	H	H	O	Me	OEt
Me	Me	H	H	2	H	H	O	Me	OPr-i
Me	Me	H	H	2	H	H	O	Me	OPh
Me	Me	H	H	2	H	H	O	Me	OCHF ₂
Me	Me	H	H	2	H	H	O	Me	Me
Me	Me	H	H	2	H	H	O	Me	CF ₃
Me	Me	H	H	2	H	H	O	Me	CN
Me	Me	H	H	2	H	H	O	OCHF ₂	F
Me	Me	H	H	2	H	H	O	OCHF ₂	Cl
Me	Me	H	H	2	H	H	O	OCHF ₂	Me

	Me	Me	H	H	2	H	H	O	OCHF ₂	CF ₃
5	Me	Me	H	H	2	H	H	O	OCHF ₂	CN
	Me	Me	H	H	2	H	H	O	CF ₃	F
	Me	Me	H	H	2	H	H	O	CF ₃	Cl
	Me	Me	H	H	2	H	H	O	CF ₃	OMe
10	Me	Me	H	H	2	H	H	O	CF ₃	OEt
	Me	Me	H	H	2	H	H	O	CF ₃	OPr-i
	Me	Me	H	H	2	H	H	O	CF ₃	OPh
	Me	Me	H	H	2	H	H	O	CF ₃	OCHF ₂
15	Me	Me	H	H	2	H	H	O	CF ₃	SMe
	Me	Me	H	H	2	H	H	O	CF ₃	SOMe
	Me	Me	H	H	2	H	H	O	CF ₃	SO ₂ Me
	Me	Me	H	H	2	H	H	O	CF ₃	SEt
20	Me	Me	H	H	2	H	H	O	CF ₃	SOEt
	Me	Me	H	H	2	H	H	O	CF ₃	SO ₂ Et
	Me	Me	H	H	2	H	H	O	CF ₃	SPr-i
	Me	Me	H	H	2	H	H	O	CF ₃	SOPr-i
25	Me	Me	H	H	2	H	H	O	CF ₃	SO ₂ Pr-i
	Me	Me	H	H	2	H	H	O	CF ₃	SPh
	Me	Me	H	H	2	H	H	O	CF ₃	SOPh
	Me	Me	H	H	2	H	H	O	CF ₃	SO ₂ Ph
30	Me	Me	H	H	2	H	H	O	CF ₃	SCHF ₂
	Me	Me	H	H	2	H	H	O	CF ₃	SOCHF ₂
	Me	Me	H	H	2	H	H	O	CF ₃	SO ₂ CHF ₂
	Me	Me	H	H	2	H	H	O	CF ₃	SCF ₃
35	Me	Me	H	H	2	H	H	O	CF ₃	SOCF ₃
	Me	Me	H	H	2	H	H	O	CF ₃	SO ₂ CF ₃
	Me	Me	H	H	2	H	H	O	CF ₃	NH ₂
	Me	Me	H	H	2	H	H	O	CF ₃	NHC(=O)Me
40	Me	Me	H	H	2	H	H	O	CF ₃	NHC(=O)Ph
	Me	Me	H	H	2	H	H	O	CF ₃	NHC(=O)CH ₂ Ph
	Me	Me	H	H	2	H	H	O	CF ₃	NHC(=O)CF ₃
	Me	Me	H	H	2	H	H	O	CF ₃	NHSO ₂ Me
45	Me	Me	H	H	2	H	H	O	CF ₃	NHSO ₂ Ph
	Me	Me	H	H	2	H	H	O	CF ₃	NHSO ₂ CHF ₂
	Me	Me	H	H	2	H	H	O	CF ₃	NHSO ₂ CF ₃
	Me	Me	H	H	2	H	H	O	CF ₃	NHMe
50	Me	Me	H	H	2	H	H	O	CF ₃	NHPh
	Me	Me	H	H	2	H	H	O	CF ₃	N(Me)C(=O)Me
	Me	Me	H	H	2	H	H	O	CF ₃	N(Me)C(=O)Ph
	Me	Me	H	H	2	H	H	O	CF ₃	N(Me)C(=O)CH ₂ Ph
	Me	Me	H	H	2	H	H	O	CF ₃	N(Me)C(=O)CF ₃
55	Me	Me	H	H	2	H	H	O	CF ₃	N(Me)SO ₂ Me
	Me	Me	H	H	2	H	H	O	CF ₃	N(Me)SO ₂ Ph

5	Me	Me	H	H	2	H	H	O	CF ₃	N(Me)SO ₂ CHF ₂
	Me	Me	H	H	2	H	H	O	CF ₃	N(Me)SO ₂ CF ₃
	Me	Me	H	H	2	H	H	O	CF ₃	N(Me) ₂
	Me	Me	H	H	2	H	H	O	CF ₃	N(Me)Ph
	Me	Me	H	H	2	H	H	O	CF ₃	Me
10	Me	Me	H	H	2	H	H	O	CF ₃	CF ₃
	Me	Me	H	H	2	H	H	O	CF ₃	CN
	Me	Me	H	H	2	H	H	O	Ph	Me
	H	H	H	H	2	H	H	O	CF ₃	Me
15	Me	H	H	H	2	H	H	O	CF ₃	Me
	Me	H	Me	H	2	H	H	O	CF ₃	Me
	Me	Me	Me	H	2	H	H	O	CF ₃	Me
	Me	Me	H	H	2	Me	H	O	CF ₃	Me
20	Me	Me	H	H	2	Et	H	O	CF ₃	Me
	Me	Me	H	H	2	Pr-i	H	O	CF ₃	Me
	Me	Me	H	H	2	Me	Me	O	CF ₃	Me
	Me	Et	H	H	2	H	H	O	CF ₃	Me
	Et	Et	H	H	2	H	H	O	CF ₃	Me
25	Me	Pr-i	H	H	2	H	H	O	CF ₃	Me
	Me	Pr	H	H	2	H	H	O	CF ₃	Me
	Me	Pr-c	H	H	2	H	H	O	CF ₃	Me
	Me	CH ₂ Pr-c	H	H	2	H	H	O	CF ₃	Me
30	-(CH ₂) ₂ -		H	H	2	H	H	O	CF ₃	Me
	-(CH ₂) ₃ -		H	H	2	H	H	O	CF ₃	Me
	-(CH ₂) ₄ -		H	H	2	H	H	O	CF ₃	Me
	-(CH ₂) ₅ -		H	H	2	H	H	O	CF ₃	Me
35	H	-(CH ₂) ₃ -	H	H	2	H	H	O	CF ₃	Me
	H	-(CH ₂) ₄ -	H	H	2	H	H	O	CF ₃	Me
	H	-(CH ₂) ₅ -	H	H	2	H	H	O	CF ₃	Me
	H	-(CH ₂) ₆ -	H	H	2	H	H	O	CF ₃	Me
40	Me	Me	H	H	2	H	H	S	Me	F
	Me	Me	H	H	2	H	H	S	Me	Cl
	Me	Me	H	H	2	H	H	S	Me	OMe
	Me	Me	H	H	2	H	H	S	Me	OEt
45	Me	Me	H	H	2	H	H	S	Me	OPr-i
	Me	Me	H	H	2	H	H	S	Me	OPh
	Me	Me	H	H	2	H	H	S	Me	OCHF ₂
	Me	Me	H	H	2	H	H	S	OCHF ₂	F
50	Me	Me	H	H	2	H	H	S	OCHF ₂	Cl
	Me	Me	H	H	2	H	H	S	OCHF ₂	Me
	Me	Me	H	H	2	H	H	S	OCHF ₂	CF ₃
	Me	Me	H	H	2	H	H	S	OCHF ₂	CN
55	Me	Me	H	H	2	H	H	S	CF ₃	F
	Me	Me	H	H	2	H	H	S	CF ₃	Cl

	Me	Me	H	H	2	H	H	S	CF ₃	OMe
5	Me	Me	H	H	2	H	H	S	CF ₃	OEt
	Me	Me	H	H	2	H	H	S	CF ₃	OPh
	Me	Me	H	H	2	H	H	S	CF ₃	OCHF ₂
	Me	Me	H	H	2	H	H	S	CF ₃	SMe
10	Me	Me	H	H	2	H	H	S	CF ₃	SOMe
	Me	Me	H	H	2	H	H	S	CF ₃	SO ₂ Me
	Me	Me	H	H	2	H	H	S	CF ₃	SEt
	Me	Me	H	H	2	H	H	S	CF ₃	SOEt
15	Me	Me	H	H	2	H	H	S	CF ₃	SO ₂ Et
	Me	Me	H	H	2	H	H	S	CF ₃	SP <i>r</i> -i
	Me	Me	H	H	2	H	H	S	CF ₃	SOP <i>r</i> -i
	Me	Me	H	H	2	H	H	S	CF ₃	SO ₂ <i>Pr</i> -i
20	Me	Me	H	H	2	H	H	S	CF ₃	SPh
	Me	Me	H	H	2	H	H	S	CF ₃	SOPh
	Me	Me	H	H	2	H	H	S	CF ₃	SO ₂ Ph
	Me	Me	H	H	2	H	H	S	CF ₃	SCHF ₂
25	Me	Me	H	H	2	H	H	S	CF ₃	SOCHF ₂
	Me	Me	H	H	2	H	H	S	CF ₃	SO ₂ CHF ₂
	Me	Me	H	H	2	H	H	S	CF ₃	SCF ₃
	Me	Me	H	H	2	H	H	S	CF ₃	SOCF ₃
30	Me	Me	H	H	2	H	H	S	CF ₃	SO ₂ CF ₃
	Me	Me	H	H	2	H	H	S	CF ₃	NH ₂
	Me	Me	H	H	2	H	H	S	CF ₃	NHC(=O)Me
	Me	Me	H	H	2	H	H	S	CF ₃	NHC(=O)Ph
35	Me	Me	H	H	2	H	H	S	CF ₃	NHC(=O)CH ₂ Ph
	Me	Me	H	H	2	H	H	S	CF ₃	NHC(=O)CF ₃
	Me	Me	H	H	2	H	H	S	CF ₃	NHSO ₂ Me
	Me	Me	H	H	2	H	H	S	CF ₃	NHSO ₂ Ph
	Me	Me	H	H	2	H	H	S	CF ₃	NHSO ₂ CHF ₂
40	Me	Me	H	H	2	H	H	S	CF ₃	NHSO ₂ CF ₃
	Me	Me	H	H	2	H	H	S	CF ₃	NHMe
	Me	Me	H	H	2	H	H	S	CF ₃	NHPh
	Me	Me	H	H	2	H	H	S	CF ₃	N(Me)C(=O)Me
45	Me	Me	H	H	2	H	H	S	CF ₃	N(Me)C(=O)Ph
	Me	Me	H	H	2	H	H	S	CF ₃	N(Me)C(=O)CH ₂ Ph
	Me	Me	H	H	2	H	H	S	CF ₃	N(Me)C(=O)CF ₃
	Me	Me	H	H	2	H	H	S	CF ₃	N(Me)SO ₂ Me
50	Me	Me	H	H	2	H	H	S	CF ₃	N(Me)SO ₂ Ph
	Me	Me	H	H	2	H	H	S	CF ₃	N(Me)SO ₂ CHF ₂
	Me	Me	H	H	2	H	H	S	CF ₃	N(Me)SO ₂ CF ₃
	Me	Me	H	H	2	H	H	S	CF ₃	N(Me) ₂
55	Me	Me	H	H	2	H	H	S	CF ₃	N(Me)Ph
	Me	Me	H	H	2	H	H	S	CF ₃	Me

5	Me	Me	H	H	2	H	H	S	CF ₃	CN
	H	H	H	H	2	H	H	S	CF ₃	Cl
	Me	H	H	H	2	H	H	S	CF ₃	Cl
	Me	H	Me	H	2	H	H	S	CF ₃	Cl
	Me	Me	Me	H	2	H	H	S	CF ₃	Cl
10	Me	Me	H	H	2	Me	H	S	CF ₃	Cl
	Me	Me	H	H	2	Et	H	S	CF ₃	Cl
	Me	Me	H	H	2	Pr-i	H	S	CF ₃	Cl
	Me	Me	H	H	2	Me	Me	S	CF ₃	Cl
15	Me	Et	H	H	2	H	H	S	CF ₃	Cl
	Et	Et	H	H	2	H	H	S	CF ₃	Cl
	Me	Pr-i	H	H	2	H	H	S	CF ₃	Cl
	Me	Pr	H	H	2	H	H	S	CF ₃	Cl
20	Me	Pr-c	H	H	2	H	H	S	CF ₃	Cl
	Me	CH ₂ Pr-c	H	H	2	H	H	S	CF ₃	Cl
	-(CH ₂) ₂ -		H	H	2	H	H	S	CF ₃	Cl
	-(CH ₂) ₃ -		H	H	2	H	H	S	CF ₃	Cl
25	-(CH ₂) ₄ -		H	H	2	H	H	S	CF ₃	Cl
	-(CH ₂) ₅ -		H	H	2	H	H	S	CF ₃	Cl
	H	-(CH ₂) ₃ -	H	H	2	H	H	S	CF ₃	Cl
30	H	-(CH ₂) ₄ -	H	H	2	H	H	S	CF ₃	Cl
	H	-(CH ₂) ₅ -	H	H	2	H	H	S	CF ₃	Cl
	H	-(CH ₂) ₆ -	H	H	2	H	H	S	CF ₃	Cl
	Me	Me	H	H	1	H	H	O	Me	F
35	Me	Me	H	H	1	H	H	O	Me	Cl
	Me	Me	H	H	1	H	H	O	Me	OMe
	Me	Me	H	H	1	H	H	O	Me	OEt
	Me	Me	H	H	1	H	H	O	Me	OPr-i
	Me	Me	H	H	1	H	H	O	Me	OPh
40	Me	Me	H	H	1	H	H	O	Me	OCHF ₂
	Me	Me	H	H	1	H	H	O	Me	Me
	Me	Me	H	H	1	H	H	O	Me	CF ₃
	Me	Me	H	H	1	H	H	O	Me	CN
45	Me	Me	H	H	1	H	H	O	OCHF ₂	F
	Me	Me	H	H	1	H	H	O	OCHF ₂	Cl
	Me	Me	H	H	1	H	H	O	OCHF ₂	Me
	Me	Me	H	H	1	H	H	O	OCHF ₂	CF ₃
	Me	Me	H	H	1	H	H	O	OCHF ₂	CN
50	Me	Me	H	H	1	H	H	O	CF ₃	F
	Me	Me	H	H	1	H	H	O	CF ₃	Cl
	Me	Me	H	H	1	H	H	O	CF ₃	OMe
	Me	Me	H	H	1	H	H	O	CF ₃	OEt
55	Me	Me	H	H	1	H	H	O	CF ₃	OPr-i
	Me	Me	H	H	1	H	H	O	CF ₃	OPh

5	Me	Me	H	H	1	H	H	O	CF ₃	OCHF ₂
	Me	Me	H	H	1	H	H	O	CF ₃	SMe
	Me	Me	H	H	1	H	H	O	CF ₃	SO ₂ Me
	Me	Me	H	H	1	H	H	O	CF ₃	SEt
	Me	Me	H	H	1	H	H	O	CF ₃	SO ₂ Et
10	Me	Me	H	H	1	H	H	O	CF ₃	SP _r -i
	Me	Me	H	H	1	H	H	O	CF ₃	SO ₂ Pr-i
	Me	Me	H	H	1	H	H	O	CF ₃	SPh
	Me	Me	H	H	1	H	H	O	CF ₃	SO ₂ Ph
	Me	Me	H	H	1	H	H	O	CF ₃	SCHF ₂
15	Me	Me	H	H	1	H	H	O	CF ₃	SO ₂ CHF ₂
	Me	Me	H	H	1	H	H	O	CF ₃	SCF ₃
	Me	Me	H	H	1	H	H	O	CF ₃	SO ₂ CF ₃
	Me	Me	H	H	1	H	H	O	CF ₃	NH ₂
20	Me	Me	H	H	1	H	H	O	CF ₃	NHC(=O)Me
	Me	Me	H	H	1	H	H	O	CF ₃	NHC(=O)Ph
	Me	Me	H	H	1	H	H	O	CF ₃	NHC(=O)CH ₂ Ph
	Me	Me	H	H	1	H	H	O	CF ₃	NHC(=O)CF ₃
25	Me	Me	H	H	1	H	H	O	CF ₃	NHSO ₂ Me
	Me	Me	H	H	1	H	H	O	CF ₃	NHSO ₂ Ph
	Me	Me	H	H	1	H	H	O	CF ₃	NHSO ₂ CHF ₂
	Me	Me	H	H	1	H	H	O	CF ₃	NHSO ₂ CF ₃
30	Me	Me	H	H	1	H	H	O	CF ₃	NHMe
	Me	Me	H	H	1	H	H	O	CF ₃	NHPh
	Me	Me	H	H	1	H	H	O	CF ₃	N(Me)C(=O)Me
	Me	Me	H	H	1	H	H	O	CF ₃	N(Me)C(=O)Ph
	Me	Me	H	H	1	H	H	O	CF ₃	N(Me)C(=O)CH ₂ Ph
35	Me	Me	H	H	1	H	H	O	CF ₃	N(Me)C(=O)CF ₃
	Me	Me	H	H	1	H	H	O	CF ₃	N(Me)SO ₂ Me
	Me	Me	H	H	1	H	H	O	CF ₃	N(Me)SO ₂ Ph
	Me	Me	H	H	1	H	H	O	CF ₃	N(Me)SO ₂ CHF ₂
40	Me	Me	H	H	1	H	H	O	CF ₃	N(Me)SO ₂ CF ₃
	Me	Me	H	H	1	H	H	O	CF ₃	N(Me) ₂
	Me	Me	H	H	1	H	H	O	CF ₃	N(Me)Ph
	Me	Me	H	H	1	H	H	O	CF ₃	Me
45	Me	Me	H	H	1	H	H	O	CF ₃	CF ₃
	Me	Me	H	H	1	H	H	O	CF ₃	CN
	Me	Me	H	H	1	H	H	O	Ph	Me
	H	H	H	H	1	H	H	O	CF ₃	Me
50	Me	H	H	H	1	H	H	O	CF ₃	Me
	Me	H	Me	H	1	H	H	O	CF ₃	Me
	Me	Me	Me	H	1	H	H	O	CF ₃	Me
	Me	Me	H	H	1	Me	H	O	CF ₃	Me
55	Me	Me	H	H	1	Et	H	O	CF ₃	Me

5	Me	Me	H	H	1	Pr-i	H	O	CF ₃	Me
	Me	Me	H	H	1	Me	Me	O	CF ₃	Me
	Me	Et	H	H	1	H	H	O	CF ₃	Me
	Et	Et	H	H	1	H	H	O	CF ₃	Me
	Me	Pr-i	H	H	1	H	H	O	CF ₃	Me
10	Me	Pr	H	H	1	H	H	O	CF ₃	Me
	Me	Pr-c	H	H	1	H	H	O	CF ₃	Me
	Me	CH ₂ Pr-c	H	H	1	H	H	O	CF ₃	Me
		-(CH ₂) ₂ -	H	H	1	H	H	O	CF ₃	Me
15		-(CH ₂) ₃ -	H	H	1	H	H	O	CF ₃	Me
		-(CH ₂) ₄ -	H	H	1	H	H	O	CF ₃	Me
		-(CH ₂) ₅ -	H	H	1	H	H	O	CF ₃	Me
	H	-(CH ₂) ₃ -	H	H	1	H	H	O	CF ₃	Me
20	H	-(CH ₂) ₄ -	H	H	1	H	H	O	CF ₃	Me
	H	-(CH ₂) ₅ -	H	H	1	H	H	O	CF ₃	Me
	H	-(CH ₂) ₆ -	H	H	1	H	H	O	CF ₃	Me
	Me	Me	H	H	1	H	H	S	Me	F
25	Me	Me	H	H	1	H	H	S	Me	Cl
	Me	Me	H	H	1	H	H	S	Me	OMe
	Me	Me	H	H	1	H	H	S	Me	OEt
	Me	Me	H	H	1	H	H	S	Me	OPr-i
30	Me	Me	H	H	1	H	H	S	Me	OPh
	Me	Me	H	H	1	H	H	S	Me	OCHF ₂
	Me	Me	H	H	1	H	H	S	OCHF ₂	F
	Me	Me	H	H	1	H	H	S	OCHF ₂	Cl
35	Me	Me	H	H	1	H	H	S	OCHF ₂	Me
	Me	Me	H	H	1	H	H	S	OCHF ₂	CF ₃
	Me	Me	H	H	1	H	H	S	OCHF ₂	CN
	Me	Me	H	H	1	H	H	S	CF ₃	F
	Me	Me	H	H	1	H	H	S	CF ₃	Cl
40	Me	Me	H	H	1	H	H	S	CF ₃	OMe
	Me	Me	H	H	1	H	H	S	CF ₃	OEt
	Me	Me	H	H	1	H	H	S	CF ₃	OPh
	Me	Me	H	H	1	H	H	S	CF ₃	OCHF ₂
45	Me	Me	H	H	1	H	H	S	CF ₃	SMe
	Me	Me	H	H	1	H	H	S	CF ₃	SO ₂ Me
	Me	Me	H	H	1	H	H	S	CF ₃	SEt
	Me	Me	H	H	1	H	H	S	CF ₃	SO ₂ Et
50	Me	Me	H	H	1	H	H	S	CF ₃	SPr-i
	Me	Me	H	H	1	H	H	S	CF ₃	SO ₂ Pr-i
	Me	Me	H	H	1	H	H	S	CF ₃	SPh
	Me	Me	H	H	1	H	H	S	CF ₃	SO ₂ Ph
55	Me	Me	H	H	1	H	H	S	CF ₃	SCHF ₂
	Me	Me	H	H	1	H	H	S	CF ₃	SO ₂ CHF ₂

5	Me	Me	H	H	1	H	H	S	CF ₃	SCF ₃
	Me	Me	H	H	1	H	H	S	CF ₃	SO ₂ CF ₃
	Me	Me	H	H	1	H	H	S	CF ₃	NH ₂
	Me	Me	H	H	1	H	H	S	CF ₃	NHC(=O)Me
	Me	Me	H	H	1	H	H	S	CF ₃	NHC(=O)Ph
10	Me	Me	H	H	1	H	H	S	CF ₃	NHC(=O)CH ₂ Ph
	Me	Me	H	H	1	H	H	S	CF ₃	NHC(=O)CF ₃
	Me	Me	H	H	1	H	H	S	CF ₃	NHSO ₂ Me
	Me	Me	H	H	1	H	H	S	CF ₃	NHSO ₂ Ph
15	Me	Me	H	H	1	H	H	S	CF ₃	NHSO ₂ CHF ₂
	Me	Me	H	H	1	H	H	S	CF ₃	NHSO ₂ CF ₃
	Me	Me	H	H	1	H	H	S	CF ₃	NHMe
	Me	Me	H	H	1	H	H	S	CF ₃	NHPh
20	Me	Me	H	H	1	H	H	S	CF ₃	N(Me)C(=O)Me
	Me	Me	H	H	1	H	H	S	CF ₃	N(Me)C(=O)Ph
	Me	Me	H	H	1	H	H	S	CF ₃	N(Me)C(=O)CH ₂ Ph
	Me	Me	H	H	1	H	H	S	CF ₃	N(Me)C(=O)CF ₃
	Me	Me	H	H	1	H	H	S	CF ₃	N(Me)SO ₂ Me
25	Me	Me	H	H	1	H	H	S	CF ₃	N(Me)SO ₂ Ph
	Me	Me	H	H	1	H	H	S	CF ₃	N(Me)SO ₂ CHF ₂
	Me	Me	H	H	1	H	H	S	CF ₃	N(Me)SO ₂ CF ₃
	Me	Me	H	H	1	H	H	S	CF ₃	N(Me) ₂
30	Me	Me	H	H	1	H	H	S	CF ₃	N(Me)Ph
	Me	Me	H	H	1	H	H	S	CF ₃	Me
	Me	Me	H	H	1	H	H	S	CF ₃	CN
	H	H	H	H	1	H	H	S	CF ₃	Cl
35	Me	H	H	H	1	H	H	S	CF ₃	Cl
	Me	H	Me	H	1	H	H	S	CF ₃	Cl
	Me	Me	Me	H	1	H	H	S	CF ₃	Cl
	Me	Me	H	H	1	Me	H	S	CF ₃	Cl
40	Me	Me	H	H	1	Et	H	S	CF ₃	Cl
	Me	Me	H	H	1	Pr-i	H	S	CF ₃	Cl
	Me	Me	H	H	1	Me	Me	S	CF ₃	Cl
	Me	Et	H	H	1	H	H	S	CF ₃	Cl
45	Et	Et	H	H	1	H	H	S	CF ₃	Cl
	Me	Pr-i	H	H	1	H	H	S	CF ₃	Cl
	Me	Pr	H	H	1	H	H	S	CF ₃	Cl
	Me	Pr-c	H	H	1	H	H	S	CF ₃	Cl
	Me	CH ₂ Pr-c	H	H	1	H	H	S	CF ₃	Cl
50		-(CH ₂) ₂ -	H	H	1	H	H	S	CF ₃	Cl
		-(CH ₂) ₃ -	H	H	1	H	H	S	CF ₃	Cl
		-(CH ₂) ₄ -	H	H	1	H	H	S	CF ₃	Cl
		-(CH ₂) ₅ -	H	H	1	H	H	S	CF ₃	Cl
55	H	-(CH ₂) ₃ -	H	H	1	H	H	S	CF ₃	Cl


5	H	-(CH ₂) ₄ -		H	1	H	H	S	CF ₃	Cl
	H	-(CH ₂) ₅ -		H	1	H	H	S	CF ₃	Cl
	H	-(CH ₂) ₆ -		H	1	H	H	S	CF ₃	Cl
10	Me	Me	H	H	0	H	H	O	Me	F
	Me	Me	H	H	0	H	H	O	Me	Cl
	Me	Me	H	H	0	H	H	O	Me	OMe
15	Me	Me	H	H	0	H	H	O	Me	OEt
	Me	Me	H	H	0	H	H	O	Me	OPr-i
	Me	Me	H	H	0	H	H	O	Me	OPh
20	Me	Me	H	H	0	H	H	O	Me	OCHF ₂
	Me	Me	H	H	0	H	H	O	Me	Me
	Me	Me	H	H	0	H	H	O	Me	CF ₃
25	Me	Me	H	H	0	H	H	O	Me	CN
	Me	Me	H	H	0	H	H	O	OCHF ₂	F
	Me	Me	H	H	0	H	H	O	OCHF ₂	Cl
30	Me	Me	H	H	0	H	H	O	OCHF ₂	Me
	Me	Me	H	H	0	H	H	O	OCHF ₂	CF ₃
	Me	Me	H	H	0	H	H	O	OCHF ₂	CN
35	Me	Me	H	H	0	H	H	O	CF ₃	F
	Me	Me	H	H	0	H	H	O	CF ₃	Cl
	Me	Me	H	H	0	H	H	O	CF ₃	OMe
40	Me	Me	H	H	0	H	H	O	CF ₃	OEt
	Me	Me	H	H	0	H	H	O	CF ₃	OPr-i
	Me	Me	H	H	0	H	H	O	CF ₃	OPh
45	Me	Me	H	H	0	H	H	O	CF ₃	OCHF ₂
	Me	Me	H	H	0	H	H	O	CF ₃	SMe
	Me	Me	H	H	0	H	H	O	CF ₃	SO ₂ Me
50	Me	Me	H	H	0	H	H	O	CF ₃	SEt
	Me	Me	H	H	0	H	H	O	CF ₃	SO ₂ Et
	Me	Me	H	H	0	H	H	O	CF ₃	SPr-i
55	Me	Me	H	H	0	H	H	O	CF ₃	SO ₂ Pr-i
	Me	Me	H	H	0	H	H	O	CF ₃	SPh
	Me	Me	H	H	0	H	H	O	CF ₃	SO ₂ Ph
	Me	Me	H	H	0	H	H	O	CF ₃	SCHF ₂
	Me	Me	H	H	0	H	H	O	CF ₃	SO ₂ CHF ₂
	Me	Me	H	H	0	H	H	O	CF ₃	SCF ₃
	Me	Me	H	H	0	H	H	O	CF ₃	SO ₂ CF ₃
	Me	Me	H	H	0	H	H	O	CF ₃	NH ₂
	Me	Me	H	H	0	H	H	O	CF ₃	NHC(=O)Me
	Me	Me	H	H	0	H	H	O	CF ₃	NHC(=O)Ph
	Me	Me	H	H	0	H	H	O	CF ₃	NHC(=O)CH ₂ Ph
	Me	Me	H	H	0	H	H	O	CF ₃	NHC(=O)CF ₃
	Me	Me	H	H	0	H	H	O	CF ₃	NHSO ₂ Me
	Me	Me	H	H	0	H	H	O	CF ₃	NHSO ₂ Ph

5	Me	Me	H	H	0	H	H	O	CF ₃	NHSO ₂ CHF ₂
	Me	Me	H	H	0	H	H	O	CF ₃	NHSO ₂ CF ₃
	Me	Me	H	H	0	H	H	O	CF ₃	NHMe
	Me	Me	H	H	0	H	H	O	CF ₃	NHPh
	Me	Me	H	H	0	H	H	O	CF ₃	N(Me)C(=O)Me
10	Me	Me	H	H	0	H	H	O	CF ₃	N(Me)C(=O)Ph
	Me	Me	H	H	0	H	H	O	CF ₃	N(Me)C(=O)CH ₂ Ph
	Me	Me	H	H	0	H	H	O	CF ₃	N(Me)C(=O)CF ₃
	Me	Me	H	H	0	H	H	O	CF ₃	N(Me)SO ₂ Me
15	Me	Me	H	H	0	H	H	O	CF ₃	N(Me)SO ₂ Ph
	Me	Me	H	H	0	H	H	O	CF ₃	N(Me)SO ₂ CHF ₂
	Me	Me	H	H	0	H	H	O	CF ₃	N(Me)SO ₂ CF ₃
	Me	Me	H	H	0	H	H	O	CF ₃	N(Me) ₂
20	Me	Me	H	H	0	H	H	O	CF ₃	N(Me)Ph
	Me	Me	H	H	0	H	H	O	CF ₃	Me
	Me	Me	H	H	0	H	H	O	CF ₃	CF ₃
	Me	Me	H	H	0	H	H	O	CF ₃	CN
25	Me	Me	H	H	0	H	H	O	Ph	Me
	H	H	H	H	0	H	H	O	CF ₃	Me
	Me	H	H	H	0	H	H	O	CF ₃	Me
	Me	H	Me	H	0	H	H	O	CF ₃	Me
	Me	Me	Me	H	0	H	H	O	CF ₃	Me
30	Me	Me	H	H	0	Me	H	O	CF ₃	Me
	Me	Me	H	H	0	Et	H	O	CF ₃	Me
	Me	Me	H	H	0	Pr-i	H	O	CF ₃	Me
	Me	Me	H	H	0	Me	Me	O	CF ₃	Me
35	Me	Et	H	H	0	H	H	O	CF ₃	Me
	Et	Et	H	H	0	H	H	O	CF ₃	Me
	Me	Pr-i	H	H	0	H	H	O	CF ₃	Me
	Me	Pr	H	H	0	H	H	O	CF ₃	Me
40	Me	Pr-c	H	H	0	H	H	O	CF ₃	Me
	Me	CH ₂ Pr-c	H	H	0	H	H	O	CF ₃	Me
		-(CH ₂) ₂ -	H	H	0	H	H	O	CF ₃	Me
		-(CH ₂) ₃ -	H	H	0	H	H	O	CF ₃	Me
45		-(CH ₂) ₄ -	H	H	0	H	H	O	CF ₃	Me
		-(CH ₂) ₅ -	H	H	0	H	H	O	CF ₃	Me
	H	-(CH ₂) ₃ -	H	0	H	H	H	O	CF ₃	Me
	H	-(CH ₂) ₄ -	H	0	H	H	H	O	CF ₃	Me
50	H	-(CH ₂) ₅ -	H	0	H	H	H	O	CF ₃	Me
	H	-(CH ₂) ₆ -	H	0	H	H	H	O	CF ₃	Me
	Me	Me	H	H	0	H	H	S	Me	F
	Me	Me	H	H	0	H	H	S	Me	Cl
	Me	Me	H	H	0	H	H	S	Me	OMe
55	Me	Me	H	H	0	H	H	S	Me	OEt

5	Me	Me	H	H	0	H	H	S	Me	OPr-i
	Me	Me	H	H	0	H	H	S	Me	OPh
	Me	Me	H	H	0	H	H	S	Me	OCHF ₂
	Me	Me	H	H	0	H	H	S	OCHF ₂	F
	Me	Me	H	H	0	H	H	S	OCHF ₂	Cl
10	Me	Me	H	H	0	H	H	S	OCHF ₂	Me
	Me	Me	H	H	0	H	H	S	OCHF ₂	CF ₃
	Me	Me	H	H	0	H	H	S	OCHF ₂	CN
	Me	Me	H	H	0	H	H	S	CF ₃	F
15	Me	Me	H	H	0	H	H	S	CF ₃	Cl
	Me	Me	H	H	0	H	H	S	CF ₃	OMe
	Me	Me	H	H	0	H	H	S	CF ₃	OEt
	Me	Me	H	H	0	H	H	S	CF ₃	OPh
20	Me	Me	H	H	0	H	H	S	CF ₃	OCHF ₂
	Me	Me	H	H	0	H	H	S	CF ₃	SMe
	Me	Me	H	H	0	H	H	S	CF ₃	SO ₂ Me
	Me	Me	H	H	0	H	H	S	CF ₃	SEt
25	Me	Me	H	H	0	H	H	S	CF ₃	SO ₂ Et
	Me	Me	H	H	0	H	H	S	CF ₃	SPr-i
	Me	Me	H	H	0	H	H	S	CF ₃	SO ₂ Pr-i
	Me	Me	H	H	0	H	H	S	CF ₃	SPh
30	Me	Me	H	H	0	H	H	S	CF ₃	SO ₂ Ph
	Me	Me	H	H	0	H	H	S	CF ₃	SCHF ₂
	Me	Me	H	H	0	H	H	S	CF ₃	SO ₂ CHF ₂
	Me	Me	H	H	0	H	H	S	CF ₃	SCF ₃
	Me	Me	H	H	0	H	H	S	CF ₃	SO ₂ CF ₃
35	Me	Me	H	H	0	H	H	S	CF ₃	NH ₂
	Me	Me	H	H	0	H	H	S	CF ₃	NHC(=O)Me
	Me	Me	H	H	0	H	H	S	CF ₃	NHC(=O)Ph
	Me	Me	H	H	0	H	H	S	CF ₃	NHC(=O)CH ₂ Ph
40	Me	Me	H	H	0	H	H	S	CF ₃	NHC(=O)CF ₃
	Me	Me	H	H	0	H	H	S	CF ₃	NHSO ₂ Me
	Me	Me	H	H	0	H	H	S	CF ₃	NHSO ₂ Ph
	Me	Me	H	H	0	H	H	S	CF ₃	NHSO ₂ CHF ₂
45	Me	Me	H	H	0	H	H	S	CF ₃	NHSO ₂ CF ₃
	Me	Me	H	H	0	H	H	S	CF ₃	NHMe
	Me	Me	H	H	0	H	H	S	CF ₃	NHPh
	Me	Me	H	H	0	H	H	S	CF ₃	N(Me)C(=O)Me
50	Me	Me	H	H	0	H	H	S	CF ₃	N(Me)C(=O)Ph
	Me	Me	H	H	0	H	H	S	CF ₃	N(Me)C(=O)CH ₂ Ph
	Me	Me	H	H	0	H	H	S	CF ₃	N(Me)C(=O)CF ₃
	Me	Me	H	H	0	H	H	S	CF ₃	N(Me)SO ₂ Me
	Me	Me	H	H	0	H	H	S	CF ₃	N(Me)SO ₂ Ph
55	Me	Me	H	H	0	H	H	S	CF ₃	N(Me)SO ₂ CHF ₂

5	Me	Me	H	H	0	H	H	S	CF ₃	N(Me)SO ₂ CF ₃
	Me	Me	H	H	0	H	H	S	CF ₃	N(Me) ₂
	Me	Me	H	H	0	H	H	S	CF ₃	N(Me)Ph
	Me	Me	H	H	0	H	H	S	CF ₃	Me
	Me	Me	H	H	0	H	H	S	CF ₃	CN
10	H	H	H	H	0	H	H	S	CF ₃	Cl
	Me	H	H	H	0	H	H	S	CF ₃	Cl
	Me	H	Me	H	0	H	H	S	CF ₃	Cl
	Me	Me	Me	H	0	H	H	S	CF ₃	Cl
15	Me	Me	H	H	0	Me	H	S	CF ₃	Cl
	Me	Me	H	H	0	Et	H	S	CF ₃	Cl
	Me	Me	H	H	0	Pr-i	H	S	CF ₃	Cl
	Me	Me	H	H	0	Me	Me	S	CF ₃	Cl
20	Me	Et	H	H	0	H	H	S	CF ₃	Cl
	Et	Et	H	H	0	H	H	S	CF ₃	Cl
	Me	Pr-i	H	H	0	H	H	S	CF ₃	Cl
	Me	Pr	H	H	0	H	H	S	CF ₃	Cl
25	Me	Pr-c	H	H	0	H	H	S	CF ₃	Cl
	Me	CH ₂ Pr-c	H	H	0	H	H	S	CF ₃	Cl
	-(CH ₂) ₂ -		H	H	0	H	H	S	CF ₃	Cl
	-(CH ₂) ₃ -		H	H	0	H	H	S	CF ₃	Cl
30	-(CH ₂) ₄ -		H	H	0	H	H	S	CF ₃	Cl
	-(CH ₂) ₅ -		H	H	0	H	H	S	CF ₃	Cl
	H	-(CH ₂) ₃ -	H	0	H	H	H	S	CF ₃	Cl
	H	-(CH ₂) ₄ -	H	0	H	H	H	S	CF ₃	Cl
35	H	-(CH ₂) ₅ -	H	0	H	H	H	S	CF ₃	Cl
	H	-(CH ₂) ₆ -	H	0	H	H	H	S	CF ₃	Cl

Table 5

									
R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Z ⁴	R ³³	R ³⁴
Me	Me	H	H	2	H	H	NMe	Cl	H
Me	Me	H	H	2	H	H	NMe	Cl	Me
Me	Me	H	H	2	H	H	NMe	Cl	Et
Me	Me	H	H	2	H	H	NMe	Cl	CF ₃
Me	Me	H	H	2	H	H	NMe	CF ₃	H
Me	Me	H	H	2	H	H	NMe	CF ₃	Me

	Me	Me	H	H	2	H	H	NMe	OCHF ₂	H
	Me	Me	H	H	2	H	H	NMe	OCHF ₂	Me
5	Me	Me	H	H	2	H	H	NMe	C(=O)Me	H
	Me	Me	H	H	2	H	H	NMe	C(=O)Me	Me
	Me	Me	H	H	2	H	H	NMe	-(CH ₂) ₃ -	
	Me	Me	H	H	2	H	H	NMe	-(CH ₂) ₄ -	
10	Me	Me	H	H	2	H	H	NEt	Cl	Me
	Me	Me	H	H	2	H	H	NEt	CF ₃	H
	Me	Me	H	H	2	H	H	NEt	CF ₃	Me
	Me	Me	H	H	2	H	H	NEt	OCHF ₂	H
15	Me	Me	H	H	2	H	H	NEt	OCHF ₂	Me
	Me	Me	H	H	2	H	H	NEt	-(CH ₂) ₃ -	
	Me	Me	H	H	2	H	H	NEt	-(CH ₂) ₄ -	
20	Me	Me	H	H	2	H	H	NPr-i	Cl	Me
	Me	Me	H	H	2	H	H	NPr-i	CF ₃	H
	Me	Me	H	H	2	H	H	NPr-i	CF ₃	Me
	Me	Me	H	H	2	H	H	NPr-i	OCHF ₂	H
25	Me	Me	H	H	2	H	H	NPr-i	OCHF ₂	Me
	Me	Me	H	H	2	H	H	NPr-i	-(CH ₂) ₃ -	
	Me	Me	H	H	2	H	H	NPr-i	-(CH ₂) ₄ -	
30	Me	Me	H	H	2	H	H	NPr	Cl	Me
	Me	Me	H	H	2	H	H	NPr	CF ₃	H
	Me	Me	H	H	2	H	H	NPr	CF ₃	Me
	Me	Me	H	H	2	H	H	NPr	OCHF ₂	H
35	Me	Me	H	H	2	H	H	NPr	OCHF ₂	Me
	Me	Me	H	H	2	H	H	NPr	-(CH ₂) ₃ -	
	Me	Me	H	H	2	H	H	NPr	-(CH ₂) ₄ -	
40	Me	Me	H	H	2	H	H	NBu-t	Cl	Me
	Me	Me	H	H	2	H	H	NBu-t	CF ₃	H
	Me	Me	H	H	2	H	H	NBu-t	CF ₃	Me
	Me	Me	H	H	2	H	H	NBu-t	OCHF ₂	H
45	Me	Me	H	H	2	H	H	NBu-t	OCHF ₂	Me
	Me	Me	H	H	2	H	H	NBu-t	-(CH ₂) ₃ -	
	Me	Me	H	H	2	H	H	NBu-t	-(CH ₂) ₄ -	
50	Me	Me	H	H	2	H	H	NCH ₂ Ph	Cl	Me
	Me	Me	H	H	2	H	H	NCH ₂ Ph	CF ₃	H
	Me	Me	H	H	2	H	H	NCH ₂ Ph	OCHF ₂	H
	Me	Me	H	H	2	H	H	NCH ₂ OMe	Cl	Me
55	Me	Me	H	H	2	H	H	NCH ₂ OMe	CF ₃	H
	Me	Me	H	H	2	H	H	NCH ₂ OMe	OCHF ₂	H

5	Me	Me	H	H	2	H	H	NCH ₂ C≡CH	Cl	Me
	Me	Me	H	H	2	H	H	NCH ₂ C≡CH	CF ₃	H
	Me	Me	H	H	2	H	H	NCH ₂ C≡CH	OCHF ₂	H
	Me	Me	H	H	2	H	H	NCH ₂ CH=CH ₂	Cl	Me
10	Me	Me	H	H	2	H	H	NCH ₂ CH=CH ₂	CF ₃	H
	Me	Me	H	H	2	H	H	NCH ₂ CH=CH ₂	OCHF ₂	H
	Me	Me	H	H	2	H	H	NCHF ₂	Cl	Me
	Me	Me	H	H	2	H	H	NCHF ₂	CF ₃	H
15	Me	Me	H	H	2	H	H	NCHF ₂	CF ₃	Me
	Me	Me	H	H	2	H	H	NCHF ₂	OCHF ₂	H
	Me	Me	H	H	2	H	H	NCHF ₂	OCHF ₂	Me
	Me	Me	H	H	2	H	H	NCHF ₂	C(=O)Me	H
20	Me	Me	H	H	2	H	H	NCHF ₂	C(=O)Me	Me
	Me	Me	H	H	2	H	H	NCHF ₂	-(CH ₂) ₃ -	
	Me	Me	H	H	2	H	H	NCHF ₂	-(CH ₂) ₄ -	
25	Me	Me	H	H	2	H	H	NPh	OMe	Me
	Me	Me	H	H	2	H	H	NPh	OEt	Me
	Me	Me	H	H	2	H	H	NPh	OCHF ₂	Me
	Me	Me	H	H	2	H	H	NPh	OCH ₂ CF ₃	Me
	Me	Me	H	H	2	H	H	NPh	CF ₃	H
30	Me	Me	H	H	2	H	H	NPh	OCH ₂ CH=CH ₂	Me
	Me	Me	H	H	2	H	H	NPh	OCH ₂ C≡CH	Me
	Me	Me	H	H	2	H	H	NPh	Cl	Me
	Me	Me	H	H	2	H	H	N(2-Cl)Ph	Cl	Me
35	Me	Me	H	H	2	H	H	N(2-F)Ph	Cl	Me
	Me	Me	H	H	2	H	H	N(2-OMe)Ph	Cl	Me
	Me	Me	H	H	2	H	H	N(2-Me)Ph	Cl	Me
	Me	Me	H	H	2	H	H	N(3-Cl)Ph	Cl	Me
	Me	Me	H	H	2	H	H	N(3-F)Ph	Cl	Me
40	Me	Me	H	H	2	H	H	N(3-OMe)Ph	Cl	Me
	Me	Me	H	H	2	H	H	N(3-Me)Ph	Cl	Me
	Me	Me	H	H	2	H	H	N(4-Cl)Ph	Cl	Me
	Me	Me	H	H	2	H	H	N(4-F)Ph	Cl	Me
	Me	Me	H	H	2	H	H	N(4-OMe)Ph	Cl	Me
45	Me	Me	H	H	2	H	H	N(4-Me)Ph	Cl	Me
	Me	Me	H	H	2	H	H	N(Thiophen-2-yl)	Cl	Me
	Me	Me	H	H	2	H	H	N(Thiophen-2-yl)	CF ₃	H
	Me	Me	H	H	2	H	H	N(Thiophen-2-yl)	OCHF ₂	H
50	Me	Me	H	H	2	H	H	NC(=O)Me	Cl	Me
	Me	Me	H	H	2	H	H	NC(=O)Me	CF ₃	H
	Me	Me	H	H	2	H	H	NC(=O)Me	OCHF ₂	H
	Me	Me	H	H	2	H	H	NC(=O)CF ₃	Cl	Me
55	Me	Me	H	H	2	H	H	NC(=O)CF ₃	CF ₃	H

	Me	Me	H	H	2	H	H	NC(=O)CF ₃	OCHF ₂	H
5	Me	Me	H	H	2	H	H	NC(=O)CH ₂ Ph	Cl	Me
	Me	Me	H	H	2	H	H	NC(=O)CH ₂ Ph	CF ₃	H
	Me	Me	H	H	2	H	H	NC(=O)CH ₂ Ph	OCHF ₂	H
	Me	Me	H	H	2	H	H	NC(=O)Ph	Cl	Me
10	Me	Me	H	H	2	H	H	NC(=O)Ph	CF ₃	H
	Me	Me	H	H	2	H	H	NC(=O)Ph	OCHF ₂	H
	Me	Me	H	H	2	H	H	NC(=O)OMe	Cl	Me
	Me	Me	H	H	2	H	H	NC(=O)OMe	CF ₃	H
15	Me	Me	H	H	2	H	H	NC(=O)OMe	OCHF ₂	H
	Me	Me	H	H	2	H	H	NC(=O)OCH ₂ Ph	Cl	Me
	Me	Me	H	H	2	H	H	NC(=O)OCH ₂ Ph	CF ₃	H
	Me	Me	H	H	2	H	H	NC(=O)OCH ₂ Ph	OCHF ₂	H
20	Me	Me	H	H	2	H	H	NC(=O)OPh	Cl	Me
	Me	Me	H	H	2	H	H	NC(=O)OPh	CF ₃	H
	Me	Me	H	H	2	H	H	NC(=O)OPh	OCHF ₂	H
	Me	Me	H	H	2	H	H	NC(=O)NHMe	Cl	Me
25	Me	Me	H	H	2	H	H	NC(=O)NHMe	CF ₃	H
	Me	Me	H	H	2	H	H	NC(=O)NHMe	OCHF ₂	H
	Me	Me	H	H	2	H	H	NC(=O)N(Me) ₂	Cl	Me
	Me	Me	H	H	2	H	H	NC(=O)N(Me) ₂	CF ₃	H
30	Me	Me	H	H	2	H	H	NC(=O)N(Me) ₂	OCHF ₂	H
	H	H	H	H	2	H	H	NPh	Cl	Me
	Me	H	H	H	2	H	H	NPh	Cl	Me
	Me	H	Me	H	2	H	H	NPh	Cl	Me
	Me	Me	H	H	2	Me	H	NPh	Cl	Me
35	Me	Me	H	H	2	Et	H	NPh	Cl	Me
	Me	Me	H	H	2	Pr-i	H	NPh	Cl	Me
	Me	Me	H	H	2	Me	Me	NPh	Cl	Me
	Me	Et	H	H	2	H	H	NPh	Cl	Me
40	Et	Et	H	H	2	H	H	NPh	Cl	Me
	Me	Pr-i	H	H	2	H	H	NPh	Cl	Me
	Me	Pr	H	H	2	H	H	NPh	Cl	Me
	Me	Pr-c	H	H	2	H	H	NPh	Cl	Me
	Me	CH ₂ Pr-c	H	H	2	H	H	NPh	Cl	Me
45		-(CH ₂) ₂ -	H	H	2	H	H	NPh	Cl	Me
		-(CH ₂) ₃ -	H	H	2	H	H	NPh	Cl	Me
		-(CH ₂) ₄ -	H	H	2	H	H	NPh	Cl	Me
		-(CH ₂) ₅ -	H	H	2	H	H	NPh	Cl	Me
50	H	-(CH ₂) ₃ -	H		2	H	H	NPh	Cl	Me
	H	-(CH ₂) ₄ -	H		2	H	H	NPh	Cl	Me
	H	-(CH ₂) ₅ -	H		2	H	H	NPh	Cl	Me
	H	-(CH ₂) ₆ -	H		2	H	H	NPh	Cl	Me
55	Me	Me	H	H	2	H	H	O	H	Me

5	Me	Me	H	H	2	H	H	O	Cl	Me
	Me	Me	H	H	2	H	H	S	H	Me
	Me	Me	H	H	2	H	H	S	Cl	Me
	Me	Me	H	H	1	H	H	NMe	Cl	H
	Me	Me	H	H	1	H	H	NMe	Cl	Me
	Me	Me	H	H	1	H	H	NMe	Cl	Et
10	Me	Me	H	H	1	H	H	NMe	Cl	CF ₃
	Me	Me	H	H	1	H	H	NMe	CF ₃	H
	Me	Me	H	H	1	H	H	NMe	CF ₃	Me
	Me	Me	H	H	1	H	H	NMe	OCHF ₂	H
15	Me	Me	H	H	1	H	H	NMe	OCHF ₂	Me
	Me	Me	H	H	1	H	H	NMe	C(=O)Me	H
	Me	Me	H	H	1	H	H	NMe	C(=O)Me	Me
	Me	Me	H	H	1	H	H	NMe	-(CH ₂) ₃ -	
20	Me	Me	H	H	1	H	H	NMe	-(CH ₂) ₄ -	
	Me	Me	H	H	1	H	H	NEt	Cl	Me
	Me	Me	H	H	1	H	H	NEt	CF ₃	H
	Me	Me	H	H	1	H	H	NEt	CF ₃	Me
25	Me	Me	H	H	1	H	H	NEt	OCHF ₂	H
	Me	Me	H	H	1	H	H	NEt	OCHF ₂	Me
	Me	Me	H	H	1	H	H	NEt	-(CH ₂) ₃ -	
	Me	Me	H	H	1	H	H	NEt	-(CH ₂) ₄ -	
30	Me	Me	H	H	1	H	H	NPr-i	Cl	Me
	Me	Me	H	H	1	H	H	NPr-i	CF ₃	H
	Me	Me	H	H	1	H	H	NPr-i	CF ₃	Me
35	Me	Me	H	H	1	H	H	NPr-i	OCHF ₂	H
	Me	Me	H	H	1	H	H	NPr-i	OCHF ₂	Me
	Me	Me	H	H	1	H	H	NPr-i	-(CH ₂) ₃ -	
	Me	Me	H	H	1	H	H	NPr-i	-(CH ₂) ₄ -	
40	Me	Me	H	H	1	H	H	NPr	Cl	Me
	Me	Me	H	H	1	H	H	NPr	CF ₃	H
	Me	Me	H	H	1	H	H	NPr	CF ₃	Me
	Me	Me	H	H	1	H	H	NPr	OCHF ₂	H
45	Me	Me	H	H	1	H	H	NPr	OCHF ₂	Me
	Me	Me	H	H	1	H	H	NPr	-(CH ₂) ₃ -	
	Me	Me	H	H	1	H	H	NPr	-(CH ₂) ₄ -	
50	Me	Me	H	H	1	H	H	NBu-t	Cl	Me
	Me	Me	H	H	1	H	H	NBu-t	CF ₃	H
	Me	Me	H	H	1	H	H	NBu-t	CF ₃	Me
	Me	Me	H	H	1	H	H	NBu-t	OCHF ₂	H
55	Me	Me	H	H	1	H	H	NBu-t	OCHF ₂	Me
	Me	Me	H	H	1	H	H	NBu-t	-(CH ₂) ₃ -	

	Me	Me	H	H	1	H	H	NBu-t	-(CH ₂) ₄ -	
5	Me	Me	H	H	1	H	H	NCH ₂ Ph	Cl	Me
	Me	Me	H	H	1	H	H	NCH ₂ Ph	CF ₃	H
	Me	Me	H	H	1	H	H	NCH ₂ Ph	OCHF ₂	H
	Me	Me	H	H	1	H	H	NCH ₂ OMe	Cl	Me
10	Me	Me	H	H	1	H	H	NCH ₂ OMe	CF ₃	H
	Me	Me	H	H	1	H	H	NCH ₂ OMe	OCHF ₂	H
	Me	Me	H	H	1	H	H	NCH ₂ C≡CH	Cl	Me
15	Me	Me	H	H	1	H	H	NCH ₂ C≡CH	CF ₃	H
	Me	Me	H	H	1	H	H	NCH ₂ C≡CH	OCHF ₂	H
	Me	Me	H	H	1	H	H	NCH ₂ CH=CH ₂	Cl	Me
	Me	Me	H	H	1	H	H	NCH ₂ CH=CH ₂	CF ₃	H
20	Me	Me	H	H	1	H	H	NCH ₂ CH=CH ₂	OCHF ₂	H
	Me	Me	H	H	1	H	H	NCHF ₂	Cl	Me
	Me	Me	H	H	1	H	H	NCHF ₂	CF ₃	H
	Me	Me	H	H	1	H	H	NCHF ₂	CF ₃	Me
25	Me	Me	H	H	1	H	H	NCHF ₂	OCHF ₂	H
	Me	Me	H	H	1	H	H	NCHF ₂	OCHF ₂	Me
	Me	Me	H	H	1	H	H	NCHF ₂	C(=O)Me	H
	Me	Me	H	H	1	H	H	NCHF ₂	C(=O)Me	Me
30	Me	Me	H	H	1	H	H	NCHF ₂	-(CH ₂) ₃ -	
	Me	Me	H	H	1	H	H	NCHF ₂	-(CH ₂) ₄ -	
	Me	Me	H	H	1	H	H	NPh	OMe	Me
	Me	Me	H	H	1	H	H	NPh	OEt	Me
35	Me	Me	H	H	1	H	H	NPh	OCHF ₂	Me
	Me	Me	H	H	1	H	H	NPh	OCH ₂ CF ₃	Me
	Me	Me	H	H	1	H	H	NPh	CF ₃	H
40	Me	Me	H	H	1	H	H	NPh	OCH ₂ CH=CH ₂	Me
	Me	Me	H	H	1	H	H	NPh	OCH ₂ C≡CH	Me
	Me	Me	H	H	1	H	H	NPh	Cl	Me
	Me	Me	H	H	1	H	H	N(2-Cl)Ph	Cl	Me
	Me	Me	H	H	1	H	H	N(2-F)Ph	Cl	Me
45	Me	Me	H	H	1	H	H	N(2-OMe)Ph	Cl	Me
	Me	Me	H	H	1	H	H	N(2-Me)Ph	Cl	Me
	Me	Me	H	H	1	H	H	N(3-Cl)Ph	Cl	Me
	Me	Me	H	H	1	H	H	N(3-F)Ph	Cl	Me
	Me	Me	H	H	1	H	H	N(3-OMe)Ph	Cl	Me
50	Me	Me	H	H	1	H	H	N(3-Me)Ph	Cl	Me
	Me	Me	H	H	1	H	H	N(4-Cl)Ph	Cl	Me
	Me	Me	H	H	1	H	H	N(4-F)Ph	Cl	Me
	Me	Me	H	H	1	H	H	N(4-OMe)Ph	Cl	Me
	Me	Me	H	H	1	H	H	N(4-Me)Ph	Cl	Me
55	Me	Me	H	H	1	H	H	N(Thiophen-2-yl)	Cl	Me

5	Me	Me	H	H	1	H	H	N(Thiophen-2-yl)	CF ₃	H
	Me	Me	H	H	1	H	H	N(Thiophen-2-yl)	OCHF ₂	H
	Me	Me	H	H	1	H	H	NC(=O)Me	Cl	Me
	Me	Me	H	H	1	H	H	NC(=O)Me	CF ₃	H
	Me	Me	H	H	1	H	H	NC(=O)Me	OCHF ₂	H
10	Me	Me	H	H	1	H	H	NC(=O)CF ₃	Cl	Me
	Me	Me	H	H	1	H	H	NC(=O)CF ₃	CF ₃	H
	Me	Me	H	H	1	H	H	NC(=O)CF ₃	OCHF ₂	H
15	Me	Me	H	H	1	H	H	NC(=O)CH ₂ Ph	Cl	Me
	Me	Me	H	H	1	H	H	NC(=O)CH ₂ Ph	CF ₃	H
	Me	Me	H	H	1	H	H	NC(=O)CH ₂ Ph	OCHF ₂	H
	Me	Me	H	H	1	H	H	NC(=O)Ph	Cl	Me
20	Me	Me	H	H	1	H	H	NC(=O)Ph	CF ₃	H
	Me	Me	H	H	1	H	H	NC(=O)Ph	OCHF ₂	H
	Me	Me	H	H	1	H	H	NC(=O)OMe	Cl	Me
	Me	Me	H	H	1	H	H	NC(=O)OMe	CF ₃	H
25	Me	Me	H	H	1	H	H	NC(=O)OMe	OCHF ₂	H
	Me	Me	H	H	1	H	H	NC(=O)OCH ₂ Ph	Cl	Me
	Me	Me	H	H	1	H	H	NC(=O)OCH ₂ Ph	CF ₃	H
	Me	Me	H	H	1	H	H	NC(=O)OCH ₂ Ph	OCHF ₂	H
30	Me	Me	H	H	1	H	H	NC(=O)OPh	Cl	Me
	Me	Me	H	H	1	H	H	NC(=O)OPh	CF ₃	H
	Me	Me	H	H	1	H	H	NC(=O)OPh	OCHF ₂	H
	Me	Me	H	H	1	H	H	NC(=O)NHMe	Cl	Me
35	Me	Me	H	H	1	H	H	NC(=O)NHMe	CF ₃	H
	Me	Me	H	H	1	H	H	NC(=O)NHMe	OCHF ₂	H
	Me	Me	H	H	1	H	H	NC(=O)N(Me) ₂	Cl	Me
	Me	Me	H	H	1	H	H	NC(=O)N(Me) ₂	CF ₃	H
40	Me	Me	H	H	1	H	H	NC(=O)N(Me) ₂	OCHF ₂	H
	H	H	H	H	1	H	H	NPh	Cl	Me
	Me	H	H	H	1	H	H	NPh	Cl	Me
	Me	H	Me	H	1	H	H	NPh	Cl	Me
45	Me	Me	H	H	1	Me	H	NPh	Cl	Me
	Me	Me	H	H	1	Et	H	NPh	Cl	Me
	Me	Me	H	H	1	Pr-i	H	NPh	Cl	Me
	Me	Me	H	H	1	Me	Me	NPh	Cl	Me
	Me	Et	H	H	1	H	H	NPh	Cl	Me
50	Et	Et	H	H	1	H	H	NPh	Cl	Me
	Me	Pr-i	H	H	1	H	H	NPh	Cl	Me
	Me	Pr	H	H	1	H	H	NPh	Cl	Me
	Me	Pr-c	H	H	1	H	H	NPh	Cl	Me
	Me	CH ₂ Pr-c	H	H	1	H	H	NPh	Cl	Me
55		-(CH ₂) ₂ -	H	H	1	H	H	NPh	Cl	Me


5

5	Me	Me	H	H	0	H	H	NPr	-(CH ₂) ₃ -	
	Me	Me	H	H	0	H	H	NPr	-(CH ₂) ₄ -	
	Me	Me	H	H	0	H	H	NBu-t	Cl	Me
10	Me	Me	H	H	0	H	H	NBu-t	CF ₃	H
	Me	Me	H	H	0	H	H	NBu-t	CF ₃	Me
	Me	Me	H	H	0	H	H	NBu-t	OCHF ₂	H
15	Me	Me	H	H	0	H	H	NBu-t	OCHF ₂	Me
	Me	Me	H	H	0	H	H	NBu-t	-(CH ₂) ₃ -	
	Me	Me	H	H	0	H	H	NBu-t	-(CH ₂) ₄ -	
20	Me	Me	H	H	0	H	H	NCH ₂ Ph	Cl	Me
	Me	Me	H	H	0	H	H	NCH ₂ Ph	CF ₃	H
	Me	Me	H	H	0	H	H	NCH ₂ Ph	OCHF ₂	H
25	Me	Me	H	H	0	H	H	NCH ₂ OMe	Cl	Me
	Me	Me	H	H	0	H	H	NCH ₂ OMe	CF ₃	H
	Me	Me	H	H	0	H	H	NCH ₂ OMe	OCHF ₂	H
30	Me	Me	H	H	0	H	H	NCH ₂ C≡CH	Cl	Me
	Me	Me	H	H	0	H	H	NCH ₂ C≡CH	CF ₃	H
	Me	Me	H	H	0	H	H	NCH ₂ C≡CH	OCHF ₂	H
35	Me	Me	H	H	0	H	H	NCH ₂ CH=CH ₂	Cl	Me
	Me	Me	H	H	0	H	H	NCH ₂ CH=CH ₂	CF ₃	H
	Me	Me	H	H	0	H	H	NCH ₂ CH=CH ₂	OCHF ₂	H
40	Me	Me	H	H	0	H	H	NCHF ₂	Cl	Me
	Me	Me	H	H	0	H	H	NCHF ₂	CF ₃	H
	Me	Me	H	H	0	H	H	NCHF ₂	CF ₃	Me
45	Me	Me	H	H	0	H	H	NCHF ₂	OCHF ₂	H
	Me	Me	H	H	0	H	H	NCHF ₂	OCHF ₂	Me
	Me	Me	H	H	0	H	H	NCHF ₂	C(=O)Me	H
50	Me	Me	H	H	0	H	H	NCHF ₂	C(=O)Me	Me
	Me	Me	H	H	0	H	H	NCHF ₂	-(CH ₂) ₃ -	
	Me	Me	H	H	0	H	H	NCHF ₂	-(CH ₂) ₄ -	
55	Me	Me	H	H	0	H	H	NPh	OMe	Me
	Me	Me	H	H	0	H	H	NPh	OEt	Me
	Me	Me	H	H	0	H	H	NPh	OCHF ₂	Me
55	Me	Me	H	H	0	H	H	NPh	OCH ₂ CF ₃	Me
	Me	Me	H	H	0	H	H	NPh	CF ₃	H
	Me	Me	H	H	0	H	H	NPh	OCH ₂ CH=CH ₂	Me
55	Me	Me	H	H	0	H	H	NPh	OCH ₂ C≡CH	Me
	Me	Me	H	H	0	H	H	NPh	Cl	Me
	Me	Me	H	H	0	H	H	N(2-Cl)Ph	Cl	Me
55	Me	Me	H	H	0	H	H	N(2-F)Ph	Cl	Me
	Me	Me	H	H	0	H	H	N(2-OMe)Ph	Cl	Me

5	Me	Me	H	H	0	H	H	N(2-Me)Ph	Cl	Me
	Me	Me	H	H	0	H	H	N(3-Cl)Ph	Cl	Me
	Me	Me	H	H	0	H	H	N(3-F)Ph	Cl	Me
	Me	Me	H	H	0	H	H	N(3-OMe)Ph	Cl	Me
	Me	Me	H	H	0	H	H	N(3-Me)Ph	Cl	Me
	Me	Me	H	H	0	H	H	N(4-Cl)Ph	Cl	Me
10	Me	Me	H	H	0	H	H	N(4-F)Ph	Cl	Me
	Me	Me	H	H	0	H	H	N(4-OMe)Ph	Cl	Me
	Me	Me	H	H	0	H	H	N(4-Me)Ph	Cl	Me
	Me	Me	H	H	0	H	H	N(Thiophen-2-yl)	Cl	Me
	Me	Me	H	H	0	H	H	N(Thiophen-2-yl)	CF ₃	H
15	Me	Me	H	H	0	H	H	N(Thiophen-2-yl)	OCHF ₂	H
	Me	Me	H	H	0	H	H	NC(=O)Me	Cl	Me
	Me	Me	H	H	0	H	H	NC(=O)Me	CF ₃	H
	Me	Me	H	H	0	H	H	NC(=O)Me	OCHF ₂	H
20	Me	Me	H	H	0	H	H	NC(=O)CF ₃	Cl	Me
	Me	Me	H	H	0	H	H	NC(=O)CF ₃	CF ₃	H
	Me	Me	H	H	0	H	H	NC(=O)CF ₃	OCHF ₂	H
25	Me	Me	H	H	0	H	H	NC(=O)CH ₂ Ph	Cl	Me
	Me	Me	H	H	0	H	H	NC(=O)CH ₂ Ph	CF ₃	H
	Me	Me	H	H	0	H	H	NC(=O)CH ₂ Ph	OCHF ₂	H
	Me	Me	H	H	0	H	H	NC(=O)Ph	Cl	Me
30	Me	Me	H	H	0	H	H	NC(=O)Ph	CF ₃	H
	Me	Me	H	H	0	H	H	NC(=O)Ph	OCHF ₂	H
	Me	Me	H	H	0	H	H	NC(=O)OMe	Cl	Me
	Me	Me	H	H	0	H	H	NC(=O)OMe	CF ₃	H
35	Me	Me	H	H	0	H	H	NC(=O)OMe	OCHF ₂	H
	Me	Me	H	H	0	H	H	NC(=O)OCH ₂ Ph	Cl	Me
	Me	Me	H	H	0	H	H	NC(=O)OCH ₂ Ph	CF ₃	H
	Me	Me	H	H	0	H	H	NC(=O)OCH ₂ Ph	OCHF ₂	H
40	Me	Me	H	H	0	H	H	NC(=O)OPh	Cl	Me
	Me	Me	H	H	0	H	H	NC(=O)OPh	CF ₃	H
	Me	Me	H	H	0	H	H	NC(=O)OPh	OCHF ₂	H
	Me	Me	H	H	0	H	H	NC(=O)NHMe	Cl	Me
45	Me	Me	H	H	0	H	H	NC(=O)NHMe	CF ₃	H
	Me	Me	H	H	0	H	H	NC(=O)NHMe	OCHF ₂	H
	Me	Me	H	H	0	H	H	NC(=O)N(Me) ₂	Cl	Me
	Me	Me	H	H	0	H	H	NC(=O)N(Me) ₂	CF ₃	H
50	Me	Me	H	H	0	H	H	NC(=O)N(Me) ₂	OCHF ₂	H
	H	H	H	H	0	H	H	NPh	Cl	Me
	Me	H	H	H	0	H	H	NPh	Cl	Me
	Me	H	Me	H	0	H	H	NPh	Cl	Me
55	Me	Me	H	H	0	Me	H	NPh	Cl	Me
	Me	Me	H	H	0	Et	H	NPh	Cl	Me

Me	Me	H	H	0	Pr-i	H	NPh	Cl	Me
Me	Me	H	H	0	Me	Me	NPh	Cl	Me
Me	Et	H	H	0	H	H	NPh	Cl	Me
Et	Et	H	H	0	H	H	NPh	Cl	Me
Me	Pr-i	H	H	0	H	H	NPh	Cl	Me
Me	Pr	H	H	0	H	H	NPh	Cl	Me
Me	Pr-c	H	H	0	H	H	NPh	Cl	Me
Me	CH ₂ Pr-c	H	H	0	H	H	NPh	Cl	Me
-(CH ₂) ₂ -		H	H	0	H	H	NPh	Cl	Me
-(CH ₂) ₃ -		H	H	0	H	H	NPh	Cl	Me
-(CH ₂) ₄ -		H	H	0	H	H	NPh	Cl	Me
-(CH ₂) ₅ -		H	H	0	H	H	NPh	Cl	Me
H	-(CH ₂) ₃ -		H	0	H	H	NPh	Cl	Me
H	-(CH ₂) ₄ -		H	0	H	H	NPh	Cl	Me
H	-(CH ₂) ₅ -		H	0	H	H	NPh	Cl	Me
H	-(CH ₂) ₆ -		H	0	H	H	NPh	Cl	Me
Me	Me	H	H	0	H	H	O	H	Me
Me	Me	H	H	0	H	H	O	Cl	Me
Me	Me	H	H	0	H	H	S	H	Me
Me	Me	H	H	0	H	H	S	Cl	Me
Me	Et	H	H	2	H	H	NH	H	H

Table 6


									
R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Z ⁵	R ³⁵	R ³⁶
Me	Me	H	H	2	H	H	NMe	H	OMe
Me	Me	H	H	2	H	H	NMe	H	OEt
Me	Me	H	H	2	H	H	NMe	H	OCHF ₂
Me	Me	H	H	2	H	H	NMe	H	OCH ₂ CF ₃
Me	Me	H	H	2	H	H	NMe		-(CH ₂) ₃ -
Me	Me	H	H	2	H	H	NMe		-(CH ₂) ₄ -
Me	Me	H	H	2	H	H	NEt		-(CH ₂) ₃ -
Me	Me	H	H	2	H	H	NEt		-(CH ₂) ₄ -
Me	Me	H	H	2	H	H	NPr-i		-(CH ₂) ₃ -
Me	Me	H	H	2	H	H	NPr-i		-(CH ₂) ₄ -
Me	Me	H	H	2	H	H	NCHF ₂		-(CH ₂) ₃ -
Me	Me	H	H	2	H	H	NCHF ₂		-(CH ₂) ₄ -
Me	Me	H	H	2	H	H	N(CH ₂) ₃ O-		H
Me	Me	H	H	2	H	H	N(CH ₂) ₄ O-		H

	Me	Me	H	H	2	H	H	N(CH ₂) ₄ -	H
	Me	Me	H	H	2	H	H	N(CH ₂) ₅ -	H
5	Me	Me	H	H	2	H	H	NPh	OMe
	Me	Me	H	H	2	H	H	NPh	OEt
	Me	Me	H	H	2	H	H	NPh	OCHF ₂
10	Me	Me	H	H	2	H	H	NPh	OCH ₂ CF ₃
	Me	Me	H	H	2	H	H	O	Me
	Me	Me	H	H	2	H	H	S	Me
	H	H	H	H	2	H	H	NPh	H
	Me	H	H	H	2	H	H	NPh	OMe
15	Me	H	Me	H	2	H	H	NPh	OEt
	Me	Me	H	H	2	Me	H	NPh	OMe
	Me	Me	H	H	2	Et	H	NPh	OEt
	Me	Me	H	H	2	Pr-i	H	NPh	OMe
20	Me	Me	H	H	2	Me	Me	NPh	OEt
	Me	Et	H	H	2	H	H	NPh	OMe
	Et	Et	H	H	2	H	H	NPh	OEt
	Me	Pr-i	H	H	2	H	H	NPh	OMe
25	Me	Pr	H	H	2	H	H	NPh	OEt
	Me	Pr-c	H	H	2	H	H	NPh	OMe
	Me	CH ₂ Pr-c	H	H	2	H	H	NPh	OMe
		-(CH ₂) ₂ -	H	H	2	H	H	NPh	OEt
30		-(CH ₂) ₃ -	H	H	2	H	H	NPh	OMe
		-(CH ₂) ₄ -	H	H	2	H	H	NPh	OEt
		-(CH ₂) ₅ -	H	H	2	H	H	NPh	OMe
35	H	-(CH ₂) ₃ -	H	2	H	H	H	NPh	OEt
	H	-(CH ₂) ₄ -	H	2	H	H	H	NPh	OMe
	H	-(CH ₂) ₅ -	H	2	H	H	H	NPh	OMe
	H	-(CH ₂) ₆ -	H	2	H	H	H	NPh	OEt
40	Me	Me	H	H	1	H	H	NMe	OMe
	Me	Me	H	H	1	H	H	NMe	OEt
	Me	Me	H	H	1	H	H	NMe	OCHF ₂
	Me	Me	H	H	1	H	H	NMe	OCH ₂ CF ₃
45	Me	Me	H	H	1	H	H	NMe	-(CH ₂) ₃ -
	Me	Me	H	H	1	H	H	NMe	-(CH ₂) ₄ -
	Me	Me	H	H	1	H	H	NEt	-(CH ₂) ₃ -
	Me	Me	H	H	1	H	H	NEt	-(CH ₂) ₄ -
50	Me	Me	H	H	1	H	H	NPr-i	-(CH ₂) ₃ -
	Me	Me	H	H	1	H	H	NPr-i	-(CH ₂) ₄ -
	Me	Me	H	H	1	H	H	NCHF ₂	-(CH ₂) ₃ -
	Me	Me	H	H	1	H	H	NCHF ₂	-(CH ₂) ₄ -
55	Me	Me	H	H	1	H	H	N(CH ₂) ₃ O-	H
	Me	Me	H	H	1	H	H	N(CH ₂) ₄ O-	H

	Me	Me	H	H	1	H	H	N(CH ₂) ₄ -	H
5	Me	Me	H	H	1	H	H	N(CH ₂) ₅ -	H
	Me	Me	H	H	1	H	H	NPh	OMe
	Me	Me	H	H	1	H	H	NPh	OEt
	Me	Me	H	H	1	H	H	NPh	OCHF ₂
10	Me	Me	H	H	1	H	H	NPh	OCH ₂ CF ₃
	Me	Me	H	H	1	H	H	O	Me
	Me	Me	H	H	1	H	H	S	Me
	H	H	H	H	1	H	H	NPh	H
15	Me	H	H	H	1	H	H	NPh	OMe
	Me	H	Me	H	1	H	H	NPh	OEt
	Me	Me	H	H	1	Me	H	NPh	OMe
	Me	Me	H	H	1	Et	H	NPh	OEt
	Me	Me	H	H	1	Pr-i	H	NPh	OMe
20	Me	Me	H	H	1	Me	Me	NPh	OEt
	Me	Et	H	H	1	H	H	NPh	OMe
	Et	Et	H	H	1	H	H	NPh	OEt
	Me	Pr-i	H	H	1	H	H	NPh	OMe
25	Me	Pr	H	H	1	H	H	NPh	OEt
	Me	Pr-c	H	H	1	H	H	NPh	OMe
	Me	CH ₂ Pr-c	H	H	1	H	H	NPh	OEt
	-(CH ₂) ₂ -		H	H	1	H	H	NPh	OMe
30	-(CH ₂) ₃ -		H	H	1	H	H	NPh	OEt
	-(CH ₂) ₄ -		H	H	1	H	H	NPh	OMe
	-(CH ₂) ₅ -		H	H	1	H	H	NPh	OEt
	H	-(CH ₂) ₃ -	H	H	1	H	H	NPh	OMe
35	H	-(CH ₂) ₄ -	H	H	1	H	H	NPh	OEt
	H	-(CH ₂) ₅ -	H	H	1	H	H	NPh	OMe
	H	-(CH ₂) ₆ -	H	H	1	H	H	NPh	OMe
40	Me	Me	H	H	0	H	H	NMe	OEt
	Me	Me	H	H	0	H	H	NMe	OCHF ₂
	Me	Me	H	H	0	H	H	NMe	OCH ₂ CF ₃
	Me	Me	H	H	0	H	H	NMe	-(CH ₂) ₃ -
45	Me	Me	H	H	0	H	H	NMe	-(CH ₂) ₄ -
	Me	Me	H	H	0	H	H	NEt	-(CH ₂) ₃ -
	Me	Me	H	H	0	H	H	NEt	-(CH ₂) ₄ -
	Me	Me	H	H	0	H	H	NPr-i	-(CH ₂) ₃ -
50	Me	Me	H	H	0	H	H	NPr-i	-(CH ₂) ₄ -
	Me	Me	H	H	0	H	H	NCHF ₂	-(CH ₂) ₃ -
	Me	Me	H	H	0	H	H	NCHF ₂	-(CH ₂) ₄ -
55	Me	Me	H	H	0	H	H	N(CH ₂) ₃ O-	H
	Me	Me	H	H	0	H	H	N(CH ₂) ₄ O-	H

5	Me	Me	H	H	0	H	H	N(CH ₂) ₄ -	H
	Me	Me	H	H	0	H	H	N(CH ₂) ₅ -	H
	Me	Me	H	H	0	H	H	NPh	OMe
	Me	Me	H	H	0	H	H	NPh	OEt
10	Me	Me	H	H	0	H	H	NPh	OCHF ₂
	Me	Me	H	H	0	H	H	NPh	OCH ₂ CF ₃
	Me	Me	H	H	0	H	H	O	Me
	Me	Me	H	H	0	H	H	S	Me
15	H	H	H	H	0	H	H	NPh	OMe
	Me	H	H	H	0	H	H	NPh	OEt
	Me	H	Me	H	0	H	H	NPh	OMe
	Me	Me	H	H	0	Me	H	NPh	OEt
20	Me	Me	H	H	0	Et	H	NPh	OMe
	Me	Me	H	H	0	Pr-i	H	NPh	OEt
	Me	Me	H	H	0	Me	Me	NPh	OMe
	Me	Et	H	H	0	H	H	NPh	OEt
25	Et	Et	H	H	0	H	H	NPh	OMe
	Me	Pr-i	H	H	0	H	H	NPh	OEt
	Me	Pr	H	H	0	H	H	NPh	OMe
	Me	Pr-c	H	H	0	H	H	NPh	OEt
30	Me	CH ₂ Pr-c	H	H	0	H	H	NPh	OMe
	-(CH ₂) ₂ -		H	H	0	H	H	NPh	OEt
	-(CH ₂) ₃ -		H	H	0	H	H	NPh	OMe
	-(CH ₂) ₄ -		H	H	0	H	H	NPh	OEt
35	-(CH ₂) ₅ -		H	H	0	H	H	NPh	OMe
	H	-(CH ₂) ₃ -	H	0	H	H	H	NPh	OEt
	H	-(CH ₂) ₄ -	H	0	H	H	H	NPh	OMe
40	H	-(CH ₂) ₅ -	H	0	H	H	H	NPh	OMe
	H	-(CH ₂) ₆ -	H	0	H	H	H	NPh	OEt
	Me	Et	H	H	2	H	H	O	H
	Me	Et	H	H	2	H	H	S	H
45	Me	Et	H	H	2	H	H	NH	H

Table 7

											
R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	R ³⁷	R ³⁸	R ³⁹	R ⁴⁰	
Me	Me	H	H	2	H	H	H	H	H	H	-

	Me	Me	H	H	2	H	H	H	H	H	N-oxide
	Me	Me	H	H	2	H	H	Cl	Ph	H	-
5	Me	Me	H	H	2	H	H	OMe	Ph	H	-
	Me	Me	H	H	2	H	H	Cl	Me	H	-
	Me	Me	H	H	2	H	H	OMe	Me	H	-
	Me	Me	H	H	2	H	H	H	CF ₃	H	-
	Me	Me	H	H	2	H	H	H	CF ₃	H	N-oxide
10	Me	Me	H	H	2	H	H	Cl	CF ₃	H	-
	Me	Me	H	H	2	H	H	CN	CF ₃	H	-
	Me	Me	H	H	2	H	H	OMe	CF ₃	H	-
15	Me	Me	H	H	2	H	H	OEt	CF ₃	H	-
	Me	Me	H	H	2	H	H	Me	Me	H	N-oxide
	Me	Me	H	H	2	H	H	Ph	Ph	H	-
	Me	Me	H	H	2	H	H	Cl	(4-Cl)Ph	H	Me
20	Me	Me	H	H	2	H	H	Cl	(4-Cl)Ph	H	-
	Me	Me	H	H	2	H	H	OMe	Cl	H	-
	Me	Me	H	H	2	H	H	Cl	(CH ₂) ₃	H	-
	Me	Me	H	H	2	H	H	Me	(CH ₂) ₃	H	-
	Me	Me	H	H	2	H	H	Cl	(CH ₂) ₄	H	-
25	Me	Me	H	H	2	H	H	Me	(CH ₂) ₄	H	-
	Me	Me	H	H	2	H	H	Cl	H	(CH ₂) ₃	-
	Me	Me	H	H	2	H	H	Me	H	(CH ₂) ₃	-
	Me	Me	H	H	2	H	H	Cl	H	(CH ₂) ₄	-
30	Me	Me	H	H	2	H	H	Me	H	(CH ₂) ₄	-
	H	H	H	H	2	H	H	H	CF ₃	H	H
	Me	H	H	H	2	H	H	H	CF ₃	H	H
35	Me	H	Me	H	2	H	H	H	CF ₃	H	H
	Me	Me	Me	H	2	H	H	H	CF ₃	H	H
	Me	Me	H	H	2	Me	H	H	CF ₃	H	H
	Me	Me	H	H	2	Et	H	H	CF ₃	H	H
40	Me	Me	H	H	2	Pr-i	H	H	CF ₃	H	H
	Me	Me	H	H	2	Me	Me	H	CF ₃	H	H
	Me	Et	H	H	2	H	H	H	CF ₃	H	H
	Et	Et	H	H	2	H	H	H	CF ₃	H	H
45	Me	Pr-i	H	H	2	H	H	H	CF ₃	H	H
	Me	Pr	H	H	2	H	H	H	CF ₃	H	H
	Me	Pr-c	H	H	2	H	H	H	CF ₃	H	H
50	Me	CH ₂ Pr-c	H	H	2	H	H	H	CF ₃	H	H
		-(CH ₂) ₂ -	H	H	2	H	H	H	CF ₃	H	H
		-(CH ₂) ₃ -	H	H	2	H	H	H	CF ₃	H	H
55		-(CH ₂) ₄ -	H	H	2	H	H	H	CF ₃	H	H

		-(CH ₂) ₅ -	H	H	2	H	H	H	CF ₃	H	H	-
	H	-(CH ₂) ₃ -		H	2	H	H	H	CF ₃	H	H	-
5	H	-(CH ₂) ₄ -		H	2	H	H	H	CF ₃	H	H	-
	H	-(CH ₂) ₅ -		H	2	H	H	H	CF ₃	H	H	-
	H	-(CH ₂) ₆ -		H	2	H	H	H	CF ₃	H	H	-
10	Me	Me	H	H	1	H	H	H	H	H	H	-
	Me	Me	H	H	1	H	H	H	H	H	H	N-oxide
	Me	Me	H	H	1	H	H	Cl	Ph	H	H	-
	Me	Me	H	H	1	H	H	OMe	Ph	H	H	-
	Me	Me	H	H	1	H	H	Cl	Me	H	H	-
15	Me	Me	H	H	1	H	H	OMe	Me	H	H	-
	Me	Me	H	H	1	H	H	H	CF ₃	H	H	-
	Me	Me	H	H	1	H	H	Cl	CF ₃	H	H	-
	Me	Me	H	H	1	H	H	CN	CF ₃	H	H	-
20	Me	Me	H	H	1	H	H	OMe	CF ₃	H	H	-
	Me	Me	H	H	1	H	H	OEt	CF ₃	H	H	-
	Me	Me	H	H	1	H	H	Me	Me	H	H	N-oxide
	Me	Me	H	H	1	H	H	Ph	Ph	H	H	-
25	Me	Me	H	H	1	H	H	Cl	(4-Cl)Ph	H	Me	-
	Me	Me	H	H	1	H	H	Cl	(4-Cl)Ph	H	H	-
	Me	Me	H	H	1	H	H	OMe	Cl	H	H	-
	Me	Me	H	H	1	H	H	Cl	(CH ₂) ₃		H	-
30	Me	Me	H	H	1	H	H	Me	(CH ₂) ₃		H	-
	Me	Me	H	H	1	H	H	Cl	(CH ₂) ₄		H	-
	Me	Me	H	H	1	H	H	Me	(CH ₂) ₄		H	-
	Me	Me	H	H	1	H	H	Cl	H	(CH ₂) ₃		-
35	Me	Me	H	H	1	H	H	Me	H	(CH ₂) ₃		-
	Me	Me	H	H	1	H	H	Cl	H	(CH ₂) ₄		-
	Me	Me	H	H	1	H	H	Me	H	(CH ₂) ₄		-
	H	H	H	H	1	H	H	H	CF ₃	H	H	-
40	Me	H	H	H	1	H	H	H	CF ₃	H	H	-
	Me	H	Me	H	1	H	H	H	CF ₃	H	H	-
	Me	Me	Me	H	1	H	H	H	CF ₃	H	H	-
45	Me	Me	H	H	1	Me	H	H	CF ₃	H	H	-
	Me	Me	H	H	1	Et	H	H	CF ₃	H	H	-
	Me	Me	H	H	1	Pr-i	H	H	CF ₃	H	H	-
	Me	Me	H	H	1	Me	Me	H	CF ₃	H	H	-
50	Me	Et	H	H	1	H	H	H	CF ₃	H	H	-
	Et	Et	H	H	1	H	H	H	CF ₃	H	H	-
	Me	Pr-i	H	H	1	H	H	H	CF ₃	H	H	-
	Me	Pr	H	H	1	H	H	H	CF ₃	H	H	-
55	Me	Pr-c	H	H	1	H	H	H	CF ₃	H	H	-

	Me	CH ₂ Pr-c	H	H	1	H	H	H	CF ₃	H	H	-
		-(CH ₂) ₂ -	H	H	1	H	H	H	CF ₃	H	H	-
5		-(CH ₂) ₃ -	H	H	1	H	H	H	CF ₃	H	H	-
		-(CH ₂) ₄ -	H	H	1	H	H	H	CF ₃	H	H	-
		-(CH ₂) ₅ -	H	H	1	H	H	H	CF ₃	H	H	-
10	H	-(CH ₂) ₃ -	H	1	H	H	H	H	CF ₃	H	H	-
	H	-(CH ₂) ₄ -	H	1	H	H	H	H	CF ₃	H	H	-
	H	-(CH ₂) ₅ -	H	1	H	H	H	H	CF ₃	H	H	-
	H	-(CH ₂) ₆ -	H	1	H	H	H	H	CF ₃	H	H	-
15	Me	Me	H	H	0	H	H	H	H	H	H	-
	Me	Me	H	H	0	H	H	H	H	H	H	N-oxide
	Me	Me	H	H	0	H	H	H	Cl	Ph	H	-
	Me	Me	H	H	0	H	H	H	OMe	Ph	H	-
	Me	Me	H	H	0	H	H	H	Cl	Me	H	-
20	Me	Me	H	H	0	H	H	H	OMe	Me	H	-
	Me	Me	H	H	0	H	H	H	H	CF ₃	H	-
	Me	Me	H	H	0	H	H	H	Cl	CF ₃	H	-
	Me	Me	H	H	0	H	H	H	CN	CF ₃	H	-
25	Me	Me	H	H	0	H	H	H	OMe	CF ₃	H	-
	Me	Me	H	H	0	H	H	H	OEt	CF ₃	H	-
	Me	Me	H	H	0	H	H	H	Me	Me	H	N-oxide
	Me	Me	H	H	0	H	H	H	Ph	Ph	H	-
30	Me	Me	H	H	0	H	H	H	Cl	(4-Cl)Ph	H	Me
	Me	Me	H	H	0	H	H	H	Cl	(4-Cl)Ph	H	-
	Me	Me	H	H	0	H	H	H	OMe	Cl	H	-
	Me	Me	H	H	0	H	H	H	Cl	(CH ₂) ₃	H	-
35	Me	Me	H	H	0	H	H	H	Me	(CH ₂) ₃	H	-
	Me	Me	H	H	0	H	H	H	Cl	(CH ₂) ₄	H	-
	Me	Me	H	H	0	H	H	H	Me	(CH ₂) ₄	H	-
	Me	Me	H	H	0	H	H	H	Cl	H	(CH ₂) ₃	-
40	Me	Me	H	H	0	H	H	H	Me	H	(CH ₂) ₃	-
	Me	Me	H	H	0	H	H	H	Cl	H	(CH ₂) ₄	-
	Me	Me	H	H	0	H	H	H	Me	H	(CH ₂) ₄	-
45	Me	Me	H	H	0	H	H	(2-Chloropyridin-3-yl)methylthio	H	H	H	-
	H	H	H	H	0	H	H	H	CF ₃	H	H	-
	Me	H	H	H	0	H	H	H	CF ₃	H	H	-
50	Me	Me	Me	H	0	H	H	H	CF ₃	H	H	-
	Me	H	Me	H	0	H	H	H	CF ₃	H	H	-
	Me	Me	H	H	0	Me	H	H	CF ₃	H	H	-
	Me	Me	H	H	0	Et	H	H	CF ₃	H	H	-
55	Me	Me	H	H	0	Pr-i	H	H	CF ₃	H	H	-

5

10

15

20

25

Me	Me	H	H	0	Me	Me	H	CF ₃	H	H	-
Me	Et	H	H	0	H	H	H	CF ₃	H	H	-
Et	Et	H	H	0	H	H	H	CF ₃	H	H	-
Me	Pr-i	H	H	0	H	H	H	CF ₃	H	H	-
Me	Pr	H	H	0	H	H	H	CF ₃	H	H	-
Me	Pr-c	H	H	0	H	H	H	CF ₃	H	H	-
Me	CH ₂ Pr-c	H	H	0	H	H	H	CF ₃	H	H	-
	-(CH ₂) ₂ -	H	H	0	H	H	H	CF ₃	H	H	-
	-(CH ₂) ₃ -	H	H	0	H	H	H	CF ₃	H	H	-
	-(CH ₂) ₄ -	H	H	0	H	H	H	CF ₃	H	H	-
	-(CH ₂) ₅ -	H	H	0	H	H	H	CF ₃	H	H	-
H	-(CH ₂) ₃ -	H	0	H	H	H	H	CF ₃	H	H	-
H	-(CH ₂) ₄ -	H	0	H	H	H	H	CF ₃	H	H	-
H	-(CH ₂) ₅ -	H	0	H	H	H	H	CF ₃	H	H	-
H	-(CH ₂) ₆ -	H	0	H	H	H	H	CF ₃	H	H	-
Me	Et	H	H	2	H	H	H	H	H	H	-

30

35


40

45

50

55

Table 8

									
R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	R ⁴¹	R ⁴²	R ⁴³
Me	Me	H	H	2	H	H	H	Cl	Cl
Me	Me	H	H	2	H	H	H	OH	Cl
Me	Me	H	H	2	H	H	H	OMe	Cl
Me	Me	H	H	2	H	H	H	OEt	Cl
Me	Me	H	H	2	H	H	H	OPr-i	Cl
Me	Me	H	H	2	H	H	H	OPr	Cl
Me	Me	H	H	2	H	H	H	OBu-t	Cl
Me	Me	H	H	2	H	H	H	OCH ₂ Pr-c	Cl
Me	Me	H	H	2	H	H	H	OCH ₂ Bu-c	Cl
Me	Me	H	H	2	H	H	H	OCH ₂ Pen-c	Cl
Me	Me	H	H	2	H	H	H	OCH ₂ Hex-c	Cl
Me	Me	H	H	2	H	H	H	OPen-c	Cl
Me	Me	H	H	2	H	H	H	OHex-c	Cl
Me	Me	H	H	2	H	H	H	OCH ₂ Ph	Cl
Me	Me	H	H	2	H	H	H	OPh	Cl
Me	Me	H	H	2	H	H	H	OCHF ₂	Cl
Me	Me	H	H	2	H	H	H	SH	Cl
Me	Me	H	H	2	H	H	H	SMe	Cl

	Me	Me	H	H	2	H	H	H	SO ₂ Me	Cl
	Me	Me	H	H	2	H	H	H	SEt	Cl
5	Me	Me	H	H	2	H	H	H	SO ₂ Et	Cl
	Me	Me	H	H	2	H	H	H	SPr-i	Cl
	Me	Me	H	H	2	H	H	H	SO ₂ Pr-i	Cl
	Me	Me	H	H	2	H	H	H	SPh	Cl
10	Me	Me	H	H	2	H	H	H	SO ₂ Ph	Cl
	Me	Me	H	H	2	H	H	H	SCHF ₂	Cl
	Me	Me	H	H	2	H	H	H	SO ₂ CHF ₂	Cl
	Me	Me	H	H	2	H	H	H	NH ₂	Cl
	Me	Me	H	H	2	H	H	H	NHMe	Cl
15	Me	Me	H	H	2	H	H	H	NMe ₂	Cl
	Me	Me	H	H	2	H	H	H	NHEt	Cl
	Me	Me	H	H	2	H	H	H	NEt ₂	Cl
	Me	Me	H	H	2	H	H	H	NHPh	Cl
20	Me	Me	H	H	2	H	H	H	N(Me)Ph	Cl
	Me	Me	H	H	2	H	H	H	CN	Cl
	Me	Me	H	H	2	H	H	H	F	Me
	Me	Me	H	H	2	H	H	H	Cl	Me
	Me	Me	H	H	2	H	H	H	OH	Me
25	Me	Me	H	H	2	H	H	H	OMe	Me
	Me	Me	H	H	2	H	H	H	OEt	Me
	Me	Me	H	H	2	H	H	H	OPr-i	Me
	Me	Me	H	H	2	H	H	H	OPr	Me
30	Me	Me	H	H	2	H	H	H	OBu-t	Me
	Me	Me	H	H	2	H	H	H	OCH ₂ Pr-c	Me
	Me	Me	H	H	2	H	H	H	OCH ₂ Bu-c	Me
	Me	Me	H	H	2	H	H	H	OCH ₂ Pen-c	Me
	Me	Me	H	H	2	H	H	H	OCH ₂ Hex-c	Me
35	Me	Me	H	H	2	H	H	H	OPen-c	Me
	Me	Me	H	H	2	H	H	H	OHex-c	Me
	Me	Me	H	H	2	H	H	H	OCH ₂ Ph	Me
	Me	Me	H	H	2	H	H	H	OPh	Me
40	Me	Me	H	H	2	H	H	H	OCHF ₂	Me
	Me	Me	H	H	2	H	H	H	SH	Me
	Me	Me	H	H	2	H	H	H	SMe	Me
	Me	Me	H	H	2	H	H	H	SO ₂ Me	Me
45	Me	Me	H	H	2	H	H	H	SEt	Me
	Me	Me	H	H	2	H	H	H	SO ₂ Et	Me
	Me	Me	H	H	2	H	H	H	SPr-i	Me
	Me	Me	H	H	2	H	H	H	SO ₂ Pr-i	Me
	Me	Me	H	H	2	H	H	H	SPh	Me
50	Me	Me	H	H	2	H	H	H	SO ₂ Ph	Me
	Me	Me	H	H	2	H	H	H	SCHF ₂	Me
	Me	Me	H	H	2	H	H	H	SO ₂ CHF ₂	Me
	Me	Me	H	H	2	H	H	H	NH ₂	Me
55	Me	Me	H	H	2	H	H	H	NHMe	Me
	Me	Me	H	H	2	H	H	H	NMe ₂	Me

	Me	Me	H	H	2	H	H	H	NHEt	Me
	Me	Me	H	H	2	H	H	H	NEt ₂	Me
5	Me	Me	H	H	2	H	H	H	NHPh	Me
	Me	Me	H	H	2	H	H	H	N(Me)Ph	Me
	Me	Me	H	H	2	H	H	H	CN	Me
	Me	Me	H	H	2	H	H	H	F	Pr-i
	Me	Me	H	H	2	H	H	H	Cl	Pr-i
10	Me	Me	H	H	2	H	H	H	OH	Pr-i
	Me	Me	H	H	2	H	H	H	OMe	Pr-i
	Me	Me	H	H	2	H	H	H	OEt	Pr-i
	Me	Me	H	H	2	H	H	H	OPr-i	Pr-i
15	Me	Me	H	H	2	H	H	H	OPr	Pr-i
	Me	Me	H	H	2	H	H	H	OBu-t	Pr-i
	Me	Me	H	H	2	H	H	H	OCH ₂ Pr-c	Pr-i
	Me	Me	H	H	2	H	H	H	OCH ₂ Bu-c	Pr-i
20	Me	Me	H	H	2	H	H	H	OCH ₂ Pen-c	Pr-i
	Me	Me	H	H	2	H	H	H	OCH ₂ Hex-c	Pr-i
	Me	Me	H	H	2	H	H	H	OPen-c	Pr-i
	Me	Me	H	H	2	H	H	H	OHex-c	Pr-i
	Me	Me	H	H	2	H	H	H	OCH ₂ Ph	Pr-i
25	Me	Me	H	H	2	H	H	H	OPh	Pr-i
	Me	Me	H	H	2	H	H	H	OCHF ₂	Pr-i
	Me	Me	H	H	2	H	H	H	SH	Pr-i
	Me	Me	H	H	2	H	H	H	SMe	Pr-i
30	Me	Me	H	H	2	H	H	H	SO ₂ Me	Pr-i
	Me	Me	H	H	2	H	H	H	SEt	Pr-i
	Me	Me	H	H	2	H	H	H	SO ₂ Et	Pr-i
	Me	Me	H	H	2	H	H	H	SPr-i	Pr-i
	Me	Me	H	H	2	H	H	H	SO ₂ Pr-i	Pr-i
35	Me	Me	H	H	2	H	H	H	SPh	Pr-i
	Me	Me	H	H	2	H	H	H	SO ₂ Ph	Pr-i
	Me	Me	H	H	2	H	H	H	SCHF ₂	Pr-i
	Me	Me	H	H	2	H	H	H	SO ₂ CHF ₂	Pr-i
40	Me	Me	H	H	2	H	H	H	NH ₂	Pr-i
	Me	Me	H	H	2	H	H	H	NHMe	Pr-i
	Me	Me	H	H	2	H	H	H	NMe ₂	Pr-i
	Me	Me	H	H	2	H	H	H	NHEt	Pr-i
45	Me	Me	H	H	2	H	H	H	NEt ₂	Pr-i
	Me	Me	H	H	2	H	H	H	NHPh	Pr-i
	Me	Me	H	H	2	H	H	H	N(Me)Ph	Pr-i
	Me	Me	H	H	2	H	H	H	CN	Pr-i
50	Me	Me	H	H	2	H	H	H	F	Pr-c
	Me	Me	H	H	2	H	H	H	Cl	Pr-c
	Me	Me	H	H	2	H	H	H	OH	Pr-c
	Me	Me	H	H	2	H	H	H	OMe	Pr-c
	Me	Me	H	H	2	H	H	H	OEt	Pr-c
55	Me	Me	H	H	2	H	H	H	OPr-i	Pr-c
	Me	Me	H	H	2	H	H	H	OPr	Pr-c

	Me	Me	H	H	2	H	H	H	OBu-t	Pr-c
	Me	Me	H	H	2	H	H	H	OCH ₂ Pr-c	Pr-c
5	Me	Me	H	H	2	H	H	H	OCH ₂ Bu-c	Pr-c
	Me	Me	H	H	2	H	H	H	OCH ₂ Pen-c	Pr-c
	Me	Me	H	H	2	H	H	H	OCH ₂ Hex-c	Pr-c
	Me	Me	H	H	2	H	H	H	OPen-c	Pr-c
10	Me	Me	H	H	2	H	H	H	OHex-c	Pr-c
	Me	Me	H	H	2	H	H	H	OCH ₂ Ph	Pr-c
	Me	Me	H	H	2	H	H	H	OPh	Pr-c
	Me	Me	H	H	2	H	H	H	OCHF ₂	Pr-c
15	Me	Me	H	H	2	H	H	H	SH	Pr-c
	Me	Me	H	H	2	H	H	H	SMe	Pr-c
	Me	Me	H	H	2	H	H	H	SO ₂ Me	Pr-c
	Me	Me	H	H	2	H	H	H	SEt	Pr-c
	Me	Me	H	H	2	H	H	H	SO ₂ Et	Pr-c
20	Me	Me	H	H	2	H	H	H	SPr-i	Pr-c
	Me	Me	H	H	2	H	H	H	SO ₂ Pr-i	Pr-c
	Me	Me	H	H	2	H	H	H	SPh	Pr-c
	Me	Me	H	H	2	H	H	H	SO ₂ Ph	Pr-c
25	Me	Me	H	H	2	H	H	H	SCHF ₂	Pr-c
	Me	Me	H	H	2	H	H	H	SO ₂ CHF ₂	Pr-c
	Me	Me	H	H	2	H	H	H	NH ₂	Pr-c
	Me	Me	H	H	2	H	H	H	NHMe	Pr-c
	Me	Me	H	H	2	H	H	H	NMe ₂	Pr-c
30	Me	Me	H	H	2	H	H	H	NHEt	Pr-c
	Me	Me	H	H	2	H	H	H	NEt ₂	Pr-c
	Me	Me	H	H	2	H	H	H	NHPh	Pr-c
	Me	Me	H	H	2	H	H	H	N(Me)Ph	Pr-c
35	Me	Me	H	H	2	H	H	H	CN	Pr-c
	Me	Me	H	H	2	H	H	H	F	CHF ₂
	Me	Me	H	H	2	H	H	H	Cl	CHF ₂
	Me	Me	H	H	2	H	H	H	OH	CHF ₂
40	Me	Me	H	H	2	H	H	H	OMe	CHF ₂
	Me	Me	H	H	2	H	H	H	OEt	CHF ₂
	Me	Me	H	H	2	H	H	H	OPr-i	CHF ₂
	Me	Me	H	H	2	H	H	H	OPr	CHF ₂
45	Me	Me	H	H	2	H	H	H	OBu-t	CHF ₂
	Me	Me	H	H	2	H	H	H	OCH ₂ Pr-c	CHF ₂
	Me	Me	H	H	2	H	H	H	OCH ₂ Bu-c	CHF ₂
	Me	Me	H	H	2	H	H	H	OCH ₂ Pen-c	CHF ₂
	Me	Me	H	H	2	H	H	H	OCH ₂ Hex-c	CHF ₂
50	Me	Me	H	H	2	H	H	H	OPen-c	CHF ₂
	Me	Me	H	H	2	H	H	H	OHex-c	CHF ₂
	Me	Me	H	H	2	H	H	H	OCH ₂ Ph	CHF ₂
	Me	Me	H	H	2	H	H	H	OPh	CHF ₂
55	Me	Me	H	H	2	H	H	H	OCHF ₂	CHF ₂
	Me	Me	H	H	2	H	H	H	SH	CHF ₂

	Me	Me	H	H	2	H	H	H	SMe	CHF ₂
	Me	Me	H	H	2	H	H	H	SO ₂ Me	CHF ₂
5	Me	Me	H	H	2	H	H	H	SEt	CHF ₂
	Me	Me	H	H	2	H	H	H	SO ₂ Et	CHF ₂
	Me	Me	H	H	2	H	H	H	SPr-i	CHF ₂
	Me	Me	H	H	2	H	H	H	SO ₂ Pr-i	CHF ₂
10	Me	Me	H	H	2	H	H	H	SPh	CHF ₂
	Me	Me	H	H	2	H	H	H	SO ₂ Ph	CHF ₂
	Me	Me	H	H	2	H	H	H	SCHF ₂	CHF ₂
	Me	Me	H	H	2	H	H	H	SO ₂ CHF ₂	CHF ₂
15	Me	Me	H	H	2	H	H	H	NH ₂	CHF ₂
	Me	Me	H	H	2	H	H	H	NHMe	CHF ₂
	Me	Me	H	H	2	H	H	H	NMe ₂	CHF ₂
	Me	Me	H	H	2	H	H	H	NHEt	CHF ₂
20	Me	Me	H	H	2	H	H	H	NEt ₂	CHF ₂
	Me	Me	H	H	2	H	H	H	NHPh	CHF ₂
	Me	Me	H	H	2	H	H	H	N(Me)Ph	CHF ₂
	Me	Me	H	H	2	H	H	H	CN	CHF ₂
	Me	Me	H	H	2	H	H	H	F	CF ₃
25	Me	Me	H	H	2	H	H	H	Cl	CF ₃
	Me	Me	H	H	2	H	H	H	OH	CF ₃
	Me	Me	H	H	2	H	H	H	OMe	CF ₃
	Me	Me	H	H	2	H	H	H	OEt	CF ₃
30	Me	Me	H	H	2	H	H	H	OPr-i	CF ₃
	Me	Me	H	H	2	H	H	H	OPr	CF ₃
	Me	Me	H	H	2	H	H	H	OBu-t	CF ₃
	Me	Me	H	H	2	H	H	H	OCH ₂ Pr-c	CF ₃
35	Me	Me	H	H	2	H	H	H	OCH ₂ Bu-c	CF ₃
	Me	Me	H	H	2	H	H	H	OCH ₂ Pen-c	CF ₃
	Me	Me	H	H	2	H	H	H	OCH ₂ Hex-c	CF ₃
	Me	Me	H	H	2	H	H	H	OPen-c	CF ₃
	Me	Me	H	H	2	H	H	H	OHex-c	CF ₃
40	Me	Me	H	H	2	H	H	H	OCH ₂ Ph	CF ₃
	Me	Me	H	H	2	H	H	H	OPh	CF ₃
	Me	Me	H	H	2	H	H	H	OCHF ₂	CF ₃
	Me	Me	H	H	2	H	H	H	SH	CF ₃
45	Me	Me	H	H	2	H	H	H	SMe	CF ₃
	Me	Me	H	H	2	H	H	H	SO ₂ Me	CF ₃
	Me	Me	H	H	2	H	H	H	SEt	CF ₃
	Me	Me	H	H	2	H	H	H	SO ₂ Et	CF ₃
50	Me	Me	H	H	2	H	H	H	SPr-i	CF ₃
	Me	Me	H	H	2	H	H	H	SO ₂ Pr-i	CF ₃
	Me	Me	H	H	2	H	H	H	SPh	CF ₃
	Me	Me	H	H	2	H	H	H	SO ₂ Ph	CF ₃
	Me	Me	H	H	2	H	H	H	SCHF ₂	CF ₃
55	Me	Me	H	H	2	H	H	H	SO ₂ CHF ₂	CF ₃
	Me	Me	H	H	2	H	H	H	NH ₂	CF ₃

	Me	Me	H	H	2	H	H	H	NHMe	CF ₃
	Me	Me	H	H	2	H	H	H	NMe ₂	CF ₃
5	Me	Me	H	H	2	H	H	H	NHEt	CF ₃
	Me	Me	H	H	2	H	H	H	NEt ₂	CF ₃
	Me	Me	H	H	2	H	H	H	NHPh	CF ₃
	Me	Me	H	H	2	H	H	H	N(Me)Ph	CF ₃
	Me	Me	H	H	2	H	H	H	CN	CF ₃
10	Me	Me	H	H	2	H	H	H	F	OMe
	Me	Me	H	H	2	H	H	H	OH	OMe
	Me	Me	H	H	2	H	H	H	OMe	OMe
	Me	Me	H	H	2	H	H	H	OEt	OMe
15	Me	Me	H	H	2	H	H	H	OPr-i	OMe
	Me	Me	H	H	2	H	H	H	OPr	OMe
	Me	Me	H	H	2	H	H	H	OBu-t	OMe
	Me	Me	H	H	2	H	H	H	OCH ₂ Pr-c	OMe
20	Me	Me	H	H	2	H	H	H	OCH ₂ Bu-c	OMe
	Me	Me	H	H	2	H	H	H	OCH ₂ Pen-c	OMe
	Me	Me	H	H	2	H	H	H	OCH ₂ Hex-c	OMe
	Me	Me	H	H	2	H	H	H	OPen-c	OMe
	Me	Me	H	H	2	H	H	H	OHex-c	OMe
25	Me	Me	H	H	2	H	H	H	OCH ₂ Ph	OMe
	Me	Me	H	H	2	H	H	H	OPh	OMe
	Me	Me	H	H	2	H	H	H	OCHF ₂	OMe
	Me	Me	H	H	2	H	H	H	SH	OMe
30	Me	Me	H	H	2	H	H	H	SMe	OMe
	Me	Me	H	H	2	H	H	H	SO ₂ Me	OMe
	Me	Me	H	H	2	H	H	H	SEt	OMe
	Me	Me	H	H	2	H	H	H	SO ₂ Et	OMe
35	Me	Me	H	H	2	H	H	H	SPr-i	OMe
	Me	Me	H	H	2	H	H	H	SO ₂ Pr-i	OMe
	Me	Me	H	H	2	H	H	H	SPh	OMe
	Me	Me	H	H	2	H	H	H	SO ₂ Ph	OMe
	Me	Me	H	H	2	H	H	H	SCHF ₂	OMe
40	Me	Me	H	H	2	H	H	H	SO ₂ CHF ₂	OMe
	Me	Me	H	H	2	H	H	H	NH ₂	OMe
	Me	Me	H	H	2	H	H	H	NHMe	OMe
	Me	Me	H	H	2	H	H	H	NMe ₂	OMe
45	Me	Me	H	H	2	H	H	H	NHEt	OMe
	Me	Me	H	H	2	H	H	H	NEt ₂	OMe
	Me	Me	H	H	2	H	H	H	NHPh	OMe
	Me	Me	H	H	2	H	H	H	N(Me)Ph	OMe
	Me	Me	H	H	2	H	H	H	CN	OMe
50	Me	Me	H	H	2	H	H	H	F	OPh
	Me	Me	H	H	2	H	H	H	OH	OPh
	Me	Me	H	H	2	H	H	H	OMe	OPh
	Me	Me	H	H	2	H	H	H	OEt	OPh
55	Me	Me	H	H	2	H	H	H	OPr-i	OPh
	Me	Me	H	H	2	H	H	H	OPr	OPh

	Me	Me	H	H	2	H	H	H	OBu-t	OPh
	Me	Me	H	H	2	H	H	H	OCH ₂ Pr-c	OPh
5	Me	Me	H	H	2	H	H	H	OCH ₂ Bu-c	OPh
	Me	Me	H	H	2	H	H	H	OCH ₂ Pen-c	OPh
	Me	Me	H	H	2	H	H	H	OCH ₂ Hex-c	OPh
	Me	Me	H	H	2	H	H	H	OPen-c	OPh
	Me	Me	H	H	2	H	H	H	OHex-c	OPh
10	Me	Me	H	H	2	H	H	H	OCH ₂ Ph	OPh
	Me	Me	H	H	2	H	H	H	OPh	OPh
	Me	Me	H	H	2	H	H	H	OCHF ₂	OPh
	Me	Me	H	H	2	H	H	H	SH	OPh
15	Me	Me	H	H	2	H	H	H	SMe	OPh
	Me	Me	H	H	2	H	H	H	SO ₂ Me	OPh
	Me	Me	H	H	2	H	H	H	SEt	OPh
	Me	Me	H	H	2	H	H	H	SO ₂ Et	OPh
	Me	Me	H	H	2	H	H	H	SPr-i	OPh
20	Me	Me	H	H	2	H	H	H	SO ₂ Pr-i	OPh
	Me	Me	H	H	2	H	H	H	SPh	OPh
	Me	Me	H	H	2	H	H	H	SO ₂ Ph	OPh
	Me	Me	H	H	2	H	H	H	SCHF ₂	OPh
25	Me	Me	H	H	2	H	H	H	SO ₂ CHF ₂	OPh
	Me	Me	H	H	2	H	H	H	NH ₂	OPh
	Me	Me	H	H	2	H	H	H	NHMe	OPh
	Me	Me	H	H	2	H	H	H	NMe ₂	OPh
30	Me	Me	H	H	2	H	H	H	NHEt	OPh
	Me	Me	H	H	2	H	H	H	NEt ₂	OPh
	Me	Me	H	H	2	H	H	H	NHPh	OPh
	Me	Me	H	H	2	H	H	H	N(Me)Ph	OPh
	Me	Me	H	H	2	H	H	H	CN	OPh
35	Me	Me	H	H	2	H	H	H	F	OCHF ₂
	Me	Me	H	H	2	H	H	H	OH	OCHF ₂
	Me	Me	H	H	2	H	H	H	OMe	OCHF ₂
	Me	Me	H	H	2	H	H	H	OEt	OCHF ₂
40	Me	Me	H	H	2	H	H	H	OPr-i	OCHF ₂
	Me	Me	H	H	2	H	H	H	OPr	OCHF ₂
	Me	Me	H	H	2	H	H	H	OBu-t	OCHF ₂
	Me	Me	H	H	2	H	H	H	OCH ₂ Pr-c	OCHF ₂
45	Me	Me	H	H	2	H	H	H	OCH ₂ Bu-c	OCHF ₂
	Me	Me	H	H	2	H	H	H	OCH ₂ Pen-c	OCHF ₂
	Me	Me	H	H	2	H	H	H	OCH ₂ Hex-c	OCHF ₂
	Me	Me	H	H	2	H	H	H	OPen-c	OCHF ₂
	Me	Me	H	H	2	H	H	H	OHex-c	OCHF ₂
50	Me	Me	H	H	2	H	H	H	OCH ₂ Ph	OCHF ₂
	Me	Me	H	H	2	H	H	H	OPh	OCHF ₂
	Me	Me	H	H	2	H	H	H	OCHF ₂	OCHF ₂
	Me	Me	H	H	2	H	H	H	SH	OCHF ₂
55	Me	Me	H	H	2	H	H	H	SMe	OCHF ₂

	Me	Me	H	H	2	H	H	H	SO ₂ Me	OCHF ₂
	Me	Me	H	H	2	H	H	H	SEt	OCHF ₂
5	Me	Me	H	H	2	H	H	H	SO ₂ Et	OCHF ₂
	Me	Me	H	H	2	H	H	H	SPr-i	OCHF ₂
	Me	Me	H	H	2	H	H	H	SO ₂ Pr-i	OCHF ₂
	Me	Me	H	H	2	H	H	H	SPh	OCHF ₂
10	Me	Me	H	H	2	H	H	H	SO ₂ Ph	OCHF ₂
	Me	Me	H	H	2	H	H	H	SCHF ₂	OCHF ₂
	Me	Me	H	H	2	H	H	H	SO ₂ CHF ₂	OCHF ₂
	Me	Me	H	H	2	H	H	H	NH ₂	OCHF ₂
15	Me	Me	H	H	2	H	H	H	NHMe	OCHF ₂
	Me	Me	H	H	2	H	H	H	NMe ₂	OCHF ₂
	Me	Me	H	H	2	H	H	H	NHEt	OCHF ₂
	Me	Me	H	H	2	H	H	H	NEt ₂	OCHF ₂
	Me	Me	H	H	2	H	H	H	NHPh	OCHF ₂
20	Me	Me	H	H	2	H	H	H	N(Me)Ph	OCHF ₂
	Me	Me	H	H	2	H	H	H	CN	OCHF ₂
	Me	Me	H	H	2	H	H	Me	F	CF ₃
	Me	Me	H	H	2	H	H	Me	Cl	CF ₃
25	Me	Me	H	H	2	H	H	Me	OH	CF ₃
	Me	Me	H	H	2	H	H	Me	OMe	CF ₃
	Me	Me	H	H	2	H	H	Me	OEt	CF ₃
	Me	Me	H	H	2	H	H	Me	OPr-i	CF ₃
30	Me	Me	H	H	2	H	H	Me	OPr	CF ₃
	Me	Me	H	H	2	H	H	Me	OBu-t	CF ₃
	Me	Me	H	H	2	H	H	Me	OCH ₂ Pr-c	CF ₃
	Me	Me	H	H	2	H	H	Me	OCH ₂ Bu-c	CF ₃
35	Me	Me	H	H	2	H	H	Me	OCH ₂ Pen-c	CF ₃
	Me	Me	H	H	2	H	H	Me	OCH ₂ Hex-c	CF ₃
	Me	Me	H	H	2	H	H	Me	OPen-c	CF ₃
	Me	Me	H	H	2	H	H	Me	OHex-c	CF ₃
	Me	Me	H	H	2	H	H	Me	OCH ₂ Ph	CF ₃
40	Me	Me	H	H	2	H	H	Me	OPh	CF ₃
	Me	Me	H	H	2	H	H	Me	OCHF ₂	CF ₃
	Me	Me	H	H	2	H	H	Me	SH	CF ₃
	Me	Me	H	H	2	H	H	Me	SMe	CF ₃
45	Me	Me	H	H	2	H	H	Me	SO ₂ Me	CF ₃
	Me	Me	H	H	2	H	H	Me	SEt	CF ₃
	Me	Me	H	H	2	H	H	Me	SO ₂ Et	CF ₃
	Me	Me	H	H	2	H	H	Me	SPr-i	CF ₃
50	Me	Me	H	H	2	H	H	Me	SO ₂ Pr-i	CF ₃
	Me	Me	H	H	2	H	H	Me	SPh	CF ₃
	Me	Me	H	H	2	H	H	Me	SO ₂ Ph	CF ₃
	Me	Me	H	H	2	H	H	Me	SCHF ₂	CF ₃
	Me	Me	H	H	2	H	H	Me	SO ₂ CHF ₂	CF ₃
55	Me	Me	H	H	2	H	H	Me	NH ₂	CF ₃
	Me	Me	H	H	2	H	H	Me	NHMe	CF ₃

	Me	Me	H	H	2	H	H	Me	NMe ₂	CF ₃
	Me	Me	H	H	2	H	H	Me	NHEt	CF ₃
5	Me	Me	H	H	2	H	H	Me	NEt ₂	CF ₃
	Me	Me	H	H	2	H	H	Me	NHPh	CF ₃
	Me	Me	H	H	2	H	H	Me	N(Me)Ph	CF ₃
	Me	Me	H	H	2	H	H	Me	CN	CF ₃
	Me	Me	H	H	2	H	H	OMe	F	CF ₃
10	Me	Me	H	H	2	H	H	OMe	Cl	CF ₃
	Me	Me	H	H	2	H	H	OMe	OH	CF ₃
	Me	Me	H	H	2	H	H	OMe	OMe	CF ₃
	Me	Me	H	H	2	H	H	OMe	OEt	CF ₃
15	Me	Me	H	H	2	H	H	OMe	OPr-i	CF ₃
	Me	Me	H	H	2	H	H	OMe	OPr	CF ₃
	Me	Me	H	H	2	H	H	OMe	OBu-t	CF ₃
	Me	Me	H	H	2	H	H	OMe	OCH ₂ Pr-c	CF ₃
20	Me	Me	H	H	2	H	H	OMe	OCH ₂ Bu-c	CF ₃
	Me	Me	H	H	2	H	H	OMe	OCH ₂ Pen-c	CF ₃
	Me	Me	H	H	2	H	H	OMe	OCH ₂ Hex-c	CF ₃
	Me	Me	H	H	2	H	H	OMe	OPen-c	CF ₃
25	Me	Me	H	H	2	H	H	OMe	OHex-c	CF ₃
	Me	Me	H	H	2	H	H	OMe	OCH ₂ Ph	CF ₃
	Me	Me	H	H	2	H	H	OMe	OPh	CF ₃
	Me	Me	H	H	2	H	H	OMe	OCHF ₂	CF ₃
30	Me	Me	H	H	2	H	H	OMe	SH	CF ₃
	Me	Me	H	H	2	H	H	OMe	SMe	CF ₃
	Me	Me	H	H	2	H	H	OMe	SO ₂ Me	CF ₃
	Me	Me	H	H	2	H	H	OMe	SEt	CF ₃
35	Me	Me	H	H	2	H	H	OMe	SO ₂ Et	CF ₃
	Me	Me	H	H	2	H	H	OMe	SPr-i	CF ₃
	Me	Me	H	H	2	H	H	OMe	SO ₂ Pr-i	CF ₃
	Me	Me	H	H	2	H	H	OMe	SPh	CF ₃
	Me	Me	H	H	2	H	H	OMe	SO ₂ Ph	CF ₃
40	Me	Me	H	H	2	H	H	OMe	SCHF ₂	CF ₃
	Me	Me	H	H	2	H	H	OMe	SO ₂ CHF ₂	CF ₃
	Me	Me	H	H	2	H	H	OMe	NH ₂	CF ₃
	Me	Me	H	H	2	H	H	OMe	NHMe	CF ₃
45	Me	Me	H	H	2	H	H	OMe	NMe ₂	CF ₃
	Me	Me	H	H	2	H	H	OMe	NHEt	CF ₃
	Me	Me	H	H	2	H	H	OMe	NEt ₂	CF ₃
	Me	Me	H	H	2	H	H	OMe	NHPh	CF ₃
	Me	Me	H	H	2	H	H	OMe	N(Me)Ph	CF ₃
50	Me	Me	H	H	2	H	H	OMe	CN	CF ₃
	Me	Me	H	H	2	H	H	SMe	F	CF ₃
	Me	Me	H	H	2	H	H	SMe	Cl	CF ₃
	Me	Me	H	H	2	H	H	SMe	OH	CF ₃
55	Me	Me	H	H	2	H	H	SMe	OMe	CF ₃
	Me	Me	H	H	2	H	H	SMe	OEt	CF ₃

	Me	Me	H	H	2	H	H	SMe	OPr-i	CF ₃
	Me	Me	H	H	2	H	H	SMe	OPr	CF ₃
5	Me	Me	H	H	2	H	H	SMe	OBu-t	CF ₃
	Me	Me	H	H	2	H	H	SMe	OCH ₂ Pr-c	CF ₃
	Me	Me	H	H	2	H	H	SMe	OCH ₂ Bu-c	CF ₃
	Me	Me	H	H	2	H	H	SMe	OCH ₂ Pen-c	CF ₃
10	Me	Me	H	H	2	H	H	SMe	OCH ₂ Hex-c	CF ₃
	Me	Me	H	H	2	H	H	SMe	OPen-c	CF ₃
	Me	Me	H	H	2	H	H	SMe	OHex-c	CF ₃
	Me	Me	H	H	2	H	H	SMe	OCH ₂ Ph	CF ₃
15	Me	Me	H	H	2	H	H	SMe	OPh	CF ₃
	Me	Me	H	H	2	H	H	SMe	OCHF ₂	CF ₃
	Me	Me	H	H	2	H	H	SMe	SH	CF ₃
	Me	Me	H	H	2	H	H	SMe	SMe	CF ₃
20	Me	Me	H	H	2	H	H	SMe	SO ₂ Me	CF ₃
	Me	Me	H	H	2	H	H	SMe	SEt	CF ₃
	Me	Me	H	H	2	H	H	SMe	SO ₂ Et	CF ₃
	Me	Me	H	H	2	H	H	SMe	SPr-i	CF ₃
25	Me	Me	H	H	2	H	H	SMe	SO ₂ Pr-i	CF ₃
	Me	Me	H	H	2	H	H	SMe	SPh	CF ₃
	Me	Me	H	H	2	H	H	SMe	SO ₂ Ph	CF ₃
	Me	Me	H	H	2	H	H	SMe	SCHF ₂	CF ₃
	Me	Me	H	H	2	H	H	SMe	SO ₂ CHF ₂	CF ₃
30	Me	Me	H	H	2	H	H	SMe	NH ₂	CF ₃
	Me	Me	H	H	2	H	H	SMe	NHMe	CF ₃
	Me	Me	H	H	2	H	H	SMe	NMe ₂	CF ₃
	Me	Me	H	H	2	H	H	SMe	NHEt	CF ₃
35	Me	Me	H	H	2	H	H	SMe	NEt ₂	CF ₃
	Me	Me	H	H	2	H	H	SMe	NHPh	CF ₃
	Me	Me	H	H	2	H	H	SMe	N(Me)Ph	CF ₃
	Me	Me	H	H	2	H	H	SMe	CN	CF ₃
40	Me	Me	H	H	2	H	H	SO ₂ Me	F	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	Cl	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	OH	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	OMe	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	OEt	CF ₃
45	Me	Me	H	H	2	H	H	SO ₂ Me	OPr-i	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	OPr	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	OBu-t	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	OCH ₂ Pr-c	CF ₃
50	Me	Me	H	H	2	H	H	SO ₂ Me	OCH ₂ Bu-c	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	OCH ₂ Pen-c	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	OCH ₂ Hex-c	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	OPen-c	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	OHex-c	CF ₃
55	Me	Me	H	H	2	H	H	SO ₂ Me	OCH ₂ Ph	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	OPh	CF ₃

	Me	Me	H	H	2	H	H	SO ₂ Me	OCHF ₂	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	SH	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	SMe	CF ₃
5	Me	Me	H	H	2	H	H	SO ₂ Me	SO ₂ Me	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	SEt	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	SO ₂ Et	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	SPr-i	CF ₃
10	Me	Me	H	H	2	H	H	SO ₂ Me	SO ₂ Pr-i	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	SPh	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	SO ₂ Ph	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	SCHF ₂	CF ₃
15	Me	Me	H	H	2	H	H	SO ₂ Me	SO ₂ CHF ₂	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	NH ₂	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	NHMe	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	NMe ₂	CF ₃
20	Me	Me	H	H	2	H	H	SO ₂ Me	NHEt	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	NEt ₂	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	NHPh	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	N(Me)Ph	CF ₃
	Me	Me	H	H	2	H	H	SO ₂ Me	CN	CF ₃
25	Me	Me	H	H	2	H	H	NH ₂	F	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	Cl	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	OH	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	OMe	CF ₃
30	Me	Me	H	H	2	H	H	NH ₂	OEt	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	OPr-i	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	OPr	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	OBu-t	CF ₃
35	Me	Me	H	H	2	H	H	NH ₂	OCH ₂ Pr-c	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	OCH ₂ Bu-c	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	OCH ₂ Pen-c	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	OCH ₂ Hex-c	CF ₃
40	Me	Me	H	H	2	H	H	NH ₂	OPen-c	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	OHex-c	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	OCH ₂ Ph	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	OPh	CF ₃
45	Me	Me	H	H	2	H	H	NH ₂	OCHF ₂	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	SH	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	SMe	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	SO ₂ Me	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	SEt	CF ₃
50	Me	Me	H	H	2	H	H	NH ₂	SO ₂ Et	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	SPr-i	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	SO ₂ Pr-i	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	SPh	CF ₃
55	Me	Me	H	H	2	H	H	NH ₂	SO ₂ Ph	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	SCHF ₂	CF ₃

	Me	Me	H	H	2	H	H	NH ₂	SO ₂ CHF ₂	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	NH ₂	CF ₃
5	Me	Me	H	H	2	H	H	NH ₂	NHMe	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	NMe ₂	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	NHEt	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	NEt ₂	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	NHPh	CF ₃
10	Me	Me	H	H	2	H	H	NH ₂	N(Me)Ph	CF ₃
	Me	Me	H	H	2	H	H	NH ₂	CN	CF ₃
	H	H	H	H	2	H	H	H	OMe	CF ₃
	H	H	H	H	2	H	H	H	OEt	CF ₃
15	Me	H	H	H	2	H	H	H	OMe	CF ₃
	Me	H	H	H	2	H	H	H	OEt	CF ₃
	Me	H	Me	H	2	H	H	H	OMe	CF ₃
	Me	H	Me	H	2	H	H	H	OEt	CF ₃
20	Me	Me	H	H	2	Me	H	H	OMe	CF ₃
	Me	Me	H	H	2	Me	H	H	OEt	CF ₃
	Me	Me	H	H	2	Et	H	H	OMe	CF ₃
	Me	Me	H	H	2	Et	H	H	OEt	CF ₃
	Me	Me	H	H	2	Pr-i	H	H	H	CF ₃
25	Me	Me	H	H	2	Pr-i	H	H	OMe	CF ₃
	Me	Me	H	H	2	Pr-i	H	H	OEt	CF ₃
	Me	Me	H	H	2	Me	Me	H	OMe	CF ₃
	Me	Me	H	H	2	Me	Me	H	OEt	CF ₃
30	Me	Et	H	H	2	H	H	H	OMe	CF ₃
	Me	Et	H	H	2	H	H	H	OEt	CF ₃
	Et	Et	H	H	2	H	H	H	OMe	CF ₃
	Et	Et	H	H	2	H	H	H	OEt	CF ₃
35	Me	Pr-i	H	H	2	H	H	H	OMe	CF ₃
	Me	Pr-i	H	H	2	H	H	H	OEt	CF ₃
	Me	Pr	H	H	2	H	H	H	OMe	CF ₃
	Me	Pr	H	H	2	H	H	H	OEt	CF ₃
40	Me	Pr-c	H	H	2	H	H	H	OMe	CF ₃
	Me	Pr-c	H	H	2	H	H	H	OEt	CF ₃
	Me	CH ₂ Pr-c	H	H	2	H	H	H	OMe	CF ₃
	Me	CH ₂ Pr-c	H	H	2	H	H	H	OEt	CF ₃
45		-(CH ₂) ₂ -	H	H	2	H	H	H	OMe	CF ₃
		-(CH ₂) ₂ -	H	H	2	H	H	H	OEt	CF ₃
		-(CH ₂) ₃ -	H	H	2	H	H	H	OMe	CF ₃
		-(CH ₂) ₃ -	H	H	2	H	H	H	OEt	CF ₃
		-(CH ₂) ₄ -	H	H	2	H	H	H	OMe	CF ₃
50		-(CH ₂) ₄ -	H	H	2	H	H	H	OEt	CF ₃
		-(CH ₂) ₅ -	H	H	2	H	H	H	OMe	CF ₃
		-(CH ₂) ₅ -	H	H	2	H	H	H	OEt	CF ₃
55	H	-(CH ₂) ₃ -	H		2	H	H	H	OMe	CF ₃
	H	-(CH ₂) ₃ -	H		2	H	H	H	OEt	CF ₃
	H	-(CH ₂) ₄ -	H		2	H	H	H	OMe	CF ₃

	H	-(CH ₂) ₄ -	H	2	H	H	H	OEt	CF ₃
	H	-(CH ₂) ₅ -	H	2	H	H	H	OMe	CF ₃
	H	-(CH ₂) ₅ -	H	2	H	H	H	OEt	CF ₃
5	H	-(CH ₂) ₆ -	H	2	H	H	H	OMe	CF ₃
	H	-(CH ₂) ₆ -	H	2	H	H	H	OEt	CF ₃
	Me	Me	H	1	H	H	H	Cl	Cl
	Me	Me	H	1	H	H	H	OH	Cl
10	Me	Me	H	1	H	H	H	OMe	Cl
	Me	Me	H	1	H	H	H	OEt	Cl
	Me	Me	H	1	H	H	H	OPr-i	Cl
	Me	Me	H	1	H	H	H	OPr	Cl
15	Me	Me	H	1	H	H	H	OBu-t	Cl
	Me	Me	H	1	H	H	H	OCH ₂ Pr-c	Cl
	Me	Me	H	1	H	H	H	OCH ₂ Bu-c	Cl
	Me	Me	H	1	H	H	H	OCH ₂ Pen-c	Cl
20	Me	Me	H	1	H	H	H	OCH ₂ Hex-c	Cl
	Me	Me	H	1	H	H	H	OPen-c	Cl
	Me	Me	H	1	H	H	H	OHex-c	Cl
	Me	Me	H	1	H	H	H	OCH ₂ Ph	Cl
	Me	Me	H	1	H	H	H	OPh	Cl
25	Me	Me	H	1	H	H	H	OCHF ₂	Cl
	Me	Me	H	1	H	H	H	SH	Cl
	Me	Me	H	1	H	H	H	SMe	Cl
	Me	Me	H	1	H	H	H	SO ₂ Me	Cl
	Me	Me	H	1	H	H	H	SEt	Cl
30	Me	Me	H	1	H	H	H	SO ₂ Et	Cl
	Me	Me	H	1	H	H	H	SPr-i	Cl
	Me	Me	H	1	H	H	H	SO ₂ Pr-i	Cl
	Me	Me	H	1	H	H	H	SPh	Cl
35	Me	Me	H	1	H	H	H	SO ₂ Ph	Cl
	Me	Me	H	1	H	H	H	SCHF ₂	Cl
	Me	Me	H	1	H	H	H	SO ₂ CHF ₂	Cl
	Me	Me	H	1	H	H	H	NH ₂	Cl
40	Me	Me	H	1	H	H	H	NHMe	Cl
	Me	Me	H	1	H	H	H	NMe ₂	Cl
	Me	Me	H	1	H	H	H	NHEt	Cl
	Me	Me	H	1	H	H	H	NEt ₂	Cl
	Me	Me	H	1	H	H	H	NHPh	Cl
45	Me	Me	H	1	H	H	H	N(Me)Ph	Cl
	Me	Me	H	1	H	H	H	CN	Cl
	Me	Me	H	1	H	H	H	F	Me
	Me	Me	H	1	H	H	H	Cl	Me
50	Me	Me	H	1	H	H	H	OH	Me
	Me	Me	H	1	H	H	H	OMe	Me
	Me	Me	H	1	H	H	H	OEt	Me
	Me	Me	H	1	H	H	H	OPr-i	Me
	Me	Me	H	1	H	H	H	OPr	Me
55	Me	Me	H	1	H	H	H	OBu-t	Me

	Me	Me	H	H	1	H	H	H	OCH ₂ Pr-c	Me
	Me	Me	H	H	1	H	H	H	OCH ₂ Bu-c	Me
	Me	Me	H	H	1	H	H	H	OCH ₂ Pen-c	Me
5	Me	Me	H	H	1	H	H	H	OCH ₂ Hex-c	Me
	Me	Me	H	H	1	H	H	H	OPen-c	Me
	Me	Me	H	H	1	H	H	H	OHex-c	Me
	Me	Me	H	H	1	H	H	H	OCH ₂ Ph	Me
10	Me	Me	H	H	1	H	H	H	OPh	Me
	Me	Me	H	H	1	H	H	H	OCHF ₂	Me
	Me	Me	H	H	1	H	H	H	SH	Me
	Me	Me	H	H	1	H	H	H	SMe	Me
15	Me	Me	H	H	1	H	H	H	SO ₂ Me	Me
	Me	Me	H	H	1	H	H	H	SEt	Me
	Me	Me	H	H	1	H	H	H	SO ₂ Et	Me
	Me	Me	H	H	1	H	H	H	SPr-i	Me
	Me	Me	H	H	1	H	H	H	SO ₂ Pr-i	Me
20	Me	Me	H	H	1	H	H	H	SPh	Me
	Me	Me	H	H	1	H	H	H	SO ₂ Ph	Me
	Me	Me	H	H	1	H	H	H	SCHF ₂	Me
	Me	Me	H	H	1	H	H	H	SO ₂ CHF ₂	Me
25	Me	Me	H	H	1	H	H	H	NH ₂	Me
	Me	Me	H	H	1	H	H	H	NHMe	Me
	Me	Me	H	H	1	H	H	H	NMe ₂	Me
	Me	Me	H	H	1	H	H	H	NHEt	Me
	Me	Me	H	H	1	H	H	H	NEt ₂	Me
30	Me	Me	H	H	1	H	H	H	NHPh	Me
	Me	Me	H	H	1	H	H	H	N(Me)Ph	Me
	Me	Me	H	H	1	H	H	H	CN	Me
	Me	Me	H	H	1	H	H	H	F	Pr-i
35	Me	Me	H	H	1	H	H	H	Cl	Pr-i
	Me	Me	H	H	1	H	H	H	OH	Pr-i
	Me	Me	H	H	1	H	H	H	OMe	Pr-i
	Me	Me	H	H	1	H	H	H	OEt	Pr-i
40	Me	Me	H	H	1	H	H	H	OPr-i	Pr-i
	Me	Me	H	H	1	H	H	H	OPr	Pr-i
	Me	Me	H	H	1	H	H	H	OBu-t	Pr-i
	Me	Me	H	H	1	H	H	H	OCH ₂ Pr-c	Pr-i
45	Me	Me	H	H	1	H	H	H	OCH ₂ Bu-c	Pr-i
	Me	Me	H	H	1	H	H	H	OCH ₂ Pen-c	Pr-i
	Me	Me	H	H	1	H	H	H	OCH ₂ Hex-c	Pr-i
	Me	Me	H	H	1	H	H	H	OPen-c	Pr-i
	Me	Me	H	H	1	H	H	H	OHex-c	Pr-i
50	Me	Me	H	H	1	H	H	H	OCH ₂ Ph	Pr-i
	Me	Me	H	H	1	H	H	H	OPh	Pr-i
	Me	Me	H	H	1	H	H	H	OCHF ₂	Pr-i
	Me	Me	H	H	1	H	H	H	SH	Pr-i
55	Me	Me	H	H	1	H	H	H	SMe	Pr-i
	Me	Me	H	H	1	H	H	H	SO ₂ Me	Pr-i

	Me	Me	H	H	1	H	H	H	SEt	Pr-i
	Me	Me	H	H	1	H	H	H	SO ₂ Et	Pr-i
5	Me	Me	H	H	1	H	H	H	SPr-i	Pr-i
	Me	Me	H	H	1	H	H	H	SO ₂ Pr-i	Pr-i
	Me	Me	H	H	1	H	H	H	SPh	Pr-i
	Me	Me	H	H	1	H	H	H	SO ₂ Ph	Pr-i
10	Me	Me	H	H	1	H	H	H	SCHF ₂	Pr-i
	Me	Me	H	H	1	H	H	H	SO ₂ CHF ₂	Pr-i
	Me	Me	H	H	1	H	H	H	NH ₂	Pr-i
	Me	Me	H	H	1	H	H	H	NHMe	Pr-i
15	Me	Me	H	H	1	H	H	H	NMe ₂	Pr-i
	Me	Me	H	H	1	H	H	H	NHEt	Pr-i
	Me	Me	H	H	1	H	H	H	NEt ₂	Pr-i
	Me	Me	H	H	1	H	H	H	NHPh	Pr-i
20	Me	Me	H	H	1	H	H	H	N(Me)Ph	Pr-i
	Me	Me	H	H	1	H	H	H	CN	Pr-i
	Me	Me	H	H	1	H	H	H	F	Pr-c
	Me	Me	H	H	1	H	H	H	Cl	Pr-c
	Me	Me	H	H	1	H	H	H	OH	Pr-c
25	Me	Me	H	H	1	H	H	H	OMe	Pr-c
	Me	Me	H	H	1	H	H	H	OEt	Pr-c
	Me	Me	H	H	1	H	H	H	OPr-i	Pr-c
	Me	Me	H	H	1	H	H	H	OPr	Pr-c
30	Me	Me	H	H	1	H	H	H	OBu-t	Pr-c
	Me	Me	H	H	1	H	H	H	OCH ₂ Pr-c	Pr-c
	Me	Me	H	H	1	H	H	H	OCH ₂ Bu-c	Pr-c
	Me	Me	H	H	1	H	H	H	OCH ₂ Pen-c	Pr-c
	Me	Me	H	H	1	H	H	H	OCH ₂ Hex-c	Pr-c
35	Me	Me	H	H	1	H	H	H	OPen-c	Pr-c
	Me	Me	H	H	1	H	H	H	OHex-c	Pr-c
	Me	Me	H	H	1	H	H	H	OCH ₂ Ph	Pr-c
	Me	Me	H	H	1	H	H	H	OPh	Pr-c
40	Me	Me	H	H	1	H	H	H	OCHF ₂	Pr-c
	Me	Me	H	H	1	H	H	H	SH	Pr-c
	Me	Me	H	H	1	H	H	H	SMe	Pr-c
	Me	Me	H	H	1	H	H	H	SO ₂ Me	Pr-c
45	Me	Me	H	H	1	H	H	H	SEt	Pr-c
	Me	Me	H	H	1	H	H	H	SO ₂ Et	Pr-c
	Me	Me	H	H	1	H	H	H	SPr-i	Pr-c
	Me	Me	H	H	1	H	H	H	SO ₂ Pr-i	Pr-c
	Me	Me	H	H	1	H	H	H	SPh	Pr-c
50	Me	Me	H	H	1	H	H	H	SO ₂ Ph	Pr-c
	Me	Me	H	H	1	H	H	H	SCHF ₂	Pr-c
	Me	Me	H	H	1	H	H	H	SO ₂ CHF ₂	Pr-c
	Me	Me	H	H	1	H	H	H	NH ₂	Pr-c
55	Me	Me	H	H	1	H	H	H	NHMe	Pr-c
	Me	Me	H	H	1	H	H	H	NMe ₂	Pr-c
	Me	Me	H	H	1	H	H	H	NHEt	Pr-c

	Me	Me	H	H	1	H	H	H	NEt ₂	Pr-c
	Me	Me	H	H	1	H	H	H	NHPh	Pr-c
5	Me	Me	H	H	1	H	H	H	N(Me)Ph	Pr-c
	Me	Me	H	H	1	H	H	H	CN	Pr-c
	Me	Me	H	H	1	H	H	H	F	CHF ₂
	Me	Me	H	H	1	H	H	H	Cl	CHF ₂
	Me	Me	H	H	1	H	H	H	OH	CHF ₂
10	Me	Me	H	H	1	H	H	H	OMe	CHF ₂
	Me	Me	H	H	1	H	H	H	OEt	CHF ₂
	Me	Me	H	H	1	H	H	H	OPr-i	CHF ₂
	Me	Me	H	H	1	H	H	H	OPr	CHF ₂
15	Me	Me	H	H	1	H	H	H	OBu-t	CHF ₂
	Me	Me	H	H	1	H	H	H	OCH ₂ Pr-c	CHF ₂
	Me	Me	H	H	1	H	H	H	OCH ₂ Bu-c	CHF ₂
	Me	Me	H	H	1	H	H	H	OCH ₂ Pen-c	CHF ₂
20	Me	Me	H	H	1	H	H	H	OCH ₂ Hex-c	CHF ₂
	Me	Me	H	H	1	H	H	H	OPen-c	CHF ₂
	Me	Me	H	H	1	H	H	H	OHex-c	CHF ₂
	Me	Me	H	H	1	H	H	H	OCH ₂ Ph	CHF ₂
25	Me	Me	H	H	1	H	H	H	OPh	CHF ₂
	Me	Me	H	H	1	H	H	H	OCHF ₂	CHF ₂
	Me	Me	H	H	1	H	H	H	SH	CHF ₂
	Me	Me	H	H	1	H	H	H	SMe	CHF ₂
30	Me	Me	H	H	1	H	H	H	SO ₂ Me	CHF ₂
	Me	Me	H	H	1	H	H	H	SEt	CHF ₂
	Me	Me	H	H	1	H	H	H	SO ₂ Et	CHF ₂
	Me	Me	H	H	1	H	H	H	SPr-i	CHF ₂
	Me	Me	H	H	1	H	H	H	SO ₂ Pr-i	CHF ₂
35	Me	Me	H	H	1	H	H	H	SPh	CHF ₂
	Me	Me	H	H	1	H	H	H	SO ₂ Ph	CHF ₂
	Me	Me	H	H	1	H	H	H	SCHF ₂	CHF ₂
	Me	Me	H	H	1	H	H	H	SO ₂ CHF ₂	CHF ₂
40	Me	Me	H	H	1	H	H	H	NH ₂	CHF ₂
	Me	Me	H	H	1	H	H	H	NHMe	CHF ₂
	Me	Me	H	H	1	H	H	H	NMe ₂	CHF ₂
	Me	Me	H	H	1	H	H	H	NHEt	CHF ₂
45	Me	Me	H	H	1	H	H	H	NEt ₂	CHF ₂
	Me	Me	H	H	1	H	H	H	NHPh	CHF ₂
	Me	Me	H	H	1	H	H	H	N(Me)Ph	CHF ₂
	Me	Me	H	H	1	H	H	H	CN	CHF ₂
50	Me	Me	H	H	1	H	H	H	F	CF ₃
	Me	Me	H	H	1	H	H	H	Cl	CF ₃
	Me	Me	H	H	1	H	H	H	OH	CF ₃
	Me	Me	H	H	1	H	H	H	OMe	CF ₃
	Me	Me	H	H	1	H	H	H	OEt	CF ₃
55	Me	Me	H	H	1	H	H	H	OPr-i	CF ₃
	Me	Me	H	H	1	H	H	H	OPr	CF ₃

	Me	Me	H	H	1	H	H	H	OBu-t	CF ₃
	Me	Me	H	H	1	H	H	H	OCH ₂ Pr-c	CF ₃
5	Me	Me	H	H	1	H	H	H	OCH ₂ Bu-c	CF ₃
	Me	Me	H	H	1	H	H	H	OCH ₂ Pen-c	CF ₃
	Me	Me	H	H	1	H	H	H	OCH ₂ Hex-c	CF ₃
	Me	Me	H	H	1	H	H	H	OPen-c	CF ₃
10	Me	Me	H	H	1	H	H	H	OHex-c	CF ₃
	Me	Me	H	H	1	H	H	H	OCH ₂ Ph	CF ₃
	Me	Me	H	H	1	H	H	H	OPh	CF ₃
	Me	Me	H	H	1	H	H	H	OCHF ₂	CF ₃
15	Me	Me	H	H	1	H	H	H	SH	CF ₃
	Me	Me	H	H	1	H	H	H	SMe	CF ₃
	Me	Me	H	H	1	H	H	H	SO ₂ Me	CF ₃
	Me	Me	H	H	1	H	H	H	SEt	CF ₃
	Me	Me	H	H	1	H	H	H	SO ₂ Et	CF ₃
20	Me	Me	H	H	1	H	H	H	SPr-i	CF ₃
	Me	Me	H	H	1	H	H	H	SO ₂ Pr-i	CF ₃
	Me	Me	H	H	1	H	H	H	SPh	CF ₃
	Me	Me	H	H	1	H	H	H	SO ₂ Ph	CF ₃
25	Me	Me	H	H	1	H	H	H	SCHF ₂	CF ₃
	Me	Me	H	H	1	H	H	H	SO ₂ CHF ₂	CF ₃
	Me	Me	H	H	1	H	H	H	NH ₂	CF ₃
	Me	Me	H	H	1	H	H	H	NHMe	CF ₃
30	Me	Me	H	H	1	H	H	H	NMe ₂	CF ₃
	Me	Me	H	H	1	H	H	H	NHEt	CF ₃
	Me	Me	H	H	1	H	H	H	NEt ₂	CF ₃
	Me	Me	H	H	1	H	H	H	NHPh	CF ₃
35	Me	Me	H	H	1	H	H	H	N(Me)Ph	CF ₃
	Me	Me	H	H	1	H	H	H	CN	CF ₃
	Me	Me	H	H	1	H	H	H	F	OMe
	Me	Me	H	H	1	H	H	H	OH	OMe
	Me	Me	H	H	1	H	H	H	OMe	OMe
40	Me	Me	H	H	1	H	H	H	OEt	OMe
	Me	Me	H	H	1	H	H	H	OPr-i	OMe
	Me	Me	H	H	1	H	H	H	OPr	OMe
	Me	Me	H	H	1	H	H	H	OBu-t	OMe
45	Me	Me	H	H	1	H	H	H	OCH ₂ Pr-c	OMe
	Me	Me	H	H	1	H	H	H	OCH ₂ Bu-c	OMe
	Me	Me	H	H	1	H	H	H	OCH ₂ Pen-c	OMe
	Me	Me	H	H	1	H	H	H	OCH ₂ Hex-c	OMe
	Me	Me	H	H	1	H	H	H	OPen-c	OMe
50	Me	Me	H	H	1	H	H	H	OHex-c	OMe
	Me	Me	H	H	1	H	H	H	OCH ₂ Ph	OMe
	Me	Me	H	H	1	H	H	H	OPh	OMe
	Me	Me	H	H	1	H	H	H	OCHF ₂	OMe
55	Me	Me	H	H	1	H	H	H	SH	OMe
	Me	Me	H	H	1	H	H	H	SMe	OMe

	Me	Me	H	H	1	H	H	H	SO ₂ Me	OMe
	Me	Me	H	H	1	H	H	H	SEt	OMe
5	Me	Me	H	H	1	H	H	H	SO ₂ Et	OMe
	Me	Me	H	H	1	H	H	H	SPr-i	OMe
	Me	Me	H	H	1	H	H	H	SO ₂ Pr-i	OMe
	Me	Me	H	H	1	H	H	H	SPh	OMe
10	Me	Me	H	H	1	H	H	H	SO ₂ Ph	OMe
	Me	Me	H	H	1	H	H	H	SCHF ₂	OMe
	Me	Me	H	H	1	H	H	H	SO ₂ CHF ₂	OMe
	Me	Me	H	H	1	H	H	H	NH ₂	OMe
	Me	Me	H	H	1	H	H	H	NHMe	OMe
15	Me	Me	H	H	1	H	H	H	NMe ₂	OMe
	Me	Me	H	H	1	H	H	H	NHEt	OMe
	Me	Me	H	H	1	H	H	H	NEt ₂	OMe
	Me	Me	H	H	1	H	H	H	NHPh	OMe
20	Me	Me	H	H	1	H	H	H	N(Me)Ph	OMe
	Me	Me	H	H	1	H	H	H	CN	OMe
	Me	Me	H	H	1	H	H	H	F	OPh
	Me	Me	H	H	1	H	H	H	OH	OPh
25	Me	Me	H	H	1	H	H	H	OMe	OPh
	Me	Me	H	H	1	H	H	H	OEt	OPh
	Me	Me	H	H	1	H	H	H	OPr-i	OPh
	Me	Me	H	H	1	H	H	H	OPr	OPh
	Me	Me	H	H	1	H	H	H	OBu-t	OPh
30	Me	Me	H	H	1	H	H	H	OCH ₂ Pr-c	OPh
	Me	Me	H	H	1	H	H	H	OCH ₂ Bu-c	OPh
	Me	Me	H	H	1	H	H	H	OCH ₂ Pen-c	OPh
	Me	Me	H	H	1	H	H	H	OCH ₂ Hex-c	OPh
35	Me	Me	H	H	1	H	H	H	OPen-c	OPh
	Me	Me	H	H	1	H	H	H	OHex-c	OPh
	Me	Me	H	H	1	H	H	H	OCH ₂ Ph	OPh
	Me	Me	H	H	1	H	H	H	OPh	OPh
	Me	Me	H	H	1	H	H	H	OCHF ₂	OPh
40	Me	Me	H	H	1	H	H	H	SH	OPh
	Me	Me	H	H	1	H	H	H	SMe	OPh
	Me	Me	H	H	1	H	H	H	SO ₂ Me	OPh
	Me	Me	H	H	1	H	H	H	SEt	OPh
45	Me	Me	H	H	1	H	H	H	SO ₂ Et	OPh
	Me	Me	H	H	1	H	H	H	SPr-i	OPh
	Me	Me	H	H	1	H	H	H	SO ₂ Pr-i	OPh
	Me	Me	H	H	1	H	H	H	SPh	OPh
50	Me	Me	H	H	1	H	H	H	SO ₂ Ph	OPh
	Me	Me	H	H	1	H	H	H	SCHF ₂	OPh
	Me	Me	H	H	1	H	H	H	SO ₂ CHF ₂	OPh
	Me	Me	H	H	1	H	H	H	NH ₂	OPh
	Me	Me	H	H	1	H	H	H	NHMe	OPh
55	Me	Me	H	H	1	H	H	H	NMe ₂	OPh
	Me	Me	H	H	1	H	H	H	NHEt	OPh

	Me	Me	H	H	1	H	H	H	NEt ₂	OPh
	Me	Me	H	H	1	H	H	H	NHPh	OPh
5	Me	Me	H	H	1	H	H	H	N(Me)Ph	OPh
	Me	Me	H	H	1	H	H	H	CN	OPh
	Me	Me	H	H	1	H	H	H	F	OCHF ₂
	Me	Me	H	H	1	H	H	H	OH	OCHF ₂
10	Me	Me	H	H	1	H	H	H	OMe	OCHF ₂
	Me	Me	H	H	1	H	H	H	OEt	OCHF ₂
	Me	Me	H	H	1	H	H	H	OPr-i	OCHF ₂
	Me	Me	H	H	1	H	H	H	OPr	OCHF ₂
15	Me	Me	H	H	1	H	H	H	OBu-t	OCHF ₂
	Me	Me	H	H	1	H	H	H	OCH ₂ Pr-c	OCHF ₂
	Me	Me	H	H	1	H	H	H	OCH ₂ Bu-c	OCHF ₂
	Me	Me	H	H	1	H	H	H	OCH ₂ Pen-c	OCHF ₂
	Me	Me	H	H	1	H	H	H	OCH ₂ Hex-c	OCHF ₂
20	Me	Me	H	H	1	H	H	H	OPen-c	OCHF ₂
	Me	Me	H	H	1	H	H	H	OHex-c	OCHF ₂
	Me	Me	H	H	1	H	H	H	OCH ₂ Ph	OCHF ₂
	Me	Me	H	H	1	H	H	H	OPh	OCHF ₂
25	Me	Me	H	H	1	H	H	H	OCHF ₂	OCHF ₂
	Me	Me	H	H	1	H	H	H	SH	OCHF ₂
	Me	Me	H	H	1	H	H	H	SMe	OCHF ₂
	Me	Me	H	H	1	H	H	H	SO ₂ Me	OCHF ₂
30	Me	Me	H	H	1	H	H	H	SEt	OCHF ₂
	Me	Me	H	H	1	H	H	H	SO ₂ Et	OCHF ₂
	Me	Me	H	H	1	H	H	H	SPr-i	OCHF ₂
	Me	Me	H	H	1	H	H	H	SO ₂ Pr-i	OCHF ₂
35	Me	Me	H	H	1	H	H	H	SPh	OCHF ₂
	Me	Me	H	H	1	H	H	H	SO ₂ Ph	OCHF ₂
	Me	Me	H	H	1	H	H	H	SCHF ₂	OCHF ₂
	Me	Me	H	H	1	H	H	H	SO ₂ CHF ₂	OCHF ₂
40	Me	Me	H	H	1	H	H	H	NH ₂	OCHF ₂
	Me	Me	H	H	1	H	H	H	NHMe	OCHF ₂
	Me	Me	H	H	1	H	H	H	NMe ₂	OCHF ₂
	Me	Me	H	H	1	H	H	H	NHEt	OCHF ₂
45	Me	Me	H	H	1	H	H	H	NEt ₂	OCHF ₂
	Me	Me	H	H	1	H	H	H	NHPh	OCHF ₂
	Me	Me	H	H	1	H	H	H	N(Me)Ph	OCHF ₂
	Me	Me	H	H	1	H	H	H	CN	OCHF ₂
50	Me	Me	H	H	1	H	H	Me	F	CF ₃
	Me	Me	H	H	1	H	H	Me	Cl	CF ₃
	Me	Me	H	H	1	H	H	Me	OH	CF ₃
	Me	Me	H	H	1	H	H	Me	OMe	CF ₃
	Me	Me	H	H	1	H	H	Me	OEt	CF ₃
55	Me	Me	H	H	1	H	H	Me	OPr-i	CF ₃
	Me	Me	H	H	1	H	H	Me	OPr	CF ₃
	Me	Me	H	H	1	H	H	Me	OBu-t	CF ₃

	Me	Me	H	H	1	H	H	Me	OCH ₂ Pr-c	CF ₃
	Me	Me	H	H	1	H	H	Me	OCH ₂ Bu-c	CF ₃
5	Me	Me	H	H	1	H	H	Me	OCH ₂ Pen-c	CF ₃
	Me	Me	H	H	1	H	H	Me	OCH ₂ Hex-c	CF ₃
	Me	Me	H	H	1	H	H	Me	OPen-c	CF ₃
	Me	Me	H	H	1	H	H	Me	OHex-c	CF ₃
10	Me	Me	H	H	1	H	H	Me	OCH ₂ Ph	CF ₃
	Me	Me	H	H	1	H	H	Me	OPh	CF ₃
	Me	Me	H	H	1	H	H	Me	OCHF ₂	CF ₃
	Me	Me	H	H	1	H	H	Me	SH	CF ₃
15	Me	Me	H	H	1	H	H	Me	SMe	CF ₃
	Me	Me	H	H	1	H	H	Me	SO ₂ Me	CF ₃
	Me	Me	H	H	1	H	H	Me	SEt	CF ₃
	Me	Me	H	H	1	H	H	Me	SO ₂ Et	CF ₃
	Me	Me	H	H	1	H	H	Me	SPr-i	CF ₃
20	Me	Me	H	H	1	H	H	Me	SO ₂ Pr-i	CF ₃
	Me	Me	H	H	1	H	H	Me	SPh	CF ₃
	Me	Me	H	H	1	H	H	Me	SO ₂ Ph	CF ₃
	Me	Me	H	H	1	H	H	Me	SCHF ₂	CF ₃
25	Me	Me	H	H	1	H	H	Me	SO ₂ CHF ₂	CF ₃
	Me	Me	H	H	1	H	H	Me	NH ₂	CF ₃
	Me	Me	H	H	1	H	H	Me	NHMe	CF ₃
	Me	Me	H	H	1	H	H	Me	NMe ₂	CF ₃
	Me	Me	H	H	1	H	H	Me	NHEt	CF ₃
30	Me	Me	H	H	1	H	H	Me	NEt ₂	CF ₃
	Me	Me	H	H	1	H	H	Me	NHPh	CF ₃
	Me	Me	H	H	1	H	H	Me	N(Me)Ph	CF ₃
	Me	Me	H	H	1	H	H	Me	CN	CF ₃
35	Me	Me	H	H	1	H	H	OMe	F	CF ₃
	Me	Me	H	H	1	H	H	OMe	Cl	CF ₃
	Me	Me	H	H	1	H	H	OMe	OH	CF ₃
	Me	Me	H	H	1	H	H	OMe	OMe	CF ₃
40	Me	Me	H	H	1	H	H	OMe	OEt	CF ₃
	Me	Me	H	H	1	H	H	OMe	OPr-i	CF ₃
	Me	Me	H	H	1	H	H	OMe	OPr	CF ₃
	Me	Me	H	H	1	H	H	OMe	OBu-t	CF ₃
45	Me	Me	H	H	1	H	H	OMe	OCH ₂ Pr-c	CF ₃
	Me	Me	H	H	1	H	H	OMe	OCH ₂ Bu-c	CF ₃
	Me	Me	H	H	1	H	H	OMe	OCH ₂ Pen-c	CF ₃
	Me	Me	H	H	1	H	H	OMe	OCH ₂ Hex-c	CF ₃
	Me	Me	H	H	1	H	H	OMe	OPen-c	CF ₃
50	Me	Me	H	H	1	H	H	OMe	OHex-c	CF ₃
	Me	Me	H	H	1	H	H	OMe	OCH ₂ Ph	CF ₃
	Me	Me	H	H	1	H	H	OMe	OPh	CF ₃
	Me	Me	H	H	1	H	H	OMe	OCHF ₂	CF ₃
55	Me	Me	H	H	1	H	H	OMe	SH	CF ₃
	Me	Me	H	H	1	H	H	OMe	SMe	CF ₃

	Me	Me	H	H	1	H	H	OMe	SO ₂ Me	CF ₃
	Me	Me	H	H	1	H	H	OMe	SEt	CF ₃
5	Me	Me	H	H	1	H	H	OMe	SO ₂ Et	CF ₃
	Me	Me	H	H	1	H	H	OMe	SP <i>r</i> -i	CF ₃
	Me	Me	H	H	1	H	H	OMe	SO ₂ Pr-i	CF ₃
	Me	Me	H	H	1	H	H	OMe	SPh	CF ₃
10	Me	Me	H	H	1	H	H	OMe	SO ₂ Ph	CF ₃
	Me	Me	H	H	1	H	H	OMe	SCHF ₂	CF ₃
	Me	Me	H	H	1	H	H	OMe	SO ₂ CHF ₂	CF ₃
	Me	Me	H	H	1	H	H	OMe	NH ₂	CF ₃
	Me	Me	H	H	1	H	H	OMe	NHMe	CF ₃
15	Me	Me	H	H	1	H	H	OMe	NMe ₂	CF ₃
	Me	Me	H	H	1	H	H	OMe	NHEt	CF ₃
	Me	Me	H	H	1	H	H	OMe	NEt ₂	CF ₃
	Me	Me	H	H	1	H	H	OMe	NHPh	CF ₃
20	Me	Me	H	H	1	H	H	OMe	N(Me)Ph	CF ₃
	Me	Me	H	H	1	H	H	OMe	CN	CF ₃
	Me	Me	H	H	1	H	H	SMe	F	CF ₃
	Me	Me	H	H	1	H	H	SMe	Cl	CF ₃
25	Me	Me	H	H	1	H	H	SMe	OH	CF ₃
	Me	Me	H	H	1	H	H	SMe	OMe	CF ₃
	Me	Me	H	H	1	H	H	SMe	OEt	CF ₃
	Me	Me	H	H	1	H	H	SMe	OP <i>r</i> -i	CF ₃
30	Me	Me	H	H	1	H	H	SMe	OP <i>r</i>	CF ₃
	Me	Me	H	H	1	H	H	SMe	OBu-t	CF ₃
	Me	Me	H	H	1	H	H	SMe	OCH ₂ Pr-c	CF ₃
	Me	Me	H	H	1	H	H	SMe	OCH ₂ Bu-c	CF ₃
	Me	Me	H	H	1	H	H	SMe	OCH ₂ Pen-c	CF ₃
35	Me	Me	H	H	1	H	H	SMe	OCH ₂ Hex-c	CF ₃
	Me	Me	H	H	1	H	H	SMe	OPen-c	CF ₃
	Me	Me	H	H	1	H	H	SMe	OHex-c	CF ₃
	Me	Me	H	H	1	H	H	SMe	OCH ₂ Ph	CF ₃
40	Me	Me	H	H	1	H	H	SMe	OPh	CF ₃
	Me	Me	H	H	1	H	H	SMe	OCHF ₂	CF ₃
	Me	Me	H	H	1	H	H	SMe	SH	CF ₃
	Me	Me	H	H	1	H	H	SMe	SMe	CF ₃
45	Me	Me	H	H	1	H	H	SMe	SO ₂ Me	CF ₃
	Me	Me	H	H	1	H	H	SMe	SEt	CF ₃
	Me	Me	H	H	1	H	H	SMe	SO ₂ Et	CF ₃
	Me	Me	H	H	1	H	H	SMe	SP <i>r</i> -i	CF ₃
50	Me	Me	H	H	1	H	H	SMe	SO ₂ Pr-i	CF ₃
	Me	Me	H	H	1	H	H	SMe	SPh	CF ₃
	Me	Me	H	H	1	H	H	SMe	SO ₂ Ph	CF ₃
	Me	Me	H	H	1	H	H	SMe	SCHF ₂	CF ₃
	Me	Me	H	H	1	H	H	SMe	SO ₂ CHF ₂	CF ₃
55	Me	Me	H	H	1	H	H	SMe	NH ₂	CF ₃
	Me	Me	H	H	1	H	H	SMe	NHMe	CF ₃

	Me	Me	H	H	1	H	H	SMe	NMe ₂	CF ₃
	Me	Me	H	H	1	H	H	SMe	NHEt	CF ₃
5	Me	Me	H	H	1	H	H	SMe	NEt ₂	CF ₃
	Me	Me	H	H	1	H	H	SMe	NHPh	CF ₃
	Me	Me	H	H	1	H	H	SMe	N(Me)Ph	CF ₃
	Me	Me	H	H	1	H	H	SMe	CN	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	F	CF ₃
10	Me	Me	H	H	1	H	H	SO ₂ Me	Cl	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	OH	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	OMe	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	OEt	CF ₃
15	Me	Me	H	H	1	H	H	SO ₂ Me	OPr-i	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	OPr	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	OBu-t	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	OCH ₂ Pr-c	CF ₃
20	Me	Me	H	H	1	H	H	SO ₂ Me	OCH ₂ Bu-c	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	OCH ₂ Pen-c	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	OCH ₂ Hex-c	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	OPen-c	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	OHex-c	CF ₃
25	Me	Me	H	H	1	H	H	SO ₂ Me	OCH ₂ Ph	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	OPh	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	OCHF ₂	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	SH	CF ₃
30	Me	Me	H	H	1	H	H	SO ₂ Me	SMe	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	SO ₂ Me	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	SEt	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	SO ₂ Et	CF ₃
35	Me	Me	H	H	1	H	H	SO ₂ Me	SPr-i	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	SO ₂ Pr-i	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	SPh	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	SO ₂ Ph	CF ₃
40	Me	Me	H	H	1	H	H	SO ₂ Me	SCHF ₂	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	SO ₂ CHF ₂	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	NH ₂	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	NHMe	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	NMe ₂	CF ₃
45	Me	Me	H	H	1	H	H	SO ₂ Me	NHEt	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	NEt ₂	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	NHPh	CF ₃
	Me	Me	H	H	1	H	H	SO ₂ Me	N(Me)Ph	CF ₃
50	Me	Me	H	H	1	H	H	SO ₂ Me	CN	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	F	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	Cl	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	OH	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	OMe	CF ₃
55	Me	Me	H	H	1	H	H	NH ₂	OEt	CF ₃

	Me	Me	H	H	1	H	H	NH ₂	OPr-i	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	OPr	CF ₃
5	Me	Me	H	H	1	H	H	NH ₂	OBu-t	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	OCH ₂ Pr-c	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	OCH ₂ Bu-c	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	OCH ₂ Pen-c	CF ₃
10	Me	Me	H	H	1	H	H	NH ₂	OCH ₂ Hex-c	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	OPen-c	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	OHex-c	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	OCH ₂ Ph	CF ₃
15	Me	Me	H	H	1	H	H	NH ₂	OPh	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	OCHF ₂	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	SH	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	SMe	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	SO ₂ Me	CF ₃
20	Me	Me	H	H	1	H	H	NH ₂	SEt	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	SO ₂ Et	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	SPr-i	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	SO ₂ Pr-i	CF ₃
25	Me	Me	H	H	1	H	H	NH ₂	SPh	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	SO ₂ Ph	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	SCHF ₂	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	SO ₂ CHF ₂	CF ₃
30	Me	Me	H	H	1	H	H	NH ₂	NH ₂	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	NHMe	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	NMe ₂	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	NHEt	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	NEt ₂	CF ₃
35	Me	Me	H	H	1	H	H	NH ₂	NHPh	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	N(Me)Ph	CF ₃
	Me	Me	H	H	1	H	H	NH ₂	CN	CF ₃
	H	H	H	H	1	H	H	H	OMe	CF ₃
40	H	H	H	H	1	H	H	H	OEt	CF ₃
	Me	H	H	H	1	H	H	H	OMe	CF ₃
	Me	H	H	H	1	H	H	H	OEt	CF ₃
	Me	H	Me	H	1	H	H	H	OMe	CF ₃
45	Me	H	Me	H	1	H	H	H	OEt	CF ₃
	Me	Me	H	H	1	Me	H	H	OMe	CF ₃
	Me	Me	H	H	1	Me	H	H	OEt	CF ₃
	Me	Me	H	H	1	Et	H	H	OMe	CF ₃
	Me	Me	H	H	1	Et	H	H	OEt	CF ₃
50	Me	Me	H	H	1	Pr-i	H	H	H	CF ₃
	Me	Me	H	H	1	Pr-i	H	H	OMe	CF ₃
	Me	Me	H	H	1	Pr-i	H	H	OEt	CF ₃
	Me	Me	H	H	1	Me	Me	H	OMe	CF ₃
55	Me	Me	H	H	1	Me	Me	H	OEt	CF ₃
	Me	Et	H	H	1	H	H	H	OMe	CF ₃

	Me	Et	H	H	1	H	H	H	OEt	CF ₃
	Et	Et	H	H	1	H	H	H	OMe	CF ₃
5	Et	Et	H	H	1	H	H	H	OEt	CF ₃
	Me	Pr-i	H	H	1	H	H	H	OMe	CF ₃
	Me	Pr-i	H	H	1	H	H	H	OEt	CF ₃
	Me	Pr	H	H	1	H	H	H	OMe	CF ₃
10	Me	Pr	H	H	1	H	H	H	OEt	CF ₃
	Me	Pr-c	H	H	1	H	H	H	OMe	CF ₃
	Me	Pr-c	H	H	1	H	H	H	OEt	CF ₃
	Me	CH ₂ Pr-c	H	H	1	H	H	H	OMe	CF ₃
	Me	CH ₂ Pr-c	H	H	1	H	H	H	OEt	CF ₃
15		-(CH ₂) ₂ -	H	H	1	H	H	H	OMe	CF ₃
		-(CH ₂) ₂ -	H	H	1	H	H	H	OEt	CF ₃
		-(CH ₂) ₃ -	H	H	1	H	H	H	OMe	CF ₃
		-(CH ₂) ₃ -	H	H	1	H	H	H	OEt	CF ₃
20		-(CH ₂) ₄ -	H	H	1	H	H	H	OMe	CF ₃
		-(CH ₂) ₄ -	H	H	1	H	H	H	OEt	CF ₃
		-(CH ₂) ₅ -	H	H	1	H	H	H	OMe	CF ₃
		-(CH ₂) ₅ -	H	H	1	H	H	H	OEt	CF ₃
25	H	-(CH ₂) ₃ -	H	H	1	H	H	H	OMe	CF ₃
	H	-(CH ₂) ₃ -	H	H	1	H	H	H	OEt	CF ₃
	H	-(CH ₂) ₄ -	H	H	1	H	H	H	OMe	CF ₃
	H	-(CH ₂) ₄ -	H	H	1	H	H	H	OEt	CF ₃
	H	-(CH ₂) ₅ -	H	H	1	H	H	H	OMe	CF ₃
30	H	-(CH ₂) ₅ -	H	H	1	H	H	H	OEt	CF ₃
	H	-(CH ₂) ₆ -	H	H	1	H	H	H	OMe	CF ₃
	H	-(CH ₂) ₆ -	H	H	1	H	H	H	OEt	CF ₃
35	Me	Me	H	H	0	H	H	H	Cl	Cl
	Me	Me	H	H	0	H	H	H	OH	Cl
	Me	Me	H	H	0	H	H	H	OMe	Cl
	Me	Me	H	H	0	H	H	H	OEt	Cl
	Me	Me	H	H	0	H	H	H	OPr-i	Cl
40	Me	Me	H	H	0	H	H	H	OPr	Cl
	Me	Me	H	H	0	H	H	H	OBu-t	Cl
	Me	Me	H	H	0	H	H	H	OCH ₂ Pr-c	Cl
	Me	Me	H	H	0	H	H	H	OCH ₂ Bu-c	Cl
45	Me	Me	H	H	0	H	H	H	OCH ₂ Pen-c	Cl
	Me	Me	H	H	0	H	H	H	OCH ₂ Hex-c	Cl
	Me	Me	H	H	0	H	H	H	OPen-c	Cl
	Me	Me	H	H	0	H	H	H	OHex-c	Cl
	Me	Me	H	H	0	H	H	H	OCH ₂ Ph	Cl
50	Me	Me	H	H	0	H	H	H	OPh	Cl
	Me	Me	H	H	0	H	H	H	OCHF ₂	Cl
	Me	Me	H	H	0	H	H	H	SH	Cl
	Me	Me	H	H	0	H	H	H	SMe	Cl
	Me	Me	H	H	0	H	H	H	SO ₂ Me	Cl
55	Me	Me	H	H	0	H	H	H	SEt	Cl

	Me	Me	H	H	0	H	H	H	SO ₂ Et	Cl
	Me	Me	H	H	0	H	H	H	SP <i>r</i> -i	Cl
5	Me	Me	H	H	0	H	H	H	SO ₂ Pr-i	Cl
	Me	Me	H	H	0	H	H	H	SPh	Cl
	Me	Me	H	H	0	H	H	H	SO ₂ Ph	Cl
	Me	Me	H	H	0	H	H	H	SCHF ₂	Cl
10	Me	Me	H	H	0	H	H	H	SO ₂ CHF ₂	Cl
	Me	Me	H	H	0	H	H	H	NH ₂	Cl
	Me	Me	H	H	0	H	H	H	NHMe	Cl
	Me	Me	H	H	0	H	H	H	NMe ₂	Cl
15	Me	Me	H	H	0	H	H	H	NHEt	Cl
	Me	Me	H	H	0	H	H	H	NEt ₂	Cl
	Me	Me	H	H	0	H	H	H	NHPh	Cl
	Me	Me	H	H	0	H	H	H	N(Me)Ph	Cl
	Me	Me	H	H	0	H	H	H	CN	Cl
20	Me	Me	H	H	0	H	H	H	F	Me
	Me	Me	H	H	0	H	H	H	Cl	Me
	Me	Me	H	H	0	H	H	H	OH	Me
	Me	Me	H	H	0	H	H	H	OMe	Me
25	Me	Me	H	H	0	H	H	H	OEt	Me
	Me	Me	H	H	0	H	H	H	OP <i>r</i> -i	Me
	Me	Me	H	H	0	H	H	H	OP <i>r</i>	Me
	Me	Me	H	H	0	H	H	H	OBu-t	Me
30	Me	Me	H	H	0	H	H	H	OCH ₂ Pr-c	Me
	Me	Me	H	H	0	H	H	H	OCH ₂ Bu-c	Me
	Me	Me	H	H	0	H	H	H	OCH ₂ Pen-c	Me
	Me	Me	H	H	0	H	H	H	OCH ₂ Hex-c	Me
	Me	Me	H	H	0	H	H	H	OPen-c	Me
35	Me	Me	H	H	0	H	H	H	OHex-c	Me
	Me	Me	H	H	0	H	H	H	OCH ₂ Ph	Me
	Me	Me	H	H	0	H	H	H	OPh	Me
	Me	Me	H	H	0	H	H	H	OCHF ₂	Me
40	Me	Me	H	H	0	H	H	H	SH	Me
	Me	Me	H	H	0	H	H	H	SMe	Me
	Me	Me	H	H	0	H	H	H	SO ₂ Me	Me
	Me	Me	H	H	0	H	H	H	SEt	Me
45	Me	Me	H	H	0	H	H	H	SO ₂ Et	Me
	Me	Me	H	H	0	H	H	H	SP <i>r</i> -i	Me
	Me	Me	H	H	0	H	H	H	SO ₂ Pr-i	Me
	Me	Me	H	H	0	H	H	H	SPh	Me
	Me	Me	H	H	0	H	H	H	SO ₂ Ph	Me
50	Me	Me	H	H	0	H	H	H	SCHF ₂	Me
	Me	Me	H	H	0	H	H	H	SO ₂ CHF ₂	Me
	Me	Me	H	H	0	H	H	H	NH ₂	Me
	Me	Me	H	H	0	H	H	H	NHMe	Me
55	Me	Me	H	H	0	H	H	H	NMe ₂	Me
	Me	Me	H	H	0	H	H	H	NHEt	Me
	Me	Me	H	H	0	H	H	H	NEt ₂	Me

	Me	Me	H	H	0	H	H	H	NHPh	Me
	Me	Me	H	H	0	H	H	H	N(Me)Ph	Me
	Me	Me	H	H	0	H	H	H	CN	Me
5	Me	Me	H	H	0	H	H	H	F	Pr-i
	Me	Me	H	H	0	H	H	H	Cl	Pr-i
	Me	Me	H	H	0	H	H	H	OH	Pr-i
	Me	Me	H	H	0	H	H	H	OMe	Pr-i
10	Me	Me	H	H	0	H	H	H	OEt	Pr-i
	Me	Me	H	H	0	H	H	H	OPr-i	Pr-i
	Me	Me	H	H	0	H	H	H	OPr	Pr-i
	Me	Me	H	H	0	H	H	H	OBu-t	Pr-i
15	Me	Me	H	H	0	H	H	H	OCH ₂ Pr-c	Pr-i
	Me	Me	H	H	0	H	H	H	OCH ₂ Bu-c	Pr-i
	Me	Me	H	H	0	H	H	H	OCH ₂ Pen-c	Pr-i
	Me	Me	H	H	0	H	H	H	OCH ₂ Hex-c	Pr-i
	Me	Me	H	H	0	H	H	H	OPen-c	Pr-i
20	Me	Me	H	H	0	H	H	H	OHex-c	Pr-i
	Me	Me	H	H	0	H	H	H	OCH ₂ Ph	Pr-i
	Me	Me	H	H	0	H	H	H	OPh	Pr-i
	Me	Me	H	H	0	H	H	H	OCHF ₂	Pr-i
25	Me	Me	H	H	0	H	H	H	SH	Pr-i
	Me	Me	H	H	0	H	H	H	SMe	Pr-i
	Me	Me	H	H	0	H	H	H	SO ₂ Me	Pr-i
	Me	Me	H	H	0	H	H	H	SEt	Pr-i
	Me	Me	H	H	0	H	H	H	SO ₂ Et	Pr-i
30	Me	Me	H	H	0	H	H	H	SPr-i	Pr-i
	Me	Me	H	H	0	H	H	H	SO ₂ Pr-i	Pr-i
	Me	Me	H	H	0	H	H	H	SPh	Pr-i
	Me	Me	H	H	0	H	H	H	SO ₂ Ph	Pr-i
35	Me	Me	H	H	0	H	H	H	SCHF ₂	Pr-i
	Me	Me	H	H	0	H	H	H	SO ₂ CHF ₂	Pr-i
	Me	Me	H	H	0	H	H	H	NH ₂	Pr-i
	Me	Me	H	H	0	H	H	H	NHMe	Pr-i
40	Me	Me	H	H	0	H	H	H	NMe ₂	Pr-i
	Me	Me	H	H	0	H	H	H	NHEt	Pr-i
	Me	Me	H	H	0	H	H	H	NEt ₂	Pr-i
	Me	Me	H	H	0	H	H	H	NHPh	Pr-i
45	Me	Me	H	H	0	H	H	H	N(Me)Ph	Pr-i
	Me	Me	H	H	0	H	H	H	CN	Pr-i
	Me	Me	H	H	0	H	H	H	F	Pr-c
	Me	Me	H	H	0	H	H	H	Cl	Pr-c
	Me	Me	H	H	0	H	H	H	OH	Pr-c
50	Me	Me	H	H	0	H	H	H	OMe	Pr-c
	Me	Me	H	H	0	H	H	H	OEt	Pr-c
	Me	Me	H	H	0	H	H	H	OPr-i	Pr-c
	Me	Me	H	H	0	H	H	H	OPr	Pr-c
	Me	Me	H	H	0	H	H	H	OBu-t	Pr-c
55	Me	Me	H	H	0	H	H	H	OCH ₂ Pr-c	Pr-c

	Me	Me	H	H	0	H	H	H	OCH ₂ Bu-c	Pr-c
	Me	Me	H	H	0	H	H	H	OCH ₂ Pen-c	Pr-c
5	Me	Me	H	H	0	H	H	H	OCH ₂ Hex-c	Pr-c
	Me	Me	H	H	0	H	H	H	OPen-c	Pr-c
	Me	Me	H	H	0	H	H	H	OHex-c	Pr-c
	Me	Me	H	H	0	H	H	H	OCH ₂ Ph	Pr-c
10	Me	Me	H	H	0	H	H	H	OPh	Pr-c
	Me	Me	H	H	0	H	H	H	OCHF ₂	Pr-c
	Me	Me	H	H	0	H	H	H	SH	Pr-c
	Me	Me	H	H	0	H	H	H	SMe	Pr-c
15	Me	Me	H	H	0	H	H	H	SO ₂ Me	Pr-c
	Me	Me	H	H	0	H	H	H	SEt	Pr-c
	Me	Me	H	H	0	H	H	H	SO ₂ Et	Pr-c
	Me	Me	H	H	0	H	H	H	SPr-i	Pr-c
	Me	Me	H	H	0	H	H	H	SO ₂ Pr-i	Pr-c
20	Me	Me	H	H	0	H	H	H	SPh	Pr-c
	Me	Me	H	H	0	H	H	H	SO ₂ Ph	Pr-c
	Me	Me	H	H	0	H	H	H	SCHF ₂	Pr-c
	Me	Me	H	H	0	H	H	H	SO ₂ CHF ₂	Pr-c
25	Me	Me	H	H	0	H	H	H	NH ₂	Pr-c
	Me	Me	H	H	0	H	H	H	NHMe	Pr-c
	Me	Me	H	H	0	H	H	H	NMe ₂	Pr-c
	Me	Me	H	H	0	H	H	H	NHEt	Pr-c
30	Me	Me	H	H	0	H	H	H	NEt ₂	Pr-c
	Me	Me	H	H	0	H	H	H	NHPh	Pr-c
	Me	Me	H	H	0	H	H	H	N(Me)Ph	Pr-c
	Me	Me	H	H	0	H	H	H	CN	Pr-c
35	Me	Me	H	H	0	H	H	H	F	CHF ₂
	Me	Me	H	H	0	H	H	H	Cl	CHF ₂
	Me	Me	H	H	0	H	H	H	OH	CHF ₂
	Me	Me	H	H	0	H	H	H	OMe	CHF ₂
40	Me	Me	H	H	0	H	H	H	OEt	CHF ₂
	Me	Me	H	H	0	H	H	H	OPr-i	CHF ₂
	Me	Me	H	H	0	H	H	H	OPr	CHF ₂
	Me	Me	H	H	0	H	H	H	OBu-t	CHF ₂
	Me	Me	H	H	0	H	H	H	OCH ₂ Pr-c	CHF ₂
45	Me	Me	H	H	0	H	H	H	OCH ₂ Bu-c	CHF ₂
	Me	Me	H	H	0	H	H	H	OCH ₂ Pen-c	CHF ₂
	Me	Me	H	H	0	H	H	H	OCH ₂ Hex-c	CHF ₂
	Me	Me	H	H	0	H	H	H	OPen-c	CHF ₂
	Me	Me	H	H	0	H	H	H	OHex-c	CHF ₂
50	Me	Me	H	H	0	H	H	H	OCH ₂ Ph	CHF ₂
	Me	Me	H	H	0	H	H	H	OPh	CHF ₂
	Me	Me	H	H	0	H	H	H	OCHF ₂	CHF ₂
	Me	Me	H	H	0	H	H	H	SH	CHF ₂
55	Me	Me	H	H	0	H	H	H	SMe	CHF ₂
	Me	Me	H	H	0	H	H	H	SO ₂ Me	CHF ₂

	Me	Me	H	H	0	H	H	H	SEt	CHF ₂
	Me	Me	H	H	0	H	H	H	SO ₂ Et	CHF ₂
5	Me	Me	H	H	0	H	H	H	SPr-i	CHF ₂
	Me	Me	H	H	0	H	H	H	SO ₂ Pr-i	CHF ₂
	Me	Me	H	H	0	H	H	H	SPh	CHF ₂
	Me	Me	H	H	0	H	H	H	SO ₂ Ph	CHF ₂
10	Me	Me	H	H	0	H	H	H	SCHF ₂	CHF ₂
	Me	Me	H	H	0	H	H	H	SO ₂ CHF ₂	CHF ₂
	Me	Me	H	H	0	H	H	H	NH ₂	CHF ₂
	Me	Me	H	H	0	H	H	H	NHMe	CHF ₂
15	Me	Me	H	H	0	H	H	H	NMe ₂	CHF ₂
	Me	Me	H	H	0	H	H	H	NHEt	CHF ₂
	Me	Me	H	H	0	H	H	H	NEt ₂	CHF ₂
	Me	Me	H	H	0	H	H	H	NHPh	CHF ₂
20	Me	Me	H	H	0	H	H	H	N(Me)Ph	CHF ₂
	Me	Me	H	H	0	H	H	H	CN	CHF ₂
	Me	Me	H	H	0	H	H	H	F	CF ₃
	Me	Me	H	H	0	H	H	H	Cl	CF ₃
	Me	Me	H	H	0	H	H	H	OH	CF ₃
25	Me	Me	H	H	0	H	H	H	OMe	CF ₃
	Me	Me	H	H	0	H	H	H	OEt	CF ₃
	Me	Me	H	H	0	H	H	H	OPr-i	CF ₃
	Me	Me	H	H	0	H	H	H	OPr	CF ₃
30	Me	Me	H	H	0	H	H	H	OBu-t	CF ₃
	Me	Me	H	H	0	H	H	H	OCH ₂ Pr-c	CF ₃
	Me	Me	H	H	0	H	H	H	OCH ₂ Bu-c	CF ₃
	Me	Me	H	H	0	H	H	H	OCH ₂ Pen-c	CF ₃
	Me	Me	H	H	0	H	H	H	OCH ₂ Hex-c	CF ₃
35	Me	Me	H	H	0	H	H	H	OPen-c	CF ₃
	Me	Me	H	H	0	H	H	H	OHex-c	CF ₃
	Me	Me	H	H	0	H	H	H	OCH ₂ Ph	CF ₃
	Me	Me	H	H	0	H	H	H	OPh	CF ₃
40	Me	Me	H	H	0	H	H	H	OCHF ₂	CF ₃
	Me	Me	H	H	0	H	H	H	SH	CF ₃
	Me	Me	H	H	0	H	H	H	SMe	CF ₃
	Me	Me	H	H	0	H	H	H	SO ₂ Me	CF ₃
45	Me	Me	H	H	0	H	H	H	SEt	CF ₃
	Me	Me	H	H	0	H	H	H	SO ₂ Et	CF ₃
	Me	Me	H	H	0	H	H	H	SPr-i	CF ₃
	Me	Me	H	H	0	H	H	H	SO ₂ Pr-i	CF ₃
	Me	Me	H	H	0	H	H	H	SPh	CF ₃
50	Me	Me	H	H	0	H	H	H	SO ₂ Ph	CF ₃
	Me	Me	H	H	0	H	H	H	SCHF ₂	CF ₃
	Me	Me	H	H	0	H	H	H	SO ₂ CHF ₂	CF ₃
	Me	Me	H	H	0	H	H	H	NH ₂	CF ₃
55	Me	Me	H	H	0	H	H	H	NHMe	CF ₃
	Me	Me	H	H	0	H	H	H	NMe ₂	CF ₃

	Me	Me	H	H	0	H	H	H	NHEt	CF ₃
	Me	Me	H	H	0	H	H	H	NEt ₂	CF ₃
5	Me	Me	H	H	0	H	H	H	NHPh	CF ₃
	Me	Me	H	H	0	H	H	H	N(Me)Ph	CF ₃
	Me	Me	H	H	0	H	H	H	CN	CF ₃
	Me	Me	H	H	0	H	H	H	F	OMe
	Me	Me	H	H	0	H	H	H	OH	OMe
10	Me	Me	H	H	0	H	H	H	OMe	OMe
	Me	Me	H	H	0	H	H	H	OEt	OMe
	Me	Me	H	H	0	H	H	H	OPr-i	OMe
	Me	Me	H	H	0	H	H	H	OPr	OMe
15	Me	Me	H	H	0	H	H	H	OBu-t	OMe
	Me	Me	H	H	0	H	H	H	OCH ₂ Pr-c	OMe
	Me	Me	H	H	0	H	H	H	OCH ₂ Bu-c	OMe
	Me	Me	H	H	0	H	H	H	OCH ₂ Pen-c	OMe
20	Me	Me	H	H	0	H	H	H	OCH ₂ Hex-c	OMe
	Me	Me	H	H	0	H	H	H	OPen-c	OMe
	Me	Me	H	H	0	H	H	H	OHex-c	OMe
	Me	Me	H	H	0	H	H	H	OCH ₂ Ph	OMe
	Me	Me	H	H	0	H	H	H	OPh	OMe
25	Me	Me	H	H	0	H	H	H	OCHF ₂	OMe
	Me	Me	H	H	0	H	H	H	SH	OMe
	Me	Me	H	H	0	H	H	H	SMe	OMe
	Me	Me	H	H	0	H	H	H	SO ₂ Me	OMe
30	Me	Me	H	H	0	H	H	H	SEt	OMe
	Me	Me	H	H	0	H	H	H	SO ₂ Et	OMe
	Me	Me	H	H	0	H	H	H	SPr-i	OMe
	Me	Me	H	H	0	H	H	H	SO ₂ Pr-i	OMe
	Me	Me	H	H	0	H	H	H	SPh	OMe
35	Me	Me	H	H	0	H	H	H	SO ₂ Ph	OMe
	Me	Me	H	H	0	H	H	H	SCHF ₂	OMe
	Me	Me	H	H	0	H	H	H	SO ₂ CHF ₂	OMe
	Me	Me	H	H	0	H	H	H	NH ₂	OMe
40	Me	Me	H	H	0	H	H	H	NHMe	OMe
	Me	Me	H	H	0	H	H	H	NMe ₂	OMe
	Me	Me	H	H	0	H	H	H	NHEt	OMe
	Me	Me	H	H	0	H	H	H	NEt ₂	OMe
45	Me	Me	H	H	0	H	H	H	NHPh	OMe
	Me	Me	H	H	0	H	H	H	N(Me)Ph	OMe
	Me	Me	H	H	0	H	H	H	CN	OMe
	Me	Me	H	H	0	H	H	H	F	OPh
	Me	Me	H	H	0	H	H	H	OH	OPh
50	Me	Me	H	H	0	H	H	H	OMe	OPh
	Me	Me	H	H	0	H	H	H	OEt	OPh
	Me	Me	H	H	0	H	H	H	OPr-i	OPh
	Me	Me	H	H	0	H	H	H	OPr	OPh
55	Me	Me	H	H	0	H	H	H	OBu-t	OPh
	Me	Me	H	H	0	H	H	H	OCH ₂ Pr-c	OPh

	Me	Me	H	H	0	H	H	H	OCH ₂ Bu-c	OPh
	Me	Me	H	H	0	H	H	H	OCH ₂ Pen-c	OPh
5	Me	Me	H	H	0	H	H	H	OCH ₂ Hex-c	OPh
	Me	Me	H	H	0	H	H	H	OPen-c	OPh
	Me	Me	H	H	0	H	H	H	OHex-c	OPh
	Me	Me	H	H	0	H	H	H	OCH ₂ Ph	OPh
10	Me	Me	H	H	0	H	H	H	OPh	OPh
	Me	Me	H	H	0	H	H	H	OCHF ₂	OPh
	Me	Me	H	H	0	H	H	H	SH	OPh
	Me	Me	H	H	0	H	H	H	SMe	OPh
15	Me	Me	H	H	0	H	H	H	SO ₂ Me	OPh
	Me	Me	H	H	0	H	H	H	SEt	OPh
	Me	Me	H	H	0	H	H	H	SO ₂ Et	OPh
	Me	Me	H	H	0	H	H	H	SPr-i	OPh
	Me	Me	H	H	0	H	H	H	SO ₂ Pr-i	OPh
20	Me	Me	H	H	0	H	H	H	SPh	OPh
	Me	Me	H	H	0	H	H	H	SO ₂ Ph	OPh
	Me	Me	H	H	0	H	H	H	SCHF ₂	OPh
	Me	Me	H	H	0	H	H	H	SO ₂ CHF ₂	OPh
25	Me	Me	H	H	0	H	H	H	NH ₂	OPh
	Me	Me	H	H	0	H	H	H	NHMe	OPh
	Me	Me	H	H	0	H	H	H	NMe ₂	OPh
	Me	Me	H	H	0	H	H	H	NHEt	OPh
	Me	Me	H	H	0	H	H	H	NEt ₂	OPh
30	Me	Me	H	H	0	H	H	H	NHPh	OPh
	Me	Me	H	H	0	H	H	H	N(Me)Ph	OPh
	Me	Me	H	H	0	H	H	H	CN	OPh
	Me	Me	H	H	0	H	H	H	F	OCHF ₂
35	Me	Me	H	H	0	H	H	H	OH	OCHF ₂
	Me	Me	H	H	0	H	H	H	OMe	OCHF ₂
	Me	Me	H	H	0	H	H	H	OEt	OCHF ₂
	Me	Me	H	H	0	H	H	H	OPr-i	OCHF ₂
40	Me	Me	H	H	0	H	H	H	OPr	OCHF ₂
	Me	Me	H	H	0	H	H	H	OBu-t	OCHF ₂
	Me	Me	H	H	0	H	H	H	OCH ₂ Pr-c	OCHF ₂
	Me	Me	H	H	0	H	H	H	OCH ₂ Bu-c	OCHF ₂
45	Me	Me	H	H	0	H	H	H	OCH ₂ Pen-c	OCHF ₂
	Me	Me	H	H	0	H	H	H	OCH ₂ Hex-c	OCHF ₂
	Me	Me	H	H	0	H	H	H	OPen-c	OCHF ₂
	Me	Me	H	H	0	H	H	H	OHex-c	OCHF ₂
	Me	Me	H	H	0	H	H	H	OCH ₂ Ph	OCHF ₂
50	Me	Me	H	H	0	H	H	H	OPh	OCHF ₂
	Me	Me	H	H	0	H	H	H	OCHF ₂	OCHF ₂
	Me	Me	H	H	0	H	H	H	SH	OCHF ₂
	Me	Me	H	H	0	H	H	H	SMe	OCHF ₂
55	Me	Me	H	H	0	H	H	H	SO ₂ Me	OCHF ₂
	Me	Me	H	H	0	H	H	H	SEt	OCHF ₂

	Me	Me	H	H	0	H	H	H	SO ₂ Et	OCHF ₂
	Me	Me	H	H	0	H	H	H	SP <i>r</i> -i	OCHF ₂
5	Me	Me	H	H	0	H	H	H	SO ₂ Pr-i	OCHF ₂
	Me	Me	H	H	0	H	H	H	SPh	OCHF ₂
	Me	Me	H	H	0	H	H	H	SO ₂ Ph	OCHF ₂
	Me	Me	H	H	0	H	H	H	SCHF ₂	OCHF ₂
10	Me	Me	H	H	0	H	H	H	SO ₂ CHF ₂	OCHF ₂
	Me	Me	H	H	0	H	H	H	NH ₂	OCHF ₂
	Me	Me	H	H	0	H	H	H	NHMe	OCHF ₂
	Me	Me	H	H	0	H	H	H	NMe ₂	OCHF ₂
15	Me	Me	H	H	0	H	H	H	NHEt	OCHF ₂
	Me	Me	H	H	0	H	H	H	NEt ₂	OCHF ₂
	Me	Me	H	H	0	H	H	H	NHPh	OCHF ₂
	Me	Me	H	H	0	H	H	H	N(Me)Ph	OCHF ₂
20	Me	Me	H	H	0	H	H	H	CN	OCHF ₂
	Me	Me	H	H	0	H	H	Me	F	CF ₃
	Me	Me	H	H	0	H	H	Me	Cl	CF ₃
	Me	Me	H	H	0	H	H	Me	OH	CF ₃
25	Me	Me	H	H	0	H	H	Me	OMe	CF ₃
	Me	Me	H	H	0	H	H	Me	OE <i>t</i>	CF ₃
	Me	Me	H	H	0	H	H	Me	OP <i>r</i> -i	CF ₃
	Me	Me	H	H	0	H	H	Me	OP <i>r</i>	CF ₃
30	Me	Me	H	H	0	H	H	Me	OBu- <i>t</i>	CF ₃
	Me	Me	H	H	0	H	H	Me	OCH ₂ Pr- <i>c</i>	CF ₃
	Me	Me	H	H	0	H	H	Me	OCH ₂ Bu- <i>c</i>	CF ₃
	Me	Me	H	H	0	H	H	Me	OCH ₂ Pen- <i>c</i>	CF ₃
	Me	Me	H	H	0	H	H	Me	OCH ₂ Hex- <i>c</i>	CF ₃
35	Me	Me	H	H	0	H	H	Me	OPen- <i>c</i>	CF ₃
	Me	Me	H	H	0	H	H	Me	OHex- <i>c</i>	CF ₃
	Me	Me	H	H	0	H	H	Me	OCH ₂ Ph	CF ₃
	Me	Me	H	H	0	H	H	Me	OPh	CF ₃
40	Me	Me	H	H	0	H	H	Me	OCHF ₂	CF ₃
	Me	Me	H	H	0	H	H	Me	SH	CF ₃
	Me	Me	H	H	0	H	H	Me	SMe	CF ₃
	Me	Me	H	H	0	H	H	Me	SO ₂ Me	CF ₃
45	Me	Me	H	H	0	H	H	Me	SE <i>t</i>	CF ₃
	Me	Me	H	H	0	H	H	Me	SO ₂ Et	CF ₃
	Me	Me	H	H	0	H	H	Me	SP <i>r</i> -i	CF ₃
	Me	Me	H	H	0	H	H	Me	SO ₂ Pr-i	CF ₃
50	Me	Me	H	H	0	H	H	Me	SPh	CF ₃
	Me	Me	H	H	0	H	H	Me	SO ₂ Ph	CF ₃
	Me	Me	H	H	0	H	H	Me	SCHF ₂	CF ₃
	Me	Me	H	H	0	H	H	Me	SO ₂ CHF ₂	CF ₃
55	Me	Me	H	H	0	H	H	Me	NH ₂	CF ₃
	Me	Me	H	H	0	H	H	Me	NHMe	CF ₃
	Me	Me	H	H	0	H	H	Me	NMe ₂	CF ₃
	Me	Me	H	H	0	H	H	Me	NHEt	CF ₃

	Me	Me	H	H	0	H	H	Me	NEt ₂	CF ₃
	Me	Me	H	H	0	H	H	Me	NHPh	CF ₃
5	Me	Me	H	H	0	H	H	Me	N(Me)Ph	CF ₃
	Me	Me	H	H	0	H	H	Me	CN	CF ₃
	Me	Me	H	H	0	H	H	OMe	F	CF ₃
	Me	Me	H	H	0	H	H	OMe	Cl	CF ₃
10	Me	Me	H	H	0	H	H	OMe	OH	CF ₃
	Me	Me	H	H	0	H	H	OMe	OMe	CF ₃
	Me	Me	H	H	0	H	H	OMe	OEt	CF ₃
	Me	Me	H	H	0	H	H	OMe	OPr-i	CF ₃
15	Me	Me	H	H	0	H	H	OMe	OPr	CF ₃
	Me	Me	H	H	0	H	H	OMe	OBu-t	CF ₃
	Me	Me	H	H	0	H	H	OMe	OCH ₂ Pr-c	CF ₃
	Me	Me	H	H	0	H	H	OMe	OCH ₂ Bu-c	CF ₃
	Me	Me	H	H	0	H	H	OMe	OCH ₂ Pen-c	CF ₃
20	Me	Me	H	H	0	H	H	OMe	OCH ₂ Hex-c	CF ₃
	Me	Me	H	H	0	H	H	OMe	OPen-c	CF ₃
	Me	Me	H	H	0	H	H	OMe	OHex-c	CF ₃
	Me	Me	H	H	0	H	H	OMe	OCH ₂ Ph	CF ₃
25	Me	Me	H	H	0	H	H	OMe	OPh	CF ₃
	Me	Me	H	H	0	H	H	OMe	OCHF ₂	CF ₃
	Me	Me	H	H	0	H	H	OMe	SH	CF ₃
	Me	Me	H	H	0	H	H	OMe	SMe	CF ₃
	Me	Me	H	H	0	H	H	OMe	SO ₂ Me	CF ₃
30	Me	Me	H	H	0	H	H	OMe	SEt	CF ₃
	Me	Me	H	H	0	H	H	OMe	SO ₂ Et	CF ₃
	Me	Me	H	H	0	H	H	OMe	SPr-i	CF ₃
	Me	Me	H	H	0	H	H	OMe	SO ₂ Pr-i	CF ₃
35	Me	Me	H	H	0	H	H	OMe	SPh	CF ₃
	Me	Me	H	H	0	H	H	OMe	SO ₂ Ph	CF ₃
	Me	Me	H	H	0	H	H	OMe	SCHF ₂	CF ₃
	Me	Me	H	H	0	H	H	OMe	SO ₂ CHF ₂	CF ₃
40	Me	Me	H	H	0	H	H	OMe	NH ₂	CF ₃
	Me	Me	H	H	0	H	H	OMe	NHMe	CF ₃
	Me	Me	H	H	0	H	H	OMe	NMe ₂	CF ₃
	Me	Me	H	H	0	H	H	OMe	NHEt	CF ₃
45	Me	Me	H	H	0	H	H	OMe	NEt ₂	CF ₃
	Me	Me	H	H	0	H	H	OMe	NHPh	CF ₃
	Me	Me	H	H	0	H	H	OMe	N(Me)Ph	CF ₃
	Me	Me	H	H	0	H	H	OMe	CN	CF ₃
50	Me	Me	H	H	0	H	H	SMe	F	CF ₃
	Me	Me	H	H	0	H	H	SMe	Cl	CF ₃
	Me	Me	H	H	0	H	H	SMe	OH	CF ₃
	Me	Me	H	H	0	H	H	SMe	OMe	CF ₃
	Me	Me	H	H	0	H	H	SMe	OEt	CF ₃
55	Me	Me	H	H	0	H	H	SMe	OPr-i	CF ₃
	Me	Me	H	H	0	H	H	SMe	OPr	CF ₃

	Me	Me	H	H	0	H	H	SMe	OBu-t	CF ₃
	Me	Me	H	H	0	H	H	SMe	OCH ₂ Pr-c	CF ₃
5	Me	Me	H	H	0	H	H	SMe	OCH ₂ Bu-c	CF ₃
	Me	Me	H	H	0	H	H	SMe	OCH ₂ Pen-c	CF ₃
	Me	Me	H	H	0	H	H	SMe	OCH ₂ Hex-c	CF ₃
	Me	Me	H	H	0	H	H	SMe	OPen-c	CF ₃
10	Me	Me	H	H	0	H	H	SMe	OHex-c	CF ₃
	Me	Me	H	H	0	H	H	SMe	OCH ₂ Ph	CF ₃
	Me	Me	H	H	0	H	H	SMe	OPh	CF ₃
	Me	Me	H	H	0	H	H	SMe	OCHF ₂	CF ₃
	Me	Me	H	H	0	H	H	SMe	SH	CF ₃
15	Me	Me	H	H	0	H	H	SMe	SMe	CF ₃
	Me	Me	H	H	0	H	H	SMe	SO ₂ Me	CF ₃
	Me	Me	H	H	0	H	H	SMe	SEt	CF ₃
	Me	Me	H	H	0	H	H	SMe	SO ₂ Et	CF ₃
20	Me	Me	H	H	0	H	H	SMe	SPr-i	CF ₃
	Me	Me	H	H	0	H	H	SMe	SO ₂ Pr-i	CF ₃
	Me	Me	H	H	0	H	H	SMe	SPh	CF ₃
	Me	Me	H	H	0	H	H	SMe	SO ₂ Ph	CF ₃
25	Me	Me	H	H	0	H	H	SMe	SCHF ₂	CF ₃
	Me	Me	H	H	0	H	H	SMe	SO ₂ CHF ₂	CF ₃
	Me	Me	H	H	0	H	H	SMe	NH ₂	CF ₃
	Me	Me	H	H	0	H	H	SMe	NHMe	CF ₃
	Me	Me	H	H	0	H	H	SMe	NMe ₂	CF ₃
30	Me	Me	H	H	0	H	H	SMe	NHEt	CF ₃
	Me	Me	H	H	0	H	H	SMe	NEt ₂	CF ₃
	Me	Me	H	H	0	H	H	SMe	NHPh	CF ₃
	Me	Me	H	H	0	H	H	SMe	N(Me)Ph	CF ₃
35	Me	Me	H	H	0	H	H	SMe	CN	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	F	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	Cl	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	OH	CF ₃
40	Me	Me	H	H	0	H	H	SO ₂ Me	OMe	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	OEt	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	OPr-i	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	OPr	CF ₃
45	Me	Me	H	H	0	H	H	SO ₂ Me	OBu-t	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	OCH ₂ Pr-c	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	OCH ₂ Bu-c	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	OCH ₂ Pen-c	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	OCH ₂ Hex-c	CF ₃
50	Me	Me	H	H	0	H	H	SO ₂ Me	OPen-c	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	OHex-c	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	OCH ₂ Ph	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	OPh	CF ₃
55	Me	Me	H	H	0	H	H	SO ₂ Me	OCHF ₂	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	SH	CF ₃

	Me	Me	H	H	0	H	H	SO ₂ Me	SMe	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	SO ₂ Me	CF ₃
5	Me	Me	H	H	0	H	H	SO ₂ Me	SEt	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	SO ₂ Et	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	SPr-i	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	SO ₂ Pr-i	CF ₃
10	Me	Me	H	H	0	H	H	SO ₂ Me	SPh	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	SO ₂ Ph	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	SCHF ₂	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	SO ₂ CHF ₂	CF ₃
15	Me	Me	H	H	0	H	H	SO ₂ Me	NH ₂	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	NHMe	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	NMe ₂	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	NHEt	CF ₃
20	Me	Me	H	H	0	H	H	SO ₂ Me	NEt ₂	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	NHPh	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	N(Me)Ph	CF ₃
	Me	Me	H	H	0	H	H	SO ₂ Me	CN	CF ₃
25	Me	Me	H	H	0	H	H	NH ₂	F	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	Cl	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	OH	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	OMe	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	OEt	CF ₃
30	Me	Me	H	H	0	H	H	NH ₂	OPr-i	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	OPr	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	OBu-t	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	OCH ₂ Pr-c	CF ₃
35	Me	Me	H	H	0	H	H	NH ₂	OCH ₂ Bu-c	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	OCH ₂ Pen-c	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	OCH ₂ Hex-c	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	OPen-c	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	OHex-c	CF ₃
40	Me	Me	H	H	0	H	H	NH ₂	OCH ₂ Ph	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	OPh	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	OCHF ₂	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	SH	CF ₃
45	Me	Me	H	H	0	H	H	NH ₂	SMe	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	SO ₂ Me	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	SEt	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	SO ₂ Et	CF ₃
50	Me	Me	H	H	0	H	H	NH ₂	SPr-i	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	SO ₂ Pr-i	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	SPh	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	SO ₂ Ph	CF ₃
55	Me	Me	H	H	0	H	H	NH ₂	SCHF ₂	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	SO ₂ CHF ₂	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	NH ₂	CF ₃

5	Me	Me	H	H	0	H	H	NH ₂	NHMe	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	NMe ₂	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	NHEt	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	NEt ₂	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	NHPh	CF ₃
	Me	Me	H	H	0	H	H	NH ₂	N(Me)Ph	CF ₃
10	Me	Me	H	H	0	H	H	NH ₂	CN	CF ₃
	H	H	H	H	0	H	H	H	OMe	CF ₃
	H	H	H	H	0	H	H	H	OEt	CF ₃
	Me	H	H	H	0	H	H	H	OMe	CF ₃
15	Me	H	H	H	0	H	H	H	OEt	CF ₃
	Me	H	Me	H	0	H	H	H	OMe	CF ₃
	Me	H	Me	H	0	H	H	H	OEt	CF ₃
	Me	Me	H	H	0	Me	H	H	OMe	CF ₃
	Me	Me	H	H	0	Me	H	H	OEt	CF ₃
20	Me	Me	H	H	0	Et	H	H	OMe	CF ₃
	Me	Me	H	H	0	Et	H	H	OEt	CF ₃
	Me	Me	H	H	0	Pr-i	H	H	H	CF ₃
	Me	Me	H	H	0	Pr-i	H	H	OMe	CF ₃
25	Me	Me	H	H	0	Pr-i	H	H	OEt	CF ₃
	Me	Me	H	H	0	Me	Me	H	OMe	CF ₃
	Me	Me	H	H	0	Me	Me	H	OEt	CF ₃
	Me	Et	H	H	0	H	H	H	OMe	CF ₃
	Me	Et	H	H	0	H	H	H	OEt	CF ₃
30	Et	Et	H	H	0	H	H	H	OMe	CF ₃
	Et	Et	H	H	0	H	H	H	OEt	CF ₃
	Me	Pr-i	H	H	0	H	H	H	OMe	CF ₃
	Me	Pr-i	H	H	0	H	H	H	OEt	CF ₃
35	Me	Pr	H	H	0	H	H	H	OMe	CF ₃
	Me	Pr	H	H	0	H	H	H	OEt	CF ₃
	Me	Pr-c	H	H	0	H	H	H	OMe	CF ₃
	Me	Pr-c	H	H	0	H	H	H	OEt	CF ₃
40	Me	CH ₂ Pr-c	H	H	0	H	H	H	OMe	CF ₃
	Me	CH ₂ Pr-c	H	H	0	H	H	H	OEt	CF ₃
	-(CH ₂) ₂ -		H	H	0	H	H	H	OMe	CF ₃
	-(CH ₂) ₂ -		H	H	0	H	H	H	OEt	CF ₃
45	-(CH ₂) ₃ -		H	H	0	H	H	H	OMe	CF ₃
	-(CH ₂) ₃ -		H	H	0	H	H	H	OEt	CF ₃
	-(CH ₂) ₄ -		H	H	0	H	H	H	OMe	CF ₃
	-(CH ₂) ₄ -		H	H	0	H	H	H	OEt	CF ₃
50	-(CH ₂) ₅ -		H	H	0	H	H	H	OMe	CF ₃
	-(CH ₂) ₅ -		H	H	0	H	H	H	OEt	CF ₃
	H	-(CH ₂) ₃ -	H	0	H	H	H	H	OMe	CF ₃
	H	-(CH ₂) ₃ -	H	0	H	H	H	H	OEt	CF ₃
	H	-(CH ₂) ₄ -	H	0	H	H	H	H	OMe	CF ₃
55	H	-(CH ₂) ₄ -	H	0	H	H	H	H	OEt	CF ₃
	H	-(CH ₂) ₅ -	H	0	H	H	H	H	OMe	CF ₃

H	-(CH ₂) ₅ -	H	0	H	H	H	· OEt	CF ₃
H	-(CH ₂) ₆ -	H	0	H	H	H	OMe	CF ₃
H	-(CH ₂) ₆ -	H	0	H	H	H	· OEt	CF ₃

5

10

15

20

25

30

35


40

45

50

55

Table 9



R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Y ¹
Me	Me	H	H	2	H	H	Pyridin-2-yl
Me	Me	H	H	2	H	H	Pyridin-2-yl 1-oxide
Me	Me	H	H	2	H	H	Pyridin-4-yl
Me	Me	H	H	2	H	H	Pyridin-4-yl 1-oxide
Me	Me	H	H	2	H	H	1,2,4-Oxadiazol-3-yl
Me	Me	H	H	2	H	H	3-Phenyl-1,2,4-oxadiazol-5-yl
Me	Me	H	H	2	H	H	3-Benzyl-1,2,4-oxadiazol-5-yl
Me	Me	H	H	2	H	H	2-Chlorothiazol-4-yl
Me	Me	H	H	2	H	H	5-Trifluoromethyl-1,3,4-thiadiazol-2-yl
Me	Me	H	H	2	H	H	1,4-Dimethylimidazol-5-yl
Me	Me	H	H	2	H	H	1-Phenyl-4-methoxycarbonyl-1,2,3-triazol-5-yl
Me	Me	H	H	2	H	H	1-Difluoromethyl-1,2,4-triazol-3-yl
Me	Me	H	H	2	H	H	1-Difluoromethyl-1,2,4-triazol-5-yl
Me	Me	H	H	2	H	H	4-Difluoromethyl-1,2,4-triazol-3-yl
Me	Me	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
Me	Me	H	H	2	H	H	4,6-Diethoxypyrimidin-2-yl
Me	Me	H	H	2	H	H	4,6-Dimethylpyrimidin-2-yl
Me	Me	H	H	2	H	H	4-Chloro-6-methylpyrimidin-2-yl
Me	Me	H	H	2	H	H	4-Methoxy-6-methylpyrimidin-2-yl
Me	Me	H	H	2	H	H	4-Difluoromethoxy-6-methylpyrimidin-2-yl
Me	Me	H	H	2	H	H	4-Phenoxy-6-methylpyrimidin-2-yl
Me	Me	H	H	2	H	H	4-Chloro-6-trifluoromethylpyrimidin-2-yl
Me	Me	H	H	2	H	H	4-Methoxy-6-trifluoromethylpyrimidin-2-yl
Me	Me	H	H	2	H	H	4-Difluoromethoxy-6-trifluoromethylpyrimidin-2-yl
Me	Me	H	H	2	H	H	4-Phenoxy-6-trifluoromethylpyrimidin-2-yl
H	H	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
Me	H	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
Me	H	Me	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
Me	Me	H	H	2	Me	H	4,6-Dimethoxypyrimidin-2-yl
Me	Me	H	H	2	Et	H	4,6-Dimethoxypyrimidin-2-yl
Me	Me	H	H	2	Pr-i	H	4,6-Dimethoxypyrimidin-2-yl
Me	Me	H	H	2	Me	Me	4,6-Dimethoxypyrimidin-2-yl
Me	Et	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
Et	Et	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl

	Me	Pr-i	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
	Me	Pr	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
5	Me	Pr-c	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
	Me	CH ₂ Pr-c	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
		-(CH ₂) ₂ -	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
		-(CH ₂) ₃ -	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
10		-(CH ₂) ₄ -	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
		-(CH ₂) ₅ -	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
	H	-(CH ₂) ₃ -	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
	H	-(CH ₂) ₄ -	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
15	H	-(CH ₂) ₅ -	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
	H	-(CH ₂) ₆ -	H	H	2	H	H	4,6-Dimethoxypyrimidin-2-yl
	Me	Me	H	H	1	H	H	Pyridin-2-yl
	Me	Me	H	H	1	H	H	Pyridin-2-yl 1-oxide
	Me	Me	H	H	1	H	H	Pyridin-4-yl
20	Me	Me	H	H	1	H	H	Pyridin-4-yl 1-oxide
	Me	Me	H	H	1	H	H	1,2,4-Oxadiazol-3-yl
	Me	Me	H	H	1	H	H	3-Phenyl-1,2,4-oxadiazol-5-yl
	Me	Me	H	H	1	H	H	3-Benzyl-1,2,4-oxadiazol-5-yl
25	Me	Me	H	H	1	H	H	2-Chlorothiazol-4-yl
	Me	Me	H	H	1	H	H	5-Trifluoromethyl-1,3,4-thiadiazol-2-yl
	Me	Me	H	H	1	H	H	1,4-Dimethylimidazol-5-yl
	Me	Me	H	H	1	H	H	1-Phenyl-4-methoxycarbonyl-1,2,3-triazol-5-yl
	Me	Me	H	H	1	H	H	1-Difluoromethyl-1,2,4-triazol-3-yl
30	Me	Me	H	H	1	H	H	1-Difluoromethyl-1,2,4-triazol-5-yl
	Me	Me	H	H	1	H	H	4-Difluoromethyl-1,2,4-triazol-3-yl
	Me	Me	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
	Me	Me	H	H	1	H	H	4,6-Diethoxypyrimidin-2-yl
35	Me	Me	H	H	1	H	H	4,6-Dimethylpyrimidin-2-yl
	Me	Me	H	H	1	H	H	4-Chloro-6-methylpyrimidin-2-yl
	Me	Me	H	H	1	H	H	4-Methoxy-6-methylpyrimidin-2-yl
	Me	Me	H	H	1	H	H	4-Difluoromethoxy-6-methylpyrimidin-2-yl
	Me	Me	H	H	1	H	H	4-Phenoxy-6-methylpyrimidin-2-yl
40	Me	Me	H	H	1	H	H	4-Chloro-6-trifluoromethylpyrimidin-2-yl
	Me	Me	H	H	1	H	H	4-Methoxy-6-trifluoromethylpyrimidin-2-yl
	Me	Me	H	H	1	H	H	4-Difluoromethoxy-6-trifluoromethylpyrimidin-2-yl
	Me	Me	H	H	1	H	H	4-Phenoxy-6-trifluoromethylpyrimidin-2-yl
45	H	H	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
	Me	H	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
	Me	H	Me	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
	Me	Me	H	H	1	Me	H	4,6-Dimethoxypyrimidin-2-yl
50	Me	Me	H	H	1	Et	H	4,6-Dimethoxypyrimidin-2-yl
	Me	Me	H	H	1	Pr-i	H	4,6-Dimethoxypyrimidin-2-yl
	Me	Me	H	H	1	Me	Me	4,6-Dimethoxypyrimidin-2-yl
	Me	Et	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
	Et	Et	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
55	Me	Pr-i	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl

	Me	Pr	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
5	Me	Pr-c	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
	Me	CH ₂ Pr-c	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
		-(CH ₂) ₂ -	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
		-(CH ₂) ₃ -	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
10		-(CH ₂) ₄ -	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
		-(CH ₂) ₅ -	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
	H	-(CH ₂) ₃ -	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
	H	-(CH ₂) ₄ -	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
	H	-(CH ₂) ₅ -	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
15	H	-(CH ₂) ₆ -	H	H	1	H	H	4,6-Dimethoxypyrimidin-2-yl
	Me	Me	H	H	0	H	H	Pyridin-2-yl
	Me	Me	H	H	0	H	H	Pyridin-2-yl 1-oxide
	Me	Me	H	H	0	H	H	Pyridin-4-yl
20	Me	Me	H	H	0	H	H	Pyridin-4-yl 1-oxide
	Me	Me	H	H	0	H	H	1,2,4-Oxadiazol-3-yl
	Me	Me	H	H	0	H	H	3-Phenyl-1,2,4-oxadiazol-5-yl
	Me	Me	H	H	0	H	H	3-Benzyl-1,2,4-oxadiazol-5-yl
	Me	Me	H	H	0	H	H	2-Chlorothiazol-4-yl
25	Me	Me	H	H	0	H	H	5-Trifluoromethyl-1,3,4-thiadiazol-2-yl
	Me	Me	H	H	0	H	H	1,4-Dimethylimidazol-5-yl
	Me	Me	H	H	0	H	H	1-Phenyl-4-methoxycarbonyl-1,2,3-triazol-5-yl
	Me	Me	H	H	0	H	H	1-Difluoromethyl-1,2,4-triazol-3-yl
30	Me	Me	H	H	0	H	H	1-Difluoromethyl-1,2,4-triazol-5-yl
	Me	Me	H	H	0	H	H	4-Difluoromethyl-1,2,4-triazol-3-yl
	Me	Me	H	H	0	H	H	4,6-Dimethoxypyrimidin-2-yl
	Me	Me	H	H	0	H	H	4,6-Diethoxypyrimidin-2-yl
	Me	Me	H	H	0	H	H	4,6-Dimethylpyrimidin-2-yl
35	Me	Me	H	H	0	H	H	4-Chloro-6-methylpyrimidin-2-yl
	Me	Me	H	H	0	H	H	4-Methoxy-6-methylpyrimidin-2-yl
	Me	Me	H	H	0	H	H	4-Difluoromethoxy-6-methylpyrimidin-2-yl
	Me	Me	H	H	0	H	H	4-Phenoxy-6-methylpyrimidin-2-yl
40	Me	Me	H	H	0	H	H	4-Chloro-6-trifluoromethylpyrimidin-2-yl
	Me	Me	H	H	0	H	H	4-Methoxy-6-trifluoromethylpyrimidin-2-yl
	Me	Me	H	H	0	H	H	4-Difluoromethoxy-6-trifluoromethylpyrimidin-2-yl
	Me	Me	H	H	0	H	H	4-Phenoxy-6-trifluoromethylpyrimidin-2-yl
45	H	H	H	H	0	H	H	4,6-Dimethoxypyrimidin-2-yl
	Me	H	H	H	0	H	H	4,6-Dimethoxypyrimidin-2-yl
	Me	H	Me	H	0	H	H	4,6-Dimethoxypyrimidin-2-yl
	Me	Me	H	H	0	Me	H	4,6-Dimethoxypyrimidin-2-yl
	Me	Me	H	H	0	Et	H	4,6-Dimethoxypyrimidin-2-yl
50	Me	Me	H	H	0	Pr-i	H	4,6-Dimethoxypyrimidin-2-yl
	Me	Me	H	H	0	Me	Me	4,6-Dimethoxypyrimidin-2-yl
	Me	Et	H	H	0	H	H	4,6-Dimethoxypyrimidin-2-yl
	Et	Et	H	H	0	H	H	4,6-Dimethoxypyrimidin-2-yl
	Me	Pr-i	H	H	0	H	H	4,6-Dimethoxypyrimidin-2-yl
55	Me	Pr	H	H	0	H	H	4,6-Dimethoxypyrimidin-2-yl

	Me	Pr-c	H	H	0	H	H	4,6-Dimethoxypyrimidin-2-yl
5	Me	CH ₂ Pr-c	H	H	0	H	H	4,6-Dimethoxypyrimidin-2-yl
		-(CH ₂) ₂ -	H	H	0	H	H	4,6-Dimethoxypyrimidin-2-yl
		-(CH ₂) ₃ -	H	H	0	H	H	4,6-Dimethoxypyrimidin-2-yl
		-(CH ₂) ₄ -	H	H	0	H	H	4,6-Dimethoxypyrimidin-2-yl
10		-(CH ₂) ₅ -	H	H	0	H	H	4,6-Dimethoxypyrimidin-2-yl
	H	-(CH ₂) ₃ -	H	0	H	H	H	4,6-Dimethoxypyrimidin-2-yl
	H	-(CH ₂) ₄ -	H	0	H	H	H	4,6-Dimethoxypyrimidin-2-yl
	H	-(CH ₂) ₅ -	H	0	H	H	H	4,6-Dimethoxypyrimidin-2-yl
15	H	-(CH ₂) ₆ -	H	0	H	H	H	4,6-Dimethoxypyrimidin-2-yl
	Me	Et	H	H	2	H	H	Pirrol-1-yl
	Me	Et	H	H	2	H	H	Oxazol-2-yl
	Me	Et	H	H	2	H	H	Thiazol-2-yl
20	Me	Et	H	H	2	H	H	Thiazol-4-yl
	Me	Et	H	H	2	H	H	1,2,3-Thiadiazol-4-yl
	Me	Et	H	H	2	H	H	1,2,3-Thiadiazol-5-yl
	Me	Et	H	H	2	H	H	1,2,4-Thiadiazol-3-yl
25	Me	Et	H	H	2	H	H	1,2,4-Thiadiazol-5-yl
	Me	Et	H	H	2	H	H	1,3,4-Thiadiazol-2-yl
	Me	Et	H	H	2	H	H	1,3,4-Thiadiazol-5-yl
	Me	Et	H	H	2	H	H	Pyridin-2-yl
30	Me	Et	H	H	2	H	H	Pyridin-3-yl
	Me	Et	H	H	2	H	H	Pyridin-4-yl
	Me	Et	H	H	2	H	H	1H-Imidazol-2-yl
	Me	Et	H	H	2	H	H	1H-Imidazol-4-yl
35	Me	Et	H	H	2	H	H	1H-Imidazol-5-yl
	Me	Et	H	H	2	H	H	1H-1,3,4-Triazol-2-yl
	Me	Et	H	H	2	H	H	1H-1,3,4-Triazol-5-yl

Table 10


$ \begin{array}{c} \text{R}^2 \quad \text{R}^3 \\ \diagup \quad \diagdown \\ \text{R}^1 - \text{C} - \text{C} - \text{R}^4 \\ \diagdown \quad \diagup \\ \text{O} \quad \text{N} \end{array} \text{S(O)}_n - \text{C} \begin{array}{l} \text{R}^6 \\ \text{Y}^1 \\ \text{R}^5 \end{array} $							
R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Y ¹
Me	Me	H	H	2	H	H	Benzimidazol-2-yl
Me	Me	H	H	2	H	H	Benzothiophen-2-yl
Me	Me	H	H	2	H	H	3-Chlorobenzothiophen-2-yl
Me	Me	H	H	2	H	H	Benzotriazol-1-yl
Me	Me	H	H	2	H	H	1-Methylindazol-4-yl
Me	Me	H	H	2	H	H	Benzothiazol-2-yl
Me	Me	H	H	2	H	H	Benzothiophen-3-yl
Me	Me	H	H	2	H	H	5-Chlorobenzothiophen-3-yl

	Me	Me	H	H	2	H	H	Benzoxazol-2-yl
5	Me	Me	H	H	2	H	H	3-Methylbenzothiophen-2-yl
	Me	Me	H	H	2	H	H	3-Bromobenzothiophen-2-yl
	Me	Me	H	H	2	H	H	Benzofuran-2-yl
	Me	Me	H	H	2	H	H	2-Methylbenzofuran-7-yl
10	Me	Me	H	H	2	H	H	3-Bromobenzofuran-2-yl
	Me	Me	H	H	2	H	H	Benzothiophen-7-yl
	Me	Me	H	H	2	H	H	1-Methylindazol-7-yl
	Me	Me	H	H	2	H	H	1-Difluoromethylindazol-7-yl
	Me	Me	H	H	2	H	H	3-Methylbenzofuran-2-yl
15	Me	Me	H	H	2	H	H	3-Chloro-1-methylindol-2-yl
	Me	Me	H	H	1	H	H	Benzimidazol-2-yl
	Me	Me	H	H	1	H	H	Benzothiophen-2-yl
	Me	Me	H	H	1	H	H	3-Chlorobenzothiophen-2-yl
20	Me	Me	H	H	1	H	H	Benzotriazol-1-yl
	Me	Me	H	H	1	H	H	1-Methylindazol-4-yl
	Me	Me	H	H	1	H	H	Benzothiazol-2-yl
	Me	Me	H	H	1	H	H	Benzothiophen-3-yl
	Me	Me	H	H	1	H	H	5-Chlorobenzothiophen-3-yl
25	Me	Me	H	H	1	H	H	Benzoxazol-2-yl
	Me	Me	H	H	1	H	H	3-Methylbenzothiophen-2-yl
	Me	Me	H	H	1	H	H	3-Bromobenzothiophen-2-yl
	Me	Me	H	H	1	H	H	Benzofuran-2-yl
30	Me	Me	H	H	1	H	H	2-Methylbenzofuran-7-yl
	Me	Me	H	H	1	H	H	3-Bromobenzofuran-2-yl
	Me	Me	H	H	1	H	H	Benzothiophen-7-yl
	Me	Me	H	H	1	H	H	1-Methylindazol-7-yl
	Me	Me	H	H	1	H	H	3-Methylbenzofuran-2-yl
35	Me	Me	H	H	1	H	H	3-Chloro-1-methylindol-2-yl
	Me	Me	H	H	0	H	H	Benzimidazol-2-yl
	Me	Me	H	H	0	H	H	Benzothiophen-2-yl
	Me	Me	H	H	0	H	H	3-Chlorobenzothiophen-2-yl
40	Me	Me	H	H	0	H	H	Benzotriazol-1-yl
	Me	Me	H	H	0	H	H	1-Methylindazol-4-yl
	Me	Me	H	H	0	H	H	Benzothiazol-2-yl
	Me	Me	H	H	0	H	H	Benzothiophen-3-yl
	Me	Me	H	H	0	H	H	5-Chlorobenzothiophen-3-yl
45	Me	Me	H	H	0	H	H	Benzoxazol-2-yl
	Me	Me	H	H	0	H	H	3-Methylbenzothiophen-2-yl
	Me	Me	H	H	0	H	H	3-Bromobenzothiophen-2-yl
	Me	Me	H	H	0	H	H	Benzofuran-2-yl
50	Me	Me	H	H	0	H	H	2-Methylbenzofuran-7-yl
	Me	Me	H	H	0	H	H	3-Bromobenzofuran-2-yl
	Me	Me	H	H	0	H	H	Benzothiophen-7-yl
	Me	Me	H	H	0	H	H	1-Methylindazol-7-yl
	Me	Me	H	H	0	H	H	3-Methylbenzofuran-2-yl
55	Me	Me	H	H	0	H	H	3-Chloro-1-methylindol-2-yl

	Me	Et	H	H	2	H	H	Benzoxazol-2-yl
	Me	Et	H	H	2	H	H	4-Chlorobenzoxazol-2-yl
5	Me	Et	H	H	2	H	H	5-Chlorobenzoxazol-2-yl
	Me	Et	H	H	2	H	H	6-Chlorobenzoxazol-2-yl
	Me	Et	H	H	2	H	H	7-Chlorobenzoxazol-2-yl
	Me	Et	H	H	2	H	H	4-Fluorobenzoxazol-2-yl
	Me	Et	H	H	2	H	H	5-Fluorobenzoxazol-2-yl
10	Me	Et	H	H	2	H	H	6-Fluorobenzoxazol-2-yl
	Me	Et	H	H	2	H	H	7-Fluorobenzoxazol-2-yl
	Me	Et	H	H	2	H	H	4-Methylbenzoxazol-2-yl
	Me	Et	H	H	2	H	H	5-Methylbenzoxazol-2-yl
15	Me	Et	H	H	2	H	H	6-Methylbenzoxazol-2-yl
	Me	Et	H	H	2	H	H	7-Methylbenzoxazol-2-yl
	Me	Et	H	H	2	H	H	4-Methoxybenzoxazol-2-yl
	Me	Et	H	H	2	H	H	5-Methoxybenzoxazol-2-yl
20	Me	Et	H	H	2	H	H	6-Methoxybenzoxazol-2-yl
	Me	Et	H	H	2	H	H	7-Methoxybenzoxazol-2-yl
	Me	Et	H	H	2	H	H	Benzothiazol-2-yl
	Me	Et	H	H	2	H	H	4-Chlorobenzothiazol-2-yl
	Me	Et	H	H	2	H	H	5-Chlorobenzothiazol-2-yl
25	Me	Et	H	H	2	H	H	6-Chlorobenzothiazol-2-yl
	Me	Et	H	H	2	H	H	7-Chlorobenzothiazol-2-yl
	Me	Et	H	H	2	H	H	4-Fluorobenzothiazol-2-yl
	Me	Et	H	H	2	H	H	5-Fluorobenzothiazol-2-yl
30	Me	Et	H	H	2	H	H	6-Fluorobenzothiazol-2-yl
	Me	Et	H	H	2	H	H	7-Fluorobenzothiazol-2-yl
	Me	Et	H	H	2	H	H	4-Methylbenzothiazol-2-yl
	Me	Et	H	H	2	H	H	5-Methylbenzothiazol-2-yl
	Me	Et	H	H	2	H	H	6-Methylbenzothiazol-2-yl
35	Me	Et	H	H	2	H	H	7-Methylbenzothiazol-2-yl
	Me	Et	H	H	2	H	H	4-Methoxybenzothiazol-2-yl
	Me	Et	H	H	2	H	H	5-Methoxybenzothiazol-2-yl
	Me	Et	H	H	2	H	H	6-Methoxybenzothiazol-2-yl
40	Me	Et	H	H	2	H	H	7-Methoxybenzothiazol-2-yl
	Me	Et	H	H	2	H	H	Qnolin-2-yl
	Me	Et	H	H	2	H	H	Qinolin-6-yl
	Me	Et	H	H	2	H	H	Quinoxalin-2-yl
	Me	Et	H	H	2	H	H	Benzofuran-2-yl
45	Me	Et	H	H	2	H	H	3-Chlorobenzofuran-2-yl
	Me	Et	H	H	2	H	H	4-Chlorobenzofuran-2-yl
	Me	Et	H	H	2	H	H	5-Chlorobenzofuran-2-yl
	Me	Et	H	H	2	H	H	6-Chlorobenzofuran-2-yl
50	Me	Et	H	H	2	H	H	7-Chlorobenzofuran-2-yl
	Me	Et	H	H	2	H	H	3-Methylbenzofuran-2-yl
	Me	Et	H	H	2	H	H	4-Methylbenzofuran-2-yl
	Me	Et	H	H	2	H	H	5-Methylbenzofuran-2-yl
55	Me	Et	H	H	2	H	H	6-Methylbenzofuran-2-yl

Me	Et	H	H	2	H	H	7-Methylbenzofuran-2-yl
Me	Et	H	H	2	H	H	3-Methoxybenzofuran-2-yl
Me	Et	H	H	2	H	H	4-Methoxybenzofuran-2-yl
Me	Et	H	H	2	H	H	5-Methoxybenzofuran-2-yl
Me	Et	H	H	2	H	H	6-Methoxybenzofuran-2-yl
Me	Et	H	H	2	H	H	7-Methoxybenzofuran-2-yl

[0046] The present compound represented by the general formula [I] can be produced according to the processes shown below; however, the compound can be produced also by other processes. <Production Process 1> Step 1 to Step 5



[0047] In the above production scheme, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and Y have the same definitions as given above; X^1 is a halogen atom; R^7 is a C1 to C4 alkyl group, an optionally substituted phenyl group or an optionally substituted benzyl group; L is a leaving group such as halogen atom, C1 to C4 alkylsulfonyl group, optionally substituted phenylsulfonyl group, optionally substituted benzylsulfonyl group or the like; and x is an integer of 1 or more.

[0048] The above production process is described below in detail on each step.

(Step 1)

[0049] A sulfide derivative represented by the general formula [5] can be produced by reacting a compound represented by the general formula [1] with a sodium hydrosulfide hydrate represented by the general formula [2] in the presence or absence of a solvent (preferably in an appropriate solvent) in the presence of a salt to produce a base of a mercaptan, represented by the general formula [3] in the reaction system, and then, without isolating the salt of a mercaptan [3], reacting the salt [3] with a halogen derivative represented by the general formula [4] [in this case, a radical-generating agent, for example, Rongalit (trade name) : $\text{CH}_2(\text{OH})\text{SO}_2\text{Na} \cdot 2\text{H}_2\text{O}$ may be added].

[0050] The reaction temperature in each reaction is any temperature between 0°C and the reflux temperature of each reaction system and is preferably 10 to 100°C . The reaction time varies depending upon the compounds used, but is 0.5 to 24 hours.

[0051] With respect to the amounts of the reagents used in each reaction, each of the compound represented by the general formula [2] and the compound represented by the general formula [4] is used in an amount of 1 to 3 equivalents relative to one equivalent of the compound represented by the general formula [1] and, when a base is used, the base is used in an amount of 0.5 to 3 equivalents.

[0052] As the solvent, there can be mentioned, for example, ethers such as dioxane, tetrahydrofuran (THF) and the like; halogenated hydrocarbons such as dichloroethane, carbon tetrachloride, chlorobenzene, dichlorobenzene and the like; amides such as N,N-dimethylacetamide, N,N-dimethylformamide, N-methyl-2-pyrrolidinone and the like; sulfur compounds such as dimethyl sulfoxide, sulfolane and the like; aromatic hydrocarbons such as benzene, toluene, xylene and the like; alcohols such as methanol, ethanol, propanol, isopropanol, butanol, tert-butanol and the like; ketones such as acetone, 2-butanone and the like; nitriles such as acetonitrile and the like; water; and mixtures thereof.

[0053] As the base, there can be mentioned, for example, metal hydrides such as sodium hydride and the like; alkali metal amides such as sodium amide, lithium diisopropylamide and the like; organic bases such as pyridine, triethylamine, 1,8-diazabicyclo[5.4.0]-7-undecene and the like; alkali metal hydroxides such as sodium hydroxide, potassium hydroxide and the like; alkaline earth metal hydroxides such as calcium hydroxide, magnesium hydroxide and the like; alkali metal carbonates such as sodium carbonate, potassium carbonate and the like; alkali metal hydrogencarbonates such as sodium hydrogencarbonate, potassium hydrogencarbonate and the like; and metal alcoholates such as sodium methoxide, potassium tertbutoxide and the like.

(Step 2)

[0054] A sulfoxide derivative represented by the general formula [6] can be produced by reacting the sulfide derivative represented by the general formula [5] with an oxidizing agent in an appropriate solvent.

[0055] The reaction temperature is any temperature between 0°C and the reflux temperature of the reaction system and is preferably 0 to 60°C . The reaction time varies depending upon the compounds used, but is 1 to 72 hours.

[0056] With respect to the amounts of the reagents used in the reaction, the oxidizing agent is used in an amount of 1 to 3 equivalents per equivalent of the compound represented by the general formula [5].

[0057] As the solvent, there can be mentioned, for example, halogenated hydrocarbons such as dichloromethane, chloroform, dichloroethane, carbon tetrachloride, chlorobenzene, dichlorobenzene and the like; ethers such as dioxane, tetrahydrofuran (THF), dimethoxyethane, diethyl ether and the like; amides such as N,N-dimethylacetamide, N,N-dimethylformamide, N-methyl-2-pyrrolidinone and the like; alcohols such as methanol, ethanol, propanol, isopropanol, butanol, tert-butanol and the like; ketones such as acetone, 2-butanone and the like; nitriles such as acetonitrile and the like; acetic acid; water; and mixtures thereof.

[0058] As the oxidizing agent, there can be mentioned, for example, organic peroxides such as m-chloroperbenzoic acid, performic acid, peracetic acid and the like; and inorganic peroxides such as hydrogen peroxide, potassium permanganate, sodium periodate and the like.

(Step 3)

[0059] A sulfone derivative represented by the general formula [7] can be produced by reacting the sulfoxide derivative represented by the general formula [6] with an oxidizing agent in an appropriate solvent.

[0060] The reaction temperature is any temperature between 0°C and the reflux temperature of the reaction system and is preferably 0 to 60°C . The reaction time varies depending upon the compounds used, but is 1 to 72 hours.

[0061] With respect to the amounts of the reagents used in the reaction, the oxidizing agent is used in an amount of 1 to 3 equivalents per equivalent of the compound represented by the general formula [6].

[0062] As the solvent and the oxidizing agent, there can be mentioned the same solvents and oxidizing agents as

in the step 2.

(Step 4)

[0063] The sulfone derivative represented by the general formula [7] can also be produced by reacting the sulfide derivative represented by the general formula [5] with an oxidizing agent of appropriate amount in an appropriate solvent without isolating the sulfoxide derivative represented by the general formula [6].

[0064] The reaction temperature is any temperature between 0°C and the reflux temperature of the reaction system and is preferably 0 to 60°C. The reaction time varies depending upon the compounds used, but is 1 to 72 hours.

[0065] With respect to the amounts of the reagents used in the reaction, the oxidizing agent is used in an amount of 1 to 3 equivalents per equivalent of the compound represented by the general formula [5].

[0066] As the solvent and the oxidizing agent, there can be mentioned the same solvents and oxidizing agents as in the step 2.

(Step 5)

[0067] The sulfide derivative represented by the general formula [5] can also be produced by reacting a compound represented by the general formula [8] with a mercaptan derivative represented by the general formula [9] in the presence or absence of a solvent (preferably in an appropriate solvent) in the presence of a base.

[0068] The reaction temperature is any temperature between 0°C and the reflux temperature of the reaction system and is preferably 10 to 100°C. The reaction time varies depending upon the compounds used, but is 0.5 to 24 hours.


[0069] With respect to the amounts of the reagents used in the reaction, the compound represented by the general formula [9] is used in an amount of 1 to 3 equivalents per equivalent of the compound represented by the general formula [8], and the base is used in an amount of 0.5 to 3 equivalents.

[0070] As the solvent, there can be mentioned, for example, ethers such as diethyl ether, dimethoxyethane, dioxane, tetrahydrofuran (THF) and the like; halogenated hydrocarbons such as dichloromethane, chloroform, carbon tetrachloride, dichloroethane, chlorobenzene, dichlorobenzene and the like; amides such as N,N-dimethylacetamide, N,N-dimethylformamide, N-methyl-2-pyrrolidinone and the like; sulfur compounds such as dimethyl sulfoxide, sulfolane and the like; aromatic hydrocarbons such as benzene, toluene, xylene and the like; alcohols such as methanol, ethanol, propanol, isopropanol, butanol, tert-butanol and the like; ketones such as acetone, 2-butanone and the like; nitriles such as acetonitrile and the like; water; and mixtures thereof.

[0071] As the base, there can be mentioned, for example, metal hydrides such as sodium hydride and the like; alkali metal amides such as sodium amide, lithium diisopropylamide and the like; organic bases such as pyridine, triethylamine, 1,8-diazabicyclo[5.4.0]-7-undecene and the like; alkali metal hydroxides such as sodium hydroxide, potassium hydroxide and the like; alkaline earth metal hydroxides such as calcium hydroxide, magnesium hydroxide and the like; alkali metal carbonates such as sodium carbonate, potassium carbonate and the like; alkali metal hydrogencarbonates such as sodium hydrogencarbonate, potassium hydrogencarbonate and the like; and metal alcoholates such as sodium methoxide, potassium tertbutoxide and the like.

[0072] A compound of the general formula [8] wherein L is a halogen atom, i.e. a compound [12] can be produced by a process shown by the following step 6. As necessary, a mixture of the compound [12] and a compound [13] is subjected to a separation and purification procedure to isolate the compound [12].

(Step 6)



[0073] In the above reaction, X¹, R¹, R², R³ and R⁴ have the same definitions as given above.

[0074] The isoxazoline compounds represented by the general formulas [12] and [13] can be produced by reacting

an olefin derivative represented by the general formula [10] with an oxime derivative represented by the general formula [11] in the presence or absence of a solvent (preferably in an appropriate solvent) in the presence of a base. When R³ and R⁴ are each a hydrogen atom, the isoxazoline compound represented by the general formula [12] can be obtained preferentially.

[0075] The reaction temperature is any temperature between 0°C and the reflux temperature of the reaction system and is preferably 10 to 80°C. The reaction time varies depending upon the compounds used, but is 0.5 hours to 2 weeks.


[0076] With respect to the amounts of the reagents used in the reaction, the compound represented by the general formula [10] is used in an amount of 1 to 3 equivalents per equivalent of the compound represented by the general formula [11].

[0077] As the solvent, there can be mentioned, for example, ethers such as ethylene glycol dimethyl ether, ethylene glycol diethyl ether, diethyl ether, dioxane, tetrahydrofuran and the like; halogenated hydrocarbons such as dichloroethane, carbon tetrachloride, chlorobenzene, dichlorobenzene and the like; aromatic hydrocarbons such as benzene, toluene, xylene and the like; acetic acid esters such as ethyl acetate, butyl acetate and the like; water; and mixtures thereof.

[0078] As the base, there can be mentioned, for example, alkali metal hydroxides such as sodium hydroxide, potassium hydroxide and the like; alkaline earth metal hydroxides such as calcium hydroxide, magnesium hydroxide and the like; alkali metal carbonates such as sodium carbonate, potassium carbonate and the like; alkali metal hydrogen-carbonates such as sodium hydrogencarbonate, potassium hydrogencarbonate and the like; alkali metal acetates such as sodium acetate, potassium acetate and the like; alkali metal fluorides such as sodium fluoride, potassium fluoride and the like; and organic bases such as pyridine, triethylamine, 1,8-diazabicyclo[5.4.0]-7-undecene and the like.

[0079] In the above production process, the compound represented by the general formula [10], used as an intermediate can be a commercial product or can be produced by a known reaction such as Wittig reaction or the like. The compound represented by the general formula [11] can be produced, for example, by a process described in Liebigs Annalen der Chemie, 985 (1989).

[0080] The compound represented by the general formula [1] can be produced from the above-shown compound represented by the general formula [12-] by the following process.




[0081] In the above reaction, X¹, R¹, R², R³, R⁴ and R⁷ have the same definitions as given above.

[0082] A compound represented by the general formula [15] can be produced by the above-described step 5; a compound represented by the general formula [16] can be produced by the above-described step 2; and the compound represented by the general formula [1] can be produced from the compound [15] by the above-described step 4 or from the compound [16] by the above-described step 3.

[0083] As the solvent, base and oxidizing agent, there can be mentioned the same solvents, bases and oxidizing agents as mentioned in the step 2, 3, 4 or 5.

[0084] A compound represented by the general formula [4] wherein R⁶ is a hydrogen atom, i.e. a compound represented by the general formula [21] can be produced by the following process.



[0085] In the above reaction, R⁵, X¹ and Y have the same definitions as given above; and R⁸ is an alkyl group.

(Step 11)

[0086] A compound represented by the general formula [20] can be produced by reacting a compound [17], [18] or [19] with a reducing agent in a solvent.

[0087] This reaction is conducted ordinarily at -60 to 150°C for 10 minutes to 24 hours.

[0088] With respect to the amounts of the reagents used in the reaction, the reducing agent is used in an amount of desirably 0.5 to 2 equivalents per equivalent of the compound [17], [18] or [19]; however, the amount can be varied appropriately depending upon the condition of the reaction.

[0089] As the reducing agent, there can be mentioned, for example, metal hydrides (e.g. diisobutyl aluminum hydride) and metal hydrogen complex compounds (e.g. sodium borohydride and lithium aluminum hydride) in production of [20] from [17]; and metal hydrides (e.g. diisobutyl aluminum hydride), metal hydrogen complex compounds (e.g. sodium borohydride and lithium aluminum hydride) and diborane in production of [20] from [18] or [19].

[0090] As the solvent, there can be mentioned, for example, ethers such as diethyl ether, tetrahydrofuran, dioxane and the like; aromatic hydrocarbons such as benzene, toluene and the like; and alcohols such as methanol, ethanol and the like.

(Step 12)

[0091] A compound represented by the general formula [21] can be produced by reacting the compound [20] with a halogenating agent in a solvent.


[0092] This reaction is conducted ordinarily at -50 to 100°C for 10 minutes to 24 hours.

[0093] With respect to the amounts of the reagents used in the reaction, the halogenating agent is used in an amount of desirably 1 to 3 equivalents per equivalent of the compound [20]; however, the amount can be varied appropriately depending upon the condition of the reaction.

[0094] As the halogenating agent, there can be mentioned, for example, hydrogen chloride, hydrogen bromide, phosphorus trichloride, phosphorus tribromide and thionyl chloride.

[0095] As the solvent, there can be mentioned, for example, halogenated hydrocarbons such as dichloroethane, carbon tetrachloride and the like; acids such as acetic acid and the like; and ethers such as tetrahydrofuran and the like.

[0096] The compound represented by the general formula [4] can be produced by the following process.



[0097] In the above reaction, R⁵, R⁶, X¹ and Y have the same definitions as given above.

[0098] The compound represented by the general formula [4] can be produced by reacting a compound [22] with a halogenating agent in a solvent in the presence or absence of a catalyst.

[0099] This reaction is conducted ordinarily at 30 to 150°C for 10 minutes to 24 hours.


[0100] With respect to the amounts of the reagents used in the reaction, the halogenating agent is used in an amount of desirably 1 to 10 equivalents relative to one equivalent of the compound [22]; however, the amount of the halogenating agent can be varied appropriately depending upon the condition of the reaction. The catalyst is used in an amount of 0.01 to 0.5 equivalent.

[0101] As the halogenating agent, there can be mentioned, for example, halogens such as bromine, chlorine and the like; N-halosuccinimides such as N-bromosuccinimide and the like; and pyridine salts such as pyridinium perbromide and the like.

[0102] As the solvent, there can be mentioned, for example, halogenated hydrocarbons such as dichloroethane, carbon tetrachloride, chlorobenzene, dichlorobenzene and the like; amides such as N,N-dimethylacetamide, N,N-dimethylformamide, N-methyl-2-pyrrolidinone and the like; sulfur compounds such as dimethyl sulfoxide, sulfolane and the like; and carboxylic acids such as formic acid, acetic acid and the like.

[0103] As the catalyst, there can be mentioned, for example, benzoyl peroxide, α, α'-azobisisobutyronitrile and a mixture thereof.

[0104] A compound represented by the general formula [4] wherein R⁵ and R⁶ are each a hydrogen atom, i.e. a compound represented by the general formula [24] can be produced by the following process.



[0105] In the above reaction, X¹ and Y have the same definitions as given above.

[0106] The compound represented by the general formula [24] can be produced by reacting a compound [23], hydrogen halide, and formaldehyde or paraformaldehyde in a solvent in the presence or absence of a Lewis acid according to the method described in Org. Synth., III, 557 (1955) or J. Am. Chem. Soc., 72, 2216 (1950), or by reacting the compound [23] with a halogenomethyl ether in a solvent in the presence of a Lewis acid according to the method described in J. Am. Chem. Soc., 97, 6155 (1975).

[0107] This reaction is conducted ordinarily at -40 to 150°C for 10 minutes to 24 hours.


[0108] With respect to the amounts of the reagents used in the reaction, the hydrogen halide, formaldehyde, paraformaldehyde, Lewis acid or halogenomethyl ether is used in an amount of desirably 1 to 2 equivalents per equivalent of the compound [23]; however, the amount of the former can be varied appropriately depending upon the condition of the reaction.

[0109] As the Lewis acid, there can be mentioned, for example, titanium tetrachloride, zinc chloride, aluminum chloride and zinc bromide.

[0110] As the hydrogen halide, there can be mentioned hydrogen chloride, hydrogen bromide and hydrogen iodide.

[0111] As the solvent, there can be mentioned, for example, halogenated hydrocarbons such as dichloroethane, carbon tetrachloride, chloroform and the like; aliphatic hydrocarbons such as hexane, heptane and the like; ethers such as dioxane, tetrahydrofuran and the like; carboxylic acids such as acetic acid and the like; carbon disulfide; and mixtures thereof.

[0112] A compound represented by the general formula [19] wherein R⁵ is a hydrogen atom, i.e. a compound represented by the general formula [25] can be produced by the following process.



[0113] In the above reaction, Y has the same definition as given above.

[0114] The compound represented by the general formula [25] can be produced by reacting the compound [23] with N,N-dimethylformamide in the presence of phosphoryl chloride, phosgene or thionyl chloride in the presence or absence of a solvent according to the Vilsmeier method described in Org. Synth., IV, 831 (1963), or by reacting the compound [23] with a dihalogenomethyl ether in a solvent in the presence of a Lewis acid and then giving rise to hydrolysis according to the method described in Chem. Ber., 93, 88 (1960).


[0115] This reaction is conducted ordinarily at -40 to 150°C for 10 minutes to 24 hours.

[0116] With respect to the amounts of the reagents used in the reaction, the phosphoryl chloride, phosgene, thionyl chloride, N,N-dimethylformamide, Lewis acid or dihalogenomethyl ether is used in an amount of desirably 1 to 2 equivalents per equivalent of the compound [23]; however, the amount of the former can be varied appropriately depending upon the condition of the reaction.

[0117] As the Lewis acid, there can be mentioned, for example, titanium tetrachloride, tin tetrachloride, zinc chloride, aluminum chloride and zinc bromide.

[0118] As the solvent, there can be mentioned, for example, halogenated hydrocarbons such as dichloroethane, carbon tetrachloride, chloroform and the like; aliphatic hydrocarbons such as hexane, heptane and the like; ethers such as dioxane, tetrahydrofuran and the like; carboxylic acids such as acetic acid and the like; amides such as N,N-dimethylformamide and the like; carbon disulfide; and mixtures thereof.

[0119] The compounds represented by the general formulas [17], [18], [19] and [20] can be produced by the following process.



[0120] In the above reaction, R⁵, R⁸ and Y have the same definitions as given above; and X² is a chlorine atom, a bromine atom or an iodine atom.

[0121] The compounds represented by the general formulas [17], [18], [19] and [20] can be produced by reacting a compound [26] with a magnesium reagent in the presence or absence of a solvent to obtain a compound [27] and then reacting the compound [27] with an electrophilic reagent according to the method described in J. Org. Chem., 65, 4618 (2000), or by reacting the compound [26] with n-butyl lithium in a solvent to obtain a compound [28] and then reacting the compound [28] with an electrophilic reagent according to the method described in Synth. Commun., 24 (2), 253 (1994).

[0122] This reaction is conducted ordinarily at -100 to 150°C for 10 minutes to 24 hours.

[0123] With respect to the amounts of the reagents used in the reaction, the amount of the magnesium reagent or the lithium reagent is desirably 1 to 5 equivalents per equivalent of the compound [26], and the amount of the electrophilic reagent is desirably 1 to 5 equivalents; however, these amounts can be varied appropriately depending upon the condition of the reaction.

[0124] As the magnesium reagent, there can be mentioned, for example, metal magnesium, isopropyl magnesium


bromide and diisopropyl magnesium.

[0125] As the lithium reagent, there can be mentioned, for example, n-butyl lithium, sec-butyl lithium and tert-butyl lithium.

[0126] As the electrophilic reagent, there can be mentioned, for example, esters such as ethyl formate, ethyl cyanoformate, ethyl acetate and the like; acid halides such as acetyl chloride, methyl chloroformate and the like; amides such as N,N-dimethylformamide and the like; aldehydes such as paraformaldehyde and the like; and carbon dioxide.

[0127] As the solvent, there can be mentioned, for example, halogenated hydrocarbons such as dichloroethane, carbon tetrachloride, chloroform and the like; aliphatic hydrocarbons such as hexane, pentane and the like; ethers such as dioxane, tetrahydrofuran and the like; and mixtures thereof.

[0128] Among compounds represented by the general formulas [4], [17], [18], [19], [20], [22], [23], [26], [29] or [34], a compound represented by the general formula [31] can be produced by the following process.



[0129] In the above reaction, Y has the same definition as given above; R⁹ is an alkyl group, a haloalkyl group, a cycloalkyl group, a cycloalkylalkyl group, an alkoxyalkyl group, an optionally substituted benzyl group, an optionally substituted heterocyclic alkyl group, an alkenyl group, an optionally substituted heterocyclic alkyl group, an alkenyl group, an alkynyl group, an alkylsulfonyl group, a haloalkylsulfonyl group, an optionally substituted aromatic heterocyclic group, an optionally substituted phenylsulfonyl group, an acyl group, a haloalkylcarbonyl group, an optionally substituted benzylcarbonyl group or an optionally substituted benzoyl group; and L¹ is a leaving group such as halogen atom, C1 to C4 alkylsulfonate group, C1 to C4 alkylsulfonyl group, optionally substituted benzylsulfonyl group, optionally substituted phenylsulfonate group, optionally substituted benzylsulfonate group or the like. When R⁹ is a haloalkyl group, L¹ is a leaving group having a higher reactivity than the halogen atom remaining after haloalkylation. For example, when R⁹ is a CHF₂ group, L¹ is a chlorine atom or a bromine atom; and when R⁹ is a CH₂CF₃ group, L¹ is a chlorine atom, a bromine atom, a p-toluenesulfonyloxy group or a methylsulfonyloxy group.

[0130] The compound represented by the general formula [31] can be produced by reacting a compound [29] with a compound [30] in a solvent in the presence of a base.


[0131] This reaction is conducted ordinarily at 0 to 120°C for 10 minutes to 24 hours.

[0132] With respect to the amounts of the reagents used in the reaction, the amount of the compound [30] is 1 to 20 equivalents per equivalent of the compound [29], and the amount of the base is 1 to 3 equivalents.

[0133] As the base, there can be mentioned, for example, alkali metal carbonates such as sodium carbonate, potassium carbonate and the like; alkali metal hydroxides such as sodium hydroxide, potassium hydroxide and the like; alkali metal hydrides such as potassium hydride, sodium hydride and the like; alkali metal alcoholates such as sodium ethoxide, sodium methoxide and the like; and organic bases such as 1,8-diazabicyclo[5.4.0]-7-undecene and the like.

[0134] As the solvent, there can be mentioned, for example, halogenated hydrocarbons such as dichloromethane, chloroform and the like; ethers such as diethyl ether, tetrahydrofuran and the like; aromatic hydrocarbons such as benzene, toluene and the like; aliphatic hydrocarbons such as hexane, heptane and the like; ketones such as acetone, methyl isobutyl ketone and the like; esters such as ethyl acetate, methyl acetate and the like; amides such as N-methylpyrrolidone, N,N-dimethylformamide and the like; sulfur compounds such as dimethyl sulfoxide, sulfolane and the like; nitriles such as acetonitrile and the like; and mixtures thereof.

[0135] Among compounds represented by the general formulas [4], [17], [18], [19], [20], [22], [23], [26], [29] or [31], a compound represented by the general formula [34] can be produced by the following process.



[0136] In the above reaction, L^1 has the same definition as given above; and R^{10} is an alkyl group, an alkyl group mono-substituted with a group selected from the substituent group β , a haloalkyl group, a cycloalkyl group, an alkenyl group, an alkynyl group, an alkylsulfinyl group, an alkylsulfonyl group, an alkylsulfonyl group mono-substituted with a group selected from the substituent group γ , a haloalkylsulfonyl group, an optionally substituted phenyl group, an optionally substituted aromatic heterocyclic group, an optionally substituted phenylsulfonyl group, an optionally substituted aromatic heterocyclicsulfonyl group, an acyl group, a haloalkylcarbonyl group, an optionally substituted benzylcarbonyl group, an optionally substituted benzoyl group, an alkoxycarbonyl group, an optionally substituted benzyloxy-carbonyl group, an optionally substituted phenoxycarbonyl group, or a carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from alkyl groups and optionally substituted phenyl group). The carbon atoms of the pyrazole ring may be substituted with 1 to 2 same or different groups selected from the substituent group α .

[0137] The compound represented by the general formula [34] can be produced by reacting a compound [32] with a compound [33] in a solvent in the presence of a base.

[0138] This reaction is conducted ordinarily at 0 to 120°C for 10 minutes to 24 hours.

[0139] With respect to the amounts of the reagents used in the reaction, the amount of the compound [33] is 1 to 20 equivalents per equivalent of the compound [32], and the amount of the base is 1 to 3 equivalents.


[0140] As the base and the solvent, there can be mentioned, for example, the same bases and solvents as mentioned in production of the compound [31] from the compound [29].

[0141] Introduction of a trifluoromethyl group into Y can be conducted according to or based on, for example, the methods described in J. Chem. Soc. Perkin Trans. 1, 8, 2293-2299 (1990); J. Fluorine Chem., 50 (3), 411-426 (1990); J. Chem. Soc. Chem. Commun., 18, 1389-1391 (1993); J. Chem. Soc. Chem. Commun., 1, 53-54 (1992); Chem. Lett., 1719-1720 (1981); Chem. Pharm. Bull., 38 (9), 2446-2458 (1990); J. Chem. Soc. Perkin Trans. 1, 921-926 (1988); Heterocycles, 37 (2), 775-782 (1994); Tetrahedron Lett., 30 (16), 2133-2136 (1989); J. Chem. Soc. Perkin Trans. 1, 2755-2761 (1980); Heterocycles, 22 (1), 117-124 (1984); Eur. J. Med. Chem. Chim. Ther., 24, 249-258 (1989); Acta Chem. Scand. Ser. B, 38 (6), 505-508 (1984); J. Fluorine Chem., 21, 495-514 (1982); J. Chem. Soc. Chem. Commun., 10, 638-639 (1988); J. Fluorine Chem., 67 (1), 5-6 (1994); J. Heterocycl. Chem., 31 (6), 1413-1416 (1994); Chem. Heterocycl. Compd., 30 (5), 576-578 (1994); J. Fluorine Chem., 78 (2), 177-182 (1996); J. Heterocycl. Chem., 34 (2), 551-556 (1997); Tetrahedron, 55 (52), 15067-15070 (1999); and Synthesis, 11, 932-933 (1980).

[0142] The -compounds represented by the general formulas [4], [17], [18], [19], [20], [21], [22], [23], [24], [25], [26], [29] and [31] can be produced according to or based on, for example, the methods described in Methoden der Organischen Chemie, E6a, 16-185 (1994) when Y is a furyl group; Methoden der Organischen Chemie, E6a, 186-555 (1994) when Y is a thienyl group; Methoden der Organischen Chemie, E6a, 556-798 (1994) when Y is a pyrrolyl group; Methoden der Organischen Chemie, E8b, 399-763 (1994) and JP-A-2000-219679 when Y is a pyrazolyl group; Methoden der Organischen Chemie, E8a, 45-225 (1993) when Y is an isoxazolyl group; Methoden der Organischen Chemie, E8a, 668-798 (1993) when Y is an isothiazolyl group; Methoden der Organischen Chemie, E8a, 891-1019 (1993) when Y is an oxazolyl group; Methoden der Organischen Chemie, E8b, 1-398 (1994) when Y is a thiazolyl group; Methoden der Organischen Chemie, E8c, 1-215 (1994) when Y is an imidazolyl group; Methoden der Organischen Chemie, E7a, 286-686 (1992) when Y is a pyridyl group; Methoden der Organischen Chemie, E9a, 557-682 (1997) when Y is a pyridazinyl group; Methoden der Organischen Chemie, E9b/1, 1-249 (1998) when Y is a pyrimidinyl group; Methoden der Organischen Chemie, E9b/1, 250-372 (1998) when Y is a pyrazinyl group; Methoden der Organischen Chemie, E9c, 530-796 (1998) when Y is a triazinyl group; Methoden der Organischen Chemie, E8d, 305-405 and 479-598 (1994) when Y is a triazolyl group; Methoden der Organischen Chemie, E8c, 397-818 (1994) when Y is an oxadiazolyl group; Methoden der Organischen Chemie, E8d, 59-304 (1994) when Y is a thiadiazolyl group; Methoden der Organischen Chemie, E6b1, 33-216 (1994) and Published International Patent Application WO-1997/29105 when Y is a benzofuryl group; Methoden der Organischen Chemie, E6b1, 217-322 (1994) when Y is a benzothienyl group; Methoden der Organischen Chemie, E6b1, 546-848 (1994), Methoden der Organischen Chemie, E6b2, 849-1336 (1994) and Published International Patent Application WO-1997/42188-A1 when Y is an indolyl group; Methoden der Organischen Chemie, E8a, 1020-1194 (1993) when Y is a benzoxazolyl group; Methoden der Organischen Chemie, E8b, 865-1062 (1994) when Y is a benzothiazolyl group; Methoden der Organischen Chemie, E8c, 216-391 (1994) when Y is a benzimidazolyl group; Methoden der Organischen Chemie, E8a, 226-348 (1993) when Y is a benzisoxazolyl group; Methoden der Organischen Chemie, E8a, 799-852 (1993) when Y is a benzisothiazolyl group; Methoden der Organischen Chemie, E8b, 764-864 (1994) when Y is an indazolyl group; Methoden der Organischen Chemie, E7a, 290-570 (1991) when Y is a quinolyl group; Methoden der Organischen Chemie, E7a, 571-758 (1991) when Y is an isoquinolyl group; Methoden der Organischen Chemie, E9a, 744-789 (1997) when Y is a phthalazinyl group; Methoden der Organischen Chemie, E9b/2, 93-265 (1998) when Y is a quinoxaliny group; Methoden der Organischen Chemie, E9b/2, 1-192 (1998) when Y is a quinazolinyl group; Methoden der Organischen Chemie, E9a, 683-743 (1997) when Y is a cinnolinyl group; and Methoden der Organischen Chemie, E8d, 406-478 (1994) when Y is a benzotriazolyl group.

<Production Process 2>

[0143]



[0144] In the above reaction, R¹, R², R³, R⁴, R⁵ and R⁶ have the same definitions as given above. The carbon atoms of the pyrazole ring may be substituted with 1 to 2 same or different groups selected from the substituent group α .

[0145] A compound of the present invention represented by the general formula [36] can be produced by reacting a compound [35] of the present invention, produced by the Production Process 1, with an acid in a solvent.

[0146] This reaction is conducted ordinarily at 0 to 120°C for 10 minutes to 24 hours.


[0147] With respect to the amounts of the reagents used in the reaction, the amount of the acid is 1 to 10 equivalents per equivalent of the compound [35]; however, the amount can be varied appropriately depending upon the condition of the reaction.

[0148] As the acid, there can be mentioned, for example, hydrochloric acid, hydrobromic acid and trifluoroacetic acid.

[0149] As the solvent, there can be mentioned, for example, halogenated hydrocarbons such as dichloroethane, carbon tetrachloride, chlorobenzene, dichlorobenzene and the like; amides such as N,N-dimethylacetamide, N,N-dimethylformamide, N-methyl-2-pyrrolidinone and the like; sulfur compounds such as dimethyl sulfoxide, sulfolane and the like; carboxylic acids such as formic acid, acetic acid and the like; and water.

<Production Process 3>

[0150]



[0151] In the above reaction, n, L¹, R¹, R², R³, R⁴, R⁵, R⁶ and R¹⁰ have the same definitions as given above. The carbon atoms of the pyrazole ring may be substituted with 1 to 2 same or different groups selected from the substituent group α .

[0152] A compound of the present invention represented by the general formula [37] can be produced by reacting the compound [36] of the present invention with the compound [33] in a solvent in the presence of a base.


[0153] With respect to the amounts of the reagents used in the reaction, the amount of the compound [33] is 1 to 3 equivalents per equivalent of the compound represented by the general formula [36] and the amount of the base is 1 to 3 equivalents.

[0154] As the solvent, there can be mentioned, for example, ethers such as dioxane, tetrahydrofuran (THF) and the like; halogenated hydrocarbons such as dichloroethane, carbon tetrachloride, chlorobenzene, dichlorobenzene and the like; amides such as N,N-dimethylacetamide, N,N-dimethylformamide, N-methyl-2-pyrrolidinone and the like; sulfur compounds such as dimethyl sulfoxide, sulfolane and the like; aromatic hydrocarbons such as benzene, toluene, xylene and the like; alcohols such as methanol, ethanol, propanol, isopropanol, butanol, tert-butanol and the like; ketones such as acetone, 2-butanone and the like; nitriles such as acetonitrile and the like; water; and mixtures thereof.

[0155] As the base, there can be mentioned, for example, metal hydrides such as sodium hydride and the like; alkali metal amides such as sodium amide, lithium diisopropylamide and the like; organic bases such as pyridine, triethylamine, 1,8-diazabicyclo[5.4.0]-7-undecene and the like; alkali metal hydroxides such as sodium hydroxide, potassium hydroxide and the like; alkaline earth metal hydroxides such as calcium hydroxide, magnesium hydroxide and the like; alkali metal carbonates such as sodium carbonate, potassium carbonate and the like; alkali metal hydrogencarbonates such as sodium hydrogencarbonate, potassium hydrogencarbonate and the like; and metal alcoholates such as sodium methoxide, potassium tertbutoxide and the like.

<Production Process 4>

[0156]



[0157] In the above reaction, R^1, R^2, R^3, R^4, R^5 and R^6 have the same definitions as given above; R^{11} is a hydrogen atom or substituent group α as mentioned above; X^3 is a chlorine atom, a fluorine atom, an alkylsulfonyl group or an optionally substituted benzylsulfonyl group; R^{12} is an alkyl group, a haloalkyl group, a cycloalkyl group, a cycloalkylalkyl group, an alkenyl group, an alkynyl group, an optionally substituted phenyl group, an optionally substituted aromatic heterocyclic group, an alkoxy carbonylalkyl group, an optionally substituted heterocyclic alkyl group or an optionally substituted benzyl group; R^{13} is an alkyl group, a haloalkyl group, an optionally substituted phenyl group, an optionally substituted aromatic heterocyclic group, an alkoxy carbonylalkyl group or an optionally substituted benzyl group; R^{14} and R^{15} may be the same or different and are each a hydrogen atom, an alkyl group, an optionally substituted phenyl group, an acyl group, a haloalkylcarbonyl group, an optionally substituted benzylcarbonyl group, an optionally substituted benzoyl group, an alkylsulfonyl group, a haloalkylsulfonyl group, an optionally substituted benzylsulfonyl group or an optionally substituted phenylsulfonyl group; and Z is an oxygen atom, a sulfur atom, $N=CR^{11a}$, $CR^{11a}=N$, $CR^{11a}=CR^{11b}$ or $N-R^{16}$ (wherein R^{16} is a hydrogen atom or has the same definition as R^{10} , and R^{11a} and R^{11b} have the same definition as R^{11}).

[0158] Compounds of the present invention represented by the general formulas [40], [42] and [44] can be produced by reacting a compound of the present invention represented by the general formula [38] with a compound [39], a compound [41] and a compound [43], respectively, in the presence or absence of a solvent and, as necessary, in the presence of a base.

[0159] This reaction is conducted ordinarily at 20 to 200°C, preferably 30 to 180°C for 10 minutes to 48 hours and, as necessary, under pressure.

[0160] With respect to the amounts of the reagents used in the reaction, the amount of the compound [39], the


compound [41] or the compound [43] is 1 to 20 equivalents per equivalent of the compound [38].

[0161] As the base used as necessary, there can be mentioned, for example, alkali metal hydroxides such as potassium hydroxide, sodium hydroxide and the like; alkali metal hydrides such as potassium hydride, sodium hydride and the like; alkali metal alcoholates such as sodium ethoxide, sodium methoxide and the like; and organic bases such as 1,8-diazabicyclo[5.4.0]-7-undecene and the like.

[0162] As the solvent, there can be mentioned, for example, halogenated hydrocarbons such as chloroform and the like; ethers such as diethyl ether, tetrahydrofuran and the like; aromatic hydrocarbons such as benzene, toluene and the like; aliphatic hydrocarbons such as hexane, heptane and the like; ketones such as acetone, methyl isobutyl ketone and the like; esters such as ethyl acetate and the like; amides such as N-methylpyrrolidone, N,N-dimethylformamide and the like; sulfur compounds such as dimethyl sulfoxide, sulfolane and the like; acetonitrile; and mixtures thereof.

<Production Process 5>

[0163]



[0164] In the above reaction, R¹, R², R³, R⁴, R⁵, R⁶, R⁸, R¹¹ and Z have the same definitions as given above.

[0165] A compound of the present invention represented by the general formula [46] can be produced by reacting a compound [45] of the present invention with an acid in a solvent.


[0166] This reaction is conducted ordinarily at 0 to 120°C for 10 minutes to 24 hours.

[0167] With respect to the amounts of the reagents used in the reaction, the amount of the acid is desirably 1 to 10 equivalents per equivalent of the compound [45]; however, the amount can be varied appropriately depending upon the condition of the reaction.

[0168] As the acid and the solvent, there can be mentioned the same acids and solvents as mentioned in the Production Process 2.

<Production Process 6>

[0169]



[0170] In the above reaction, Y, R¹, R², R³, R⁴, R⁵, R⁶, R⁹ and L¹ have the same definitions as given above. Y may be substituted with 1 to 5 same or different groups selected from the substituent group α.

[0171] A compound represented by the general formula [48] according to the present invention can be produced by reacting a compound [47] of the present invention with the compound [30] in a solvent in the presence of a base.

[0172] This reaction is conducted ordinarily at 0 to 150°C for 10 minutes to 24 hours.


[0173] With respect to the amounts of the reagents used in the reaction, the amount of the acid is desirably 1 to 1.2

equivalents per equivalent of the compound [47]; however, the amount can be varied appropriately depending upon the condition of the reaction.

[0174] As the base and the solvent, there can be mentioned the same bases and solvents as mentioned in the Production Process 3.

<Production Process 7>

[0175]



[0176] In the above reaction, Y, R¹, R², R³, R⁴, and R⁶ have the same definitions as given above; and R¹⁷ is an alkyl group, an optionally substituted benzyl group or an optionally substituted phenyl group. Y may be substituted with 1 to 5 same or different groups selected from the substituent group α .

[0177] A compound represented by the general formula [50] according to the present invention can be produced by hydrolyzing a compound [49] of the present invention in water or a mixed solvent of water and other solvent in the presence or absence of a base.

[0178] This reaction is conducted ordinarily at 0 to 100°C for 10 minutes to 24 hours.


[0179] With respect to the amounts of the reagents used in the reaction, the amount of the base, when used, is desirably 1 to 2 equivalents per equivalent of the compound [49]; however, the amount can be varied appropriately depending upon the condition of the reaction.

[0180] As the base, there can be mentioned, for example, inorganic bases such as potassium carbonate, sodium hydride, sodium hydroxide and the like; and organic bases such as 1,8-diazabicyclo[5.4.0]-7-undecene and the like.

[0181] As the other solvent mixed with water, there can be mentioned, for example, alcohols such as methanol, ethanol and the like; ethers such as tetrahydrofuran and the like; ketones such as acetone, methyl isobutyl ketone and the like; amides such as N,N-dimethylformamide and the like; sulfur compounds such as dimethyl sulfoxide, sulfolane and the like; acetonitrile; and mixtures thereof.

<Production Process 8>

[0182]



[0183] In the above reaction, Y, R¹, R², R³, R⁴, R⁵, R⁶ and R⁸ have the same definitions as given above; and R¹⁸ is an alkyl group. Y may be substituted with 1 to 5 same or different groups selected from the substituent group α .

[0184] A compound represented by the general formula [53] according to the present invention can be produced by reacting a compound [51] of the present invention with a compound [52] in a solvent in the presence of a base.

[0185] This reaction is conducted ordinarily at 0 to 100°C for 10 minutes to 24 hours.

[0186] With respect to the amounts of the reagents used in the reaction, the amount of the hydrochloride or sulfate of the compound [52] is desirably 1 to 5 equivalents per equivalent of the compound [51] and the amount of the base


is desirably 1 to 10 equivalents; however, these amounts can be varied appropriately depending upon the condition of the reaction.

[0187] As the base, there can be mentioned, for example, metal carbonates such as potassium carbonate, sodium carbonate and the like; metal acetates such as potassium acetate, sodium acetate and the like; and organic bases such as triethylamine, dimethylamine, 1,8-diazabicyclo[5.4.0]-7-undecene and the like.

[0188] As the solvent, there can be mentioned, for example, alcohols such as methanol, ethanol and the like; ethers such as tetrahydrofuran and the like; amides such as N,N-dimethylformamide and the like; water; and mixtures thereof.

<Production Process 9>

[0189]



[0190] In the above reaction, Y, R¹, R², R³, R⁴, R⁵ and R⁶ have the same definitions as given above; and R¹⁹ and R²⁰ are each a hydrogen atom or an alkyl group. Y may be substituted with 1 to 5 same or different groups selected from the substituent group α.

[0191] A compound represented by the general formula [57] according to the present invention can be produced by reacting the compound [50] of the present invention with thionyl chloride in the presence or absence of a solvent to obtain a compound [55] and then reacting the compound [55] with a compound [56] in the presence or absence of a solvent.

[0192] The reaction from the compound [50] to the compound [55] is conducted ordinarily at 0 to 100°C for 10 minutes to 24 hours.

[0193] With respect to the amounts of the reagents used in the reaction, the amount of thionyl chloride [54] is desirably 1 to 100 equivalents per equivalent of the compound [50] but it can be varied appropriately depending upon the condition of the reaction.

[0194] As the solvent, there can be mentioned, for example, halogenated hydrocarbons such as dichloromethane, chloroform and the like; ethers such as diethyl ether, tetrahydrofuran and the like; and aromatic hydrocarbons such as benzene, toluene and the like.


[0195] The reaction from the compound [55] to the compound [57] is conducted ordinarily at 0 to 100°C for 10 minutes to 24 hours.

[0196] With respect to the amounts of the reagents used in the reaction, the amount of the compound [56] is desirably 1 to 100 equivalents per equivalent of the compound [55] but it can be varied appropriately depending upon the condition of the reaction.

[0197] As the solvent, there can be mentioned, for example, the same solvents as used in the reaction from the compound [50] to the compound [55].

<Production Process 10>

[0198]



[0199] In the above reaction, Z, R¹, R², R³, R⁴, R⁵, R⁶, R¹¹ and X³ have the same definitions as given above.

[0200] A compound represented by the general formula [59] according to the present invention can be produced by reacting the compound [38] of the present invention with a compound [58] in a solvent.


[0201] This reaction is conducted ordinarily at 0 to 100°C for 10 minutes to 24 hours.

[0202] With respect to the amounts of the reagents used in the reaction, the amount of the compound [58] is desirably 1 to 2 equivalents per equivalent of the compound [38] but it can be varied appropriately depending upon the condition of the reaction.

[0203] As the solvent, there can be mentioned, for example, ethers such as dioxane, tetrahydrofuran (THF) and the like; halogenated hydrocarbons such as dichloroethane, carbon tetrachloride, chlorobenzene, dichlorobenzene and the like; amides such as N,N-dimethylacetamide, N,N-dimethylformamide, N-methyl-2-pyrrolidinone and the like; sulfur compounds such as dimethyl sulfoxide, sulfolane and the like; ketones such as acetone, 2-butanone and the like; nitriles such as acetonitrile and the like; water; and mixtures thereof.

<Production Process 11>

[0204]



[0205] In the above reaction, Y, R¹, R², R³, R⁴, R⁵ and R⁶ have the same definitions as given above; and R²¹ is an alkyl group, a haloalkyl group, a cycloalkyl group, a cycloalkylalkyl group, an alkenyl group, an alkynyl group, an alkoxycarbonylalkyl group, an optionally substituted heteroalkyl group or an optionally substituted benzyl group. Y may be substituted with 1 to 5 same or different groups selected from the substituent group α.

[0206] A compound represented by the general formula [61] according to the present invention can be produced by reacting the compound [47] of the present invention with a compound [60] in the presence of an azo compound and triphenylphosphine in a solvent according to a known method [Synthesis, 1-28 (1981)].

[0207] This reaction is conducted ordinarily at 0 to 100°C for 10 minutes to 24 hours.

[0208] With respect to the amounts of the reagents used in the reaction, the amounts of the compound [60], the azo compound and triphenylphosphine are desirably each 1 to 1.5 equivalents per equivalent of the compound [47] but the amounts can be varied appropriately depending upon the condition of the reaction.


[0209] As the solvent, there can be mentioned, for example, ethers such as dioxane, tetrahydrofuran (THF) and the like; halogenated hydrocarbons such as dichloroethane, carbon tetrachloride, chlorobenzene, dichlorobenzene and the like; amides such as N,N-dimethylacetamide, N,N-dimethylformamide, N-methyl-2-pyrrolidinone and the like; sulfur compounds such as dimethyl sulfoxide, sulfolane and the like; aromatic hydrocarbons such as benzene, toluene, xylene

and the like; acetonitrile; and mixtures thereof.

[0210] As the azo compound, there can be mentioned, for example, diethyl azodicarboxylate and diisopropyl azodicarboxylate.

<Production Process 12>

[0211]



[0212] In the above reaction, X^3 , n , R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and Z have the same definitions as given above; and m is an integer of 1 to 4. The carbon atom of the 3-position of the pyrazole ring may be substituted with a group selected from the substituent group α .

[0213] A compound represented by the general formula [63] according to the present invention can be produced by reacting a compound [62] of the present invention in the presence of a base in a solvent.

[0214] This reaction is conducted ordinarily at 0 to 120°C for 10 minutes to 24 hours.

[0215] With respect to the amounts of the reagents used in the reaction, the amount of the base is desirably 1 to 3 equivalents per equivalent of the compound represented by the general formula [62] but the amount can be varied appropriately depending upon the condition of the reaction.

[0216] As the base and the solvent, there can be mentioned the same bases and solvents as mentioned in the Production Process 3.

[0217] Incidentally, the sulfide compound mentioned in the Production Process 2 or the Production Processes 4 to 11 can be converted into a sulfoxide compound or a sulfone compound by oxidation according to the method described in the Production Process 1. Furthermore, the sulfide compound mentioned in the Production Process 2 or the Production Processes 4 to 11 wherein substituent Y is substituted by C1 to C10 alkylthio group, C1 to C10 alkylthio group mono-substituted with a group selected from the substituent group γ or C1 to C4 haloalkylthio group, can be converted into a sulfoxide compound or a sulfone compound according to the method described in the Production Process 1, by adding equi-molar to excess amount of an oxidizing agent to the sulfide compound; oxidizing the substituent substituted to substituent Y (C1 to C10 alkylthio group, C1 to C10 alkylthio group mono-substituted with a group selected from the substituent group γ or C1 to C4 haloalkylthio group) at the same time, and convert these substituent into a sulfoxide group or a sulfone group.

[0218] Then, specific description is made on the production process of the present compound, the production method of the present herbicide and the application of the present herbicide by way of Examples. Description is also made on the production process of each intermediate of the present compound.

<Example 1>

Production of 3-(5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0001)

[0219] 2.1 g of sodium hydrosulfide hydrate (purity: 70%, 26.2 mmoles) was added to a solution of 2.3 g (13.1 mmoles) of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline dissolved in 20 ml of N,N -dimethylformamide. The mixture was stirred for 2 hours. Thereto were added 1.8 g (13.1 mmoles) of anhydrous potassium carbonate, 2.0 g (13.1 mmoles) of Rongalit and 3.6 g (10.5 mmoles) of 4-bromomethyl-5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazole. The resulting mixture was stirred at room temperature for 15 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 2.7 g (yield: 65.5%) of 3-(5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as white crystals.

tals (melting point: 89 to 90°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)] :

7.55-7.50 (5H,m), 4.33 (2H,s), 2.83 (2H,s),

1.45 (6H,s)

<Example 2>

Production of 3-(5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0002)

[0220] 0.63 g of m-chloroperbenzoic acid (purity: 70%, 2.6 mmoles) was added, with ice-cooling, to a solution of 0.4 g (1.0 mmoles) of 3-(5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 15 ml of chloroform. The mixture was stirred at room temperature for 22 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogen-carbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting crystals were washed with hexane to obtain 0.4 g (yield: 83.2%) of 3-(5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline as white crystals (melting point: 132 to 133°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.60-7.51 (5H,m), 4.37 (2H,s), 3.14 (2H,s) 1.53 (6H,s)

<Example 3>

Production of 3-(5-chloro-1-methyl-3-phenyl-1H-pyrazol-4-ylmethylsulfinyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0003)

[0221] 0.87 g of m-chloroperbenzoic acid (purity: 70%, 3.54 mmoles) was added, with ice-cooling, to a solution of 0.85 g (2.53 mmoles) of 3-(5-chloro-1-methyl-3-phenyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 30 ml of chloroform. The mixture was stirred at room temperature for 1 hour to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogen-carbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.48 g (yield: 53.9%) of 3-(5-chloro-1-methyl-3-phenyl-1H-pyrazol-4-ylmethylsulfinyl)-5,5-dimethyl-2-isoxazoline as a transparent viscous substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.63-7.60 (2H,m), 7.48-7.37 (3H,m), 4.29 (2H,q), 3.91 (3H,s), 3.12 (1H,d), 2.79 (1H,d), 1.41 (3H,s), 1.35 (3H,s)

<Example 4>

Production of 5,5-dimethyl-3-(5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline (present compound No. 3-0021)

[0222] 9.3 g of sodium hydrosulfide hydrate (purity: 70%, 116.3 mmoles) was added to a solution of 18.7 g (105.7 mmoles) of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline (present compound No. 2-1) dissolved in 300 ml of N,N-dimethylformamide. The mixture was stirred for 2 hours. The reaction system was ice-cooled. Thereto was added a solution of 30.3 g (93.8 mmoles) of 4-bromomethyl-5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazole dissolved in 200 ml of N,N-dimethylformamide. The mixture was stirred at 0°C for 30 minutes to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 13.11 g (yield: 37.4%) of 5,5-dimethyl-3-(5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.65-7.39 (5H,m), 4.24 (2H,s), 2.81 (2H,s), 1.43 (6H,s)

<Example 5>

Production of 5,5-dimethyl-3-(5-ethylthio-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline (present compound No. 3-0022)

[0223] 0.2 g (4.0 mmoles) of sodium hydroxide and 1 ml of water were added to a solution of 0.25 g (4.0 mmoles) of ethanethiol dissolved in 10 ml of N,N-dimethylformamide. The mixture was stirred at room temperature for 30 minutes. Thereto was added a solution of 0.5 g (1.4 mmoles) of 5,5-dimethyl-3-(5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 5 ml of N,N-dimethylformamide. The resulting mixture was stirred for 1 hour to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.6 g (yield: 100%) of 5,5-dimethyl-3-(5-ethylthio-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.62-7.47 (5H,m), 4.44 (2H,s), 2.83 (2H,s), 2.50 (2H,q), 1.45 (6H,s), 1.02 (3H,t)

<Example 6>

Production of 5,5-dimethyl-3-(5-ethylsulfonyl-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-isoxazoline (present compound No. 3-0004)

[0224] 1.7 g of m-chloroperbenzoic acid (purity: 70%, 6.7 mmoles) was added, with ice-cooling, to a solution of 0.6 g (1.3 mmoles) of 5,5-dimethyl-3-(5-ethylthio-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 10 ml of chloroform. The mixture was stirred at room temperature for 16 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogen-carbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting crystals were washed with hexane to obtain 0.6 g (yield: 93.0%) of 5,5-dimethyl-3-(5-ethylsulfonyl-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-isoxazoline as light yellow crystals (melting point: 158 to 160°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.58-7.54 (5H,m), 5.16 (2H,s), 3.18 (2H,s), 3.15 (2H,q), 1.55 (6H,s), 1.24 (3H,t)

<Example 7>

Production of 5,5-dimethyl-3-(5-dimethylamino-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline (present compound 3-0023)

[0225] 0.8 g (6.7 mmoles) of a 40% aqueous dimethylamine solution was added to a solution of 0.5 g (1.3 mmoles) of 5,5-dimethyl-3-(5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 10 ml of N,N-dimethylformamide. The mixture was stirred at 100°C for 9 hours in a sealed tube. Thereto was added 3.0 g (26.6 mmoles) of a 40% aqueous dimethylamine solution, and the resulting mixture was stirred for 9 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous sodium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.4 g (yield: 80.6%) of 5,5-dimethyl-3-(5-dimethylamino-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.58-7.38 (5H,m), 4.35 (2H,s), 2.82 (2H,s), 2.77 (6H,s), 1.45 (6H,s)

<Example 8>

Production of 5,5-dimethyl-3-(5-dimethylamino-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-isoxazoline (present compound 3-0005)

[0226] 0.7 g of m-chloroperbenzoic acid (purity: 70%, 2.7 mmoles) was added, with ice-cooling, to a solution of 0.4

g (1.1 mmoles) of 5,5-dimethyl-3-(5-dimethylamino-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 10 ml of chloroform. The mixture was stirred at room temperature for 20 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting crystals were washed with hexane to obtain 0.2 g (yield: 52.0%) of 5,5-dimethyl-3-(5-dimethylamino-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-isoxazoline as a white powder (melting point: 150 to 151°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.61-7.38 (5H,m), 4.75 (2H,s), 3.13 (2H,s), 2.76 (6H,s), 1.53 (6H,s)

<Example 9>

Production of 3-(1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0006)

[0227] 21.8 g of sodium hydrosulfide (purity: 70%, 272.5 mmoles) was added to a solution of 24.1 g (136.0 mmoles) of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline dissolved in 200 ml of N,N-dimethylformamide. The mixture was stirred for 1 hour. Thereafter were added 18.8 g (136.2 mmoles) of anhydrous potassium carbonate and 21.0 g (136.2 mmoles) of Rongalit. The resulting mixture was stirred for 2 hours. Thereafter was added, with ice-cooling, 40 g (125 mmoles) of 4-bromomethyl-1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazole. The resulting mixture was stirred at room temperature for 2 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 23.0 g (yield: 57.1%) of 3-(1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as light pink crystals (melting point: 79.0 to 81.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.24 (2H,s), 2.80 (2H,s), 1.71 (9H,s), 1.43 (6H,s)

<Example 10>

Production of 3-(5-chloro-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0007)

[0228] 19.8 g (53.4 mmoles) of 3-(1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline was added to 170 ml of a 25% hydrogen bromide-acetic acid solution. The mixture was stirred at 40 to 50°C for 2 hours to give rise to a reaction. After the completion of the reaction was confirmed, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 12.0 g (yield: 60.6%) of 3-(5-chloro-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as light yellow crystals (melting point: 120.0 to 122.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)] :

4.26 (2H,s), 2.81 (2H,s), 1.44 (6H,s)

<Example 11>

Production of 3-(5-chloro-1-difluoromethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0008) and 3-(3-chloro-1-difluoromethyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound 3-0009)

[0229] 3.1 g (22.5 mmoles) of anhydrous potassium carbonate was added to a solution of 2.3 g (7.3 mmoles) of 3-(5-chloro-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 50 ml of N,N-dimethylformamide. Thereinto was blown chlorodifluoromethane. The resulting mixture was stirred at 130 to 140°C for 3 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium

chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.69 g (yield: 25.8%) of 3-(5-chloro-1-difluoromethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as light yellow crystals (melting point: 41.0 to 42.0°C) and 0.54 g (yield: 20.2%) of 3-(3-chloro-1-difluoromethyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as a white powder (melting point: 89.0 to 90.0°C).

3-(5-Chloro-1-difluoromethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.22 (1H,t), 4.25 (2H,s), 2.80 (2H,s), 0.44 (6H,s) 3-(3-Chloro-1-difluoromethyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.19 (1H, t), 4.28 (2H,s), 2.80 (2H,s), 1.44 (6H,s)

<Example 12>

Production of 3-(5-chloro-1-difluoromethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0010)

[0230] 1.4 g of m-chloroperbenzoic acid (purity: 70%, 8.1 mmoles) was added, with ice-cooling, to a solution of 0.69 g (1.9 mmoles) of 3-(5-chloro-1-difluoromethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting solid was washed with n-hexane to obtain 0.4 g (yield: 53.3%) of 3-(5-chloro-1-difluoromethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline as a white powder (melting point: 126.0 to 127.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.26 (1H,t), 4.68 (2H,s), 3.11 (2H,s), 1.53 (6H,s)

<Example 13>

Production of 3-(3-chloro-1-difluoromethyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0011)

[0231] 1.1 g of m-chloroperbenzoic acid (purity: 70%, 6.4 mmoles) was added, with ice-cooling, to a solution of 0.54 g (1.5 mmoles) of 3-(3-chloro-1-difluoromethyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting solid was washed with n-hexane to obtain 0.47 g (yield: 79.7%) of 3-(3-chloro-1-difluoromethyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline as a white powder (melting point: 136.0 to 137.0°C).

¹R-NMR [CDCl₃/TMS, δ (ppm)]:

7.23 (1H,t), 4.71 (2H,s), 3.11 (2H,s), 1.53 (6H,s)

<Example 14>

Production of 5,5-dimethyl-3-(3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline (present compound No. 3-0024)

[0232] 3.1 g of sodium hydrosulfide hydrate (purity: 70%, 22.0 mmoles) was added to a solution of 3.3 g (17.3 mmoles) of 5,5-dimethyl-3-ethylsulfonyl-2-isoxazoline dissolved in 10 ml of N,N-dimethylformamide. The mixture was stirred for 2 hours. Thereto were added 3.1 g (22.0 mmoles) of anhydrous potassium carbonate, 2.7 g (17.5 mmoles) of Rongalit and 4.0 g (17.5 mmoles) of 4-chloromethyl-3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazole. The resulting mixture

was stirred at room temperature for 2 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 2.8 g (yield: 52.0%) of 5,5-dimethyl-3-(3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline.

<Example 15>

Production of 5,5-dimethyl-3-(3-hydroxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline (present compound No. 3-0025)

[0233] To 20 ml of a 25% hydrogen bromide acetic acid solution was added 3.3 g (10.6 mmoles) of 5,5-dimethyl-3-(3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline. The mixture was stirred at 50°C for 3 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the solvent contained therein. The residue was poured into water. The resulting crystals were collected by filtration, washed with water and dried to obtain 3.1 g (yield: 96.0%) of intended 5,5-dimethyl-3-(3-hydroxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline.

<Example 16>

Production of 5,5-dimethyl-3-(3-ethoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline (present compound No. 3-0026)

[0234] 0.20 g (1.3 mmoles) of anhydrous potassium carbonate and 0.20 g (1.5 mmoles) of ethyl iodide were added to a solution of 0.30 g (1.0 mmoles) of 5,5-dimethyl-3-(3-hydroxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 10 ml of N,N-dimethylformamide. The mixture was stirred at 50°C for 3 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.30 g (yield: 92.0%) of intended 5,5-dimethyl-3-(3-ethoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline.

<Example 17>

Production of 5,5-dimethyl-3-(3-ethoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-isoxazoline (present compound No. 3-0012)

[0235] 0.68 g of m-chloroperbenzoic acid (purity: 70%, 2.76 mmoles) was added, with ice-cooling, to a solution of 0.30 g (0.92 mmoles) of 5,5-dimethyl-3-(3-ethoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 10 ml of chloroform. The mixture was stirred at room temperature for 5 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting crystals were washed with hexane to obtain 0.24 g (yield: 73.0%) of 5,5-dimethyl-3-(3-ethoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-isoxazoline as white crystals (melting point: 124 to 125°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.50 (2H,s), 4.27 (2H,q), 3.86 (3H,s), 3.04 (2H,s), 1.49 (6H,s), 1.39 (3H,t)

<Example 18>

Production of 5,5-dimethyl-3-(5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline (present compound No. 3-0027)

[0236] 19.3 g of sodium hydrosulfide (purity: 70%, 344.6 mmoles) was added to a solution of 21.3 g (120.3 mmoles) of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline dissolved in 200 ml of N,N-dimethylformamide. The mixture was stirred for 1 hour. Thereto were added 16.7 g (121.0 mmoles) of anhydrous potassium carbonate and 18.6 g (120.7 mmoles)

of Rongalit. The resulting mixture was stirred for 2 hours. Thereto was added, with ice-cooling, 31.4 g (120.3 mmoles) of 4-bromomethyl-5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazole. The resulting mixture was stirred at room temperature for 2 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 29.0 g (yield: 90.3%) of 5,5-dimethyl-3-(5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.24 (2H,s), 3.90 (3H,s), 2.78 (2H,s), 1.42 (6H,s)

<Example 19>

Production of 5,5-dimethyl-3-(5-methoxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline (present compound No. 3-0028)

[0237] 0.77 g (4.0 mmoles) of sodium methoxide (a 28% methanol solution) was added to a solution of 0.5 g (1.6 mmoles) of 5,5-dimethyl-3-(5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 20 ml of methanol. The mixture was stirred for 4 hours under refluxing, to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.5 g (yield: 96.7%) of 5,5-dimethyl-3-(5-methoxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.26 (2H,s), 4.07 (3H,s), 3.72 (3H,s), 2.80 (2H,s), 1.43 (6H,s)

<Example 20>

Production of 5,5-dimethyl-3-(5-methoxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-isoxazoline (present compound No. 3-0013)

[0238] 1.3 g of m-chloroperbenzoic acid (purity: 70%, 7.5 mmoles) was added, with ice-cooling, to a solution of 0.5 g (1.5 mmoles) of 5,5-dimethyl-3-(5-methoxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting solid was washed with n-hexane to obtain 0.31 g (yield: 58.2%) of 5,5-dimethyl-3-(5-methoxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-isoxazoline as a white powder (melting point: 113.0 to 114.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.60 (2H,s), 4.11 (3H,s), 3.79 (3H,s), 3.10 (2H,s), 1.51(6H,s)

<Example 21>

Production of 3-(5-(2-chlorophenoxy)-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0029)

[0239] 0.2 g (8.3 mmoles) of sodium hydride (purity: 60%) was added, with ice-cooling, to a solution of 0.44 g (3.4 mmoles) of 2-chlorophenol dissolved in 30 ml of N,N-dimethylformamide. The mixture was stirred for 1 hour. Thereto was added 0.7 g (2.2 mmoles) of 5,5-dimethyl-3-(5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline. The resulting mixture was stirred at 120 to 130°C for 5 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.63 g (yield: 66.7%) of 3-(5-(2-chlorophenoxy)-1-methyl-3-trifluoromethyl-1H-pyrazol-4-yl-

methylthio)-5,5-dimethyl-2-isoxazoline as a yellow oily substance.

<Example 22>

Production of 3-(5-(2-chlorophenoxy)-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0014)

[0240] 1.0 g of m-chloroperbenzoic acid (purity: 70%, 5.8 mmoles) was added, with ice-cooling, to a solution of 0.63 g (1.5 mmoles) of 3-(5-(2-chlorophenoxy)-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting solid was washed with n-hexane to obtain 0.31 g (yield: 45.7%) of 3-(5-(2-chlorophenoxy)-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline as a white powder (melting point: 67.0 to 70.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.50-6.91 (4H,m), 4.45 (2H,s), 3.71 (3H,s), 3.03 (2H,s), 1.47 (6H,s)

<Example 23>

Production of 3-(5-cyclopentyloxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0030)

[0241] To a solution of 0.43 g (1.6 mmoles) of triphenylphosphine dissolved in 10 ml of benzene were added 0.14 g (1.6 mmoles) of cyclopentanol, 0.5 g (1.6 mmoles) of 5,5-dimethyl-3-(5-hydroxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline and 0.7 g (1.6 mmoles) of diethyl azodicarboxylate (a 40% toluene solution). The mixture was stirred at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting organic layer was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.52 g (yield: 85.2%) of 3-(5-cyclopentyloxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as a colorless transparent oily substance.

<Example 24>

Production of 3-(5-cyclopentyloxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0015)

[0242] 0.85 g of m-chloroperbenzoic acid (purity: 70%, 4.9 mmoles) was added, with ice-cooling, to a solution of 0.52 g (1.4 mmoles) of 3-(5-(cyclopentyloxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting solid was washed with n-hexane to obtain 0.2 g (yield: 35.5%) of 3-(5-cyclopentyloxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline as a white powder (melting point: 113.0 to 114.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

5.03 (1H,br), 4.60 (2H,s), 3.73 (3H,s), 3.05 (2H,s), 1.88-1.70 (8H,m), 1.50 (6H,s)

<Example 25>

Production of 3-(5-cyano-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0031)

[0243] 0.2 g (4.0 mmoles) of sodium cyanide was added to a solution of 0.5 g (1.6 mmoles) of 5,5-dimethyl-3-(5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 30 ml of N,N-dimethylformamide. The mixture was stirred at 40°C for 1 hour to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.9 g of crude 3-(5-cyano-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.30 (2H,s), 4.08 (3H,s), 2.81 (2H,s), 1.43 (6H,s)

<Example 26>

Production of 3-(5-cyano-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0016)

[0244] 2.1 g of m-chloroperbenzoic acid (purity: 70%, 12.2 mmoles) was added, with ice-cooling, to a solution of 0.9 g of 3-(5-cyano-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (crude compound) dissolved in 50 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting solid was washed with n-hexane to obtain 0.43 g (yield: 76.4%) of 3-(5-cyano-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline as a white powder (melting point: 105.0 to 108.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.73 (2H,s), 4.16 (3H,s), 3.14 (2H,s), 1.53 (6H,s)

<Example 27>

Production of 3-(3,5-dichloro-1-ethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0032)

[0245] 0.6 g of sodium hydrosulfide (purity: 70%, 10.7 mmoles) was added to a solution of 0.7 g (3.7 mmoles) of 5,5-dimethyl-3-ethylsulfonyl-2-isoxazoline dissolved in 30 ml of N,N-dimethylformamide. The mixture was stirred for 1 hour. Thereto were added 0.51 g (3.7 mmoles) of anhydrous potassium carbonate and 0.56 g (3.6 mmoles) of Rongalit. The resulting mixture was stirred for 2 hours. Thereto was added, with ice-cooling, 0.9 g (3.5 mmoles) of 4-bromomethyl-3,5-dichloro-1-ethyl-1H-pyrazole. The resulting mixture was stirred at room temperature for 2 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.8 g (yield: 70.8%) of 3-(3,5-dichloro-1-ethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as a colorless transparent oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.14 (2H,s), 4.14 (2H, q), 2.81 (2H,s), 1.43 (6H,s), 1.42 (3H,t)

<Example 28>

Production of 3-(3,5-dichloro-1-ethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0017)

[0246] 2.0 g of m-chloroperbenzoic acid (purity: 70%, 11.6 mmoles) was added, with ice-cooling, to a solution of 0.8 g (2.6 mmoles) of 3-(3,5-dichloro-1-ethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting solid was washed with n-hexane to obtain 0.41 g (yield: 46.6%) of 3-(3,5-dichloro-1-ethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline as a white powder (melting point: 105.0 to 107.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.48 (2H,s), 4.19 (2H,q), 3.05 (2H,s), 1.51 (6H,s), 1.45 (3H,t)

<Example 29>

Production of 3-(5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0020)

[0247] 1.2 g of sodium hydrosulfide hydrate (purity: 70%, 15.0 mmoles) was added to a solution of 1.9 g (10.0 mmoles) of 5,5-dimethyl-3-ethylsulfonyl-2-isoxazoline dissolved in 30 ml of N,N-dimethylformamide. The mixture was stirred for 2 hours. Thereto were added 2.1 g (15.0 mmoles) of anhydrous potassium carbonate, 2.3 g (15.0 mmoles) of Rongalit and 2.6 g (10.0 mmoles) of 4-bromomethyl-5-chloro-3-difluoromethyl-1-methyl-1H-pyrazole. The resulting mixture was stirred at room temperature for 15 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 2.1 g (yield: 68.0%) of 3-(5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as a colorless viscous liquid ($n_D^{20} = 1.5183$).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

6.70 (1H,t, J=54.2 Hz), 4.24 (2H,s), 3.86 (3H,s), 2.80 (2H,s), 1.42 (6H,s)

<Example 30>

Production of 3-(5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0018)

[0248] 3.6 g of m-chloroperbenzoic acid (purity: 70%, 14.5 mmoles) was added, with ice-cooling, to a solution of 1.8 g (5.8 mmoles) of 3-(5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 15 ml of chloroform. The mixture was stirred at room temperature for 22 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting crystals were washed with hexane to obtain 1.7 g (yield: 85.9%) of 3-(5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline as white crystals (melting point: 78 to 79°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

6.80 (1H,t, J=54.8 Hz), 4.60 (2H,s), 3.91 (3H,s), 3.08 (2H,s), 1.51 (6H,s)

<Example 31>

Production of 5,5-dimethyl-3-(5-methyl-3-trifluoromethylisoxazol-4-ylmethylthio)-2-isoxazoline (present compound No. 4-0003)

[0249] 0.4 g of sodium hydrosulfide hydrate (purity: 70%, 4.6 mmoles) was added to a solution of 0.4 g (2.3 mmoles) of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline dissolved in 10 ml of N,N-dimethylformamide. The mixture was stirred for 2 hours. Thereto were added 0.3 g (2.3 mmoles) of potassium carbonate, 0.4 g (2.3 mmoles) of Rongalit and 0.5 g (1.8 mmoles) of 4-bromomethyl-5-methyl-3-trifluoromethylisoxazole. The resulting mixture was stirred at room temperature for 14 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.4 g (yield : 70.0%) of 5,5-dimethyl-3-(5-methyl-3-trifluoromethylisoxazol-4-ylmethylthio)-2-isoxazoline.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.11 (2H,s), 2.77 (2H,s), 2.54 (3H,s), 1.42 (6H,s)

<Example 32>

Production of 5,5-dimethyl-3-(5-methyl-3-trifluoromethylisoxazol-4-ylmethylsulfonyl)-2-isoxazoline (present compound No. 4-0001)

[0250] 0.8 g of m-chloroperbenzoic acid (purity: 70%, 3.2 mmoles) was added, with ice-cooling, to a solution of 0.4 g (1.3 mmoles) of 5,5-dimethyl-3-(5-methyl-3-trifluoromethylisoxazol-4-ylmethylthio)-2-isoxazoline dissolved in 10 ml of chloroform. The mixture was stirred at room temperature for 4 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting crystals were washed with hexane to obtain 0.4 g (yield: 95.0%) of 5,5-dimethyl-3-(5-methyl-3-trifluoromethylisoxazol-4-ylmethylsulfonyl)-2-isoxazoline as white crystals (melting point: 135 to 136°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.54 (2H,s), 3.11 (2H,s), 2.61 (3H,s), 1.52 (6H,s)

<Example 33>

Production of [(5-chloro-3-methyl-isothiazol-4-yl)-methylthio]-5,5-dimethyl-2-isoxazoline (present compound No. 4-0004)

[0251] 0.82 g of sodium hydrosulfide (purity: 70%, 10.00 mmoles) was added at the room temperature to a solution of 0.89 g (5.00 mmoles) of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline dissolved in 10 ml of N,N-dimethylformamide. The mixture was stirred for 2 hours. Thereto were added 0.70 g (5.00 mmoles) of anhydrous potassium carbonate, 0.78 g (5.00 mmoles) of Rongalit and 0.91 g (5.00 mmoles) of 5-chloro-4-chloromethyl-3-methylisothiazole. The resulting mixture was stirred at room temperature overnight to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 1.38 g (yield: quantitative) of [(5-chloro-3-methyl-isothiazol-4-yl)-methylthio]-5,5-dimethyl-2-isoxazoline.

<Example 34>

Production of [(5-chloro-3-methyl-isothiazol-4-yl)-methylsulfonyl]-5,5-dimethyl-2-isoxazoline (present compound No. 4-0002)

[0252] 2.96 g of m-chloroperbenzoic acid (purity: 70%, 12.00 mmoles) was added, with ice-cooling, to a solution of 1.38 g (5.00 mmoles) of [(5-chloro-3-methyl-isothiazol-4-yl)-methylthio]-5,5-dimethyl-2-isoxazoline dissolved in 20 ml

of chloroform. The mixture was stirred for 1 hour and then at room temperature for overnight to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 0.65 g (yield: 47.0%) of [(5-chloro-3-methyl-isothiazol-4-yl)-methylsulfonyl]-5,5-dimethyl-2-isoxazoline as a light yellow powder (melting point: 113 to 114°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.89 (1H,s), 4.67 (2H,s), 3.05 (2H,s), 2.59 (3H,s), 1.51 (6H,s)

<Example 35>

Production of 5,5-dimethyl-3-[2,5-dimethyl-4-(1-methoxyiminoethyl)-thiophen-3-ylmethylthio]-2-isoxazoline (present compound No. 2-0002)

[0253] 0.57 g (6.8 mmoles) of O-methylhydrozylamine hydrochloride and 0.56 g (6.8 mmoles) of sodium acetate were added to a solution of 1.0 g (3.4 mmoles) of 3-(4-acetyl-2,5-dimethylthiophen-3-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 50 ml of ethanol. The mixture was stirred for 5 hours under refluxing, to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.4 g (36.4%) of 5,5-dimethyl-3-[2,5-dimethyl-4-(1-methoxyiminoethyl)-thiophen-3-ylmethylthio]-2-isoxazoline as a yellow oily substance.

¹R-NMR [CDCl₃/TMS, δ (ppm)]:

4.21 (2H,s), 3.95 (3H,s), 2.76 (2H,s), 2.38 (3H,s), 2.34 (3H,s), 2.13 (3H,s), 1.42 (6H,s)

<Example 36>

Production of 5,5-dimethyl-3-[2,5-dimethyl-4-(1-methoxyiminoethyl)-thiophen-3-ylmethylsulfonyl]-2-isoxazoline (present compound No. 2-0001)

[0254] 0.61 g of m-chloroperbenzoic acid (purity: 70%, 3.5 mmoles) was added, with ice-cooling, to a solution of 0.4 g (1.2 mmoles) of 5,5-dimethyl-3-[2,5-dimethyl-4-(1-methoxyiminoethyl)-thiophen-3-ylmethylthio]-2-isoxazoline dissolved in 30 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.35 g (80%) of 5,5-dimethyl-3-[2,5-dimethyl-4-(1-methoxyiminoethyl)-thiophen-3-ylmethylsulfonyl]-2-isoxazoline as white crystals (melting point: 95.0 to 96.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.79 (2H,s), 3.95 (3H,s), 2.93 (2H,s), 2.42 (3H,s), 2.37 (3H,s), 2.17 (3H,s), 1.47 (6H,s)

<Example 37>

Production of 5,5-dimethyl-3-(4-trifluoromethyl-pyridin-3-ylmethylthio)-2-isoxazoline (present compound No. 7-0003)

[0255] 0.26 g of sodium hydrosulfide (purity: 70%, 4.6 mmoles) was added to a solution of 0.3 g (1.6 mmoles) of 5,5-dimethyl-3-ethylsulfonyl-2-isoxazoline dissolved in 20 ml of N,N-dimethylformamide. The mixture was stirred for 1 hour. Thereto were added 0.22 g (1.6 mmoles) of anhydrous potassium carbonate and 0.25 g (1.6 mmoles) of Rongalit. The resulting mixture was stirred for 2 hours. Thereto was added, with ice-cooling, 0.3 g (1.3 mmoles) of 3-bromomethyl-4-trifluoromethyl-pyridine. The resulting mixture was stirred at room temperature for 2 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent:

hexane-ethyl acetate mixed solvent) to obtain 0.45 g (yield: 98.9%) of 5, 5-dimethyl-3-(4-trifluoromethyl-pyridin-3-yl-methylthio)-2-isoxazoline as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.98 (1H,s) 8.70 (1H,d), 7.51 (1H,d), 4.47 (2H,s), 2.79 (2H,s), 1.43 (6H,s)

<Example 38>

Production of 5,5-dimethyl-3-(4-trifluoromethyl-pyridin-3-ylmethylsulfonyl)-2-isoxazoline (present compound No. 7-0001) and 5,5-dimethyl-3-(4-trifluoromethyl-pyridine-N-oxide-3-ylmethylsulfonyl)-2-isoxazoline (present compound No. 7-0002)

[0256] 0.77 g of m-chloroperbenzoic acid (purity: 70%, 4.5 mmoles) was added, with ice-cooling, to a solution of 0.45 g (1.6 mmoles) of 5,5-dimethyl-3-(4-trifluoromethyl-pyridin-3-ylmethylthio)-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.06 g (yield: 12.0%) of 5,5-dimethyl-3-(4-trifluoromethyl-pyridin-3-ylmethylsulfonyl)-2-isoxazoline as light yellow crystals (melting point: 77.0 to 80.0°C) and 0.12 g (yield: 23.1%) of 5,5-dimethyl-3-(4-trifluoromethyl-pyridine-N-oxide-3-ylmethylsulfonyl)-2-isoxazoline as white crystals (melting point: 114.0 to 116.0°C). 5,5-Dimethyl-3-(4-trifluoromethyl-pyridin-3-ylmethylsulfonyl)-2-isoxazoline

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.98 (1H,s), 8.84 (1H,d) 7.64 (1H,d), 4.92 (2H,s), 3.09 (2H,s), 1.52 (6H,s)

5,5-Dimethyl-3-(4-trifluoromethyl-pyridine-N-oxide-3-ylmethylsulfonyl)-2-isoxazoline

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.50 (1H,s) 8.25 (1H,d), 7.59 (1H,d), 4.81 (2H,s), 3.12 (2H,s), 1.53 (6H,s)

<Example 39>

Production of 5,5-dimethyl-[(4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methylthio]-2-isoxazoline (present compound No. 8-0002)

[0257] 0.32 g of sodium hydrosulfide (purity: 70%, 4.00 mmoles) was added, at room temperature, to a solution of 0.35 g (2.00 mmoles) of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline dissolved in 10 ml of dimethylformamide. The mixture was stirred for 2 hours. To the reaction mixture were added 0.28 g (2.00 mmoles) of anhydrous potassium carbonate, 0.31 g (2.00 mmoles) of Rongalit and 0.45 g (2.00 mmoles) of 5-chloromethyl-4-methoxy-6-trifluoromethylpyrimidine. The resulting mixture was stirred at room temperature for 2 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 0.55 g (yield: 85.9%) of 5,5-dimethyl-[(4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methylthio]-2-isoxazoline.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.81 (1H,s), 4.44 (2H,d), 4.12 (3H,s), 2.81 (2H,s), 1.45 (6H,s)

<Example 40>

Production of 5,5-dimethyl-[(4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methylsulfonyl]-2-isoxazoline (present compound No. 8-0001)

[0258] 1.05 g of m-chloroperbenzoic acid (purity: 70%, 4.28 mmoles) was added, with ice-cooling, to a solution of 0.55 g (1.71 mmoles) of 5,5-dimethyl-[(4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methylthio]-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 4 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and then dried over anhy-

drous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 0.45 g (yield: 75.0%) of 5,5-dimethyl-[(4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methylsulfonyl]-2-isoxazoline as white feather-like crystals (melting point: 175 to 176°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.89 (1H,s), 5.00 (2H,d), 4.11 (3H,s), 3.11 (2H,s), 1.53 (6H,s)

<Example 41>

Production of 3-(5,5-dimethyl-2-isoxazolin-3-ylthiomethyl)-2-trifluoromethyl-6,7-dihydro-5H-pyrazolo[5,1-b][1,3]oxazine (present compound No. 3-0033)

[0259] A solution of 0.82 g (2.3 mmoles) of 3-[5-chloro-1-(3-hydroxypropyl)-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio]-5,5-dimethyl-2-isoxazole dissolved in 5 ml of N,N-dimethylformamide was dropwise added to a suspension of 0.11 g (2.8 mmoles) of sodium hydride in 15 ml of N,N-dimethylformamide. After the completion of the dropwise addition, the resulting mixture was stirred at room temperature for 30 minutes, then heated to 100°C, and stirred for 1 hour to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous citric acid solution and an aqueous sodium chloride solution, and then dried over magnesium sulfate. The resulting solution was subjected to vacuum distillation to obtain 0.77 g (yield: 100%) of 3-(5,5-dimethyl-2-isoxazolin-3-ylthiomethyl)-2-trifluoromethyl-6,7-dihydro-5H-pyrazolo[5,1-b][1,3]oxazine.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.37 (2H,t), 4.19 (2H,t), 4.15 (2H,s), 2.80 (2H,s), 2.31 (2H,m), 1.42 (6H,s)

<Example 42>

Production of 3-(5,5-dimethyl-2-isoxazolin-3-ylsulfonylmethyl)-2-trifluoromethyl-6,7-dihydro-5H-pyrazolo[5,1-b][1,3]oxazine (present compound No. 3-0019)

[0260] 1.25 g of m-chloroperbenzoic acid (purity: 70%, 5.1 mmoles) was added, with ice-cooling, to a solution of 0.77 g (2.3 mmoles) of 3-(6,7-dihydro-3-trifluoromethyl-5H-pyrazolo[5,1-b][1,3]oxazin-4-yl-methylthio)-5,5-dimethyl-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 0.36 g (yield: 43%) of 3-(5,5-dimethyl-2-isoxazolin-3-ylsulfonylmethyl)-2-trifluoromethyl-6,7-dihydro-5H-pyrazolo[5,1-b][1,3]oxazine as a white powder (melting point: 151.0 to 152.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.47 (2H,s), 4.40 (2H,t), 4.23 (2H,t), 3.09 (2H,s), 2.34 (2H,m), 1.50 (6H,s)

[0261] Compound numbers shown in Tables 11 to 20 are referred to in the Examples.

Table 11




												
Compound No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Z ₁	R ²²	R ²³	R ²⁴	Melting point(°C) or refractive index (n _D ²⁰)
1-0001	Me	Me	H	H	2	H	H	S	Me	H	H	66-68
1-0002	Me	Me	H	H	2	H	H	S	Cl	Me	H	87-88
1-0003	Me	Me	H	H	2	H	H	S	H	H	Me	95-97
1-0004	Me	Me	H	H	2	H	H	S	Cl	H	H	70-72
1-0005	Me	Me	H	H	2	H	H	S	H	H	Cl	118-119
1-0006	Me	Me	H	H	2	H	H	O	H	H	H	Impossible to measure
1-0007	Me	Me	H	H	2	H	H	O	H	H	C(=O)OMe	124-125

Table 12



Compound No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Z ²	R ²⁵	R ²⁶	R ²⁷	Melting point(°C) or refractive index (n _D ²⁰)
2-0001	Me	Me	H	H	2	H	H	S	Me	C(=NOMe)Me	Me	95-96
2-0002	Me	Me	H	H	0	H	H	S	Me	C(=NOMe)Me	Me	
2-0003	Me	Me	H	H	2	H	H	S	H	H	H	99-101
2-0004	Me	Me	H	H	2	H	H	S	H	Ome	H	96-97
2-0005	Me	Me	H	H	2	H	H	S	Cl	H	Cl	125-127
2-0006	Me	Me	H	H	2	H	H	S	Cl	Cl	Cl	158-160
2-0007	Me	Me	H	H	2	H	H	S	Me	Me	Me	117-117
2-0008	Me	Me	H	H	2	H	H	S	Me	C(=O)Me	Me	146-148
2-0009	Me	Me	H	H	2	H	H	S	Ph	C(=O)Me	Me	1.5730
2-0010	Me	Me	H	H	2	H	H	S	Ph	C(=NOMe)Me	Me	129-131
2-0011	Me	Me	H	H	2	H	H	S	Cl	C(=O)Ome	Cl	157-158
2-0012	Me	Me	H	H	2	H	H	S	Cl	C(=O)NHMe	Cl	178-180
2-0013	Me	Me	H	H	2	H	H	O	H	H	H	58-61
2-0014	Me	Me	H	H	2	H	H	O	Me	H	Cl	180-181

Table 13

											
Compound No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	R ²⁹	R ²⁸	R ³⁰	Melting point(°C) or refractive index (n _D ²⁰)
3-0001	Me	Me	H	H	0	H	H	CF ₃	Ph	Cl	89-90
3-0002	Me	Me	H	H	2	H	H	CF ₃	Ph	Cl	132-133
3-0003	Me	Me	H	H	1	H	H	Ph	Me	Cl	Impossible to measure
3-0004	Me	Me	H	H	2	H	H	CF ₃	Ph	SO ₂ Et	158-160
3-0005	Me	Me	H	H	2	H	H	CF ₃	Ph	N(Me) ₂	150-151
3-0006	Me	Me	H	H	0	H	H	CF ₃	Bu-t	Cl	79-81
3-0007	Me	Me	H	H	0	H	H	CF ₃	H	Cl	120-122
3-0008	Me	Me	H	H	0	H	H	CF ₃	CHF ₂	Cl	41-42
3-0009	Me	Me	H	H	0	H	H	Cl	CHF ₂	CF ₃	89-90
3-0010	Me	Me	H	H	2	H	H	CF ₃	CHF ₂	Cl	126-127
3-0011	Me	Me	H	H	2	H	H	Cl	CHF ₂	CF ₃	136-137
3-0012	Me	Me	H	H	2	H	H	OEt	Me	CF ₃	124-125
3-0013	Me	Me	H	H	2	H	H	CF ₃	Me	OMe	113-114
3-0014	Me	Me	H	H	2	H	H	CF ₃	Me	O(2-Cl)Ph	67-70
3-0015	Me	Me	H	H	2	H	H	CF ₃	Me	OPen-c	113-114
3-0016	Me	Me	H	H	2	H	H	CF ₃	Me	CN	105-108
3-0017	Me	Me	H	H	2	H	H	Cl	Et	Cl	105-107
3-0018	Me	Me	H	H	2	H	H	CHF ₂	Me	Cl	78-79
3-0019	Me	Me	H	H	2	H	H	CF ₃	-(CH ₂) ₃ O-		151-152
3-0020	Me	Me	H	H	0	H	H	CHF ₂	Me	Cl	1.5183
3-0021	Me	Me	H	H	0	H	H	CF ₃	Ph	F	
3-0022	Me	Me	H	H	0	H	H	CF ₃	Ph	SEt	
3-0023	Me	Me	H	H	0	H	H	CF ₃	Ph	N(Me) ₂	
3-0024	Me	Me	H	H	0	H	H	OMe	Me	CF ₃	
3-0025	Me	Me	H	H	0	H	H	OH	Me	CF ₃	
3-0026	Me	Me	H	H	0	H	H	OEt	Me	CF ₃	
3-0027	Me	Me	H	H	0	H	H	CF ₃	Me	F	

	3-0028	Me	Me	H	H	0	H	H	CF ₃	Me	OMe	
	3-0029	Me	Me	H	H	0	H	H	CF ₃	Me	O(2-Cl)Ph	
5	3-0030	Me	Me	H	H	0	H	H	CF ₃	Me	OPen-c	
	3-0031	Me	Me	H	H	0	H	H	CF ₃	Me	CN	
	3-0032	Me	Me	H	H	0	H	H	Cl	Et	Cl	
	3-0033	Me	Me	H	H	0	H	H	CF ₃	-(CH ₂) ₃ O-		
10	3-0034	Me	Me	H	H	2	H	H	CF ₃	H	Cl	138-140
	3-0035	Me	Me	H	H	2	H	H	H	Me	Cl	105-106
	3-0036	Me	Me	H	H	2	H	H	Me	Me	Me	148-150
15	3-0037	Me	Me	H	H	2	H	H	Me	Me	Cl	99-101
	3-0038	Me	Me	H	H	2	H	H	Cl	Me	Cl	143-145
	3-0039	Me	Me	H	H	2	H	H	CF ₃	Me	Cl	115-116
	3-0040	Me	Me	H	H	2	H	H	Cl	Me	CF ₃	120-122
20	3-0041	Me	Me	H	H	2	H	H	CF ₃	Me	F	79-82
	3-0042	Me	Me	H	H	2	H	H	CF ₃	Me	OH	90-92
	3-0043	Me	Me	H	H	2	H	H	OMe	Me	CF ₃	125-126
25	3-0044	Me	Me	H	H	2	H	H	CF ₃	Me	OEt	92-94
	3-0045	Me	Me	H	H	2	H	H	CF ₃	Me	OPr-i	69-71
	3-0046	Me	Me	H	H	2	H	H	CF ₃	Me	OPr	82-83
	3-0047	Me	Me	H	H	2	H	H	CF ₃	Me	OBu-t	86-89
30	3-0048	Me	Me	H	H	2	H	H	CF ₃	Me	OBu	61-62
	3-0049	Me	Me	H	H	2	H	H	CF ₃	Me	OHex-c	124-125
	3-0050	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ Pr-c	93-94
	3-0051	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ Pen-c	112-113
35	3-0052	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ Hex-c	56-59
	3-0053	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ C≡CH	92-93
	3-0054	Me	Me	H	H	2	H	H	CF ₃	Me	OCHF ₂	129-130
40	3-0055	Me	Me	H	H	2	H	H	OCHF ₂	Me	CF ₃	Impossible to measure
	3-0056	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ CHF ₂	89-91
	3-0057	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ CF ₃	93-95
45	3-0058	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ CN	1.4872
	3-0059	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ Ph	79-81
	3-0060	Me	Me	H	H	2	H	H	CF ₃	Me	OPh	122-123
	3-0061	Me	Me	H	H	2	H	H	CF ₃	Me	O(3-Cl)Ph	Impossible to measure
50	3-0062	Me	Me	H	H	2	H	H	CF ₃	Me	O(3-OMe)Ph	1.5059
	3-0063	Me	Me	H	H	2	H	H	CF ₃	Me	O(4-Cl)Ph	68-69
	3-0064	Me	Me	H	H	2	H	H	CF ₃	Me	O(4-Me)Ph	132-133
55	3-0065	Me	Me	H	H	2	H	H	CF ₃	Me	O(4-OMe)Ph	115-117

	3-0066	Me	Me	H	H	2	H	H	CF ₃	Me	OC(=O)Me	130-131
	3-0067	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ Me	168-169
5	3-0068	Me	Me	H	H	2	H	H	CF ₃	Me	SEt	100-102
	3-0069	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ Et	107-108
	3-0070	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ Ph	166-168
10	3-0071	Me	Me	H	H	2	H	H	CF ₃	Me	Me	105-107
	3-0072	Me	Me	H	H	2	H	H	Ph	Me	Cl	127-129
	3-0073	Me	Me	H	H	2	H	H	CF ₃	Et	Cl	111-112
	3-0074	Me	Me	H	H	2	H	H	Cl	Et	CF ₃	112-114
15	3-0075	Me	Me	H	H	2	H	H	CF ₃	Pr-i	Cl	157-158
	3-0076	Me	Me	H	H	2	H	H	Cl	Pr-i	CF ₃	135-136
	3-0077	Me	Me	H	H	2	H	H	CF ₃	Pr	Cl	89-90
	3-0078	Me	Me	H	H	2	H	H	Cl	Pr	CF ₃	111-113
20	3-0079	Me	Me	H	H	2	H	H	CF ₃	Bu-t	H	101-103
	3-0080	Me	Me	H	H	2	H	H	CF ₃	Bu-t	Cl	118-119
	3-0081	Me	Me	H	H	2	H	H	CF ₃	Bu-s	Cl	110-112
25	3-0082	Me	Me	H	H	2	H	H	Cl	Bu-s	CF ₃	110-111
	3-0083	Me	Me	H	H	2	H	H	CF ₃	Bu-i	Cl	96-98
	3-0084	Me	Me	H	H	2	H	H	Cl	Bu-i	CF ₃	140-141
	3-0085	Me	Me	H	H	2	H	H	CF ₃	Bu	Cl	89-90
30	3-0086	Me	Me	H	H	2	H	H	Cl	Bu	CF ₃	108-110
	3-0087	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Ph	Cl	132-133
	3-0088	Me	Me	H	H	2	H	H	Cl	CH ₂ Ph	CF ₃	118-120
	3-0089	Me	Me	H	H	2	H	H	CF ₃	Pen-c	Cl	130-131
35	3-0090	Me	Me	H	H	2	H	H	Cl	Pen-c	CF ₃	147-148
	3-0091	Me	Me	H	H	2	H	H	CF ₃	Hex-c	Cl	151-152
	3-0092	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	Cl	93-95
40	3-0093	Me	Me	H	H	2	H	H	Cl	CH ₂ Pr-c	CF ₃	129-130
	3-0094	Me	Me	H	H	2	H	H	CF ₃	1-cyclopropylethyl	Cl	87-89
	3-0095	Me	Me	H	H	2	H	H	Cl	1-cyclopropylethyl	CF ₃	121-123
	3-0096	Me	Me	H	H	2	H	H	CF ₃	CH ₂ (2-Methylcyclopropyl)	Cl	102-103
45	3-0097	Me	Me	H	H	2	H	H	Cl	CH ₂ (2-Methylcyclopropyl)	CF ₃	118-119
	3-0098	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Bu-c	Cl	94-96
	3-0099	Me	Me	H	H	2	H	H	Cl	CH ₂ Bu-c	CF ₃	141-142
50	3-0100	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pen-c	Cl	127-129
	3-0101	Me	Me	H	H	2	H	H	Cl	CH ₂ Pen-c	CF ₃	146-149
	3-0102	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Hex-c	Cl	152-154
	3-0103	Me	Me	H	H	2	H	H	Cl	CH ₂ Hex-c	CF ₃	115-117
55	3-0104	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH=CH ₂	Cl	78-80

	3-0105	Me	Me	H	H	2	H	H	Cl	CH ₂ CH=CH ₂	CF ₃	105-106
	3-0106	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C≡CH	Cl	73-74
5	3-0107	Me	Me	H	H	2	H	H	Cl	CH ₂ C≡CH	CF ₃	108-109
	3-0108	Me	Me	H	H	2	H	H	CF ₃	CHMeC≡CH	Cl	95-96
	3-0109	Me	Me	H	H	2	H	H	Cl	CHMeC≡CH	CF ₃	116-118
10	3-0110	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C≡CMe	Cl	114-115
	3-0111	Me	Me	H	H	2	H	H	Cl	CH ₂ C≡CMe	CF ₃	115-116
	3-0112	Me	Me	H	H	2	H	H	CF ₃	CHF ₂	OMe	72-74
	3-0113	Me	Me	H	H	2	H	H	OMe	CHF ₂	CF ₃	108-109
15	3-0114	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CHF ₂	Cl	99-100
	3-0115	Me	Me	H	H	2	H	H	Cl	CH ₂ CHF ₂	CF ₃	107-109
	3-0116	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CF ₃	Cl	135-136
	3-0117	Me	Me	H	H	2	H	H	Cl	CH ₂ CF ₃	CF ₃	112-115
20	3-0118	Me	Me	H	H	2	H	H	CF ₃	CH ₂ OMe	Cl	87-89
	3-0119	Me	Me	H	H	2	H	H	Cl	CH ₂ OMe	CF ₃	125-128
	3-0120	Me	Me	H	H	2	H	H	CF ₃	CH ₂ OEt	Cl	97-98
25	3-0121	Me	Me	H	H	2	H	H	Cl	CH ₂ OEt	CF ₃	128-129
	3-0122	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ OH	Cl	79-81
	3-0123	Me	Me	H	H	2	H	H	Cl	CH ₂ CH ₂ OH	CF ₃	93-94
	3-0124	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ OMe	Cl	102-104
30	3-0125	Me	Me	H	H	2	H	H	Cl	CH ₂ CH ₂ OMe	CF ₃	118-119
	3-0126	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ OEt	Cl	56-59
	3-0127	Me	Me	H	H	2	H	H	Cl	CH ₂ CH ₂ OEt	CF ₃	118-119
	3-0128	Me	Me	H	H	2	H	H	CF ₃	CH ₂ SMe	Cl	103-105
35	3-0129	Me	Me	H	H	2	H	H	Cl	CH ₂ SMe	CF ₃	128-129
	3-0130	Me	Me	H	H	2	H	H	CF ₃	CH ₂ SO ₂ Me	Cl	157-159
	3-0131	Me	Me	H	H	2	H	H	Cl	CH ₂ SO ₂ Me	CF ₃	165-166
40	3-0132	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ SO ₂ Me	Cl	155-157
	3-0133	Me	Me	H	H	2	H	H	Cl	CH ₂ CH ₂ SO ₂ Me	CF ₃	166-168
	3-0134	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CN	Cl	128-129
	3-0135	Me	Me	H	H	2	H	H	Cl	CH ₂ CN	CF ₃	117-118
45	3-0136	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=O)OEt	Cl	127-129
	3-0137	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=O)OEt	CF ₃	143-145
	3-0138	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=O)NH ₂	Cl	173-174
50	3-0139	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=O)NH ₂	CF ₃	182-183
	3-0140	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=O)N(Me) ₂	Cl	142-143
	3-0141	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=O)N(Me) ₂	CF ₃	181-182
	3-0142	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=O)Me	Cl	148-149
55	3-0143	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=O)Me	CF ₃	163-164

EP 1 364 946 A1


	3-0144	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ C(=O)Me	Cl	89-91
	3-0145	Me	Me	H	H	2	H	H	Me	Ph	Me	140-141
5	3-0146	Me	Me	H	H	2	H	H	Me	Ph	Cl	124-125
	3-0147	Me	Me	H	H	2	H	H	Et	Ph	Cl	112-113
	3-0148	Me	Me	H	H	2	H	H	Pr	Ph	Cl	122-123
	3-0149	Me	Me	H	H	2	H	H	Pr-i	Ph	Cl	116-117
10	3-0150	Me	Me	H	H	2	H	H	Bu-t	Ph	Cl	100-102
	3-0151	Me	Me	H	H	2	H	H	CF ₃	Ph	H	111-112
	3-0152	Me	Me	H	H	2	H	H	CF ₃	Ph	Me	129-132
15	3-0153	Me	Me	H	H	2	H	H	CF ₃	Ph	CF ₃	112-113
	3-0154	Me	Me	H	H	2	H	H	CF ₃	Ph	F	90-91
	3-0155	Me	Me	H	H	2	H	H	CF ₃	Ph	OMe	104-106
	3-0156	Me	Me	H	H	2	H	H	CF ₃	Ph	OEt	129-131
20	3-0157	Me	Me	H	H	2	H	H	CF ₃	Ph	OPr-i	86-88
	3-0158	Me	Me	H	H	2	H	H	CF ₃	Ph	OPr	117-118
	3-0159	Me	Me	H	H	2	H	H	CF ₃	Ph	OBu-t	105-108
	3-0160	Me	Me	H	H	2	H	H	CF ₃	Ph	OCHF ₂	90-92
25	3-0161	Me	Me	H	H	2	H	H	CF ₃	Ph	SO ₂ Me	167-168
	3-0162	Me	Me	H	H	2	H	H	CF ₃	Ph	CN	113-115
	3-0163	Me	Me	H	H	2	H	H	CF ₃	(2-Cl)Ph	Cl	153-154
30	3-0164	Me	Me	H	H	2	H	H	CF ₃	(3-Cl)Ph	Cl	106-107
	3-0165	Me	Me	H	H	2	H	H	CF ₃	(4-Cl)Ph	Cl	142-143
	3-0166	Me	Me	H	H	2	H	H	CF ₃	(4-F)Ph	Cl	135-138
	3-0167	Me	Me	H	H	2	H	H	CF ₃	(4-OMe)Ph	Cl	136-138
35	3-0168	Me	Me	H	H	2	H	H	CF ₃	(4-Me)Ph	Cl	129-130
	3-0169	Me	Me	H	H	2	H	H	CF ₃	(4-NO ₂)Ph	Cl	145-147
	3-0170	Me	Me	H	H	2	H	H	CF ₃	(4-CN)Ph	Cl	91-93
40	3-0171	Me	Me	H	H	2	H	H	CF ₃	(4-C(=O)Me)Ph	Cl	133-135
	3-0172	Me	Me	H	H	2	H	H	CF ₃	(4-C(=O)OMe)Ph	Cl	121-124
	3-0173	Me	Me	H	H	2	H	H	CF ₃	Pyrimidin-2-yl	Cl	148-150
	3-0174	Me	Me	H	H	2	H	H	CF ₃	4,6-Dimethoxypyrimidin-2-yl	Cl	117-118
45	3-0175	Me	Me	H	H	2	H	H	CF ₃	SO ₂ Me	Cl	146-148
	3-0176	Me	Me	H	H	2	H	H	CF ₃	SO ₂ Ph	Cl	145-148
	3-0177	Me	Me	H	H	2	H	H	CF ₃	C(=O)Me	Cl	130-131
50	3-0178	Me	Me	H	H	2	H	H	CF ₃	C(=O)Ph	Cl	114-117
	3-0179	Me	Me	H	H	2	H	H	CF ₃	C(=O)OMe	Cl	104-106
	3-0180	Me	Et	H	H	2	H	H	CF ₃	Me	Cl	108-110
	3-0181	Me	Me	H	H	0	H	H	CHF ₂	Me	Cl	1.5183
55	3-0182	Me	Me	H	H	0	H	H	Ph	Me	Cl	76-77

3-0183	Me	Me	H	H	0	H	H	CF ₃	Bu-t	OMe	1.4831
3-0184	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C(=O)NH ₂	Cl	179-180
3-0185	Me	Me	H	H	0	H	H	Me	Ph	Cl	58-60

Table 14

R ²	R ³	R ⁴	n	R ⁵	R ⁶	Z ³	R ³¹	R ³²	Melting point(°C) or refractive index (n _D ²⁰)
Me	H	H	2	H	H	O	CF ₃	Me	135-136
Me	H	H	2	H	H	S	Me	Cl	113-114
Me	H	H	0	H	H	O	CF ₃	Me	
Me	H	H	0	H	H	S	Me	Cl	
Me	H	H	2	H	H	O	Me	Me	178-179
Me	H	H	2	H	H	O	CF ₃	OEt	89-91
Me	H	H	2	H	H	O	Ph	Me	81-83
Me	H	H	2	H	H	S	Me	OEt	109-111

Table 15



Compound No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Z ⁴	R ³³	R ³⁴	Melting point(°C) or refractive index (n _D ²⁰)
5-0001	Me	Me	H	H	2	H	H	NMe	Cl	Me	114-115
5-0002	Me	Me	H	H	2	H	H	NMe	Cl	Et	107-108
5-0003	Me	Me	H	H	2	H	H	NMe	CF ₃	H	142-143
5-0004	Me	Me	H	H	2	H	H	NCHF ₂	-(CH ₂) ₄ -		123-125
5-0005	Me	Me	H	H	2	H	H	NPh	Oet	Me	1.5397
5-0006	Me	Me	H	H	2	H	H	NPh	OCHF ₂	Me	1.5339
5-0007	Me	Me	H	H	2	H	H	NPh	CF ₃	H	99-101
5-0008	Me	Me	H	H	2	H	H	NPh	OCH ₂ CH=CH ₂	Me	87-90
5-0009	Me	Me	H	H	1	H	H	NPh	OCH ₂ CH=CH ₂	Me	1.5702

Table 16


											
Compound No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Z ⁵	R ³⁵	R ³⁶	Melting point(°C) or refractive index (n _D ²⁰)
6-0001	Me	Me	H	H	2	H	H	NCHF ₂	-(CH ₂) ₄ -		Impossible to measure
6-0002	Me	Me	H	H	2	H	H	NPh	H	Oet	107-108
6-0003	Me	Me	H	H	2	H	H	NPh	H	OCHF ₂	1.5383
6-0004	Me	Me	H	H	2	H	H	O	Me	H	100-102
6-0005	Me	Me	H	H	0	H	H	NCHF ₂	-(CH ₂) ₄ -		1.5264

Table 17


													
Compound No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	R ³⁷	R ³⁸	R ³⁹	R ⁴⁰		Melting point(°C) or refractive index (n _D ²⁰)
7-0001	Me	Me	H	H	2	H	H	H	CF ₃	H	H	-	77-80
7-0002	Me	Me	H	H	2	H	H	H	CF ₃	H	H	N-oxide	114-116
7-0003	Me	Me	H	H	0	H	H	H	CF ₃	H	H	-	
7-0004	Me	Me	H	H	2	H	H	H	H	H	H	-	130-131
7-0005	Me	Me	H	H	2	H	H	H	H	H	H	N-oxide	166-168
7-0006	Me	Me	H	H	2	H	H	Cl	Ph	H	H	-	118-120
7-0007	Me	Me	H	H	2	H	H	OMe	Ph	H	H	-	105-106
7-0008	Me	Me	H	H	2	H	H	Cl	Me	H	H	-	115-116
7-0009	Me	Me	H	H	2	H	H	OMe	Me	H	H	-	134-135
7-0010	Me	Me	H	H	2	H	H	Me	Me	H	H	N-oxide	198-199
7-0011	Me	Me	H	H	2	H	H	Ph	Ph	H	H	-	161-162
7-0012	Me	Me	H	H	1	H	H	H	H	H	H	-	97-99
7-0013	Me	Me	H	H	0	H	H	(2-Chloropyridin-3-yl)methylthio	H	H	H	-	154-155

Table 18


										
R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	R ⁴¹	R ⁴²	R ⁴³	Melting point(°C) or refractive index (n _D ²⁰)
Me	Me	H	H	2	H	H	H	OMe	CF ₃	175-176
Me	Me	H	H	0	H	H	H	OMe	CF ₃	
Me	Me	H	H	2	H	H	H	Cl	Cl	119-120
Me	Me	H	H	2	H	H	H	OEt	CF ₃	94-95
Me	Me	H	H	2	H	H	H	OMe	OMe	186-187
Me	Me	H	H	2	H	H	Me	OMe	CF ₃	143-144
Me	Me	H	H	2	H	H	OMe	OMe	CF ₃	144-145
Me	Me	H	H	2	H	H	SMe	OMe	CF ₃	160-162
Me	Me	H	H	2	H	H	SO ₂ Me	OMe	CF ₃	144-146
Me	Me	H	H	2	H	H	NH ₂	OMe	CF ₃	208-209
Me	Me	H	H	2	Pr-i	H	H	H	CF ₃	112-113
Me	Me	H	H	0	Pr-i	H	H	H	CF ₃	1.4986

Table 19



									
Compound No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Y ¹	Melting point(°C) or refractive index (n _D ²⁰)
9-0001	Me	Me	H	H	2	H	H	Pyridin-2-yl	116-118
9-0002	Me	Me	H	H	2	H	H	Pyridin-2-yl 1-oxide	140-143
9-0003	Me	Me	H	H	2	H	H	Pyridin-4-yl	133-136
9-0004	Me	Me	H	H	2	H	H	Pyridin-4-yl 1-oxide	110-113
9-0005	Me	Me	H	H	2	H	H	1,2,4-Oxadiazol-3-yl	Impossible to measure
9-0006	Me	Me	H	H	2	H	H	3-Phenyl-1,2,4-oxadiazol-5-yl	153-154
9-0007	Me	Me	H	H	2	H	H	3-Benzyl-1,2,4-oxadiazol-5-yl	108-109
9-0008	Me	Me	H	H	2	H	H	2-Chlorothiazol-4-yl	110-112
9-0009	Me	Me	H	H	2	H	H	1,4-Dimethylimidazol-5-yl	163-164
9-0010	Me	Me	H	H	1	H	H	Pyridin-2-yl	81-82
9-0011	Me	Me	H	H	1	H	H	Pyridin-4-yl	94-96
9-0012	Me	Me	H	H	1	H	H	1,4-Dimethylimidazol-5-yl	138-140
9-0013	Me	Me	H	H	0	H	H	1,4-Dimethylimidazol-5-yl	1.5427

Table 20

									
Compound No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Y ¹	Melting point(°C) or refractive index (n _D ²⁰)
10-0001	Me	Me	H	H	2	H	H	Benzimidazol-2-yl	171-174
10-0002	Me	Me	H	H	2	H	H	Benzothiophen-2-yl	181-183
10-0003	Me	Me	H	H	2	H	H	3-Chlorobenzothiophen-2-yl	109-112
10-0004	Me	Me	H	H	2	H	H	Benzotriazol-1-yl	206-207
10-0005	Me	Me	H	H	2	H	H	1-Methylindazol-4-yl	128-130
10-0006	Me	Me	H	H	2	H	H	Benzothiazol-2-yl	142-143
10-0007	Me	Me	H	H	2	H	H	Benzothiophen-3-yl	188-191
10-0008	Me	Me	H	H	2	H	H	5-Chlorobenzothiophen-3-yl	129-130
10-0009	Me	Me	H	H	2	H	H	Benzoxazol-2-yl	127-129
10-0010	Me	Me	H	H	2	H	H	3-Methylbenzothiophen-2-yl	161-163
10-0011	Me	Me	H	H	2	H	H	3-Bromobenzothiophen-2-yl	118-119
10-0012	Me	Me	H	H	2	H	H	Benzofuran-2-yl	123-124
10-0013	Me	Me	H	H	2	H	H	2-Methylbenzofuran-7-yl	135-137
10-0014	Me	Me	H	H	2	H	H	3-Bromobenzofuran-2-yl	107-108
10-0015	Me	Me	H	H	2	H	H	Benzothiophen-7-yl	95-97
10-0016	Me	Me	H	H	2	H	H	1-Methylindazol-7-yl	89-90
10-0017	Me	Me	H	H	2	H	H	3-Methylbenzofuran-2-yl	111-112
10-0018	Me	Me	H	H	2	H	H	3-Chloro-1-methylindol-2-yl	162-165

(Production Examples of intermediates)

<Reference Example 1>

Production of 3-chloro-5,5-dimethyl-2-isoxazoline

[0262] 534.0 g (4.0 moles) of N-chlorosuccinimide was gradually added, at 65 to 70°C, to a solution of 182.7 g (2.05 moles) of glyoxylic acid aldoxime dissolved in 2 liters of 1,2-dimethoxyethane. The mixture was refluxed for 1 hour with heating. Thereto were added, with ice-cooling, 1,440.0 g (14.4 moles) of potassium hydrogencarbonate and 10 ml of water. Then, 360.0 g (6.4 moles) of 2-methylpropene was added. The resulting mixture was stirred at room temperature for 24 hours to give rise to a reaction. The reaction mixture was poured into water, followed by extraction with diisopropyl ether. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and

then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 107.7 g (yield: 40.0%) of 3-chloro-5,5-dimethyl-2-isoxazoline as a yellow viscous liquid.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

2.93 (2H,s), 1.47 (6H,s)

<Reference Example 2>

Production of 3-chloro-5-ethyl-5-methyl-2-isoxazoline

[0263] 61.9 g (463.4 mmoles) of N-chlorosuccinimide was gradually added, at 60°C, to a solution of 20.6 g (231.7 mmoles) of glyoxylic acid aldoxime dissolved in 500 ml of 1,2-dimethoxyethane. After the addition, the mixture was refluxed for 10 minutes with heating. Thereto were added, with ice-cooling, 50 ml (463.4 mmoles) of 2-methyl-1-butene, 98.9 g (1,622 mmoles) of potassium hydrogencarbonate and 10 ml of water. The resulting mixture was stirred for 12 hours to give rise to a reaction. The reaction mixture was poured into water, followed by extraction with n-hexane. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 13.9 g (yield: 40.6%) of 3-chloro-5-ethyl-5-methyl-2-isoxazoline as a yellow viscous liquid.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

2.91 (2H, ABq, J=17.0, Δv=46.1 Hz), 1.73 (2H,q), 1.42 (3H,s), 0.96 (3H,t)

<Reference Example 3>

Production of 3-benzylthio-5,5-dimethyl-2-isoxazoline

[0264] 3.2 g (23.2 mmoles) of anhydrous potassium carbonate and 3.0 g (22.5 mmoles) of 3-chloro-5,5-dimethyl-2-isoxazoline were added, in a nitrogen atmosphere, to a solution of 2.8 g (22.5 mmoles) of benzylmercaptan dissolved in 50 ml of N,N-dimethylformamide. The mixture was stirred at 100°C for 2 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 3.1 g (yield: 62.0%) of 3-benzylthio-5,5-dimethyl-2-isoxazoline as a yellow oily substance (n_D²⁰ = 1.5521).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.24-7.39 (5H,m), 4.26 (2H,s), 2.77 (2H,s), 1.40 (6H,s)

<Reference Example 4>

Production of 3-(2,6-difluorobenzylsulfinyl)-5-ethyl-5-methyl-2-isoxazoline

[0265] 4.6 g of m-chloroperbenzoic acid (purity: 70%, 18.8 mmoles) was added, with ice-cooling, to a solution of 4.1 g (15.0 mmoles) of 3-(2,6-difluorobenzylthio)-5-ethyl-5-methyl-2-isoxazoline dissolved in 50 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After the completion of the reaction the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 1.5 g (yield: 34.8%) of 3-(2,6-difluorobenzylsulfinyl)-5-ethyl-5-methyl-2-isoxazoline as a white powder (melting point: 30°C or less).

¹H-NMR [CDCl₃/TMS, δ (ppm)] :

7.39-7.28 (1H,m), 7.03-6.94 (2H,m), 4.38 (2H,s), 3.04 (1H, ABq, J=17.2, Δv=85.7 Hz), 3.12 (1H,s), 1.75 (2H,m), 1.44 (3H,S)+1.41 (3H,s), 0.97 (3H,m)

<Reference Example 5>

Production of 3-(2,6-difluorobenzylsulfonyl)-5-ethyl-5-methyl-2-isoxazoline

[0266] 1.0 g of m-chloroperbenzoic acid (purity: 70%, 4.1 mmoles) was added, with ice-cooling, to a solution of 0.8

g (2.8 mmoles) of 3-(2,6-difluorobenzylsulfinyl)-5-ethyl-5-methyl-2-isoxazoline dissolved in 50 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.6 g (yield: 75.0%) of 3-(2, 6-difluorobenzylsulfonyl)-5-ethyl-5-methyl-2-isoxazoline as a white powder (melting point: 64 to 65°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.36-7.46 (1H,m), 6.98-7.04 (2H,m), 4.73 (2H,s), 3.04 (2H, ABq, J=17.2, Δv=51.1 Hz), 1.77 (2H,q), 1.46 (3H,s), 0.97 (3H,t)

<Reference Example 6>

Production of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline

[0267] 1.0 kg of an aqueous sodium methanethiolate solution (content: 15%, 2.14 mmoles) was dropwise added, with ice-cooling, to a solution of 143.0 g (1.07 moles) of 3-chloro-5,5-dimethyl-2-isoxazoline dissolved in 500 ml of N, N-dimethylformamide. The mixture was stirred at room temperature for 12 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 115.0 g (yield: 74.1%) of 5,5-dimethyl-3-methylthio-2-isoxazoline. This residue (741.2 mmoles) was dissolved in 1 liter of chloroform. Thereto was added, with ice-cooling, 392.0 g of m-chloroperbenzoic acid (purity: 70%, 1.59 moles). The resulting mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After the completion of the reaction, the separated m-chloroperbenzoic acid was removed by filtration. The resulting filtrate was washed with an aqueous sodium hydrogensulfite solution, water, an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was washed with diisopropyl ether to obtain 77.6 g (yield: 59.1%) of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline as a white powder (melting point: 82 to 84°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

3.26 (3H,s), 3.12 (2H,s), 1.51 (6H,s)

<Reference Example 7>

Production of 5,5-dimethyl-3-ethylthio-2-isoxazoline

[0268] 1,500 ml of an aqueous solution containing 560.0 g (9.0 moles) of ethyl mercaptan and 360.0 g (9.0 moles) of sodium hydroxide was added to a solution containing 3-chloro-5,5-dimethyl-2-isoxazoline. The mixture was stirred at 60 to 70°C for 16 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 270.0 g of crude 5,5-dimethyl-3-ethylthio-2-isoxazoline as a dark red oily substance.

<Reference Example 8>

Production of 5,5-dimethyl-3-ethylsulfonyl-2-isoxazoline

[0269] 270.0 g (1.7 moles) of crude oily 5,5-dimethyl-3-ethylthio-2-isoxazoline was dissolved in 1.0 liter of chloroform. Thereto was added, with ice-cooling, 1,050 g of m-chloroperbenzoic acid (purity: 70%, 6.1 moles). The resulting mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the separated m-chloroperbenzoic acid was removed by filtration. The resulting filtrate was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was washed with n-hexane to obtain 133.6 g (yield: 65.4%) of 5,5-dimethyl-3-ethylsulfonyl-2-isoxazoline as a white powder.

<Reference Example 9>

Production of 1-phenyl-3-trifluoromethyl-1H-pyrazol-5-ol

[0270] 20 g (184.9 mmoles) of phenylhydrazine and 4 ml of concentrated hydrochloric acid were added to a solution of 34.1 g (184.9 mmoles) of ethyl trifluoroacetoacetate dissolved in 500 ml of ethanol. The mixture was refluxed for 1 hour with heating, to give rise to a reaction. After the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the most part of the solvent contained therein. The residue was mixed with water to precipitate crystals. The crystals were collected by filtration, washed with water until the filtrate became neutral, and dried to obtain 37.1 g (yield: 87.9%) of 1-phenyl-3-trifluoromethyl-1H-pyrazol-5-ol as ochreous crystals.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.68-7.41 (5H,m), 5.86 (1H,s), 3.71 (1H,s)

<Reference Example 10>

Production of 5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde

[0271] 33.6 g (219.1 mmoles) of phosphorus oxychloride was added to 7.7 g (105.2 mmoles) of N,N-dimethylformamide with ice-cooling. Thereto was added, at room temperature, 20 g (87.7 mmoles) of 1-phenyl-3-trifluoromethyl-1H-pyrazol-5-ol. The resulting mixture was refluxed for 1 hour with heating, to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water with ice-cooling, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 19.1 g (yield: 79.1%) of 5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde as white crystals.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

10.06 (1H,s), 7.57 (5H,s)

<Reference Example 11>

Production of (5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol

[0272] A solution of 0.21 g (5.5 mmoles) of lithium aluminum hydride dissolved in 70 ml of THF was cooled to -30°C. Thereto was gradually added a solution of 3 g (10.9 mmoles) of 5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde dissolved in 30 ml of tetrahydrofuran. The resulting mixture was stirred at -30°C for 30 minutes to give rise to a reaction. After the completion of the reaction, ethyl acetate was added, followed by stirring. Then, water was added, followed by stirring for a while. The reaction mixture was filtered under vacuum. The filtrate was extracted with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 3.0 g (yield: 99.9%) of (5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol as white crystals.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.54-7.51 (5H,m), 4.71 (2H,d) 1.79 (1H,b)

<Reference Example 12>

Production of 4-bromomethyl-5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazole

[0273] A solution of 3.0 g (10.9 mmoles) of (5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol dissolved in 60 ml of diethyl ether was cooled to -10°C. Thereto was added 1.0 g (3.8 mmoles) of phosphorus tribromide. The mixture was stirred at room temperature for 1 hour to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 3.6 g (yield: 95.8%) of 4-bromomethyl-5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazole as white crystals.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.58-7.48 (5H,m), 4.48 (2H,s)

<Reference Example 13>

Production of 5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde

[0274] 10.5 g (180.2 mmoles) of potassium fluoride was added to a solution of 33.0 g (120.1 mmoles) of 5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde dissolved in 500 ml of dimethyl sulfoxide. The mixture was stirred at 100°C for 2 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 26.5 g (yield: 85.0%) of 5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde.

¹H-NMR [CDCl₃/TMS, δ (ppm)] :
9.96 (1H,s), 7.68-7.51 (5H,m)

<Reference Example 14>

Production of (5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol

[0275] To a solution of 1.6 g (41.0 mmoles) of sodium borohydride dissolved in 300 ml of methanol was added, with ice-cooling, a solution of 26.5 g (102.5 mmoles) of 5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde dissolved in 200 ml of methanol. The resulting mixture was stirred at 0°C for 30 minutes to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 28.5 g (yield: 100%) of (5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:
7.65-7.41 (5H,m), 4.68 (2H,d), 1.73 (1H,t)

<Reference Example 15>

Production of 4-bromomethyl-5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazole

[0276] A solution of 27.5 g (105.7 mmoles) of (5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol dissolved in 300 ml of diethyl ether was cooled to 0°C. Thereto was added 10.0 g (37.0 mmoles) of phosphorus tribromide. The mixture was stirred at room temperature for 2 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 30.3 g (yield: 88.8%) of 4-bromomethyl-5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazole.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:
7.66-7.42 (5H,m), 4.44 (2H,s)

<Reference Example 16>

Production of 1-tert-butyl-3-trifluoromethyl-1H-pyrazol-5-ol

[0277] 373.8 g (3.0 moles) of tert-butylhydrazine hydrochloride and 50 ml of concentrated hydrochloric acid were added to a solution of 552.3 g (3.0 moles) of ethyl trifluoroacetoacetate dissolved in 1,500 ml of ethanol. The mixture was refluxed for 2 days with heating, to give rise to a reaction. After the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the most part of the solvent contained therein. The residue was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was washed with n-hexane to obtain 369.0 g (yield: 59.1%) of 1-tert-butyl-3-trifluoromethyl-1H-pyrazol-5-ol as a white powder.

<Reference Example 17>

Production of 1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde

[0278] 462.0 g (3.0 moles) of phosphorus oxychloride was added to 87.7 g (1.2 moles) of N,N-dimethylformamide with ice-cooling. Thereto was added, at room temperature, 208.2 g (1.0 moles) of 1-tert-butyl-3-trifluoromethyl-1H-pyrazol-5-ol. The resulting mixture was refluxed for 10 hours with heating, to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with water, a 5% aqueous sodium hydroxide solution and water in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 131.5 g (yield: 21.7%) of 1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde as white crystals.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

9.97 (1H,d) 1.76 (9H,s)

<Reference Example 18>

Production of (1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol

[0279] A solution of 39.9 g (156.9 mmoles) of (1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde dissolved in 300 ml of methanol was cooled to 0°C. Thereto was gradually added 6.5 g (172.6 mmoles) of sodium borohydride. The mixture was stirred at room temperature for 3 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 37.7 g (yield: 93.6%) of (1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol.

¹H-NMR [CDCl₃/TMS, δ (ppm)] :

4.60 (2H,d), 1.72 (9H,s), 1.58 (1H,t)

<Reference Example 19>

Production of 4-bromomethyl-1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazole

[0280] A solution of 9.2 g (35.7 mmoles) of (1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol dissolved in 100 ml of diethyl ether was cooled to -10°C. Thereto was added 11.6 g (42.9 mmoles) of phosphorus tribromide. The mixture was stirred at room temperature overnight to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 10.0 g (yield: 87.3%) of 4-bromomethyl-1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazole.

<Reference Example 20>

Production of (1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazol-4-yl)-methanethiol

[0281] 43.5 g (136.1 mmoles) of 4-bromomethyl-1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazole was added to a solution of 21.8 g of sodium hydrosulfide hydrate (purity: 70%, 272.2 mmoles) dissolved in 300 ml of N,N-dimethylformamide. The mixture was stirred at room temperature overnight to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 32.3 g (yield: 87.0%) of (1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazol-4-yl)-methanethiol.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

3.65 (2H,d), 1.90 (1H,t), 1.70 (9H,s)

<Reference Example 21>

Production of 1-tert-butyl-5-methoxy-3-trifluoromethyl-1H-pyrazole

[0282] 15.0 g (108.4 mmoles) of anhydrous potassium carbonate and 19.3 g (135.5 mmoles) of methyl iodide were added, at room temperature, to a solution of 18.8 g (90.3 mmoles) of 1-tert-butyl-3-trifluoromethyl-1H-pyrazol-5-ol dissolved in 100 ml of N,N-dimethylformamide. The mixture was stirred for 15 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 20.0 g (yield: 99.8%) of 1-tert-butyl-5-methoxy-3-trifluoromethyl-1H-pyrazole.

<Reference Example 22>

Production of 1-tert-butyl-4-chloromethyl-5-methoxy-3-trifluoromethyl-1H-pyrazole

[0283] 5.4 g of paraformaldehyde (180.2 mmoles in terms of formaldehyde) and 20 ml of concentrated hydrochloric acid were added to a solution of 20.0 g (90.1 mmoles) of 1-tert-butyl-5-methoxy-3-trifluoromethyl-1H-pyrazole dissolved in 90 ml of acetic acid. The mixture was stirred at 60°C for 30 minutes to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diisopropyl ether. The resulting organic layer was washed with water and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 21.7 g (yield: 89.0%) of 1-tert-butyl-4-chloromethyl-5-methoxy-3-trifluoromethyl-1H-pyrazole.

<Reference Example 23>

Production of 3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazole

[0284] 10.0 g (72.3 mmoles) of anhydrous potassium carbonate and 12.8 g (90.3 mmoles) of methyl iodide were added, at room temperature, to a solution of 10.0 g (60.2 mmoles) of 3-hydroxy-1-methyl-5-trifluoromethyl-1H-pyrazole dissolved in 50 ml of N,N-dimethylformamide. The mixture was stirred for 15 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 9.8 g (yield: 90.7%) of 3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazole.

<Reference Example 24>

Production of 4-chloromethyl-3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazole

[0285] 0.45 g of paraformaldehyde (15.0 mmoles in terms of formaldehyde) and 5 ml of concentrated hydrochloric acid were added to a solution of 1.00 g (5.6 mmoles) of 3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazole dissolved in 25 ml of acetic acid. The mixture was stirred at 80°C for 2 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water and neutralized with potassium carbonate, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.83 g (yield: 65.0%) of 4-chloromethyl-3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazole.

<Reference Example 25>

Production of 5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde

[0286] 42.0 g (711.9 mmoles) of potassium fluoride was added to a solution of 60.4 g (282.7 mmoles) of 5-chloro-1-methyl-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde dissolved in 700 ml of dimethyl sulfoxide. The mixture was stirred at 120 to 140°C for 5 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The

resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 36.8 g (yield: 66.0%) of 5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde.

<Reference Example 26>

Production of (5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol

[0287] To a solution of 3.9 g (102.6 mmoles) of sodium borohydride dissolved in 500 ml of methanol was added, with ice-cooling, a solution of 36.8 g (187.6 mmoles) of 5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde dissolved in 200 ml of methanol. The resulting mixture was stirred at 0°C for 30 minutes to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 35.4 g (yield: 95.4%) of (5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol.

<Reference Example 27>

Production of 4-bromomethyl-5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazole

[0288] A solution of 35.4 g (178.7 mmoles) of 5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazole-4-methanol dissolved in 500 ml of diethyl ether was cooled to -30°C. Thereto was added 54.0 g (199.5 mmoles) of phosphorus tribromide. The mixture was stirred at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 31.4 g (yield: 80.8%) of 4-bromomethyl-5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazole.

<Reference Example 28>

Production of (ethoxycarbonyl)malondialdehyde

[0289] 12.6 g of sodium hydride (purity: 60%, 525.0 mmoles) was washed with diethyl ether by decantation several times and then made into a solution in 500 ml of diethyl ether. Thereto were added, in a nitrogen current at 0 to 10°C, 194 g (2.6 moles) of ethyl formate and 50 g (262.0 mmoles) of ethyl 3,3-diethoxy-propionate. The resulting mixture was stirred at room temperature for 15 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by washing with diethyl ether. The resulting aqueous layer was allowed to have a pH of 1 with hydrochloric acid, followed by extraction with dichloromethane. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 37.6 g (yield: 100%) of crude (ethoxycarbonyl)malondialdehyde as a dark red oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

9.09 (2H,s), 5.26 (1H,s), 4.27 (2H,q), 1.28 (3H,t)

<Reference Example 29>

Production of ethyl 1H-pyrazole-4-carboxylate

[0290] 6.2 g (193 mmoles) of hydrazine was added, with ice-cooling, to a solution of 27.6 g (192 mmoles) of (ethoxycarbonyl)malondialdehyde dissolved in 150 ml of ethanol. The mixture was stirred at room temperature for 17 hours to give rise to a reaction. The reaction mixture was subjected to vacuum distillation to remove the ethanol contained therein. The residue was purified by silica gel column chromatography (developing solvent: dichloromethane-ethyl acetate mixed solvent) to obtain 19.4 g (72.4%) of ethyl 1H-pyrazole-4-carboxylate as yellow crystals.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.08 (2H,s), 5.30 (1H,s) 4.31 (2H,q), 1.36 (3H,t)

<Reference Example 30>

Production of ethyl 1-ethyl-1H-pyrazole-4-carboxylate

[0291] 3.7 g (26.8 mmoles) of anhydrous potassium carbonate and 4.2 g (26.6 mmoles) of ethyl iodide were added to a solution of 1.5 g (10.7 mmoles) of ethyl 1H-pyrazole-4-carboxylate dissolved in 50 ml of N,N-dimethylformamide. The mixture was stirred at room temperature for 20 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 1.6 g (yield: 88.9%) of ethyl 1-ethyl-1H-pyrazole-4-carboxylate as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.90 (2H,s), 4.28 (2H,q), 4.18 (2H,q), 1.51 (3H,t), 1.35 (3H,t)

<Reference Example 31>

Production of ethyl 3,5-dichloro-1-ethyl-1H-pyrazole-4-carboxylate

[0292] In a glass sealed tube were placed 1.6 g (9.5 mmoles) of ethyl 1-ethyl-1H-pyrazole-4-carboxylate and 5.1 g (38.3 mmoles) of N-chlorosuccinimide. There were allowed to react at 160°C for 6 hours. After the completion of the reaction, the reaction mixture was cooled to room temperature, washed with carbon tetrachloride and chloroform, and filtered under vacuum. The resulting filtrate (organic layer) was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 1.0 g (yield: 44.2%) of ethyl 3,5-dichloro-1-ethyl-1H-pyrazole-4-carboxylate as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.36 (2H,q), 4.21 (2H,q), 1.44 (3H,t), 1.38 (3H,t)

<Reference Example 32>

Production of (3,5-dichloro-1-ethyl-1H-pyrazol-4-yl)methanol

[0293] A solution of 0.16 g (4.2 mmoles) of lithium aluminum hydride dissolved in 70 ml of tetrahydrofuran was cooled to -50°C. Thereto was gradually added dropwise a solution of 1.0 g (4.2 mmoles) of ethyl 3,5-dichloro-1-ethyl-1H-pyrazole-4-carboxylate dissolved in 30 ml of tetrahydrofuran. The mixture was stirred at -50°C for 3 hours to give rise to a reaction. After confirmation of the completion of the reaction, ethyl acetate was added, followed by stirring for a while. Water was added, followed by stirring for a while. The resulting mixture was filtered under vacuum. The filtrate was extracted with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.82 g (yield: 100%) of (3,5-dichloro-1-ethyl-1H-pyrazol-4-yl)methanol as a brown oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.52 (2H,s), 4.16 (2H,q), 1.43 (3H,t)

<Reference Example 33>

Production of 4-bromomethyl-3,5-dichloro-1-ethyl-1H-pyrazole

[0294] A solution of 0.82 g (4.2 mmoles) of (3,5-dichloro-1-ethyl-1H-pyrazol-4-yl)methanol dissolved in 50 ml of diethyl ether was cooled to -30°C. Thereto was added 1.3 g (4.8 mmoles) of phosphorus tribromide. The mixture was stirred at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.9 g (yield: 81.8%) of 4-bromomethyl-3,5-dichloro-1-ethyl-1H-pyrazole as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.33 (2H,s), 4.13 (2H,q), 1.43 (3H,t)

<Reference Example 34>

5 Production of 3-difluoromethyl-1-methyl-1H-pyrazol-5-ol

[0295] 8.3 g (180.6 mmoles) of methylhydrazine and 5 ml of concentrated hydrochloric acid were added to a solution of 30.0 g (180.6 mmoles) of ethyl difluoroacetoacetate dissolved in 200 ml of ethanol. The mixture was refluxed for 2 days with heating, to give rise to a reaction. After the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the most part of the solvent contained therein. The residue was poured into water. The mixture was allowed to have a pH of 4 using citric acid and extracted with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 8.9 g (yield: 33.3%) of 3-difluoromethyl-1-methyl-1H-pyrazol-5-ol.

<Reference Example 35>

20 Production of 5-chloro-3-difluoromethyl-1-methyl-1H-pyrazole-4-carbaldehyde

[0296] 41.6 g (270.1 mmoles) of phosphorus oxychloride was added, with ice-cooling, to 7.9 g (108.0 mmoles) of N,N-dimethylformamide. Thereto was added, at room temperature, 8.0 g (54.0 mmoles) of 3-difluoromethyl-1-methyl-1H-pyrazol-5-ol. The mixture was refluxed for 4 hours with heating, to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with water, a 5% aqueous sodium hydroxide solution and water in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 7.7 g (yield: 73.3%) of 5-chloro-3-difluoromethyl-1-methyl-1H-pyrazole-4-carbaldehyde as white crystals.

30 ¹H-NMR [CDCl₃/TMS, δ (ppm)]:
9.96 (1H,s), 6.90 (1H,t, J=53.6 Hz), 3.93 (3H,s)

<Reference Example 36>

35 Production of (5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-yl)-methanol

[0297] A solution of 7.2 g (37.0 mmoles) of 5-chloro-3-difluoromethyl-1-methyl-1H-pyrazole-4-carbaldehyde dissolved in 100 ml of methanol was cooled to 0°C. Thereto was gradually added 2.1 g (55.5 mmoles) of sodium borohydride. The mixture was stirred at room temperature for 3 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 3.8 g (yield: 52.1%) of (5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-yl)-methanol.

45 ¹H-NMR [CDCl₃/TMS, δ (ppm)]:
6.70 (1H,t, J=40.8 Hz), 4.63 (2H,s), 3.86 (3H,s), 1.79 (1H,br)

<Reference Example 37>

50 Production of 4-bromomethyl-5-chloro-3-difluoromethyl-1-methyl-1H-pyrazole

[0298] A solution of 2.0 g (10.0 mmoles) of (5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-yl)-methanol dissolved in 50 ml of diethyl ether was cooled to -10°C. Thereto was added 1.0 g (3.5 mmoles) of phosphorus tribromide. The mixture was stirred at room temperature overnight to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into ice water, followed by extraction with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 2.6 g (yield: 100.0%) of 4-bromomethyl-5-chloro-3-difluoromethyl-1-methyl-1H-pyrazole.

<Reference Example 38>

Production of trifluoroacetaldehyde oxime etherate

- 5 **[0299]** 24.1 g (347.0 mmoles) of hydroxylamine hydrochloride and 160 ml of water were added to a solution of 50.0 g (347.0 mmoles) of trifluoroacetaldehyde hemiethyl acetal dissolved in 80 ml of methanol. Thereto was dropwise added, with ice-cooling, 80.0 g of a 50% aqueous sodium hydroxide solution (1.7 moles). After the completion of the dropwise addition, the resulting mixture was stirred at room temperature for 6 hours to give rise to a reaction. After the completion of the reaction, 10% hydrochloric acid was added for pH adjustment to 6. The resulting mixture was extracted with diethyl ether. The extract was subjected to vacuum distillation to remove the solvent contained therein. The residue was subjected to distillation to obtain 24.7 g (yield: 38.0%) of trifluoroacetaldehyde oxime etherate.

<Reference Example 39>

- 15 Production of trifluoroacetohydroximoyl bromide etherate

- [0300]** A solution of 38.8 g (218.0 mmoles) of N-bromosuccinimide dissolved in 125 ml of N,N-dimethylformamide was added, with ice-cooling, to a solution of 24.7 g (131.7 mmoles) of trifluoroacetaldehyde oxime etherate dissolved in 50 ml of N,N-dimethylformamide. The mixture was stirred at room temperature for 3 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was subjected to distillation to obtain 33.3 g (yield: 95.0%) of trifluoroacetohydroximoyl bromide etherate as a brown oily substance.
- 25 ¹H-NMR [CDCl₃/TMS, δ (ppm)]:
9.30 (1H,s)

<Reference Example 40>

- 30 Production of 4-ethoxycarbonyl-5-methyl-3-trifluoromethylisoxazole

- [0301]** 2.8 g (51.3 mmoles) of sodium methoxide was added to a solution of 6.7 g (51.3 mmoles) of ethyl acetoacetate dissolved in 80 ml of methanol. Thereto was added, with ice-cooling, a solution of 5.0 g (18.8 mmoles) of trifluorohydroximoyl bromide etherate dissolved in 20 ml of methanol. The resulting mixture was stirred at room temperature for 3 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the solvent contained therein. Water was added to the residue, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 2.9 g (yield: 69.0%) of 4-ethoxycarbonyl-5-methyl-3-trifluoromethylisoxazole as a colorless oily substance.
- 35 ¹H-NMR [CDCl₃/TMS, δ (ppm)]:
4.36 (2H,q), 2.77 (3H,s), 1.37 (3H,t)

- 45 <Reference Example 41>

Production of (5-methyl-3-trifluoromethylisoxazol-4-yl)-methanol

- [0302]** A solution of 0.16 g (4.2 mmoles) of lithium aluminum hydride dissolved in 15 ml of THF was cooled to 0°C. Thereto was gradually added a solution of 0.93 g (4.2 mmoles) of 4-ethoxycarbonyl-5-methyl-3-trifluoromethylisoxazole dissolved in 15 ml of THF. The mixture was stirred at 0°C for 1 hour to give rise to a reaction. After the completion of the reaction, ethyl acetate was added, followed by stirring for a while. Water was added, followed by stirring for a while. The reaction mixture was filtered under vacuum. The filtrate was extracted with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.5 g (yield: 60.0%) of (5-methyl-3-trifluoromethylisoxazol-4-yl)-methanol.
- 55 ¹H-NMR [CDCl₃/TMS, δ (ppm)]:
4.60 (2H,d), 2.54 (3H,s), 1.66 (1H, br)

<Reference Example 42>

Production of 4-bromomethyl-5-methyl-3-trifluoromethylisoxazole

[0303] A solution of 0.45 g (2.5 mmoles) of (5-methyl-3-trifluoromethylisoxazol-4-yl)-methanol dissolved in 10 ml of diethyl ether was cooled to 0°C. Thereto was added 0.2 g (8.9 mmoles) of phosphorus tribromide. The mixture was stirred at room temperature for 1 hour to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.5 g (yield: 74.0%) of 4-bromomethyl-5-methyl-3-trifluoromethylisoxazole.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.31 (2H,d), 2.51 (3H,s)

<Reference Example 43>

Production of (5-chloro-3-methyl-isothiazol-4-yl)-methanol

[0304] A solution of 2.06 g (10.0 mmoles) of ethyl 5-chloro-3-methyl-isothiazole-4-carboxylate dissolved in 10 ml of THF was dropwise added, at -30°C, to a solution of 0.42 g (11.0 mmoles) of lithium aluminum hydride dissolved in 10 ml of THF. The mixture was stirred at the same temperature for 1 hour to give rise to a reaction. After confirmation of the completion of the reaction, ethyl acetate was added to the reaction mixture. The resulting mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 1.50 g (yield: 91.5%) of (5-chloro-3-methylisothiazol-4-yl)-methanol.

<Reference Example 44>

Production of 4-chloromethyl-5-chloro-3-methylisothiazole

[0305] 3.26 g (27.44 mmoles) of thionyl chloride was added, at room temperature, to a solution of 1.50 g (9.15 mmoles) of (5-chloro-3-methyl-isothiazol-4-yl)-methanol dissolved in 10 ml of chloroform. The mixture was stirred for 3 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the solvent contained therein, to obtain 1.67 g (yield: quantitative) of 4-chloromethyl-5-chloro-3-methylisothiazole.

<Reference Example 45>

Production of methyl 4-trifluoromethylnicotinate

[0306] 6.7 g (48.6 mmoles) of anhydrous potassium carbonate and 6.9 g (48.6 mmoles) of methyl iodide were added to a solution of 4.6 g (24.1 mmoles) of 4-trifluoromethylnicotinic acid dissolved in 70 ml of N,N-dimethylformamide. The mixture was stirred at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 2.77 g (yield: 56.1%) of methyl 4-trifluoromethylnicotinate as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

9.11 (1H,s) 8.92 (1H,d), 7.64 (1H,d), 3.99 (3H,s)

<Reference Example 46>

Production of (4-trifluoromethylpyridin-3-yl)-methanol

[0307] A solution of 0.37 g (9.7 mmoles) of lithium aluminum hydride dissolved in 100 ml of THF was cooled to -50°C. Thereto was gradually added dropwise a solution of 2.0 g (9.8 mmoles) of methyl 4-trifluoromethylnicotinate dissolved

in 30 ml of THF. The mixture was stirred at -50°C for 3 hours to give rise to a reaction. After confirmation of the completion of the reaction, ethyl acetate was added, followed by stirring for a while. Water was added, followed by stirring for a while. The reaction mixture was filtered under vacuum. The filtrate was extracted with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.6 g (yield: 35.3%) of (4-trifluoromethylpyridin-3-yl)-methanol as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

9.00 (1H,s), 8.73 (1H,d), 7.51 (1H,d), 4.95 (2H,s)

<Reference Example 47>

Production of 3-bromomethyl-4-trifluoromethylpyridine

[0308] A solution of 0.6 g (3.4 mmoles) of (4-trifluoromethylpyridin-3-yl)-methanol dissolved in 50 ml of diethyl ether was cooled to -30°C. Thereto was added 1.4 g (5.2 mmoles) of phosphorus tribromide. The mixture was stirred at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.61 g (yield: 75.3%) of 3-bromomethyl-4-trifluoromethylpyridine as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.88 (1H,s), 8.73 (1H,d), 7.54 (1H,d), 4.63 (2H,s)

<Reference Example 48>

Production of 5-bromo-4-hydroxy-6-trifluoromethylpyrimidine

[0309] 77.5 g (945.0 mmoles) of anhydrous sodium acetate was added, at room temperature, to a solution of 49.2 g (300.0 mmoles) of 4-hydroxy-6-trifluoromethylpyrimidine dissolved in 600 ml of acetic acid. Thereto was gradually added 50.3 g (315 mmoles) of bromine at 45°C. The resulting mixture was stirred at the same temperature for 3 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the solvent contained therein. The residue was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was washed with n-hexane to obtain 38.9 g (yield: 53.4%) of 5-bromo-4-hydroxy-6-trifluoromethylpyrimidine.

<Reference Example 49>

Production of 5-bromo-4-chloro-6-trifluoromethylpyrimidine

[0310] 24.3 g (100.0 mmoles) of 5-bromo-4-hydroxy-6-trifluoromethylpyrimidine was suspended in 18.5 g (120.0 mmoles) of phosphorus oxychloride. The mixture was stirred at 100°C for 2 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water gradually, followed by extraction with chloroform. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 21.5 g (yield: 82.4%) of 5-bromo-4-chloro-6-trifluoromethylpyrimidine.

<Reference Example 50>

Production of 5-bromo-4-methoxy-6-trifluoromethylpyrimidine

[0311] 16.7 ml of sodium methoxide (a 28% methanol solution, 86.4 mmoles) was added, at room temperature, to a solution of 21.5 g (82.2 mmoles) of 5-bromo-4-chloro-6-trifluoromethylpyrimidine dissolved in 100 ml of methanol. The mixture was stirred to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the solvent contained therein. The residue was poured into

water, followed by extraction with chloroform. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was washed with n-hexane to obtain 19.2 g (yield: 91.0%) of 5-bromo-4-methoxy-6-trifluoromethylpyrimidine.

<Reference Example 51>

Production of 5-bromo-4-ethoxy-6-trifluoromethylpyrimidine

[0312] 0.94 g (13.77 mmoles) of sodium ethoxide was added, at room temperature, to a solution of 3.00 g (11.48 mmoles) of 5-bromo-4-chloro-6-trifluoromethylpyrimidine dissolved in 50 ml of ethanol. The mixture was stirred to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the solvent contained therein. The residue was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 2.44 g (yield: 82.9%) of 5-bromo-4-ethoxy-6-trifluoromethylpyrimidine.

<Reference Example 52>

Production of 4-methoxy-6-trifluoromethylpyrimidine-5-carbaldehyde

[0313] 30.0 ml of n-butyllithium (a 1.6 moles/liter n-hexane solution, 48.0 mmoles) was gradually added, at -65 to -60°C, to a solution of 10.3 g (40.0 mmoles) of 5-bromo-4-methoxy-6-trifluoromethylpyrimidine dissolved in 100 ml of tetrahydrofuran. The mixture was stirred for 30 minutes. Thereto was added 3.6 g (48.0 mmoles) of ethyl formate at the same temperature. The resulting mixture was stirred at the same temperature for 3 hours to give rise to a reaction. The reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 1.3 g (yield: 15.8%) of 4-methoxy-6-trifluoromethylpyrimidine-5-carbaldehyde.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

10.41 (1H,q), 8.91 (1H,s), 4.18 (3H,s)

<Reference Example 53>

Production of 4-ethoxy-6-trifluoromethylpyrimidine-5-carbaldehyde

[0314] A solution of 5.76 g (21.3 mmoles) of 5-bromo-4-ethoxy-6-trifluoromethylpyrimidine dissolved in 250 ml of THF was cooled to -78°C. Thereto was dropwise added 22.6 ml of n-butyllithium (a 1.6 moles/liter n-hexane solution, 36.1 mmoles). The mixture was stirred for 40 minutes. Thereto was added 2.7 g (45.1 mmoles) of methyl formate. The resulting mixture was stirred for 1.5 hours to give rise to a reaction. After the completion of the reaction, an aqueous ammonium chloride solution was added. The mixture was extracted with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 3.82 g (yield: 81.6%) of 4-ethoxy-6-trifluoromethylpyrimidine-5-carbaldehyde.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

10.41 (1H,s), 8.95 (1H,s), 4.63 (2H,q), 1.48 (3H,t)

<Reference Example 54>

Production of (4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methanol

[0315] 0.24 g (6.3 mmoles) of sodium borohydride was gradually added, at room temperature, to a solution of 1.3 g (6.3 mmoles) of 4-methoxy-6-trifluoromethylpyrimidine-5-carbaldehyde dissolved in 30 ml of methanol. The mixture was stirred for 3 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water

and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 0.42 g (yield: 32.1%) of (4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methanol

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.93 (1H,s), 4.81 (2H,s), 4.13 (3H,s), 2.26 (1H,br)

<Reference Example 55>

Production of (4-ethoxy-6-trifluoromethylpyrimidin-5-yl)-methanol

[0316] A solution of 3.82 g (17.2 mmoles) of 4-ethoxy-6-trifluoromethylpyrimidine-5-carbaldehyde dissolved in 50 ml of methanol was added, with ice-cooling, to a solution of 1.7 g (45.7 mmoles) of sodium borohydride dissolved in 50 ml of methanol. The mixture was stirred at 0°C for 1 hour to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 3.77 g (yield: 97.8%) of (4-ethoxy-6-trifluoromethylpyrimidin-5-yl)-methanol

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.80 (1H,s), 4.81 (2H,s), 4.59 (2H,q), 2.28 (1H,b), 1.48 (3H,t)

<Reference Example 56>

Production of 5-chloromethyl-4-methoxy-6-trifluoromethylpyrimidine

[0317] 1.19 g (10.1 mmoles) of thionyl chloride was added, at room temperature, to a solution of 0.42 g (2.02 mmoles) of (4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methanol. The mixture was stirred for 3 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.45 g (yield: quantitative) of 5-chloromethyl-4-methoxy-6-trifluoromethylpyrimidine.

<Reference Example 57>

Production of 5-bromomethyl-4-ethoxy-6-trifluoromethylpyrimidine

[0318] A solution of 3.77 g (17.0 mmoles) of (4-ethoxy-6-trifluoromethylpyrimidin-5-yl)-methanol dissolved in 50 ml of diethyl ether was cooled to 0°C. Thereto was added 2.0 g (7.2 mmoles) of phosphorus tribromide. The mixture was stirred at room temperature for 1 hour. The resulting salt was dissolved using methanol. The resulting mixture was stirred for 1 hour to give rise to a reaction. The reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain crude 5-bromomethyl-4-ethoxy-6-trifluoromethylpyrimidine.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.79 (1H,s), 4.61 (2H,q), 4.55 (2H,s), 1.49 (3H,t)

<Reference Example 58>

Production of (2-chloro-4-methylpyridin-3-yl)methanol

[0319] A solution of 1.9 g (10.0 mmoles) of methyl 2-chloro-4-methylnicotinate dissolved in 5.0 ml of THF was gradually added, at -65 to -60°C, to a suspension of 0.4 g (10.0 mmoles) of lithium aluminum hydride in 30 ml of tetrahydrofuran. The mixture was stirred for 30 minutes and at -20°C for 1 hour to give rise to a reaction. The reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 0.6 g (yield: 38.2%) of (2-chloro-4-methylpyridin-3-yl)methanol.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.19 (1H,d), 7.08 (1H,d), 4.85 (2H,s), 2.49 (3H,s)

<Reference Example 59>

Production of 3-acetyl-4-chloromethyl-2,5-dichlorothiophene

[0320] 33 ml of titanium tetrachloride (a 2 moles/liter dichloromethane solution, 66.0 mmoles) was dropwise added, at 10°C with ice-cooling, to a solution of 5.0 g (32.4 mmoles) of 3-acetyl-2,5-dichlorothiophene dissolved in 26 ml (323.0 mmoles) of chloromethyl methyl ether. The mixture was stirred at room temperature for 2 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into ice water, followed by extraction with chloroform. The resulting organic layer was washed with sodium bicarbonate, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane/ethyl acetate = 9/1) to obtain 2.6 g (yield: 39.7%) of 3-acetyl-4-chloromethyl-2,5-dichlorothiophene as yellow crystals.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.70 (2H,s), 2.56 (3H,s), 2.54 (3H,s), 2.39 (3H,s)

<Reference Example 60>

Production of 3-bromo-2-bromomethylbenzofuran

[0321] 2.7 g (15.3 mmoles) of N-bromosuccinimide and 0.4 g (2.7 mmoles) of azobisisobutyronitrile were added to a solution of 2.8 g (13.3 mmoles) of 3-bromo-2-methylbenzofuran dissolved in 30 ml of monochlorobenzene. The mixture was stirred at 80°C for 30 minutes to give rise to a reaction. After confirmation of the disappearance of the raw materials, the reaction mixture was cooled to room temperature. The insolubles were removed by filtration. The filtrate was subjected to vacuum distillation to remove the solvent contained therein. The residue was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 3.0 g (yield: 79.0%) of 3-bromo-2-bromomethylbenzofuran.

<Reference Example 61>

Production of ethyl 1-difluoromethyl-1H-pyrazole-4-carboxylate

[0322] 6.0 g (43.5 mmoles) of anhydrous potassium carbonate was added to a solution of 3.0 g (21.4 mmoles) of ethyl 1H-pyrazole-4-carboxylate dissolved in 100 ml of N,N-dimethylformamide. Thereinto was blown chlorodifluoromethane. The resulting mixture was stirred at 130 to 140°C for 3 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 1.67 g (yield: 41.0%) of ethyl 1-difluoromethyl-1H-pyrazole-4-carboxylate as a colorless transparent oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.32 (1H,s), 8.04 (1H,s), 7.20 (1H,t), 4.32 (2H,q), 1.37 (3H,t)

[0323] The herbicide of the present invention contains, as the active ingredient, an isoxazoline derivative represented by the general formula [I] or a salt thereof.

[0324] In using the compound of the present invention as a herbicide, the present compound may be used by itself. It can also be used in the form of a powder, a wettable powder, an emulsifiable concentrate, a flowable, fine granules, granules, etc. by mixing with a carrier, a surfactant, a dispersant, an adjuvant, etc. all generally used in formulation.

[0325] As the carrier used in formulation, there can be mentioned, for example, solid carriers such as talc, bentonite, clay, kaolin, diatomaceous earth, white carbon, vermiculite, calcium carbonate, slaked lime, siliceous sand, ammonium sulfate, urea and the like; and liquid carriers such as isopropyl alcohol, xylene, cyclohexane, methylnaphthalene and the like.

[0326] As the surfactant and the dispersant, there can be mentioned, for example, metal salts of alkylbenzenesulfonic acids, metal salts of dinaphthylmethanedisulfonic acid, salts of alcohol sulfates, alkylarylsulfonic acid salts, ligninsulfonic acid salts, polyoxyethylene glycol ether, polyoxyethylene alkyl aryl ethers, monoalkylates of polyoxyethylene sorbitan and the like. As the adjuvant, there can be mentioned, for example, carboxymethyl cellulose, polyethylene

glycol and gum arabic. The present herbicide, when used, is diluted to an appropriate concentration and sprayed or applied directly.

[0327] The herbicide of the present invention can be used by spraying on foliage, application to soil, application on water surface, etc. The amount of the active ingredient used is determined appropriately so as to meet the application purpose. The content of the active ingredient is appropriately determined according to the purpose. When the present compound is formulated as a powder or granules, the content is in a range of 0.01% to 10% by weight, preferably 0.05% to 5% by weight. When the present compound is made into an emulsifiable concentrate or a wettable powder, the amount is appropriately determined in a range of 1 to 50% by weight, 5 to 30% by weight. When the present compound is made into a flowable, the amount is appropriately determined in a range of 1 to 40% by weight, preferably 5 to 30% by weight.

[0328] The amount of the present herbicide used varies depending upon the kind of the compound used, the target weed, the tendency of weed emergence, the environmental conditions, the type of the herbicide used, etc. When the present herbicide is used per se as in the case of a powder or granules, the amount is appropriately selected in a range of 1 g to 50 kg, preferably 10 g to 10 kg per 1 hectare in terms of the active ingredient. When the present herbicide is used in a liquid form as in the case of an emulsifiable concentrate, a wettable powder or a flowable, the amount is appropriately selected in a range of 0.1 to 50,000 ppm, preferably 10 to 10,000 ppm.

[0329] The compound of the present invention may be mixed as necessary with an insecticide, a fungicide, other herbicide, a plant growth-regulating agent, a fertilizer, etc.

[0330] Next, formulation from the present compound is described specifically by showing typical examples of formulation. The kinds of compounds and additives and their compounding ratios are not restricted to those shown below and can be varied widely. In the following description, "parts" refer to parts by weight.

<Formulation 1> Wettable powder

[0331] 10 parts of a compound (3-0006) were mixed with 0.5 part of polyoxyethylene octylphenyl ether, 0.5 part of a sodium salt of a β -naphthalenesulfonic acid-formalin condensate, 20 parts of diatomaceous earth and 69 parts of clay. The mixture was mixed and pulverized to obtain a wettable powder.

<Formulation 2> Flowable

[0332] 20 parts of a coarsely ground compound (3-0006) were dispersed in 69 parts of water. Thereto were added 4 parts of a polyoxyethylene styryl phenylether sulfate, 7 parts of ethylene glycol and 200 ppm, relative to the herbicide produced, of Silicone AF-118N (a product of Asahi Chemical Industry, Co. Ltd.). The resulting mixture was stirred for 30 minutes using a high-speed stirrer and then ground using a wet grinder to obtain a flowable.

<Formulation 3> Emulsion

[0333] To 30 parts of a compound (3-0006) were added 60 parts of an equal volume mixture of xylene and isophorone and 10 parts of a surfactant mixture of a polyoxyethylene sorbitan alkylate, a polyoxyethylene alkylaryl polymer and an alkylaryl sulfonate. The resulting mixture was stirred sufficiently to obtain an emulsifiable concentrate.

<Formulation 4> Granules

[0334] There were mixed 10 parts of a compound (3-0006), 80 parts of an extender which was a 1:3 mixture of talc and bentonite, 5 parts of white carbon and 5 parts of a surfactant mixture of a polyoxyethylene sorbitan alkylate, a polyoxyethylene alkylaryl polymer and an alkylaryl sulfonate. To the mixture were added 10 parts of water. The resulting mixture was kneaded sufficiently to form a paste. The paste was extruded through the eyes (diameter: 0.7 mm) of a sieve. The extrudate was dried and cut into a length of 0.5 to 1 mm to obtain granules.

[0335] Next, Application Examples of the present compound are described to show the effect of the present compound.

<Application Example 1> Test for herbicidal effect by paddy field soil treatment

[0336] A paddy field soil was filled in a plastic pot of 100 cm² and subjected to puddling. Then, seeds of *Echinochloa oryzicola* Vasing. and *Monochoria vaginalis* (Murm. f.) Presl var. *plantaginea* (Roxb.) Solms-Laub. were sowed and water was filled in a depth of 3 cm. Next day, wettable powders produced in accordance with the Formulation 1 were diluted with water and dropped on the water surface. The application amount of each wettable powder was 1,000 g per 1 hectare in terms of the active ingredient. Then, breeding was made in a greenhouse, and the herbicidal effect of

EP 1 364 946 A1

each wettable powder was examined at the 21st day from the treatment in accordance with the standard shown in Table 21. The results are shown in Table 22.

Table 21

Index	Herbicidal effect (extent of growth inhibition) or phytotoxicity
5	A herbicidal effect or phytotoxicity of 90%
4	A herbicidal effect or phytotoxicity of 70% to less than 90%
3	A herbicidal effect or phytotoxicity of 50% to less than 70%
2	A herbicidal effect or phytotoxicity of 30% to less than 50%
1	A herbicidal effect or phytotoxicity of 10% to less than 30%
0	A herbicidal effect or phytotoxicity of 0% to less than 10%

Table 22

Compound No.	Active ingredient (g/ha)	Echinochloa oryzicola Vasing	Monochoria vaginalis (Burm.F.) Presl var. plantaginea (Roxb.) Solms-Laub.
1-0001	1000	5	5
1-0002	1000	5	5
1-0003	1000	5	5
1-0004	1000	5	5
1-0005	1000	5	5
2-0001	1000	5	5
2-0003	1000	5	5
2-0004	1000	5	5
2-0005	1000	5	5
2-0006	1000	5	5
2-0008	1000	5	5
2-0011	1000	5	5
2-0012	1000	5	5
3-0002	1000	5	5
3-0004	1000	5	5
3-0009	1000	5	5
3-0013	1000	5	5
3-0014	1000	5	5
3-0015	1000	5	5
3-0016	1000	5	5
3-0034	1000	5	5
3-0035	1000	5	5
3-0037	1000	5	5
3-0038	1000	5	5
3-0039	1000	5	5
3-0040	1000	5	5
3-0041	1000	5	5
3-0044	1000	5	5
3-0047	1000	5	5
3-0049	1000	5	5
3-0051	1000	5	5
3-0054	1000	5	5
3-0059	1000	5	5

EP 1 364 946 A1

Table 22 (continued)

	Compound No.	Active ingredient (g/ha)	Echinochloa oryzicola Vasing	Monochoria vaginalis (Burm.F.) Presl var. plantaginea (Roxb.) Solms-Laub.
5	3-0060	1000	5	5
	3-0061	1000	5	5
	3-0070	1000	5	5
	3-0072	1000	5	5
10	3-0073	1000	5	5
	3-0074	1000	5	5
	3-0081	1000	5	5
	3-0082	1000	5	5
15	3-0083	1000	5	5
	3-0084	1000	5	5
	3-0085	1000	5	5
	3-0086	1000	5	5
	3-0087	1000	5	5
20	3-0088	1000	5	5
	3-0089	1000	5	5
	3-0090	1000	5	5
	3-0091	1000	5	5
25	3-0100	1000	5	5
	3-0101	1000	5	5
	3-0102	1000	5	5
	3-0103	1000	5	5
	3-0114	1000	5	5
30	3-0115	1000	5	5
	3-0117	1000	5	5
	3-0118	1000	5	5
	3-0119	1000	5	5
35	3-0120	1000	5	5
	3-0121	1000	5	5
	3-0124	1000	5	5
	3-0125	1000	5	5
	3-0126	1000	5	5
40	3-0127	1000	5	5
	3-0128	1000	5	5
	3-0129	1000	5	5
	3-0130	1000	5	5
	3-0131	1000	5	5
45	3-0134	1000	5	5
	3-0135	1000	5	5
	3-0137	1000	5	5
	3-0139	1000	5	5
50	3-0144	1000	5	5
	3-0153	1000	5	5
	3-0156	1000	5	5
	3-0160	1000	5	5
	3-0173	1000	5	5
55	3-0174	1000	5	5
	3-0176	1000	5	5
	3-0177	1000	5	5

EP 1 364 946 A1

Table 22 (continued)

	Compound No.	Active ingredient (g/ha)	Echinochloa oryzicola Vasing	Monochoria vaginalis (Burm.F.) Presl var. plantaginea (Roxb.) Solms-Laub.
5	3-0178	1000	5	5
	3-0180	1000	5	5
	4-0001	1000	5	5
	4-0002	1000	5	5
10	4-0005	1000	5	5
	4-0007	1000	5	5
	4-0008	1000	5	5
	5-0001	1000	5	5
15	5-0002	1000	5	5
	5-0003	1000	5	5
	5-0005	1000	5	5
	5-0006	1000	5	5
	5-0007	1000	5	5
20	6-0003	1000	5	5
	6-0004	1000	5	5
	7-0004	1000	5	5
	7-0006	1000	5	5
25	7-0008	1000	5	5
	7-0009	1000	5	5
	8-0001	1000	5	5
	8-0012	1000	5	5
30	9-0001	1000	5	5
	9-0003	1000	5	5
	9-0005	1000	5	5
	9-0006	1000	5	5
	9-0008	1000	5	5
35	10-0002	1000	5	4
	10-0003	1000	5	5
	10-0004	1000	5	5
	10-0005	1000	5	5
	10-0006	1000	5	5
40	10-0008	1000	5	5
	10-0009	1000	5	5
	10-0011	1000	5	5
	10-0012	1000	5	5
45	10-0013	1000	5	5
	10-0014	1000	5	5
	10-0015	1000	5	5
	10-0016	1000	5	5
	10-0017	1000	5	5
50	10-0018	1000	5	5

<Application Example 2> Test for herbicidal effect by upland field soil treatment

55 [0337] An upland field soil was filled in a plastic pot of 80 cm². Seeds of Echinochloa crus-galli (L.) Beauv. var. crus-galli and Setaria viridis (L.) Beauv. were sowed, followed by covering with the same soil. Wettable powders produced in accordance with the Formulation 1 were diluted with water and sprayed uniformly on the soil surface using a small sprayer, in an amount of 1,000 liters per 1 hectare so that the amount of each active ingredient became 1,000 g per 1

EP 1 364 946 A1

hectare. Then, breeding was made in a greenhouse, and the herbicidal effect of each wettable powder was examined at the 21st day from the treatment in accordance with the standard shown in Table 21. The results are shown in Table 23.

Table 23

	Compound No.	Active ingredient (g/ha)	Echinochloa crus-galli (L.) Beauv. Var. crus-galli	Setaria viridis (L.) Beauv.
5	1-0001	1000	5	5
	1-0002	1000	5	5
10	1-0003	1000	5	5
	1-0004	1000	5	5
	1-0005	1000	5	5
	1-0006	1000	5	4
15	2-0001	1000	5	5
	2-0003	1000	5	5
	2-0004	1000	5	5
	2-0005	1000	5	5
	2-0006	1000	5	4
20	2-0007	1000	4	4
	2-0008	1000	5	5
	2-0011	1000	5	4
	2-0012	1000	5	5
25	3-0002	1000	5	5
	3-0004	1000	5	5
	3-0006	1000	4	4
	3-0008	1000	5	5
	3-0009	1000	5	5
30	3-0012	1000	5	5
	3-0013	1000	5	5
	3-0015	1000	5	5
	3-0016	1000	5	5
	3-0017	1000	5	5
35	3-0018	1000	5	5
	3-0019	1000	5	5
	3-0020	1000	5	5
	3-0034	1000	5	5
40	3-0035	1000	5	5
	3-0036	1000	5	5
	3-0037	1000	5	5
	3-0038	1000	5	5
	3-0039	1000	5	5
45	3-0040	1000	5	5
	3-0041	1000	5	5
	3-0043	1000	5	5
	3-0044	1000	5	5
50	3-0047	1000	5	5
	3-0048	1000	5	5
	3-0049	1000	5	5
	3-0050	1000	5	5
55	3-0053	1000	5	5
	3-0054	1000	5	5
	3-0054	1000	5	5
	3-0056	1000	5	5

EP 1 364 946 A1

Table 23 (continued)

Compound No.	Active ingredient (g/ha)	Echinochloa crus-galli (L.) Beauv. Var. crus-galli	Setaria viridis (L.) Beauv.
5	3-0059	1000	5
	3-0060	1000	5
	3-0063	1000	5
	3-0070	1000	4
	3-0072	1000	5
10	3-0073	1000	5
	3-0074	1000	5
	3-0081	1000	5
	3-0082	1000	5
15	3-0083	1000	5
	3-0084	1000	5
	3-0085	1000	5
	3-0086	1000	5
	3-0087	1000	5
20	3-0088	1000	4
	3-0091	1000	5
	3-0114	1000	5
	3-0115	1000	5
25	3-0117	1000	5
	3-0118	1000	5
	3-0119	1000	5
	3-0120	1000	5
	3-0121	1000	5
30	3-0124	1000	5
	3-0125	1000	5
	3-0126	1000	5
	3-0127	1000	5
35	3-0128	1000	5
	3-0129	1000	5
	3-0130	1000	5
	3-0131	1000	5
	3-0134	1000	5
40	3-0135	1000	5
	3-0136	1000	5
	3-0137	1000	5
	3-0138	1000	4
45	3-0139	1000	5
	3-0142	1000	5
	3-0143	1000	5
	3-0144	1000	5
	3-0153	1000	5
50	3-0156	1000	5
	3-0173	1000	5
	3-0174	1000	5
	3-0180	1000	5
55	4-0001	1000	5
	4-0001	1000	4
	4-0002	1000	5
	4-0005	1000	5

EP 1 364 946 A1

Table 23 (continued)

Compound No.	Active ingredient (g/ha)	Echinochloa crus-galli (L.) Beauv. Var. crus-galli	Setaria viridis (L.) Beauv.
5	4-0006	1000	5
	4-0007	1000	5
	4-0008	1000	5
	5-0001	1000	5
	5-0002	1000	5
10	5-0003	1000	5
	5-0005	1000	4
	5-0006	1000	5
	5-0007	1000	5
15	6-0001	1000	5
	6-0003	1000	5
	6-0004	1000	5
	7-0002	1000	5
	7-0004	1000	4
20	7-0006	1000	5
	7-0007	1000	4
	7-0008	1000	5
	7-0009	1000	5
25	8-0001	1000	5
	8-0004	1000	5
	8-0005	1000	4
	8-0007	1000	5
30	9-0001	1000	5
	9-0005	1000	4
	9-0006	1000	4
	9-0007	1000	4
	9-0008	1000	5
35	10-0003	1000	5
	10-0004	1000	5
	10-0005	1000	5
	10-0006	1000	4
	10-0009	1000	5
40	10-0012	1000	4
	10-0013	1000	5
	10-0014	1000	5
	10-0015	1000	5
45	10-0016	1000	4
	10-0017	1000	5
	10-0018	1000	5

<Application Example 3> Test for herbicidal effect by upland foliage treatment

[0338] A sand was filled in a plastic pot of 80 cm². Seeds of Echinochloa crus-galli (L.) Beauv. var. crus-galli and Setaria viridis (L.) Beauv. were sowed. Breeding was made in a greenhouse for 2 weeks. Wettable powders produced in accordance with the Formulation 1 were diluted with water and sprayed on the whole foliage of plants from above the plants using a small sprayer in an amount of 1,000 liters per 1 hectare so that the amount of each active ingredient became 1,000 g per 1 hectare. Then, breeding was made in the greenhouse, and the herbicidal effect of each wettable powder was examined at the 14th day from the treatment in accordance with the standard shown in Table 21. The results are shown in Table 24.

EP 1 364 946 A1


Table 24

	Compound No. Compound No.	Active ingredient (g/ha)	Echinochloa crus-galli (L.) Beauv. Var. crus-galli	Setaria viridis (L.) Beauv.
5	1-0001	1000	5	4
	1-0004	1000	5	4
	2-0001	1000	5	4
	2-0003	1000	5	4
10	2-0004	1000	5	4
	2-0008	1000	5	5
	2-0011	1000	5	4
	3-0008	1000	4	4
15	3-0010	1000	5	4
	3-0011	1000	5	4
	3-0013	1000	5	5
	3-0015	1000	5	4
	3-0035	1000	4	4
20	3-0036	1000	4	4
	3-0037	1000	5	4
	3-0038	1000	5	5
	3-0039	1000	5	5
25	3-0044	1000	5	4
	3-0049	1000	4	4
	3-0073	1000	5	4
	3-0074	1000	5	4
	3-0076	1000	5	4
30	3-0077	1000	5	4
	3-0081	1000	4	4
	3-0082	1000	4	4
	3-0083	1000	4	4
35	3-0084	1000	4	4
	3-0085	1000	4	4
	3-0086	1000	4	4
	3-0092	1000	4	4
	3-0104	1000	5	4
40	3-0105	1000	5	4
	3-0106	1000	5	4
	3-0107	1000	5	5
	3-0115	1000	5	4
	3-0118	1000	5	4
45	3-0119	1000	5	4
	3-0120	1000	5	5
	3-0144	1000	5	5
	4-0002	1000	5	4
50	4-0005	1000	5	4
	5-0001	1000	5	4
	5-0002	1000	5	5
	5-0003	1000	5	4
	5-0007	1000	5	5
55	6-0004	1000	5	4
	7-0008	1000	5	5
	7-0009	1000	4	4

Industrial Applicability

[0340] The herbicide of the present invention has high safety to crops, particularly to rice, wheat, barley, corn, grain sorghum, soybean, cotton, sugar beat, etc.

1. An isoxazoline derivative represented by the following general formula [I] or a pharmaceutically acceptable salt thereof:



R³ and R⁴ may be the same or different and are each a hydrogen atom, a C1 to C10 alkyl group or a C3 to C8 cycloalkyl group; or R³ and R⁴ may be bonded to each other to form a C3 to C7 spiro ring together with the carbon atoms to which they bond; or R¹, R², R³ and R⁴ may form a 5- to 8-membered ring together with the carbon atoms to which they bond;

R⁵ and R⁶ may be the same or different and are each a hydrogen atom or a C1 to C10 alkyl group;

213

stituent group α , the two groups may form, together with the atoms to which they bond, a 5- to 8-membered ring which may be substituted with 1 to 4 halogen atoms; the hetero atom of the heterocyclic group, when it is a nitrogen atom, may be oxidized to become N-oxide;

n is an integer of 0 to 2.

[Substituent group α]

Hydroxyl group; thiol group; halogen atoms; C1 to C10 alkyl groups; C1 to C10 alkyl groups each mono-substituted with a group selected from the following substituent group β , C1 to C4 haloalkyl groups; C3 to C8 cycloalkyl groups; C1 to C10 alkoxy groups; C1 to C10 alkoxy groups each mono-substituted with a group selected from the following substituent group γ ; C1 to C4 haloalkoxy groups; C3 to C8 cycloalkoxy groups; C3 to C8 cycloalkyl C1 to C3 alkyloxy groups; C1 to C10 alkylthio groups; C1 to C10 alkylthio groups each mono-substituted with a group selected from the substituent group γ ; C1 to C4 haloalkylthio groups; C2 to C6 alkenyl groups; C2 to C6 alkenyloxy groups; C2 to C6 alkynyl groups; C2 to C6 alkynyloxy groups; C1 to C10 alkylsulfinyl groups; C1 to C10 alkylsulfinyl groups each mono-substituted with a group selected from the substituent group γ ; C1 to C10 alkylsulfonyl groups; C1 to C10 alkylsulfonyl groups each mono-substituted with a group selected from the substituent group γ ; C1 to C4 haloalkylsulfinyl groups; C1 to C10 alkylsulfonyloxy groups each mono-substituted with a group selected from the substituent group γ ; C1 to C4 haloalkylsulfonyloxy groups; optionally substituted phenyl group; optionally substituted phenoxy group; optionally substituted phenylthio group; optionally substituted aromatic heterocyclic groups; optionally substituted aromatic heterocyclic oxy groups; optionally substituted aromatic heterocyclic thio groups; optionally substituted phenylsulfinyl groups; optionally substituted phenylsulfonyl groups; optionally substituted aromatic heterocyclic sulfonyl groups; optionally substituted phenylsulfonyloxy groups; acyl groups; C1 to C4 haloalkylcarbonyl groups; optionally substituted benzylcarbonyl group; optionally substituted benzoyl group; carboxyl group; C1 to C10 alkoxy carbonyl groups; optionally substituted benzyloxy carbonyl group; optionally substituted phenoxy carbonyl group; cyano group; carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups and optionally substituted phenyl group); C1 to C6 acyloxy groups; C1 to C4 haloalkylcarbonyloxy groups; optionally substituted benzylcarbonyloxy group; optionally substituted benzoyloxy group; nitro group; and amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, optionally substituted phenyl group, C1 to C6 acyl groups, C1 to C4 haloalkylcarbonyl groups, optionally substituted benzylcarbonyl group, optionally substituted benzoyl group, C1 to C10 alkylsulfonyl group, C1 to C4 haloalkylsulfonyl groups, optionally substituted benzylsulfonyl group, and optionally substituted phenylsulfonyl group).

[Substituent group β]

Hydroxyl group; C3 to C8 cycloalkyl groups (which may be substituted with halogen atom or alkyl group); C1 to C10 alkoxy groups; C1 to C10 alkylthio groups; C1 to C10 alkylsulfonyl groups; C1 to C10 alkoxy carbonyl groups; C2 to C6 haloalkenyl groups; amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, C1 to C6 acyl groups; C1 to C4 haloalkylcarbonyl groups, C1 to C10 alkylsulfonyl groups and C1 to C4 haloalkylsulfonyl groups); carbamoyl group (its nitrogen atom may be substituted with same or different C1 to C10 alkyl groups); C1 to C6 acyl groups; C1 to C4 haloalkylcarbonyl groups; C1 to C10 alkoxyimino groups; cyano group; optionally substituted phenyl group; and optionally substituted phenoxy group.

[Substituent group γ]

C1 to C10 alkoxy carbonyl groups; optionally substituted phenyl group; optionally substituted aromatic heterocyclic groups; cyano group; and carbamoyl group (its nitrogen atom may be substituted with same or different C1 to C10 alkyl groups).

2. An isoxazoline derivative according to Claim 1, wherein the substituent group α on the heterocycle which may be substituted with 0 to 6 same or different groups, includes hydroxyl group; halogen atoms; C1 to C10 alkyl groups; C1 to C10 alkyl groups each mono-substituted with a group selected from the substituent group β , C1 to C4 haloalkyl groups; C3 to C8 cycloalkyl groups; C1 to C10 alkoxy groups; C1 to C10 alkoxy groups each mono-substituted with a group selected from the substituent group γ ; C1 to C4 haloalkoxy groups; C3 to C8 cycloalkoxy groups; C3 to C8 cycloalkyl C1 to C3 alkyloxy groups; C1 to C10 alkylthio groups; C1 to C10 alkylthio groups each mono-substituted with a group selected from the substituent group γ ; C1 to C4 haloalkylthio groups; C2 to C6 alkenyl groups; C2 to C6 alkenyloxy groups; C2 to C6 alkynyl groups; C2 to C6 alkynyloxy groups; C1 to C10 alkylsulfonyl groups; C1 to C4 haloalkylsulfonyl groups; optionally substituted phenyl group; optionally substituted phenoxy group; optionally substituted phenylthio group; optionally substituted aromatic heterocyclic groups; optionally substituted aromatic heterocyclic oxy groups; optionally substituted aromatic heterocyclic thio groups; optionally substituted phenylsulfonyl groups; optionally substituted aromatic heterocyclic sulfonyl groups; C1 to C6 acyl groups; C1 to C4 haloalkylcarbonyl groups; optionally substituted benzylcarbonyl group; optionally substituted benzoyl

group; carboxyl group; C1 to C10 alkoxy carbonyl groups; cyano group; carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups and optionally substituted phenyl group); nitro group; and amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, optionally substituted phenyl group, C1 to C6 acyl groups, C1 to C4 haloalkyl carbonyl groups, optionally substituted benzyl carbonyl group, optionally substituted benzoyl group, C1 to C10 alkylsulfonyl groups, C1 to C4 haloalkylsulfonyl groups, optionally substituted benzylsulfonyl group, and optionally substituted phenylsulfonyl group); when the heterocyclic group is substituted at the two adjacent positions with two alkyl groups, two alkoxy groups, an alkyl group and an alkoxy group, an alkyl group and an alkylthio group, an alkyl group and an alkylsulfonyl group, an alkyl group and a monoalkylamino group, or an alkyl group and a dialkylamino group, all selected from the substituent group α , the two groups may form, together with the atoms to which they bond, a 5- to 8-membered ring which may be substituted with 1 to 4 halogen atoms.

3. An isoxazoline derivative according to Claim 2, wherein the substituent group α on the heterocycle which may be substituted with 0 to 6 same or different groups, includes halogen atoms; C1 to C10 alkyl groups; C1 to C4 haloalkyl groups; C1 to C10 alkoxy C1 to C3 alkyl groups; C3 to C8 cycloalkyl groups (which may be substituted with halogen atom or alkyl group); C1 to C10 alkoxy groups; C1 to C4 haloalkoxy groups; C3 to C8 cycloalkyl C1 to C3 alkyloxy groups; optionally substituted phenoxy group; C1 to C10 alkylthio groups; C1 to C10 alkylsulfonyl groups; acyl groups; C1 to C4 haloalkyl carbonyl groups; C1 to C10 alkoxy carbonyl groups; cyano group and carbamoyl group (its nitrogen atom may be substituted with same or different C1 to C10 alkyl groups).
4. An isoxazoline derivative according to any of Claim 1, 2 or 3, wherein R¹ and R² may be the same or different and are each a methyl group or an ethyl group; and R³, R⁴, R⁵ and R⁶ are each a hydrogen atom.
5. An isoxazoline derivative according to any of Claim 1, 2, 3 or 4, wherein Y is a 5- or 6-membered aromatic heterocyclic group having a hetero atom selected from a nitrogen atom, an oxygen atom and a sulfur atom.
6. An isoxazoline derivative according to Claim 5, wherein Y is a thienyl group, a pyrazolyl group, an isoxazolyl group, an isothiazolyl group, a pyridyl group or a pyrimidinyl group.
7. An isoxazoline derivative according to Claim 6, wherein Y is a thiophen-3-yl group, a pyrazol-4-yl group, a pyrazol-5-yl group, an isoxazol-4-yl group, an isothiazol-4-yl group, a pyridyn-3-yl group or a pyrimidin-5-yl group.
8. An isoxazoline derivative according to Claim 7, wherein Y is a thiophen-3-yl group and the thiophene ring is substituted with the substituent group α at the 2- and 4-positions.
9. An isoxazoline derivative according to Claim 7, wherein Y is a pyrazol-4-yl group and the pyrazole ring is substituted at the 3- and 5-positions with the substituent group α and at the 1-position with a hydrogen atom, a C1 to C10 alkyl group, a C1 to C10 alkyl group mono-substituted with a group selected from the substituent group β , a C1 to C4 haloalkyl group, a C3 to C8 cycloalkyl group, a C2 to C6 alkenyl group, a C2 to C6 alkynyl group, a C1 to C10 alkylsulfinyl group, a C1 to C10 alkylsulfonyl group, a C1 to C10 alkylsulfonyl group mono-substituted with a group selected from the substituent group γ , a C1 to C4 haloalkylsulfonyl group, an optionally substituted phenyl group, an optionally substituted aromatic heterocyclic group, an optionally substituted phenylsulfonyl group, an optionally substituted aromatic heterocyclic sulfonyl group, an acyl group, a C1 to C4 haloalkyl carbonyl group, an optionally substituted benzyl carbonyl group, an optionally substituted benzoyl group, a C1 to C10 alkoxy carbonyl group, an optionally substituted benzyloxycarbonyl group, an optionally substituted phenoxycarbonyl group, a carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups and optionally substituted phenyl group), or an amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, optionally substituted phenyl group, acyl groups, C1 to C4 haloalkyl carbonyl groups, optionally substituted benzyl carbonyl group, optionally substituted benzoyl group, C1 to C10 alkylsulfonyl groups, C1 to C4 haloalkylsulfonyl groups, optionally substituted benzylsulfonyl group and optionally substituted phenylsulfonyl group).
10. An isoxazoline derivative according to Claim 7, wherein Y is a pyrazol-5-yl group and the pyrazole ring is substituted at the 4-position with the substituent group α and at the 1-position with a hydrogen atom, a C1 to C10 alkyl group, a C1 to C10 alkyl group mono-substituted with a group selected from the substituent group β , a C1 to C4 haloalkyl group, a C3 to C8 cycloalkyl group, a C2 to C6 alkenyl group, a C2 to C6 alkynyl group, a C1 to C10 alkylsulfinyl group, a C1 to C10 alkylsulfonyl group, a C1 to C10 alkylsulfonyl group mono-substituted with a group selected from the substituent group γ , a C1 to C4 haloalkylsulfonyl group, an optionally substituted phenyl group, an op-

tionally substituted aromatic heterocyclic group, an optionally substituted phenylsulfonyl group, an optionally substituted aromatic heterocyclic sulfonyl group, an acyl group, a C1 to C4 haloalkylcarbonyl group, an optionally substituted benzylcarbonyl group, an optionally substituted benzoyl group, a C1 to C10 alkoxy carbonyl group, an optionally substituted benzyloxycarbonyl group, an optionally substituted phenoxycarbonyl group, a carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups and optionally substituted phenyl group), or an amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, optionally substituted phenyl group, acyl groups, C1 to C4 haloalkylcarbonyl groups, optionally substituted benzylcarbonyl group, optionally substituted benzoyl group, C1 to C10 alkylsulfonyl groups, C1 to C4 haloalkylsulfonyl groups, optionally substituted benzylsulfonyl group and optionally substituted phenylsulfonyl group).

11. An isoxazoline derivative according to Claim 7, wherein Y is an isoxazol-4-yl group and the isoxazole ring is substituted with the substituent group α at the 3- and 5-positions.
12. An isoxazoline derivative according to Claim 7, wherein Y is an isothiazol-4-yl group and the isothiazole ring is substituted with the substituent group α at the 3- and 5-positions.
13. An isoxazoline derivative according to Claim 7, wherein Y is a pyridin-3-yl group and the pyridine ring is substituted with the substituent group α at the 2- and 4-positions.
14. An isoxazoline derivative according to Claim 7, wherein Y is a pyrimidin-5-yl group and the pyrimidine ring is substituted with the substituent group α at the 4- and 6-positions.
15. An isoxazoline derivative according to any of Claims 1 to 14, wherein n is an integer of 2.
16. An isoxazoline derivative according to any of Claims 1 to 14, wherein n is an integer of 1.
17. An isoxazoline derivative according to any of Claims 1 to 14, wherein n is an integer of 0.
18. A herbicide containing, as the active ingredient, an isoxazoline derivative set forth in any of Claims 1 to 17 or a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/01015

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl ⁷ C07D261/10, 261/12, 413/12, 498/04, 513/04, 413/14, 417/12, 261/20, 487/04, A01N43/80 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int.Cl ⁷ C07D261/10, 261/12, 413/12, 498/04, 513/04, 413/14, 417/12, 261/20, 487/04, A01N43/80 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Jitsuyo Shinan Koho 1992-1996 Jitsuyo Shinan Toroku Koho 1996-2002 Kokai Jitsuyo Shinan Koho 1971-2002 Toroku Jitsuyo Shinan Koho 1994-2002 Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS (STN) , REGISTRY (STN)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y A	JP 9-328483 A (Sankyo Co., Ltd.), 22 December, 1997 (22.12.97), Claims (Family: none)	1-8, 11, 13 15-18 9, 10, 12, 14
Y	WO 00/50410 A1 (Nippon Soda Co.), 31 August, 2000 (31.08.00), Full text & AU 200026912 A & JP 2000-297080 A	1-3, 5-8, 11, 13 15-18
Y	WO 99/23094 A1 (Nippon Soda Co.), 14 May, 1999 (14.05.99), Full text & EP 1031537 A1 & BR 9814832 A & AU 9896505 A & US 6147031 A & CN 1278259 A & JP 11-240872 A	1-8, 11, 13, 15-18
☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family	
Date of the actual completion of the international search 09 April, 2002 (09.04.02)		Date of mailing of the international search report 23 April, 2002 (23.04.02)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1998)