



(11) **EP 1 379 508 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

- (15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Description Paragraph(s) 18, 19, 22, 23, 67, 85, 87, 88, 101, 105, 135, 136, 145, 149, 150, 153, 154, 166, 167, 170
Claims EN 1, 7, 23
- (51) Int Cl.:
C07D 215/02 (2006.01)
- (86) International application number:
PCT/US2002/005979
- (87) International publication number:
WO 2002/067865 (06.09.2002 Gazette 2002/36)
- (48) Corrigendum issued on:
09.02.2011 Bulletin 2011/06
- (45) Date of publication and mention of the grant of the patent:
01.09.2010 Bulletin 2010/35
- (21) Application number: **02802229.1**
- (22) Date of filing: **28.02.2002**

(54) **N-(ARYL)-2-ARYLETHENESULFONAMIDES AND THERAPEUTIC USES THEREOF**
N-(ARYL)-2-ARYLETHENSULFONAMIDE UND IHRE THERAPEUTISCHEN ANWENDUNGEN
N-(ARYL)-2-ARYLETHENESULFONAMIDES ET LEUR USAGE THERAPEUTIQUE

- (84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE TR
- (30) Priority: **28.02.2001 US 271985 P**
- (43) Date of publication of application:
14.01.2004 Bulletin 2004/03
- (73) Proprietors:
• **Temple University - Of The Commonwealth System of Higher Education Philadelphia, PA 19122 (US)**
• **Onconova Therapeutics, Inc. Newtown, PA 18940 (US)**
- (72) Inventors:
• **REDDY, E., Premkumar Villanova, PA 19085 (US)**
• **REDDY, M., V., Ramana Upper Darby, PA 19082-5429 (US)**
• **BELL, Stanley, C. Narberth, PA 19072 (US)**
- (74) Representative: **Ouzman, Beverley Nicola Claire et al**
Murgitroyd & Company
Scotland House
165-169 Scotland Street
Glasgow G5 8PL (GB)
- (56) References cited:
WO-A-00/65913 WO-A-95/24190
US-A- 2 531 367 US-A- 4 258 058
US-A- 5 780 483 US-A- 6 121 304
US-B1- 6 191 170
- **DATABASE WPI Section Ch, Week 199236 Derwent Publications Ltd., London, GB; Class B05, AN 1992-295365 XP002352169 & JP 04 202173 A (ZERIA SHINYAKU KOGYO KK) 22 July 1992 (1992-07-22)**
 - **PUROHIT, D. M. ET AL: "Synthesis and antimicrobial activity of sulfonamides, imidazolinones, N',N''-diarylphosphinic amide derivatives with potent "dichloran" moiety" INDIAN JOURNAL OF HETEROCYCLIC CHEMISTRY , 8(1), 67-70 CODEN: IJCHEI; ISSN: 0971-1627, 1998, XP008054977**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 379 508 B9

- HARTIG, SIEGFRIED: "Derivatives of .beta.-styrenesulfonic acid" JOURNAL FUER PRAKTISCHE CHEMIE (LEIPZIG) , 33(3-4), 215-24 CODEN: JPCEAO; ISSN: 0021-8383, 1966, XP008054975
- MANECKE, GEORG ET AL: "Polymere Hydrochinonsulfonamide. 1. Mitt.: Monomere und Modellverbindungen" MAKROMOLEKULARE CHEMIE , 145, 53-66 CODEN: MACEAK; ISSN: 0025-116X, 1971, XP008054976
- OKUMA, KENTARO ET AL: "Reactions of (aryloxy)oxosulfonium ylides with carbonyl compounds" JOURNAL OF ORGANIC CHEMISTRY , 49(8), 1402-7 CODEN: JOCEAH; ISSN: 0022-3263, 1984, XP002352158
- GODFREY, CHRISTOPHER R. A. ET AL: "A novel route to unsymmetrical stilbene derivatives via intramolecular free radical ipso substitution reactions" TETRAHEDRON LETTERS , 39(7), 723-726 CODEN: TELEAY; ISSN: 0040-4039, 1998, XP002352159
- DEMILO, ALBERT B. ET AL: "Insect chemosterilants. Sulfonamides" JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY , 22(2), 197-9 CODEN: JAFCAU; ISSN: 0021-8561, 1974, XP002352160
- TRUCE, WILLIAM E. ET AL: "Cyclopropanesulfonic acid esters and amides" JOURNAL OF ORGANIC CHEMISTRY , 33(10), 3849-51 CODEN: JOCEAH; ISSN: 0022-3263, 1968, XP002352161

Description**Field of the Invention**

5 [0001] The invention relates to compositions and compounds for the treatment of proliferative disorders, including but not limited to cancer. The invention relates to the field of protecting normal cells and tissues from anticipated, planned or inadvertent exposure to ionizing radiation.

Background of the Invention

10 α - β -Unsaturated Sulfonamides

[0002] Cancer remains a leading cause of mortality in the United States and in the world. To be useful, a new chemotherapeutic agent should have a wide spectrum of activity and significant therapeutic index. Styrene- ω -sulfonanilide has been prepared by reacting styrylsulfonyl chloride with aniline (Bordwell et al., J. Amer. Chem. Soc. 68:139, 1946). This and certain other styrene- ω -sulfonanilides have been prepared by Knoevenagel-type synthesis as possible chemosterilants against the common house fly *Musca domestica* L. (Oliver et al., Synthesis 321-322, 1975).

[0003] U.S. Patent 4,035,421 to Snyder, Jr. describes the preparation of N-(3,4-dichlorophenyl)-2-phenylethanesulfonamide and its use as an antibacterial agent.

20 [0004] WO 95/24190 describes the use of tyrosine kinase inhibitors for treating cell proliferative disorders.

[0005] The styrene- ω -sulfonanilides 3'-hydroxy-4-nitrostyrene- β -sulfonanilide, 3'-hydroxy-2-nitrostyrene- β -sulfonanilide and 5'-hydroxy-2'-methyl-4-nitrostyrene- β -sulfonanilide were utilized as intermediates in the preparation of certain stilbenes by Waldau et al. Angew. Chem., Int. Ed. Engl. 11(9):826-8 (1972). The styrene- ω -sulfonanilides 3'-hydroxy-3-nitrostyrene- β -sulfonanilide and 5'-hydroxy-2'-methyl-4-nitrostyrene- β -sulfonanilide have been utilized in the preparation of stilbenes used as dyes (DE 2118493 - Farbenfab AG).

25 [0006] Aswarthamma et al., Chimica Acta Turcica 24:7-10 (1996) disclose the preparation of certain *trans*-(1-aryl-(2-anilinesulphonyl)ethylenes. No biological activity is set forth for the compounds. Touarti et al., J. Soc. Alger. Chim. 6(1): 39-52 (1996) disclose the preparation of certain α , β -unsaturated sulfonamides for inhibition of coniferyl alcohol dehydrogenase (CADH).

30 [0007] Except for the isolated teaching of antibacterial activity of N-(3,4-dichlorophenyl)-2-phenylethanesulfonamide, no useful pharmaceutical activity has been proposed for the limited numbers of α , β -unsaturated sulfonamides known to the prior art. In particular, no anti-cell proliferation or anticancer utility has been proposed for this class of compounds.

[0008] New cell antiproliferative agents, and anticancer therapeutics in particular, are needed which are useful in inhibiting proliferation of and/or killing cancer cells. In particular, such agents are needed which are selective in the killing of proliferating cells such as tumor cells, but not normal cells. Antineoplastic agents are needed which are effective against a broad range of tumor types.

Ionizing Radiation Health Risks

40 [0009] Ionizing radiation has an adverse effect on cells and tissues, primarily through cytotoxic effects. In humans, exposure to ionizing radiation occurs primarily through therapeutic techniques (such as anticancer radiotherapy) or through occupational and environmental exposure.

[0010] A major source of exposure to ionizing radiation is the administration of therapeutic radiation in the treatment of cancer or other proliferative disorders. Depending on the course of treatment prescribed by the treating physician, multiple doses may be received by a subject over the course of several weeks to several months.

45 [0011] Therapeutic radiation is generally applied to a defined area of the subject's body which contains abnormal proliferative tissue, in order to maximize the dose absorbed by the abnormal tissue and minimize the dose absorbed by the nearby normal tissue. However, it is difficult (if not impossible) to selectively administer therapeutic ionizing radiation to the abnormal tissue. Thus, normal tissue proximate to the abnormal tissue is also exposed to potentially damaging doses of ionizing radiation throughout the course of treatment. There are also some treatments that require exposure of the subject's entire body to the radiation, in a procedure called "total body irradiation", or "TBI." The efficacy of radiotherapeutic techniques in destroying abnormal proliferative cells is therefore balanced by associated cytotoxic effects on nearby normal cells. Because of this, radiotherapy techniques have an inherently narrow therapeutic index which results in the inadequate treatment of most tumors. Even the best radiotherapeutic techniques may result in incomplete tumor reduction, tumor recurrence, increasing tumor burden, and induction of radiation resistant tumors.

55 [0012] Numerous methods have been designed to reduce normal tissue damage while still delivering effective therapeutic doses of ionizing radiation. These techniques include brachytherapy, fractionated and hyperfractionated dosing, complicated dose scheduling and delivery systems, and high voltage therapy with a linear accelerator. However, such

techniques only attempt to strike a balance between the therapeutic and undesirable effects of the radiation, and full efficacy has not been achieved.

5 [0013] For example, one treatment for subjects with metastatic tumors involves harvesting their hematopoietic stem cells and then treating the subject with high doses of ionizing radiation. This treatment is designed to destroy the subject's tumor cells, but has the side effect of also destroying their normal hematopoietic cells. Thus, a portion of the subject's bone marrow (containing the hematopoietic stem cells), is removed prior to radiation therapy. Once the subject has been treated, the autologous hematopoietic stem cells are returned to their body.

10 [0014] However, if tumor cells have metastasized away from the tumor's primary site, there is a high probability that some tumor cells will contaminate the harvested hematopoietic cell population. The harvested hematopoietic cell population may also contain neoplastic cells if the subject suffers from a cancer of the bone marrow such as the various French-American-British (FAB) subtypes of acute myelogenous leukemias (AML), chronic myeloid leukemia (CML), or acute lymphocytic leukemia (ALL). Thus, the metastasized tumor cells or resident neoplastic cells must be removed or killed prior to reintroducing the stem cells to the subject. If any living tumorigenic or neoplastic cells are reintroduced into the subject, they can lead to a relapse.

15 [0015] Prior art methods of removing tumorigenic or neoplastic cells from harvested bone marrow are based on a whole-population tumor cell separation or killing strategy, which typically does not kill or remove all of the contaminating malignant cells. Such methods include leukopheresis of mobilized peripheral blood cells, immunoaffinity-based selection or killing of tumor cells, or the use of cytotoxic or photosensitizing agents to selectively kill tumor cells. In the best case, the malignant cell burden may still be at 1 to 10 tumor cells for every 100,000 cells present in the initial harvest (Lazarus et al. J. of Hematotherapy, 2(4):457-66, 1993).

20 [0016] Thus, there is needed a purging method designed to selectively destroy the malignant cells present in the bone marrow, while preserving the normal hematopoietic stem cells needed for hematopoietic reconstitution in the transplantation subject.

25 [0017] Exposure to ionizing radiation can also occur in the occupational setting. Occupational doses of ionizing radiation may be received by persons whose job involves exposure (or potential exposure) to radiation, for example in the nuclear power and nuclear weapons industries. Military personnel stationed on vessels powered by nuclear reactors, or soldiers required to operate in areas contaminated by radioactive fallout, risk similar exposure to ionizing radiation. Occupational exposure may also occur in rescue and emergency personnel called in to deal with catastrophic events involving a nuclear reactor or radioactive material. Other sources of occupational exposure may be from machine parts, plastics, and solvents left over from the manufacture of radioactive medical products, smoke alarms, emergency signs, and other consumer goods. Occupational exposure may also occur in persons who serve on nuclear powered vessels, particularly those who tend the nuclear reactors, in military personnel operating in areas contaminated by nuclear weapons fallout, and in emergency personnel who deal with nuclear accidents. Environmental exposure to ionizing radiation may also result from nuclear weapons detonations (either experimental or during wartime), discharges of actinides from nuclear waste storage and processing and reprocessing of nuclear fuel, and from naturally occurring radioactive materials such as radon gas or uranium. There is also increasing concern that the use of ordnance containing depleted uranium results in low-level radioactive contamination of combat areas.

30 [0018] Radiation exposure from any source can be classified as acute (a single large exposure) or chronic (a series of small low-level, or continuous low-level exposures spread over time). Radiation sickness generally results from an acute exposure of a sufficient dose, and presents with a characteristic set of symptoms that appear in an orderly fashion, including hair loss, weakness, vomiting, diarrhea, skin burns and bleeding from the gastrointestinal tract and mucous membranes. Genetic defects, sterility and cancers (particularly bone marrow cancer) often develop over time. Chronic exposure is usually associated with delayed medical problems such as cancer and premature ageing. An acute total body exposure of 125,000 millirem may cause radiation sickness. Localized doses such as those used in radiotherapy may not cause radiation sickness, but may result in the damage or death of exposed normal cells.

35 [0019] For example, an acute total body radiation dose of 100,000 - 125,000 millirem (equivalent to 1 Gy) received in less than one week would result in observable physiologic effects such as skin burns or rashes, mucosal and GI bleeding, nausea, diarrhea and/or excessive fatigue. Longer term cytotoxic and genetic effects such as hematopoietic and immunocompetent cell destruction, hair loss (alopecia), gastrointestinal, and oral mucosal sloughing, venoocclusive disease of the liver and chronic vascular hyperplasia of cerebral vessels, cataracts, pneumonites, skin changes, and an increased incidence of cancer may also manifest over time. Acute doses of less than 10,000 millirem (equivalent to 0.1 Gy) typically will not result in immediately observable biologic or physiologic effects, although long term cytotoxic or genetic effects may occur.

40 [0020] A sufficiently large acute dose of ionizing radiation, for example 500,000 to over 1 million millirem (equivalent to 5 - 10 Gy), may kill a subject immediately. Doses in the hundreds of thousands of millirems may kill within 7 to 21 days from a condition called "acute radiation poisoning." Reportedly, some of the Chernobyl firefighters died of acute radiation poisoning, having received acute doses in the range of 200,000 - 600,000 millirem (equivalent to 2 - 6 Gy). Acute doses below approximately 200,000 millirem do not result in death, but the exposed subject will likely suffer long-

EP 1 379 508 B9

term cytotoxic or genetic effects as discussed above.

[0021] Acute occupational exposures usually occur in nuclear power plant workers exposed to accidental releases of radiation, or in fire and rescue personnel who respond to catastrophic events involving nuclear reactors or other sources of radioactive material. Suggested limits for acute occupational exposures in emergency situations were developed by the Brookhaven National Laboratories, and are given in Table 1.

Table 1: Acute Occupational Exposure Limits for Emergency Operations

Whole Body Conditions for Dose Limit	Activity Required	Conditions for Exposure
10,000 millirem*	Protect property	Voluntary, when lower dose not practical
25,000 millirem	Lifesaving Operation; Protect General Public	Voluntary, when lower dose not practical
>25,000 millirem	Lifesaving operation; Protect large population	Voluntary, when lower dose not practical, and the risk has been clearly explained
*100,000 millirem equals one sievert (Sv). For penetrating radiation such as gamma radiation, one Sv equals approximately one Gray (Gy). Thus, the dosage in Gy can be estimated as 1 Gy for every 100,000 millirem.		

[0022] A chronic dose is a low level (*i.e.*, 100 - 5000 millirem) incremental or continuous radiation dose received over time. Examples of chronic doses include a whole body dose of ~5000 millirem per year, which is the dose typically received by an adult working at a nuclear power plant. By contrast, the Atomic Energy Commission recommends that members of the general public should not receive more than 100 millirem per year. Chronic doses may cause long-term cytotoxic and genetic effects, for example manifesting as an increased risk of a radiation-induced cancer developing later in life. Recommended limits for chronic exposure to ionizing radiation are given in Table 2.

Table 2: Annual Chronic Occupational Radiation Exposure Limits

Organ or Subject	Annual Occupational Dose in millirem
Whole Body	5000
Lens of the Eye	15,000
Hands and wrists	50,000
Any individual organ	50,000
Pregnant worker	500 /9 months
Minor (16-18) receiving training	100

[0023] By way of comparison, Table 3 sets forth the radiation doses from common sources.

Table 3: Radiation Dosages From Common Sources

Sources	Dose In Millirem
Television	<1/yr
Gamma Rays, Jet Cross Country	1
Mountain Vacation - 2 week	3
Atomic Test Fallout	5
U.S. Water, Food & Air (Average)	30/yr
Wood	50/yr
Concrete	50/yr
Brick	75/yr

(continued)

Sources	Dose In Millirem
Chest X-Ray	100
Cosmic Radiation (Sea Level)	40/yr (add 1 millirem/100 ft elev.)
Natural Background San Francisco	120/yr
Natural Background Denver	50/yr
Atomic Energy Commission Limit For Workers	5000/yr
Complete Dental X-Ray	5000
Natural Background at Pocos de Caldras, Brazil	7000/yr
Whole Body Diagnostic X-Ray	100,000
Cancer Therapy	500,000 (localized)
Radiation Sickness-Nagasaki	125,000 (single doses)
LD ₅₀ Nagasaki & Hiroshima	400,000-500,000 (single dose)

[0024] Chronic doses of greater than 5000 millirem per year (0.05 Gy per year) may result in long-term cytotoxic or genetic effects similar to those described for persons receiving acute doses. Some adverse cytotoxic or genetic effects may also occur at chronic doses of significantly less than 5000 millirem per year. For radiation protection purposes, it is assumed that any dose above zero can increase the risk of radiation-induced cancer (*i.e.*, that there is no threshold). Epidemiologic studies have found that the estimated lifetime risk of dying from cancer is greater by about 0.04% per rem of radiation dose to the whole body.

[0025] While anti-radiation suits or other protective gear may be effective at reducing radiation exposure, such gear is expensive, unwieldy, and generally not available to the public. Moreover, radioprotective gear will not protect normal tissue adjacent a tumor from stray radiation exposure during radiotherapy. What is needed, therefore, is a practical way to protect subjects who are scheduled to incur, or are at risk for incurring, exposure to ionizing radiation. In the context of therapeutic irradiation, it is desirable to enhance protection of normal cells while causing tumor cells to remain vulnerable to the detrimental effects of the radiation. Furthermore, it is desirable to provide systemic protection from anticipated or inadvertent total body irradiation, such as may occur with occupational or environmental exposures, or with certain therapeutic techniques.

[0026] Pharmaceutical radioprotectants offer a cost-efficient, effective and easily available alternative to radioprotective gear. However, previous attempts at radioprotection of normal cells with pharmaceutical compositions have not been entirely successful. For example, cytokines directed at mobilizing the peripheral blood progenitor cells confer a myelo-protective effect when given prior to radiation (Neta et al., *Semin. Radiat. Oncol.* 6:306-320, 1996), but do not confer systemic protection. Other chemical radioprotectors administered alone or in combination with biologic response modifiers have shown minor protective effects in mice, but application of these compounds to large mammals was less successful, and it was questioned whether chemical radioprotection was of any value (Maisin, J.R., Bacq and Alexander Award Lecture. "Chemical radioprotection: past, present, and future prospects", *Int J. Radiat Biol.* 73:443-50, 1998). Pharmaceutical radiation sensitizers, which are known to preferentially enhance the effects of radiation in cancerous tissues, are clearly unsuited for the general systemic protection of normal tissues from exposure to ionizing radiation.

[0027] Therapeutic agents are needed to protect subjects who have incurred, or are at risk for incurring exposure to ionizing radiation. In the context of therapeutic irradiation, it is desirable to enhance protection of normal cells while causing tumor cells to remain vulnerable to the detrimental effects of the radiation. Furthermore, it is desirable to provide systemic protection from anticipated or inadvertent total body irradiation, such as may occur with occupational or environmental exposures, or with certain therapeutic techniques.

Summary of the Invention

[0028] It is an object of the invention to provide compounds and compositions. The biologically active compounds are in the form of N-(aryl)-2-arylethanesulfonamides, and pharmaceutically acceptable salts thereof.

[0029] It is an object of the invention to provide compounds and compositions for the treatment of cancer and other proliferative diseases.

[0030] It is an object of the invention to provide compounds which are selective in killing tumor cells but not normal cells.

[0031] It is an object of the invention to provide compounds and compositions for inducing neoplastic cells to selectively

undergo apoptosis.

[0032] It is an object of the invention to provide compounds and compositions for protecting normal cells and tissues from the cytotoxic and genetic effects of exposure to ionizing radiation, in subjects who have incurred or are at risk for incurring exposure to ionizing radiation. The exposure to ionizing radiation may occur in controlled doses during the treatment of cancer and other proliferative disorders, or may occur in uncontrolled doses beyond the norm accepted for the population at large during high risk activities or environmental exposures.


[0033] In another aspect, a radioprotectant N-(aryl)-2-arylethanesulfonamide compound for treating cancer or other proliferative disorders is provided, for administration prior to administering an effective amount of ionizing radiation, wherein the radioprotective N-(aryl)-2-arylethanesulfonamide compound induces a temporary radioresistant phenotype in the subject's normal tissue.

[0034] In a further aspect, the invention provides a radioprotective N-(aryl)-2-arylethanesulfonamide for safely increasing the dosage of therapeutic ionizing radiation used in the treatment of cancer or other proliferative disorders, wherein an effective amount of at least one radioprotective N-(aryl)-2-arylethanesulfonamide compound is administered prior to administration of the therapeutic ionizing radiation, which radioprotective compound induces a temporary radioresistant phenotype in the subject's normal tissue.

[0035] In yet a further aspect, the invention provides a radioprotective N-(aryl)-2-arylethanesulfonamide for treating individuals who have incurred or are at risk for incurring remediable radiation damage from exposure to ionizing radiation. In one embodiment, an effective amount of at least one radioprotective N-(aryl)-2-arylethanesulfonamide compound is administered to the subject before the subject incurs remediable radiation damage from exposure to ionizing radiation. In another embodiment, an effective amount of at least one radioprotective N-(aryl)-2-arylethanesulfonamide compound is administered to the subject after the subject incurs remediable radiation damage from exposure to ionizing radiation.

[0036] In yet another embodiment, the invention provides an N-(aryl)-2-arylethanesulfonamide compound for purging bone marrow of neoplastic cells (such as leukemic cells) or tumor cells which have metastasized into the bone marrow. In this embodiment, bone marrow cells from an individual afflicted with a proliferative disorder are harvested, an effective amount of at least one N-(aryl)-2-arylethanesulfonamide compound is used to treat the harvested bone marrow cells, and the treated bone marrow cells are subjected to an effective amount of ionizing radiation. The harvested cells are then returned to the body of the afflicted individual.

[0037] In another aspect, the invention is directed to novel compounds of formula I:



wherein:

Q_1 is selected from the group consisting of substituted and unsubstituted aryl, and substituted and unsubstituted heteroaryl;

Q_2 is selected from the group consisting of substituted aryl, and substituted and unsubstituted heteroaryl;

R is selected from the group consisting of hydrogen and (C₁-C₆)alkyl;

wherein the substituents for the substituted aryl and substituted heteroaryl groups comprising Q_1 are independently selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, cyano, carboxy, hydroxy, (C₂-C₆)hydroxyalkyl, phosphonato, amino, (C₁-C₆)acylamino, sulfamyl, acetoxy, di(C₁-C₆)alkylamino(C₂-C₆)alkoxy, trifluoromethyl and



wherein:

X is oxygen or sulfur,

R₅ is selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)heteroalkyl, substituted phenyl, and unsubstituted phenyl, and

R₆ is selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)heteroalkyl, substituted aryl, unsubstituted aryl, substituted heteroaryl, unsubstituted heteroaryl, substituted aryl(C₁-C₃)alkyl, unsubstituted aryl-(C₁-C₃)alkyl and (C₁-C₆)alkoxycarbonyl(C₁-C₆)alkylenyl;

wherein the substituents for the substituted aryl and substituted heteroaryl groups comprising Q₂, and the substituents for the substituted aryl and substituted heteroaryl groups comprising or included within R₅ and R₆, are independently selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, nitro, cyano, carboxy, carboxy(C₁-C₃)alkoxy, hydroxy, (C₂-C₆)hydroxyalkyl, phosphonato, amino, (C₁-C₆)acylamino, sulfamyl, acetoxy, di(C₁-C₆)alkylamino(C₂-C₆ alkoxy) and trifluoromethyl;

provided that when R is hydrogen:

(a) when Q₁ is unsubstituted phenyl, Q₂ is other than dimethoxyphenyl, 2-methylphenyl, 2-chlorophenyl, 4-chlorophenyl, 4-N,N-dimethylaminophenyl, 4-methylphenyl, 4-methoxyphenyl, 4-nitrophenyl, 3-methoxy-4-hydroxyphenyl, unsubstituted pyrenyl, unsubstituted benzodioxolyl, and unsubstituted 2-thienyl;

(b) when Q₁ is 3-hydroxyphenyl, Q₂ is other than nitrophenyl;

(c) when Q₁ is 2-methyl-5-hydroxyphenyl, Q₂ is other than 4-nitrophenyl; and

(d) when Q₁ is unsubstituted 2-pyridyl, Q₂ is other than 3-methoxy-4-hydroxyphenyl;

or a pharmaceutically acceptable salt thereof.

[0038] In a further sub-embodiment, novel compounds of formula I are provided wherein Q₁ and Q₂ are independently selected from the group consisting of substituted aryl and substituted heteroaryl; R is defined as above; the substituents for the substituted aryl and substituted heteroaryl groups comprising Q₁ are defined as above; the substituents for the substituted aryl and substituted heteroaryl groups comprising Q₂, and the substituents for the substituted aryl and substituted heteroaryl groups comprising or included within R₅ and R₆, are defined as above, provided, when R is hydrogen:

(i) Q₁ may not be dinitrophenyl;

(ii) Q₂ may not be dinitrophenyl; and


(iii) when Q₂ is mononitrophenyl:

Q₁ is other than substituted phenyl, or

Q₁ is substituted phenyl wherein at least the 4-position is substituted, and the substituent is other than hydroxy;

or a pharmaceutically acceptable salt thereof.


[0039] According to another embodiment, the invention is directed to a process for preparing a novel compound as defined above, the process comprising reacting a compound of the formula **B**:



with a compound of the formula C




in a nonprotic solvent in the presence of a base to form a compound of the formula:




10 wherein R, Q₁ and Q₂ are defined as above. Compound **B** may be prepared by reacting a compound of the formula **A**, Q₂-CH=CH₂, with sulfonyl chloride in the presence of a nonprotic solvent.


15 **[0040]** According to another embodiment, the invention is directed to an alternative process for preparing a novel compound as defined above, said process comprising reacting a compound of the formula **G**



25 with a compound of the formula H




35 in the presence of a basic catalyst to form a compound of the formula:



wherein R, Q₁ and Q₂ are defined as above.

50 **[0041]** Compounds of formula **G** may be prepared by reacting a compound of the formula **E**, ClSO₂-CH₂-C(O)OR', with a compound of formula **C** (as defined above) in a nonprotic solvent in the presence of a base to form a compound of the formula **F**,



10 and then treating the formula **F** compound with a base capable of hydrolyzing the ester function thereof to an acid to form compound **G**; wherein **R** and Q_1 are defined as above, and R' is methyl or ethyl.

[0042] According to another embodiment of the invention, pharmaceutical compositions are provided comprising a pharmaceutically acceptable carrier and a compound according to formula I wherein

Q_1 , Q_2 and **R** are as defined as above;

15 wherein the substituents for the substituted aryl and substituted heteroaryl groups comprising Q_1 are independently selected from the group consisting of halogen, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, cyano, carboxy, hydroxy, (C_2-C_6) hydroxyalkyl, phosphonato, amino, (C_1-C_6) acylamino, sulfamyl, acetoxyl, di (C_1-C_6) alkylamino (C_2-C_6) alkoxy, trifluoromethyl; and



25 wherein:

X is oxygen or sulfur,

R_5 is selected from the group consisting of hydrogen, (C_1-C_6) alkyl, (C_2-C_6) heteroalkyl, substituted phenyl, and unsubstituted phenyl, and

30 R_6 is selected from the group consisting of hydrogen, (C_1-C_6) alkyl, (C_2-C_6) heteroalkyl, substituted aryl, unsubstituted aryl, substituted heteroaryl, unsubstituted heteroaryl, substituted aryl (C_1-C_3) alkyl, unsubstituted aryl- (C_1-C_3) alkyl and (C_1-C_6) alkoxycarbonyl (C_1-C_6) alkylenyl;

35 wherein the substituents for the substituted aryl and substituted heteroaryl groups comprising Q_2 , and the substituents for the substituted aryl and substituted heteroaryl groups comprising or included within R_5 and R_6 , are independently selected from the group consisting of halogen, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, nitro, cyano, carboxy, carboxy (C_1-C_3) alkoxy, hydroxy, (C_2-C_6) hydroxyalkyl, phosphonato, amino, (C_1-C_6) acylamino, sulfamyl, acetoxyl, di (C_1-C_6) alkylamino (C_2-C_6) alkoxy) and trifluoromethyl;

40 provided, when **R** is hydrogen and Q_2 is unsubstituted phenyl, then Q_1 must be other than 3,4-dichlorophenyl, more particularly other than 3,4-dihalophenyl, even more particularly other than dihalophenyl; or a pharmaceutically acceptable salt thereof.

[0043] According to another embodiment of the invention, an N-(aryl)-2-arylethanesulfonamide compound for treating a proliferative disorder in an individual is provided.

45 **[0044]** According to another embodiment of the invention, an N-(aryl)-2-arylethanesulfonamide compound for inducing apoptosis of tumor cells in an individual afflicted with cancer is provided.

[0045] According to another embodiment of the invention, an N-(aryl)-2-arylethanesulfonamide compound for reducing or eliminating the effects of ionizing radiation on normal cells in a subject who has incurred or is at risk for incurring exposure to ionizing radiation is provided. An effective amount of the at least one N-(aryl)-2-arylethanesulfonamide compound is administered to the subject prior to or after exposure to ionizing radiation.

50 **[0046]** An N-(aryl)-2-arylethanesulfonamide compound for safely increasing the dosage of therapeutic ionizing radiation used in the treatment of cancer or other proliferative disorders is also provided. An effective amount of the at least one radioprotective N-(aryl)-2-arylethanesulfonamide compound is administered prior to administration of the therapeutic ionizing radiation, which radioprotective compound induces a temporary radioresistant phenotype in the normal tissue of the subject.

55 **[0047]** An N-(aryl)-2-arylethanesulfonamide compound for treating a subject who has incurred or is at risk for incurring remediable radiation damage from exposure to ionizing radiation is also provided, wherein an effective amount of at least one radioprotective N-(aryl)-2-arylethanesulfonamide compound is administered prior to or after incurring remedial radiation damage from exposure to ionizing radiation.

[0048] For all of the aforementioned uses, the administered compound is a compound according to formula I wherein:

Q₁ and Q₂ and R are as defined as above;

wherein the substituents for the substituted aryl and substituted heteroaryl groups comprising Q₁ are independently selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, cyano, carboxy, hydroxy, (C₂-C₆)hydroxyalkyl, phosphonato, amino, (C₁-C₆)acylamino, sulfamyl, acetoxy, di(C₁-C₆)alkylamino(C₂-C₆)alkoxy, trifluoromethyl and



wherein:

X is oxygen or sulfur,

R₅ is selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)heteroalkyl, substituted phenyl, and unsubstituted phenyl, and

R₆ is selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)heteroalkyl, substituted aryl, unsubstituted aryl, substituted heteroaryl, unsubstituted heteroaryl, substituted aryl(C₁-C₃)alkyl, unsubstituted aryl-(C₁-C₃)alkyl and (C₁-C₆)alkoxycarbonyl(C₁-C₆)alkylenyl;

wherein the substituents for the substituted aryl and substituted heteroaryl groups comprising Q₂, and the substituents for the substituted aryl and substituted heteroaryl groups comprising or included within R₅ and R₆, are independently selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, nitro, cyano, carboxy, carboxy(C₁-C₃)alkoxy, hydroxy, (C₂-C₆)hydroxyalkyl, phosphonato, amino, (C₁-C₆)acylamino, sulfamyl, acetoxy, di(C₁-C₆)alkylamino(C₂-C₆alkoxy) and trifluoromethyl;

or a pharmaceutically acceptable salt thereof.

[0049] The term "acyl" means a radical of the general formula -C(=O)-R, wherein -R is hydrogen, hydrocarbyl, amino or alkoxy. Examples include for example, acetyl (-C(=O)CH₃), propionyl (-C(=O)CH₂CH₃), benzoyl (-C(=O)C₆H₅), phenylacetyl (-C(=O)CH₂C₆H₅), carboethoxy (-CO₂Et), and dimethylcarbamoyl (-C(=O)N(CH₃)₂).

[0050] The term "aromatic" refers to a carbocycle or heterocycle having one or more polyunsaturated rings having aromatic character (4n + 2) delocalized π (pi) electrons).

[0051] The term "(C₂-C₆)acylamino" means a radical containing a two to six carbon straight or branched chain acyl group attached to a nitrogen atom via the acyl carbonyl carbon. Examples include -NHC(O)CH₂CH₂CH₃ and -NHC(O)CH₂CH₂CH₂CH₂CH₃.

[0052] The term "alkyl", by itself or as part of another substituent means, unless otherwise stated, a straight or branched chain hydrocarbon radical, including di- and multi-radicals, having the number of carbon atoms designated (*i.e.* (C₁-C₆) means one to six carbons) and includes straight or branched chain groups. Most preferred is (C₁-C₃)alkyl, ethyl or methyl.

[0053] The term "alkoxy" employed alone or in combination with other terms means, unless otherwise stated, an alkyl group having the designated number of carbon atoms, as defined above, connected to the rest of the molecule via an oxygen atom, such as, for example, methoxy, ethoxy, 1-propoxy, 2-propoxy and the higher homologs and isomers. Preferred are (C₁-C₃)alkoxy, ethoxy or methoxy.

[0054] The term "alkylenyl" by itself or as part of another substituent means a divalent radical derived from a straight or branched chain alkane having the indicated number of carbon atoms, as exemplified by the four-carbon radical -CH₂CH₂CH₂CH₂-.

[0055] The term "alkenyl" employed alone or in combination with other terms, means, unless otherwise stated, a stable straight chain or branched monounsaturated or diunsaturated hydrocarbon group having the stated number of carbon atoms. Examples include vinyl, propenyl (allyl), crotyl, isopentenyl, butadienyl, 1,3-pentadienyl, 1,4-pentadienyl, and the higher homologs and isomers. A divalent radical derived from an alkene is exemplified by -CH=CH-CH₂-.

[0056] The term "carboxy(C₁-C₃)alkoxy" means a radical in which the carboxy group -COOH is attached to a carbon of a straight or branched chain alkoxy group containing one to three carbon atoms. The radical thus contains up to four carbon atoms. Examples include HOC(O)CH₂CH₂CH₂O- and HOC(O)CH₂CH₂O-.

[0057] The term "heteroalkyl" by itself or in combination with another term means, unless otherwise stated, a stable straight or branched chain radical consisting of the stated number of carbon atoms and one or two heteroatoms selected from the group consisting of O, N, and S, and wherein the nitrogen and sulfur atoms may be optionally oxidized and the

nitrogen heteroatom may be optionally quaternized. The heteroatom(s) may be placed at any position of the heteroalkyl group, including between the rest of the heteroalkyl group and the fragment to which it is attached, as well as attached to the most distal carbon atom in the heteroalkyl group. Examples include: $-O-CH_2-CH_2-CH_3$, $-CH_2-CH_2-CH_2-OH$, $-CH_2-CH_2-NH-CH_3$, $-CH_2-S-CH_2-CH_3$, and $-CH_2-CH_2-S(O)-CH_3$. Up to two heteroatoms may be consecutive, such as, for example, $-CH_2-NH-OCH_3$.

[0058] The term "heteroalkenyl" by itself or in combination with another term means, unless otherwise stated, a stable straight or branched chain monounsaturated or diunsaturated hydrocarbon radical consisting of the stated number of carbon atoms and one or two heteroatoms selected from the group consisting of O, N, and S, and wherein the nitrogen and sulfur atoms may optionally be oxidized and the nitrogen heteroatom may optionally be quaternized. Up to two heteroatoms may be placed consecutively. Examples include $-CH=CH-O-CH_3$, $-CH=CH-CH_2-OH$, $-CH_2-CH=N-OCH_3$, $-CH=CH-N(CH_3)-CH_3$, and $-CH_2-CH=CH-CH_2-SH$.

[0059] The term "hydroxyalkyl" means an alkyl radical wherein one or more of the carbon atoms is substituted with hydroxy. Examples include $-CH_2CH(OH)CH_3$ and $-CH_2CH_2OH$. The terms "halo" or "halogen" by themselves or as part of another substituent mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom.

[0060] The term "(C₁-C₆)alkoxycarbonyl(C₁-C₆)alkylenyl" means a group of the formula $CH_3(CH_2)_pOC(O)(CH_2)_q-$ wherein p is an integer from zero to five and q is an integer from one to six.

[0061] The term "di(C₁-C₆)alkylamino(C₂-C₆)alkoxy" means $(alkyl)_2N(CH_2)_nO-$ wherein the two alkyl chains connected to the nitrogen atom independently contain from one to six carbon atoms, preferably from one to three carbon atoms, and n is an integer from 2 to 6. Preferably, n is 2 or 3. Most preferably, n is 2, and the alkyl groups are methyl, that is, the group is the dimethylaminoethoxy group, $(CH_3)_2NCH_2CH_2O-$.

[0062] The term "phosphonato" means the group $-PO(OH)_2$.

[0063] The term "sulfamyl" means the group $-SO_2NH_2$.

[0064] The term "aryl" employed alone or in combination with other terms, means, unless otherwise stated, a carbocyclic aromatic system containing one or more rings (typically one, two or three rings) wherein such rings may be attached together in a pendent manner or may be fused. Examples include phenyl; anthracyl; and naphthyl, particularly 1-naphthyl and 2-naphthyl.

[0065] The term "aryl-(C₁-C₃)alkyl" means a radical wherein a one to three carbon alkylene chain is attached to an aryl group, *e.g.*, $-CH_2CH_2$ -phenyl. Similarly, the term "heteroaryl-(C₁-C₃)alkyl" means a radical wherein a one to three carbon alkylene chain is attached to a heteroaryl group, *e.g.*, $-CH_2CH_2$ -pyridyl. The term "substituted aryl-(C₁-C₃)alkyl" means an aryl-(C₁-C₃)alkyl radical in which the aryl group is substituted. The term "substituted heteroaryl-(C₁-C₃)alkyl" means a heteroaryl-(C₁-C₃)alkyl radical in which the heteroaryl group is substituted.

[0066] The term "heteroaryl" by itself or as part of another substituent means, unless otherwise stated, an unsubstituted or substituted, stable, mono- or multicyclic heterocyclic aromatic ring system which consists of carbon atoms and from one to four heteroatoms selected from the group consisting of N, O, and S, and wherein the nitrogen and sulfur heteroatoms may be optionally oxidized, and the nitrogen atom may be optionally quaternized. The heterocyclic system may be attached, unless otherwise stated, at any heteroatom or carbon atom which affords a stable structure.

[0067] Examples of such heteroaryls include benzimidazolyl, particularly 2-benzimidazolyl; benzofuryl, particularly 3-, 4-, 5-, 6- and 7-benzofuryl; 2-benzothiazolyl and 5-benzothiazolyl; benzothienyl, particularly 3-, 4-, 5-, 6-, and 7-benzothienyl; 4-(2-benzoxazolyl); furyl, particularly 2- and 3-furyl; isoquinolyl, particularly 1- and 5-isoquinolyl; isoxazolyl, particularly 3-, 4- and 5-isoxazolyl; imidazolyl, particularly 2-, 4- and 5-imidazolyl; indolyl, particularly 3-, 4-, 5-, 6- and 7-indolyl; oxazolyl, particularly 2-, 4- and 5-oxazolyl; purinyl; pyrrolyl, particularly 2-pyrrolyl, 3-pyrrolyl; pyrazolyl, particularly 3- and 5-pyrazolyl; pyrazinyl, particularly 2-pyrazinyl; pyridazinyl, particularly 3- and 4-pyridazinyl; pyridyl, particularly 2-, 3- and 4-pyridyl; pyrimidinyl, particularly 2- and 4-pyrimidinyl; quinoxalinyl, particularly 2- and 5-quinoxalinyl; quinolinyl, particularly 2- and 3-quinolinyl; 5-tetrazolyl; thiazolyl; particularly 2-thiazolyl, 4-thiazolyl and 5-thiazolyl; thienyl, particularly 2- and 3-thienyl; and 3-(1,2,4-triazolyl). The aforementioned listing of heteroaryl moieties is intended to be representative, not limiting. In another embodiment of the invention, Q₁ is independently selected from the group consisting of substituted and unsubstituted aryl, and substituted and unsubstituted heteroaryl, provided that Q₁ is not 2-thiazolyl. In a further embodiment of the invention, Q₁ is independently selected from the group consisting of substituted and unsubstituted aryl, and substituted and unsubstituted heteroaryl, provided that Q₁ is not 2-thiazolyl, 4-thiazolyl or 5-thiazolyl.

[0068] The term "substituted" means that an atom or group of atoms has replaced hydrogen as the substituent attached to another group. For aryl and heteroaryl groups, by "substituted" is meant any level of substitution, namely mono-, di-, tri-, tetra-, or penta-substitution. The substituents are independently selected.

[0069] For purposes of this disclosure, the term aryl in the expression "N-(aryl)-2-arylethanesulfonamide" is deemed to include both "aryl" and "heteroaryl" radicals, either substituted or unsubstituted, unless otherwise indicated.

[0070] The term "subject" or "individual" includes human beings and non-human animals. With respect to the disclosed radioprotective methods, these terms further refer to an organism which is scheduled to incur, is at risk for incurring, or has incurred, exposure to ionizing radiation.

[0071] As used herein, "ionizing radiation" is radiation of sufficient energy that, when absorbed by cells and tissues,

induces formation of reactive oxygen species and DNA damage. This type of radiation includes X-rays, gamma rays, and particle bombardment (*e.g.*, neutron beam, electron beam, protons, mesons and others), and is used for medical testing and treatment, scientific purposes, industrial testing, manufacturing and sterilization, weapons and weapons development, and many other uses. Radiation is typically measured in units of absorbed dose, such as the rad or gray (Gy), or in units of dose equivalence, such as the rem or sievert (Sv). The relationship between these units is given below:

rad and gray (Gy)	rem and sievert (Sv)
1 rad = 0.01 Gy	1 rem = 0.01 Sv

[0072] The Sv is the Gy dosage multiplied by a factor that includes tissue damage done. For example, penetrating ionizing radiation (*e.g.*, gamma and beta radiation) have a factor of about 1, so 1 Sv = ~1 Gy. Alpha rays have a factor of 20, so 1 Gy of alpha radiation = 20 Sv.

[0073] By "effective amount of ionizing radiation" is meant an amount of ionizing radiation effective in killing, or reducing the proliferation, of abnormally proliferating cells in a subject. As used with respect to bone marrow purging, "effective amount of ionizing radiation" means an amount of ionizing radiation effective in killing, or in reducing the proliferation, of malignant cells in a bone marrow sample removed from a subject.

[0074] By "acute exposure to ionizing radiation" or "acute dose of ionizing radiation" is meant a dose of ionizing radiation absorbed by a subject in less than 24 hours. The acute dose may be localized, as in radiotherapy techniques, or may be absorbed by the subject's entire body. Acute doses are typically above 10,000 millirem (0.1 Gy), but may be lower.

[0075] By "chronic exposure to ionizing radiation" or "chronic dose of ionizing radiation" is meant a dose of ionizing radiation absorbed by a subject over a period greater than 24 hours. The dose may be intermittent or continuous, and may be localized or absorbed by the subject's entire body. Chronic doses are typically less than 10,000 millirem (0.1 Gy), but may be higher.

[0076] By "effective amount of radioprotective N-(aryl)-2-arylethanesulfonamide compound" is meant an amount of compound effective to reduce or eliminate the toxicity associated with radiation in normal cells of the subject, and also to impart a direct cytotoxic effect to abnormally proliferating cells in the subject. As used with respect to bone marrow purging, "effective amount of the radioprotective N-(aryl)-2-arylethanesulfonamide compound" means an amount of compound effective to reduce or eliminate the toxicity associated with radiation in bone marrow removed from a subject, and also to impart a direct cytotoxic effect to malignant cells in the bone marrow removed from the subject.

[0077] By "at risk of incurring exposure to ionizing radiation" is meant that a subject may advertently (such as by scheduled radiotherapy sessions) or inadvertently be exposed to ionizing radiation in the future. Inadvertent exposure includes accidental or unplanned environmental or occupational exposure.

Description of the Figures

[0078]

Fig. 1 is a plot of the growth inhibition effect of (E)-4-methoxystyryl-N-4-fluorophenyl sulfonamide on the breast tumor cell line BT20, as a function of concentration.

Fig. 2 is a Western blot of BT20 cells and normal human lung fibroblast cells (HFL-1) treated with 20 micromolar (E)-4-methoxystyryl-N-4-fluorophenyl sulfonamide or vehicle (DMSO) and probed with antibody which recognizes both full length and cleaved poly(ADP-ribose)polymerase (PARP). The 83 kDa cleavage product is a marker for apoptosis.

Detailed Description of the Invention

[0079] According to the present invention, N-(aryl)-2-arylethanesulfonamides and pharmaceutically acceptable salts thereof selectively inhibit proliferation of cancer cells, and kill various tumor cell types without killing normal cells. Cells are killed at concentrations where normal cells may be temporarily growth-arrested but not killed.

[0080] The N-(aryl)-2-arylethanesulfonamides compounds of the invention have been shown to inhibit the proliferation of tumor cells, and for some compounds, induce cell death. Cell death results from the induction of apoptosis. The compounds are believed effective against a broad range of tumor types, including but not limited to the following: breast, prostate, ovarian, lung, colorectal, brain (i.e, glioma) and renal. The compounds are also effective against leukemic cells.

[0081] The N-(aryl)-2-arylethanesulfonamides compounds are also believed useful in the treatment of non-cancer proliferative disorders, including but not limited to the following: hemangiomas in new born, secondary progressive multiple sclerosis, chronic progressive myelodegenerative disease, neurofibromatosis, ganglioneuromatosis, keloid for-

mation, Paget's disease of the bone, fibrocystic disease of the breast, Peyronie's fibrosis, Dupuytren's fibrosis, restenosis and cirrhosis.

[0082] The N-(aryl)-2-arylethanesulfonamides also protect normal cells and tissues from the effects of acute and chronic exposure to ionizing radiation.


[0083] Subjects may be exposed to ionizing radiation when undergoing therapeutic irradiation for the treatment of the above proliferative disorders. The N-(aryl)-2-arylethanesulfonamides are effective in protecting normal cells during therapeutic irradiation of abnormal tissues. The compounds are also believed useful in protecting normal cells during radiation treatment for leukemia, especially in the purging of malignant cells from autologous bone marrow grafts with ionizing radiation.

[0084] According to the invention, therapeutic ionizing radiation may be administered to a subject on any schedule and in any dose consistent with the prescribed course of treatment, as long as the N-(aryl)-2-arylethanesulfonamide radioprotectant compound is administered prior to the radiation. The course of treatment differs from subject to subject, and those of ordinary skill in the art can readily determine the appropriate dose and schedule of therapeutic radiation in a given clinical situation.


[0085] In some embodiments of the invention, R in formula I is selected from hydrogen and (C₁-C₆)alkyl, particularly (C₁-C₃)alkyl, and even more particularly ethyl or methyl. In other embodiments, Q₁ is substituted or unsubstituted phenyl, particularly mono-, di- or tri-substituted phenyl and Q₂ is substituted phenyl, particularly mono-, di- or tri-substituted phenyl. In certain embodiments, at least one of Q₁ and Q₂ is at least trisubstituted, at least tetrasubstituted, or even pentasubstituted.

[0086] According to another embodiment of the invention of formula I, Q₁ is optionally substituted phenyl and Q₂ is substituted phenyl. In some embodiments, the substituents are selected from halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, hydroxy and sulfamyl, and for Q₂ can be nitro. In certain sub-embodiments, at least one of Q₁ or Q₂ is substituted in at least the 4-position, or both of Q₁ and Q₂ are substituted at the 4-position. According to certain other sub-embodiments, the substitutions are selected from the group consisting of halogen, (C₁-C₆)alkyl and (C₁-C₆)alkoxy.

[0087] According to another sub-embodiment of the invention, a compound has the formula II:



wherein R₁ is selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, cyano, carboxy, hydroxy, (C₂-C₆)hydroxyalkyl, phosphonato, amino, (C₁-C₆)acylamino, sulfamyl, acetoxy, di(C₁-C₆)alkylamino(C₂-C₆)alkoxy and trifluoromethyl; R is defined as above; and R₂, R₃ and R₄ are independently selected from the group consisting of (C₁-C₆)alkoxy. A preferred pattern of substitution for R₂/R₃/R₄ is 2,4,6, that is, the compound has the formula IIa:



wherein R₁, R₂, R₃ and R₄ are defined as for formula II.


[0088] In some embodiments of formula II and IIa, R₁ is selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, hydroxy and sulfamyl.

[0089] According to another sub-embodiment, Q₁ is optionally substituted phenyl and Q₂ is substituted phenyl, and at least one of Q₁ or Q₂ is at least tetrasubstituted. In other embodiments, at least one of Q₁ and Q₂ is pentasubstituted, e.g. particularly with halogen, most preferably with fluorine.

[0090] Compounds having a carbon-carbon double bond are characterized by cis-trans isomerism. Such compounds are named according to the Cahn-Ingold-Prelog system, the IUPAC 1974 Recommendations, Section E: Stereochemistry, in Nomenclature of Organic Chemistry, John Wiley & Sons, Inc., New York, NY, 4th ed., 1992, p. 127-138. Steric relations around a double bond are designated as "Z" or "E". The compounds of the present invention have the "E" configuration.


[0091] The N-(aryl)-2-arylethenesulfonamides may be prepared by one of two methods. In the synthesis methods to follow, reference to "aryl" is intended to include substituted and unsubstituted aryl, and also substituted and unsubstituted heteroaryl.

[0092] According to Scheme 1, the arylolethene **A**, where Q₂ is substituted aryl, is reacted with sulfonyl chloride in the presence of a nonprotic solvent to form the corresponding arylolethene sulfonyl chloride **B**. Appropriate solvents for this reaction include, for example, dimethylformamide, chloroform and benzene. The arylolethene sulfonyl chloride **B** is then reacted in a nonprotic solvent in the presence of a base with the N-aryl compound **C** wherein Q₁ is substituted or unsubstituted aryl, to obtain the desired N-(aryl)-2-arylethenesulfonamide of formula I. The arylolethene sulfonyl chloride is highly reactive with N-aryl compound **C** and HCl is a byproduct of the reaction. The base is present in the solvent to serve as a scavenger for the produced HCl. The same compound may serve as both the nonprotic solvent and the base. Such dual-function solvents include, for example, pyridine, substituted pyridines, trimethylamine and triethylamine.



25 **[0093]** According to Scheme 2, a Knoevenagel-type condensation according to Oliver et al., Synthesis 321-322 (May 1975) is utilized, relying on the condensation of an arylaminosulfonylacetic acid intermediate **G** with an appropriate aryl aldehyde **H**.

30 **Scheme 2: Synthesis of compounds of formula I.**



[0094] A methyl (or ethyl) β -chlorosulfonylacetate intermediate **E** is prepared from methyl (or ethyl) bromoacetate (R' = methyl or ethyl). To do this, methyl (or ethyl) bromoacetate is reacted with sodium sulfite to form the sodium sulfoacetate intermediate $Na_2OSO_2CH_2CO_2R'$. Potassium sulfite may be used as a substitute for sodium sulfite. The sodium sulfoacetate intermediate is then reacted with a chlorinating agent, preferably PCl_5 , to form the methyl (or ethyl) β -chlorosulfonylacetate intermediate **E**. Reaction of intermediate **E** with the aromatic amine **C** yields the arylaminosulfonylacetate intermediate **F**. The latter reaction is conducted in a nonprotic solvent in the presence of a base. The same compound may serve as both the nonprotic solvent and the base. Such dual-function solvents include, for example, pyridine, substituted pyridines, trimethylamine and triethylamine. The arylaminosulfonylacetate **F** is then converted to the corresponding arylaminosulfonylacetic acid compound **G** by any base capable of hydrolyzing the ester function of **F** to an acid. Such bases include KOH and NaOH, for example. In the final step, the arylaminosulfonylacetic acid compound is condensed with arylaldehyde **H** in the presence of a basic catalyst via a Knoevenagel reaction and decarboxylation of an intermediate. Basic catalysts include, for example, pyridine and benzylamine. The reaction yields the desired N-(aryl)-2-arylethanesulfonamide of formula I.

[0095] The following are more detailed procedures for the preparation of the formula I compounds, according to either Scheme 1 (General Procedure 1) or Scheme 2 (General Procedure 2).

General Procedure 1

A. Synthesis of (E)-Q₂-CH=CH-SO₂Cl.

[0096] To a stirred solution of an arylolefin **A** (0.1mol) in dimethyl formamide (30mL), sulfonyl chloride (0.2mol) is added dropwise for 30 minutes under nitrogen atmosphere. After the addition is complete, the solution is stirred further for 5 hours under nitrogen atmosphere. The reaction mixture is then slowly poured into cold water (250mL) and the precipitated material is extracted with diethyl ether. Evaporation of the dried ethereal layer yields the corresponding sulfonyl chloride (E)-Q₂-CH=CH-SO₂Cl (**B**).

B. Condensation of (E)-Q₂-CH=CH-SO₂Cl with arylamine.

[0097] The N-aryl compound **C** (10 mmol) and sulfonyl chloride **B** (10mmol) are dissolved in 15mL of pyridine under nitrogen. The mixture is stirred for 6 hours at room temperature, and the solvent is removed at aspirator pressure. Water (100mL) is added to the residue and the product is filtered. Recrystallization of the product gives pure N-(aryl)-2-arylethanesulfonamide of formula I.

General Procedure 2

A. Synthesis of sodium ethyl or methyl sulfoacetate

[0098] A solution of ethyl or methyl bromoacetate (0.1 mol) in ethanol (50mL) is added dropwise to a stirred cold solution of sodium sulfite (0.1mol) in water (100mL). After the addition is complete, the mixture is heated briefly to 50°C and then concentrated to dryness. The solid residue is extracted with boiling 2:1 acetic acid/ethyl acetate (200mL) and the hot solution is filtered and chilled overnight. The sodium methyl or ethyl sulfoacetate obtained as a white solid is collected by filtration.

B. Synthesis of ethyl or methyl arylaminosulfonylacetate

[0099] Sodium methyl or ethyl sulfoacetate (0.1 mol) and phosphorus (V) chloride (0.11mol) are separately pulverized and then combined in a flask equipped with a condenser and drying tube. After swirling a few minutes, an exothermic reaction occurs. After the reaction subsides, the flask is warmed on a steam bath for 1 hour and then phosphoryl chloride is removed *in vacuo*. A portion of benzene is added and the resulting solution is filtered and evaporated to yield ethyl or methyl β -chlorosulfonyl acetate (**E**) solution. Benzene (50mL) is added to this clear oil and the solution is stirred and cooled. To this solution, an N-aryl compound **C** and triethylamine (10mL) in benzene (50mL) is added dropwise. After the addition is complete, the mixture is warmed gently for 5 minutes, then is cooled and filtered. The filtrate is washed with water, dilute hydrochloric acid, aqueous sodium hydrogen carbonate and aqueous sodium chloride. After drying the solvent is removed to give crude ethyl or methyl arylaminosulfonylacetate **F**. Recrystallization from benzene yields a pure compound.

C. Synthesis of arylaminosulfonylacetic acid

[0100] The ethyl or methyl arylaminosulfonylacetate **F** (0.1mol) is refluxed for 2.5 hours in a solution of potassium hydroxide (15g) in water (100mL) and ethanol (40mL). Charcoal is added and the solution is heated to boiling for 5 minutes, filtered, acidified with hydrochloric acid and extracted with ether. The ether extract is washed with water, dried and evaporated to give the crude arylaminosulfonylacetic acid **G** which is purified by recrystallization from benzene.


D. Condensation of arylaminosulfonylacetic acid with arylaldehyde

[0101] A solution of the arylaminosulfonylacetic acid **G** (10mmol), arylaldehyde **H** (10mmol), pyridine (1mL) and ammonium acetate (250mg) are refluxed for 22 hours in toluene with azeotropic removal of water. The solution is cooled, washed with water, dilute hydrochloric acid and aqueous sodium hydrogen carbonate and then is extracted with 10% potassium hydroxide. The two-phase aqueous extract is washed with ether and acidified with hydrochloric acid. Evaporation of the ether extract yields crude N-(aryl)-2-arylethenesulfonamide of formula I. Recrystallization from appropriate solvent provides an analytical sample.


N-(aryl)-2-arylethenesulfonamides according to formula I wherein the aryl nucleus of Q_1 is substituted with the group



wherein X is sulfur or oxygen and R_5 and R_6 are defined as above, are prepared according to a variation of the above procedure. Accordingly, an aryl intermediate which is substituted with at least one amino and at least one nitro group, preferably a phenyl intermediate of the formula III




wherein R is defined as above, and wherein the phenyl ring may be further substituted as described above, is reacted with the sulfonyl chloride **B** as in Scheme I to obtain the desired N-(nitroaryl)-2-arylethenesulfonamide IV:



[0102] The nitro group is reduced to an amino group by hydrogenation with a catalyst of palladium on carbon, for example, to form amino intermediate V:


5



10

[0103] Alternatively, the amine intermediate Va is prepared:

15




20

25

wherein R_5 is selected from the group consisting of hydrogen, (C_1-C_6) alkyl, (C_2-C_6) heteroalkyl and substituted or unsubstituted phenyl. The amino group is then made to react with a compound of formula VI


30



35

wherein X is oxygen or sulfur, and R_6 is selected from the group consisting of hydrogen, (C_1-C_6) alkyl, (C_2-C_6) heteroalkyl, substituted aryl, unsubstituted aryl, substituted heteroaryl, unsubstituted heteroaryl, substituted aryl- (C_1-C_3) alkyl, unsubstituted aryl- (C_1-C_3) alkyl and (C_1-C_6) alkoxycarbonyl (C_1-C_6) alkylenyl, to form the urea/thiourea derivative of formula VII:

40



45

50

[0104] The reaction may be carried out, for example, by dissolving the compound of formula VI in deionized water and adding an approximately equimolar amount thereof to the intermediate V or Va dissolved in an appropriate solvent, such as glacial acetic acid. The reaction mixture is stirred at room temperature for three hours. The reaction mixture is then poured into deionized water and extracted 3 times with ethyl acetate. The resulting combined organic layers are

55

washed with saturated NaHCO_3 and saturated brine. The solvent is dried over MgSO_4 , filtered, and concentrated under reduced pressure. The resulting solid is recrystallized from hot ethyl acetate/hexane, for example, to give the sulfonamide VII.

[0105] The compounds of the present invention may take the form of pharmaceutically acceptable salts. The term "pharmaceutically acceptable salts", embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically acceptable acid addition salts may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic and sulfonic classes of organic acids, example of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, 4-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, 2-hydroxyethanesulfonic, toluenesulfonic, sulfanilic, cyclohexylaminosulfonic, stearic, alginic, beta-hydroxybutyric, salicylic, galactaric and galacturonic acid. Suitable pharmaceutically acceptable base addition salts of compounds of formula I include metallic salts made from calcium, magnesium, potassium, sodium and zinc or organic salts made from N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. All of these salts may be prepared by conventional means from the corresponding compound of formula I by reacting, for example, the appropriate acid or base with the compound of formula I.

[0106] The compounds of the invention may be administered to individuals (mammals, including animals and humans) afflicted with cancer.

[0107] The compounds are also useful in the treatment of non-cancer proliferative disorders, that is, proliferative disorders which are characterized by benign indications. Such disorders may also be known as "cytoproliferative" or "hyperproliferative" in that cells are made by the body at an atypically elevated rate. Such disorders include, but are not limited to, the following: hemangiomas in new born, secondary progressive multiple sclerosis, chronic progressive myelodegenerative disease, neurofibromatosis, ganglioneuromatosis, keloid formation, Paget's disease of the bone, fibrocystic disease of the breast, Peyronie's fibrosis, Dupuytren's fibrosis, restenosis and cirrhosis.

[0108] For treating proliferative disorders, the specific dose of compound according to the invention to obtain therapeutic benefit will, of course, be determined by the particular circumstances of the individual patient including, the size, weight, age and sex of the patient, the nature and stage of the disease, the aggressiveness of the disease, and the route of administration. For example, a daily dosage of from about 0.05 to about 50 mg/kg/day may be utilized. Higher or lower doses are also contemplated.

[0109] For radioprotective administration, the specific dose and schedule of N-(aryl)-2-arylethanesulfonamide to obtain the radioprotective benefit will, of course, be determined by the particular circumstances of the individual patient including, the size, weight, age and sex of the patient, the nature and stage of the disease being treated, the aggressiveness of the disease, and the route of administration, and the specific toxicity of the radiation. For example, a daily dosage of from about 0.01 to about 150 mg/kg/day may be utilized, more preferably from about 0.05 to about 50 mg/kg/day. Particularly preferred are doses from about 1.0 to about 10.0 mg/kg/day, for example, a dose of about 7.0 mg/kg/day. The dose may be given over multiple administrations, for example, two administrations of 3.5 mg/kg. Higher or lower doses are also contemplated.

[0110] For radioprotective administration, the N-(aryl)-2-arylethanesulfonamide should be administered far enough in advance of the therapeutic radiation such that the compound is able to reach the normal cells of the subject in sufficient concentration to exert a radioprotective effect on the normal cells. The compound may be administered as much as about 24 hours, preferably no more than about 18 hours, prior to administration of the radiation. In one embodiment, the N-(aryl)-2-arylethanesulfonamide is administered at least about 6-12 hours before administration of the therapeutic radiation. Most preferably, the compound is administered once at about 18 hours and again at about 6 hours before the radiation exposure. One or more N-(aryl)-2-arylethanesulfonamides may be administered simultaneously, or different N-(aryl)-2-arylethanesulfonamides may be administered at different times during the treatment.

[0111] Where the therapeutic radiation is administered in serial fashion, it is preferable to intercalate the administration of one or more N-(aryl)-2-arylethanesulfonamides within the schedule of radiation treatments. As above, different N-(aryl)-2-arylethanesulfonamides may be administered either simultaneously or at different times during the treatment. Preferably, an about 24 hour period separates administration of the radioprotective compound and the therapeutic radiation. More preferably, the administration of the radioprotective N-(aryl)-2-arylethanesulfonamide and the therapeutic radiation is separated by about 6 to 18 hours. This strategy will yield significant reduction of radiation-induced side effects without affecting the anticancer activity of the therapeutic radiation.

[0112] For example, therapeutic radiation at a dose of 0.1 Gy may be given daily for five consecutive days, with a two day rest, for a total period of 6 - 8 weeks. One or more N-(aryl)-2-arylethanesulfonamides may be administered to the subject 18 hours previous to each round of radiation. It should be pointed out, however, that more aggressive treatment schedules, *i.e.*, delivery of a higher dosage, is contemplated according to the present invention due to the protection of

the normal cells afforded by the N-(aryl)-2-arylethanesulfonamides. Thus, the radioprotective effect of the compound increases the therapeutic index of the therapeutic radiation, and may permit the physician to safely increase the dosage of therapeutic radiation above presently recommended levels without risking increased damage to the surrounding normal cells and tissues.

5 [0113] The N-(aryl)-2-arylethanesulfonamides of the invention are further useful in protecting normal bone marrow cells from radiologic treatments designed to destroy hematologic neoplastic cells or tumor cells which have metastasized into the bone marrow. Such cells include, for example, myeloid leukemia cells. The appearance of these cells in the bone marrow and elsewhere in the body is associated with various disease conditions, such as the French-American-British (FAB) subtypes of acute myelogenous leukemias (AML), chronic myeloid leukemia (CML), and acute lymphocytic leukemia (ALL). CML, in particular, is characterized by abnormal proliferation of immature granulocytes (e.g., neutrophils, eosinophils, and basophils) in the blood, bone marrow, spleen, liver, and other tissues and accumulation of granulocytic precursors in these tissues. The subject who presents with such symptoms will typically have more than 20,000 white blood cells per microliter of blood, and the count may exceed 400,000. Virtually all CML patients will develop "blast crisis", the terminal stage of the disease during which immature blast cells rapidly proliferate, leading to death.

10 [0114] Other subjects suffer from metastatic tumors, and require treatment with total body irradiation (TBI). Because TBI will also kill the subject's hematopoietic cells, a portion of the subject's bone marrow is removed prior to irradiation for subsequent reimplantation. However, metastatic tumor cells are likely present in the bone marrow, and reimplantation often results in a relapse of the cancer within a short time.

15 [0115] Subjects presenting with neoplastic diseases of the bone marrow or metastatic tumors may be treated by removing a portion of the bone marrow (also called "harvesting"), purging the harvested bone marrow of malignant stem cells, and reimplanting the purged bone marrow. Preferably, the subject is treated with radiation or some other anti-cancer therapy before the autologous purged bone marrow is reimplanted.

20 [0116] Thus, the compounds of the invention can be used to reduce the number of malignant cells in bone marrow, wherein a portion of the subject's (previously removed) bone marrow, has an effective amount of at least one N-(aryl)-2-arylethanesulfonamide administered thereto and the treated bone marrow is irradiated with a sufficient dose of ionizing radiation such that malignant cells in the bone marrow are killed. As used herein, "malignant cell" means any uncontrollably proliferating cell, such as a tumor cell or neoplastic cell. The N-(aryl)-2-arylethanesulfonamides protect the normal hematopoietic cells present in the bone marrow from the deleterious effects of the ionizing radiation. The N-(aryl)-2-arylethanesulfonamides also exhibit a direct killing effect on the malignant cells. The number of malignant cells in the bone marrow is significantly reduced prior to reimplantation, thus minimizing the occurrence of a relapse.

25 [0117] Preferably, each N-(aryl)-2-arylethanesulfonamide is administered in a concentration from about 0.25 to about 100 micromolar; more preferably, from about 1.0 to about 50 micromolar; in particular from about 2.0 to about 25 micromolar. Particularly preferred concentrations are 0.5, 1.0 and 2.5 micromolar and 5, 10 and 20 micromolar. Higher or lower concentrations may also be used.

30 [0118] The N-(aryl)-2-arylethanesulfonamides may be added directly to the harvested bone marrow, but are preferably dissolved in an organic solvent such as dimethylsulfoxide (DMSO). Pharmaceutical formulations of N-(aryl)-2-arylethanesulfonamides such as are described in more detail below may also be used.

35 [0119] Preferably, the N-(aryl)-2-arylethanesulfonamide is added to the harvested bone marrow about 20 hours prior to radiation exposure, preferably no more than about 24 hours prior to radiation exposure. In one embodiment, the N-(aryl)-2-arylethanesulfonamide is administered to the harvested bone marrow at least about 6 hours before radiation exposure. One or more N-(aryl)-2-arylethanesulfonamides may be administered simultaneously, or different N-(aryl)-2-arylethanesulfonamides may be administered at different times. Other dosage regimens are also contemplated.

40 [0120] If the subject is to be treated with ionizing radiation prior to reimplantation of the purged bone marrow, the subject may be treated with one or more N-(aryl)-2-arylethanesulfonamides prior to receiving the ionizing radiation dose, as described above.

45 [0121] A subject may also be exposed to ionizing radiation from occupation or environmental sources, as discussed in the background section. For purposes of the invention, the source of the radiation is not as important as the type (i.e., acute or chronic) and dose level absorbed by the subject. It is understood that the following discussion encompasses ionizing radiation exposures from both occupational and environmental sources.

50 [0122] Subjects suffering from effects of acute or chronic exposure to ionizing radiation that are not immediately fatal are said to have remediable radiation damage. Such remediable radiation damage can be reduced or eliminated by the compounds and methods of the present invention.

55 [0123] An acute dose of ionizing radiation which may cause remediable radiation damage includes a localized or whole body dose, for example, between about 10,000 millirem (0.1 Gy) and about 1,000,000 millirem (10 Gy) in 24 hours or less, preferably between about 25,000 millirem (0.25 Gy) and about 200,000 (2 Gy) in 24 hours or less, and more preferably between about 100,000 millirem (1 Gy) and about 150,000 millirem (1.5 Gy) in 24 hours or less.

[0124] A chronic dose of ionizing radiation which may cause remediable radiation damage includes a whole body dose of about 100 millirem (.001 Gy) to about 10,000 millirem (0.1 Gy), preferably a dose between about 1000 millirem

(.01 Gy) and about 5000 millirem (.05 Gy) over a period greater than 24 hours, or a localized dose of 15,000 millirem (0.15 Gy) to 50,000 millirem (0.5 Gy) over a period greater than 24 hours.

5 [0125] The invention therefore provides the use of at least one N-(aryl)-2-arylethanesulfonamide compound for the preparation of a medicament for treating individuals who have incurred remediable radiation damage from acute or chronic exposure to ionizing radiation, comprising reducing or eliminating the cytotoxic effects of radiation exposure on normal cells and tissues. The compound is intended to be administered in as short a time as possible following radiation exposure, for example between 0 - 6 hours following exposure.

10 [0126] Remediable radiation damage may take the form of cytotoxic and genotoxic (*i.e.*, adverse genetic) effects in the subject. The compounds of the present invention are also useful for reducing or eliminating the cytotoxic and genotoxic effects of radiation exposure on normal cells and tissues, wherein an effective amount of at least one radioprotective N-(aryl)-2-arylethanesulfonamide compound is administered prior to acute or chronic radiation exposure. The N-(aryl)-2-arylethanesulfonamide may be administered, for example about 24 hours prior to radiation exposure, preferably no more than about 18 hours prior to radiation exposure. In one embodiment, the N-(aryl)-2-arylethanesulfonamide is administered at least about 6 hours before radiation exposure. Most preferably, the N-(aryl)-2-arylethanesulfonamide is administered at about 18 and again at about 6 hours before the radiation exposure. One or more N-(aryl)-2-arylethanesulfonamides may be administered simultaneously, or different N-(aryl)-2-arylethanesulfonamides may be administered at different times.

15 [0127] When multiple acute exposures are anticipated, the radioprotective may be administered multiple times. For example, if fire or rescue personnel must enter contaminated areas multiple times, N-(aryl)-2-arylethanesulfonamides may be administered prior to each exposure. Preferably, an about 24 hour period separates administration of the compound and the radiation exposure. More preferably, the administration of N-(aryl)-2-arylethanesulfonamide and the radiation exposure is separated by about 6 to 18 hours. It is also contemplated that a worker in a nuclear power plant may be administered an effective amount of N-(aryl)-2-arylethanesulfonamide prior to beginning each shift, to reduce or eliminate the effects of exposure to ionizing radiation.

20 [0128] If a subject is anticipating chronic exposure to ionizing radiation, the N-(aryl)-2-arylethanesulfonamide may be administered periodically throughout the duration of anticipated exposure. For example, a nuclear power plant worker or a soldier operating in a forward area contaminated with radioactive fallout may be given the radioprotective compound every 24 hours, preferably every 6-18 hours, in order to mitigate the effects of radiation damage. Likewise, N-(aryl)-2-arylethanesulfonamide compound may be periodically administered to civilians living in areas contaminated by radioactive fallout until the area is decontaminated or the civilians are removed to a safer environment.

25 [0129] As used herein, "administered" means the act of making the N-(aryl)-2-arylethanesulfonamide compound available to the subject such that a pharmacologic effect is obtained. For administration of drug for radioprotection, the pharmacologic effect may manifest as the absence of expected physiologic or clinical symptoms at a certain level of radiation exposure. One skilled in the art may readily determine the presence or absence of radiation-induced effects, by well-known laboratory and clinical methods. The N-(aryl)-2-arylethanesulfonamide compound may thus be administered by any route which is sufficient to bring about the desired radioprotective effect in the patient.

30 [0130] The N-(aryl)-2-arylethanesulfonamide compounds may be administered for therapeutic effect by any route, for example enteral (*e.g.*, oral, rectal, intranasal, etc.) and parenteral administration. Parenteral administration includes, for example, intravenous, intramuscular, intraarterial, intraperitoneal, intravaginal, intravesical (*e.g.*, into the bladder), intradermal, topical or subcutaneous administration. Also contemplated within the scope of the invention is the instillation of drug in the body of the patient in a controlled formulation, with systemic or local release of the drug to occur at a later time. For example, a depot of N-(aryl)-2-arylethanesulfonamide may be administered to the patient more than 24 hours before the administration of radiation. Preferably, at least a portion of the compound is retained in the depot and not released until an about 6-18 hour window prior to the radiation exposure. For anticancer use, the drug may similarly be localized in a depot for controlled release to the circulation, or local site of tumor growth.

35 [0131] The compounds of the invention may be administered in the form of a pharmaceutical composition, in combination with a pharmaceutically acceptable carrier. The active ingredient in such formulations may comprise from 0.1 to 99.99 weight percent. By "pharmaceutically acceptable carrier" is meant any carrier, diluent or excipient which is compatible with the other ingredients of the formulation and not deleterious to the recipient.

40 [0132] The active agent is preferably administered with a pharmaceutically acceptable carrier selected on the basis of the selected route of administration and standard pharmaceutical practice. The active agent may be formulated into dosage forms according to standard practices in the field of pharmaceutical preparations. See Alphonso Gennaro, ed., Remington's Pharmaceutical Sciences, 18th Ed., (1990) Mack Publishing Co., Easton, PA. Suitable dosage forms may comprise, for example, tablets, capsules, solutions, parenteral solutions, troches, suppositories, or suspensions.

45 [0133] For parenteral administration, the active agent may be mixed with a suitable carrier or diluent such as water, an oil (particularly a vegetable oil), ethanol, saline solution, aqueous dextrose (glucose) and related sugar solutions, glycerol, or a glycol such as propylene glycol or polyethylene glycol. Solutions for parenteral administration preferably contain a water-soluble salt of the active agent. Stabilizing agents, antioxidizing agents and preservatives may also be

added. Suitable antioxidizing agents include sulfite, ascorbic acid, citric acid and its salts, and sodium EDTA. Suitable preservatives include benzalkonium chloride, methyl- or propyl-paraben, and chlorbutanol. The composition for parenteral administration may take the form of an aqueous or nonaqueous solution, dispersion, suspension or emulsion.

[0134] For oral administration, the active agent may be combined with one or more solid inactive ingredients for the preparation of tablets, capsules, pills, powders, granules or other suitable oral dosage forms. For example, the active agent may be combined with at least one excipient such as fillers, binders, humectants, disintegrating agents, solution retarders, absorption accelerators, wetting agents, absorbents or lubricating agents. According to one tablet embodiment, the active agent may be combined with carboxymethylcellulose calcium, magnesium stearate, mannitol and starch, and then formed into tablets by conventional tableting methods.

[0135] The practice of the invention is illustrated by the following non-limiting examples. In each of Examples 1-17, the starting (E)-Q₂-CH=CH-SO₂Cl compound was made according to part A of General Procedure 1, above. The synthesized compounds are tabulated in Table 4.




Table 4

Example #	X	Y	R
1*	4-Cl	H	H
2	3-F, 4-OCH ₃	4-Cl	H
3	4-F	4-Cl	H
4	4-F	4-OCH ₃	H
5	3-F, 4-OCH ₃	4-OCH ₃	H
6*	4-F	H	H
7	H	4-OCH ₃	CH ₃
8	3-Cl	4-Cl	H
9	2-Cl	4-Cl	H
10	4-F	4-OCH ₃	H
11	4-Cl	4-F	H
12	2,4,6-(OCH ₃) ₃	4-OCH ₃	H
13	2,3,4,5,6-F ₅	4-OCH ₃	H
14*	2,3,4,5,6-F ₅	H	H
15	2,3,4,5,6-F ₅	4-F	H
16*	4-SO ₂ NH ₂	H	H
17	4-SO ₂ NH ₂	4-OCH ₃	H
* Comparative			

Example 1: (E)-styryl-N-4-chlorophenyl sulfonamide - Comparative

[0136] A solution of (E)-styrylsulfonyl chloride (10mmol) and 4-chloroaniline (10mmol) was subjected to General Pro-

cedure 1, part B. The title compound, melting point 107-109°C was obtained in 56% yield.

Example 2: (E)-4-chlorostyryl-N-3-fluoro-4-methoxyphenyl sulfonamide

5 [0137] A solution of (E)-4-chlorostyrylsulfonyl chloride (10mmol) and 3-fluoro-4-methoxyaniline (10mmol) was subjected to General Procedure 1, part B. The title compound, melting point 101-102°C was obtained in 58% yield.

Example 3: (E)-4-chlorostyryl-N-4-fluorophenyl sulfonamide

10 [0138] A solution of (E)-4-chlorostyrylsulfonyl chloride (10mmol) and 4-fluoroaniline (10mmol) was subjected to General Procedure 1, part B. The title compound, melting point 105-107°C, was obtained in 68.5% yield.

Example 4: (E)-4-methoxystyryl-N-4-fluorophenyl sulfonamide,

15 [0139] A solution of (E)-4-methoxystyrylsulfonyl chloride (10mmol) and 4-fluoroaniline (10mmol) was subjected to General Procedure 1, part B. The title compound, melting point 115-117°C, was obtained 86.4% yield.

Example 5: (E)-4-methoxystyryl-N-3-fluoro-4-methoxyphenyl sulfonamide

20 [0140] A solution of (E)-4-methoxystyrylsulfonyl chloride (10mmol) and 3-fluoro-4-methoxyaniline (10mmol) was subjected to General Procedure 1, part B. The title compound, melting point 151-153°C, was obtained 80% yield.

Example 6: (E)-styryl-N-4-fluorophenyl sulfonamide - Comparative

25 [0141] A solution of (E)-styrylsulfonyl chloride (10mmol) and 4-fluoroaniline (10mmol) was subjected to General Procedure 1, part B. The title compound, melting point 83-85°C, was obtained 58.7% yield.

Example 7: (E)-4-methoxystyryl-N-methyl-N-phenyl sulfonamide

30 [0142] A solution of (E)-4-methoxystyrylsulfonyl chloride (10mmol) and N-methyl aniline (10mmol) was subjected to General Procedure 1, part B. The title compound, melting point 126-129°C, was obtained 80.9% yield.

Example 8: (E)-4-chlorostyryl-N-3-chlorophenyl sulfonamide

35 [0143] A solution of (E)-4-chlorostyrylsulfonyl chloride (10mmol) and 3-chloroaniline (10mmol) was subjected to General Procedure 1, part B. The title compound, melting point 118-120°C, was obtained in 31.8% yield.

Example 9: (E)-4-chlorostyryl-N-2-chlorophenyl sulfonamide

40 [0144] A solution of (E)-4-chlorostyrylsulfonyl chloride (10mmol) and 2-chloroaniline (10mmol) was subjected to General Procedure 1, part B. The title compound, melting point 107-109°C, was obtained in 57.8% yield.

Example 10: (E)-4-methoxystyryl-N-4-fluorophenyl sulfonamide.

45 [0145] A solution of (E)-4-methoxystyrylsulfonyl chloride (10mmol) and 4-fluoroaniline (10mmol) was subjected to General Procedure 1, part B. The title compound was obtained in 60.3% yield.

Example 11: (E)-4-fluorostyryl-N-4-chlorophenyl sulfonamide.

50 [0146] A solution of (E)-4-fluorostyrylsulfonyl chloride (10mmol) and 4-chloroaniline (10mmol) was subjected to General Procedure 1, part B. The title compound, melting point 126-128°C; was obtained in 81.7% yield.

Example 12: (E)-4-methoxystyryl-N-2,4,6-trimethoxyphenyl sulfonamide.

55 [0147] A solution of (E)-4-methoxystyrylsulfonyl chloride (10mmol) and 2,4,6-trimethoxyaniline (10mmol) was subjected to General Procedure 1, part B. The title compound, melting point 103-106°C, was obtained in 78.7% yield.

Example 13: (E)-4-methoxystyryl-N-2,3,4,5,6-pentafluorophenyl sulfonamide.

[0148] A solution of (E)-4-methoxystyrylsulfonyl chloride (10mmol) and 2,3,4,5,6-pentafluoroaniline (10mmol) was subjected to General Procedure 1, part B. The title compound, melting point 58-60°C, was obtained in 41% yield.

Example 14: (E)-styryl-N-2,3,4,5,6-pentafluorophenyl sulfonamide - Comparative

[0149] A solution of (E)-styrylsulfonyl chloride (10mmol) and 2,3,4,5,6-pentafluoroaniline (10mmol) was subjected to General Procedure 1, part B. The title compound, melting point 145-148°C, was obtained in 34.8% yield.

Example 15: (E)-4-fluorostyryl-N-2,3,4,5,6-pentafluorophenyl sulfonamide.

[0150] A solution of (E)-4-fluorostyrylsulfonyl chloride (10mmol) and 2,3,4,5,6-pentafluoroaniline (10mmol) was subjected to General Procedure 1, part B. The title compound, melting point 182-184°C, was obtained in 36.1% yield.


Example 16: (E)-styryl-N-4-sulfamylphenyl sulfonamide. - Comparative

[0151] A solution of (E)-styrylsulfonyl chloride (10mmol) and sulfanilamide (10mmol) was subjected to General Procedure 1, part B. The title compound, melting point 171-173°C, was obtained in 80% yield.

Example 17: (E)-4-methoxystyryl-N-4-sulfamylphenyl sulfonamide.

[0152] A solution of (E)-4-methoxystyrylsulfonyl chloride (10mmol) and sulfanilamide (10mmol) was subjected to General Procedure 1, part B. The title compound, melting point 181-183°C, was obtained in 46.2% yield.

[0153] The following additional compounds, tabulated in Tables 5 and 6, are made by subjecting a solution containing a 10 mmolar concentration of the two indicated reactants to either General Procedure 1, part B (Examples Nos. 18, 19, 21-23, 26, 27, 30, 33, 36, 39, 41, 44, 47-50, 55, 59, 61, 64 and 65) or General Procedure 2, part D (Examples Nos. 20, 24, 25, 28, 29, 31, 32, 34, 35, 37, 38, 40, 42, 43, 45, 46, 51-54, 56-58, 60, 62, 63, 66 and 67).

**Table 5**

Ex.	X	Y	Reactants	Product
18	2-Cl	2,3,4,5,6-F ₅	pentafluorostyrylsulfonyl chloride and 2-chloroaniline	(E)-pentafluorostyryl-N-2-chlorophenyl sulfonamide
19	4-F	2,3,4,5,6-F ₅	pentafluorostyrylsulfonyl chloride and 4-fluoroaniline sulfonamide	(E)-pentafluorostyryl-N-4-fluorophenyl sulfonamide
20	4-Br	2,3,4,5,6-F ₅	4-bromophenylaminosulfonylacetic acid and pentafluorobenzaldehyde	(E)-pentafluorostyryl-N-4-bromophenyl sulfonamide
21	2-F,4-Cl	2,3,4,5,6-F ₅	pentafluorostyrylsulfonyl chloride and 2-fluoro-4-chloroaniline	(E)-pentafluorostyryl-N-2-fluoro-4-chlorophenyl sulfonamide

EP 1 379 508 B9

(continued)

Ex.	X	Y	Reactants	Product
5 22	4-OCH ₃	2,3,4,5,6-F ₅	pentafluorostyrylsulfonyl chloride and 4-methoxyaniline	(E)-pentafluorostyryl-N-4-methoxyphenyl sulfonamide
23	3-F,4-OCH ₃	2,3,4,5,6-F ₅	pentafluorostyrylsulfonyl chloride and 3-fluoro-4-methoxyaniline	(E)-pentafluorostyryl-N-3-fluoro-4-methoxyphenyl sulfonamide
10 24	2,3,4-(OCH ₃) ₃	2,3,4,5,6-F ₅	2,3,4-trimethoxyphenylaminosulfonylacetic acid and pentafluorobenzaldehyde	(E)-pentafluorostyryl-N-2,3,4-trimethoxyphenyl sulfonamide
15 25	4-OH	2,3,4,5,6-F ₅	4-hydroxyphenylaminosulfonylacetic acid and pentafluorobenzaldehyde	(E)-pentafluorostyryl-N-4-hydroxyphenyl sulfonamide
26	4-NO ₂	2,3,4,5,6-F ₅	pentafluorostyrylsulfonyl chloride and 4-nitroaniline	(E)-pentafluorostyryl-N-4-nitrophenyl sulfonamide
20 27	4-SO ₂ NH ₂	2,3,4,5,6-F ₅	pentafluorostyrylsulfonyl chloride and 4-sulfanilamide	(E)-pentafluorostyryl-N-4-sulfamylphenyl sulfonamide
28	3-F,4-OCH ₃	2,3,4,5,6-F ₅	3-fluoro-4-methoxyphenylaminosulfonylacetic acid and pentafluorobenzaldehyde	(E)-pentafluorostyryl-N-3-fluoro-4-methoxyphenyl sulfonamide
25 29	2,4,6-(OCH ₃) ₃	2,3,4,5,6-F ₅	2,4,6-trimethoxyphenylaminosulfonylacetic acid and pentafluorobenzaldehyde	(E)-pentafluorostyryl-N-2,4,6-trimethoxyphenyl sulfonamide
30 30	2,3,4,5,6-F ₅	2,3,4,5,6-F ₅	pentafluorostyrylsulfonyl chloride and pentafluoroaniline	(E)-pentafluorostyryl-N-pentafluorophenyl sulfonamide
31	2-Cl	2,4,6-(OCH ₃) ₃	2-chlorophenylaminosulfonylacetic acid and 2,4,6-trimethoxybenzaldehyde	(E)-2,4,6-trimethoxystyryl-N-2-chlorophenyl sulfonamide
35 32	4-F	2,4,6-(OCH ₃) ₃	4-bromophenylaminosulfonylacetic acid and 2,4,6-trimethoxybenzaldehyde	(E)-2,4,6-trimethoxystyryl-N-4-fluorophenyl sulfonamide
33	4-Br	2,4,6-(OCH ₃) ₃	2,4,6-trimethoxystyrylsulfonyl chloride and 4-bromoaniline	(E)-2,4,6-trimethoxystyryl-N-4-bromophenyl sulfonamide
40 34	2-F,4-Cl	2,4,6-(OCH ₃) ₃	2-fluoro-4-chlorophenylaminosulfonylacetic acid and 2,4,6-trimethoxybenzaldehyde	(E)-2,4,6-trimethoxystyryl-N-2-fluoro-4-chlorophenyl sulfonamide
45 35	4-OCH ₃	2,4,6-(OCH ₃) ₃	4-methoxyphenylaminosulfonylacetic acid and 2,4,6-trimethoxybenzaldehyde	(E)-2,4,6-trimethoxystyryl-N-4-methoxyphenyl sulfonamide
36	3-F,4-OCH ₃	2,4,6-(OCH ₃) ₃	2,4,6-trimethoxystyrylsulfonyl chloride and 3-fluoro-4-methoxyaniline	(E)-2,4,6-trimethoxystyryl-N-3-fluoro-4-methoxyphenyl sulfonamide
50 37	2,3,4-(OCH ₃) ₃	2,4,6-(OCH ₃) ₃	2,3,4-trimethoxyphenylaminosulfonylacetic acid and 2,4,6-trimethoxybenzaldehyde	(E)-2,4,6-trimethoxystyryl-N-2,3,4-trimethoxyphenyl sulfonamide
55 38	4-OH	2,4,6-(OCH ₃) ₃	4-hydroxyphenylaminosulfonylacetic acid and 2,4,6-trimethoxybenzaldehyde	(E)-2,4,6-trimethoxystyryl-N-4-hydroxyphenyl sulfonamide

(continued)

Ex.	X	Y	Reactants	Product
39	4-H ₂ PO ₄	2,4,6-(OCH ₃) ₃	2,4,6-trimethoxystyrylsulfonyl chloride and 4-phosphonatoaniline	(E)-2,4,6-trimethoxystyryl-N-4-phosphonatophenyl sulfonamide
40	4-NO ₂	2,4,6-(OCH ₃) ₃	4-nitrophenylaminosulfonylacetic acid and 2,4,6-trimethoxybenzaldehyde	(E)-2,4,6-trimethoxystyryl-N-4-nitrophenyl sulfonamide
41	2,4,6-(OCH ₃) ₃	2,4,6-(OCH ₃) ₃	2,4,6-trimethoxystyrylsulfonyl chloride and 2,4,6-trimethoxyaniline	(E)-2,4,6-trimethoxystyryl-N-2,4,6-trimethoxyphenyl sulfonamide
42	2,3,4,5,6-F ₅	2,4,6-(OCH ₃) ₃	2,3,4,5,6-pentafluorophenylaminosulfonylacetic acid and 2,4,6-trimethoxybenzaldehyde	(E)-2,4,6-trimethoxystyryl-N-pentafluorophenyl sulfonamide
43	4-SO ₂ NH ₂	2,4,6-(OCH ₃) ₃	4-sulfamylphenylaminosulfonylacetic acid and 2,4,6-trimethoxybenzaldehyde	(E)-2,4,6-trimethoxystyryl-N-4-sulfamylphenyl sulfonamide
44	2,3,4,5,6-F ₅	2,4,6-(OCH ₃) ₃	2,4,6-trimethoxystyrylsulfonyl chloride and pentafluoroaniline	(E)-2,4,6-trimethoxystyryl-N-pentafluoro sulfonamide
45	4-OCH ₃	3,4,5-(OCH ₃) ₃	4-methoxyphenylaminosulfonylacetic acid and 3,4,5-trimethoxybenzaldehyde	(E)-3,4,5-trimethoxystyryl-N-4-methoxyphenyl sulfonamide
46	3-F,4-OCH ₃	3,4,5-(OCH ₃) ₃	3-fluoro-4-methoxyphenylaminosulfonylacetic acid and 3,4,5-trimethoxybenzaldehyde	(E)-3,4,5-trimethoxystyryl-N-3-fluoro-4-methoxyphenyl sulfonamide
47	2,3,4-(OCH ₃) ₃	3,4,5-(OCH ₃) ₃	3,4,5-trimethoxystyrylsulfonyl chloride and 2,3,4-trimethoxyaniline	(E)-3,4,5-trimethoxystyryl-N-2,3,4-trimethoxyphenyl sulfonamide




Table 6

Ex.	Q ₁	Q ₂	Reactants	Product
48	2-benzothiazolyl	pentafluorophenyl	pentafluorostyryl-sulfonyl chloride and 2-aminobenzathiazole	(E)-pentafluorostyryl-N-2-benzothiazolylsulfonamide
49	2-benzoxazolyl	2,4,6-trimethoxyphenyl	2,4,6-trimethoxystyrylsulfonyl chloride and 2-aminobenzoxazole	(E)-2,4,6-trimethoxystyryl-N-2-benzoxazolylsulfonamide
50	2-thiazolyl	4-methoxyphenyl	4-methoxystyrylsulfonyl chloride and 2-aminothiazole	(E)-4-methoxystyryl-N-2-thiazolylsulfonamide
51	3-indolyl	pentafluorophenyl	3-indolylaminosulfonylacetic acid and pentafluorobenzaldehyde	(E)-pentafluorostyryl-N-3-indolyl sulfonamide
52	4-indolyl	2,4,6-trimethoxyphenyl	4-indolylaminosulfonylacetic acid and 2,4,6-trimethoxybenzaldehyde	(E)-2,4,6-trimethoxystyryl-N-4-indolyl sulfonamide
53	5-indolyl	3-fluoro-4-methoxyphenyl	5-indolylaminosulfonylacetic acid and 3-fluoro-4-methoxybenzaldehyde	(E)-3-fluoro-4-methoxystyryl-N-5-indolyl sulfonamide
54	3-isoxazolyl	3-indolyl	3-isoxazolylaminosulfonylacetic acid and 3-indolylcarbaldehyde	(E)-N-3-isoxazolyl-3-indolyl ethenesulfonamide
55	3-quinolinyl	4-pyridinyl	3-Aminoquinoline and 4-pyridine ethenyl sulfonylchloride	(E)-N-3-quinolinyl-4-pyridyl ethenesulfonamide
56	3-(1,2,3-triazolyl)	pentafluorophenyl	3-(1,2,3-triazolyl)aminosulfonylacetic acid and pentafluorobenzaldehyde	(E)-N-1,2,3-triazolyl-2,3,4,5,6-pentafluorostyrylsulfonamide
57	2,4,6-trimethoxyphenyl	5-indolyl	2,4,6-trimethoxyphenylaminosulfonyl-acetic acid and 5-indolylcarboxaldehyde	(E)-N-2,4,6-trimethoxyphenyl-5-indolyl ethenesulfonamide
58	2-imidazolyl	pentafluorophenyl	2-imidazolylaminosulfonylacetic acid and pentafluorobenzaldehyde	(E)-N-2-imidazolyl-2,3,4,5,6-pentafluorostyrylsulfonamide
59	5-isothiazolyl	2-pyridinyl	5-aminoisothiazole and 2-pyridene ethenyl sulfonylchloride	(E)-N-5-thiazolyl-2-pyridyl ethenesulfonamide
60	3-pyrazolyl	2,4,6-trimethoxyphenyl	3-pyrazolylaminosulfonylacetic acid and 2,4,6-trimethoxybenzaldehyde	(E)-N-3-pyrazolyl-2,4,6-trimethoxystyryl sulfonamide
61	4-cyano-3-pyrazolyl	pentafluorophenyl	4-cyano-3-pyrazolylaminosulfonylacetic acid and pentafluorobenzaldehyde	(E)-N-4-cyano-3-pyrazolyl-pentafluorostyrylsulfonamide

(continued)

Ex.	Q ₁	Q ₂	Reactants	Product
62	4-methoxyphenyl	3-quinolinyl	4-methoxyphenylaminosulfonylacetic acid and 3-quinolinylcarboxaldehyde	(E)-N-4-methoxyphenyl-3-quinolinyl ethenesulfonamide
63	3-fluoro-4-methoxyphenyl	5-indolyl	3-fluoro-4-methoxyphenylaminosulfonyl-acetic acid and 5-indolylcarboxaldehyde	(E)-N-3-fluoro-4-methoxyphenyl-5-indolyl ethenesulfonamide
64	2,3,4-trimethoxyphenyl	3-pyrazolyl	2,3,4-trimethoxyphenylaminosulfonyl-acetic acid and 3-pyrazolecarboxaldehyde	(E)-N-2,3,4-trimethoxyphenyl-3-pyrazolyl ethenesulfonamide
65	4-hydroxyphenyl	4-pyridinyl	4-hydroxyaniline and 4-pyridine ethenyl sulfonylchloride	(E)-N-4-hydroxyphenyl-4-pyridyl ethenesulfonamide
66	pentafluorophenyl	4-quinolinyl	pentafluorophenylaminosulfonylacetic acid and 4-quinolinylcarboxaldehyde	(E)-N-pentafluorophenyl-4-quinolinyl ethenesulfonamide
67	2,4,6-trimethoxyphenyl	2-imidazolyl	2,4,6-trimethoxyphenylaminosulfonyl-acetic acid and 2-imidazolylcarboxaldehyde	(E)-N-2,4,6-trimethoxyphenyl-2-imidazolyl ethenesulfonamide

Example 68: Effect of N-(Aryl)-2-Arylethanesulfonamides on Tumor Cell Lines.

[0154] The effect of the N-(aryl)-2-arylethanesulfonamides on normal fibroblasts and on tumor cells was determined by the assay described by Latham et al., *Oncogene* 12:827-837 (1996). Normal diploid lung human fibroblasts (HFL-1) or tumor cells (prostate, colorectal, breast, glial, pancreatic ovarian or leukemic) were plated in 6-well dishes at a cell density of 1.0×10^5 cells per 35-mm² well. The plated cells were treated 24 hours later with various concentrations of N-(aryl)-2-arylethanesulfonamide dissolved in dimethyl sulfoxide (DMSO). The total number of viable cells was determined 96 hours later by trypsinizing the wells and counting the number of viable cells, as determined by trypan blue exclusion, using a hemacytometer. Each compound tested (Exs. 1-17) inhibited cell proliferation when tested at a concentration of 30 micromolar. Some compounds inhibited proliferation at lower concentrations. Certain compounds induced cell death, at concentrations from 1 to 10 micromolar. Normal HFL cells were treated with the same compounds under the same conditions of concentration and time. The normal cells displayed growth inhibition but no appreciable cell death.

Example 69: Determination of GI₅₀.

[0155] A dose response curve was plotted for the tumor growth inhibitory effect of (E)-4-methoxystyryl-N-4-fluorophenyl sulfonamide on the estrogen-unresponsive breast cell line BT-20 (Fig. 1). The GI₅₀ (the concentration of drug resulting in 50% net loss of growth inhibition) was determined as 10 μ M. In contrast, the normal fibroblast line HFL-1 exhibited a GI₅₀ of 50 μ M. The (E)-4-methoxystyryl-N-4-fluorophenyl sulfonamide GI₅₀ for the same compound against the cell lines listed in Table 7 was determined in the same manner.

Table 7: GI₅₀ for growth inhibition effect of (E)-4-methoxystyryl-N-4-fluorophenyl sulfonamide on various cell lines

CELL LINE	TUMOR TYPE	GI ₅₀ (μ M)
DU145	Prostate	10
PC-3	Prostate	20
LNCAP	Prostate	20
DLD-1	Colo-rectal	15
HCT-116	Colo-rectal	20
COLO-320	Colo-rectal	5
BT20	Breast	10
SK-br-3	Breast	30
U87	Glioblastoma	20
MIA-PaCa-2	Pancreatic	7
SK-ov-3	Ovarian	15
CEM	Leukemic	30
HFL-1	Normal diploid lung	50

[0156] The GI₅₀ of (E)-4-methoxystyryl-N-3-fluoro-4-methoxyphenyl sulfonamide was determined for the following cell lines: BT20, DU145, H157 and DLD-1. The GI₅₀ for each cell line was in essential agreement with the GI₅₀ determined for (E)-4-methoxystyryl-N-4-fluorophenyl sulfonamide in the same cell line.

Example 70: Induction of Apoptosis in Tumor Cells

[0157] The following assay demonstrates the apoptotic activity of the compounds of the invention against tumor cells.

[0158] The caspases and the ICE-family proteases are cysteine proteases which are activated during apoptosis (Patel et al., *FASEB* 10:587-597, 1996). The cleavage of poly(ADP-ribose) polymerase (PARP), which is a target of caspase-3, apopain, and several other activated proteases, is a widely used and accepted marker for apoptosis (Nicholson et al., *Nature* 376(6533):37-43, 1995; Lippke et al., *J. Biol. Chemistry* 271:1825, 1996). For this assay, BT20 cells, an estrogen receptor negative breast carcinoma, and HFL-1 cells, normal lung fibroblasts, were treated with either (E)-4-

methoxystyryl-N-4-fluorophenyl sulfonamide at a final concentration of 20 μ M or dimethyl sulfoxide (DMSO) for 96 hours. The cells were then lysed in RIPA buffer and 100 μ g of total cellular protein from each sample was resolved on a 10% SDS-polyacrylamide gel. The proteins were then Western blotted onto PROTRAN filter paper (S/S) and the filter was then probed with antibody (Boehringer Mannheim) specific for PARP. This antibody recognizes both the 116 kDa full length PARP and the 83 kDa cleaved product. The results, set forth in Fig. 2, show that a 96 hour treatment with the test compound specifically activated caspases in the treated breast carcinoma cell line and not in the normal cell line. The western blot clearly shows that only the test compound-treated BT20 cells had the presence of the 83 kDa PARP cleavage product. The HFL-1 cells treated in a similar manner showed no cleavage of the full length PARP. The BT20 cells treated with DMSO for the same amount of time also had no activation of the apoptotic pathway. These results show that the compounds of the invention selectively kill cancer cells by activating the apoptotic pathway as indicated by the activation of the cysteine proteases, a molecular marker for apoptosis. Cells which are not tumorigenic do not undergo apoptosis but become growth arrested at concentrations significantly higher than the concentration necessary for tumor cell death.

Example 71: Radioprotective Effects of N-Aryl-2-Arylethanesulfonamides on Cultured Normal Cells

[0159] The radioprotective effects of (a) styryl-N-phenylsulfonamide, (b) 4-methoxystyryl-N-fluorophenylsulfonamide; (c) styryl-N-methyl-N-phenylsulfonamide and (d) 4-methoxystyryl-N-2,4,6-trimethoxyphenylsulfonamide were evaluated on cultured normal cells as follows.

[0160] HFL-1 cells, which are normal diploid lung fibroblasts, were plated into 24 well dishes at a cell density of 3000 cells per 10 mm² in DMEM completed with 10% fetal bovine serum and antibiotics. The test compounds were added to the cells 24 hours later in select concentrations from 2.5 μ M and 10.0 μ M, inclusive, using DMSO as a solvent. Control cells were treated with DMSO alone. The cells were exposed to the test compound or DMSO for 24 hours.

[0161] The cells were then irradiated with 10 Gy (gray) of ionizing radiation (IR) using a J.L. Shepherd Mark I, Model 30-1 Irradiator equipped with ¹³⁷cesium as a source. After irradiation, the medium on the test and control cells was removed and replaced with fresh growth medium without the test compounds or DMSO. The irradiated cells were incubated for 96 hours and then duplicate wells were trypsinized and replated onto 100 mm² tissue culture dishes. The replated cells were grown under normal conditions with one change of fresh medium for 2 weeks. The number of colonies from each 100 mm² culture dish, which represents the number of surviving cells, was determined by staining the dishes as described below.

[0162] In order to visualize and count the colonies derived from the clonal outgrowth of individual protected cells, the medium was removed and the plates were washed one time with room temperature phosphate buffered saline. The cells were stained with a 1:10 diluted Modified Geimsa staining solution (Sigma) for 20 minutes. The stain was removed, and the plates were washed with tap water. The plates were air dried, the number of colonies from each plate was counted and the average from duplicate plates was determined. Each compound provided radioprotective activity of between 4- and 6-fold at the concentrations tested. Fold protection was determined by dividing the average number of colonies from the test plates by the average number of colonies counted on the control plates.

Example 72: Protection of Mice from Radiation Toxicity by Pre-Treatment with N-Aryl-2-Arylethanesulfonamides.

[0163] C57 black mice age 10-12 weeks (Taconic) are divided into treatment groups of 10 mice each and given intraperitoneal injections of 200 micrograms of N-aryl-2-arylethanesulfonamide dissolved in DMSO (a 10 mg/Kg dose, based on 20g mice). The injections are given 18 and 6 hours before irradiation with 8 Gy gamma radiation. A control group of 10 animals receives 8 Gy gamma radiation alone. Mortality of control and experimental groups is assessed for 40 days after irradiation.

Example 73: Radioprotective Effect of N-Aryl-2-Arylethanesulfonamides in Mice When Given After Radiation Exposure.

[0164] C57 B6/J mice age 10-12 weeks (Taconic) are divided into treatment groups and one control group of 10 mice each. Each treatment group receives intraperitoneal injections of 200 micrograms of N-aryl-2-arylethanesulfonamide dissolved in DMSO (a 10 mg/Kg dose, based on 20g mice) 15 minutes after irradiation with 8 Gy gamma radiation. The control group receives 8 Gy gamma radiation alone. Mortality of control and treatment groups are assessed for 40 days after irradiation.

Example 74: Effect of Exposure to Ionizing Radiation on Normal and Malignant Hematopoietic Progenitor Cell Growth After Pretreatment with N-Aryl-2-Arylethanesulfonamides.

5 [0165] The effect of ionizing radiation on normal and malignant hematopoietic progenitor cells which are pretreated with N-aryl-2-arylethanesulfonamides is investigated by assessing cloning efficiency and development of the pretreated cells after irradiation.

10 [0166] To obtain hematopoietic progenitor cells, human bone marrow cells (BMC) or peripheral blood cells (PB) are obtained from normal healthy, or acute or chronic myelogenous leukemia (AML, CML), volunteers by Ficoll-Hypaque density gradient centrifugation, and are partially enriched for hematopoietic progenitor cells by positively selecting CD34⁺ cells with immunomagnetic beads (DynaI A.S., Oslo, Norway). The CD34⁺ cells are suspended in supplemented alpha medium and incubated with mouse anti-HPCA-I antibody in 1:20 dilution, 45 minutes, at 4°C with gentle inverting of tubes. Cells are washed x 3 in supplemented alpha medium, and then incubated with beads coated with the Fc fragment of goat anti-mouse IgG₁ (75 µl of immunobeads/10⁷ CD34⁺ cells). After 45 minutes of incubation (4°C), cells adherent to the beads are positively selected using a magnetic particle concentrator as directed by the manufacturer.

15 [0167] 2 x 10⁴ CD34⁺ cells are incubated in 5 ml polypropylene tubes (Fisher Scientific, Pittsburgh, PA) in a total volume of 0.4 ml of Iscove's modified Dulbecco's medium (IMDM) containing 2% human AB serum and 10 mM Hepes buffer. An N-aryl-2-arylethanesulfonamide, for example styryl-N-phenylsulfonamide; 4-methoxystyryl-N-fluorophenylsulfonamide; styryl-N-methyl-N-phenylsulfonamide; or 4-methoxystyryl-N-2,4,6-trimethoxyphenylsulfonamide at three different concentrations (2.5 µM, 5.0 µM and 10.0 µM) in DMSO are added separately to the cells. Control cells received 20 DMSO alone. The cells are incubated for 20 - 24 hours and irradiated with 5 Gy or 10 Gy of ionizing radiation. Immediately after irradiation, the medium is removed and replaced with fresh medium without the test compound or DMSO. Twenty-four hours after irradiation, the treatment and control cells are prepared for plating in plasma clot or methylcellulose cultures. Cells (1x 10⁴ CD34⁺ cells per dish) were not washed before plating.

25 [0168] Assessment of the cloning efficiency and development of the treated hematopoietic progenitor cells are carried out essentially as reported in Gewirtz et al., Science 242, 1303-1306 (1988).

Example 75: Bone Marrow Purging with Ionizing Radiation After Pretreatment with N-Aryl-2-Arylethanesulfonamides.


30 [0169] Bone marrow is harvested from the iliac bones of a subject under general anesthesia in an operating room using standard techniques. Multiple aspirations are taken into heparinized syringes. Sufficient marrow is withdrawn so that the subject will be able to receive about 4 x 10⁸ to about 8 x 10⁸ processed marrow cells per kg of body weight. Thus, about 750 to 1000 ml of marrow is withdrawn. The aspirated marrow is transferred immediately into a transport medium (TC-199, Gibco, Grand Island, New York) containing 10,000 units of preservative-free heparin per 100 ml of medium. The aspirated marrow is filtered through three progressively finer meshes to obtain a cell suspension devoid of cellular aggregates, debris and bone particles. The filtered marrow is then processed further into an automated cell separator (e.g., Cobe 2991 Cell Processor) which prepares a "buffy coat" product, (i.e., leukocytes devoid of red cells and platelets). The buffy coat preparation is then placed in a transfer pack for further processing and storage. It may be stored until purging in liquid nitrogen using standard procedures. Alternatively, purging can be carried out immediately, then the purged marrow may be stored frozen in liquid nitrogen until it is ready for transplantation.

35 [0170] The purging procedure is carried out as follows. Cells in the buffy coat preparation are adjusted to a cell concentration of about 2 x 10⁷/ml in TC-199 containing about 20% autologous plasma. An N-aryl-2-arylethanesulfonamide; for example 2.5 to 10 micromolar of either styryl-N-phenylsulfonamide; 4-methoxystyryl-N-fluorophenylsulfonamide, styryl-N-methyl-N-phenylsulfonamide; or 4-methoxystyryl-N-2,4,6-trimethoxyphenylsulfonamide in DMSO is added to the transfer packs containing the cell suspension and incubated in a 37°C water bath for 20 - 24 hours with gentle shaking. The transfer packs are then exposed to 5 - 10 Gy ionizing radiation. Recombinant human hematopoietic growth factors, e.g., rH IL-3 or rH GM-CSF, may be added to the suspension to stimulate growth of hematopoietic neoplasms and thereby increase their sensitivity to ionizing radiation.

40 [0171] The cells may then either be frozen in liquid nitrogen or washed once at 4°C in TC-199 containing about 20% autologous plasma. Washed cells are then infused into the subject. Care must be taken to work under sterile conditions wherever possible and to maintain scrupulous aseptic techniques at all times.

Claims

55 1. A compound of the formula I:



I


wherein:

Q₁ is selected from the group consisting of substituted and unsubstituted aryl, and substituted and unsubstituted heteroaryl;

Q₂ is selected from the group consisting of substituted aryl, and substituted and unsubstituted heteroaryl;

R is selected from the group consisting of hydrogen and (C₁-C₆)alkyl;

wherein the substituents for the substituted aryl and substituted heteroaryl groups comprising Q₁ are independently selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, cyano, carboxy, hydroxy, (C₂-C₆)hydroxyalkyl, phosphonato, amino, (C₁-C₆)acylamino, sulfamyl, acetoxy, di(C₁-C₆)alkylamino(C₂-C₆)alkoxy, trifluoromethyl and



wherein:

X is oxygen or sulfur,

R₅ is selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)heteroalkyl, substituted phenyl, and unsubstituted phenyl, and

R₆ is selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)heteroalkyl, substituted aryl, unsubstituted aryl, substituted heteroaryl, unsubstituted heteroaryl, substituted aryl(C₁-C₃)alkyl, unsubstituted aryl-(C₁-C₃)alkyl and (C₁-C₆)alkoxycarbonyl(C₁-C₆)alkylenyl;

wherein the substituents for the substituted aryl and substituted heteroaryl groups comprising Q₂, and the substituents for the substituted aryl and substituted heteroaryl groups comprising or included within R₅ and R₆, are independently selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, nitro, cyano, carboxy, carboxy(C₁-C₃)alkoxy, hydroxy, (C₂-C₆)hydroxyalkyl, phosphonato, amino, (C₁-C₆)acylamino, sulfamyl, acetoxy, di(C₁-C₆)alkylamino(C₂-C₆ alkoxy) and trifluoromethyl; provided that when R is hydrogen:

(a) when Q₁ is unsubstituted phenyl, Q₂ is other than dimethoxyphenyl, 2-methylphenyl, 2-chlorophenyl, 4-chlorophenyl, 4-N,N-dimethylaminophenyl, 4-methylphenyl, 4-methoxyphenyl, 4-nitrophenyl, 3-methoxy-4-hydroxyphenyl, unsubstituted pyrenyl, unsubstituted benzodioxolyl, and unsubstituted 2-thienyl;

(b) when Q₁ is 3-hydroxyphenyl, Q₂ is other than nitrophenyl;

(c) when Q₁ is 2-methyl-5-hydroxyphenyl, Q₂ is other than 4-nitrophenyl;

and

(d) when Q₁ is unsubstituted 2-pyridyl, Q₂ is other than 3-methoxy-4-hydroxyphenyl;

or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1, provided:

(a) when Q₁ is unsubstituted phenyl, Q₂ is other than dialkoxyphenyl, 2-alkylphenyl, 2-halophenyl, 4-halophenyl, 4-N,N-dialkylaminophenyl, 4-alkylphenyl, 4-alkoxyphenyl, 4-nitrophenyl, 3-alkoxy-4-hydroxyphenyl, unsubsti-

tuted phenyl, unsubstituted pyrenyl, unsubstituted benzodioxolyl, unsubstituted 1-naphthyl and unsubstituted 2-thienyl;

(b) when Q₁ is 3-hydroxyphenyl, Q₂ is other than nitrophenyl;

(c) when Q₁ is 2-methyl-5-hydroxyphenyl, Q₂ is other than 4-nitrophenyl; and

(d) when Q₁ is unsubstituted 2-pyridyl, Q₂ is other than 3-methoxy-4-hydroxyphenyl;

or a pharmaceutically acceptable salt thereof.

3. A compound according to claim 2, provided:

when R is hydrogen:

(i) Q₁ may not be dinitrophenyl;

(ii) Q₂ may not be dinitrophenyl; and


(iii) when Q₂ is mononitrophenyl:

Q₁ is other than substituted phenyl, or

Q₁ is substituted phenyl wherein at least the 4-position is substituted, and the substituent is other than hydroxy;

or a pharmaceutically acceptable salt thereof.

4. A compound of formula I, for use in medicine:




wherein:

Q₁ is selected from the group consisting of substituted and unsubstituted aryl, and substituted and unsubstituted heteroaryl;

Q₂ is selected from the group consisting of substituted aryl, and substituted and unsubstituted heteroaryl;

R is selected from the group consisting of hydrogen and (C₁-C₆)alkyl;

wherein the substituents for the substituted aryl and substituted heteroaryl groups comprising Q₁ are independently selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, cyano, carboxy, hydroxy, (C₂-C₆)hydroxyalkyl, phosphonato, amino, (C₁-C₆)acylamino, sulfamyl, acetoxy, di(C₁-C₆)alkylamino(C₂-C₆)alkoxy, trifluoromethyl and



wherein:

X is oxygen or sulfur,

R₅ is selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)heteroalkyl, substituted phenyl, and

unsubstituted phenyl, and

R₆ is selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)heteroalkyl, substituted aryl, unsubstituted aryl, substituted heteroaryl, unsubstituted heteroaryl, substituted aryl(C₁-C₃)alkyl, unsubstituted aryl-(C₁-C₃)alkyl and (C₁-C₆)alkoxycarbonyl(C₁-C₆)alkylenyl;

wherein the substituents for the substituted aryl and substituted heteroaryl groups comprising Q₂, and the substituents for the substituted aryl and substituted heteroaryl groups comprising or included within R⁵ and R⁶, are independently selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, nitro, cyano, carboxy, carboxy(C₁-C₃)alkoxy, hydroxy, (C₂-C₆)hydroxyalkyl, phosphonato, amino, (C₁-C₆)acylamino, sulfamyl, acetoxy, di(C₁-C₆)alkylamino(C₂-C₆ alkoxy) and trifluoromethyl;

provided, when R is hydrogen and Q₂ is unsubstituted phenyl, then Q₁ must be other than dihalophenyl;

or a pharmaceutically acceptable salt thereof.

5. A compound according to claim 1 or 4 wherein Q₁ is optionally substituted phenyl and Q₂ is substituted phenyl, or a pharmaceutically acceptable salt thereof.

6. A compound according to claim 5 wherein at least one of Q₁ and Q₂ is substituted in at least the 4-position, or a pharmaceutically acceptable salt thereof.

7. A compound according to claim 6 wherein the substituents are independently selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, nitro, hydroxy and sulfamyl, or a pharmaceutically acceptable salt thereof.

8. A compound according to claim 7 wherein the compound is:

(E)-4-methoxystyryl-N-methyl-N-phenyl sulfonamide,
(E)-4-chlorostyryl-N-3-chlorophenyl sulfonamide,
(E)-4-chlorostyryl-N-2-chlorophenyl sulfonamide,

or a pharmaceutically acceptable salt thereof.

9. A compound according to claim 8 wherein Q₁ and Q₂ are both independently substituted in at least the 4-position, or a pharmaceutically acceptable salt thereof.

10. A compound according to claim 9 wherein the substituents are independently selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, nitro, hydroxy and sulfamyl, or a pharmaceutically acceptable salt thereof.

11. A compound according to claim 10 wherein the compound is:

(E)-4-methoxystyryl-N-4-sulfamylphenyl sulfonamide,
(E)-4-chlorostyryl-N-3-fluoro-4-methoxyphenyl sulfonamide,
(E)-4-chlorostyryl-N-4-fluorophenyl sulfonamide,
(E)-4-methoxystyryl-N-4-fluorophenyl sulfonamide,
(E)-4-methoxystyryl-N-3-fluoro-4-methoxyphenyl sulfonamide,
(E)-4-methoxystyryl-N-4-fluorophenyl sulfonamide,
(E)-4-fluorostyryl-N-4-chlorophenyl sulfonamide,
(E)-4-methoxystyryl-N-2,4,6-trimethoxyphenyl sulfonamide,


or a pharmaceutically acceptable salt thereof.

12. A compound according to claim 6 of the formula:

5

10

15



II

20


wherein R is hydrogen or (C₁-C₆)alkyl; R₁ is selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, cyano, carboxy, hydroxy, (C₂-C₆)hydroxyalkyl, phosphonato, amino, (C₁-C₆)acylamino, sulfamyl, acetoxy, di(C₁-C₆)alkylamino(C₂-C₆)alkoxy and trifluoromethyl; and R₂, R₃ and R₄, are independently selected from (C₁-C₆)alkoxy, or a pharmaceutically acceptable salt thereof.

13. A compound according to claim 12 of the formula:

25

30

35



IIa

wherein R, R₁, R₂, R₃ and R₄ are defined as in claim 12, or a pharmaceutically acceptable salt thereof.

40

14. A compound according to claim 13 wherein the compound is (E)-2,4,6-trimethoxystyryl-N-4-methoxyphenyl sulfonamide, or a pharmaceutically acceptable salt thereof.

45

15. A compound according to claim 5 wherein at least one of Q₁ and Q₂ is penta-substituted with halogen, or a pharmaceutically acceptable salt thereof.

16. A compound according to claim 15 wherein at least one of Q₁ and Q₂ is pentafluorophenyl, or a pharmaceutically acceptable salt thereof.

50

17. A compound according to claim 16 wherein the compound is:


(E)-4-methoxystyryl-N-2,3,4,5,6-pentafluorophenyl sulfonamide,
(E)-4-fluorostyryl-N-2,3,4,5,6-pentafluorophenyl sulfonamide,

55

or a pharmaceutically acceptable salt thereof.

18. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and at least one compound, or pharmaceutically acceptable salt thereof, according to any preceding claim.

19. Use of a compound according to the formula:



wherein:

15 Q₁ is selected from the group consisting of substituted and unsubstituted aryl, and substituted and unsubstituted heteroaryl;

Q₂ is selected from the group consisting of substituted aryl, and substituted and unsubstituted heteroaryl;

R is selected from the group consisting of hydrogen and (C₁-C₆)alkyl;

20 wherein the substituents for the substituted aryl and substituted heteroaryl groups comprising Q₁ are independently selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, cyano, carboxy, hydroxy, (C₂-C₆)hydroxyalkyl, phosphonato, amino, (C₁-C₆)acylamino, sulfamyl, acetoxy, di(C₁-C₆)alkylamino(C₂-C₆)alkoxy, trifluoromethyl and



wherein:

X is oxygen or sulfur,

35 R₅ is selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)heteroalkyl, substituted phenyl, and unsubstituted phenyl, and

R₆ is selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)heteroalkyl, substituted aryl, unsubstituted aryl, substituted heteroaryl, unsubstituted heteroaryl, substituted aryl(C₁-C₃)alkyl, unsubstituted aryl-(C₁-C₃)alkyl and (C₁-C₆)alkoxycarbonyl(C₁-C₆)alkylenyl;

40 wherein the substituents for the substituted aryl and substituted heteroaryl groups comprising Q₂, and the substituents for the substituted aryl and substituted heteroaryl groups comprising or included within R₅ and R₆, are independently selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, nitro, cyano, carboxy, carboxy(C₁-C₃)alkoxy, hydroxy, (C₂-C₆)hydroxyalkyl, phosphonato, amino, (C₁-C₆)acylamino, sulfamyl, acetoxy, di(C₁-C₆)alkylamino(C₂-C₆ alkoxy) and trifluoromethyl;

45 or a pharmaceutically acceptable salt thereof;


for preparation of a medicament for treatment of a proliferative disorder.

20. A use according to claim 19 wherein the proliferative disorder is selected from the group consisting of cancer, hemangiomas in newborn, secondary progressive multiple sclerosis, chronic progressive myelodegenerative disease, neurofibromatosis, ganglioneuromatosis, keloid formation, Paget's disease of the bone, fibrocystic disease of the breast, Peyronie's fibrosis, Dupuytren's fibrosis, restenosis and cirrhosis.

21. A use according to claim 20 wherein the proliferative disorder is cancer, and the cancer is selected from the group consisting of ovarian, breast, prostate, lung, renal, colorectal and brain cancers, or the cancer is a leukemia.

22. A use according to claim 20 wherein the proliferative disorder is cancer, for inducing apoptosis of tumor cells.

23. Use of a compound according to the formula




wherein:

15 Q_1 is selected from the group consisting of substituted and unsubstituted aryl, and substituted and unsubstituted heteroaryl;

Q_2 is selected from the group consisting of substituted aryl, and substituted and unsubstituted heteroaryl;

R is selected from the group consisting of hydrogen and (C₁-C₆)alkyl;

20 wherein the substituents for the substituted aryl and substituted heteroaryl groups comprising Q_1 are independently selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, cyano, carboxy, hydroxy, (C₂-C₆)hydroxyalkyl, phosphonato, amino, (C₁-C₆)acylamino, sulfamyl, acetoxy, di(C₁-C₆)alkylamino(C₂-C₆)alkoxy, trifluoromethyl and



wherein:

X is oxygen or sulfur,

35 R_5 is selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)heteroalkyl, substituted phenyl, and unsubstituted phenyl, and

R_6 is selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₂-C₆)heteroalkyl, substituted aryl, unsubstituted aryl, substituted heteroaryl, unsubstituted heteroaryl, substituted aryl(C₁-C₃)alkyl, unsubstituted aryl-(C₁-C₃)alkyl and (C₁-C₆)alkoxycarbonyl(C₁-C₆)alkylenyl;

40 wherein the substituents for the substituted aryl and substituted heteroaryl groups comprising Q_2 , and the substituents for the substituted aryl and substituted heteroaryl groups comprising or included within R_5 and R_6 , are independently selected from the group consisting of halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, nitro, cyano, carboxy, carboxy(C₁-C₃)alkoxy, hydroxy, (C₂-C₆)hydroxyalkyl, phosphonato, amino, (C₁-C₆)acylamino, sulfamyl, acetoxy, di(C₁-C₆)alkylamino(C₂-C₆)alkoxy and trifluoromethyl;

45 or a pharmaceutically acceptable salt thereof;


for the preparation of a medicament for reducing or eliminating the effects of ionizing radiation on normal cells in a subject who has incurred or is at risk for incurring exposure to ionizing radiation, or in bone marrow cells from such a subject.

50 **24.** A use according to claim 23, wherein the ionizing radiation is therapeutic ionizing radiation for the treatment of a proliferative disorder.

25. A use according to claim 24 wherein the proliferative disorder is cancer.

55 **26.** A process for preparing a compound according to claim 1, comprising reacting a compound of the formula Q_2 -CH=CH-SO₂Cl with a compound of the formula Q_1 -NRH in a nonprotic solvent in the presence of a base to form a compound of the formula:

5



10


wherein:

Q₁, Q₂, and R are defined as in claim 1.

15

27. A process for preparing a compound according to claim 1, comprising reacting a compound of the formula


20



25

with a compound of the formula Q₂-C(O)H in a nonprotic solvent in the presence of a base to form a compound formula:

30



35

wherein:

40


Q₁, Q₂, and R are defined as in claim 1.

Patentansprüche

45

1. Eine Verbindung der Formel I:

50



55

wobei:

Q₁ aus der Gruppe, bestehend aus substituiertem und nichtsubstituiertem Aryl und substituiertem und nicht-substituiertem Heteroaryl, ausgewählt ist;

Q₂ aus der Gruppe, bestehend aus substituiertem Aryl und substituiertem und nichtsubstituiertem Heteroaryl, ausgewählt ist;

R aus der Gruppe, bestehend aus Wasserstoff und (C₁-C₆)-Alkyl, ausgewählt ist;

wobei die Substituenten für die substituierten Aryl- und substituierten Heteroarylgruppen, die Q₁ beinhalten, unabhängig aus der Gruppe, bestehend aus Halogen, (C₁-C₆)-Alkyl, (C₁-C₆)-Alkoxy, Cyan, Carboxy, Hydroxy, (C₂-C₆)-Hydroxyalkyl, Phosphonato, Amino, (C₁-C₆)-Acylamino, Sulfamyl, Acetoxy, Di-(C₁-C₆)-alkylamino-(C₂-C₆)-alkoxy, Trifluormethyl und Folgendem, ausgewählt sind:



wobei:

X Sauerstoff oder Schwefel ist,

R₅ aus der Gruppe, bestehend aus Wasserstoff, (C₁-C₆)-Alkyl, (C₂-C₆)-Heteroalkyl, substituiertem Phenyl und nichtsubstituiertem Phenyl, ausgewählt ist und

R₆ aus der Gruppe, bestehend aus Wasserstoff, (C₁-C₆)-Alkyl, (C₂-C₆)-Heteroalkyl, substituiertem Aryl, nicht-substituiertem Aryl, substituiertem Heteroaryl, nichtsubstituiertem Heteroaryl, substituiertem Aryl-(C₁-C₃)-alkyl, nichtsubstituiertem Aryl-(C₁-C₃)-alkyl und (C₁-C₆)-Alkoxy-carbonyl-(C₁-C₆)-alkylenyl, ausgewählt ist;

wobei die Substituenten für die substituierten Aryl- und substituierten Heteroarylgruppen, die Q₂ beinhalten, und die Substituenten für die substituierten Aryl- und substituierten Heteroarylgruppen, die R₅ und R₆ beinhalten oder darin umfasst sind, unabhängig aus der Gruppe, bestehend aus Halogen, (C₁-C₆)-Alkyl, (C₁-C₆)-Alkoxy, Nitro, Cyan, Carboxy, Carboxy-(C₁-C₃)-alkoxy, Hydroxy, (C₂-C₆)-Hydroxyalkyl, Phosphonato, Amino, (C₁-C₆)-Acylamino, Sulfamyl, Acetoxy, Di-(C₁-C₆)-alkylamino-(C₂-C₆)-alkoxy) und Trifluormethyl, ausgewählt sind;

vorausgesetzt dass, wenn R Wasserstoff ist:

(a) wenn Q₁ nichtsubstituiertes Phenyl ist, Q₂ anders als Dimethoxyphenyl, 2-Methylphenyl, 2-Chlorphenyl, 4-Chlorphenyl, 4-N,N-Dimethylaminophenyl, 4-Methylphenyl, 4-Methoxyphenyl, 4-Nitrophenyl, 3-Methoxy-4-hydroxyphenyl, nichtsubstituiertes Pyrenyl, nichtsubstituiertes Benzodioxolyl und nichtsubstituiertes 2-Thienyl ist;

(b) wenn Q₁ 3-Hydroxyphenyl ist, Q₂ anders als Nitrophenyl ist;

(c) wenn Q₁ 2-Methyl-5-hydroxyphenyl ist, Q₂ anders als 4-Nitrophenyl ist und

(d) wenn Q₁ nichtsubstituiertes 2-Pyridyl ist, Q₂ anders als 3-Methoxy-4-hydroxyphenyl ist;

oder ein pharmazeutisch zulässiges Salz davon.

2. Verbindung gemäß Anspruch 1, vorausgesetzt, dass:

(a) wenn Q₁ nichtsubstituiertes Phenyl ist, Q₂ anders als Dialkoxyphenyl, 2-Alkylphenyl, 2-Halogenphenyl, 4-Halogenphenyl, 4-N,N-Dialkylaminophenyl, 4-Alkylphenyl, 4-Alkoxyphenyl, 4-Nitrophenyl, 3-Alkoxy-4-hydroxyphenyl, nichtsubstituiertes Phenyl, nichtsubstituiertes Pyrenyl, nichtsubstituiertes Benzodioxolyl, nichtsubstituiertes 1-Naphthyl and nichtsubstituiertes 2-Thienyl ist;

(b) wenn Q₁ 3-Hydroxyphenyl ist, Q₂ anders als Nitrophenyl ist;

(c) wenn Q₁ 2-Methyl-5-hydroxyphenyl ist, Q₂ anders als 4-Nitrophenyl ist und

(d) wenn Q₁ nichtsubstituiertes 2-Pyridyl ist, Q₂ anders als 3-Methoxy-4-hydroxyphenyl ist;

oder ein pharmazeutisch zulässiges Salz davon.

3. Verbindung gemäß Anspruch 2, vorausgesetzt, dass:

wenn R Wasserstoff ist:


- 5 (i) Q₁ möglicherweise anders als Dinitrophenyl ist;
 (ii) Q₂ möglicherweise anders als Dinitrophenyl ist und
 (iii) wenn Q₂ Mononitrophenyl ist:

10 Q₁ anders als substituiertes Phenyl ist oder
 Q₁ substituiertes Phenyl ist, wobei mindestens die 4-Position substituiert ist und der Substituent anders als Hydroxy ist;
 oder ein pharmazeutisch zulässiges Salz davon.

4. Verbindung der Formel I zur Verwendung in der Medizin:

15

20



25

30

wobei:

Q₁ aus der Gruppe, bestehend aus substituiertem und nichtsubstituiertem Aryl und substituiertem und nicht-substituiertem Heteroaryl, ausgewählt ist;

35

Q₂ aus der Gruppe, bestehend aus substituiertem Aryl und substituiertem und nichtsubstituiertem Heteroaryl, ausgewählt ist;

R aus der Gruppe, bestehend aus Wasserstoff und (C₁-C₆)-Alkyl, ausgewählt ist;

wobei die Substituenten für die substituierten Aryl- und substituierten Heteroarylgruppen, die Q₁ beinhalten, unabhängig aus der Gruppe, bestehend aus Halogen, (C₁-C₆)-Alkyl, (C₁-C₆)-Alkoxy, Cyan, Carboxy, Hydroxy, (C₂-C₆)-Hydroxyalkyl, Phosphonato, Amino, (C₁-C₆)-Acylamino, Sulfamyl, Acetoxy, Di-(C₁-C₆)-alkylamino-(C₂-C₆)-alkoxy, Trifluormethyl und Folgendem, ausgewählt sind:

40

45



50

wobei:

X Sauerstoff oder Schwefel ist,

R₅ aus der Gruppe, bestehend aus Wasserstoff, (C₁-C₆)-Alkyl, (C₂-C₆)-Heteroalkyl, substituiertem Phenyl und nichtsubstituiertem Phenyl, ausgewählt ist und

55

R₆ aus der Gruppe, bestehend aus Wasserstoff, (C₁-C₆)-Alkyl, (C₂-C₆)-Heteroalkyl, substituiertem Aryl, nicht-substituiertem Aryl, substituiertem Heteroaryl, nichtsubstituiertem Heteroaryl, substituiertem Aryl-(C₁-C₃)-Alkyl, nichtsubstituiertem Aryl-(C₁-C₃)-alkyl und (C₁-C₆)-Alkoxy-carbonyl-(C₁-C₆)-alkylenyl, ausgewählt ist;
 wobei die Substituenten für die substituierten Aryl- und substituierten Heteroarylgruppen, die Q₂ beinhalten,

und die Substituenten für die substituierten Aryl- und substituierten Heteroarylgruppen, die R₅ und R₆ beinhalten oder darin umfasst sind, unabhängig aus der Gruppe, bestehend aus Halogen, (C₁-C₆)-Alkyl, (C₁-C₆)-Alkoxy, Nitro, Cyan, Carboxy, Carboxy-(C₁-C₃)-alkoxy, Hydroxy, (C₂-C₆)-Hydroxyalkyl, Phosphonato, Amino, (C₁-C₆)-Acylamino, Sulfamyl, Acetoxy, Di-(C₁-C₆)-alkylamino-(C₂-C₆-alkoxy) und Trifluormethyl, ausgewählt sind;

vorausgesetzt dass, wenn R Wasserstoff und Q₂ nichtsubstituiertes Phenyl ist, Q₁ dann anders als Dihalogenphenyl sein muss;

oder ein pharmazeutisch zulässiges Salz davon.

5
10 **5.** Verbindung gemäß Anspruch 1 oder 4, wobei Q₁ optional substituiertes Phenyl ist und Q₂ substituiertes Phenyl ist, oder ein pharmazeutisch zulässiges Salz davon.

6. Verbindung gemäß Anspruch 5, wobei mindestens eines von Q₁ und Q₂ an mindestens der 4-Position substituiert ist, oder ein pharmazeutisch zulässiges Salz davon.

15 **7.** Verbindung gemäß Anspruch 6, wobei die Substituenten unabhängig aus der Gruppe, bestehend aus Halogen, (C₁-C₆)-Alkyl, (C₁-C₆)-Alkoxy, Nitro, Hydroxy und Sulfamyl, ausgewählt sind, oder ein pharmazeutisch zulässiges Salz davon.

20 **8.** Verbindung gemäß Anspruch 7, wobei die Verbindung Folgende ist:

(E)-4-Methoxystyryl-N-methyl-N-phenyl-sulfonamid,

(E)-4-Chlorstyryl-N-3-chlorphenyl-sulfonamid,

(E)-4-Chlorstyryl-N-2-chlorphenyl-sulfonamid

25
oder ein pharmazeutisch zulässiges Salz davon.

9. Verbindung gemäß Anspruch 8, wobei sowohl Q₁ als auch Q₂ unabhängig an mindestens der 4-Position substituiert sind, oder ein pharmazeutisch zulässiges Salz davon.

30 **10.** Verbindung gemäß Anspruch 9, wobei die Substituenten unabhängig aus der Gruppe, bestehend aus Halogen, (C₁-C₆)-Alkyl, (C₁-C₆)-Alkoxy, Nitro, Hydroxy und Sulfamyl, ausgewählt sind, oder ein pharmazeutisch zulässiges Salz davon.

35 **11.** Verbindung gemäß Anspruch 10, wobei die Verbindung Folgende ist:

(E)-4-Methoxystyryl-N-4-sulfamylphenyl-sulfonamid,

(E)-4-Chlorstyryl-N-3-fluor-4-methoxyphenyl-sulfonamid,

(E)-4-Chlorstyryl-N-4-fluorphenyl-sulfonamid,

40 (E)-4-Methoxystyryl-N-4-fluorphenyl-sulfonamid,

(E)-4-Methoxystyryl-N-3-fluor-4-methoxyphenyl-sulfonamid,

(E)-4-Methoxystyryl-N-4-fluorphenyl-sulfonamid,

(E)-4-Fluorstyryl-N-4-chlorphenyl-sulfonamid,

(E)-4-Methoxystyryl-N-2,4,6-trimethoxyphenyl-sulfonamid

45
oder ein pharmazeutisch zulässiges Salz davon.

12. Verbindung gemäß Anspruch 6 der Formel:


50

55

5

10

15



II

20

wobei R Wasserstoff oder (C₁-C₆)-Alkyl ist; R₁ aus der Gruppe, bestehend aus Halogen, (C₁-C₆)-Alkyl, (C₁-C₆)-Alkoxy, Cyan, Carboxy, Hydroxy, (C₂-C₆)-Hydroxyalkyl, Phosphonato, Amino, (C₁-C₆)-Acylamino, Sulfamyl, Acetoxy, Di-(C₁-C₆)-alkylamino-(C₂-C₆)-alkoxy und Trifluormethyl, ausgewählt ist und R₂, R₃ und R₄ unabhängig aus (C₁-C₆)-Alkoxy ausgewählt sind, oder ein pharmazeutisch zulässiges Salz davon.


25

13. Verbindung gemäß Anspruch 12 der Formel:

30

35

40



IIa

45

wobei R, R₁, R₂, R₃ und R₄ gemäß Anspruch 12 definiert sind, oder ein pharmazeutisch zulässiges Salz davon.

50

14. Verbindung gemäß Anspruch 13, wobei die Verbindung (E)-2,4,6-Trimethoxystyryl-N-4-methoxyphenyl-sulfonamid ist, oder ein pharmazeutisch zulässiges Salz davon.

15. Verbindung gemäß Anspruch 5, wobei mindestens eines von Q₁ und Q₂ mit Halogen pentasubstituiert ist, oder ein pharmazeutisch zulässiges Salz davon.

55

16. Verbindung gemäß Anspruch 15, wobei mindestens eines von Q₁ und Q₂ Pentafluorphenyl ist, oder ein pharmazeutisch zulässiges Salz davon.

17. Verbindung gemäß Anspruch 16, wobei die Verbindung Folgende ist:


EP 1 379 508 B9

(E)-4-Methoxystyryl-N-2,3,4,5,6-pentafluorphenyl-sulfonamid,
(E)-4-Fluorstyryl-N-2,3,4,5,6-pentafluorphenyl-sulfonamid

oder ein pharmazeutisch zulässiges Salz davon.

18. Eine pharmazeutische Zusammensetzung, die einen pharmazeutisch zulässigen Träger und mindestens eine Verbindung beinhaltet, oder ein pharmazeutisch zulässiges Salz davon gemäß einem der vorhergehenden Ansprüche.

19. Eine Verwendung einer Verbindung gemäß folgender Formel:




wobei:

Q₁ aus der Gruppe, bestehend aus substituiertem und nichtsubstituiertem Aryl und substituiertem und nichtsubstituiertem Heteroaryl, ausgewählt ist;

Q₂ aus der Gruppe, bestehend aus substituiertem Aryl und substituiertem und nichtsubstituiertem Heteroaryl, ausgewählt ist;

R aus der Gruppe, bestehend aus Wasserstoff und (C₁-C₆)-Alkyl, ausgewählt ist;

wobei die Substituenten für die substituierten Aryl- und substituierten Heteroarylgruppen, die Q₁ beinhalten, unabhängig aus der Gruppe, bestehend aus Halogen, (C₁-C₆)-Alkyl, (C₁-C₆)-Alkoxy, Cyan, Carboxy, Hydroxy, (C₂-C₆)-Hydroxyalkyl, Phosphonato, Amino, (C₁-C₆)-Acylamino, Sulfamyl, Acetoxy, Di-(C₁-C₆)-alkylamino-(C₂-C₆)-alkoxy, Trifluormethyl und Folgendem, ausgewählt sind:



wobei:

X Sauerstoff oder Schwefel ist,

R₅ aus der Gruppe, bestehend aus Wasserstoff, (C₁-C₆)-Alkyl, (C₂-C₆)-Heteroalkyl, substituiertem Phenyl und nichtsubstituiertem Phenyl, ausgewählt ist und

R₆ aus der Gruppe, bestehend aus Wasserstoff, (C₁-C₆)-Alkyl, (C₂-C₆)-Heteroalkyl, substituiertem Aryl, nichtsubstituiertem Aryl, substituiertem Heteroaryl, nichtsubstituiertem Heteroaryl, substituiertem Aryl-(C₁-C₃)-alkyl, nichtsubstituiertem Aryl-(C₁-C₃)-alkyl und (C₁-C₆)-Alkoxy-carbonyl-(C₁-C₆)-alkylenyl, ausgewählt ist;

wobei die Substituenten für die substituierten Aryl- und substituierten Heteroarylgruppen, die Q₂ beinhalten, und die Substituenten für die substituierten Aryl- und substituierten Heteroarylgruppen, die R₅ und R₆ beinhalten oder darin umfasst sind, unabhängig aus der Gruppe, bestehend aus Halogen, (C₁-C₆)-Alkyl, (C₁-C₆)-Alkoxy, Nitro, Cyan, Carboxy, Carboxy-(C₁-C₃)-alkoxy, Hydroxy, (C₂-C₆)-Hydroxyalkyl, Phosphonato, Amino, (C₁-C₆)-Acylamino, Sulfamyl, Acetoxy, Di-(C₁-C₆)-alkylamino-(C₂-C₆)-alkoxy) und Trifluormethyl, ausgewählt sind;

oder eines pharmazeutisch zulässigen Salzes davon


zur Zubereitung eines Medikaments zur Behandlung einer proliferativen Störung.

20. Verwendung gemäß Anspruch 19, wobei die proliferative Störung aus der Gruppe, bestehend aus Krebs, Haemangiomas bei Neugeborenen, sekundärer progressiver Multipler Sklerose, chronisch progressiver myelodegenerativer Krankheit, Neurofibromatosis generalisata, Ganglioneuromatose, Keloidbildung, Paget-Krankheit, fibrözytischer Mastopathie, Peyronie-Krankheit, Dupuytren-Krankheit, Restenose und Zirrhose, ausgewählt ist.

21. Verwendung gemäß Anspruch 20, wobei die proliferative Störung Krebs ist und der Krebs aus der Gruppe, bestehend aus Eierstock-, Brust-, Prostata-, Lungen-, Nieren-, Kolorektal- und Gehirnkrebs, ausgewählt ist oder der Krebs Leukämie ist.

22. Verwendung gemäß Anspruch 20, wobei die proliferative Störung Krebs zum Induzieren der Apoptose von Tumorzellen ist.

23. Eine Verwendung einer Verbindung gemäß folgender Formel:




wobei:

Q₁ aus der Gruppe, bestehend aus substituiertem und nichtsubstituiertem Aryl und substituiertem und nichtsubstituiertem Heteroaryl, ausgewählt ist;

Q₂ aus der Gruppe, bestehend aus substituiertem Aryl und substituiertem und nichtsubstituiertem Heteroaryl, ausgewählt ist;

R aus der Gruppe, bestehend aus Wasserstoff und (C₁-C₆)-Alkyl, ausgewählt ist;

wobei die Substituenten für die substituierten Aryl- und substituierten Heteroarylgruppen, die Q₁ beinhalten, unabhängig aus der Gruppe, bestehend aus Halogen, (C₁-C₆)-Alkyl, (C₁-C₆)-Alkoxy, Cyan, Carboxy, Hydroxy, (C₂-C₆)-Hydroxyalkyl, Phosphonato, Amino, (C₁-C₆)-Acylamino, Sulfamyl, Acetoxy, Di-(C₁-C₆)-alkylamino-(C₂-C₆)-alkoxy, Trifluormethyl und Folgendem, ausgewählt sind:



wobei:

X Sauerstoff oder Schwefel ist,

R₅ aus der Gruppe, bestehend aus Wasserstoff, (C₁-C₆)-Alkyl, (C₂-C₆)-Heteroalkyl, substituiertem Phenyl und nichtsubstituiertem Phenyl, ausgewählt ist und

R₆ aus der Gruppe, bestehend aus Wasserstoff, (C₁-C₆)-Alkyl, (C₂-C₆)-Heteroalkyl, substituiertem Aryl, nichtsubstituiertem Aryl, substituiertem Heteroaryl, nichtsubstituiertem Heteroaryl, substituiertem Aryl-(C₁-C₃)-alkyl, nichtsubstituiertem Aryl-(C₁-C₃)-alkyl und (C₁-C₆)-Alkoxy-carbonyl-(C₁-C₆)-alkylenyl, ausgewählt ist;

wobei die Substituenten für die substituierten Aryl- und substituierten Heteroarylgruppen, die Q₂ beinhalten, und die Substituenten für die substituierten Aryl- und substituierten Heteroarylgruppen, die R₅ und R₆ beinhalten

EP 1 379 508 B9

oder darin umfasst sind, unabhängig aus der Gruppe, bestehend aus Halogen, (C₁-C₆)-Alkyl, (C₁-C₆)-Alkoxy, Nitro, Cyan, Carboxy, Carboxy-(C₁-C₃)-alkoxy, Hydroxy, (C₂-C₆)-Hydroxyalkyl, Phosphonato, Amino, (C₁-C₆)-Acylamino, Sulfamyl, Acetoxy, Di-(C₁-C₆)-alkylamino-(C₂-C₆-alkoxy) und Trifluormethyl, ausgewählt sind;


5 oder eines pharmazeutisch zulässigen Salzes davon zur Zubereitung eines Medikaments zum Reduzieren oder Beseitigen der Wirkungen ionisierender Strahlung auf normale Zellen in einem Patienten, der Aussetzung gegenüber ionisierender Strahlung erlitten hat oder Gefahr läuft, diese zu erleiden, oder in Knochenmarkzellen eines solchen Patienten.

10 **24.** Verwendung gemäß Anspruch 23, wobei die ionisierende Strahlung therapeutische ionisierende Strahlung zur Behandlung einer proliferativen Störung ist.

25. Verwendung gemäß Anspruch 24, wobei die proliferative Störung Krebs ist.

15 **26.** Ein Prozess zur Zubereitung einer Verbindung gemäß Anspruch 1, der das Reagierenlassen einer Verbindung der Formel Q₂-CH=CH-SO₂Cl mit einer Verbindung der Formel Q₁-NRH in einem nicht protischen Lösungsmittel in der Gegenwart einer Base, um eine Verbindung der folgenden Formel zu bilden, beinhaltet:

20




30

wobei:

Q₁, Q₂ und R gemäß Anspruch 1 definiert sind.

35 **27.** Ein Prozess zum Zubereiten einer Verbindung gemäß Anspruch 1, der das Reagierenlassen einer Verbindung folgender Formel:


40



50

mit einer Verbindung der Formel Q₂-C(O)H in einem nicht protischen Lösungsmittel in der Gegenwart einer Base, um eine Verbindung folgender Formel zu bilden, beinhaltet:

55




15 wobei:

20 Q₁, Q₂ und R gemäß Anspruch 1 definiert sind.

25 **Revendications**

30 1. Un composé de la formule I :




45 où :

Q₁ est sélectionné dans le groupe consistant en aryle substitué et non substitué, et hétéroaryle substitué et non substitué ;

Q₂ est sélectionné dans le groupe consistant en aryle substitué, et hétéroaryle substitué et non substitué ;

R est sélectionné dans le groupe consistant en hydrogène et alkyle(C₁-C₆) ;

où les substituants pour les groupes aryle substitué et hétéroaryle substitué comprenant Q₁ sont indépendamment sélectionnés dans le groupe consistant en halogène, alkyle(C₁-C₆), alcoxy(C₁-C₆), cyano, carboxy, hydroxy, hydroxyalkyle(C₂-C₆), phosphonato, amino, acyl(C₁-C₆)amino, sulfamyle, acétoxy, dialkyl(C₁-C₆)aminoalcoxy(C₂-C₆), trifluorométhyle et



où :

X est de l'oxygène ou du soufre,

R₅ est sélectionné dans le groupe consistant en hydrogène, alkyle(C₁-C₆), hétéroalkyle(C₂-C₆), phényle substitué, et phényle non substitué, et

R₆ est sélectionné dans le groupe consistant en hydrogène, alkyle(C₁-C₆), hétéroalkyle(C₂-C₆), aryle substitué, aryle non substitué, hétéroaryle substitué, hétéroaryle non substitué, arylalkyle(C₁-C₃) substitué, aryl-alkyle (C₁-C₃) non substitué et alcoxy(C₁-C₆)carbonylalkylényle(C₁-C₆) ;

où les substituants pour les groupes aryle substitué et hétéroaryle substitué comprenant Q₂, et les substituants pour les groupes aryle substitué et hétéroaryle substitué comprenant ou inclus dans R₅ et R₆, sont indépendamment sélectionnés dans le groupe consistant en halogène, alkyle(C₁-C₆), alcoxy(C₁-C₆), nitro, cyano, carboxy, carboxyalcoxy(C₁-C₃), hydroxy, hydroxyalkyle(C₂-C₆), phosphonato, amino, acyl(C₁-C₆)amino, sulfamyle, acétoxy, dialkyl(C₁-C₆)amino(alcoxy C₂-C₆) et trifluorométhyle ; à condition que lorsque R est de l'hydrogène :

(a) lorsque Q₁ est du phényle non substitué, Q₂ est autre que du diméthoxyphényle, 2-méthylphényle, 2-chlorophényle, 4-chlorophényle, 4-N,N-diméthylaminophényle, 4-méthylphényle, 4-méthoxyphényle, 4-nitrophényle, 3-méthoxy-4-hydroxyphényle, pyrényle non substitué, benzodioxolyle non substitué, et 2-thiényle non substitué ;

(b) lorsque Q₁ est du 3-hydroxyphényle, Q₂ est autre que du nitrophényle ;

(c) lorsque Q₁ est du 2-méthyl-5-hydroxyphényle, Q₂ est autre que du 4-nitrophényle ; et

(d) lorsque Q₁ est du 2-pyridyle non substitué, Q₂ est autre que du 3-méthoxy-4-hydroxyphényle ;

ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.

2. Un composé selon la revendication 1, à condition que :

(a) lorsque Q₁ est du phényle non substitué, Q₂ est autre que du dialcoxyphényle, 2-alkylphényle, 2-halophényle, 4-halophényle, 4-N,N-dialkylaminophényle, 4-alkylphényle, 4-alcoxyphényle, 4-nitrophényle, 3-alcoxy-4-hydroxyphényle, phényle non substitué, pyrényle non substitué, benzodioxolyle non substitué, 1-naphthyle non substitué et 2-thiényle non substitué ;

(b) lorsque Q₁ est du 3-hydroxyphényle, Q₂ est autre que du nitrophényle ;

(c) lorsque Q₁ est du 2-méthyl-5-hydroxyphényle, Q₂ est autre que du 4-nitrophényle ; et

(d) lorsque Q₁ est du 2-pyridyle non substitué, Q₂ est autre que du 3-méthoxy-4-hydroxyphényle ;

ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.

3. Un composé selon la revendication 2, à condition que :

lorsque R est de l'hydrogène :

(i) Q₁ peut ne pas être du dinitrophényle ;

(ii) Q₂ peut ne pas être du dinitrophényle ; et


(iii) lorsque Q₂ est du mononitrophényle :

Q₁ est autre que du phényle substitué, ou

Q₁ est du phényle substitué où au moins la position 4 est substituée, et le substituant est autre que de l'hydroxy ;

ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.

4. Un composé de la formule I, destiné à être utilisé en médecine :



I

où :

Q_1 est sélectionné dans le groupe consistant en aryle substitué et non substitué, et hétéroaryle substitué et non substitué ;

Q_2 est sélectionné dans le groupe consistant en aryle substitué, et hétéroaryle substitué et non substitué ;

R est sélectionné dans le groupe consistant en hydrogène et alkyle(C_1-C_6) ;

où les substituants pour les groupes aryle substitué et hétéroaryle substitué comprenant Q_1 sont indépendamment sélectionnés dans le groupe consistant en halogène, alkyle(C_1-C_6), alcoxy(C_1-C_6), cyano, carboxy, hydroxy, hydroxyalkyle(C_2-C_6), phosphonato, amino, acyl(C_1-C_6)amino, sulfamyle, acétoxy, dialkyl(C_1-C_6)aminoalcoxy(C_2-C_6), trifluorométhyle et



où :

X est de l'oxygène ou du soufre,

R_5 est sélectionné dans le groupe consistant en hydrogène, alkyle(C_1-C_6), hétéroalkyle(C_2-C_6), phényle substitué, et phényle non substitué, et

R_6 est sélectionné dans le groupe consistant en hydrogène, alkyle(C_1-C_6), hétéroalkyle(C_2-C_6), aryle substitué, aryle non substitué, hétéroaryle substitué, hétéroaryle non substitué, arylalkyle(C_1-C_3) substitué, aryl-alkyle(C_1-C_3) non substitué et alcoxy(C_1-C_6)carbonylalkylényle(C_1-C_6) ;

où les substituants pour les groupes aryle substitué et hétéroaryle substitué comprenant Q_2 , et les substituants pour les groupes aryle substitué et hétéroaryle substitué comprenant ou inclus dans R_5 et R_6 , sont indépendamment sélectionnés dans le groupe consistant en halogène, alkyle(C_1-C_6), alcoxy(C_1-C_6), nitro, cyano, carboxy, carboxyalcoxy(C_1-C_3), hydroxy, hydroxyalkyle(C_2-C_6), phosphonato, amino, acyl(C_1-C_6)amino, sulfamyle, acétoxy, dialkyl(C_1-C_6)amino(alcoxy C_2-C_6) et trifluorométhyle ;

à condition que, lorsque R est de l'hydrogène et Q_2 est du phényle non substitué, alors Q_1 doit être autre que du dihalophényle ;

ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.

5. Un composé selon la revendication 1 ou la revendication 4 où Q_1 est de manière facultative du phényle substitué et Q_2 est du phényle substitué, ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.
6. Un composé selon la revendication 5 où au moins soit Q_1 , soit Q_2 est substitué à la position 4 au moins, ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.
7. Un composé selon la revendication 6 où les substituants sont indépendamment sélectionnés dans le groupe consistant en halogène, alkyle(C_1-C_6), alcoxy(C_1-C_6), nitro, hydroxy et sulfamyle, ou un sel acceptable d'un point de

vue pharmaceutique de celui-ci.

8. Un composé selon la revendication 7 où le composé est :

- 5 (E)-4-méthoxystyryl-N-méthyl-N-phényl sulfonamide,
 (E)-4-chlorostyryl-N-3-chlorophényl sulfonamide,
 (E)-4-chlorostyryl-N-2-chlorophényl sulfonamide,

ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.

9. Un composé selon la revendication 8 où Q₁ et Q₂ sont tous deux indépendamment substitués à la position 4 au moins, ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.


10. Un composé selon la revendication 9 où les substituants sont indépendamment sélectionnés dans le groupe consistant en halogène, alkyle(C₁-C₆), alcoxy(C₁-C₆), nitro, hydroxy et sulfamyle, ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.

11. Un composé selon la revendication 10 où le composé est :

- 20 (E)-4-méthoxystyryl-N-4-sulfamylphényl sulfonamide,
 (E)-4-chlorostyryl-N-3-fluoro-4-méthoxyphényl sulfonamide,
 (E)-4-chlorostyryl-N-4-fluorophényl sulfonamide,
 (E)-4-méthoxystyryl-N-4-fluorophényl sulfonamide,
 (E)-4-méthoxystyryl-N-3-fluoro-4-méthoxyphényl sulfonamide,
 25 (E)-4-méthoxystyryl-N-4-fluorophényl sulfonamide,
 (E)-4-fluorostyryl-N-4-chlorophényl sulfonamide,
 (E)-4-méthoxystyryl-N-2,4,6-triméthoxyphényl sulfonamide,

ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.

12. Un composé selon la revendication 6 de la formule :



II


où R est de l'hydrogène ou de l'alkyle(C₁-C₆) ; R₁ est sélectionné dans le groupe consistant en halogène, alkyle (C₁ C₆), alcoxy(C₁-C₆), cyano, carboxy, hydroxy, hydroxyalkyle(C₂-C₆), phosphonato, amino, acyl(C₁-C₆)amino, sulfamyle, acétoxy, dialkyl(C₁-C₆)aminoalcoxy(C₂-C₆) et trifluorométhyle ; et R₂, R₃ et R₄ sont indépendamment sélectionnés parmi l'alcoxy(C₁-C₆), ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.

13. Un composé selon la revendication 12 de la formule :

5

10

15



IIa

20

où R, R₁, R₂, R₃ et R₄ sont définis tel que dans la revendication 12, ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.

25

14. Un composé selon la revendication 13 où le composé est (E)-2,4,6-triméthoxystyryl-N-4-méthoxyphényl sulfonamide, ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.

15. Un composé selon la revendication 5 où au moins soit Q₁, soit Q₂ est penta-substitué par de l'halogène, ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.

30

16. Un composé selon la revendication 15 où au moins soit Q₁, soit Q₂ est du pentafluorophényle, ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.

17. Un composé selon la revendication 16 où le composé est :

35

(E)-4-méthoxystyryl-N-2,3,4,5,6-pentafluorophényl sulfonamide,
(E)-4-fluorostyryl-N-2,3,4,5,6-pentafluorophényl sulfonamide,

ou un sel acceptable d'un point de vue pharmaceutique de celui-ci.

40


18. Une composition pharmaceutique comprenant un support acceptable d'un point de vue pharmaceutique et au moins un composé, ou sel acceptable d'un point de vue pharmaceutique de celui-ci, selon n'importe quelle revendication précédente.

45

19. Utilisation d'un composé selon la formule :

50

55



I


où :

Q_1 est sélectionné dans le groupe consistant en aryle substitué et non substitué, et hétéroaryle substitué et non substitué ;

Q_2 est sélectionné dans le groupe consistant en aryle substitué, et hétéroaryle substitué et non substitué ;

R est sélectionné dans le groupe consistant en hydrogène et alkyle(C_1-C_6) ;

où les substituants pour les groupes aryle substitué et hétéroaryle substitué comprenant Q_1 sont indépendamment sélectionnés dans le groupe consistant en halogène, alkyle(C_1-C_6), alcoxy(C_1-C_6), cyano, carboxy, hydroxy, hydroxyalkyle(C_2-C_6), phosphonato, amino, acyl(C_1-C_6)amino, sulfamyle, acétoxy, dialkyl(C_1-C_6)aminoalcoxy(C_2-C_6), trifluorométhyle et



où :

X est de l'oxygène ou du soufre,

R_5 est sélectionné dans le groupe consistant en hydrogène, alkyle(C_1-C_6), hétéroalkyle(C_2-C_6), phényle substitué, et phényle non substitué, et

R_6 est sélectionné dans le groupe consistant en hydrogène, alkyle(C_1-C_6), hétéroalkyle(C_2-C_6), aryle substitué, aryle non substitué, hétéroaryle substitué, hétéroaryle non substitué, arylalkyle(C_1-C_3) substitué, aryl-alkyle(C_1-C_3) non substitué et alcoxy(C_1-C_6)carbonylalkylényle(C_1-C_6) ;

où les substituants pour les groupes aryle substitué et hétéroaryle substitué comprenant Q_2 , et les substituants pour les groupes aryle substitué et hétéroaryle substitué comprenant ou inclus dans R_5 et R_6 , sont indépendamment sélectionnés dans le groupe consistant en halogène, alkyle(C_1-C_6), alcoxy(C_1-C_6), nitro, cyano, carboxy, carboxyalcoxy(C_1-C_3), hydroxy, hydroxyalkyle(C_2-C_6), phosphonato, amino, acyl(C_1-C_6)amino, sulfamyle, acétoxy, dialkyl(C_1-C_6)amino(alcoxy C_2-C_6) et trifluorométhyle ;

ou un sel acceptable d'un point de vue pharmaceutique de celui-ci ;


pour la préparation d'un médicament destiné au traitement d'un trouble prolifératif.

20. Une utilisation selon la revendication 19 où le trouble prolifératif est sélectionné dans le groupe consistant en cancer, hémangiomatose néonatale, sclérose en plaques progressive secondaire, trouble myélo-dégénératif progressif chronique, neurofibromatose, ganglioneuromatose, formation de chéloïdes, maladie osseuse de Paget, maladie fibrokystique du sein, fibrose de La Peyronie, fibrose de Dupuytren, resténose et cirrhose.

21. Une utilisation selon la revendication 20 où le trouble prolifératif est le cancer, et le cancer est sélectionné dans le groupe consistant en cancer de l'ovaire, cancer du sein, cancer de la prostate, cancer du poumon, cancer du rein, cancer colorectal et cancer du cerveau, ou le cancer est une leucémie.

22. Une utilisation selon la revendication 20 où le trouble prolifératif est le cancer, destinée à induire l'apoptose des cellules tumorales.

23. Utilisation d'un composé selon la formule



I


où :

Q₁ est sélectionné dans le groupe consistant en aryle substitué et non substitué, et hétéroaryle substitué et non substitué ;

Q₂ est sélectionné dans le groupe consistant en aryle substitué, et hétéroaryle substitué et non substitué ;

R est sélectionné dans le groupe consistant en hydrogène et alkyle(C₁-C₆) ;

où les substituants pour les groupes aryle substitué et hétéroaryle substitué comprenant Q₁ sont indépendamment sélectionnés dans le groupe consistant en halogène, alkyle(C₁-C₆), alcoxy(C₁-C₆), cyano, carboxy, hydroxy, hydroxyalkyle(C₂-C₆), phosphonato, amino, acyl(C₁-C₆)amino, sulfamyle, acétoxy, dialkyl(C₁-C₆)aminoalcoxy(C₂-C₆), trifluorométhyle et



où :

X est de l'oxygène ou du soufre,

R₅ est sélectionné dans le groupe consistant en hydrogène, alkyle(C₁-C₆), hétéroalkyle(C₂-C₆), phényle substitué, et phényle non substitué, et

R₆ est sélectionné dans le groupe consistant en hydrogène, alkyle(C₁-C₆), hétéroalkyle(C₂-C₆), aryle substitué, aryle non substitué, hétéroaryle substitué, hétéroaryle non substitué, arylalkyle(C₁-C₃) substitué, aryl-alkyle(C₁-C₃) non substitué et alcoxy(C₁-C₆)carbonylalkylényle(C₁-C₆) ;

où les substituants pour les groupes aryle substitué et hétéroaryle substitué comprenant Q₂, et les substituants pour les groupes aryle substitué et hétéroaryle substitué comprenant ou inclus dans R₅ et R₆, sont indépendamment sélectionnés dans le groupe consistant en halogène, alkyle(C₁-C₆), alcoxy(C₁-C₆), nitro, cyano, carboxy, carboxyalcoxy(C₁-C₃), hydroxy, hydroxyalkyle(C₂-C₆), phosphonato, amino, acyl(C₁-C₆)amino, sulfamyle, acétoxy, dialkyl(C₁-C₆)amino(alcoxy C₂-C₆), et trifluorométhyle ;

ou un sel acceptable d'un point de vue pharmaceutique de celui-ci ;

pour la préparation d'un médicament destiné à réduire ou éliminer les effets du rayonnement ionisant sur des cellules normales chez un sujet qui a subi ou qui risque de subir une exposition à un rayonnement ionisant, ou dans des cellules de moelle osseuse provenant d'un tel sujet.

24. Une utilisation selon la revendication 23, où le rayonnement ionisant est un rayonnement ionisant thérapeutique destiné au traitement d'un trouble prolifératif.


25. Une utilisation selon la revendication 24 où le trouble prolifératif est le cancer.

26. Un processus destiné à préparer un composé selon la revendication 1, comprenant faire entrer en réaction un composé de la formule Q₂-CH=CH-SO₂Cl avec un composé de la formule Q₁-NRH dans un solvant non protique

en présence d'une base afin de former un composé de la formule :

5

10



15

I

où :


20

Q₁, Q₂ et R sont définis tel que dans la revendication 1.

27. Un processus destiné à préparer un composé selon la revendication 1, comprenant faire entrer en réaction un composé de la formule

25

30




35

avec un composé de la formule Q₂-C(O)H dans un solvant non protique en présence d'une base afin de former un composé de la formule :

40

45



50

I

où :

55

Q₁, Q₂ et R sont définis tel que dans la revendication 1.




Figure 1




Figure 2

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 4035421 A, Snyder, Jr. [0003]
- WO 9524190 A [0004]
- DE 2118493 [0005]

Non-patent literature cited in the description

- **Bordwell et al.** *J. Amer. Chem. Soc.*, 1946, vol. 68, 139 [0002]
- **Oliver et al.** *Synthesis*, 1975, 321-322 [0002]
- **Waldau et al.** *Angew. Chem., Int. Ed. Engl.*, 1972, vol. 11 (9), 826-8 [0005]
- **Aswarthamma et al.** *Chimica Acta Turcica*, 1996, vol. 24, 7-10 [0006]
- **Touarti et al.** *J. Soc. Alger. Chim.*, 1996, vol. 6 (1), 39-52 [0006]
- **Lazarus et al.** *J. of Hematotherapy*, 1993, vol. 2 (4), 457-66 [0015]
- **Neta et al.** *Semin. Radiat. Oncol.*, 1996, vol. 6, 306-320 [0026]
- **Maisin, J.R. ; Alexander Award.** Chemical radio-protection: past, present, and future prospects. *Int J. Radiat Biol.*, 1998, vol. 73, 443-50 [0026]
- Section E: Stereochemistry, in *Nomenclature of Organic Chemistry*. John Wiley & Sons, Inc, 1974, 127-138 [0090]
- **Oliver et al.** *Synthesis*, May 1975, 321-322 [0093]
- Remington's Pharmaceutical Sciences. Mack Publishing Co, 1990 [0132]
- **Latham et al.** *Oncogene*, 1996, vol. 12, 827-837 [0154]
- **Patel et al.** *FASEB*, 1996, vol. 10, 587-597 [0158]
- **Nicholson et al.** *Nature*, 1995, vol. 376 (6533), 37-43 [0158]
- **Lippke et al.** *J. Biol. Chemistry*, 1996, vol. 271, 1825 [0158]
- **Gewirtz et al.** *Science*, 1988, vol. 242, 1303-1306 [0168]