

(19)



(11)

EP 1 409 016 B9

(12)

CORRECTED NEW EUROPEAN PATENT SPECIFICATION

(15) Correction information:

Corrected version no 1 (W1 B2)
Corrections, see
Claims EN 1

(51) Int Cl.:

A61K 39/395 ^(2006.01) **A61K 39/00** ^(2006.01)
G01N 33/53 ^(2006.01) **C07K 16/28** ^(2006.01)
A61P 19/08 ^(2006.01)

(48) Corrigendum issued on:

28.10.2020 Bulletin 2020/44

(86) International application number:

PCT/US2002/020181

(45) Date of publication and mention of the opposition decision:

15.07.2020 Bulletin 2020/29

(87) International publication number:

WO 2003/002713 (09.01.2003 Gazette 2003/02)

(45) Mention of the grant of the patent:

14.04.2010 Bulletin 2010/15

(21) Application number: **02749660.3**

(22) Date of filing: **25.06.2002**

(54) **ANTIBODIES TO OPGL**

ANTIKÖRPER GEGEN OPGL

ANTICORPS OPGL

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**

- **DAVIS, Geoffrey, C.**
Burlingame, CA 94010 (US)

(30) Priority: **26.06.2001 US 301172 P**

(74) Representative: **Dörries, Hans Ulrich et al**
df-mp Dörries Frank-Molnia & Pohlman
Patentanwälte Rechtsanwälte PartG mbB
Theatinerstrasse 16
80333 München (DE)

(43) Date of publication of application:

21.04.2004 Bulletin 2004/17

(60) Divisional application:

09156995.4 / 2 087 908
10181323.6 / 2 270 052
10185344.8 / 2 295 081
18202340.8 / 3 492 100

(56) References cited:

WO-A-00/15807 **WO-A1-01/62932**
WO-A1-98/46751 **WO-A2-00/37504**
WO-A2-02/053596 **WO-A2-03/086289**
US-A- 5 886 152 **US-A- 5 891 438**
US-A- 6 015 938 **US-A1- 2002 015 055**

(73) Proprietors:

- **Amgen Fremont Inc.**
Thousand Oaks, CA 91320-1799 (US)
- **Amgen Inc.**
Thousand Oaks, CA 91320-1799 (US)

- **SHEPHERD P.; DEAN C.:** 'Monoclonal Antibodies: A Practical Approach', 2000, OXFORD UNIVERSITY PRESS * page 58 - page 65 *
- **TSUKII K. ET AL:** 'Osteoclast differentiation factor mediates an essential signal for bone resorption induced by 1 alpha,25-dihydroxyvitamin D3, prostaglandin E2, or parathyroid hormone in the microenvironment of bone' BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS vol. 246, no. 2, 19 May 1998, pages 337 - 341, XP002169940

(72) Inventors:

- **BOYLE, William, J.**
Malibu, CA 90265 (US)
- **MARTIN, Francis, H.**
Newbury Park, CA 91320 (US)
- **CORVALAN, Jose, R.**
Foster City, CA 94404 (US)

EP 1 409 016 B9

- NAGAI M.; KYAKUMOTO S.; SATO N.: 'Cancer cells responsible for humoral hypercalcemia express mRNA encoding a secreted form of ODF/TRANCE that induces osteoclast formation' **BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS** vol. 269, 16 March 2000, pages 532 - 536, XP002956545
- OYAJOBIB.O. ET AL: 'Therapeutic efficacy of a soluble receptor activator of nuclear factor kappaB-IgG Fc fusion protein in suppressing bone resorption and hypercalcemia in a model of humoral hypercalcemia of malignancy' **CANCER RESEARCH** vol. 61, no. 6, 15 March 2001, pages 2572 - 2578, XP002301285
- KINPARA K. ET AL: 'Osteoclast differentiation factor in human osteosarcoma cell line' **JOURNAL OF IMMUNOASSAY** vol. 21, no. 4, 2000, pages 327 - 340, XP002956542
- DE SOUZA ET AL: 'Effect of platelet-activating factor antagonists (BN-52021, WEB-2170, and BB-882) on bacterial translocation in acute pancreatitis' **JOURNAL OF GASTROINTESTINAL SURGERY** vol. 5, no. 4, 2001, pages 364 - 370
- VAN DER VELDEN ET AL: 'Glucocorticoids: mechanisms of action and anti-inflammatory potential in asthma' **JOURNAL OF GASTROINTESTINAL SURGERY** vol. 5, no. 4, 2001, pages 229 - 237
- Thomas R.J. et al: "Breast Cancer Cells Interact With Osteoblasts to Support Osteoclast Formation", **Endocrinology** ., vol. 140, no. 10, 1 October 1999 (1999-10-01), pages 4451-4458,
- KINPARA K. ET AL.: "Osteoclast differentiation factor in human osteosarcoma cell line", **JOURNAL OF IMMUNOASSAY**., vol. 21, no. 4, 1 January 2000 (2000-01-01), pages 327-340, CH ISSN: 0197-1522
- Jean-Jacques Body et al: "A Study of the Biological Receptor Activator of Nuclear factor-kappaB Ligand Inhibitor, Denosumab, in Patients With Multiple Myeloma or Bone Metastases From Breast Cancer", **Clin Cancer Res** ., vol. 12, no. 4, 15 February 2006 (2006-02-15), pages 1221-1228,
- Daniel Schramek et al: "Osteoclast differentiation factor RANKL controls development of progesterin-driven mammary cancer.", **Nature**, vol. 468, no. 7320, 4 November 2010 (2010-11-04), pages 98-102, London ISSN: 0028-0836, DOI: 10.1038/NATURE09387
- LACEY D.L. ET AL.: "Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation", **CELL**, vol. 93., no. 02., 17 April 1998 (1998-04-17), pages 165-176., AMSTERDAM, NL ISSN: 0092-8674, DOI: 10.1016/S0092-8674(00)81569-X
- Clarke S.H. et al: "Inter- and intraclonal diversity in the antibody response to influenza hemagglutinin.", **J Exp Med**, vol. 161, no. 4, 1 January 1985 (1985-01-01), pages 687-704,
- GARDNEROVA M. et al: "THE USE OF TNF FAMILY LIGANDS AND RECEPTORS AND AGENTS WHICH MODIFY THEIR INTERACTION AS THERAPEUTIC AGENTS", **CURRENT DRUG TARGETS**, vol. 01, no. 04, 1 December 2000 (2000-12-01), pages 327-364, US ISSN: 1389-4501, DOI: 10.2174/1389450003349092
- NAGAI MASAZUMI et al: "Cancer cells responsible for humoral hypercalcemia express mRNA encoding a secreted form of ODF/TRANCE that induces osteoclast formation", **Biochemical and Biophysical Research Communications**NL, vol. 269, no. 2, 16 March 2000 (2000-03-16), pages 532-536, AMSTERDAM, NL ISSN: 0006-291X, DOI: 10.1006/bbrc.2000.2314
- OYAJOBIB.O. ET AL: "Therapeutic efficacy of a soluble receptor activator of nuclear factor kappaB-IgG Fc fusion protein in suppressing bone resorption and hypercalcemia in a model of humoral hypercalcemia of malignancy.", **Cancer research**, vol. 61, no. 6, 15 March 2001 (2001-03-15) , pages 2572-2578, US ISSN: 0008-5472
- Kostenuik P.J.: "Osteoprotegerin and RANKL regulate bone resorption, density, geometry and strength", **CURRENT OPINION IN PHARMACOLOGY**, vol. 5, no. 6, 1 December 2005 (2005-12-01), pages 618-625, NL ISSN: 1471-4892, DOI: 10.1016/j.coph.2005.06.005
- Record (CN) 615258-40-7, for "Denosumab", Database CAS Registry, (pub. Am. Chem. Soc., 2011), retrieved via STN, June 24, 2011

Description

FIELD OF THE INVENTION

5 **[0001]** The present invention relates to antibodies that bind osteoprotegerin ligand (OPGL). Compositions and methods for the treatment of bone diseases, such as osteoporosis, bone loss from arthritis, Paget's disease, and osteopenia, are also described.

BACKGROUND OF THE INVENTION

10 **[0002]** Bone tissue provides support for the body and includes mineral (including calcium and phosphorous), a matrix of collagenous and noncollagenous proteins, and cells. Living bone tissue exhibits a dynamic equilibrium between formation of bone, which is called deposition, and breakdown of bone, which is called resorption. Three types of cells found in bone, osteocytes, osteoblasts and osteoclasts, are involved in this equilibrium. Osteoblasts promote formation of bone tissue whereas osteoclasts are associated with resorption. Resorption, or the dissolution of bone matrix and mineral, is a fast and efficient process compared to bone formation and can release large amounts of mineral from bone. Osteoclasts are involved in the regulation of the normal remodeling of skeletal tissue and in resorption induced by hormones. For instance, resorption is stimulated by the secretion of parathyroid hormone in response to decreasing concentrations of calcium ion in extracellular fluids. In contrast, inhibition of resorption is a function of calcitonin. In addition, metabolites of vitamin D alter the responsiveness of bone to parathyroid hormone and calcitonin.

20 **[0003]** Osteoprotegerin ligand (OPGL), which is a member of the TNF family of cytokines, promotes formation of osteoclasts through binding to the receptor activator of NF- κ B (RANK, also called osteoclast differentiation and activation receptor, or ODAR). Osteoprotegerin (OPG), on the other hand, inhibits the formation of osteoclasts by sequestering OPGL and preventing OPGL association with ODAR. Thus, the amount of OPGL associated with ODAR correlates with the equilibrium between bone deposition and resorption.

25 **[0004]** After skeletal maturity, the amount of bone in the skeleton reflects the balance (or imbalance) of bone formation and bone resorption. Peak bone mass occurs after skeletal maturity prior to the fourth decade. Between the fourth and fifth decades, the equilibrium shifts and bone resorption dominates. The inevitable decrease in bone mass with advancing years starts earlier in females than males and is distinctly accelerated after menopause in some females (principally those of Caucasian and Asian descent).

30 **[0005]** Osteopenia is a condition relating generally to any decrease in bone mass to below normal levels. Such a condition may arise from a decrease in the rate of bone synthesis or an increase in the rate of bone destruction or both. A common form of osteopenia is primary osteoporosis, also referred to as postmenopausal and senile osteoporosis. This form of osteoporosis is a consequence of the universal loss of bone with age and is often a result of increase in bone resorption with a normal rate of bone formation. Many white females in the United States develop symptomatic osteoporosis. A direct relationship exists between osteoporosis and the incidence of hip, femoral, neck and inter-trochanteric fracture in women 45 years and older. Elderly males may develop symptomatic osteoporosis between the ages of 50 and 70. Osteoporosis may, in certain instances, result from increased levels or activity of OPGL. Thus, it would be useful to have molecules that can regulate the activity of OPGL in osteoclastogenesis.

35 **[0006]** Several factors have been identified which may contribute to postmenopausal and senile osteoporosis. They include alteration in hormone levels accompanying aging and inadequate calcium consumption attributed to decreased intestinal absorption of calcium and other minerals. Certain treatments have included hormone therapy or dietary supplements in an attempt to retard the process. More recently, anti-resorptive agents such as bisphosphonates and selective estrogen receptor modifiers (SERMs) have emerged for the prevention and treatment of reduced bone mass. Thus, it may be useful to combine those treatments with molecules that can regulate the activity of OPGL in treating certain osteopenic disorders.

40 **[0007]** Tsukii et al. (Biochemical and Biophysical Research Communications (1998), vol. 246, pages 336-341) disclose a polyclonal antibody directed against murine ODF (= OPGL, RAHKL) which is effective in inhibiting bone resorption *in vitro*.

SUMMARY OF THE INVENTION

45 **[0008]** The present invention is defined in the claims.

50 **[0009]** Described herein is an antibody, comprising a heavy chain and a light chain, wherein the heavy chain comprises an amino acid sequence as set forth in SEQ ID NO: 2, and the light chain comprises an amino acid sequence as set forth in SEQ ID NO: 4.

[0010] In certain embodiments, the invention provides for an antibody, comprising a heavy chain and a light chain, wherein the heavy chain comprises a variable region comprising an amino acid sequence as set forth in SEQ ID NO:

13, and wherein the light chain comprises a variable region comprising an amino acid sequence as set forth in SEQ ID NO: 14.

[0011] The disclosure provides for an antibody comprising a heavy chain and a light chain, wherein the heavy chain comprises an amino acid sequence as set forth in SEQ ID NO: 2 or a fragment thereof.

[0012] In certain embodiments, the disclosure provides for an antibody comprising a heavy chain and a light chain, wherein the heavy chain comprises a variable region comprising an amino acid sequence as set forth in SEQ ID NO: 13 or a fragment thereof

[0013] The disclosure provides for an antibody comprising a heavy chain and a light chain, wherein the light chain comprises an amino acid sequence as set forth in SEQ ID NO: 4 or a fragment thereof.

[0014] In certain embodiments, the disclosure provides for an antibody comprising a heavy chain and a light chain, wherein the light chain comprises a variable region comprising an amino acid sequence as set forth in SEQ ID NO: 14 or a fragment thereof.

[0015] In certain embodiments, the disclosure provides for an antibody, comprising a heavy chain and a light chain, (a) wherein the heavy chain comprises a first variable region, and wherein the first variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 13, and (b) wherein the light chain comprises a second variable region, and wherein the second variable region comprises a sequence that has at least 90% identity to the amino acid sequence set forth in SEQ ID NO: 14, and (c) wherein the antibody interacts with an osteoprotegerin ligand (OPGL).

[0016] In certain embodiments, the first variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 13, and the second variable region comprises a sequence that has at least 95% identity to the amino acid sequence set forth in SEQ ID NO: 14.

[0017] In certain embodiments, the first variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 13, and the second variable region comprises a sequence that has at least 99% identity to the amino acid sequence set forth in SEQ ID NO: 14.

[0018] The disclosure provides for a heavy chain, comprising an amino acid sequence as set forth in SEQ ID NO:2 or a fragment thereof. In certain embodiments, the invention provides for a heavy chain comprising an amino acid sequence as set forth in SEQ ID NO: 13.

[0019] The disclosure provides for a light chain, comprising an amino acid sequence as set forth in SEQ ID NO:4 or a fragment thereof. In certain embodiments, the invention provides for a light chain comprising an amino acid sequence as set forth In SEQ ID NO: 14.

[0020] In certain embodiments of the disclosure, single chain antibodies are provided. In certain embodiments of the invention, single chain Fv antibodies are provided. In certain embodiments of the invention, Fab antibodies are provided. In certain embodiments of the invention, Fab' antibodies are provided. In certain embodiments of the invention, (Fab')₂ antibodies are provided.

[0021] In certain embodiments, a pharmaceutical composition comprising an antibody of the invention is provided. In certain embodiments, the pharmaceutical composition may comprise a therapeutically effective amount of an antibody to OPGL.

[0022] In certain embodiments, the pharmaceutical composition may comprise an antibody to OPGL and at least one therapeutic agent selected from a bone morphogenic factor, transforming growth factor- β (TGF- β), an interleukin 1 (IL-1) inhibitor, IL-1ra, KineretTM, anakinra, a TNF α inhibitor, a soluble TNF α receptor, EnbrelTM, etanercept, an anti-TNF α antibody, RemicadeTM, infliximab, a D2E7 antibody, a parathyroid hormone, an analog of a parathyroid hormone, a parathyroid hormone related protein, an analog of a parathyroid hormone related protein, a prostaglandin, a bisphosphonate, an alendronate, fluoride, calcium, a non-steroidal anti-inflammatory drug (NSAID), a COX-2 inhibitor, CelebrexTM, celecoxib, VioxxTM, rofecoxib, an immunosuppressant, methotrexate, leflunomide, a serine protease inhibitor, a secretory leukocyte protease inhibitor (SLPI), an IL-6 inhibitor, an antibody to IL-6, an IL-8 inhibitor, an antibody to IL-8, an IL-18 inhibitor, an IL-18 binding protein, an IL-18 antibody, an interleukin-1 converting enzyme (ICE) modulator, a fibroblast growth factor (FGF), an FGF modulator, a PAF antagonist, a keratinocyte growth factor (KGF), a KGF-related molecule, a KGF modulator; a matrix metalloproteinase (MMP) modulator, a nitric oxide synthase (NOS) modulator; a modulator of glucocorticoid receptor, a modulator of glutamate receptor, a modulator of lipopolysaccharide (LPS) levels, a noradrenaline, a noradrenaline mimetic, and a noradrenaline modulator.

[0023] In certain aspects of the disclosure, the antibody of the invention is for a method of treating an osteopenic disorder, comprising administering a pharmaceutically effective amount of the antibody. In certain aspects of the disclosure, the pharmaceutical composition of the invention is for a method of treating an osteopenic disorder comprising administering the pharmaceutical composition.

[0024] In certain aspects of the disclosure, the pharmaceutical composition is for a method of treating an inflammatory condition with attendant bone loss in a patient comprising administering the pharmaceutical composition.

[0025] In certain aspects of the disclosure, the pharmaceutical composition is for a method of treating an autoimmune condition with attendant bone loss in a patient comprising administering the pharmaceutical composition.

[0026] In certain aspects of the disclosure, the pharmaceutical composition is for a method of treating rheumatoid arthritis in a patient, comprising administering the pharmaceutical composition of the invention.

[0027] In certain aspects of the disclosure, a method of detecting the level of OPGL in a biological sample is provided, comprising contacting the sample with an antibody of the invention.

5

BRIEF DESCRIPTION OF THE FIGURES

[0028]

- 10 Figure 1 shows a cDNA sequence encoding the α OPGL-1 antibody heavy chain (SEQ ID NO: 1)
 Figure 2 shows the amino acid sequence of the α OPGL-1 antibody heavy chain (SEQ ID NO: 2).
 Figure 3 shows a cDNA sequence encoding the α OPGL-1 antibody light chain (SEQ ID NO: 3).
 Figure 4 shows the amino acid sequence of the α OPGL-1 antibody light chain (SEQ ID NO: 4).
 Figure 5 shows a schematic diagram of the α OPGL-1 kappa light chain expression plasmid α OPGL-1-Kappa/
 15 pDSRa19.
 Figure 6 shows a schematic diagram of the α OPGL-1 IgG2 heavy chain expression plasmid, α OPGL-1-IgG2/pDSRa19.
 Figure 7 shows dose-dependent binding of α OPGL-1 to OPGL-coated EIA plates.
 Figure 8 shows specific binding of α OPGL-1 to membrane-bound OPGL.
 20 Figure 9 shows inhibition of α OPGL-1 binding to OPGL-coated EIA plates by soluble OPGL.
 Figure 10 shows specific binding of α OPGL-1 to OPGL-coated EIA plates.
 Figure 11 shows dose-dependent inhibition of osteoclast formation by α OPGL-1.
 Figure 12 shows dose-dependent inhibition of OPGL binding to ODAR by α OPGL-1.
 Figure 13 shows the mean serum concentration time profiles after administering a single dose of α OPGL-1 to
 25 Cynomolgus Monkeys.
 Figure 14 shows the mean percent change in serum N-Tx concentration after administering a single dose of α OPGL-1 to Cynomolgus Monkeys.
 Figure 15 shows the mean percent change in urine N-Tx concentration after administering a single dose of α OPGL-1 to Cynomolgus Monkeys.
 30 Figure 16 shows antibody positive and negative serum-concentration time profiles after administering a single dose of α OPGL-1 to Cynomolgus Monkeys.
 Figure 17 shows the amino acid sequence of the α OPGL-1 antibody heavy chain variable region (SEQ ID NO: 13).
 Figure 18 shows the amino acid sequence of the α OPGL-1 antibody light chain variable region (SEQ ID NO: 14).
 Figure 19 shows a cell culture process for production of α OPGL-1.
 35 Figure 20 shows the serum calcium percent change after administering a single dose of α OPGL-1 to Cynomolgus monkeys.
 Figure 21 shows the mean serum Alkaline Phosphatase percent change after administering a single dose of α OPGL-1 to Cynomolgus monkeys.

40 DETAILED DESCRIPTION OF CERTAIN PREFERRED EMBODIMENTS

[0029] The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described.

45 Definitions

[0030] Standard techniques may be used for recombinant DNA, oligonucleotide synthesis, and tissue culture and transformation (e.g., electroporation, lipofection). Enzymatic reactions and purification techniques may be performed according to manufacturer's specifications or as commonly accomplished in the art or as described herein. The foregoing
 50 techniques and procedures may be generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification. See e.g., Sambrook et al. *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)). Unless specific definitions are provided, the nomenclatures utilized in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal
 55 and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques may be used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

[0031] As utilized in accordance with the present disclosure, the following terms, unless otherwise indicated, shall be

understood to have the following meanings:

[0032] The term "isolated polynucleotide" as used herein shall mean a polynucleotide of genomic, cDNA, or synthetic origin or some combination thereof, which by virtue of its origin the "isolated polynucleotide" (1) is not associated with all or a portion of a polynucleotide in which the "isolated polynucleotide" is found in nature, (2) is linked to a polynucleotide which it is not linked to in nature, or (3) does not occur in nature as part of a larger sequence.

[0033] The term "isolated protein" referred to herein means a protein encoded by cDNA, recombinant RNA, or synthetic origin or some combination thereof, which (1) is free of at least some proteins with which it would normally be found, (2) is essentially free of other proteins from the same source, e.g., from the same species, (3) is expressed by a cell from a different species, or (4) does not occur in nature.

[0034] The term "polypeptide" is used herein as a generic term to refer to native proteins, or sequences that have deletions, additions, and/or substitutions of one or more amino acids of the native sequence. The term "polypeptide" also encompasses α OPGL-1 (as described below, SEQ ID NO: 2 and SEQ ID NO: 4), or sequences that have deletions, additions, and/or substitutions of one or more amino acid of α OPGL-1. According to certain embodiments, the invention comprises the human heavy chain immunoglobulin molecule represented by Figure 2 (SEQ ID NO: 2) and the human light chain immunoglobulin molecule represented by Figure 4 (SEQ ID NO: 4).

[0035] The term "naturally-occurring" as used herein as applied to an object refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be isolated from a source in nature and which has not been intentionally modified by man in the laboratory or otherwise is naturally-occurring.

[0036] The term "operably linked" as used herein refers to components that are in a relationship permitting them to function in their intended manner. For example, a control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

[0037] The term "control sequence" as used herein refers to polynucleotide sequences which may effect the expression and processing of coding sequences to which they are ligated. The nature of such control sequences may differ depending upon the host organism. According to certain embodiments, control sequences for prokaryotes may include promoter, ribosomal binding site, and transcription termination sequence. According to certain embodiments, control sequences for eukaryotes may include promoters and transcription termination sequence. In certain embodiments, "control sequences" can include leader sequences and/or fusion partner sequences.

[0038] The term "polynucleotide" as referred to herein means a polymeric form of nucleotides of at least 10 bases in length. In certain embodiments, the bases may be ribonucleotides or deoxyribonucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA.

[0039] The term "oligonucleotide" referred to herein includes naturally occurring, and modified nucleotides linked together by naturally occurring, and/or non-naturally occurring oligonucleotide linkages. Oligonucleotides are a polynucleotide subset generally comprising a length of 200 bases or fewer. In certain embodiments, oligonucleotides are 10 to 60 bases in length. In certain embodiments, oligonucleotides are 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 bases in length. Oligonucleotides may be single stranded or double stranded, e.g. for use in the construction of a gene mutant. Oligonucleotides of the invention may be sense or antisense oligonucleotides.

[0040] The term "naturally occurring nucleotides" includes deoxyribonucleotides and ribonucleotides. The term "modified nucleotides" includes nucleotides with modified or substituted sugar groups and the like. The term "oligonucleotide linkages" includes oligonucleotides linkages such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate, phosphoroamidate, and the like. See, e.g., LaPlanche et al. Nucl. Acids Res. 14:9081 (1986); Stec et al. J. Am. Chem. Soc. 106:6077 (1984); Stein et al. Nucl. Acids Res. 16:3209 (1988); Zon et al. Anti-Cancer Drug Design 6:539 (1991); Zon et al. Oligonucleotides and Analogues: A Practical Approach, pp. 87-108 (F. Eckstein, Ed., Oxford University Press, Oxford England (1991)); Stec et al. U.S. Pat. No. 5,151,510; Uhlmann and Peyman Chemical Reviews 90:543 (1990). An oligonucleotide can include a label for detection.

[0041] Identity and similarity of related and polypeptides can be readily calculated by known methods. Such methods include, but are not limited to, those described in Computational Molecular Biology, Lesk, A.M., ed., Oxford University Press, New York (1988); Biocomputing: informatics and Genome Projects, Smith, D.W., ed., Academic Press, New York (1993); Computer Analysis of Sequence Data, Part 1, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey (1994); Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press (1987); Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M. Stockton Press, New York (1991); and Carillo et al., SIAM J. Applied Math., 48:1073 (1988).

[0042] Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity are described in publicly available computer programs. Preferred computer program methods to determine identity between two sequences include, but are not limited to, the GCG program package, including GAP (Devereux et al., Nucl. Acid. Res., 12:387 (1984); Genetics Computer Group, University of Wisconsin, Madison, WI, BLASTP, BLASTN, and FASTA (Altschul et al., J. Mol. Biol., 215:403-410 (1990)). The BLASTX program is publicly available from the National Center for Biotechnology Information (NCBI) and other sources (BLAST *Manual*,

Altschul *et al.* NCB/NLM/NIH Bethesda, MD 20894; Altschul *et al.*, *supra* (1990)). The well-known Smith Waterman algorithm may also be used to determine identity.

[0043] Certain alignment schemes for aligning two amino acid sequences may result in the matching of only a short region of the two sequences, and this small aligned region may have very high sequence identity even though there is no significant relationship between the two full-length sequences. Accordingly, in certain embodiments, the selected alignment method (GAP program) will result in an alignment that spans at least 50 contiguous amino acids of the target polypeptide.

[0044] For example, using the computer algorithm GAP (Genetics Computer Group, University of Wisconsin, Madison, WI), two polypeptides for which the percent sequence identity is to be determined are aligned for optimal matching of their respective amino acids (the "matched span", as determined by the algorithm). In certain embodiments, a gap opening penalty (which is calculated as 3X the average diagonal; the "average diagonal" is the average of the diagonal of the comparison matrix being used; the "diagonal" is the score or number assigned to each perfect amino acid match by the particular comparison matrix) and a gap extension penalty (which is usually 1/10 times the gap opening penalty), as well as a comparison matrix such as PAM 250 or BLOSUM 62 are used in conjunction with the algorithm. In certain embodiments, a standard comparison matrix (see Dayhoff *et al.*, Atlas of Protein Sequence and Structure, 5(3)(1978) for the PAM 250 comparison matrix; Henikoff *et al.*, Proc. Natl. Acad. Sci USA, 89:10915-10919 (1992) for the BLOSUM 62 comparison matrix) is also used by the algorithm.

[0045] In certain embodiments, the parameters for a polypeptide sequence comparison include the following:

Algorithm: Needleman *et al.*, J. Mol. Biol., 48:443-453 (1970);
 Comparison matrix: BLOSUM 62 from Henikoff *et al.*, *supra* (1992);
 Gap Penalty: 12
 Gap Length Penalty: 4
 Threshold of Similarity: 0

[0046] The GAP program may be useful with the above parameters. In certain embodiments, the aforementioned parameters are the default parameters for polypeptide comparisons (along with no penalty for end gaps) using the GAP algorithm.

[0047] As used herein, the twenty conventional amino acids and their abbreviations follow conventional usage. See Immunology-A Synthesis (2nd Edition, E. S. Golub and D. R. Gren, Eds., Sinauer Associates, Sunderland, Mass. (1991)). Stereoisomers (e.g., D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as α -, α -disubstituted amino acids, N-alkyl amino acids, lactic acid, and other unconventional amino acids may also be suitable components for polypeptides of the present invention. Examples of unconventional amino acids include: 4-hydroxyproline, γ -carboxyglutamate, ϵ -N,N,N-trimethyllysine, ϵ -N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, σ -N-methylarginine, and other similar amino acids and imino acids (e.g., 4-hydroxyproline). In the polypeptide notation used herein, the left-hand direction is the amino terminal direction and the right-hand direction is the carboxy-terminal direction, in accordance with standard usage and convention.

[0048] Similarly, unless specified otherwise, the left-hand end of single-stranded polynucleotide sequences is the 5' end; the left-hand direction of double-stranded polynucleotide sequences is referred to as the 5' direction. The direction of 5' to 3' addition of nascent RNA transcripts is referred to as the transcription direction; sequence regions on the DNA strand having the same sequence as the RNA and which are 5' to the 5' end of the RNA transcript are referred to as "upstream sequences"; sequence regions on the DNA strand having the same sequence as the RNA and which are 3' to the 3' end of the RNA transcript are referred to as "downstream sequences".

[0049] Conservative amino acid substitutions may encompass non-naturally occurring amino acid residues, which are typically incorporated by chemical peptide synthesis rather than by synthesis in biological systems. These include peptidomimetics and other reversed or inverted forms of amino acid moieties.

[0050] Naturally occurring residues may be divided into classes based on common side chain properties:

- 1) hydrophobic: norleucine, Met, Ala, Val, Leu, Ile;
- 2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;
- 3) acidic: Asp, Glu;
- 4) basic: His, Lys, Arg;
- 5) residues that influence chain orientation: Gly, Pro; and
- 6) aromatic: Trp, Tyr, Phe.

[0051] For example, non-conservative substitutions may involve the exchange of a member of one of these classes for a member from another class. Such substituted residues may be introduced into regions of the human antibody that are homologous with non-human antibodies, or into the non-homologous regions of the molecule.

[0052] In making such changes, according to certain embodiments, the hydrophobic index of amino acids may be considered. Each amino acid has been assigned a hydrophobic index on the basis of its hydrophobicity and charge characteristics. They are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

[0053] The importance of the hydrophobic amino acid index in conferring interactive biological function on a protein is understood in the art. Kyte et al., J. Mol. Biol., 157:105-131 (1982). It is known that certain amino acids may be substituted for other amino acids having a similar hydrophobic index or score and still retain a similar biological activity. In making changes based upon the hydrophobic index, in certain embodiments, the substitution of amino acids whose hydrophobic indices are within ± 2 is included. In certain embodiments, those which are within ± 1 are included, and in certain embodiments, those within ± 0.5 are included.

[0054] It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity, particularly where the biologically functional protein or peptide thereby created is intended for use in immunological embodiments, as in the present case. In certain embodiments, the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with its immunogenicity and antigenicity, i.e., with a biological property of the protein.

[0055] The following hydrophilicity values have been assigned to these amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0 \pm 1); glutamate (+3.0 \pm 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5 \pm 1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5) and tryptophan (-3.4). In making changes based upon similar hydrophilicity values, in certain embodiments, the substitution of amino acids whose hydrophilicity values are within ± 2 is included, in certain embodiments, those which are within ± 1 are included, and in certain embodiments, those within ± 0.5 are included. One may also identify epitopes from primary amino acid sequences on the basis of hydrophilicity. These regions are also referred to as "epitopic core regions."

[0056] Exemplary amino acid substitutions are set forth in Table 1.

Table 1: Amino Acid Substitutions

| Original Residues | Exemplary Substitutions | Preferred Substitutions |
|-------------------|---|-------------------------|
| Ala | Val, Leu, Ile | Val |
| Arg | Lys, Gln, Asn | Lys |
| Asn | Gln | Gln |
| Asp | Glu | Glu |
| Cys | Ser, Ala | Ser |
| Gln | Asn | Asn |
| Glu | Asp | Asp |
| Gly | Pro, Ala | Ala |
| His | Asn, Gln, Lys, Arg | Arg |
| Ile | Leu, Val, Met, Ala, Phe, Norleucine | Leu |
| Leu | Norleucine, Ile, Val, Met, Ala, Phe | Ile |
| Lys | Arg, 1,4 Diamino-butyric Acid, Gln, Asn | Arg |
| Met | Leu, Phe, Ile | Leu |
| Phe | Leu, Val, Ile, Ala, Tyr | Leu |
| Pro | Ala | Gly |
| Ser | Thr, Ala, Cys | Thr |
| Thr | Ser | Ser |
| Trp | Tyr, Phe | Tyr |
| Tyr | Trp, Phe, Thr, Ser | Phe |

(continued)

| Original Residues | Exemplary Substitutions | Preferred Substitutions |
|-------------------|-------------------------------------|-------------------------|
| Val | Ile, Met, Leu, Phe, Ala, Norleucine | Leu |

[0057] A skilled artisan will be able to determine suitable variants of the polypeptide as set forth herein using well-known techniques. In certain embodiments, one skilled in the art may identify suitable areas of the molecule that may be changed without destroying activity by targeting regions not believed to be important for activity. In certain embodiments, one can identify residues and portions of the molecules that are conserved among similar polypeptides. In certain embodiments, even areas that may be important for biological activity or for structure may be subject to conservative amino acid substitutions without destroying the biological activity or without adversely affecting the polypeptide structure.

[0058] Additionally, one skilled in the art can review structure-function studies identifying residues in similar polypeptides that are important for activity or structure. In view of such a comparison, one can predict the importance of amino acid residues in a protein that correspond to amino acid residues which are important for activity or structure in similar proteins. One skilled in the art may opt for chemically similar amino acid substitutions for such predicted important amino acid residues.

[0059] One skilled in the art can also analyze the three-dimensional structure and amino acid sequence in relation to that structure in similar polypeptides. In view of such information, one skilled in the art may predict the alignment of amino acid residues of an antibody with respect to its three dimensional structure. In certain embodiments, one skilled in the art may choose not to make radical changes to amino acid residues predicted to be on the surface of the protein, since such residues may be involved in important interactions with other molecules. Moreover, one skilled in the art may generate test variants containing a single amino acid substitution at each desired amino acid residue. The variants can then be screened using activity assays known to those skilled in the art. Such variants could be used to gather information about suitable variants. For example, if one discovered that a change to a particular amino acid residue resulted in destroyed, undesirably reduced, or unsuitable activity, variants with such a change may be avoided. In other words, based on information gathered from such routine experiments, one skilled in the art can readily determine the amino acids where further substitutions should be avoided either alone or in combination with other mutations.

[0060] A number of scientific publications have been devoted to the prediction of secondary structure. See Moulton J., *Curr. Op. in Biotech.*, 7(4):422-427 (1996), Chou et al., *Biochemistry*, 13(2):222-245 (1974); Chou et al., *Biochemistry*, 113(2):211-222 (1974); Chou et al., *Adv. Enzymol. Relat. Areas Mol. Biol.*, 47:45-148 (1978); Chou et al., *Ann. Rev. Biochem.*, 47:251-276 and Chou et al., *Biophys. J.*, 26:367-384 (1979). Moreover, computer programs are currently available to assist with predicting secondary structure. One method of predicting secondary structure is based upon homology modeling. For example, two polypeptides or proteins which have a sequence identity of greater than 30%, or similarity greater than 40% often have similar structural topologies. The recent growth of the protein structural database (PDB) has provided enhanced predictability of secondary structure, including the potential number of folds within a polypeptide's or protein's structure. See Holm et al., *Nucl. Acid. Res.*, 27(1):244-247 (1999). It has been suggested (Brenner et al., *Curr. Op. Struct. Biol.*, 7(3):369-376 (1997)) that there are a limited number of folds in a given polypeptide or protein and that once a critical number of structures have been resolved, structural prediction will become dramatically more accurate.

[0061] Additional methods of predicting secondary structure include "threading" (Jones, D., *Curr. Opin. Struct. Biol.*, 7(3):377-87 (1997); Sippl et al., *Structure*, 4(1):15-19 (1996)), "profile analysis" (Bowie et al., *Science*, 253:164-170 (1991); Gribskov et al., *Meth. Enzym.*, 183:146-159 (1990); Gribskov et al., *Proc. Nat. Acad. Sci.*, 84(13):4355-4358 (1987)), and "evolutionary linkage" (See Holm, *supra* (1999), and Brenner, *supra* (1997)).

[0062] In certain embodiments, antibody variants include glycosylation variants wherein the number and/or type of glycosylation site has been altered compared to the amino acid sequences of the parent polypeptide. In certain embodiments, protein variants comprise a greater or a lesser number of N-linked glycosylation sites than the native protein. An N-linked glycosylation site is characterized by the sequence: Asn-X-Ser or Asn-X-Thr, wherein the amino acid residue designated as X may be any amino acid residue except proline. The substitution of amino acid residues to create this sequence provides a potential new site for the addition of an N-linked carbohydrate chain. Alternatively, substitutions which eliminate this sequence will remove an existing N-linked carbohydrate chain. Also provided is a rearrangement of N-linked carbohydrate chains wherein one or more N-linked glycosylation sites (typically those that are naturally occurring) are eliminated and one or more new N-linked sites are created. Additional preferred antibody variants include cysteine variants wherein one or more cysteine residues are deleted from or substituted for another amino acid (e.g., serine) as compared to the parent amino acid sequence. Cysteine variants may be useful when antibodies must be refolded into a biologically active conformation such as after the isolation of insoluble inclusion bodies. Cysteine variants generally have fewer cysteine residues than the native protein, and typically have an even number to minimize interactions resulting from unpaired cysteines.

[0063] According to certain embodiments, amino acid substitutions are those which: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter binding affinities, and/or (4) confer or modify other physicochemical or functional properties on such polypeptides. According to certain embodiments, single or multiple amino acid substitutions (in certain embodiments, conservative amino acid substitutions) may be made in the naturally-occurring sequence (in certain embodiments, in the portion of the polypeptide outside the domain(s) forming intermolecular contacts). In certain embodiments, a conservative amino acid substitution typically may not substantially change the structural characteristics of the parent sequence (e.g., a replacement amino acid should not tend to break a helix that occurs in the parent sequence, or disrupt other types of secondary structure that characterizes the parent sequence). Examples of art-recognized polypeptide secondary and tertiary structures are described in *Proteins, Structures and Molecular Principles* (Creighton, Ed., W. H. Freeman and Company, New York (1984)); *Introduction to Protein Structure* (C. Branden and J. Tooze, eds., Garland Publishing, New York, N.Y. (1991)); and Thornton et al. *Nature* 354:105 (1991).

[0064] The term "polypeptide fragment" as used herein refers to a polypeptide that has an amino-terminal and/or carboxy-terminal deletion. In certain embodiments, fragments are at least 5 to 467 amino acids long. It will be appreciated that in certain embodiments, fragments are at least 5, 6, 8, 10, 14, 20, 50, 70, 100, 150, 200, 250, 300, 350, 400, or 450 amino acids long.

[0065] Peptide analogs are commonly used in the pharmaceutical industry as non-peptide drugs with properties analogous to those of the template peptide. These types of non-peptide compound are termed "peptide mimetics" or "peptidomimetics". Fauchere, J. *Adv. Drug Res.* 15:29 (1986); Veber and Freidinger *TINS* p.392 (1985); and Evans et al. *J. Med. Chem.* 30:1229 (1987). Such compounds are often developed with the aid of computerized molecular modeling. Peptide mimetics that are structurally similar to therapeutically useful peptides may be used to produce a similar therapeutic or prophylactic effect. Generally, peptidomimetics are structurally similar to a paradigm polypeptide (i.e., a polypeptide that has a biochemical property or pharmacological activity), such as human antibody, but have one or more peptide linkages optionally replaced by a linkage selected from; $-\text{CH}_2 \text{NH}-$, $-\text{CH}_2 \text{S}-$, $-\text{CH}_2 -\text{CH}_2-$, $-\text{CH}=\text{CH}-$ (cis and trans), $-\text{COCH}_2-$, $-\text{CH}(\text{OH})\text{CH}_2-$, and $-\text{CH}_2 \text{SO}-$, by methods well known in the art. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (e.g., D-lysine in place of L-lysine) may be used in certain embodiments to generate more stable peptides. In addition, constrained peptides comprising a consensus sequence or a substantially identical consensus sequence variation may be generated by methods known in the art (Rizo and Gierasch *Ann. Rev. Biochem.* 61:387 (1992); for example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

[0066] "Antibody" or "antibody peptide(s)" refer to an intact antibody, or a binding fragment thereof that competes with the intact antibody for specific binding. In certain embodiments, binding fragments are produced by recombinant DNA techniques. In certain embodiments, binding fragments are produced by enzymatic or chemical cleavage of intact antibodies. Binding fragments include, but are not limited to, Fab, Fab', F(ab')₂, Fv, and single-chain antibodies.

[0067] The term "heavy chain" includes any polypeptide having sufficient variable region sequence to confer specificity for an OPGL. The term "light chain" includes any polypeptide having sufficient variable region sequence to confer specificity for an OPGL. A full-length heavy chain includes a variable region domain, V_H, and three constant region domains, C_H1, C_H2, and C_H3. The V_H domain is at the amino-terminus of the polypeptide, and the C_H3 domain is at the carboxy-terminus. The term "heavy chain", as used herein, encompasses a full-length heavy chain and fragments thereof. A full-length light chain includes a variable region domain, V_L, and a constant region domain, C_L. Like the heavy chain, the variable region domain of the light chain is at the amino-terminus of the polypeptide. The term "light chain", as used herein, encompasses a full-length light chain and fragments thereof. A Fab fragment is comprised of one light chain and the C_H1 and variable regions of one heavy chain. The heavy chain of a Fab molecule cannot form a disulfide bond with another heavy chain molecule. A Fab' fragment contains one light chain and one heavy chain that contains more of the constant region, between the C_H1 and C_H2 domains, such that an interchain disulfide bond can be formed between two heavy chains to form a F(ab')₂ molecule. The Fv region comprises the variable regions from both the heavy and light chains, but lacks the constant regions. Single-chain antibodies are Fv molecules in which the heavy and light chain variable regions have been connected by a flexible linker to form a single polypeptide chain which forms an antigen-binding region. Single chain antibodies are discussed in detail in WO 88/01649 and U.S. Patent Nos. 4,946,778 and 5,260,203.

[0068] A bivalent antibody other than a "multispecific" or "multifunctional" antibody, in certain embodiments, typically is understood to have each of its binding sites identical.

[0069] An antibody substantially inhibits adhesion of a ligand to a receptor when an excess of antibody reduces the quantity of receptor bound to counterreceptor by at least about 20%, 40%, 60%, 80%, 85%, or more (as measured in an *in vitro* competitive binding assay).

[0070] The term "epitope" includes any polypeptide determinant capable of specific binding to an immunoglobulin or T-cell receptor. In certain embodiments, epitope determinants include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl, or sulfonyl, and, in certain embodiments, may have specific three

dimensional structural characteristics, and/or specific charge characteristics. An epitope is a region of an antigen that is bound by an antibody. In certain embodiments, an antibody is said to specifically bind an antigen when it preferentially recognizes its target antigen in a complex mixture of proteins and/or macromolecules. In certain embodiments, an antibody is said to specifically bind an antigen when the dissociation constant is $\leq 1 \mu\text{M}$, in certain embodiments, when the dissociation constant is $\leq 100 \text{ nM}$, and in certain embodiments, when the dissociation constant is $\leq 10 \text{ nM}$.

[0071] The term "agent" is used herein to denote a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials.

[0072] As used herein, the terms "label" or "labeled" refers to incorporation of a detectable marker, e.g., by incorporation of a radiolabeled amino acid or attachment to a polypeptide of biotin moieties that can be detected by marked avidin (e.g., streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or colorimetric methods). In certain embodiments, the label or marker can also be therapeutic. Various methods of labeling polypeptides and glycoproteins are known in the art and may be used. Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes or radionuclides (e.g., ^3H , ^{14}C , ^{15}N , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I), fluorescent labels (e.g., FITC, rhodamine, lanthanide phosphors), enzymatic labels (e.g., horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase), chemiluminescent, biotinyl groups, predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In certain embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance.

[0073] The term "biological sample", as used herein, includes, but is not limited to, any quantity of a substance from a living thing or formerly living thing. Such living things include, but are not limited to, humans, mice, monkeys, rats, rabbits, and other animals. Such substances include, but are not limited to, blood, serum, urine, cells, organs, tissues, bone, bone marrow, lymph nodes, and skin.

[0074] The term "osteopenic disorder" includes, but is not limited to, osteoporosis, osteopenia, Paget's disease, lytic bone metastases, periodontitis, rheumatoid arthritis, and bone loss due to immobilization. In addition to these bone disorders, certain cancers are known to increase osteoclast activity and induce bone resorption, such as breast, prostate, and multiple myeloma. These cancers are now known to produce factors that result in the over-expression of OPGL in the bone, and lead to increased osteoclast numbers and activity.

[0075] The term "pharmaceutical agent or drug" as used herein refers to a chemical compound or composition capable of inducing a desired therapeutic effect when properly administered to a patient.

[0076] The term "modulator," as used herein, is a compound that changes or alters the activity or function of a molecule. For example, a modulator may cause an increase or decrease in the magnitude of a certain activity or function of a molecule compared to the magnitude of the activity or function observed in the absence of the modulator. In certain embodiments, a modulator is an inhibitor, which decreases the magnitude of at least one activity or function of a molecule. Certain exemplary activities and functions of a molecule include, but are not limited to, binding affinity, enzymatic activity, and signal transduction. Certain exemplary inhibitors include, but are not limited to, proteins, peptides, antibodies, peptibodies, carbohydrates or small organic molecules. Peptibodies are described, e.g., in WO01/83525.

[0077] As used herein, "substantially pure" means an object species is the predominant species present (i.e., on a molar basis it is more abundant than any other individual species in the composition). In certain embodiments, a substantially purified fraction is a composition wherein the object species comprises at least about 50 percent (on a molar basis) of all macromolecular species present. In certain embodiments, a substantially pure composition will comprise more than about 80%, 85%, 90%, 95%, or 99% of all macromolar species present in the composition. In certain embodiments, the object species is purified to essential homogeneity (contaminant species cannot be detected in the composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species.

[0078] The term patient includes human and animal subjects.

[0079] In this application, the use of the singular includes the plural unless specifically stated otherwise. In this application, the use of "or" means "and/or" unless stated otherwise. Furthermore, the use of the term "including", as well as other forms, such as "includes" and "included", is not limiting. Also, terms such as "element" or "component" encompass both elements and components comprising one unit and elements and components that comprise more than one subunit unless specifically stated otherwise.

[0080] Osteoprotegerin Ligand (OPGL), a member of the tumor necrosis factor (TNF) family of cytokines, is involved in the formation of osteoclasts. Increased osteoclast activity correlates with a number of osteopenic disorders, including post-menopausal osteoporosis, Paget's disease, lytic bone metastases, and rheumatoid arthritis. Therefore, a reduction in OPGL activity may result in a decrease in osteoclast activity and may reduce the severity of osteopenic disorders. As disclosed herein, antibodies directed to OPGL may be used to treat osteopenic disorders, including but not limited to, those mentioned above.

[0081] In certain embodiments of the present invention, there is provided a fully human monoclonal antibody against human osteoprotegerin ligand (OPGL). In certain embodiments, amino acid sequences comprising, heavy and light chain immunoglobulin molecules, particularly sequences corresponding to the variable regions are provided. In certain

embodiments, sequences corresponding to complementarity determining regions (CDR's), specifically from CDR1 through CDR3, are provided. According to certain embodiments, a hybridoma cell line expressing such an immunoglobulin molecule and monoclonal antibody is also provided. In certain embodiments, purified human monoclonal antibody against human OPGL is provided.

5 **[0082]** The ability to clone and reconstruct megabase-sized human loci in yeast artificial chromosomes (YACs) and to introduce them into the mouse germline provides an approach to elucidating the functional components of very large or crudely mapped loci as well as generating useful models of human disease. Furthermore, the utilization of such technology for substitution of mouse loci with their human equivalents could provide unique insights into the expression and regulation of human gene products during development, their communication with other systems, and their involvement in disease induction and progression.

10 **[0083]** An important practical application of such a strategy is the "humanization" of the mouse humoral immune system. Introduction of human immunoglobulin (Ig) loci into mice in which the endogenous Ig genes have been inactivated offers the opportunity to study the mechanisms underlying programmed expression and assembly of antibodies as well as their role in B-cell development. Furthermore, such a strategy could provide a source for production of fully human monoclonal antibodies (MAbs). In certain embodiments, fully human antibodies are expected to minimize the immunogenic and allergic responses intrinsic to mouse or mouse-derivatized Mabs, and thus, in certain embodiments, increase the efficacy and safety of the administered antibodies. Fully human antibodies may be used in the treatment of chronic and recurring human diseases, such as osteoporosis, inflammation, autoimmunity, and cancer, which may involve repeated antibody administrations.

20 **[0084]** One can engineer mouse strains deficient in mouse antibody production with large fragments of the human Ig loci in anticipation that such mice would produce human antibodies in the absence of mouse antibodies. Large human Ig fragments may preserve the large variable gene diversity as well as the proper regulation of antibody production and expression. By exploiting the mouse machinery for antibody diversification and selection and the lack of immunological tolerance to human proteins, the reproduced human antibody repertoire in these mouse strains may yield high affinity antibodies against any antigen of interest, including human antigens. Using the hybridoma technology, antigen-specific human MAbs with the desired specificity may be produced and selected.

25 **[0085]** In certain embodiments, one may use constant regions from species other than human along with the human variable region(s).

30 Naturally Occurring Antibody Structure

[0086] Naturally occurring antibody structural units typically comprise a tetramer. Each such tetramer typically is composed of two identical pairs of polypeptide chains, each pair having one full-length "light" (in certain embodiments, about 25 kDa) and one full-length "heavy" chain (in certain embodiments, about 50-70 kDa). The amino-terminal portion of each chain typically includes a variable region of about 100 to 110 or more amino acids that typically is responsible for antigen recognition. The carboxy-terminal portion of each chain typically defines a constant region that may be responsible for effector function. Human light chains are typically classified as kappa and lambda light chains. Heavy chains are typically classified as mu, delta, gamma, alpha, or epsilon, and define the antibody's isotype as IgM; IgD, IgG, IgA, and IgE, respectively. IgG has several subclasses, including, but not limited to, IgG1, IgG2, IgG3, and IgG4. IgM has subclasses including, but not limited to, IgM1 and IgM2. IgA is similarly subdivided into subclasses including, but not limited to, IgA1 and IgA2. Within full-length light and heavy chains, typically, the variable and constant regions are joined by a "J" region of about 12 or more amino acids, with the heavy chain also including a "D" region of about 10 more amino acids. See, e.g., *Fundamental Immunology* Ch 7 (Paul, W., ed., 2nd ed. Raven Press, N Y. (1989)). The variable regions of each light/heavy chain pair typically form the antigen binding site.

45 **[0087]** The variable regions typically exhibit the same general structure of relatively conserved framework regions (FR) joined by three hyper variable regions, also called complementarity determining regions or CDRs. The CDRs from the two chains of each pair typically are aligned by the framework regions, which may enable binding to a specific epitope. From N-terminal to C-terminal, both light and heavy chain variable regions typically comprise the domains FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The assignment of amino acids to each domain is typically in accordance with the definitions of Kabat Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md. (1987 and 1991)), or Chothia & Lesk *J. Mol. Biol.* 196:901-917 (1987); Chothia et al. *Nature* 342:877-883 (1989).

Bispecific or Bifunctional Antibodies

55 **[0088]** A bispecific or bifunctional antibody typically is an artificial hybrid antibody having two different heavy/light chain pairs and two different binding sites. Bispecific antibodies may be produced by a variety of methods including, but not limited to, fusion of hybridomas or linking of Fab' fragments. See, e.g., Songsivilai & Lachmann *Clin. Exp. Immunol.* 79: 315-321 (1990), Kostelny et al. *J. Immunol.* 148:1547-1553 (1992).

Preparation of Antibodies

5 [0089] According to certain embodiments, certain antibodies specifically binding to OPGL are encompassed by the invention. In certain embodiments, the antibodies may be produced by immunization with full-length OPGL, soluble forms of OPGL, or a fragment thereof. In certain embodiments, the antibodies of the invention may be polyclonal or monoclonal, and/or may be recombinant antibodies. In certain embodiments, antibodies of the invention are human antibodies prepared, for example, by immunization of transgenic animals capable of producing human antibodies (see, for example, PCT Published Application No. WO 93/12227).

10 [0090] In certain embodiments, the complementarity determining regions (CDRs) of the light and heavy chain variable regions of α OPGL-1 may be grafted to framework regions (FRs) from the same, or another, species. In certain embodiments, the CDRs of the light and heavy chain variable regions of α OPGL-1 may be grafted to consensus human FRs. To create consensus human FRs, in certain embodiments, FRs from several human heavy chain or light chain amino acid sequences are aligned to identify a consensus amino acid sequence. In certain embodiments, the FRs of the α OPGL-1 heavy chain or light chain are replaced with the FRs from a different heavy chain or light chain. In certain
15 embodiments, rare amino acids in the FRs of the heavy and light chains of α OPGL-1 are not replaced, while the rest of the FR amino acids are replaced. Rare amino acids are specific amino acids that are in positions in which they are not usually found in FRs. In certain embodiments, the grafted variable regions from α OPGL-1 may be used with a constant region that is different from the constant region of α OPGL-1. In certain embodiments, the grafted variable regions are part of a single chain Fv antibody. CDR grafting is described, e.g., in U.S. Patent Nos. 6,180,370, 5,693,762, 5,693,761,
20 5,585,089, and 5,530,101.

[0091] According to certain embodiments, antibodies of the invention are prepared through the utilization of a transgenic mouse that has a substantial portion of the human antibody producing genome inserted but that is rendered deficient in the production of endogenous, murine, antibodies. Such mice, then, are capable of producing human immunoglobulin molecules and antibodies and are deficient in the production of murine immunoglobulin molecules and antibodies.
25 Technologies utilized for achieving this result are disclosed in the patents, applications, and references disclosed in the specification, herein. In certain embodiments, one may employ methods such as those disclosed in PCT Published Application No. WO 98/24893. See also Mendez et al. Nature Genetics 15:146-156 (1997).

[0092] According to certain embodiments, fully human monoclonal antibodies specific for OPGL are produced as follows. Transgenic mice containing human immunoglobulin genes are immunized with the antigen of interest. Lymphatic
30 cells (such as B-cells) from the mice that express antibodies are obtained. Such recovered cells are fused with a myeloid-type cell line to prepare immortal hybridoma cell lines, and such hybridoma cell lines are screened and selected to identify hybridoma cell lines that produce antibodies specific to the antigen of interest. In certain embodiments, the production of a hybridoma cell line that produces antibodies specific to OPGL is provided.

[0093] In certain embodiments, antibodies of the invention are produced by hybridoma lines AMG 6.1, AMG 6.4, AMG
35 6.5, AMG 7.1, and AMG 7.2. In certain embodiments, antibodies of the invention are produced by hybridoma lines AMG 6.1, AMG 6.4, and AMG 6.5. In certain embodiments, the antibodies of the invention bind to OPGL with a dissociation constant (Kd) of between approximately 0.23 and 0.29 nM. In certain embodiments of the invention, the antibodies bind to OPGL with a Kd of less than 0.23 nM.

[0094] In certain embodiments, the antibodies of the invention are of the IgG2 isotype. In certain embodiments of the
40 invention, the antibodies comprise a human kappa light chain and a human IgG2 heavy chain. In certain embodiments, the antibodies of the invention have been cloned for expression in mammalian cells. In certain embodiments, the variable regions of the antibodies are ligated to a constant region other than the constant region for the IgG2 isotype.

[0095] In certain embodiments, conservative modifications to the heavy and light chains of α OPGL-1 (and corresponding modifications to the encoding nucleotides) will produce antibodies to OPGL having functional and chemical characteristics similar to those of α OPGL-1. In contrast, substantial modifications in the functional and/or chemical characteristics
45 of α OPGL-1 may be accomplished by selecting substitutions in the amino acid sequence of the heavy and light chains that differ significantly in their effect on maintaining (a) the structure of the molecular backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

[0096] For example, a "conservative amino acid substitution" may involve a substitution of a native amino acid residue
50 with a nonnative residue such that there is little or no effect on the polarity or charge of the amino acid residue at that position. Furthermore, any native residue in the polypeptide may also be substituted with alanine, as has been previously described for "alanine scanning mutagenesis."

[0097] Desired amino acid substitutions (whether conservative or non-conservative) can be determined by those skilled
55 in the art at the time such substitutions are desired. In certain embodiments, amino acid substitutions can be used to identify important residues of α OPGL-1, or to increase or decrease the affinity of the antibodies to OPGL described herein.

[0098] In certain embodiments, antibodies of the present invention can be expressed in cell lines other than hybridoma cell lines. In certain embodiments, sequences encoding particular antibodies can be used for transformation of a suitable

mammalian host cell. According to certain embodiments, transformation can be by any known method for introducing polynucleotides into a host cell, including, for example packaging the polynucleotide in a virus (or into a viral vector) and transducing a host cell with the virus (or vector) or by transfection procedures known in the art, as exemplified by U.S. Pat. Nos. 4,399,216, 4,912,040, 4,740,461, and 4,959,455. In certain embodiments, the transformation procedure used may depend upon the host to be transformed. Methods for introduction of heterologous polynucleotides into mammalian cells are well known in the art and include, but are not limited to dextran-mediated transfection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei.

[0099] Mammalian cell lines available as hosts for expression are well known in the art and include, but are not limited to, many immortalized cell lines available from the American Type Culture Collection (ATCC), including but not limited to Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), and a number of other cell lines. In certain embodiments, cell lines may be selected through determining which cell lines have high expression levels and produce antibodies with constitutive OPGL binding properties.

[0100] According to certain embodiments, antibodies of the invention are useful for detecting OPGL in biological samples. In certain embodiments, this allows the identification of cells or tissues which produce the protein. In certain embodiments, antibodies which bind to OPGL and block interaction with other binding compounds may have therapeutic use in modulating osteoclast differentiation and bone resorption. In certain embodiments, antibodies to OPGL may block OPGL binding to ODAR, which may result in a block in the signal transduction cascade and loss of NF- κ B mediated transcription activation. Assays for measuring NF- κ B-mediated transcription activation using, e.g., a luciferase reporter assay, are known to those skilled in the art.

[0101] Methods are disclosed of treating a bone disorder comprising administering a therapeutically effective amount of an antibody to OPGL. Methods are disclosed of treating a bone disorder comprising administering a therapeutically effective amount of an antibody to OPGL and another therapeutic agent. In certain such methods, the additional therapeutic agent is administered in a therapeutically effective amount. In certain methods, the bone disorder is a disorder characterized by a net bone loss, including but not limited to, osteopenia and osteolysis. In certain methods, treatment with an antibody to OPGL is used to suppress the rate of bone resorption. Therefore, in certain methods, treatment may be used to reduce the rate of bone resorption where the resorption rate is above normal, or to reduce bone resorption to below normal levels in order to compensate for below normal levels of bone formation. In certain methods, antibodies can be tested for binding to OPGL in the absence or presence of OPG and examined for their ability to inhibit OPGL-mediated osteoclastogenesis and/or bone resorption.

[0102] Conditions which may be treated according to certain aspects of the disclosure include, but are not limited to, the following:

Osteoporosis, including, but not limited to, primary osteoporosis, endocrine osteoporosis (including, but not limited to, hyperthyroidism, hyperparathyroidism, Cushing's syndrome, and acromegaly), hereditary and congenital forms of osteoporosis (including, but not limited to, osteogenesis imperfecta, homocystinuria, Menkes' syndrome, Riley-Day syndrome), and osteoporosis due to immobilization of extremities;

Paget's disease of bone (osteitis deformans) in adults and juveniles;

Osteomyelitis, i.e., an infectious lesion in bone, leading to bone loss;

Hypercalcemia, including, but not limited to, hypercalcemia resulting from solid tumors (including, but not limited to, breast, lung and kidney) and hematologic malignancies (including, but not limited to, multiple myeloma, lymphoma and leukemia), idiopathic hypercalcemia, and hypercalcemia associated with hyperthyroidism and renal function disorders;

Osteopenia, including but not limited to, osteopenia following surgery, osteopenia induced by steroid administration, osteopenia associated with disorders of the small and large intestine, and osteopenia associated with chronic hepatic and renal diseases;

Osteonecrosis, i.e., bone cell death, including, but not limited to, osteonecrosis associated with traumatic injury, osteonecrosis associated with Gaucher's disease, osteonecrosis associated with sickle cell anemia, osteonecrosis associated with systemic lupus erythematosus, osteonecrosis associated with rheumatoid arthritis, osteonecrosis associated with periodontal disease, osteonecrosis associated with osteolytic metastasis, and osteonecrosis associated with other condition; and

Loss of cartilage and joint erosion associated with rheumatoid arthritis.

[0103] In certain aspects of the disclosure, an antibody to OPGL may be for use alone or with at least one additional therapeutic agents for the treatment of bone disorders. In certain aspects of the disclosure, an antibody to OPGL is for use in conjunction with a therapeutically effective amount of an additional therapeutic agent. Exemplary therapeutic agents that may be administered with an antibody to OPGL include, but are not limited to, the bone morphogenic factors

designated BMP-1 through BMP-12; transforming growth factor- β (TGF- β) and TGF- β family members; interleukin-1 (IL-1) inhibitors, including, but not limited to, IL-1ra and derivatives thereof and Kineret™ anakinra TNF α inhibitors, including, but not limited to, soluble TNF α receptors, Enbrel™, etanercept anti-TNF α antibodies, Remicade™, infliximab and D2E7 antibodies; parathyroid hormone and analogs thereof; parathyroid related protein and analogs thereof; E series prostaglandins; bisphosphonates (such as alendronate and others); bone-enhancing minerals such as fluoride and calcium; non-steroidal anti-inflammatory drugs (NSAIDs), including, but not limited to, COX-2 inhibitors, such as Celebrex™ celecoxib, and Vioxx™, rofecoxib, immunosuppressants, such as methotrexate or leflunomide; serine protease inhibitors, including, but not limited to, secretory leukocyte protease inhibitor (SI.PI); IL-6 inhibitors (including, but not limited to, antibodies to IL-6), IL-8 inhibitors (including, but not limited to, antibodies to IL-8); IL-18 inhibitors (including, but not limited to, IL-18 binding protein and IL-18 antibodies); Interleukin-1 converting enzyme (ICE) modulators; fibroblast growth factors FGF-1 to FGF-10 and FGF modulators; PAF antagonists; keratinocyte growth factor (KGF), KGF-related molecules, and KGF modulators; matrix metalloproteinase (MMP) modulators; Nitric oxide synthase (NOS) modulators, including, but not limited to, modulators of inducible NOS; modulators of glucocorticoid receptor; modulators of glutamate receptor; modulators of lipopolysaccharide (LPS) levels; and noradrenaline and modulators and mimetics thereof.

[0104] In certain aspects of the disclosure, an antibody to OPGL is for use with particular therapeutic agents to treat various inflammatory conditions, autoimmune conditions, or other conditions with attendant bone loss. In certain aspects of the disclosure, in view of the condition and the desired level of treatment, two, three, or more agents may be administered. In certain aspects of the disclosure, such agents may be provided together by inclusion in the same formulation. In certain aspects of the disclosure, such agents and an antibody to OPGL may be provided together by inclusion in the same formulation. In certain aspects of disclosure, such agents may be provided together by inclusion in a treatment kit. In certain aspects of the disclosure, such agents and an antibody to OPGL may be provided together by inclusion in a treatment kit. In certain aspects of the disclosure, such agents may be provided separately. In certain aspects of the disclosure, when administered by gene therapy, the genes encoding protein agents and/or an antibody to OPGL may be included in the same vector. In certain aspects of the disclosure, the genes encoding protein agents and/or an antibody to OPGL may be under the control of the same promoter region. In certain aspects of the disclosure, the genes encoding protein agents and/or an antibody to OPGL may be in separate vectors.

[0105] In certain aspects of the disclosure, the present invention is directed to an antibody to OPGL and at least one interleukin-1 (IL-1) inhibitor for use in therapies and methods of treatment using such therapies. In certain aspects of the disclosure, an antibody to OPGL and an IL-1 inhibitor and at least one additional molecule described herein are for use in a therapy. In certain aspects of the disclosure, IL-1 inhibitors and/or TNF- α inhibitors are for use in conjunction with an antibody to OPGL in methods of treatment. In certain aspects of the disclosure, an antibody to OPGL in combination with IL-1 inhibitors and/or TNF- α inhibitors is for use for treatment of conditions such as asthma, rheumatoid arthritis, and multiple sclerosis.

[0106] Interleukin-1 (IL-1) is an anti-inflammatory cytokine. In certain instances, IL-1 is a mediator in many diseases and medical conditions. In certain instances, IL-1 is manufactured by cells of the macrophage/monocyte lineage. In certain instances, IL-1 is produced in two forms: IL-1 alpha (IL-1 α) and IL-1 beta (IL-1 β).

[0107] A disease or medical condition is considered to be an "interleukin-1 mediated disease" if the spontaneous or experimental disease or medical condition is associated with elevated levels of IL-1 in bodily fluids or tissue and/or if cells or tissues taken from the body produce elevated levels of IL-1 in culture. In certain aspects of the disclosure, such interleukin-1 mediated diseases are also recognized by the following additional two conditions: (1) pathological findings associated with the disease or medical condition can be mimicked experimentally in animals by administration of IL-1 or upregulation of expression of IL-1; and (2) a pathology induced in experimental animal models of the disease or medical condition can be inhibited or abolished by treatment with agents that inhibit the action of IL-1. In certain aspects of the disclosure, one or more of the above conditions are met in an IL-1-mediated disease. In certain aspects of the disclosure, all three of the conditions are met in an IL-1-mediated disease.

[0108] Acute and chronic interleukin-1 (IL-1) -mediated diseases include, but are not limited to, the following: acute pancreatitis; amyotrophic lateral sclerosis (ALS, or Lou Gehrig's disease); Alzheimer's disease; cachexia/anorexia, including, but not limited to, AIDS-induced cachexia; asthma and other pulmonary diseases; atherosclerosis; autoimmune vasculitis; chronic fatigue syndrome; Clostridium associated illnesses, including, but not limited to, Clostridium-associated diarrhea; coronary conditions and indications, including, but not limited to, congestive heart failure, coronary restenosis, myocardial infarction, myocardial dysfunction (e.g., related to sepsis), and coronary artery bypass graft; cancer, including, but not limited to, leukemias, including, but not limited to, multiple myeloma leukemia and myelogenous (e.g., AML and CML), and tumor metastasis; diabetes (including, but not limited to, insulin-dependent diabetes); endometriosis; fever; fibromyalgia; glomerulonephritis; graft versus host disease and/or transplant rejection; hemohorrhagic shock; hyperalgesia; inflammatory bowel disease; inflammatory conditions of a joint, including, but not limited to, osteoarthritis, psoriatic arthritis, and rheumatoid arthritis; inflammatory eye disease, including, but not limited to, those associated with, for example, corneal transplant; ischemia, including, but not limited to, cerebral ischemia (including, but not limited to, brain injury as a result of, e.g., trauma, epilepsy, hemorrhage or stroke, each of which may lead to neurodegeneration);

Kawasaki's disease; learning impairment; lung diseases (including, but not limited to, acute respiratory distress syndrome, or ARDS); multiple sclerosis; myopathies (e.g., muscle protein metabolism, including, but not limited to, muscle protein metabolism in sepsis); neurotoxicity (including, but not limited to, such condition induced by HIV); osteoporosis; pain, including, but not limited to, cancer-related pain; Parkinson's disease; periodontal disease; pre-term labor; psoriasis; reperfusion injury; septic shock; side effects from radiation therapy; temporal mandibular joint disease; sleep disturbance; uveitis; and an inflammatory condition resulting from, e.g., strain, sprain, cartilage damage, trauma, orthopedic surgery, infection, or other disease processes.

[0109] In certain aspects of the disclosure, an IL-1 inhibitor may be any protein or molecule capable of specifically preventing activation of cellular receptors to IL-1, which may result from any number of mechanisms. Exemplary mechanisms include, but are not limited to, downregulating IL-1 production, binding free IL-1, interfering with IL-1 binding to its receptor, interfering with formation of the IL-1 receptor complex (i.e., association of IL-1 receptor with IL-1 receptor accessory protein), and interfering with modulation of IL-1 signaling after binding to its receptor.

[0110] Certain interleukin-1 inhibitors include, but are not limited to, IL-1 receptor antagonists, including, but not limited to, Kineret™, anakinra, IL-1ra, IL-1ra variants, and IL-1ra derivatives, which are collectively termed "IL-1ra proteins;" anti-IL-1 receptor monoclonal antibodies (see, e.g., EP 623674); IL-1 binding proteins, including, but not limited to, soluble IL-1 receptors (see, e.g., U. S. Pat No. 5,492,888, U. S. Pat No. 5,488,032, and U. S. Pat No. 5,464,937, U. S. Pat. No. 5,319,071, and U.S. Pat No. 5,180,812); anti-IL-1 monoclonal antibodies (see, e.g., WO 9501997, WO 9402627, WO 9006371, U.S. Pat No. 4,935,343, EP 364778, EP 267611 and EP 220063); IL-1 receptor accessory proteins and antibodies thereto (see, e.g., WO 96/23067 and WO 99/37773); inhibitors of interleukin-1 beta converting enzyme (ICE) or caspase 1 (see, e.g., WO 99/46248, WO 99/47545, and WO 99/47154), which may be used to inhibit IL-1 beta production and secretion; interleukin-1 beta protease inhibitors; and other compounds and proteins that block *in vivo* synthesis or extracellular release of IL-1.

[0111] Exemplary IL-1 inhibitors are disclosed, e.g., in US Pat. Nos. 5,747,444; 5,359,032; 5,608,035; 5,843,905; 5,359,032; 5,866,576; 5,869,660; 5,869,315; 5,872,095; 5,955,480; 5,965,564; International (WO) patent applications 98/21957, 96/09323, 91/17184, 96/40907, 98/32733, 98/42325, 98/44940, 98/47892, 98/56377, 99/03837, 99/06426, 99/06042, 91/17249, 98/82733, 98/17661, 97/08174, 95/34326, 99/36426, 99/36415; European (EP) patent applications 534978 and 894795; and French patent application FR 2762514.

[0112] Interleukin-1 receptor antagonist (IL-1ra) is a human protein that acts as a natural inhibitor of interleukin-1 and is a member of the IL-1 family, which includes IL-1 α and IL-1 β . Certain receptor antagonists, including IL-1ra and variants and derivatives thereof, as well as methods of making and using them, are described in U.S. Patent No. 5,075,222; WO 91/08285; WO 91/17184; AU 9173636; WO 92/16221; WO 93/21946; WO 94/06457; WO 94/21275; FR 2706772; WO 94/21235; DE 4219626, WO 94/20517; WO 96/22793; WO 97/28828; and WO 99/36541. In certain aspects of the disclosure, an IL-1 receptor antagonist may be glycosylated. In certain aspects of the disclosure, an IL-1 receptor antagonist may be non-glycosylated.

[0113] Three forms of IL-1ra and variants thereof are described in U.S. Pat. No 5,075,222 (the '222 patent). The first form, called "IL-1I" in the '222 patent, is characterized as a 22-23 kD molecule on SDS-PAGE with an approximate isoelectric point of 4.8, which elutes from a Mono Q FPLC column at around 52 mM NaCl in Tris buffer, pH 7.6. The second form, IL-1ra β , is characterized as a 22-23 kD protein, which elutes from a Mono Q column at 48 mM NaCl. Both IL-1ra α and IL-1ra β are glycosylated. The third form, IL-1rax, is characterized as a 20 kD protein, which elutes from a Mono Q column at 48 mM NaCl and is non-glycosylated. The '222 patent also describes certain methods for isolating genes that code for the inhibitors, cloning those genes in suitable vectors, transforming and transfecting those genes into certain cell types, and expressing those genes to produce the inhibitors.

[0114] In certain aspects of the disclosure, deletions, insertions, and/or substitutions (individually or collectively referred to as "variant(s)") are made within the amino acid sequences of IL-1ra. In certain aspects of the disclosure, an IL-1ra variant is biologically active (e.g., possesses the ability to inhibit IL-1).

[0115] In certain aspects of the disclosure, the present invention is directed to an antibody to OPGL and at least one TNF α inhibitor for use in therapies and methods of treatment using such therapies. In certain aspects of the disclosure, an antibody to OPGL and a TNF α inhibitor and at least one additional molecule described herein is for use in a therapy.

[0116] Certain diseases and medical conditions are mediated by TNF and may be categorized as inflammatory conditions. As used herein, a "TNF-mediated disease" includes, but is not limited to, a disease or medical condition that is associated with elevated levels of TNF in bodily fluids or tissue and/or in which cells or tissues taken from the body produce elevated levels of TNF in culture. In certain aspects of the disclosure, a disease is a TNF-mediated disease if (1) pathological findings associated with the disease or medical condition can be mimicked experimentally in animals by the administration or upregulation of expression of TNF and/or (2) a pathology induced in experimental animal models of the disease or medical condition can be inhibited or abolished by treatment with agents that inhibit the action of TNF.

[0117] Certain acute and chronic TNF-mediated diseases include, but are not limited to: cachexia and anorexia; cancer, including, but not limited to, leukemia; chronic fatigue syndrome; coronary conditions and/or indications, including, but not limited to, congestive heart failure, coronary restenosis, myocardial infarction, myocardial dysfunction (including but

not limited to, such condition related to sepsis), and coronary artery bypass graft; depression; diabetes, including, but not limited to, juvenile onset Type 1 diabetes, diabetes mellitus, and insulin resistance (including, but not limited to, insulin resistance associated with obesity); endometriosis, endometritis, and related conditions; fibromyalgia and analgesia; graft versus host rejection; hyperalgesia; inflammatory bowel diseases, including, but not limited to, Crohn's disease and *Clostridium difficile*-associated diarrhea; ischemia, including, but not limited to, cerebral ischemia, which includes, but is not limited to, brain injury as a result of trauma, epilepsy, hemorrhage, and/or stroke; lung disease, including, but not limited to, adult respiratory distress syndrome, asthma, and pulmonary fibrosis; multiple sclerosis; neuroinflammatory diseases; ocular diseases and conditions, including, but not limited to, corneal transplant, ocular degeneration and uveitis; pain, including, but not limited to, cancer-related pain; pancreatitis; periodontal diseases; Pityriasis rubra pilaris (PRP); prostatitis, including bacterial and non-bacterial prostatitis, and related conditions; psoriasis and related conditions; pulmonary fibrosis; reperfusion injury; rheumatic diseases, including, but not limited to, rheumatoid arthritis, osteoarthritis, juvenile arthritis (including, but not limited to, juvenile rheumatoid arthritis), seronegative polyarthritis, ankylosing spondylitis, Reiter's syndrome and reactive arthritis, Still's disease, psoriatic arthritis, enteropathic arthritis, polymyositis, dermatomyositis, scleroderma, systemic sclerosis, vasculitis (e.g., Kawasaki's disease), cerebral vasculitis, Lyme disease, staphylococcal-induced ("septic") arthritis, Sjögren's syndrome, rheumatic fever, polychondritis and polymyalgia rheumatica and giant cell arteritis); septic shock; side effects from radiation therapy; systemic lupus erythematosus (SLE); temporal mandibular joint disease; thyroiditis; and tissue transplantation and/or an inflammatory condition, e.g., resulting from strain, sprain, cartilage damage, trauma, orthopedic surgery, infection (e.g., HIV, *Clostridium difficile* and related species) or other disease process.

[0118] In certain embodiments, TNF inhibitors may act by at least one of downregulating or inhibiting TNF production, binding free TNF, interfering with TNF binding to its receptor, and interfering with modulation of TNF signaling after binding to its receptor. The term "TNF inhibitor" includes, but is not limited to, solubilized TNF receptors, including, but not limited to, soluble tumor necrosis factor receptor type I (sTNF-RI; also called the p55 receptor), soluble tumor necrosis factor receptor type II (also called the p75 receptor), and Enbrel™ etanercept; antibodies to TNF, including, but not limited to, Remicade™, infliximab and D2E7 (see, e.g., U.S. Patent Nos. 6,090,382 and 6,258,562); antibodies to TNF receptor; sTNF-RI (see, e.g., WO 98/24463), etanercept (Enbrel™), Avakine™; inhibitors of TNF- α converting enzyme (TACE); and other molecules that affect TNF activity.

[0119] Exemplary TNF- α inhibitors are described, e.g., in European patent applications EP 308 378; EP 422 339; EP 393 438; EP 398 327; EP 412 486; EP 418 014, EP 417 563, EP 433 900; EP 464 533; EP 512 528; EP 526 905; EP 568 928; EP 607 776, which describes the use of leflunomide for inhibition of TNF- α ; EP 663 210; EP 542 795; EP 818 439; EP 664 128; EP 542 795; EP 741 707; EP 874 819 ; EP 882 714; EP 880 970; EP 648 783; EP 731 791; EP 895 988; EP 550 376; EP 882 714; EP 853 083; EP 550 376; EP 943 616; EP 939 121; EP 614 984 ; EP 853 083; U.S. Patent Nos. 5,136,021; 5,929,117; 5,948,638; 5,807,862; 5,695,953; 5,834,435; 5,817,822; 5,830,742; 5,834,435; 5,851,556; 5,853,977; 5,359,037; 5,512,544; 5,695,953; 5,811,261; 5,633,145; 5,863,926; 5,866,616; 5,641,673; 5,869,677; 5,869,511; 5,872,146; 5,854,003; 5,856,161; 5,877,222; 5,877,200; 5,877,151; 5,886,010; 5,869,660; 5,859,207; 5,891,883; 5,877,180; 5,955,480; 5,955,476; 5,955,435; 5,994,351; 5,990,119; 5,952,320; 5,962,481; International patent applications WO 90/13575, WO 91/03553, WO 92/01002, WO 92/13095, WO 92/16221, WO 93/07863, WO 93/21946, WO 93/19777, WO 95/34326, WO 96/28546, WO 98/27298, WO 98/30541, WO 96/38150, WO 96/38150, WO 97/18207, WO 97/15561, WO 97/12902, WO 96/25861, WO 96/12735, WO 96/11209, WO 98/39326, WO 98/39316, WO 98/38859, WO 98/39315, WO 98/42659, WO 98/39329, WO 98/43959, WO 98/45268, WO 98/47863, WO 96/33172, WO 96/20926, WO 97/37974, WO 97/37973, WO 97/47599, WO 96/35711, WO 98/51665, WO 98/43946, WO 95/04045, WO 98/56377, WO 97/12244, WO 99/00364, WO 99/00363, WO 98/57936, WO 99/01449, WO 99/01139, WO 98/56788, WO 98/56756, WO 98/53842, WO 98/52948, WO 98/52937, WO 99/02510, WO 97/43250, WO 99/06410, WO 99/06042, WO 99/09022, WO 99/08688, WO 99/07679, WO 99/09965, WO 99/07704, WO 99/06041, WO 99/37818, WO 99/37625, WO 97/11668, WO 99/50238, WO 99/47672, WO 99/48491; Japanese patent applications 10147531, 10231285, 10259140, and 10130149, 10316570, 11001481, and 127,800/1991; German application no. 19731521; and British application nos. 2 218 101, 2 326 881, 2 246 569.

[0120] EP 393 438 and EP 422 339 describe the amino acid and nucleic acid sequences of a soluble TNF receptor type I (also known as sTNFR-I or 30kDa TNF inhibitor) and a soluble TNF receptor type II (also known as sTNFR-II or 40kDa TNF inhibitor), which are collectively termed "sTNFRs". EP 393 438 and EP 422 339 also describe modified forms of sTNFR-1 and sTNFR-II, including, but not limited to fragments, functional derivatives, and variants. Furthermore, EP 393 438 and EP 422 339 describe methods for isolating genes that code for the inhibitors, cloning the genes into suitable vectors, transforming or transfecting the genes into certain cell types, and expressing the genes to produce the inhibitors.

[0121] sTNFR-I and sTNFR-II are members of the nerve growth factor/TNF receptor superfamily of receptors, which includes the nerve growth factor receptor (NGF), the B cell antigen CD40, 4-1 BB, the rat T-cell antigen MRC OX40, the fas antigen, and the CD27 and CD30 antigens (Smith et al. (1990) Science, 248:1019-1023). A conserved feature of that group of cell surface receptors is a cysteine-rich extracellular ligand binding domain, which can be divided into four

repeated motifs of about forty amino acids that contain 4-6 cysteine residues at positions that are well conserved (Smith et al. (1990), *supra*).

[0122] EP 393 438 teaches a 40kDa TNF inhibitor $\Delta 51$ and a 40kDa TNF inhibitor $\Delta 53$, which are truncated versions of the full-length recombinant 40kDa TNF inhibitor protein. $\Delta 51$ and $\Delta 53$ have 51 or 53 amino acids, respectively, deleted from the carboxyl terminus of the mature protein.

[0123] Published PCT Application No. WO 98/01555 describes truncated forms of sTNFR-I and sTNFR-II that do not contain the fourth domain (amino acid residues Thr¹²⁷-Asn¹⁶¹ of sTNFR-I and amino acid residues Pro¹⁴¹-Thr¹⁷⁹ of sTNFR-II); a portion of the third domain (amino acid residues Asn¹¹¹-Cys¹²⁶ of sTNFR-I and amino acid residues Pro¹²³-Lys¹⁴⁰ of sTNFR-II); and, optionally, do not contain a portion of the first domain (amino acid residues Asp¹-Cys¹⁹ of sTNFR-I and amino acid residues Leu¹-Cys³² of sTNFR-II). In certain aspects of the disclosure, the truncated sTNFRs include the proteins represented by the formula R₁-[Cys¹⁹-Cys¹⁰³]-R₂ and R₄-[Cys³⁷-Cys¹¹⁵]-R₅. These proteins are truncated forms of sTNFR-I and sTNFR-II, respectively.

[0124] As used herein, "R₁-[Cys¹⁹-Cys¹⁰³]-R₂" represents one or more proteins wherein [Cys¹⁹-Cys¹⁰³] is residues 19 through 103 of sTNFR-I, the sequence of which is provided in Figure 1 of WO 98/01555; wherein R₁ represents a methionylated or nonmethionylated amine group of Cys¹⁹ or one or more amino-terminal amino acid residues selected from Cys¹⁸ to Asp¹; and wherein R₂ represents a carboxy group of Cys¹⁰³ or one or more carboxy-terminal amino acid residues selected from Phe¹⁰⁴ to Leu¹¹⁰.

[0125] Exemplary truncated sTNFR-I's for use with the present invention include, but are not limited to, sTNFR-I 2.6D/C105, sTNFR-I 2.6D/C106, sTNFR-I 2.6D/N105, sTNFR-I 2.3D/d8, sTNFR-I 2.3D/d18, sTNFR-I 2.3D/d15, either methionylated or nonmethionylated, and variants and derivatives thereof. Certain exemplary truncated sTNFR-I's are described, e.g., in published PCT Application No. WO 98/01555.

[0126] As used herein, "R₃-[Cys³²-Cys¹¹⁵]-R₄" represents one or more proteins wherein [Cys³²-Cys¹¹⁵] is residues Cys³² through Cys¹¹⁵ of sTNFR-II, the sequence of which is provided in Figure 8 of WO 98/01555; wherein R₃ represents a methionylated or nonmethionylated amine group of Cys³² or one or more amino-terminal amino acid residues selected from Cys³¹ to Leu¹; and wherein R₄ represents a carboxy group of Cys¹¹⁵ or one or more carboxy-terminal amino acid residues selected from Ala¹¹⁶ to Arg¹²².

[0127] In certain aspects of the disclosure, the present invention is directed to an antibody to OPGL and at least one serine protease inhibitor for use in a therapy and methods of treatment using such therapies. In certain embodiments, an antibody to OPGL and a serine protease inhibitor and at least one additional molecule described herein is for use in a therapy.

[0128] Endogenous proteolytic enzymes may degrade invading organisms, antigen-antibody complexes, and certain tissue proteins that are no longer necessary or useful. Infective agents may introduce additional proteolytic enzymes into the organism. Protease inhibitors may regulate both endogenous and invading proteolytic enzymes.

[0129] In certain aspects of the disclosure, naturally occurring protease inhibitors serve to control endogenous proteases by limiting their reactions locally and temporally. In certain aspects of the disclosure, protease inhibitors may inhibit proteases introduced into the body by infective agents. In certain instances, tissues that are particularly prone to proteolytic attack and infection, including, but not limited to, those of the respiratory tract, are rich in protease inhibitors.

[0130] Protease inhibitors comprise approximately 10% of the human plasma proteins. At least eight inhibitors have been isolated from this source and characterized in the literature. These include, but are not limited to, alpha 2-macroglobulin (alpha 2M), alpha 1-protease inhibitor (alpha 1PI), alpha 1-antichymotrypsin (alpha 1Achy), alpha 1-antitrypsin (alpha 1AC), and inter-alpha-trypsin inhibitor (I-alpha-I).

[0131] In certain instances, a disturbance of the protease/protease inhibitor balance can lead to protease-mediated tissue destruction, including, but not limited to, emphysema, arthritis, glomerulonephritis, periodontitis, muscular dystrophy, tumor invasion, and various other pathological conditions. In certain situations, e.g. severe pathological processes such as sepsis or acute leukemia, the amount of free proteolytic enzymes present may increase due to the release of enzyme from secretory cells.

[0132] Furthermore, in certain instances, a diminished regulating inhibitor capacity of the organism may also cause alterations in the protease/protease inhibitor balance. A nonlimiting example of such a diminished regulating inhibitor capacity is an alpha 1-protease inhibitor deficiency, which is correlated with the development of pulmonary emphysema.

[0133] In certain instances, serious damage to the organism can occur when such aberrant conditions are present unless measures can be taken to control the proteolytic enzymes. Therefore, protease inhibitors have been sought that can be administered to an organism to control proteolytic enzymes.

[0134] Leukocyte elastase, trypsin, cathepsin G, and pancreatic elastase are nonlimiting examples of a class of proteases known as serine proteases.

[0135] In certain instances, leukocyte elastase, when released extracellularly, degrades connective tissue and other valuable proteins. While a normally functioning organism degrades a certain amount of connective tissue and other proteins, the presence of an excessive amount of leukocyte elastase may be associated with various pathological states, including, but not limited to, emphysema and rheumatoid arthritis. In certain aspects of the disclosure, to counteract the

effects of leukocyte elastase when it is present in amounts greater than normal, a protease inhibitor has been sought which is specific for leukocyte elastase. Such a protease inhibitor may be useful if it were capable of being isolated or prepared in a purified form and in sufficient quantities to be pharmaceutically useful.

5 **[0136]** Certain leukocyte elastase inhibitors are described, e.g., in Schiessler et al., "Acid-Stable Inhibitors of Granulocyte Neutral Proteases in Human Mucous Secretions: Biochemistry and Possible Biological Function", in *Neutral Proteases of Human Polymorphonuclear Leucocytes*, Havemann et al. (eds), Urban and Schwarzenberg, Inc. (1978), and in Travis and Salvesen, *Ann. Rev. Biochem.* 52: 655-709 (1983).

[0137] In certain instances, trypsin initiates degradation of certain soft organ tissue, such as pancreatic tissue, during a variety of acute conditions, including, but not limited to, pancreatitis. A trypsin inhibitor may be useful if it could be isolated and prepared in a purified form and in sufficient quantities to be pharmaceutically useful.

10 **[0138]** Cathepsin G is another protease present in leukocytes. Cathepsin G is capable of degrading a variety of proteins *in vitro*, including those of the complement pathway. Pancreatic elastase is another protease that may have a role in pancreatitis. Thus, inhibitors for these proteases may also be of pharmaceutical value.

[0139] In certain aspects of the disclosure, the substrate specificity and sensitivity to different inhibitors of serine proteases are believed to result from changes in only a few amino acid residues. By analogy, it may be possible to conceive of a class of serine protease inhibitors in which changes in a relatively few amino acids will result in inhibition of different proteases. In certain aspects of the disclosure, a member of this class inhibits every serine protease.

15 **[0140]** An exemplary serine protease inhibitor is secretory leukocyte protease inhibitor (SLPI) and fragments and analogs thereof. Exemplary serine protease inhibitors also include, but are not limited to, anti-leukoprotease (ALP), mucous protease inhibitor (MPI), human seminal plasma inhibitor-I (HUSH-I), bronchial mucus inhibitor (BMI), and cervical mucus inhibitor (CUSI). In certain aspects of the disclosure, a serine protease inhibitors also may be LPS modulator. See, e.g., Jin et al. (1997), *Cell* 88(3): 417-26. In certain embodiments, these molecules are well-suited for use in conditions leading to bone loss because they are preferentially directed to the cartilage.

20 **[0141]** Exemplary serine protease inhibitors are described, e.g., in U.S. Pat. No. 4,760,130; U.S. Pat. No. 5,900,400; and U.S. Pat. No. 5,633,227. The molecules disclosed in the foregoing references as well as any variants or analogues thereof are collectively termed "serine protease inhibitors."

[0142] IL-18 is a pro-inflammatory cytokine that was found to induce interferon- γ and was previously named interferon gamma inducing factor (IGIF). In certain instances, IL-1 has been shown to upregulate IL-18 production, and IL-18 induces production of a number of proinflammatory cytokines, including IL-6 and MMP-1. See, e.g., Dinarello et al. (1998), *J. Leukocyte Biol.* 63: 658-84. In certain instances, caspase 1 is also important for IL-18 production. Experiments also suggest that TNF- α regulates IL-18 production, and that simultaneous inhibition of TNF- α and IL-18 protects against liver toxicity. See, e.g., Faggioni et al. (2000), *PNAS* 97: 2367-72.

30 **[0143]** IL-18 acts *in vivo* through a receptor system reminiscent of the IL-1 system. IL-18 interacts with a cell surface receptor (IL-18R), which interacts with an accessory protein (IL-18RACp). IL-18-mediated signaling proceeds upon formation of the complex of IL-18, IL-18R, and IL-18RACp. A natural inhibitor for IL-18 is IL-18bp. In certain embodiments, IL-18bp acts as a "decoy receptor" by binding to IL-18 molecules and preventing interaction with IL-18R.

35 **[0144]** In certain aspects of the disclosure, the present invention is directed to an antibody to OPGL and at least one IL-18 inhibitor for use in a therapy and methods of treatment using such therapies. In certain aspects of the disclosure, an antibody to OPGL and an IL-18 inhibitor and at least one additional molecule described herein is for use in a therapy. Exemplary conditions that may be treated according to certain aspects of the disclosure include, but are not limited to, inflammation, autoimmune diseases, IL-1 mediated diseases, and TNF-mediated diseases. Exemplary conditions that may be treated with an antibody to OPGL and at least one IL-18 inhibitor according to certain aspects of the disclosure include, but are not limited to, arthritis, including, but not limited to rheumatoid arthritis; systemic lupus erythematosus (SLE); graft versus host disease (GvHD); hepatitis; sepsis; and the loss of bone and cartilage accompanying these diseases.

45 **[0145]** Exemplary IL-18 inhibitors include, but are not limited to, antibodies that bind to IL-18; antibodies that bind to IL-18R; antibodies that bind to IL-18RACp; IL-18bp; IL-18R fragments (e.g., a solubilized extracellular domain of the IL-18 receptor); peptides that bind to IL-18 and reduce or prevent its interaction with IL-18R; peptides that bind to IL-18R and reduce or prevent its interaction with IL-18 or with IL-18RACp; peptides that bind to IL-18RACp and reduce or prevent its interaction with IL-18R; and small molecules that reduce or prevent IL-18 production or the interaction between any of IL-18, IL-18R, and IL-18RACp.

50 **[0146]** Certain IL-18 inhibitors are described, e.g., in US Pat. No. 5,912,324, issued July 14, 1994; EP 0 962 531, published Dec. 8, 1999; EP 712 931, published Nov. 15, 1994; US Pat. No. 5,914,253, issued July 14, 1994; WO 97/24441, published July 10, 1997; US Pat. No. 6,060,283, issued May 9, 2000; EP 850 952, published Dec. 26, 1996; EP 864 585, published Sep. 16, 1998; WO 98/41232, published Sep. 24, 1998; US Pat. No. 6,054,487, issued April 25, 2000; WO 99/09063, published Aug 14, 1997; WO 99/22760, published Nov. 3, 1997; WO 99/37772, published Jan. 23, 1998; WO 99/37773, published March 20, 1998; EP 0 974 600, published Jan. 26, 2000; WO 00/12555, published Mar. 9, 2000; Japanese patent application JP 111,399/94, published Oct. 31, 1997; Israel patent application IL 121554

A0. published Feb. 8, 1998.

[0147] In certain aspects of the disclosure, an antibody to OPGL may be used with at least one therapeutic agent for inflammation. In certain aspects of the disclosure, an antibody to OPGL may be used with at least one therapeutic agent for an immune disorder. Exemplary therapeutic agents for inflammation and immune disorders include, but are not limited to, corticosteroids, including, but not limited to, prednisolone; nonsteroidal anti-inflammatory drugs (NSAIDs), including, but not limited to, cyclooxygenase type 1 (COX-1) and cyclooxygenase type 2 (COX-2) inhibitors; disease modifying antirheumatic drugs (DMARDs); including, but not limited to, methotrexate, hydroxychloroquine, chloroquine, cyclosporine, gold compounds (such as auranofin, aurothiomalate and aurothioglucose), and leflunomide; type IV phosphodiesterase inhibitors, including, but not limited to, Rolipram and Pentoxifylline; Tacrolimus (FK-506); Sirolimus (rapamycin); mycophenolic acid; 5-lipoxygenase inhibitors, including, but not limited to, Zileuton; modulators of interleukin-6 (IL-6); small molecule modulators of 38 kDa mitogen-activated protein kinase (p38-MAPK); small molecule modulators of intracellular molecules involved in inflammation pathways, wherein such intracellular molecules include, but are not limited to, jnk, IKK, NF- κ B, ZAP70, and Ick. Certain exemplary therapeutic agents for inflammation are described, e.g., in C.A. Dinarello and L.L. Moldawer Proinflammatory and Anti-inflammatory Cytokines in Rheumatoid Arthritis: A Primer for Clinicians Third Edition (2001) Amgen Inc. Thousand Oaks, CA. Certain exemplary therapeutic agents for inflammation and autoimmune diseases include, but are not limited to, interferon gamma (IFN- γ) modulators; modulators of OX40/OX40L (including soluble forms of OX40); modulators of 4-1 BB/4-1 BB ligand (including soluble forms of 4-1 BB); and modulators of B cell-T cell costimulatory pathways, including, but not limited to, modulators of the receptor ligand pairs CD28/B7, CD40/CD40L, ICOS/B7RP1, and AGP-3/TACI/BAFFR (AGP-3 binds to both TACI and BAFFR receptors). Certain exemplary modulators of B cell-T cell costimulatory pathways include, but are not limited to, inhibitors of CD28, B7.1, and B7.2 (including soluble forms of B7.1 or B7.2 and soluble forms of CTLA4, both of which may be fused to a heterologous peptide or protein which reduces or prevents degradation and/or increases half-life, reduces toxicity, reduces immunogenicity, or increases biological activity of a therapeutic protein by increasing solubility or circulating half-life); inhibitors of CD40 and CD40L (including soluble forms of CD40 which may be fused to a heterologous peptide or protein); inhibitors of ICOS and B7RP1 (including soluble forms of ICOS which may be fused to a heterologous peptide or protein) and inhibitors of AGP-3, TACI and BAFFR (including soluble forms of TACI and BAFFR). ICOS, B7RP1 and inhibitors thereof are described, e.g., in WO00/46240. AGP-3, TACI and BAFFR and inhibitors thereof are described, e.g., in WO00/47740, WO01/85872, WO02/15273, WO98/39361, and von Bulow and Bram (1997) Science 278:138-140.

[0148] In certain aspects of the disclosure, an antibody to OPGL is for use to treat bone loss, including, but not limited to, bone loss resulting from osteolytic destruction of bone caused by malignant or metastatic tumors. In certain aspects of the disclosure, an antibody to OPGL is for use to treat bone loss associated with cancer. Exemplary cancers include, but are not limited to, breast, prostate, thyroid, kidney, lung, esophageal, rectal, bladder, cervical, ovarian, and liver cancers, as well as cancer of the gastrointestinal tract. In certain aspects of the disclosure, an antibody to OPGL is for use to treat bone loss associated with, e.g., certain hematological malignancies, including, but not limited to, multiple myeloma and lymphoma, including Hodgkin's Disease.

[0149] In certain aspects of the disclosure, an antibody to OPGL is for administration alone. In certain aspects of the disclosure, an antibody to OPGL is for administration with at least one other therapeutic agent, including, but not limited to, at least one other cancer therapy agent. Exemplary cancer therapy agents include, but are not limited to, radiation therapy and chemo-therapy. In certain aspects of the disclosure, chemotherapy may involve treatment with one or more of the following: anthracyclines, taxol, tamoxifene, doxorubicin, 5-fluorouracil, and other drugs known in the art. In certain embodiments, a cancer therapy agent is a luteinizing hormone-releasing hormone (LHRH) antagonist. In certain embodiments, a LHRH antagonist is a peptide antagonist.

[0150] In certain aspects of the disclosure, an LHRH antagonist comprises the peptide: Ao-D-Nal-4-CI-Phe-O-Pal-Ser-N-Me-Tyr-D-Asn-Leu-Lys(iPr)-Pro-D-Ala-NH₂ (SEQ ID NO: 20), where Nal is 3-(2-naphthyl)alaninyl; 4-CI-Phe is (4'-chlorophenyl) alaninyl; Pal is 3-(3'-pyridyl)alaninyl; and Lys(iPr) is N-epsilon-2-propyl-lysinylnyl.

[0151] In certain aspects of the disclosure, an LHRH antagonist is an LHRH antagonist decapeptide. Certain exemplary decapeptides are described, e.g., in U.S. Patent No. 5,843,901.

[0152] Exemplary therapeutic antibodies according to certain embodiments include, but are not limited to, mouse, mouse-human chimeric, CDR-grafted, humanized and fully human antibodies, and synthetic antibodies, including, but not limited to, those selected by screening antibody libraries. Exemplary antibodies include, but are not limited to, those which bind to cell surface proteins Her2, CDC20, CDC33, mucin-like glycoprotein, and epidermal growth factor receptor (EGFR) present on tumor cells, and optionally induce a cytostatic and/or cytotoxic effect on tumor cells displaying these proteins. Exemplary antibodies also include HERCEPTIN™ (trastuzumab), which may be used to treat breast cancer and other forms of cancer, and RITUXAN™ (rituximab), ZEVALIN™ (ibrutinomab tiuxetan), and LYMPHOCIDE™ (epratuzumab), which may be used to treat non-Hodgkin's lymphoma and other forms of cancer. Certain exemplary antibodies also include ERBITUX™ (IMC-C225), BEXXAR™ (iodine 131 tositumomab), and Campath.

[0153] In certain aspects of the disclosure, cancer therapy agents are polypeptides which selective induce apoptosis in tumor cells, including, but not limited to, the TNF-related polypeptide TRAIL. In certain aspects of the disclosure, an

antibody to OPGL. may be administered at least one of prior to, concurrent with, and subsequent to treatment with a cancer therapy agent. In certain aspects of the disclosure, an antibody to OPGL may be administered prophylactically to prevent or mitigate the onset of bone loss by metastatic cancer. In certain aspects of the disclosure, an antibody to OPGL may be administered for the treatment of an existing condition of bone loss due to metastasis.

5 **[0154]** In certain aspects of the disclosure, an antibody to OPGL is for use to prevent and/or treat bone loss associated with multiple myeloma and/or to prevent and/or treat the disease itself. Multiple myeloma is a B cell derived tumor that may result in significant morbidity and/or mortality. In certain instances, a clinical manifestation of multiple myeloma is focal bone loss, which may be due to increased osteoclast activation in localized regions. Many myeloma patients present with bone lesions visible by radiological analysis and suffer from skeletal pain. In certain instances, patients with myeloma are susceptible to pathological fractures of involved bone, which may occur either spontaneously or due to injury. In certain instances, the skeletal lesions that occur during myeloma not only lead to bone fractures, but also deformity and occasionally nerve compression, particularly in the vertebral spine. In some patients, a pathological increase in serum calcium (hypercalcemia) occurs, and may cause significant problems during disease treatment. In certain aspects of the disclosure, an antibody to OPGL may be administered to patients to reduce or block bone resorption and release of calcium, which may reduce the risk of fractures and spinal deformity.

10 **[0155]** In certain instances, myeloma cells do not directly participate in bone destruction, but instead produce extra-cellular signals that lead to osteoclast differentiation and activation. In certain instances, osteoclasts produce high levels of the cytokine IL-6, particularly when they become activated. IL-6 is a B-cell growth factor, and contributes to the growth of both murine and human myeloma cells *in vitro*. Myeloma cells may also either directly or indirectly produce OPGL, which may result in local bone lysis surrounding the myeloma cells embedded in bone marrow spaces. In certain instances, the normal osteoclasts adjacent to the myeloma cell in turn produce IL-6, which may lead to local expansion of the tumor cells. Myeloma cells expand in a clonal fashion and may occupy bone spaces that are created by inappropriate bone resorption.

15 **[0156]** It has been observed that OPG administration in rodents induces rapid death of the osteoclast population (see, e.g., Lacey et al. (2000) Am. J. Pathol. 157:435-448). A reduction in the number of osteoclasts may counteract the effect of increased IL-6 production by those cells and may therefore affect the growth and survival of myeloma cells within trabecular bone. Thus, in certain aspects of the disclosure, administration of an antibody to OPGL to a myeloma patient may not only block the hyper resorption of bone, but may also affect the expansion and survival of the tumor itself.

20 **[0157]** B-cells express the receptor for OPGL, ODAR. Myeloma cells also express ODAR, and in addition may produce OPGL. In certain instances, the expression of both OPGL and ODAR in the same cell population may create an autocrine stimulus that affects survival of the myeloma cell. Thus, in certain aspects of the disclosure, administration of an antibody to OPGL may reduce tumor cell survival, thereby decreasing or eliminating the tumor burden seen in myeloma patients.

25 **[0158]** In certain embodiments, the invention provides for pharmaceutical compositions comprising a therapeutically effective amount of an antibody to OPGL together with a pharmaceutically acceptable diluent, carrier, solubilizer, emulsifier, preservative and/or adjuvant.

30 **[0159]** In certain embodiments, the invention provides for a pharmaceutical composition; the pharmaceutical composition may comprise a therapeutically effective amount of an antibody to OPGL of the invention and a therapeutically effective amount of at least one additional therapeutic agent, together with a pharmaceutically acceptable diluent, carrier, solubilizer, emulsifier, preservative and/or adjuvant. In certain embodiments, the at least one additional therapeutic agent may be selected from bone morphogenic factors designated BMP-1 through BMP-12; transforming growth factor- β (TGF- β) and TGF- β family members; interleukin-1 (IL-1) inhibitors, including, but not limited to, IL-1ra and derivatives thereof and Kineret™; anakinra TVF α inhibitors, including, but not limited to, a soluble TNF α receptor, Enbrel™, etanercept anti-TNF α antibodies. Remicade™, infliximab and D2E7 antibody; parathyroid hormone and analogs thereof, parathyroid related protein and analogs thereof; E series prostaglandins; bisphosphonates (such as alendronate and others); bone-enhancing minerals such as fluoride and calcium; non-steroidal anti-inflammatory drugs (NSAIDs), including COX-2 inhibitors, such as Celebrex™, celecoxib and Vioxx™; rofecoxib, immunosuppressants, such as methotrexate or leflunomide; serine protease inhibitors such as secretory leukocyte protease inhibitor (SLPI); IL-6 inhibitors (e.g., antibodies to IL-6), IL-8 inhibitors (e.g., antibodies to IL-8); IL-18 inhibitors (e.g., IL-18 binding protein or IL-18 antibodies); Interleukin-1 converting enzyme (ICE) modulators; fibroblast growth factors FGF-1 to FGF-10 and FGF modulators; PAF antagonists; keratinocyte growth factor (KGF), KGF-related molecules, or KGF modulators; matrix metalloproteinase (MMP) modulators; Nitric oxide synthase (NOS) modulators, including modulators of inducible NOS; modulators of glucocorticoid receptor; modulators of glutamate receptor; modulators of lipopolysaccharide (LPS) levels; and noradrenaline and modulators and mimetics thereof.

35 **[0160]** In certain embodiments, acceptable formulation materials preferably are nontoxic to recipients at the dosages and concentrations employed.

40 **[0161]** In certain embodiments, the pharmaceutical composition may contain formulation materials for modifying, maintaining or preserving, for example, the pH, osmolarity, viscosity, clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption or penetration of the composition. In certain embodiments, suitable formulation

materials include, but are not limited to, amino acids (such as glycine, glutamine, asparagine, arginine or lysine); anti-microbials; antioxidants (such as ascorbic acid, sodium sulfite or sodium hydrogen-sulfite); buffers (such as borate, bicarbonate, Tris-HCl, citrates, phosphates or other organic acids); bulking agents (such as mannitol or glycine); chelating agents (such as ethylenediamine tetraacetic acid (EDTA)); complexing agents (such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin or hydroxypropyl-beta-cyclodextrin); fillers; monosaccharides; disaccharides; and other carbohydrates (such as glucose, mannose or dextrans); proteins (such as serum albumin, gelatin or immunoglobulins); coloring, flavoring and diluting agents; emulsifying agents; hydrophilic polymers (such as polyvinylpyrrolidone); low molecular weight polypeptides; salt-forming counterions (such as sodium); preservatives (such as benzalkonium chloride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid or hydrogen peroxide); solvents (such as glycerin, propylene glycol or polyethylene glycol); sugar alcohols (such as mannitol or sorbitol); suspending agents; surfactants or wetting agents (such as pluronics, PEG, sorbitan esters, polysorbates such as polysorbate 20, polysorbate 80, triton, tromethamine, lecithin, cholesterol, tyloxapal); stability enhancing agents (such as sucrose or sorbitol); tonicity enhancing agents (such as alkali metal halides, preferably sodium or potassium chloride, mannitol sorbitol); delivery vehicles; diluents; excipients and/or pharmaceutical adjuvants. (Remington's Pharmaceutical Sciences, 18th Edition, A.R. Gennaro, ed., Mack Publishing Company (1990).

[0162] In certain embodiments, an antibody to OPGL and/or a therapeutic molecule is linked to a half-life extending vehicle known in the art. Such vehicles include, but are not limited to, the Fc domain, polyethylene glycol, and dextran. Such vehicles are described, e.g., in U.S. Application Serial No. 09/428,082 and published PCT Application No. WO 99/25044.

[0163] In certain embodiments, the optimal pharmaceutical composition will be determined by one skilled in the art depending upon, for example, the intended route of administration, delivery format and desired dosage. See, for example, *Remington's Pharmaceutical Sciences, supra*. In certain embodiments, such compositions may influence the physical state, stability, rate of *in vivo* release and rate of *in vivo* clearance of the antibodies of the invention.

[0164] In certain embodiments, the primary vehicle or carrier in a pharmaceutical composition may be either aqueous or non-aqueous in nature. For example, in certain embodiments, a suitable vehicle or carrier may be water for injection, physiological saline solution or artificial cerebrospinal fluid, possibly supplemented with other materials common in compositions for parenteral administration. In certain embodiments, neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. In certain embodiments, pharmaceutical compositions comprise Tris buffer of about pH 7.0-8.5, or acetate buffer of about pH 4.0-5.5, which may further include sorbitol or a suitable substitute therefor. In certain embodiments, a composition comprising an antibody to OPGL, with or without at least one additional therapeutic agents, may be prepared for storage by mixing the selected composition having the desired degree of purity with optional formulation agents (*Remington's Pharmaceutical Sciences, supra*) in the form of a lyophilized cake or an aqueous solution. Further, in certain embodiments, a composition comprising an antibody to OPGL, with or without at least one additional therapeutic agents, may be formulated as a lyophilizate using appropriate excipients such as sucrose.

[0165] In certain embodiments, the pharmaceutical compositions of the invention can be selected for parenteral delivery. In certain embodiments, the compositions may be selected for inhalation or for delivery through the digestive tract, such as orally. The preparation of such pharmaceutically acceptable compositions is within the skill of the art.

[0166] In certain embodiments, the formulation components are present in concentrations that are acceptable to the site of administration. In certain embodiments, buffers are used to maintain the composition at physiological pH or at a slightly lower pH, typically within a pH range of from about 5 to about 8.

[0167] In certain embodiments, when parenteral administration is contemplated, a therapeutic composition may be in the form of a pyrogen-free, parenterally acceptable aqueous solution comprising the desired antibody to OPGL, with or without additional therapeutic agents, in a pharmaceutically acceptable vehicle. In certain embodiments, a vehicle for parenteral injection is sterile distilled water in which the antibody to OPGL, with or without at least one additional therapeutic agent, is formulated as a sterile, isotonic solution, properly preserved. In certain embodiments, the preparation can involve the formulation of the desired molecule with an agent, such as injectable microspheres, bio-erodible particles, polymeric compounds (such as polylactic acid or polyglycolic acid), beads or liposomes, that may provide for the controlled or sustained release of the product which may then be delivered via a depot injection. In certain embodiments, hyaluronic acid may also be used, and may have the effect of promoting sustained duration in the circulation. In certain aspects of the disclosure, implantable drug delivery devices may be used to introduce the desired molecule.

[0168] In certain embodiments, a pharmaceutical composition may be formulated for inhalation. In certain embodiments, an antibody to OPGL, with or without at least one additional therapeutic agent, may be formulated as a dry powder for inhalation. In certain embodiments, an inhalation solution comprising an antibody to OPGL, with or without at least one additional therapeutic agent, may be formulated with a propellant for aerosol delivery. In certain embodiments, solutions may be nebulized. Pulmonary administration is further described in PCT application no. PCT/US94/001875, which describes pulmonary delivery of chemically modified proteins.

[0169] In certain embodiments, it is contemplated that formulations may be administered orally. In certain embodiments, an antibody to OPGL, with or without at least one additional therapeutic agents, that is administered in this fashion may

be formulated with or without those carriers customarily used in the compounding of solid dosage forms such as tablets and capsules. In certain embodiments, a capsule may be designed to release the active portion of the formulation at the point in the gastrointestinal tract when bioavailability is maximized and pre-systemic degradation is minimized. In certain embodiments, at least one additional agent can be included to facilitate absorption of the antibody to OPGL and/or any additional therapeutic agents. In certain embodiments, diluents, flavorings, low melting point waxes, vegetable oils, lubricants, suspending agents, tablet disintegrating agents, and binders may also be employed.

[0170] In certain embodiments, a pharmaceutical composition may involve an effective quantity of antibodies to OPGL, with or without at least one additional therapeutic agents, in a mixture with non-toxic excipients which are suitable for the manufacture of tablets. In certain embodiments, by dissolving the tablets in sterile water, or another appropriate vehicle, solutions may be prepared in unit-dose form. In certain embodiments, suitable excipients include, but are not limited to, inert diluents, such as calcium carbonate, sodium carbonate or bicarbonate, lactose, or calcium phosphate; or binding agents, such as starch, gelatin, or acacia; or lubricating agents such as magnesium stearate, stearic acid, or talc.

[0171] Additional pharmaceutical compositions will be evident to those skilled in the art, including formulations involving antibodies to OPGL, with or without at least one additional therapeutic agents, in sustained- or controlled-delivery formulations. In certain embodiments, techniques for formulating a variety of other sustained- or controlled-delivery means, such as liposome carriers, bio-erodible microparticles or porous beads and depot injections, are also known to those skilled in the art. See for example, PCT Application No. PCT/US93/00829 which describes the controlled release of porous polymeric microparticles for the delivery of pharmaceutical compositions. In certain embodiments, sustained-release preparations may include semipermeable polymer matrices in the form of shaped articles, e.g. films, or microcapsules. Sustained release matrices may include polyesters, hydrogels, polylactides (U.S. 3,773,919 and EP 058,481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman et al., Biopolymers, 22:547-556 (1983)), poly (2-hydroxyethylmethacrylate) (Langer et al., J. Biomed. Mater. Res., 15:167-277 (1981) and Langer, Chem. Tech., 12: 98-105 (1982)), ethylene vinyl acetate (Langer *et al.*, *supra*) or poly-D(-)-3-hydroxybutyric acid (EP 133,988). In certain embodiments, sustained release compositions may also include liposomes, which can be prepared by any of several methods known in the art. See e.g., Eppstein et al., Proc. Natl. Acad. Sci. USA, 82:3688-3692 (1985); EP 036,676; EP 088,046 and EP 143,949.

[0172] The pharmaceutical composition to be used for *in vivo* administration typically is sterile. In certain embodiments, this may be accomplished by filtration through sterile filtration membranes. In certain embodiments, where the composition is lyophilized, sterilization using this method may be conducted either prior to or following lyophilization and reconstitution. In certain embodiments, the composition for parenteral administration may be stored in lyophilized form or in a solution. In certain embodiments, parenteral compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[0173] In certain embodiments, once the pharmaceutical composition has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, or as a dehydrated or lyophilized powder. In certain embodiments, such formulations may be stored either in a ready-to-use form or in a form (e.g., lyophilized) that is reconstituted prior to administration.

[0174] The present disclosure includes kits for producing a single-dose administration unit. In certain aspects of the disclosure, the kits may each contain both a first container having a dried protein and a second container having an aqueous formulation. In certain aspects of the disclosure, kits containing single and multi-chambered pre-filled syringes (e.g., liquid syringes and lyosyringes) are included.

[0175] In certain aspects of the disclosure, the effective amount of a pharmaceutical composition comprising an antibody to OPGL, with or without at least one additional therapeutic agent, to be employed therapeutically will depend, for example, upon the therapeutic context and objectives. One skilled in the art will appreciate that the appropriate dosage levels for treatment, will thus vary depending, in part, upon the molecule delivered, the indication for which the antibody to OPGL, with or without at least one additional therapeutic agent, is being used, the route of administration, and the size (body weight, body surface or organ size) and/or condition (the age and general health) of the patient. In certain aspects of the disclosure, the clinician may titer the dosage and modify the route of administration to obtain the optimal therapeutic effect. In certain aspects of the disclosure, a typical dosage may range from about 0.1 $\mu\text{g}/\text{kg}$ to up to about 100 mg/kg or more, depending on the factors mentioned above. In certain aspects of the disclosure, the dosage may range from 0.1 $\mu\text{g}/\text{kg}$ up to about 100 mg/kg; or 1 $\mu\text{g}/\text{kg}$ up to about 100 $\mu\text{g}/\text{kg}$; or 5 $\mu\text{g}/\text{kg}$ up to about 100 mg/kg.

[0176] In certain aspects of the disclosure, the frequency of dosing will take into account the pharmacokinetic parameters of the antibody to OPGL and/or any additional therapeutic agents in the formulation used. In certain aspects of the disclosure, a clinician will administer the composition until a dosage is reached that achieves the desired effect. In certain aspects of the disclosure, the composition may therefore be administered as a single dose, or as two or more doses (which may or may not contain the same amount of the desired molecule) over time, or as a continuous infusion via an implantation device or catheter. Further refinement of the appropriate dosage is routinely made by those of ordinary skill in the art and is within the ambit of tasks routinely performed by them. In certain aspects of the disclosure, appropriate dosages may be ascertained through use of appropriate dose-response data.

[0177] In certain aspects of the disclosure, the route of administration of the pharmaceutical composition is in accord with known methods, e.g. orally, through injection by intravenous, intraperitoneal, intracerebral (intra-parenchymal), intracerebroventricular, intramuscular, intra-ocular, intraarterial, intraportal, or intralesional routes; by sustained release systems or by implantation devices. In certain aspects of the disclosure, the compositions may be administered by bolus injection or continuously by infusion, or by implantation device.

[0178] In certain aspects of the disclosure, the composition may be administered locally via implantation of a membrane, sponge or another appropriate material onto which the desired molecule has been absorbed or encapsulated. In certain aspects of the disclosure, where an implantation device is used, the device may be implanted into any suitable tissue or organ, and delivery of the desired molecule may be via diffusion, timed-release bolus, or continuous administration.

[0179] In certain aspects of the disclosure, it may be desirable to use a pharmaceutical composition comprising an antibody to OPGL, with or without at least one additional therapeutic agent, in an *ex vivo* manner. In such instances, cells, tissues and/or organs that have been removed from the patient are exposed to a pharmaceutical composition comprising an antibody to OPGL, with or without at least one additional therapeutic agent, after which the cells, tissues and/or organs are subsequently implanted back into the patient.

[0180] In certain aspects of the disclosure, an antibody to OPGL and/or any additional therapeutic agents can be delivered by implanting certain cells that have been genetically engineered, using methods such as those described herein, to express and secrete the polypeptides. In certain aspects of the disclosure, such cells may be animal or human cells, and may be autologous, heterologous, or xenogeneic. In certain aspects of the disclosure, the cells may be immortalized. In certain aspects of the disclosure, in order to decrease the chance of an immunological response, the cells may be encapsulated to avoid infiltration of surrounding tissues. In certain aspects of the disclosure, the encapsulation materials are typically biocompatible, semi-permeable polymeric enclosures or membranes that allow the release of the protein product(s) but prevent the destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissues.

EXAMPLES

[0181] The following examples, including the experiments conducted and results achieved are provided for illustrative purposes only and are not to be construed as limiting the present invention.

Example 1

Cloning the α OPGL-1 Heavy and Light Chains

[0182] CHO cells expressing the full-length human OPGL cDNA are used to immunize transgenic mice containing human immunoglobulin genes. Lymph nodes from the immunized mice are fused to murine myeloma cells to generate hybridomas. Supernatants from the hybridoma lines are tested in an ELISA assay for antibodies that react with human OPGL. Anti-OPGL expressing hybridoma lines AMG 6.5, AMG 6.4, and AMG 6.1 are found to express antibodies with high affinity for OPGL (Kd's of 0.28 nM, 0.29 nM, and 0.23 nM, respectively), and AMG 6.5 is selected for cloning. Heavy and light chain cDNA clones from AMG 6.5 and AMG 6.4 are identical, and AMG 6.5 is used to clone the α OPGL-1 light chain cDNA, while AMG 6.4 is used to clone the α OPGL-1 heavy chain cDNA.

Cloning of the α OPGL-1 light chain

[0183] The α OPGL-1 kappa light chain variable region is obtained using PCR amplification methods from first strand cDNA prepared from AMG 6.5 total RNA. First strand cDNA is prepared from AMG 6.5 total RNA using a random primer with an extended 5'-adapter (5'-GGCCGGATAGGCCTCACNNNNNT-3' (SEQ ID NO: 15)) and the materials and methods provided by the Gibco SuperScript II™ Pre-amplification System for First Strand cDNA Synthesis kit (cat. no. 18089-011). The oligonucleotides below are used for the PCR:

5' kappa RACE primer:

5' - GAT GAC CCA GTC TCC AGC CAC CCT G - 3' (SEQ ID NO: 5)

3' kappa RACE primer

5' - AAG GGT CAG AGG CCA AAG GAT GG - 3' (SEQ ID NO: 6)

[0184] The amplified DNAs, are cloned into pCRII-TOPO (Invitrogen) and the resulting plasmids are sequenced. The

kappa chain consensus sequence is used to design primers for PCR amplification of the full-length α OPGL-1 kappa chain. The 5' α OPGL-1 kappa primer incorporates a *Xba*I site (TCTAGA) for cloning and a "CCACC" Kozak sequence before the initiator Met codon. The 3' α OPGL-1 kappa primer incorporates a *Sa*I site (GTTCGAC) following the stop codon for cloning.

5' α OPGL-1 kappa primer:

5'-CAA CTC TAG A CC ACC ATG GAA ACC CCA GCG-3' (SEQ ID NO: 7)

***Xba*I Site Kozak M E T P A (SEQ ID NO: 16)**

3' α OPGL-1 kappa primer:

5'-TTT GAC GTC GAC TTA TCA ACA CTC TCC CCT GTT GAA G -3' (SEQ ID NO: 8)

***Sa*I Site * * C E G R N F (SEQ ID NO: 17)**

[0185] The full-length α OPGL-1 kappa chain cDNA clone is obtained using the AMG 6.5 first strand cDNA, described above, by PCR amplification with the 5' and 3' α OPGL-1 kappa primers. The PCR reaction generates a 738 bp fragment encoding the 235 amino acid residues (including the 20 amino acid kappa chain signal sequence) of the α OPGL-1 kappa chain (Figure 4, SEQ ID NO: 4). Following purification using a QIAquick PCR Purification kit (Qiagen cat. no.28104), this fragment is used to construct the kappa light chain expression vector.

[0186] The 738 bp full-length kappa fragment generated above is cut with *Xba*I and *Sa*I, is purified using the Promega Wizard DNA Clean-Up System (Promega cat no. A7100), and is cloned into pDSR α 19 to generate plasmid α OPGL-1-kappa/pDSR α 19 (Figure 5). pDSR α 19 has been described previously (see WO 90/14363 (see, e.g., Figure 12)). Briefly, to make pDSR α 19, pDSR α 2 is modified in the following ways: the FSH polyA is shortened by approximately 1400 base pairs, to 885 base pairs, and now ends at the *Nde*I site; the dihydrofolate reductase (DHFR) promoter now contains 209 base pairs, having been shortened from the 5' end by approximately 1 kilobase; and an approximately 550 base pair *Bgl*III fragment from the DHFR polyA sequence is deleted.

[0187] The α OPGL-1 kappa light chain expression clone is sequenced to confirm that it coded for the same peptide that is identified in the AMG 6.5 hybridoma. The final expression vector, α OPGL-1-kappa/pDSR α 19, is 5476 bp and contains the 7 functional regions shown in Table 2.

Table 2: Features of α OPGL-1-kappa/DSR α 19

Plasmid Base Pair Number.

| | |
|--------------|--|
| 2 to 881 | A transcription termination/polyadenylation signal from the α -subunit of the bovine pituitary glycoprotein hormone (α -FSH) (Goodwin, et al, Nucleic Acids Res. 1983 11: 6873-82; Genbank Accession Number X00004) |
| 882 to 2027 | A mouse dihydrofolate reductase (DHFR) minigene containing the endogenous mouse DHFR promoter, the cDNA coding sequences, and the DHFR transcription termination/polyadenylation signals (Gasser et al, Proc Natl Acad Sci U S A. 1982 79:6522-6.; Nunberg et al, Cell 1980 19: 355-64; Setzer et al, J Biol Chem. 1982 257: 5143-7; McGrogan et al, J Biol Chem. 1985 260: 2307-14) |
| 2031 to 3947 | pBR322 sequences containing the ampicillin resistance marker gene and the origin for replication of the plasmid in <i>E. coli</i> (Genbank Accession Number J01749) |
| 3949 to 4292 | An SV40 early promoter, enhancer and origin of replication (Takebe et al, Mol Cell Biol. 1988 8: 466-72., Genbank Accession Number J02400) |
| 4299 to 4565 | A translational enhancer element from the HTLV-1 LTR domain (Seiki et al, Proc Natl Acad Sci U S A. 1983 80: 3618-22, Genbank Accession Number J02029) |
| 4574 to 4730 | An intron from the SV40 16S, 19S splice donor/acceptor signals (Okayama and Berg, Mol Cell Biol. 1983 3: 280-9, Genbank Accession Number J02400) |
| 4750 to 5476 | The α OPGL-1 Kappa light chain cDNA between the <i>Xba</i> I and <i>Sa</i> I sites |

[0188] A circular plasmid map of the vector is shown in Figure 5.

Cloning of the α OPGL-1 heavy chain

5 [0189] The α OPGL-1 IgG2 heavy chain is cloned from AMG 6.4 hybridoma double-stranded cDNA produced with the Clontech Marathon™ cDNA Amplification Kit (cat. no. K1802-1). Amplification of AMG 6.4 heavy chain cDNA is accomplished by 5' and 3' rapid amplification of cDNA ends (RACE) techniques performed with human germline IgG2 heavy chain constant region specific primers (shown below) and RACE primers and other materials and methods provided in the Marathon™ cDNA amplification kit.

10 5' IgG2 RACE primer
5'- GGC ACG GTC ACC ACG CTG CTG AG -3' (SEQ ID NO: 9)
3' IgG2 RACE primer
5'- CCT CCA CCA AGG GCC CAT CGG TCT -3' (SEQ ID NO: 10)

15 [0190] The 600 bp 5' RACE product and 1200 bp 3' RACE product are cloned into pCR2.1 (Invitrogen) and are sequenced. This sequence information is used to design α OPGL-1 heavy chain specific primers for the cloning of the full-length sequence. The heavy chain 5' primer (5' α OPGL-1 IgG2 Primer) is directed against the sense strand and has a HindIII site and consensus Kozak sequence before the natural start site. The heavy chain 3' primer (3' α OPGL-1 IgG2 Primer) is an antisense primer that contains a Sall site and stop codon after the last amino acid of the heavy chain IgG2 sequence.

20 5' α OPGL-1 IgG2 Primer:

5'- CAGAAGCTTGACCACC ATG GAG TTT GGG CTG AGC TGG CTT TTT CTT GTG GC -3'

25 (SEQ ID NO: 11)

HindIII Kozak M E F G L S W L F L V A

(SEQ ID NO: 18)

30 3' α OPGL-1 IgG2 Primer:

5'- GCA TGTCGAC TTA TCA TTT ACC CGG AGA CAG GGA GAG -3' (SEQ ID NO: 12)

35 *Sall* * * K G P S L S L (SEQ ID NO: 19)

[0191] The double-stranded cDNA described above is used to generate the full-length heavy chain cDNA by PCR amplification with the 5'- and 3'- α OPGL-1 IgG2 primers. The PCR generates a 1433 bp fragment encoding the 467 amino acid residues (including the 19 amino acid IgG signal sequence) of the α OPGL-1 IgG2 heavy chain protein (Figure 2, SEQ ID NO: 2). Following purification using a QIAquick PCR Purification kit (Qiagen cat. no.28104), this fragment is used to construct the heavy chain expression vector as follows.

40 [0192] DNA encoding the full-length IgG2 heavy fragment generated above is cut with *HindIII* and *Sall*, purified using a QIAquick Gel Extraction kit (Qiagen cat. no.28704), and this fragment is cloned into pDSRa19. The resulting expression plasmid is named α OPGL-1-IgG2/pDSRa19 (Figure 6). All vector components are identical to α OPGL-1-kappa/pDSRa19 vector, described above, except the α OPGL-1 IgG2 heavy chain cDNA replaces the α OPGL-1 kappa light chain cDNA between the *XbaI* and *Sall* sites. The α OPGL-1 IgG2 heavy chain expression clone is sequenced to confirm that it coded for the same polypeptide that is identified in the AMG 6.4 hybridoma.

50 **Example 2**

α OPGL-1 expression in CHO cells

[0193] Stable expression of α OPGL-1 antibody is achieved by co-transfection of α OPGL-1-kappa/pDSRa19 and α OPGL-1-IgG2/pDSRa19 plasmids into dihydrofolate reductase deficient (DHFR-) Chinese hamster ovary cells (CHO AM-1/D, U.S. Patent No. 6,210,924) followed by isolation and testing of individual clones.

[0194] A 100 mm tissue culture dish is plated with 1.5×10^6 AM-1/D cells grown in CHO d⁻ medium (DMEM-high glucose, 10% fetal bovine serum, 1% penicillin/ streptomycin/glutamine, 1 X NaPyruvate, 1% nonessential amino acids (NEAA))

(Gibco®) and 1% ht supplement (Gibco®)) the day before transfections (Day 0). On day one, 400 µl of serum-free RPMI 1640 medium (Gibco®) is aliquoted into a 12x75 mm polypropylene tube. Twenty four microliters of *TransIT*®-LT1 reagent (Mirus Corporation) is added dropwise to the medium and the mixture is allowed to incubate at room temperature for 10 minutes. A total of 15 µg of linearized plasmid DNA (7.5 µg of αOPGL-1-kappa/pDSRα19 and 7.5 µg of αOPGL-1 -IgG2/pDSRα19, digested with Pvu1) is then added dropwise to the mixture and is incubated at room temperature for 10 minutes.

[0195] The CHO d- medium is removed from the cells, which are washed with 10 ml of Dulbecco's phosphate buffered saline (Gibco®). Six milliliters of serum-free MEM medium supplemented with HT, L-glu, NEAA, and Na pyruvate (Gibco®) is added to the cells. The DNA/LT1 complex is added dropwise to the plates, which are gently rocked back and forth to distribute the DNA evenly over the cells. After 6 hours in a tissue culture incubator, the medium is replaced with fresh CHO d- medium. Forty eight hours later the cells are split to ten 100 mm culture dishes in CHO select medium (DMEM high glucose, 10% dialyzed fetal bovine serum (FBS), 1% penicillin /streptomycin /glutamine, 1% nonessential amino acids and 1X Na pyruvate) (Gibco®). Medium is changed twice weekly until colonies appeared.

[0196] After 10-14 days, colonies are picked using 5 mm cloning discs (Labcore®) soaked in 1x trypsin-EDTA (Gibco®) and are cultured in 24 well tissue culture plates with CHO select medium. When the cells become confluent, serum free media (CHO select medium minus FBS) is added and is then collected 48 hours later. These conditioned media are analyzed for antibody expression by Western blot with horse radish peroxidase (HRP)-conjugated goat anti-human IgG Fc antibody (Pierce, Rockford, IL) to detect the αOPGL-1 heavy chain, and goat anti-human kappa chain antibody (Pierce, Rockford, IL) followed by HRP-conjugated rabbit anti-goat IgG(H+L) antibody (Pierce, Rockford, IL) to detect the αOPGL-1 light chain. The highest expressing clones are expanded and stored in liquid nitrogen.

Example 3

Production of αOPGL-1

Preparation and Creation of Cell Line 125Q

[0197] CHO cells producing αOPGL-1 are cloned by two rounds of limiting dilution in 96 well plates under serum-free conditions. The clones are selected based on production and growth characteristics in various suspension vessels. EIAs are performed to select the clone that produces the highest level of αOPGL-1. Growth characteristics, including doubling times and densities are then measured by growing the clones in 100 ml, 250 ml, 500 ml, 1 L, and 3 L spinner flasks, as well as in 3L Aplikon bioreactors. The clone with the fastest doubling time that reaches the highest density in culture is selected, and is designated Cell Line 125Q. When the clone has expanded to yield sufficient cells to freeze 360 ampules at approximately 1x10⁷ cells/mL, cells are resuspended in a cryopreservative, serum-free medium (90% VM-Soy Batch Medium (see Table 3 for details) supplemented with 10 ml/L nonessential amino acids and 10 ml/L L-Glutamine (Gibco/LTI/Invitrogen), and 10% dimethyl sulfoxide (JT Baker)) and frozen. Ampules are stored in a limited access facility and are submerged in liquid nitrogen in liquid nitrogen dewars.

[0198] Based on growth and production in small-scale spinners and larger scale bioreactors, Cell Line 125Q is chosen as the cell line for manufacturing of αOPGL-1.

Cell Culture

[0199] αOPGL-1 is produced by expression in Cell Line 125Q, a clonal line of CHO cells that expresses αOPGL-1 from plasmids αOPGL-1-kappa/pDSRα19 and αOPGL-1 -IgG2/pDSRα19. The cell culture process for αOPGL-1 is shown in Figure 19. For each production run, cells from a vial of Cell Line 125Q are initially grown in 50 mL of VM-Soy Batch Medium (see Table 3 for composition) supplemented with 10 ml/L nonessential amino acids and 10 ml/L L-glutamine (Gibco/LTI/Invitrogen) (VM-Soy Supp) in 125 ml erlenmeyer shakers at 100 rpm for 5 days. The entire culture is then used to inoculate VM-Soy Supp in a 500 ml spinner flask to 3x10⁵ viable cells/ml (3E5 vc/ml), and is grown with 70 rpm spinning for 3-4 days. The entire culture from the 500 ml spinner flask is then used to inoculate VM-Soy Supp in a 3L spinner flask to 3E5 vc/ml, and is grown with 70 rpm spinning for 3-4 days.

[0200] The culture from the 3L spinner flask is then split to two 3 L spinner flasks at 3E5 vc/ml in VM-Soy Supp without phenol red and grown under the same conditions. These spinner flask cultures are then used to inoculate four additional spinner flasks at 3E5 vc/ml in VM-Soy Supp without phenol red, and grown under the same conditions. Four liters of culture from the four 3L spinner flasks is used to inoculate 10 L of VM-Soy Supp without phenol red in a 20 L bioreactor, and the bioreactor is run in fed-batch mode for 7-10 days. In fed-batch mode, a nutrient feed containing concentrated media components ("Feed", as set forth below in Table 3) is added to maintain cell growth and culture viability.

[0201] The entire culture from the 20L bioreactor is then used to inoculate 70 L of VM-Soy Supp without phenol red in a 150L bioreactor, and the bioreactor is run in fed-batch mode for 9-10 days. Finally, the entire culture from the 150L

EP 1 409 016 B9

bioreactor is used to inoculate approximately 880 L of VM-Soy (without supplement or phenol red) in a 2000L bioreactor, and the bioreactor is run in fed-batch mode. The rate of feed during fed-batch mode is determined such that the glucose level in the culture is maintained at 0.6 g/L for each bioreactor. The cell density and glucose concentration are measured daily and the rate of feed adjusted accordingly.

[0202] Production in the 2000L bioreactor lasts for approximately two weeks during which time α OPGL-1 is constitutively produced by the cells and secreted into the cell culture medium.

[0203] The production reactor is controlled at set pH, temperature, and dissolved oxygen level: pH is 7.0 and is controlled by carbon dioxide gas and sodium carbonate addition; dissolved oxygen is 120 mmHg and is controlled by air, nitrogen, and oxygen gas flows. Cells are maintained at 37°C throughout the process. All gases are passed through membrane filters of pore size 0.22 μ m or less.

[0204] At the end of production, the cell broth is fed into a disk stack centrifuge and the culture supernatant is separated from the cells. The centrate is further clarified through a Cuno 90SP depth filter followed by a 0.2 μ m Posidyne filter (Pall Co.). The clarified conditioned media is then concentrated by tangential flow ultrafiltration (UF) using 50kD NMWL membranes (Millipore Biomax 50). The conditioned media is concentrated 15- to 30- fold. The resulting concentrated conditioned medium (CCM) is then either processed through purification or frozen for purification at a later date. The production process is summarized in Figure 19.

Cell Culture Medium

[0205] The cell culture medium for use throughout the entire cell culture process is based on Dulbecco's Modified Eagle's Medium /Ham's Nutrient F12 (DMEM/F12, 1:1), and contains supplemental levels of amino acids, additional nutrients and salts, a soy hydrolysate and recombinant human insulin (Nucellin[®]Zn, Eli Lilly). The components are listed in Table 3. This media is referred to as VM-Soy. Media solutions are filtered through membrane filters of 0.2 μ m pore size prior to use.

Table 3. Cell Culture Media Components

| COMPONENTS FOR BASAL Media and FEEDS | | |
|--|----------------------------------|--------------------|
| COMPONENT | VMSoy Batch Medium (mg/L) | FEED (mg/L) |
| DMEM/F12 COMPONENTS | | |
| Inorganic salts | | |
| CaCl ₂ (anhyd.) | 116.60 | 233.2 |
| CuSO ₄ .5H ₂ O | 0.0026 | 0.0052 |
| Fe(NO ₃) ₃ .9H ₂ O | 0.1000 | 0.2 |
| FeSO ₄ .7H ₂ O | 0.8340 | 1.668 |
| KCl | 311.80 | 623.6 |
| MgCl ₂ (anhyd.) | 57.280 | 114.56 |
| MgSO ₄ (anhyd.) | 97.680 | 195.36 |
| NaCl | 905.990 | 1811.98 |
| NaH ₂ PO ₄ .H ₂ O | 125.00 | 250 |
| Na ₂ HPO ₄ | 142.040 | 284.08 |
| ZnSO ₄ .7H ₂ O | 0.8640 | 1.728 |
| Other Components | | |
| D-Glucose | 3151.00 | 12302 |
| Na Hypoxanthine | 5.40 | 10.8 |
| Linoleic acid | 0.090 | 0.18 |
| Lipoic acid | 0.2060 | 0.412 |
| Phenol Red | 8.10 | 16.2 |
| Putrescine.2HCl | 0.1620 | 0.324 |
| Sodium Pyruvate | 110.00 | 220 |
| Amino acids | | |
| L-Alanine | 26.70 | 53.4 |
| L-Arginine HCl | 295.00 | 590 |
| L-Asparagine.H ₂ O | 45.00 | 90 |

EP 1 409 016 B9

(continued)

| | | | |
|----|----------------------------------|----------|---------|
| | Amino acids | | |
| 5 | L-Aspartic acid | 39.90 | 79.8 |
| | L-Cysteine.HCl.H ₂ O | 35.120 | 70.24 |
| | L-Cystine.2HCl | 62.580 | 125.16 |
| | L-Glutamic acid | 44.10 | 88.2 |
| | L-Glutamine | 657.00 | 1314 |
| 10 | Glycine | 52.50 | 105 |
| | L-Histidine.HCl.H ₂ O | 62.950 | 125.9 |
| | L-Isoleucine | 108.940 | 217.88 |
| | L-Leucine | 118.10 | 236.2 |
| | L-Lysine HCl | 182.50 | 365 |
| 15 | L-Methionine | 34.480 | 68.96 |
| | L-Phenylalanine | 70.960 | 141.92 |
| | L-Proline | 57.50 | 115 |
| | L-Serine | 73.50 | 147 |
| 20 | L-Threonine | 106.90 | 213.8 |
| | L-Tryptophan | 18.040 | 36.08 |
| | L-Tyrosine.2Na.2H ₂ O | 111.580 | 223.16 |
| | L-Valine | 105.70 | 211.4 |
| | Vitamins | | |
| 25 | Biotin | 0.0073 | 0.0146 |
| | D-Ca Pantothenate | 4.480 | 8.96 |
| | Choline Chloride | 17.960 | 35.92 |
| | Folic Acid | 5.30 | 10.6 |
| 30 | i-Inositol | 25.20 | 50.4 |
| | Niacinamide | 4.040 | 8.08 |
| | Pyridoxal HCl | 4.00 | 8 |
| | Pyridoxine HCl | 0.0620 | 0.124 |
| | Riboflavin | 0.4380 | 0.876 |
| 35 | Thiamine HCl | 4.340 | 8.68 |
| | Thymidine | 0.3635 | 0.727 |
| | Vitamin B12 | 1.360 | 2.72 |
| | ADDITIONAL COMPONENTS | | |
| 40 | Nucellin Zn, (rhu insulin) | 5.00 | 15 |
| | Selenous Acid | 0.0050 | 0.015 |
| | Ethanolamine | 0.0012 | 0.0037 |
| | Triiodothyronine | 0.000040 | 0.00012 |
| 45 | Hydrocortisone | 0.020 | 0.06 |
| | Ferric Citrate | 122.450 | 122.450 |
| | Pluronic F-68 | 1000.00 | 500 |
| | Soy Hydrolysate | 6000.00 | 6000.00 |
| | NaHCO ₃ | 3000.00 | 3000.00 |
| 50 | NaCl | 3500.00 | |

Purification Process

55 **[0206]** αOPGL-1 expressed in CHO cells is secreted into the extracellular medium. A series of steps may be used to generate pure material. The process uses hydrophobic charge induction, cation exchange, and hydrophobic interaction chromatography along with a low pH step and viral filter. These procedures are described below.

A. Hydrophobic Charge Induction Chromatography (HCIC)

5 [0207] This chromatography step removes the majority of host cell proteins and DNA. The concentrated conditioned media (CCM) is filtered through a Cuno 30SP filter and then through a Cuno VR07 charged cellulose-based filter, and then loaded on to an MEP HyperCel resin. After loading, the column is washed with equilibration buffer (20 mM Tris pH 7.2). The antibody is eluted from the resin using a low pH buffer (20 mM Sodium Acetate, pH 5.0). As it is eluted from the column, the product is collected based on the absorbance at 280 nm of the column effluent.

B. Viral Inactivation

10 [0208] The MEP pool is titrated to pH 3.7 and is held for approximately 60 minutes to inactivate potentially contaminating retrovirus. Following the hold step, the pH is adjusted to approximately 6.0.

C. Viral Filtration

15 [0209] The pH-adjusted pool is filtered through a Millipore Viresolve NFR filter or equivalent. The antibody flows through the filter while potentially contaminating viruses ≥ 50 nm are retained.

D. Cation Exchange Chromatography (CEX)

20 [0210] The antibody is further purified by cation exchange chromatography using SP Sepharose HP (Amersham Pharmacia) or equivalent. The cation exchange chromatography step removes additional CHO cell proteins, DNA, lower molecular weight proteins, and aggregated forms of α OPGL-1. The viral filtered pool is loaded onto the cation exchange resin. After loading, the column is washed with equilibration buffer (20 mM NaMES pH 6.2). The antibody is then eluted with a linear gradient of increasing salt (20 mM NaMES pH 6.2, 0 M NaCl to 20 mM NaMES pH 6.2, 0.3 M NaCl). As it is eluted from the column, the product is collected based on the absorbance at 280 nm of the column effluent.

E. Hydrophobic Interaction Chromatography (HIC)

30 [0211] The antibody is further purified by hydrophobic interaction chromatography using Phenyl Toyopearl 650S (Tosoh Biosep) or equivalent. The hydrophobic interaction chromatography step is used as a polishing step and removes additional CHO cell proteins, DNA, lower molecular weight proteins, and aggregated forms of α OPGL-1. The cation exchange pool is conditioned before loading onto the column by addition of ammonium sulfate to a conductivity of >105 mS/cm at 15-25°C. After loading, the column is washed with the equilibration buffer (1 M Potassium Phosphate pH 8). The antibody is then eluted with a linear gradient of decreasing salt concentration (1 M Potassium Phosphate, 0mM Tris pH 8 to 0M Potassium Phosphate, 20 mM Tris pH 8). As it is eluted from the column, the product is collected based on the absorbance at 280 nm of the column effluent.

F. Concentration and Diafiltration

40 [0212] The HIC column pool is concentrated and diafiltered into formulation buffer by tangential flow ultrafiltration using 50kD NMWL membranes (Millipore Biomax 50). The formulation buffer includes 10 mM Acetate, 5% Sorbitol, pH 5.2 and α OPGL-1 is at 30 mg/mL.

Final Filtration and Storage

45 [0213] The purified bulk is passed through a 0.22 μ m PVDF filter (Millipore), is sampled, and stored at approximately -30°C in a secured freezer.

Example 4Binding Specificity of α OPGL-1

50 [0214] Antibodies that are produced in CHO cells that are transfected with the two expression vectors as discussed in Examples 1 and 2 may be used in the following examples 4, 5, and 6.

[0215] Human OPG binds and neutralizes OPGL in rats, mice and cynomolgus monkeys, as well as in humans. α OPGL-1 binds human OPGL with high affinity but does not bind significantly to murine OPGL (Table 4).

Table 4: Affinity of α OPGL-1 to Cell Membrane Expressed OPGL of Human, Cynomolgus Monkey, or Mouse Sequence.

| OPGL Species | α OPGL-1 ED ₅₀ (ng/ml) |
|--------------|--|
| Human | 16 |
| Cynomolgus | 19 |
| Mouse | No Specific Binding |

OPGL of these species is expressed in CHO cells as the full-length, membrane-bound protein. Binding of α OPGL-1 to the cell surface expressed OPGL is assessed by FACS analysis of cells incubated with α OPGL-1 and a FITC-labeled secondary antibody to human IgG2. α OPGL-1 binds human and cynomolgus OPGL but there is no specific binding to mouse OPGL.

[0216] In addition, human OPG has been reported to show weak binding to tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) (Truneh et al, 2000), a related member of the TNF family, which shows DNA and amino acid sequence homology to OPGL (Lacey et al., 1998). However, OPG does not detectably bind to other TNF-related proteins such as TNF α , TNF β , or CD40 ligand.

[0217] α OPGL-1 binds specifically to OPGL on EIA plates (Figure 7). Recombinant soluble OPGL (2 μ g/ml) is coated onto 96-well EIA plates at room temperature for 16 to 24 hours. After blocking with 1% BSA in PBS, varying concentrations of α OPGL-1 (approximately 2 ng/ml to 1000 ng/ml) diluted in 1% BSA/PBS are added to the wells and the plates are incubated for about 2 hours at room temperature. Bound antibody is detected with goat anti-Human IgG (Fab')-HRP using TMB-H₂O₂ (tetramethylbenzidine and hydrogen peroxide) substrate cocktail. The absorbance is read at 450nm and 650nm.

[0218] α OPGL-1 binds specifically OPGL expressed on the surface of transfected cells (Figure 8). α OPGL-1 (100 ng/ml) diluted in FACS Buffer (PBS, 0.1%BSA, 0.01% Sodium Azide) is preincubated with varying concentrations of OPGL, TNF α , TNF β , TRAIL, or CD40 ligand (approximately 0.1 ng/ml to 1000 ng/ml), and is then added to approximately 200,000 CHO REN 218-9 cells, which are CHO cells stably expressing membrane-bound OPGL on the cell surface. After 1 hour at 2-8°C, unbound antibody is removed by centrifugation and washing. Cells are then incubated for 30 minutes at 2-8°C with FITC-labeled F(ab')₂ Goat anti-Human IgG (Fc γ fragment specific). After centrifugation and washing, cell surface fluorescence is measured using flow cytometry. Figure 8 shows that binding of α OPGL-1 to CHO REN 218-9 cells is specific, and is competitively reduced by addition of soluble OPGL, but not by addition of TNF α , TNF β , TRAIL, or CD40 ligand.

[0219] In competition experiments, α OPGL-1 binding to OPGL on EIA plates is inhibited by addition of exogenous OPGL (Figure 9), but not by the addition of TNF α , TNF β , TRAIL, or CD40 ligand (Figure 10). This procedure is performed in substantially the same manner as above, for binding of α OPGL-1 to OPGL on EIA plates, except a constant concentration of α OPGL-1 (100ng/mL) is preincubated with varying concentrations of soluble OPGL or other ligands (approximately 1 ng/ml to 1000 ng/ml for each) before it is added to the OPGL-coated plates.

Example 5

α OPGL-1 Neutralizing Activity

Inhibition of Osteoclast Formation

[0220] RAW 264.7 (ATCC No. TIB-71, Manassas, VA) is a murine macrophage cell line that was derived from an Abelson murine leukemia virus-induced tumor. RAW 264.7 cells will differentiate to osteoclast-like cells in the presence of OPGL. The basic assay for generation of osteoclasts in culture from RAW cells in the presence of OPGL has been described in detail in Simonet et al (1997) Cell 89 p. 309, and Lacey et al (1998) Cell 93 p. 165.

[0221] RAW cells are stimulated by ligand to differentiate into osteoclast-like cells, and the differentiation can be measured by TRAP activity, a property of osteoclasts. Thus, the effect of α OPGL-1 on osteoclastogenesis can be measured.

[0222] RAW cells are incubated for 4 days in the presence of a constant amount of OPGL (40 ng/ml) and varying amounts of α OPGL-1 (6.3 ng/ml to 200 ng/ml) in cell culture medium (DMEM, 10% FBS, 0.292 mg/ml L-Glut, 100 units/ml Penicillin G, 100 μ g/ml Streptomycin sulfate). At the end of 4 days, the cells are stained for tartrate-resistant acid phosphatase (TRAP) activity by permeabilization and acidification, followed by treatment with para-nitrophenylphosphate for 5 minutes. Briefly, the media is aspirated off of the cells, and 100 μ l of citrate buffer (410 ml 0.1M citric acid, 590 ml 0.1 M citrate, trisodium salt, 1 mL triton X-100) is added to each well and the plates are incubated for 3 to 5 minutes at

room temperature. One hundred microliters of PNPP solution is then added (157.8 mg acid phosphatase reagent (Sigma 104-100), 7.2 ml tartrate solution (Sigma cat. no. 387-3), and 22.8 ml citrate buffer), and plates are incubated for 3 to 5 minutes at room temperature. The reaction is terminated by addition of 50 μ l 0.5 M NaOH solution.

[0223] TRAP will convert para-nitrophenylphosphate to para-nitrophenol, which can be quantitated by optical density measurement at 405 nm. The TRAP activity, which is a surrogate marker for osteoclast development, therefore correlates with the optical density at 405 nm. A plot of optical density versus α OPGL-1 concentration is shown in Figure 11, and demonstrates that α OPGL-1 inhibits osteoclast formation in this assay.

Inhibition of OPGL Binding to its Receptor

[0224] The potency of α OPGL-1 is demonstrated by its ability to block the binding of OPG ligand to its cognate receptor, the osteoclast differentiation and activating receptor (ODAR, also known as RANK). This assay uses homogeneous time resolved fluorescent resonance (HTRF) to detect binding of α OPGL-1 to Europium-conjugated osteoprotegerin ligand (Eu-OPGL). If α OPGL-1 inhibits Eu-OPGL binding to ODAR, fluorescent output will decrease, and the amount of α OPGL-1 present will be inversely related to the amount of fluorescence.

[0225] OPGL is labeled with europium, which emits light at 620 nm when excited with 337 nm light. ODAR is fused to FLAG and to Fc, and the Fc-ODAR-FLAG fusion protein is labeled with an anti-FLAG antibody linked to allophycocyanin (APC), a fluorophore which emits 665 nm light when excited by light at 620 nm. Therefore, when Eu-labeled OPG ligand binds to the Fc-ODAR-FLAG/anti-FLAG-APC complex, the tertiary complex will emit 665 nm light when excited with light at 337 nm.

[0226] Eu-OPGL at 0.05 μ g/ml is preincubated with various concentrations (0.1 to 150 ng/ml) of α OPGL-1 in assay buffer (50 mM Tris pH 8, 100 mM NaCl, 0.05% NaN₃, 0.1% BSA, and 0.05% Tween 20) at room temperature for approximately one hour (Preincubation mix). A mixture of Fc-ODAR-FLAG (1 μ g/ml) and anti-FLAG-APC (2.5 μ g/ml) is also prepared in assay buffer and incubated at room temperature for one hour (Fluorochrome mix). Equal volumes of Preincubation mix and Fluorochrome mix are then combined and incubated at room temperature for 3 hours. The fluorescence is measured by reading plates on the Packard Discovery HTRF microplate analyzer using an excitation wavelength of 337 nm and an emission wavelength of 665 nm.

[0227] When α OPGL-1 is preincubated with Eu-OPG ligand and is then mixed with Fc-ODAR-FLAG/anti-FLAG-APC, the fluorescence intensity at 665 nm decreases in a dose-dependent manner, as shown in Figure 12, demonstrating that α OPGL162 can effectively inhibit OPGL binding to ODAR.

Example 6

Pharmacokinetics in Cynomolgus Monkeys

[0228] Six male and six female cynomolgus monkeys, not greater than 4.5 years of age and weighing 2 to 4 kg are assigned to 4 dose groups. Group 1 consists of 3 males and 3 females. Groups 2, 3, and 4 each consists of 1 male and 1 female. Animals in Group 1 are administered a single SC dose of 1 mg/kg α OPGL-1, while animals in Groups 2, 3 and 4 are administered single IV doses of 0.1, 1.0, or 10.0 mg/kg of α OPGL-1, respectively.

[0229] Animals are dosed with α OPGL-1 expressed from transfected chinese hamster ovary (CHO) cells. Serum samples are taken for determination of α OPGL-1 levels, antibody analysis, and analysis of the bone turnover marker serum N-telopeptide (serum N-Tx), alkaline phosphatase (ALP), and serum calcium (serum Ca). Urine is also collected for analysis of N-telopeptide (urine N-Tx) and creatinine.

[0230] The serum concentration-time profiles following IV administration are characterized by a tri-phasic distribution (Figure 13). Initially, there is a rapid distribution phase, followed by a significantly slower plateau phase, which appears to be concentration-dependent. The third observed phase is a rapid elimination phase.

[0231] Non-compartmental analysis of complete serum concentration-time profiles using WinNonlin Professional (v1.5), and exponential analysis of the data up to 14 days after test article administration and above 10,000 ng/mL using SAAM II (v1.1.2) are utilized to investigate the pharmacokinetics of α OPGL-1 in monkeys. The initial volume of distribution from all IV doses averages 28.9 ml/kg, similar to plasma volume. Steady state volume (V_{ss}) of distribution averages 39 ml/kg across all IV doses. Exponential analysis indicates that α OPGL-1 has an average distribution half-life ($t_{1/2\alpha}$) of 6.02 hours, an extended secondary phase with a half-life ($t_{1/2\beta}$) that increases with dose from 86.9 hours at a dose of 0.1 mg/kg to a maximum of 444 hours at a dose of 10.0 mg/kg. Terminal elimination half-life ($t_{1/2z}$) estimated non-compartmentally averages 31 hours across all IV dose groups. Clearance (CL, CUF) of α OPGL-1 is found to be non-linear, with animals receiving IV doses of 10 mg/kg having an average clearance (0.120 ml/hr/kg) that is 3.3-fold lower than those receiving 0.1 mg/kg (0.401 ml/hr/kg).

[0232] After administering subcutaneously, absorption is slow, with average peak concentrations (C_{max}) of 11,600 ng/ml at 132 hr. There is high variability in the range of exposure after SC administration, resulting in an average clearance

of 0.387 ± 0.281 ml/hr/kg and mean residence time of 202 ± 80.1 hours. Average bioavailability is 89%.

[0233] The preceding data are summarized in Table 5.

Table 5: Mean (\pm SD) Non-Compartmental Pharmacokinetic Parameters^a in Cynomolgus Monkeys After Administering a Single Dose of α OPGL-1 IV and SC

| Non-Compartmental Parameter Estimates | | | | | | |
|---------------------------------------|----------|------------------|-------|-----------|-----------|-----------------|
| Parameter | Unit | 1.0 mg/kg | | 0.1 mg/kg | 1.0 mg/kg | 10 mg/kg |
| | | SC (n=6) | | IV (n=2) | IV (n=2) | IV (n=2) |
| | | Mean | SD | Mean | Mean | Mean |
| T _{max} | hr | 132 | 60.2 | 0 | 0 | 0 |
| C _{max} | ng/ml | 11600 | 3410 | 4330 | 38200 | 326000 |
| t _{1/2,z} | hr | 34.9 | 11.1 | 30.7 | 31.4 | ND ^b |
| AUC _(0-∞) | μg*hr/ml | 3520 | 1750 | 253 | 3950 | 99900 |
| CL, CL/F | ml/hr/kg | 0.387 | 0.281 | 0.401 | 0256 | 0.120 |
| MRT | hr | 202 | 80.1 | 84.8 | 124 | 519 |
| V _{ss} | ml/kg | N/A ^c | N/A | 33.7 | 31.7 | 55.9 |

^aValues are reported to 3 significant figures
^bNot Determined, PK samples end during plateau (p) phase hence terminal phase is not observed
^cNot Applicable

[0234] α OPGL-1 causes a rapid decrease in serum N-Tx levels within 24 hours post dose (Figure 14). The average time of maximum effect is observed to occur between 12 hours and 7 days post dose as IV doses increase from 0.1 to 10 mg/kg, and between 12 hours and 11 days in animals receiving a SC dose of 1.0 mg/kg. Maximum effect increases with dose from approximately 80 to 91% over the dose range of 0.1 to 1 mg/kg. However, at higher doses no further suppression is observed with maximum inhibition of 91%. Mean levels of serum N-Tx return to baseline by day 28 after administering 0.1 mg/kg IV and by day 70 after administering 1 mg/kg SC. Urine N-Tx shows similar trends to those of serum N-Tx, except that all groups return to baseline values by study day 105 (Figure 15).

[0235] Suppression of serum Ca increases with dose to a mean nadir of 31.6% below the baseline average seven days after IV administration of 10.0 mg/kg. All other dose groups have mean decreases in serum Ca of less than 26.4% from their baseline averages. By day 17 all serum Ca levels in treated animals return to within 10% of their baseline averages (Figure 20).

[0236] As bone resorption and formation are intimately linked, changes in bone formation markers (ALP) are also observed with a much slower decline in ALP levels and a more prolonged suppression than the formation marker, N-Tx (Figure 21). Observing bone resorption markers decrease prior to bone formation markers (ALP) following dosing with α OPGL-1 confirms that the α OPGL-1 is a bone anti-resorptive agent.

[0237] The majority of animals (9 of 12) develop antibodies to α OPGL-1. The incidence of antibodies to α OPGL-1 is not dose or route dependent. It is not possible to assess the effect of antibodies to α OPGL-1 on α OPGL-1 pharmacokinetics above 0.1 mg/kg when no dose group has both antibody negative and positive animals. At 0.1 mg/kg IV, the majority of α OPGL-1 is cleared prior to antibody development and therefore, effects on α OPGL-1 disposition are not observed (Figure 16).

SEQUENCE LISTING

[0238]

<110> Boyle, William J
 Martin, Francis H
 Corvalan, Jose R
 Davis, C. Geoffrey

<120> Antibodies to OPGL

<130> 06843.0049-00000

EP 1 409 016 B9

<150> 60/301,172

<151> 2001-06-26

<160> 20

5

<170> PatentIn version 3.1

<210> 1

<211> 1426

10

<212> DNA

<213> Mus musculus

<400> 1

15

```

aagcttgacc accatggagt ttgggctgag ctggcttttt. cttgtggcta ttttaaagg 60
tgtccagtgt gaggtgcagc tgttggagtc tgggggagge ttggtacagc ctgggggggc 120
cctgagactc tctgtgcag cctctggatt cacctttage agctatgcca tgagctgggt 180
20 ccgccaggct ccaggaagg ggctggagtg ggtctcaggt attactggga gtggtggtag 240
tacatactac gcagactccg tgaagggccg gttcaccatc tccagagaca attccaagaa 300
cacgctgtat ctgcaaatga acagcctgag agccgaggac acggccgtat attactgtgc 360

```

25

```

gaaagatcca gggactacgg tgattatgag ttggttcgac ccctggggcc agggaaccct 420
ggtcaccgtc tctcagcct ccaccaaggg cccatcggtc tccccctgg cgcctgtctc 480
30 caggagcacc tccgagagca cagcggccct gggctgcctg gtcaaggact acttccccga 540
accggtgacg gtgtcgtgga actcaggcgc tctgaccagc ggcgtgcaca ccttcccagc 600
tgtcctacag tctcaggac tctactccct cagcagcgtg gtgaccgtgc cctccagcaa 660
35 cttcggcaec cagacctaca cctgcaacgt agatcacaag cccagcaaca ccaaggtgga 720
caagacagtt gagcgcaaat gttgtgtcga gtgccaccog tgcccagcac cacctgtggc 780
aggaccgtca gtcttctct tcccccaaa acccaaggac accctcatga tctcccggac 840
40 ccctgaggtc acgtgcgtgg tgggtggacgt gagccacgaa gaccccagg tccagttcaa 900
ctggtaoctg gacggcgtgg aggtgcataa tgccaagaca aagccacggg aggagcagtt 960
caacagcacg ttccgtgtgg tcaçcgtcct caccgttgtg caccaggact ggctgaacgg 1020
45 caaggagtac aagtgcagg tctccaacaa aggcctccca gccccatcg agaaaacat 1080
ctccaaaacc aaagggcagc cccgagaacc acaggtgtac accctgcccc catcccggga 1140
ggagatgacc aagaaccagg tcagcctgac ctgcctggtc aaaggcttct accccagcga 1200
50 catcgcctg gagtgggaga gcaatgggca gccggagaac aactacaaga ccacacctcc 1260
catgctggac tccgacggt ccttcttct ctacagcaag ctcaccgtgg acaagagcag 1320
gtggcagcag gggaaacgtct tctcatgctc cgtgatgcat gaggctctgc acaaccaacta 1380
cacgcagaag agcctctccc tgtctccggg taaatgataa gtcgac 1426

```

55

<210> 2

<211> 467

<212> PRT

EP 1 409 016 B9

<213> Mus musculus

<400> 2

5 Met Glu Phe Gly Leu Ser Trp Leu Phe Leu Val Ala Ile Leu Lys Gly
1 . 5 10 15

10 Val Gln Cys Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln
20 25 30

15 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
35 40 45

15

20

25

30

35

40

45

50

55

EP 1 409 016 B9

Ser Ser Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 5
 Glu Trp Val Ser Gly Ile Thr Gly Ser Gly Gly Ser Thr Tyr Tyr Ala
 65 70 75 80
 10
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn
 85 90 95
 15
 Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
 100 105 110
 Tyr Tyr Cys Ala Lys Asp Pro Gly Thr Thr Val Ile Met Ser Trp Phe
 115 120 125
 20
 Asp Pro Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr
 130 135 140
 Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser
 145 150 155 160
 25
 Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu
 165 170 175
 30
 Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His
 180 185 190
 Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser
 195 200 205
 35
 Val Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys
 210 215 220
 40
 Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Glu
 225 230 235 240
 Arg Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro Pro Val Ala
 245 250 255
 45
 Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met
 260 265 270
 50
 Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His

55

EP 1 409 016 B9

tctagaccac catggaacc ccagcgcagc ttctcttctt cctgctactc tggctcccag 60
 ataccaccgg agaaattgtg ttgacgcagt ctccaggcac cctgtctttg tctccagggg 120
 5 aaagagccac cctctcctgt agggccagtc agagtgttcg cggcaggtae ttagcctggt 180
 accagcagaa acctggccag gctcccagge tctcatcta tggcgcaccc agcagggcca 240
 ctggcatccc agacagggtc agtggcagtg ggtctgggac agacttcaet ctcaccatca 300
 10 gcagactgga gctgaagat tttgcagtgt tttactgtca gcagtatggt agttcacctc 360
 ggacgttcgg ccaagggacc aaggtggaaa tcaaacgaac tgtggctgca ccatctgtct 420
 tcatcttccc gccatctgat gagcagttga aatctggaac tgcctctggt gtgtgcctgc 480
 15 tgaataactt ctatcccaga gaggccaaag tacagtggaa ggtggataac gccctccaat 540
 cgggtaactc ccaggagagt gtcacagagc aggacagcaa ggacagcacc tacagcctca 600
 gcagcacctt gacgtgagc aaagcagact acgagaaaca caaagtctac gcctgcgaag 660
 20 tcacccatca gggcctgagc tegcccgta caaagagctt caacagggga gagtgttgat 720
 aagtcgac 728

<210> 4

<211> 235

25 <212> PRT

<213> Mus musculus

<400> 4

30 Met Glu Thr Pro Ala Gln Leu Leu Phe Leu Leu Leu Leu Trp Leu Pro
 1 5 10 15
 Asp Thr Thr Gly Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser
 35 20 25 30
 Leu Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser
 40 35 40 45
 Val Arg Gly Arg Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala
 50 55 60
 Pro Arg Leu Leu Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro

45

50

55

EP 1 409 016 B9

| | | | | | | | | | | | | | | | | |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 65 | | | | | 70 | | | | | | 75 | | | | 80 |
| 5 | Asp | Arg | Phe | Ser | Gly | Ser | Gly | Ser | Gly | Thr | Asp | Phe | Thr | Leu | Thr | Ile |
| | | | | | 85 | | | | | 90 | | | | | 95 | |
| | Ser | Arg | Leu | Glu | Pro | Glu | Asp | Phe | Ala | Val | Phe | Tyr | Cys | Gln | Gln | Tyr |
| 10 | | | | 100 | | | | | 105 | | | | | 110 | | |
| | Gly | Ser | Ser | Pro | Arg | Thr | Phe | Gly | Gln | Gly | Thr | Lys | Val | Glu | Ile | Lys |
| | | | 115 | | | | | 120 | | | | | 125 | | | |
| 15 | Arg | Thr | Val | Ala | Ala | Pro | Ser | Val | Phe | Ile | Phe | Pro | Pro | Ser | Asp | Glu |
| | | 130 | | | | | 135 | | | | | 140 | | | | |
| | Gln | Leu | Lys | Ser | Gly | Thr | Ala | Ser | Val | Val | Cys | Leu | Leu | Asn | Asn | Phe |
| 20 | 145 | | | | | 150 | | | | | 155 | | | | | 160 |
| | Tyr | Pro | Arg | Glu | Ala | Lys | Val | Gln | Trp | Lys | Val | Asp | Asn | Ala | Leu | Gln |
| | | | | | 165 | | | | | 170 | | | | | 175 | |
| 25 | Ser | Gly | Asn | Ser | Gln | Glu | Ser | Val | Thr | Glu | Gln | Asp | Ser | Lys | Asp | Ser |
| | | | | 180 | | | | | 185 | | | | | 190 | | |
| | Thr | Tyr | Ser | Leu | Ser | Ser | Thr | Leu | Thr | Leu | Ser | Lys | Ala | Asp | Tyr | Glu |
| | | | 195 | | | | | 200 | | | | | 205 | | | |
| 30 | Lys | His | Lys | Val | Tyr | Ala | Cys | Glu | Val | Thr | His | Gln | Gly | Leu | Ser | Ser |
| | | 210 | | | | | 215 | | | | | 220 | | | | |
| 35 | Pro | Val | Thr | Lys | Ser | Phe | Asn | Arg | Gly | Glu | Cys | | | | | |
| | 225 | | | | | 230 | | | | | 235 | | | | | |

40 <210> 5
 <211> 25
 <212> DNA
 <213> Artificial Sequence

45 <220>
 <223> 5' kappa RACE primer

<400> 5
 gatgaccag tctccagcca ccctg 25

50 <210> 6
 <211> 23
 <212> DNA
 <213> Artificial Sequence

55 <220>
 <223> 3' kappa RACE primer

<400> 6
 aagggtcaga ggccaaagga tgg 23

EP 1 409 016 B9

<210> 7
 <211> 30
 <212> DNA
 <213> Artificial Sequence
 5
 <220>
 <223> 5' anti-OPGL-1 kappa primer
 <400> 7
 10 caactctaga ccaccatgga aaccccagcg 30
 <210> 8
 <211> 37
 <212> DNA
 15 <213> Artificial Sequence
 <220>
 <223> 3' anti-OPGL-1 kappa primer
 20 <400> 8
 ttgacgtcg acttatcaac actctcccct gttgaag 37
 <210> 9
 <211> 23
 25 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> 5' IgG2 RACE primer
 30 <400> 9
 ggcacggtca ccacgctgct gag 23
 <210> 10
 <211> 24
 35 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> 3' IgG2 RACE primer
 40 <400> 10
 cctccaccaa gggcccatcg gtct 24
 <210> 11
 <211> 51
 <212> DNA
 <213> Artificial Sequence
 50 <220>
 <223> 5' anti-OPGL-1 IgG2 Primer
 <400> 11
 55 cagaagcttg accaccatgg agtttgggct gagctggctt tttctgtgg c 51
 <210> 12
 <211> 37
 <212> DNA

EP 1 409 016 B9

<213> Artificial Sequence

<220>

<223> 3' anti-OPGL-1 IgG2 Primer

5

<400> 12

gcatgtcgac ttatcattta cccggagaca gggagag 37

<210> 13

10

<211> 122

<212> PRT

<213> Mus musculus

<400> 13

15

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

20

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

25

Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Gly Ile Thr Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
50 55 60

30

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

35

Ala Lys Asp Pro Gly Thr Thr Val Ile Met Ser Trp Phe Asp Pro Trp
100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

40

<210> 14

<211> 108

<212> PRT

<213> Mus musculus

45

<400> 14

50

55

EP 1 409 016 B9

Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

5 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Arg Gly Arg
 20 25 30

10 Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu
 35 40 45

Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser
 50 55 60

15 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu
 65 70 75 80

20 Pro Glu Asp Phe Ala Val Phe Tyr Cys Gln Gln Tyr Gly Ser Ser Pro
 85 90 95

25 Arg Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 15
 <211> 24
 <212> DNA
 <213> Artificial Sequence

30 <220>
 <223> random primer
 <220>
 <221> misc-feature
 <222> (18)..(23)
 <223> N is A, C, G, or T

35 <400> 15
 ggccggatag gcctcacnnn nnnt 24

40 <210> 16
 <211> 5
 <212> PRT
 <213> Artificial Sequence

45 <220>
 <223> translation of portion of 5' anti-OPGL-1 kappa primer
 <400> 16

50 Met Glu Thr Pro Ala
 1 5

55 <210> 17
 <211> 6
 <212> PRT
 <213> Artificial Sequence

<220>

EP 1 409 016 B9

<223> translation of portion of 3' anti-OPGL-1 kappa primer

<400> 17

5
Cys Glu Gly Arg Asn Phe
1 5

<210> 18

<211> 12

10 <212> PRT

<213> Artificial Sequence

<220>

<223> translation of portion of 5' anti-OPGL-1 IgG2 Primer

15

<400> 18

20 Met Glu Phe Gly Leu Ser Trp Leu Phe Leu Val Ala
1 5 10

<210> 19

<211> 7

<212> PRT

25 <213> Artificial Sequence

<220>

<223> translation of portion of 3' anti-OPGL-1 IgG2 Primer

30 <400> 19

35 Lys Gly Pro Ser Leu Ser Leu
1 5

<210> 20

<211> 10

<212> PRT

<213> Artificial Sequence

40

<220>

<223> LHRH antagonist peptide

<220>

<221> MISC_FEATURE

45 <222> (1)..(1)

<223> Xaa is Ac-D-Nal, where Nal is 3-(2-naphthyl)alaninyl

<220>

<221> MISC_FEATURE

50 <222> (2)..(2)

<223> Xaa is (4'-chlorophenyl)alaninyl

<220>

<221> MISC_FEATURE

55 <222> (3) .. (3)

<223> Xaa is D-Pal, where Pal is 3-(3'-pyridyl)alaninyl

<220>

<221> MISC_FEATURE
 <222> (5) .. (5')
 <223> Xaa is N-methyl tyrosine

5 <220>
 <221> MISC_FEATORE
 <222> (6) .. (6)
 <223> Xaa is D-asparagine

10 <220>
 <221> MISC_FEATURE
 <222> (8)..(B)
 <223> Xaa is N-epsilon-2-propyl-lysinyI

15 <220>
 <221> MISC_FEATURE
 <222> (10)..(10)
 <223> Xaa is D-alanine-NH2

20 <400> 20

Xaa Xaa Xaa Ser Xaa Xaa Leu Xaa Pro Xaa
 1 5 10

25

Claims

- 30 1. An antibody, comprising a heavy chain and a light chain, wherein:
- the heavy chain comprises:
 an amino acid sequence as set forth in SEQ ID NO: 13; and
 the light chain comprises:
 an amino acid sequence as set forth in SEQ ID NO: 14; and
- 35 wherein the antibody binds to an osteoprotegerin ligand (OPGL) and inhibits binding of OPGL to an osteoclast differentiation and activation receptor (ODAR).
- 40 2. The antibody of claim 1, wherein the heavy chain and the light chain are connected by a flexible linker to form a single-chain antibody.
- 45 3. The antibody of claim 2, which is a single-chain Fv antibody.
4. The antibody of claim 1, which is a Fab antibody.
5. The antibody of claim 1, which is a Fab' antibody.
6. The antibody of claim 1, which is a (Fab')₂ antibody.
- 50 7. The antibody of claim 1, which is fully human.
8. An antibody of claim 1 comprising a heavy chain and a light chain, wherein the heavy chain comprises an amino acid sequence of SEQ ID NO:2 from residue 20 to residue 467, and wherein the light chain comprises an amino acid sequence of SEQ ID NO:4 from residue 21 to residue 235.
- 55 9. An antibody of claim 1 comprising a heavy chain and a light chain, wherein the heavy chain consists of an amino acid sequence of SEQ ID NO:2 from residue 20 to residue 467, and wherein the light chain consists of an amino acid sequence of SEQ ID NO:4 from residue 21 to residue 235.

10. A pharmaceutical composition comprising an antibody of any of claims 1 to 9.

Patentansprüche

- 5
1. Antikörper, der eine schwere Kette und eine leichte Kette umfasst, wobei:
- die schwere Kette umfasst:
eine Aminosäuresequenz, wie in SEQ ID NO: 13 dargestellt; und
10 die leichte Kette umfasst:
eine Aminosäuresequenz, wie in SEQ ID NO: 14 dargestellt; und
- wobei der Antikörper an einen Osteoprotegerin-Liganden (OPGL) bindet und die OPGL-Bindung an einen Rezeptor zur Differenzierung und Aktivierung von Osteoklasten (ODAR) hemmt.
- 15
2. Antikörper nach Anspruch 1, wobei die schwere Kette und die leichte Kette über einen flexiblen Linker verbunden sind, um einen Einzelkettenantikörper zu bilden.
- 20
3. Antikörper nach Anspruch 2, der ein Fv-Einzelkettenantikörper ist.
4. Antikörper nach Anspruch 1, der ein Fab-Antikörper ist.
5. Antikörper nach Anspruch 1, der ein Fab'-Antikörper ist.
- 25
6. Antikörper nach Anspruch 1, der ein (Fab')₂-Antikörper ist.
7. Antikörper nach Anspruch 1, der vollständig human ist.
8. Antikörper nach Anspruch 1, der eine schwere Kette und eine leichte Kette umfasst, wobei die schwere Kette eine Aminosäuresequenz der SEQ ID NO: 2 von Rest 20 bis Rest 467 umfasst und wobei die leichte Kette eine Amino-
30 säuresequenz der SEQ ID NO: 4 von Rest 21 bis Rest 235 umfasst.
9. Antikörper nach Anspruch 1, der eine schwere Kette und eine leichte Kette umfasst, wobei die schwere Kette aus einer Aminosäuresequenz der SEQ ID NO: 2 von Rest 20 bis Rest 467 besteht und wobei die leichte Kette aus
35 einer Aminosäuresequenz der SEQ ID NO: 4 von Rest 21 bis Rest 235 besteht.
10. Pharmazeutische Zusammensetzung, die einen Antikörper nach einem der Ansprüche 1 bis 9 umfasst.

40 Revendications

1. Un anticorps comprenant une chaîne lourde et une chaîne légère, où :
- la chaîne lourde comprend :
45 une séquence d'acides aminés telle que définie dans SEQ ID NO: 13, et
la chaîne légère comprend :
une séquence d'acides aminés telle que définie dans SEQ ID NO: 14, et
- où l'anticorps se lie à un ligand de l'ostéoprotégérine (OPGL) et inhibe la liaison de l'OPGL à un récepteur d'activation
50 et de différenciation des ostéoclasts (ODAR).
2. L'anticorps selon la revendication 1, où la chaîne lourde et la chaîne légère sont reliées par un lieur souple de façon à former un anticorps à chaîne unique.
- 55
3. L'anticorps selon la revendication 2, qui est un anticorps Fv à chaîne unique.
4. L'anticorps selon la revendication 1, qui est un anticorps Fab.

EP 1 409 016 B9

5. L'anticorps selon la revendication 1, qui est un anticorps Fab'.
6. L'anticorps selon la revendication 1, qui est un anticorps (Fab')₂.
- 5 7. L'anticorps selon la revendication 1, qui est entièrement humain.
8. Un anticorps selon la revendication 1 comprenant une chaîne lourde et une chaîne légère, où la chaîne lourde comprend une séquence d'acides aminés de SEQ ID NO: 2 allant du résidu 20 au résidu 467 et où la chaîne légère comprend une séquence d'acides aminés de SEQ ID NO: 4 allant du résidu 21 au résidu 235.
- 10 9. Un anticorps selon la revendication 1 comprenant une chaîne lourde et une chaîne légère, où la chaîne lourde se compose d'une séquence d'acides aminés de SEQ ID NO: 2 allant du résidu 20 au résidu 467 et où la chaîne légère se compose d'une séquence d'acides aminés de SEQ ID NO: 4 allant du résidu 21 au résidu 235.
- 15 10. Une composition pharmaceutique contenant un anticorps selon l'une quelconque des revendications 1 à 9.

20

25

30

35

40

45

50

55

FIGURE 1

 α OPGL-1 Heavy Chain cDNA

DNA sequence of heavy chain expression plasmid beginning at HindIII site, through Sall site. Start codon begins at nt 14, stop codon begins at nt 1415.

```

1  AAGCTTGACC ACCATGGAGT TTGGGCTGAG CTGGCTTTTT CTTGTGGCTA TTTTAAAAGG
61  TGTCAGTGT GAGGTGCAGC TGTTGGAGTC TGGGGGAGGC TTGTACAGC CTGGGGGGTC
121 CCTGAGACTC TCCTGTGCAG CCTCTGGATT CACCTTTAGC AGCTATGCCA TGAGCTGGGT
181 CCGCCAGGCT CCAGGGAAGG GGCTGGAGTG GGTCTCAGGT ATTACTGGGA GTGGTGGTAG
241 TACATACTAC GCAGACTCCG TGAAGGGCCG GTTCACCATC TCCAGAGACA ATTCCAAGAA
301 CACGCTGTAT CTGCAAATGA ACAGCCTGAG AGCCGAGGAC ACGGCCGTAT ATTACTGTGC
361 GAAAGATCCA GGGACTACGG TGATTATGAG TTGGTTCGAC CCCTGGGGCC AGGGAACCTT
421 GGTCAACGTC TCCTCAGCCT CCACCAAGGG CCCATCGGTC TTCCCCCTGG CGCCCTGCTC
481 CAGGAGCACC TCCGAGAGCA CAGCGGCCCT GGGCTGCCTG GTCAAGGACT ACTTCCCCGA
541 ACCGGTGACG GTGTCTGGA ACTCAGGCGC TCTGACCAGC GGCCTGCACA CCTTCCCAGC
601 TGTCCTACAG TCCTCAGGAC TCTACTCCCT CAGCAGCGTG GTGACCGTGC CCTCCAGCAA
661 CTTCGGCACC CAGACCTACA CCTGCAACGT AGATCACAAG CCCAGCAACA CCAAGGTGGA
721 CAAGACAGTT GAGCGCAAAT GTTGTGTCGA GTGCCACCG TGCCCAGCAC CACCTGTGGC
781 AGGACCGTCA GTCTTCTCT TCCCCCAA ACCCAAGGAC ACCCTCATGA TCTCCCGGAC
841 CCCTGAGGTC ACGTGCCTGG TGGTGGACGT GAGCCACGAA GACCCCGAGG TCCAGTTCAA
901 CTGGTACGTG GACGGCGTGG AGGTGCATAA TGCCAAGACA AAGCCACGGG AGGAGCAGTT
961 CAACAGCACG TTCCGTGTGG TCAGCGTCT CACCGTTGTG CACCAGGACT GGCTGAACGG
1021 CAAGGAGTAC AAGTGCAAGG TCTCCAACAA AGGCCTCCCA GCCCCATCG AGAAAACCAT
1081 CTCAAAACC AAAGGGCAGC CCCGAGAACC ACAGGTGTAC ACCCTGCCCC CATCCCGGGA
1141 GGAGATGACC AAGAACCAGG TCAGCCTGAC CTGCCTGGTC AAAGGTTCT ACCCCAGCGA
1201 CATCGCCGTG GAGTGGGAGA GCAATGGGCA GCCGGAGAAC AACTACAAGA CCACACCTCC
1261 CATGCTGGAC TCCGACGGCT CTTCTTCTCT CTACAGCAAG CTCACCGTGG ACAAGAGCAG
1321 GTGGCAGCAG GGAACGTCT TCTCATGCTC CGTGATGCAT GAGGCTCTGC ACAACCACTA
1381 CACGCAGAAG AGCCTCTCCC TGCTCCGGG TAAATGATAA GTCGAC (SEQ ID NO: 1)

```

FIGURE 2

 α OPGL-1 Heavy Chain Amino Acid Sequence

The IgG2 signal peptide is underlined, the variable region is in capital letters and is not underlined, and the constant region is in lower case.

```

1 MEFGLSWLFL VAILKGVOCE VQLLESGGGL VQPGGSLRLS CAASGFTFSS YAMSWVRQAP
61 GKGLEWVSGI TGGGGSTYYA DSVKGRFTIS RDNSKNTLYL QMNSLRAEDT AVYYCAKDPG
121 TTVIMSWFDP WGQGTLTVS Sastkgpsvf plapcsrsts estaalgclv kdyfpepvtv
181 swnsgaltsg vhtfpavlqs sglyslssvv tvpssnfgtq tytcnvdhkp sntkvdkve
241 rkccvecppc pappvagpsv flfppkpkdt lmisrtpevt cvvvdvshed pevqfnwyvd
301 gvevhnaktk preeqfnstf rvsvltvvh qdwlngkeyk ckvsnkglpa piektisktk
361 gqprepvyt lppsreemtk nqvsltclvk gfypsdiave wesngqpenn ykttppmlds
421 dgsfflyskl tvdksrwqqg nvfscsvmhe alhnhytqks lsispqk (SEQ ID NO: 2)

```

FIGURE 3 **α OPGL-1 kappa Light Chain cDNA**

DNA sequence of kappa chain expression plasmid sequence from XbaI site through Sall site. Start codon begins at nt 12; stop codon begins at nt 717

```

1 TCTAGACCAC CATGGAAACC CCAGCGCAGC TTCTCTTCCT CCTGCTACTC TGGCTCCCAG
61 ATACCACCGG AGAAATTGTG TTGACGCAGT CTCCAGGCAC CCTGTCTTTG TCTCCAGGGG
121 AAAGAGCCAC CCTCTCCTGT AGGGCCAGTC AGAGTGTTTCG CGGCAGGTAC TTAGCCTGGT
181 ACCAGCAGAA ACCTGGCCAG GCTCCCAGGC TCCTCATCTA TGGTGCATCC AGCAGGGCCA
241 CTGGCATCCC AGACAGGTTT AGTGGCAGTG GGTCTGGGAC AGACTTCACT CTCACCATCA
301 GCAGACTGGA GCCTGAAGAT TTTGCACTGT TTTACTGTCA GCAGTATGGT AGTTCACCTC
361 GGACGTTTCGG CCAAGGGACC AAGGTGGAAA TCAAACGAAC TGTGGCTGCA CCATCTGTCT
421 TCATCTTCCC GCCATCTGAT GAGCAGTTGA AATCTGGAAC TGCCTCTGTT GTGTGCCTGC
481 TGAATAACTT CTATCCCAGA GAGGCCAAAG TACAGTGGAA GGTGGATAAC GCCCTCCAAT
541 CGGGTAAGTC CCAGGAGAGT GTCACAGAGC AGGACAGCAA GGACAGCACC TACAGCCTCA
601 GCAGCACCCCT GACGCTGAGC AAAGCAGACT ACGAGAAACA CAAAGTCTAC GCCTGCGAAG
661 TCACCCATCA GGCCTGAGC TCGCCCGTCA CAAAGAGCTT CAACAGGGGA GAGTGTGAT
721 AAGTCGAC (SEQ ID NO: 3)

```

FIGURE 4 **α OPGL-1 kappa Light Chain Amino Acid Sequence**

The kappa signal peptide is underlined, the variable region is in capital letters and not underlined, and the constant region is in lower case.

```

1 METPAQLLFL LLLWLPDTTG EIVLTQSPGT LSLSPGERAT LSCRASQSVR
51 GRYLAWYQK PGQAPRLLIY GASSRATGIP DRFSGSGSGT DFTLTISRLE
101 PEDFAVFYCQ QYGSSPRTFG QGTKVEIKrt vaapsvfifp psdeqlksgt
151 asvvcilnnf ypreakvqwk vdnalqsgns qesvteqdk dstyslsstl
201 tiskadyekh kvyacevthq glsspvtksf nrgec (SEQ ID NO: 4)

```

FIGURE 5

Circular Plasmid Map of α OPGL-1-kappa/pDSR α 19




FIGURE 6

Circular Plasmid Map of α OPGL-1-IgG2/pDSRa19






FIGURE 7

 α OPGL-1 Binds to Soluble OPG Ligand Coated on the EIA Plate in a Dose-dependent Manner

96-well EIA plates are coated with recombinant soluble OPG. Varying concentrations of α OPGL-1 are added to the wells and incubated for about 2hrs at room temperature. Bound antibody is detected with goat anti-Human IgG (Fab')-Horse Radish Peroxidase. The absorbance is read at 450 nm and 650 nm.


FIGURE 8

 α OPGL-1 Binds Specifically to Membrane-bound OPGL

α OPGL-1 binds to OPGL expressed on a cell surface of transfected CHO REN 218-9 cells in a dose-dependent manner. This binding is competed by exogenously added human OPGL but not by TNF- α , TNF- β , TRAIL or CD40 ligand. α OPGL-1 (100 ng/ml) is pre-incubated with varying concentrations of soluble OPGL or other ligands and is then incubated with CHO REN 218-9 cells expressing OPGL on the surface. Cells are then incubated for 30 minutes at 2-8°C with FITC-labeled F(ab')₂ Goat anti-Human IgG, Fc γ Fragment Specific. After centrifugation and washing cell surface fluorescence is measured using flow cytometry.

FIGURE 9


α OPGL-1 Binds to Soluble OPGL



α OPGL-1 binding to OPGL on an EIA plate is reduced competitively by exogenously added soluble OPGL.

Figure 10

α OPGL-1 does not bind to TNF family members TNF α , TNF β , TRAIL, or CD40 ligand



α OPGL-1 binding to OPGL on an EIA plate is not reduced by exogenously added TNF- α , TNF- β , TRAIL or CD40 Ligand.

FIGURE 11

Dose-dependent Inhibition by α OPGL-1 of OPG Ligand-induced TRAP Activity in RAW 264.7 Cells




FIGURE 12

Dose-dependent Inhibition by α OPGL-1 of Europium-labeled OPG Ligand Binding to ODAR-FLAG/anti-FLAG-APC




FIGURE 13

Mean (\pm SD) Serum-Concentration Time Profiles Following Single Dose Administration of α OPGL-1 to Cynomolgus Monkeys at Doses of 0.1, 1 and 10.0 mg/kg IV (n=2/dose) and 1.0 mg/kg SC (n=6/dose)




FIGURE 14

Mean (\pm SD) Percent Change in Serum N-Tx Concentration After Administering a Single Dose of α OPGL-1 IV (n=2/dose) or SC (n=6) to Cynomolgus Monkeys at Doses of 0.1, 1 and 10.0 mg/kg.




FIGURE 15

Mean (\pm SD) Percent Change in Urine N-Tx Concentration Following Single Dose IV (n=2/dose) or SC (n=6) Administration of α OPGL-1 to Cynomolgus Monkeys at Doses of 0.1, 1 and 10.0 mg/kg




FIGURE 16

**Antibody Positive (open black symbols) and Negative (closed red symbols)
Animal Serum-Concentration Time Profiles After Administering a Single Dose of
 α OPGL-1 IV to Cynomolgus Monkeys at a Dose of 0.1 mg/kg**




FIGURE 17

α OPGL-1 Heavy Chain Variable Region Amino Acid Sequence

1 EVQLLES GGG LVQP GGS LRL SCAAS GFTFS SYAM SWRQA
41 PGK GLEW VSG ITG SGG STYY ADSV KGRFTI SRD NSKNTLY
81 LQM NSLRAED TAVYY CAKDP GTTV IMSWFD PWG QGTLVTV
121 SS (SEQ ID NO: 13)

FIGURE 18

α OPGL-1 Light Chain Variable Region Amino Acid Sequence

1 EIVLTQSPGT LSLSPGERAT LSCRASQSVR GRYLAWYQQK
41 PGQAPRLLIY GASSRATGIP DRFSGSGSGT DFTLTISRLE
81 PEDFAVFCYQ QYGSSPRTFG QGTKVEIK (SEQ ID NO: 14)

FIGURE 19

Cell Culture Process Flow Chart




FIGURE 20

Mean Serum Calcium Percent Change From Baseline After Administering a Single Dose of α OPGL-1 to Cynomolgus Monkeys





FIGURE 21

Mean Serum Alkaline Phosphatase Percent Change From Baseline After Administering a Single Dose of α OPGL-1 to Cynomolgus Monkeys



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 5151510 A, Stec [0040]
- WO 8801649 A [0067]
- US 4946778 A [0067]
- US 5260203 A [0067]
- WO 0183525 A [0076]
- WO 9312227 A [0089]
- US 6180370 B [0090]
- US 5693762 A [0090]
- US 5693761 A [0090]
- US 5585089 A [0090]
- US 5530101 A [0090]
- WO 9824893 A [0091]
- US 4399216 A [0098]
- US 4912040 A [0098]
- US 4740461 A [0098]
- US 4959455 A [0098]
- EP 623674 A [0110]
- US 5492888 A [0110]
- US 5488032 A [0110]
- US 5464937 A [0110]
- US 5319071 A [0110]
- US 5180812 A [0110]
- WO 9501997 A [0110]
- WO 9402627 A [0110]
- WO 9006371 A [0110]
- US 4935343 A [0110]
- EP 364778 A [0110]
- EP 267611 A [0110]
- EP 220063 A [0110]
- WO 9623067 A [0110]
- WO 9937773 A [0110] [0146]
- WO 9946248 A [0110]
- WO 9947545 A [0110]
- WO 9947154 A [0110]
- US 5747444 A [0111]
- US 5359032 A [0111]
- US 5608035 A [0111]
- US 5843905 A [0111]
- US 5866576 A [0111]
- US 5869660 A [0111] [0119]
- US 5869315 A [0111]
- US 5872095 A [0111]
- US 5955480 A [0111] [0119]
- US 5965564 A [0111]
- WO 9821957 A [0111]
- WO 9609323 A [0111]
- WO 9117184 A [0111] [0112]
- WO 9640907 A [0111]
- WO 9832733 A [0111]
- WO 9842325 A [0111]
- WO 9844940 A [0111]
- WO 9847892 A [0111]
- WO 9856377 A [0111] [0119]
- WO 9903837 A [0111]
- WO 9906426 A [0111]
- WO 9906042 A [0111] [0119]
- WO 9117249 A [0111]
- WO 9882733 A [0111]
- WO 9817661 A [0111]
- WO 9708174 A [0111]
- WO 9534326 A [0111] [0119]
- WO 9936426 A [0111]
- WO 9936415 A [0111]
- EP 534978 A [0111]
- EP 894795 A [0111]
- FR 2762514 [0111]
- US 5075222 A [0112] [0113]
- WO 9108285 A [0112]
- AU 9173636 [0112]
- WO 9216221 A [0112] [0119]
- WO 9321946 A [0112] [0119]
- WO 9406457 A [0112]
- WO 9421275 A [0112]
- FR 2706772 [0112]
- WO 9421235 A [0112]
- DE 4219626 [0112]
- WO 9420517 A [0112]
- WO 9622793 A [0112]
- WO 9728828 A [0112]
- WO 9936541 A [0112]
- US 6090382 A [0118]
- US 6258562 B [0118]
- WO 9824463 A [0118]
- EP 308378 A [0119]
- EP 422339 A [0119] [0120]
- EP 393438 A [0119] [0120] [0122]
- EP 398327 A [0119]
- EP 412486 A [0119]
- EP 418014 A [0119]
- EP 417563 A [0119]
- EP 433900 A [0119]
- EP 464533 A [0119]
- EP 512528 A [0119]
- EP 526905 A [0119]
- EP 568928 A [0119]
- EP 607776 A [0119]
- EP 663210 A [0119]
- EP 542795 A [0119]

EP 1 409 016 B9

- EP 818439 A [0119]
- EP 664128 A [0119]
- EP 741707 A [0119]
- EP 874819 A [0119]
- EP 882714 A [0119]
- EP 880970 A [0119]
- EP 648783 A [0119]
- EP 731791 A [0119]
- EP 895988 A [0119]
- EP 550376 A [0119]
- EP 853083 A [0119]
- EP 943616 A [0119]
- EP 939121 A [0119]
- EP 614984 A [0119]
- US 5136021 A [0119]
- US 5929117 A [0119]
- US 5948638 A [0119]
- US 5807862 A [0119]
- US 5695953 A [0119]
- US 5834435 A [0119]
- US 5817822 A [0119]
- US 5830742 A [0119]
- US 5851556 A [0119]
- US 5853977 A [0119]
- US 5359037 A [0119]
- US 5512544 A [0119]
- US 5811261 A [0119]
- US 5633145 A [0119]
- US 5863926 A [0119]
- US 5866616 A [0119]
- US 5641673 A [0119]
- US 5869677 A [0119]
- US 5869511 A [0119]
- US 5872146 A [0119]
- US 5854003 A [0119]
- US 5856161 A [0119]
- US 5877222 A [0119]
- US 5877200 A [0119]
- US 5877151 A [0119]
- US 5886010 A [0119]
- US 5859207 A [0119]
- US 5891883 A [0119]
- US 5877180 A [0119]
- US 5955476 A [0119]
- US 5955435 A [0119]
- US 5994351 A [0119]
- US 5990119 A [0119]
- US 5952320 A [0119]
- US 5962481 A [0119]
- WO 9013575 A [0119]
- WO 9103553 A [0119]
- WO 9201002 A [0119]
- WO 9213095 A [0119]
- WO 9307863 A [0119]
- WO 9319777 A [0119]
- WO 9628546 A [0119]
- WO 9827298 A [0119]
- WO 9830541 A [0119]
- WO 9638150 A [0119]
- WO 9718207 A [0119]
- WO 9715561 A [0119]
- WO 9712902 A [0119]
- WO 9625861 A [0119]
- WO 9612735 A [0119]
- WO 9611209 A [0119]
- WO 9839326 A [0119]
- WO 9839316 A [0119]
- WO 9838859 A [0119]
- WO 9839315 A [0119]
- WO 9842659 A [0119]
- WO 9839329 A [0119]
- WO 9843959 A [0119]
- WO 9845268 A [0119]
- WO 9847863 A [0119]
- WO 9633172 A [0119]
- WO 9620926 A [0119]
- WO 9737974 A [0119]
- WO 9737973 A [0119]
- WO 9747599 A [0119]
- WO 9635711 A [0119]
- WO 9851665 A [0119]
- WO 9843946 A [0119]
- WO 9504045 A [0119]
- WO 9712244 A [0119]
- WO 9900364 A [0119]
- WO 9900363 A [0119]
- WO 9857936 A [0119]
- WO 9901449 A [0119]
- WO 9901139 A [0119]
- WO 9856788 A [0119]
- WO 9856756 A [0119]
- WO 9853842 A [0119]
- WO 9852948 A [0119]
- WO 9852937 A [0119]
- WO 9902510 A [0119]
- WO 9743250 A [0119]
- WO 9906410 A [0119]
- WO 9909022 A [0119]
- WO 9908688 A [0119]
- WO 9907679 A [0119]
- WO 9909965 A [0119]
- WO 9907704 A [0119]
- WO 9906041 A [0119]
- WO 9937818 A [0119]
- WO 9937625 A [0119]
- WO 9711668 A [0119]
- WO 9950238 A [0119]
- WO 9947672 A [0119]
- WO 9948491 A [0119]
- JP 10147531 B [0119]
- JP 10231285 B [0119]
- JP 10259140 B [0119]
- JP 10130149 B [0119]
- JP 10316570 B [0119]
- JP 11001481 B [0119]
- JP 3127800 A [0119]

- DE 19731521 [0119]
- GB 2218101 A [0119]
- GB 2326881 A [0119]
- GB 2246569 A [0119]
- WO 9801555 A [0123] [0124] [0125]
- WO 98101555 A [0126]
- US 4760130 A [0141]
- US 5900400 A [0141]
- US 5633227 A [0141]
- US 5912324 A [0146]
- EP 0962531 A [0146]
- EP 712931 A [0146]
- US 5914253 A [0146]
- WO 9724441 A [0146]
- US 6060283 A [0146]
- EP 850952 A [0146]
- EP 864585 A [0146]
- WO 9841232 A [0146]
- US 6054487 A [0146]
- WO 9909063 A [0146]
- WO 9922760 A [0146]
- WO 9937772 A [0146]
- EP 0974600 A [0146]
- WO 0012555 A [0146]
- JP 6111399 A [0146]
- IL 121554 A0 [0146]
- WO 0046240 A [0147]
- WO 0047740 A [0147]
- WO 0185872 A [0147]
- WO 0215273 A [0147]
- WO 9839361 A [0147]
- US 5843901 A [0151]
- US 428082 [0162]
- WO 9925044 A [0162]
- US 94001875 W [0168]
- US 9300829 W [0171]
- US 3773919 A [0171]
- EP 058481 A [0171]
- EP 133988 A [0171]
- EP 036676 A [0171]
- EP 088046 A [0171]
- EP 143949 A [0171]
- WO 9014363 A [0186]
- US 6210924 B [0193]
- WO 60301172 A [0238]

Non-patent literature cited in the description

- **TSUKII et al.** *Biochemical and Biophysical Research Communications*, 1998, vol. 246, 336-341 [0007]
- **SAMBROOK et al.** *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press [0030]
- **LAPLANCHE et al.** *Nucl. Acids Res.*, 1986, vol. 14, 9081 [0040]
- **STEC et al.** *J. Am. Chem. Soc.*, 1984, vol. 106, 6077 [0040]
- **STEIN et al.** *Nucl. Acids Res.*, 1988, vol. 16, 3209 [0040]
- **ZON et al.** *Anti-Cancer Drug Design*, 1991, vol. 6, 539 [0040]
- **ZON et al.** *Oligonucleotides and Analogues: A Practical Approach*. Oxford University Press, 1991, 87-108 [0040]
- **UHLMANN ; PEYMAN.** *Chemical Reviews*, 1990, vol. 90, 543 [0040]
- *Computational Molecular Biology*. Oxford University Press, 1988 [0041]
- *Biocomputing: informatics and Genome Projects*. Academic Press, 1993 [0041]
- *Computer Analysis of Sequence Data*. Humana Press, 1994 [0041]
- **HEINJE, G.** *Sequence Analysis in Molecular Biology*. Academic Press, 1987 [0041]
- *Sequence Analysis Primer*. M. Stockton Press, 1991 [0041]
- **CARILLO et al.** *SIAM J. Applied Math.*, 1988, vol. 48, 1073 [0041]
- **DEVEREUX et al.** *Nucl. Acid. Res.*, 1984, vol. 12, 387 [0042]
- **ALTSCHUL et al.** *J. Mol. Biol.*, 1990, vol. 215, 403-410 [0042]
- **DAYHOFF et al.** *Atlas of Protein Sequence and Structure*, 1978, vol. 5 (3 [0044]
- **HENIKOFF et al.** *Proc. Natl. Acad. Sci USA*, 1992, vol. 89, 10915-10919 [0044]
- **NEEDLEMAN et al.** *J. Mol. Biol.*, 1970, vol. 48, 443-453 [0045]
- *Immunology-A Synthesis*. Sinauer Associates, 1991 [0047]
- **KYTE et al.** *J. Mol. Biol.*, 1982, vol. 157, 105-131 [0053]
- **MOULT J.** *Curr. Op. in Biotech.*, 1996, vol. 7 (4), 422-427 [0060]
- **CHOU et al.** *Biochemistry*, 1974, vol. 13 (2), 222-245 [0060]
- **CHOU et al.** *Biochemistry*, 1974, vol. 113 (2), 211-222 [0060]
- **CHOU et al.** *Adv. Enzymol. Relat. Areas Mol. Biol.*, 1978, vol. 47, 45-148 [0060]
- **CHOU et al.** *Ann. Rev. Biochem.*, vol. 47, 251-276 [0060]
- **CHOU et al.** *Biophys. J.*, 1979, vol. 26, 367-384 [0060]
- **HOLM et al.** *Nucl. Acid. Res.*, 1999, vol. 27 (1), 244-247 [0060]
- **BRENNER et al.** *Curr. Op. Struct. Biol.*, 1997, vol. 7 (3), 369-376 [0060]
- **JONES, D.** *Curr. Opin. Struct. Biol.*, 1997, vol. 7 (3), 377-87 [0061]
- **SIPPL et al.** *Structure*, 1996, vol. 4 (1), 15-19 [0061]

- **BOWIE et al.** *Science*, 1991, vol. 253, 164-170 [0061]
- **GRIBSKOV et al.** *Meth. Enzym.*, 1990, vol. 183, 146-159 [0061]
- **GRIBSKOV et al.** *Proc. Nat. Acad. Sci.*, 1987, vol. 84 (13), 4355-4358 [0061]
- Proteins, Structures and Molecular Principles. W. H. Freeman and Company, 1984 [0063]
- Introduction to Protein Structure. Garland Publishing, 1991 [0063]
- **THORNTON.** *Nature*, 1991, vol. 354, 105 [0063]
- **FAUCHERE.** *J. Adv. Drug Res.*, 1986, vol. 15, 29 [0065]
- **VEBER ; FREIDINGER.** *TINS*, 1985, 392 [0065]
- **EVANS et al.** *J. Med. Chem.*, 1987, vol. 30, 1229 [0065]
- **RIZO ; GIERASCH.** *Ann. Rev. Biochem.*, 1992, vol. 61, 387 [0065]
- Fundamental Immunology. Raven Press, 1989 [0086]
- Kabat Sequences of Proteins of Immunological Interest. National Institutes of Health, 1987 [0087]
- **CHOTHIA ; LESK.** *J. Mol. Biol.*, 1987, vol. 196, 901-917 [0087]
- **CHOTHIA et al.** *Nature*, 1989, vol. 342, 877-883 [0087]
- **SONGSIVILAI ; LACHMANN.** *Clin. Exp. Immunol.*, 1990, vol. 79, 315-321 [0088]
- **KOSTELNY et al.** *J. Immunol.*, 1992, vol. 148, 1547-1553 [0088]
- **MENDEZ et al.** *Nature Genetics*, 1997, vol. 15, 146-156 [0091]
- **SMITH et al.** *Science*, 1990, vol. 248, 1019-1023 [0121]
- Acid-Stable Inhibitors of Granulocyte Neutral Proteases in Human Mucous Secretions: Biochemistry and Possible Biological Function. **SCHIESSLER et al.** Neutral Proteases of Human Polymorphonuclear Leucocytes. Urban and Schwarzenberg, Inc, 1978 [0136]
- **TRAVIS ; SALVESEN.** *Ann. Rev. Biochem.*, 1983, vol. 52, 655-709 [0136]
- **JIN et al.** *Cell*, 1997, vol. 88 (3), 417-26 [0140]
- **DINARELLO et al.** *J. Leukocyte Biol.*, 1998, vol. 63, 658-84 [0142]
- **FAGGIONI et al.** *PNAS*, 2000, vol. 97, 2367-72 [0142]
- **C.A. DINARELLO ; L.L. MOLDAWER.** Proinflammatory and Anti-inflammatory Cytokines in Rheumatoid Arthritis: A Primer for Clinicians. Amgen Inc, 2001 [0147]
- **BULOW ; BRAM.** *Science*, 1997, vol. 278, 138-140 [0147]
- **LACEY et al.** *Am. J. Pathol.*, 2000, vol. 157, 435-448 [0156]
- Remington's Pharmaceutical Sciences. Mack Publishing Company, 1990 [0161]
- **SIDMAN et al.** *Biopolymers*, 1983, vol. 22, 547-556 [0171]
- **LANGER et al.** *J. Biomed. Mater. Res.*, 1981, vol. 15, 167-277 [0171]
- **LANGER.** *Chem. Tech.*, 1982, vol. 12, 98-105 [0171]
- **EPPSTEIN et al.** *Proc. Natl. Acad. Sci. USA*, 1985, vol. 82, 3688-3692 [0171]
- **GOODWIN et al.** *Nucleic Acids Res.*, 1983, vol. 11, 6873-82 [0187]
- **GASSER et al.** *Proc Natl Acad Sci U S A.*, 1982, vol. 79, 6522-6 [0187]
- **NUNBERG.** *Cell*, 1980, vol. 19, 355-64 [0187]
- **SETZER et al.** *J Biol Chem.*, 1982, vol. 257, 5143-7 [0187]
- **MCGROGAN et al.** *J Biol Chem.*, 1985, vol. 260, 2307-14 [0187]
- **TAKEBE et al.** *Mol Cell Biol.*, 1988, vol. 8, 466-72 [0187]
- **SEIKI et al.** *Proc Natl Acad Sci U S A.*, 1983, vol. 80, 3618-22 [0187]
- **OKAYAMA ; BERG.** *Mol Cell Biol.*, 1983, vol. 3, 280-9 [0187]
- **SIMONET et al.** *Cell*, 1997, vol. 89, 309 [0220]
- **LACEY et al.** *Cell*, 1998, vol. 93, 165 [0220]