

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 452 586 A8

(12)

CORRECTED EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

Note: Bibliography reflects the latest situation

(15) Correction information:

Corrected version no 1 (W1 A1)
INID code(s) 54,84

(51) Int Cl.7: **C12N 5/06**, A61K 35/28,
A61P 25/00

(48) Corrigendum issued on:

27.04.2005 Bulletin 2005/17

(86) International application number:

PCT/JP2002/011294

(43) Date of publication:

01.09.2004 Bulletin 2004/36

(87) International publication number:

WO 2003/038075 (08.05.2003 Gazette 2003/19)

(21) Application number: **02775435.7**

(22) Date of filing: **30.10.2002**

(84) Designated Contracting States:

DE FR GB IT SE

(72) Inventors:

- **HONMOU, Osamu**
Sapporo-shi, Hokkaido 064-0801 (JP)
- **HAMADA, Hirofumi**
Sapporo-shi, Hokkaido 064-0959 (JP)

(30) Priority: **30.10.2001 WOPCT/JP01/09510**

03.04.2002 WOPCT/JP02/03344

(74) Representative: **VOSSIUS & PARTNER**

Siebertstrasse 4
81675 München (DE)

(71) Applicant: **Renomedix Institute Inc.**

Sapporo-shi, Hokkaido 060-0031 (JP)

(54) **METHOD OF INDUCING DIFFERENTIATION OF MESODERMAL STEM CELLS, ES CELLS OR IMMORTALIZED CELLS INTO NERVOUS SYSTEM CELLS**

(57) Mesodermal stem cells or ES cells, prepared from the mononuclear cell fraction isolated from bone marrow fluid or umbilical blood, were found to differentiate into neural stem cells, neurons, or glial cells when cultured in a basal culture medium. In addition, the differentiation of the mesodermal stem cells or ES cells into neural cells was promoted through the addition of an ischemic brain extract to the above-mentioned basal culture medium. Furthermore, the neural cells obtained using the above-described method for inducing differentiation were revealed to have neural regeneration po-

tency in a brain infarction model, a dementia model, a spinal cord injury model and a demyelination model.

In addition, according to the present invention, mesodermal stem cells can be differentiated into neural cells by immortalizing the mesodermal stem cells by highly expressing or activating an immortalization gene in the mesodermal stem cells and culturing the cells under an appropriate condition. The methods of the present invention are very useful in the medical field of neural regeneration.

EP 1 452 586 A8