



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 461 331 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

Note: Bibliography reflects the latest situation

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see page(s) 2, 4, 6

(48) Corrigendum issued on:
28.06.2006 Bulletin 2006/26

(45) Date of publication and mention
of the grant of the patent:
08.03.2006 Bulletin 2006/10

(21) Application number: **02793596.4**

(22) Date of filing: **09.12.2002**

(51) Int Cl.:
C07D 319/06 (2006.01)

(86) International application number:
PCT/NL2002/000876

(87) International publication number:
WO 2003/059901 (24.07.2003 Gazette 2003/30)

(54) **PROCESS FOR THE PREPARATION OF 2-(6-SUBSTITUTED-1,3-DIOXANE-4-YL) ACETIC ACID DERIVATIVES**

PROZESS FÜR DIE HERSTELLUNG VON 2-(6-SUBSTITUIERTEN-1,3-DIOXAN-4-YL)
ESSIGSÄURE DERIVATEN

PROCEDE DE PREPARATION DE DERIVES D'ACIDE 2-(6-SUBSTITUE-1,3-DIOXANE-4-YL)
ACETIQUE

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
IE IT LI LU MC NL PT SE SI SK TR**
Designated Extension States:
AL LT LV MK RO

(30) Priority: **27.12.2001 EP 01000794**

(43) Date of publication of application:
29.09.2004 Bulletin 2004/40

(60) Divisional application:
05014484.9

(73) Proprietor: **AstraZeneca UK Limited**
London,
Greater London W1K 1LN (GB)

(72) Inventor: **HOF, Robert, Patrick**
NL-5981 GD Panningen (NL)

(74) Representative: **Bill, Kevin**
AstraZeneca
R&D Headquarters
Global Intellectual Property, Patents
SE-151 85 Södertälje (SE)


(56) References cited:
EP-A- 1 024 139

EP 1 461 331 B9


Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] The invention relates to a process for the preparation of a 2-(6-substituted-1,3-dioxane-4-yl) acetic acid derivative according to formula 1,



wherein R¹, R² and R³ are each independently a C1-4 alkylgroup or wherein R¹ and R² together with the C-atom to which they are bound form a 5- or 6-membered cycloalkyl and wherein Y stands for R^A-CO- or for R^B-SO₂- with R^A, R^B are chosen from the group of alkyl or aryl with 1-12 C-atoms, from its corresponding 2-(6-substituted-1,3-dioxane-4-yl) acetic acid derivative according to formula 2,



wherein R¹, R² and R³ are as defined above and wherein X stands for a halogen, in the presence of a phase transfer catalyst and an oxyating agent.

[0002] Such a process is known from EP 1 024 139, wherein the preparation of a compound according to formula 1 from a compound according to formula 2 is achieved in the presence of a quaternary ammonium salt (phase transfer catalyst) and a carboxylic acid salt (acyloxylating agent).

[0003] It is the object of the invention to provide an alternative process for the preparation of a 2-(6-substituted-1,3-dioxane-4-yl) acetic acid derivative according to formula 1 from its corresponding 2-(6-substituted-1,3-dioxane-4-yl) acetic acid derivative according to formula 2.

[0004] This is achieved according to the invention by using a quaternary phosphonium ion according to formula 3 as a phase transfer catalyst,




wherein R⁴, R⁵, R⁶, R⁷ each independently stand for an alkyl, cycloalkyl, aralkyl or aryl with 1 to 12 C-atoms, and an ion according to formula 4,




wherein Y is as defined above, as an oxyating agent. The reaction has a high yield.

[0005] The quaternary phosphonium ion according to formula 3 used as a phase transfer catalyst and the ion according to formula 4 used as an oxyating agent may be present in a quaternary phosphonium salt according to formula 5,



wherein Y, R⁴, R⁵, R⁶ and R⁷ are as defined above. The phosphonium salt according to formula 5 can be used as both a phase transfer catalyst and as an oxyating agent. The quarternary phosphonium salt according to formula 5 can be prepared according to methods known to the person skilled in the art (e.g. analogous to the preparation of tetra-n-butylammoniumacetate as described in US 5 278 313).

[0006] In a preferred embodiment of the invention, the phase transfer catalyst and the oxyating agent are not present in the same molecule. In this embodiment, a quarternary phosphonium salt according to formula 3a,



wherein R⁴, R⁵, R⁶ and R⁷ are as defined above and wherein A stands for the anion of the quarternary phosphonium salt and is selected from the group of halogens, for example Cl, Br, I is used as a phase transfer catalyst and the acid salt according to formula 4a,



wherein Y is as defined above, wherein M stands for alkali metal or an alkaline metal, for example Li, K, Na, Mg, Ca, Ba and wherein n represents an integer of 1 or 2, depending on the valence of M is used as an oxyating agent. Preferred is M is K or Na.

[0007] In the process according to the invention, halogens X are preferably Cl, Br or I, more preferably Cl.

[0008] In the process according to the invention, R¹, R² and R³ are preferably a C1-4 alkyl group, more preferably R¹ and R² are a methyl or an ethyl group, more preferably a methyl group. R³ is preferably a methyl or a butyl, most preferably a t-butyl.

[0009] In the process according to the invention Y groups are preferably represented by R^A-CO- or R^B-SO₂-, wherein R^A, R^B are chosen from the group of C₁-C₄ alkyl or aryl with 6-10 C-atoms. In a preferred embodiment, Y is chosen from the group of acyl, more preferably acetyl (with R^A is CH₃), benzenesulphonyl (with R^B is benzene), more preferably nitro substituted benzenesulphonyl (with R^B is p-nitrobenzene), tosyl (with R^B is p-methyl-benzene) or mesyl (with R^B is methyl).

[0010] In the process according to the invention, preferably a phosphonium salt according to formula 3a or according to formula 5, with at least three out of four R groups are the same (e.g. R⁴, R⁵, R⁶ are butyl and R⁷ is methyl or R⁴, R⁵, R⁶ are phenyl and R⁷ is butyl), more preferably a phosphonium salt with all four R groups are the same, is used.

[0011] R^A and R^B and R⁴, R⁵, R⁶, R⁷, -in case R⁴, R⁵, R⁶, R⁷ are aryl or aralkyl-, may be substituted for example with substituents chosen from the group of halogens, alkoxy (e.g. methoxy or ethoxy) with 1-6 C-atoms, alkyl with 1-6 C atoms, (e.g. methyl if R^B is toluene) or nitro, preferably only R^A or R^B are substituted.

[0012] The quarternary phosphonium salt according to formula 3a is preferably used in a molar equivalent amount of 0.01 to 1.0, more preferably 0.05 to 0.7, most preferably 0.1 to 0.5 relative to the amount of compound according to formula 2.

[0013] The quarternary phosphonium salt according to formula 5 is preferably used in a molar equivalent amount of 0.8 to 5, preferably 1 to 3, most preferably 1 to 1.5.

[0014] The acid salt according to formula 4a is preferably used in a molar equivalent amount of 1 to 5 relative to the amount of compound according to formula 2 present. More preferably, a molar equivalent amount of acid salt according to formula 4a of 1 to 4, most preferably 2 to 3, is used.

[0015] The solvents suitable for use in the present invention are a various number of organic solvents, which are known to the person skilled in the art. Organic solvents, which may be used are hydrocarbon series solvents, for example benzene, toluene, cyclohexane, etc; ether series solvents, for example diethyl ether tetrahydrofuran, 1,4-dioxane, methyl-

10

15

Example 1

20

25


30


40

45

50

55





10 wherein R¹, R² and R³ are as defined above and wherein X stands for a halogen, in the presence of a phase transfer catalyst and an oxyating agent, **characterized in that** a quarternary phosphonium ion according to formula 3,




20 wherein R⁴, R⁵, R⁶, R⁷ each independently stand for an alkyl, cycloalkyl, aralkyl or aryl with 1 to 12 C-atoms, is used as a phase transfer catalyst and an ion according to formula 4,



25 wherein Y is as defined above, is used as an oxyating agent;

wherein R^A and R^B and R⁴, R⁵, R⁶, R⁷, -in case R⁴, R⁵, R⁶, R⁷ are aryl or aralkyl-, may be substituted with substituents chosen from the group of halogens, alkoxy with 1-6 C-atoms, alkyl with 1-6 C atoms, or nitro.

- 30
2. Process according to claim 1, **characterized in that** R^A, R^B are chosen from the group of C₁-C₄ alkyl or aryl with 6-10 C-atoms.
 3. Process according to any of claims 1-2, **characterized in that** as a phase transfer catalyst a quarternary phosphonium salt according to formula 3a,



wherein R⁴, R⁵, R⁶ and R⁷ are as defined above and wherein A stands for a halogen, is used and **in that** an acid salt according to formula 4a,



wherein Y is as defined above, wherein M stands for alkali metal or an alkaline metal, is used as an oxyating agent.


- 50
4. Process according to claim 3, **characterized in that** the quarternary phosphonium salt according to formula 3a is used in a molar equivalent amount of 0.05 to 0.7 relative to the amount of compound according to formula 2.
 5. Process according to claim 4, **characterized in that** the quarternary phosphonium salt according to formula 3a is used in a molar equivalent amount of 0.1 to 0.5 relative to the amount of compound according to formula 2.
 - 55 6. Process according to any of claims 1-5, **characterized in that** the process is carried out at a temperature between 100 and 160°C.
 7. Process according to any of claims 1-6, **characterized in that** the process is carried out at a temperature between

110 and 150°C.

8. Process according to any of claims 1-7, **characterized in that** the compound according to formula 1 is tert-butyl 2-
 5 { (4R,6S)-2,2 dimethyl-6-[(methylcarbonyloxy)methyl]-1,3-dioxan-4-yl}acetate and **in that** the compound according to formula 2 is tert-butyl 2-[(4R,6S)-6-(chloromethyl)-2,2-dimethyl-1,3-dioxan-4-yl]acetate.


Patentansprüche

1. Verfahren zur Herstellung eines 2-(1,3-Dioxan-4-yl)essigsäurederivats mit 6-substituierter Dioxanylgruppe gemäß Formel 1




(1)

worin R¹, R² und R³ jeweils unabhängig voneinander für eine C₁₋₄-Alkylgruppe stehen oder R¹ und R² gemeinsam mit dem C-Atom, an das sie gebunden sind, 5- oder 6-gliedriges Cycloalkyl bilden und Y für R^A-CO- oder R^B-SO₂- steht, wobei R^A und R^B aus der Gruppe bestehend aus Alkyl oder Aryl mit 1-12 C-Atomen ausgewählt sind, aus dem entsprechenden 2-(1,3-Dioxan-4-yl)essigsäurederivat mit 6-substituierter Dioxanylgruppe gemäß Formel 2



(2)

worin R¹, R² und R³ die oben angegebene Bedeutung besitzen und X für Halogen steht, in Gegenwart eines Phasentransferkatalysators und eines Oxylierungsmittels, **dadurch gekennzeichnet, daß** man als Phasentransferkatalysator ein quaternäres Phosphoniumion gemäß Formel 3




(3)

worin R⁴, R⁵, R⁶ und R⁷ jeweils unabhängig voneinander für Alkyl, Cycloalkyl, Aralkyl oder Aryl mit 1 bis 12 C-Atomen stehen, und als Oxylierungsmittel ein Ion gemäß Formel 4

OY⁻ (4)

worin Y die oben angegebene Bedeutung besitzt, verwendet;
 wobei R^A und R^B und R⁴, R⁵, R⁶ und R⁷ in dem Fall, daß R⁴, R⁵, R⁶ und R⁷ für Aryl oder Aralkyl stehen - durch Substituenten aus der Gruppe bestehend aus Halogenen, Alkoxy mit 1-6 C-Atomen, Alkyl mit 1-6 C-Atomen oder Nitro substituiert sind.

2. Verfahren nach Anspruch 1, **dadurch gekennzeichnet, daß** man R^A und R^B aus der Gruppe bestehend aus C_1 - C_4 -Alkyl oder Aryl mit 6-10 C-Atomen auswählt.
3. Verfahren nach einem der Ansprüche 1-2, **dadurch gekennzeichnet, daß** man als Phasentransferkatalysator ein quaternäres Phosphoniumsalz gemäß Formel 3a



worin R^4 , R^5 , R^6 und R^7 die oben angegebene Bedeutung besitzen und A für Halogen steht, und als Oxylierungsmittel ein Säuresalz gemäß Formel 4a




worin Y die oben angegebene Bedeutung besitzt und M für Alkalimetall oder ein alkalisches Metall steht, verwendet.


4. Verfahren nach Anspruch 3, **dadurch gekennzeichnet, daß** man das quaternäre Phosphoniumsalz gemäß Formel 3a in einer Moläquivalentmenge von 0,05 bis 0,7, bezogen auf die Menge an Verbindung der Formel 2, verwendet.
5. Verfahren nach Anspruch 4, **dadurch gekennzeichnet, daß** man das quaternäre Phosphoniumsalz gemäß Formel 3a in einer Moläquivalentmenge von 0,1 bis 0,5, bezogen auf die Menge an Verbindung der Formel 2, verwendet.
6. Verfahren nach einem der Ansprüche 1-5, **dadurch gekennzeichnet, daß** man es bei einer Temperatur zwischen 100 und 160°C durchführt.
7. Verfahren nach einem der Ansprüche 1-6, **dadurch gekennzeichnet, daß** man es bei einer Temperatur zwischen 110 und 150°C durchführt.
8. Verfahren nach einem der Ansprüche 1-7, **dadurch gekennzeichnet, daß** es sich bei der Verbindung gemäß Formel 1 um 2-[(4R,6S)-2,2-Dimethyl-6-[(methylcarbonyloxy)methyl]-1,3-dioxan-4-yl]essigsäure-tert.-butylester und bei der Verbindung gemäß Formel 2 um 2-[(4R,6S)-6-(Chlormethyl)-2,2-dimethyl-1,3-dioxan-4-yl]essigsäure-tert.-butylester handelt.

Revendications

1. Procédé pour la préparation d'un dérivé d'acide 2-((6-substituant)-1,3-dioxane-4-yl)acétique selon la formule 1,




dans laquelle R^1 , R^2 et R^3 sont chacun indépendamment un groupe alkyle en C1-4 ou dans laquelle R^1 et R^2 avec l'atome C auquel ils sont liés forment un cycloalkyle à 5 ou 6 éléments et dans laquelle Y représente R^A -CO- ou R^B -SO₂-, R^A et R^B étant choisis entre un alkyle ou un aryle ayant 1-12 atomes de C, à partir de son dérivé d'acide 2-((6-substituant)-1,3-dioxane-4-yl)acétique correspondant selon la formule 2,



(2)

dans laquelle R^1 , R^2 et R^3 sont tels que définis ci-dessus et dans laquelle X représente un halogène, en présence d'un catalyseur de transfert de phase et d'un agent oxydant, **caractérisé en ce que** qu'on utilise comme catalyseur de transfert de phase un ion phosphonium quaternaire selon la formule 3,



(3)


dans laquelle R^4 , R^5 , R^6 , R^7 représentent chacun indépendamment un alkyle, un cycloalkyle, un aralkyle ou un aryle ayant 1 à 12 atomes de C et **en ce qu'**on utilise comme agent oxydant un ion selon la formule 4,



dans laquelle Y est tel que défini ci-dessus ;

dans lequel R^A et R^B et R^4 , R^5 , R^6 , R^7 , -dans le cas où R^4 , R^5 , R^6 , R^7 sont un aryle ou un aralkyle-, peuvent être substitués par des substituants choisis entre des halogènes, un alcoxy ayant 1-6 atomes de C, un alkyle ayant 1-6 atomes de C ou un nitro.

2. Procédé selon la revendication 1, **caractérisé en ce que** R^A , R^B sont choisis entre un alkyle en C_1 - C_4 ou un aryle ayant 6-10 atomes de C.
3. Procédé selon l'une quelconque des revendications 1-2, **caractérisé en ce qu'**on utilise comme catalyseur de transfert de phase un sel de phosphonium quaternaire selon la formule 3a,



(3a)


dans laquelle R^4 , R^5 , R^6 et R^7 sont tels que définis ci-dessus et dans laquelle A représente un halogène et **en ce qu'**on utilise comme agent oxydant un sel d'acide selon la formule 4a,



dans laquelle Y est tel que défini ci-dessus, dans laquelle M représente un métal d'un alcali ou un métal alcalin.


4. Procédé selon la revendication 3, **caractérisé en ce que** le sel de phosphonium quaternaire selon la formule 3a est utilisé en quantité en équivalent molaire de 0,05 à 0,7 par rapport à la quantité de composé selon la formule 2.

5. Procédé selon la revendication 4, **caractérisé en ce qu'on** utilise le sel de phosphonium quaternaire selon la formule 3a en quantité en équivalent molaire de 0,1 à 0,5 par rapport à la quantité de composé selon la formule 2.
6. Procédé selon l'une quelconque des revendications 1-5, **caractérisé en ce qu'on** effectue le procédé à une température comprise entre 100 et 160°C.
7. Procédé selon l'une quelconque des revendications 1-6, **caractérisé en ce qu'on** effectue le procédé à une température comprise entre 110 et 150°C.
8. Procédé selon l'une quelconque des revendications 1-7, **caractérisé en ce que** le composé selon la formule 1 est le 2-[(4R,6S)-2,2-diméthyl-6-[(méthylcarbonyloxy)méthyl]-1,3-dioxan-4-yl]acétate de tert-butyle et **en ce que** le composé selon la formule 2 est le 2-[(4R,6S)-6-(chlorométhyl)-2,2-diméthyl-1,3-dioxan-4-yl]acétate de tert-butyle.




(1)

Formula 1




(2)

Formula 2



(3)

Formula 3




(3a)

Formula 3a




(4)

Formula 4



(4a)

Formula 4a



(5)

Formula 5