

(19)



(11)

**EP 1 478 339 B9**

(12)

**CORRECTED EUROPEAN PATENT SPECIFICATION**

(15) Correction information:

**Corrected version no 1 (W1 B1)**  
**Corrections, see**  
**Claims EN 1, 4, 15, 19, 21, 29**

(51) Int Cl.:

**A61K 9/08** (2006.01) **A61K 47/18** (2006.01)  
**A61P 35/00** (2006.01)

(48) Corrigendum issued on:

**19.08.2009 Bulletin 2009/34**

(86) International application number:

**PCT/US2003/005018**

(45) Date of publication and mention of the grant of the patent:

**04.06.2008 Bulletin 2008/23**

(87) International publication number:

**WO 2003/072082 (04.09.2003 Gazette 2003/36)**

(21) Application number: **03709188.1**

(22) Date of filing: **20.02.2003**

(54) **PHARMACEUTICAL FORMULATIONS OF ANTINEOPLASTIC AGENTS, IN PARTICULAR TEMOZOLOMIDE, PROCESSES OF MAKING AND USING THE SAME**

PHARMAZEUTISCHE ZUBEREITUNGEN VON ANTINEOPLASTISCHEN WIRKSTOFFEN, INSBESONDERE TEMOZOLOMID, VERFAHREN ZU IHRER HERSTELLUNG UND DEREN VERWENDUNG

FORMULATIONS PHARMACEUTIQUES D'AGENTS ANTI-NÉOPLASTIQUES, NOTAMMENT LA TEMOZOLOMIDE, ET PROCÉDÉS DE FABRICATION ET D'UTILISATION ASSOCIÉS

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LI LU MC NL PT SE SI SK TR**

Designated Extension States:

**AL LT LV MK RO**

• **WITCHEY-LAKSHMANAN, Leonore, C.**  
**Piscataway, NJ 08854 (US)**

(30) Priority: **22.02.2002 US 359198 P**

(74) Representative: **Adams, Harvey Vaughan John et al**

**Mathys & Squire LLP**  
**120 Holborn**  
**London**  
**EC1N 2SQ (GB)**

(43) Date of publication of application:

**24.11.2004 Bulletin 2004/48**

(56) References cited:

**EP-A- 0 653 210** **US-A- 4 455 256**  
**US-A- 4 675 183** **US-B1- 6 251 886**

(60) Divisional application:

**08102060.4 / 1 938 798**

(73) Proprietor: **Schering Corporation**

**Kenilworth, NJ 07033-0530 (US)**

• **DATABASE WPI Section Ch, Week 198906**  
**Derwent Publications Ltd., London, GB; Class B05, AN 1989-042233 XP002243408 & JP 63 313721 A (TEYSAN SEIYAKU KK), 21 December 1988 (1988-12-21)**

(72) Inventors:

- **UGWU, Sydney**  
**Gurnee, IL 60031 (US)**
- **RADHAKRISHNAN, Vinay**  
**Laurence Harbor, NJ 08879 (US)**
- **IHNAT, Peter, M.**  
**Brooklyn, NY 11222 (US)**

• **DATABASE WPI Section Ch, Week 199009**  
**Derwent Publications Ltd., London, GB; Class D13, AN 1990-065752 XP002243409 & SU 1 479 049 A (BIOLAR IND COMPLEX), 15 May 1989 (1989-05-15)**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 1 478 339 B9**

**Description**

**[0001]** This application claims priority from U.S. provisional patent application, Serial No. 60/359,198 filed February 22, 2002.

FIELD OF THE INVENTION

**[0002]** The present invention pertains to pharmaceutical formulations comprising antineoplastic agents, such as Temozolomide, and dissolution enhancing agents.

BACKGROUND OF INVENTION

**[0003]** Antineoplastic agents are useful in cancer therapies against a wide array of cancer and other diseases. Temozolomide is one such antineoplastic agent. U.S. Patent 6,096,759 lists a variety of antineoplastic agents including Temozolomide, the disclosure of which is incorporated herein by reference.

**[0004]** Temozolomide is known for its anti-tumor effects. For example, one study showed that clinical responses were achieved in 17% of patients having advanced melanoma (Newlands ES, et al. Br J Cancer 65 (2) 287-2981 (1992)). In another study, a clinical response was achieved in 21 % of patients with advanced melanoma (Journal of Clinical Oncology, 13(4) 910-913 (1995)). Treatment of gliomas in adults with Temozolomide is also known (Eur. J. Cancer 29A 940 (1993)). Treatment of the following cancers in adults with Temozolomide has also been disclosed: metastatic melanoma; malignant melanoma, high grade glioma, glioblastoma and other brain cancers; lung cancer; breast cancer; testicular cancer; colon and rectal cancers; carcinomas; sarcomas; lymphomas; leukemias; anaplastic astrocytoma; and mycosis fungoides.

**[0005]** Temozolomide is a water-insoluble compound. Temozolomide has been administered in patients as micronized suspensions, as disclosed in U.S. Patent 6,251,886. However, suspension formulations are not desirable because they may lead to clogged veins.

**[0006]** Storage of pharmaceutical and biological agents, especially antineoplastic agents, as a frozen solution can cause the active ingredient therein to rapidly deteriorate.

**[0007]** Lyophilization, also known as freeze-drying, is a process whereby water is sublimed from a composition after it is frozen. In this process, pharmaceutical and biological agents that are relatively unstable in an aqueous solution over a period of time can be placed into dosage containers in an easily processed liquid state, dried without the use of damaging heat and stored in a dried state for extended periods. Most pharmaceutical and biological agents, including antineoplastic agents, require additional ingredients to protect the active ingredient during lyophilization. In addition, it can be difficult to reconstitute a lyophilized antineoplastic agent into an aqueous solution.

**[0008]** Accordingly there is an increased need for formulations containing antineoplastic agents, such as Temozolomide, which are water soluble, stable and/or suitable for lyophilization, long term storage and reconstitution of the lyophilized formulation into an aqueous solution.

**[0009]** Furthermore, there is an increased need for administering to a patient an antineoplastic agent, such as Temozolomide, as a water soluble and stable formulation.

SUMMARY OF THE INVENTION

**[0010]** This invention relates to pharmaceutical formulations comprising at least one antineoplastic agent, processes of making the same, processes of lyophilization of the pharmaceutical formulations, lyophilized powders and articles of manufacture thereof, pharmaceutical formulations comprising the lyophilized powder reconstituted in at least one aqueous diluent, and processes of treating or preventing diseases comprising administering the pharmaceutical formulation to an animal in need thereof.

**[0011]** One aspect of the invention relates to a pharmaceutical formulation comprising at least one antineoplastic agent or a pharmaceutically acceptable salt thereof, at least one aqueous diluent, and at least one dissolution enhancing agent sufficient to substantially dissolve said antineoplastic agent(s), wherein said dissolution enhancing agent is urea, L-histidine, L-threonine, L-asparagine, L-serine, L-glutamine or mixtures thereof.

**[0012]** Another aspect of the invention relates to a process for making the pharmaceutical formulation of the invention. This process comprises the steps of dissolving at least one dissolution enhancing agent in at least one aqueous diluent, and adding at least one antineoplastic agent or a pharmaceutically acceptable salt thereof.

**[0013]** Another aspect of the invention relates to a lyophilized powder produced by lyophilization of the pharmaceutical formulation of the invention.

**[0014]** Another aspect of the invention relates to an article of manufacture comprising a container containing the lyophilized powder of the invention:

**[0015]** Another aspect of the invention relates to a pharmaceutical formulation suitable for administration to a patient, wherein the formulation is prepared by reconstituting (resolubilizing) the lyophilized powder of the invention in a volume of water or other aqueous diluent.

**[0016]** Another aspect of the invention relates to a process for treating or preventing diseases in patients comprising administering a therapeutically effective amount of the pharmaceutical formulation of the invention to a patient in need thereof.

**[0017]** Other aspects of this invention relate to and disclose pharmaceutical formulations of Temozolomide, a process of making the same, a lyophilized powder of said formulation and articles of manufacture thereof, a pharmaceutical formulation comprising the lyophilized powder reconstituted in water or other aqueous diluents, and a process of treating or preventing diseases (such as, for example, cancer) comprising administering the pharmaceutical formulation to a patient in need thereof.

#### DETAILED DESCRIPTION

**[0018]** The pharmaceutical formulation of the invention comprises at least one antineoplastic agent or a pharmaceutically acceptable salt thereof, at least one aqueous diluent, and at least one dissolution enhancing agent sufficient to substantially dissolve the antineoplastic agent in the aqueous diluent(s). The percentage of the antineoplastic agent which is dissolved in the pharmaceutical formulation can range from about 50% to about 100%, preferably from about 75% to about 100%, and more preferably about 100%.

**[0019]** The dissolution enhancing agent is urea, L-histidine, L-threonine, L-asparagine, L-serine, L-glutamine or mixtures thereof. The dissolution enhancing agent increases the rate in which the antineoplastic agent dissolves in the aqueous diluent(s). The time to it takes to complete dissolution of at least one antineoplastic agent with a dissolution agent in at least one aqueous diluent in a 25 mg vial can range from about 30 seconds to about 90 seconds, preferably from about 30 seconds to about 60 seconds, more preferably about 30 seconds.

**[0020]** When urea is used in the pharmaceutical formulation as the dissolution enhancing agent, its weight percent (wt%) in the pharmaceutical formulation can range from about 4 wt% to about 60 wt%, preferably from about 8 wt% to about 30 wt%, more preferably from about 12 wt% to about 22 wt%.

**[0021]** When L-histidine, L-threonine, L-asparagine, L-serine, L-glutamine or mixtures thereof are used in the pharmaceutical formulation as the dissolution enhancing agent(s), its wt% in the pharmaceutical formulation can range from about 2 wt% to about 60 wt%, preferably from about 4 wt% to about 40 wt%, more preferably from about 8 wt% to about 20 wt%.

**[0022]** When L-histidine is the only amino acid used in the pharmaceutical formulation as the dissolution enhancing agent, its wt% in the pharmaceutical formulation can range preferably from about 1 wt% to about 30 wt%, more preferably from about 2 wt% to about 20 wt%, and most preferably from about 4 wt% to about 10 wt%.

**[0023]** Non-limiting examples of useful antineoplastic agents include Temozolomide (commercially available under the trademark TEMODAR® from Schering-Plough Corporation, Kenilworth, New Jersey), Uracil Mustard, Chlormethine, Cyclophosphamide, Ifosfamide, Melphalan, Chlorambucil, Pipobroman, Triethylenemelamine, Triethylenethiophosphoramine, Busulfan, Carmustine, Lomustine, Streptozocin, Dacarbazine, Methotrexate, 5-Fluorouracil, Floxuridine, Cytarabine, 6-Mercaptopurine, 6-Thioguanine, Fludarabine phosphate, Pentostatine, Gemcitabine, Vinblastine, Vincristine, Vindesine, Bleomycin, Dactinomycin, Daunorubicin, Doxorubicin, Epirubicin, Idarubicin, Paclitaxel, Mithramycin, Deoxycoformycin, Mitomycin-C, L-Asparaginase, Interferons, Etoposide, Teniposide 17.alpha.-Ethinylestradiol, Diethylstilbestrol, Testosterone, Prednisone, Fluoxymesterone, Dromostanolone propionate, Testolactone, Megestrolacetate, Tamoxifen, Methylprednisolone, Methyltestosterone, Prednisolone, Triamcinotone, Chlorotrianisene, Hydroxyprogesterone, Aminoglutethimide, Estramustine, Medroxyprogesteroneacetate, Leuprolide, Flutamide, Toremifene, Goserelin, Cisplatin, Carboplatin, Hydroxyurea, Amsacrine, Procarbazine, Mitotane, Mitoxantrone, Levamisole, Navelbene, Anastrozole, Letrozole, Capecitabine, Reloxafine, Droloxafine, Hexamethylmelamine, and mixtures thereof.

**[0024]** In a preferred embodiment, at least one of the antineoplastic agents is Temozolomide.

**[0025]** In another preferred embodiment, the antineoplastic agent is a therapeutically effective amount of Temozolomide.

**[0026]** The wt% of the antineoplastic agent in the pharmaceutical formulation can range from about 1 wt% to about 50 wt%, preferably from about 2 wt% to about 30 wt%, more preferably from about 4 wt% to about 16 wt%.

**[0027]** In another embodiment, the pharmaceutical formulation further comprises at least one excipient. Non-limiting examples of suitable excipients include polysorbates, polyethylene glycols (PEG), propylene glycols, polysorbates or suitable mixtures thereof. The excipient is used to increase the solubility of the antineoplastic agent.

**[0028]** The average molecular weight of polysorbates can range from about 500g/mole to about 1900g/mole, preferably from about 800g/mole to about 1600g/mole, more preferably from about 1000g/mole to about 1400g/mole. Non-limiting examples of polysorbates include polysorbate 20, polysorbate 21, polysorbate 40, polysorbate 60, polysorbate 61, polysorbate 65, polysorbate 81, polysorbate 85, and polysorbate 120. Preferred polysorbates include polysorbate 20,

polysorbate 80, and mixtures thereof.

**[0029]** The average molecular weight of PEG can range from about 200g/mole to about 600g/mole, preferably from about 200g/mole to about 500g/mole, more preferably from about 200g/mole to about 400g/mole. Non-limiting examples include PEG 200, PEG 300, PEG 400, PEG 540, and PEG 600.

**[0030]** Propylene glycol is a small molecule with a molecular weight of about 76.1 g/mole.

**[0031]** When an excipient is used in the pharmaceutical formulation, its wt% in the pharmaceutical formulation can range from about 1 wt% to about 50 wt%, preferably from about 2 wt% to about 30 wt%, more preferably from about 4 wt% to about 16 wt%.

**[0032]** In another embodiment, the pharmaceutical formulation further comprises at least one bulking agent. Non-limiting examples of suitable bulking agents which can be included in the pharmaceutical formulation include mannitol, lactose, sucrose, sodium chloride, trehalose, dextrose, starch, hydroxyethylstarch (hetastarch), cellulose, cyclodextrins, glycine, or mixtures thereof.

**[0033]** In a preferred embodiment, the bulking agent in the pharmaceutical formulation is mannitol.

**[0034]** When a bulking agent is used in the pharmaceutical formulation, its wt% in the pharmaceutical formulation can range from about 20 wt% to about 80 wt%, preferably from about 35 wt% to about 65 wt%, more preferably from about 40 wt% to about 56 wt%.

**[0035]** In another embodiment, the pharmaceutical formulation further comprises at least one buffer.

**[0036]** Non-limiting examples of suitable buffers which can be included in the pharmaceutical formulation include citrate buffers, lithium lactate, sodium lactate, potassium lactate, calcium lactate, lithium phosphate, sodium phosphate, potassium phosphate, calcium phosphate, lithium maleate, sodium maleate, potassium maleate, calcium maleate, lithium tartarate, sodium tartarate, potassium tartarate, calcium tartarate, lithium succinate, sodium succinate, potassium succinate, calcium succinate, lithium acetate, sodium acetate, potassium acetate, calcium acetate, or mixtures thereof.

**[0037]** Preferably, a buffer used in the pharmaceutical formulation is at least one citrate buffer. Non-limiting examples of suitable citrate buffers include lithium citrate monohydrate, sodium citrate monohydrate, potassium citrate monohydrate, calcium citrate monohydrate, lithium citrate dihydrate, sodium citrate dihydrate, potassium citrate dihydrate, calcium citrate dihydrate, lithium citrate trihydrate, sodium citrate trihydrate, potassium citrate trihydrate, calcium citrate trihydrate, lithium citrate tetrahydrate, sodium citrate tetrahydrate, potassium citrate tetrahydrate, calcium citrate tetrahydrate, lithium citrate pentahydrate, sodium citrate pentahydrate, potassium citrate pentahydrate, calcium citrate pentahydrate, lithium citrate hexahydrate, sodium citrate hexahydrate, potassium citrate hexahydrate, calcium citrate hexahydrate, lithium citrate heptahydrate, sodium citrate heptahydrate, potassium citrate heptahydrate, or calcium citrate heptahydrate.

**[0038]** When a buffer is used in the pharmaceutical formulation, its wt% in the pharmaceutical formulation can range from about 5 wt% to about 60 wt%, preferably from about 10 wt% to about 40 wt%, more preferably from about 15 wt% to about 28 wt%.

**[0039]** In another embodiment, the pharmaceutical formulation further comprises a pH adjuster. Non-limiting examples of suitable pH adjusters which can be included in the pharmaceutical formulation are hydrochloric acid, sodium hydroxide, citric acid, phosphoric acid, lactic acid, tartaric acid, succinic acid, or mixtures thereof.

**[0040]** A preferred pH adjuster for the pharmaceutical formulation is hydrochloric acid.

**[0041]** When a pH adjuster is used in the pharmaceutical formulation, its wt% in the pharmaceutical formulation can range from about 1 wt% to about 20 wt%, preferably from about 2 wt% to about 12 wt%, more preferably from about 4 wt% to about 8 wt%.

**[0042]** The pH of the pharmaceutical formulation preferably ranges from about 2.5 to about 6.0, more preferably from about 3.0 to about 4.5, and most preferably from about 3.8 to about 4.2.

**[0043]** The pharmaceutical formulation and the lyophilized powders thereof can be stored in containers commonly used in the pharmaceutical industry, which can include plastic containers or glass containers such as standard USP Type I borosilicate glass containers. For example, the container used can be a syringe or vial.

**[0044]** Another aspect of the invention relates to a process for making the pharmaceutical formulation of the invention. This process comprises the steps of dissolving at least one dissolution enhancing agent in at least one aqueous diluent, and adding at least one antineoplastic agent or a pharmaceutically acceptable salt thereof, preferably in that order. In the ideal embodiment, the antineoplastic agent is added after the dissolution enhancing agent is completely dissolved. The dissolution enhancing agent can be urea, L-histidine, L-threonine, L-asparagine, L-serine, L-glutamine or mixtures thereof. The volume of aqueous diluent(s) preferably comprises at least 80% of the total volume.

**[0045]** In another embodiment, the process further comprises adding at least one bulking agent; adding at least one buffer; and adding at least one pH adjuster to form a solution; preferably in that order, and filtering the solution.

**[0046]** In another embodiment, the process further comprises filling the filtered solution into a lyophilization container and lyophilizing the solution contained within the lyophilization container to produce a lyophilized powder. "Lyophilized powders" for purposes of this invention is meant to include all lyophilized forms including lyophilized cakes.

**[0047]** The moisture content of the lyophilized powders can range from to about 0.1 % to about 3%, preferably from about 0.2% to about 0.8%, more preferably from about 0.2% to about 0.6%. Moisture content can be measured by a

moisture analyzer; many suitable moisture analyzers are commercially available.

**[0048]** Lyophilization is a process whereby water is sublimed from a formulation after it is frozen. In this process, pharmaceutical and biological agents that are relatively unstable in an aqueous solution over a period of time can be placed into dosage containers in an easily processed liquid state, dried without the use of damaging heat, and stored in a dried state for extended periods.

**[0049]** Another aspect of the invention relates to lyophilized powders produced by lyophilization of the pharmaceutical formulation of the invention.

**[0050]** Another aspect of the invention relates to an article of manufacture comprising a container containing the lyophilized powder produced by the lyophilization of the pharmaceutical formulation of the invention. Suitable containers are discussed above. In a preferred embodiment, the article of manufacture contains a therapeutically effective amount of the antineoplastic agent(s) in a lyophilized powder.

**[0051]** In another embodiment, the article of manufacture further comprises a volume of at least one aqueous diluent for reconstitution of the lyophilized powder. Reconstitution time generally takes from about 30 seconds to about 60 seconds.

**[0052]** Another aspect of the invention relates to a pharmaceutical formulation suitable for administration to a patient, said formulation prepared by reconstituting (resolubilizing) the lyophilized powder of the invention in a volume of at least one aqueous diluent.

**[0053]** The lyophilized formulations of the pharmaceutical formulations can be diluted or reconstituted prior to administration with a suitable aqueous diluent(s) to obtain a finished concentration. For example, a concentration of from about 0.5 mg/ml to about 5 mg/ml, preferably from about 1 mg/ml to about 3 mg/ml, and more preferably from about 2 mg/ml to about 3 mg/ml is suitable for transfer to an infusion bag for use by a patient in need of an antineoplastic agent such as Temozolomide.

**[0054]** Another aspect of the invention relates to a process for treating or preventing disease in a patient comprising administering a therapeutically effective amount of the pharmaceutical formulation of the invention to a patient in need thereof. The pharmaceutical formulation in this aspect of the invention can be a formulation which are the lyophilized powders reconstituted with water or other aqueous diluents, or the formulations which are not prepared by reconstituting the lyophilized powders. Non-limiting examples of diseases which can be treated or prevented include carcinoma, sarcoma, melanoma, glioma, glioblastoma, brain cancer, lung cancer, thyroid follicular cancer, pancreatic cancer, breast cancer, anaplastic astrocytoma, bladder cancer, myelodysplasia, prostate cancer, testicular cancer, colon and rectal cancer, lymphoma, leukemia, or mycosis fungoides.

**[0055]** The dosage regimen utilizing the pharmaceutical formulations of the invention is selected based upon consideration of a variety of factors, including species, age, weight, sex and medical condition of the patient; the specific disease to be treated, the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular active ingredient or salt thereof employed. An ordinarily skilled physician can readily determine and prescribe the effective amount of antineoplastic agent required to prevent, counter, or arrest the progress of the disease condition. For example, the adult dosage of temozolomide for an adult is generally about 150mg/m<sup>2</sup> of body surface area.

**[0056]** The pharmaceutical formulation of the invention can be used for treating or preventing one or more diseases such as carcinoma, sarcoma, melanoma, glioma, glioblastoma, brain cancer, lung cancer, thyroid follicular cancer, pancreatic cancer, breast cancer, bladder cancer, myelodysplasia, anaplastic astrocytoma, prostate cancer, testicular cancer, anaplastic astrocytoma, colon and rectal cancer, lymphoma, leukemia, and mycosis fungoides.

**[0057]** The pharmaceutical formulation, its lyophilized powder, and the pharmaceutical formulation formed by reconstituting the lyophilized powder with at least one aqueous diluent can provide enhanced chemical stability. Enhanced chemical stability of the pharmaceutical formulation means the pharmaceutical formulation is stable in solution for at least 60 hours at room temperature (about 25°C) and ambient light conditions. Enhanced chemical stability of the lyophilized powder means the lyophilized powder is suitable for storage from about 4°C to about 40°C preferably for about 12 months or more. Enhanced chemical stability of the pharmaceutical formulation formed by reconstituting the lyophilized powder with water or other aqueous diluent means the reconstituted lyophilized powder is stable in solution for about 48 hours or more at room temperature and ambient light conditions. One advantage of the stability of the lyophilized powder is extended pharmaceutical product shelf life. Extended pharmaceutical shelf life offers significant economic advantages.

**[0058]** In yet another embodiment, the present invention discloses stable pharmaceutical formulations comprising temozolomide and at least one dissolution enhancing agent sufficient to substantially dissolve temozolomide in at least one aqueous diluent. The dissolution enhancing agent can be urea, L-histidine, L-threonine, L-asparagine, L-serine, L-glutamine, or mixtures thereof. The pharmaceutical formulation comprising Temozolomide may have at least one other ingredient such as, for example, a bulking agent, buffer or pH adjuster, which are described above both as to their nature and amounts. Such pharmaceutical formulations can have the stability as discussed above. The invention additionally describes a process of preparing such stable pharmaceutical formulations.

[0059] In a further embodiment, the above-described formulations comprising Temozolomide can be lyophilized into a lyophilized powder and stored in a suitable container or such article of manufacture in a condition suitable to be reconstituted later at an appropriate time in water or other aqueous diluent(s) for administration to a patient in need of treatment as described above. The lyophilized powders can have long term storage stability as discussed above.

[0060] The stable pharmaceutical formulations and lyophilized forms comprising Temozolomide are described in more detail in the EXAMPLES section below.

[0061] The invention is therefore advantageous in that it allows the formation of a stable aqueous solution containing an antineoplastic agent. Other advantages include the ability of the pharmaceutical formulation to be lyophilized and stored as a lyophilized powder suitable to be reconstituted as an aqueous solution, which solution is suitable to be administered to a patient in need thereof.

[0062] The term "aseptic" means preventing microbial contamination.

[0063] The term "aqueous diluent(s)" means aqueous fluids suitable for injection into a patient. Non-limiting examples of aqueous diluents include water, normal saline, 5% dextrose solution, and other fluids suitable for injection into a patient, preferably suitable for intravenous injection into a patient.

[0064] Pharmaceutically acceptable salts suitable as acid addition salts as well as salts providing the anion of the quaternary salt include those prepared from acids such as hydrochloric, hydrobromic, phosphoric, sulfuric, maleic, citric, acetic, tartaric, succinic, oxalic, malic, glutamic, pamoic and the like, and other acids related to the pharmaceutically acceptable salts listed in Journal of Pharmaceutical Science, 66, 2 (1977).

[0065] The term "therapeutically effective amount" shall mean that amount of active ingredient that will elicit the biological or medical response of a tissue, system or animal that is being sought by a researcher or clinician.


[0066] The term "extended pharmaceutical shelf life" is intended to mean a shelf life for pharmaceutical products from about 12 months to about 18 months wherein there is a loss of no greater than 10% of the active agent when stored at recommended storage conditions. The active agent for this invention is intended to mean the antineoplastic agent.

[0067] The term "patient" is intended to mean animals, mammals, humans, monkeys, rodents, domestic and farm animals.

[0068] The term "therapeutically effective amount" is intended to mean an amount of a therapeutic agent of the composition, such as temozolomide or other antineoplastic agents described above, that will have an antineoplastic effect on a tissue, system, animal or mammal that is being sought by the administrator (such as a researcher, doctor or veterinarian), which includes alleviation of the symptoms of the condition or disease being treated and the prevention, slowing or halting of progression of the neoplastic condition.

[0069] The term "weight percent" ("wt%") for purposes of this invention is calculated on a basis of total weight of the pharmaceutical formulation.

[0070] The term "Temozolomide" is intended to mean a compound having the formula:



[0071] The synthesis of Temozolomide is well known. See, for example, Stevens et al., J. Med. Chem, 27,196-201 (1984), and Wang et al., J. Chem. Soc., Chem. Commun., 1687-1688 (1994) which are incorporated herein by reference.

## EXAMPLES

[0072] The following examples have been set forth below as a guide to the practitioner and are not meant in any way to limit the scope of the present invention.

EXAMPLE 1: The pharmaceutical formulation of the invention is generally prepared by the following procedure:

[0073]

1. Urea or an amino acid with Polysorbate a bulking agent, and a buffer, is charged and dissolved in at least one

## EP 1 478 339 B9

aqueous diluent, wherein the amino acid is L-histidine, L-threonine, L-asparagine, L-serine, L-glutamine or mixtures thereof.

2. An antineoplastic agent is charged and dissolved into the solution from step 1. Dissolution of the antineoplastic agent is completed by mixing.

3. Water is added to the solution from step 2 to bring the batch to a volume with a desired solution density.

4. The solution from step 3 is aseptically filtered.

**EXAMPLE 2:** \* For batches A-D in examples 2-5, the actual amount of Temozolomide to be charged is adjusted according to the purity of the drug substance batch using the following formula:

[0074]

$$\text{grams Temozolomide} = 2.50 \times 100/\% \text{ Purity.}$$

### Sample Calculation:

[0075] Temozolomide drug substance = 97.0% pure.

$2.50 \times 100/97.0 = 2.58$  grams of Temozolomide to be charged for a 1-Liter batch.

\*\*For batches A-D in Examples 2-5, grams of concentrated Hydrochloric Acid (HCl) to be charged will be calculated as follows:

$$\text{grams of concentrated HCL} = 100 \times (\text{grams of HCl required})/\%w/w \text{ Concentrated HCl}$$

### Sample Calculation:

[0076] Concentrated HCl = 38.0% w/w

$$100 \times 1.48/38.0 = 3.895 \text{ grams Concentrated HCl}$$

[0077] 1 L of Batch A was prepared by the procedure described below.

### Batch A

Ingredients	mg/ml	wt%
*Temozolomide	2.50	8
L-Threonine	4.00	13
Polysorbate 80	3.00	9
Sodium Citrate Dihydrate	5.88	19
Mannitol	15.0	48
**Hydrochloric acid	1.48	5
Water for injection qs ad	1.00 ml	

### PROCESS OF MANUFACTURING 1-LITER OF BATCH A PRIOR TO LYOPHILIZATION:

[0078]

1. 4.00 g of L- Threonine, 3.00 g of polysorbate 80, 15 g of mannitol, 5.88 g of sodium citrate dihydrate, and 1.48 g of Hydrochloric acid, in that order, were charged and dissolved in water with agitation. The amount of water was about 80% of the total volume (batch volume).

## EP 1 478 339 B9

2. \*2.58 g of Temozolomide was charged and dissolved with agitation into the solution from Step 1. Complete dissolution of Temozolomide may require up to 2 hours of mixing.

3. \*\*Water is added to bring the batch to the final volume with a solution density of  $1.008 \pm 0.002$  g/mL at 25°C. The solution was mixed for at least 15 minutes.

4. The solution was aseptically filtered through a sterilized 0.22  $\mu$ m filter (Millipore, GVWP, Durapore), which was washed and tested for integrity, into a sterilized, stainless steel pressure vessel or equivalent. For a batch of 10 liters or less, a 293-mm membrane filter or equivalent was used. For batch sizes greater than 10 Liters, a 0.22  $\mu$ m cartridge filter (MCGL40S Millidisk, Durapore GVWP) was used.

The compounded batch can be stored at room temperature (19 - 25°C) for up to 8 hours in a sealed, sterilized, stainless steel pressure vessel, and then refiltered following storage.

5. The solution from step 4 was aseptically filled into 20-mL Type 1 flint glass vials in aliquots of  $10.7 \pm 0.2$  mL. The vials were washed and sterilized prior to being filled.

6. 20-mm Daikyo D-713 lyo-shape rubber stoppers, which were washed, siliconized and sterilized, were aseptically inserted into the glass vials from step 5 in the lyophilization position.

**EXAMPLE 3:** 1 L of Batch B was prepared by the procedure described above except that 2 g of L-Histidine was used instead of 4.00 g of L-Threonine, and 2.08 g of hydrochloric acid was used instead of 1.48 g of hydrochloric acid in step 1.

### [0079]

**Batch B**

Ingredients	mg/ml	wt%
*Temozolomide	2.50	8%
L-Histidine	2.00	7%
Polysorbate 80	3.00	10%
Sodium Citrate dihydrate	5.88	19%
Mannitol	15.0	49
**Hydrochloric acid	2.08	7
Water for injection qs ad	1.00 ml	

**EXAMPLE 4:** 1L of Batch C was prepared by the procedure described above except that 4 g of L-Asparagine was used instead of 4.00 g of L-Threonine in step 1.

### [0080]

**Batch C**

Ingredients	mg/ml (except for water)	wt%
*Temozolomide	2.50	8
L-Asparagine	4.00	13
Polysorbate 80	3.00	9
Sodium Citrate dihydrate	5.88	19
Mannitol	15.0	48
**Hydrochloric acid	1.48	5
Water for injection qs ad	1.00 ml	

**EXAMPLE 5:** 1 L of Batch D was prepared by the procedure described above except that 5 g of Urea was used instead of 4.00 g of L-Threonine and 3.00 g of Polysorbate 80 in step 1.

### [0081]



**EP 1 478 339 B9**

**Batch D**

Ingredients	mg/ml	wt%
*Temozolomide	2.50	9
Urea	5.00	17
Sodium Citrate dihydrate	5.88	20
Mannitol	15.0	51
**Hydrochloric acid	1.48	5
Water for injection qs ad	1.00 ml	

**[0082]** After lyophilization of batches A-D, the resulting powder was stored and, over a period of one week, four weeks, eight weeks and twelve weeks, the samples were reconstituted with water for analysis. The results are present in Table 1 below.

**Table 1**

Batch #	Stability Time Point/ Condition	% Initial Assay 1	% Initial Assay 2	Mean % Initial (n=2)	Moisture (%)	Ph	Reconstitution Time (see)
A	Initial	100.00	100.00	100.00	0.4	3.75	30
A	1 week, LTC/LTR	100.00	99.63	99.82	0.4	3.77	30
A	1 week, LTC/LTR Control	100.11	99.99	100.05	0.4	3.77	30
A	4 week, 4	100.18	99.62	99.9	0.4	3.83	30
A	4 week, 25H	100.24	99.13	99.69	0.3	3.84	30
A	4 week, 40	99.17	98.11	98.64	0.4	3.84	30
A	8 week, 25H	99.43	98.82	99.125	0.4	3.81	30
A	12 week, 4	99.37	99.59	99.48	0.3	3.83	30
A	12 week, 25H	99.24	99.22	99.23	0.4	3.82	30
B	Initial	100.00	100.00	100.00	1.6	3.96	30
B	1 week, LTC/LTR	100.65	100.41	100.53	1.3	6.97	30
B	1 week, LTC/LTR Control	99.63	100.27	99.95	1.6	3.97	30
B	4 week, 4	100.45	100.43	100.44	2.1	4.00	30
B	4 week, 25H	99.89	99.04	99.47	2.0	4.00	30
B	4 week, 40	99.86	99.66	99.76	2.0	4.00	30
B	8 week, 25H	99.91	100.15	100.03	2.0	4.04	30
B	12 week, 4	100.16	100.00	100.08	2.0	4.01	30
B	12 week, 25H	98.84	100.2	99.52	2.0	4.00	30
C	Initial	100.00	100.00	100.00	0.3	4.00	30
C	1 week, LTC/LTR	97.59	103.68	100.635	0.4	4.04	30

**EP 1 478 339 B9**

(continued)

	Batch #	Stability Time Point/ Condition	% Initial Assay 1	% Initial Assay 2	Mean % Initial (n=2)	Moisture (%)	Ph	Reconstitution Time (see)
5	C	1 week. LTC/LTR Control	98.14	99.55	98.845	0.5	4.04	30
	C	4 week. 4	97.09	97.47	97.28	0.4	4.03	30
10	C	4 week, 25H	98.49	97.18	97.835	0.5	4.03	30
	C	4 week. 40	97.96	97.28	97.62	0.6	4.06	90
	C	8 week. 25H	98.42	97.56	97.99	0.6	4.04	30
	C	12 week, 4	98.27	96.94	97.605	0.4	4.02	30
15	C	12 week, 25H	98.67	97.77	98.22	0.6	4.02	30
	D	Initial	100.00	100.00	100.00	0.4	4	30
	D	1 week. LTC/LTR	103.87	98.41	101.14	0.3	4.03	30
20	D	1 week, LTCILTR Control	100.46	98.95	99.705	0.4	4.02	30
	D	4 week, 4	101.06	99.3	100.18	0.4	4.02	30
25	D	4 week. 25H	101.69	98.97	100.33	0.4	4.02	30
	D	4 week. 40	101.49	98.95	100.22	0.7	4.03	30
	D	8 week. 25H	100.86	98.78	99.82	0.5	4.01	30
	D	12 week, 4	101.3	99.28	100.29	0.4	4	30
30	D	12 week, 25H	101.37	98.59	99.98	0.5	4	30

**[0083]** The results show that the temozolamide in batches A-D was stable over one, four, eight and twelve week periods.

**EXAMPLE 6: Procedure for Lyophilization**

**[0084]** The 20-mL glass vials from Examples 2-5, which were filled with the solution from one of Batches A-D, were placed into trays prior to lyophilization. Lyophilization was then carried out by the following procedure. A Lyostar Lyophilizer, which is manufactured by FTS systems, was used for the lyophilization.

1. Lyophilizer shelves were cooled to  $-50 \pm 5^{\circ}\text{C}$ .
2. The trays of filled vials were aseptically loaded onto the lyophilizer shelves.
3. The lyophilizer shelves were held at a temperature of about  $50 \pm 5^{\circ}\text{C}$  for 4.5 hours.
4. The lyophilizer shelves were warmed to about  $-22 \pm 2^{\circ}\text{C}$  in 1.5 hours and the product was maintained at a temperature of about  $-22 \pm 2^{\circ}\text{C}$  for at least 6 hours.
5. The shelves were cooled to about  $-50^{\circ}\text{C}$  in 3 hours and held at about  $-50 \pm 5^{\circ}\text{C}$  for 3 hours.
6. The condenser temperature was lowered to about  $-45^{\circ}\text{C}$  or below, and then the chamber was evacuated to about  $100 \mu\text{m Hg}$ .
7. Once the chamber reached  $100 \mu\text{m Hg}$  pressure, full vacuum ( $50\text{-}70 \mu\text{m Hg}$ ) was applied and held for about 2 hours with the shelf temperature at about  $-50 \pm 5^{\circ}\text{C}$ .
8. The pressure was changed to about  $150 \mu\text{m Hg}$  and held for 30 minutes.
9. The shelves were heated to about  $5^{\circ}\text{C}$  in 1 hour and 45 minutes. The shelf temperature was then maintained at about  $5^{\circ}\text{C}$  for about 6 hours at approximately  $150 \mu\text{m Hg}$  pressure.
10. The shelves were cooled to  $-2^{\circ}\text{C}$  in about 3 hours and the shelf temperature was maintained at about  $-2 \pm 2^{\circ}\text{C}$  for 32 hours at approximately  $150 \mu\text{m Hg}$  pressure.
11. The shelves were heated to about  $45^{\circ}\text{C}$  in 8 hours and maintained at a temperature of about  $45 \pm 2^{\circ}\text{C}$  for about 5 hours at about  $150 \mu\text{m Hg}$  pressure. The product temperature was then kept above  $-10^{\circ}\text{C}$ .
12. The shelves were cooled to about  $4^{\circ}\text{C}$  at a chamber pressure of about  $150 \mu\text{m}$  and maintained at about  $4^{\circ}\text{C}$

for a minimum of about 1 hour.

13. The chamber was vented with sterile filtered nitrogen to about 933 mBar.

14. The vials were stoppered inside the lyophilizer.

15. The chamber was vented with sterile filtered nitrogen to atmospheric pressure.

16. The vials were removed from the lyophilizer and crimped with 20-mm aluminum seals.

17. The vials were stored at about 2° to about 8°C until inspection was completed.

**[0085]** It will be appreciated by those skilled in the art that changes could be made to the embodiments described above, including chemical and stereochemical changes, without departing from the broad inventive concept thereof. It is understood, therefore, that this invention is not limited to the particular embodiments disclosed, but it is intended to cover modifications which are within the spirit and scope of the invention, as defined by the appended claims.

## Claims

1. A pharmaceutical formulation comprising Temozolomide of a pharmaceutically acceptable salt thereof, at least one aqueous diluent, and at least one dissolution enhancing agent sufficient to substantially dissolve said Temozolomide, wherein said dissolution enhancing agent is selected from the group consisting of urea, L-histidine, L-threonine, L-asparagine, L-serine, L-glutamine and mixtures thereof.
2. The pharmaceutical formulation according to claim 1, further comprising an excipient selected from the group consisting of polysorbate, polyethylene glycol, propylene glycol, polypropylene glycol, and mixtures thereof.
3. The pharmaceutical formulation according to claim 2, wherein said excipient is selected from the group consisting of polysorbate 20, polysorbate 80, and mixtures thereof.
4. The pharmaceutical formulation according to claim 1, further comprising at least one bulking agent.
5. The pharmaceutical formulation according to claim 4, wherein said bulking agent is selected from the group consisting of mannitol, lactose, sucrose, sodium chloride, trehalose, dextrose, starch, hetastarch, cellulose, cyclodextrins, glycine, and mixtures thereof.
6. The pharmaceutical formulation according to claim 5, wherein said bulking agent is mannitol.
7. The pharmaceutical formulation according to claim 1, further comprising at least one buffer.
8. The pharmaceutical formulation according to claim 7, wherein said buffer is selected from the group consisting of lithium citrate monohydrate, sodium citrate monohydrate, potassium citrate monohydrate, calcium citrate monohydrate, lithium citrate dihydrate, sodium citrate dihydrate, potassium citrate dihydrate, calcium citrate dihydrate, lithium citrate trihydrate, sodium citrate trihydrate, potassium citrate trihydrate, calcium citrate trihydrate, lithium citrate tetrahydrate, sodium citrate tetrahydrate, potassium citrate tetrahydrate, calcium citrate tetrahydrate, lithium citrate pentahydrate, sodium citrate pentahydrate, potassium citrate pentahydrate, calcium citrate pentahydrate, lithium citrate hexahydrate, sodium citrate hexahydrate, potassium citrate hexahydrate, calcium citrate hexahydrate, lithium citrate heptahydrate, sodium citrate heptahydrate, potassium citrate heptahydrate, calcium citrate heptahydrate, lithium lactate, sodium lactate, potassium lactate, calcium lactate, lithium phosphate, sodium phosphate, potassium phosphate, calcium phosphate, lithium maleate, sodium maleate, potassium maleate, calcium maleate, lithium tartarate, sodium tartarate, potassium tartarate, calcium tartarate, lithium succinate, sodium succinate, potassium succinate, calcium succinate, lithium acetate, sodium acetate, potassium acetate, calcium acetate, and mixtures thereof.
9. The pharmaceutical formulation according to claim 8, wherein said buffer is selected from the group consisting of lithium citrate monohydrate, sodium citrate monohydrate, potassium citrate monohydrate, calcium citrate monohydrate, lithium citrate dihydrate, sodium citrate dihydrate, potassium citrate dihydrate, calcium citrate dihydrate, lithium citrate trihydrate, sodium citrate trihydrate, potassium citrate trihydrate, calcium citrate trihydrate, lithium citrate tetrahydrate, sodium citrate tetrahydrate, potassium citrate tetrahydrate, calcium citrate tetrahydrate, lithium citrate pentahydrate, sodium citrate pentahydrate, potassium citrate pentahydrate, calcium citrate pentahydrate, lithium citrate hexahydrate, sodium citrate hexahydrate, potassium citrate hexahydrate, calcium citrate hexahydrate, lithium citrate heptahydrate, sodium citrate heptahydrate, potassium citrate heptahydrate, calcium citrate heptahydrate and

mixtures thereof.

10. The pharmaceutical formulation according to claim 1, further comprising a pH adjuster.

5 11. The pharmaceutical formulation according to claim 10, wherein said pH adjuster is selected from the group consisting of hydrochloric acid, sodium hydroxyde, citric acid, phosphoric acid, lactic acid, tartaric acid, succinic acid, and mixture thereof.

10 12. The pharmaceutical formulation according to claim 11, wherein said pH adjuster is hydrochloric add.

13. The pharmaceutical formulation according to claim 1, wherein the pH of said formulation ranges from about 2.5 to about 6.0.

15 14. The pharmaceutical formulation according to claim 13. wherein the pH of said formulation ranges from about 3.0 to about 4.5.

15. The pharmaceutical formulation according to claim 14, wherein the pH of said formulation ranges from about 3.8 to about 4.2.

20 16. The pharmaceutical formulation according to claim 1 wherein said aqueous diluent is selected from the group consisting of water, normal saline, 5% dextrose solution, and mixtures thereof.

25 17. The pharmaceutical formulation according to claim 1, wherein said dissolution enhancing agent is urea and wherein said pharmaceutical formulation further comprises hydrochloric acid, at least one citrate buffer, and mannitol.

30 18. The pharmaceutical formulation according to claim 17, wherein said Temozolomide is present in an amount ranging from about 1 wt% to about 50 wt%, said hydrochloric acid is present in an amount ranging from about 1 wt% to about 20 wt%, said citrate buffer(s) is present in an amount ranging from about 5 wt% to about 60 wt%, said urea is present in an amount ranging from about 4 wt% to about 60 wt%, and said mannitol is present in an amount ranging from about 10 wt% to about 85 wt%.

35 19. The pharmaceutical formulation according to claim 1, wherein said dissolution enhancing agent is selected from the group consisting of L-histidine, L-threonine, L-asparagine, L-serine, L-glutamine and mixture thereof, and wherein said pharmaceutical formulation further comprises polysorbate, hydrochloric acid, at least one citrate buffer, mannitol, and water.

40 20. The pharmaceutical formulation according to claim 19, wherein said Temozolomide is present in an amount ranging from 1 wt% to about 50%, said hydrochloric acid is present in an amount ranging from about 1 wt% to about 20 wt%, said citrate buffer (s) is present in an amount ranging from about 5 wt% to about 60wt%, said polysorbate is present in an amount ranging from about 1 wt% about 50 wt%, said dissolution enhancing agent is present in an amount ranging from about 2 wt% to about 60 wt%, and said mannitol to present in an amount ranging from about 15 wt% to about 85 wt%.

45 21. A process of making a pharmaceutical formulation comprising the step of:

(i) dissolving at least one dissolution enhancing agent in at least one aqueous diluent wherein said dissolution enhancing agent is selected from the group consisting of urea, L-histidine, L-threonine, L-asparagine, L-serine, and L-glutamine; and

(ii) adding Temozolomide or a pharmaceutically acceptable salt thereof.

50

22. The process according to claim 21 further comprising:

a) adding at least one bulking agent;

b) adding at least one buffer;

55

c) adding at least one pH adjuster to form a solution; and

d) filtering said solution.

23. The process according to claim 22, further comprising lyophilizing said solution from step (d) to form a lyophilized

powder.

24. The lyophilized powder, produced by the process of claim 23.
- 5 25. An article of manufacture comprising a container containing the lyophilized powder of claim 24.
26. The article of manufacture according to claim 25, wherein said container is a syringe or vial.
- 10 27. The article of manufacture according to claim 25, further comprising a volume of at least one aqueous diluent suitable for reconstitution of said lyophilized powder.
28. A pharmaceutical formulation suitable for administration to a patient, said formulation prepared by reconstituting the lyophilized powder of claim 24 in volume of at least one aqueous diluent.
- 15 29. A pharmaceutical formulation comprising Temozolomide or a pharmaceutically acceptable salt thereof, at least one aqueous diluent, and at least one dissolution enhancing agent sufficient to substantially dissolve said Temozolomide, wherein said dissolution enhancing agent is selected from the group consisting of urea, L-histidine, L-threonine, L-asparagine L-aerine, and L-glutamine, for use as a medicament.
- 20 30. The pharmaceutical formulation of claim 29 for use in treating or preventing carcinoma, sarcoma, melanoma, glioma, glioblastoma, brain cancer, lung cancer, thyroid follicular cancer, pancreatic cancer, breast cancer, bladder cancer, myelodysplasia, prostate cancer, testicular cancer, anaplastic astrocytoma, colon and rectal cancer, lymphoma, leukemia or mycosis fungoides.
- 25 31. The pharmaceutical formulation of claim 29 for administration to an animal.
32. The pharmaceutical formulation of claim 31, wherein said animal is a mammal.
33. The pharmaceutical formulation of claim 32, wherein said mammal is a human.
- 30 34. The pharmaceutical formulation of claim 28 for use as a medicament.
- 35 35. The pharmaceutical formulation of claim 34, for use in treating or preventing carcinoma, sarcoma, melanoma, glioma, glioblastoma, brain cancer, lung cancer, thyroid follicular cancer, pancreatic cancer, breast cancer, bladder cancer, myelodysplasia, prostate cancer, testicular cancer, anaplastic astrocytoma, colon and rectal cancer, lymphoma, leukemia, or mycosis fungoides.
36. The pharmaceutical formulation of claim 34, for administration to an animal.
- 40 37. The pharmaceutical formulation of claim 36, wherein said animal is a mammal.
38. The pharmaceutical formulation of claim 37, wherein said animal is a human.
- 45 39. The pharmaceutical formulation of claim 29, wherein the dissolution enhancing agent is L-threonine.

#### Patentansprüche

- 50 1. Pharmazeutische Formulierung, die Temozolomid oder ein pharmazeutisch annehmbares Salz davon, mindestens ein wässriges Verdünnungsmittel und mindestens ein Auflösungsverbesserungsmittel enthält, das ausreicht, um das Temozolomid im Wesentlichen zu lösen, wobei das Auflösungsverbesserungsmittel ausgewählt ist aus der Gruppe bestehend aus Harnstoff, L-Histidin, L-Threonin, L-Asparagin, L-Serin, L-Glutamin und Mischungen davon.
- 55 2. Pharmazeutische Formulierung nach Anspruch 1, die ferner einen Hilfsstoff ausgewählt aus der Gruppe bestehend aus Polysorbat, Polyethylenglykol, Propylenglykol, Polypropylenglykol und Mischungen davon enthält.
3. Pharmazeutische Formulierung nach Anspruch 2, bei der der Hilfsstoff ausgewählt ist aus der Gruppe bestehend aus Polysorbat 20, Polysorbat 80 und Mischungen davon.

## EP 1 478 339 B9

4. Pharmazeutische Formulierung nach Anspruch 1, die ferner mindestens ein Füllmittel enthält.
5. Pharmazeutische Formulierung nach Anspruch 4, bei der das Füllmittel ausgewählt ist aus der Gruppe bestehend aus Mannit, Lactose, Sucrose, Natriumchlorid, Trehalose, Dextrose, Stärke,  $\beta$ -Stärke, Cellulose, Cyclodextrinen, Glycin und Mischungen davon.
6. Pharmazeutische Formulierung nach Anspruch 5, bei der das Füllmittel Mannit ist.
7. Pharmazeutische Formulierung nach Anspruch 1, die ferner mindestens einen Puffer enthält.
8. Pharmazeutische Formulierung nach Anspruch 7, bei der der Puffer ausgewählt ist aus der Gruppe bestehend aus Lithiumcitratmonohydrat, Natriumcitratmonohydrat, Kaliumcitratmonohydrat, Calciumcitratmonohydrat, Lithiumcitratdihydrat, Natriumcitratdihydrat, Kaliumcitratdihydrat, Calciumcitratdihydrat, Lithiumcitrattrihydrat, Natriumcitrattrihydrat, Kaliumcitrattrihydrat, Calciumcitrattrihydrat, Lithiumcitratтетetrahydrat, Natriumcitratтетetrahydrat, Kaliumcitratтетetrahydrat, Calciumcitratтетetrahydrat, Lithiumcitratpentahydrat, Natriumcitratpentahydrat, Kaliumcitratpentahydrat, Calciumcitratpentahydrat, Lithiumcitrathexahydrat, Natriumcitrathexahydrat, Kaliumcitrathexahydrat, Calciumcitrathexahydrat, Lithiumcitratheptahydrat, Natriumcitratheptahydrat, Kaliumcitratheptahydrat, Calciumcitratheptahydrat, Lithiumlactat, Natriumlactat, Kaliumlactat, Calciumlactat, Lithiumphosphat, Natriumphosphat, Kaliumphosphat, Calciumphosphat, Lithiummaleat, Natriummaleat, Kaliummaleat, Calciummaleat, Lithiumtartrat, Natriumtartrat, Kaliumtartrat, Calciumtartrat, Lithiumsuccinat, Natriumsuccinat, Kaliumsuccinat, Calciumsuccinat, Lithiumacetat, Natriumacetat, Kaliumacetat, Calciumacetat und Mischungen davon.
9. Pharmazeutische Zusammensetzung nach Anspruch 8, bei der der Puffer ausgewählt ist aus der Gruppe bestehend aus Lithiumcitratmonohydrat, Natriumcitratmonohydrat, Kaliumcitratmonohydrat, Calciumcitratmonohydrat, Lithiumcitratdihydrat, Natriumcitratdihydrat, Kaliumcitratdihydrat, Calciumcitratdihydrat, Lithiumcitrattrihydrat, Natriumcitrattrihydrat, Kaliumcitrattrihydrat, Calciumcitrattrihydrat, Lithiumcitratтетetrahydrat, Natriumcitratтетetrahydrat, Kaliumcitratтетetrahydrat, Calciumcitratтетetrahydrat, Lithiumcitratpentahydrat, Natriumcitratpentahydrat, Kaliumcitratpentahydrat, Calciumcitratpentahydrat, Lithiumcitrathexahydrat, Natriumcitrathexahydrat, Kaliumcitrathexahydrat, Calciumcitrathexahydrat, Lithiumcitratheptahydrat, Natriumcitratheptahydrat, Kaliumcitratheptahydrat, Calciumcitratheptahydrat und Mischungen davon.
10. Pharmazeutische Formulierung nach Anspruch 1, die ferner mindestens ein Mittel zum Einstellen des pH-Werts enthält.
11. Pharmazeutische Formulierung nach Anspruch 10, bei der das Mittel zur Einstellung des pH-Werts ausgewählt ist aus der Gruppe bestehend aus Salzsäure, Natriumhydroxid, Citronensäure, Phosphorsäure, Milchsäure, Weinsäure, Bernsteinsäure und Mischungen davon.
12. Pharmazeutische Formulierung nach Anspruch 11, bei der das Mittel zur Einstellung des pH-Werts Salzsäure ist.
13. Pharmazeutische Formulierung nach Anspruch 1, bei der der pH-Wert der Formulierung im Bereich von etwa 2,5 bis etwa 6,0 liegt.
14. Pharmazeutische Formulierung nach Anspruch 13, bei der der pH-Wert der Formulierung im Bereich von etwa 3,0 bis etwa 4,5 liegt.
15. Pharmazeutische Formulierung nach Anspruch 14, bei der der pH-Wert der Formulierung im Bereich von etwa 3,8 bis etwa 4,2 liegt.
16. Pharmazeutische Formulierung nach Anspruch 1, bei der das wässrige Verdünnungsmittel ausgewählt ist aus der Gruppe bestehend aus Wasser, normaler Salzlösung, 5 % Dextroslösung und Mischungen davon.
17. Pharmazeutische Formulierung nach Anspruch 1, bei der das Auflösungsverbesserungsmittel Harnstoff ist und bei der die pharmazeutische Formulierung ferner Salzsäure, mindestens einen Citratpuffer und Mannit enthält.
18. Pharmazeutische Formulierung nach Anspruch 17, bei der das Temozolomid in einer Menge im Bereich von etwa 1 Gew.% bis 50 Gew.% vorhanden ist, die Salzsäure in einer Menge im Bereich von etwa 1 Gew.% bis etwa 20 Gew.% vorhanden ist, der (die) Citratpuffer in einer Menge im Bereich von etwa 5 Gew.% bis etwa 60 Gew.%

## EP 1 478 339 B9

vorhanden ist (sind), der Harnstoff in einer Menge im Bereich von etwa 4 Gew.% bis etwa 60 Gew.% vorhanden ist, und das Mannit in einer Menge im Bereich von etwa 10 Gew.% bis etwa 85 Gew.% vorhanden ist.

- 5 19. Pharmazeutische Formulierung nach Anspruch 1, bei der das Auflösungsverbesserungsmittel ausgewählt ist aus der Gruppe bestehend aus L-Histidin, L-Threonin, L-Asparagin, L-Serin, L-Glutamin und Mischungen davon, und wobei die pharmazeutische Formulierung ferner Polysorbat, Salzsäure, mindestens einen Citratpuffer, Mannit und Wasser enthält.
- 10 20. Pharmazeutische Formulierung nach Anspruch 19, bei der das Temozolomid in einer Menge im Bereich von etwa 1 Gew.% bis etwa 50 Gew.% vorhanden ist, die Salzsäure in einer Menge im Bereich von etwa 1 Gew.% bis etwa 20 Gew.% vorhanden ist, der (die) Citratpuffer in einer Menge im Bereich von etwa 5 Gew.% bis etwa 60 Gew.% vorhanden ist (sind), das Polysorbat in einer Menge im Bereich von etwa 1 Gew.% bis etwa 50 Gew.% vorhanden ist, das Auflösungsverbesserungsmittel in einer Menge im Bereich von etwa 2 Gew.% bis etwa 60 Gew.% vorhanden ist, und das Mannit in einer Menge im Bereich von etwa 15 Gew.% bis etwa 85 Gew.% vorhanden ist.
- 15 21. Verfahren zur Herstellung einer pharmazeutischen Formulierung, bei dem
- (i) mindestens ein Auflösungsverbesserungsmittel in mindestens einem wässrigen Verdünnungsmittel gelöst wird, wobei das Auflösungsverbesserungsmittel ausgewählt ist aus der Gruppe bestehend aus Harnstoff, L-Histidin, L-Threonin, L-Asparagin, L-Serin und L-Glutamin, und
- 20 (ii) Temozolomid oder ein pharmazeutisch annehmbares Salz davon zugegeben wird.
22. Verfahren nach Anspruch 21, bei dem ferner
- 25 a) mindestens ein Füllmittel zugegeben wird;  
b) mindestens ein Puffer zugegeben wird;  
c) mindestens ein Mittel zum Einstellen des pH-Werts zugegeben wird, um eine Lösung zu bilden, und  
d) die Lösung filtriert wird.
- 30 23. Verfahren nach Anspruch 22, bei dem ferner die Lösung aus Stufe (d) lyophilisiert wird, um ein lyophilisiertes Pulver zu bilden.
24. Lyophilisiertes Pulver, das nach dem Verfahren gemäß Anspruch 23 hergestellt ist.
- 35 25. Fertigungsgegenstand, der einen Behälter umfasst, welcher das lyophilisierte Pulver gemäß Anspruch 24 enthält.
26. Fertigungsgegenstand nach Anspruch 25, bei dem der Behälter eine Spritze oder Ampulle ist.
27. Fertigungsgegenstand nach Anspruch 25, der ferner ein Volumen an mindestens einem wässrigen Verdünnungsmittel enthält, das zum Rekonstituieren des lyophilisierten Pulvers geeignet ist.
- 40 28. Pharmazeutische Formulierung, die zur Verabreichung an einen Patienten geeignet ist, welche durch Rekonstituieren des lyophilisierten Pulvers gemäß Anspruch 24 in einem Volumen von mindestens einem wässrigen Verdünnungsmittel hergestellt ist.
- 45 29. Pharmazeutische Formulierung, die Temozolomid oder ein pharmazeutisch annehmbares Salz davon, mindestens ein wässriges Verdünnungsmittel und mindestens ein Auflösungsverbesserungsmittel enthält, das ausreicht, um das Temozolomid im Wesentlichen zu lösen, wobei das Auflösungsverbesserungsmittel ausgewählt ist aus der Gruppe bestehend aus Harnstoff, L-Histidin, L-Threonin, L-Asparagin, L-Serin, L-Glutamin, zur Verwendung als
- 50 Medikament.
30. Pharmazeutische Formulierung nach Anspruch 29 zur Verwendung bei der Behandlung oder Verhinderung von Karzinom, Sarkom, Melanom, Gliom, Glioblastom, Hirnkrebs, Lungenkrebs, Schilddrüsenfollikelkrebs, Pankreas-
- 55 krebs, Brustkrebs, Blasenkrebs, Myelodysplasie, Prostatakrebs, Hodenkrebs, anaplastischem Astrozytom, Kolon- und Rektumkrebs, Lymphom, Leukämie oder Mycosis fungoides.
31. Pharmazeutische Formulierung nach Anspruch 29 zur Verabreichung an ein Tier.

32. Pharmazeutische Formulierung nach Anspruch 31, bei der das Tier ein Säuger ist.
33. Pharmazeutische Formulierung nach Anspruch 32, bei der der Säuger ein Mensch ist.
- 5 34. Pharmazeutische Formulierung nach Anspruch 28 zur Verwendung als Medikament.
35. Pharmazeutische Formulierung nach Anspruch 34 zur Verwendung bei der Behandlung oder Verhinderung von Karzinom, Sarkom, Melanom, Gliom, Glioblastom, Hirnkrebs, Lungenkrebs, Schilddrüsenfollikelkrebs, Pankreas-  
 10 krebs, Brustkrebs, Blasenkrebs, Myelodysplasie, Prostatakrebs, Hodenkrebs, anaplastischem Astrozytom, Kolon- und Rektumkrebs, Lymphom, Leukämie oder Mycosis fungoides.
36. Pharmazeutische Formulierung nach Anspruch 34 zur Verabreichung an ein Tier.
37. Pharmazeutische Formulierung nach Anspruch 36, bei der das Tier ein Säuger ist.
- 15 38. Pharmazeutische Formulierung nach Anspruch 37, bei der das Tier ein Mensch ist.
39. Pharmazeutische Formulierung nach Anspruch 29, bei der das Auflösungsverbesserungsmittel L-Threonin ist.

20 **Revendications**

1. Formulation pharmaceutique comprenant du témozolomide ou de l'un de ses sels pharmacologiquement admissi-  
 25 bles, au moins un diluant aqueux, et au moins un agent favorisant la dissolution, suffisant pour que ledit témozolomide se dissolve sensiblement, pour laquelle ledit agent favorisant la dissolution est choisi dans l'ensemble constitué par l'urée, la L-histidine, la L-thréonine, la L-asparagine, la L-sérine et la L-glutamine, ainsi que leurs mélanges.
2. Formulation pharmaceutique conforme à la revendication 1, qui comprend en outre un excipient choisi dans l'en-  
 30 semble constitué par les polysorbates, polyéthylèneglycols, propylèneglycol et polypropylèneglycols, ainsi que les mélanges de tels corps.
3. Formulation pharmaceutique conforme à la revendication 2, dans laquelle ledit excipient est choisi parmi du poly-  
 sorbate 20, du polysorbate 80, et leurs mélanges.
- 35 4. Formulation pharmaceutique conforme à la revendication 1, qui comprend en outre au moins un diluant.
5. Formulation pharmaceutique conforme à la revendication 4, dans laquelle ledit diluant est choisi dans l'ensemble  
 constitué par les mannitol, lactose, saccharose, chlorure de sodium, tréhalose, dextrose, amidon, amidon bêta,  
 40 cellulose, cyclodextrines et glycine, ainsi que leurs mélanges.
6. Formulation pharmaceutique conforme à la revendication 5, dans laquelle ledit diluant est du mannitol.
7. Formulation pharmaceutique conforme à la revendication 1, qui comprend en outre au moins un agent tampon.
- 45 8. Formulation pharmaceutique conforme à la revendication 7, dans laquelle ledit agent tampon est choisi dans l'en-  
 semble constitué par les suivants : citrate de lithium monohydraté, citrate de sodium monohydraté, citrate de po-  
 tassium monohydraté, citrate de calcium monohydraté, citrate de lithium dihydraté, citrate de sodium dihydraté,  
 citrate de potassium dihydraté, citrate de calcium dihydraté, citrate de lithium trihydraté, citrate de sodium trihydraté,  
 citrate de potassium trihydraté, citrate de calcium trihydraté, citrate de lithium tétrahydraté, citrate de sodium tétra-  
 50 hydraté, citrate de potassium tétrahydraté, citrate de calcium tétrahydraté, citrate de lithium pentahydraté, citrate  
 de sodium pentahydraté, citrate de potassium pentahydraté, citrate de calcium pentahydraté, citrate de lithium  
 hexahydraté, citrate de sodium hexahydraté, citrate de potassium hexahydraté, citrate de calcium hexahydraté,  
 citrate de lithium heptahydraté, citrate de sodium heptahydraté, citrate de potassium heptahydraté, citrate de calcium  
 55 heptahydraté, lactate de lithium, lactate de sodium, lactate de potassium, lactate de calcium, phosphate de lithium,  
 phosphate de sodium, phosphate de potassium, phosphate de calcium, maléate de lithium, maléate de sodium,  
 maléate de potassium, maléate de calcium, tartrate de lithium, tartrate de sodium, tartrate de potassium, tartrate  
 de calcium, succinate de lithium, succinate de sodium, succinate de potassium, succinate de calcium, acétate de  
 lithium, acétate de sodium, acétate de potassium, acétate de calcium, et leurs mélanges.



## EP 1 478 339 B9

9. Formulation pharmaceutique conforme à la revendication 8, dans laquelle ledit agent tampon est choisi dans l'ensemble constitué par les suivants : citrate de lithium monohydraté, citrate de sodium monohydraté, citrate de potassium monohydraté, citrate de calcium monohydraté, citrate de lithium dihydraté, citrate de sodium dihydraté, citrate de potassium dihydraté, citrate de calcium dihydraté, citrate de lithium trihydraté, citrate de sodium trihydraté, citrate de potassium trihydraté, citrate de calcium trihydraté, citrate de lithium tétrahydraté, citrate de sodium tétrahydraté, citrate de potassium tétrahydraté, citrate de calcium tétrahydraté, citrate de lithium pentahydraté, citrate de sodium pentahydraté, citrate de potassium pentahydraté, citrate de calcium pentahydraté, citrate de lithium hexahydraté, citrate de sodium hexahydraté, citrate de potassium hexahydraté, citrate de calcium hexahydraté, citrate de lithium heptahydraté, citrate de sodium heptahydraté, citrate de potassium heptahydraté, citrate de calcium heptahydraté, et leurs mélanges.
10. Formulation pharmaceutique conforme à la revendication 1, qui comprend en outre au moins un agent d'ajustement de pH.
11. Formulation pharmaceutique conforme à la revendication 10, dans laquelle ledit agent d'ajustement de pH est choisi dans l'ensemble constitué par l'acide chlorhydrique, l'hydroxyde de sodium, l'acide citrique, l'acide phosphorique, l'acide lactique, l'acide tartrique et l'acide succinique, ainsi que leurs mélanges.
12. Formulation pharmaceutique conforme à la revendication 11, dans laquelle ledit agent d'ajustement de pH est de l'acide chlorhydrique.
13. Formulation pharmaceutique conforme à la revendication 1, dont le pH vaut d'environ 2,5 à environ 6,0.
14. Formulation pharmaceutique conforme à la revendication 13, dont le pH vaut d'environ 3,0 à environ 4,5.
15. Formulation pharmaceutique conforme à la revendication 14, dont le pH vaut d'environ 3,8 à environ 4,2.
16. Formulation pharmaceutique conforme à la revendication 1, dans laquelle ledit diluant aqueux est choisi parmi de l'eau, une solution physiologique salée, une solution à 5 % de dextrose, et leurs mélanges.
17. Formulation pharmaceutique conforme à la revendication 1, dans laquelle ledit agent favorisant la dissolution est de l'urée, et laquelle formulation pharmaceutique comprend en outre de l'acide chlorhydrique, au moins un agent tampon citrate, et du mannitol.
18. Formulation pharmaceutique conforme à la revendication 17, dans laquelle ledit témozolomide se trouve en une proportion pondérale d'environ 1 à environ 50 %, ledit acide chlorhydrique se trouve en une proportion pondérale d'environ 1 à environ 20 %, ledit ou lesdits agent(s) tampon(s) citrate(s) se trouve(nt) en une proportion pondérale d'environ 5 à environ 60 %, ladite urée se trouve en une proportion pondérale d'environ 4 à environ 60 %, et ledit mannitol se trouve en une proportion pondérale d'environ 10 à environ 85 %.
19. Formulation pharmaceutique conforme à la revendication 1, dans laquelle ledit agent favorisant la dissolution est choisi dans l'ensemble constitué par la L-histidine, la L-thréonine, la L-asparagine, la L-sérine et la L-glutamine, ainsi que leurs mélanges, et laquelle formulation pharmaceutique comprend en outre un polysorbate, de l'acide chlorhydrique, au moins un agent tampon citrate, du mannitol et de l'eau.
20. Formulation pharmaceutique conforme à la revendication 19, dans laquelle ledit témozolomide se trouve en une proportion pondérale d'environ 1 à environ 50 %, ledit acide chlorhydrique se trouve en une proportion pondérale d'environ 1 à environ 20 %, ledit ou lesdits agent(s) tampon(s) citrate(s) se trouve(nt) en une proportion pondérale d'environ 5 à environ 60 %, ledit polysorbate se trouve en une proportion pondérale d'environ 1 à environ 50 %, ledit agent favorisant la dissolution se trouve en une proportion pondérale d'environ 2 à environ 60 %, et ledit mannitol se trouve en une proportion pondérale d'environ 15 à environ 85 %.
21. Procédé de fabrication d'une formulation pharmaceutique, qui comporte les étapes suivantes :
- dissoudre au moins un agent favorisant la dissolution dans au moins un diluant aqueux, lequel agent favorisant la dissolution est choisi dans l'ensemble formé par l'urée, la L-histidine, la L-thréonine, la L-asparagine, la L-sérine et la L-glutamine ;
  - et y ajouter le témozolomide ou de l'un de ses sels pharmacologiquement admissibles.

## EP 1 478 339 B9

22. Procédé conforme à la revendication 21, qui comporte en outre les étapes suivantes :
- a) ajouter au moins un diluant ;
  - b) ajouter au moins un agent tampon ;
  - c) ajouter au moins un agent d'ajustement de pH, de manière à obtenir une solution ;
  - d) et filtrer cette solution.
23. Procédé conforme à la revendication 22, qui comporte en outre le fait de lyophiliser ladite solution issue de l'étape (d), de manière à obtenir une poudre lyophilisée.
24. Poudre lyophilisée obtenue selon un procédé conforme à la revendication 23.
25. Article manufacturé qui comporte un récipient contenant une poudre lyophilisée conforme à la revendication 24.
26. Article manufacturé conforme à la revendication 25, dans lequel ledit récipient est une seringue ou un flacon.
27. Article manufacturé conforme à la revendication 25, qui comporte en outre un certain volume d'au moins un diluant aqueux, approprié pour la reconstitution d'une solution à partir de ladite poudre lyophilisée.
28. Formulation pharmaceutique appropriée pour être administrée à un patient, laquelle formulation est préparée par reconstitution d'une solution à partir d'une poudre lyophilisée conforme à la revendication 24 et d'un certain volume d'au moins un diluant aqueux.
29. Formulation pharmaceutique comprenant du témozolomide ou de l'un de ses sels pharmacologiquement admissibles, au moins un diluant aqueux, et au moins un agent favorisant la dissolution, suffisant pour que ledit témozolomide se dissolve sensiblement, cet agent favorisant la dissolution étant choisi parmi l'urée, la L-histidine, la L-thréonine, la L-asparagine, la L-sérine et la L-glutamine, et conçue pour servir de médicament.
30. Formulation pharmaceutique conforme à la revendication 29, conçue pour servir à traiter ou prévenir un carcinome, un sarcome, un mélanome, un gliome, un glioblastome, un cancer du cerveau, un cancer pulmonaire, un cancer des vésicules thyroïdiennes, un cancer du pancréas, un cancer du sein, un cancer de la vessie, une myélodisplasie, un cancer de la prostate, un cancer du testicule, un glioblastome multiforme, un cancer colo-rectal, un lymphome, une leucémie ou un mycosis fongoïde.
31. Formulation pharmaceutique conforme à la revendication 29, conçue pour être administrée à un animal.
32. Formulation pharmaceutique conforme à la revendication 31, ledit animal étant un mammifère.
33. Formulation pharmaceutique conforme à la revendication 32, ledit mammifère étant un humain.
34. Formulation pharmaceutique conforme à la revendication 28, conçue pour servir de médicament.
35. Formulation pharmaceutique conforme à la revendication 34, conçue pour servir à traiter ou prévenir un carcinome, un sarcome, un mélanome, un gliome, un glioblastome, un cancer du cerveau, un cancer pulmonaire, un cancer des vésicules thyroïdiennes, un cancer du pancréas, un cancer du sein, un cancer de la vessie, une myélodisplasie, un cancer de la prostate, un cancer du testicule, un glioblastome multiforme, un cancer colo-rectal, un lymphome, une leucémie ou un mycosis fongoïde.
36. Formulation pharmaceutique conforme à la revendication 34, conçue pour être administrée à un animal.
37. Formulation pharmaceutique conforme à la revendication 36, ledit animal étant un mammifère.
38. Formulation pharmaceutique conforme à la revendication 37, ledit mammifère étant un humain.
39. Formulation pharmaceutique conforme à la revendication 29, dans laquelle l'agent favorisant la dissolution est de la L-thréonine.

**REFERENCES CITED IN THE DESCRIPTION**

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

**Patent documents cited in the description**

- US 35919802 P [0001]
- US 6096759 A [0003]
- US 6251886 B [0005]

**Non-patent literature cited in the description**

- **NEWLANDS ES et al.** *Br J Cancer*, 1992, vol. 65 (2), 287-2981 [0004]
- *Journal of Clinical Oncology*, 1995, vol. 13 (4), 910-913 [0004]
- *Eur. J. Cancer*, 1993, vol. 29A, 940 [0004]
- *Journal of Pharmaceutical Science*, 1977, vol. 66, 2 [0064]
- **STEVENS et al.** *J. Med. Chem*, 1984, vol. 27, 196-201 [0071]
- **WANG et al.** *J. Chem. Soc., Chem. Commun.*, 1994, 1687-1688 [0071]