



(11) **EP 1 554 576 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

(15) Correction information:
Corrected version no 2 (W2 B1)
Corrections, see
Description Paragraph(s) 7, 12-17

(48) Corrigendum issued on:
20.08.2008 Bulletin 2008/34

(45) Date of publication and mention
of the grant of the patent:
03.10.2007 Bulletin 2007/40

(21) Application number: **02792319.2**

(22) Date of filing: **02.12.2002**

(51) Int Cl.:
G01N 33/53 (2006.01) **G01N 33/537** (2006.01)
G01N 33/566 (2006.01) **G01N 33/557** (2006.01)

(86) International application number:
PCT/US2002/038530

(87) International publication number:
WO 2003/048730 (12.06.2003 Gazette 2003/24)

(54) **IDENTIFICATION OF HIGH AFFINITY MOLECULES BY LIMITED DILUTION SCREENING**

IDENTIFIZIERUNG HOCHAFFINER MOLEKÜLE DURCH SCREENING MIT BEGRENZTER
VERDÜNNUNG

IDENTIFICATION DE MOLECULES DE HAUTE AFFINITE PAR DEPISTAGE PAR DILUTIONS
LIMITEES

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
IE IT LI LU MC NL PT SE SI SK TR**

(30) Priority: **03.12.2001 US 337250 P**

(43) Date of publication of application:
20.07.2005 Bulletin 2005/29

(73) Proprietor: **Amgen Fremont Inc.**
Fremont, CA 94555 (US)

(72) Inventors:
• **BABCOOK, John**
Vancouver, British Columbia V6R 2R2 (CA)
• **KANG, Jaspal, Singh**
Surrey, British Columbia V4N 5H6 (CA)
• **JIA, Xiao-Chi**
San Mateo, CA 94403 (US)
• **ZHANG, Li**
Fremont, CA 94555 (US)

(74) Representative: **Brants, Johan P.E. et al**
De Clercq, Brants & Partners cv
Edgard Gevaertdreef 10a
9830 Sint-Martens-Latem (BE)

(56) References cited:
WO-A-03/048729 US-A- 5 324 633
US-A- 5 800 815 US-A- 5 929 212
US-B1- 6 207 160

- **WATKINS J D ET AL: "Determination of the relative affinities of antibody fragments expressed in Escherichia coli by enzyme-linked immunosorbent assay." ANALYTICAL BIOCHEMISTRY. 1 NOV 1997, vol. 253, no. 1, 1 November 1997 (1997-11-01), pages 37-45, XP002367211 New York ISSN: 0003-2697**
- **BATRA ET AL.: 'Recombinant anti-erbB2 immunotoxins containing pseudomonas exotoxin' PROC. NATL. ACAD. SCI. USA vol. 89, July 1992, pages 5867 - 5871, XP000990722**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

DescriptionField of the Invention

[0001] Embodiments of the invention relate to techniques for determining the binding affinities of molecules for a molecule binding partner. More specifically, embodiments of the invention relate to methods of determining the relative binding affinity of antibodies to a target antigen by contacting each antibody with limited dilutions of antigen.

Background of the Invention

[0002] During an immune response B-cells are stimulated to generate antibodies that specifically bind molecules that activate the immune system. The genetic recombination events involved with the synthesis of antibodies enables animals to create an immense diversity of antibodies each with its own characteristics such as binding specificity and binding affinity. Antibodies are currently being used for diverse diagnostic, imaging, or therapeutic applications. Antibodies that can be successfully used for diagnostic, imaging or therapeutic applications often require special characteristics. In particular, the kinetic properties of the antibody often dictate their utility. In general, the effectiveness or suitability of an antibody shows a strong positive correlation with the antibody's, affinity for its target. As higher affinity antibodies are able to bind their target faster and remain bound to the target longer, they generally are effective at lower concentrations. Monoclonal antibodies (mAbs) can be generated by a variety of techniques including hybridoma technology, which involves fusing antibody-producing B-cells with cancerous myeloma cells. (Kohler & Milstein (1975) Nature 256: 52-53). The resulting hybridoma cell is an "immortalized" cell that secretes an antibody of single specificity (monoclonal antibody). An alternative method for monoclonal antibody generation is termed the Selected Lymphocyte Antibody Method (SLAM), described in U.S. Patent No. 5,627,052 entitled "Methods for the production of proteins with a desired function" and in "A novel strategy for generating monoclonal antibodies from single, isolated lymphocytes producing antibodies of defined specificities" Babcook et al. (1996) Proc Natl Acad Sci U S A. 93:7843-8. This method allows one to first identify B-cells that are making antibodies with desired characteristics such as binding specificity, function, and optimal kinetics. The selected B-cells are then isolated and the antibody variable domain genes encoding the binding portion of the antibody are rescued by molecular techniques, such as the polymerase chain reaction (PCR), and cloned into expression vectors having an antibody constant region domain. The expression vectors are then transfected into cells, such as CHO or NS0 cells, to produce secreted antibody molecules. This process allows the virtually unlimited production of the desired antibody. Other methodologies such as phage display or the viral immortalization (including Epstein Barr Vins (EBV)) of B-cells have also been used successfully in the generation of monoclonal antibodies.

[0003] During the SLAM process, millions of B-cells are screened for the specificity, function and kinetic characteristics of the antibodies they produce. For a selected target, thousands of target-specific B-cells might be identified. Examples of known methods of screening include screening for antibodies which (1) bind to natively expressed antigen on the cell surface, (2) bind cell lines and induce apoptosis, (3) bind cells and either induce proliferation or block proliferation, and (4) bind to ligand and block binding of the ligand to receptor and vice versa. To determine formal affinity measurements of large panels of antibodies is a time-consuming, expensive, and inefficient process for identifying optimal antibodies. Accordingly, what is needed is a simple, rapid, and accurate mechanism for screening antibodies to determine their relative binding characteristics.

[0004] Watkins *et al.* have described the determination of the relative affinities of antibody fragments expressed in *Escherichia coli* by enzyme-linked immunosorbent assay (ELISA). (Watkins et al. (1997) Anal Biochem 253:37-45). U.S. Patent No. 5,800,815 relates to compositions and methods for treating inflammation and other pathological conditions using novel blocking P-selectin antibodies.

Summary of the Invention

[0005] The present invention includes a method for kinetically ranking each antibody in an antibody set by preparing a set of diluted antigen preparations and thereafter measuring the binding of each antibody in the set of antibodies to the diluted antigen preparations. During the binding assay, antibody and antigen are combined, the binding reaction is allowed to go to equilibrium, and after equilibrium is achieved, each antibody is preferably labeled with a detectable label, and the strength each antibodies label signal is related to the binding affinity of the antibody for a diluted antigen preparation. A comparison of each antibody's relative affinity for each particular antigen preparation can thereby be performed. Antibodies that bind to the more dilute antigen preparations have a higher relative affinity for the antigen, whereas antibodies that only bind to more concentrated antigen preparations have a relatively lower binding affinity to the antigen.

[0006] One embodiment of the invention is a method of determining the relative binding affinities of antibodies to an antigen. This embodiment includes: determining the limiting concentration of an antigen that maximizes the detectable

range of antibody affinities for said antigen; providing a set of antibody solutions, wherein the concentration of antibodies in each antibody solution is known; incubating the limiting concentration of antigen with each member of the set of antibody solutions; measuring the relative degree to which the antibodies in each member of the set of antibody solutions binds to the limiting concentration of antigen; and ranking the antibodies in each member of said set of antibody solutions based on their binding affinity to said different dilutions in order to determine the relative binding affinities of each of said different antibodies to said antigen.

[0007] Another embodiment of the invention is a method of determining the relative binding affinities of a set of antibodies to an antigen which includes: providing an antigen solution comprising a first dilution of the antigen, wherein said antigen is labeled with a first label, and a second dilution of the antigen, wherein said antigen is labeled with a second label; providing a set of antibody solutions wherein each antibody solution has substantially the same relative concentration of said antibody; measuring the binding of the antibody to the first dilution of said antigen; measuring the binding of the antibody to the second dilution of the antigen; and ranking the antibody in the set of antibody solutions which bind to a relatively more dilute antigen preparation as having a higher relative affinity and the antibody that substantially only bind to relatively more concentrated antigen preparations as having a relatively lower binding affinity to the antigen.

[0008] Still another embodiment is a method of determining the relative binding affinities of a set of molecules to a molecule binding partner. This method includes: determining the limiting concentration of a molecule binding partner that maximizes the detectable range of molecule affinities for said molecule binding partner; providing a set of molecule solutions, wherein the concentration of molecules in each molecule solution is known; incubating the limiting concentration of molecule binding partners with each member of the set of molecule solutions; measuring the relative degree to which the molecules in each member of the set of molecule solutions binds to the limiting concentration of molecule binding partners; and ranking the molecules in each member of said set of molecule solutions based on their binding affinity to said different dilutions in order to determine the relative binding affinities of each of said different molecules to said molecule binding partners.

Detailed Description

[0009] Embodiments of the invention relate to methods for rapidly determining the differential binding properties within a set of antibodies. Accordingly, rapid identification of optimal antibodies for binding to a target can be determined. Any set of antibodies raised against a particular target antigen may bind to a variety of epitopes on the antigen. In addition, antibodies might bind to one particular epitope with varying affinities. Embodiments of the invention provide methods for determining how strongly or weakly an antibody binds to a particular epitope in relation to other antibodies generated against the antigen.

[0010] One embodiment of the invention is provided by preparing a set of diluted antigen preparations and thereafter measuring the binding of each antibody in a set of antibodies to the diluted antigen preparations. A comparison of each antibody's relative affinity for a particular concentration of antigen can thereby be performed. Accordingly, this method discerns which antibodies bind to the more dilute concentration of antigen, or to the more concentrated antigen preparations, as part of a comparative assay for the relative affinity of each antibody in a set.

[0011] Another embodiment of the invention is provided by preparing a set of diluted antibody preparations and thereafter measuring the binding of an antigen to each of the diluted antibody preparations. A comparison of each antibody's relative affinity for a particular antigen can thereby be performed. Accordingly, this method discerns whether a particular concentration of an antigen binds to the more dilute concentration of antibody preparations, or to the more concentrated antibody preparations, as part of a comparative assay for the relative affinity of each antibody in a set.

[0012] Although a process is disclosed in which an antibody's relative affinity can be determined, a similar protocol can be foreseen for the identification of high affinity antibody fragments, protein ligands, small molecules or any other molecule with affinity toward another. Thus, the invention is not limited to only analyzing binding of antibodies to antigens.

[0013] One embodiment of the invention provides a method for analyzing the kinetic properties of antibodies to allow ranking and selection of antibodies with desired kinetic properties. Affinity, as defined herein, reflects the relationship between the rate at which one molecule binds to another molecule (association constant, K_{on}) and the rate at which dissociation of the complex occurs (dissociation constant, K_{off}). When an antibody and target are combined under suitable conditions, the antibody will associate with the target antigen. At some point the ratio of the amount of antibody binding and releasing from its target reaches an equilibrium. This equilibrium is referred to as the "affinity constant" or just "affinity".

[0014] When binding reactions having identical concentrations of antibody and target molecule are compared, reactions containing higher affinity antibodies will have more antibodies bound to the target at equilibrium than reactions containing antibodies of lower affinity.

[0015] In assays where the binding of one molecule to another is measured by the formation of complexes which generate a signal, the amount of signal is proportional to the concentrations of the molecules as well as to the affinity of the interaction. For purposes of the present disclosure, assays are employed to measure formation of complexes

between antibodies and their targets (on antigens), where signals being measured in such assays may be proportional to the concentrations of antibody or antibodies, concentration of target antigen, and the affinity of the interaction. Suitable assay methods for measuring formation of antibody-target complexes include enzyme linked immunosorbent assays (ELISA), fluorescence-linked immunosorbent assays (including Luminex systems, fluorometric microvolume assay technology (FMAT) and fluorescence activated cell sorting (FACS) systems), radioisotopic assay (RIA) as well as others which can be chosen by one of skill in the art.

[0016] Another aspect of the present invention includes methods for kinetically ranking antibodies by affinity based on the signal strength of an assay such as an assay listed above, when the target or antigen is provided at limiting concentrations. Antibody and antigen are combined, the binding reaction is allowed to go to equilibrium, and after equilibrium is achieved, an assay is performed to determine the amount of antibody bound to the target or antigen. According to one aspect of the present invention, the amount of bound antibody detected by the assay is directly proportional to the affinity of the antibody for the target or antigen. At very low concentrations of antigen, some antibodies of low affinity will not generate a detectable signal due to an insufficient amount of bound antibody. At the same very concentrations of antigen, antibodies of moderate affinity will generate low signals, and antibodies with high affinity will generate strong signals.

[0017] During a standard assay using moderate to high concentrations of target, a collection of different antibodies having different affinities for the same target antigen may generate signals of equal or similar intensity. However, as the amount of antigen is diluted, it becomes possible to discern differences in affinity among the antibodies. Using limiting concentrations of target antigen in the assay in accordance with the teachings of the present disclosure, it is possible to establish a kinetic ranking of a collection of antibodies against the same target antigen.

[0018] Under conditions of limiting amounts of antigen, a collection of antibodies against the same antigen will give a range of signals from high to low or no signal, even though in the original assay using high to moderate levels of antigen, some of these antibodies may have produced signals of similar apparent strength. Antibodies can thus be affinity-ranked by their signal intensity in a limiting antigen assay carried out in accordance with the teachings of the present disclosure.

[0019] Another aspect of the invention is a method of determining antibodies with higher affinities than currently known and characterized antibodies. This method involves using the characterized antibodies as kinetic standards. A plurality of test antibodies are then measured against the kinetic standard antibodies to determine those antibodies that bind to more dilute antigen preparations than to the standard antibodies. A plurality of test antibodies is then measured against the kinetic standard antibody to determine those antibodies which have more antibody bound to a given dilute preparation of antigen. This allows the rapid discovery of antibodies that have a higher affinity for antigen in comparison to the kinetic standard antibodies.

[0020] In one preferred embodiment, an ELISA is used in a limiting antigen assay in accordance with the present disclosure.

[0021] It has been empirically determined that supernatants of cultured B-cells generally secrete antibodies in a concentration range from 20 ng/ml to 800 ng/ml. Because there is often a limited amount of supernatant from these cultures, B-cell culture supernatants are typically diluted 10-fold for most assays, giving a working concentration of from 2 ng/ml- 80 ng/ml for use in affinity determination assays. In one aspect of the invention, the appropriate concentration of target antigen used to coat ELISA plates was determined by using a reference solution from a monoclonal antibody at a concentration of 100 ng/ml. This number could change depending on the concentration range of test antibodies and the affinity of the reference antibody, such that the concentration of target antigen required to give half-maximal signal in a ELISA-based measurement of antibody/antigen binding can be empirically determined. This determination is discussed in more detail below.

[0022] Antigen at an empirically determined optimal coating concentration was used in affinity measurement assays to discern the antibodies produced by various B-cell cultures that gave an ELISA value higher than a reference monoclonal antibody. According to the methods of the present invention, the only way to obtain a higher signal than that obtained using the reference antibody is if (1) the antibody is of higher affinity than the reference antibody or (2) the antibody has the same affinity but is present in a higher concentration than the reference monoclonal antibody. As disclosed previously, antibodies in B-cell culture supernatants are usually at concentrations of between 20-800 ng/ml and are diluted to a working concentration of between 2 to 80 ng/ml. In one embodiment, test antibodies at a concentration of between 2 to 80 ng/ml are used in assays having a reference antibody concentration of 100 ng/ml. The signal achieved from the test antibodies is compared to that of the 100 ng/ml reference antibody. If antibodies within the test group are found to have a higher signal, then the antibody is assumed to be of a higher affinity than the reference antibody.

[0023] In another embodiment, antibodies generated from hybridomas were ranked using a limiting kinetic antigen assay in an ELISA-based protocol. The binding affinities for these antibodies was confirmed by quantifying and kinetically ranked the antibodies using a Biacore system. As is known, the Biacore system gives formal kinetic values for the binding coefficient between each antibody and the antigen. It was determined that the kinetic ranking of antibodies using the limiting antigen assay as taught by the present disclosure closely correlated with the formal kinetic values for these antibodies as determined by the Biacore method, as shown in Table 5 (Example 5).

[0024] Briefly, the Biacore technology uses surface plasmon resonance (SPR) to measure the decay of antibody from antigen at various concentrations of antigen and at a known concentration of antibody. For example, chips are loaded with antibody, washed, and the chip is exposed to a solution of antigen to load the antibodies with antigen. The chip is then continually washed with a solution without antigen. An initial increase in SPR is seen as the antibody and antigen complex forms, followed by decay as the antigen-antibody complex dissociates. This decay in signal is directly proportional to antibody affinity. Similarly this method could run the reverse assay with limited concentrations of antibody coated on the chip.

[0025] Using the Luminex (MiraiBio, Inc., Alameda, CA) technology antibodies are assayed for how they bound a plurality of different antigen coated beads. In this assay each bead set is preferably coated with a different concentration of antigen. As the Luminex reader has the ability to multiplex all the beads sets, the bead sets are combined and antibody binding to each of the different bead sets are determined. The behavior of antibodies on the differentially coated beads can then be tracked. Once normalized for antibody concentration, then antibodies which maintain a high degree of binding as one moves from non-antigen limiting concentrations to limited antigen concentrations correlate well to high affinity. Advantageously, these differential shifts can be used to relatively rank antibody affinities. For example, samples with smaller shifts correspond to higher affinity antibodies and antibodies with larger shifts correspond to lower affinity antibodies.

Table 1

Comparison of Affinity Rankings Between Biacore and Luminex Methods				
BiaCore Affinity Measurements				Luminex rank
ka (M ⁻¹ s ⁻¹)	kd (s ⁻¹)	Biacore Med-res KD (nM)	Rank	
9.9 x 10 ⁵	9.3 x 10 ⁻³	9.4	1	1
2.7 x 10 ⁵	4.2 x 10 ⁻³	16	2	14
3.1 x 10 ⁵	5.6 x 10 ⁻³	18	3	57
8.2 x 10 ⁵	2.7 x 10 ⁻²	33	4	83
1.4 x 10 ⁶	6.2 x 10 ⁻²	42	5	116
2.9 x 10 ⁵	1.6 x 10 ⁻²	54	6	123

[0026] In another embodiment of the invention, a series of limited concentrations of the antibody being tested are compared to a standard solution of antibody. Such a method using limiting concentrations of antibody would appear to be a "reverse" of the method using limiting antigen concentrations, but it provides a similar mechanism for rapidly screening a set of antibodies to determine each antibody's relative affinity for the target antigen. Other plates that are, or can be, chemically modified to allow covalent or passive coating can also be used. One of skill in the relevant art can devise further modifications of the methods presented herein to carry out an assay using limiting antibody dilution to screen and kinetically rank test antibodies.

Determining optimal bound antigen concentration

[0027] Embodiments of the limiting antigen assay method are practiced using a method by which antigen is bound or attached to a stationary surface prior to subsequent manipulations. The surface is preferably part of a vessel in which subsequent manipulations may occur; more preferably, the surface is in a flask or test tube, even more preferably the surface is in the well of a microtiter plate such as a 96-well plate, a 384-well plate, or a 864-well plate. Alternately, the surface to which antigen is bound may be part of a surface such as a slide or bead, where the surface with bound antigen may be manipulated in subsequent antibody binding and detection steps. Preferably, the process by which the antigen is bound or attached to the surface does not interfere with the ability of antibodies to recognize and bind to the target antigen.

[0028] In one embodiment, the surface is coated with streptavidin and the antigen is biotinylated. In a particularly preferred embodiment, the plate is a microtiter plate, preferably a 96-well plate, having streptavidin coating at least one surface in each well, and the antigen is biotinylated. Most preferably, the plate is Sigma SA 96-well plate and the antigen is biotinylated with Pierce EZ-link Sulpho-NHS Biotin (Sigma-Aldrich Canada, Oakville Ontario, CANADA). Alternative methods of biotinylation which attach the biotin molecule to other moieties can also be used.

[0029] In the unlikely event that an antigen cannot be biotinylated, alternative surfaces to which antigen can be bound can be substituted. For example, the Costar® Universal-BIND™ surface, which is intended to covalently immobilize

biomolecules via an abstractable hydrogen using UV illumination resulting in a carbon-carbon bond. (Coming Life Sciences, Coming, NY). Plates, for example, Costar® Universal-BIND™ 96-well plates, may be used. One of skill in the art can modify subsequent manipulations in the event that the use of alternate surfaces such as Costar® Universal-BIND™ increases the time of the assay and/or requires the use of more antigen.

[0030] In one embodiment of the present invention, a "checkerboard" assay design is used to find optimal concentration of bound antigen. One example is shown below in Table 2. The following description includes a disclosure of the steps to determine the optimal coating concentration of biotinylated antigen using 96-well plates coated with streptavidin. This disclosure is intended merely to illustrate one way to practice various aspects of the present invention. The scope of the present invention is not limited to the methods of the assay described above and below, as one of skill in the art can practice the methods of the present invention using a wide variety of materials and manipulations. Methods including but not limited to; expression of antigen on cells (transient or stable), using phage which express different copy number of antigen per phage.

Antigen dilution and distribution.

[0031] An antigen to be tested is selected. Such an antigen may be, for example, any antigen that might provide a therapeutic target by antibodies. For example, tumor markers, cell surface molecules, Lymphokines, chemokines, pathogen associated proteins, and immunomodulators are non-limiting examples of such antigens.

[0032] A solution of antigen at an initial concentration, preferably about 1 ug/ml, is diluted in a series of stepwise dilutions. Diluted samples are then placed on surfaces such as in the wells of a microtiter plate, and replicates of each sample are also distributed on surfaces. Antigen solutions may contain blocking agents if desired. In a preferred embodiment, serial dilutions of antigen are distributed across the columns of a 96-well plate. Specifically, a different antigen dilution is placed in each column, with replicate samples in each row of the column. In a 96-well plate, replicates of each dilution are placed in rows A-H under each column. Although the standard dilutions vary from antigen to antigen, the typical dilution series starts at 1 ug/ml and is serially diluted 1:2 to a final concentration of about 900 pg/ml.

[0033] In one embodiment, biotinylated antigen is diluted from a concentration of 1 µg/ml to 900 pg/ml horizontally across a 96 well plate. While a preferred blocking buffer is a phosphate buffered saline (PBS) Milk solution, others buffers such as bovine serum albumin (BSA) diluted in PBS can be substituted. In another embodiment, biotinylated antigen is diluted from a concentration of 1 ug/ml to 900 pg/ml in 1% skim milk/ 1X PBS pH 7.4, and pipetted into the wells of columns 1 to 11 of a Sigma SA (streptavidin) microtiter plate, with 8 replicates of each dilution placed in rows A-H of each column. Column 12 is left blank, serving as the "antibody-only" control. The final volume in each well is 50 ul. Antigen is incubated on the surface (e.g., in the wells of the plate) for a suitable amount of time for the antigen to become attached to the surface; incubation time, temperature, and other conditions can be determined from manufacturer's instructions and/or standard protocols for the surface being used. After incubation, excess antigen solution is removed. If needed, plates are then blocked with a suitable blocking solution containing, e.g., skim milk, powdered milk, BSA, gelatin, detergent, or other suitable blocking agents, to prevent nonspecific binding during subsequent steps.

[0034] Plates with biotinylated antigen are then incubated for a suitable amount of time for antigen to bind or attach to the surface. Biotinylated antigen in a Sigma SA plate is incubated at room temperature for 30 minutes. Excess biotinylated antigen solution is then removed from the plate. In this embodiment, blocking is not necessary because Sigma SA plates are pre-blocked.

[0035] In another embodiment using Costar® Universal-BIND™ plates, antigen is passively adsorbed overnight at 4 degrees C in 1X PBS pH 7.4, 0.05% azide. Generally, if Costar® Universal-BIND™ plates are used, the initial concentration of antigen is a somewhat higher concentration, preferably 2-4 ug/ml. The next morning, excess antigen solution is removed from Costar® Universal-BIND™ plate or plates, preferably by "flicking", and each plate is exposed to UV light at 365 nm for four (4) minutes. Each plate is then blocked with 1% skim milk / 1X PBS pH 7.4 at 100 ul of blocking solution per well, for 30 minutes.

[0036] After incubation with antigen and removal of excess antigen solution, and blocking, if necessary, plates are washed four times (4X) with tap water. Plates may be washed by hand, or a microplate washer or other suitable washing tool may be used.

Reference antibody dilution and distribution.

[0037] A reference antibody that recognizes and binds to the antigen is then added. The reference antibody is preferably a monoclonal antibody, but can alternatively be polyclonal antibodies, natural ligands or soluble receptors, antibody fragments or small molecules.

[0038] A solution of reference antibody, also known as anti-antigen antibody, at an initial concentration, preferably about 1 ug/ml, is diluted in a series of stepwise dilutions. Diluted samples are placed on surfaces such as in the wells of a microtiter plate, and replicates of each sample are also distributed on surfaces. Serial dilutions of reference antibody

are distributed across the rows of a 96-well plate. Specifically, each reference antibody dilution is placed in a row, with replicate samples placed in each column of the row. In a 96-well plate, a different dilution of reference antibody is placed in each row, with replicates of each dilution placed in each column across each row starting at an initial concentration of about 1 ug/ml progressively and diluted 1:2 seven times for a series of seven wells. An ending concentration of about 30 ng/ml is used as the standard solution series. Solutions of reference antibody are incubated with bound antigen under suitable conditions determined by the materials and reagents being used, preferably about 24 hours at room temperature. One of skill in the art can determine whether incubation for longer or shorter times, or at higher or lower temperatures would be suitable for a particular embodiment.

Optional Step: Incubation with shaking.

[0039] If desired, the plate may be tightly wrapped and incubation of the reference antibody with bound antigen may be carried out with shaking to promote mixing and more efficient binding. Plates containing reference antibody and bound antigen may be incubated overnight with shaking, for example as provided by a Lab Line Microplate Shaker at setting 3.

Add detection antibody.

[0040] Plates are washed to remove unbound reference antibody, preferably about five times (5X) with water. Next, a labeled detection antibody that recognizes and binds to the reference antibody is added, and the solution is incubated to permit binding of the detection antibody to the reference antibody. The detection antibody may be polyclonal or monoclonal. The detection antibody may be labeled in any manner that allows detection of antibody bound to the reference antibody. The label may be an enzymatic label such as alkaline phosphatase or horseradish peroxidase (HRP), or a non-enzymatic label such as biotin or digoxigenin, or may be a radioactive label such as ^{32}P , ^3H , or ^{14}C , or may be any other label suitable for the assay based on reagents, materials, and detection methods available.

[0041] Following labeling, 50 ul of goat anti-Human IgG Fc HRP polyclonal antibody (Pierce Chemical Co, Rockford IL, catalog number 31416) at a concentration of 0.5 ug/ml in 1 % skim milk, 1X PBS pH 7.4 is added to each well of a microtiter plate. The plate is then incubated for 1 hr at room temperature.

[0042] Excess solution containing detection antibody is removed, and plates are washed with water repeatedly, preferably at least five times, in order to remove all unbound detection antibody.

Measurement of bound detection antibody.

[0043] The amount of detection antibody bound to reference antibody is determined by using the appropriate method for measuring and quantifying the amount of label present. Depending on the label chosen, methods of measuring may include measuring enzymatic activity against added substrate, measuring binding to a detectable binding partner (e.g., for biotin) scintillation counting to measure radioactivity, or any other suitable method to be determined by one of skill in the relevant art.

[0044] In the embodiment described above using goat anti-Human IgG Fc HRP polyclonal antibody as the detection antibody, 50 ul of the chromogenic HRP substrate tetramethylbenzidine (TMB) is added to each well. The substrate solution is incubated for about 30 minutes at room temperature. The HRP/TMB reaction is stopped by adding 50 ul of 1M phosphoric acid to each well.

Quantification.

[0045] The amount of bound label is then quantified by the appropriate method, such as spectrophotometric measurement of formation of reaction products or binding complexes, or calculation of the amount of radioactive label detected. Under the conditions disclosed here, the amount of label measured in this step is a measure of the amount of labeled detection antibody bound to the reference antibody.

[0046] In the embodiment described above using goat anti-Human IgG Fc HRP polyclonal antibody and TMB substrate, the amount of detection antibody bound to reference antibody is quantified by reading the absorbance (optical density or "OD") at 450 nm of each well of the plate.

Data analysis to determine optimal antigen concentration.

[0047] A known reference antibody concentration is chosen, and the results from wells having the chosen antibody concentration and different amounts of antigen are examined. The antigen concentration that produces the desired signal strength, or standard signal, is chosen as the optimal antigen concentration for subsequent experiments. The

standard signal may be empirically determined according to the conditions and materials used in a particular embodiment, because the standard signal will serve as a reference point for comparing signals from other reactions. For a detection method that produces a chromogenic product, a desirable standard signal is one that falls within the most dynamic region of the ELISA reader or other detector and may be an optical density (OD) of between about 0.4 and 1.6 OD units and for this system preferably about 1.0 OD units, although it is possible to achieve signals ranging from 0.2 to greater than 3.0 OD units. Any OD value may be chosen as the standard signal, although an OD value of about 1.0 OD units permits a accurate measurement of a range of test signals above and below 1.0 OD units, and further permits easy comparison with other test signals and reference signals. The concentration of antigen identified as the concentration that produces the standard signal will be used in subsequent experiments to screen and kinetically rank antibodies.

[0048] In a preferred embodiment using a 96-well plate, a reference antibody concentration of 100 ng/ml is chosen. It is possible, depending on the sensitivity and antibody concentrations employed in the system, to use other reference antibody concentrations. The signals from the detection antibody reaction in the wells in all columns of the row containing 100 ng/ml antibody are then examined to find the antigen concentration that produces an OD value of about 1.0. In the preferred embodiment described above using goat anti-Human IgG Fc HRP polyclonal antibody and TMB substrate, the wells in the row containing 100 ng/ml antibody are examined to determine which antigen concentration produces a reaction which, when absorbance is measured at 450 nm, has an OD value of about 1.0. This concentration of antigen will then be used for the subsequent experiments to screen and kinetically rank antibodies. A similar approach for identifying optimal antigen densities was used for the Luminex bead based system.

Screening antibodies using limiting antigen concentrations

Coat surfaces at optimized antigen concentration

[0049] The surface or surfaces being used to carry out antibody screening are coated with antigen at the optimal concentration as previously determined. In a preferred embodiment, the surfaces are wells of a 96-well streptavidin plate such as a Sigma SA plate, and biotinylated antigen at optimal concentration is added to the wells. In a more preferred embodiment, 50 µl of antigen in a solution of 1% skim milk, 1X PBS pH 7.4, and plates are incubated for 30 minutes. In another preferred embodiment, unmodified antigen is added to Costar® Universal-BIND™ plates, and incubation and UV-mediated antigen binding are carried out according to manufacturer's instructions and/or standard protocols, as described above.

[0050] After incubation with antigen solution for a suitable amount of time, plates are washed to remove unbound antigen, preferably at least four times (4X).

Addition of test antibodies to be screened and ranked

[0051] Antibodies to be screened and ranked by the limiting antigen assay are called test antibodies. Test antibodies may be recovered from the solution surrounding antibody-producing cells. Preferably, test antibodies are recovered from the media of antibody-producing B cell cultures, hybridoma supernatants, antibody or antibody fragments expressed from any type of cell, more preferably from the supernatant of B cell cultures. Solutions containing test antibodies, for example B cell culture supernatants, generally do not require additional processing; however, additional steps to concentrate, isolate, or purify test antibodies would also be compatible with the disclosed methods.

[0052] Each solution containing test antibodies is diluted to bring the concentration within a desirable range and samples are added to a surface having attached antigen. Typically, a desirable concentration range for test antibodies has a maximum concentration lower than the concentration of reference antibody used to select the optimal antigen concentration as described above. One aspect of the present invention provides that a test antibody would produce a signal higher than that of the reference antibody for the same antigen concentration if the test antibody (a) has a higher affinity for the antigen, or (b) has a similar affinity but is present in higher concentration than the reference antigen. Thus, when test antibodies are used at concentrations lower than the concentration of the reference antibody used to select the antigen concentration used in the screening assay, only a test antibody having higher affinity for the antigen would produce a higher signal than the reference antibody signal.

[0053] In one embodiment in which a reference antibody concentration of 100 ng/ml is used to select the optimal antigen concentration (as described above), B cell culture supernatants having an empirically determined test antibody concentration range of between about 20 ng/ml to 800 ng/ml are typically diluted ten-fold to produce a working assay test antibody concentration of between about 2 ng/ml to 80 ng/ml. Preferably, at least two duplicate samples of each diluted B cell culture supernatant are tested. Preferably, the diluted B cell culture supernatants are added to wells of a microtiter plate, where the wells are coated with antigen at an optimal concentration previously determined using antigen and a reference antibody.

[0054] A positive control should be included as part of the screening, wherein the reference antibody used to optimize

the assay by determining optimal antigen concentration is diluted and reacted with the antigen. The positive control provides a set of measurements useful both as an internal control and also to compare with previous optimization results in order to confirm, assure, and demonstrate that results from a screening of test antibodies are comparable with the expected results of the positive control, and are consistent with previous optimization results.

[0055] In one embodiment, each B cell culture supernatant to be tested is diluted 1:10 in 1% skim milk / 1X PBS pH 7.4 / .05% azide, and 50 ul is added to each of two antigen-coated wells of a 96-well plate, such that 48 different samples are present in each 96-well plate. A positive control comprising a dilution series of the reference antibody is preferably added to wells of about one-half a 96-well plate, to provide confirmation and to demonstrate that results of the screening of test antibodies in B cell culture supernatants run in parallel with the positive control are internally consistent and also consistent with previous optimization results.

[0056] Test antibodies are incubated with antigen under suitable conditions. Reference antibodies used as positive controls are incubated in parallel under the same conditions. In one preferred embodiment, plates are wrapped tightly, for example with plastic wrap or paraffin film, and incubated with shaking for 24 hours at room temperature.

Add detection antibody to test antibodies

[0057] Plates are washed to remove unbound test antibodies, preferably about five times (5X) with water. Next, a labeled detection antibody that recognizes and binds to the test antibody is added, and the solution is incubated to permit binding of the detection antibody to the test antibody. Detection antibody is also added to the positive control, to confirm the interaction between the reference antibody and detection antibody. The detection antibody may be polyclonal or monoclonal. The detection antibody may be labeled in any matter that allows detection of antibody bound to the reference antibody. The label may be an enzymatic label such as alkaline phosphatase or horseradish peroxidase (HRP), or a non-enzymatic label such as biotin or digoxigenin, or a radioactive label such as ^{32}P , ^3H , or ^{14}C , or fluorescence, or it may be any other label suitable for the assay based on reagents, materials, and detection methods available.

[0058] In one embodiment, using human test antibodies, 50 ul of goat anti-Human IgG Fc HRP polyclonal antibody (Pierce Chemical Co, Rockford IL, catalog number 31416) at a concentration of 0.5 ug/ml in 1% skim milk, 1X PBS pH 7.4 is added to each well of microtiter plates containing test antibodies and reference antibodies (as a positive control). The plate is then incubated for 1 hr at room temperature.

[0059] Excess solution containing detection antibody is removed, and plates are washed with water repeatedly, preferably at least five times, in order to remove all unbound detection antibody.

Measurement of bound detection antibody.

[0060] The amount of detection antibody bound to test antibody (and bound to reference antibody of the control) is determined by using the appropriate method for measuring and quantifying the amount of label present. Depending on the label chosen, methods of measuring may include measuring enzymatic activity against added substrate, measuring binding to a detectable binding partner (e.g., for biotin) scintillation counting to measure radioactivity, or any other suitable method to be determined by one of skill in the relevant art.

[0061] In the method described above, using goat anti-Human IgG Fc HRP polyclonal antibody as the detection antibody, 50 ul of the chromogenic HRP substrate tetramethylbenzidine (TMB) is added to each well. The antibody-substrate solution is incubated for about 30 minutes at room temperature. The HRP/TMB reaction is stopped by adding 50 ul of 1M phosphoric acid to each well.

Quantification.

[0062] The amount of bound label is then quantified by the appropriate method, such as the spectrophotometric measurement of formation of reaction products or binding complexes, or calculation of the amount of radioactive label detected. In accordance with one aspect of the present invention, the amount of label provides a measure of the amount of labeled detection antibody bound to the test antibody (or, in the positive control, bound to the reference antibody). In accordance with another aspect of the present invention, the amount of label provides a measure of the amount of test antibody bound to antigen. Thus, detecting and quantifying the amount of label provides a means of measuring the binding of test antibody to the test antigen. By comparing the standard signal with the signal that quantifies the amount of test antibody bound to antigen, it is possible to identify test antibodies with higher affinities by searching for test antibodies which give a higher signal than the reference.

[0063] In the method described above using goat anti-Human IgG Fc HRP polyclonal antibody and TMB substrate, the amount of detection antibody bound to test antibody (and reference antibody in the positive control) is quantified by reading the absorbance (optical density, OD) at 450 nm of each well of each plate.

Data analysis to identify and rank antibodies of interest.

[0064] The results from each test antibody are averaged and the standard range is determined. In a preferred embodiment wherein two samples of each test antibody are assayed using a HRP-labeled detection antibody, OD values at 450 nm are averaged and the standard deviation is calculated. The average OD values of test antibodies are compared against the OD value of the standard signal. Values from the positive control assays are also calculated and examined for reliability of the assay.

[0065] Test antibodies are kinetically ranked by considering the average OD value and the range of the OD's between replicates. The average OD value provides a measure of the affinity of the test antibody for the antigen, where affinity is determined by comparison with the standard signal, or the OD value of the reference antibody in the positive control. The range provides a measure of reliability of the assay, where a narrow range indicates that the OD values are likely to be accurate measurements of the amount of test antibody bound to the antigen, and a wide range indicates that the OD values may not be accurate measurements of binding. Acceptable standard deviations are typically OD's of between 5-15% of each other. Test antibodies giving the highest OD values, where the standard deviation of the average value is low, are given the highest kinetic ranking.

[0066] In one embodiment, wherein the standard signal is 1.0 OD units, any test antibody with both an average OD of greater than 1.0 OD units, and an acceptably low standard deviation, is considered to have a higher affinity for the antigen than the affinity of the reference antibody.

[0067] In another embodiment, Luminex based assays using differentially antigen coated beads were used. In this assay antibodies were ranked based on how they bound antigen at higher then at lower antigen densities.

Example 1Determination of optimal antigen concentration*Antigen Preparation*

[0068] Parathyroid hormone (PTH) was biotinylated using Pierce EZ-Link Sulpho-NHS biotin according the manufacturer's directions (Pierce EZ-link Sulpho-NHS Biotin, (Pierce Chemical Co., Rockford, IL, Catalogue number 21217). When the antigen could not be biotinylated, Costar UV plates were substituted. The use of Costar UV plates increased the time of the assay and generally required the use of considerably more antigen.

Checkerboard ELISA

[0069] An assay laid out in a "checkerboard" arrangement was carried out as described below to determine optimal coating concentration of the antigen. The assay was performed using streptavidin-coated 96-well plates (Sigma SA microtiter plates, Sigma-Aldrich Chemicals, St Louis MO, Catalogue number-M5432) as follows.

[0070] The parathyroid hormone (PTH) antigen was biotinylated using Pierce EZ-link Sulpho-NHS biotin ((Pierce Chemical Co, Rockford IL, catalog number 21217) according to manufacturer's instructions. Biotinylated antigen diluted in 1% skim milk/ 1X PBS pH 7.4 in a series of stepwise dilutions from a beginning concentration of 500 ng/ml to a final concentration of 0.5ng/ml. Diluted biotinylated antigen was distributed horizontally across a 96-well Sigma SA microtiter plate (Sigma Aldrich Chemicals, catalogue M-5432), placing 50 ul of each dilution in wells of each of columns 1 through 11, with replicates in each well of rows A-H under each column. No antigen was added to column 12. The plate was incubated at room temperature for 30 minutes. No blocking step was performed because Sigma SA plates are pre-blocked.

[0071] The plate was washed four times with tap water. Plates were washed by hand, or using a microplate washer when available.

[0072] An anti-PTH antibody with known affinity was used as a reference antibody. Anti-PTH antibody 15g2 was diluted 1% skim milk / 1X PBS pH 7.4 / 0.05% to final initial dilution of 1 ug/ml was serially diluted 1:2, 7 wells to an ending concentration 15 ng/ml and 50 ul of each dilution was distributed in each well of row A to row G, with replicates in each well of columns 1-12. No antibody was added to row H. Plates containing the antigen and reference antibody were incubated at room temperature for approximately 24 hours.

[0073] The plate was wrapped tightly ("air tight") with plastic wrap or paraffin film, and incubated overnight with shaking using a Lab Line Titer Plate Shaker at setting 3.

[0074] The plates were washed five times (5X) with water to remove unbound reference antibody. Bound reference antibody was detected by adding fifty microliters (50 ul) of 0.5 ug/ml goat anti-Human IgG Fc HRP polyclonal antibody (Pierce Chemical Co, Rockford IL, catalog number 31416) in 1% skim milk / 1X PBS pH 7.4 to each well and incubating the plate 1 hr at room temperature. (Gt anti-Human Fc HRP- Pierce catalogue number-31416).

[0075] The plate was washed at least five times (5X) with water to remove unbound goat anti-Human IgG Fc HRP

polyclonal antibody

[0076] Fifty microliters (50 μ l) of the HRP substrate TMB (Kirkegaard & Perry Laboratories, Inc, Gaithersburg, MD) was added to each well and the plate was incubated for one-half hour at room temperature. The HRP-TMB reaction was stopped by adding 50 μ l of 1M phosphoric acid to each well. Optical density (absorbance) at 450 nm was measured for each well of the plate.

Data analysis

[0077] Table 2 shows the results from the reference assay using PTH as the antigen and 15g2 anti-PTH as the reference antibody. OD measurements from the row of samples corresponding to the reference antibody concentration of 100 ng/ml were examined to find the antigen concentration that gives an OD of approximately 1.0. This concentration was determined to be approximately 15 ng/ml PTH. This concentration of antigen was considered the optimal antigen concentration and will be used for the subsequent experiments.

Table 2

Optical Density Measurements of Test Antibodies Bound to
Various Concentrations of PTH

		PTH Concentration (ng/mL)											
		500.00	250.00	125.00	62.50	31.25	15.63	7.81	3.91	1.95	0.98	0.49	0.00
Reference antibody concentration (ng/mL)	1000	3.218	3.273	3.075	3.103	2.521	1.910	1.269	0.885	0.438	0.329	0.256	0.086
	500	3.199	3.133	3.144	3.068	2.608	1.928	1.283	0.708	0.424	0.293	0.224	0.062
	250	3.130	3.274	3.208	2.945	2.393	1.634	1.182	0.543	0.295	0.201	0.156	0.055
	125	3.190	3.194	3.177	2.733	2.116	1.251	0.863	0.444	0.489	0.178	0.147	0.067
	62.5	3.187	3.262	2.952	2.137	1.678	0.946	0.515	0.295	0.179	0.126	0.103	0.055
	31.3	3.148	3.001	2.628	1.767	1.168	0.604	0.336	0.199	0.131	0.098	0.127	0.063
	15.6	2.998	2.792	2.099	1.245	0.736	0.371	0.189	0.127	0.093	0.073	0.070	0.056
	0	0.114	0.121	0.089	0.088	0.069	0.068	0.054	0.052	0.054	0.057	0.058	0.063

Example 2

Limiting Antigen Assay of Test Antibodies

[0078] SA microtiter plates were coated with biotinylated antigen PTH at the optimal concentration of 15 ng/ml as determined in Example 1. Fifty microliters (50 μ l) of biotinylated antigen at a concentration of 15 ng/ml in 1% skim milk / 1X PBS pH 7.4 was added to each well, in a dilution pattern as described in Example 1. The plate was incubated for 30 minutes.

[0079] Plates were washed four times (4X) with water, and a B-cell culture supernatant containing test antibodies diluted 1:10 in 1% skim milk / 1X PBS pH 7.4 / .05% azide, and 50 μ l of each sample was added to each of two wells. Forty-eight (48) different samples were added per 96 well plate. On a separate plate, reference antibody 15g2 anti-PTH at the concentration used to determine the optimal antigen concentration was diluted out at least half a plate. This provided a positive control to assure that results from assays of test antibodies are comparable with optimization results.

[0080] Plates were wrapped tightly with plastic wrap or paraffin film, and incubated with shaking for 24 hours at room temperature.

[0081] On the following day, all plates were washed five times (5X) and 50 μ l goat anti-Human IgG Fc HRP polyclonal antibody at a concentration of 0.5 μ g/ml in 1% milk, 1X PBS pH 7.4 was added to each well. The plates were incubated for 1 hour at room temperature.

[0082] Plates were washed at least five times (5X with tap water). Fifty microliters (50 μ l) of HRP substrate TMB was added to each well, and the plate were incubated for 30 minutes. The HRP-TMB reaction was stopped by adding 50 μ l of 1M phosphoric acid to each well. Optical density (absorbance) at 450 nm was measured for each well of the plate.

Data Analysis

[0083] OD values of test antibodies were averaged and the range was calculated. Antibodies with the highest signal and acceptably low standard deviation were selected as antibodies having a higher affinity for the antigen than did the reference antibody.

[0084] Table 3 shows the results of a limiting antigen dilution assay using PTH as a ligand. Antibodies are ranked according to their relative affinity for various PTH antigens, and identified by their well number.

Table 3

Affinity Ranking of Test Antibodies to Limited Dilution of PTH								
Well	Limiting Ag OD	Limiting Ag Rank	Primary OD	Secondary OD	PTH (1-84)	PTH (7-84)	PTH (17-44)	Rat PTH (1-84)
292A10	2.747	1	0.992	ND	1.40	1.95	3.26	0.62
302A7	1.376	2	0.317	ND	0.35	0.36	2.66	0.19
253D10	1.009	3	0.954	0.511	0.79	1.10	2.10	1.18
263C8	0.693	5	0.372	0.286	1.75	1.98	3.29	1.34
245B10	0.644	6	0.622	0.580	0.84	0.32	0.12	0.19
238F8	0.566	7	0.667	0.541	1.05	1.34	2.79	1.19
228E3	0.504	8	0.560	0.259	0.48	0.80	3.12	1.40
262H1	0.419	9	0.461	0.274	0.86	1.20	2.45	0.36
161G7	0.411	10	0.409	0.212	0.49	0.90	1.88	0.84
331H6	0.322	11	0.312	ND	0.52	0.45	2.40	0.24
287E7	0.261	12	0.682	ND	0.71	0.13	0.36	1.03
315D8	0.221	13	0.441	ND	0.14	0.17	0.29	0.31
279E6	0.213	14	0.379	ND	0.31	0.10	0.17	0.19
250G6	0.178	15	0.560	0.248	0.44	0.66	1.77	0.19
244H11	0.175	16	0.405	0.556	0.50	0.86	0.98	0.31
313D5	0.170	17	0.664	ND	0.12	0.29	0.43	0.30
339F5	0.120	18	0.319	ND	0.40	0.21	0.11	0.25
279D2	0.114	19	0.353	ND	0.31	0.11	0.27	0.18
307H1	0.084	20	0.401	ND	0.10	0.14	0.30	0.42
308A1	0.079	21	0.312	ND	0.19	0.22	0.30	0.45
322F2	ND	22	1.870	ND	1.01	0.15	0.34	1.41

Example 3**Dilutions of Antibodies Against Interleukin-8 (IL-8)**

[0085] The proper coating concentration of IL-8 was determined as described above to determine a concentration of IL-8 that resulted in an OD of approximately 1. The optimal concentration was then incubated with a variety of anti-IL-8 antibody supernatants derived from XenoMouse animals immunized with IL-8. Table 4 illustrates typical results and ranking of antibodies screened for their affinity for IL-8. The columns "primary OD" and "secondary OD" refer to primary and secondary binding screen OD's achieved when non-limited amounts of IL-8 were used in the binding ELISA. OD values reported in the limited antigen section refer to an average of two binding ELISA's done at limited antigen. As shown by Table 4, the top three antibodies are able retain their binding to antigen even at the limited concentrations. Other antibodies which also achieved high OD's in the primary and secondary non-limited antigen binding ELISA were not able to achieve the same signal when antigen concentrations were limiting.

EP 1 554 576 B9

Table 4

Affinity Ranking of Test Antibodies to Limited Dilution of IL-8						
Clone Number				Limited Ag		
plate	well	Primary OD	Secondary OD	Average	St dev.	Limited Ag Rank
36	C6	1.95	3.023	1.32	4%	1
6	G11	2.021	1.403	0.90	9%	2
50	B1	1.818	2.398	0.82	14%	3
41	C11	1.83	3.218	0.81	19%	4
53	G5	1.128	2.521	0.80	1 %	5
44	B8	2.09	2.707	0.78	2%	6
51	G10	1.408	1.652	0.78	2%	7
53	E1	1.992	3.035	0.72	12%	8
38	C1	2.571	2.945	0.71	3%	9
32	F3	2.339	3.322	0.66	13%	10
13	F10	1.505	1.833	0.66	5%	11
41	D2	2.997	2.944	0.66	5%	12
53	C2	1.56	1.869	0.64	22%	13
14	E2	1.255	1.875	0.57	25%	14
54	C3	2.131	2.486	0.51	12%	15
50	F3	0.572	1.635	0.51	26%	16
55	E8	1.031	1.917	0.50	10%	17
42	E5	3.07	3.147	0.49	4%	18
6	E7	0.637	1.545	0.49	22%	19
7	E10	1.794	1.953	0.48	18%	20
8	B2	1.725	1.777	0.48	5%	21
48	E6	2.103	3.004	0.48	25%	22
33	A1	2.623	2.351	0.47	17%	23
51	F5	2.062	2.838	0.45	15%	24
51	B1	1.778	2.631	0.45	0%	25
44	A5	2.473	2.55	0.44	5%	26
6	G4	2.117	1.505	0.41	7%	27
43	G4	0.991	1.943	0.41	2%	28
47	E3	1.049	2.222	0.40	16%	29
46	F11	1.641	1.843	0.39	9%	30
43	F4	0.744	1.449	0.39	7%	31
54	H1	1.465	1.584	0.38	25%	32
44	F4	2.05	2.573	0.38	13%	33
49	G11	1.334	2.019	0.37	6%	34
11	C10	1.169	1.498	0.37	3%	35

EP 1 554 576 B9

(continued)

Affinity Ranking of Test Antibodies to Limited Dilution of IL-8						
Clone Number				Limited Ag		
plate	well	Primary OD	Secondary OD	Average	St dev.	Limited Ag Rank
41	B12	1.107	1.347	0.37	3%	36
46	F2	0.865	1.15	0.37	11%	37
52	E11	0.961	2.034	0.37	5%	38
7	B6	2.039	1.802	0.33	6%	39
39	F6	1.434	1.196	0.33	6%	40
10	E5	0.886	1.262	0.33	6%	41
36	C12	1.078	1.991	0.33	10%	42
44	B9	1.469	1.683	0.32	4%	43
8	H1	1.338	1.316	0.31	2%	44
52	F3	1.289	1.204	0.28	16%	45
45	A4	1.136	1.302	0.28	13%	46
25	A11	1.199	1.17	0.27	25%	47
51	C12	0.955	1.148	0.26	11 %	48
6	E5	1.41	1.138	0.24	8%	49
39	H3	0.471	1.155	0.23	6%	50
14	E3	1.958	1.255	0.22	15%	51
3	D1	2.254	3.497	0.21	24%	52
33	F4	1.323	1.408	0.21	24%	53
51	A12	0.555	1.522	0.19	17%	54
5	G1	2.205	2.274	0.17	4%	55
35	C9	1.217	1.249	0.17	4%	56
6	B10	1.006	1.145	0.17	8%	57
39	B4	1.326	1.62	0.17	8%	58
5	G3	1.192	1.387	0.17	29%	59
35	F10	1.307	1.777	0.17	29%	60
17	E11	0.839	1.805	0.17	15%	61
3	D3	0.605	1.351	0.16	5%	62
31	A1	1.557	1.826	0.16	17%	63
28	C5	1.373	1.942	0.16	5%	64
14	F5	1.441	1.482	0.15	25%	65
43	D8	0.714	1.501	0.15	22%	66
29	D5	1.326	1.322	0.14	23%	67
32	F11	1.36	1.284	0.48	71%	68
7	D4	0.874	2.333	0.44	34%	69
47	G11	0.811	1.209	0.42	76%	70
39	G2	0.676	1.157	0.42	32%	71

EP 1 554 576 B9

(continued)

Affinity Ranking of Test Antibodies to Limited Dilution of IL-8						
Clone Number				Limited Ag		
plate	well	Primary OD	Secondary OD	Average	St dev.	Limited Ag Rank
15	G4	2.046	2.461	0.39	41%	72
31	G12	1.902	1.929	0.36	44%	73
41	C2	1.201	2.522	0.33	34%	74
7	E11	1.402	1.719	0.32	50%	75
40	A4	1.786	1.427	0.32	50%	76
45	E12	1.986	2.887	0.26	54%	77
2	B10	1.871	1.389	0.22	38%	78
7	H8	1.516	1.171	0.22	45%	79
28	C3	1.246	1.182	0.15	52%	80

Table 4A

Affinity Measurement of Reference Antibody 1		
Reference antibody 1		
Conc. ng/ml	Limited Ag OD	St. Dev.
125.00	1.52	1 %
62.50	1.38	2%
31.25	1.25	12%
15.63	1.13	28%
7.81	0.80	2%
3.91	0.78	18%
1.95	0.67	0%
0.98	0.73	8%
0.49	0.53	18%
0.24	0.39	17%

Table 4B

Affinity Measurement of Reference Antibody 2		
Reference antibody 2		
Conc. ng/ml	Limited Ag OD	St. Dev.
125.00	0.52	23%
62.50	0.38	11%
31.25	0.34	1%
15.63	0.42	43%
7.81	0.54	13%
3.91	0.46	30%

EP 1 554 576 B9

(continued)

Affinity Measurement of Reference Antibody 2		
Reference antibody 2		
Conc. ng/ml	Limited Ag OD	St. Dev.
1.95	0.54	9%
0.98	0.34	9%
0.49	0.49	32%
0.24	0.55	38%

Example 4

Affinity Ranking

Preparation of Antigens

[0086] In order to increase the effective throughput of the antibody affinity ranking process, we labeled different concentrations of an antigen with different colored beads. In this example, beads from the Luminex system were used. As is known, each bead, when activated, emits light of a varying wavelength. When put in a Luminex reader, the identity of each bead can be readily ascertained.

[0087] In this example, a different color of strepavidin luminex bead was bound to each of four concentrations of biotinylated antigen (1 μ g/ml, 100ng/ml, 30ng/ml, and 10ng/ml). Thus, each concentration of the antigen was represented by a different color bead. The four concentrations were then mixed into a single solution containing all four color-bound concentrations.

[0088] All of the antibody samples were then diluted to the same concentration (~500ng/ml) using Luminex quantitation results or a one-point quantitation by Luminex. A serial dilution (1:5) of all of the samples was then performed so a total of four dilution points were obtained, while preferably diluting enough sample for two plates: a quantitation plate and the ranking plate.

Ranking of Antibodies

[0089] In order to rank the antibodies, ~ 2000 of each mixture of luminex bead-antigen samples was loaded into each well of the luminex plate, and then the well was aspirated. Then 50 μ l of each antibody sample (24 samples total) was loaded into each well and left overnight while shaking in 4°C. The plates were washed three times (3X) with washing buffer. Detection with a fluorescent anti-human antibody (hIgG-Phycoerythrin (PE) (1:500 dilution)) that bound 50 μ l/well was then performed while shaking at room temperature for 20 min. The plates were then washed three times (3X) with washing buffer. The plates were re-suspended in 80 μ l blocking buffer. Next, the plates were loaded in the Luminex.

Data Analysis

[0090] Because each well held four different concentrations of the same antigen, that could be distinguished based on color, it was possible to rapidly rank binding affinities of the different antibodies. For example, antibodies that had very strong binding affinity for the antigen bound to even the weakest dilution of antibody. This could be measured by analyzing the amount of fluorescent anti-human antibody bound to the colored bead attached to the weakest antigen concentration. Alternatively, antibodies that did not bind strongly might be only detected as binding with the 1 μ g/ml and 100ng/ml antigen concentrations, but not the 30ng/ml or 10ng/ml concentrations.

[0091] Data analysis was performed using SoftMax Pro for the quantitation data. The Luminex signal of samples tested at several concentrations were compared. The samples were then ranked accordingly.

Example 5

Comparison of Limiting Antigen Output Compared to Absolute Biacore KD Measurements

[0092] The following kinetic ranking technique was performed by ELISA and compared to formal Biacore kinetics. Below in Table 5 is a comparison of a typical limited antigen output as compared to absolute Biacore derived KD

measurements. In short, 68 antibodies were ranked (relative to each other) using limited antigen ranked. From the 68 antibodies 17 were scaled up to sufficient quantities for formal affinity measurements using Biacore technology.

Table 5

Comparison of Affinity Measurement Based on Limited Dilutions with Biacore Affinity Measurements		
Sample ID	Limited Antigen Ranking	Biacore Affinity (nM)
A	1	1.9
B	3	1.9
C	4	1.3
D	5	6.9
E	7	3.3
F	10	17.7
G	11	28.9
H	12	3.8
I	13	4.4
J	23	11.2
K	28	57.8
L	30	29.2
M	34	1667
N	46	115.2
O	47	305.1
P	51	1000
Q	60	33.1

Data Analysis

[0093] As can be seen overall there is a high degree of correlation between high limited antigen rank and the formal K_D . In the case of antibodies which do not correlate well, there are a number of reasons why such discrepancies could exist. For example, although antigen is coated on ELISA plates at a low density avidity effects cannot completely be ruled out. In addition, it is possible that, when coating assay material for the limited antigen ranking technique, certain epitopes could be masked or altered. In Biacore analysis, if antigen is flowed over an antibody coated chip, these epitopes on the antigen could be presented in a different conformation and, therefore, seen at a different relative concentration. This could, in turn, could result in a different kinetic ranking between the two methods.

[0094] It is also possible that an antibody with lower Biacore derived affinities may give a high limited antigen rank due to a much higher than average concentration of antigen specific antibody being present in the test sample. This could, in turn, lead to an artificially high limited antigen score.

[0095] Importantly, the limited antigen kinetics method did allow a rapid determination of relative affinity and it identified the antibodies with the highest formal affinity of the tested antibodies in this panel. Further, as the limited antigen kinetic relative ranking method is easily scalable to interrogate 1000's of antibodies at early stages of antibody generation it offers significant advantage over other technologies which do not offer similar advantages of scale.

Claims

1. A method of determining the relative binding affinities of antibodies to an antigen, comprising:

providing a reference antibody that binds to a target antigen;
determining the limiting concentration of the target antigen for the reference antibody, wherein the limiting concentration-maximizes the detectable range of antibody affinities within a predetermined concentration range

for a set of test antibody solutions;
 providing a the set of test antibody solutions, wherein the concentration of antibodies in each test antibody solution is within the predetermined concentration range;
 incubating the limiting concentration of antigen with each test antibody solution in the set of test antibody solutions;
 measuring the relative degree to which the antibodies in each test antibody solution binds to the limiting concentration of antigen; and
 ranking the antibodies in each test antibody solution based on their binding affinity to said limited concentration of antigen in order to determine the relative binding affinity of each test antibody to said antigen.

2. The method of Claim 1, wherein the test antibodies are selected from the group consisting of: mammalian, human, humanized, non-human primate, mouse, rat, rabbit, goat, horse, guinea pig, sheep, cow, and chimeric antibodies.
3. The method of Claim 1, wherein the test antibodies are recombinant antibodies.
4. The method of Claim 1, wherein the test antibodies are antibody fragments.
5. The method of Claim 1, wherein measuring the relative degree comprises an assay selected from a group consisting of: an enzyme linked immunosorbent assay (ELISA), a fluorescence-linked immunosorbent assay, a radioisotopic assay (RIA), and a surface plasmon resonance (SPR) based assay.
6. The method of Claim 1, wherein the concentration of each test antibody is between 10 ng/ml and 100 ng/ml.
7. The method of Claim 1, wherein said concentration of each test antibody is between 10 ng/ml and 50 ng/ml.
8. The method of Claim 1, wherein said concentration of each test antibody is between 10 ng/ml and 25 ng/ml.
9. The method of Claim 1, wherein said set of test antibody solutions comprises a set of supernatants comprising monoclonal antibodies secreted from a set of hybridomas.
10. The method of Claim 1, wherein said set of test antibody solutions is between 100 and 5000 antibody solutions.
11. The method of Claim 1, wherein said set of test antibody solutions is between 2500 and 5000 antibody solutions.
12. The method of Claim 1, wherein each test antibody solution is a polyclonal antibody solution.
13. The method of Claim 1, wherein each test antibody solution is a monoclonal antibody solution.

Patentansprüche

1. Verfahren zur Bestimmung der relativen Bindungsaffinitäten von Antikörpern zu einem Antigen, umfassend:

das Bereitstellen eines Referenzantikörpers, der an ein Zielantigen bindet;
 das Bestimmen der Grenzkonzentration des Zielantigens für den Referenzantikörper, wobei die Grenzkonzentration den Nachweisbereich von Antikörperaffinitäten innerhalb eines vorbestimmten Konzentrationsbereichs für einen Satz von Testantikörperlösungen maximiert;
 das Bereitstellen eines Satzes von Testantikörperlösungen, wobei die Antikörperkonzentration in jeder Testantikörperlösung innerhalb des vorbestimmten Konzentrationsbereichs liegt;
 das Inkubieren der Antigengrenzkonzentration mit jeder Testantikörperlösung im Satz von Testantikörperlösungen;
 das Messen des relativen Grads, zu dem die Antikörper in jeder Testantikörperlösung an die Antigengrenzkonzentration binden; und
 das Einordnen der Antikörper in jeder Testantikörperlösung auf der Grundlage ihrer Bindungsaffinität zur Antikörpergrenzkonzentration in eine Rangfolge, um die relative Bindungsaffinität eines jeden Testantikörpers an das Antigen zu bestimmen.

2. Verfahren nach Anspruch 1, wobei die Testantikörper aus der aus Säugetierantikörpern, humanen und humanisierten

Antikörpern, nicht-humanen Primatenantikörpern, Maus-, Ratten-, Kaninchen-, Ziegen-, Pferd-, Meerschweinchen-, Schaf- und Rindantikörpern und chimären Antikörpern bestehenden Gruppe ausgewählt sind.

3. Verfahren nach Anspruch 1, wobei die Testantikörper rekombinante Antikörper sind.

4. Verfahren nach Anspruch 1, wobei die Antikörper Antikörperfragmente sind.

5. Verfahren nach Anspruch 1, wobei das Messen des relativen Grads einen Assay umfasst, der aus einer aus einem enzymgekoppelten Immunadsorptionstest (ELISA-Assay), einem fluoreszenzgekoppelten Immunadsorptionsassay, einem radioisotopischen Assay (RIA) und einem auf Oberflächenplasmonenresonanz (SPR) basierenden Assay bestehenden Gruppe ausgewählt ist.

6. Verfahren nach Anspruch 1, wobei die Konzentration eines jeden Testantikörpers zwischen 10 ng/ml und 100 ng/ml beträgt.

7. Verfahren nach Anspruch 1, wobei die Konzentration eines jeden Testantikörpers zwischen 10 ng/ml und 50 ng/ml beträgt.

8. Verfahren nach Anspruch 1, wobei die Konzentration eines jeden Testantikörpers zwischen 10 ng/ml und 25 ng/ml beträgt.

9. Verfahren nach Anspruch 1, wobei der Satz von Testantikörperlösungen einen Satz von Überständen umfasst, die monoklonale Antikörper umfassen, die aus einem Satz von Hybridomen sekretiert sind.

10. Verfahren nach Anspruch 1, worin der Satz von Testantikörperlösungen aus zwischen 100 und 5000 Antikörperlösungen besteht.

11. Verfahren nach Anspruch 1, worin der Satz von Testantikörperlösungen aus zwischen 2500 und 5000 Antikörperlösungen besteht.

12. Verfahren nach Anspruch 1, worin jede Testantikörperlösung eine polyklonale Antikörperlösung ist.

13. Verfahren nach Anspruch 1, worin jede Testantikörperlösung eine monoklonale Antikörperlösung ist.

Revendications

1. Procédé de détermination des affinités de liaison relatives d'anticorps à un antigène, qui comprend:

la fourniture d'un anticorps de référence qui se lie à un antigène cible ;
la détermination de la concentration limite de l'antigène cible pour l'anticorps de référence, dans lequel la concentration limite maximise la plage détectable d'affinités de l'anticorps à l'intérieur d'une plage de concentrations prédéterminée pour un jeu de solutions d'anticorps de test ;
la fourniture du jeu de solutions d'anticorps de test, dans lequel la concentration en anticorps dans chaque solution d'anticorps de test se trouve à l'intérieur de la plage de concentrations prédéterminée ;
l'incubation de la concentration limite d'antigène avec chaque solution d'anticorps de test dans le jeu de solutions d'anticorps de test ;
la mesure du degré relatif auquel les anticorps dans chaque solution d'anticorps de test se lient à la concentration limite d'antigène ; et
la notation des anticorps dans chaque solution d'anticorps de test en se basant sur leur affinité de liaison à ladite concentration limite d'antigène de manière à déterminer l'affinité de liaison relative de chaque anticorps de test au dit antigène.

2. Procédé selon la revendication 1, dans lequel les anticorps de test sont choisis dans le groupe constitué par des anticorps : de mammifère, humains, humanisés, de primate non humain, de souris, de rat, de lapin, de chèvre, de cheval, de cobaye, de mouton, de vache, et chimériques.

3. Procédé selon la revendication 1, dans lequel les anticorps de test sont des anticorps recombinants.

EP 1 554 576 B9

4. Procédé selon la revendication 1, dans lequel les anticorps de test sont des fragments d'anticorps.
5. Procédé selon la revendication 1, dans lequel la mesure du degré relatif comprend un dosage choisi dans un groupe constitué par: un dosage immunoenzymatique (ELISA), un dosage immunofluorescent (FLISA), un dosage radioisotopique (RIA), et un dosage basé sur la résonnance plasmonique de surface (SPR).
6. Procédé selon la revendication 1, dans lequel la concentration de chaque anticorps de test est comprise entre 10 ng/mL et 100 ng/mL.
7. Procédé selon la revendication 1, dans lequel ladite concentration de chaque anticorps de test est comprise entre 10 ng/mL et 50 ng/mL.
8. Procédé selon la revendication 1, dans lequel ladite concentration de chaque anticorps de test est comprise entre 10 ng/mL et 25 ng/mL.
9. Procédé selon la revendication 1, dans lequel ledit jeu de solutions d'anticorps de test comprend un jeu de surnageants comprenant des anticorps monoclonaux sécrétés par un jeu d'hybridomes.
10. Procédé selon la revendication 1, dans lequel ledit jeu de solution d' anticorps de test comprend de 100 à 5000 solutions d'anticorps.
11. Procédé selon la revendication 1, dans lequel ledit jeu de solution d'anticorps de test comprend de 2500 à 5000 solutions d'anticorps.
12. Procédé selon la revendication 1, dans lequel chaque solution d'anticorps de test est une solution d'anticorps polyclonaux.
13. Procédé selon la revendication 1, dans lequel chaque solution d'anticorps de test est une solution d'anticorps monoclonaux.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 5627052 A [0002]
- US 5800815 A [0004]

Non-patent literature cited in the description

- KOHLER ; MILSTEIN. *Nature*, 1975, vol. 256, 52-53 [0002]
- BABCOOK et al. *Proc Natl Acad Sci U S A.*, 1996, vol. 93, 7843-8 [0002]
- WATKINS et al. *Anal Biochem*, 1997, vol. 253, 37-45 [0004]