

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 557 157 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

27.07.2005 Bulletin 2005/30

(51) Int Cl.7: **A61K 7/13**

(21) Application number: **04001401.1**

(22) Date of filing: **23.01.2004**

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR**

Designated Extension States:

AL LT LV MK

(72) Inventors:

- **Lorenz, Heribert**
64401 Gross-Bieberau (DE)
- **Kaffenberger, Klaus**
64665 Alsbach-Hähnlein (DE)

(71) Applicant: **KPSS-Kao Professional Salon**

Services GmbH

64297 Darmstadt (DE)

(54) **Composition for the dyeing of human hair**

(57) Hair dyeing composition on the basis of an oxidation dyestuff precursor reacting with peroxide, comprising at least one developing and/or coupling substance selected from the group

- 3-methyl-p-aminophenol and/or 2-methyl-p-aminophenol,
- 3-(N-methyl-N-hydroxyethyl amino)-phenol, 3-morpholinophenol, 3-(N-hydroxyethyl amino)-phenol and/or 3-(N-hydroxypropyl amino)-phenol, and
- p-phenylenediamine, p-toluylenediamine, 2,6-dimethyl-p-phenylenediamine, 2-hydroxyme-

thyl-p-phenylenediamine, 2-hydroxyethyl-p-phenylenediamine, 2-n-propyl-p-phenylenediamine, 2-isopropyl-p-phenylenediamine, N-(β -hydroxypropyl)-p-phenylenediamine, N,N-bis-(β -hydroxyethyl)-p-phenylenediamine, N-methoxyethyl-p-phenylenediamine and/or 5-chloro-2-hydroxyethyl-p-phenylenediamine or the water-soluble salts thereof.

EP 1 557 157 A1

Description

[0001] The present invention concerns a composition for the dyeing of human hair on the basis of an oxidation dyestuff precursor system reacting with peroxide which provides long-lasting, intensive colors either used as such, or which can be used to obtain further shades in combination with additional developing and/or coupling agents and which does not damage the hair even upon repeated application within short intervals.

[0002] The developing substances still most frequently used in hair dyeing compositions are 1,4-diamineobenzene (p-phenylenediamine) and 1-methyl-2,5-diamineobenzene (p-toluylenediamine). Although incorporation of these substances largely fulfills the user's color wishes, there are still shades that cannot be completely achieved by the use thereof.

[0003] Proposals have also been made to close this gap by the use of alternative developing substances. To a limited degree this is possible with the use of tetraaminopyrimidine or 2-(2,5-diamineophenyl)ethanol (see EP-B 400 330); however, it is then necessary to accept reduced color intensity in other shades.

[0004] A further satisfactory solution of this problem is disclosed in EP-A 615 743, with the use of 2-(2'-hydroxyethyl amino)-5-aminotoluene or the water-soluble salts thereof, and 3,4-diamineo-5-hydroxypyrazole as components of oxidation hair dye compositions.

[0005] However, to the present it has not been possible to achieve strong colorations in the range of brown, violet and brown-violet by this means.

[0006] The invention starts from the task of counteracting this deficiency and providing an oxidation dyestuff composition which achieves intensive, glossy colorations, especially in the range of brown, violet and brown-violet, and which leaves the hair without damage even upon repeated application within short periods of time.

[0007] This task is solved when such a hair dyeing composition comprises an oxidation dyestuff system reacting with peroxide which is selected from

- a) 3-methyl-p-aminophenol and/or 2-methyl-p-aminophenol,
- b) 3-(N-methyl-N-hydroxyethyl amino)-phenol, 3-morpholinophenol, 3-(N-hydroxyethyl amino)-phenol and/or 3-(N-hydroxypropyl amino)-phenol, and
- c) p-phenylenediamine, p-toluylenediamine, 2,6-dimethyl-p-phenylenediamine, 2-hydroxy methyl-p-phenylenediamine, 2-hydroxyethyl-p-phenylenediamine, 2-n-propyl-p-phenylenediamine, 2-isopropyl-p-phenylenediamine, N-(β-hydroxypropyl)-p-phenylenediamine, N,N-bis-(β-hydroxyethyl)-p-phenylenediamine, N-methoxyethyl-p-phenylenediamine and/or 5-chloro-2-hydroxyethyl-p-phenylenediamine or the water-soluble salts thereof.

[0008] After oxidation with peroxide, use of these compositions on the basis of a customary carrier provides very expressive, intensive, long-lasting hair colorations, especially in the range of brown, violet and brown-violet, which can be varied to achieve further shades by the addition of the respective further developing and coupling substances.

[0009] In addition to the named developing and coupling substances it is also possible to incorporate further substances of this type.

[0010] Further suitable coupling substances are, for example, 1-methoxy-2-amino-4-(β-hydroxyethyl amino)benzene, 2-amino-3-hydroxypyridine, 3-amino-2-methylamino-6-methoxypyridine, resorcinol, 2-methyl resorcinol, 4-chlororesorcinol, 2-amino-4-chlorophenol, 1,3-diaminobenzene, 1,6-dihydroxynaphthaline, 1,7-dihydroxynaphthaline, 2-aminophenol and/or 3-aminophenol.

[0011] The total concentration of the developing substances customarily ranges between about 0.05 % and 5 %, preferably 0.1 % and 4 %, in particular 0.25 % to 3 % by weight, calculated to the total hair dyeing composition (excluding the oxidizing agent), whereby these figures are always related to the proportion of free base. The preferred weight proportion of the developing substances to the coupling substances ranges between about 1 : 8 to 8 : 1, preferably about 1 : 5 to 5 : 1, in particular 1 : 2 to 2 : 1.

[0012] In the hair dyeing compositions according to the invention, the coupling substance(s) as reaction partners of the developing substance(s) are present in approximately the same molecular proportions as the developing substances, i.e. in amounts from 0.05 % to 5.0 %, preferably 0.1 % to 4 %, in particular 0.5 % to 3 % by weight, calculated to the total composition (excluding the oxidizing agent), whereby these figures are always related to the proportion of free base.

[0013] If desired, the compositions according to the invention can also contain so-called shading agents for precise adjustment of the desired shade, in particular direct-acting dyestuffs.

[0014] Such shading agents are, for example, nitro dyestuffs such as 2-amino-4,6-dinitrophenol, 2-amino-4-nitrophenol, 2-amino-6-chloro-4-nitrophenol, etc., preferably in amounts from about 0.05 % to 2.5 %, in particular 0.1 % to 1 % by weight of the dyestuff composition (excluding the oxidizing agent).

[0015] The hair dyeing compositions according to the invention can comprise the basic substances and additives customarily found in such compositions, conditioning agents, etc., known as state of the art and described, for example,

in the monograph by K. Schrader, "Grundlagen und Rezepturen der Kosmetika", 2nd Ed. (Hüthig Buch Verlag, Heidelberg, 1989), pp. 782 to 815. They can be prepared as solutions, creams, gels or also in the form of aerosol products; suitable carrier material compositions are known as state of the art.

[0016] For application, the oxidation dyestuff precursor is mixed with an oxidizing agent. The preferred oxidizing agent is hydrogen peroxide, for example in a concentration of 2 % to 6 % by weight.

[0017] However, the use of other peroxides such as urea peroxide and melamine peroxide is also possible.

[0018] The pH-value of the ready-to-use hair dyeing composition, i.e. after mixing with peroxide, can be in a slightly acidic range, i.e. from 5.5 to 6.9, as well as in the neutral or alkaline range, i.e. between pH 7.1 and 10.

[0019] In the following, various Examples are used to illustrate the invention.

Carrier

[0020]

Stearyl alcohol	8.0 (% by wt.)
Coco fatty acid monoethanolamide	4.5
1,2-Propanediol mono/distearate	1.3
Coco fatty alcohol polyglycol ether	4.0
Sodium lauryl sulfate	1.0
Oleic acid	2.0
1,2-Propanediol	1.5
Na-EDTA	0.5
Sodium sulfite	1.0
Protein hydrolyzate	0.5
Ascorbic acid	0.2
Perfume	0.4
Ammonia, 25%	1.0
Ammonium chloride	0.5
Panthenol	0.8
Water	ad 100.00

[0021] The oxidation dyestuff combinations according to the invention were incorporated into this carrier, whereby the water content was reduced accordingly.

[0022] The colorations were carried out on wool patches and strands of bleached human hair by application of a 1:1 mixture of a dyestuff precursor and a 6 % hydrogen peroxide solution (pH-value of the mixture: 9.8) with twenty minutes processing at room temperature, subsequent rinsing and drying.

[0023] The following colorations were achieved:

Example 1:

[0024]

0.19 (% by wt.)	3-Methyl-p-aminophenol
0.25	3-(N-methyl-N-hydroxyethyl aminophenol
0.33	p-Toluylenediamine sulfate
<u>Coloration:</u>	Light gold-brown

Example 2:

[0025]

0.19 (% by wt.)	3-Methyl-p-aminophenol
0.27	2-Morpholinophenol
0.16	p-Phenylenediamine

EP 1 557 157 A1

(continued)

<u>Coloration:</u>	Brown gray-violet
--------------------	-------------------

Example 3:

[0026]

0.19	(% by wt.)	3-Methyl-p-aminophenol
0.23		3-(N-hydroxyethyl amino)-phenol
0.38		2-Hydroxyethyl-p-phenylenediamine sulfate
<u>Coloration:</u>		Magenta-brown

Example 4:

[0027]

0.19	(% by wt.)	3-Methyl-p-aminophenol
0.25		3-(N-hydroxypropyl amino)-phenol
0.47		N,N-bis-(β -hydroxyethyl)-p-phenylenediamine sulfate
<u>Coloration:</u>		Gray-violet

Example 5:

[0028]

0.19	(% by wt.)	2-Methyl-p-aminophenol
0.27		2-Morpholinophenol
0.38		2-Hydroxyethyl-p-phenylenediamine sulfate
<u>Coloration:</u>		Red-violet (purple)

Example 6:

[0029]

0.14	(% by wt.)	3-Methyl-p-aminophenol
0.35		3-(N-hydroxyethyl amino)-phenol
0.28		2-Hydroxyethyl-p-phenylenediamine sulfate
<u>Coloration:</u>		Magenta-brown

Example 7:

[0030]

0.14	(% by wt.)	3-Methyl-p-aminophenol
0.17		3-(N-hydroxyethyl amino)-phenol
0.28		2-Hydroxyethyl-p-phenylenediamine sulfate
0.12		Resorcinol
<u>Coloration:</u>		Hazelnut blond

Claims

1. Hair dyeing composition on the basis of an oxidation dyestuff precursor reacting with peroxide, comprising at least

one developing and/or coupling substance selected from the group

- a) 3-methyl-p-aminophenol and/or 2-methyl-p-aminophenol,
- b) 3-(N-methyl-N-hydroxyethyl amino)-phenol, 3-morpholinophenol, 3-(N-hydroxyethyl amino)-phenol and/or 3-(N-hydroxypropyl amino)-phenol, and
- c) p-phenylenediamine, p-toluylenediamine, 2,6-dimethyl-p-phenylenediamine, 2-hydroxy methyl-p-phenylenediamine, 2-hydroxyethyl-p-phenylenediamine, 2-n-propyl-p-phenylenediamine, 2-isopropyl-p-phenylenediamine, N-(β -hydroxypropyl)-p-phenylenediamine, N,N-bis-(β -hydroxyethyl)-p-phenylenediamine, N-methoxyethyl-p-phenylenediamine and/or 5-chloro-2-hydroxyethyl-p-phenylenediamine or the water-soluble salts thereof.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 04 00 1401

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	EP 1 374 841 A (KPSS KAO PROFESSIONAL SALON SE) 2 January 2004 (2004-01-02) * claims *	1	A61K7/13
A	EP 1 374 840 A (KPSS KAO PROFESSIONAL SALON SE) 2 January 2004 (2004-01-02) * claims *	1	
A	EP 1 297 818 A (KPSS KAO PROFESSIONAL SALON SE) 2 April 2003 (2003-04-02) * claims 1,2; examples 4,5 *	1	
A	US 6 264 703 B1 (COOPE JANET LYNN) 24 July 2001 (2001-07-24) * claims 1,4,5 *	1	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			A61K
The present search report has been drawn up for all claims			
Place of search Berlin		Date of completion of the search 21 June 2004	Examiner Boeker, R
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

1
EPO FORM 1503 03/02 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 04 00 1401

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

21-06-2004

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1374841	A	02-01-2004	DE 10228244 A1	22-01-2004
			EP 1374841 A2	02-01-2004
			US 2003233716 A1	25-12-2003

EP 1374840	A	02-01-2004	DE 10228245 A1	22-01-2004
			EP 1374840 A2	02-01-2004
			US 2003233715 A1	25-12-2003

EP 1297818	A	02-04-2003	EP 1297818 A2	02-04-2003

US 6264703	B1	24-07-2001	CA 2334288 A1	10-08-2001
			EP 1129688 A1	05-09-2001
			JP 2001226239 A	21-08-2001
