



(11) **EP 1 595 950 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

(15) Correction information:
Corrected version no 2 (W2 B1)
Corrections, see
Description Paragraph(s) 36, 38

(48) Corrigendum issued on:
10.06.2009 Bulletin 2009/24

(45) Date of publication and mention
of the grant of the patent:
29.10.2008 Bulletin 2008/44

(21) Application number: **04705917.5**

(22) Date of filing: **28.01.2004**

(51) Int Cl.:
G01N 33/68 (2006.01) **C12Q 1/25** (2006.01)
C12Q 1/26 (2006.01) **C12Q 1/48** (2006.01)

(86) International application number:
PCT/JP2004/000787

(87) International publication number:
WO 2004/074477 (02.09.2004 Gazette 2004/36)

(54) **METHOD OF MEASURING HOMOCYSTEINE**
VERFAHREN ZUR MESSUNG VON HOMOCYSTEIN
PROCEDE DE MESURE D'HOMOCYSTEINE

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR

(30) Priority: **19.02.2003 JP 2003041585**

(43) Date of publication of application:
16.11.2005 Bulletin 2005/46

(73) Proprietor: **Alfresa Pharma Corporation**
Osaka-shi, Osaka 540-8575 (JP)

(72) Inventors:
• **ESAKI, Nobuyoshi**
Ohtsu-shi, Shiga 5202279 (JP)
• **YOSHIMURA, Tohru,**
Room 403, Corpo Momoyama
Kyoto-shi,
Kyoto 6120066 (JP)
• **MATSUYAMA, Naoto**
Takatsuki-shi, Osaka 5691041 (JP)
• **MIZUNO, Koji**
Ibaraki-shi, Osaka 5670805 (JP)
• **BOGAKI, Takayuki**
Higashinada-ku Kobe-shi, Hyogo 6580082 (JP)
• **MINETOKI, Toshitaka**
Suma-ku Kobe-shi, Hyogo 6540123 (JP)
• **OZEKI, Kenji**
Nishinomiya-shi, Hyogo 6638184 (JP)

(74) Representative: **Arth, Hans-Lothar et al**
ABK Patent Attorneys
Jasminweg 9
14052 Berlin (DE)

(56) References cited:
EP-A2- 1 213 358 WO-A1-00/34771
WO-A1-02/02802 WO-A1-93/15220
WO-A1-95/30151 WO-A1-98/07872
JP-A- 2001 017 198 JP-A- 2002 360 295
JP-A- 2003 144 195 US-A- 5 998 191
US-A1- 2002 123 088

- **MATSUYAMA N ET AL: "New enzymatic colorimetric assay for total homocysteine" CLINICAL CHEMISTRY, AMERICAN ASSOCIATION FOR CLINICAL CHEMISTRY, WASHINGTON, DC, US, vol. 47, no. 12, December 2001 (2001-12), pages 2155-2157, XP002346199 ISSN: 0009-9147**
- **ASANO Y ET AL: "Enzymes acting on peptides containing D-amino acid." JOURNAL OF BIOSCIENCE AND BIOENGINEERING 2000, vol. 89, no. 4, 2000, pages 295-306, XP002415780 ISSN: 1389-1723**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 595 950 B9

- DUCROS V ET AL: "Methods for homocysteine analysis and biological relevance of the results"
JOURNAL OF CHROMATOGRAPHY B:
BIOMEDICAL SCIENCES & APPLICATIONS,
ELSEVIER, AMSTERDAM, NL, vol. 781, no. 1-2, 5
December 2002 (2002-12-05), pages 207-226,
XP004394155 ISSN: 1570-0232

Description

Technical Field

5 **[0001]** The present invention relates to a method of detecting or measuring homocysteine in a sample. More specifically, the present invention relates to a method of measuring homocysteine including the step of removing a D-amino acid present in a sample in advance.

Background Art

10 **[0002]** As a method of measuring homocysteine, a method of reacting homocysteine in a sample with homocysteine methyltransferase and D-methionine methylsulfonium, and then detecting the produced D-methionine with D-amino acid oxidase has been reported (see WO 02/02802). However, it is known that biological samples contain D-amino acids such as D-alanine and D-serine in small amounts, and these amino acids increase in the case of renal diseases or the like (e.g., see Fukushima, T., Biol. Pharm. Bull., 1995, Vol. 18, No. 8, pp. 1130-1132). It is believed that D-alanine and D-serine, which can be a substrate of D-amino acid oxidase, leads to a positive reading in the method of measuring homocysteine. Therefore, as described in WO 02/02802, in order to avoid the influence of endogenous D-amino acids that are originally present in a sample, it is necessary to subtract a value obtained by measurement by the same operation except that homocysteine methyltransferase is not contained from a measured value in the case of containing this enzyme. That is to say, it is necessary to provide a sample blank for each individual sample to measure the amount of endogenous D-amino acids.

Disclosure of Invention

25 **[0003]** It is an object of the present invention to provide a method of measuring homocysteine that is not affected by endogenous D-amino acids, that is, that does not require sample blanks.

[0004] As a result of in-depth research in order to achieve the above object, it became possible to provide a method of measuring homocysteine that is not affected by endogenous D-amino acids, that is, that does not require sample blanks by leading D-alanine and/or D-serine to the outside of the reaction system of the homocysteine measurement by enzyme actions.

[0005] The present invention provides a method for detecting or measuring homocysteine in a sample, and the method includes the steps of: (a) reacting a D-amino acid present in a sample with a D-amino acid converting enzyme to convert the D-amino acid into a substance that does not serve as a substrate of D-amino acid oxidase or D-amino acid acetyltransferase; (b) reducing homocysteine in the sample with a thiol compound;

35 (c) reacting the reduced homocysteine with a methyltransferase and D-methionine methylsulfonium to newly produce D-methionine; and

(d) reacting the D-methionine produced in step (c) with the D-amino acid oxidase in the presence of an SH reagent to produce hydrogen peroxide, and color-developing the produced hydrogen peroxide by using an oxidative color-developing agent, or

40 (d') reacting the D-methionine produced in step (c) with the D-amino acid acetyltransferase in the presence of acetyl coenzyme A thereby producing coenzyme A; reacting the produced coenzyme A with acyl coenzyme A synthetase thereby producing acyl coenzyme A; and reacting the produced acyl coenzyme A with acyl coenzyme A oxidase in the presence of an SH reagent to produce hydrogen peroxide; and color-developing the produced hydrogen peroxide by using an oxidative color-developing agent, wherein the thiol compound is selected from dithiothreitol, mercaptoethanol, N-acetylcysteine, dithioerythritol and thioglycolic acid and wherein the SH-reagent is selected from Ellman's reagent, mercapto forming agents, p-mercuri benzoic acid, alkylating agents, iodoacetic acid and N-ethylmaleimide.

45 **[0006]** In a preferred embodiment, the step (a) is a step of reacting D-alanine present in a sample with D-alanyl-D-alanine ligase (which may be referred to "Ddl" hereinafter) in a presence of adenosine triphosphate to convert the D-alanine into D-alanyl-D-alanine and/or a step of reacting D-serine present in a sample with D-serine dehydratase (which may be referred to "Dsd" hereinafter) to convert the D-serine into pyruvic acid.

[0007] In a preferred embodiment, the methyltransferase is homocysteine methyltransferase, and the methyl donor is D-methionine methylsulfonium.

[0008] In a preferred embodiment, in the step (d), the produced hydrogen peroxide is detected or measured by color-development using peroxidase and an oxidative color-developing agent.

55 **[0009]** The present invention also provides a reagent kit for measuring homocysteine comprising D-alanyl-D-alanine ligase and/or D-serine dehydratase; a thiol compound; a methyltransferase; D-methionine methylsulfonium; D-amino acid oxidase or D-amino acid acetyltransferase and acylcoenzyme A synthetase and acylcoenzyme A oxidase; an SH reagent; and an oxidative color-developing agent wherein the thiol compound is selected from dithiothreitol, mercap-

toethanol, N-acetylcysteine, dithioerythritol and thioglycolic acid and wherein the SH-reagent is selected from Ellman's reagent, mercapto forming agents, p-mercuri benzoic acid, alkylating agents, iodoacetic acid and N-ethylmaleimide.

Brief Description of Drawings

[0010]

Fig. 1 is a schematic diagram of the reaction in a method of measuring homocysteine using homocysteine transferase and D-methionine methylsulfonium.

Fig. 2 is a schematic diagram showing construction of expression vector pKdIA.

Fig. 3 is a schematic diagram showing construction of expression vector pKdIB.

Fig. 4 is a graph showing a relationship between the concentration of D-alanyl-D-alanine ligase and the sensitivity of the homocysteine measurement.

Fig. 5 is a graph showing a relationship between the concentration of D-serine dehydratase and the sensitivity of the homocysteine measurement.

Fig. 6 are graphs showing a correlation between the concentration of the measured homocysteine according to (a) a conventional single channel method, (b) a conventional double channel method, and (c) the method of the present invention and the concentration measured by a HPLC method.

Best Mode for Carrying Out the Invention

Principle of homocysteine measurement:

[0011] The method of measuring homocysteine of the present invention is based on the principle that homocysteine in a sample is subjected to a reduction treatment with a thiol compound, and is reacted with a methyltransferase in the presence of a methyl donor (first process), and then the produced D-amino acid or D-amino acid derivative is measured (second process). For example, as shown in Fig. 1, when D-methionine produced in the first process is reacted with D-amino acid oxidase in the second process, hydrogen peroxide is produced, and the hydrogen peroxide can be led to an oxidative color-developing agent commonly used in the presence of an SH reagent so as to be determined colorimetrically. When the D-methionine is reacted with D-amino acid acetyltransferase, the produced coenzyme A is led to hydrogen peroxide using acyl coenzyme A synthetase [EC 6.2.1.3] and acyl coenzyme A oxidase [EC 1.3.3.6], and this can be determined in the same manner.

[0012] The method of detecting or measuring homocysteine of the present invention is characterized in that in order to eliminate the influence of endogenous D-amino acids when measuring D-amino acids, first, a D-amino acid present in a sample is reacted with a D-amino acid converting enzyme to convert the D-amino acid to a substance that does not serve as a substrate of D-amino acid oxidase or D-amino acid acetyltransferase.

[0013] More specifically, the method of detecting or measuring homocysteine of the present invention includes the steps of:

- (a) reacting a D-amino acid present in a sample with a D-amino acid converting enzyme to convert the D-amino acid to a substance that does not serve as a substrate of D-amino acid oxidase or D-amino acid acetyltransferase;
- (b) reducing homocysteine in the sample with a thiol compound;
- (c) reacting the reduced homocysteine with a methyltransferase and D-methionine methylsulfonium to newly produce D-methionine; and
- (d) reacting the D-methionine produced in step (c) with the D-amino acid oxidase in the presence of an SH reagent to produce hydrogen peroxide, and color-developing the produced hydrogen peroxide by using an oxidative color-developing agent, or
- (d') reacting the D-methionine produced in step (c) with the D-amino acid acetyltransferase in the presence of acetyl coenzyme A thereby producing coenzyme A; reacting the produced coenzyme A with acyl coenzyme A synthetase thereby producing acyl coenzyme A; and reacting the produced acyl coenzyme A with acyl coenzyme A oxidase in the presence of an SH reagent to produce hydrogen peroxide; and color-developing the produced hydrogen peroxide by using an oxidative color-developing agent, wherein the thiol compound is selected from dithiothreitol, mercaptoethanol, N-acetylcysteine, dithioerythritol and thioglycolic acid and wherein the SH-reagent is selected from Ellman's reagent, mercapto forming agents, p-mercuri benzoic acid, alkylating agents, iodoacetic acid and N-ethylmaleimide.

[0014] Any sample can be used as a sample to be subjected to detect or measure homocysteine by the method of the present invention, as long as it is believed to contain homocysteine. The homocysteine can be present in the form

of, not only reduced homocysteine, but also oxidized homocysteine that is bound to another molecule by a disulfide bond such as a complex with a protein, a homocysteine dimer and a homocysteine-cysteine dimer. For example, serum, plasma, blood, urine and a dilution thereof can be used.

Step (a):

[0015] As the D-amino acid converting enzyme that is used in the method of the present invention, any D-amino acid converting enzyme can be used, as long as it can react with a D-amino acid to convert the D-amino acid into a substance that does not serve as a substrate of D-amino acid oxidase or D-amino-acid acetyltransferase so as to lead it to the outside of the reaction system of the homocysteine measurement. In the present invention, a D-amino acid converting enzyme that can react with D-alanine and/or D-serine is preferable. As an enzyme that reacts with D-alanine, D-alanyl-D-alanine ligase [EC 6.3.2.4], D-alanine hydroxymethyltransferase [EC 2.1.2.7], D-alanine- γ -glutamyl-transferase [EC 2.3.2.14], or the like can be used. As an enzyme that reacts with D-serine, D-serine dehydratase [EC 4.3.1.18], diaminopropionate ammonia-lyase [EC 4.3.1.18], or the like can be used. These enzymes can be used alone or in combination if necessary. As the enzyme that reacts with D-alanine, D-alanyl-D-alanine ligase is preferably used, and as the enzyme that reacts with D-serine, D-serine dehydratase is preferably used.

[0016] Any D-alanyl-D-alanine ligase (Ddl) derived from any sources can be used in the method of the present invention, enzymes, as long as it can condense two D-alanine molecules to produce D-alanyl-D-alanine. For example, Ddls derived from almost all bacteria can be utilized. Enzymes derived from *E. coli* are preferable. "E. coli" in this specification refers to *Escherichia coli* described in Bergey's Manual of Determinative Bacteriology, the 8th edition (edited by R.E. Buchanan and N.E. Gibbons, The Williams & Wilkins Company, Baltimore, pp. 295-296, 1974) and its variants and modified forms.

[0017] As Ddl used in the method of the present invention, enzymes derived from *E. coli* having an amino acid sequence estimated from the base sequence of GenBank Accession No. J05319 described in Biochemistry 30:1673-1682 (1991) or GenBank Accession No. AE000118 REGION: 18688., 19608 described in Journal of Bacteriology 167: 809-817 (1986) are preferably used. Furthermore, enzymes derived from microorganisms belonging to the genus of *Bacillus*, *Enterococcus*, *Lactobacillus* and the like also can be used. The sequence can be modified in some amino acids (e.g., addition, deletion or Substitution of one or more amino acids), as long as the Ddl activity does not disappear. The Ddl can be obtained by any methods well-known to those skilled in the art, for example, by a method of preparing a crude enzyme from the *E. coli* (e.g., an *E. coli* strain that has been transformed by introducing ddl gene obtained from bacteria or the like) and then purifying it by, for example, various chromatography techniques.

[0018] The D-serine dehydratase (Dsd) used in the method of the present invention is an enzyme also called D-serine ammonia-lyase, D-serine dehydrase, D-hydroxyamino acid dehydratase, D-serine hydrase, or D-serine deaminase. Dsd derived from any sources can be used, as long as it can deaminate D-serine to produce pyruvic acid. Enzymes derived from *E. coli* having an amino acid sequence estimated from the base sequence of GenBank Accession No. J01603 described in J. Bacteriol. 154(3), 1508-1512 (1983) are preferably used. Furthermore, Dsds derived from microorganisms belonging to the genus of *Pseudomonas*, *Bacillus*, *Salmonella*, *Fusobacterium*, *Vibrio*, *Shigella*, *Ralstonia* and the like can be used. The sequence can be modified in some amino acids (e.g., addition, deletion or substitution of one or more amino acids), as long as the Dsd activity does not disappear. The Dsd can be obtained by any methods well-known to those skilled in the art, for example, by a method of preparing a crude enzyme from *E. coli* (e.g., an *E. coli* strain that has been transformed by introducing dsd gene obtained from bacteria or the like) and then purifying it by, for example, various chromatography techniques.

[0019] The ddl gene and the dsd gene can be obtained by a method commonly used by those skilled in the art, based on the estimated amino acid sequences described above. For example, methods performing plaque hybridization, colony hybridization, PCR or the like can be employed, where a part or all of the genes encoding Ddl or Dsd or genes containing these sequences are used as probes. The gene source of Ddl or Dsd is not limited to *E. coli*, but other species of bacteria can be used as well.

[0020] In this specification, "ddl gene" refers to a DNA chain or DNA sequence that encodes a polypeptide having the Ddl activity that exhibits the characteristics of Ddl, and "dsd gene" refers to a DNA chain or DNA sequence that encodes a polypeptide having the Dsd activity that exhibits the characteristics of Dsd. In both cases, a polypeptide having a modification (e.g., addition, deletion or substitution) in the amino acid sequence that does not affect the activity of the enzyme as described above may be encoded. Moreover, a plurality of sequences may encode the same polypeptide, for example, due to degeneracy. Furthermore, the ddl gene or the dsd gene may be derived from natural sources or may be totally synthesized or semi-synthesized chemically.

[0021] The obtained ddl gene or the dsd gene is, for example, ligated to an expression vector that can multiply and is introduced into a host such as *E. coli*. The expression vector used herein may be any expression vector, as long as it can be usually used for *E. coli*, and for example, ColE1, pCR1, pBR322, pMB9 and the like can be preferably used.

[0022] In order to express a large amount of DNA encoding Ddl or Dsd in *E. coli*, or to increase the expression amount thereof, a promoter for controlling transcription and translation may be incorporated into a 5' upstream region of the DNA

chain of the vector, and/or a terminator may be incorporated into a 3' downstream region thereof. Such a promoter and/or a terminator may be derived from a *ddl* gene or *dsd* gene itself, derived from a gene that is well known such as β -galactosidase gene, or obtained by artificially modifying these known genes. Therefore, expression vectors in which such control sequences are incorporated are preferably used as the expression vector, and examples thereof include

pTrc99A, pKK223-3 (which are manufactured by Amersham Pharmacia Biotech), and pET-3, pET-11 (which are manufactured by Stratagene), although there is no limitation to these vectors.

[0023] Any microorganisms can be used as a host for expressing Ddl or Dsd, but bacteria are preferable and *E. coli* is more preferable. A transformant for expressing Ddl or Dsd can be produced by a method known in the field of genetic engineering, for example, by a rubidium chloride method (J. Mol. Biol., 166:557, 1983). The thus obtained transformant of *E. coli* whose ability of expressing Ddl or Dsd is increased is cultured so that Ddl or Dsd can be obtained.

[0024] The Ddl or Dsd obtained in the above-described manner may be used alone or in combination.

[0025] When Ddl is applied to a reagent for measuring homocysteine, the produced D-alanyl-D-alanine can not be a substrate of D-amino acid oxidase, so that it is possible to eliminate the D-alanine contained in a sample. There is no limitation regarding the amount of enzyme used, as long as it has a concentration that can eliminate the D-alanine in the sample, and for example, the amount can be 0.01 to 100 U/mL, and preferably is 0.1 to 10 U/mL. In the present invention, the amount of Ddl that produces 1 μ mol of D-alanyl-D-alanine per minute at 37°C, using D-alanine as the substrate is defined as 1 unit. There is also no limitation regarding the amounts of ATP and Mg ions to be used that is necessary to exhibit the activities, as long as they have a concentration that can eliminate D-alanine, and for example, for ATP, an amount of 0.1 to 10 mM, and for Mg ions, an amount of 0.1 to 20 mM are preferable. This treatment of a sample with enzymes can be performed alone, or can be performed simultaneously with the following steps (b) and (c).

[0026] When Dsd is used as a reagent for measuring homocysteine, D-serine can be removed from the reaction system by converting D-serine contained in a sample into pyruvic acid. There is no limitation regarding the amount of enzyme used, as long as it has a concentration that can eliminate D-serine in the sample, and for example, the amount can be 0.001 to 10 U/mL, and preferably is 0.01 to 1 U/mL. In the present invention, the amount of Dsd that degrades 1 μ mol of D-serine per minute at 37°C is defined as 1 unit. This treatment of a sample with enzymes can be performed alone, or can be performed simultaneously with the following steps (b) and (c).

Step (b):

[0027] The step (b) in the method of the present invention is a step of reducing homocysteine in various forms in a sample with a thiol compound into a reduced homocysteine.

[0028] The thiol compound used in the method of present invention is selected from dithiothreitol, mercaptoethanol, N-acetylcysteine, dithioerythritol, and thioglycolic acid. Any concentration of the thiol compound may be employed as the concentration of the thiol compound, as long as it is in a range that allows oxidized homocysteine to reduced homocysteine. Preferably, the concentration is 0.1 mM or more in terms of thiol groups, and more preferably 1 mM or more.

Step (c):

[0029] The step (c) in the method of the present invention is a step of reacting the reduced homocysteine made in the step (b) above with a methyltransferase and a methyl donor so as to newly produce D-amino acid. In the present invention, a methyltransferase using homocysteine as a methyl acceptor is preferably used, and D-methionine methylsulfonium is used as a methyl donor. That is, the reduced homocysteine in the sample is reacted with a methyltransferase and D-methionine methylsulfonium to produce D-methionine.

[0030] There is no limitation regarding the methyltransferase, as long as it reacts with D-methionine methylsulfonium to catalyze the production of D-methionine. Examples of the methyltransferase include homocysteine methyltransferase [EC 2.1.1.10], 5-methyltetrahydrofolic acid-homocysteine S-methyltransferase [EC 2.1.1.13], and 5-methyl-tetrahydropteroyltriglutamic acid-homocysteine S-methyltransferase [EC 2.1.1.14]. Preferably, homocysteine methyltransferase [EC 2.1.1.10] can be used. This enzyme produces D-methionine, using D-methionine methylsulfonium as a methyl donor, in spite of its low specificity, as reported by G. Grue-Sorensen et al (J. Chem. Soc. Perkin Trans. I 1091-7 (1984)). The homocysteine methyltransferase to be used may be derived from any sources, as long as it uses D-methionine methylsulfonium as a methyl donor. For example, enzymes derived from bacteria, yeasts, rats or the like can be used. In the present invention, the amount of the homocysteine methyltransferase that produces 1 μ mol of D-methionine per minute at 37°C, using homocysteine and D-methionine methylsulfonium as the substrates, is defined as 1 unit.

Step (d) or (d')

[0031] The step (d) or (d') in the method of the present invention is a step of reacting the D-amino acid produced in the step (c) with D-amino acid oxidase or D-amino acid acetyltransferase in the presence of an SH reagent to produce

hydrogen peroxide, and color-developing the produced hydrogen peroxide by using an oxidative color-developing agent.

[0032] When D-amino acid is reacted with D-amino acid oxidase [EC 1.4.3.3], hydrogen peroxide is produced. This is led to an oxidative color-developing agent commonly used in the presence of an SH reagent so as to be determined colorimetrically. The D-amino acid oxidase may be derived from any sources. For example, it may be derived from animal organs, bacteria, or fungi. Preferably, those derived from porcine kidney can be used. In the present invention, the amount of the D-amino acid oxidase that converts 1 μ mol of D-alanine into pyruvic acid per minute at 37°C is defined as 1 unit.

[0033] When D-amino acid is reacted with D-amino acid acetyltransferase [EC 2.3.1.36], coenzyme A is produced. Coenzyme A is reacted with acyl-coenzyme A synthetase [EC 6.2.1.3] and acyl-coenzyme A oxidase [EC 1.3.3.6], sequentially, then hydrogen peroxide is produced. The produced hydrogen peroxide can be quantitatively determined in the same manner. The D-amino acid acetyltransferase may be derived from any sources. For example, those derived from yeasts can be used.

[0034] The SH reagent is selected from Ellman's reagent, a mercapto forming agent such as p-mercuri benzoic acid, and an alkylating agent such as iodoacetic acid and N-ethylmaleimide, as described in Biochemistry Dictionary (3rd edition, p.182, Tokyo Kagaku Doujin, 1998). Preferably, an alkylating agent, and more preferably a maleimide compound, and most preferably, N-ethylmaleimide can be used.

[0035] The produced hydrogen peroxide can develop the color of an ordinary oxidative color-developing agent with peroxidase. As the oxidative color-developing agent, various Trinder reagents can be used in combination with a coupler reagent. This method is called the Trinder method, and is commonly used in the field of clinical chemical analysis, which is not described herein in detail. It is preferable to use 4-aminoantipyrine as the coupler reagent and to use ADOS [N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3-methoxyaniline], DAOS [N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3,5-dimethoxyaniline], HDAOS [N-(2-hydroxy-3-sulfopropyl)-3,5-dimethoxyaniline], MAOS [N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3,5-dimethylaniline], TOOS [N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3-methylaniline] or the like as the Trinder reagent. Furthermore, a leuco-type color-developing agent such as o-tolidine, o-dianisidine, DA-67 [10-(carboxymethylaminocarbonyl)-3,7-bis(dimethylamino)phenothiazine sodium, manufactured by Wako Pure Chemical Industries Ltd.], and TPM-PS [N,N,N',N'',N''',N''-hexa(3-sulfopropyl)-4,4',4''-tri-aminotriphenylmethane hexasodium salt, Dojindo Laboratories], which do not require the coupler reagent, can be used as well. In particular, DA-67 and TPM-PS have a mole absorption coefficient larger than that of the Trinder reagent, so that the determination can be performed with higher sensitivity.

Reagent kit for homocysteine measurement:

[0036] The present invention provides a reagent kit for homocysteine measurement including (a) D-alanyl-D-alanine ligase and/or D-serine hydratase, (b) a thiol compound, (c) a methyltransferase and D-methionine methylsulfonium, and (d) D-amino acid oxidase or D-amino acid acetyltransferase + acyl-coenzyme A synthetase + acyl-coenzyme A oxidase, an SH reagent, and an oxidative color-developing agent wherein the thiol compound is selected from dithiothreitol, mercaptoethanol, N-acetylcysteine, dithioerythritol and thioglycolic acid and wherein the SH-reagent is selected from Ellman's reagent, mercapto forming agents, p-mercuri benzoic acid, alkylating agents, iodoacetic acid and N-ethylmaleimide. In general, these (a) to (d) are provided separately, but (a) to (c) may be provided as a reagent for measurement that is previously prepared by mixing in a buffer solution.

[0037] Hereinafter, the present invention will be described more specifically by way of examples, but the present invention is not limited by the following examples.

[Example 1] Preparation of recombinant D-alanyl-D-alanine ligase (A) (Dd1A) derived from E. coli.

(1-1) Synthesis of probes and obtainment of ddIA gene

[0038] Synthetic primers shown in SEQ ID NOS: 1 and 2, which include EcoRI and PstI recognition sites, respectively, were synthesized based on the base sequence information of the ddIA gene from E. coli encoding an enzyme having DdIA activity (Biochemistry 30:1673-1682 (1991), GenBank Accession No. J05319). PCR was performed in a buffer solution (2 μ l of KOD DNA polymerase (manufactured by TOYOBO Co., Ltd.; hereinafter, referred to as "KOD"), 10 μ l of 10 x KOD buffer solution, 10 μ l of dNTP mixture, 5 μ l of DMSO and 5 μ l of distilled water), using these primers in an amount of 3 nmol (100 pmol/ μ l, 30 μ l) each, and using 2 μ l of chromosomal DNA of E. coli JM109 as a template to give 1.09 kb DNA containing the DdIA structural gene.

(1-2) Preparation of plasmid including ddIA gene

[0039] The thus obtained ddIA gene was ligated to a SmaI fragment of E. coli vector pUC19 having DNA replication origin of E. coli ColE1 and ampicillin resistance gene, and introduced in E. coli JM109 by the rubidium chloride method

(J. Mol. Biol., 166:557, 1983) to give a transformant having a recombinant plasmid containing the *ddlA* gene. All the restriction enzymes used in the Examples were obtained from TAKARA BIO INC.

[0040] Using the recombinant plasmid in the transformant, the base sequence was determined by using 377 Automate Sequencing System (manufactured by PerkinElmer Co., Ltd.) based on the dideoxy terminator method using fluorescent-labeled primers. The *ddlA* has a structural gene region with 1095 bases, and encodes 364 amino acids, which was completely consistent with the above-described base sequence information on the *ddlA* gene.

(1-3) Preparation of expression vector and transformant including *ddlA* gene

[0041] The above-described recombinant plasmid was digested with restriction enzymes *EcoRI* and *PstI* to give a DNA fragment, and then the 1.09 kb DNA fragment containing the *ddlA* DNA was purified by agarose gel electrophoresis. *E. coli* vector pKK233*3 (manufactured by Amersham Pharmacia Biotech) having ampicillin resistance gene was digested with restriction enzymes *EcoRI* and *PstI*, and ligated to the obtained 1.09 kb DNA fragment to give an expression vector pKdIA (see Figure 2). This expression vector pKdIA is induced to express DdIA by isopropyl- β -D-thiogalactopyranoside (hereinafter, referred to as "IPTG").

[0042] The obtained expression vector pKdIA was introduced into *E. coli* JM109 by the rubidium chloride method, and one in which DdIA is expressed was selected to give a transformant JM109-*ddlA*-3.

(1-4) Confirmation of activity of expressed DdIA

[0043] The obtained transformant JM109-*ddlA*-3 was cultured in 3 ml of LB liquid medium (1% yeast extract, 2% bactopectone, and 2% glucose) containing ampicillin at 37°C for about four hours with shaking. A 0.3 ml aliquot was added to 10 ml of LB liquid medium and cultured at 37°C for three hours with shaking, and IPTG (TAKARA BIO INC.) was added thereto such that the final concentration became 1 mM, and was cultured further for four hours. The culture was centrifuged at 8000 rpm for 15 minutes to collect cells. Then, after the cells were washed with 1 ml of 100 mM Bis-tris-HCl buffer solution (pH 7.4) once, they were suspended in a solubilization solution (100 mM Bis-tris-HCl buffer solution (pH 7.4), 1 mM EDTA, 5 mM MgCl₂, and 100 μ g/ml lysozyme) in a volume ten times that of the cells. The Suspension was treated twice with an ultrasonic generator (UD-200 manufactured by TOMY SEIKO Co., Ltd.) at scale 1 for 10 seconds to disrupt the cells. A supernatant was obtained by centrifugation at 15000 rpm for 10 minutes, and this was used as a sample for measuring the Ddl activity. As a control, a transformant of *E. coli* JM109 strain with non-recombinant pKK223-3 that had been treated in the same manner was used.

[0044] The enzyme activity was measured in the following manner. First, a reagent for measuring the activity (100 μ l of 20 mM D-alanine, 20 mM ATP, 100 mM HEPES, 40 mM MgCl₂, and 50 μ l of 40 mM KCl) was added to the disrupted cell solution (50 μ l), and allowed to react at 37°C for one hour. Then, 2 μ l of the reaction mixture was spotted onto a silica gel thin layer, and then developed using ethanol : 25% aqueous ammonia = 74 : 26 (w/w) as an eluent in a closed vessel. After development, a ninhydrin solution (0.1 M citric acid buffer solution saturated with n-butanol, which contains 0.2% of ninhydrin) was sprayed to detect the produced D-alanyl-D-alanine. The D-alanyl-D-alanine was produced in the disrupted recombinant cell solution in a significantly larger amount compared to the disrupted control cell solution.

(1-5) Preparation of DdIA

[0045] The recombinant *E. coli* obtained in the above steps (1-3), which has high production ability of DdIA, was inoculated in 200 ml of LB medium (1% yeast extract, 2% bactopectone, and 2% glucose) containing ampicillin and was cultured at 37°C for 15 hours. Then, the culture was inoculated to 1.8 L of LB medium containing ampicillin in a jar fermenter with a volume of 5 L, and cultured at 37°C for 100 minutes with aeration and stirring. To the culture, IPTG was added such that the final concentration became 1 mM, and cultured further for 4 hours. The culture was centrifuged at 8000 rpm for 10 minutes to collect cells. The cells were suspended in a buffer solution (20 mM Bis-tris-HCl buffer solution (pH 7.4), 1 mM EDTA, and 5 mM MgCl₂) in a volume nine times that of the cells. The suspension was treated with an ultrasonic generator (UD-200 manufactured by TOMY SEIKO CO., Ltd.) to disrupt the cells. Then, the suspension was centrifuged at 15000 rpm for 10 minutes to remove the debris and thus a crude enzyme solution was obtained.

[0046] Ammonium sulfate was added to the obtained crude enzyme solution so as to be 5% saturation under ice cooling with stirring, and thereafter was allowed to stand for 30 minutes, and then was centrifuged at 14000 rpm. Ammonium sulfate was added to the obtained supernatant so as to be 45% saturation under ice cooling with stirring, and thereafter allowed to stand for 30 minutes, and then centrifuged at 14000 rpm. The obtained precipitate was dialyzed overnight against a buffer solution (20 mM Bis-tris-HCl buffer solution (pH 7.4), 1 mM EDTA, and 5 mM MgCl₂).

[0047] Then, the dialyzed crude enzyme solution was purified by column chromatography (adsorption: a buffer solution A (20 mM Bis-tris-HCl buffer solution (pH 7.4), 1 mM EDTA, and 5 mM MgCl₂), elution: a buffer solution A - 0.0 to 0.6 M sodium chloride gradient) using Q-Sepharose FF (manufactured by Amersham Pharmacia Biotech). The Ddl active

fractions were collected and further purified by gel filtration column chromatography with Sephacryl S-100 (manufactured by Amersham Pharmacia Biotech). The obtained active fractions were subjected to SDS-polyacryl amide gel electrophoresis and stained by Coomassie brilliant blue, which confirmed that the DdlA was purified to substantially a single band.

[Example 2] Preparation of recombinant D-alanyl-D-alanine ligase (B) (DdlB) derived from *E. coli*

(2-1) Synthesis of probe and acquisition of ddIB gene

[0048] Synthetic primers shown in SEQ ID NOS: 3 and 4 were synthesized, based on the base sequence information of the ddIB gene of *E. coli* encoding an enzyme having the DdlB activity (Journal of Biochemistry 167:809-817 (1986), GenBank Accession No. AE000118 REGION: 18688.. 19608). PCR was performed in a buffer solution (2 μ l of KOD, 10 \times 1 of 10 \times KOD buffer solution, 10 μ l of dNTP mixture, 5 μ l of DMSO and 5 μ l of distilled water), using these primers in an amount of 3 nmol (100 pmol/ μ l, 30 μ l) each, and using 2 μ l of chromosome DNA of *E. coli* JM109 as a template to give 0.92 kb DNA containing the DdlB structural gene.

(2-2) Preparation of plasmid including ddIB gene

[0049] The thus obtained ddIB gene was ligated to a SmaI fragment of an *E. coli* vector pUC19 having DNA replication origin of *E. coli* ColEI and an ampicillin resistance gene, and introduced into *E. coli* JM109 by the rubidium chloride method (J. Mol. Biol., 166:557, 1983) to give a transformant having a recombinant plasmid containing the ddIB gene.

[0050] Using the recombinant plasmid in the transformant, the base sequence was determined by using 377 Automate Sequencing System (manufactured by PerkinElmer Co., Ltd.) based on the dideoxy terminator method using fluorescent-labeled primers. The ddIB has a structural gene region with 921 bases, and encodes 306 amino acids, which was completely consistent with the base sequence information on the ddIB gene.

(2-3) Preparation of expression vector and transformant including ddIB gene

[0051] An *E. coli* vector pKK233-3 (manufactured by Amersham Pharmacia Biotech) was digested with restriction enzymes EcoRI, and blunt-ended using DNA Blunting Kit (manufactured by TAKARA BIO INC.), then dephosphorylated using alkaline phosphatase (manufactured by TAKARA BIO INC.), and a fragment of the DdlB structural gene obtained in (2-1) was ligated thereto to give an expression vector pKdlB (see Figure 3). This expression vector pKdlB is induced to express DdlB by IPTG.

[0052] The obtained expression vector pKdlB was introduced into *E. coli* JM109 by the rubidium chloride method, and one in which DdlB is expressed was selected to give a transformant JM109-ddIB-1.

(2-4) Preparation of DdlB

[0053] The recombinant *E. coli* obtained in the above steps (2-3), which has high production capacity of DdlB, was inoculated in 200 ml of LB medium (1% yeast extract, 2% bactopectone, and 2% glucose) containing ampicillin and was preliminarily cultured at 37°C for 15 hours. The culture was inoculated to 1.8 L of LB medium containing ampicillin in a jar fermenter with a volume of 5 L, and cultured at 37°C for 100 minutes with aeration and stirring. To the culture, IPTG was added such that the final concentration became 1 mM and the culture was cultured further for 4 hours. The culture was centrifuged at 8000 rpm for 10 minutes to collect the cells. The cells were suspended in a buffer solution (20 mM Bis-tris-HCl buffer solution (pH 7.2), 1 mM EDTA, and 5 mM MgCl₂) in a volume nine times that of the cells. The suspension was treated with an ultrasonic generator (UD-200 manufactured by TOMY SEIKO CO., Ltd.) to disrupt the cells. Then, the suspension was centrifuged at 15000 rpm for 10 minutes to remove the debris and thus a crude enzyme solution was obtained.

[0054] Ammonium sulfate was added to the obtained crude enzyme solution so as to be 25% saturation under ice cooling with stirring, and thereafter was allowed to stand for 30 minutes, and then centrifuged at 14000 rpm. Ammonium sulfate was added to the obtained supernatant so as to be 50% saturation under ice cooling with stirring, and thereafter allowed to stand for 30 minutes, and then centrifuged at 14000 rpm. The obtained precipitate was dialyzed overnight against a buffer solution (20 mM Bis-tris-HCl buffer solution (pH 7.2), 1 mM EDTA, and 5 mM MgCl₂).

[0055] Then, the dialyzed crude enzyme solution was purified by column chromatography (adsorption: a buffer solution A (20 mM Bis-tris-HCl buffer solution (pH 7.2), 1 mM EDTA, and 5 mM MgCl₂), elution: a buffer solution A - 0.0 to 0.6 M sodium chloride gradient) using Q-Sepharose FF (manufactured by Amersham Pharmacia Biotech). The Ddl active fractions were collected and further purified by gel filtration column chromatography with Sephacryl S-100 (manufactured by Amersham Pharmacia Biotech). The obtained active fractions were subjected to SDS-polyacryl amide gel electrophoresis and stained by Coomassie brilliant blue, which confirmed that the DdlB was purified to substantially a single band.

[Example 3] Preparation of recombinant D-serine dehydratase (Dsd) derived from *E. coli*

(3-1) Preparation of expression vector and transformant including *dsd* gene

[0056] The following two synthetic primers (SEQ ID NOS: 5 and 6) were synthesized, based on the base sequence information of the *dsd* gene of *E. coli* encoding an enzyme having the Dsd activity.

[0057] PCR was performed, using these primers and using genome DNA of *E. coli* as a template. The obtained DNA fragment was digested with *EcoRI* and *HindIII*, and ligated to a vector PUC118 that had been digested with the same restriction enzymes. This was introduced into *E. coli* JM109 by the rubidium chloride method to give a transformant having a recombinant plasmid containing the *dsd* gene.

(3-2) Preparation of Dsd

[0058] The Dsd expressing cell strain obtained in the above-described manner was cultured in about 10 L of LB medium containing 100 µg/ml of ampicillin. Enzyme purification was performed according to the method described in J. Biol. Chem., 263, 16926-16933, 1988. The cell paste was suspended in SPE lysis buffer (20% sucrose, 20 mM EDTA, 30 mM potassium phosphate, 0.5 mg/mL lysozyme, pH7.8) in almost the same volume. After incubation, water containing a protease inhibitor and pyridoxal 5'-phosphorate (PLP) was added thereto for lysis. The lysate was cooled with ice and a buffer solution B (1M potassium phosphate, 800 µM PLP, 50 mM EDTA, 10 mM DTT, pH7.5) was added thereto in almost the same volume, and the cell debris was removed by centrifugation at 18000 rpm for 30 minutes. The obtained supernatant was adjusted to pH 7.3, and nucleic acid was precipitated by 1% Polymyxin P and then centrifuged to give supernatant. Ammonium sulfate was added thereto and dissolved therein so as to be 70% saturation. Then, the solution was stirred for 40 minutes and then centrifuged at 18000 rpm for two hours. The obtained precipitate was suspended in a small amount of buffer solution C (10mM potassium phosphate, 80 µM PLP, 1 mM EDTA, 1 mM DTT, pH 7.2) and dialyzed overnight against the same buffer solution.

[0059] Then, the dialyzed crude enzyme solution was purified by column chromatography using DEAE-Toyopearl (manufactured by Tosoh Corporation). For adsorption, the buffer solution C was used, and elution was performed using 0 to 200 mM KCl gradient in this buffer solution. The Dsd active fractions were collected and precipitated with 70% saturated ammonium sulfate and recovered. The recovered fractions were dissolved in a small amount of the buffer solution C, and dialyzed overnight against the same buffer solution to remove ammonium sulfate. Furthermore, they were dialyzed against a buffer solution D (1mM potassium phosphate, 1 mM DTT, pH7.0) for 3 to 4 hours.

[0060] The dialyzed partially purified enzyme solution was further purified by hydroxyapatite column chromatography (Gigapite, manufactured by TOAGOSEI CO., LTD., available from SEIKAGAKU CORPORATION). This enzyme solution was applied to a column previously equilibrated with the buffer solution D, washed with the same buffer solution in a volume three times the column, and eluted with a buffer solution E (10 mM potassium phosphate, 80 µM PLP, 1 mM EDTA, pH7.8). A buffer solution in a volume of 1/10 was added to each eluted fraction, and Dsd active fractions were collected, precipitated with 70% saturated ammonium sulfate and recovered. The obtained precipitate was suspended in a small amount of buffer solution F (100 mM potassium phosphate, 80 µM PLP, 1 mM EDTA, 1 mM DTT, pH7.8), and dialyzed against this buffer solution to give a purified enzyme.

(3-3) Measurement of Dsd activity

[0061] The Dsd activity was determined by measuring coupling of the pyruvic acid produced from D-serine with lactate dehydrogenase in the presence of NADH. More specifically, 0.01 to 0.1 U Dsd was added to 100 mM D-serine, 0.5 mM NADH and 5 U lactate dehydrogenase at 37 °C to initiate a reaction, and a decrease of the absorbance at 340 nm was traced.

[Example 4] Prevention of influence of D-alanine by DdIA, DdIB and Dsd

[0062] Samples (1 to 5) were prepared in the following manner:

- Sample 1: Normal control blood serum SERACLEAR HE (manufactured by Azwell Co. Ltd.)
- Sample 2: Addition of 25 µM of homocystine (corresponding to 50 µM of homocysteine) to the sample 1
- Sample 3: Addition of 100 µM of D-alanine to the sample 2
- Sample 4: Addition of 500 µM of D-serine to the sample 2
- Sample 5: Addition of 100 µM of D-alanine and 500 µM of D-serine to the sample 2

[0063] First reagents (I to V) and a second reagent were prepared in the following manner:

First reagent:

Reagent I: 50 mM Bicine (pH 8.0), 123 U/L homocysteine methyltransferase (derived from bacteria), 5.6 mM dithiothreitol, 0.06 mM D-methionine methylsulfonium, 1 mM zinc bromide, 0.3 mM DA-67, 1 mM ATP, 1 mM magnesium chloride

Reagent II: Addition of 0.2285 mg protein/mL of DdlB to the reagent I

Reagent III: Addition of 2.0875mg protein/mL of DdlA to the reagent I

Reagent IV: Addition of 1 U/mL of Dsd to the reagent I

Reagent V: Addition of 0.2285 mg protein/mL of DdlB and 1U / mL of Dsd to the reagent I at the same time

Second reagent:

A reagent containing 50 mM citric acid (pH 5.6), 23 mM NEM, 6.4 U/mL D-amino acid oxidase derived from porcine kidney, and 5.5 U/mL peroxidase

[0064] Measurement was performed using Hitachi 7170 in the following manner. Any one of the first reagents in an amount of 180 μ L was added to each of the samples 1 to 5 in an amount of 15 μ L, and allowed to react at 37°C for 5 minutes. Then, 120 μ L of the second reagent was added thereto, and further allowed to react at 37°C for 5 minutes. The absorbance change (dominant wavelength 660 nm, secondary wavelength 750nm) from the detection point 16th to 34th was measured.

[0065] First, the absorbance change for the samples 1 and 2 were measured with the reagents I to V, respectively, and the measurement sensitivity with respect to the homocysteine added to the samples in each case was taken as 100 %. Then, the absorbance change for the samples 3 to 5 were measured with the reagents I to V. The measurement sensitivities of the samples 3 to 5 were shown in Table 1.

Table 1

		Sample 3	Sample 4	Sample 5
		D-Ala	D-Ser	D-Ala+D-Ser
Reagent I	Control	148%	123%	169%
Reagent II	D-alanyl-D-alanine ligase (B) (DdlB)	112%	119%	128%
Reagent III	D-alanyl-D-alanine ligase (A) (DdlA)	103%	117%	117%
Reagent IV	D-serine dehydratase (Dsd)	148%	100%	147%
Reagent V	DdlB + Dsd	112%	101%	112%

[0066] As seen from these results, in the control, because of the influence of D-alanine and D-serine, the measurement value was high, whereas when DdlB or DdlA was contained in the measurement reagent, the influence of D-alanine in the sample was reduced, and when Dsd was contained, the influence of D-serine in the sample was reduced. Furthermore, it is evident that when both are used simultaneously, the influence of D-alanine and D-serine can be reduced at the same time.

[Example 5] Elimination of D-alanine and D-serine by Ddl and Dsd

[0067] Samples (four types) were prepared in the following manner:

Sample 1: Normal control serum SERACLEAR HE (manufactured by Azwell Co. Ltd.)

Sample 2: Addition of 25 μ M of homocystine (corresponding to 50 μ M of homocysteine) to the sample 1.

Sample 6: Addition of 200 μ M of D-alanine to the sample 2

Sample 7: Addition of 1000 μ M of D-serine to the sample 2

[0068] First reagents and a second reagent were prepared in the following manner:

First reagents:

A reagent in which 0, 0.073, 0.145, 0.29 or 0.58 U/mL of DdlB or 0, 0.125, 0.25 or 0.5 U/ml of Dsd was added

to a reagent containing 50 mM Bicine (pH 8.0), 123 U/L homocysteine methyltransferase (derived from bacteria), 5.6 mM dithiothreitol, 0.06 mM D-methionine methylsulfonium, 1 mM zinc bromide, 0.3 mM DA-67, 5 mM ATP, and 10 mM magnesium chloride

Second reagent:

A reagent containing 50 mM citric acid (pH 5.6), 23 mM NEM, 6.4 U/mL D-amino acid oxidase derived from porcine kidney, and 5.5 U/mL peroxidase

[0069] Measurement was performed using Hitachi 7170 in the following manner. First, 180 μ L of first reagent was added to 15 μ L of the sample, and allowed to react at 37°C for 5 minutes. Then, 120 μ L of the second reagent was added thereto, and further allowed to react at 37°C for 5 minutes. The absorbance change (dominant wavelength 660 nm, secondary wavelength 750nm) from the detection point 16th to 34th was measured.

[0070] The results are shown in Figs. 4 and 5.

[0071] Fig. 4 shows the concentration of DdlB on the horizontal axis, and the relative sensitivity at the time of homocysteine measurement on the vertical axis. It was confirmed that, by using about 0.5 U/mL of DdlB, the influence of 200 mM D-alanine was substantially prevented.

[0072] Fig. 5 shows the concentration of Dsd on the horizontal axis, and the relative sensitivity at the time of homocysteine measurement on the vertical axis. By using about 0.2 U/mL of Dsd, the influence of 1000 mM D-serine was substantially prevented.

[Example 6] Prevention of influence of D-amino acid in a sample by Ddl and Dsd

[0073] Twelve samples of EDTA plasma were used as samples. As the standard, a sample in which 50 μ M D-methionine was added to a control blood serum was used.

[0074] Three first reagents (i to iii) and a second reagent (used in common) were prepared in the following manner:

First reagents:

Reagent i: 50 mM Bicine (pH 8.0), 126 U/L homocysteine methyltransferase (derived from bacteria), 5.6 mM dithiothreitol, 0.06 mM D-methionine methylsulfonium, 1 mM zinc bromide, 0.3 mM DA-67

Reagent ii: excluding the homocysteine methyltransferase from the reagent i

Reagent iii: Addition of 5 mM ATP, 10 mM magnesium chloride, 0.58 U/mL DdlB, and 1 U/mL Dsd to the reagent i

Second reagent:

A reagent containing 50 mM citric acid (pH 5.6), 23 mM NEM, 6.4 U/mL D-amino acid oxidase derived from porcine kidney, and 5.5 U/mL peroxidase

[0075] Measurement was performed using Hitachi 7170 in the following manner. Any one of the first reagents in an amount of 180 μ L was added to 15 μ L of the sample, and allowed to react at 37°C for 5 minutes. Then, 120 μ L of the second reagent was added thereto, and further allowed to react at 37°C for 5 minutes. The absorbance change (dominant wavelength 660 nm, secondary wavelength 750nm) from the detection point 16th to 34th was measured. The homocysteine concentration in the sample was obtained based on the absorbance change of the Standard. The value obtained as a result of measurement with only the reagent i was taken as the value according to the conventional single Channel method, the value obtained as a result of taking a difference between measurements with the reagent i and the reagent ii was taken as the value according to the conventional double Channel method, and the value obtained as a result of measurement with the reagent iii was taken as the value according to the method of the present invention. Each of these values was compared with the value measured according to the HPLC method.

[0076] As seen from Fig. 6, compared with the conventional Single Channel method, the method of the present invention has an improved correlation with the HPLC method, and exhibits a comparable correlation to the conventional double Channel method.

Industrial Applicability

[0077] According to the present invention, homocysteine can be measured without being affected by endogenous D-amino acid and without taking a sample blank. That is to say, homocysteine can be measured accurately by simple operations.

SEQUENCE LISTING

[0078]

5 <110> Azwell Inc.

 <120> A method of measuring homocysteine

 <130> P2-03A03191

10 <160> 6

 <170> Patent in version 3.1

15 <210> 1

 <211> 26

 <212> DNA

 <213> Artificial Sequence

20 <220>

 <223> primer

 <400> 1

25 gcgaattcat ggaaaaactg cgggta 26

 <210> 2

 <211> 26

 <212> DNA

 <213> Artificial Sequence

30 <220>

 <223> primer

 <400> 2

35 ggctgcagtt acattgtggt ttcaa 26

 <210> 3

 <211> 21

 <212> DNA

40 <213> Artificial Sequence

 <220>

 <223> primer

45 <400> 3

 atgactgata aaatcgcggt c 21

 <210> 4

 <211> 21

50 <212> DNA

 <213> Artificial Sequence

 <220>

 <223> primer

55 <400> 4

 ttagtccgcc agttccagaa t 21

<210> 5
 <211> 31
 <212> DNA
 <213> Artificial Sequence

5

<220>
 <223> primer

<400> 5
 ggtgaattca aagagacgta ctatggaaaa c 31

10

<210> 6
 <211> 30
 <212> DNA
 <213> Artificial Sequence

15

<220>
 <223> primer

<400> 6
 ggtaagcttt tgcgatgctg cgttgaaacg 30

20

Claims

25

1. A method for detecting or measuring homocysteine in a sample, comprising:

(a) reacting a D-amino acid present in a sample with a D-amino acid converting enzyme to convert the D-amino acid into a substance that does not serve as a substrate of D-amino acid oxidase or D-amino acid acetyltransferase;

30

(b) reducing homocysteine in the sample with a thiol compound;

(c) reacting the reduced homocysteine with a methyltransferase and D-methionine methylsulfonium to newly produce D-methionine;

and

35

(d) reacting the D-methionine produced in step (c) with the D-amino acid oxidase in the presence of an SH reagent to produce hydrogen peroxide, and color-developing the produced hydrogen peroxide by using an oxidative color-developing agent,

or

(d') reacting the D-methionine produced in step (c) with the D-amino acid acetyltransferase in the presence of acetyl coenzyme A thereby producing coenzyme A; reacting the produced coenzyme A with acyl coenzyme A synthetase thereby producing acyl coenzyme A; and reacting the produced acyl coenzyme A with acyl coenzyme A oxidase in the presence of an SH reagent to produce hydrogen peroxide; and color-developing the produced hydrogen peroxide by using an oxidative color-developing agent,

40

wherein the thiol compound is selected from dithiothreitol, mercaptoethanol, N-acetylcysteine, dithioerythritol and thioglycolic acid and,

45

wherein the SH-reagent is selected from Ellman's reagent, mercaptido forming agents, p-mercuri benzoic acid, alkylating agents, iodoacetic acid and N-ethylmaleimide.

2. The method of claim 1, wherein the step (a) is a step of reacting D-alanine present in a sample with D-alanyl-D-alanine ligase in the presence of adenosine triphosphate to convert the D-alanine into D-alanyl-D-alanine and/or a step of reacting D-serine present in a sample with D-serine dehydratase to convert the D-serine into pyruvic acid.

50

3. The method of claim 2, wherein the methyltransferase is homocysteine methyltransferase.

55

4. The method of claim 2 or 3, wherein in the step (d) or (d'), the produced hydrogen peroxide is detected or measured by color-development using peroxidase and an oxidative color-developing agent.

5. A reagent kit for measuring homocysteine comprising D-alanyl-D-alanine ligase and/or D-serine dehydratase; a thiol compound; a methyltransferase; D-methionine methylsulfonium; D-amino acid oxidase, or D-amino acid acetyltransferase and acyl coenzyme A synthetase and acyl coenzyme A oxidase; an SH reagent; and an oxidative color-developing agent,
 wherein the thiol compound is selected from dithiothreitol, mercaptoethanol, N-acetylcysteine, dithioerythritol and thioglycolic acid and,
 wherein the SH-reagent is selected from Ellman's reagent, mercaptido forming agents, p-mercuri benzoic acid, alkylating agents, iodoacetic acid and N-ethylmaleimide.

Patentansprüche

1. Ein Verfahren zum Nachweis oder Messen von Homocystein in einer Probe, umfassend:

- (a) Umsetzen einer D-Aminosäure, die in einer Probe vorhanden ist, mit einem D-Aminosäure-umsetzenden Enzym zur Umsetzung der D-Aminosäure zu einer Substanz, die nicht als ein Substrat der D-Aminosäure-Oxidase oder D-Aminosäure-Acetyltransferase dient;
 (b) Reduzieren von Homocystein in der Probe mit einer Thiolverbindung;
 (c) Umsetzen des reduzierten Homocysteins mit einer Methyltransferase und D-Methioninmethylsulfonium zur neuen Erzeugung von D-Methionin;
 und
 (d) Umsetzen des in Schritt (c) erzeugten D-Methionins mit der D-Aminosäure-Oxidase in Gegenwart eines SH-Reagenz zur Erzeugung von Wasserstoffperoxid; und Farbentwickeln des erzeugten Wasserstoffperoxids unter Verwendung eines oxidativen Farbentwicklungsmittels,
 oder
 (d') Umsetzen des in Schritt (c) erzeugten D-Methionins mit der D-Aminosäure-Acetyltransferase in Gegenwart von Acetyl-Coenzym A, wodurch Coenzym A erzeugt wird; Umsetzen des erzeugten Coenzym A mit Acyl-Coenzym-A-Synthetase, wodurch Acyl-Coenzym A erzeugt wird; und Umsetzen des erzeugten Acyl-Coenzym A mit Acyl-Coenzym-A-Oxidase in Gegenwart eines SH-Reagenz zur Erzeugung von Wasserstoffperoxid; und Farbentwickeln des erzeugten Wasserstoffperoxids unter Verwendung eines oxidativen Farbentwicklungsmittels,

wobei die Thiolverbindung aus Dithiothreitol, Mercaptoethanol, N-Acetylcystein, Dithioerythritol und Thioglycolsäure ausgewählt ist, und

wobei das SH-Reagenz aus Ellmanns-Reagenz, Mercaptid-bildenden Mitteln, p-Quecksilberbenzoesäure, Alkylierungsmitteln, Iodessigsäure und N-Ethylmaleimid ausgewählt ist.

2. Das Verfahren nach Anspruch 1, wobei der Schritt (a) ein Umsetzungsschritt von D-Alanin, das in einer Probe vorhanden ist, mit D-Alanyl-D-alanin-Ligase in Gegenwart von Adenosintriphosphat zur Umsetzung des D-Alanin zu D-Alanyl-D-alanin und/oder ein Umsetzungsschritt von D-Serin, das in einer Probe vorhanden ist, mit D-Serin-Dehydratase zur Umsetzung des D-Serin zu Brenztraubensäure ist.

3. Das Verfahren nach Anspruch 2, wobei die Methyltransferase Homocystein-Methyltransferase ist.

4. Das Verfahren nach Anspruch 2 oder 3, wobei in dem Schritt (d) oder (d') das erzeugte Wasserstoffperoxid durch Farbentwicklung unter Verwendung von Peroxidase und eines oxidativen Farbentwicklungsmittels nachgewiesen oder gemessen wird.

5. Ein Reagenz-Kit zum Messen von Homocystein umfassend D-Alanyl-D-alanin-Ligase und/oder D-Serin-Dehydratase; eine Thiolverbindung; eine Methyltransferase; D-Methioninmethylsulfonium; D-Aminosäure-Oxidase; oder D-Aminosäure-Acetyltransferase und Acyl-Coenzym-A-Synthetase und Acyl-Coenzym-A-Oxidase; ein SH-Reagenz; und ein oxidatives Farbentwicklungsmittel,
 wobei die Thiolverbindung aus Dithiothreitol, Mercaptoethanol, N-Acetylcystein, Dithioerythritol und Thioglycolsäure ausgewählt ist, und
 wobei das SH-Reagenz aus Ellmanns-Reagenz, Mercaptid-bildenden Mitteln, p-Quecksilberbenzoesäure, Alkylierungsmitteln, Iodessigsäure und N-Ethylmaleimid ausgewählt ist.

Revendications

1. Procédé permettant de détecter ou de mesurer de l'homocystéine dans un échantillon, comprenant:

- (a) la réaction d'un acide D-aminé présent dans un échantillon avec une enzyme de conversion d'acide D-aminé afin de convertir l'acide D-aminé en une substance qui ne sert pas de substrat d'oxydase d'acide D-aminé ou d'acétyltransférase d'acide D-aminé ;
 (b) la réduction de l'homocystéine dans l'échantillon au moyen d'un composé de thiol ;
 (c) la réaction de l'homocystéine réduite avec une méthyltransférase et du D-méthionine méthyl sulfonium pour permettre la production nouvelle de D-méthionine ;
 et
 (d) la réaction de la D-méthionine produite dans l'étape (c) avec l'oxydase d'acide D-aminé en présence d'un réactif de SH afin de produire du peroxyde d'hydrogène, et le développement couleur du peroxyde d'hydrogène produit en utilisant un agent oxydant de développement couleur,
 ou
 (d') la réaction de la D-méthionine produite dans l'étape (c) avec l'acétyltransférase d'acide D-aminé en présence d'acétyl-coenzyme A, produisant ainsi du coenzyme A ; la réaction du coenzyme A produit avec de la syntétase d'acyle-coenzyme A, produisant ainsi de l'acyle-coenzyme A ; et la réaction de l'acyle-coenzyme A produit avec l'oxydase d'acyle-coenzyme A en présence d'un réactif de SH afin de produire du peroxyde d'hydrogène ; et le développement couleur du peroxyde d'hydrogène produit en utilisant un agent oxydant de développement couleur,

dans lequel le composé de thiol est choisi parmi le dithiothréitol, le mercaptoéthanol, la N-acetylcystéine, le dithioerythritol et l'acide thioglycolique, et

dans lequel le réactif de SH est choisi parmi le réactif d'Ellman, les agents formant du mercaptide, l'acide benzoïque de p-mercure, les agents alkylants, l'acide iodoacétique et le N-éthylmaléimide.

2. Procédé selon la revendication 1, dans lequel l'étape (a) est une étape de réaction de D-alanine présente dans un échantillon avec de la ligase de D-alanyl-D-alanine en présence d'adénosine triphosphate afin de convertir la D-alanine en D-alanyl-D-alanine, et/ou une étape de réaction de D-sérine présente dans un échantillon avec de la déshydratase de D-sérine afin de convertir la D-sérine en acide pyruvique.

3. Procédé selon la revendication 2, dans lequel la méthyltransférase est de l'homocystéine méthyltransférase.

4. Procédé selon la revendication 2 ou 3, dans lequel, dans l'étape (d) ou (d'), le peroxyde d'hydrogène produit est détecté ou mesuré par développement couleur en utilisant de la peroxydase et un agent oxydant de développement couleur.

5. Trousse de réactif permettant de mesurer de l'homocystéine, comprenant de la ligase de D-alanyl-D-alanine et/ou de la déshydratase de D-sérine ; un composé de thiol, une méthyltransférase; du D-méthionine méthyl sulfonium; de l'oxydase d'acide D-aminé, ou de l'acétyltransférase d'acide D-aminé et de la syntétase d'acyle-coenzyme A et de l'oxydase d'acyle-coenzyme A; un réactif de SH; et un agent oxydant de développement couleur, dans laquelle le composé de thiol est choisi parmi le dithiothréitol, le mercaptoéthanol, la N-acetylcystéine, le dithioerythritol et l'acide thioglycolique, et dans laquelle le réactif de SH est choisi parmi le réactif d'Ellman, les agents formant du mercaptide, l'acide benzoïque de p-mercure, les agents alkylants, l'acide iodoacétique et le N-éthylmaléimide.

Fig.1




Fig. 2




Fig. 3




Fig. 4




Fig. 5





Fig. 6



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 0202802 A [0002] [0002]

Non-patent literature cited in the description

- **FUKUSHIMA, T.** *Biol. Pharm. Bull.*, 1995, vol. 18 (8), 1130-1132 [0002]
- Bergey's Manual of Determinative Bacteriology. The Williams & Wilkins Company, 1974, 295-296 [0016]
- *Biochemistry*, 1991, vol. 30, 1673-1682 [0017] [0038]
- *Journal of Bacteriology*, 1986, vol. 167, 809-817 [0017]
- *J. Bacteriol.*, 1983, vol. 154 (3), 1508-1512 [0018]
- *J. Mol. Biol.*, 1983, vol. 166, 557 [0023] [0039] [0049]
- *Biochemistry Dictionary*. Tokyo Kagaku Doujin, 1998, 182 [0034]
- *Journal of Biochemistry*, 1986, vol. 167, 809-817 [0048]
- *J. Biol. Chem.*, 1988, vol. 263, 16926-16933 [0058]