



(11)

EP 1 609 875 B9

(12)

CORRECTED EUROPEAN PATENT SPECIFICATION

(15) Correction information:
Corrected version no 1 (W1 B1)
 Corrections, see
 Description Paragraph(s) 70

(51) Int Cl.:
C12Q 1/68 (2006.01) **C12P 19/34 (2006.01)**
C07H 15/12 (2006.01) **G01N 33/53 (2006.01)**

(48) Corrigendum issued on:
02.03.2011 Bulletin 2011/09

(45) Date of publication and mention
 of the grant of the patent:
03.11.2010 Bulletin 2010/44

(21) Application number: **05020832.1**

(22) Date of filing: **28.01.1992**

(54) **DNA typing with short tandem repeat polymorphisms and identification of polymorphic short tandem repeats**

DNS "Typing" mit kurzen tandem-repetitiven Polymorphismen und Identifikation von kurzen polymorphen Tandem-Wiederholungen

Typage de l'ADN au moyen des polymorphismes de courts segments à répétition en tandem et identification de ces courts segments à répétition en tandem polymorphiques

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU MC NL SE

(30) Priority: **31.01.1991 US 647655**

(43) Date of publication of application:
28.12.2005 Bulletin 2005/52

(60) Divisional application:
08005776.3 / 1 967 593

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
92906069.7 / 0 639 228

(73) Proprietor: **BAYLOR COLLEGE OF MEDICINE**
Houston, TX 77030 (US)

(72) Inventors:
 • **Caskey, Thomas C.**
Houston, TX 77005 (US)
 • **Edwards, Albert O.**
Houston, TX 77021 (US)

(74) Representative: **Brasnett, Adrian Hugh et al**
Mewburn Ellis LLP
33 Gutter Lane
London
EC2V 8AS (GB)

(56) References cited:
EP-A- 0 266 787 WO-A-90/04040

- **STANDEN G R ET AL: "FAMILY STUDIES IN VON WILLEBRAND'S DISEASE BY ANALYSIS OF RESTRICTION FRAGMENT LENGTH POLYMORPHISMS AND AN INTRAGENIC VARIABLE NUMBER TANDEM REPEAT VNTR SEQUENCE" BRITISH JOURNAL OF HAEMATOLOGY, vol. 76, no. 2, 1990, pages 242-249, XP002453978 ISSN: 0007-1048**
- **ZULIANI G ET AL: "TETRANUCLEOTIDE REPEAT POLYMORPHISM IN THE APOLIPOPROTEIN B GENE" NUCLEIC ACIDS RESEARCH, vol. 18, no. 14, 1990, page 4299, XP002453979 ISSN: 0305-1048**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

- ZULIANI G ET AL: "TETRANUCLEOTIDE REPEAT POLYMORPHISM IN THE APOLIPOPROTEIN C-III GENE" NUCLEIC ACIDS RESEARCH, vol. 18, no. 14, 1990, page 4299, XP002453980 ISSN: 0305-1048
- POLYMEROPoulos M H ET AL: "TETRANUCLEOTIDE REPEAT POLYMORPHISM AT THE HUMAN AROMATASE CYTOCHROME P-450 GENE CYP 19" NUCLEIC ACIDS RESEARCH, vol. 19, no. 1, 1991, page 195, XP002453981 ISSN: 0305-1048
- ZULIANI G ET AL: "Tetranucleotide repeat polymorphism in the LPL gene" NUCLEIC ACIDS RESEARCH, OXFORD UNIVERSITY PRESS, SURREY, GB, vol. 18, no. 16, 25 August 1990 (1990-08-25), page 4958, XP002429804 ISSN: 0305-1048
- VAN AMSTEL H K P ET AL: "TETRANUCLEOTIDE REPEAT POLYMORPHISM IN THE VWF GENE" NUCLEIC ACIDS RESEARCH, vol. 18, no. 16, 1990, page 4957, XP002453982 ISSN: 0305-1048
- YANDELL D W ET AL: "DETECTION OF DNA SEQUENCE POLYMORPHISMS BY ENZYMATIC AMPLIFICATION AND DIRECT GENOMIC SEQUENCING" AMERICAN JOURNAL OF HUMAN GENETICS, vol. 45, no. 4, 1989, pages 547-555, XP002453990 ISSN: 0002-9297

DescriptionField of the Invention

5 **[0001]** The present invention relates generally to Kits for the detection of short tandem repeat sequence polymorphisms. More particularly, it relates to kits for detecting short tandem repeat sequences which show polymorphisms in the number of repeats for the detection or identification of medical and forensic samples, paternity, sample origin and tissue origin.

Background of the Invention

10 **[0002]** The volume of crime committed in the United States has risen with the increase of population and expansion of population centers. A large portion of violent crimes involve the creation of body fluid evidence having the potential of providing significant information about the perpetrator of a particular offense. Although the forensic science community has made tremendous effort in using this evidence, the results have historically been limited and are not useful in many situations when dealing with human remains and crime scene evidence. Identification by genetically inherited markers has long been seen as a possibility that would overcome most of the problems that are encountered when identification is not accomplished by fingerprints, forensic odontology, medical records or other methods. The establishment of a genetically inherited method that could be used for identification would have tremendous impact on investigation of the violent crimes of sexual assault and murder, identification of human remains and missing persons, and disputed parentage.

15 **[0003]** Methods enabling the matching of unidentified tissue samples to specific individuals would have wide application in the criminal justice system and the forensic sciences. With the possible exception of monozygotic twins, each individual in the human population has a unique genetic composition which could be used to specifically identify each individual. This phenomenon presents the theoretical possibility of using DNA sequence variation to determine whether a forensic sample was derived from any given individual.

20 **[0004]** Genetic marker systems, including blood groups and isoenzymes, have been used by forensic and medical serologists to provide estimates of individuality ranging from 1:1000 to 1:1,000,000 using 10 to 15 markers. Numbers in this range are often not available since a large percentage of the evidence does not yield results for ten genetic marker systems. Forensic scientists, investigators and the court system have been using inclusions as low as 1:5 to 1:100 in a population to bolster their case against defendants.

25 **[0005]** The fields of forensic and medical serology, paternity testing, and tissue and sample origin has been altered by the use of DNA sequence variation, e.g., satellite sequences and variable number of tandem repeats (VNTRs) or AMP-FLPs, in the crime laboratory, the court, hospitals and research and testing labs. Inclusion probabilities stated by the laboratories performing the analyses in such cases often exceed 1:1,000,000. The first implementation of DNA typing in forensics, was Jeffreys' use of a multilocus DNA probe "fingerprint" that identified a suspect in a murder case occurring in England. In the United States, DNA profiling has been established using a battery of unlinked highly polymorphic single locus VNTR probes. The use of these batteries of probes permits the development of a composite DNA profile for an individual. These profiles can be compared to ethnic databases using the principles of Hardy-Weinberg to determine the probability of the match between suspect and unknown forensic samples.

30 **[0006]** The application of VNTRs to gene mapping, population genetics, and personal identification has been limited by the low frequency and asymmetric distribution of these repeats in the genome and by the inability to precisely determine alleles with Southern hybridization-based detection schemes. The inability to make precise allele determinations complicates the application of VNTRs to personal identification. Binning protocols have been devised in which all alleles occurring within a region of the gel are treated as the same allele for genotype calculations. Since the allele distribution appears continuous because of the limited resolving power of Southern gels, heterozygotes with alleles of similar size may be scored as homozygotes. These features have led to claims that VNTR loci are not in Hardy-Weinberg equilibrium, and therefore the method for calculating the significance of a match is not agreed upon.

35 **[0007]** Although these methods have markedly improved the power of the forensic and medical scientist to distinguish between individuals, they suffer from a number of shortcomings including a lack of sensitivity, the absence of internal controls, expense, time intensity, relatively large sample size, an inability to perform precise allele identification and problems with identifying degraded DNA samples.

40 **[0008]** Medical and forensic studies have also employed the polymerase chain reaction (PCR) to examine variation in the HLA locus. PCR has also been used to amplify short VNTRs or AMP-FLPs. The use of PCR addresses some of the problems of sensitivity and sample degradation, however, the HLA typing system, still has some problems. A simpler, more powerful technique is needed which makes use of the most recent advances in DNA technology.

45 **[0009]** The present invention involves the novel application of these advances to medical and forensic science. In the present invention novel classes of highly polymorphic, primarily trimeric and tetrameric, short or simple tandem repeats (STRs) which are present within the human genome have been identified. These STRs have characteristics suitable for

inclusion in a DNA profiling assay. This assay incorporates internal or external standards, provides higher sensitivity, requires shorter analysis time, lowers expense, and enables precise identification of alleles. The STRs are amplified with great fidelity and the allele patterns are easily interpreted. Amplification of highly polymorphic tandemly reiterated sequences may be the most cost effective and powerful method available to the medical and forensic community.

5 [0010] The DNA profiling assay of the present invention has features which represent significant improvements over existing technology and brings increased power and precision to DNA profiling for criminal justice, paternity testing, and other forensic and medical uses.

Summary of the Invention

10 [0011] The present invention provides a kit for a DNA profiling assay comprising a container having oligonucleotide primer pairs for amplifying a short tandem repeat sequence, comprising primers with the sequence of SEQ ID 13 and SEQ ID 14.

15 [0012] In a further embodiment the present invention provides a kit for a DNA profiling assay comprising a container having oligonucleotide primer pairs for amplifying a short tandem repeat sequence, comprising primers with the sequence of SEQ ID 13 and SEQ ID 23.

[0013] The present invention also provides a respective kit further comprising a labelled standard.

20 [0014] According to one embodiment, these kits further comprise a container having reagents for multiplex polymerase chain reaction.

Description of the Drawings

[0015]

25 Figure 1 illustrates the strategy used to determine the sequence flanking a STR.

Figure 2 shows the development of a polymorphic STR locus.

30 Figure 3 shows examples of products from the multiplex and single PCR assays used in generating multilocus genotype data. Figure 3A shows the multiplex PCR(mPCR) of HUMHPRTB[AGAT]_n (top) and HUMFABP[AAT]_n. Figure 3B shows the mPCR of HUMRENA4[ACAG]_n (top) and HUMTH01[AATG]_n. Figure 3C shows the single PCR of HUMARA[AGC]_n.

35 Figures 4A to 4E plot the relative allele frequency distributions. Allele counts were used to calculate and plot the relative frequencies of alleles at: 4A HUMRENA4[ACAG]_n; 4B HUMW01[AATG]_n; 4C HUMARA[AGC]_n; 4D HUM-PRTB[AGAT]_n; and 4E HUMFABP[AAT]_n.

40 Figure 5 shows the results of a fluorescent DNA typing assay. The analysis software scales the intensities of the fluorescent profiles relative to the strongest signal.

45 [0016] The drawings and figures are not necessarily to scale and certain features of the invention may be exaggerated in scale or shown in schematic form in the interest of clarity and conciseness.

Detailed Description of Invention

50 [0017] As used herein, the term "short tandem repeat" (STR) refers to all sequences between 2 and 7 nucleotides long which are tandemly reiterated within the human organism. The STRs can be represented by the formula $(A_wG_xT_yC_z)_n$ where A, G, T and C represent the nucleotides which can be in any order; w, x, y and z represent the number of each nucleotide in the sequence and range between 0 and 7 with the sum of w+x+y+z ranging between 2 and 7; and n represents the number of times the sequence is tandemly repeated and is between about 5 and 50. Most of the useful polymorphisms usually occur when the sum of w+x+y+z ranges between 3 and 7 and n ranges between 5 and 40. For dimeric repeat sequences n usually ranges between 10 and 40.

55 [0018] As used herein "non-duplicative" sequence means the sequence and its complement. It is represented in its lowest alphabetical form as shown in Table 1. For example (ACT) represents ACT, CTA, TAC, AGT, TAG and GTA. Each representative sequence can represent a maximum of two times the number of nucleotides in the sequence.

[0019] As used herein "flanking sequence" refers to the nucleotide sequence on either side of the STR. "Unique flanking sequences" are those flanking sequences which are only found at one location within the genome.

[0020] The term "oligonucleotide primers" as used herein defines a molecule comprised of more than three deoxyri-

bonucleotides or ribonucleotides. Its exact length will depend on many factors relating to the ultimate function and use of the oligonucleotide primer, including temperature, source of the primer and use of the method. The oligonucleotide primer can occur naturally, as in a purified restriction digest, or be produced synthetically. The oligonucleotide primer is capable of acting as an initiation point for synthesis when placed under conditions which induce synthesis of a primer extension product complementary to a nucleic acid strand. The conditions can include the presence of nucleotides and an inducing agent such as a DNA polymerase at a suitable temperature and pH. In the preferred embodiment, the primer is a single-stranded oligodeoxyribonucleotide of sufficient length to prime the synthesis of an extension product from a specific sequence in the presence of an inducing agent. Sensitivity and specificity of the oligonucleotide primers are determined by the primer length and uniqueness of sequence within a given sample of template DNA. In the present invention the oligonucleotide primers are usually about greater than 15 mer and in the preferred embodiment are about 20 to 30 mer in length.

[0021] Each pair of primers is selected to detect a different STR. Each primer of each pair herein is selected to be substantially complementary to a different strand in the flanking sequence of each specific STR sequence to be amplified. Thus one primer of each pair is sufficiently complementary to hybridize with a part of the sequence in the sense strand and the other primer of each pair is sufficiently complementary to hybridize with a different part of the same sequence in the antisense strand. Although the primer sequence need not reflect the exact sequence of the template, the more closely the 3' end reflects the exact sequence, the better the binding during the annealing stage.

[0022] As used herein the term "extension product" refers to the nucleotide sequence which is synthesized from the 3' end of the oligonucleotide primer and which is complementary to the strand to which the oligonucleotide primer is bound.

[0023] As used herein the term "differentially labeled" indicates that each extension product can be distinguished from all others because it has a different label attached and/or is of a different size and/or binds to a specifically labeled oligonucleotide. One skilled in the art will recognize that a variety of labels are available. For example, these can include radioisotopes, fluorescers, chemiluminescers, stains, enzymes and antibodies. Various factors affect the choice of the label. These include the effect of the label on the rate of hybridization and binding of the primer to the DNA, the sensitivity of the label, the ease of making the labeled primer, probe or extension products, the ability to automate, the available instrumentation, convenience and the like. For example, differential radioisotope labelling could include ^{32}P , ^{3}H and ^{14}C ; differential fluorescers labelling could include fluorescein-5-isothiocyanate, tetramethylrhodamine-5- (and-6) isothiocyanate, Texas Red and NBD aminoheanoic acid; or a mixture of different labels such as radioisotopes, fluorescers and chemiluminescers.

[0024] Each specific, different DNA sequence, which is to be detected herein is derived from genomic DNA. The source of the genomic DNA to be tested can be any medical or forensic sample. Examples of medical and forensic samples include blood, semen, vaginal swabs, tissue, hair, saliva, urine and mixtures of body fluids. These samples can be fresh, old, dried and/or partially-degraded. The samples can be collected from evidence at the scene of a crime.

[0025] The term "forensic sample" as used herein means using the technology for legal problems including but not limited to criminal, paternity testing and mixed-up samples. The term "medical sample" as used herein means use of the technology for medical problems including but not limited to research, diagnosis, and tissue and organ transplants.

[0026] As used herein the term "polymorphism" refers to the genetic variation seen in the tandem repeats or flanking sequences. One example of this polymorphism is in the number of times the 3 to 7 nucleotide sequence is repeated.

[0027] As used herein the term "multiplex polymerase chain reaction" (mPCR) refers to a novel variation of PCR. It is a procedure for simultaneously performing PCR on greater than two different sequences. The mPCR reaction comprises: creating said extracted DNA to form single stranded complementary strands, adding a plurality of labeled paired oligonucleotide primers, each paired primer specific for a different short tandem repeat sequence, one primer of each pair substantially complementary to a part of the sequence in the sense strand and the other primer of each pair substantially complementary to a different part of the same sequence in the complementary antisense strand, annealing the plurality of paired primers to their complementary sequences, simultaneously extending said plurality of annealed primers from the 3' terminus of each primer to synthesize an extension product complementary to the strands annealed to each primer, said extension products, after separation from their complement, serving as templates for the synthesis of an extension product for the other primer of each pair, separating said extension products from said templates to produce single stranded molecules, amplifying said single stranded molecules by repeating at least once said annealing, extending and separating steps. In the mPCR process the preferred method for three loci includes: (1) primers composed of similar GC base compositions and lengths; (2) longer extension times up to 8 fold the normally utilized times; and (3) minimization of the number of PCR cycles performed to achieve detection for example approximately 23-25 cycles.

[0028] The mPCR reaction is optimized for each reaction. In some mPCR reactions the optimization further includes more enzyme than one normally adds to a PCR reaction.

[0029] As used herein, the term "match probability" refers to the chance that two unrelated persons will have the same combined genotype at the examined loci.

[0030] A DNA profiling assay for detecting polymorphisms in a short tandem repeat may comprise the steps of : extracting DNA from a sample to be tested; amplifying the extracted DNA; and identifying said amplified extension

products for each different sequence, wherein each different sequence has a differential label. Although a variety of known amplification procedures are known, the preferred embodiment employs PCR or mPCR.

[0031] In one embodiment of the present invention an external standard is used. In a preferred embodiment an internal standard is used. The internal standard is composed of labeled alleles of the STR loci of interest. One skilled in the art will recognize that the choice of standard and the intervals chosen will depend on the label and the desired resolution. For example, in a DNA profiling assay STR alleles from a DNA sample can be localized with greater than 1 bp resolution using an internal standard marker every 3-5 bp.

[0032] In a preferred assay, short tandem repeat sequences of nucleotides characterized by the formula $(A_w G_x T_y C_z)_n$ wherein A, G, T and C represent the nucleotides, w, x, y and z represent the number of each nucleotide and range from 0 to 7 and the sum of w+x+y+z ranges from 3 to 7 and n represents the repeat number and ranges from 5 to 40 are used. In another preferred assay, the sum of w+x+y+z is either 3 or 4. In the preferred embodiment of the profiling assay, at least two STRs are assayed simultaneously.

[0033] The DNA profiling assay may be automated. This automation can be achieved by a variety of methods. One method is to use an automated DNA label analyzer capable of distinguishing simultaneously different fluorescers, radioactive labels or chemiluminescers during the identifying step. One skilled in the art will readily recognize that a variety of instrumentation meets this requirement. One example of such an analyzer used in the preferred assay is the Applied Biosystems 370A Fluorescent DNA sequence device ("370A device") which has the capability to distinguish between four different fluors during electrophoresis.

[0034] A method of detecting polymorphic STRs for use in the DNA profiling assay, may comprise the steps of: determining possible non-duplicative nucleotide sequences of the formula $(A_w G_x T_y C_z)_n$, wherein A, G, T and C represent each respective nucleotide and w, x, y and z represent the number of each nucleotide in the sequence and range between 0 and 7 with the sum of w+x+y+z ranging between 2 and 7; searching for and identifying $(A_w G_x T_y C_z)_n$ in databases containing known genetic sequences and identifying the $(A_w G_x T_y C_z)_n$ sequence of said genetic sequence and its flanking sequence, wherein n represents the number of tandem repeats of the sequence and is at least about 5; extracting each identified sequence and its flanking sequence; identifying the extracted sequences which have unique flanking sequences; synthesizing oligonucleotide primer pairs to the unique flanking sequences; performing a PCR with the primer pairs on DNA samples from a test population; and examining the extension products of the PCR to detect polymorphic STRs.

[0035] The present invention provides a kit for DNA profiling assays as defined above. The kit is comprised of a container having oligonucleotide primer pairs for amplifying a STR. In the preferred embodiment the number of STR primer pairs is selected such that the genotype frequency (p) is at least 10^{-6} . This usually requires 6-10 STR primer pairs.

[0036] A further addition to the kit can be a container having labelled standards. An additional enhancement to the kit is the addition of reagents for mPCR.

[0037] In examples, all percentages are by weight, if for solids and by volume if for liquids, and all temperatures are in degrees Celsius unless otherwise noted.

Example 1

Computer Identification of STR loci

[0038] STRs were identified by searching all human sequences in the GenBank DNA sequence repository for the presence of all possible classes of dimeric, trimeric, and tetrameric STRs. One skilled in the art will readily recognize that a similar search using repeats of 5 to 7 nucleotides can be used to identify STRs of 5 to 7 nucleotides in length. The possible non-duplicative nucleotide sequences used in this search are given by their lowest alphabetical representation.

Table 1: Possible Non-Duplicative STRs

(AA) (AC) (AG) (AT) (CC) (CG)
(AAC) (AAG) (AAT) (ACC) (ACG) (ACT) (AGC) (AGG)

(ATC) (CCG) (AAAC) (AAAG) (AAAT) (AACC) (AACG) (AACT)
 (AAGC) (AAGG) (AAGT) (AATC) (AATG) (AATT) (ACAG) (ACAT)
 5 (AGAT) (ACCC) (ACCG) (ACCT) (ACGC) (ACGG) (ACGT) (ACTC)
 (ACTG) (ACTT) (AGCC) (AGCG) (AGCT) (AGGC) (AGGG) (ATCC)
 (ATCG) (ATGC) (CCCG) (CCGG)

10 [0039] As shown in Table 2, the computer search identified a considerable number of STRs. The dimeric search was set to identify sequences in which the STR was repeated at least 10 times and the trimeric and tetrameric search was set to identify sequences which were repeated at least 5 times.

15

MONOMER LENGTH	DINUCLEOTIDE (10 OR MORE)	TRINUCLEOTIDE (5 OR MORE)	TETRANUCLEOTIDE (5 OR MORE)
FRACTION OBSERVED	5/6	10/10	16/34
TOTAL NUMBER	217	101	67

20

[0040] The fraction observed refers to the number of different classes of STRs observed out of the total number of possible STR classes. All classes of trimeric repeats were present and about half of the possible tetrameric sequences were represented.

25 [0041] Approximately 50% of STRs studied were polymorphic. Trimeric and tetrameric STRs have features of polymorphic markers useful for the physical and genetic mapping of the human genome and personal identification in the medical and forensic sciences.

30 Example 2

30 Molecular Biological Identification of STR Loci

35 [0042] In addition to the procedure for identifying STRs in the GenBank, other methods are available to identify additional STR loci. For example, oligonucleotide probes for the possible 50 unique dimeric, trimeric and tetrameric STR sequences can be synthesized and used to screen total human DNA libraries. In one example recombinant bacteriophage lambda of the human X chromosome were plated at a density of 255 plaque forming units per 15 cm plate. Plaque lifts made from the plates are hybridized to ^{32}P 5' end-labeled oligonucleotides of the STR motifs. Standard hybridization methods were used. Oligonucleotides were labeled according to standard protocols.

40 [0043] With nucleotide sequences up to above 100 bp the conditions for hybridization may be estimated using the following formula:

$$T_i = T_m - 15^\circ\text{C}$$

45

$$T_m = 16.6 \log[M] + 0.41[P_{gc}] + 81.5 - P_m \cdot B/L - 0.65[P_f]$$

50

where: M is the molar concentration of Na^+ , to a maximum of 0.5 (1 X SSC contains 0.165 M Na^+);

P_{gc} is the percent of G or C bases in the oligonucleotide and is 1-16 between 30 and 70;

P_m is the percent of mismatched bases, if known;

55 P_f is the percent of formamide in the buffer;

B is 675 for synthetic probes up to 100 bases;

L is the length of the probe in bases.

[0044] The formula was used to arrive at the conditions in Table 3:

Table 3

Oligo	(ml)2X Hyb Mix	Pf(%)	Formamide(ml)	H2O(ml)	V(ml)
<u>Id sequence</u>					
1152 [AATC] _{7.5}	12.5	10%	2.5	10	25
1154 [AGAT] _{7.5}	12.5	10	2.5	10	25
1525 [AAT] ₁₀	12.5	0	0	12.5	25
1526 [AATG] _{7.5}	12.5	10	2.5	10	25
1528 [ACAG] _{7.5}	12.5	30	7.5	5	25

[0045] The 2 X hybridization mix contained 37.5 ml of 20 X SSC, 15 ml of 50 X Denhardts, 7.5 ml of 20% SDS, and 15 ml of H₂O. Hybridizations were performed at 42°C.

[0046] These conditions were used to determine the frequency of each STR shown on the X chromosome using recombinant lambda from an X chromosome genomic library picked to a grid. (Table 4).

TABLE 4. The frequency of trimeric and tetrameric STRs.

STR	Positive bacteriophage(%)	Frequency (kb/STR)
[AAT]	5	300
[AATC]	5	300
[AATG]	3	500
[ACAG]	3	500
[AGAT]	3	500

[0047] A total of 1020 recombinant bacteriophage were hybridized to radiolabeled 30 bp oligonucleotides (e.g., [AATC]_{7.5}). Calculations were based on an average insert size of 15 kb in the library. These hybridization results and the results from the GenBank studies suggest the presence of approximately 400 million STRs in the human genome. The X-chromosome results have been extended to the entire human genome by utilizing the complete genomic phage lambda library. Thus identification of sufficient STRs to extend the DNA typing assay to very high levels of individualization (e.g., one in a billion) is feasible.

Example 3

Determination of DNA sequence flanking STRs

[0048] Clones containing STRs, for example M13, lambda, cosmid and YAC, can be identified by any procedure which allows hybridization to one of the core oligonucleotides. Most of the hybridization methods are usually laborious for determining the sequence of the unique DNA segments flanking both sides of the STR. In the present invention a strategy called STR-PCR was used. This strategy is shown in schematic form in (Figure 1).

[0049] The STR-PCR strategy was based upon the method of Riley, et al, Nucleic Acids Res. 18: 2887 (1990). The Riley method was designed to amplify the ends of YAC molecules from total yeast genomic DNA. This method was adapted to amplify the DNA segments flanking STRs, and coupled to direct DNA sequencing of the products via a solid-phased-DNA sequencing technology. The procedure involves the following steps: (1) Blunt ends are generated to flank both sides of a STR in a cloned DNA segment by digestion with a single restriction enzyme. Multiple enzymes can be used, separately, to generate a flanking sequence length in the range of 100 to 150 bp. (2) A linker which contains a region of non-complementary DNA is ligated to the population of blunt ended molecules. (3) The flanking sequences are amplified in separate reactions. The left end is amplified with the anchored PCR primer and a primer of one strand of the STR. The right end is amplified with the same anchored PCR primer and a primer of the other strand of the STR. The STR primers may be biotinylated (*). (4) The final biotinylated (*) PCR product. (5) The biotinylated strand may be captured with avidin coated beads. And (6) The flanking sequence may be obtained by extension from the sequencing primer in the presence of dideoxynucleotides.

[0050] Results from using the STR-PCR strategy are shown in Figure 2. Amplification of the DNA sequence flanking both sides of an (AGAT) STR from two recombinant bacteriophage is shown in Figure 2A. While Figure 2B shows direct DNA sequencing of single stranded template following capture and strand separation of the biotinylated amplification products of λ AE[AGAT]-2 with avidin coated magnetic beads. Figure 2C demonstrates the use of oligonucleotides

complementary to the sequence flanking the STR to amplify the STR locus in a family.

[0051] Oligonucleotide primers which generate a blunt ended linker upon annealing were synthesized. Examples of these oligonucleotides are SEQ ID Nos: 1, 2, 3 and 4. Oligonucleotides (SEQ ID Nos: 1 and 2) form the double stranded linker, oligonucleotide (SEQ ID No: 3) is the PCR primer for the anchor and oligonucleotide (SEQ ID No: 4) is the DNA sequencing primer.

[0052] Oligonucleotide (SEQ ID No: 1) was phosphorylated and annealed to (SEQ ID No: 2) to form the double stranded linker by standard protocol. In this procedure 10 μ L of one of the linker oligonucleotides (SEQ. ID No: 1) (100 μ M; 1 nanomole) is combined with about 10 μ L of 32 P-gamma-ATP (10mM); about 5 μ L of PNK buffer; about 3 μ L of T4 PNKinase (30U); about 22 μ L of H₂O for about a 50 μ L final volume (at about 37°C, for about 40 min.; and at about 65°C, for about 5 min). Then about 10 μ L of the other linker oligonucleotide (SEQ ID No: 2) (100 μ M) is added. This mixture is held at about 95°C for about 5 min. The mixture is then slowly cooled to room temperature.

[0053] Although the Riley method taught that both oligonucleotides should be phosphorylated, the present invention has discovered that it is sufficient and possibly better to phosphorylate only the first oligonucleotide.

[0054] The next procedure was to ligate the double stranded linker (PCR anchor) to recombinant bacteriophage lambda DNA. First each clone was digested with frequent cutting restriction enzymes which give blunt ends. For example: Alul, HaeIII, and Rsal. Second, the linker was ligated to the blunt ended DNA. Third, the flanking segments were amplified. In this procedure the sample was digested with sufficient enzyme to cut the DNA (about 0.5 μ L Enzyme) in about 1.5 μ L of One Phor All Plus Restriction Buffer (Pharmacia) about 250 ng Lambda Phage DNA (50 kb) and sufficient H₂O to bring the final volume to about 15 μ L. The temperature was held at about 37°C until the DNA was cut.

[0055] After the digestion, a cocktail for ligations is added. Although a variety of cocktails are known, the present invention used: up to about 2.0 μ L of Annealed Oligonucleotides with about 0.05 μ L of 1M DTT, about 3.0 μ L of 0.1 M rATP, about 0.25 μ L of T4 DNA Ligase (Pharmacia), about 1.5 μ L of One Phor All Plus Restriction Buffer and about 9.7 μ L of H₂O. The mixture was held at about 15°C for at least about 2 hours. Longer times, for example overnight, may give better results.

[0056] Next, the flanking segments were amplified. The amplification mix included about 3 μ L of STR-PCR Primer (10 X Cetus; 10 μ M), about 3 μ L of Anchor-PCR Primer (same as with STR-PCR primer), about 3 μ L of dNTP mix (10 X Cetus; 2 mM), about 3 μ L of PCR Buffer (10 X Cetus, without Mg⁺⁺), about 3.0 μ L of 0.01 M MgCl₂, about 1 μ L of DNA, about 14.2 μ L of H₂O, and about 0.3 μ L of AmpliTaq (Cetus).

[0057] The mixture was heated for about 2 min at about 95°C, then the PCR assay on the mixture included about 25-30 cycles of about 45 sec at about 95°C, then about 30 sec at about 60°C, then about 1 min at about 72°C. Finally, the mixture was held at about 72°C for about 10 min, then transferred to about 4°C. In this procedure the STR-PCR was performed separately for both strands of the STR. Control of the concentration of Mg⁺⁺ appeared to be important.

[0058] The amplified products were sequenced. If the STR primers are biotinylated (for example, by the Aminolink 2 methodology of Applied Biosystems, Inc.) the products were captured with avidin coated beads (Dynal). The unwanted DNA strand was removed. The preferred conditions for isolation for the amplified products was as follows: about 25 μ L of M280 Beads from Dynal were mixed with about 25 μ L of PCR Product. This was held for about 30 min at about 25°C on a rotating wheel. The supernatant was removed and about 150 μ L of 0.15 M NaOH was added. This was held for 5 min. at about 25°C. The supernatant is then removed, the remaining material is washed at least once with H₂O and resuspended in about 7 μ L of H₂O for DNA sequencing. Any standard DNA sequencing reactions can be used. In the present invention the sequencing was performed as for any single stranded template.

Example 4

Frequency of Polymorphic Variation of STRs and examples

[0059] Seventeen STRs present either within the human HPRT locus or in human sequences in the GenBank database were assayed for variation in the human population. Nine were polymorphic.

[0060] Amplifications were performed with Perkin-Elmer-Cetus thermocyclers, AmpliTaq enzyme, and recommended buffer conditions in a volume of about 15 μ L. Amplification conditions were about 95°C for about 45 sec., then about 60°C for about 30 sec., then about 72°C for about 30 sec. Approximately 23-28 cycles were run. Amplified products were radiolabeled by inclusion of 2 μ Ci 32 P-dCTP (3000 Ci/mmol) in the PCR. The HUMHPRTB [AGAT]_n and HL1MFABP [AAT]_n loci, and the HUMRENA4 [ACAG]_n and HUMTH01 [AATG]_n loci, were studied as a multiplex PCR of two loci. Approximately 50 ng of genomic DNA was used in the PCRs. PCR products were diluted 2:5 in formamide, denatured at about 95°C for about 2 min., and loaded onto a DNA sequencing gel (about 6% (39:1) acrylamide:bisacrylamide, with about 7 M urea, and about 0.04% TEMED). Control reactions without added DNA were included in every set of amplifications. The amplification products ranged in size from between about 100 to 350 bp. This allowed precise determination of allele lengths.

[0061] The GenBank data for locus name, approximate repeat sequence, Primer SEQ ID No:, number of alleles

observed, number of chromosomes studied and average predicted heterozygote frequencies are shown in Table 6.

TABLE 6: Example STR's

	Locus and STR Sequence	PCR Primer SEQ. ID NO:	Alleles Detected	Chromosomes Studied	Heterozygotes (%)	Chromosomes
5	HUMFABP [AAT] ₈₋₁₅	5,6	8	314	52-78	4
10	HUMARA [AGC] ₁₂₋₃₀	7,8	17	228	87-91	X
15	HUMGPP3A09 [AAGG] ₉	9,10	4	24	31	N.D.
20	HUMERP[c: AATG] ₂	11,12	2	20	19	7
25	[ACTC] ₄ [c: AATG] ₅					
30	HUMTH01 [AATG] ₆₋₁₂	13,14	7	320	72-78	11
35	HUMTNFAB [AATG] ₅	15,16	4	24	31	6
40	HUMRENA4 [ACAG] ₇₋₁₂	17,18	6	310	34-48	1
45	HUMHPRTB [AGAT] ₉₋₁₆	19,20	8	227	69-78	X
50	HUMSTRX1 [AGAT] ₁₃	21,22	11	44	60-84	X

[0062] In Table 6 the features of 9 polymorphic STR loci are shown. The range of heterozygote frequencies represents the values obtained for the least to most polymorphic racial group. Alleles from loci shown with a range of reiteration numbers (for example, HUMFABP[AAT]₈₋₁₅) were sequenced to enable precise association of the number of tandem reiterations with specific alleles. The reiteration number of the GenBank clone is given for loci at which the range in the number of repeats is unknown. The lowest alphabetical representation of each STR motif is used, with the reverse complement (c:) indicated where appropriate for compound STR loci. Variability in the human population was assayed with a radioactive PCR assay.

Example 5

Examples of data from the radioactive PCR assay for 5 STR loci

[0063] Genotype data for five STR loci were determined in two multiplex and one single PCR (Figure 3). Both DNA strands of the amplified products are radiolabeled and the alleles of different loci have distinct appearances based on the relative mobilities of the two DNA strands. HUMHPRTB [AGAT]_n and HUMTH01 [AATG]_n alleles appear as closely spaced doublet bands, while HUMRENA4 [ACAG]_n and HUMFABP [AAT]_n alleles usually appear as singlets. HUMARA [AGC]_n alleles appear as widely spaced doublets, such that adjacent alleles overlap. The faster strand of HUMHPRTB [AGAT]_n alleles usually appear as a doublet, due to incomplete addition of an extra, non-complementary base to the 3' end of the product. The relative mobilities of the strands are influenced by the composition of the polyacrylamide gel. The data for Figure 3 was selected from the population surveys as a fair representation of the clarity with which allele designations were made. The autoradiograms were overexposed to illustrate the faint artifactual bands differing in the number of repeats which are thought to arise during the PCR.

[0064] Representative alleles from each of the polymorphic STRs were sequenced. The results show that the variation in size is a function of the number of repeats.

Example 6

Population Genetics of STRs in four human ethnic groups

[0065] Trimeric and tetrameric STRs represent a rich source of highly polymorphic markers in the human genome.

Analysis of a multilocus genotype survey of 40 or more individuals in U.S. Black, White, Hispanic, and Asian populations at five STR loci located on chromosomes 1, 4, 11, and X was performed. The heterozygote frequencies of the loci ranged from 0.34 to 0.91 and the number of alleles from 6 to 17 for the 20 race and locus combinations. Relative allele frequencies exhibited differences between races and unimodal, bimodal, and complex distributions. Genotype data from the loci 5 were consistent with Hardy-Weinberg equilibrium by three tests and population sub-heterogeneity within each ethnic group was not detected by two additional tests. No mutations were detected in a total of 860 meiosis for two to five loci studied in various kindreds. An indirect estimate of the mutation rates give values from 2.5×10^{-5} to 15×10^{-5} for the 10 five loci. Higher mutation rates appear to be associated with more tandem repeats of the core motif. The most frequent genotype for all five loci combined appears to have a frequency of 6.51×10^{-4} . Together, these results suggest that trimeric and tetrameric STR loci are ideal markers for understanding the mechanism of production of new mutations at hypervariable DNA regions and are suitable for application to personal identification in the medical and forensic sciences.

TABLE 7: Polymorphic Short Tandem Repeat Loci Studied

Locus and STR	Accession Number	Chromosome	PCR Primer SEQ. ID. NO:	Product Length (bp)	Gene
HUMFABP [AAT] _n	M18079	4q31	5,6	199-220	Intestinal Fatty Acid Binding Protein
HUMARA[AGC] _n	M21748	Xcen-q13	7,8	261-312	Androgen Receptor
HTMTMTH01 [AATG] _n	D00269	11p15.5	13,14	183-207	Tyrosine hydroxylase
HUMRENA4 [ACAG] _n	M10151	1q32	17,18	251-271	Renin
HUMHPRTB [AGAT] _n	M26434	Xq26	19,20	263-299	Hypoxanthine Phosphoribosyl transferase

30 Samples

[0066] DNA was extracted from blood samples obtained at local blood banks from unrelated volunteer donors. Blood bank personnel visually designated donors as Black, White, or Other. Hispanics and Orientals were identified on the basis of surname. A total of 40 individuals in each of these four ethnic groups were studied. Genotype data in 40 families 35 (10 French, 27 Utah/Mormon, 2 Venezuelan, and 1 Amish) was determined with HUMHPRTB (AGAT)_n and HUMFABP (AAT)_n. Five STR loci were studied in additional families for a minimum of 31 meiosis.

STR Loci

[0067] The STR loci are designated by their GenBank locus name and the lowest alphabetical representation of the 40 44 possible unique trimeric and tetrameric repeat motifs. For example, HUMHPRTB (AGAT)_n refers to the polymorphic (CTAT) STR located in intron 3 of the human hypoxanthine phosphoribosyltransferase (HPRT) gene. The loci studied, their GenBank accession numbers, chromosomal assignments, amplification primers, and range of product sizes (based on the GenBank sequence) are given in Table 7.

[0068] In Table 8 the alleles are numbered according to the number of tandem repeats present in the amplification products. The number of repeated motifs was determined by direct DNA sequencing of amplified products or by subcloning into M13 for sequencing. The repeat number of subcloned fragments was verified relative to the original genomic DNA source by amplification of the cloned segment.

50 Computations and Statistics

[0069] A variety of standard population genetics tests were employed to evaluate the heterozygote frequencies, allele frequencies and random association of alleles at different loci. These tests included measurements of standard errors, G-statistics for the likelihood-ratio test, binomial distributions, Hardy-Weinberg equilibrium and the summary statistic (S_k^2). 55

Relative Allele Frequencies

[0070] Allele frequencies and their standard errors were calculated from the genotypes of approximately 40 individuals for the 20 combinations of five STR loci and four populations (Table 8).

5

TABLE 8 Allele Frequencies and Their Standard Errors at Five STR Loci in Four Populations

Allele frequencies (%) and standard errors (%) in Allele ^a						
	Whites	Blacks	Hispanics	Asians	Pooled ^b	
LOCUS - HUMHPRTB[AGAT] _n						
10	7	0.4±0.4	---	---	---	0.3±0.3
	9	0.4±0.4	1.6±1.6	---	---	0.5±0.4
	10	0.4±0.4	1.6±1.6	---	---	0.5±0.4
15	11	12.1±2.2	3.2±2.2	8.9±3.8	11.3±4.4	10.1±1.5
	12	34.4±3.2	29.0±5.8	39.3±6.5	26.4±6.1	33.2±2.4
	13	33.0±3.1	30.6±5.9	39.3±6.5	39.6±6.7	34.4±2.4
	14	14.7±2.4	21.0±5.2	5.4±30	13.2±4.7	14.2±1.8
20	15	2.2±1.0	11.3±4.0	7.1±34	9.4±4.0	5.3±1.1
	16	2.2±1.0	1.6±1.6	---	---	1.5±0.6
	(n ^d)	224	62	56	53	395
LOCUS - HUMTH01[AATG] _n						
25	6	26.2±4.9	12.5±3.7	21.3±4.6	8.8±3.2	17.2±2.1
	7	8.8±3.2	32.5±5.2	30.0±5.1	23.7±4.8	23.8±2.4
	8	11.3±3.5	21.3±4.6	6.3±2.7	3.8±2.1	10.6±1.7
	9	16.2±4.1	21.3±4.6	13.7±3.9	47.5±5.6	24.7±2.4
	10	36.2±5.4	12.5±3.7	28.7±5.1	7.5±2.9	21.3±2.3
30	11	1.3±1.2	---	---	7.5±2.9	2.2±2.3
	12	---	---	---	1.3±1.2	0.3±0.3
	(n)	80	80	80	80	320
LOCUS - HUMRENA4[ACAG] _n						
35	7	---	2.5±1.7	---	---	0.6±0.5
	8	80.3±4.6	71.2±5.1	76.2±4.8	69.7±5.3	74.4±2.5
	9	---	3.8±2.1	---	---	1.0±0.6
	10	7.9±3.1	11.3±3.5	6.3±2.7	19.7±4.6	11.2±1.8
	11	11.8±3.7	10.0±3.4	12.5±3.7	3.9±2.2	9.6±1.7
40	12	---	1.3±1.2	5.0±2.4	6.6±2.8	3.2±1.0
	(n)	76	80	80	76	312
LOCUS - HUMFABP[AAT] _n						
45	8	---	2.5±1.7	1.3±1.2	---	0.6±0.3
	9	0.3±0.3	27.5±5.0	1.3±1.2	5.1±2.5	5.2±1.0
	10	49.7±2.9	32.5±5.2	55.0±5.6	66.7±5.3	50.4±2.2
	11	17.4±2.2	2.5±1.7	8.8±3.2	6.4±2.8	12.3±1.4
	12	3.4±1.0	5.0±2.4	2.5±1.7	2.6±1.8	3.4±0.8
	13	24.8±2.5	12.5±3.7	25.0±4.8	19.2±4.4	22.2±1.8
	14	4.4±1.2	16.3±4.1	6.2±2.7	---	5.8±1.0
50	15	---	1.3±1.2	---	---	0.2±0.2
	(n)	298	80	80	78	536
	0					
LOCUS - HUMARA[AGC] _n						
55	13	---	1.6±1.6	---	---	0.4±0.4
	15	---	---	---	5.7±3.2	1.3±0.8
	16	---	1.6±1.6	---	---	0.4±0.4
	17	1.7±1.7	17.7±4.9	---	1.9±1.9	5.7±1.5

(continued)

Allele frequencies (%) and standard errors (%) in Allele ^a						
	Whites	Blacks	Hispanics	Asians	Pooled ^b	
5	18	1.7±1.7	16.1±4.7	9.3±3.9	3.8±2.6	7.9±1.8
	19	10.2±3.9	8.1±3.5	9.3±3.9	9.4±4.0	9.2±1.9
	20	15.3±4.7	9.7±3.8	3.7±2.6	5.7±3.2	8.8±1.9
	21	16.9±4.9	8.1±3.5	18.5±5.3	18.9±5.4	15.4±2.4
10	22	8.5±3.6	17.7±4.9	9.3±3.9	17.0±5.2	13.6±2.3
	23	15.3±4.7	4.8±2.7	9.3±3.9	15.1±4.9	11.0±2.1
	24	20.3±5.2	6.5±3.1	13.0±4.6	3.8±2.6	10.5±2.0
	25	6.8±3.3	1.6±1.6	11.1±4.3	5.7±3.2	6.1±1.6
	26	3.4±2.4	---	5.6±3.1	11.3±4.3	4.8±1.4
15	27	---	1.6±1.6	1.8±1.8	1.9±1.9	1.3±0.8
	28	---	3.2±2.2	1.8±1.8	---	1.3±0.8
	29	---	1.6±1.6	5.6±3.1	---	1.8±0.9
	30	---	---	1.8±1.8	---	0.4±0.4
20	(n)	59	62	54	53	228

^aAllelic designations refer to the number of repeats of the core sequence motif indicated in the locus column.

^bAlleles from all four racial groups were pooled for this column.

^cA dash indicates the absence of that allele in the respective sample.

^dRefers to the number of chromosomes samples.

[0071] Frequencies of some specific alleles (for example, allele 7 of HLTMT01 [AATG]_n and allele 17 of HUMARA [AGC]_n) are clearly variable across the four racial groups. Allele frequency distributions by race are given in Figure 4. With the exception of HUNFHPRTB (AGAT)_n, which is unimodal and symmetrical, the allele frequency distributions appear bimodal or more complex. The most common allele, however, appears to be the same for some loci (for example, HUMRENA4 (ACAG)_n), while at other loci predominant alleles do not coincide between races (for example, HUMARA (AGC)_n).

35 Most Frequent Genotypes

[0072] The frequencies of the most common genotypes of a DNA typing system reflect the utility of that assay in practice. The most frequent genotypes for the five STR loci have frequencies from 0.048 to 0.645 in the 20 STR-race combinations (Table 9). The most common genotypes for all five loci combined (p) have frequencies from 1.40 X 10⁻⁴ to 6.54 X 10⁻⁴ in the four racial groups.

TABLE 9 Most Frequent Genotype at the Five STR Loci in Four Populations

Locus	Whites	Blacks	Hispanics	Asians	Pooled
HUMHPRTB[AGAT] _a	0.227	0.177	0.309	0.209	0.228
HUMTH01[AATC] _a	0.190	0.138	0.172	0.225	0.118
HUMRENA4[ACAG] _a	0.645	0.507	0.581	0.486	0.553
HUMFABP1[AAT] _a	0.247	0.179	0.303	0.445	0.254
HUMARA[AGC] _n	0.069	0.063	0.048	0.064	0.042
Combined (P)	4.74X10 ⁻⁴	1.40X10 ⁻⁴	4.49X10 ⁻⁴	6.51X10 ⁻⁴	1.69X10 ⁻⁴
P ²	2.25X10 ⁻⁷	1.96X10 ⁻⁸	2.02X10 ⁻⁷	4.24X10 ⁻⁷	2.52X10 ⁻⁸

[0073] The match probability (P²) for the most common genotype of all five loci combined was 4.24 X 10⁻⁷. The frequencies of the least common genotypes for all five loci combined were on the order of 10⁻¹⁷. The probability calculations in Table 15 are only relevant for female individuals since two of the loci are X-linked. The most common male genotypes for all five loci combined have frequencies from 6.78 X 10⁻⁴ to 36.4 X 10⁻⁴ in the four racial groups.

[0074] The best markers for individualization in medicine and forensic science may be those with symmetrical and similar allele frequency distributions. Choice of the proper ethnic database appears less critical at such loci for the four ethnic populations we have studied.

5 [0075] The faint artifactual bands which are thought to arise during the PCR, assist in genotype determination relative to external standards. It is possible to count between lanes from allele to allele on overexposed autoradiograms such that even widely separated alleles differing by approximately 6 repeat units were accurately scored. The use of mPCR with fluorescent labels and internal standards improves upon this accuracy.

10 [0076] The data demonstrate that genotype data from trimeric and tetrameric tandem repeats were accurately and efficiently obtained via multiplex PCR. The fidelity with which trimeric and tetrameric STRs were amplified compared to the dimeric STRs make these new class of polymorphic markers well suited for application to DNA typing in forensic science and medicine.

Example 7

15 Fluorescent DNA profiling assay with internal standards

[0077] DNA typing is a powerful technique for determining the relationship, if any, between two genomic DNA samples. Applications for DNA typing include personal identification in paternity testing and forensic science, and sample source determinations in transplantation, prenatal diagnosis, and pedigree validation. Several features of polymorphic STRs 20 suggest that they could form the basis of a powerful and simple DNA typing assay. The small size of the amplified units allows several loci to be easily amplified simultaneously by mPCR, and analyzed with precise allele identification on DNA sequencing gels. The precision, sensitivity, and speed of detecting alleles with PCR offers special opportunities for the study of forensic specimens. For example, trimeric and tetrameric STRs show excellent fidelity of amplification indicating that the genotyping fingerprints can be easily interpreted and are amenable to automation. Fluorescent DNA 25 fragment detection can be used for internal size standards and precise allele quantitation.

30 [0078] For genetic typing, alleles from three chromosomally unlinked STR loci were amplified simultaneously in a mPCR (Figure 5). One primer from each of the three amplification primer sets is differentially labeled with one of the four fluorescent dyes used with the DNA sequencing device. In the ABI 370A system, one dye is reserved for the internal standards, while three dyes are available for the amplification products of STR loci. Theoretically, any given region of the sequencing gel can contain internal standards as well as alleles from three unlinked STR loci. Used to full potential the approach has enormous personal identification power of high accuracy.

35 [0079] Amplification incorporates a fluorescent label into one end, and a *Mlul* site into the other end of each product in the mPCR (Figure 5). Following amplification of the STR loci from a genomic DNA sample, residual activity of the *T. aquaticus* polymerase is destroyed and a homogeneous fragment length is achieved for each allele by digestion with *Mlul*. The treated multiplex products are then mixed with internal standards and loaded onto a sequencing gel for analysis 40 on an ABI 370A.

[0080] Internal standards were generated by pooling amplification products from individuals of known genotype such that the molar ratios of each allele observed were approximately equal. The pooled alleles were diluted, reamplified, and treated with *Mlul*. This scheme for generating internal standard size markers insures a virtually unlimited supply of 45 standards.

[0081] The combination of a quantitative detection system and mPCR enabled additional levels of internal control and precision. Using mPCR products synthesized under standardized amplification conditions, the fluorescent intensity of specific alleles at different loci was related. Because of the relationship between alleles of different loci, it was possible to distinguish between homozygosity and hemizygosity at a given locus (Figure 5). While failure of allele amplification 50 can occur by primer binding site polymorphism, the null was detectable by quantitation. This quantitative capacity removes the doubt which has been cast on the use of VNTRs due to the observation of homozygosity excess in population studies. The quantitative nature of the allele identification also facilitated the analysis of mixed body samples which occurs in forensics, prenatal diagnosis, the detection of chromosomal aneuploidy, and true somatic mosaicism seen in patients with chromosomal abnormalities and following bone marrow transplantation.

[0082] The average individualization potential (P_I) of the three loci together was one in 500 individuals. The combined genotype frequencies (three loci) of the individuals in panels A and B were 0.00026 and 0.0085 assuming Hardy-Weinberg equilibrium. The addition of three more loci will give a P_I of approximately one in 200,000, while the addition of six more loci will give a P_I of one in 90 million. Multiplex PCRs of this complexity have been done. Eight and nine genetic site mPCR for the hypoxanthine phosphoribosyltransferase and dystrophin genes are known.

[0083] Oligonucleotides were synthesized on an Applied Biosystems (ABI) 380B DNA synthesizer. Underivatized oligonucleotides were not purified after deprotection and lyophilization. ABI Aminolink 2 chemistry was used to derivatize oligonucleotides for biotin and fluorescent labeling, after which they were ethanol precipitated and purified by polyacrylamide gel electrophoresis. The fluorescent dyes (Molecular Probes, Eugene, OR) used in the assay were (i) NBD

aminoheanoic acid for all internal standard markers, (ii) 5-(and-6)-carboxyfluorescein succinimidyl ester for the HUMTH01 [AATG]_n and HUMHPRTB [AGAT]_n loci; and (iii) Texas Red™ sulfonyl chloride for the HUMFABP [AAT]_n locus. The primer sets had the first primer derivatized and the second primer containing an *M*lul restriction site. The primers used were: HUMTH01 (AATG)_n, (SEQ. ID NOS: 13,23); ;HUMFABP(ATT)_n (SEQ ID NOS; 5, 24); HUMHPRTB (AGAT)_n (SEQ ID. NOS; 19, 25). Simultaneous amplification with all six primers was performed with 25 cycles by denaturing at about 95°C for about 45 sec., annealing at about 60°C for about 30 sec., and extending at about 72°C for about 30 sec. using Perkin-Elmer-Cetus thermocyclers, amplitaq, and buffer conditions. The concentration of primers in the multiplex were about 0.06 μ M for HUMTH01 [AATG]_n, about 1.6 μ M for HUMFABP [AAT]_n, and about 0.56 μ M for HUMHPRTB [AGAT]_n. Following amplification, the products were phenol extracted, ethanol precipitated, and digested with *M*lul. The digested multiplex products were then combined with the internal size standards and electrophoresed through a 6.5% polyacrylamide, 8.3 M urea gel at 1300-1500 V, 24 mA and 32 W at a temperature of about 46°C. Internal size standards were prepared by amplifying specific alleles from individuals of known genotype. The products were quantitated, combined to give near equimolarity, diluted approximately 5000 fold, and reamplified with approximately 12 cycles.

[0084] In Figure 5A, fluorescent profiles of the internal standard cocktails when combined and electrophoresed in a single lane of an ABI 370A DNA sequencing device are shown. In Figure 5B the internal standards were combined with the amplification products of a multiplex PCR composed of (left to right) the HUMTH01 (AATG)_n, (I), HUMFABP (AAT)_n (II), and HUMHPRTB (AGAT)_n (III) loci. The individual shown is heterozygous for all three markets. In Figure 5C multiplex amplification from an individual homozygous at the HUMFABP (AAT)_n locus and hemizygous at HUMHPRTB (AGAT)_n is shown.

Example 8

Alternative detection schemes: radioactive, silver stain, intercalation

[0085] Since some forensic laboratories may not have access to fluorescent detection devices, the STR markers can be detected with non-denaturing and denaturing electrophoretic systems using alternative labeling and detection strategies. For example, radioactive and silver staining detection methods, and ethidium bromide staining methods are all applicable. The 6-15 STR loci are sorted into 4 to 5 separate mPCR reactions, each containing 2-4 loci. The three loci are selected such that the amplification products run in non-overlapping regions of the gel (i.e., the base pair lengths of the alleles from different loci do not overlap). Alleles from unknown samples are identified with reference to external standards in adjacent lanes (the same cocktails used in the fluorescent detection scheme (Figure 5) can be employed.

Example 9

Species Specificity

[0086] The species specificity of amplification of all STR loci can be determined. Primate DNAs, for example human, baboon, chimpanzee, gorilla, and various bacterial and yeast strains are compared. Additionally, *Drosophila*, common farm animals, common household pets, and common human flora are also examined. There is no difficulty in obtaining the samples since only 10 μ g of DNA are needed to perform over 100 studies. The high similarity of sequence between humans and other primates suggests that some of the loci amplify genomes from non-human primates. It is important to document which loci can be amplified from which species for optimal deployment of the method in the forensic arena. Amplification is not seen in non-primates.

Example 10

Kits

[0087] The kit includes a container having a oligonucleotide primer pair for amplifying short tandem repeats. The kit can also include standards. One kit includes standards and three oligonucleotide primer pairs. In a preferred embodiment the kit includes sufficient oligonucleotide primer pairs needed to perform mPCR for at least 6-10 loci. The kit can further include the reagents, and established protocols for using the kit. These kits provide for efficient and effective transfer and distribution of the method to the forensic community. The oligonucleotides and reaction mixtures in these kits can be stored at -70°C for extended periods of time. This facilitates mass production and quality control of the reagents needed to provide accurate reagents at a reasonable cost.

Example 11

Novel STR Sequence

5 [0088] A novel short tandem repeat sequence (SEQ ID. NO. 26) was identified from a lambda clone containing the X chromosome library by screening with a 30 base pair oligonucleotide of the sequence AGAT tandemly repeated. This locus was identified as HUMSTRX1. The sequence flanking the AGAT repeat was amplified and sequenced. Oligonucleotides were designed to amplify the AGAT repeat SEQ ID. NOS. 21 and 22.

10 [0089] The number of AGAT repeats is variable. The exact sequence length was inferred by length polymorphic short tandem repeat sequences with verification of the end sequences. The STR is between approximately base 153 and 203 of SEQ ID. NO. 26. The sense primer is between bases 61 and 84 and corresponds to SEQ ID. NO. 22 and the antisense primer is the reverse complement of the sequence between base 346 and 369 and corresponds to SEQ ID. NO. 21.

Example 12

15 Results and Benefits Expected

20 [0090] The novel methodology of the present invention provides the most powerful technique to date for the characterization of blood and other body fluids. The increase in credible evidence that this assay produces should result in an increase in the conviction rate for such violent crimes as sexual assault and murder. More importantly, many innocent suspects will be categorically cleared of false accusations. Additional applications would result in increased investigative power in the identification of missing persons, abducted children, military personnel, and human remains from natural and physical disasters. The sensitivity of body fluid identification methods would also be increased well beyond current limits. This would provide obvious benefits in the number of cases in which useful evidence was available.

25 [0091] Another significant improvement over the DNA technology currently used is the profound decrease in the amount of time required to provide results and the amount of labor required to produce the results. The time of actual testing for the new methodology is only 10% of the time required for existing DNA profiling techniques. Furthermore, existing technology is of limited investigative use in sexual assaults and homicides, due to the length of time required to obtain results.

30 [0092] The STR DNA profiling assay enables precise allele determination. The analysis of databases for locus stability, population heterogeneity, population allele frequencies, the Mendelian inheritance are greatly simplified. The collection of data from the fluorescent STR profiling assay lends itself to automation, thereby reducing the chance of operator error. Defined discrete allele designations promote generation of a national databank.

35 [0093] The loci developed, the profiling assay methods, and the population studies are of interest to the general scientific community. DNA profiling has direct application in the medical diagnostic and research laboratories for verifying specimen identity.

SEQUENCE LISTING

40 [0094]

(1) GENERAL INFORMATION:

45 (i) APPLICANT: Albert O. Edwards and
Charles Thomas caskey

(ii) TITLE OF INVENTION: DNA profiling with short tandem repeat polymorphisms and identification of polymorphic STRs

50 (iii) NUMBER OF SEQUENCES: 26

(iv) CORRESPONDENCE ADDRESS:

55 (A) ADDRESSEE: Fulbright & Jaworski Patent Department

(B) STREET: 1301 McKinney, Suite 5100

(C) CITY: Houston

(D) STATE: Texas

(E) COUNTRY: U.S.A.

5 (F) ZIP: 77010-3095

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Disk, 3.5 inch (1.44MB)

10 (B) COMPUTER: IBM PC/AT

(C) OPERATING SYSTEM: MS-DOS

15 (D) SOFTWARE: BASIC

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

20 (B) FILING DATE:

(C) CLASSIFICATION:

25 (vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER:

30 (B) FILING DATE:

(viii) ATTORNEY/AGENT INFORMATION:

(A) AME: THOMAS D. PAUL

35 (B) REGISTRATION NUMBER: 32,714

(C) REFERENCE/DOCKET NUMBER: D-5217

(ix) TELECOMMUNICATION INFORMATION:

40 (A) TELEPHONE: (713) 651-5325

(B) TELEFAX: (713) 651-5246

45 (C) TELEX: WESTERN UNION 762829

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

50 (A) LENGTH: 52

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

55 (ii) MOLECULE TYPE: Synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

ACTGCAGAGA	CGCTGTCTGT	CGAAGGTAAG	40
GAACGGACGA	GAGAAGGGAG	AG	52

5

(3) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 52
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

15 (ii) MOLECULE TYPE: Synthetic DNA
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CTCTCCCTTC	TCGAATCGTA	ACCGTTCGTA	40
CGAGAACATCGC	TGTCTCTGCA	GT	52

(4) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 38
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

30 (ii) MOLECULE TYPE: Synthetic DNA
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

35 GCCGGATCCC	GAATCGTAAC	CGTTCGTACG	30
AGAACATCGC		38	

(40 (5) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

45 (A) LENGTH: 24
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

50 (ii) MOLECULE TYPE: Synthetic DNA
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:
 TACGAGAACATC GCTGTCTCTG CAGT 24

(6) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

55 (A) LENGTH: 25
 (B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

5 (ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:
GTAGTATCAG TTTCATAGGG TCACC 25

(7) INFORMATION FOR SEQ ID NO: 6:

10 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear
15 (ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:
CAGTTCGTTT CCATTGTCTG TCCG 24

20 (8) INFORMATION FOR SEQ ID NO: 7:

25 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear
30 (ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:
TCCAGAAATCT GTTCCAGAGC GTGC 24

35 (9) INFORMATION FOR SEQ ID NO: 8:

40 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear
45 (ii) MOLECULE TYPE: Genomic DNA (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:
GCTGTGAAGG TTGCTGTTCC TCAT 24

(10) INFORMATION FOR SEQ ID NO: 9:

50 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear
55 (ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:
TGTGAGTCCC AGTTGCCAGT CTAC 24

(11) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

5 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

10 (ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:
ACTGGTCACC TTGGAAAGTG GCAT 24

(12) INFORMATION FOR SEQ ID NO: 11:

15 (i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

25 (ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11: TGAGGGCTGT ATGGAATACG TTCA 24

(13) INFORMATION FOR SEQ ID NO: 12:

30 (i) SEQUENCE CHARACTERISTICS:

35 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:
CAAGCACCAA GCTGAGCAAA CAGA 24

(14) INFORMATION FOR SEQ ID NO: 13:

40 (i) SEQUENCE CHARACTERISTICS:

45 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:
GTGGGCTGAA AAGCTCCGAT TAT 23

(15) INFORMATION FOR SEQ ID NO: 14:

55 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:
5 ATTCAAAGGG TATCTGGGCT CTGG 24

(16) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

15 (ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:
GGAGAGACAG GATGTCTGGC ACAT 24

20 (17) INFORMATION FOR SEQ ID NO: 16:

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

30 (ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:
CCATCTCTCT CCTTAGCTGT CATA 24

(18) INFORMATION FOR SEQ ID NO: 17:

35 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
40 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:
45 AGAGTACCTT CCCTCCTCTA CTCA 24

(19) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:

50 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

55 (ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:
CTCTATGGAG CTGGTAGAAC CTGA 24

(20) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

5 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

10 (ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:
ATGCCACAGA TAATACACAT CCCC 24

(21) INFORMATION FOR SEQ ID NO: 20:

15 (i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

25 (ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:
CTCTCCAGAA TAGTTAGATG TAGG 24

(22) INFORMATION FOR SEQ ID NO: 21:

30 (i) SEQUENCE CHARACTERISTICS:

35 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:
CTCCTTGTGG CCTTCCTTAA ATGG 24

40 (23) INFORMATION FOR SEQ ID NO: 22:

(i) SEQUENCE CHARACTERISTICS:

45 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:
CTTCTCCAGC ACCCAAGGAA GTCA 24

(24) INFORMATION FOR SEQ ID NO: 23:

55 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32
(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

5 (ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:
TTACGCGTAT TCAAAGGGTA TCTGGGCTCT GG 32

(25) INFORMATION FOR SEQ ID NO: 24:

10 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

15 (ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:
TTACGCGTCT CGGACAGTAT TCAGTTCGTT TC 32

20 (26) INFORMATION FOR SEQ ID NO: 25:

25 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 34
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

30 (ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:
TTACGCGTTC TCCAGAATAG TTAGATGTAG GTAT 34

(27) INFORMATION FOR SEQ ID NO: 26:

35 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 504
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

40 (ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

45 50

55

5	TGTTTGT	GT	TTTTTTAAA	TCTGTTCTCA	40
	TTGGTGTTC	TG	CTTCTCCAGC	ACCCAAGGAA	80
	GTCACCA	TATTGTCCT	TAGTCCTGTG	TTTCTTAGCT	120
	GGTCTGCC	CTTCTCTCA	CTTTTCAGAG	TCAGATAGAT	160
	GATAGATAGA	TAGATAGATA	GATAGATAGA	TAGATAGATA	200
	GATATAAAA	GATAAATAGA	TAGTCTCTAT	ACAGATATAG	240
	ATGTATATCA	TTCCAAGTTT	TAGCTGTATT	TGCAGGAGGA	280
	AACGAGAAGT	ATGTCTACTT	CTTTTCCCTCT	GGAACCAAGAG	320
	GTTTCCCCTT	TGTTACCTTG	TTTTTCCATT	TAAGGAAGGC	360
	CACAAGGAGA	ACCATGAATC	TGCTCTAATT	GATTTTTACG	400
	AAAGGAAGGA	AAAAAAGACA	GGAAATAATT	CACAATCTCC	440
10	AACTCTTCA	TCGTAATTAG	TGAGTGACAA	GTAGTTGTA	480
	ACACTCTCAG	TGTATCTTGA	TAAT		504

15

SEQUENCE LISTING

[0095]

20 (1) GENERAL INFORMATION:

(i) APPLICANT: Albert O. Edwards and
Charles Thomas Caskey

25 (ii) TITLE OF INVENTION: DNA profiling with short tandem repeat polymorphisms and identification of polymorphic STRs

(iii) NUMBER OF SEQUENCES: 26

30 (iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Fulbright & Jaworski Patent Department

(B) STREET: 1301 McKinney, Suite 5100

35 (C) CITY: Houston

(D) STATE: Texas

40 (E) COUNTRY: U.S.A.

(F) ZIP: 77010-3095

(v) COMPUTER READABLE FORM:

45 (A) MEDIUM TYPE: Disk,3.5 inch (1.44MB)

(B) COMPUTER: IBM PC/AT

50 (C) OPERATING SYSTEM: MS-DOS

(D) SOFTWARE: BASIC

(vi) CURRENT APPLICATION DATA:

55 (A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

5 (A) APPLICATION NUMBER:
(B) FILING DATE:

(viii) ATTORNEY/AGENT INFORMATION:

10 (A) AME: THOMAS D. PAUL
(B) REGISTRATION NUMBER: 32,714
(C) REFERENCE/DOCKET NUMBER: D-5217

(ix) TELECOMMUNICATION INFORMATION:

15 (A) TELEPHONE: (713) 651-5325
(B) TELEFAX: (713) 651-5246
(C) TELEX: WESTERN UNION 762829

20 (2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 52
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Synthetic DNA

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

ACTGCAGAGA CGCTGTCTGT CGAAGGTAAG 40

35 GAACGGACGA GAGAAGGGAG AG 52

(3) INFORMATION FOR SEQ ID NO: 2:

40 (i) SEQUENCE CHARACTERISTICS:

45 (A) LENGTH: 52
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Synthetic DNA

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

50 CTCTCCCTTC TCGAATCGTA ACCGTTCGTA 40

CGAGAATCGC TGTCTCTGCA GT 52

55 (4) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 38
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

5

- (ii) MOLECULE TYPE: Synthetic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

10 GCCGGATCCC GAATCGTAAC CGTCGTACG 30
 AGAACATCGC 38

15 (5) INFORMATION FOR SEQ ID NO: 4:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

20 (ii) MOLECULE TYPE: Synthetic DNA
25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:
 TACGAGAATC GCTGTCTCTG CAGT 24

(6) INFORMATION FOR SEQ ID NO: 5:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

- (ii) MOLECULE TYPE: Genomic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:
 GTAGTATCAG TTTCATAGGG TCACC 25

40

(7) INFORMATION FOR SEQ ID NO: 6:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

- (ii) MOLECULE TYPE: Genomic DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:
 CAGTCGTTT CCATTGTCTG TCCG 24

50

(8) INFORMATION FOR SEQ ID NO: 7:

55

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24

- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

5 (ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:
TCCAGAACATCT GTTCCAGAGC GTGC 24

(9) INFORMATION FOR SEQ ID NO: 8:

10 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: Nucleic Acid
15 (C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:
20 GCTGTGAAGG TTGCTGTTCC TCAT 24

(10) INFORMATION FOR SEQ ID NO: 9:

25 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
30 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:
TGTGAGTCCC AGTTGCCAGT CTAC 24

35 (11) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
40 (B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:
ACTGGTCACC TTGGAAAGTG GCAT 24

(12) INFORMATION FOR SEQ ID NO: 11:

50 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
55 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

TGAGGGCTGT ATGGAATACG TTCA 24

(13) INFORMATION FOR SEQ ID NO: 12:

5 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

10 (ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:
CAAGCACCAA GCTGAGCAAA CAGA 24

15 (14) INFORMATION FOR SEQ ID NO: 13:

20 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

25 (ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:
GTGGGCTGAA AAGCTCCCGAT TAT 23

30 (15) INFORMATION FOR SEQ ID NO: 14:

35 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

40 (ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:
ATTCAAAGGG TATCTGGGCT CTGG 24

(16) INFORMATION FOR SEQ ID NO: 15:

45 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

50 (ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:
GGAGAGACAG GATGTCTGGC ACAT 24

55 (17) INFORMATION FOR SEQ ID NO: 16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

5

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:
CCATCTCT CCTTAGCTGT CATA 24

10

(18) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:

15

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

20

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:
AGAGTACCTT CCCTCCTCTA CTCA 24

25

(19) INFORMATION FOR SEQ ID NO: 18:

30

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

35

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:
CTCTATGGAG CTGGTAGAAC CTGA 24

35

(20) INFORMATION FOR SEQ ID NO: 19:

40

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

45

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:
ATGCCACAGA TAATACACAT CCCC 24

50

(21) INFORMATION FOR SEQ ID NO: 20:

55

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:
CTCTCCAGAA TAGTTAGATG TAGG 24

(22) INFORMATION FOR SEQ ID NO: 21:

5 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: Nucleic Acid
10 (C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:
15 CTCCTTGTGG CCTTCCTTAA ATGG 24

(23) INFORMATION FOR SEQ ID NO: 22:

20 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
25 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:
CTTCTCCAGC ACCCAAGGAA GTCA 24

30 (24) INFORMATION FOR SEQ ID NO: 23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32
35 (B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:
TTACGCGTAT TCAAAGGGTA TCTGGGCTCT GG 32

(25) INFORMATION FOR SEQ ID NO: 24:

45 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32
(B) TYPE: Nucleic Acid
50 (C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:
TTACGCGTCT CGGACAGTAT TCAGTTCGTT TC 32

55 (26) INFORMATION FOR SEQ ID NO: 25:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 34
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

5
 (ii) MOLECULE TYPE: Genomic DNA
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:
 TTACGCGTTC TCCAGAATAG TTAGATGTAG GTAT 34

10 (27) INFORMATION FOR SEQ ID NO: 26:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 504
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear
 20 (ii) MOLECULE TYPE: Genomic DNA
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

25	TGTTTGT	TTTTGTTGT	TTTTTTAAA	TCTGTTCTCA	40
	TTGGTGT	TGTTTGCTGC	CTTCTCCAGC	ACCCAAGGAA	80
	GTCACCACCA	TATTGTTCC	TAGCCTGTG	TTTCTTAGCT	120
	GGTCTGCCTT	CTTCTCTCCA	CTTTTCAGAG	TCAGATAGAT	160
30	GATAGATAGA	TAGATAGATA	GATAGATAGA	TAGATAGATA	200
	GATATAAAAAA	GATAAATAGA	TAGTCTCTAT	ACAGATATAG	240
	ATGTATATCA	TTCCAAGTT	TAGCTGTATT	TGCAGGAGGA	280
35	AACGAGAAGT	ATGTCTACTT	CTTTTCCTCT	GGAACCAGAG	320
	GTTTCCCCTT	TGTTACCTTG	TTTTCCATT	TAAGGAAGGC	360
	CACAAGGAGA	ACCATGAATC	TGCTCTAATT	GATTTTACG	400
40	AAAGGAAGGA	GAAAAAGACA	GGAAATAATT	CACAATCTCC	440
	AACTCTTCA	TCGTAATTAG	TGAGTGACAA	GTAGTTGTA	480
	ACACTCTCAG	TGTATCTTGA	TAAT		504

45

SEQUENCE LISTING

[0096]

50 (1) GENERAL INFORMATION:

(i) APPLICANT: Albert O. Edwards and Charles Thomas caskey
 55 (ii) TITLE OF INVENTION: DNA profiling with short tandem repeat polymorphisms and identification of polymorphic STRs
 (iii) NUMBER OF SEQUENCES: 26

(iv) CORRESPONDENCE ADDRESS:

- (A) ADDRESSEE: Fulbright & Jaworski Patent Department
- 5 (B) STREET: 1301 McKinney, Suite 5100
- (C) CITY: Houston
- 10 (D) STATE: Texas
- (E) COUNTRY: U.S.A.
- (F) ZIP: 77010-3095

15 (v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Disk,3.5 inch (1.44MB)
- 20 (B) COMPUTER: IBM PC/AT
- (C) OPERATING SYSTEM: MS-DOS
- (D) SOFTWARE: BASIC

25 (vi) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER:
- 30 (B) FILING DATE:
- (C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

35 (A) APPLICATION NUMBER:

- (B) FILING DATE:

40 (viii) ATTORNEY/AGENT INFORMATION:

- (A) AME: THOMAS D. PAUL
- (B) REGISTRATION NUMBER: 32,714
- 45 (C) REFERENCE/DOCKET NUMBER: D-5217

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: (713) 651-5325
- 50 (B) TELEFAX: (713) 651-5246
- (C) TELEX: WESTERN UNION 762829

55 (2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 52
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

5

(ii) MOLECULE TYPE: Synthetic DNA
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

10	ACTGCAGAGA	CGCTGTCTGT	CGAAGGTAAG	40
	GAACGGACGA	GAGAAGGGAG	AG	52

(3) INFORMATION FOR SEQ ID NO: 2:

15

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 52
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Synthetic DNA
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

25

	CTCTCCCTTC	TCGAATCGTA	ACCGTTCGTA	40
	CGAGAACATCGC	TGTCTCTGCA	GT	52

30

(4) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 38
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

40

(ii) MOLECULE TYPE: Synthetic DNA
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

45	GCCGGATCCC	GAATCGTAAC	CGTTCGTACG	30
	AGAACATCGC		38	

(5) INFORMATION FOR SEQ ID NO: 4:

50

(i) SEQUENCE CHARACTERISTICS:

55

(A) LENGTH: 24
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Synthetic DNA
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

TACGAGAATC GCTGTCTCTG CAGT

24

(6) INFORMATION FOR SEQ ID NO: 5:

5 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

10 (ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:
GTAGTATCAG TTTCATAGGG TCACC 25

15 (7) INFORMATION FOR SEQ ID NO: 6:

20 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

25 (ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:
CAGTTCGTTT CCATTGTCTG TCCG 24

30 (8) INFORMATION FOR SEQ ID NO: 7:

35 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

40 (ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:
TCCAGAAATCT GTTCCAGAGC GTGC 24

(9) INFORMATION FOR SEQ ID NO: 8:

45 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

50 (ii) MOLECULE TYPE: Genomic DNA (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GCTGTGAAGG TTGCTGTTCC TCAT 24

(10) INFORMATION FOR SEQ ID NO: 9:

55 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24

- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

5 (ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:
TGTGAGTCCC AGTTGCCAGT CTAC 24

10 (11) INFORMATION FOR SEQ ID NO: 10:

- (i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:
20 ACTGGTCACC TTGGAAAGTG GCAT 24

(12) INFORMATION FOR SEQ ID NO: 11:

- (i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:
30 TGAGGGCTGT ATGGAATACG TTCA 24

35 (13) INFORMATION FOR SEQ ID NO: 12:

- (i) SEQUENCE CHARACTERISTICS:

40 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:
45 CAAGCACCAA GCTGAGCAAA CAGA 24

(14) INFORMATION FOR SEQ ID NO: 13:

50 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

GTGGGCTGAA AAGCTCCGAT TAT 23

(15) INFORMATION FOR SEQ ID NO: 14:

5 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

10 (ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:
ATTCAAAGGG TATCTGGGCT CTGG 24

15 (16) INFORMATION FOR SEQ ID NO: 15:

20 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

25 (ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:
GGAGAGACAG GATGTCTGGC ACAT 24

30 (17) INFORMATION FOR SEQ ID NO: 16:

35 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

40 (ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:
CCATCTCTCT CCTTAGCTGT CATA 24

(18) INFORMATION FOR SEQ ID NO: 17:

45 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

50 (ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:
AGAGTACCTT CCCTCCTCTA CTCA 24

55 (19) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

5
(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:
CTCTATGGAG CTGGTAGAAC CTGA 24

10 (20) INFORMATION FOR SEQ ID NO: 19:

- (i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:
ATGCCACAGA TAATACACAT CCCC 24

(21) INFORMATION FOR SEQ ID NO: 20:

25 (i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:
CTCTCCAGAA TAGTTAGATG TAGG 24

35 (22) INFORMATION FOR SEQ ID NO: 21:

- (i) SEQUENCE CHARACTERISTICS:

40 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:
CTCCTTGTGG CCTTCCTTAA ATGG 24

(23) INFORMATION FOR SEQ ID NO: 22:

50 (i) SEQUENCE CHARACTERISTICS:

55 (A) LENGTH: 24
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:
CTTCTCCAGC ACCCAAGGAA GTCA 24

(24) INFORMATION FOR SEQ ID NO: 23:

5 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32
(B) TYPE: Nucleic Acid
10 (C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:
15 TTACGCGTAT TCAAAGGGTA TCTGGGCTCT GG 32

(25) INFORMATION FOR SEQ ID NO: 24:

20 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32
(B) TYPE: Nucleic Acid
25 (C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:
TTACGCGTCT CGGACAGTAT TCAGTTCGTT TC 32

30 (26) INFORMATION FOR SEQ ID NO: 25:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 34
35 (B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:
TTACGCGTTC TCCAGAATAG TTAGATGTAG GTAT 34

(27) INFORMATION FOR SEQ ID NO: 26:

45 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 504
(B) TYPE: Nucleic Acid
50 (C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

	TGTTTGT	GTTTGTTGT	TTTTTTAAA	TCTGTTCTCA	40
	TTGGTGT	TGTTTGCTGC	CTTCTCCAGC	ACCCAAGGAA	80
5	GTCACCACCA	TATTGTTCTT	TAGTCCTGTG	TTTCTTAGCT	120
	GGTCTGCCCT	CTTCTCTCCA	CTTTTCAGAG	TCAGATAGAT	160
	GATAGATAGA	TAGATAGATA	GATAGATAGA	TAGATAGATA	200
	GATATAAAA	GATAAATAGA	TAGTCTCTAT	ACAGATATAG	240
	ATGTATATCA	TTCCAAGTTT	TAGCTGTATT	TGCAGGAGGA	280
	AACGAGAAGT	ATGTCCTACTT	CTTTTCTCTT	GGAACCAGAG	320
10	GTTTCCCTT	TGTTACCTTG	TTTTTCCATT	TAAGGAAGGC	360
	CACAAGGAGA	ACCATGAATC	TGCTCTAATT	GATTTTTACG	400
	AAAGGAAGGA	GAAAAAAGACA	GGAAATAATT	CACAATCTCC	440
	AACTCTTCA	TCGTAATTAG	TGAGTGACAA	GTAGTTTGTA	480
	ACACTCTCAG	TGTATCTTGA	TAAT		504

15

Claims

20 1. A kit for a DNA profiling assay comprising a container having oligonucleotide primer pairs for amplifying a short tandem repeat sequence, comprising primers with the sequence of SEQ ID 13 and SEQ ID 14.

2. A kit for a DNA profiling assay comprising a container having oligonucleotide primer pairs for amplifying a short tandem repeat sequence, comprising primers with the sequence of SEQ ID 13 and SEQ ID 23.

25 3. A kit according to claim 1 or 2 further comprising a labelled standard.

4. A kit according to one of claims 1 to 3, further comprising a container having reagents for multiplex polymerase chain reaction.

30

Patentansprüche

35 1. Set für einen DNA-Fingerprinting-Test, umfassend einen Behälter mit Oligonucleotidprimerpaaren zum Amplifizieren einer kurzen Tandemwiederholungssequenz, die Primer mit der Sequenz von Seq.-ID Nr. 13 und Seq.-ID Nr. 14 umfasst.

2. Set für einen DNA-Fingerprinting-Test, umfassend einen Behälter mit Oligonucleotidprimerpaaren zum Amplifizieren einer kurzen Tandemwiederholungssequenz, die Primer mit der Sequenz von Seq.-ID Nr. 13 und Seq.-ID Nr. 23 umfasst.

40 3. Set nach Anspruch 1 oder 2, das weiters einen markierten Standard umfasst.

4. Set nach einem der Ansprüche 1 bis 3, das weiters einen Behälter mit Reagenzien für die Multiplex-Polymerasekettenreaktion umfasst.

45

Revendications

50 1. kit pour un essai de profilage d'ADN comprenant un contenant ayant des paires d'amorces d'oligonucléotides pour l'amplification avec une courte séquence de répétition tandem, comprenant des amorces ayant la séquence de SEQ ID 13 et SEQ ID 14.

2. Kit pour un essai de profilage d'ADN comprenant un contenant ayant des paires d'amorces d'oligonucléotides pour l'amplification d'une courte séquence de répétition tandem, comprenant des amorces avec la séquence de SEQ ID 13 et SEQ ID 23.

55 3. Kit selon la revendication 1 ou 2, comprenant en outre un standard marqué.

4. Kit selon l'une quelconque des revendications 1 à 3, comprenant en outre un contenant ayant des réactifs pour une réaction en chaîne par polymérase multiplexe.

5

10

15

20

25

30

35

40

45

50

55




FIGURE 1




Figure 2A




Figure 2B



Figure 2C




Figure 3A



Figure 3B




Figure 3C

Frequency Distribution of HUMFABP[AAT] Alleles




FIGURE 4A




FIGURE 4B




FIGURE 4C




FIGURE 4D




FIGURE 4E




FIG. 5A




FIG. 5B




FIG. 5C

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Non-patent literature cited in the description

- **Riley et al.** *Nucleic Acids Res*, 1990, vol. 18, 2887
[0049]