

(19)



(11)

EP 1 641 801 B9

(12)

CORRECTED EUROPEAN PATENT SPECIFICATION

(15) Correction information:

Corrected version no 1 (W1 B1)
Corrections, see
Claims EN

(51) Int Cl.:

C07D 487/08 ^(2006.01) **C07D 451/02** ^(2006.01)
C07D 451/14 ^(2006.01) **A61K 31/46** ^(2006.01)

(48) Corrigendum issued on:

25.02.2009 Bulletin 2009/09

(86) International application number:

PCT/US2004/014671

(45) Date of publication and mention
of the grant of the patent:

16.07.2008 Bulletin 2008/29

(87) International publication number:

WO 2004/101562 (25.11.2004 Gazette 2004/48)

(21) Application number: **04751862.6**

(22) Date of filing: **11.05.2004**

(54) **BRIDGED N-ARYLSULFONYLPIPERIDINES AS GAMMA-SECRETASE INHIBITORS**

VERBRÜCKTE N-ARYLSULFONYLPIPERIDINE ALS GAMMA-SEKRETASEINHIBITOREN

N-ARYLSULFONYLPIPERIDINES PONTEES UTILISEES COMME INHIBITEURS DE LA
GAMMA-SECRETASE

(84) Designated Contracting States:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PL PT RO SE SI SK TR
Designated Extension States:
AL HR LT LV MK

(74) Representative: **Adams, Harvey Vaughan John et al**

Mathys & Squire LLP
120 Holborn
London
EC1N 2SQ (GB)

(30) Priority: **13.05.2003 US 470146 P**

(43) Date of publication of application:

05.04.2006 Bulletin 2006/14

(56) References cited:

WO-A-00/50391

(73) Proprietor: **Schering Corporation**
Kenilworth, NJ 07033-0530 (US)

- **MOMOSE, TAKEFUMI ET AL: "Reaction of N-benzenesulfonyl-9-azabicyclo[3.3.1]nona-2,6-diene with diborane: failure of cyclic hydroboration in the intramolecularly faced diene system" HETEROCYCLES, 6(4), 469-74 CODEN: HTCYAM; ISSN: 0385-5414, 1977, XP009041109**

(72) Inventor: **JOSIEN, Hubert**
Hoboken, NJ 07030 (US)

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 641 801 B9

Description

CROSS REFERENCE TO RELATED APPLICATION

[0001] This application claims the benefit of U.S. Provisional Application No. 60/470,146, filed May 13, 2003.

BACKGROUND OF THE INVENTION


[0002] WO 00/50391, published August 13, 2000, discloses compounds having a sulfonamide moiety that are useful for the treatment and prevention of Alzheimer's disease and other diseases relating to the deposition of amyloid protein.

[0003] In view of the present interest in the treatment or prevention of neurodegenerative diseases, such as Alzheimer's disease, a welcome contribution to the art would be compounds for use in such treatment or prevention. This invention provides such a contribution.

SUMMARY OF THE INVENTION

[0004] Disclosed herein are novel compounds as inhibitors of gamma secretase, methods for preparing such compounds, pharmaceutical compositions comprising one or more such compounds, methods for preparing pharmaceutical formulations comprising one or more such compounds, and methods of treatment, prevention, inhibition or amelioration of one or more diseases associated with gamma secretase using such compounds or pharmaceutical compositions.

[0005] Disclosed herein is a compound, and/or pharmaceutically acceptable salts or solvates of said compound, said compound having the general structure shown in Formula I:

Formula I

wherein:

Ar is aryl or heteroaryl;

X is O, NH, or NR⁶;

m is 0, 1, 2 or 3, provided that when m = 0 no double bond is present and when

m > 0 a double bond may or may not be present or a triple bond may or may not be present;

n is 0, 1, 2 or 3;

R¹ is 1 to 3 substituents independently selected from the group consisting of: hydrogen, alkyl, cycloalkyl, halogen, -CF₃, -OCF₃, -NR⁶R⁷, -CN, -NO₂, -NH₂, -alkylene-NR⁶R⁷, -alkylene-O-alkyl, -C(O)O-alkyl, -C(O)NR⁶R⁷, -N(R⁶)C(O)alkyl, -N(R⁶)C(O)cycloalkyl, -N(R⁶)C(O)aryl,

$-N(R^6)C(O)heteroaryl$, $-N(R^6)C(O)O-alkyl$, $-N(R^6)C(O)NR^6R^7$, $-N(R^6)S(O)_2alkyl$, $-OH$, $-O-alkyl$, and $-O-cycloalkyl$;


R^2 is hydrogen, $-alkyl$, $-cycloalkyl$, $-alkylene-cycloalkyl$, $-cycloalkyl$, $-NR^6R^7$, $-alkylene-NR^6R^7$, $-alkylene-O-alkyl$, $-O-alkyl$, $-heteroaryl$, or $-aryl$;

R^3 is 1 to 6 substituents, which can be the same or different, each being independently selected from hydrogen, halogen, $-alkyl$, $-cycloalkyl$, $-OH$, $-OCF_3$, $-CF_3$, $-O-alkyl$, $-O-cycloalkyl$, or $-NR^6R^7$;


R^4 is hydrogen, $-alkyl$, $-cycloalkyl$, $-aryl$ or $-heteroaryl$; or R^3 and R^4 can be joined together to form a 3 to 6 member ring;

R^5 is $-NR^6R^7$, $-N(R^6)-alkylene-NR^6R^7$, $-alkyl$, $-cycloalkyl$, $-aryl$, $-heteroaryl$, $-cycloalkyl$, $-alkylene-aryl$, $-alkylene-heteroaryl$, $-cycloalkylene-aryl$, $-cycloalkylene-heteroaryl$, $-heterocycloalkyl-aryl$ or $-heterocycloalkyl-heteroaryl$; or

R^5 is selected from the group consisting of:



R^6 and R^7 can be the same or different, each being independently selected from the group consisting of: hydrogen, $-aryl$, $-heteroaryl$, $-alkyl$, $-cycloalkyl$, $-alkylene-aryl$, $-alkylene-heteroaryl$,



r is 0, 1, 2, 3 or 4;

s is 0, 1, 2 or 3;

R^8 is hydrogen, $-OH$, $-alkyl$, $-C(O)NR^6R^7$, $-C(O)O-alkyl$, or $aryl$, wherein said alkyl moiety may be unsubstituted or substituted with 1 to 4 hydroxy groups, and said aryl group may be unsubstituted or substituted with at least one substituent selected from the group consisting of $alkyl$, $cycloalkyl$, $halogen$, $-CF_3$, $-OCF_3$, $-NR^6R^7$, $-CN$, $-NO_2$, $-NH_2$, $-alkylene-NR^6R^7$, $-alkylene-O-alkyl$, $-C(O)O-alkyl$, $-C(O)NR^6R^7$, $-N(R^6)C(O)alkyl$; $-N(R^6)C(O)cycloalkyl$, $-N(R^6)C(O)aryl$, $-N(R^6)C(O)heteroaryl$; $-N(R^6)C(O)O-alkyl$, $-N(R^6)C(O)NR^6R^7$, $-N(R^6)S(O)_2alkyl$, $-OH$; $-O-alkyl$, and $-O-cycloalkyl$;

R^9 is selected from the group consisting of: hydrogen, $-alkyl$; $-cycloalkyl$, $-alkylene-aryl$, $-alkylene-heteroaryl$, $-alkyl-cycloalkyl$, $-C(O)O-alkyl$, $-alkylene-O-alkylene-OH$, $-aryl$ and $-heteroaryl$, wherein each-said alkyl and cycloalkyl moieties may be independently substituted or unsubstituted and if substituted then substituted with 1 to 4 hydroxy groups or at least one $-NR^6R^7$ group;

each R^{10} is independently selected from hydrogen or alkyl wherein said alkyl group may be substituted or

each R^{11} and R^{12} unsubstituted and if substituted then R^{10} is substituted with 1 to 4 hydroxy groups;
 are independently selected from hydrogen, halogen, -alkyl, -cycloalkyl, -OH, -OCF₃, -CF₃, -O-alkyl, -O-cycloalkyl, -NR⁶R⁷ or aryl; or
 R^{11} and R^{12} , together with the ring carbon atoms to which they are shown attached in Formula I, form a cycloalkyl ring, and wherein the cycloalkyl ring formed thereby can be unsubstituted or substituted with an alkyl, alkoxy, aryl, or halogen;
 p is 0, 1, 2, 3 or 4; and
 q is 0, 1 or 2.

[0006] Unless specifically stated, the moieties alkyl, aryl, heteroaryl, cycloalkyl, alkylene and heterocycloalkyl (also referred to as heterocyclyl) in the above-noted definitions can be unsubstituted or each be optionally independently substituted with one or more suitable groups which can be the same or different and are independently selected. Non-limiting examples of such suitable groups as substituents are provided in the definitions that are in the Detailed Description section below. The possibility of such non-substitution as well as independent optional substitution in the moieties above is well within the scope of this disclosure.

[0007] This disclosure teaches a pharmaceutical composition comprising a therapeutically effective amount of one or more compounds of Formula I and one or more pharmaceutical acceptable carriers.

[0008] This disclosure teaches a method for inhibiting gamma-secretase comprising administering a therapeutically effective amount of one or more compounds of Formula I to a patient in need of such inhibition.

[0009] This disclosure teaches a method of treating one or more neurodegenerative diseases comprising administering a therapeutically effective amount of one or more compounds of Formula I to a patient in need of treatment.

[0010] This disclosure teaches a method of inhibiting the deposition of amyloid protein (e.g., amyloid β protein) in, on, or around neurological tissue (e.g., the brain) comprising administering a therapeutically effective amount of one or more compounds of Formula I to a patient in need of treatment.

[0011] This disclosure teaches a method of treating Alzheimer's disease comprising administering a therapeutically effective amount of one or more compounds of Formula I to a patient in need of treatment

DETAILED DESCRIPTION OF THE INVENTION

[0012] In one aspect, the present invention relates to a compound, or pharmaceutically acceptable salts or solvates of said compound, said compound having the structure set out in the appended claim 1. Other aspects of the invention are set out in the appended claims.

[0013] In another preferred embodiment, m is 0 or 1.

[0014] In another preferred embodiment, n is 1 or 2.


[0015] In another preferred embodiment, R^2 is hydrogen.

[0016] In another preferred embodiment, R^3 is hydrogen.

[0017] In another preferred embodiment, R^4 is hydrogen or CH₃.

[0018] In an additional preferred embodiment, R^1 is 4-halo, more preferably 4-chloro.

[0019] In an additional preferred embodiment, R^5 is:




[0020] In an additional preferred embodiment, one double bond exists within a bicyclic ring in the structure represented by Formula I.

[0021] In an additional preferred embodiment, m = 1 and n = 2.

[0022] In an additional preferred embodiment, m = 1 and n = 1.

[0023] In an additional preferred embodiment, m = 0 and n = 1.

[0024] In an additional preferred embodiment, R^5 is



[0025] In an additional preferred embodiment, R^{11} and R^{12} are both H.


[0026] In an additional preferred embodiment;

R^2 is H;

R^3 is H;


R^4 is H; and


R^5 is:



[0027] In an additional preferred embodiment, m is 1, no double or triple bond is present, and R^{11} and R^{12} , together with the ring carbon atoms to which they are shown attached in Formula I, form a cyclopropyl ring.

[0028] Some preferred compounds of the invention include:





5

10

15

20

25

30


35


40

45

50

55





[0029] Except where stated otherwise, the following definitions apply throughout the present specification and claims. These definitions apply regardless of whether a term is used by itself or in combination with other terms. Hence the definition of "alkyl" applies to "alkyl" as well as to the "alkyl" portions of "heteroarylalkyl", "-N(R⁶)C(O)O-alkyl, etc.

[0030] As used above, and throughout the specification, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

"Patient" includes both human and animals.

"Mammal" means humans and other mammalian animals.

"Alkyl" means an aliphatic hydrocarbon group which may be straight or branched and comprising about 1 to about 20 carbon atoms in the chain. Preferred alkyl groups contain about 1 to about 12 carbon atoms in the chain. More preferred alkyl groups contain about 1 to about 6 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkyl chain. "Lower alkyl" means a group having about 1 to about 6 carbon atoms in the chain which may be straight or branched. The term "substituted alkyl" means that the alkyl group may be substituted by one or more substituents which may be the same or different, each substituent being independently selected from the group consisting of halo, alkyl, aryl, cycloalkyl, cyano, hydroxy, alkoxy, alkylthio, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)₂, carboxy and -C(O)O-alkyl. Non-limiting examples of suitable alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, n-pentyl, heptyl, nonyl, decyl, fluoromethyl, trifluoromethyl and cyclopropylmethyl.

"Alkenyl" means an aliphatic hydrocarbon group containing at least one carbon-carbon double bond and which may be straight or branched and comprising about 2 to about 15 carbon atoms in the chain. Preferred alkenyl groups have about 2 to about 12 carbon atoms in the chain; and more preferably about 2 to about 6 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkenyl chain. "Lower alkenyl" means about 2 to about 6 carbon atoms in the chain which may be straight or branched. The term "substituted alkenyl" means that the alkenyl group may be substituted by one or more substituents which may be the same or different, each substituent being independently selected from the group consisting of halo, alkyl, aryl, cycloalkyl, cyano, and alkoxy. Non-limiting examples of suitable alkenyl groups include ethenyl, propenyl, n-butenyl, 3-methylbut-2-enyl, n-pentenyl, octenyl and decenyl.

"Alkynyl" means an aliphatic hydrocarbon group containing at least one carbon-carbon triple bond and which may be straight or branched and comprising about 2 to about 15 carbon atoms in the chain. Preferred alkynyl groups have about 2 to about 12 carbon atoms in the chain; and more preferably about 2 to about 4 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkynyl chain. "Lower alkynyl" means about 2 to about 6 carbon atoms in the chain which may be straight or branched.

Non-limiting examples of suitable alkynyl groups include ethynyl, propynyl, 2-butylnyl, 3-methylbutynyl, n-pentylnyl, and decynyl. The term "substituted alkynyl" means that the alkynyl group may be substituted by one or more substituents which may be the same or different, each substituent being independently selected from the group consisting of alkyl, aryl and cycloalkyl.

"Alkylene" means a difunctional group obtained by removal of a hydrogen atom from an alkyl group that is defined above. Non-limiting examples of alkylene include methylene, ethylene and propylene.

"Aryl" means an aromatic monocyclic or polycyclic ring system comprising about 6 to about 14 carbon atoms, preferably about 6 to about 10 carbon atoms. The aryl group can be optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined herein. Non-limiting examples of suitable aryl groups include phenyl and naphthyl. "Aryl" is sometimes referred to herein by the abbreviation "ar".

"Heteroaryl" means an aromatic monocyclic or polycyclic ring system comprising about 5 to about 14 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the ring atoms is an element other than carbon, for example nitrogen, oxygen or sulfur, alone or in combination. Preferred heteroaryls contain about 5 to about 6 ring atoms. The "heteroaryl" can be optionally substituted by one or more "ring system substituents" which may be the same or different, and are as defined herein. The prefix aza, oxa or thia before the heteroaryl root name means that at least a nitrogen, oxygen or sulfur atom respectively, is present as a ring atom. A nitrogen atom of a heteroaryl can be optionally oxidized to the corresponding N-oxide. Non-limiting examples of suitable heteroaryls include pyridyl, pyrazinyl, furanyl, thienyl, pyrimidinyl, isoxazolyl, isothiazolyl, oxazolyl, thiazolyl, pyrazolyl, furazanyl, pyrrolyl, pyrazolyl, triazolyl, 1,2,4-thiadiazolyl, pyrazinyl, pyridazinyl, quinoxalinyl, phthalazinyl, imidazo[1,2-a]pyridinyl, imidazo[2,1-b]thiazolyl, benzofurazanyl, indolyl, azaindolyl, benzimidazolyl, benzothienyl, quinolinyl, imidazolyl, thienopyridyl, quinazolinyl, thienopyrimidyl, pyrrolopyridyl, imidazopyridyl, isoquinolinyl, benzoazaindolyl, 1,2,4-triazinyl, benzothiazolyl and the like.

"Aralkyl" means an aryl-alkyl- group in which the aryl and alkyl are as previously described. Preferred aralkyls comprise a lower alkyl group. Non-limiting examples of suitable aralkyl groups include benzyl, 2-phenethyl and naphthalenylmethyl. The bond to the parent moiety is through the alkyl.

"Alkylaryl" means an alkyl-aryl- group in which the alkyl and aryl are as previously described. Preferred alkylaryls comprise a lower alkyl group. Non-limiting examples of suitable alkylaryl groups include o-tolyl, p-tolyl and xylyl. The bond to the parent moiety is through the aryl.


"Cycloalkyl" means a non-aromatic mono- or polycyclic ring system comprising about 3 to about 10 carbon atoms, preferably about 5 to about 10 carbon atoms. Preferred cycloalkyl rings contain about 5 to about 7 ring atoms. The cycloalkyl can be optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined above. Non-limiting examples of suitable monocyclic cycloalkyls include cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl and the like. Non-limiting examples of suitable polycyclic cycloalkyls include 1-decalin, norbornyl, adamantyl and the like. The abbreviation "cy" represents cyclohexyl.

"Halo" means fluoro, chloro, bromo, or iodo groups. Preferred are fluoro, chloro or bromo, and more preferred are fluoro and chloro.

"Halogen" means fluorine, chlorine, bromine, or iodine. Preferred are fluorine, chlorine or bromine, and more preferred are fluorine and chlorine.

"Haloalkyl" means an alkyl as defined above wherein one or more hydrogen atoms on the alkyl is replaced by a halo group defined above.

"Ring system substituent" means a substituent attached to an aromatic or non-aromatic ring system which, for example, replaces an available hydrogen on the ring system. Ring system substituents may be the same or different, each being independently selected from the group consisting of aryl, heteroaryl, aralkyl, alkylaryl, aralkenyl, heteroaralkyl, alkylheteroaryl, heteroaralkenyl, hydroxy, hydroxyalkyl, alkoxy, aryloxy, aralkoxy, acyl, aroyl, halo, nitro, cyano, carboxy, alkoxy carbonyl, aryloxy carbonyl, aralkoxy carbonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, alkylsulfinyl, arylsulfinyl, heteroarylsulfinyl, alkylthio, arylthio, heteroarylthio, aralkylthio, heteroaralkylthio, cycloalkyl, cycloalkenyl, heterocyclyl, heterocyclenyl, Y_1Y_2N- , $Y_1Y_2N-alkyl-$, $Y_1Y_2NC(O)-$ and $Y_1Y_2NSO_2-$, wherein Y_1 and Y_2 may be the same or different and are independently selected from the group consisting of hydrogen, alkyl, aryl, and aralkyl. "Ring system substituent" also means a cyclic ring of 3 to 7 ring atoms of which 1-2 may be a heteroatom, attached to an aryl, heteroaryl, heterocyclyl or heterocyclenyl ring by simultaneously substituting two ring hydrogen atoms on said aryl, heteroaryl, heterocyclyl or heterocyclenyl ring. Non-limiting examples include:



and the like.

"Cycloalkenyl" means a non-aromatic mono or multicyclic ring system comprising about 3 to about 10 carbon atoms, preferably about 5 to about 10 carbon atoms which contains at least one carbon-carbon double bond. Preferred cycloalkenyl rings contain about 5 to about 7 ring atoms. The cycloalkenyl can be optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined above. Non-limiting examples of suitable monocyclic cycloalkenyls include cyclopentenyl, cyclohexenyl, cycloheptenyl, and the like. Non-limiting example of a suitable multicyclic cycloalkenyl is norbornenyl.

"Heterocyclenyl" means a non-aromatic monocyclic or multicyclic ring system comprising about 3 to about 10 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the atoms in the ring system is an element other than carbon, for example nitrogen, oxygen or sulfur atom, alone or in combination, and which contains at least one carbon-carbon double bond or carbon-nitrogen double bond. There are no adjacent oxygen and/or sulfur atoms present in the ring system. Preferred heterocyclenyl rings contain about 5 to about 6 ring atoms. The prefix aza, oxa or thia before the heterocyclenyl root name means that at least a nitrogen, oxygen or sulfur atom respectively is present as a ring atom. The heterocyclenyl can be optionally substituted by one or more ring system substituents, wherein "ring system substituent" is as defined above. The nitrogen or sulfur atom of the heterocyclenyl can be optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Non-limiting examples of suitable monocyclic azaheterocyclenyl groups include 1,2,3,4-tetrahydropyridine, 1,2-dihydropyridyl, 1,4-dihydropyridyl, 1,2,3,6-tetrahydropyridine, 1,4,5,6-tetrahydropyrimidine, 2-pyrrolinyl, 3-pyrrolinyl, 2-imidazolyl, 2-pyrazolyl, and the like. Non-limiting examples of suitable oxaheterocyclenyl groups include 3,4-dihydro-2H-pyran, dihydrofuranyl, fluorodihydrofuranyl, and the like. Non-limiting example of a suitable multicyclic oxaheterocyclenyl group is 7-oxabicyclo[2.2.1]heptenyl. Non-limiting examples of suitable monocyclic thiaheterocyclenyl rings include dihydrothiophenyl, dihydrothiopyranyl, and the like.

"Heterocyclyl" (or heterocycloalkyl) means a non-aromatic saturated monocyclic or multicyclic ring system comprising about 3 to about 10 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the atoms in the ring system is an element other than carbon, for example nitrogen, oxygen or sulfur, alone or in combination. There are no adjacent oxygen and/or sulfur atoms present in the ring system. Preferred heterocyclyls contain about 5 to about 6 ring atoms. The prefix aza, oxa or thia before the heterocyclyl root name means that at least a nitrogen, oxygen or sulfur atom respectively is present as a ring atom. The heterocyclyl can be optionally substituted by one or more "ring system substituents" which may be the same or different, and are as defined herein. The nitrogen or sulfur atom of the heterocyclyl can be optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Non-limiting examples of suitable monocyclic heterocyclyl rings include piperidyl, pyrrolidinyl, piperazinyl, morpholinyl, thiomorpholinyl, thiazolidinyl, 1,3-dioxolanyl, 1,4-dioxanyl, tetrahydrofuranyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, and the like.

"Arylcycloalkenyl" means a group derived from a fused aryl and cycloalkenyl in which the aryl and the cycloalkenyl ring share two ring carbon atoms. Preferred arylcycloalkenyls are those wherein aryl is phenyl and the cycloalkenyl consists of about 5 to about 6 ring atoms. The arylcycloalkenyl can be optionally substituted by one or more ring system substituents, wherein "ring system substituent" is as defined above. Non-limiting examples of suitable arylcycloalkenyls include 1,2-dihydronaphthalene, indene, and the like. The bond to the parent moiety is through a non-aromatic carbon atom.

"Cycloalkenylaryl" means a group derived from a fused arylcycloalkenyl as defined above. Non-limiting examples of suitable cycloalkenylaryls are as described herein for a arylcycloalkenyl, except that the bond to the parent moiety is through an aromatic carbon atom.

"Arylcycloalkyl" means a group derived from a fused aryl and cycloalkyl in which the aryl and the cycloalkyl ring share two ring carbon atoms. Preferred arylcycloalkyls are those wherein aryl is phenyl and the cycloalkyl consists of about 5 to about 6 ring atoms. The arylcycloalkyl can be optionally substituted by one or more ring system substituents, wherein "ring system substituent" is as defined above. Non-limiting examples of suitable arylcycloalkyls include 1,2,3,4-tetrahydronaphthyl, and the like. The bond to the parent moiety is through a non-aromatic carbon

atom.

"Cycloalkylaryl" means a group derived from a fused arylcycloalkyl as defined above. Non-limiting examples of suitable cycloalkylaryls are as described herein for an arylcycloalkyl group, except that the bond to the parent moiety is through an aromatic carbon atom.

"Heteroarylcyaloalkyl" means a group derived from a fused heteroaryl and cycloalkyl in which the heteroaryl and cycloalkyl share two ring atoms. The shared ring atoms may both be carbon, or one of the shared ring atoms may be a heteroatom capable of bonding to at least 3 other atoms, such as nitrogen. Preferred heteroarylcyaloalkyls are those wherein the heteroaryl thereof consists of about 5 to about 6 ring atoms and the cycloalkyl consists of about 5 to about 6 ring atoms. The prefix aza, oxa or thia before heteroaryl means that at least a nitrogen, oxygen or sulfur atom is present respectively as a ring atom. The heteroarylcyaloalkyl can be optionally substituted by one or more ring system substituents, wherein "ring system substituent" is as defined above. The nitrogen atom of the heteroaryl portion of the heteroarylcyaloalkyl can be optionally oxidized to the corresponding N-oxide. Non-limiting examples of suitable heteroarylcyaloalkyls include 5,6,7,8-tetrahydroquinolinyl, 5,6,7,8-tetrahydroisoquinolinyl, 5,6,7,8-tetrahydroquinoxalinyl, 5,6,7,8-tetrahydroquinazolyl, 4,5,6,7-tetrahydro-1 H- benzimidazolyl, 4,5,6,7-tetrahydrobenzoxazolyl, 1 H-4-oxa-1,5-diazanaphthalen-2-onyl, 1,3-dihydroimidazole-[4,5]-pyridin-2-onyl, and the like. The bond to the parent moiety is through a non-aromatic carbon atom.

"Cycloalkylheteroaryl" means a group derived from a fused heteroarylcyaloalkyl as defined above. Non-limiting examples of suitable cycloalkylheteroaryls are as described herein for heteroarylcyaloalkyl, except that the bond to the parent moiety is through an aromatic carbon atom.

"Aralkenyl" means an aryl-alkenyl- group in which the aryl and alkenyl are as previously described. Preferred aralkenyls contain a lower alkenyl group. Non-limiting examples of suitable aralkenyl groups include 2-phenethenyl and 2-naphthylethenyl. The bond to the parent moiety is through the alkenyl.

"Aralkynyl" means an aryl-alkynyl- group in which the aryl and alkynyl are as previously described. Preferred aralkynyls contain a lower alkynyl group. The bond to the parent moiety is through the alkynyl. Non-limiting examples of suitable aralkynyl groups include phenacetylenyl and naphthylacetylenyl.

"Heteroaralkyl" means a heteroaryl-alkyl- group in which the heteroaryl and alkyl are as previously described. Preferred heteroaralkyls contain a lower alkyl group. Non-limiting examples of suitable aralkyl groups include pyridylmethyl, 2-(furan-3-yl)ethyl and quinolin-3-ylmethyl. The bond to the parent moiety is through the alkyl.

"Heteroaralkenyl" means an heteroaryl-alkenyl- group in which the heteroaryl and alkenyl are as previously described. Preferred heteroaralkenyls contain a lower alkenyl group. Non-limiting examples of suitable heteroaralkenyl groups include 2-(pyrid-3-yl)ethenyl and 2-(quinolin-3-yl)ethenyl. The bond to the parent moiety is through the alkenyl.

"Heteroaralkynyl" means an heteroaryl-alkynyl- group in which the heteroaryl and alkynyl are as previously described. Preferred heteroaralkynyls contain a lower alkynyl group. Non-limiting examples of suitable heteroaralkynyl groups include pyrid-3-ylacetylenyl and quinolin-3-ylacetylenyl. The bond to the parent moiety is through the alkynyl.

"Hydroxyalkyl" means a HO-alkyl- group in which alkyl is as previously defined. Preferred hydroxyalkyls contain lower alkyl. Non-limiting examples of suitable hydroxyalkyl groups include hydroxymethyl and 2-hydroxyethyl.

"Acyl" means an H-C(O)-, alkyl-C(O)-, alkenyl-C(O)-, Alkynyl-C(O)-, cycloalkyl-C(O)-, cycloalkenyl-C(O)-, or cycloalkynyl-C(O)- group in which the various groups are as previously described. The bond to the parent moiety is through the carbonyl. Preferred acyls contain a lower alkyl. Non-limiting examples of suitable acyl groups include formyl, acetyl, propanoyl, 2-methylpropanoyl, butanoyl and cyclohexanoyl.

"Aroyl" means an aryl-C(O)- group in which the aryl group is as previously described. The bond to the parent moiety is through the carbonyl. Non-limiting examples of suitable groups include benzoyl and 1- and 2-naphthoyl.

"Heteroaroyl" means a heteroaryl-C(O)- group in which the heteroaryl group is as previously described. Non-limiting examples of suitable groups include nicotinoyl and pyrrol-2-ylcarbonyl. The bond to the parent moiety is through the carbonyl.

"Alkoxy" means an alkyl-O- group in which the alkyl group is as previously described. Non-limiting examples of suitable alkoxy groups include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy and heptoxy. The bond to the parent moiety is through the ether oxygen.

"Aryloxy" means an aryl-O- group in which the aryl group is as previously described. Non-limiting examples of suitable aryloxy groups include phenoxy and naphthoxy. The bond to the parent moiety is through the ether oxygen.

"Aralkyloxy" means an aralkyl-O- group in which the aralkyl group is as previously described. Non-limiting examples of suitable aralkyloxy groups include benzyloxy and 1- or 2-naphthalenemethoxy. The bond to the parent moiety is through the ether oxygen.

"Alkylamino" means an -NH₂ or -NH₃⁺ group in which one or more of the hydrogen atoms on the nitrogen is replaced by an alkyl group as defined above.

"Arylamino" means an -NH₂ or -NH₃⁺ group in which one or more of the hydrogen atoms on the nitrogen is replaced by an aryl group as defined above.

"Alkylthio" means an alkyl-S- group in which the alkyl group is as previously described. Non-limiting examples of

suitable alkylthio groups include methylthio, ethylthio, i-propylthio and heptylthio. The bond to the parent moiety is through the sulfur.

"Arylthio" means an aryl-S- group in which the aryl group is as previously described. Non-limiting examples of suitable arylthio groups include phenylthio and naphthylthio. The bond to the parent moiety is through the sulfur.

"Aralkylthio" means an aralkyl-S- group in which the aralkyl group is as previously described. Non-limiting example of a suitable aralkylthio group is benzylthio. The bond to the parent moiety is through the sulfur.

"Alkoxycarbonyl" means an alkyl-O-CO- group. Non-limiting examples of suitable alkoxycarbonyl groups include methoxycarbonyl and ethoxycarbonyl. The bond to the parent moiety is through the carbonyl.

"Aryloxycarbonyl" means an aryl-O-C(O)- group. Non-limiting examples of suitable aryloxycarbonyl groups include phenoxycarbonyl and naphthoxycarbonyl. The bond to the parent moiety is through the carbonyl.

"Aralkoxycarbonyl" means an aralkyl-O-C(O)- group. Non-limiting example of a suitable aralkoxycarbonyl group is benzyloxycarbonyl. The bond to the parent moiety is through the carbonyl.

"Alkylsulfonyl" means an alkyl-S(O)₂- group. Preferred groups are those in which the alkyl group is lower alkyl. The bond to the parent moiety is through the sulfonyl.

"Alkylsulfinyl" means an alkyl-S(O)- group. Preferred groups are those in which the alkyl group is lower alkyl. The bond to the parent moiety is through the sulfinyl.

"Arylsulfonyl" means an aryl-S(O)₂- group. The bond to the parent moiety is through the sulfonyl.

"Arylsulfinyl" means an aryl-S(O)- group. The bond to the parent moiety is through the sulfinyl.


[0031] The term "optionally substituted" means optional substitution with the specified groups, radicals or moieties, in available position or positions.

[0032] With reference to the number of moieties (e.g., substituents, groups or rings) in a compound, unless otherwise defined, the phrases "one or more" and "at least one" mean that there can be as many moieties as chemically permitted, and the determination of the maximum number of such moieties is well within the knowledge of those skilled in the art.


[0033] Those skilled in the art will appreciate that the term "neurodegenerative disease" has its commonly accepted medical meaning and describes diseases and conditions resulting from abnormal function of neurons, including neuronal death and abnormal release of neurotransmitters or neurotoxic substances. In this instance it also includes all diseases resulting from abnormal levels of beta amyloid protein. Examples of such diseases include, but are not limited to, Alzheimer's disease, age-related dementia, cerebral or systemic amyloidosis, hereditary cerebral hemorrhage with amyloidosis, and Down's syndrome.

[0034] Lines drawn into the ring systems indicate that the indicated bond may be attached to any of the substitutable ring carbon atoms.


[0035] As is well known in the art, a bond drawn from a particular atom wherein no moiety is depicted at the terminal end of the bond indicates a methyl group bound through that bond to the atom. For example:



represents




5



10

represents


15



20

and

25




30

35

40

represents



45




50

55


[0036] Those skilled in the art will appreciate a bicyclic ring structure, two examples of which are represented by the diagrams shown below. For convenience and description, carbon atoms can be numbered accordingly.




[0037] A dotted line within a ring structure, an example of which is shown in the diagram below, represents an optional bond or optional double bond. For example, with an optional bond,




the structure above could represent the compound below:



[0038] Further, the double or triple bond may form the bonds between any two available elements on any ring in the structure. For example, with a double bond forming,



the structure above could represent the structure below:



[0039] As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

[0040] Prodrugs and solvates of the compounds of formula I, including the compounds of the invention, are also contemplated herein. The term "prodrug", as employed herein, denotes a compound that is a drug precursor which, upon administration to a subject, undergoes chemical conversion by metabolic or chemical processes to yield a compound of formula I or a salt and/or solvate thereof. A discussion of prodrugs is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems (1987) Volume 14 of the A.C.S. Symposium Series, and in Bioreversible Carriers in Drug Design, (1987) Edward B. Roche, ed., American Pharmaceutical Association and Pergamon Press, both of which are incorporated herein by reference thereto.

[0041] "Solvate" means a physical association of a compound of formula I, including a compound of this invention, with one or more solvent molecules. This physical association involves varying degrees of ionic and covalent bonding, including hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of the crystalline solid. "Solvate" encompasses both solution-phase and isolatable solvates. Non-limiting examples of suitable solvates include ethanulates, methanulates, and the like. "Hydrate" is a solvate wherein the solvent molecule is H₂O.

[0042] "Effective amount" or "therapeutically effective amount" is meant to describe of a compound of formula I, e.g. a compound of the invention, or a composition containing such a compound, effective in inhibiting gamma secretase and thus producing the desired therapeutic effect.

[0043] The compounds of formula I, including compounds of the invention, form salts. Reference to a compound of formula I herein is understood to include reference to salts thereof, unless otherwise indicated. The term "salt(s)", as employed herein, denotes acidic salts formed with inorganic and/or organic acids, as well as basic salts formed with inorganic and/or organic bases. In addition, when a compound of formula I contains both a basic moiety, such as, but not limited to a pyridine or imidazole, and an acidic moiety, such as, but not limited to a carboxylic acid, zwitterions ("inner salts") may be formed and are included within the term "salt(s)" as used herein. Pharmaceutically acceptable (i.e., non-toxic, physiologically acceptable) salts are preferred, although other salts are also useful. Salts of the compounds of the formula I may be formed, for example, by reacting a compound of formula I with an amount of acid or base, such as an equivalent amount, in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

[0044] Exemplary acid addition salts include acetates, adipates, alginates, ascorbates, aspartates, benzoates, benzenesulfonates, bisulfates, borates, butyrates, citrates, camphorates, camphorsulfonates, cyclopentanepropionates, digluconates, dodecylsulfates, ethanesulfonates, fumarates, glucoheptanoates, glycerophosphates, hemisulfates, heptanoates, hexanoates, hydrochlorides, hydrobromides, hydroiodides, 2-hydroxyethanesulfonates, lactates, maleates, methanesulfonates, 2-naphthalenesulfonates, nicotinate, nitrates, oxalates, pectinates, persulfates, 3-phenylpropionates, phosphates, picrates, pivalates, propionates, salicylates, succinates, sulfates, sulfonates (such as those mentioned herein), tartarates, thiocyanates, toluenesulfonates (also known as tosylates,) undecanoates, and the like. Additionally, acids which are generally considered suitable for the formation of pharmaceutically useful salts from basic pharmaceutical compounds are discussed, for example, by S. Berge et al, Journal of Pharmaceutical Sciences (1977) 66 (1) 1-19; P. Gould, International J. of Pharmaceutics (1986) 33 201-217; Anderson et al, The Practice of Medicinal Chemistry (1996), Academic Press, New York; and in The Orange Book (Food & Drug Administration, Washington, D.C. on their website). These disclosures are incorporated herein by reference thereto.

[0045] Exemplary basic salts include ammonium salts, alkali metal salts such as sodium, lithium, and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases (for example, organic amines) such as benzathines, dicyclohexylamines, hydrabamines (formed with N,N-bis(dehydroabietyl)ethylenediamine), N-methyl-D-glucamines, N-methyl-D-glucamides, t-butyl amines, and salts with amino acids such as arginine, lysine and the like. Basic nitrogen-containing groups may be quarternized with agents such as lower alkyl halides (e.g. methyl, ethyl,

propyl, and butyl chlorides, bromides and iodides), dialkyl sulfates (e.g. dimethyl, diethyl, dibutyl, and diamyl sulfates), long chain halides (e.g. decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides), aralkyl halides (e.g. benzyl and phenethyl bromides), and others.

[0046] All such acid salts and base salts are intended to be pharmaceutically acceptable salts within the scope of this disclosure and all acid and base salts are considered equivalent to the free forms of the corresponding compounds for purposes of this disclosure.

[0047] Compounds of formula I, and salts, solvates and prodrugs thereof, may exist in their tautomeric form (for example, as an amide or or imino ether). All such tautomeric forms are contemplated herein as part of this disclosure.

[0048] All stereoisomers (for example, geometric isomers, optical isomers and the like) of the compounds of this disclosure (including those of the salts, solvates and prodrugs of the compounds as well as the salts and solvates of the prodrugs), such as those which may exist due to asymmetric carbons on various substituents, including enantiomeric forms (which may exist even in the absence of asymmetric carbons), rotameric forms, atropisomers, and diastereomeric forms, are contemplated within the scope of this disclosure. Individual stereoisomers of the compounds of this disclosure including compounds of the invention may, for example, be substantially free of other isomers, or may be admixed, for example, as racemates or with all other, or other selected, stereoisomers. The chiral centers in compounds of this disclosure, including compounds of the present invention can have the S or R configuration as defined by the *IUPAC* 1974 Recommendations. The use of the terms "salt", "solvate" "prodrug" and the like, is intended to equally apply to the salt, solvate and prodrug of enantiomers, stereoisomers, rotamers, tautomers, racemates or prodrugs of the compounds disclosed herein.

[0049] Polymorphic forms of the compounds of Formula I, e.g. compounds of the invention, and of the salts, solvates and/or prodrugs of the compounds of Formula I, are intended to be included in this disclosure.

[0050] This disclosure is also directed to pharmaceutical compositions which comprise at least one compound of Formula I, e.g. at least one compound of the present invention, or a pharmaceutically acceptable salt or solvate of said compound and at least one pharmaceutically acceptable carrier. Pharmaceutical compositions can comprise one or more of the compounds of Formula I.

[0051] For preparing pharmaceutical compositions from the compounds described by this disclosure, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets and suppositories. The powders and tablets may be comprised of from about 5 to about 95 percent active compound. Suitable solid carriers are known in the art, e.g. magnesium carbonate, magnesium stearate, talc, sugar or lactose. Tablets, powders, cachets and capsules can be used as solid dosage forms suitable for oral administration. Examples of pharmaceutically acceptable carriers and methods of manufacture for various compositions may be found in A. Gennaro (ed.), *Remington's Pharmaceutical Sciences*, 18th Edition, (1990), Mack Publishing Co., Easton, Pennsylvania.

[0052] Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injection or addition of sweeteners and opacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal administration.

[0053] Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier, such as an inert compressed gas, e.g. nitrogen.

[0054] Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

[0055] The compounds of the disclosure, including compounds of the invention may also be deliverable transdermally. The transdermal compositions can take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type as are conventional in the art for this purpose. ,

[0056] Preferably, the pharmaceutical preparation is in a unit dosage form. In such form, the preparation is subdivided into suitably sized unit doses containing appropriate quantities of the active compound, e.g., an effective amount to achieve the desired purpose.

[0057] The quantity of active compound in a unit dose of preparation may be varied or adjusted from about 0.01 mg to about 1000 mg, preferably from about 0.01 mg to about 750 mg, more preferably from about 0.01 mg to about 500mg, and most preferably from about 0.01 mg to about 250 mg, according to the particular application.

[0058] The actual dosage employed may be varied depending upon the requirements of the patient and the severity of the condition being treated. Determination of the proper dosage regimen for a particular situation is within the skill of the art. For convenience, the total daily dosage may be divided and administered in portions during the day as required.

[0059] The amount and frequency of administration of the compounds of this disclosure, e.g. the compounds of this invention, and/or the pharmaceutically acceptable salts thereof will be regulated according to the judgment of the attending clinician considering such factors as age, condition and size of the patient as well as severity of the symptoms being treated. A typical recommended daily dosage regimen for oral administration can range from about 0.04 mg/day to about 4000 mg/day, in one to four divided doses.

[0060] Another aspect of this disclosure is a kit comprising a therapeutically effective amount of at least one compound

of Formula I, or a pharmaceutically acceptable salt or solvate of said compound and a pharmaceutically acceptable carrier, vehicle or diluent.

EXAMPLES

[0061] The invention disclosed herein is exemplified by the following preparations and examples which should not be construed to limit the scope of the disclosure. Alternative mechanistic pathways and analogous structures will be apparent to those skilled in the art.

[0062] Where NMR data are presented, ¹H spectra were obtained on either a Varian VXR-200 (200 MHz, ¹H), Varian Gemini-300 (300 MHz) or XL-400 (400 MHz) and are reported as ppm down field from Me₄Si with number of protons, multiplicities, and coupling constants in Hertz indicated parenthetically. Where LC/MS data are presented, analyses was performed using an Applied Biosystems API-100 mass spectrometer and Shimadzu SCL-10A LC column: Alltech platinum C18, 3 micron, 33mm x 7mm ID; gradient flow: 0 min - 10% CH₃CN in water, 5 min - 95% CH₃CN in water, 7 min - 95% CH₃CN in water, 7.5 min - 10% CH₃CN in water, 9 min. The retention time and observed parent ion are given.

[0063] The following solvents and reagents may be referred to by their abbreviations in parenthesis:

AcOEt: represents ethyl acetate;

AcOH: represents acetic acid;

AIBN represents: 2,2'-azobisisobutyronitrile;

DCM: represents dichloromethane;

DCE: represents dichloroethylene;

DIBALH: represents diisobutylaluminum hydride;

DEAD: represents diethylazodicarboxylate;

DMF: represents dimethylformamide;

EDCI: represents 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide;

Et₂O: represents diethyl ether;

EtOAc: represents ethyl acetate;

HOBT represents: 1-hydroxybenzotriazole;

LDA represents: lithium diisopropylamide;

MS: represents mass spectrometry;

Me: represents methyl;

MeOH: represents methanol;

NMO represents: N-methylmorpholine N-oxide;

NaOH: represents sodium hydroxide;

NMR: represents nuclear magnetic resonance;

HRMS: represents high resolution mass spectrometry;

OTBDMS: represents t-butyldimethylsilyloxy (or t-butyldimethylsilyl ether);

OTBDPS: represents t-butyldiphenylsilyloxy (or t-butyldiphenylsilyl ether);

P: represents a protecting group;

Ph: represents phenyl;

RT: represents room temperature.

TBAF: represents tetrabutylammonium fluoride;

TBDMS: represents t-butyldimethylsilyl;

TBDMSCl: represents t-butyldimethylsilyl chloride;

TBDPSCl: represents t-butyldiphenylsilylchloride;

TBS: represents t-butylsilyl;

TFA: represents trifluoroacetic acid;

THF: represents tetrahydrofuran;


TMS: represents trimethylsilane;

TMSCl represents: trimethylsilyl chloride.

[0064] Compounds of Formula I can be prepared by various methods well known to those skilled in the art, and by the methods described below. The following methods are typical:

Method 1


[0065] In Method 1, compounds of Formula I having the structure A are prepared.



[0066] The protecting group in alcohol **1** is removed following standard conditions to lead to an amine that is further converted to alcohol **2** by reaction with a sulfonyl halide. Alcohol **2** can be converted to a variety of compounds of type **A** using methods well-known to those skilled in the art. For example, carbamates can be prepared by reaction of **2** with 4-nitrophenylchloroformate followed by reaction of the resulting carbonate with a primary or secondary amine. Alternatively, esters can be prepared by reaction of **2** with either an acid halide or a carboxylic acid in the presence of a suitable coupling reagent such as EDCI and HOBT.

Method 2


[0067] In Method 2, compounds of Formula I having the structure **B** are prepared.



[0068] Cyclic amine **3** is converted to a sulfonamide by reaction with a sulfonyl halide, then converted to enol ether **4** using standard conditions such as deprotonation with LDA followed by reaction with TMSCl. Oxidation of enol ether **4** with an appropriate mixture of reagents such as osmium tetroxide and NMO followed by protection of the alcohol using a typical protecting group such as tertbutyldimethylsilyl chloride then gives ketone **5**. Ketone **5** can be des-oxygenated to the alkylene **7** using a variety of methods known to those skilled in the art including but not limited to Wolf-Kishner type reactions and radical des-oxygenation reactions. For example, reaction of ketone **5** with an appropriate reducing agent such as sodium borohydride gives an alcohol which is then converted to thiocarbonate **6** using phenylchlorothioformate. Thiocarbonate **6** is then reduced to alkylene **7** with radical-induced reducing conditions such as AIBN with tributyltin hydride. The alcohol protecting group in alkylene **7** is then removed and the alcohol is converted to compounds of type **B** following methods described in Method 1.

Method 3


[0069] In Method 3, compounds of Formula I having the structure **C** are prepared.



[0070] In Method 3, the alcohol functionality of methyl 2-(hydroxymethyl)-6-pyridinecarboxylate **8** is protected using a typical protecting group such as tertbutyldimethylsilyl chloride and the pyridine is then hydrogenated using a suitable catalyst such as platinum oxide, preferably in the presence of acetic acid, to give amine **9**. Cyclic amine **9** is further converted to a sulfonamide by reaction with a sulfonyl halide, and then is reduced with a hydride reagent such as diisobutylaluminum hydride to give aldehyde **10**. A Wittig reaction is then performed on aldehyde **10** and the alcohol protecting group is removed under standard conditions such as TBAF to give alcohol **11**. The alcohol **11** is then oxidized to an aldehyde using an oxidative reagent such as Dess-Martin periodinate and the intermediate aldehyde is reacted with a metal-alkene such as a Grignard alkene to provide bis-alkene alcohol **12**. A ring-closure-metathesis with a reagent such as $RuCl_2(CHPh)[PCy_3]_2$ (Grubbs's second generation catalyst) is then carried out on bis-alkene alcohol **12** to obtain bicyclic alcohol **13**. Alcohol **13** is finally converted to compounds of type **C** following methods described in Method 1.

Method 4


[0071] In Method 4, compounds of Formula I having the structure **D** are prepared.



[0072] Alkene alcohol **13** is hydrogenated using a suitable catalyst such as platinum oxide, and the resulting alkyl alcohol is converted to compounds of type **D** using methods described in Method 1.

Preparative Example 1:

[0073]

**Step 1:**

[0074] To a solution of methyl 2-(hydroxymethyl)-6-pyridinecarboxylate (44.5 g, 0.26 mol) in DCE (500 mL) was added triethylamine (44 mL, 0.31 mol) followed by TBDMSCl (44 g, 0.29 mol), and then the reaction was heated at 70°C for 2 days. After concentration, the residue was purified over silica gel (eluted with hexane/AcOEt 90:10 to 50:50) to afford 68.5 g of TBS-protected pyridine ester.

Step 2:

[0075] A solution of the TBS-protected pyridine ester of Step 1 (68 g, 0.24 mmol) and platinum (IV) oxide (6 g, 0.026 mmol) in MeOH (500 mL) and AcOH (50 mL) was hydrogenated at 40 psi overnight. The mixture was filtered over Celite

and concentrated. The residue was diluted with aqueous 1 N NaOH, extracted with DCM and AcOEt, dried over Na₂SO₄ and concentrated to provide 66 g of TBS-protected piperidine ester.

Step 3:

[0076] A solution of the TBS-protected piperidine ester of Step 2 (63 g, 0.22 mol), triethylamine (100 mL, 0.66 mol) and 4-chlorobenzenesulfonyl chloride (93 g, 0.44 mol) in DCE (500 mL) was heated at 40 °C overnight. After concentration, the residue was purified over silica gel (eluted with hexane/AcOEt 97:3 to 90:10 then DCM/AcOEt 90:10) to give 89 g of TBS-protected piperidinesulfonamide ester.

Step 4:

[0077] To a solution of the TBS-protected piperidinesulfonamide ester of Step 3 (23.5 g, 50.9 mmol) in anhydrous DCM (250 mL) at -78 °C was slowly added DIBAH 1 N in toluene (55 mL, 55 mmol) and the reaction mixture was stirred for 1 h at this temperature. The final mixture was then quenched at -78 °C with 15% Rochelle salt solution (300 mL). The reaction was then diluted with DCM, Celite was added and the mixture was stirred for 1 h at RT then filtered, extracted with DCM, AcOEt, dried over Na₂SO₄ and concentrated. The residue was purified over silica gel (eluted with hexanes/AcOEt 95:5 to 80:20) to provide 16.37 g of TBS-protected piperidinesulfonamide aldehyde.

Step 5:

[0078] To a solution of methyltriphenylphosphonium bromide (14.65 g, 41 mmol) in THF (150 mL) at -78 °C was added n-BuLi 2.5 N in hexanes (16.4 mL, 41 mmol) then the reaction was warmed to 0 °C. It was then cooled to -78 °C and cannulated into a solution of the TBS-protected piperidinesulfonamide aldehyde of Step 4 (16.35 g, 37.8 mmol) in THF (100 mL) at -78 °C. The reaction was stirred for 30 min at -78 °C then for 30 min at RT. The final mixture was concentrated, diluted with aqueous 5% NaHCO₃ and DCM, extracted with DCM, dried over Na₂SO₄ and concentrated. Purification on silica gel (eluted with hexanes/AcOEt 95:5 to 80:20) afforded 9.81 g of TBS-protected alkene.

Step 6:

[0079] A solution of the TBS-protected alkene of Step 5 (9.81 g, 22.8 mmol) and TBAF 1 N in THF (45 mL, 45 mmol) in THF (50 mL) was stirred at 50 °C for 1 h then concentrated. The residue was taken up in water, extracted with DCM, AcOEt, dried over Na₂SO₄ and concentrated. The residue was purified over silica gel (eluted with hexanes/AcOEt 80:20 to 60:40) to give 7.00 g of alkene alcohol.

Step 7:

[0080] To a solution of the alkene alcohol of Step 5 (7.00 g, 22.2 mmol) in DCM (100 mL) was added Dess-Martin periodinate (18.8 g, 44.4 mmol) and NaHCO₃ (5.59 g, 66.6 mmol). The solution was stirred for 3 h at RT then diluted with 5% Na₂S₂O₃ (400 mL), extracted with DCM, dried over Na₂SO₄ and concentrated. The residue was purified over silica gel (eluted with hexanes/AcOEt 95:5 to 80:20) to give 4.78 g of alkene aldehyde.

Step 8:

[0081] To a solution of the alkene aldehyde of Step 7 (3.68 g, 11.73 mmol) in THF (60 mL) at -40 °C was added vinyl magnesium bromide 1 N in hexanes (14 mL, 14 mmol) and the reaction was allowed to warm to -20 °C. After 30 min at this temperature, the reaction was quenched with sat. aqueous NH₄Cl, extracted with DCM, AcOEt, dried over Na₂SO₄ and concentrated. The residue was purified over silica gel (eluted with hexanes/AcOEt 93:7 to 50:50) to provide, in order of elution, 540 mg of starting alkene aldehyde, 675 mg of bis-alkene alcohol isomer A (12a), and 2.63 g of bis-alkene alcohol isomer B (12b).

Step 9:

[0082] A solution of the bis-alkene alcohol isomer B of Step 8 (2.55 g, 7.46 mmol) in DCM (350 mL) was deaerated *in vacuo* then put under nitrogen. Bis(tricyclohexylphosphine)benzylidene ruthenium (IV) dichloride (930 mg, 1.11 mmol) was then added and the reaction was stirred at 50 °C overnight. The final solution was then concentrated and purified over silica gel (eluted with hexanes/AcOEt 85:15 to 40:60) to provide, in order of elution, 435 mg of starting material and 1.31 g of azabicyclo[3.3.1]nonane alkene alcohol.

Step 10:

[0083] A solution of the azabicyclo[3.3.1]nonane alkene alcohol of Step 9 (1.40 g, 4.46 mmol) and platinum (IV) oxide (215 mg) in AcOEt (30 mL) was hydrogenated at atmospheric pressure for 2 hours. The final mixture was filtered over Celite and concentrated. The residue was then purified over silica gel (eluted with hexanes/AcOEt 90:10 to 60:40) to afford 1.24 g of azabicyclo[3.3.1]nonane alcohol.

Step 11:



[0084] To a solution of the azabicyclo[3.3.1]nonane alcohol of Step 10 (170 mg, 0.54 mmol) in DCE (2.7 mL) was added p-nitrophenylchloroformate (217 mg, 1.10 mmol) followed by triethylamine (0.4 ml) and the reaction was stirred at RT overnight. After concentration, the residue was purified over silica gel (eluted with DCM) to obtain 192 mg of azabicyclo[3.3.1]nonane carbonate.

Step 12:





[0085] To a solution of azabicyclo[3.3.1]nonane carbonate (200 mg, 0.40 mmol) in DCE was added N-(2-hydroxyethyl)piperazine (0.1 ml, 0.80 mmol) and the reaction was heated at 50°C overnight. The mixture was diluted with aqueous 0.5 NaOH, extracted with DCM and AcOEt and dried over Na₂SO₄. The residue obtained after concentration was purified over silica gel (eluted with DCM/MeOH 9:1) to give 40 mg of product of formula 14. ¹H NMR (CDCl₃ 300 MHz) δ 7.76 (d, J = 8.4 Hz, 2H), 7.43 (d, J = 8.4 Hz, 2H), 4.77 (m, 1 H), 4.28 (m, 1H), 3.95 (m, 1 H), 3.55-3.70 (m, 4H), 3.35-3.55 (m, 2H), 2.40-2.65 (m, 8H), 1.45-1.90 (m, 9H); HRMS (MH⁺) 472.1994.

[0086] Following similar procedures to those of Preparative Example 1 and using bis-alkene alcohol isomer A or bis-alkene alcohol isomer B (of Step 8) in the synthesis, the compounds in Table 1 were prepared:





TABLE 1

Formula No.	COMPOUND	Mass Spec (M ⁺); retention time (min)
15		510.1; 4.81
16		510.1; 4.78





(continued)

Formula No.	COMPOUND	Mass Spec (M ⁺); retention time (min)
17		482.1; 4.45
18		500.1; 4.23
19		496.3; 3.27
20		485.1; 4.70





(continued)

Formula No.	COMPOUND	Mass Spec (M ⁺); retention time (min)
21		470.3; 3.21
22		486.1; 4.23
23		504.1; 5.23
24		553.3; 3.22

(continued)


Formula No.	COMPOUND	Mass Spec (M ⁺); retention time (min)
25		526.3; 3.28
26		442.1 ; 4.43
27		485.3; 3.17
28		443.2; 3.00

(continued)

Formula No.	COMPOUND	Mass Spec (M ⁺); retention time (min)
29		478.3; 3.18
30		439.2; 2.83
31		439.2; 2.70
32		492.3; 3.09

Preparative Example 2:

[0087]

**Formula 33**

[0088] The alcohol of Preparative Example 1, Step 9 above, was subjected to conditions similar to those described in Preparative Example 1, Step 11 and Step 12, using 4-piperidinopiperidine instead of N-(2-hydroxyethyl) piperazine in the last step, to afford the desired product of formula 33. ¹H NMR (CDCl₃ 300 MHz) δ 7.78 (d, J = 8.4 Hz, 2H), 7.43 (d, J = 8.4 Hz, 2H), 5.95-6.05 (m, 2H), 4.86 (br s, 1 H), 4.10-4.45 (m, 3H), 3.75-4.05 (m, 1H), 2.40-2.85 (m, 8H), 1.30-2.00 (m, 15H); MS (ES) *m/e* 508.1 (M+H)⁺; retention time: 5.35 min.


[0089] Following similar procedures to those of Preparative Example 2 and using bis-alkene alcohol isomer A in the synthesis, the following compound in Table 2 was prepared:

TABLE 2

Formula No.	COMPOUND	Mass Spec (M ⁺); retention time (min)
34		508.1; 5.52

Preparative Example 3:

[0090]

**Step 1:**

[0091] A mixture of exo-7-benzyl-7-aza-bicyclo[2.2.1]heptan-2-ol (1.86 g, 9.1 mmol), ammonium formate (3.47 g, 55 mmol) and Pd/C 10% (200 mg) in MeOH (20 mL) was stirred for 2 h at 50°C. The solution was filtered over Celite, rinsed with MeOH and then 1 N HCl until a pH of 1 was obtained. The solution is then diluted with water, washed with Et₂O, and concentrated. The remaining solid (4.8 g) was taken up in 1 N NaOH (100 mL) and treated with 4-chlorobenzenesulfonyl chloride (10.55 g, 50 mmol) in DCM (40 mL) overnight. The mixture was diluted with water and DCM, extracted with DCM, dried over Na₂SO₄ and concentrated. Purification over silica gel (eluting with DCM/AcOEt 90:10 to 80:20) afforded 1.87 g of sulfonamide alcohol.

Step 2:





[0092] The product of Step 1 was subjected to conditions similar to the conditions described in Preparative Example 1, Step 11 and Step 12 above, using 4-piperidinopiperidine in the last step, to afford the desired product of formula 35. ¹H NMR (CDCl₃ 300 MHz) δ 7.81 (d, J = 8.4 Hz, 2H), 7.42 (d, J = 8.4 Hz, 2H), 4.64 (m, 1 H), 3.90-4.25 (m, 4H), 2.55-2.75 (m, 2H), 2.30-2.55 (m, 5H), 1.65-1.95 (m, 6H), 1.50-1.60 (m, 4H), 1.30-1.45 (m, 6H); MS (ES) *m/e* 482.3 (M+H)⁺; retention time: 3.71 min.

[0093] Following similar procedures to those of Preparative Example 3, the compounds of Table 3 were prepared:


TABLE 3

Formula No.	COMPOUND	Mass Spec (M ⁺); retention time (min)
36		482.3; 4.06


(continued)

Formula No.	COMPOUND	Mass Spec (M ⁺); retention time (min)
37		428.2; 3.64
38		444.2; 3.38
39		416.1; 3.94
40		442.1; 4.48

(continued)

Formula No.	COMPOUND	Mass Spec (M ⁺); retention time (min)
41		430.1; 4.41

Preparative Example 4:**[0094]**



Step 1:

[0095] A solution of nor-tropinone hydrobromide (6.05 g, 29.5 mmol), 4-chlorobenzenesulfonyl chloride (6.65 g, 31.5 mmol), K_2CO_3 (9.70 g, 70 mmol) and tetrabutylammonium bromide (200 mg) in DCM (70 mL) and water (70 mL) was stirred at RT overnight. The mixture was extracted with DCM, dried over Na_2SO_4 , concentrated and purified over silica gel (eluted with DCM/AcOEt 95:5) to give 9.04 g of nor-tropinone sulfonamide.

Step 2:

[0096] To a solution of LDA 2N in heptanes (16.1 mL, 32.2 mmol) in THF (120 mL) at $-78^\circ C$ was slowly added a solution of the nor-tropinone sulfonamide of Step 1 (8.06 g, 26.9 mmol) in THF (160 mL) over 10 min. After 35 min at $-78^\circ C$, chlorotrimethylsilane (6.8 mL) was added, the reaction was stirred 20 min at $-78^\circ C$, saturated aqueous $NaHCO_3$ was added and the mixture was warmed to RT. After extraction with Et_2O , drying over Na_2SO_4 , and concentration, the residue was purified over silica gel (eluted with hexanes/DCM 1:1 to DCM) to afford 3.88 g of nor-tropinone sulfonamide enol ether.

Step 3:

[0097] To a solution of the nor-tropinone sulfonamide enol ether of Step 2 (3.88 g, 10.43 mmol) and NMO (1.51 g, 12.9 mmol) in THF (40 mL) and water (15 mL) was added OsO₄ (100 mg, 0.40 mmol) at 0°C, and then the reaction was allowed to stir at RT overnight. After concentration, the solution was diluted with AcOEt and aqueous 10% Na₂SO₃, extracted with AcOEt, dried over Na₂SO₄ and concentrated. The residue was purified over silica gel (eluted with DCM/AcOEt 90:10 to 80:20) to afford 1.51 g of α-hydroxyketone sulfonamide.

Step 4:

[0098] A solution of the α-hydroxyketone sulfonamide of Step 3 (1.50 g, 4.75 mmol), TBDMSCI (860 mg, 5.70 mmol), and imidazole (390 mg, 5.70 mmol) in DMF (6 mL) was stirred overnight at 40°C. The cooled solution was diluted with water and Et₂O, extracted with Et₂O, dried over Na₂SO₄ and concentrated. The residue was purified over silica gel (eluted with hexane/DCM 1:1 to DCM) to provide 1.53 g of O-TBS ketone sulfonamide.

Step 5:

[0099] To a solution of the O-TBS ketone sulfonamide of Step 4 (1.53 g, 3.56 mmol) in MeOH (10 mL) at 0°C was slowly added NaBH₄ (135 mg, 3.56 mmol) and the solution was stirred for 2 h at this temperature. The final mixture was concentrated, diluted with water, extracted with DCM and AcOEt, and dried over Na₂SO₄. The residue obtained after concentration was purified over silica gel (eluted with DCM to DCM/AcOEt 9:1) to afford 1.51 g of O-TBS alcohol sulfonamide.

Step 6:

[0100] To a solution of the O-TBS alcohol sulfonamide of Step 5 (1.10 g, 2.55 mmol) in THF (10 mL) at -78°C was added n-BuLi 2.5 N in hexanes (1.10 mL, 2.80 mmol) followed, 1 min later, by phenylthionochloroformate (460 μL, 3.32 mmol). The reaction mixture was stirred 30 min at -78°C then allowed to warm to RT and stirred for 1 h. The final solution was treated with 5% NaHCO₃, extracted with DCM (3x), AcOEt (1x), dried over Na₂SO₄ and concentrated. The residue was purified over silica gel (eluting with hexane/DCM 60:40 to DCM) to provide 1.34 g of O-TBS thiocarbonate sulfonamide.

Step 7:

[0101] A solution of the O-TBS thiocarbonate sulfonamide of Step 6 (1.34 g, 2.36 mmol), tributyltin hydride (1.35 mL, 5.0 mmol) and AIBN (107 mg, 0.65 mmol) was stirred under reflux for 6 h then concentrated. Purification of the crude over silica gel (eluted with hexane/DCM 60:40 to DCM) afforded 514 mg of O-TBS sulfonamide with other components.

Step 8:


[0102] A solution of the O-TBS sulfonamide of Step 7 (514 mg, 1.23 mmol) and TBAF 1 N in THF (2 mL, 2.0 mmol) in THF (3 mL) was stirred for 2 h at 40°C then concentrated. Workup with aqueous saturated NH₄Cl, DCM, and AcOEt afforded a residue that was purified over silica gel (eluted with DCM/AcOEt 88:12 to 85:15) to give 105 mg of alcohol sulfonamide.


Step 9:

[0103] The product of Step 8 was subjected to conditions similar to the conditions described in Example 1, Step 11 and Step 12, using N-(2-hydroxyethyl)-piperazine in the last step, to afford the desired product of formula 42. ¹H NMR (CDCl₃ 300 MHz) δ 7.77 (d, J = 8.4 Hz, 2H), 7.43 (d, J = 8.4 Hz, 2H), 4.59 (br s, 1 H), 4.28 (br d, 2H), 3.30-3.70 (m, 6H), 2.45-2.65 (m, 7H), 2.06 (m, 1H), 1.40-1.85 (m, 7H); MS (ES) m/e 458.1 (M+H)⁺; retention time: 3.65 min.

[0104] Following similar procedures to those of Preparative Example 4, the compound in Table 4 was prepared:

TABLE 4

Formula No.	COMPOUND	Mass Spec (M^+); retention time (min)
3		496.1; 4.36

Preparative Example 5:**[0105]****Step 1:**

[0106] A solution of the product alcohol of Preparative Example 1, Step 10 (450 mg, 1.42 mmol) and Dess-Martin periodinate (15% in DCM, 4.80 g, 1.70 mmol) in DCM (5 mL) was stirred at RT for 3 h. The mixture was treated with 5% sodium thiosulfate, extracted with DCM, dried over Na_2SO_4 and concentrated. The residue was purified over silica gel (eluted with hexanes/AcOEt 90:10 to 60:40) to give 460 mg of ketone.

Step 2:

[0107] To a solution of the ketone of Step 1 (150 mg, 0.48 mmol) in THF (3 mL) at -20°C was slowly added MeMgBr


3N (190 μ L, 0.56 mmol), and then the reaction was allowed to warm to 0°C. After 1 h, the final solution was poured into saturated NH_4Cl and extracted with DCM, AcOEt, dried over Na_2SO_4 and concentrated. The residue was purified over silica gel (eluted with hexanes/AcOEt 90:10 to 60:40) to provide 125 mg of alcohol.

Step 3:

[0108] The product of Step 2 was subjected to conditions similar to the conditions described in Preparative Example 1, Step 11 and Step 12, using N-(2-hydroxyethyl)-piperazine in the last step, to afford the desired product of formula 44. ^1H NMR (CDCl_3 300 MHz) δ 7.79 (d, J = 8.4 Hz, 2H), 7.46 (d, J = 8.4 Hz, 2H), 4.43 (m, 1 H), 4.08 (m, 1H), 3.60-3.70 (m, 3H), 3.40-3.55 (m, 4H), 2.40-2.70 (m, 8H), 1.40-2.20 (m, 9H), 1.54 (s, 3H); MS (ES) m/e 486.1 ($\text{M}+\text{H}^+$); retention time: 4.06 min.


[0109] Following similar procedures to those of Preparative Example 5, the compound of Table 5 was prepared:

TABLE 5

Formula No.	COMPOUND	Mass Spec (M^+); retention time (min)
45		524.1; 4.58

Preparative Example 6:

[0110]



Step 1:



[0111] To a solution of diethylzinc (1 N in hexanes, 5 mL, 5 mmol) in DCE (20 mL) at -20°C was slowly added chloriodomethane (370 μ L, 5 mmol), and the reaction was stirred for 5 min at this temperature. A solution of the product, the alcohol, of Preparative Example 1, Step 9 (740 mg, 2.36 mmol) in DCE (3 mL) was then added and the reaction was stirred for 1 h at -10°C and for 1 h at RT. The final mixture was quenched into sat NH_4Cl , extracted with DCM, dried over Na_2SO_4 and concentrated. The crude product was purified over silica gel (eluted with hexanes/AcOEt 80:20 to 50:50) to give 512 mg of cyclopropyl alcohol.

Step 2:

[0112] The product of Step 1 was subjected to conditions similar to the conditions described in Preparative Example 1, Step 11 and Step 12, using N-(2-hydroxyethyl)-piperazine in the last step, to afford the desired product of formula 46. ¹H NMR (CDCl₃ 300 MHz) δ 7.74 (d, J = 8.4 Hz, 2H), 7.45 (d, J = 8.4 Hz, 2H), 4.91 (d, J = 7.5 Hz, 1H), 4.30 (br s, 1H), 3.90 (m, 1 H), 3.63 (m, 2H), 3.30-3.55 (m, 4H), 2.40-2.60 (m, 8H), 1.90-2.10 (m, 1H), 1.35-1.70 (m, 8H), 0.98 (m, 1H), 0.50-0.62 (m, 2H); MS (ES) *m/e* 484.1 (M+H)⁺; retention time: 4.28 min.

[0113] Following similar procedures to those of Preparative Example 6, the compounds in Table 6 were prepared:

TABLE 6

Formula No.	COMPOUND	Mass Spec (M ⁺); retention time (min)
47		454.1; 4.44
48		522.1; 4.95

ASSAY:

[0114] The pharmacological properties of the compounds of this invention may be confirmed by a number of pharmacological assays. The exemplified pharmacological assays which are described later have been carried out with the compounds according to the invention and their salts.

[0115] Gamma secretase activity was determined as described by Zhang et al. (Biochemistry, 40 (16), 5049 -5055, 2001), which is herein incorporated by reference. Activity is expressed either as a percent inhibition or as the concentration of compound producing 50% inhibition of enzyme activity.

[0116] *Reagents.* Antibodies W02, G2-10, and G2-1 were obtained from Dr. Konrad Beyreuther (University of Heidelberg, Heidelberg, Germany). W02 recognizes residues 5-8 of Aβ peptide, while G2-10 and G2-11 recognize the specific C-terminal structure of Aβ 40 and Aβ 42, respectively. Biotin-4G8 was purchased from Senetec (St. Louis, MO). All tissue culture reagents used in this work were from Life Technologies, Inc., unless otherwise specified. Pepstatin A was purchased from Roche Molecular Biochemicals; DFK167 was from Enzyme Systems Products (Livermore, CA).

[0117] *cDNA Constructs, Tissue Culture, and Cell Line Construction.* The construct SPC99-Lon, which contains the first 18 residues and the C-terminal 99 amino acids of APP carrying the London mutation, has been described (Zhang, L., Song, L., and Parker, E. (1999) J. Biol. Chem. 274, 8966-8972). Upon insertion into the membrane, the 17 amino acid signal peptide is processed, leaving an additional leucine at the N-terminus of Aβ. SPC99-Lon was cloned into the pcDNA4/TO vector (Invitrogen) and transfected into 293 cells stably transfected with pcDNA6/TR, which is provided in the T-REx system (Invitrogen). The transfected cells were selected in Dulbecco's modified Eagle's media (DMEM)

supplemented with 10% fetal bovine serum, 100 units/mL penicillin, 100 g/mL streptomycin, 250 g/mL zeocin, and 5 g/mL blasticidin (Invitrogen). Colonies were screened for A β production by inducing C99 expression with 0.1 g/mL tetracycline for 16-20 h and analyzing conditioned media with a sandwich immunoassay (see below). One of the clones, designated as pTRE.15, was used in these studies.

[0118] *Membrane Preparation.* C99 expression in cells was induced with 0.1 g/mL tetracycline for 20 h. The cells were pretreated with 1 M phorbol 12-myristate 13-acetate (PMA) and 1 M brefeldin A (BFA) for 5-6 h at 37 C before harvesting. The cells were washed 3 times with cold phosphate-buffered saline (PBS) and harvested in buffer A containing 20 mM Hepes (pH 7.5), 250 mM sucrose, 50 mM KCl, 2 mM EDTA, 2 mM EGTA, and Complete protease inhibitor tablets (Roche Molecular Biochemicals). The cell pellets were flash-frozen in liquid nitrogen and stored at -70 C before use.

[0119] To make membranes, the cells were resuspended in buffer A and lysed in a nitrogen bomb at 600 psi. The cell lysate was centrifuged at 1500g for 10 min to remove nuclei and large cell debris. The supernatant was centrifuged at 100000g for 1 h. The membrane pellet was resuspended in buffer A plus 0.5 M NaCl, and the membranes were collected by centrifugation at 200000g for 1 h. The salt-washed membrane pellet was washed again in buffer A and centrifuged at 100000g for 1 h. The final membrane pellet was resuspended in a small volume of buffer A using a Teflon-glass homogenizer. The protein concentration was determined, and membrane aliquots were flash-frozen in liquid nitrogen and stored at -70°C.


[0120] *γ -Secretase Reaction and A β Analysis.* To measure γ -secretase activity, membranes were incubated at 37°C for 1 h in 50 μ L of buffer containing 20 mM Hepes (pH 7.0) and 2 mM EDTA. At the end of the incubation, A β 40 and A β 42 were measured using an electrochemiluminescence (ECL)-based immunoassay. A β 40 was identified with antibody pairs TAG-G2-10 and biotin-W02, while A β 42 was identified with TAG-G2-11 and biotin-4G8. The ECL signal was measured using an ECL-M8 instrument (IGEN International, Inc.) according to the manufacturer's instructions. The data presented were the means of the duplicate or triplicate measurements in each experiment. The characteristics of γ -secretase activity described were confirmed using more than five independent membrane preparations.

[0121] The compound of Example 16 had an IC₅₀ of about 16 nM. The compounds of Examples 18, 19, 20, 33 and 48 had an IC₅₀ within the range of about 50 nM to about 0.1 μ M. The compounds of Examples 14, 17, 21, 22, 23, 24, 25, 26, 27, 28, 29, 34 and 43 had an IC₅₀ within the range of about 0.1 to about 1.0 μ M. The compounds of Examples 15, 35, 36, 37 and 40 had an IC₅₀ within the range of about 1 to about 10 μ M.

[0122] While the present invention has been described in conjunction with the specific embodiments set forth above, many alternatives, modifications and variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present invention.

Claims

1. A compound, or pharmaceutically acceptable salts and/or solvates of said compound, said compound having the general structure shown in Formula I:

Formula I

wherein:

Ar is phenyl;

X is O;

m is 0, 1, 2 or 3, provided that when m = 0 no double bond is present and when m > 0 a double bond may or may not be present or a triple bond may or may not be present;

n is 0, 1, 2 or 3;

R¹ is 1 to 3 substituents independently selected from the group consisting of halogen;


R² is hydrogen, -alkyl, -cycloalkyl, -alkylene-cycloalkyl, -cycloalkyl, -NR⁶R⁷, -alkylene-NR⁶R⁷, -alkylene-O-alkyl, -O-alkyl, -heteroaryl, or -aryl;

R³ is 1 to 6 substituents, which can be the same or different, each being independently selected from hydrogen, halogen, -alkyl, -cycloalkyl, -OH, -OCF₃, -CF₃, -O-alkyl, -O-cycloalkyl, or -NR⁶R⁷;


R⁴ is hydrogen, -alkyl, -cycloalkyl, -aryl or -heteroaryl; or R³ and R⁴ can be joined together to form a 3 to 6 member ring;

R⁵ is -NR⁶R⁷, -N(R⁶)-alkylene-NR⁶R⁷; or

R⁵ is selected from the group consisting of:



R^6 and R^7 can be the same or different, each being independently selected from the group consisting of: hydrogen, -aryl, heteroaryl, -alkyl, -cycloalkyl, -alkylene-aryl, -alkylene-heteroaryl,



r is 0, 1, 2, 3 or 4;

s is 0, 1, 2 or 3;

R^8 is hydrogen, -OH, alkyl, -C(O)NR⁶R⁷, -C(O)O-alkyl, or aryl, wherein said alkyl moiety may be unsubstituted or substituted with 1 to 4 hydroxy groups, and said aryl group may be unsubstituted or substituted with at least one substituent selected from the group consisting of alkyl, cycloalkyl, halogen, -CF₃, -OCF₃, -NR⁶R⁷, -CN, -NO₂, -NH₂, -alkylene-NR⁶R⁷, -alkylene-O-alkyl, -C(O)O-alkyl, -C(O)NR⁶R⁷, -N(R⁶)C(O)alkyl, -N(R⁶)C(O)cycloalkyl, -N(R⁶)C(O)aryl, -N(R⁶)C(O)heteroaryl, -N(R⁶)C(O)O-alkyl, -N(R⁶)C(O)NR⁶R⁷, -N(R⁶)S(O)₂alkyl, -OH, -O-alkyl, and -O-cycloalkyl;

R^9 is selected from the group consisting of: hydrogen, alkyl, -cycloalkyl, -alkylene-aryl, -alkylene-heteroaryl, -alkyl-cycloalkyl, -C(O)O-alkyl, -alkylene-O-alkylene-OH, -aryl and -heteroaryl, wherein each said alkyl and cycloalkyl moieties may be independently substituted or unsubstituted, and if substituted then substituted with 1 to 4 hydroxy groups or at least one -NR⁶R⁷ group;

each R^{10} is independently selected from hydrogen or alkyl wherein said alkyl group may be substituted or unsubstituted and if substituted then R^{10} is substituted with 1 to 4 hydroxy groups;

each R^{11} and R^{12} are each independently selected from hydrogen, alkyl, or aryl; or

R^{11} and R^{12} , together with the ring carbon atoms to which they are shown attached in Formula I, form a cycloalkyl ring, and wherein the cycloalkyl ring formed thereby can be unsubstituted or substituted with an alkyl, alkoxy, aryl, or halogen;

p is 0, 1, 2, 3 or 4; and


q is 0, 1 or 2.

2. The compound according to Claim 1, wherein R^2 is H.

3. The compound according to Claim 1, wherein R^3 is H.

4. The compound according to Claim 1, wherein R^4 is hydrogen.

5. The compound according to Claim 1, wherein R^5 is: NR⁶R⁷, -NR⁶-alkylene-NR⁶R⁷,




6. The compound according to claim 1, wherein m is 0 or 1.

7. The compound according to claim 1, wherein n is 1 or 2.

8. The compound according to Claim 1, wherein R^4 is $-CH_3$.


9. The compound according to claim 1, wherein R^1 is 4-chloro.

10. The compound according to claim 5, wherein R^5 is:



11. The compound according to Claim 1, wherein:

R^2 is H;
 R^3 is H;
 R^4 is H; and
 R^5 is:




12. The compound according to Claim 11 wherein one double bond is present within a bicyclic ring in the structure represented by Formula I.

13. The compound according to Claim 11, wherein $m = 1$ and $n = 2$.

14. The compound according to Claim 11, wherein $m = 1$ and $n = 1$.

15. The compound according to Claim 11, wherein $m = 0$ and $n = 1$.

16. The compound according to Claim 11, wherein R^5 is



17. A compound selected from the group consisting of:

5

10

15

20

25

30









35

40

45

50

55



5

10

15

20

25

30


35


40

45

50


55






and


18. A compound or a pharmaceutically acceptable salt and/or solvate thereof, wherein the compound has the following structure:




19. A compound or a pharmaceutically acceptable salt and/or solvate thereof wherein the compound has the following structure:




20. A compound or a pharmaceutically acceptable salt and/or solvate thereof, wherein the compound has the following structure:




21. A compound or a pharmaceutically acceptable salt and/or solvate thereof, wherein the compound has the following structure:



22. A compound or a pharmaceutically acceptable salt and/or solvate thereof, wherein the compound has the following structure:



23. A compound or a pharmaceutically acceptable salt and/or solvate thereof, wherein the compound has the following structure:



24. A pharmaceutical composition comprising at least one compound of Claim 1 and at least one pharmaceutically acceptable carrier.

25. Use of one or more compounds of Claim 1 for the manufacture of a medicament for inhibiting gamma-secretase in a patient in need of such treatment.

26. Use of one or more compounds of Claim 1 for the manufacture of a medicament for treating one or more neurodegenerative diseases in a patient in need of such treatment.


27. Use of one or more compounds of Claim 1 for the manufacture of a medicament for inhibiting the deposition of beta amyloid protein in a patient in need of such inhibition.

28. Use of one more compounds of Claim 1 for the manufacture of a medicament for treating Alzheimer's in a patient

in need of such treatment.

Patentansprüche

1. Verbindung oder pharmazeutisch annehmbare Salze und/oder Solvate der Verbindung, wobei die Verbindung die in Formel I gezeigte allgemeine Struktur hat:



Formel I

worin:

Ar Phenyl ist,

X 0 ist,

m 0, 1, 2 oder 3 ist, mit der Maßgabe, dass keine Doppelbindung vorhanden ist, wenn m = 0 ist, und eine Doppelbindung vorhanden sein kann oder nicht oder eine Dreifachbindung vorhanden sein kann oder nicht, wenn m > 0 ist,

n 0, 1, 2 oder 3 ist,

R¹ 1 bis 3 Substituenten ist, die unabhängig ausgewählt sind aus der Gruppe bestehend aus Halogen,

R² Wasserstoff, -Alkyl, -Cycloalkyl, -Alkylen-cycloalkyl, -Cycloalkyl, -NR⁶R⁷, -Alkylen-NR⁶R⁷, -Alkylen-O-alkyl, -O-Alkyl, -Heteroaryl oder -Aryl ist,


R³ 1 bis 6 Substituenten ist, die gleich oder unterschiedlich sein können, wobei jeder unabhängig ausgewählt ist aus Wasserstoff, Halogen, -Alkyl, -Cycloalkyl, -OH, -OCF₃, -CF₃, -O-Alkyl, -O-Cycloalkyl oder -NR⁶R⁷,

R⁴ Wasserstoff, -Alkyl, -Cycloalkyl, -Aryl oder -Heteroaryl ist, oder


R³ und R⁴ unter Bildung eines 3- bis 6-gliedrigen Rings verbunden sein können,

R⁵ -NR⁶R⁷, -N(R⁶)-Alkylen-NR⁶R⁷ ist, oder

R⁵ ausgewählt ist aus der Gruppe bestehend aus:



R^6 und R^7 gleich oder unterschiedlich sein können und jeweils unabhängig ausgewählt sind aus der Gruppe bestehend aus Wasserstoff, -Aryl, Heteroaryl, -Alkyl, -Cycloalkyl, -Alkylen-aryl, -Alkylen-heteroaryl,



r 0, 1, 2, 3 oder 4 ist;

s 0, 1, 2 oder 3 ist,

R^8 Wasserstoff, -OH, Alkyl, -C(O)NR⁶R⁷, -C(O)O-Alkyl oder Aryl ist, wobei die Alkyleinheit unsubstituiert oder mit 1 bis 4 Hydroxygruppen substituiert sein kann, und die Arylgruppe unsubstituiert oder mit mindestens einem Substituenten ausgewählt aus der Gruppe bestehend aus Alkyl, Cycloalkyl, Halogen, -CF₃, -OCF₃, -NR⁶R⁷, -CN, -NO₂, -NH₂, -Alkylen-NR⁶R⁷, -Alkylen-O-alkyl, -C(O)O-Alkyl, -C(O)NR⁶R⁷, -N(R⁶)C(O)Alkyl, -N(R⁶)C(O)Cycloalkyl, -N(R⁶)C(O)Aryl, -N(R⁶)C(O)Heteroaryl, -N(R⁶)C(O)O-Alkyl, -N(R⁶)C(O)NR⁶R⁷, -N(R⁶)S(O)₂alkyl, -OH, -O-Alkyl und -O-Cycloalkyl substituiert sein kann;

R^9 ausgewählt ist aus der Gruppe bestehend aus: Wasserstoff, Alkyl, -Cycloalkyl, -Alkylen-aryl, -Alkylen-heteroaryl, -Alkyl-cycloalkyl, -C(O)O-Alkyl, -Alkylen-O-alkylen-OH, -Aryl und -Heteroaryl, wobei jede der Alkyl- und Cycloalkyleinheiten unabhängig substituiert oder unsubstituiert sein kann, und wenn sie substituiert ist, dann mit 1 bis 4 Hydroxygruppen oder mindestens einer -NR⁶R⁷ Gruppe substituiert sein kann;

jedes R^{10} unabhängig ausgewählt ist aus Wasserstoff oder Alkyl, wobei die Alkylgruppe substituiert oder unsubstituiert sein kann, und falls sie substituiert ist, dann R^{10} mit 1 bis 4 Hydroxygruppen substituiert ist,

jedes R^{11} und R^{12} unabhängig ausgewählt ist aus Wasserstoff, Alkyl oder Aryl, oder

R^{11} und R^{12} zusammen mit den Ringkohlenstoffatomen, an die sie in Formel I gebunden gezeigt sind, einen Cycloalkylring bilden, und wobei der dadurch gebildete Cycloalkylring unsubstituiert oder mit Alkyl, Alkoxy, Aryl oder Halogen substituiert sein kann,

p 0, 1, 2, 3 oder 4 ist, und


q ist 0, 1 oder 2.

2. Verbindung nach Anspruch 1, bei der R^2 H ist.

3. Verbindung nach Anspruch 1, bei der R^3 H ist.

4. Verbindung nach Anspruch 1, bei der R^4 Wasserstoff ist.

5. Verbindung nach Anspruch 1, bei der R^5 NR⁶R⁷, -NR⁶-Alkylen-NR⁶R⁷,



ist.


6. Verbindung nach Anspruch 1, bei der m 0 oder 1 ist.

7. Verbindung nach Anspruch 1, bei der n 1 oder 2 ist.

8. Verbindung nach Anspruch 1, bei der R⁴ -CH₃ ist.

9. Verbindung nach Anspruch 1, bei der R¹ 4-Chlor ist.


10. Verbindung nach Anspruch 5, bei der R⁵



ist.

11. Verbindung nach Anspruch 1, worin:

R² H ist,
R³ H ist,
R⁴ H ist, und
R⁵



ist.


12. Verbindung nach Anspruch 11, bei der in einem bicyclischen Ring in der durch Formel I wiedergegebenen Struktur eine Doppelbindung vorhanden ist.

13. Verbindung nach Anspruch 11, bei der m 1 ist und n 2 ist.

14. Verbindung nach Anspruch 11, bei der m 1 ist und n 1 ist.

15. Verbindung nach Anspruch 11, bei der m 0 ist und n 1 ist.

16. Verbindung nach Anspruch 11, bei der R⁵




ist.

17. Verbindung ausgewählt aus der Gruppe bestehend aus:

5

10

15




1


20

25

30



2




3


35

40

45




4






5


50

55







55




und



18. Verbindung oder pharmazeutisch annehmbares Salz und/oder Solvat davon, wobei die Verbindung die folgende Struktur hat:



19. Verbindung oder pharmazeutisch annehmbares Salz und/oder Solvat davon, wobei die Verbindung die folgende Struktur hat:




20. Verbindung oder pharmazeutisch annehmbares Salz und/oder Solvat davon, wobei die Verbindung die folgende Struktur hat:

5

10

15




20

21. Verbindung oder pharmazeutisch annehmbares Salz und/oder Solvat davon, wobei die Verbindung die folgende Struktur hat:

25

30

35




40

22. Verbindung oder pharmazeutisch annehmbares Salz und/oder Solvat davon, wobei die Verbindung die folgende Struktur hat:


45

50

55




23. Verbindung oder pharmazeutisch annehmbares Salz und/oder Solvat davon, wobei die Verbindung die folgende Struktur hat:



24. Pharmazeutische Zusammensetzung, die mindestens eine Verbindung nach Anspruch 1 und mindestens einen pharmazeutisch annehmbaren Träger enthält.
25. Verwendung von einer oder mehreren Verbindungen gemäß Anspruch 1 zur Herstellung eines Medikaments zum Inhibieren von γ -Sekretase bei einem Patienten, der dieser Behandlung bedarf.
26. Verwendung von einer oder mehreren Verbindungen gemäß Anspruch 1 zur Herstellung eines Medikaments zur Behandlung von einer oder mehreren neurodegenerativen Erkrankungen bei einem Patienten, der dieser Behandlung bedarf.
27. Verwendung von einer oder mehreren Verbindungen gemäß Anspruch 1 zur Herstellung eines Medikaments zum Inhibieren der Ablagerung von β -Amyloidprotein bei einem Patienten, der dieser Inhibierung bedarf.
28. Verwendung von einer oder mehreren Verbindungen gemäß Anspruch 1 zur Herstellung eines Medikaments zur Behandlung von Morbus Alzheimer bei einem Patienten, der dieser Behandlung bedarf.

Revendications


1. Composé, ou sel et/ou solvat pharmacologiquement admissible d'un tel composé, lequel composé présente la structure générale représentée par la formule I :




Formule I

dans laquelle :

- Ar représente un groupe phényle ;
- X représente un atome d'oxygène ;
- l'indice m vaut 0, 1, 2 ou 3, étant entendu que, si m est nul, il n'y a pas de double liaison, et si m est supérieur à zéro, il peut y avoir ou non une double liaison ou une triple liaison ;
- l'indice n vaut 0, 1, 2 ou 3 ;
- R¹ représente 1 à 3 substituants choisis indépendamment parmi les atomes d'halogène ;
- R² représente un atome d'hydrogène, un groupe alkyle, cycloalkyle, cycloalkyl-alkyle, cycloalkyle, alcoxy-alkyle, alcoxy, hétéroaryle ou aryle, ou un groupe symbolisé par -NR⁶R⁷ ou -alkylène-NR⁶R⁷ ;
- R³ représente 1 à 6 substituants, qui peuvent être identiques ou différents et qui sont choisis, chacun indépendamment, parmi les atomes d'hydrogène et d'halogène, les groupes alkyle, cycloalkyle, alcoxy et cycloalcoxy, et les groupes symbolisés par -OH, -CF₃, -OCF₃ ou -NR⁶R⁷ ;
- R⁴ représente un atome d'hydrogène ou un groupe alkyle, cycloalkyle, hétéroaryle ou aryle ;
- ou bien R³ et R⁴ représentent des entités qui forment conjointement un cycle de 3 à 6 chaînons ;
- R⁵ représente un groupe symbolisé par -NR⁶R⁷ ou -N(R⁶)-alkylène-NR⁶R⁷, ou un groupe choisi parmi ceux de formules suivantes :



- R⁶ et R⁷ représentent des entités qui peuvent être identiques ou différentes et qui sont choisies, chacune indépendamment, parmi un atome d'hydrogène, les groupes aryle, hétéroaryle, alkyle, cycloalkyle, aryl-alkyle et hétéroaryl-alkyle, et les groupes représentés par les formules



- l'indice r vaut 0, 1, 2, 3 ou 4 ;

- l'indice s vaut 0, 1, 2 ou 3 ;

- R^8 représente un atome d'hydrogène, un groupe alkyle ou aryle, ou un groupe symbolisé par -OH, -C(O)NR⁶R⁷ ou -C(O)O-alkyle, ledit groupe ou fragment alkyle pouvant ne porter aucun substituant ou porter 1 à 4 substituants hydroxyle, et ledit groupe aryle pouvant ne porter aucun substituant ou porter au moins un substituant choisi dans l'ensemble constitué par les atomes d'halogène, les groupes alkyle, cycloalkyle, alcoxy-alkyle, hydroxyle, alcoxy et cycloalcoxy, et les groupes symbolisés par -CF₃, -OCF₃, -NR⁶R⁷, -CN, -NO₂, -NH₂, -alkylène-NR⁶R⁷, -C(O)NR⁶R⁷, -C(O)O-alkyle, -N(R⁶)C(O)-alkyle, -N(R⁶)C(O)-cyclo-alkyle, -N(R⁶)C(O)-aryle, -N(R⁶)C(O)-hétéroaryle, -N(R⁶)C(O)O-alkyle, -N(R⁶)C(O)NR⁶R⁷ ou -N(R⁶)S(O)₂-alkyle ;

- R^9 représente une entité choisie dans l'ensemble constitué par un atome d'hydrogène et les groupes alkyle, cycloalkyle, aryl-alkyle, hétéroaryl-alkyle, cycloalkyl-alkyle, alcoxy-carbonyle, hydroxy-alcoxy-alkyle, aryle et hétéroaryle, chacun desdits groupes ou fragments alkyle et cycloalkyle pouvant indépendamment porter un ou des substituant(s) ou n'en porter aucun, et s'il en porte, porter 1 à 4 substituants hydroxy ou au moins un substituant symbolisé par -NR⁶R⁷ ;

- chaque symbole R^{10} représente une entité choisie indépendamment dans l'ensemble constitué par un atome d'hydrogène et les groupes alkyle, lesquels groupes alkyle peuvent porter un ou des substituant(s) ou n'en porter aucun, et s'ils en portent, porter 1 à 4 substituants hydroxy ;

- chaque symbole R^{11} ou R^{12} représente une entité choisie indépendamment dans l'ensemble constitué par un atome d'hydrogène et les groupes alkyle et aryle ;

- ou les symboles R^{11} et R^{12} représentent des entités qui forment, conjointement avec les atomes de carbone de cycle auxquels elles sont liées d'après la formule I, un groupe cycloalkyle qui peut ne porter aucun substituant ou porter un substituant alkyle, alcoxy, aryle ou halogéno ;

- l'indice p vaut 0, 1, 2, 3 ou 4 ;


- et l'indice q vaut 0, 1 ou 2.

2. Composé conforme à la revendication 1, dans lequel R^2 représente un atome d'hydrogène.

3. Composé conforme à la revendication 1, dans lequel R^3 représente un atome d'hydrogène.


4. Composé conforme à la revendication 1, dans lequel R^4 représente un atome d'hydrogène.

5. Composé conforme à la revendication 1, dans lequel R^5 représente un groupe symbolisé par -NR⁶R⁷ ou -N(R⁶)-alkylène-NR⁶R⁷, ou un groupe de formule




6. Composé conforme à la revendication 1, dans lequel l'indice m vaut 0 ou 1.

7. Composé conforme à la revendication 1, dans lequel l'indice n vaut 1 ou 2.
8. Composé conforme à la revendication 1, dans lequel R^4 représente un groupe méthyle.
9. Composé conforme à la revendication 1, dans lequel R^1 représente un atome de chlore placé en position 4.
10. Composé conforme à la revendication 5, dans lequel R^5 représente un groupe de formule



11. Composé conforme à la revendication 1, dans lequel :

- R^2 représente un atome d'hydrogène ;
- R^3 représente un atome d'hydrogène ;
- R^4 représente un atome d'hydrogène ;
- et R^5 représente un groupe de formule





12. Composé conforme à la revendication 11, dans lequel il y a une double liaison au sein d'un groupe bicyclique, dans la structure représentée par la formule I.
13. Composé conforme à la revendication 11, dans lequel l'indice m vaut 1 et l'indice n vaut 2.
14. Composé conforme à la revendication 11, dans lequel l'indice m vaut 1 et l'indice n vaut 1.
15. Composé conforme à la revendication 11, dans lequel l'indice m vaut 0 et l'indice n vaut 1.
16. Composé conforme à la revendication 11, dans lequel R^5 représente un groupe de formule :



17. Composé choisi dans l'ensemble formé par les suivants :



5

10





15

20





25

30



35

40



45

50

55

5

10

15

20

25

30


35

40

45

50

55



5

10

15


20

25

30

35

40



45


18. Composé, ou sel et/ou solvat pharmacologiquement admissible d'un tel composé, lequel composé présente la structure suivante :

50

55

5

10




15

19. Composé, ou sel et/ou solvat pharmacologiquement admissible d'un tel composé, lequel composé présente la structure suivante :

20

25




30

20. Composé, ou sel et/ou solvat pharmacologiquement admissible d'un tel composé, lequel composé présente la structure suivante :

35

40



45


21. Composé, ou sel et/ou solvat pharmacologiquement admissible d'un tel composé, lequel composé présente la structure suivante :

50

55

5

10




15

- 22.** Composé, ou sel et/ou solvat pharmacologiquement admissible d'un tel composé, lequel composé présente la structure suivante :

20

25




30

- 23.** Composé, ou sel et/ou solvat pharmacologiquement admissible d'un tel composé, lequel composé présente la structure suivante :

35

40



45

- 24.** Composition pharmaceutique comprenant au moins un composé conforme à la revendication 1 et au moins un véhicule pharmacologiquement admissible.

50

- 25.** Emploi d'un ou de plusieurs composés conformes à la revendication 1 en vue de la fabrication d'un médicament conçu pour inhiber la gamma-sécrétase chez un patient qui a besoin d'un tel traitement.

55

- 26.** Emploi d'un ou de plusieurs composés conformes à la revendication 1 en vue de la fabrication d'un médicament conçu pour traiter une ou plusieurs maladies neurodégénératives chez un patient qui a besoin d'un tel traitement.

- 27.** Emploi d'un ou de plusieurs composés conformes à la revendication 1 en vue de la fabrication d'un médicament conçu pour inhiber la formation de dépôts de protéine bêta-amyloïde chez un patient qui a besoin d'une telle inhibition.

EP 1 641 801 B9

- 28.** Emploi d'un ou de plusieurs composés conformes à la revendication 1 en vue de la fabrication d'un médicament conçu pour traiter la maladie d'Alzheimer chez un patient qui a besoin d'un tel traitement.

5

10

15

20

25

30

35

40

45

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 47014603 P [0001]
- WO 0050391 A [0002]

Non-patent literature cited in the description

- Pro-drugs as Novel Delivery Systems. **T. HIGUCHI ; V. STELLA**. A.C.S. Symposium Series, and in Biorversible Carriers in Drug Design. American Pharmaceutical Association and Pergamon Press, 1987, vol. 14 [0040]
- **S. BERGE et al.** *Journal of Pharmaceutical Sciences*, 1977, vol. 66 (1), 1-19 [0044]
- **P. GOULD**. *International J. of Pharmaceutics*, 1986, vol. 33, 201-217 [0044]
- **ANDERSON et al.** *The Practice of Medicinal Chemistry*. Academic Press, 1996 [0044]
- Food & Drug Administration. *The Orange Book* [0044]
- Remington's Pharmaceutical Sciences. Mack Publishing Co, 1990 [0051]
- **ZHANG et al.** *Biochemistry*, 2001, vol. 40 (16), 5049-5055 [0115]
- **ZHANG, L. ; SONG, L. ; PARKER, E.** *J. Biol. Chem.*, 1999, vol. 274, 8966-8972 [0117]