



(11) **EP 1 675 474 B9**

(12) **CORRECTED NEW EUROPEAN PATENT SPECIFICATION**

(15) Correction information:
Corrected version no 1 (W1 B2)
Corrections, see
Description Paragraph(s) 1, 89, 90, 94, 95,
96, 125

(51) Int Cl.:
A23K 50/42 ^(2016.01) **A23K 40/00** ^(2016.01)
A23K 20/163 ^(2016.01) **A23K 20/116** ^(2016.01)
A61K 9/00 ^(2006.01)

(48) Corrigendum issued on:
05.12.2018 Bulletin 2018/49

(86) International application number:
PCT/EP2004/008538

(45) Date of publication and mention
of the opposition decision:
07.12.2016 Bulletin 2016/49

(87) International publication number:
WO 2005/013714 (17.02.2005 Gazette 2005/07)

(45) Mention of the grant of the patent:
22.10.2008 Bulletin 2008/43

(21) Application number: **04763629.5**

(22) Date of filing: **29.07.2004**

(54) **PALATABLE DUCTILE CHEWABLE VETERINARY COMPOSITION**

PLASTISCHES KAUBARES TIERARZNEIPRÄPARAT MIT HOHER PALATABILITÄT
COMPOSITION VETERINAIRE DUCTILE A MACHER SAPIDE

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PL PT RO SE SI SK TR

(74) Representative: **Sexton, Jane Helen**
J A Kemp
14 South Square
Gray's Inn
London WC1R 5JJ (GB)

(30) Priority: **30.07.2003 EP 03017252**

(43) Date of publication of application:
05.07.2006 Bulletin 2006/27

(56) References cited:
EP-A- 1 247 456 WO-A1-02/00202
WO-A1-2004/016252 GB-A- 2 300 103
US-A- 4 393 085 US-A- 5 824 336
US-A1- 2001 036 464 US-B1- 6 500 463

(73) Proprietors:
• **Novartis Tiergesundheit AG**
4058 Basel (CH)
Designated Contracting States:
BE BG CH CY CZ DE DK EE ES FI FR GB GR HU
IE IT LI LU MC NL PL PT RO SE SI SK TR
• **Novartis Pharma GmbH**
1230 Wien (AT)
Designated Contracting States:
AT

- **PATENT ABSTRACTS OF JAPAN** vol. 018, no. 371 (C-1224), 13 July 1994 (1994-07-13) & JP 06 100602 A (ASAHI CHEM IND CO LTD), 12 April 1994 (1994-04-12)
- **'Starch 1500', 1999, COLORCON** deel 'Partially pregelatinized maize'

(72) Inventor: **ISELE, Ute**
79115 Freiburg (DE)

EP 1 675 474 B9

Description

[0001] The present invention relates to an easy-to-use, safe, efficacious, and stable veterinary formulation consisting of a highly palatable ductile chewable veterinary composition comprising an effective amount of one or more ingredients that are active against animal pests, pathogens or animal diseases; of meat flavoring; of partially gelatinized starch; of a softener; and of up to 9% of water. Further, the invention relates to a process for producing said highly palatable ductile chewable veterinary composition by cold extrusion. In a preferred embodiment, the highly palatable ductile chewable veterinary composition controls animal pests like endo-parasites, such as worms, and simultaneously ecto-parasites, such as biting insects like fleas, on pets. The pesticidally effective ingredient is dispensed as the animal chews the product.

Field and background of the Invention

[0002] Veterinary products can be administered to warm-blooded animals in very different ways depending on their mode of action and their ability to be taken up either by the treated animal or the target pest. Thus veterinary products can be administered, for example, topically as pour-on or spot-on formulations, in form of shampoos, showers, as a dip, bath or spray, in form of a collar, and in many variants of these application forms. They can also be administered systemically, for example, orally, parenterally and in certain cases even transdermally. Examples of systemic administration forms are: via injection, as a tablet, capsule, bolus, drink, feed additive and the like. Each of these administration forms can have advantages or disadvantages depending on the actual situation and the animal that is in need of such a treatment. Treatment of herd animals, like horses, cattle, sheep or poultry usually requires different administration methods than for the treatment of single animals, such as pets like dogs and cats.

[0003] One very convenient and easy to manage administration form for human patients is the oral uptake of a medicament. This would also be very desirable in the field of veterinary medicine but here the animal holder or veterinarian is confronted with the natural behavior of the animal and oral treatment can be a real challenge.

[0004] Many attempts have been made to design the ideal oral application form that is really accepted and voluntarily taken up by the animal but most of these application forms need still to be improved.

[0005] For example, EP-A-1247456 describes a palatable pharmaceutical composition for oral administration to companion animals comprising a pharmaceutically effective amount of a pharmaceutically active agent in combination with a palatability improving agent and a pharmaceutically acceptable carrier, wherein the palatability improving agent is non-meat and non-fish derived, and is present in amounts sufficient to make the pharmaceutical composition palatable to the companion animal. The subject matter of the present invention differs from that disclosed in said reference in that partially gelatinized starch is used in the present composition. The effect of this difference is that the product may be produced by cold extrusion.

[0006] GB-A-2300103 describes a palatable dog biscuits and their production via an extrusion process. The resulting product comprises partially gelatinized starch and creatine. However, while addressing the problem of destruction of the active ingredient, the document does not provide a means of cold extrusion. Rather the extrusion occurs at relatively high temperatures up to 130°C.

[0007] The present inventors recognized that in the field of animal health the dosage form and especially the palatability of the dosage form, i.e. the natural acceptance of the drug plays the decisive role. The underlying problems are outlined hereinafter.

[0008] While, in humans, medicaments may be administered in a wide variety of application forms, such as tablets, coated tablets, emulsions, injection solutions, suppositories and the like, because the discipline and the desire to recover in human patients can be relied upon, in the case of animals practical problems are soon encountered, since a few application forms, such as the usage of suppositories, either have to be dispensed with all together or other forms, such as injections, must only be carried out by the veterinarian.

[0009] In general, humans do not like to visit the doctor. The same is true for animal keepers who would need advice from a veterinarian. In general, the animal keeper prefers to use those treatment methods that he can carry out himself without having involved a veterinarian. Among the preferred treatment methods, which an animal keeper can carry out himself, e.g. following the veterinarian's instructions, is the oral administration of medicaments.

[0010] Treating humans with medicines is generally not problematic, because the human patient follows the advice of the doctor or reads the directions on the leaflet in the package and complies with them since this is in his own interest, and because the manufacturer usually prepares the tablet, capsule or coated tablet in a form which is appropriate for oral consumption and has been tailored for human patients.

[0011] However, as soon as a pharmaceutical active ingredient has a taste which is unpleasant to the animal, whether because it is bitter or has some other unpleasant taste or is simply alien to the animal, the animal refuses to take it orally. This inborn behavior occurs to varying degrees among the different species of animals, and essentially depends on their conventional eating habits. Unfortunately, only a few active ingredients have a neutral taste, so that the problem being discussed here is almost always present.

[0012] In the case of a human patient, an unpleasant tasting active ingredient can be masked relatively easily, e.g. by coating it with a neutral-tasting or sweet layer. Everybody has come across gelatin capsules or tablets coated with sugar or lacquer at some time or other. It is easy to instruct the human patient to take the preparation without chewing.

[0013] An animal must have a natural willingness to take a medicinal preparation orally, which means that the medical preparation must taste well and be palatable. Of course, an individual animal or a few animals can also be forced to take a medicament, by making it swallow or by injecting it. However, such forced methods are not only unacceptable to large animal operations but also to single dogs and cats which tend to bite or scrape if they are not willing to be treated. This is why animal treatment can be very labor-intensive or can require the intervention of a veterinarian and this ultimately leads to increase of costs.

[0014] Therefore, for pets but equally for animals that are kept on a large scale, simple and safe oral application forms are required, which can be easily administered by the animal keeper, which lead to reliable results, and which are affordable.

[0015] The chewable composition according to the present invention is not only suitable for replacing the treatment with a tablet or capsule. These chewables can also easily be mixed with conventional non-medicated feed pellets if herds of animals have to be treated.

[0016] Due to their excellent palatability the chewables according to the present invention are taken up by animals without causing any acceptance problems. Their handling is easy and safe, and can be adapted to the need either of an individual animal like a cat or a dog or to a herd animals like sheep and cows.

[0017] When reviewing the administration of capsules and coated tablets to animals, it has been shown that these application forms are rather unsuitable for animal medicine, since in the case of herd animals they can only be used in a controlled manner with considerable effort on a daily basis, and in the case of pets, such as dogs and cats, lead to particular acceptance problems. As already mentioned above, the eating habits of animals generally play a decisive role when using oral application forms. Thus, most important is an attractive taste and the palatability.

[0018] In the case of dogs, it has been observed that they gnaw at solid food, e.g. on bones, and gulp down other food, either in the form of large scraps or wet formulated food, almost unchewed. If a tablet or coated tablet is mixed with the wet formulated feed, varying results are obtained. In a few cases, the tablet is not noticed by the dog at all and is simply gulped down, and in other cases it remains uneaten in the dog bowl. In contrast to dogs, cats are considerably more fastidious in their eating habits. Only in the rarest cases can a tablet or coated tablet be mixed with the formulated food, without them noticing it immediately and rejecting it. Although cats also do not exactly chew their food, they generally break it down with a few small bites. They thereby damage the protective coating of a tablet or capsule and release the unpleasant tasting active ingredient. Attempts to mix the active ingredient directly with the feed likewise fail, because either the degree of dilution is insufficient to neutralize the unpleasant taste or the active ingredient breaks down too rapidly when in contact with the feed. For the same reasons, mixtures of feed, active ingredient and excipients, which should stimulate the appetite of dogs and cats, similarly do not have a successful outcome with cats. Whereas the test animals rush eagerly to a placebo which has a corresponding appetite stimulant, i.e. a tablet consisting of feed, flavoring and other excipients, but no active ingredient, the test animals reject the same combination as soon as active ingredient is added. Clearly, a different technical solution must be found to the existing problem with animals.

[0019] Of course, any other active ingredient which is suitable for animals can be administered according to the present invention, but especially those active ingredients that have the taste disadvantages mentioned initially and are therefore not willingly taken orally by animals.

[0020] Basically, a diversity of individual active ingredients or mixtures of active ingredients may be considered, e.g., those acting against external (ecto) or internal (endo) parasites or active ingredients acting against animal diseases including viral or bacterial infections, behavioral disorders, such as hypo- or hyper-activity, inflammatory diseases, and auto-immune diseases. Thus, the active ingredient can be a pesticide or a medicament or a mixture of both.

[0021] It should be kept in mind that the present invention deals with an optimized application form for veterinary compositions rather than with the treatment of animals with a specific class of active ingredients. On the contrary, the present invention provides an easy-to-use, safe, powerful, and stable veterinary formulation consisting of a highly palatable ductile chewable veterinary composition, which allows to administer orally almost each and any active ingredient to a warm-blooded animal, provided that this active ingredient or mixture of active ingredients is at the administered dose physiologically acceptable to the animal, does not display unacceptable side effects and, what is most important, exhibits after oral uptake systemic activity. This means that the main prerequisite for the active ingredient is that after oral administration it is taken up by the body fluids, including blood and lymph, and transported to the animal pest, the pathogen or the diseased organ where it can exhibit its activity. Thus, any active ingredient or class of active ingredients mentioned hereinafter is nothing but a non-limiting example of suitable active ingredients. The application form of the present invention is actually not limited to existing active ingredients but also suitable for each and any active ingredient developed in the future provided that the future active ingredient meets the main characteristics explained hereinbefore.

[0022] The highly palatable ductile chewable veterinary composition of the present invention is in principle a medicated food product and everybody working in this area is aware of the technical problems that arise in context with the production

of medicated feed. For example, stability of the active ingredient is very crucial. It is a matter of fact that many potent active compounds are somewhat unstable (temperature-sensitive), above all when in contact with feed material, especially close contact to vegetable and animal materials, during conventional extrusion of feed pellets, result in considerable losses of active ingredient.

[0023] For example, when feed pellets are prepared via extrusion, the dried organic starting material of animal or vegetable origin is ground, is intimately mixed with the active ingredient, that is to say is substantially homogenized, and then is moistened with water or steam and is compressed into pellets at elevated temperatures and under pressures of around 100 kbar. However, said high pressures and the permanent high temperatures in the range of 60-100°C are disadvantageous and do not only dramatically reduce the viscosity of the pellets but result in a considerable loss of active ingredient.

[0024] Whereas most active ingredients in pure form or in contact with carriers that are routinely used in the production of tablets or capsules withstand such relatively high temperatures per se very well and can be stored in pure form or as tablets or capsules at room temperature for months or years without any measurable loss of active ingredient, they decompose relatively rapidly under pressure and in intimate contact with animal or vegetable fibers in feedstuffs and under the prevailing elevated temperatures. It appears that contact with the fibers actually catalyses the decomposition process. Even when the elevated-pressure and elevated-temperature phase is kept as short as technically possible and the finished pellets are immediately cooled down to room temperature directly after the compression process, a quarter to a third of the active ingredient is nevertheless lost. Even though in the rare cases where the degradation products do not have disadvantageous effects on the animals treated, the unavoidable loss of active ingredient inevitably results in a considerable increase in the cost of the final product. Thus extrusion processes can lead to very undesirable effects.

[0025] For the reasons mentioned, therefore, much effort has been directed at stabilizing temperature-sensitive active ingredients so that they withstand the elevated temperatures and pressures during pellet preparation without loss of active substance and also, when in the form of the finished pellets, have a long-term storage stability suitable for practical purposes.

[0026] Unsuccessful attempts at such stabilization include, for example, (1) reduction of the active ingredient surface area by means of compression into granules, a very great variety of granule sizes having been tried; (2) sealing of the said active ingredient granules in a very great variety of protective layers, for example gelatin or various sugars and coatings; (3) enclosure of the active ingredient within porous materials such as, for example, various celluloses, starches, silicic acids or zeolites, with or without additional protective layers; and (4) chemical modification of the basic macrocyclic structure of the active ingredient. Although in a few cases chemical modification has resulted in improved stability of the compound per se, it has simultaneously resulted in loss of activity.

[0027] However, none of those attempts has resulted in an appreciably smaller loss of active ingredient on compression into feed pellets or in measurably improved storage stability.

[0028] Moreover, success has now been achieved, surprisingly, in providing the user with the user-friendly, easy-to-use, safe, powerful, stable, and especially highly palatable chewable veterinary composition of the present invention.

[0029] Astonishingly, it is now possible to provide a product that not only withstands the extrusion process undamaged but also survives for a outstanding long storage period.

[0030] Therefore, it is highly surprising and was absolutely unpredictable that even so the chewable veterinary composition of the present invention contains a relatively high amount of meat material, this has obviously when combined with the appropriate amount of partially gelatinized starch, no adverse effect on the stability of the active ingredient. It actually turned out that the chewable veterinary composition of the present invention is a very stable product that can be stored at room temperature over many months without significant loss or degradation of active ingredient. Tests with stored material demonstrate that the palatability is not decreased and the efficacy of the active ingredient stays at a high level.

[0031] Moreover, investigations of the kinetic behavior demonstrate another surprising effect. It could not have been foreseen that the administration of the chewable veterinary composition of the present invention could lead to absolutely the same level of bioavailability as the administration of tablets or capsules. Thus, the present invention provides a safe, easy to use and stable product that is at least as efficacious as conventional oral application forms, like tablets or capsules.

[0032] Many biocides and veterinary medicines that may be now be incorporated into the chewable veterinary composition and be used according to the present invention, have been known to skilled specialists for a long time but the conventional oral dosage forms are not satisfactory because they are not attractive for animals and show the disadvantages discussed above.

[0033] With the chewable veterinary composition of the present invention one can combat all kinds of parasites. External parasites, also called ecto-parasites, are understood to be parasites which normally live on the animal, i.e. on the animal's skin or in the fur. Enclosed are biting insects, such as mosquitoes, blowfly, fleas or lice, or members of the order *Acarina*, e.g. mites or ticks. Suitable products against external parasites include insecticides and acaricides. It does not matter what their mode of action actually is. They can be e.g. chitin synthesis inhibitors, growth regulators; juvenile hormones; adulticides. They can be broad-band insecticides, broad-band acaricides. The active ingredient can

be a killer or a deterrent or repellent. It can affect e.g. only adult stages or juvenile stages of the parasite or may affect any stage. The only prerequisite is that the active ingredient acts systemically. This means that it is not decomposed after oral uptake but transported by the body fluids to the skin or organ where the parasite uses to live.

[0034] If the active ingredient is an acaricide one can, for example, select a systemically acting acaricide from one of the following well-known classes of acaricides including: antibiotic acaricides such as abamectin, doramectin, eprinomectin, ivermectin, milbemectin, nikkomycins, selamectin, tetranactin, and thuringiensin; bridged diphenyl acaricides such as azobenzene, benzoximate, benzyl benzoate, bromopropylate, chlorbenside, chlorfenethol, chlorfenson, chlorfensulphide, chlorobenzilate, chloropropylate, dicofol, diphenyl sulfone, dofenapyn, fenson, fentrifanil, fluorbenside, proclonol, tetradifon, and tetrasul; carbamate acaricides such as benomyl, carbanolate, carbaryl, carbofuran, fenothiocarb, methiocarb, metolcarb, promacyl, and propoxur; oxime carbamate acaricides such as aldicarb, butocarboxim, oxamyl, thio-carboxime, and thiofanox; dinitrophenol acaricides such as binapacryl, dinex, dinobuton, dinocap, dinocap-4, dinocap-6, dinoceton, dinopenton, dinosulfon, dinoterbon, and DNOC; formamidine acaricides such as amitraz, chlordimeform, chloromebuform, formetanate, and formparanate, mite growth regulators such as clofentezine, dofenapyn, fluazuron, flubenzimine, flucycloxuron, flufenoxuron, and hexythiazox; organochlorine acaricides such as bromocyclen, cam-phechlor, dienochlor, and endosulfan; organotin acaricides such as azocyclotin, cyhexatin, and fenbutatin oxide; pyrazole acaricides such as acetoprole, Fipronil and analogues and derivatives thereof, tebufenpyrad, and vaniliprole; pyrethroid acaricides including: pyrethroid ester acaricides like acrinathrin, bifenthrin, cyhalothrin, cypermethrin, alpha-cypermethrin, fenpropathrin, fenvalerate, flucythrinate, flumethrin, fluvalinate, tau-fluvalinate, and permethrin, and pyrethroid ether acaricides like halfenprox; quinoxaline acaricides such as chinomethionat and thioquinox; sulfite ester acaricides such as propargite; tetrionic acid acaricides such as spiroadiclofen; and form unclassified acaricides such acequinocyl, amid-oflumet, arsenous oxide, chloromethiuron, closantel, crotamiton, diafenthion, dichlofluanid, disulfiram, fenazaflor, fenazaquin, fenpyroximate, fluacrypyrim, fluenetil, mesulfen, MNAF, nifluridide, pyridaben, pyrimidifen, sulfiram, sulfluramid, sulfur and triarathene.

[0035] Suitable insecticides acting either as adulticides or insect growth regulators (IGRs) can be chosen from a variety of well-known different chemical classes such as chlorinated hydrocarbons, organophosphates, carbamates, pyrethroids, formamidines, borates, phenylpyrazoles, and macrocyclic lactones (previously known as avermectins). Prominent representatives of adulticides/insect killers are imidacloprid, fenthion, fipronil, allethrin, resmethrin, fenvalerate, permethrin, malathion and derivatives thereof. Insect adulticides kill the insect in almost any development stage either by contact or as a stomach poison. Widely used representatives of insect growth regulators (IGRs) are, for example benzoylphenylureas such as diflubenzuron, lufenuron, noviflumuron, hexaflumuron, triflumuron, and teflubenzuron or substances like fenoxycarb, pyriproxifen, methoprene, kinoprene, hydroprene, cyromazine, buprofezin, pymetrozine and derivatives thereof. Insect growth inhibitors or insect growth regulators (any of which is commonly known as an IGR) are products or materials that interrupt or inhibit the life cycle of a pest.

[0036] It goes without saying that the highly palatable ductile chewable veterinary composition according to the present invention is also very suitable for administering active ingredients that combat internal parasites (endo-parasites) such as worms living in the blood or in organs of the animal. Thus, the active ingredient can be an anthelmintic (dewormer).

[0037] Anthelmintics (dewormers) are a heterogeneous group of drugs but they are selectively toxic to worms. The drugs can achieve this by either inhibiting the metabolic process vital to the parasite, or by causing the parasite to be exposed to higher concentration of drug than are the hosts cells, which means that one makes use of the existence of an advantageous therapeutic window. Anthelmintics can affect the target parasite during treatment by interfering with the integrity of parasite cells, inhibiting neuromuscular transmission and coordination, or mechanisms which protect against host immunity, that ultimately lead to the starvation, neuromuscular paralysis, death and expulsion of the parasite. Anthelmintics are commonly administered by drench, paste, orally, or by injection. The drugs are absorbed into the blood stream and widely diffused. They are metabolized in the liver and excreted in feces and urine. In the animal health field anthelmintics are used widely used against roundworms, lungworms, tapeworms, intestinal worms, whipworms, hookworms, pinworms, trichinella (trichinosis), and other less common organisms, liver flukes and other less common organisms in a broad range of animals such as beef, cattle, swine, goats, horses, and pets like cats and dogs. The activity spectrum of anthelmintics for dogs and cats embraces Trematodes such as *Alaria alata* and *Opisthorchis tenuicollis*; Cestodes such as *Taenia hydatigena*, *Taenia pisiformis*, *Taenia ovis*, *Hydatigena Taenia taeniaeformis*, *Echinococcus granulosus*, *Echinococcus multilocularis*, *Dipylidium caninum*, *Diphyliobothrium latum*, *Multiceps multiceps*, *Multiceps serialis*, *Mesocestoides lineatus*, and *Mesocestoides corti*; and Nematodes such as *Ancylostoma caninum*, *Uncinaria stenocephala*, *Toxocara canis*, *Toxocara cati*, *Toxascaris leonina*, *Strongyloides stercoralis*, *Filaroides osieri*, *Capillaria aerophila*, *Capillaria plica*, *Capillaria hepatica*, *Trichinella spiralis*, *Angiostrongylus vasorum*, *Trichuris vulpis*, *Spirocerca lupi*, *Dirofilaria immitis*, *Ancylostoma tubaeforme*, and *Aelurostrongylus abstrusus*.

[0038] Internal parasites within the present invention include all species of worm infestation (helminthes) but also bacteria and viruses causing bacterial and viral infections, in particular those that infest the organs or parts of the body, such as the lungs, heart, alimentary tract or extremities, or which spread through the whole organism.


[0039] The anthelmintic can be selected from endo-parasiticides and endectocides including one of the following well-

known groups of dewormers such as macrocyclic lactones (sometimes called simply macrolides), benzimidazoles, pro-benzimidazoles, imidazothiazoles, tetrahydropyrimidines, organophosphates and piperazines.


[0040] A most preferred group of anthelmintics consists of the more modern natural or chemically modified macrocyclic lactones (macrolides), such as avermectins, milbemycins and derivatives thereof, including prominent representatives such as Ivermectin, Doramectin, Moxidectin, Selamectin, Emamectin, Eprinomectin, Milbemectin, Abamectin, Milbemycin oxime, Nemadectin, and a derivative thereof, in free form or in the form of a physiologically acceptable salt.

[0041] The macrocyclic lactones are most preferred because they exhibit a broad spectrum of activity. Most of them exhibit ecto and in parallel endo-parasitocidal activity. Therefore, they are also called endectocides. Macrocyclic lactones bind to glutamated chlorine channel causing in the first instance paralysis and later on the death of the parasite.

[0042] In the context of the invention, a preferred group of macrocyclic lactones is represented by compounds of formula (I)




wherein X is -C(H)(OH)-; -C(O)-; or -C(=N-OH)-; Y is -C(H₂)-; =C(H)-; -C(H)(OH)-; or -C(=N-OCH₃)-; R₁ is hydrogen or one of radicals



R₄ is hydroxyl, -NH-CH₃ or -NH-OCH₃; R₂ is hydrogen, -CH₃, -C₂H₅, -CH(CH₃)-CH₃, -CH(CH₃)-C₂H₅, -C(CH₃)=CH-CH(CH₃)₂ or cyclohexyl; and if the bond between atoms 22 and 23 represents a double bond the carbon atom in 23-position is unsubstituted so that Y is =C(H)-, or if is the bond between atoms 22 and 23 is a single bond the carbon atom in 23-position is unsubstituted or substituted by hydroxy or by the group =N-O-CH₃ so that Y is -C(H₂)-; -C(H)(OH)-; or -C(=N-OCH₃)-; in free form or in the form of a physiologically acceptable salt.


[0043] Typical and especially preferred representatives of compounds of formula (I) are:

- 1) Ivermectin is 22,23-Dihydroabamectin; 22,23-dihydroavermectin B 1; or 22,23-dihydro C-076B 1, wherein X is -C(H)(OH)-; Y is -C(H₂)-; R₁ is the radical



R_2 is either $-\text{CH}(\text{CH}_3)-\text{CH}_3$ or $-\text{CH}(\text{CH}_3)-\text{C}_2\text{H}_5$ and the bond between atoms 22 and 23 represents a single bond. Ivermectin is known from US-4,199,569.


2) Doramectin is 25-Cyclohexyl-5- O- demethyl-25-de(1-methylpropyl) avermectin A 1a, wherein X is $-\text{C}(\text{H})(\text{OH})-$; Y is $=\text{C}(\text{H})-$; R_1 is the radical



R_2 is cyclohexyl and the bond between atoms 22 and 23 represents a double bond. Doramectin is known from US-5,089,480.


3) Moxidectin, is [6R, 23E, 25S (E)]-5-O- Demethyl-28-deoxy-25-(1,3-dimethyl-1-butenyl)-6,28-epoxy-23-(methoxyimino) milbemycin B, wherein X is $-\text{C}(\text{H})(\text{OH})-$; Y is $-\text{C}(=\text{N}-\text{OCH}_3)-$; R_1 is hydrogen; R_2 is $-\text{C}(\text{CH}_3)=\text{CH}-\text{CH}(\text{CH}_3)_2$; and the bond between atoms 22 and 23 represents a single bond. Moxidectin, is known from EP-0,237,339 and US-4,916,154.

4) Selamectin is 25-cyclohexyl-25-de(1-methylpropyl)-5-deoxy-22,23-dihydro-5-(hydroxyimino) avermectin B1 monosaccharide and thus a compound of formula (i), wherein X is $-\text{C}(=\text{N}-\text{OH})-$; Y is $-\text{C}(\text{H}_2)-$; R_1 is the radical




R_2 is cyclohexyl; and the bond between atoms 22 and 23 represents a single bond. Selamectin is known e.g. from: ECTOPARASITE ACTIVITY OF SELAMECTIN; A novel endectocide for dogs and cats. A Pfizer Symposium, held in conjunction with The 17th international Conference of the World Association for the Advancement of Veterinary Parasitology, 19 August 1999. Copenhagen, Denmark.

5) Emamectin is (4primeprime R)-5- O -demethyl-4primeprimedeoxy-4primeprime-(methlamino) avermectin A 1a and (4primeprime R)-5- O -demethyl-25-de(1-methylpropyl)-4primeprime-deoxy-4primeprime-(methylamino)-25-(1-methylethyl) avermectin A 1a (9:1), wherein X is $-\text{C}(\text{H})(\text{OH})-$; Y is $=\text{C}(\text{H})-$; R_1 is -



R_2 is $-\text{CH}(\text{CH}_3)-\text{CH}_3$, or $-\text{CH}(\text{CH}_3)-\text{C}_2\text{H}_5$, and the bond between atoms 22 and 23 represents a double bond. Emamectin is known from US-4,874,749.


6) Eprinomectin is (4primeprime R)-4primeprime- epi- (acetyl amino)-4primeprime-deoxy avermectin B 1, wherein X is $-\text{C}(\text{H})(\text{OH})-$; Y is $=\text{C}(\text{H})-$; R_1 is the radical



R_2 is $-\text{CH}(\text{CH}_3)-\text{CH}_3$, or $-\text{CH}(\text{CH}_3)-\text{C}_2\text{H}_5$, and the bond between atoms 22 and 23 represents a double bond. Eprinomectin is known from US-4,427,663.

7) Milbemectin is (6R, 25R)-5-O-demethyl-28-deoxy-6,28-epoxy-25-methylmilbemycin, wherein X is $-\text{C}(\text{H})(\text{OH})-$; Y is $-\text{C}(\text{H}_2)-$; R_1 is hydrogen; R_2 is $-\text{CH}_3$, or $-\text{C}_2\text{H}_5$; and the bond between atoms 22 and 23 represents a single bond. Milbemectin is known from US-3,950,360.


8) Abamectin is Avermectin B 1 which is also named 5- O- demethyl avermectin A 1 a and 5-O- demethyl-25-de(1-methylpropyl)-25-(1-methylethyl) avermectin A 1 a (4:1), wherein X is $-\text{C}(\text{H})(\text{OH})-$; Y is $=\text{C}(\text{H})-$; R_1 is the radical



R_2 is $-\text{CH}(\text{CH}_3)-\text{CH}_3$, or $-\text{CH}(\text{CH}_3)-\text{C}_2\text{H}_5$; and the bond between atoms 22 and 23 represents a double bond. Abamectin is known from US-4,310,519.

9) Milbemycin oxim is milbemycin A 4 5-oxime; milbemycin A 3 5-oxime, wherein X is $-\text{C}(\text{H})(\text{OH})-$; Y is $-\text{C}(\text{H}_2)-$; R_1 is hydrogen; R_2 is $-\text{CH}(\text{CH}_3)-\text{CH}_3$, or $-\text{CH}(\text{CH}_3)-\text{C}_2\text{H}_5$, and the bond between atoms 22 and 23 represents a single bond. Milbemycin oxim is known from US-4, 547, 520.

10) The compound of the formula (I) wherein X is $-\text{C}(\text{H})(\text{OH})-$; Y is $-\text{C}(\text{H}_2)-$; R_1 is the radical



R_2 is $-\text{CH}_3$ or C_2H_5 , and the bond between atoms 22 and 23 represents a single bond. This compound is known from WO 01/83500.

11) Nemadectin is antibiotic S-541 A; also named [6 R, 23 S, 25 S,(E)]-5-O- Demethyl-28-deoxy-25-(1,3-dimethyl-1-butenyl)-6,28-epoxy-23-hydroxymilbemycin B; wherein X is $=\text{CH}-\text{OH}$; Y is $-\text{C}(\text{H}_2)-$; R_1 is hydrogen; R_2 is $-\text{C}(\text{CH}_3)=\text{CH}-\text{CH}(\text{CH}_3)_2$, and the bond between atoms 22 and 23 represents a single bond. Nemadectin is known from US-4,869,901.

[0044] The compounds specifically mentioned under items 1-11 hereinbefore, are preferred embodiments of the present invention and can be used either alone or in combination with another endo-parasiticide, ecto-parasiticide or

endecticide.

[0045] Benzimidazoles, benzimidazole carbamate and pro-benzimidazoles interfere with energy metabolism by inhibition of polymerization of microtubules and include very potent compounds such as thiabendazole, mebendazole, fenbendazole, oxfendazole, oxibendazole, albendazole, luxabendazole, netobimin, parbendazole, flubendazole, cyclobendazole, febantel, thiophanate and derivatives thereof.

[0046] Imidazothiazoles are cholinergic agonists and include highly active compounds such as tetramisole, levamisole, and derivatives thereof.

Tetrahydropyrimidines act are also cholinergic agonists and include highly active compounds such as morantel, pyrantel, and derivatives thereof.

Organophosphates are inhibitors of cholinesterase. This class includes potent compounds such as dichlorvos, haloxon, trichlorfon, and derivatives thereof.

Piperazines exhibit anticholinergic action and block neuromuscular transmission. This class includes highly active compounds such as piperazine and derivatives thereof. Salicylanilide selected from closantel, tribromsalan, dibromsalan, oxychlozanide, clioxanide, rafoxanide, brotianide, bromoxanide and derivatives thereof.

[0047] Within the present invention the anthelmintic (dewormer) a preferred embodiment consist of a combination of a macrocyclic lactone and an anthelmintic selected from the group consisting of Albendazole, Clorsulon, Cydectin, Diethylcarbamazine, Febantel, Fenbendazole, Haloxon, Levamisole, Mebendazole, Morantel, Oxyclozanide, Oxibendazole, Oxfendazole, Oxfendazole, Oxamniquine, Pyrantel, Piperazine, Praziquantel, Thiabendazole, Tetramisole, Trichlorfon, Thiabendazole, and derivatives thereof. Most preferred is Praziquantel. In order to broaden the activity spectrum towards ecto-parasites said anthelmintic combination can contain in addition to the dewormers a parasitocidally effective amount of an insecticide, acaricide or an insecticide and an acaricide. Of course one could also add an antibiotic for treating bacterial disease.

[0048] All of the suitable parasitocides mentioned hereinbefore are known. Most of them are described in THE MERCK INDEX 1999 by Merck & Co Inc, Whitehouse Station, NJ, USA; published on CD-ROM by Chapman & Hill/CRC, 1999, Hampden Data Service Ltd. and in the literature specifically mentioned in THE MERCK INDEX 1999.

[0049] Suitable antimicrobial active ingredients are, e.g. various penicillins, tetracyclines, sulfonamides, cephalosporins, cephamycins, aminoglycosids, trimethoprim, dimetridazoles, erythromycin, framycetin, fruazolidone, various pleuromutilins such as thiamulin, valnemulin, various macrolides, streptomycin and substances acting against protozoa, e.g. clopidol, salinomycin, monensin, halofuginone, narasin, robenidine, etc.

[0050] Behavioral disorders include e.g. separation worry or travel sickness of dogs and cats. A suitable compound acting against behavioral disorders is e.g. clomipramine.

[0051] The chewable combination according to the present invention may also contain an active ingredient for the treatment of disfunctions or hypo-activity.

[0052] Dysfunction or hypo-activity is understood to include functions like autoimmune disorders, which deviate from the norm, whether through inborn or acquired damage to individual organs or tissue. This complex also includes rheumatic diseases, pathological changes to joints, bones or internal organs, and much more. A prominent representative of compounds that can be used in this complex area is cyclosporine and derivatives thereof. The term "animal disease" even includes different types of cancer and metastasis progression in connective tissues that are common in animals. In this field bisphosphonates like coledronate, clodronate, etidronate, pamidronate and alendronate play an important role. Said bisphosphonates can also be administered in the treatment or prophylaxis of ulcers, rheumatoid arthritis and other arthritides, and periodontitis. Another suitable class of active ingredients encompasses anti-inflammatory agents such as benzenesulfonamides like Deracoxib, which is extremely suitable for the control of pain and inflammation associated with osteoarthritis. Further anti-inflammatory agents are diclofenac and derivatives thereof.

[0053] In the present invention, the administration problems depicted in connection with conventional oral dosage forms, like tablets and capsules, can be very easily resolved and chewable products can be prepared, which are taken orally by the animals without causing any problems. The animals actually take the chewable veterinary composition voluntarily.

Summary of the invention

[0054] The present invention overcomes the disadvantages and shortcomings of the prior art by providing an easy-to-use, safe, powerful, and stable veterinary formulation consisting of a highly palatable ductile chewable veterinary composition which is produced by an extrusion process wherein the product is extruded at or near room temperature, and where the extruder is cooled down below room temperature, preferably to 5-10°C. The palatable ductile chewable composition constitutes a veterinary composition and is administered orally. The composition is capable of killing endoparasites and ecto-parasites and/or can be used for treating prophylactic or curative animal diseases, and it is useful for the treatment of any warm-blooded non-human animal, including herd animals, like horses, cattle, sheep or poultry and preferably pets like dogs and cats.

[0055] The highly palatable ductile chewable veterinary composition of the invention is composed of an organic composition which contains an effective amount of one or more active ingredients, preferably an effective amount of a mono, binary or ternary mixture of organic compounds capable of controlling ecto-parasites, endo-parasites or bacterial or viral pathogens or a combination of ecto-parasites, endo-parasites, bacterial or viral pathogens. Depending on the mode of action the highly palatable ductile chewable veterinary composition of the invention contains a parasitically or anti-pathogenically effective amount of one or more active ingredients. The expression "parasitically effective amount" refers to that amount of active ingredient in the composition which will fully control the target parasite which means that 95-100%, preferably 98-100% or close to 100% of the parasites are killed and the active ingredient is nevertheless well tolerated. The expression "anti-pathogenically effective amount" refers to that amount of active ingredient in the composition which will efficiently cure a bacterial, viral or behavioral disease or if administered prophylactically will suppress the outbreak of such a disease. The highly palatable ductile chewable veterinary composition of the invention comprises an effective amount of one or more ingredients that are active against animal pests, pathogens or animal diseases; it further comprises meat flavoring and partially gelatinized starch, and it comprises of a softener; and of up to 9 % (w/w), preferably 3-7 % (w/w), most preferred 4-6 % (w/w), of water. It is essential that that during the extrusion process the extruder is cooled down below room temperature. A temperature range of 5-10°C is ideal.

[0056] Each of the following paragraphs defines a preferred embodiment of the present invention:

[0057] A highly palatable ductile chewable veterinary composition that comprises (A) an effective amount of one or more ingredients that are active against animal pests, pathogens or animal diseases; (B) meat flavoring; (C) partially gelatinized starch; (D) a softener; and (E) optionally up to 9 % (w/w) water.

[0058] A highly palatable ductile chewable veterinary composition as defined above capable of controlling endo-parasites and simultaneously ecto-parasites of non-human animals.

[0059] A highly palatable ductile chewable veterinary composition as defined above wherein the animal disease comprises bacterial infections, viral infections, behavioral disorders, inflammatory diseases, and auto-immune diseases.

[0060] A highly palatable ductile chewable veterinary composition as defined above comprising 20 to 30 % (w/w) of a natural meat flavoring.

[0061] A highly palatable ductile chewable veterinary composition as defined above wherein the natural meat flavoring comprises 20 to 55 % (w/w) fat.

[0062] A highly palatable ductile chewable veterinary composition as defined above comprising 25 to 70 % (w/w) of partially gelatinized starch.

[0063] A highly palatable ductile chewable veterinary composition as defined above wherein the partially gelatinized starch comprises 12 to 17 % (w/w) of gelatinized starch.

[0064] A highly palatable ductile chewable veterinary composition as defined above comprising 10 to 20 % (w/w), preferably about 11-15 % (w/w), of a softener, based upon the weight of the partially gelatinized starch.

[0065] A highly palatable ductile chewable veterinary composition as defined above wherein the softener is selected from the group consisting of glycerol, polyethylene glycol and polypropylene glycol.

[0066] A highly palatable ductile chewable veterinary composition as defined above comprising up to 9 % (w/w), preferably 3 to 7 % (w/w), more preferably 4 to 6% (w/w) of water.

[0067] A highly palatable ductile chewable veterinary composition as defined above comprising 1 to 10 % (w/w), preferably 3 to 7 % (w/w) of a sweetener.

[0068] A highly palatable ductile chewable veterinary composition as defined above comprising 0 to 3.5 % (w/w), preferably 0.01 to 0.5 % (w/w) of an antioxidant.

[0069] A highly palatable ductile chewable veterinary composition as defined above comprising 0 to 5 % (w/w), preferably 0.05 to 2 % (w/w) of a coloring agent.

[0070] A highly palatable ductile chewable veterinary composition as defined above comprising 0 to 4% (w/w) of sodium chloride.


[0071] A highly palatable ductile chewable veterinary composition as defined above comprising an parasitically effective amount of an ecto-parasiticide, an endo-parasiticide, an endectocide or of a combination of a parasiticide selected from the group consisting of an ecto-parasiticide, an endo-parasiticide and an endectocide.

[0072] A highly palatable ductile chewable veterinary composition as defined above wherein the ecto-parasiticide is active against insects, members of the order Acarina or insects and members of the order Acarina.


[0073] A highly palatable ductile chewable veterinary composition as defined above wherein the ecto-parasiticide is an insecticide which is either an insect adulticides or insect growth regulators.

[0074] A highly palatable ductile chewable veterinary composition as defined above comprising an parasitically effective amount of an endo-parasiticide or endectocide selected from the group consisting of macrocyclic lactones, benzimidazoles, pro-benzimidazoles, imidazothiazoles, tetrahydropyrimidines, organophosphates and piperazines.

[0075] A highly palatable ductile chewable veterinary composition as defined above comprising an effective amount of a natural or chemically modified macrocyclic lactone of formula (I)




wherein X is -C(H)(OH)-; -C(O)-; or -C(=N-OH)-; Y is -C(H₂)-; =C(H)-; -C(H)(OH)-; or -C(=N-OCH₃)-; R₁ is hydrogen or one of radicals



R₄ is hydroxyl, -NH-CH₃ or -NH-OCH₃; R₂ is hydrogen, -CH₃, -C₂H₅, -CH(CH₃)-CH₃, -CH(CH₃)-C₂H₅, -C(CH₃)=CH-CH(CH₃)₂ or cyclohexyl; and if the bond between atoms 22 and 23 represents a double bond the carbon atom in 23-position is unsubstituted so that Y is =C(H)-, or if the bond between atoms 22 and 23 is a single bond the carbon atom in 23-position is unsubstituted or substituted by hydroxy or by the group =N-O-CH₃ so that Y is -C(H₂)-; -C(H)(OH)-; or -C(=N-OCH₃)-; in free form or in the form of a physiologically acceptable salt.

[0076] A highly palatable ductile chewable veterinary composition as defined above comprising an effective amount of a natural or chemically modified macrocyclic lactone of formula (I) wherein X is -C(H)(OH)-; Y is -C(H₂)-; R₁ is the radical



R₂ is -CH₃ or C₂H₅, and the bond between atoms 22 and 23 represents a single bond.

[0077] A highly palatable ductile chewable veterinary composition as defined above wherein the animal pests are external animal parasites or internal animal parasites or both.

[0078] A highly palatable ductile chewable veterinary composition as defined above wherein the macrocyclic lactone is selected from the group consisting of avermectins, milbemycins and derivatives thereof, in free form or in the form of a physiologically acceptable salt.

[0079] A highly palatable ductile chewable veterinary composition as defined above wherein the macrocyclic lactone is selected from the group consisting of Ivermectin, Doramectin, Moxidectin, Selamectin, Emamectin, Eprinomectin, Milbemectin, Abamectin, Milbemycin oxime, Nemadectin, and a derivative thereof, in free form or in the form of a physiologically acceptable salt.

[0080] A highly palatable ductile chewable veterinary composition as defined above comprising an effective amount of a macrocyclic lactone in combination with an effective amount of an anthelmintic selected from the group consisting of Albendazole, Clorsulon, Cydectin, Diethylcarbamazine, Febantel, Fenbendazole, Haloxon, Levamisole, Mebendazole,

Morantel, Oxyclozanide, Oxibendazole, Oxfendazole, Oxfendazole, Oxamniquine, Pyrantel, Piperazine, Praziquantel, Thiabendazole, Tetramisole, Trichlorfon, Thiabendazole, and a derivative thereof.

[0081] A highly palatable ductile chewable veterinary composition as defined above comprising additionally an effective amount of an insecticide, acaricide or an insecticide and an acaricide.

[0082] A highly palatable ductile chewable veterinary composition as defined above comprising an effective amount of milbemycin oxime and praziquantel.

[0083] A highly palatable ductile chewable veterinary composition as defined above comprising an effective amount of lufenuron, praziquantel and milbemycin oxime.

[0084] A highly palatable ductile chewable veterinary composition as defined above comprising an effective amount of cyclosporin.

[0085] A highly palatable ductile chewable veterinary composition as defined above comprising an effective amount of an antimicrobial selected from the group consisting of a penicillin, tetracycline, sulfonamide, cephalosporin, cephamycin, aminoglucosid, trimethoprim, dimetridazole, erythromycin, framycetin, fruazolidone, pleuromutilin, streptomycin and a compound that is active against protozoa.

[0086] A highly palatable ductile chewable veterinary composition as defined above comprising an effective amount of compound that is active against behavioral including separation worry or travel sickness of dogs and cats.

[0087] A highly palatable ductile chewable veterinary composition as defined above wherein the active ingredient or a different chemical class is an insecticide or acaricide.

[0088] A highly palatable ductile chewable veterinary composition as defined above wherein the insecticide is selected from the group consisting of insect killers and insect growth regulators.

[0089] Yet another preferred embodiment of the present invention is a process for the production of a highly palatable ductile chewable veterinary composition as defined above, comprising (i) feeding the hopper of an extruder with an effective amount of one or more ingredients that are active against animal pests, pathogens or animal diseases; meat flavoring; partially gelatinized starch; a softener; and up to 9% (w/w) of water, (ii) cooling constantly down the mixture of active ingredients and carriers so that the temperature of the extrudate in the extruder does during the whole extrusion process at no time exceed 40°C, (iii) pressing the extrudate through a die that is decisive for the shape of the chewable product, and (iv) cutting the extrudate that leaves the extruder into equal pieces.

[0090] Yet another preferred embodiment of the present invention is a process as defined above wherein the hopper of the extruder is fed continuously and simultaneously with pre-mixture (1) and pre-mixture (2), wherein pre-mixture (1) consist of a homogenized mixture of one or more active ingredients and partially gelatinized starch, and pre-mixture (2) consists of a homogenized mixture of meat flavoring, a softener and optionally of a carrier selected from the group consisting of a sweetener, softener, an antioxidant, a coloring agent and sodium chloride.

[0091] Yet another preferred embodiment of the present invention is a process as defined above wherein the extruder is cooled down below room temperature.

[0092] Another preferred embodiment of the present invention is the use of (A) an effective amount of one or more ingredients that are active against animal pests, pathogens or animal diseases; (B) meat flavoring; (C) partially gelatinized starch; (D) a softener; (E) up to 9% water; and an active ingredient suitable for combating animal pests, pathogens or animal diseases for the preparation of a highly palatable ductile chewable veterinary composition.

[0093] Another preferred embodiment of the present invention is the use as defined above, comprising 20 to 30 % (w/w) of a natural meat flavoring.

[0094] Another preferred embodiment of the present invention is the use as defined above, wherein the natural meat flavoring comprises 20 to 55 % (w/w) fat.

[0095] Another preferred embodiment of the present invention is the use as defined above, comprising 25 to 70 % (w/w) of partially gelatinized starch.

[0096] Another preferred embodiment of the present invention is the use as defined above, wherein the partially gelatinized starch comprises 12 to 17 % (w/w) of gelatinized starch.

[0097] Another preferred embodiment of the present invention is the use as defined above, comprising 10 to 20 % (w/w) of a softener, based upon the weight of the partially gelatinized starch.

[0098] Another preferred embodiment of the present invention is the use as defined above, wherein the softener is selected from the group consisting of glycerol, polyethylene glycol and polypropylene glycol.

[0099] Another preferred embodiment of the present invention is the use as defined above, comprising 3 to 7 % (w/w) of water.

[0100] Another preferred embodiment of the present invention is the use as defined above, wherein the animal pests are external animal parasites or internal animal parasites or both.

[0101] Another preferred embodiment of the present invention is the use as defined above, comprising 1 to 10 % (w/w) of a sweetener.

[0102] Another preferred embodiment of the present invention is the use as defined above, comprising 0 to 3.5 % (w/w) of an antioxidant.

[0103] Another preferred embodiment of the present invention is the use as defined above, comprising 0 to 5 % (w/w) of a coloring agent.

[0104] Another preferred embodiment of the present invention is the use as defined above, comprising 0 to 4 % (w/w) of sodium chloride.


[0105] Another preferred embodiment of the present invention is the use as defined above, comprising an parasitically effective amount of an ecto-parasiticide, an endo-parasiticide, an endectocide or of a combination of a parasiticide selected from the group consisting of an ecto-parasiticide, an endo-parasiticide and an endectocide.

[0106] Another preferred embodiment of the present invention is the use as defined above, wherein the ecto-parasiticide is active against insects, members of the order Acarina or insects and members of the order Acarina.


[0107] Another preferred embodiment of the present invention is the use as defined above, wherein the ecto-parasiticide is an insecticide which is either an insect adulticides or insect growth regulators.

[0108] Another preferred embodiment of the present invention is the use as defined above, comprising an parasitically effective amount of an endo-parasiticide or endectocide selected from the group consisting of macrocyclic lactones, benzimidazoles, pro-benzimidazoles, imidazothiazoles, tetrahydropyrimidines, organophosphates and piperazines.

[0109] Another preferred embodiment of the present invention is the use as defined above, comprising an effective amount of a natural or chemically modified macrocyclic lactone of formula (I)




wherein X is -C(H)(OH)-; -C(O)-; or -C(=N-OH)-; Y is -C(H₂)-; =C(H)-; -C(H)(OH)-; or -C(=N-OCH₃)-; R₁ is hydrogen or one of radicals



R₄ is hydroxyl, -NH-CH₃ or -NH-OCH₃; R₂ is hydrogen, -CH₃, -C₂H₅, -CH(CH₃)-CH₃, -CH(CH₃)-C₂H₅, -C(CH₃)=CH-CH(CH₃)₂ or cyclohexyl; and if the bond between atoms 22 and 23 represents a double bond the carbon atom in 23-position is unsubstituted so that Y is =C(H)-, or if the bond between atoms 22 and 23 is a single bond the carbon atom in 23-position is unsubstituted or substituted by hydroxy or by the group =N-O-CH₃ so that Y is -C(H₂)-; -C(H)(OH)-; or -C(=N-OCH₃)-; in free form or in the form of a physiologically acceptable salt.

[0110] Another preferred embodiment of the present invention is the use as defined above, wherein the macrocyclic lactone is a compound of the formula (I) wherein X is -C(H)(OH)-; Y is -C(H₂)-; R₁ is the radical



R_2 is $-CH_3$ or C_2H_5 , and the bond between atoms 22 and 23 represents a single bond.

10 **[0111]** Another preferred embodiment of the present invention is the use as defined above, wherein the endecticide is a macrocyclic lactone is selected from the group consisting of avermectins, milbemycins and derivatives thereof, in free form or in the form of a physiologically acceptable salt.

15 **[0112]** Another preferred embodiment of the present invention is the use as defined above, wherein the macrocyclic lactone is selected from the group consisting of Ivermectin, Doramectin, Moxidectin, Selamectin, Emamectin, Eprinomectin, Milbemectin, Abamectin, Milbemycin oxime, Nemadectin, and a derivative thereof, in free form or in the form of a physiologically acceptable salt.

20 **[0113]** Another preferred embodiment of the present invention is the use as defined above, comprising an effective amount of a macrocyclic lactone in combination with an effective amount of an anthelmintic selected from the group consisting of Albendazole, Clorsulon, Cydectin, Diethylcarbamazine, Febantel, Fenbendazole, Haloxon, Levamisole, Mebendazole, Morantel, Oxyclozanide, Oxibendazole, Oxfendazole, Oxfendazole, Oxamniquine, Pyrantel, Piperazine, Praziquantel, Thiabendazole, Tetramisole, Trichlorfon, Thiabendazole, and a derivative thereof.

[0114] Another preferred embodiment of the present invention is the use as defined above, comprising in addition to an endo-parasiticide or an endecticide an effective amount of an insecticide, acaricide or an insecticide and an acaricide.

[0115] Another preferred embodiment of the present invention is the use as defined above, comprising an effective amount of milbemycin oxime and praziquantel.

25 **[0116]** Another preferred embodiment of the present invention is the use as defined above, comprising an effective amount of lufenuron, praziquantel and milbemycin oxime.

[0117] Another preferred embodiment of the present invention is the use as defined above, comprising an effective amount of cyclosporin.

30 **[0118]** Another preferred embodiment of the present invention is the use as defined above, comprising an effective amount of an antimicrobial selected from the group consisting of a penicillin, tetracycline, sulfonamide, cephalosporin, cephamycin, aminoglycosid, trimethoprim, dimetridazole, erythromycin, framycetin, fruazolidone, pleuromutilin, streptomycin and a compound that is active against protozoa.

[0119] Another preferred embodiment of the present invention is the use as defined above, comprising an effective amount of compound that is active against behavioral including separation worry or travel sickness of dogs and cats.

35 Detailed Description of the Invention

40 **[0120]** Even so meat flavoring is actually not the main component of the highly palatable ductile chewable veterinary composition it plays the major role for the present invention. It has surprisingly been recognized that the desired high palatability that is necessary for achieving reliable and well-reproducible results strictly depends on the amount of meat flavoring in the final composition. The meat flavoring is either a natural product consisting of dried powdered meat derived, for example, from domestic animals and productive livestock, e.g. pigs, horses, cattle, sheep, goats and poultry including chicken, duck, goose and turkey. It further surprisingly turned out that the natural content of fat in said natural meat powder of 20-55% (w/w) is very important not only for achieving the desired high palatability and excellent taste for the animal but also for achieving the desired softness of the final chewable product.

45 In context with the present invention, a "meat flavoring" shall refer to natural dried and powdered meat as well as to artificial meat flavorings, which are well-known from the food industry. It has however been recognized that artificial meat flavorings are only suitable for the present invention if they already contain 20-55% (w/w) fat or if this amount of fat is added to the artificial flavoring. Fat that can be added to artificial meat flavorings can be chosen either from animal fats or preferably from plant fats including vegetable oils. However, if vegetable oils are used it is advantageous to use hardened/saturated oils. Unsaturated oils are usually liquid at room temperature and result in products that do not show the desired ductility/softness. They are usually too soft. Preferred is the use of saturated/hardened oils/fats that are generally solid at room temperature and lead to chewable compositions showing the desired ductility.

55 Fats and oils contain many different fatty acids which affect the body in varying ways. Most simply, they are classified as saturated or unsaturated. Saturated fats sometimes are also called hardened fats. It is the saturated fat found in many animal products. Saturated fats are generally solid at room temperature. They are mainly of animal origin but can also be isolated from plants. Typical examples stemming from plants are cocoa butter and coconut and palm oils. These products are often used in store-bought baked goods, non-dairy whipped toppings, cream substitutes, most peanut

butter and some margarines. Typical sources of saturated fat are: Animal Fat; Coconut Oil; Meat Fat; Bacon Fat; Cream; Palm Kernel Oil; Beef Fat; Palm Oil; Butter; Ham Fat; Pork Fat; Chicken Fat and Skin; Hardened Fat or Oil; Turkey Fat and Skin; Hydrogenated Vegetable Oil; Cocoa Butter; Lamb Fat; and Coconut.

Natural and artificial meat flavoring is commercially obtainable from various producers. Natural meat flavoring is, for example, obtainable from:

IDF (International Dehydrated Food) INTERNATIONAL DEHYDRATED FOODS, INC.

P.O. Box 10347 Springfield, Missouri 65808, USA 800/641-6509 or 417/881-7820

ADF (American Dehydrated Food), American Dehydrated Foods, Inc., P.O. Box 4087 3801 East Sunshine, Springfield, Missouri 65809

IFF (International Flavour and Fragrance), IFF Global Headquarters, 521 West 57th Street, New York, NY 10019, United States

Proliant, Proliant Inc. - U.S. Office, 2325 North Loop Drive

Ames, IA 50010 USA

Some examples of sources of artificial meat flavoring are.

Kemin, Worldwide Headquarters • 2100 Maury Street, Box 70 • Des Moines, Iowa 50301-0070 USA.

McCormick, 226 Schilling Circle

Hunt Valley, MD 21031

Givaudan, Givaudan Flavors Corp.

(Flavours creation, sales & production) 1199 Edison Drive

Cincinnati, Ohio 45216

Haarman und Reimer

[0121] Within the present invention the expression "soft" is used to characterize a product that is not as hard and crunchy as, for example, a cornflake and on the other hand is not as ductile as, for example, a marshmallow. The desired ductility/hardness lies somewhere in between. If measured with a commercially available TEXTURE ANALYSER that is commercially available from Stable Micro Systems(TA-XT2 iHR/25), the texture (softness/hardness) of the chewables lies ideally between 6 - 12 N.

[0122] Hard and crunchy products are especially disadvantageous if one intends to treat older dogs and cats because most of these old animals suffer from periodontal disease (Pyorrhea). This disease involves the inflammation and degeneration of tissues that surround and support the teeth. These include the gingiva, alveolar bone, periodontal ligament, and cementum. Periodontitis or the loss of supporting bone is the latest stage of this progressive disorder and is the major cause of tooth loss in old dogs and cats. Animals suffering from periodontitis avoid eating hard and crunchy products because they cause them pain.

[0123] The second important feature of the present invention is the use of partially gelatinized starch. This starch contains 10-20% (w/w), preferably about 13-17% (w/w), most preferably 13-17% (w/w) pre-gelatinized starch. This is important as non-gelatinized and completely pre-gelatinized starch do not result in the desired ductility of the final product.

[0124] Starches exhibit thermal stability to about 121 °C. Starches are carbohydrates of a general formula $(C_6H_{10}O_5)_n$ and are derived from corn, wheat, oats, rice, potatoes, yucca and similar plants and vegetables. They consist of about 27% linear polymer (amylose) and about 73% branched polymer (amylopectin). The two polymers are intertwined within starch granules. Granules are insoluble in cold water, but soaking in hot water or under steam pressure ruptures their covering and the polymers hydrate into a colloidal suspension. This product is a pregelatinized starch and has been used in mugs for many years. Thus, pregelatinized starch is water-soluble starch that has undergone irreversible changes by heating in water or steam. Many suitable pregelatinized starches are commercially available. For example, Naster® Instant which is a pregelatinized Pea Starch with a high gel strength. Due to its high amylose level, it has some remarkable properties. It shows an excellent stability to high temperatures, shearing and to variations in pH and is ideal for use in cold processes.

[0125] A further important component is the softener, which keeps the moisture within the composition and allows to store the final product for weeks and months. It does not become hard and does not dry out.

[0126] If the mentioned meat flavoring, partially gelatinized starch and the basic component containing the active ingredient does not contain moisture one should add water during the extrusion process. This has an impact on the flexibility on the chewable veterinary composition of the present invention. It turned out that it is advantageous to adjust the moisture content of the product so that the final product contains water at a concentration equal to or lower than 9 % (w/w), preferably about 3 to 7 % (w/w), more preferably 4 to 6 % (w/w).

[0127] For the present invention not only the proportion of meat flavoring and partially pre-gelatinized starch is extremely important but also the production process as such has an influence on the final product. The highly palatable ductile

chewable veterinary composition of the present invention is the outcome of a special extrusion process. As such, extrusion is a very common thermoforming process widely used in the food industry for the production of customary feed pellets. However, in order to achieve the chewables according to the present invention, i.e. a highly palatable ductile product one has to modify the process and secure that the extrudate is not heated during the whole extrusion process because this leads to hard and crunchy products, to losses of active ingredient, and especially to a decrease of the palatability. Actually, production in the desired manner can easily be achieved. The highly palatable ductile chewable veterinary composition of the present invention is conveniently carried out in an injection molding machine or extruder. A mixture comprising an effective amount of one or more ingredients that are active against animal pests, pathogens or animal diseases; meat flavoring partially gelatinized starch; a softener; and up to 9% (w/w) of water is fed through the hopper onto a rotating, reciprocating screw. The material moves along the screw towards the tip. During this process, its temperature is cooled down constantly by means of external coolers around the outside of the barrel and by the shearing action of the screw. The cooling process is controlled so that the temperature of the extrudate during the whole extrusion process does not exceed peak temperature of 40°C. Starting in the feeding zone and continuing in the compression zone, the extrudate should not reach temperatures higher than said 40°C. It has been found that ideally the product is extruded at or near room temperature, and the extruder is cooled down below room temperature, preferably to 5-10°C. [0128] It is then conveyed through the metering zone, where homogenization occurs, to the end of the screw. The homogenized material at the tip is then pressed through a form-determining die to obtain shaped articles of the desired size. The simplest way is cutting the extrudate that leaves the extruder into equal pieces of the desired size. 'Desired' means that each piece contains the appropriate amount of active ingredient. Thus, for example, for big dogs one would produce bigger pieces than for young cats. The amount of active ingredient has to be adapted to the bodyweight of the animal that has to be treated.

[0129] This cooling during the extrusion process is extremely important because extruding this kind of a material without cooling can easily lead to temperatures inside the extruder in the range of 100-200°C. High extrusion temperatures however transform the matrix of the extrudate in an undesirable manner. The reason for this being that the starch is heated above the melting and glass transition temperatures of its components so that they undergo endothermic transitions. As a consequence a melting and disordering of the molecular structure of the starch granules takes place, so that an essentially destructurized starch is obtained. Instead of a ductile and rather soft product one obtains a hard or crunchy product which is not only refused by the animals but does not contain a reproducible amount of active ingredient. Many active ingredients are not even stable enough at these relatively high temperatures and are at least partially degraded. As a result the biological activity of the product is reduced, and undesired degradation products can be formed that can cause undesired side effects or lead to allergic reactions. Cooling the extruder to temperatures near room temperature, preferably to 5-10°C, suppresses all these undesired effects and leads to a perfect product that is not only highly palatable but can surprisingly be stored for months without being degraded.

[0130] The highly palatable ductile chewable veterinary composition of the present invention may also contain a sweetener for further improving the palatability. Any natural sugar can be used including confectioners sugar, maltitol, xylitol, sorbitol, mannitol, lactose, dextrose, saccharose, glucose or fructose, or any mixture thereof. In addition, artificial sweeteners known in the art, including saccharin, aspartame and Acesulfame-K, may also be used. The sweetener is preferably present in an amount of from 1 to 10 % (w/w), preferably between from about 3 to about 7 % (w/w) based on the sweetening power of sucrose. The sweetener serves as a palatability enhancer due to its organoleptic properties. Enhancement of the palatability can be especially achieved in those cases where the active ingredient is extremely bitter or exhibits a taste that is absolutely not accepted by the animal.

[0131] The veterinary composition of the present invention may also contain an antioxidant even so it turned out that in most of the cases this is not necessary. However, in certain cases the antioxidant serves as a preservative which increases the stability of ingredients that are not stable if exposed for a longer period of time to oxygen. The term "antioxidant" represents the three groups of antioxidants, true antioxidants, such as Tenox 2, Tenox PG, Tenox s-1, BHA (2-t-butyl-4-methoxyphenol), and BHT (2,6-di-t-butyl-4-methylphenol), sodium metabisulfite reducing agents and antioxidant synergists, such as tocopherols (alpha, beta, or delta-tocopherol, tocopherol esters, alpha-tocopherol acetate), alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, citric acid, edetic acid and its salts, lecithin and tartaric acid. Further suitable antioxidants are resveratrol, quercetin, benzoic acid, Trolox (N-acetylcysteine, 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid), dimethyl thiourea (DMTU), hesperetin, tetrahydrocurcumin, tetrahydrodemethoxycurcumin, and monothioglycerol. Said antioxidant being added in concentrations ranging from 0 to 3.5 % (w/w), preferably 0.01 to 0.5 % (w/w). Preferred antioxidant are Tenox 2 and BHA (2-t-butyl-4-methoxyphenol).

[0132] Sodium chloride may also be added, up to about 4% (w/w), to further improve the palatability of the product and to bind moisture. For certain animals sodium chloride serves like a palatability enhancer.

[0133] The highly palatable ductile chewable veterinary composition of the present invention further may contain a softener. The softener for use in the invention serves as a humectant which enhances the flexibility of the pet chew and retains moisture so that the texture of the pet chew is maintained at ambient temperatures. Typically, the softener is present in the highly palatable ductile chewable veterinary composition of the present invention at concentrations from

about 10-20 % (w/w) and preferably about 11-15% (w/w) based upon the weight of the partially gelatinized starch. Suitable softeners include alcohols such as sorbitol, mannitol, hexanol, pentanol and polyols (such as glycerine, propylene glycol, polyethylene glycol, and polypropylene glycol).

[0134] The highly palatable ductile chewable veterinary composition of the present invention may further contain a coloring agent that lead to a better-looking product. The coloring agent can be selected from the group of azo dyes, organic or inorganic pigments, or coloring agents of natural origin, preferably from oxides of iron or titanium. Said coloring agent being added in concentrations ranging from 0 to 5 % (w/w), preferably 0.05 to 2 % (w/w). A preferred coloring agent is ferric oxide, that is normally used in an amount around 0.1 % (w/w).

Technical Equipment

[0135] The extruder used for the production of the chewables according to the present invention is a co-rotating twin screw extruder type BCTG-62/28D with a screw diameter of 62 mm and a screw length of L/D ratio of 28D with a pelleting device from Bühler AG; Industriestrasse; CH-9240 Uzwil; Switzerland. The feeder (delivering the dry blend into the extruder) is obtainable from K-Tron, Switzerland. It is a loss in weight feeder: type K2-ML-T 35 twin screw feeder equipped with a AC (twin auger) screw. In addition two pumps are used to deliver glycerin and water separately. In addition, two chillers are used to maintain an extruder temperature below 10°C.

Measurement of the softness/hardness of the chewables

[0136] The texture (softness/hardness) of the chewy is measured by using a TEXTURE ANALYSER type: TA-XT2 iHR/25 that is commercially available from Stable Micro Systems Ltd. (Headquater: Stable Micro Systems Ltd., Vienna Court, Lammas Road; UK). One measures the peak force (in N) necessary to push a sphere with a velocity of 1 mm/sec to penetrate 2mm into the chewable. The sphere has a diameter of 8 mm. The texture of the chewables is measured to determine when the chewables are hard enough to be packaged into bins, before they are packaged into blisters. Texture after 24 hours: Typical values lay between 8 and 20 N.

Examples

Example 1: 2-Way-formulation containing Milbemycin oxime and Praziquantel

[0137]

Ingredients	Amount	Percentage [w/w]
Active ingredient No.1 Milbemycin oxime	1.975 kg	0.395%
Active ingredient No. 2 Praziquantel	19.000 kg	3.800%
Excipients		
Pre-gelatinized starch	205.025 kg	41.005%
Natural chicken flavor	150.000 kg	30.000%
Sugar	25.000 kg	5.000%
Sodium chloride powder	7.500 kg	1.500%
Ferric oxide	0.500 kg	0.100%
Total solids	409.000 kg	81.800%
Water	20.000 kg	4.000 %
Glycerin	70.000 kg	14.000%
Tenox 2	1.000 kg	0.200%
Total liquids	91.000 kg	18.200%
Total batch size	500.000 kg	100.000%

[0138] The 2-Way-formulation containing Milbemycin oxime and Praziquantel is produced as follows:

1. Pre-blend Milbemycine oxime (1.975 kg) and iron oxide (0.500 kg) with 13 kg of pre-gelatinized Starch using a V-blender for 5 min.
2. Vacuum transfer through a 10 mesh screen into a bin blender

EP 1 675 474 B9

3. Vacuum transfer praziquantel, confectionery sugar, sodium chloride, chicken flavor and the rest of pregelatinized starch through a 10 mesh screen into a bin blender and blend for 20 min
4. Weigh water into the water tank and weigh glycerin into the glycerin tank and mix with tenox.
5. Start extruder:
6. Cooling unit temperature is set at 5°C
7. Dry blend (mixture of step 3) is fed into BCTG extruder through K-tron feeding device.
8. Glycerin/Tenox 2 mixture is pumped into the extruder
9. Water is pumped into the extruder
10. Extruder velocity (rpm) is adjusted according feeding velocity of the dry blend
11. Cutting device of the extruder is adjusted to get the appropriate weights of the chewables
12. After extrusion the chewables are transported via conveyer to a hopping conveyer and finally
13. Filled into boxes of not more than 3 inches
14. Curing of the chewables for approx. 24 hours at ambient temperature and relative humidity < 60%.
15. After curing chewables are packaged into blister packages

[0139] Extruder settings for the different chewy sizes of 0.6 g, 1.5 g, 3.0 g and 6.0 g

Weights [g]	dry blend feed rate [kg/h]	glycerin addition [kg/h]	water addition [kg/h]	Extruder [rpm]	cutting speed
0.6	120	20.52	5.76	85	2200
1.5	140	23.94	6.72	100	2000
3.0	170	29.07	8.16	120	1250
6	200	34.2	9.6	140	700

Example 2: 3-Way-formulation containing Milbemycin oxime, Praziquantel and Lufenuron

[0140]

Ingredients	Amount	Percentage [w/w]
Active ingredient No. 1 Milbemycin oxime	1.975 kg	0.395%
Active ingredient No. 2 Praziquantel	19.000 kg	3.800%
Active ingredient No. 3 Lufenuron	38.340 kg	7.667%
Excipients		
Pre-gelatinized starch	159.690 kg	31.938%
Powdered Cooked Beef	142.000 kg	30.000%
Stock Aid Bacon Flavour	25.000 kg	5.000%
Sugar	25.000 kg	5.000%
Sodium chloride powder	7.500 kg	1.500%
Ferric oxide	0.500 kg	0.100%
Total solids	419.000 kg	83.800%
Waster	20.000 kg	4.000%
Glycerin	60.000 kg	12.000%
Tenox 2	1.000 kg	0.200%
Total liquids	81.000 kg	16.200%
Total batch size	500.000 kg	100.000%

[0141] The 3-Way-formulation containing Milbemycin oxime, Praziquantel and Lufenuron is produced along the same lines as described for the 2-Way-formulation in Example 1.

Example 3: 1-Way-formulation containing Cyclosporin

[0142]

EP 1 675 474 B9

	Ingredients	Amount	Percentage [w/w]
	Active ingredient		
5	Cyclosporin	43.40 g	8.7%
	Pre-gelatinized starch	183.60 g	36.7%
	Natural chicken flavor	150.00 g	30.0%
	Sugar	25.0 g	5.0 %
	Ferric oxide	0.50 g	0.1 %
10	Total solids	410.00 g	82.0%
	Water	20.00 g	4.0 %
	Glycerin	70.00 g	14.0 %
15	Total liquids	90.00 g	18.0%
	Total batch size	500.00 g	100.00%

[0143] The 1-Way-formulation containing Cyclosporin is produced along the same lines as described for the 2-Way-formulation in Example 1.

Example 4: Palatability (acceptance) test for different flavored chewables of a 3-way formulation with 100 dogs and 100 cats

[0144] 100 male and female dogs of different breeds and age are tested. The dogs are divided in 4 groups of 25 dogs of the same bodyweight. The testing person offers once a day to each dog one test-chewable which is adapted to the bodyweight of the dog. In a first instance the chewable is offered by hand for 60 seconds. If the dog does not take the formulation it is offered the dog in his empty bowl. The dog has again 60 seconds to take the formulation, if not, it is paced in his/her mouth. If the dog/cat spits it out it is reported as not accepted. In general not more than 5 to 6 different formulations are tested on consecutive days. Each formulation is packaged separately and labeled so that they can be clearly identified. An analogous test is carried out with 100 cats.

Chewable to be tested	Flavor	Total acceptance (dogs) [%]
2-Way-formulation containing Milbemycin oxime and Praziquantel	Natural bacon	96
3-Way-formulation containing Milbemycin oxime, Praziquantel and Lufenuron	Natural bacon	93
3-Way-formulation containing Milbemycin oxime, Praziquantel and Lufenuron	Natural beef	94
3-Way-formulation containing Milbemycin oxime, Praziquantel and Lufenuron	Natural chicken	95
3-Way-formulation containing Milbemycin oxime, Praziquantel and Lufenuron	Artificial bacon	94
3-Way-formulation containing Milbemycin oxime, Praziquantel and Lufenuron	Artificial beef	95

[0145] The analogues test in cats shows results in absolutely the same range.

Example 5: Stability tests

[0146] Samples of the highly palatable ductile chewable veterinary composition of the presnet invention are tested for stability under stressing conditions in order to simulate different temperatures and humidity conditions. The samples are tested at 25°C/60 rh, 30°C 60% rh and 40°C 75%rh. They are kept in incubators and are analyzed after 3, 6, 9 and 12 months with regard to active ingredient content. The analyze of the content of active ingredient of all samples tested at 25°C/60 rh and 30°C 60% rh for 12 months show no difference in comparison with identical samples kept in the refrigerator

at - 25°C for the same period of time. Chewables kept 12 months at 40°C/75%rh show also good stability results which indicate that they would lead to a shelf live for at least 12 months if stored under normal conditions, i.e. 25°C or 30°C and 40-70% rh. No significant difference is seen with regard to the stability of 2-way-formulations containing Milbemycin oxime and Praziquantel and 3-way-formulation containing Milbemycin oxime, Praziquantel and Lufenuron.

Claims

1. A highly palatable ductile chewable veterinary composition comprising (A) an effective -amount of one or more ingredients that are active against animal pests, pathogens or animal diseases; (B) meat flavoring; (C) partially gelatinized starch; (D) a softener; and (E) up to 9% water.
2. A chewable veterinary composition according to claim 1 wherein the animal disease comprises viral or bacterial infections, behavioral disorders, inflammatory diseases, and auto-immune diseases.
3. A chewable veterinary composition according to claim 1 comprising 20 to 30 % (w/w) of a natural meat flavoring.
4. A chewable veterinary composition according to claim 4 wherein the natural meat flavoring comprises 20 to 55 % (w/w) fat.
5. A chewable veterinary composition according to claim 1 comprising 25 to 70 % (w/w) of partially gelatinized starch.
6. A chewable veterinary composition according to claim 5 wherein the partially gelatinized starch comprises 12 to 17 % (w/w) of gelatinized starch.
7. A chewable veterinary composition according to claim 1 comprising 10 to 20 % (w/w) of a softener, based upon the weight of the partially gelatinized starch.
8. A chewable veterinary composition according to claim 7 wherein the softener is selected from the group consisting of glycerol, polyethylene glycol and polypropylene glycol.
9. A chewable veterinary composition according to claim 1 comprising 4 to 6 % (w/w) of water.
10. A chewable veterinary composition according to claim 1 wherein the animal pests are external animal parasites or internal animal parasites or both.
11. A chewable veterinary composition according to claim 1 comprising 1 to 10 % (w/w) of a sweetener.
12. A chewable veterinary composition according to claim 1 comprising 0 to 3.5 % (w/w) of an antioxidant.
13. A chewable veterinary composition according to claim 1 comprising 0 to 5 % (w/w) of a coloring agent.
14. A chewable veterinary composition according to claim 1 comprising 0 to 4% (w/w) of sodium chloride.
15. A chewable veterinary composition according to claim 1 comprising an parasitically effective amount of an ecto-parasiticide, an endo-parasiticide, an endectocide or of a combination of a parasiticide selected from the group consisting of an ecto-parasiticide, an endo-parasiticide and an endectocide.
16. A chewable veterinary composition according to claim 15 wherein the ecto-parasiticide is active against insects, members of the order Acarina or insects and members of the order Acarina.
17. A chewable veterinary composition according to claim 16 wherein the ecto-parasiticide is an insecticide which is either an insect adulticides or insect growth regulators.
18. A chewable veterinary composition according to claim 15 comprising an parasitically effective amount of an endo-parasiticide or endectocide selected from the group consisting of macrocyclic lactones, benzimidazoles, pro-benzimidazoles, imidazothiazoles, tetrahydropyrimidines, organophosphates and piperazines.

10



20



35

40



50

55

55

23. A chewable veterinary composition according to claim 18 comprising an effective amount of a macrocyclic lactone in combination with an effective amount of an anthelmintic selected from the group consisting of Albendazole, Clorsulon, Cydectin, Diethylcarbamazine, Febantel, Fenbendazole, Haloxon, Levamisole, Mebendazole, Morantel, Oxytoclozanide, Oxibendazole, Oxfendazole, Oxfendazole, Oxamniquine, Pyrantel, Piperazine, Praziquantel, Thiabendazole, Tetramisole, Trichlorfon, Thiabendazole, and a derivative thereof.
24. A chewable veterinary composition according to claim 18 comprising additionally an effective amount of an insecticide, acaricide or an insecticide and an acaricide.
25. A chewable veterinary composition according to claim 1 comprising an effective amount of milbemycin oxime and praziquantel.
26. A chewable veterinary composition according to claim 1 comprising an effective amount of lufenuron, praziquantel and milbemycin oxime.
27. A chewable veterinary composition according to claim 1 comprising an effective amount of cyclosporin.
28. A chewable veterinary composition according to claim 1 comprising an effective amount of an antimicrobial selected from the group consisting of a penicillin, tetracycline, sulfonamide, cephalosporin, cephamycin, aminoglycosid, trimethoprim, dimetridazole, erythromycin, framycetin, fruazolidone, pleuromutilin, streptomycin and a compound that is active against protozoa.
29. A chewable veterinary composition according to claim 1 comprising an effective amount of compound that is active against behavioral including separation worry or travel sickness of dogs and cats.
30. Process for the production of a highly palatable ductile chewable veterinary composition of claim 1, comprising (i) feeding the hopper of an extruder with an effective amount of one or more ingredients that are active against animal pests, pathogens or animal diseases; meat flavoring; partially gelatinized starch; a softener; and up to 9% (w/w) of water, (ii) cooling constantly down the mixture of active ingredients and carriers so that the temperature of the extrudate that leaves the tip of the extruder does during the whole extrusion process at no time exceed 40°C, (iii) pressing the extrudate through a die that is decisive for the shape of the chewable product, and (iv) cutting the extrudate that leaves the extruder into equal pieces.
31. Process according to claim 30 wherein the hopper of the extruder is fed continuously and simultaneously with pre-mixture (1) and pre-mixture (2), wherein pre-mixture (1) consist of a homogenized mixture of one or more active ingredients and partially gelatinized starch, and pre-mixture (2) consists of a homogenized mixture of meat flavoring, a softener and optionally of a carrier selected from the group consisting of a sweetener, softener, an antioxidant, a coloring agent and sodium chloride.
32. Process according to claim 30 wherein the extruder is cooled down below room temperature.
33. Use of (A) an effective amount of one or more ingredients that are active against animal pests, pathogens or animal diseases; (B) meat flavoring; (C) partially gelatinized starch; (D) a softener; (E) up to 9% water; and an active ingredient suitable for combating animal pests, pathogens or animal diseases for the preparation of a highly palatable ductile chewable veterinary composition.
34. Use according to claim 33 comprising 20 to 30 % (w/w) of a natural meat flavoring.
35. Use according to claim 34 wherein the natural meat flavoring comprises 20 to 55 % (w/w) fat.
36. Use according to claim 33 comprising 25 to 70 % (w/w) of partially gelatinized starch.
37. Use according to claim 36 wherein the partially gelatinized starch comprises 12 to 17 % (w/w) of gelatinized starch.
38. Use according to claim 33 comprising 10 to 20 % (w/w) of a softener, based upon the weight of the partially gelatinized starch.
39. Use according to claim 37 wherein the softener is selected from the group consisting of glycerol, polyethylene glycol

and polypropylene glycol.

40. Use according to claim 33 comprising 3 to 7 % (w/w) of water.

41. Use according to claim 33 wherein the animal pests are external animal parasites or internal animal parasites or both.

42. Use according to claim 33 comprising 1 to 10 % (w/w) of a sweetener.

43. Use according to claim 33 comprising 0 to 3.5 % (w/w) of an antioxidant.

44. Use according to claim 33 comprising 0 to 5 % (w/w) of a coloring agent.

45. Use according to claim 33 comprising 0 to 4% (w/w) of sodium chloride.


46. Use according to claim 33 comprising an parasitically effective amount of an ecto-parasiticide, an endo-parasiticide, an endectocide or of a combination of a parasiticide selected from the group consisting of an ecto-parasiticide, an endo-parasiticide and an endectocide.

47. Use according to claim 33 wherein the ecto-parasiticide is active against insects, members of the order Acarina or insects and members of the order Acarina.

48. Use according to claim 33 wherein the ecto-parasiticide is an insecticide which is either an insect adulticides or insect growth regulators.

49. Use according to claim 33 comprising an parasitically effective amount of an endo-parasiticide or endecticide selected from the group consisting of macrocyclic lactones, benzimidazoles, pro-benzimidazoles, imidazothiazoles, tetrahydropyrimidines, organophosphates and piperazines.

50. Use according to claim 33 comprising an effective amount of a natural or chemically modified macrocyclic lactone of formula (I)



wherein X is -C(H)(OH)-; -C(O)-; or -C(=N-OH)-; Y is -C(H₂)-; =C(H)-; -C(H)(OH)-; or -C(=N-OCH₃)-; R₁ is hydrogen or one of radicals



15

- 20




30


- 35

Patentansprüche

- 5 1. Plastisches kaubares Tierarzneipräparat mit hoher Palatabilität, umfassend (A) eine wirksame Menge von einem oder mehreren Bestandteilen, die gegen Tierschädlinge, Pathogene oder Tierkrankheiten wirksam sind; (B) Fleischgeschmack bzw. Fleischaroma; (C) teilweise gelierte Stärke; (D) einen Weichmacher; und (E) bis zu 9 % Wasser.
2. Kaubares Tierarzneipräparat nach Anspruch 1, worin die Tierkrankheit virale oder bakterielle Infektionen, Verhaltensstörungen, entzündliche Krankheiten und Autoimmunkrankheiten umfasst.
- 10 3. Kaubares Tierarzneipräparat nach Anspruch 1, umfassend 20 bis 30 % (Gewicht / Gewicht) von einem natürlichen Fleischgeschmack bzw. Fleischaroma.
4. Kaubares Tierarzneipräparat nach Anspruch 3, worin der natürliche Fleischgeschmack bzw. Fleischaroma 20 bis 55 % (Gewicht / Gewicht) Fett umfasst.
- 15 5. Kaubares Tierarzneipräparat nach Anspruch 1, umfassend 25 bis 70 % (Gewicht / Gewicht) von teilweise gelierter Stärke.
- 20 6. Kaubares Tierarzneipräparat nach Anspruch 5, worin die teilweise gelierte Stärke 12 bis 17 % (Gewicht / Gewicht) von gelierter Stärke umfasst.
7. Kaubares Tierarzneipräparat nach Anspruch 1, umfassend 10 bis 20 % (Gewicht / Gewicht) von einem Weichmacher, bezogen auf das Gewicht der teilweise gelierten Stärke.
- 25 8. Kaubares Tierarzneipräparat nach Anspruch 7, worin der Weichmacher aus der Gruppe, bestehend aus Glycerin, Polyethylenglycol und Polypropylenglycol, ausgewählt ist.
9. Kaubares Tierarzneipräparat nach Anspruch 1, umfassend 4 bis 6 % (Gewicht / Gewicht) Wasser.
- 30 10. Kaubares Tierarzneipräparat nach Anspruch 1, worin die Tierschädlinge äußere Tierparasiten oder innere Tierparasiten oder beides sind.
11. Kaubares Tierarzneipräparat nach Anspruch 1, umfassend 1 bis 10 % (Gewicht / Gewicht) von einem Süßungsmittel.
- 35 12. Kaubares Tierarzneipräparat nach Anspruch 1, umfassend 0 bis 3,5 % (Gewicht / Gewicht) von einem Antioxidationsmittel.
13. Kaubares Tierarzneipräparat nach Anspruch 1, umfassend 0 bis 5 % (Gewicht / Gewicht) von einem färbenden Mittel.
- 40 14. Kaubares Tierarzneipräparat nach Anspruch 1, umfassend 0 bis 4 % (Gewicht / Gewicht) Natriumchlorid.
15. Kaubares Tierarzneipräparat nach Anspruch 1, umfassend eine parasitizid wirksame Menge von einem Ecto-Parasitizid, einem Endo-Parasitizid, einem Endectozid, oder von einer Kombination von einem Parasitizid, ausgewählt aus der Gruppe, bestehend aus einem Ecto-Parasitizid, einem Endo-Parasitizid und einem Endectozid.
- 45 16. Kaubares Tierarzneipräparat nach Anspruch 15, worin das Ecto-Parasitizid gegen Insekten, Mitglieder der Gattung Acarina oder Insekten und Mitglieder der Gattung Acarina wirksam ist.
17. Kaubares Tierarzneipräparat nach Anspruch 16, worin das Ecto-Parasitizid ein Insektizid ist, das entweder ein Insekten-Adultizid oder Insekten-Wachstumsregulatoren ist.
- 50 18. Kaubares Tierarzneipräparat nach Anspruch 15, umfassend eine parasitizid wirksame Menge von einem Endo-Parasitizid oder Endectozid, ausgewählt aus der Gruppe, bestehend aus macrocyclischen Lactonen, Benzimidazolen, Pro-Benzimidazolen, Imidazothiazolen, Tetrahydropyrimidinen, Organophosphaten und Piperazinen.
- 55 19. Kaubares Tierarzneipräparat nach Anspruch 18, umfassend eine wirksame Menge von einem natürlichen oder chemisch modifizierten macrocyclischen Lacton der Formel (I)




worin X -C(H)(OH)-; -C(O)-; oder -C(=N-OH)- darstellt; Y -C(H₂)-; =C(H)-; -C(H)(OH)-; oder -C(=N-OCH₃)- darstellt; R₁ Wasserstoff oder einen der Reste



darstellt;

R₄ Hydroxyl, -NH-CH₃ oder -NH-OCH₃ darstellt; R₂ Wasserstoff, -CH₃, -C₂H₅, -CH(CH₃)-CH₃, -CH(CH₃)-C₂H₅, -C(CH₃)=CH-CH(CH₃)₂ oder Cyclohexyl darstellt; und, wenn die Bindung zwischen Atomen 22 und 23 eine Doppelbindung wiedergibt, ist das Kohlenstoffatom in der 23-Position unsubstituiert, sodass Y =C(H)- darstellt, oder, wenn die Bindung zwischen Atomen 22 und 23 eine Einfachbindung darstellt, ist das Kohlenstoffatom in 23-Position unsubstituiert oder mit Hydroxy oder mit der Gruppe =N-O-CH₃ substituiert, sodass Y -C(H₂)-; -C(H)(OH)-; oder -C(=N-OCH₃)- darstellt; in freier Form oder in der Form von einem physiologisch verträglichen Salz.

- 20.** Kaubares Tierarzneipräparat nach Anspruch 19, worin das macrocyclische Lacton eine Verbindung der Formel (I) ist, worin X -C(H)(OH)- darstellt; Y -C(H₂)- darstellt; R₁ den Rest



darstellt;

R₂ -CH₃ oder C₂H₅ darstellt, und die Bindung zwischen Atomen 22 und 23 eine Einfachbindung wiedergibt.


- 21.** Kaubares Tierarzneipräparat nach Anspruch 19, worin das macrocyclische Lacton aus der Gruppe, bestehend aus Avermectinen, Milbemycinen und Derivaten davon, in freier Form oder in der Form von einem physiologisch verträglichen Salz, ausgewählt ist.

- 22.** Kaubares Tierarzneipräparat nach Anspruch 21, worin das macrocyclische Lacton aus der Gruppe, bestehend aus Ivermectin, Doramectin, Moxidectin, Selamectin, Emamectin, Eprinomectin, Milbemectin, Abamectin, Milbemycin-oxim, Nemadectin und einem Derivat davon, in freier Form oder in der Form von einem physiologisch verträglichen

Salz ausgewählt ist.


23. Kaubares Tierarzneipräparat nach Anspruch 18, umfassend eine wirksame Menge von einem macrocyclischen Lacton in Kombination mit einer wirksamen Menge von einem Anthelmintikum, ausgewählt aus der Gruppe, bestehend aus Albendazol, Clorsulon, Cydectin, Diethylcarbamazin, Febantel, Fenbendazol, Haloxon, Levamisol, Mebendazol, Morantel, Oxyclozanid, Oxibendazol, Oxfendazol, Oxfendazol, Oxamniquin, Pyrantel, Piperazin, Praziquantel, Thiabendazol, Tetramisol, Trichlorfon, Thiabendazol und einem Derivat davon.
24. Kaubares Tierarzneipräparat nach Anspruch 18, umfassend zusätzlich eine wirksame Menge von einem Insektizid, Akarizid oder einem Insektizid und einem Akarizid.
25. Kaubares Tierarzneipräparat nach Anspruch 1, umfassend eine wirksame Menge von Milbemycinoxim und Praziquantel.
26. Kaubares Tierarzneipräparat nach Anspruch 1, umfassend eine wirksame Menge Lufenuron, Praziquantel und Milbemycinoxim.
27. Kaubares Tierarzneipräparat nach Anspruch 1, umfassend eine wirksame Menge von Cyclosporin.
28. Kaubares Tierarzneipräparat nach Anspruch 1, umfassend eine wirksame Menge von einem antimikrobiellen Mittel, ausgewählt aus der Gruppe, bestehend aus einem Penicillin, Tetracyclin, Sulfonamid, Cephalosporin, Cephamecin, Aminoglucosid, Trimethoprim, Dimetridazol, Erythromycin, Framycetin, Fruazolidon, Pleuromutilin, Streptomycin und einer Verbindung, die gegen Protozoen wirksam ist.
29. Kaubares Tierarzneipräparat nach Anspruch 1, umfassend eine wirksame Menge einer Verbindung, die gegen Verhaltenskrankheit, einschließlich Trennungsangst oder Reisekrankheit, von Hunden und Katzen wirksam ist.
30. Verfahren für die Herstellung eines plastischen kaubaren Tierarzneipräparats mit hoher Palatabilität nach Anspruch 1, umfassend (i) Beschicken des Trichters von einem Extruder mit einer wirksamen Menge von einem oder mehreren Bestandteilen, die gegen Tierschädlinge, Pathogene oder Tierkrankheiten wirksam sind; Fleischgeschmack bzw. Fleischaroma; teilweise gelierte Stärke; einem Weichmacher; und bis zu 9 % (Gewicht / Gewicht) Wasser, (ii) konstantes Herunterkühlen des Gemisches von Wirkbestandteilen und Trägern, sodass die Temperatur des Extrudats, das die Spitze des Extruders verlässt, während des gesamten Extrusionsvorgangs zu keiner Zeit 40°C übersteigt, (iii) Pressen des Extrudats durch eine Düse, die für die Form des kaubaren Produkts entscheidend ist, und (iv) Schneiden des Extrudats, das den Extruder verlässt, in gleiche Stücke.
31. Verfahren nach Anspruch 30, wobei der Trichter von dem Extruder kontinuierlich und gleichzeitig mit einer Vormischung (1) und Vormischung (2) beschickt wird, wobei Vormischung (1) aus einem homogenisierten Gemisch von einem oder mehreren Wirkbestandteilen und teilweise gelierte Stärke besteht, und Vormischung (2) aus einem homogenisierten Gemisch von Fleischgeschmack bzw. Fleischaroma, einem Weichmacher und gegebenenfalls einem Träger, ausgewählt aus der Gruppe, bestehend aus einem Süßungsmittel, Weichmacher, einem Antioxidationsmittel, einem färbenden Mittel und Natriumchlorid, besteht.
32. Verfahren nach Anspruch 30, wobei der Extruder unter Raumtemperatur heruntergekühlt wird.
33. Verwendung von (A) einer wirksamen Menge von einem oder mehreren Bestandteilen, die gegen Tierschädlinge, Pathogene oder Tierkrankheiten wirksam sind; (B) Fleischgeschmack bzw. Fleischaroma; (C) teilweise gelierte Stärke; (D) einem Weichmacher; (E) bis zu 9 % Wasser; und einem Wirkbestandteil, der zum Bekämpfen von Tierschädlingen, Pathogenen oder Tierkrankheiten geeignet ist, für die Herstellung von einem plastischen kaubaren Tierarzneipräparat mit hoher Palatabilität.
34. Verwendung nach Anspruch 33, umfassend 20 bis 30 % (Gewicht / Gewicht) von einem natürlichen Fleischgeschmack bzw. Fleischaroma.
35. Verwendung nach Anspruch 34, wobei der natürliche Fleischgeschmack bzw. Fleischaroma 20 bis 55 % (Gewicht / Gewicht) Fett umfasst.
36. Verwendung nach Anspruch 33, umfassend 25 bis 70 % (Gewicht / Gewicht) von teilweise gelierte Stärke.

37. Verwendung nach Anspruch 36, wobei die teilweise gelierte Stärke 12 bis 17 % (Gewicht / Gewicht) gelierte Stärke umfasst.
38. Verwendung nach Anspruch 33, umfassend 10 bis 20 % (Gewicht / Gewicht) von einem Weichmacher, bezogen auf das Gewicht der teilweise gelierten Stärke.
39. Verwendung nach Anspruch 37, wobei der Weichmacher aus der Gruppe, bestehend aus Glycerin, Polyethylenglycol und Polypropylenglycol, ausgewählt ist.
40. Verwendung nach Anspruch 33, umfassend 3 bis 7 % (Gewicht / Gewicht) Wasser.
41. Verwendung nach Anspruch 33, wobei die Tierschädlinge äußere Tierparasiten oder innere Tierparasiten oder beides sind.
42. Verwendung nach Anspruch 33, umfassend 1 bis 10 % (Gewicht / Gewicht) von einem Süßungsmittel.
43. Verwendung nach Anspruch 33, umfassend 0 bis 3,5 % (Gewicht / Gewicht) von einem Antioxidationsmittel.
44. Verwendung nach Anspruch 33, umfassend 0 bis 5 % (Gewicht / Gewicht) von einem färbenden Mittel.
45. Verwendung nach Anspruch 33, umfassend 0 bis 4 % (Gewicht / Gewicht) von Natriumchlorid.
46. Verwendung nach Anspruch 33, umfassend eine parasitizid wirksame Menge von einem Ecto-Parasitizid, einem Endo-Parasitizid, einem Endectozid, oder einer Kombination von einem Parasitizid, ausgewählt aus der Gruppe, bestehend aus einem Ecto-Parasitizid, einem Endo-Parasitizid und einem Endectozid.
47. Verwendung nach Anspruch 33, wobei das Ecto-Parasitizid gegen Insekten, Mitglieder der Gattung Acarina oder Insekten und Mitglieder der Gattung Acarina wirksam ist.
48. Verwendung nach Anspruch 33, wobei das Ecto-Parasitizid ein Insektizid ist, das entweder ein Insekten-Adultizid oder Insekten-Wachstumsregulatoren ist.
49. Verwendung nach Anspruch 33, umfassend eine parasitizid wirksame Menge von einem Endo-Parasitizid oder Endectizid, ausgewählt aus der Gruppe, bestehend aus macrocyclischen Lactonen, Benzimidazolen, Pro-Benzimidazolen, Imidazothiazolen, Tetrahydropyrimidinen, Organophosphaten und Piperazinen.
50. Verwendung nach Anspruch 33, umfassend eine wirksame Menge von einem natürlichen oder chemisch modifizierten macrocyclischen Lacton der Formel (I)



worin X -C(H)(OH)-; -C(O)-; oder -C(=N-OH)- darstellt; Y -C(H₂)-; =C(H)-; -C(H)(OH)-; oder -C(=N-OCH₃)- darstellt;


R₁ Wasserstoff oder einen der Reste



darstellt;

R₄ Hydroxyl, -NH-CH₃ oder -NH-OCH₃ darstellt; R₂ Wasserstoff, -CH₃, -C₂H₅, -CH(CH₃)-CH₃, -CH(CH₃)-C₂H₅, -C(CH₃)=CH-CH(CH₃)₂ oder Cyclohexyl darstellt; und, wenn die Bindung zwischen Atomen 22 und 23 eine Doppelbindung wiedergibt, ist das Kohlenstoffatom in der 23-Position unsubstituiert, sodass Y =C(H)- darstellt, oder, wenn die Bindung zwischen Atomen 22 und 23 eine Einfachbindung darstellt, ist das Kohlenstoffatom in 23-Position unsubstituiert oder mit Hydroxy oder mit der Gruppe =N-O-CH₃ substituiert, sodass Y -C(H₂)-; -C(H)(OH)-; oder -C(=N-OCH₃)- darstellt; in freier Form oder in der Form von einem physiologisch verträglichen Salz.

51. Verwendung nach Anspruch 50, wobei das macrocyclische Lacton eine Verbindung der Formel (I) ist, worin X -C(H)(OH)- darstellt; Y -C(H₂)- darstellt; R₁ den Rest



darstellt;

R₂ -CH₃ oder C₂H₅ darstellt, und die Bindung zwischen Atomen 22 und 23 eine Einfachbindung wiedergibt.

52. Verwendung nach Anspruch 46, wobei das Endectizid ein macrocyclisches Lacton ist, das aus der Gruppe, bestehend aus Avermectinen, Milbemycinen und Derivaten davon, in freier Form oder in der Form von einem physiologisch verträglichen Salz, ausgewählt ist.
53. Verwendung nach Anspruch 50, wobei das makrocyclische Lacton aus der Gruppe, bestehend aus Ivermectin, Doramectin, Moxidectin, Selamectin, Emamectin, Eprinomectin, Milbemectin, Abamectin, Milbemycinoxim, Nemalectin und einem Derivat davon, in freier Form oder in der Form von einem physiologisch verträglichen Salz ausgewählt ist.
54. Verwendung nach Anspruch 46, umfassend eine wirksame Menge von einem macrocyclischen Lacton in Kombination mit einer wirksamen Menge von einem Anthelmintikum, ausgewählt aus der Gruppe, bestehend aus Albendazol, Clorsulon, Cydectin, Diethylcarbamazin, Febantel, Fenbendazol, Haloxon, Levamisol, Mebendazol, Morantel, Oxytoclozanid, Oxibendazol, Oxfendazol, Oxfendazol, Oxamniquin, Pyrantel, Piperazin, Praziquantel, Thiabendazol, Tetramisol, Trichlorfon, Thiabendazol und einem Derivat davon.
55. Verwendung nach Anspruch 46, umfassend zusätzlich zu einem Endo-Parasitizid oder einem Endectizid eine wirksame Menge von einem Insektizid, Akarizid oder einem Insektizid und einem Akarizid.
56. Verwendung nach Anspruch 33, umfassend eine wirksame Menge von Milbemycinoxim und Praziquantel.
57. Verwendung nach Anspruch 33, umfassend eine wirksame Menge Lufenuron, Praziquantel und Milbemycinoxim.
58. Verwendung nach Anspruch 33, umfassend eine wirksame Menge von Cyclosporin.

59. Verwendung nach Anspruch 33, umfassend eine wirksame Menge von einem antimikrobiellen Mittel, ausgewählt aus der Gruppe, bestehend aus einem Penicillin, Tetracyclin, Sulfonamid, Cephalosporin, Cephamecin, Aminoglycosid, Trimethoprim, Dimetridazol, Erythromycin, Framycetin, Fruazolidon, Pleuromutilin, Streptomycin und einer Verbindung, die gegen Protozoen wirksam ist.

60. Verwendung nach Anspruch 33, umfassend eine wirksame Menge einer Verbindung, die gegen Verhaltenskrankheit, einschließlich Trennungsangst oder Reisekrankheit, von Hunden und Katzen wirksam ist.

Revendications

1. Composition vétérinaire à mâcher ductile hautement appétente comprenant (A) une quantité efficace d'un ou plusieurs ingrédients qui sont actifs contre des animaux nuisibles, pathogènes ou maladies animales ; (B) un arôme de viande ; (C) de l'amidon partiellement gélatinisé ; (D) un assouplissant ; et (E) jusqu'à 9% d'eau.

2. Composition vétérinaire à mâcher selon la revendication 1, dans laquelle la maladie animale comprend des infections virales ou bactériennes, des troubles de comportement, des maladies inflammatoires et des maladies auto-immunes.

3. Composition vétérinaire à mâcher selon la revendication 1 comprenant 20 à 30% (en poids) d'un arôme naturel de viande.

4. Composition vétérinaire à mâcher selon la revendication 4, dans laquelle l'arôme naturel de viande comprend 20 à 55% (en poids) de graisse.

5. Composition vétérinaire à mâcher selon la revendication 1 comprenant 25 à 70% (en poids) d'amidon partiellement gélatinisé.

6. Composition vétérinaire à mâcher selon la revendication 5, dans laquelle l'amidon partiellement gélatinisé comprend 12 à 17% (en poids) d'amidon gélatinisé.

7. Composition vétérinaire à mâcher selon la revendication 1 comprenant 10 à 20% (en poids) d'un assouplissant, basé sur le poids de l'amidon partiellement gélatinisé.

8. Composition vétérinaire à mâcher selon la revendication 7, dans laquelle l'assouplissant est choisi dans le groupe constitué de glycérol, polyéthylène glycol et polypropylène glycol.

9. Composition vétérinaire à mâcher selon la revendication 1 comprenant 4 à 6% (en poids) d'eau.

10. Composition vétérinaire à mâcher selon la revendication 1, dans laquelle les animaux nuisibles sont des parasites externes des animaux ou des parasites internes des animaux ou les deux.

11. Composition vétérinaire à mâcher selon la revendication 1 comprenant 1 à 10% (en poids) d'un édulcorant.

12. Composition vétérinaire à mâcher selon la revendication 1 comprenant 0 à 3,5% (en poids) d'un antioxydant.

13. Composition vétérinaire à mâcher selon la revendication 1 comprenant 0 à 5% (en poids) d'un agent colorant.

14. Composition vétérinaire à mâcher selon la revendication 1 comprenant 0 à 4% (en poids) de chlorure de sodium.


15. Composition vétérinaire à mâcher selon la revendication 1 comprenant une quantité efficace de manière antiparasitaire d'un ectoparasiticide, d'un endoparasiticide, d'un endectocide ou d'une combinaison d'un parasiticide choisi dans le groupe constitué d'un ectoparasiticide, d'un endoparasiticide et d'un endectocide.

16. Composition vétérinaire à mâcher selon la revendication 15, dans laquelle l'ectoparasiticide est actif contre des insectes, des membres de l'ordre des acariens ou des insectes et des membres de l'ordre des acariens.


17. Composition vétérinaire à mâcher selon la revendication 16, dans laquelle l'ectoparasiticide est un insecticide qui est soit un adulticide d'insectes soit des régulateurs de croissances des insectes.

- 18.** Composition vétérinaire à mâcher selon la revendication 15 comprenant une quantité efficace de manière antiparasitaire d'un endoparasiticide ou d'un endectocide choisi dans le groupe constitué de lactones macrocycliques, benzimidazoles, pro-benzimidazoles, imidazothiazoles, tétrahydropyrimidines, organophosphates et pipérazines.

19. Composition vétérinaire à mâcher selon la revendication 18 comprenant une quantité efficace d'un lactone macrocyclique naturel ou chimiquement modifié de formule (I)




dans laquelle X est -C(H)(OH)-; -C(O)-; ou -C(=N-OH)-; Y est -C(H₂)-; =C(H)-; -C(H)(OH)-; ou -C(=N-OCH₃)-; R₁ est de l'hydrogène ou un des radicaux



R₄ est de l'hydroxyle, -NH-CH₃ ou -NH-OCH₃; R₂ est de l'hydrogène, -CH₃, -C₂H₅, -CH(CH₃)-CH₃, -CH(CH₃)-C₂H₅, -C(CH₃)=CH-CH(CH₃)₂ ou du cyclohexyle ; et si la liaison entre les atomes 22 et 23 représente une liaison double l'atome de carbone en position 23 est non substitué de sorte que Y est =C(H)-, ou si la liaison entre les atomes 22 et 23 est une liaison l'atome de carbone en position 23 est non substitué ou substitué par de l'hydroxy ou par le groupe =N-O-CH₃ de sorte que Y est -C(H)₂- ; -C(H)(OH)- ; ou -C(=N-OCH₃)- ; sous forme libre ou sous forme d'un sel physiologiquement acceptable.

20. Composition vétérinaire à mâcher selon la revendication 19, dans laquelle le lactone macrocyclique est un composé de formule (I) dans laquelle X est -C(H)(OH)- ; Y est -C(H₂)- ; R₁ est le radical



R₂ est -CH₃ ou C₇H₅, et la liaison entre les atomes 22 et 23 représente une liaison simple.


21. Composition vétérinaire à mâcher selon la revendication 19, dans laquelle le lactone macrocyclique est choisi dans le groupe constitué d'ivermectines, de milbémycines et de dérivés de celles-ci, sous forme libre ou sous forme d'un sel physiologiquement acceptable.

- 22.** Composition vétérinaire à mâcher selon la revendication 21, dans laquelle le lactone macrocyclique est choisi dans le groupe constitué d'ivermectine, de doramectine, de moxidectine, de sélamectine, d'enamectine, d'eprinomectine, de milbémycines, d'abamectine, d'oxime de milbémycine, de nemadectine et d'un dérivé de celles-ci, sous forme


libre ou sous forme d'un sel physiologiquement acceptable.

23. Composition vétérinaire à mâcher selon la revendication 18 comprenant une quantité efficace d'un lactone macrocyclique en combinaison avec une quantité efficace d'un anthelminthique choisi dans le groupe constitué d'abendazole, de clorsulon, de cydectine, de diéthylcarbamazine, de fébantel, de fenbendazole, d'haloxon, de lévamisole, de mébendazole, de morantel, d'oxyclozanide, d'oxybendazole, d'oxfendazole, d'oxfendazole, d'oxamniquine, de pyrantel, de pipérazine, de praziquantel, de thiabendazole, de tétramisole, de trichlorfon, de thiabendazole et d'un dérivé de ceux-ci.
24. Composition vétérinaire à mâcher selon la revendication 18 comprenant en outre une quantité efficace d'un insecticide, d'un acaricide ou d'un insecticide et d'un acaricide.
25. Composition vétérinaire à mâcher selon la revendication 1 comprenant une quantité efficace d'oxime de milbémycine et de praziquantel.
26. Composition vétérinaire à mâcher selon la revendication 1 comprenant une quantité efficace de lufénuron, de praziquantel et d'oxime de milbémycine.
27. Composition vétérinaire à mâcher selon la revendication 1 comprenant une quantité efficace de cyclosporine.
28. Composition vétérinaire à mâcher selon la revendication 1 comprenant une quantité efficace d'un antimicrobien choisi dans le groupe constitué d'une pénicilline, d'une tétracycline, d'un sulfonamide, d'une céphalosporine, d'une céphamycine, d'un aminoside, d'un triméthoprime, d'un dimétridazole, d'une érythromycine, d'un framycétine, d'une fruazolidone, d'une pleuromutiline, d'une streptomycine et d'un composé qui est actif contre les protozoaires.
29. Composition vétérinaire à mâcher selon la revendication 1 comprenant une quantité efficace d'un composé qui est actif contre les troubles de comportement, y compris l'anxiété de séparation ou le mal de transport des chiens et chats.
30. Procédé pour la production d'une composition vétérinaires à mâcher ductile hautement appétente selon la revendication 1 comprenant (i) l'introduction dans la trémie d'une extrudeuse d'une quantité efficace d'un ou plusieurs ingrédients qui sont actifs contre les animaux nuisibles, pathogènes ou maladies animales ; d'arôme de viande ; d'amidon partiellement gélatinisé ; d'un assouplissant ; et de jusqu'à 9% (en poids) d'eau, (ii) le refroidissement constant du mélange d'ingrédients actifs et de supports de sorte que la température de l'extrudat qui quitte la pointe de l'extrudeuse n'excède à aucun moment durant tout le procédé 40°C, (iii) la pression de l'extrudat à travers d'un moule qui est décisif pour la forme du produit à mâcher, et (iv) le découpage de l'extrudat qui quitte l'extrudeuse en plusieurs pièces égales.
31. Procédé selon la revendication 30, dans lequel la trémie de l'extrudeuse est alimentée continuellement et simultanément avec un pré-mélange (1) et un pré-mélange (2), où le pré-mélange (1) est composé d'un mélange homogénéisé d'un ou plusieurs ingrédients actifs et d'amidon partiellement gélatinisé, et le pré-mélange (2) est composé d'un mélange homogénéisé d'arôme de viande, d'un assouplissant et facultativement d'un support choisi dans le groupe constitué de d'un édulcorant, d'un assouplissant, d'un antioxydant, d'un agent colorant et de chlorure de sodium.
32. Procédé selon la revendication 30, dans lequel l'extrudeuse est refroidie en dessous de la température ambiante.
33. Utilisation (A) d'une quantité efficace d'un ou plusieurs ingrédients qui sont actifs contre des animaux nuisibles, de pathogènes ou de maladies animales ; (B) d'un arôme de viande ; (C) d'un amidon partiellement gélatinisé ; (D) d'un assouplissant ; (E) de jusqu'à 9% d'eau ; et d'un ingrédient actif approprié pour combattre des animaux nuisibles, des pathogènes ou des maladies animales pour la préparation d'une composition vétérinaire à mâcher ductile hautement appétente.
34. Utilisation selon la revendication 33 comprenant 20 à 30% (en poids) d'un arôme naturel de viande.
35. Utilisation selon la revendication 34, dans laquelle l'arôme naturel de viande comprend 20 à 55% (en poids) de graisse.
36. Utilisation selon la revendication 33 comprenant 25 à 70% (en poids) d'amidon partiellement gélatinisé.

37. Utilisation selon la revendication 36, dans laquelle l'amidon partiellement gélatinisé comprend 12 à 17% (en poids) d'amidon gélatinisé.
38. Utilisation selon la revendication 33 comprenant 10 à 20% (en poids) d'un assouplissant, basé sur le poids de l'amidon partiellement gélatinisé.
39. Utilisation selon la revendication 37, dans laquelle l'assouplissant est choisi dans le groupe constitué de glycérol, polyéthylène glycol et polypropylène glycol.
40. Utilisation selon la revendication 33 comprenant 3 à 7% (en poids) d'eau.
41. Utilisation selon la revendication 33, dans laquelle les animaux nuisibles sont des parasites externes des animaux ou des parasites internes des animaux ou les deux.
42. Utilisation selon la revendication 33 comprenant 1 à 10% (en poids) d'un édulcorant.
43. Utilisation selon la revendication 33 comprenant 0 à 3,5% (en poids) d'un antioxydant.
44. Utilisation selon la revendication 33 comprenant 0 à 5% (en poids) d'un agent colorant.
45. Utilisation selon la revendication 33 comprenant 0 à 4% (en poids) de chlorure de sodium.
46. Utilisation selon la revendication 33 comprenant une quantité efficace de manière antiparasitaire d'un ectoparasiticide, d'un endoparasiticide, d'un endectocide ou d'une combinaison d'un parasiticide choisi dans le groupe constitué de d'un ectoparasiticide, d'un endoparasiticide et d'un endectocide.
47. Utilisation selon la revendication 33, dans laquelle l'ectoparasiticide est actif contre des insectes, des membres de l'ordre des acariens ou des insectes et des membres de l'ordre des acariens.
48. Utilisation selon la revendication 33, dans laquelle l'ectoparasiticide est un insecticide qui est soit un adulticide d'insectes soit des régulateurs de croissances des insectes.
49. Utilisation selon la revendication 33 comprenant une quantité efficace de manière antiparasitaire d'un endoparasiticide ou d'un endectocide choisi dans le groupe constitué de lactones macrocycliques, benzimidazoles, pro-benzimidazoles, imidazothiazoles, tétrahydropyrimidines, organophosphates et pipérazines.
50. Utilisation selon la revendication 33 comprenant une quantité efficace d'un lactone macrocyclique naturel ou chimiquement modifié de formule (I)




dans laquelle X est -C(H)(OH)- ; -C(O)- ; ou -C(=N-OH)- ; Y est -C(H2)- ; =C(H)- ; -C(H)(OH)- ; ou -C(=N-OCH3)- ; R₁ est de l'hydrogène ou un des radicaux



15

R_4 est de l'hydroxyle, $-NH-CH_3$ ou $-NH-OCH_3$; R_2 est de l'hydrogène, $-CH_3$, $-C_2H_5$, $-CH(CH_3)-CH_3$, $-CH(CH_3)-C_2H_5$, $-C(CH_3)=CH-CH(CH_3)_2$ ou du cyclohexyle ; et si la liaison entre les atomes 22 et 23 représente une liaison double l'atome de carbone en position 23 est non substitué de sorte que Y est $=C(H)-$, ou si la liaison entre les atomes 22 et 23 est une liaison l'atome de carbone en position 23 est non substitué ou substitué par de l'hydroxy ou par le groupe $=N-O-CH_3$ de sorte que Y est $-C(H_2)-$; $-C(H)(OH)-$; ou $-C(=N-OCH_3)-$; sous forme libre ou sous forme d'un sel physiologiquement acceptable.

- 20
51. Utilisation selon la revendication 50, dans laquelle le lactone macrocyclique est un composé de formule (I) dans laquelle X est $-C(H)(OH)-$; Y est $-C(H_2)-$; R_1 est le radical



R_2 est $-CH_3$ ou C_2H_5 , et la liaison entre les atomes 22 et 23 représente une liaison simple.

- 30
52. Utilisation selon la revendication 46, dans laquelle l'endectocide est un lactone macrocyclique est choisi dans le groupe constitué d'ivermectines, de milbémycines et de dérivés de celles-ci, sous forme libre ou sous forme d'un sel physiologiquement acceptable.
- 35
53. Utilisation selon la revendication 50, dans laquelle le lactone macrocyclique est choisi dans le groupe constitué d'ivermectine, de doramectine, de moxidectine, de sélamectine, d'enamectine, d'epinomectine, de milbémectine, d'abamectine, d'oxime de milbémycine, de nemadectine et d'un dérivé de celles-ci, sous forme libre ou sous forme d'un sel physiologiquement acceptable.
- 40
54. Utilisation selon la revendication 46 comprenant une quantité efficace d'un lactone macrocyclique en combinaison avec une quantité efficace d'un anthelminthique choisi dans le groupe constitué d'abendazole, de clorsulon, de cydectine, de diéthylcarbamazine, de fébantel, de fenbendazole, d'haloxon, de lévamisole, de mébendazole, de morantel, d'oxyclozanide, d'oxybendazole, d'oxfendazole, d'oxfendazole, d'oxamniquine, de pyrantel, de pipérazine, de praziquantel, de thiabendazole, de tétramisole, de trichlorfon, de thiabendazole et d'un dérivé de ceux-ci.
- 45
55. Utilisation selon la revendication 46 comprenant en plus d'un endoparasiticide ou un endectocide une quantité efficace d'un insecticide, d'un acaricide ou d'un insecticide et d'un acaricide.
- 50
56. Utilisation selon la revendication 33 comprenant une quantité efficace d'oxime de milbémycine et de praziquantel.
57. Utilisation selon la revendication 33 comprenant une quantité efficace de lufénuron, de praziquantel et d'oxime de milbémycine.
58. Utilisation selon la revendication 33 comprenant une quantité efficace de cyclosporine.
59. Utilisation selon la revendication 33 comprenant une quantité efficace d'un antimicrobien choisi dans le groupe constitué d'une pénicilline, d'une tétracycline, d'un sulfonamide, d'une céphalosporine, d'une céphamycine, d'un aminoside, d'un triméthoprime, d'un dimétridazole, d'une érythromycine, d'un framycétine, d'une fruazolidone, d'une pleuromutiline, d'une streptomycine et d'un composé qui est actif contre les protozoaires.

- 60.** Utilisation selon la revendication 33 comprenant une quantité efficace d'un composé qui est actif contre les troubles de comportement, y compris l'anxiété de séparation ou le mal de transport des chiens et chats.

5

10

15

20

25

30

35

40

45

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- EP 1247456 A [0005]
- GB 2300103 A [0006]
- US 4199569 A [0043]
- US 5089480 A [0043]
- EP 0237339 A [0043]
- US 4916154 A [0043]
- US 4874749 A [0043]
- US 4427663 A [0043]
- US 3950360 A [0043]
- US 4310519 A [0043]
- WO 0183500 A [0043]
- US 4869901 A [0043]

Non-patent literature cited in the description

- *The 17th international Conference of the World Association for the Advancement of Veterinary Parasitology*, 19 August 1999 [0043]
- THE MERCK INDEX. Merck & Co Inc, 1999 [0048]
- CD-ROM. Chapman & Hill/CRC, 1999, Hampden Data Service Ltd, 1999 [0048]
- THE MERCK INDEX, 1999 [0048]