



(11) **EP 1 755 596 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Claims EN 1, 9, 10

(51) Int Cl.:
C07D 213/56 ^(2006.01) **A61K 31/44** ^(2006.01)

(86) International application number:
PCT/US2005/015333

(48) Corrigendum issued on:
10.02.2016 Bulletin 2016/06

(87) International publication number:
WO 2005/108349 (17.11.2005 Gazette 2005/46)

(45) Date of publication and mention
of the grant of the patent:
12.08.2015 Bulletin 2015/33

(21) Application number: **05744537.1**

(22) Date of filing: **03.05.2005**

(54) **PROCESS FOR PREPARING ATAZANAVIR BISULFATE AND NOVEL FORMS**

VERFAHREN ZUR HERSTELLUNG VON ATAZANAVIRBISULFAT UND NEUE FORMEN

PROCEDE POUR LA PREPARATION DE BISULFATE D'ATAZANAVIR ET DE NOUVELLES FORMES

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU MC NL PL PT RO SE SI SK TR**
Designated Extension States:
HR LV MK YU

(30) Priority: **04.05.2004 US 568043 P**
07.09.2004 US 607533 P
02.05.2005 US 119558

(43) Date of publication of application:
28.02.2007 Bulletin 2007/09

(60) Divisional application:
13175944.1 / 2 669 273
15180557.9 / 2 980 074

(73) Proprietor: **Bristol-Myers Squibb Holdings Ireland
6312 Steinhausen (CH)**

(72) Inventors:
• **KIM, Soojin**
Demarest, NJ 07627 (US)

- **LOTZ, Bruce, T.**
Yardville, NJ 08620 (US)
- **MALLEY, Mary, F.**
Lawrenceville, NJ 08648 (US)
- **GOUGOUTAS, Jack, Z.**
Princeton, NJ 08540 (US)
- **DAVIDOVICH, Martha**
East Brunswick, NJ 08816 (US)
- **SRIVASTAVA, Sushil, K.**
Dayton, NJ 08810 (US)

(74) Representative: **Vossius & Partner**
Patentanwälte Rechtsanwälte mbB
Siebertstrasse 3
81675 München (DE)

(56) References cited:
US-A- 5 849 911 US-A- 6 087 383

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

REFERENCE TO OTHER APPLICATIONS


- 5 [0001] The present application takes priority from U.S. provisional application Nos. 60/568,043, filed May 4, 2004, and 60/607,533, filed September 7, 2004.

FIELD OF THE INVENTION

- 10 [0002] The present invention relates to a process for preparing the HIV protease inhibitor atazanavir bisulfate and novel forms thereof.

BACKGROUND OF THE INVENTION

- 15 [0003] U.S. Patent No. 5,849,911 to Fässler et al. discloses a series of azapeptide HIV protease inhibitors (which includes atazanavir) which have the structure



25 wherein

R₁ is lower alkoxy carbonyl,

R₂ is secondary or tertiary lower alkyl or lower alkylthio-lower alkyl,

R₃ is phenyl that is unsubstituted or substituted by one or more lower alkoxy radicals, or C₄-C₈ cycloalkyl,


30 R₄ is phenyl or cyclohexyl each substituted in the 4-position by unsaturated heterocyclyl that is bonded by way of a ring carbon atom, has from 5 to 8 ring atoms, contains from 1 to 4 hetero atoms selected from nitrogen, oxygen, sulfur, sulfinyl (-SO-) and sulfonyl (-SO₂-) and is unsubstituted or substituted by lower alkyl or by phenyl-lower alkyl,

R₅, independently of R₂, has one of the meanings mentioned for R₂, and

35 R₆, independently of R₁, is lower alkoxy carbonyl, or a salt thereof, provided that at least one salt-forming group is present which includes various pharmaceutically acceptable acid addition salts thereof.

[0004] Several methods for preparing the azapeptides are provided including preparation of a compound where R₁ and R₆, and R₂ and R₅ are in each case two identical radicals, wherein a diamino compound of the structure


40 (a)



45

is condensed with an acid of the structure

50 (b)




55


or with a reactive acid derivative thereof, wherein R₁' and R₂' are as defined for R₁ and R₆, and for R₂ and R₅,

respectively.


[0005] In forming atazanavir employing the above method the diamino compound (a) which will have the structure




is prepared by coupling the epoxide




with a hyrazinocarbamate




in the presence of isopropyl alcohol to form the protected diamine



which is treated with hydrochloric acid in the presence a solvent such as tetrahydrofuran to form the diamine (a)




15 [0006] The diamine is isolated and used in the next coupling step where it is reacted with an acid (b)



25 or a reactive ester thereof employing a coupling agent such as O-(1,2-dihydro-2-oxo-1-pyridyl)-N,N,N¹,N¹-tetramethyluronium-tetrafluoro-borate (TPTU).

[0007] It has been found that the diamine free base is unstable and therefore undesirable for use in preparing the free base of atazanavir.

[0008] U.S. Patent No. 6,087,383 to Singh et al. discloses the bisulfate salt of the azapeptide HIV protease inhibitor known as atazanavir which has the structure



(also referred to as atazanavir bisulfate or atazanavir sulfate).

45 [0009] Example 3 of Singh et al. describes the preparation of atazanavir bisulfate in the form of Type-II crystals which are a hydrated hygroscopic and crystalline form and Type-I crystals which appear to be an anhydrous/desolvated crystalline form.

BRIEF DESCRIPTION OF THE INVENTION

50 [0010] In accordance with the present invention, novel forms of atazanavir bisulfate are provided which includes Pattern C material and Form E3. Pattern C material is preferred.

[0011] In addition, in accordance with the present invention, a process is provided for preparing atazanavir bisulfate in the form of Form A crystals (bulk drug) (which are referred to as Type I crystals in Example 3 of U.S. Patent No. 6,087,383 to Singh et al). The Form A crystals prepared by the process of the invention have a desired substantially consistent particle size distribution and substantially consistent mean particle size, and are employed in the conversion to Pattern C material, a partially crystalline material, which is formulated with various excipients to prepare the drug product.

55 [0012] The process of the invention for preparing Form A crystals of atazanavir bisulfate salt employs a modified cubic crystallization technique wherein sulfuric acid is added at an increasing rate according to a cubic equation (as described

hereinafter), and includes the steps of reacting a solution of atazanavir free base in an organic solvent which is acetone or a mixture of acetone and N-methyl pyrrolidone with a first portion of concentrated sulfuric acid in an amount to react with less than about 15%, preferably less than about 12%, by weight of the atazanavir free base, adding seeds of atazanavir bisulfate Form A crystals to the reaction mixture, and as crystals of atazanavir bisulfate form, adding additional concentrated sulfuric acid in multiple stages at increasing rates according to a cubic equation to effect formation of Form A crystals.


[0013] In addition, in accordance with the present invention, a process is provided for preparing a form of atazanavir which is derived from and includes atazanavir bisulfate, and which is referred to as Pattern C material. Pattern C may be produced by suspending crystals of Form A in water and drying. Alternatively, Pattern C material may be formed by subjecting crystals of Form A to high relative humidity of greater than about 95% RH (water vapor) for at least 24 hours. Pattern C material may also be formed by wet granulating the atazanavir bisulfate or a combination of atazanavir bisulfate and excipients and drying the wet granulation.

[0014] In a preferred embodiment, Form A crystals are mixed with formulating excipients such as one or more bulking agents, for example lactose, one or more disintegrants, such as croscopovidone, and wet granulated to directly form Pattern C material in admixture with the excipients.


[0015] Further in accordance with the present invention, a new form of atazanavir bisulfate is provided, namely, Form E3 which is a highly crystalline form of the triethanolate solvate of atazanavir bisulfate.

[0016] Form E3 is prepared by slurrying atazanavir free base in ethanol, treating the slurry with concentrated sulfuric acid, heating and seeding the resulting solution with ethanol wet E3 crystals, treating the mixture with heptane (or other solvent such as toluene or hexane), filtering and drying.

[0017] Still further in accordance with the present invention, a process is provided for preparing Form A crystals of atazanavir bisulfate which includes the steps of preparing a triamine salt of the structure



(preferably the HCl (3 moles) salt) and without isolating the triamine salt, reacting the triamine salt with an active ester, preferably of the structure



in the presence of a base and organic solvent to form atazanavir free base which, without isolating, is converted to the atazanavir bisulfate via a modified cubic crystallization technique as described herein.

[0018] In addition, an atazanavir bisulfate composition is described which includes atazanavir bisulfate as Form A crystals or Pattern C material, and a pharmaceutically acceptable carrier therefor. The pharmaceutically acceptable carrier may include fillers, binders, disintegrants, lubricants, and other conventional excipients.

[0019] The various forms of atazanavir bisulfate according to the invention may be characterized using various techniques, the operation of which are well known to those of ordinary skill in the art. The forms may be characterized and distinguished using single crystal X-ray diffraction, which is based on unit cell measurements of a single crystal of a form at a fixed analytical temperature. A detailed description of unit cells is provided in Stout & Jensen, X-Ray Structure Determination: A Practical Guide, Macmillan Co., New York (1968), Chapter 3, which is herein incorporated by reference. Alternatively, the unique arrangement of atoms in spatial relation within the crystalline lattice may be characterized according to the observed fractional atomic coordinates. Another means of characterizing the crystalline structure is by

powder X-ray diffraction analysis in which the experimental or observed diffraction profile is compared to a simulated profile representing pure powder material, both run at the same analytical temperature, and measurements for the subject form characterized as a series of 2θ values.

[0020] Other means of characterizing the form may be used, such as solid state nuclear magnetic resonance (SSNMR), differential scanning calorimetry (DSC) and thermal gravimetric analysis (TGA). These parameters may also be used in combination to characterize the subject form.

[0021] Form A crystals may be characterized by unit cell parameters substantially equal to the following:

Cell dimensions:

[0022]

$a = 9.86(5) \text{ \AA}$
 $b = 29.245(6) \text{ \AA}$
 $c = 8.327(2) \text{ \AA}$
 $\alpha = 93.56(2)^\circ$
 $\beta = 114.77(3)^\circ$
 $\gamma = 80.49(3)^\circ$

Space group 1

Molecules/asymmetric unit 2

wherein the crystalline form is at about +22°C.

[0023] Form A may be characterized by fractional atomic coordinates substantially as listed in Table 3 and the crystal structure substantially as shown in Figure 2.

[0024] Form A may be characterized by simulated and observed powder X-ray diffraction patterns substantially as shown in Figure 1.

[0025] Form A may be characterized by a differential scanning calorimetry (DSC) thermogram having an endotherm with peak onset at about 165.6°C substantially as shown in Figure 3.

[0026] Form A may be characterized by a thermal gravimetric analysis (TGA) curve having a negligible weight loss up to about 100°C to 150°C substantially as shown in Figure 4.

[0027] Form A may be characterized by the solid-state NMR (SSNMR) chemical shifts substantially as shown in Table 4 and by the spectrum substantially as shown in Figure 5.

[0028] Form A may be characterized by fractional atomic coordinates substantially as listed in Table 5.

[0029] Form A salt may be characterized by moisture-sorption isotherms with about 0.1% weight gain in the range from 25 to 75% RH at 25°C.

[0030] In one aspect of the present invention, Pattern C may be characterized by the observed powder X-ray diffraction pattern substantially as shown in Figure 6.

[0031] In a different aspect of the present invention, Pattern C may be characterized by a differential scanning calorimetry thermogram substantially as shown in Figure 7 having an endotherm typically in the range from about 76.7 to about 96.6°C and from about 156.8 to about 165.9°C.

[0032] In a different aspect of the present invention, Pattern C may be characterized by a thermal gravimetric analysis curve having a weight loss of about 2.4% at about 125°C and a weight loss of about 4.4% up to about 190°C substantially as shown in Figure 8.

[0033] In accordance with the present invention, Form E3 may be characterized by crystallographic data as shown in Table 5, substantially equal to the following:

$a = 10.749 \text{ \AA}$
 $b = 13.450(4) \text{ \AA}$
 $c = 9.250(2) \text{ \AA}$
 $\alpha = 98.33(2)^\circ$
 $\beta = 95.92(3)^\circ$
 $\gamma = 102.82(3)^\circ$

Space group P1

Molecules/asymmetric unit 1

when the crystalline form is at about -23°C.

[0034] In a different aspect of the present invention, Form E3 may be characterized by fractional atomic coordinates substantially as listed in Table 6.

[0035] In a different aspect of the present invention, Form E3 may be characterized by simulated and observed powder X-ray diffraction patterns substantially as shown in Figure 9.

[0036] In a different aspect of the present invention, Form E3 may be characterized by a differential scanning calorimetry thermogram having an endotherm typically within the range from about 89.4 to about 96.6°C substantially as shown in Figure 11.

[0037] In a different aspect of the present invention, Form E3 may be characterized by a thermal gravimetric analysis curve having a weight loss of about 14.7% at about 150°C substantially as shown in Table 8.

[0038] In a different aspect of the invention, Form E3 may be characterized by the crystal structure substantially as shown in Figure 10.

BRIEF DESCRIPTION OF THE FIGURES

[0039]

Figure 1 shows calculated (simulated) (22°C) and observed (experimental at room temperature) powder X-ray diffraction patterns ($\text{CuK}\alpha$ $\lambda = 1.5418 \text{ \AA}$) of Form A;

Figure 2 shows the crystal structure of Form A;

Figure 3 shows a differential scanning calorimetry (DSC) thermogram of Form A;

Figure 4 shows a thermal gravimetric analysis curve (TGA) of Form A;

Figure 5 shows a C-13 solid state NMR of Form A;

Figure 6 shows an observed (experimental at room temperature) powder X-ray diffraction pattern ($\text{CuK}\alpha$ $\lambda = 1.5418 \text{ \AA}$) of Pattern C;

Figure 7 shows a differential scanning calorimetry thermogram of Pattern C;

Figure 8 shows a thermal gravimetric analysis curve of Pattern C;

Figure 9 shows calculated (simulated) (22°C) and observed (experimental at room temperature) powder X-ray diffraction patterns ($\text{CuK}\alpha$ $\lambda = 1.5418 \text{ \AA}$) of Form E3;

Figure 10 shows the crystal structure of Form E3; and

Figure 11 shows a differential scanning calorimetry (DSC) thermogram of Form E3, and a thermal gravimetric analysis curve of Form E3.

DETAILED DESCRIPTION OF THE INVENTION

[0040] The present invention provides, at least in part, forms of atazanavir bisulfate, namely, Form E3 and Pattern C, as novel materials, in particular in pharmaceutically acceptable form. The term "pharmaceutically acceptable", as used herein, refers to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem complications commensurate with a reasonable benefit/risk ratio. In certain preferred embodiments, crystalline forms of free base I and salts thereof are in substantially pure form. The term "substantially pure", as used herein, means a compound having a purity greater than about 90% including, for example, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, and about 100%.

[0041] As used herein "polymorph" refers to crystalline forms having the same chemical composition but different spatial arrangements of the molecules, atoms, and/or ions forming the crystal.

[0042] As used herein "solvate" refers to a crystalline form of a molecule, atom, and/or ions that further contains molecules of a solvent or solvents incorporated into the crystalline structure. The solvent molecules in the solvate may be present in a regular arrangement and/or a non-ordered arrangement. The solvate may contain either a stoichiometric or nonstoichiometric amount of the solvent molecules. For example, a solvate with a nonstoichiometric amount of solvent molecules may result from partial loss of solvent from the solvate.

[0043] Samples of the crystalline forms may be provided with substantially pure phase homogeneity, indicating the presence of a dominant amount of a single crystalline form and optionally minor amounts of one or more other crystalline forms. The presence of more than one crystalline form in a sample may be determined by techniques such as powder X-ray diffraction (PXRD) or solid state nuclear magnetic resonance spectroscopy (SSNMR). For example, the presence of extra peaks in the comparison of an experimentally measured PXRD pattern with a simulated PXRD pattern may indicate more than one crystalline form in the sample. The simulated PXRD may be calculated from single crystal X-ray data. see Smith, D.K., "A FORTRAN Program for Calculating X-Ray Powder Diffraction Patterns," Lawrence Radiation Laboratory, Livermore, California, UCRL-7196 (April 1963). Preferably, the crystalline form has substantially pure phase homogeneity as indicated by less than 10%, preferably less than 5 %, and more preferably less than 2 % of the total peak area in the experimentally measured PXRD pattern arising from the extra peaks that are absent from the simulated PXRD pattern. Most preferred is a crystalline form having substantially pure phase homogeneity with less than 1% of

the total peak area in the experimentally measured PXRD pattern arising from the extra peaks that are absent from the simulated PXRD pattern.

[0044] Procedures for the preparation of crystalline forms are known in the art. The crystalline forms may be prepared by a variety of methods, including for example, crystallization or recrystallization from a suitable solvent, sublimation, growth from a melt, solid state transformation from another phase, crystallization from a supercritical fluid, and jet spraying. Techniques for crystallization or recrystallization of crystalline forms from a solvent mixture include, for example, evaporation of the solvent, decreasing the temperature of the solvent mixture, crystal seeding a supersaturated solvent mixture of the molecule and/or salt, freeze drying the solvent mixture, and addition of antisolvents (countersolvents) to the solvent mixture.

[0045] Crystals of drugs, including polymorphs, methods of preparation, and characterization of drug crystals are discussed in Solid-State Chemistry of Drugs, S.R. Bym, R.R. Pfeiffer, and J.G. Stowell, 2nd Edition, SSCI, West Lafayette, Indiana (1999).

[0046] For crystallization techniques that employ solvent, the choice of solvent or solvents is typically dependent upon one or more factors, such as solubility of the compound, crystallization technique, and vapor pressure of the solvent. Combinations of solvents may be employed, for example, the compound may be solubilized into a first solvent to afford a solution, followed by the addition of an antisolvent to decrease the solubility of the compound in the solution and to afford the formation of crystals. An antisolvent is a solvent in which the compound has low solubility. Suitable solvents for preparing crystals include polar and nonpolar solvents.

[0047] In one method to prepare crystals, atazanavir bisulfate is suspended and/or stirred in a suitable solvent to afford a slurry, which may be heated to promote dissolution. The term "slurry", as used herein, means a saturated solution of atazanavir bisulfate or a salt thereof, which may also contain an additional amount of atazanavir bisulfate or salt thereof to afford a heterogeneous mixture of atazanavir bisulfate or salt thereof and a solvent at a given temperature. Suitable solvents in this regard include, for example, polar aprotic solvents, and polar protic solvents, and mixtures of two or more of these as disclosed herein.

[0048] Seed crystals may be added to any crystallization mixture to promote crystallization. As will be clear to the skilled artisan, seeding is used as a means of controlling growth of a particular crystalline form or as a means of controlling the particle size distribution of the crystalline product. Accordingly, calculation of the amount of seeds needed depends on the size of the seed available and the desired size of an average product particle as described, for example, in "Programmed cooling of batch crystallizers," J.W. Mullin and J. Nyvlt, Chemical Engineering Science (1971) 26:369-377. In general, seeds of small size are needed to effectively control the growth of crystals in the batch. Seeds of small size may be generated by sieving, milling, or micronizing of larger crystals, or by micro-crystallization of solutions. Care should be taken that milling or micronizing of crystals does not result in any change in crystallinity from the desired crystal form (i.e. change to amorphous or to another polymorph).

[0049] A cooled mixture may be filtered under vacuum, and the isolated solids may be washed with a suitable solvent, such as cold recrystallization solvent, and dried under a nitrogen purge to afford the desired crystalline form. The isolated solids may be analyzed by a suitable spectroscopic or analytical technique, such as SSNMR, DSC, PXRD, or the like, to assure formation of the preferred crystalline form of the product. The resulting crystalline form is typically produced in an amount of greater than about 70 weight % isolated yield, but preferably greater than 90 weight % based on the weight of atazanavir bisulfate originally employed in the crystallization procedure. The product may be comilled or passed through a mesh screen to delump the product, if necessary.

[0050] Crystalline forms may be prepared directly from the reaction medium of the final process step for preparing atazanavir bisulfate. This may be achieved, for example, by employing in the final process step a solvent or mixture of solvents from which atazanavir bisulfate may be crystallized. Alternatively, crystalline forms may be obtained by distillation or solvent addition techniques. Suitable solvents for this purpose include any of those solvents described herein, including protic polar solvents such as alcohols, and aprotic polar solvents such as ketones.

[0051] By way of general guidance, the reaction mixture may be filtered to remove any undesired impurities, inorganic salts, and the like, followed by washing with reaction or crystallization solvent. The resulting solution may be concentrated to remove excess solvent or gaseous constituents. If distillation is employed, the ultimate amount of distillate collected may vary, depending on process factors including, for example, vessel size, stirring capability, and the like. By way of general guidance, the reaction solution may be distilled to about {fraction (1/10)} the original volume before solvent replacement is carried out. The reaction may be sampled and assayed to determine the extent of the reaction and the wt % product in accordance with standard process techniques. If desired, additional reaction solvent may be added or removed to optimize reaction concentration. Preferably, the final concentration is adjusted to about 50 wt % at which point a slurry typically results.

[0052] It may be preferable to add solvents directly to the reaction vessel without distilling the reaction mixture. Preferred solvents for this purpose are those which may ultimately participate in the crystalline lattice as discussed above in connection with solvent exchange. Although the final concentration may vary depending on desired purity, recovery and the like, the final concentration of free base I in solution is preferably about 4% to about 7%. The reaction mixture may

be stirred following solvent addition and simultaneously warmed. By way of illustration, the reaction mixture may be stirred for about 1 hour while warming to about 70°C. The reaction is preferably filtered hot and washed with either the reaction solvent, the solvent added or a combination thereof. Seed crystals may be added to any crystallization solution to initiate crystallization.

[0053] The various forms described herein may be distinguishable from one another through the use of various analytical techniques known to one of ordinary skill in the art. Such techniques include, but are not limited to, solid state nuclear magnetic resonance (SSNMR) spectroscopy, X-ray powder diffraction (PXRD), differential scanning calorimetry (DSC), and/or thermogravimetric analysis (TGA).

[0054] One of ordinary skill in the art will appreciate that an X-ray diffraction pattern may be obtained with a measurement error that is dependent upon the measurement conditions employed. In particular, it is generally known that intensities in a X-ray diffraction pattern may fluctuate depending upon measurement conditions employed and the shape or morphology of the crystal. It should be further understood that relative intensities may also vary depending upon experimental conditions and, accordingly, the exact order of intensity should not be taken into account. Additionally, a measurement error of diffraction angle for a conventional X-ray diffraction pattern is typically about 0.2% or less, preferably about 0.1% (as discussed hereinafter), and such degree of measurement error should be taken into account as pertaining to the aforementioned diffraction angles. Consequently, it is to be understood that the crystal forms of the instant invention are not limited to the crystal forms that provide X-ray diffraction patterns completely identical to the X-ray diffraction patterns depicted in the accompanying Figures disclosed herein. Any crystal forms that provide X-ray diffraction patterns substantially identical to those disclosed in the accompanying Figures fall within the scope of the present invention. The ability to ascertain substantial identities of X-ray diffraction patterns is within the purview of one of ordinary skill in the art.

[0055] The term "Form" as used herein with respect to Form A and Form E3 refers to a homogeneous crystal structure.

[0056] The term "Pattern" as used herein with respect to Pattern C material refers to a characteristic x-ray diffraction pattern.

[0057] The term "atazanavir bisulfate" as employed herein refers to atazanavir bisulfate as well as atazanavir sulfate.

[0058] In carrying out the process of the invention for preparing Form A crystals of atazanavir bisulfate salt, a modified cubic crystallization technique is employed wherein atazanavir free base is dissolved in an organic solvent in which the atazanavir bisulfate salt is substantially insoluble and which is acetone, or a mixture of acetone and N-methyl pyrrolidone, to provide a solution having a concentration of atazanavir free base within the range from about 6.5 to about 9.7% by weight, preferably from about 6.9 to about 8.1% by weight atazanavir free base.

[0059] The solution of atazanavir free base is heated at a temperature within the range from 35 to 55°C, preferably from 40 to 50°C, and reacted with an amount of concentrated sulfuric acid (containing from about 95 to about 100% H₂SO₄) to react with less than 15%, preferably from 5 to less than 12%, more preferably from 8 to 10% by weight of the total atazanavir free base. Thus, the starting solution of atazanavir free base will be initially reacted with less than 15%, preferably from 5 to 12%, by weight of the total amount of sulfuric acid to be employed. During the reaction, the reaction mixture is maintained at a temperature within the range from 35 to 55°C, preferably from 40 to 50°C.

[0060] The reaction is allowed to continue for a period from about 12 to about 60 minutes, preferably from about 15 to about 30 minutes.

[0061] The reaction mixture is seeded with crystals of Form A atazanavir bisulfate employing an amount of seeds within the range from 0.1 to 80% by weight, preferably from 3 to 8% by weight, based on the weight of atazanavir free base remaining in the reaction mixture while maintaining the reaction mixture at a temperature within the range from 35 to 55°C, preferably from 40 to 50°C.

[0062] The reaction is allowed to continue until crystallization begins. Thereafter, sulfuric acid is added in multiple stages at an increasing rate according to the cubic equation as described below to form atazanavir bisulfate which upon drying produces Form A crystals.

[0063] The crystal particle size and morphology of the atazanavir bisulfate salt formed are dependent on the addition rate of the sulfuric acid, which determines the crystallization rate. It has been found that a modified "cubic" crystallization technique (acid added at an increasing rate according to a cubic equation) provides relatively larger, more well defined atazanavir bisulfate crystals, along with a narrower particle size range and fewer fines, than a constant addition rate crystallization. The slow initial acid flow rate has been shown to favor crystal growth over secondary nucleation. Thus, as the surface area increases with particle size, the seed bed is able to accept the increasing acid flow rate without inducing secondary nucleation. The slow initial addition rate allows time for the crystals to grow larger, increasing the mean size. The cubic crystallization provides a less compressible filter cake, which aids in effective cake deliquoring and washing, as well as giving a more easily dried product with fewer hard lumps than the constant addition rate crystallized product.

[0064] The cubic crystallization method employed is a temperature controlled crystallization derived from Mullin, "Crystallization, 3rd Ed.", 1993, Butterworth-Heinemann, Pubs. and is defined by the following simplified equation:

$$T = T_{\max} - (T_{\max} - T_{\min}) \times \left[\frac{\text{time}}{\text{time}_{\text{total}}} \right]^3 \quad (1)$$

where

$T_{\max}$ = Starting temperature for crystallization

$T_{\min}$ = Ending temperature for crystallization

time = Elapsed time in crystallization

time_{total} = Total crystallization time.

[0065] Since the crystallization of atazanavir bisulfate is controlled by the addition rate of sulfuric acid, the temperature variable is replaced with acid volume in Equation (1). In this equation, the variable representing the minimum volume is removed.

$$V_{\text{time}} = V_{\text{total}} \times \left[\frac{\text{time}}{\text{time}_{\text{total}}} \right]^3 \quad (2)$$

where

V_{time} = Volume of sulfuric acid added during elapsed time period

V_{total} = Total volume of acid representing the 90% charge

time = Elapsed time in crystallization

time_{total} = Total crystallization time.

Equation (2) is referred to as "the cubic equation."

[0066] By controlling the crystallization rate using this expression, nucleation is controlled within acceptable limits as the system maintains a constant low level of supersaturation.

[0067] Form A crystals are identified by powder x-ray diffraction pattern and crystal structure as shown in Figures 1 and 2, respectively.

[0068] The Form A crystals of atazanavir bisulfate or the Pattern C material as well as Form E3 prepared as described above are the final atazanavir bisulfate and can be employed as drug products for administration to patients.

[0069] In accordance with the process of the invention, Pattern C material may be prepared by exposing Form A crystals to water followed by drying.

[0070] In another process in accordance with the present invention, Pattern C material may be formed by exposing crystals of Form A to high relative humidity of greater than about 95% RH, preferably from about 95 to about 100% RH (water vapor), for at least 24 hours, preferably from about 24 to about 48 hours.

[0071] In another embodiment of the invention, Pattern C material is prepared by wet granulating atazanavir bisulfate Form A to produce granules of atazanavir bisulfate and then drying the granules.

[0072] In carrying out the wet granulation process, the atazanavir bisulfate will be granulated in water and dried at a temperature within the range from about 40 to about 80°C, preferably within the range from about 50 to about 60°C. The drying step will be preferably carried out for at least about 2 hours, up to about 20 hours, preferably from about 8 to about 10 hours.

[0073] The Pattern C material may also be formed by wet granulating atazanavir bisulfate Form A in the presence of conventional pharmaceutical excipients, for example, one or more bulking agents, preferably lactose, one or more disintegrants, preferably croscopovidone, and drying as described above to form Pattern C material in admixture with the excipients.

[0074] It is the Pattern C material, Form A or Form E3, preferably Pattern C material, which is formulated for administration in the treatment of diseases caused by viruses as described hereinafter.


[0075] Pattern C material is characterized by its powder x-ray diffraction pattern as shown in Figure 3.

[0076] The Form E3 is prepared by slurrying atazanavir free base in ethanol, treating the slurry with concentrated sulfuric acid employing a molar ratio of acid: free base with the range from about 1:1 to about 1.1:1, heating the resulting solution at from about 30 to about 40°C, seeding the solution with ethanol wet E3 crystals of atazanavir sulfate, treating the mixture with heptane (or other solvent such as hexane or toluene), filtering, and drying to yield atazanavir bisulfate Form E3 (triethanol solvate).


[0077] The seeding step will employ an amount of seeds to effect formation of E3 crystals, for example a molar ratio of atazanavir bisulfate E-3 seeds:free base within the range from about 0.02:1 to about 0.04:1.

[0078] Form E3 is identified by powder x-ray diffraction pattern as shown in Figure 7 and crystal structure as shown in Figure 6.


[0079] In accordance with the present invention, the atazanavir in the form of its free base is prepared by treating a solution of a protected triamine salt of the structure




(where PG represents a protecting group such as t-butyloxycarbonyl (Boc) or trifluoroacetyl, preferably Boc, with an acid, preferably hydrochloric acid (where Boc is used), or a base (where trifluoroacetyl is used) in the presence of an organic solvent such as methylene chloride, tetrahydrofuran, or methanol, which solvent is preferably methylene chloride, at a temperature within the range from about 25 to about 50°C, preferably from about 30 to about 40°C, to form the triamine acid salt, preferably the hydrogen chloride salt of the structure



and without isolating the triamine acid salt, reacting the triamine acid salt with an active ester of an acid of the structure




preferably the active ester of the structure




in the presence of a base such as K_2HPO_4 , diisopropylethylamine, N-methylmorpholine, sodium carbonate, or potassium

carbonate, preferably K_2HPO_4 , in the presence of an organic solvent such as methylene chloride, a mixture of ethyl acetate and butyl acetate, acetonitrile or ethyl acetate, preferably methylene chloride, at a temperature within the range from 25 to 50°C, preferably from 30 to 40°C to form atazanavir free base.

[0080] The protected triamine starting material is prepared by reacting the epoxide



where PG is preferably Boc such as N-(tert-butyloxycarbonyl)-2(S)-amino-1-phenyl-3(R)-3,4-epoxy-butane, with the hydrazine carbamate



where PG is preferably Boc in the presence of isopropyl alcohol or other alcohol such as ethanol or butanol.

[0081] Atazanavir bisulfate is useful for administration to a warm-blooded animal, especially a human being, for the treatment or prevention of a disease that is responsive to inhibition of a retroviral protease, especially a retroviral aspartate protease, such as HIV-1 or HIV-II gag protease, for example a retroviral disease, such as AIDS or its preliminary stages.

[0082] Atazanavir bisulfate, especially Pattern C material, Form A or Form E3, preferably Pattern C material or Form A, may be used for treating diseases caused by viruses, especially by retroviruses, especially AIDS or its preliminary stages, wherein a therapeutically effective amount of atazanavir bisulfate Pattern C material, Form A or Form E3 is administered in a dose that is effective in the treatment of said disease especially to a warm-blooded animal, for example a human being, who on account of one of the mentioned diseases, especially AIDS or its preliminary stages, requires such treatment. The preferred dose to be administered to warm-blooded animals, for example human beings of approximately 70 kg body weight, is from about 3 mg to about 1.5 g, preferably from about 10 mg to about 1.25 g, for example from about 50 mg to about 600 mg per person per day, divided preferably into 1 to 4 single doses which may, for example, be of the same size. Usually, children receive half of the adult dose. It is preferably administered orally.

[0083] Atazanavir bisulfate Pattern C material, Form A or Form E3 is employed for the above described pharmaceutical uses. Suitable compositions containing Pattern C material or Form A or Form E3 for oral administration include tablets, powders, capsules, and elixirs. About 10 to 600 mg of active ingredient is compounded with physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, flavoring, etc., in a unit dose form as called for by accepted pharmaceutical practice.

[0084] Pharmaceutical compositions for oral administration can be obtained by combining the active ingredient with solid carriers, if desired granulating a resulting mixture, and processing the mixture, if desired or necessary, after the addition of appropriate excipients, into tablets, dragée cores, capsules or powders for oral use. It is also possible for the active ingredients to be incorporated into plastic carriers that allow the active ingredients to diffuse or be released in measured amounts.

[0085] The bulking agents or fillers will be present in the pharmaceutical compositions of the invention in an amount within the range from about 0 to about 95% by weight and preferably from about 10 to about 85% by weight of the composition. Examples of bulking agents or fillers suitable for use herein include, but are not limited to, cellulose derivatives such as microcrystalline cellulose or wood cellulose, lactose, sucrose, starch, pregelatinized starch, dextrose, mannitol, fructose, xylitol, sorbitol, corn starch, modified corn starch, inorganic salts such as calcium carbonate, calcium phosphate, dicalcium phosphate, calcium sulfate, dextrin/dextrates, maltodextrin, compressible sugars, and other known bulking agents or fillers, and/or mixtures of two or more thereof, preferably lactose.

[0086] A binder will be optionally present in the pharmaceutical compositions of the invention in an amount within the range from about 0 to about 20% weight, preferably from about 1 to about 10% by weight of the composition. Examples of binders suitable for use herein include, but are not limited to, hydroxypropyl cellulose, corn starch, pregelatinized starch, modified corn starch, polyvinyl pyrrolidone (PVP) (molecular weight ranging from about 5,000 to about 80,000, preferably about 40,000), hydroxypropylmethyl cellulose (HPMC), lactose, gum acacia, ethyl cellulose, cellulose acetate, as well as a wax binder such as carnauba wax, paraffin, spermaceti, polyethylenes or microcrystalline wax, as well as other conventional binding agent and/or mixtures by two or more thereof, preferably hydroxypropyl cellulose.

[0087] The disintegrant will be optionally present in the pharmaceutical composition of the invention in an amount within the range from about 0 to about 20% by weight, preferably from about 0.25 to about 15% by weight of the

composition. Examples of disintegrants suitable for use herein include, but are not limited to, croscarmellose sodium, crospovidone, potato starch, pregelatinized starch, corn starch, sodium starch glycolate, microcrystalline cellulose, or other known disintegrant, preferably croscarmellose sodium.

[0088] The lubricant will be optionally present in the pharmaceutical composition of the invention in an amount within the range from about 0.1 to about 4% by weight, preferably from about 0.2 to about 2% by weight of the composition. Examples of tableting lubricants suitable for use herein include, but are not limited to, magnesium stearate, zinc stearate, calcium stearate, talc, carnauba wax, stearic acid, palmitic acid, sodium stearyl fumarate or hydrogenated vegetable oils and fats, or other known tableting lubricants, and/or mixtures of two or more thereof, preferably magnesium stearate.

[0089] Capsules are hard gelatin capsules and also soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The hard gelatin capsules may include the active ingredient in the form of granules, for example with fillers, such as lactose, binders, such as starches, crospovidone and/or glidants, such as talc or magnesium stearate, and if desired with stabilizers. In soft gelatin capsules the active ingredient is preferably dissolved or suspended in suitable oily excipients, such as fatty oils, paraffin oil or liquid polyethylene glycols, it likewise being possible for stabilizers and/or antibacterial agents to be added.


[0090] The following examples represent preferred embodiments of the invention.

EXAMPLE 1

1-[4-(Pyridin-2-yl)phenyl]-5(S)-2,5-bis[[N-(methoxycarbonyl)-L-tert-leucyl]amino]-4-(S)-hydroxy-6-phenyl-2-azahexane, Bisulfate salt (Form A) (Atazanavir bisulfate - Form A)


A.

[0091]



(1-[4-(Pyridin-2-yl)phenyl]-5(S)-2,5-bis[tert-butyloxycarbonyl]amino]-4(S)-hydroxy-6-phenyl-2-azahexane.3HCl (Triamine.3HCl Salt))

[0092] To a 1000 mL, 3-neck, round-bottom flask fitted with mechanical stirrer, nitrogen inlet and temperature probe was added the protected triamine 1-[4-(pyridin-2-yl)phenyl]-5(S)-2,5-bis[tert-butyloxycarbonyl]amino]-4(S)-hydroxy-6-phenyl-2-azahexane



(100 g, 0.178 mol), and CH_2Cl_2 (500 mL; 5 mL/g of protected triamine input) (prepared as described in Z. Xu et al., Process Research and Development for an Efficient Synthesis of the HIV Protease Inhibitor BMS-232,632, Organic


Process Research and Development, 6, 323-328 (2002)) and the resulting slurry was agitated while maintaining the temperature at from about 5 to about 22°C.

[0093] Concentrated hydrochloric acid (68 mL, 0.82 mole, 4.6 eq.) was added to the reaction mixture at a rate such that the temperature of the reaction mixture remained between 5 and 30°C. The reaction mixture was heated to 30 to 40°C and agitated until the reaction was judged complete by HPLC assay.


[0094] Water was added (70-210 mL, 0.7-2.1 mL/g protected triamine input) to the reaction mixture, the reaction mixture was agitated for 15 minutes and the phases were allowed to separate. The upper, product (triamine.3HCl salt)-rich aqueous oil was transferred to an addition funnel.

B.

[0095]



(Active Ester of N-methoxycarbonyl-L-tert-leucine)



[0096] To a 3000 mL, 3-neck round bottom flask fitted with mechanical stirrer, addition funnel, nitrogen inlet, and temperature probe was added N-methoxycarbonyl-L-tert-leucine (77.2g, 0.408 mol, 2.30 eq.), 1-hydroxybenzotriazole (HOBT) (60.8 g, 0.450 mol, 2.53 eq.), and N-ethyl N'-dimethylaminopropyl carbodiimide (EDAC) (82.0 g, 0.430 mol, 2.42 eq.), followed by CH₂Cl₂ (880 mL; 8.8 mL/g of protected triamine input) and the mixture was stirred at ambient temperature (18-25°C) until formation of the active ester is complete, as judged by HPLC.

C. 1-[4-(Pyridin-2-yl)phenyl]-5(S)-2,5-bis{[N-(methoxycarbonyl)-L-tert-leucinyl]amino}-4(S)-hydroxy-6-phenyl-2-azahexane (atazanavir free base)

[0097] Anhydrous dibasic potassium phosphate (K₂HPO₄; 226 g., 1.30 mol, 7.30 eq. wrt protected triamine) was dissolved in 1130 mL of water (11.3 mL/g of protected amine; 5 mL/g of K₂HPO₄).

[0098] The K₂HPO₄ solution was added to the active ester solution prepared in Part B. To the stirred active ester/aqueous K₂HPO₄ mixture was slowly added the aqueous solution of Part A hydrogen chloride salt over a period of 1.5 to 2.0 h while maintaining agitation and a pot temperature between 5 and 20°C.

[0099] After the addition of the solution of the Part A hydrogen chloride salt was complete, the reaction mixture (coupling reaction) was heated to 30-40°C and agitated until the coupling reaction was judged complete by HPLC assay.

[0100] The coupling mixture was cooled to 15 to 20°C and the lower, product rich organic phase was separated from the upper, spent aqueous phase.

[0101] The product rich organic phase was washed with 1M NaH₂PO₄ (880 mL; pH=1.5; 8.8 mL/g of protected triamine input; 5 mole eq. wrt protected triamine), the phases were allowed to separate, and the spent aqueous phase was removed.

[0102] The washed product rich organic phase was stirred with 0.5 N NaOH (800 mL; 8 mL/g of protected triamine input) until HPLC assay of the rich organic phase showed the active esters to be below 0.3 I.I. each. The phases were allowed to separate and the spent aqueous phase was removed.

[0103] The rich organic phase was washed with 5% NaH₂PO₄ (450 mL, 4.5 mL/g of protected triamine input; pH=4.3), the phases were allowed to separate and the spent aqueous phase was removed.

[0104] The rich organic phase was washed with 10 w/v% NaCl (475 mL, 4.75 mL/g of protected triamine input) and

the spent aqueous phase was removed.

[0105] The concentration of title free base in solution was 120 to 150 mg/mL with an in-process calculated yield of 95-100 mol%.

D. Solvent Exchange from CH₂Cl₂ into Acetone/N-Methylpyrrolidone

[0106] To the rich Part C free base solution in a 3000 mL, 3-neck round-bottom flask fitted with mechanical stirrer, temperature probe, and distillation condenser, was added N-methylpyrrolidone (148 mL; 1.25 mL/g of Part C free base based on in-process quantification assay). The solution was concentrated to ca. 360 mL (2.5-3.5 mL/g of Part C free base) using a jacket temperature of 70°C or less; 500 mL of acetone (4-5 mL/g of Part C free base) was added to the concentrated solution and the mixture was distilled to a volume of about 400 mL or less.

[0107] The acetone addition and distillation were repeated until in-process assay indicated the CH₂Cl₂ level had reached the target endpoint. At crystallization volume, the CH₂Cl₂ content in the rich organic solution was 0.77 v/v %. Acetone was added to the concentrated free base solution to reach a total solution of 16 mL/g of free base. The bath temperature was maintained at 40-50°C to prevent crystallization of free base. The solution was polish filtered through a 10-micron or finer filter while maintaining the temperature at 40 to 50°C. The polish filter was rinsed with acetone (125 mL, 1.0 mL/g of free base) and the rinse was added to the rich free base acetone/N-methylpyrrolidone solution which was used in the next step.

E. 1-[4-(Pyridin-2-yl)phenyl]-5(S)-2,5-bis[[N-(methoxycarbonyl)-L-tert-leuciny]amino]-4(S)-hydroxy-6-phenyl-2-azaheptane bisulfate salt

[0108] About 10% (2 g) of the total charge of concentrated sulfuric acid (19 g, 1.10 eq.) was added to the free base acetone/N-methylpyrrolidone solution of Part D, while maintaining the temperature at 40-50°C, via subsurface addition.

[0109] The reaction mixture was seeded with 5.0 wt % (wrt calculated free base in solution) of bisulfate salt. The seeded mixture was agitated at 40-50°C for at least 30 minutes during which time the bisulfate salt began crystallizing as evidenced by the mixture increasing in opacity during this time.

[0110] The remaining sulfuric acid (17.8 g) was added over ca. 5 h in five stages according to the following protocol, defined by a cubic equation, while keeping the temperature at 40-50°C.

[0111] The rate of each addition stage was determined according to the cubic equation described hereinbefore and is shown in the table below.

TABLE 1

Stage	mL/kg/h	mL(H ₂ SO ₄)/h	g(H ₂ SO ₄)/h	Duration (min)
1	4.62	0.579	1.065	60
2	6.93	0.868	1.597	60
3	16.55	2.073	3.814	60
4	30.26	3.790	6.974	60
5	48.47	6.071	11.171	23

[0112] After addition of H₂SO₄ was complete, the slurry was cooled to 20-25°C for at least 1 h with agitation. The slurry was agitated at 20-25°C for at least 1 h. The bisulfate salt was filtered and the mother liquor was recycled as needed to effect complete transfer. The filter cake was washed with acetone (5-10 mL/g of free base; 1200 mL acetone). The bisulfate salt was dried at NMT 55°C under vacuum until the LOD <1% to produce a crystalline material.

[0113] The crystalline product was analyzed by PXRD, DSC and TGA patterns and SSNMR spectrum and found to be (non-solvated) Form A crystals of the title bisulfate (see Figures 1 to 5).

TABLE 2

Table of Crystallographic Data Form A											
T°C	a(Å)	b(Å)	c(Å)	α°	β°	γ°	V(Å ³)	Z'	sg	dcalc	R
+22	9.861(5)	29.245(6)	8.327(2)	93.56(2)	114.77(3)	80.49(3)	2150(2)	2	P1	1.240	0.06
T= temp(°C) for the crystallographic data. Z' = number of drug molecules per asymmetric unit											

TABLE 3

Table Of Fractional Parameters and Their Estimated Standard Deviations for Form A				
Atom	x	y	z	B(A ²)
S1	0.3230(4)	0.5467(1)	0.5608(5)	8.0(1)
O100	0.431(1)	0.5060(3)	0.649(1)	11.1(3)
O102	0.335(1)	0.5498(4)	0.383(1)	12.0(4)
O103	0.360(1)	0.5877(4)	0.655(2)	12.0(4)
O104	0.176(1)	0.5384(4)	0.528(1)	11.8(4)
S51	0.6177(4)	0.4505(1)	0.4003(5)	7.2(1)
O150	0.596(1)	0.4430(4)	0.564(1)	12.5(4)
O152	0.518(1)	0.4921(4)	0.317(1)	13.8(4)
O153	0.588(1)	0.4121(3)	0.289(2)	12.2(4)
O154	0.768(1)	0.4587(4)	0.454(1)	12.1(4)
O4	0.6985(7)	0.1753(3)	0.6456(9)	5.7(2)
O7	0.1687(8)	0.1941(3)	0.3411(9)	6.5(2)
O11	-0.0352(7)	0.2482(3)	0.0308(8)	5.7(2)
O14	0.2280(7)	0.1769(3)	-0.233(1)	6.1(2)
O15	0.0399(8)	0.1335(3)	-0.330(1)	6.4(2)
O17	0.6169(7)	0.2821(3)	0.963(1)	7.1(2)
O18	0.3750(7)	0.2905(3)	0.9136(9)	6.2(2)
N2	0.5015(9)	0.2182(3)	0.902(1)	4.5(2)
N5	0.4642(8)	0.1647(3)	0.6001(9)	4.2(2)
N9	0.2317(9)	0.2788(3)	0.256(1)	5.1(2)
N10	0.1820(9)	0.2760(3)	0.069(1)	4.6(2)
N13	-0.0148(8)	0.2083(3)	-0.280(1)	4.6(2)
N39	-0.087(1)	0.5265(3)	0.272(1)	6.1(3)
C1	0.491(1)	0.2627(4)	0.924(1)	5.5(3)
C3	0.6381(9)	0.1908(3)	0.892(1)	4.0(2)
C4	0.600(1)	0.1764(4)	0.702(1)	4.6(3)
C6	0.420(1)	0.1551(4)	0.403(1)	5.1(3)
C7	0.295(1)	0.1936(4)	0.297(1)	5.1(3)
C8	0.357(1)	0.2400(4)	0.346(2)	5.4(3)
C11	0.051(1)	0.2592(4)	-0.028(1)	4.9(3)
C12	0.024(1)	0.2531(4)	-0.223(1)	4.5(3)
C14	0.094(1)	0.1732(4)	-0.280(1)	4.7(3)
C16	0.146(2)	0.0943(5)	-0.342(2)	10.9(5)
C19	0.616(1)	0.3313(4)	0.996(2)	8.1(4)
C20	0.701(1)	0.1485(4)	1.025(1)	5.8(3)
C21	0.842(1)	0.1219(5)	1.007(2)	7.9(4)

EP 1 755 596 B9

(continued)

Table Of Fractional Parameters and Their Estimated Standard Deviations for Form A				
Atom	x	y	z	B(A ²)
C22	0.583(2)	0.1160(5)	0.997(2)	8.0(4)
C23	0.748(2)	0.1713(5)	1.215(1)	8.2(4)
C24	0.365(1)	0.1079(4)	0.356(2)	6.6(4)
C25	0.484(1)	0.0691(4)	0.470(1)	6.5(3)
C26	0.643(2)	0.0684(5)	0.520(2)	8.4(5)
C27	0.753(2)	0.0293(6)	0.622(2)	11.4(6)
C28	0.709(3)	-0.0044(7)	0.691(3)	15.0(9)
C29	0.553(2)	-0.0032(5)	0.644(2)	14.2(7)
C30	0.441(2)	0.0343(5)	0.534(2)	10.8(4)
C31	0.291(1)	0.3229(4)	0.311(2)	5.7(3)
C32	0.177(1)	0.3650(4)	0.259(1)	5.4(3)
C33	0.224(1)	0.4064(4)	0.262(2)	6.3(3)
C34	0.122(1)	0.4487(5)	0.233(2)	6.9(4)
C35	-0.031(1)	0.4469(4)	0.189(1)	4.8(3)
C36	-0.081(1)	0.4043(4)	0.180(1)	5.6(3)
C37	0.019(1)	0.3629(4)	0.218(1)	5.4(3)
C38	-0.136(1)	0.4918(4)	0.170(1)	5.3(3)
C40	-0.170(1)	0.5683(4)	0.279(2)	7.8(4)
C41	-0.318(2)	0.5736(5)	0.158(2)	9.1(5)
C42	-0.376(2)	0.5403(5)	0.035(2)	9.0(5)
C43	-0.283(1)	0.4964(5)	0.039(2)	8.1(4)
C44	-0.096(1)	0.2937(4)	-0.345(1)	6.2(3)
C45	-0.258(1)	0.2901(5)	-0.366(2)	8.5(4)
C46	-0.085(2)	0.2890(6)	-0.530(2)	10.8(5)
C47	-0.057(2)	0.3393(5)	-0.265(2)	8.9(5)
O54	0.2347(7)	0.8167(3)	0.8392(8)	5.3(2)
O57	0.7713(8)	0.7950(3)	1.0561(9)	5.9(2)
O61	0.9725(7)	0.7436(3)	0.9141(8)	5.3(2)
O64	0.7062(7)	0.8164(3)	0.427(1)	5.9(2)
O65	0.8911(8)	0.8598(2)	0.535(1)	6.1(2)
O67	0.3150(8)	0.7090(3)	1.184(1)	6.4(2)
O68	0.5587(9)	0.6986(3)	1.377(l)	6.6(2)
N52	0.4313(9)	0.7713(3)	1.271(1)	4.9(2)
N55	0.4709(8)	0.8265(3)	1.0332(9)	4.2(2)
N60	0.7555(8)	0.7179(3)	0.728(1)	4.6(2)
N63	0.9491(8)	0.7852(3)	0.601(1)	4.4(2)
N89	1.026(1)	0.4719(3)	0.711(1)	6.0(3)

EP 1 755 596 B9

(continued)

Table Of Fractional Parameters and Their Estimated Standard Deviations for Form A				
Atom	x	y	z	B(A ²)
C51	0.442(1)	0.7247(4)	1.282(1)	5.4(3)
C53	0.296(1)	0.7996(4)	1.141(1)	5.1(3)
C54	0.3347(9)	0.8159(3)	0.989(1)	4.1(3)
C56	0.519(1)	0.8353(4)	0.887(1)	4.7(3)
C57	0.644(1)	0.7959(4)	0.886(1)	4.5(3)
C58	0.587(1)	0.7494(4)	0.854(1)	5.2(3)
C61	0.884(1)	0.7334(4)	0.766(1)	4.2(3)
C62	0.914(1)	0.7392(4)	0.603(1)	4.4(3)
C64	0.839(1)	0.8196(4)	0.513(1)	4.6(3)
C66	0.785(2)	0.8996(5)	0.433(3)	12.1(7)
C69	0.323(1)	0.6588(4)	1.202(2)	8.8(5)
C70	0.237(1)	0.8409(4)	1.232(1)	5.6(3)
C71	0.092(1)	0.8701(5)	1.080(2)	7.6(4)
C72	0.352(1)	0.8744(4)	1.328(2)	7.1(4)
C73	0.187(1)	0.8195(6)	1.362(1)	8.9(4)
C74	0.570(1)	0.8825(4)	0.907(2)	6.4(3)
C75	0.450(1)	0.9206(4)	0.919(1)	6.3(3)
C76	0.296(2)	0.9236(5)	0.813(2)	8.1(4)
C77	0.188(2)	0.9614(6)	0.826(2)	11.2(5)
C78	0.244(2)	0.9942(6)	0.960(2)	15.2(7)
C79	0.405(3)	0.9935(6)	1.062(2)	13.9(7)
C80	0.504(2)	0.9552(4)	1.043(2)	9.3(5)
C81	0.644(1)	0.6672(4)	0.832(2)	6.2(3)
C82	0.762(1)	0.6266(3)	0.839(1)	4.7(3)
C83	0.723(1)	0.5934(4)	0.696(2)	6.1(3)
C84	0.822(1)	0.5547(4)	0.695(2)	5.9(3)
C85	0.967(1)	0.5478(4)	0.828(1)	5.0(3)
C86	1.009(1)	0.5783(4)	0.971(2)	6.6(4)
C87	0.908(1)	0.6184(4)	0.971(2)	6.4(4)
C88	1.076(1)	0.5070(4)	0.827(1)	5.5(3)
C90	1.111(1)	0.4326(4)	0.690(2)	7.4(4)
C91	1.258(2)	0.4262(5)	0.792(2)	17.8(4)
C92	1.324(2)	0.4578(5)	0.918(2)	8.7(5)
C93	1.230(1)	0.4994(5)	0.936(2)	6.9(4)
C94	1.038(1)	0.7005(4)	0.584(1)	4.8(3)
C95	1.196(1)	0.7055(4)	0.717(2)	6.7(4)
C96	1.021(2)	0.7049(5)	0.392(2)	8.9(4)

(continued)

Table Of Fractional Parameters and Their Estimated Standard Deviations for Form A				
Atom	x	y	z	B(A2)
C97	0.998(1)	0.6536(4)	0.614(2)	7.6(4)
N59	0.7084(8)	0.7114(3)	0.866(1)	5.1(2)
H391	0.047	0.523	0.383	6.0*
H891	0.931	0.477	0.646	5.8*
H15'	0.491	0.471	0.600	3.8*
H15"	0.440	0.512	0.322	4.6*

[0114] Most hydrogens have been omitted; only the hydrogens on N9 and the acid are included.

[0115] Anisotropically refined atoms are given in the form of the isotropic equivalent displacement parameter defined as: $(4/3) * [a^2 B(1,1) + b^2 B(2,2) + c^2 B(3,3) + ab(\cos \gamma) B(1,2) + ac(\cos \beta) B(1,3) + bc(\cos \alpha) B(2,3)]$.

[0116] Form A is characterized by a differential scanning calorimetry thermogram having an endotherm typically within the range from about 165.6°C to about 200.9°C as shown in Figure 3.

[0117] Form A is also characterized by a thermal gravimetric analysis curve having a negligible weight loss up to about 100 to 150°C.

[0118] The crystals produced by cubic crystallization where H₂SO₄ is added at an increasing rate according to the cubic equation described above were relatively larger and more well-defined, and had a narrower particle size range and fewer fines, than crystals obtained employing constant addition rate crystallization.

[0119] The filter cake obtained using the cubic crystallization technique was less compressible than that obtained using constant addition rate crystallization, which aided in effective cake deliquoring and washing and produced a homogeneous product.

TABLE 4

Carbon-13 SSNMR Chemical Shifts for Form A, measured relative to TMS (tetramethyl silane)	
δ /ppm	
26.9	
27.5	
33.9	
37.7	
49.2	
53.5	
62.7	
63.3	
66.0	
69.2	
69.5	
122.6	
123.7	
125.3	
126.1	
127.6	
128.5	

(continued)

Carbon-13 SSNMR Chemical Shifts for Form A, measured relative to TMS (tetramethyl silane)	
δ /ppm	
129.4	
131.1	
134.4	
138.8	
139.7	
140.6	
143.2	
143.9	
149.9	
150.3	
153.9	
159.3	
172.0	

EXAMPLE 2**Atazanavir Bisulfate - Pattern C Material****Method A:**

[0120] Form A crystals of atazanavir bisulfate (prepared as described in Example 1) (25.33 g) were suspended in 200 mL of water and the mixture was stirred mechanically to produce a thick gel which was dried.

[0121] The dried mixture was ground with a spatula to produce Pattern C material. A powder X-ray diffraction pattern of Pattern C material is shown in Figure 6.

Method B:

[0122] Form A crystals of atazanavir bisulfate was wet granulated using a sufficient amount of water (about 40% w/w) in a suitable mixer-granulator. The wet mass was dried in an oven. The product was sized using a suitable screen. The x-ray diffraction pattern of the resultant product is consistent with Pattern C material as shown in Figure 6.

[0123] Pattern C is characterized by the differential scanning calorimetry thermogram shown in Figure 7 having an endotherm typically in the range from about 76.7 to about 96.6°C and from about 156.8 to about 165.9°C.

[0124] Pattern C is also characterized by a thermal gravimetric analysis curve having a weight loss of about 2.4% at about 125°C and about 4.4% at about 190°C as shown in Figure 8.

EXAMPLE 3**Atazanavir Bisulfate - Form E3 (Triethanol Solvate)**

[0125] Atazanavir free base (prepared as described in Example 1, Part C) (3.0 g, 4.26 mmol) was slurried in dry, 200 proof ethanol (20.25 mL, 6.75 mL/g of free base) in a 100 mL, 3-neck round-bottom flask fitted with a mechanical stirrer, temperature probe, and a pressure-equalizing liquid addition funnel.

[0126] Concentrated H₂SO₄ (0.25 mL, 0.46 g, 4.69 mmol, 1.1 eq.) was added to the slurry of atazanavir free base which was maintained at 20-25°C. The resulting solution (KF of 0.2 to 1.0% water) was polish filtered (Whatman #1 paper), the filter rinsed with 2.25 mL of absolute ethanol and the rinse added to the filtered solution. The solution was heated to 37°C and seeded with 10 mg of amorphous atazanavir bisulfate derived from Form E3 crystals (by exposing Form E3 crystals to ambient temperature), and the mixture was agitated for 15 min. Heptane (380 mL, 8.25 mL/g of free

EP 1 755 596 B9

base) was added over 1 hour. The resulting crystallization mixture was agitated for 8 h at 15-25°C. Crystallized atazanavir bisulfate was filtered on a Büchner funnel. The product cake was washed with 184 mL (4 mL/g of free base) of 1:1 ethanol: heptane. The product cake was washed with 46 mL (1 mL/g of free base) of heptane. The resulting product was dried under vacuum at 40-50°C until it had an LOD = 0.97%. The yield of product was 47.7 g (0.0594 mol, 74.3 mol %) of atazanavir bisulfate Form E3 (triethanol solvate) with HPLC HI=100.0 (see Figures 9 and 10).

TABLE 5

Table of Crystallographic Data Form E3											
T°C	a(Å)	b(Å)	c(Å)	α°	β°	γ°	V(Å ³)	Z'	sg	dcalc	R
-23	10.749(5)	13.450(4)	9.250(2)	98.33(2)	95.92(3)	102.82(3)	1277(2)	1	P1	1.223	0.06
T= temp(°C) for the crystallographic data. Z' = number of drug molecules per asymmetric unit											

TABLE 6

Table of Fractional Parameters and Their Estimated Standard Deviations for Form E3					
Atom	x	y	z	B(A2)	Occupany if not equal to 1
S99	0.5568(1)	0.0760(1)	0.5936(1)	3.45(2)	
O1	0.4200(5)	0.5541(4)	0.8496(5)	6.9(1)	
O2	0.2889(5)	0.6016(4)	1.0066(6)	8.1(1)	
O4	0.7004(4)	0.4509(3)	1.0233(4)	4.23(8)	
O8	0.2913(4)	0.2932(3)	1.1074(4)	4.23(8)	
O12	0.1057(4)	0.1088(3)	0.9299(4)	4.16(8)	
O15'	0.329(1)	-0.0602(9)	1.064(1)	4.8(3)*	.3
O15"	0.324(2)	-0.156(1)	1.003(2)	3.2(3)*	.17
O15	0.3312(7)	-0.1150(6)	1.0380(8)	4.9(1)*	.53
O16	0.1810(5)	-0.1433(3)	1.1819(4)	5.7(1)	
O86	0.391(1)	0.6646(7)	0.6196(9)	11.5(4)	
O89	0.3714(7)	0.5646(5)	0.3408(6)	6.5(2)	
O90	0.7502(4)	0.2721(3)	0.8957(5)	4.99(9)	
O95	0.4984(4)	0.0446(3)	0.7188(4)	4.50(8)	
O96	0.6644(4)	0.0315(3)	0.5660(4)	4.83(8)	
O97	0.4651(4)	0.0667(3)	0.4636(4)	5.08(9)	
O98	0.6112(5)	0.1957(3)	0.6332(5)	5.9(1)	
N2	0.4938(5)	0.6229(3)	1.0921(5)	4.8(1)	
N5	0.5365(4)	0.4385(3)	1.1609(4)	3.16(8)	
N10	0.2952(4)	0.2239(3)	0.8056(4)	3.17(8)	
N11	0.2716(4)	0.1163(3)	0.7961(4)	3.08(8)	
N14	0.1336(5)	-0.0874(4)	0.9743(5)	4.9(1)	
N38	-0.2764(4)	0.0574(3)	0.2878(4)	3.24(8)	
C1	0.4011(6)	0.5893(4)	0.9712(7)	5.3(1)	
C3	0.6225(5)	0.6026(4)	1.0813(5)	3.9(1)	

EP 1 755 596 B9

(continued)

Table of Fractional Parameters and Their Estimated Standard Deviations for Form E3					
Atom	x	y	z	B(A ²)	Occupancy if not equal to 1
C4	0.6231(5)	0.4896(3)	1.0873(5)	3.19(9)	
C6	0.5220(5)	0.3284(3)	1.1691(5)	3.14(9)	
C8	0.4026(5)	0.2632(3)	1.0653(5)	3.21(9)	
C9	0.4165(5)	0.2747(4)	0.9050(5)	3.6(1)	
C12	0.1740(5)	0.0661(4)	0.8596(5)	3.4(1)	
C13	0.1592(5)	-0.0523(4)	0.8367(5)	3.8(1)	
C15	0.2248(6)	-0.1124(5)	1.0627(6)	4.6(1)	
C17	0.2720(9)	-0.1732(6)	1.2842(7)	7.3(2)	
C18	0.1818(9)	0.5715(9)	0.894(1)	11.2(3)	
C19	0.7292(7)	0.6818(4)	1.1928(7)	5.8(2)	
C20	0.725(1)	0.7914(6)	1.169(1)	10.7(3)	
C21	0.8613(9)	0.6645(8)	1.165(1)	10.5(3)	
C22	0.710(1)	0.6694(7)	1.3507(8)	10.2(3)	
C23	0.5158(5)	0.3135(4)	1.3298(5)	3.8(1)	
C24	0.6305(6)	0.3765(4)	1.4359(5)	4.0(1)	
C25	0.7519(7)	0.3708(6)	1.4192(7)	6.1(2)	
C26	0.8581(7)	0.4279(9)	1.5213(9)	7.9(2)	
C27	0.8398(8)	0.4935(6)	1.6375(8)	8.6(2)	
C28	0.715(1)	0.5002(6)	1.6576(7)	8.0(2)	
C29	0.6112(8)	0.4430(5)	1.5589(6)	6.0(2)	
C30	0.3043(5)	0.2519(4)	0.6582(5)	3.6(1)	
C31	0.1813(5)	0.2051(4)	0.5532(5)	3.4(1)	
C32	0.0645(5)	0.2123(4)	0.5934(5)	3.9(1)	
C33	-0.0489(5)	0.1725(4)	0.4957(5)	3.8(1)	
C34	-0.0441(5)	0.1243(4)	0.3503(5)	3.16(9)	
C35	0.0756(5)	0.1176(4)	0.3097(5)	3.9(1)	
C36	0.1867(5)	0.1568(4)	0.4095(5)	3.9(1)	
C37	-0.1615(5)	0.0853(4)	0.2417(4)	3.11(9)	
C39	-0.3885(5)	0.0247(4)	0.1969(5)	3.9(1)	
C40	-0.3891(5)	0.0200(4)	0.0470(5)	4.2(1)	
C41	-0.2737(6)	0.0469(4)	-0.0057(5)	4.1(1)	
C42	-0.1596(5)	0.0781(4)	0.0890(5)	3.7(1)	
C43	0.0488(6)	-0.1114(4)	0.7094(6)	4.6(1)	
C44	-0.0819(7)	-0.0958(6)	0.7378(9)	6.8(2)	
C45	0.0496(9)	-0.2266(5)	0.6929(9)	7.8(2)	
C46	0.0797(8)	-0.0738(5)	0.5667(7)	6.2(2)	
C84	0.569(1)	0.7880(9)	0.725(1)	6.3(3)	

(continued)

Table of Fractional Parameters and Their Estimated Standard Deviations for Form E3					
Atom	x	y	z	B(A2)	Occupancy if not equal to 1
C85	0.448(1)	0.7726(9)	0.673(2)	8.4(4)	
C87	0.204(1)	0.449(1)	0.405(2)	10.6(4)	
C88	0.240(1)	0.517(1)	0.316(1)	8.6(3)	
C91	0.8826(7)	0.2919(5)	0.8896(8)	5.8(2)	
C92	0.9613(7)	0.3439(6)	1.035(1)	7.8(2)	
H381	-0.275	0.053	0.403	3.2	
H891	0.397	0.602	0.446	6.6	
H981	0.658	0.219	0.717	6.6	

[0127] Most hydrogens have been omitted; only the hydrogens on N9 and the acid are included.

[0128] Anisotropically refined atoms are given in the form of the isotropic equivalent displacement parameter defined as: $(4/3) * [a^2 B(1,1) + b^2 B(2,2) + c^2 B(3,3) + ab(\cos \gamma) B(1,2) + ac(\cos \beta) B(1,3) + bc(\cos \alpha) B(2,3)]$.

[0129] Form E3 is characterized by the differential scanning calorimetry thermogram having an endotherm typically within the range from about 89.4 to about 96.6 as shown in Figure 11.

[0130] Form E3 is also characterized by a thermal gravimetric analysis curve having a weight loss of about 14.7% at about 150°C as shown in Figure 11.

EXAMPLE 4

[0131] Atazanavir bisulfate Pattern C capsule formulations having the following compositions were prepared as described below.

Ingredient	Stock Granulation ^a (% w/w)	50-mg Capsule (mg/Capsule)	100-mg Capsule (mg/Capsule)	200-mg Capsule (mg/Capsule)
Atazanavir bisulfate	63.2	56.84 ^b	113.67 ^b	227.34 ^b
Lactose, Monohydrate, NF	30.4	27.33 ^c	54.69 ^c	109.35 ^c
Crospovidone, NF	6.0	5.39	10.79	21.58
Magnesium Stearate, NF	0.4	0.36 ^d	0.72 ^d	1.44 ^d
Purified Water, USP or Water for Injection, USP	q.s. ^e	q.s. ^e	q.s. ^e	q.s. ^e
Size #4 Capsule	-	1 Each	-	-
Size #2 Capsule	-	-	1 Each	-
Size #0 Capsule	-	-	-	1 Each
Total Fill Weight	100.0	89.9	179.9	359.7

^a Atazanavir bisulfate Stock Granulation for Capsules (55.5% w/w as the Free Base) was used to manufacture the 50 mg, 100 mg, and 200 mg capsules.

^b This amount is expressed in terms of atazanavir bisulfate at 100% potency, and is equal to 55.5% w/w as the Free Base.

^c The amount of lactose, hydrous will vary depending on the purity of atazanavir bisulfate and the amount of magnesium stearate used.

^d The amount of magnesium stearate used may vary from 0.4% w/w to 0.8% w/w.

^e This is used for processing only and is removed by drying.

[0132] The stock granulation of atazanavir bisulfate was prepared as follows, in which Pattern C material was formed.

[0133] Atazanavir bisulfate Form A, lactose hydrous, and a portion of crospovidone (3 % by weight of total crospovidone present) were mixed in a planetary mixer. The resulting blend was wet granulated with purified water to convert Form A to Pattern C material. The wet granulation was dried in a tray dryer and was sized using a hammer mill. The remaining crospovidone was added to the milled granulation and the mixture was mixed in a PK V-blender. Magnesium stearate was added and the mixture was mixed until a substantially uniform stock granulation was formed.

[0134] The appropriate weight of stock granulations were filled into capsules to produce 50 mg, 100 mg and 200 mg capsules containing atazanavir bisulfate.

EXAMPLE 5

[0135] Atazanavir bisulfate Form A material powder for oral use formulation having the following composition is prepared as described below.

Ingredients	Amount (% w/w)
Atazanavir Bisulfate Form A	3.79
Aspartame, NF	10.00
Sucrose, NF	81.21
Orange vanilla flavor	5.00

[0136] Atazanavir bisulfate Form A is mixed with aspartame, orange vanilla flavor and sucrose in a suitable mixer. The mixture is milled using a hammer mill, followed by a second mixing operation to obtain a uniform mixture. The product is filled into high density polyethylene bottles.

Claims

1. A process for preparing atazanavir bisulfate in the form of Form A crystals, which comprises reacting a solution of atazanavir free base in an organic solvent, which is acetone or mixture of acetone and N-methylpyrrolidone with a first portion of concentrated sulfuric acid in an amount to react with less than about 15% by weight of the atazanavir free base, adding seeds of Form A crystals of atazanavir bisulfate to the reaction mixture, as crystals of atazanavir bisulfate form, adding additional concentrated sulfuric acid in multiple stages at an increasing rate according to the following equation

$$V_{\text{time}} = V_{\text{total}} \times \left(\frac{\text{time}}{\text{time}_{\text{total}}} \right)^3$$

where

V_{time} = Volume of sulfuric acid added during elapsed time period

V_{total} = Total volume of acid representing the 90% charge

time = Elapsed time in crystallization

$\text{time}_{\text{total}}$ = Total crystallization time or total time for acid charging

to effect formation of atazanavir bisulfate crystals, and drying the atazanavir bisulfate to form Form A crystals.

2. The process as defined in Claim 1 wherein the solution of atazanavir free base is initially reacted with from 5 to 15% by weight of the total amount of sulfuric acid employed.
3. The process as defined in Claim 1 wherein the solution of atazanavir free base is initially reacted with from 8 to 12% by weight of the total amount of sulfuric acid employed.
4. The process as defined in Claim 1 wherein the atazanavir free base is reacted with the first portion of sulfuric acid

at a temperature within the range from 35 to 55°C.

5 5. The process as defined in Claim 1 wherein the solution of atazanavir free base is heated to a temperature within the range from 35 to 55°C before it is reacted with sulfuric acid.

10 6. The process as defined in Claim 1 wherein the reaction mixture of atazanavir free base and sulfuric acid is seeded with from 0.1 to 80 weight % of Form A crystals based on the weight of atazanavir free base.

7. The process as defined in Claim 1 wherein the seeded reaction mixture is heated at a temperature within the range from 35 to 55°C.

8. The process as defined in Claim 1 wherein the organic solvent for the atazanavir free base is a mixture of acetone and N-methyl pyrrolidone

15 9. A process for preparing the atazanavir bisulfate Pattern C material, which comprises subjecting Form A crystals of atazanavir bisulfate to a high relative humidity of at least about 95 % RH for at least 24 hours and then drying.

10. A process for preparing atazanavir bisulfate Pattern C material, which comprises

20 (a) reacting a solution of atazanavir free base in an organic solvent, which is acetone or a mixture of acetone and N-methylpyrrolidone, with a first portion of concentrated sulfuric acid in an amount to react with less than about 15 % by weight of the atazanavir free base, adding seeds of Form A crystals of atazanavir bisulfate to the reaction mixture, as crystals of atazanavir bisulfate form, adding additional concentrated sulfuric acid in multiple stages at an increasing rate according to the following equation

$$V_{\text{time}} = V_{\text{total}} \times \left(\frac{\text{time}}{\text{time}_{\text{total}}} \right)^3$$

where

V_{time} = Volume of sulfuric acid added during elapsed time period

V_{total} = Total volume of acid representing the 90% charge

time = Elapsed time in crystallization

$\text{time}_{\text{total}}$ = Total crystallization time or total time for acid charging


to effect formation of atazanavir bisulfate crystals, and drying the atazanavir bisulfate to form Form A crystals;

40 (b) suspending the crystals of Form A atazanavir bisulfate from step (a) in water and drying the suspension to form Pattern C material; or


(c) subjecting crystals of Form A atazanavir bisulfate from step (a) to high relative humidity of greater than 95% RH for at least 24 hours to form Pattern C material or

45 (d) mixing Form A crystals from step (a) with one or more formulating excipients and wet granulating the resulting mixture to directly form Pattern C material in admixture with the excipients.


11. A process according to claim 1 for preparing atazanavir bisulfate




in the form of Form A crystals, which comprises preparing a triamine salt of the structure



and without isolating the triamine salt, reacting the triamine salt with an active ester of an acid of the structure



and a base in the presence of an organic solvent to form a solution of the atazanavir free base of the structure



and converting the free base to the corresponding bisulfate salt

by treating a solution of free base in methylene chloride with N-methyl pyrrolidone and acetone, heating the above mixture to remove methylene chloride and treating the above mixture with sulfuric acid to form the bisulfate salt of the free base wherein the sulfuric acid is added at an increasing rate according to the following equation

55

$$V_{\text{time}} = V_{\text{total}} \times \left(\frac{\text{time}}{\text{time}_{\text{total}}} \right)^3$$

where


V_{time} = Volume of sulfuric acid added during elapsed time period

V_{total} = Total volume of acid representing the 90% charge


time = Elapsed time in crystallization

time_{total} = Total crystallization time or total time for acid charging.

12. The process as defined in Claim 11 where the triamine salt is the hydrochloride salt



13. The process as defined in Claim 11 wherein the active ester of the acid has the structure



14. The process as defined in Claim 11 wherein the base is an alkali metal hydroxide, an alkaline earth metal hydroxide, an alkali metal carbonate, an alkaline earth metal carbonate, an alkali metal phosphate, an alkaline earth metal phosphate or an organic base.


15. The process, as defined in Claim 14 wherein the base is NaOH, KOH, Mg(OH)₂, K₂HPO₄, MgCO₃, Na₂CO₃, K₂CO₃, triethylamine, diisopropylethylamine or N-methylmorpholine and the organic solvent is methylene chloride, ethyl acetate, dichloroethane, tetrahydrofuran, acetonitrile or N,N-dimethylformamide.

16. The process as defined in Claim 11 wherein the triamine salt and the active ester are reacted at a temperature within the range from 30 to 40°C.


17. The process as defined in Claim 16 wherein the triamine salt and the active ester are reacted in the presence of K₂HPO₄ as the base and methylene chloride as the solvent.

18. The process as defined in Claim 11 including the step of seeding the mixture of free base, acetone and N-methylpyrrolidone with crystals of atazanavir bisulfate.


19. A process for preparing atazanavir bisulfate




15 which comprises preparing a triamine hydrogen chloride salt of the structure



30 reacting the triamine hydrogen chloride salt with an active ester of structure



40 and K_2HPO_4 in the presence of methylene chloride to form a solution of the free base of the structure



55 in methylene chloride, and converting the free base to the corresponding bisulfate salt via a cubic crystallization technique wherein the sulfuric acid is added at an increasing rate according to the following equation

$$V_{\text{time}} = V_{\text{total}} \times \left(\frac{\text{time}}{\text{time}_{\text{total}}} \right)^3$$

where

V_{time} = Volume of sulfuric acid added during elapsed time period

V_{total} = Total volume of acid representing the 90% charge

time = Elapsed time in crystallization

time_{total} = Total crystallization time or total time for acid charging.

20. Form E3 of atazanavir bisulfate.

21. The compound as defined in Claim 20 prepared as a triethanolate solvate of atazanavir bisulfate.

22. The compound as defined in Claim 20 as **characterized by** a powder X-ray diffraction pattern substantially in accordance with that shown in Figure 9.

23. The compound as defined in Claim 20 having the crystal structure substantially shown in Figure 10.

24. The compound as defined in Claim 20 which is **characterized by** fractional atomic coordinates substantially as listed in Table 6.

25. The compound as defined in Claim 20 which is **characterized by** crystallographic data substantially equal to the following:

Cell dimensions:

$a = 10.749(5) \text{ \AA}$

$b = 13.450(4) \text{ \AA}$

$c = 9.250(2) \text{ \AA}$

$\alpha = 98.33(2)^\circ$

$\beta = 95.92(3)^\circ$

$\gamma = 102.82(3)^\circ$

Space group P1

Molecules/asymmetric unit 1

wherein said crystalline form is at about -23°C.

26. The compound as defined in Claim 20 which is **characterized by** a differential scanning calorimetry thermogram data substantially in accordance with that shown in Figure 11.

27. The compound as defined in Claim 20 which is **characterized by** a thermal gravimetric analysis curve substantially in accordance with that shown in Figure 11.

Patentansprüche

1. Verfahren zur Herstellung von Atazanavirbisulfat in der Form von Form-A-Kristallen, umfassend Umsetzen einer Lösung freier Atazanavirbase in einem organischen Lösungsmittel, das Aceton oder ein Gemisch aus Aceton und N-Methylpyrrolidon ist, mit einem ersten Teil konzentrierter Schwefelsäure in einer Menge zur Umsetzung mit weniger als etwa 15 Gew.-% der freien Atazanavirbase, Zugabe von Keimen von Form-A-Kristallen von Atazanavirbisulfat zum Reaktionsgemisch, während sich Kristalle von Atazanavirbisulfat bilden, Zugabe zusätzlicher konzentrierter Schwefelsäure in mehreren Stufen mit einer zunehmenden Rate gemäß der folgenden Gleichung

$$V_{\text{Zeit}} = V_{\text{gesamt}} \times \left(\frac{\text{Zeit}}{\text{Zeit}_{\text{gesamt}}} \right)^3$$

worin

V_{Zeit} = Schwefelsäurevolumen, zugegeben während verstrichenen Zeitraums, V_{gesamt} = Säuregesamtvolumen,

darstellend die 90%ige Ladung,

Zeit = bei Kristallisation verstrichene Zeit,

Zeit_{gesamt} = Gesamtkristallisationszeit oder Gesamtzeit für Säurebeladung,

um die Bildung von Atazanavirbisulfatkristallen zu bewirken, und Trocknen des Atazanavirbisulfats, um Form-A-Kristalle zu bilden.

2. Verfahren nach Anspruch 1, wobei die Lösung freier Atazanavirbase anfangs mit 5 bis 15 Gew.-% der verwendeten Schwefelsäure-Gesamtmenge umgesetzt wird.

3. Verfahren nach Anspruch 1, wobei die Lösung freier Atazanavirbase anfangs mit 8 bis 12 Gew.-% der verwendeten Schwefelsäure-Gesamtmenge umgesetzt wird.

4. Verfahren nach Anspruch 1, wobei die freie Atazanavirbase mit dem ersten Teil Schwefelsäure bei einer Temperatur innerhalb des Bereichs von 35 bis 55 °C umgesetzt wird.

5. Verfahren nach Anspruch 1, wobei die Lösung freier Atazanavirbase auf eine Temperatur innerhalb des Bereichs von 35 bis 55 °C erhitzt wird, bevor sie mit Schwefelsäure umgesetzt wird.

6. Verfahren nach Anspruch 1, wobei das Reaktionsgemisch aus freier Atazanavirbase und Schwefelsäure mit 0,1 bis 80 Gew.-% von Form-A-Kristallen, bezogen auf das Gewicht freier Atazanavirbase, geimpft wird.

7. Verfahren nach Anspruch 1, wobei das geimpfte Reaktionsgemisch bei einer Temperatur innerhalb des Bereichs von 35 bis 55 °C erhitzt wird.

8. Verfahren nach Anspruch 1, wobei das organische Lösungsmittel für die freie Atazanavirbase ein Gemisch aus Aceton und N-Methylpyrrolidon ist.

9. Verfahren zur Herstellung des Atazanavirbisulfat-Muster-C-Materials, das Aussetzen von Form-A-Kristallen von Atazanavirbisulfat gegenüber einer hohen relativen Feuchtigkeit von zumindest etwa 95% rF zumindest 24 Stunden lang und dann Trocknen umfasst.

10. Verfahren zur Herstellung von Atazanavirbisulfat-Muster-C-Material, das umfasst

(a) Umsetzen einer Lösung freier Atazanavirbase in einem organischen Lösungsmittel, das Aceton oder ein Gemisch aus Aceton und N-Methylpyrrolidon ist, mit einem ersten Teil konzentrierter Schwefelsäure in einer Menge zur Umsetzung mit weniger als etwa 15 Gew.-% der freien Atazanavirbase, Zugabe von Keimen von Form-A-Kristallen von Atazanavirbisulfat zum Reaktionsgemisch, während sich Kristalle von Atazanavirbisulfat bilden, Zugabe zusätzlicher konzentrierter Schwefelsäure in mehreren Stufen mit einer zunehmenden Rate gemäß der folgenden Gleichung

$$V_{\text{Zeit}} = V_{\text{gesamt}} \times \left(\frac{\text{Zeit}}{\text{Zeit}_{\text{gesamt}}} \right)^3$$

worin

V_{Zeit} = Schwefelsäurevolumen, zugegeben während verstrichenen Zeitraums, V_{gesamt} = Säuregesamtvolumen, darstellend die 90%ige Ladung,

Zeit = bei Kristallisation verstrichene Zeit,

Zeit_{gesamt} = Gesamtkristallisationszeit oder Gesamtzeit für Säurebeladung,

um die Bildung von Atazanavirbisulfatkristallen zu bewirken, und Trocknen des Atazanavirbisulfats, um Form-A-Kristalle zu bilden;


(b) Suspendieren der Kristalle von Form-A-Atazanavirbisulfat aus Schritt (a) in Wasser und Trocknen der Suspension, um Muster-C-Material zu bilden; oder

(c) Aussetzen von Kristallen von Form-A-Atazanavirbisulfat aus Schritt (a) gegenüber hoher relativer Feuchtigkeit von mehr als 95% rF zumindest 24 Stunden lang, um Muster-C-Material zu bilden; oder


(d) Mischen von Form-A-Kristallen aus Schritt (a) mit einem oder mehreren Formulierungsexzipienten und Nassgranulieren des sich ergebenden Gemischs, um Muster-C-Material in Beimischung mit den Exzipienten

direkt zu bilden.


11. Verfahren nach Anspruch 1 zur Herstellung von Atazanavirbisulfat




in der Form von Form-A-Kristallen, umfassend Herstellen eines Triaminsalzes der Struktur



und, ohne Isolieren des Triaminsalzes, Umsetzen des Triaminsalzes mit einem aktiven Ester einer Säure der Struktur



und einer Base in Gegenwart eines organischen Lösungsmittels zwecks Bildung einer Lösung der freien Atazanavirbase der Struktur



und Umwandeln der freien Base zum entsprechenden Bisulfatsalz, durch Behandeln einer Lösung freier Base in Methylenchlorid mit N-Methylpyrrolidon und Aceton, Erhitzen des obigen Gemischs, um Methylenchlorid zu entfernen, und Behandeln des obigen Gemischs mit Schwefelsäure, um das Bisulfatsalz der freien Base zu bilden, wobei die Schwefelsäure zugegeben wird mit einer zunehmenden Rate gemäß der folgenden Gleichung

$$V_{\text{Zeit}} = V_{\text{gesamt}} \times \left(\frac{\text{Zeit}}{\text{Zeit}_{\text{gesamt}}} \right)^3$$


worin

V_{Zeit} = Schwefelsäurevolumen, zugegeben während verstrichenen Zeitraums, V_{gesamt} = Säuregesamtvolumen, darstellend die 90%ige Ladung,

Zeit = bei Kristallisation verstrichene Zeit,


$\text{Zeit}_{\text{gesamt}}$ = Gesamtkristallisationszeit oder Gesamtzeit für Säurebeladung.

12. Verfahren nach Anspruch 11, wobei das Triaminsalz das Hydrochloridsalz



ist.

13. Verfahren nach Anspruch 11, wobei der aktive Ester der Säure die Struktur



aufweist.

14. Verfahren nach Anspruch 11, wobei die Base ein Alkalimetallhydroxid, ein Erdalkalimetallhydroxid, ein Alkalimetallcarbonat, ein Erdalkalimetallcarbonat, ein Alkalimetallphosphat, ein Erdalkalimetallphosphat oder eine organische Base ist.


15. Verfahren nach Anspruch 14, wobei die Base NaOH, KOH, $\text{Mg}(\text{OH})_2$, K_2HPO_4 , MgCO_3 , Na_2CO_3 , K_2CO_3 , Triethylamin, Diisopropylethylamin oder N-Methylmorpholin ist und das organische Lösungsmittel Methylenchlorid, Ethylacetat, Dichlorethan, Tetrahydrofuran, Acetonitril oder N,N-Dimethylformamid ist.

16. Verfahren nach Anspruch 11, wobei das Triaminsalz und der aktive Ester bei einer Temperatur innerhalb des Bereichs von 30 bis 40 °C umgesetzt werden.


17. Verfahren nach Anspruch 16, wobei das Triaminsalz und der aktive Ester in Gegenwart von K_2HPO_4 als der Base und Methylenchlorid als dem Lösungsmittel umgesetzt werden.

18. Verfahren nach Anspruch 11, einschließlich den Schritt des Impfens des Gemischs aus freier Base, Aceton und N-Methylpyrrolidon mit Kristallen von Atazanavirbisulfat.


19. Verfahren zur Herstellung von Atazanavirbisulfat




umfassend Herstellen eines Triaminwasserstoffchloridsalzes der Struktur



Umsetzen des Triaminwasserstoffchloridsalzes mit einem aktiven Ester der Struktur



und K_2HPO_4 in Gegenwart von Methylenchlorid zwecks Bildung einer Lösung der freien Base der Struktur



in Methylenchlorid, und Umwandeln der freien Base zum entsprechenden Bisulfatsalz via eine Technik zur kubischen Kristallisation, wobei die Schwefelsäure zugegeben wird mit einer zunehmenden Rate gemäß der folgenden Gleichung

$$V_{\text{Zeit}} = V_{\text{gesamt}} \times \left(\frac{\text{Zeit}}{\text{Zeit}_{\text{gesamt}}} \right)^3$$

worin

V_{Zeit} = Schwefelsäurevolumen, zugegeben während verstrichenen Zeitraums,

V_{gesamt} = Säuregesamtvolumen, darstellend die 90%ige Ladung,
 Zeit = bei Kristallisation verstrichene Zeit,
 $\text{Zeit}_{\text{gesamt}}$ = Gesamtkristallisationszeit oder Gesamtzeit für Säurebeladung.

20. Form E3 von Atazanavirbisulfat.

21. Verbindung nach Anspruch 20, hergestellt als Triethanolatsolvat von Atazanavirbisulfat.

22. Verbindung nach Anspruch 20 gemäß Charakterisierung durch ein Pulverröntgenbeugungsmuster, das im Wesentlichen mit dem in Figur 9 dargestellten übereinstimmt.

23. Verbindung nach Anspruch 20 mit der im Wesentlichen in Figur 10 dargestellten Kristallstruktur.

24. Verbindung nach Anspruch 20, die durch fraktionelle Atomkoordinaten im Wesentlichen gemäß Auflistung in Tabelle 6 gekennzeichnet ist.

25. Verbindung nach Anspruch 20, die durch kristallographische Daten gekennzeichnet ist, die im Wesentlichen gleich den folgenden sind:

Zelldimensionen:

$$a = 10,749(5) \text{ \AA}$$

$$b = 13,450(4) \text{ \AA}$$

$$c = 9,250(2) \text{ \AA}$$

$$\alpha = 98,33(2)^\circ$$

$$\beta = 95,92(3)^\circ$$

$$\gamma = 102,82(3)^\circ$$

Raumgruppe P1

Moleküle/asymmetrische Einheit 1

wobei die kristalline Form bei etwa -23 °C ist.

26. Verbindung nach Anspruch 20, die durch Dynamische-Differenz-Kalorimetrie-Thermogrammdaten gekennzeichnet ist, die im Wesentlichen mit den in Figur 11 dargestellten übereinstimmen.

27. Verbindung nach Anspruch 20, die durch eine Thermogravimetrieanalysekurve gekennzeichnet ist, die im Wesentlichen mit der in Figur 11 dargestellten übereinstimmt.

Revendications

1. Procédé de préparation de bisulfate d'atazanavir sous forme de cristaux de Forme A, lequel comprend mettre une solution de base libre d'atazanavir dans un solvant organique, qui est de l'acétone ou un mélange d'acétone et de N-méthylpyrrolidone, à réagir avec une première portion d'acide sulfurique concentré en une quantité pour réagir avec moins d'environ 15 % en poids de la base libre d'atazanavir, ajouter des semences de cristaux de Forme A de bisulfate d'atazanavir au mélange de réaction, au fur et à mesure que des cristaux de bisulfate d'atazanavir se forment, ajouter de l'acide sulfurique concentré additionnel par échelons multiples à un rythme croissant conformément à l'équation suivante:

$$V_{\text{temps}} = V_{\text{total}} \times \left(\frac{\text{temps}}{\text{temps}_{\text{total}}} \right)^3$$

dans laquelle:

V_{temps} = Volume d'acide sulfurique ajouté pendant la période de temps écoulé

V_{total} = Volume total d'acide représentant la charge de 90 %

Temps = Temps écoulé dans la cristallisation

Temps_{total} = Temps total de cristallisation ou temps total de chargement d'acide
afin d'effectuer la formation de cristaux de bisulfate d'atazanavir et sécher le bisulfate d'atazanavir pour former des cristaux de Forme A.

2. Procédé selon la revendication 1, dans lequel la solution de base libre d'atazanavir est initialement mise à réagir avec de 5 à 15 % en poids de la quantité totale d'acide sulfurique utilisé.
3. Procédé selon la revendication 1, dans lequel la solution de base libre d'atazanavir est initialement mise à réagir avec de 8 à 12 % en poids de la quantité totale d'acide sulfurique utilisé.
4. Procédé selon la revendication 1, dans lequel la base libre d'atazanavir est mise à réagir avec la première portion d'acide sulfurique à une température dans la plage de 35 à 55 °C.
5. Procédé selon la revendication 1, dans lequel la solution de base libre d'atazanavir est chauffée à une température dans la plage de 35 à 55 °C avant d'être mise à réagir avec l'acide sulfurique.
6. Procédé selon la revendication 1, dans lequel le mélange de réaction de base libre d'atazanavir et d'acide sulfurique estensemencé avec de 0,1 à 80 % en poids de cristaux de Forme A sur la base du poids total de la base libre d'atazanavir.
7. Procédé selon la revendication 1, dans lequel le mélange de réactionensemencé est chauffé à une température dans la plage de 35 à 55 °C.
8. Procédé selon la revendication 1, dans lequel le solvant organique pour la base libre d'atazanavir est un mélange d'acétone et de N-méthylpyrrolidone.
9. Procédé de préparation de la matière de bisulfate d'atazanavir à Diagramme C, qui comprend soumettre des cristaux de Forme A de bisulfate d'atazanavir à une humidité relative élevée d'au moins environ 95 % HR pendant au moins 24 heures et puis sécher.
10. Procédé de préparation d'une matière de bisulfate d'atazanavir à Diagramme C, qui comprend:

(a) mettre une solution de base libre d'atazanavir dans un solvant organique, qui est de l'acétone ou un mélange d'acétone et de N-méthylpyrrolidone, à réagir avec une première portion d'acide sulfurique concentré en une quantité pour réagir avec moins d'environ 15 % en poids de la base libre d'atazanavir, ajouter des semences de cristaux de Forme A de bisulfate d'atazanavir au mélange de réaction, au fur et à mesure que des cristaux de bisulfate d'atazanavir se forment, ajouter de l'acide sulfurique concentré additionnel par échelons multiples à un rythme croissant conformément à l'équation suivante:

$$V_{\text{temps}} = V_{\text{total}} \times \left(\frac{\text{temps}}{\text{temps}_{\text{total}}} \right)^3$$

dans laquelle:

V_{temps} = Volume d'acide sulfurique ajouté pendant la période de temps écoulé

V_{total} = Volume total d'acide représentant la charge de 90 %

Temps = Temps écoulé dans la cristallisation

Temps_{total} = Temps total de cristallisation ou temps total de chargement d'acide

afin d'effectuer la formation de cristaux de bisulfate d'atazanavir et sécher le bisulfate d'atazanavir pour former des cristaux de Forme A;


(b) mettre les cristaux de Forme A de bisulfate d'atazanavir de l'étape (a) en suspension dans de l'eau et sécher la suspension pour former une matière à Diagramme C; ou bien

(c) soumettre les cristaux de Forme A de bisulfate d'atazanavir de l'étape (a) à une humidité relative élevée supérieure à 95 % HR pendant au moins 24 heures pour former une matière à Diagramme C; ou bien


(d) mélanger les cristaux de Forme A de l'étape (a) avec un ou plusieurs excipients de formulation et granuler

par voie humide le mélange résultant pour former directement une matière à Diagramme C mélangée avec les excipients.


11. Procédé selon la revendication 1 de préparation de bisulfate d'atazanavir




sous forme de cristaux de Forme A, lequel comprend préparer un sel de triamine de la structure



et, sans isoler le sel de triamine, mettre le sel de triamine à réagir avec un ester actif d'un acide de la structure



et une base en présence d'un solvant organique pour former une solution de la base libre d'atazanavir de la structure



et convertir la base libre en le sel bisulfate correspondant en traitant une solution de base libre dans du chlorure de méthylène avec de la N-méthylpyrrolidone et de l'acétone, chauffer le mélange ci-dessus pour retirer le chlorure de méthylène et traiter le mélange ci-dessus avec de l'acide sulfurique pour former le sel bisulfate de la base libre, où l'acide sulfurique est ajouté à un rythme croissant conformément à l'équation suivante:

$$V_{\text{temps}} = V_{\text{total}} \times \left(\frac{\text{temps}}{\text{temps}_{\text{total}}} \right)^3$$

dans laquelle:


V_{temps} = Volume d'acide sulfurique ajouté pendant la période de temps écoulé

V_{total} = Volume total d'acide représentant la charge de 90 %


Temps = Temps écoulé dans la cristallisation

$\text{Temps}_{\text{total}}$ = Temps total de cristallisation ou temps total de chargement d'acide.

12. Procédé selon la revendication 11, dans lequel le sel de triamine est le sel chlorhydrate



13. Procédé selon la revendication 11, dans lequel l'ester actif de l'acide a la structure



14. Procédé selon la revendication 11, dans lequel la base est un hydroxyde de métal alcalin, un hydroxyde de métal alcalino-terreux, un carbonate de métal alcalin, un carbonate de métal alcalino-terreux, un phosphate de métal alcalin, un phosphate de métal alcalino-terreux ou une base organique.


15. Procédé selon la revendication 14, dans lequel la base est du NaOH, KOH, $\text{Mg}(\text{OH})_2$, K_2HPO_4 , MgCO_3 , Na_2CO_3 , K_2CO_3 , triéthylamine, diisopropyléthylamine ou de la N-méthylmorpholine et le solvant organique est du chlorure de méthylène, acétate d'éthyle, dichloroéthane, tétrahydrofurane, acétonitrile ou du N,N-diméthylformamide.

16. Procédé selon la revendication 11, dans lequel le sel de triamine et l'ester actif sont mis à réagir à une température dans la plage de 30 à 40 °C.


17. Procédé selon la revendication 16, dans lequel le sel de triamine et l'ester actif sont mis à réagir en présence de K_2HPO_4 en tant que base et de chlorure de méthylène en tant que solvant.

18. Procédé selon la revendication 11, comprenant l'étape consistant à ensemercer le mélange de base libre, d'acétone et de N-méthylpyrrolidone avec des cristaux de bisulfate d'atazanavir.


19. Procédé de préparation de bisulfate d'atazanavir




15 qui comprend préparer un sel chlorhydrate de triamine de la structure



mettre le sel chlorhydrate de triamine à réagir avec un ester actif de la structure



35 et du K_2HPO_4 en présence de chlorure de méthylène pour former une solution de la base libre de la structure



50 dans du chlorure de méthylène et convertir la base libre en le sel bisulfate correspondant par une technique de cristallisation cubique, dans lequel l'acide sulfurique est ajouté à un rythme croissant conformément à l'équation suivante:

55

$$V_{\text{temps}} = V_{\text{total}} \times \left(\frac{\text{temps}}{\text{temps}_{\text{total}}} \right)^3$$

dans laquelle:

V_{temps} = Volume d'acide sulfurique ajouté pendant la période de temps écoulé
 V_{total} = Volume total d'acide représentant la charge de 90 %
 Temps = Temps écoulé dans la cristallisation
 $\text{Temps}_{\text{total}}$ = Temps total de cristallisation ou temps total de chargement d'acide.

20. Forme E3 de bisulfate d'atazanavir.

21. Composé selon la revendication 20, préparé comme un solvate triéthanolate de bisulfate d'atazanavir.

22. Composé selon la revendication 20, tel que **caractérisé par** le diagramme de diffraction des rayons X sur poudre conformément à celui présenté sur la Figure 9.

23. Composé selon la revendication 20, ayant la structure cristalline sensiblement présentée sur la Figure 10.

24. Composé selon la revendication 20, qui est **caractérisé par** des coordonnées atomiques fractionnaires sensiblement telles que présentées dans la Table 6.

25. Composé selon la revendication 20, qui est **caractérisé par** des données cristallographiques sensiblement égales aux suivantes:

Dimensions cellulaires:

$a = 10,749(5) \text{ \AA}$

$b = 13,450(4) \text{ \AA}$

$c = 9,250(2) \text{ \AA}$

$\alpha = 98,33(2)^\circ$

$\beta = 95,92(3)^\circ$

$\gamma = 102,82(3)^\circ$

Groupe spatial P1.

Molécules/unité asymétrique 1

dans lequel la forme cristalline est à environ -23°C .

26. Composé selon la revendication 20, qui est **caractérisé par** des données de thermogramme de calorimétrie différentielle par balayage sensiblement conformes à celles présentées sur la Figure 11.

27. Composé selon la revendication 20, qui est **caractérisé par** une courbe d'analyse thermogravimétrique sensiblement conforme à celle présentée sur la Figure 11.

FIG. 1




FIG. 2

Crystal structure of atazanavir Form A





FIG. 3


DSC



DSC Form A


FIG. 4

TGA



TGA Form A

FIG. 5



Carbon-13 SSNMR of FORM A

FIG. 6

Powder x-ray diffraction pattern of atazanavir Pattern C





FIG. 7


DSC



DSC Pattern C

FIG. 8

TGA



TGA Pattern C

FIG. 9

Observed (top) and Simulated (Bottom) Powder X-ray
Diffraction Patterns for Form E3




FIG. 10

Crystal structure of atazanavir Form E3





FIG. 11



DSC and TGA Form E3

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 56804304 P [0001]
- US 60607533 B [0001]
- US 5849911 A, Fässler [0003]
- US 6087383 A, Singh [0008] [0011]

Non-patent literature cited in the description

- **STOUT ; JENSEN.** X-Ray Structure Determination: A Practical Guide. Macmillan Co, 1968 [0019]
- **SMITH, D.K.** A FORTRAN Program for Calculating X-Ray Powder Diffraction Patterns. Lawrence Radiation Laboratory, April 1963 [0043]
- **S.R. BYM ; R.R. PFEIFFER ; J.G. STOWELL.** Solid-State Chemistry of Drugs. SSCI, 1999 [0045]
- **J.W. MULLIN ; J. NYVLT.** Programmed cooling of batch crystallizers. *Chemical Engineering Science*, 1971, vol. 26, 369-377 [0048]
- Crystallization. Butterworth-Heineman, 1993 [0064]
- **Z. XU et al.** Process Research and Development for an Efficient Synthesis of the HIV Protease Inhibitor BMS-232,632. *Organic Process Research and Development*, 2002, vol. 6, 323-328 [0092]