



(11) **EP 1 763 364 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Description Paragraph(s) 176

(48) Corrigendum issued on:
03.07.2013 Bulletin 2013/27

(45) Date of publication and mention
of the grant of the patent:
11.04.2012 Bulletin 2012/15

(21) Application number: **05723323.1**

(22) Date of filing: **22.02.2005**

(51) Int Cl.:
A61K 39/385 (2006.01) **A61K 39/39** (2006.01)
C07K 14/47 (2006.01) **A61K 39/00** (2006.01)
C07K 16/18 (2006.01)

(86) International application number:
PCT/US2005/005285

(87) International publication number:
WO 2005/081872 (09.09.2005 Gazette 2005/36)

(54) **METHODS AND COMPOSITIONS COMPRISING SUPRAMOLECULAR CONSTRUCTS**
VERFAHREN UND ZUSAMMENSETZUNGEN MIT SUPRAMOLEKULAREN KONSTRUKTEN
PROCEDES ET COMPOSITIONS CONTENANT DES CONSTRUCTIONS SUPRAMOLECULAIRES

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU MC NL PL PT RO SE SI SK TR
Designated Extension States:
AL BA HR LV MK YU

(30) Priority: **20.02.2004 US 783975**
04.10.2004 US 958211

(43) Date of publication of application:
21.03.2007 Bulletin 2007/12

(60) Divisional application:
12153152.9 / 2 465 533

(73) Proprietor: **AC Immune S.A.**
1015 Lausanne (CH)

(72) Inventors:
• **NICOLAU, Claude Yves**
Newton MA Massachusetts 02459 (US)
• **GREFERATH, Ruth, Dr.**
77694 Kehl (DE)
• **HICKMAN, David**
1025 Saint-Sulpice (CH)

(74) Representative: **Vossius & Partner**
Siebertstrasse 4
81675 München (DE)

(56) References cited:
WO-A1-2005/014036 WO-A2-99/41279
US-A- 4 565 696 US-A1- 2004 242 845

US-B1- 6 521 211

- **JANSSEN A P C A ET AL: "Peptide-targeted PEG-liposomes in anti-angiogenic therapy"**
INTERNATIONAL JOURNAL OF PHARMACEUTICS, ELSEVIER BV, NL, vol. 254, 1 January 2003 (2003-01-01), pages 55-58, XP002497490 ISSN: 0378-5173
- **NICOLAU C ET AL: "A LIPOSOME-BASED THERAPEUTIC VACCINE AGAINST BETA-AMYLOID PLAQUES ON THE PANCREAS OF TRANSGENIC NORBA MICE" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE, WASHINGTON, DC, US, vol. 99, no. 4, 19 February 2002 (2002-02-19), pages 2332-2337, XP009062068 ISSN: 0027-8424**
- **ALLEN T M ET AL: "LIPOSOMES CONTAINING SYNTHETIC LIPID DERIVATIVES OF POLY (ETHYLENE GLYCOL) SHOW PROLONGED CIRCULATION HALF-LIVES IN VIVO"**
BIOCHIMICA ET BIOPHYSICA ACTA, ELSEVIER, NL, vol. 1066, 1 January 1991 (1991-01-01), pages 29-36, XP000672933 ISSN: 0006-3002
- **PAWLAK-ROBLIN C ET AL: "Inhibition of multidrug resistance by immunisation with synthetic P-glycoprotein-derived peptides"**
EUROPEAN JOURNAL OF CANCER, PERGAMON PRESS, OXFORD, GB, vol. 40, no. 4, 1 March 2004 (2004-03-01), pages 606-613, XP004488461 ISSN: 0959-8049

EP 1 763 364 B9

- GIOIA DE L ET AL: "CONFORMATIONAL POLYMORPHISM OF THE AMYLOIDOGENIC AND NEUROTOXIC PEPTIDE HOMOLOGOUS TO RESIDUES 106-126 OF THE PRION PROTEIN" JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY OF BIOCHEMICAL BIOLOGISTS, BIRMINGHAM, US, vol. 269, no. 11, 18 March 1994 (1994-03-18), pages 7859-7862, XP000887214 ISSN: 0021-9258
- MUHS ANDREAS ET AL: "Liposomal vaccines with conformation-specific amyloid peptide antigens define immune response and efficacy in APP transgenic mice" June 2007 (2007-06), PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, VOL. 104, NR. 23, PAGE(S) 9810-9815, XP002542049 ISSN: 0027-8424 * the whole document *
- MUHS ET AL: "Improved memory capacity of amyloid precursor protein transgenic mice through passive administration of a monoclonal antibody inducing a conformational shift of amyloid-beta" ALZHEIMER'S & DEMENTIA: THE JOURNAL OF THE ALZHEIMER'S ASSOCIATION, ELSEVIER, NEW YORK, NY, US, vol. 2, no. 3, 01-06-06, 1 July 2006 (2006-07-01), page S21, XP005547955 ISSN: 1552-5260
- GATOULLAT ET AL: "Immunization with liposome-anchored pegylated peptides modulates doxorubicin sensitivity in P-glycoprotein-expressing P388 cells" CANCER LETTERS, NEW YORK, NY, US, vol. 257, no. 2, 11 October 2007 (2007-10-11), pages 165-171, XP022294900 ISSN: 0304-3835

Description

[0001] The present invention is related to methods and compositions for eliciting high immune responses. In particular, the present invention includes novel compositions and methods for yielding conformationally sensitive antibodies.

[0002] The immune system is a complex response system of the body that involves many different kinds of cells that have differing activities. Activation of one portion of the immune system usually causes a variety of responses due to unwanted activation of other related portions of the system. Currently, there are no satisfactory methods or compositions for producing a specifically desired response by targeting the specific components of the immune system.

[0003] The immune system is a complex interactive system of the body that involves a wide variety of components, including cells, and cellular factors, which interact with stimuli from both inside the body and outside the body. Aside from its direct action, the immune system's response is also influenced by other systems of the body including the nervous, respiratory, circulatory, and digestive systems.

[0004] One of the better-known aspects of the immune system is its ability to respond to foreign antigens presented by invading organisms, cellular changes within the body, or from vaccination. Some of the first kinds of cells that respond to such activation of the immune system are phagocytes and natural killer cells. Phagocytes include among other cells, monocytes, macrophages, and polymorphonuclear neutrophils. These cells generally bind to the foreign antigen, internalize it and often times destroy it. They also produce soluble molecules that mediate other immune responses, such as inflammatory responses. Natural killer cells can recognize and destroy certain virally- infected embryonic and tumor cells. Other factors of the immune response include complement pathways, which are capable of responding independently to foreign antigens or acting in concert with cells or antibodies.

[0005] Generally, it is thought that the response to antigens involves both humoral responses and cellular responses. Humoral immune responses are mediated by non-cellular factors that are released by cells and which may or may not be found free in the plasma or intracellular fluids. A major component of a humoral response of the immune system is mediated by antibodies produced by B lymphocytes. Cell-mediated immune responses result from the interactions of cells, including antigen presenting cells and B lymphocytes (B cells) and T lymphocytes (T cells).

[0006] One of the most widely employed aspects of the immune response capabilities is the production of monoclonal antibodies. The advent of monoclonal antibody (Mab) technology in the mid 1970s provided a valuable new therapeutic and diagnostic tool. For the first time, researchers and clinicians had access to unlimited quantities of uniform antibodies capable of binding to a predetermined antigenic site and having various immunological effector functions. Currently, the techniques for production of monoclonal antibodies are well known in the art. However there remains a continuing need for specialized antibodies. In essence, what is desired is the ability to produce customized antibodies. The need is especially great in the area of combating infectious disease where pathogens have acquired resistance to commonly used antibiotics. In addition, there is a need for antibiotics, for addressing pathological conditions resulting from cause other than an infectious agent.

[0007] Alzheimer's Disease (AD) is a neurological disorder primarily thought to be caused by the build of amyloid plaques caused by abnormal deposit of proteins in the brain. Scientific evidence demonstrates that AD results from an increase in the production or accumulation of beta-amyloid protein in plaques that leads to nerve cell death. Loss of nerve cells in strategic brain areas, in turn, causes reduction in the neurotransmitters and impairment of memory. The proteins principally responsible for the plaque build up include amyloid precursor protein (APP) and two presenilins (presenilin I and presenilin II). The degradation of APPs likely increases their propensity to aggregate in plaques. There is a need for specific antibodies that can target and diffuse amyloid plaque formation.

[0008] The symptoms of AD manifest slowly and the first symptom may only be mild forgetfulness. In this stage, individuals may forget recent events, activities, the names of familiar people or things and may not be able to solve simple math problems. As the disease progresses, symptoms are more easily noticed and become serious enough to cause people with AD or their family members to seek medical help. Mid-stage symptoms of AD include forgetting how to do simple tasks such as grooming, and problems develop with speaking, understanding, reading, or writing. Later stage AD patients may become anxious or aggressive, may wander away from home and ultimately need total care.

[0009] Presently, the only definite way to diagnose AD is to identify plaques and tangles in brain tissue in an autopsy after death of the individual. Therefore, doctors can only make a diagnosis of "possible" or "probable" AD while the person is still alive. Using current methods, physicians can diagnose AD correctly up to 90 percent of the time using several tools to diagnose "probable" AD. Physicians ask questions about the person's general health, past medical problems, and the history of any difficulties the person has carrying out daily activities. Behavioral tests of memory, problem solving, attention, counting, and language provide information on cognitive degeneration and medical tests—such as tests of blood, urine, or spinal fluid, and brain scans can provide some further information.

[0010] The management of AD consists of medication-based and non-medication based treatments. Treatments aimed at changing the underlying course of the disease (delaying or reversing the progression) have so far been largely unsuccessful. Medicines that restore the deficit (defect), or malfunctioning, in the chemical messengers of the nerve cells (neurotransmitters), such as the cholinesterase inhibitors (ChEIs), have been shown to improve symptoms. Med-

ications are also available to address the psychiatric manifestations of AD.

[0011] Cholinesterase inhibitors, such as Tacrine and Rivastigmine, are currently the only class of agents that are approved by the FDA for the treatment of AD. These agents are medicines that restore the defect, or malfunctioning, in the chemical neurotransmission in the brain. ChEIs impede the enzymatic degradation of neurotransmitters thereby increasing the amount of chemical messengers available to transmit the nerve signals in the brain.

[0012] For some people in the early and middle stages of the disease, the drugs tacrine (COGNEX[®], Morris Plains, NJ), donepezil (ARICEPT[®], Tokyo, JP), rivastigmine (EXELON[®], East Hanover, NJ), or galantamine (REMINYL[®], New Brunswick, NJ) may help prevent some symptoms from becoming worse for a limited time. Another drug, memantine (NAMENDA[®], New York, NY), has been approved for treatment of moderate to severe AD. Also, some medicines may help control behavioral symptoms of AD such as sleeplessness, agitation, wandering, anxiety, and depression. Treating these symptoms often makes patients more comfortable and makes their care easier for caregivers. Unfortunately, despite significant treatment advances showing that this class of agents is consistently better than a placebo, the disease continues to progress despite treatment, and the average effect on mental functioning has only been modest. ChEIs also have side effects that include gastrointestinal dysfunction, liver toxicity and weight loss.

[0013] Advances in the understanding of the brain abnormalities that occur in AD are hoped to provide the framework for new targets of treatment that are more focused on altering the course and development of the disease. Many compounds, including anti-inflammatory agents, are being actively investigated. Clinical trials using specific cyclooxygenase inhibitors (COX-2), such as rofecoxib and celecoxib, are also underway.

[0014] Another factor to consider when developing new drugs is the ease of use for the target patients. Oral drug delivery—specifically tablets, capsules and softgels—account for 70% of all dosage forms consumed because of patient convenience. Drug developers agree that patients prefer oral delivery rather than subjecting themselves to injections or other, more invasive forms of medicinal administration. Formulations resulting in low dosing intervals (i.e. once a day or sustained release) are also preferable. The ease of administering antibiotics in oral dosage forms results in an increase of patient compliance during treatment.

[0015] The prior art discloses different compositions and methods for treating diseases (e.g. cancer) or neurological disease (e.g. AD).

[0016] A peptide targeted PEG-liposome used in anti-angiogenic therapy is described by Janssen et al., wherein cyclic RGD-peptides or ATWLPPR-peptides coupled to the surface of PEG-liposomes causes binding of these liposome to endothelial cells in vitro and RGD-peptide targeted liposomes demonstrate specific binding to the tumor endothelium in vivo (Janssen et al., *International Journal of Pharmaceutics*, 2003, 254, 55-58).

[0017] One known method for inhibiting amyloid fibrillogenesis is disclosed in WO99/41279, wherein the method comprises contacting tissue with a composition comprising an amyloidogenic peptide that has been blocked.

[0018] Nicolau et al. teaches an immunogenic composition comprising an amyloid antigenic peptide modified with palmitic acid reconstituted in a liposome and the use thereof for reducing plaques in the pancreas of transgenic NOBRA mice (Nicolau et al., *PNAS*, 2002, 99(4), 2332-2337).

[0019] Allen et al. discloses synthetic lipid derivatives of poly(ethylene glycol) (PEG) which have been synthesized and tested for their ability to decrease uptake of liposomes into the mononuclear phagocyte system (MPS, reticuloendothelial system) in mice and to prolong circulation half-lives of liposomes (AllXen et al., *Biochimica et Biophysica acta*, 1991, 1066, 29-36). More particularly, poly(ethylene glycol) covalently attached to bovine serum albumine is reported to decrease the immunogenicity of this protein.

[0020] What is needed are effective methods and compositions for generation of highly specific and highly effective antibodies. Preferably such antibodies would recognize specific epitopes on various antigens such as amyloid protein, prion protein or P₁₇₀ glycoprotein.

[0021] What is also needed therefore, are effective compositions and methods for addressing the complications associated with neurological disease associated with amyloid plaque formation such as Alzheimer's disease. In particular what is need are specialized antibodies capable of counteracting the physiological manifestations of the disease such as the formation plaques associated with aggregation of fibers of the amyloid peptide in its beta sheet conformation.

SUMMARY OF THE INVENTION

[0022] The present invention includes novel methods and compositions for eliciting highly specific and highly effective antibodies. Unlike currently available products the present invention provides unique methods and compositions resulting in antibodies having the ability to recognize specific epitopes from a range of antigens.

[0023] The present invention satisfies the long felt need for compositions that enable the generation of antibodies that specifically recognize epitopes such as those of amyloid protein or prion protein.

[0024] The present invention comprises unique antigen presentation that results in enhanced exposure and ultimately antibodies with a higher degree of conformational sensitivity.

[0025] In one embodiment, the invention relates to a composition comprising an antigenic construct wherein the

antigenic construct comprises

- a) an antigenic peptide or active fragment thereof, wherein the antigenic peptide or active fragment thereof is modified to have a polyethylene glycol covalently attached, one at each terminus, and
 b) wherein the antigenic construct is reconstituted in liposomes; to enhance antigenicity;

[0026] In another embodiment, the invention relates to the composition of the invention comprising an antigenic construct wherein the antigenic construct comprises

- a) an antigenic peptide or active fragment thereof, wherein the antigenic peptide or active fragment thereof is modified to have a pegylated lysine covalently attached, one at each terminus, and
 b) wherein the antigenic construct is reconstituted in liposomes to enhance antigenicity.

[0027] In one embodiment the invention includes compositions comprising antigenic constructs comprising a peptide sequence, covalently attached to pegylated amino acid (such as pegylated lysine)- one at each terminus.

[0028] Accordingly, it is an object of the present invention to provide methods and compositions for eliciting specific and effective immune responses.

[0029] It is another object of the present invention to provide methods and compositions for treating and preventing the occurrence or spread of disease.

[0030] It is a further object of the present invention to provide methods and compositions for preventing, treating or reducing disease by eliciting an active cellular and humoral response in the host.

[0031] Yet another object of the present invention is to provide methods and compositions for reducing and preventing the occurrence of neurological disorders.

[0032] Also disclosed are methods and compositions for reducing and preventing the occurrence of hyperproliferative disorders.

[0033] Yet another object of the present invention to provide methods and compositions for therapeutic immunological intervention in neurological disorders.

[0034] Further disclosed are methods and compositions for vaccinating a human or animal against selected infectious organisms, and methods and compositions for passively immunizing a human or animal against selected infectious organisms.

[0035] Another object of the present invention is to provide an antigenic construct and compositions that are antigenic and elicit an immune response against pathological manifestation in humans or animals.

[0036] Yet another object of the present invention is to provide an antigenic construct and compositions that are antigenic and elicit an immune response against pathological manifestation in humans or animals, wherein such pathological manifestation comprises abnormalities such as amyloid plaques.

[0037] Further disclosed is an antigenic construct and compositions that are antigenic and elicit an immune response against infectious organisms in humans or animals.

[0038] Another object of the present invention is to provide vaccine compositions comprising antigenic constructs that are non-immunogenic in a human or animal to be immunized with the composition; and carriers wherein the antigenic peptide is uniquely presented on the surface of the carrier such that resulting antibodies are highly specific and have a greater degree of conformational sensitivity when administered into the human or animal.

[0039] Yet another object of the present invention is to provide methods and compositions comprising modified antigenic moieties to increase an individual's response to disease and disorders.

[0040] Another object of the present invention is to provide vaccine compositions comprising antigenic constructs wherein peptides are modified to enhance antigenic effect.

[0041] Yet another object of the present invention is to provide vaccine compositions comprising antigenic constructs comprising peptides modified to enhance antigenic effect wherein such peptides are modified via pegylation (using polyethylene glycol or modified polyethylene glycol), or modified via other methods such by poly-amino acids (e.g. poly-glycine, poly-histidine), poly-saccharides (e.g. polygalacturonic acid, polylactic acid, polyglycolide, chitin, chitosan), synthetic polymers (polyamides, polyurethanes, polyesters) or co-polymers (poly(methacrylic acid) and N-(2-hydroxy) propyl methacrylamide) and the like.

[0042] It is yet another object of the present invention to provide immunogenic compositions wherein the carrier for the antigenic peptide comprises modified liposomes.

[0043] It is another object of the present invention to provide immunogenic compositions wherein the carrier for the antigenic peptide comprises a colloidal metal.

[0044] Another object of the present invention is to provide immunogenic compositions wherein the carrier for the antigenic peptide comprises a baculovirus-derived vesicle.

[0045] It is still another object of the present invention to provide immunogenic compositions in combination with

pharmaceutically acceptable adjuvants to stimulate the immune response.

[0046] Yet another object of the present invention is to provide immunogenic compositions that may be administered intramuscularly, intravenously, transdermally, orally, or subcutaneously.

[0047] These and other objects, features and advantages of the present invention will become apparent after a review of the following detailed description of the disclosed embodiment and the appended claims.

Figure 1 provides a schematic showing chemically modified β -amyloid antigen.

Figure 2 provides representative schematic showing liposome reconstituted with a chemically modified amyloid-antigen.

Figure 3 provides a schematic showing a multiple P₁₇₀ antigen.

Figure 4 provides synthetic peptides, homologous to different segments of PrP^o used to investigate their influence on the viability of primary rat hippocampal neurons.

Figure 5 provides a schematic of the peptides derived from the A β sequences 4-11 (SEQ ID NO: 2), 1-16 (SEQ ID NO: 5), 22-35 (SEQ ID NO: 3) and 29-40 (SEQ ID NO: 4).

Figure 6 provides a schematic showing the general synthetic approaches to antigens derived from peptides sequences with or without internal His or Lys residues.

Figure 7 provides the results of ELISA conducted with 1:5000 diluted sera from pegylated amyloid/ liposomes/ lipid A immunized C57BL/6 mice. PEG-A β ₁₋₁₆ (- -black), PEG-AP₁₋₁₆ + ALUM (- -grey), PEG-AP₄₋₁₁ (—grey). Means of the values of 10 mice per antigen; means of values from 2 mice are shown for A β ₁₋₁₆ +ALUM. As a control mean values of 12 palmitoylated A β ₁₋₁₆(- - bright grey) injected animals are shown (published 2002).

Figure 8 provides the results of assays evaluating solubilization of A β ₁₋₄₂ fibers by sera of PEG-A β ₄₋₁₁ immunized C57BL/6 mice. Thioflavin fluorescence emission intensity correlates with the amount of fibrillar amyloid present in solution. A β ₁₋₄₂ fibers formation during 7 days at 37°C in PBS, pH = 7.1. Sera were added on day 7 and incubated for 24hrs. Bars 1-9 represent solubilization experiments made with sera of vaccinated animals. Means of 4 samples + SD are shown.

Figure 9 provides the results of solubilization assay of A β ₁₋₄₂ fibers by supernatants of hybridoma clones from palm-A β ₁₋₁₆ immunized C57BL/6 mice. A β ₁₋₄₂ fibers formation during 7 days at 37°C in PBS, pH = 7.1. Supernatants were incubated for 24 hrs. sfr medium = medium without FCS. The hybridoma clones were grown in serum free medium for 1 day. Means of 4 samples + SD are shown.

Figure 10 provides ¹³C-¹³C correlation spectrum of amyloid fibres made of the amyloid β -peptide labeled at ¹⁰Tyr and ¹²Val.

Figure 11 provides projection of the ¹³C-¹³C correlation spectrum of A β -peptide fibers (A) and after incubation with the antibody for 12 days (B).

Figure 12 provides NMR spectra data for assessing the effect of monoclonal antibodies on amyloid beta fibers.

Figure 13 provides a graph showing comparative data for pegylated and palmitoylated antigens.

Figure 14 provides a graph showing comparative data for pegylated beta amyloid (1-16, 4-11, 22-35, 1-15) and palmitoylated beta amyloid (1-16).

[0048] The present invention may be understood more readily by reference to the following detailed description of specific embodiments included herein. Although the present invention has been described with reference to specific details of certain embodiments, thereof, it is not intended that such details should be regarded as limitations upon the scope of the invention.

[0049] We report here a method of eliciting high immune responses, of great specificity yielding conformationally sensitive antibodies. These antibodies recognize specific epitopes on a wide variety of antigens including but not limited to, amyloid protein, prion protein, P₁₇₀ glycoprotein. More specifically, we report here the concept of modifying peptides, such as amyloid peptides, to elicit an improved immunogenic response. In certain embodiments, the peptides are modified via pegylation.

Definitions

[0050] The terms "polypeptide", "peptide", and "protein", as used herein, are interchangeable and are defined to mean a biomolecule composed of two or more amino acids linked by a peptide bond.

[0051] The term "peptides," are chains of amino acids (typically L-amino acids) whose alpha carbons are linked through peptide bonds formed by a condensation reaction between the carboxyl group of the alpha carbon of one amino acid and the amino group of the alpha carbon of another amino acid. The terminal amino acid at one end of the chain (i.e., the amino terminal) has a free amino group, while the terminal amino acid at the other end of the chain (i.e., the carboxy terminal) has a free carboxyl group. As such, the term "amino terminus" (abbreviated N-terminus) refers to the free alpha-amino group on the amino acid at the amino terminal of the peptide, or to the alpha-amino group (imino group

when participating in a peptide bond) of an amino acid at any other location within the peptide. Similarly, the term "carboxy terminus" (abbreviated C-terminus) refers to the free carboxyl group on the amino acid at the carboxy terminus of a peptide, or to the carboxyl group of an amino acid at any other location within the peptide.

5 [0052] Typically, the amino acids making up a peptide are numbered in order, starting at the amino terminal and increasing in the direction toward the carboxy terminal of the peptide. Thus, when one amino acid is said to "follow" another, that amino acid is positioned closer to the carboxy terminal of the peptide than the preceding amino acid.

[0053] The term "residue" is used herein to refer to an amino acid that is incorporated into a peptide by an amide bond. As such, the amino acid may be a naturally occurring amino acid or, unless otherwise limited, may encompass known analogs of natural amino acids that function in a manner similar to the naturally occurring amino acids (i.e., amino acid mimetics). Moreover, an amide bond mimetic includes peptide backbone modifications well known to those skilled in the art.

10 [0054] The phrase "consisting essentially of" is used herein to exclude any elements that would substantially alter the essential properties of the peptides to which the phrase refers. Thus, the description of a peptide "consisting essentially of ..." excludes any amino acid substitutions, additions, or deletions that would substantially alter the biological activity of that peptide.

[0055] Furthermore, one of skill will recognize that, as mentioned above, individual substitutions, deletions or additions which alter, add or delete a single amino acid or a small percentage of amino acids (typically less than 5%, more typically less than 1%) in an encoded sequence are conservatively modified variations where the alterations result in the substitution of an amino acid with a chemically similar amino acid. Conservative substitution tables providing functionally similar amino acids are well known in the art. The following six groups each contain amino acids that are conservative substitutions for one another:

- 1) Alanine (A), Serine (S), Threonine (T);
- 2) Aspartic acid (D), Glutamic acid (E);
- 25 3) Asparagine (N), Glutamine (Q);
- 4) Arginine (R), Lysine (K);
- 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and
- 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W).

30 [0056] The phrases "isolated" or "biologically pure" refer to material which is substantially or essentially free from components which normally accompany it as found in its native state. Thus, the peptides described herein do not contain materials normally associated with their in situ environment. Typically, the isolated, immunogenic peptides described herein are at least about 80% pure, usually at least about 90%, and preferably at least about 95% as measured by band intensity on a silver stained gel.

35 [0057] Protein purity or homogeneity may be indicated by a number of methods well known in the art, such as polyacrylamide gel electrophoresis of a protein sample, followed by visualization upon staining. For certain purposes high resolution will be needed and HPLC or a similar means for purification utilized.

[0058] When the immunogenic peptides are relatively short in length (i.e., less than about 50 amino acids), they are often synthesized using standard chemical peptide synthesis techniques.

40 [0059] Solid phase synthesis in which the C-terminal amino acid of the sequence is attached to an insoluble support followed by sequential addition of the remaining amino acids in the sequence is a preferred method for the chemical synthesis of the immunogenic peptides described herein. Techniques for solid phase synthesis are known to those skilled in the art.

[0060] Alternatively, the immunogenic peptides described herein are synthesized using recombinant nucleic acid methodology. Generally, this involves creating a nucleic acid sequence that encodes the peptide, placing the nucleic acid in an expression cassette under the control of a particular promoter, expressing the peptide in a host, isolating the expressed peptide or polypeptide and, if required, renaturing the peptide. Techniques sufficient to guide one of skill through such procedures are found in the literature.

50 [0061] Once expressed, recombinant peptides can be purified according to standard procedures, including ammonium sulfate precipitation, affinity columns, column chromatography, gel electrophoresis and the like. Substantially pure compositions of about 50 to 95% homogeneity are preferred, and 80 to 95% or greater homogeneity are most preferred for use as therapeutic agents.

[0062] One of skill in the art will recognize that after chemical synthesis, biological expression or purification, the immunogenic peptides may possess a conformation substantially different than the native conformations of the constituent peptides. In this case, it is often necessary to denature and reduce the antiproliferative peptide and then to cause the peptide to re-fold into the preferred conformation. Methods of reducing and denaturing proteins and inducing re-folding are well known to those of skill in the art.

55 [0063] Antigenicity of the purified protein may be confirmed, for example, by demonstrating reaction with immune

serum, or with antisera produced against the protein itself.

[0064] The terms "a", "an" and "the" as used herein are defined to mean "one or more" and include the plural unless the context is inappropriate.

[0065] The terms "detecting" or "detected" as used herein mean using known techniques for detection of biologic molecules such as immunochemical or histological methods and refer to qualitatively or quantitatively determining the presence or concentration of the biomolecule under investigation.

[0066] By "isolated" is meant a biological molecule free from at least some of the components with which it naturally occurs.

[0067] The terms "antibody" or "antibodies" as used herein include monoclonal antibodies, polyclonal, chimeric, single chain, bispecific, simianized, and humanized antibodies as well as Fab fragments, including the products of an Fab immunoglobulin expression library.

[0068] The term "antigen" refers to an entity or fragment thereof which can induce an immune response in a mammal. The term includes immunogens and regions responsible for antigenicity or antigenic determinants.

[0069] As used herein, the term "soluble" means partially or completely dissolved in an aqueous solution.

[0070] Also as used herein, the term "immunogenic" refers to substances which elicit or enhance the production of antibodies, T-cells and other reactive immune cells directed against an immunogenic agent and contribute to an immune response in humans or animals.

[0071] An immune response occurs when an individual produces sufficient antibodies, T-cells and other reactive immune cells against administered immunogenic compositions of the present invention to moderate or alleviate the disorder to be treated.

[0072] The term "carrier" as used herein means a structure in which antigenic peptide or supramolecular construct can be incorporated into or can be associated with, thereby presenting or exposing antigenic peptides or part of the peptide to the immune system of a human or animal. The term "carrier" further comprises methods of delivery wherein supramolecular antigenic construct compositions comprising the antigenic peptide may be transported to desired sites by delivery mechanisms. One example of such a delivery system utilizes colloidal metals such as colloidal gold.

[0073] In addition, the term "carrier" further comprises delivery mechanisms known to those skilled in the art including, but not limited to, keyhole limpet hemocyanin (KLH), bovine serum albumin (BSA) and other adjuvants. It is also to be understood that the supramolecular antigenic construct compositions of the present invention can further comprise adjuvants, preservatives, diluents, emulsifiers, stabilizers, and other components that are known and used in vaccines of the prior art. Any adjuvant system known in the art can be used in the composition of the present invention. Such adjuvants include, but are not limited to, Freund's incomplete adjuvant, Freund's complete adjuvant, polydispersed β -(1,4) linked acetylated mannan ("Acemannan"), TITERMAX[®] (polyoxyethylene-polyoxypropylene copolymer adjuvants from CytRx Corporation), modified lipid adjuvants from Chiron Corporation, saponin derivative adjuvants from Cambridge Biotech, killed *Bordetella pertussis*, the lipopolysaccharide (LPS) of gram-negative bacteria, large polymeric anions such as dextran sulfate, and inorganic gels such as alum, aluminum hydroxide, or aluminum phosphate.

[0074] Carrier proteins that can be used in the supramolecular antigenic construct compositions of the present invention include, but are not limited to, maltose binding protein "MBP"; bovine serum albumin "BSA"; keyhole limpet hemocyanin "KLH"; ovalbumin; flagellin; thyroglobulin; serum albumin of any species; gamma globulin of any species; syngeneic cells; syngeneic cells bearing Ia antigens; and polymers of D- and/or L- amino acids.

[0075] Further, the term "effective amount" refers to the amount of antigenic/immunogenic composition which, when administered to a human or animal, elicits an immune response. The effective amount is readily determined by one of skill in the art following routine procedures.

[0076] For example, supramolecular antigenic construct compositions may be administered parenterally or orally in a range of approximately 1.0 μ g to 10.0 mg per patient, though this range is not intended to be limiting. The actual amount of the composition required to elicit an immune response will vary for each individual patient depending on the immunogenicity of the composition administered and on the immune response of the individual. Consequently, the specific amount administered to an individual will be determined by routine experimentation and based upon the training and experience of one skilled in the art.

[0077] The compositions of the present invention are used to produce antibodies directed against antigenic peptides. Resulting antibodies are administered to individuals to passively immunize them against a variety of diseases or disorders, including but not limited to, Alzheimer's disease, multidrug resistant cancer or prion disease.

[0078] The immunogenic compositions of the present invention may comprise liposomes made by reconstituting liposomes in the presence of purified or partially purified or modified antigenic peptides. Additionally, peptide fragments may be reconstituted into liposomes. The present invention also includes antigenic peptide fragments modified so as to increase their antigenicity. For example, antigenic moieties and adjuvants may be attached to or admixed with the peptide. Examples of antigenic moieties and adjuvants include, but are not limited to, lipophilic muramyl dipeptide derivatives, nonionic block polymers, aluminum hydroxide or aluminum phosphate adjuvant, and mixtures thereof.

[0079] The present invention further encompasses antigenic peptides modified with hydrophobic moieties, such as

palmitic acid, that facilitate insertion into the hydrophobic lipid bilayer of a carrier. Hydrophobic moieties of the present invention may be fatty acids, triglycerides and phospholipids wherein the fatty acid carbon backbone has at least 10 carbon atoms. Most preferable are lipophilic moieties having fatty acids with a carbon backbone of at least approximately 14 carbon atoms and up to approximately 24 carbon atoms. The most preferred hydrophobic moieties have a carbon backbone of at least 14 carbon atoms. Examples of hydrophobic moieties include, but are not limited to, palmitic acid, stearic acid, myristic acid, lauric acid, oleic acid, linoleic acid, and linolenic acid. The most preferred hydrophobic moiety is palmitic acid.

[0080] The supramolecular antigenic construct compositions of the present invention are administered to a human or animal to induce immunity to antigenic agents such as infectious organisms. The immunized human or animal develops circulating antibodies against the infectious organism, thereby reducing or inactivating its ability to stimulate disease.

[0081] The supramolecular antigenic construct compositions of the present invention are also used to produce a panel of monoclonal or polyclonal antibodies that are specific for various disorders, including for example, Alzheimer's disease. Antibodies are made by methods well known to those of ordinary skill in the art.

[0082] The compositions of the present invention are administered to a human or animal by any appropriate means, preferably by injection. For example, a modified antigenic peptide reconstituted in liposomes is administered by subcutaneous injection. Whether internally produced or provided from external sources, the circulating antibodies bind to antigen and reduce or inactivate its ability to stimulate disease.

[0083] Liposomes that can be used in the compositions of the present invention include those known to one skilled in the art. Any of the standard lipids useful for making liposomes may be used. Standard bilayer and multi-layer liposomes may be used to make compositions of the present invention. While any method of making liposomes known to one skilled in the art may be used, the most preferred liposomes are made according to the method of Alving et al., *Infect. Immun.* 60:2438-2444, 1992. The liposome can optionally contain an adjuvant. A preferred adjuvant is detoxified lipid A, such as monophosphoryl or diphosphoryl lipid A.

[0084] When the vesicles are liposomes, the antigenic peptide generally has a hydrophobic tail that inserts into the liposome membrane as it is formed. Additionally, antigenic peptides can be modified to contain a hydrophobic tail so that it can be inserted into the liposome. For example, antigenic peptide may be exposed on the surface of previously formed liposomes by chemical attachment or electroinsertion.

[0085] The antibodies provided herein are monoclonal or polyclonal antibodies having binding specificity for infectious organisms or antigenic peptides representative of various disorders such as Alzheimer's disease, multi drug resistant cancer and prion diseases.

[0086] The monoclonal antibody is prepared by immunizing an animal, such as a mouse or rabbit, with supramolecular antigenic construct compositions of the present invention. Spleen cells are harvested from the immunized animals and hybridomas generated by fusing sensitized spleen cells with a myeloma cell line, such as murine SP2/O myeloma cells (ATCC, Manassas, VA). The cells are induced to fuse by the addition of polyethylene glycol. Hybridomas are chemically selected by plating the cells in a selection medium containing hypoxanthine, aminopterin and thymidine (HAT).

[0087] Hybridomas are subsequently screened for the ability to produce monoclonal antibodies against specific diseases or disorders. Hybridomas producing antibodies of interest are cloned, expanded and stored frozen for future production. The preferred hybridoma produces a monoclonal antibody having the IgG isotype, more preferably the IgG1 isotype.

[0088] The polyclonal antibody is prepared by immunizing animals, such as mice or rabbits with supramolecular antigenic construct compositions of the present invention described above. Blood sera is subsequently collected from the animals, and antibodies in the sera screened for binding reactivity against target agents.

[0089] Either the monoclonal antibody or the polyclonal antibody, or both may be labeled directly with a detectable label for identification a target agent in a biological sample as described below. Labels for use in immunoassays are generally known to those skilled in the art and include enzymes, radioisotopes, and fluorescent, luminescent and chromogenic substances including colored particles, such as colloidal gold and latex beads. The antibodies may also be bound to a solid phase to facilitate separation of antibody-antigen complexes from non-reacted components in an immunoassay. Exemplary solid phase substances include, but are not limited to, microtiter plates, test tubes, magnetic, plastic or glass beads and slides. Methods for coupling antibodies to solid phases are well known to those skilled in the art.

[0090] Alternatively, the antibody may be labeled indirectly by reaction with labeled substances that have an affinity for immunoglobulin, such as protein A or G or second antibodies. The antibody may be conjugated with a second substance and detected with a labeled third substance having an affinity for the second substance conjugated to the antibody. For example, the antibody may be conjugated to biotin and the antibody-biotin conjugate detected using labeled avidin or streptavidin. Similarly, the antibody may be conjugated to a hapten and the antibody-hapten conjugate detected using labeled anti-hapten antibody. These and other methods of labeling antibodies and assay conjugates are well known to those skilled in the art.

[0091] In a preferred embodiment, the antibody is labeled indirectly by reactivity with a second antibody that has been labeled with a detectable label. The second antibody is preferably one that binds to antibodies of the animal from which

the monoclonal antibody is derived. In other words, if the monoclonal antibody is a mouse antibody, then the labeled, second antibody is an anti-mouse antibody. For the monoclonal antibody to be used in the assay described below, this label is preferably an antibody-coated bead, particularly a magnetic bead. For the polyclonal antibody to be employed in the immunoassay described herein, the label is preferably a detectable molecule such as a radioactive, fluorescent or an electrochemiluminescent substance.

Formulations

[0092] The naturally occurring or synthetic protein, peptide, or protein fragment, containing all or an active portion of an immunogenic protein or peptide can be prepared in a physiologically acceptable formulation, such as in a pharmaceutically acceptable carrier, using known techniques. For example, the protein, peptide or protein fragment is combined with a pharmaceutically acceptable excipient to form a therapeutic composition.

[0093] Alternatively, the gene for the protein, peptide, or protein fragment, containing all or an active portion of the immunogenic peptide, may be delivered in a vector for continuous administration using gene therapy techniques. The vector may be administered in a vehicle having specificity for a target site, such as a tumor.

[0094] The compositions of the present invention may be administered in the form of a solid, liquid or aerosol. Examples of solid compositions include pills, creams, and implantable dosage units. Pills may be administered orally. Therapeutic creams may be administered topically. Implantable dosage units may be administered locally, for example, at a tumor site, or may be implanted for systematic release of the therapeutic composition, for example, subcutaneously. Examples of liquid compositions include formulations adapted for injection intramuscularly, subcutaneously, intravenously, intra-arterially, and formulations for topical and intraocular administration. Examples of aerosol formulations include inhaler formulations for administration to the lungs.

[0095] The compositions may be administered by standard routes of administration. In general, the composition may be administered by topical, oral, rectal, nasal or parenteral (for example, intravenous, subcutaneous, or intramuscular) routes. In addition, the composition may be incorporated into sustained release matrices such as biodegradable polymers, the polymers being implanted in the vicinity of where delivery is desired, for example, at the site of a tumor. The method includes administration of a single dose, administration of repeated doses at predetermined time intervals, and sustained administration for a predetermined period of time.

[0096] A sustained release matrix, as used herein, is a matrix made of materials, usually polymers which are degradable by enzymatic or acid/base hydrolysis or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes and body fluids. The sustained release matrix desirably is chosen by biocompatible materials such as liposomes, polylactides (polylactide acid), polyglycolide (polymer of glycolic acid), polylactide co-glycolide (copolymers of lactic acid and glycolic acid), polyanhydrides, poly(ortho)esters, polypeptides, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides, nucleic acids, polyamino acids, amino acids such phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene, polyvinylpyrrolidone and silicone. A preferred biodegradable matrix is a matrix of one of either polylactide, polyglycolide, or polylactide co-glycolide (co-polymers of lactic acid and glycolic acid).

[0097] The dosage of the composition will depend on the condition being treated, the particular composition used, and other clinical factors such as weight and condition of the patient, and the route of administration.

[0098] The composition may be administered in combination with other compositions and procedures for the treatment of diseases. For example, unwanted cell proliferation may be treated conventionally with surgery, radiation or chemotherapy in combination with the administration of the composition, and additional doses of the composition may be subsequently administered to the patient to stabilize and inhibit the growth of any residual unwanted cell proliferation.

Supramolecular Antigenic Constructs

[0099] The supramolecular antigenic constructs of the present invention generally comprise peptides modified to enhance antigenic effect wherein such peptides are modified via pegylation (using polyethylene glycol or modified polyethylene glycol), or modified via other methods such by palmitic acid, poly-amino acids (eg poly-glycine, poly-histidine), poly-saccharides (eg polygalacturonic acid, polylactic acid, polyglycolide, chitin, chitosan), synthetic polymers (polyamides, polyurethanes, polyesters) or co-polymers (eg. poly(methacrylic acid) and N-(2-hydroxy) propyl methacrylamide) and the like.

[0100] In certain embodiments, the supramolecular antigenic constructs of the present invention comprise a peptide sequence, covalently attached to pegylated lysine- one at each terminus. The length of the PEG (polyethyleneglycol) chain may vary from 8 to 150000. The free PEG terminus is covalently attached to a molecule of phosphatidylethanolamine (where the fatty acid can be: myristic, palmitic, stearic, oleic etc. or combination thereof). This antigenic construct may be reconstituted in liposomes consisting of phospholipids and cholesterol (phosphatidylethanol amine, phosphatidyl glycerol, cholesterol in varied molar ratios. Other phospholipids can be used. Lipid A is used at a concentration of

approximately 40 $\mu\text{g}/\mu\text{mole}$ of phospholipids.

[0101] In certain embodiments, the antigenic constructs comprise a peptide having the amino acid sequence of β -amyloid. The peptides may also comprise or correspond to whole amyloid beta peptide and active fragments thereof. Additionally, peptides useful for the present invention further comprise $\text{A}\beta_{4-11}$ (SEQ ID NO: 2), $\text{A}\beta_{22-35}$ (SEQ ID NO: 3), and $\text{A}\beta_{29-40}$ (SEQ ID NO: 4) and $\text{A}\beta_{1-16}$ (SEQ ID NO: 5); and active fragments thereof.


[0102] Also disclosed is an antigenic construct comprising peptide sequences that comprise the extracellular loops 1, 4 and 6 of the P170 glycoprotein. In certain other embodiments, the antigenic construct comprises peptide sequences that comprise amino acid sequences 109-129 of the prion protein.

[0103] The present invention further discloses monoclonal antibodies raised against an antigenic construct reconstituted in liposome, wherein, for example, the peptide sequence comprises an amino acid sequence from amyloid protein. Additionally, monoclonal antibodies raised against antigenic construct wherein the peptide sequence is an/or several amino acid sequences from the P- glycoprotein (P_{170}) extracellular loops are also disclosed in the present application.

[0104] Also disclosed in the present invention are monoclonal antibodies raised against a antigenic construct wherein the peptide sequences comprise an amino acid sequence selected from protein of interest. More specifically, for example, the invention discloses monoclonal antibodies raised against an antigenic construct reconstituted in liposome wherein the peptide sequence is an amino acid sequence selected from 13-amyloid protein (4 -10, or 1 - 8, or 8 - 16, etc.) which does not induce cerebral bleeding in transgenic mice for human Alzheimer's disease. The invention further includes monoclonal antibodies sensitive to the conformational characteristics of antigenic peptides. Specific protocols for the manufacture of the antibodies of the present invention and specific information regarding the characterization of such antibodies are provided in the Examples below.

Amyloid

[0105] The 7 amino acid sequence: FRHDSGY (SEQ ID NO:1) of β -amyloid was synthesized. One lysine was attached covalently at each end of the sequence (1). The lysines, prior to attachment to the above sequence were reacted with a chain of Polyethyleneglycol (PEG, $n = 8-2000$). Polyethyleneglycol chains bound to lysine at one end are covalently attached to a molecule of dioleoyl-phosphatidyl choline ethanolamine (or any fatty acid - phosphatidylcholine) as described (2).



[0106] The chemically modified antigen is then reconstituted in liposomes consisting of phospholipids and cholesterol (3). Examples of suitable liposomes include, but are not limited to, DOPG, DOPEA, Chol. (Lipid A was at the concentration of 40 $\mu\text{g}/\mu\text{mole}$ phospholipid.) A representative schematic showing liposome reconstituted with a chemically modified amyloid-antigen is shown in Figure 2.

[0107] The supramolecular antigenic constructs of the present invention have vast advantages over the palmitoylated antigens, reconstituted in liposomes. Primarily, the long PEG chains ($n = 8 - 5000$) enhance significantly the exposure and accessibility of the peptide sequence. Antigen presentation is improved and the conformation sensitivity of the elicited antibodies is enhanced. Another advantage of the present invention is that peptide sequences in different conformations may be used. The increased distance between the sequence and surface of the liposome makes sure that the surface does not interact with the sequence, thus, possibly influencing its conformation. Also, antigenicity of the construct becomes significantly higher than that of palmitoylated sequences reconstituted in liposomes. High titers of antibodies comprised between 1:5000 and 1:10000 are obtained in mice, within a few weeks. Additionally, the affinity of the antibodies for the antigen is significantly increased. In the case of the amyloid sequence FRHDSGY (SEQ ID NO: 1), the antibody elicited by ip or iv injection of the construct are efficiently solubilizing $\text{A}\beta_{1-40}$ and $\text{A}\beta_{1-42}$ fibers, protecting *in vitro* PC12 cells against apoptosis and metabolic inhibition (MTT reduction) induced by $\text{A}\beta_{1-42}$ and $\text{A}\beta_{1-40}$ fibers.

[0108] In one embodiment of the present invention, the FRHDSGY (SEQ ID NO: 1) sequence of the amyloid protein is used, however any other amyloid protein sequence can be substituted. Monoclonal antibodies obtained from mice immunized with the described construct display, besides the *in vitro* properties mentioned above for the polyclonal antibodies, biological activity in APP[V717I] FVB transgenic mice for human Alzheimer's Disease. Significant levels of memory restoration and of curiosity awakening in these mice are observed. The mAb does not induce bleeding in the brain of the immunized, transgenic mice.

[0109] Though not wishing to be bound by the following theory, based on *in vitro* studies of the interaction of anti-

amyloid mAb (against the 1-16 sequence, generated by the methods of the present invention) mainly of fiber solubilization and of CD spectra, it appears that the antibodies bind preferentially to β -amyloid in its α -helix conformation. This would explain the amyloid fiber solubilization effect in thermodynamic terms. Since the antibody, by binding preferentially to the α -helix, removes the α -helix amyloid from the equilibrium:



thereby increasing amounts of β -amyloid, in β -sheet conformation undergo conformational transition to the soluble α -helix form in order to re-establish the equilibrium. The stoichiometric observations made, support the idea of the mAbs directly influencing the conformation equilibrium.

[0110] As Selkoe (2002) has elaborated, Alzheimer's Disease appears as a synaptic failure. In the earlier stages of the disease memory loss may originate in such failure. It is thought that soluble oligomers $A\beta_{1-40}$, for example, might be able to block the synapse. The monoclonal antibodies, generated by methods of the present invention, bind to soluble oligomers $A\beta_{1-40}$. Measurement of conductivity of synapses in the presence and absence of the antibodies permits the determination of the action of the antibody on synapses, in the presence of soluble oligomers.

[0111] The inventors of the present invention checked the activity of a number of mAbs obtained with the epitopes such as $A\beta_{4-11}$ (SEQ ID NO: 2), $A\beta_{22-35}$ (SEQ ID NO: 3), and AP_{29-40} (SEQ ID NO: 4) embedded in an supramolecular construct (see Figure 5). The sequence 4-11 was determined to be the epitope to the mAb elicited by the palmitoylated $A\beta$ 1-16 antigen (SEQ ID NO: 5).

[0112] According to the methods of the present invention, new and uniquely modified peptide antigens were used in order to raise mAbs:

Residue 22-35: EDVGSNKGAIIGLM (SEQ ID NO:3)

[0113] Junctions between extracellular and transmembrane (TM) domains have been found to be targeted by inhibitory antibodies (such as Herceptin-Trastuzumab anti-HER2/neu antibodies) and, in multispanning TM proteins, to form pockets that are targeted by small molecular weight inhibitors (Dragic et al., 2000). Though not wishing to be bound by the following theory, this sequence is likely to be crucial for the oligomerization capacity of $A\beta_{1-42}$ and $A\beta_{1-40}$, as it represents the transition between polar and hydrophobic regions (wherein the phrase "extracellular sequence" is used to refer to the extracellular sequences in the $A\beta_{1-42}$ amyloidogenic sequence). The sequence contains the first two GXXXGXXXG motifs of the $A\beta_{1-42}$ and $A\beta_{1-40}$ sequences. GXXXG are key inducers of oligomerization of hydrophobic sequences (Russ and Engelmann, 2000). Interestingly, the first GXXXG motif is predicted to be extracellular, while the following two are predicted to be placed in the membrane. Though not wishing to be bound by the following theory, it may be assumed, by analogy, that oligomerization of $A\beta$ peptides is specifically triggered by the GXXXG motifs.

Residue 29-40: GAIIGLMVGGW (SEQ ID NO:4)

[0114] The hydrophobic sequence of $A\beta_{1-42}$ and $A\beta_{1-40}$ contain the motif GXXXGXXXGG, which has been found to induce strong oligomerization of hydrophobic sequences (Eilers et al., 2002; Leeds et al., 2001; Lemmon et al., 1994; Russ and Engelmann, 1999; Russ and Engelmann, 2000; Smith and Bormann, 1995). This motif is viewed as a prime target for therapeutic approaches; since it must play a major role in all pathogenic processes that lead to $A\beta_{1-42}$ and $A\beta_{1-40}$ formation, oligomerization and accumulation. In the intact sequence of APP, it is likely that this motif caps the downstream sequence that will need to unfold for γ -secretase to process, as was shown for the SREBP cleavage (Ye et al., 2000). This sequence has not previously been identified by anybody as being important for amyloid oligomerization. As taught herein, the supramolecular modified (preferably pegylated) antigens of the present invention have higher antigenicity and the antibodies elicited by them have higher affinities. Beside $A\beta_{1-16}$, supramolecular constructs of the present invention also include peptides represented by $A\beta_{4-11}$ (SEQ ID NO: 2), $A\beta_{22-35}$ (SEQ ID NO: 3), $A\beta_{29-40}$ (SEQ ID NO: 4) for use in vaccines.

[0115] Methodologies for the mono-pegylation of peptides at the N- α -position are known and widely used. Site-specific mono-pegylation at internal, N- or C- terminal amino-acid residues of medium sized peptides has also been described following either solid-phase or peptide-grafting approaches. However, solid-phase synthetic approaches to di-pegylated peptides have been shown to be severely hampered by steric hindrance and upon starting this project no efficient synthetic methodologies were reported for such compounds. Furthermore, peptides derivatised site-specifically at the N- and C-termini with both a PEG and lipid moiety have not previously been reported. Herein the present inventors describe a novel methodology for the synthesis of such $A\beta$ peptide conjugates.

[0116] In arriving at the present invention several approaches were attempted most of which were unsuccessful. For example, the initial approach to the synthesis focused upon the on-resin grafting of lipid-PEG conjugates containing distal amine groups, to side-chain protected peptides ($A\beta_{4-11, 1-16, 22-35}$ and $29-40$) containing terminal Glutamic acid residues. No coupling products were observed under a wide variety of reaction conditions. As described in Example 2 and shown in Figure 5, the supramolecular constructs described herein were generally synthesized using Fmoc/tBu amino acid side-chain protections.

[0117] This novel approach to the synthesis of N- and C- terminal lipid-PEG β -amyloid antigens using protected

peptides is applicable to a wide variety of peptide sequences including for example multi-drug resistance protein P-glycoprotein.

[0118] In an effort to evaluate the efficacy of the antigenic peptides described herein, experiments were conducted to compare the immunogenicity of PEGylated and palmitoylated antigens using ELISA and disaggregation assays (see Example B, and Figure 7). The ELISA data showed that liposomal PEG-A β_{1-16} is significantly more immunogenic than palmitoylated A β_{1-16} . Additional ALUM did not enhance the immunogenicity of PEG-A β_{1-16} in the mice. The antibody response induced by PEG-A β_{4-11} was slower in comparison to PEG-A β_{1-16} .

[0119] In summary therefore, provided are novel monoclonal antibodies against supramolecular antigens exposing different amyloid sequences. In particular, original synthetic pathways were devised in order to bind covalently two polyethylene glycol (n=70) chains to selected amyloid sequences. At the free end of the PEG chain, phosphatidyl ethanol amine was covalently attached. Though not wishing to be bound by the following theory, it is believed that its function is to anchor the pegylated amyloid sequence in the bilayer of liposomes. Pegylation is shown herein to increase the immunogenicity of the antigens as compared to palmitoylation. Affinity studies, epitope determination, induction of conformational transition by these monoclonal antibodies are being conducted presently in our laboratory. The unique modification methodology of the present invention is applicable to a variety of peptides and can ultimately be employed in therapeutic formulations and vaccines for diseases and disorders including, but not limited to Alzheimer's disease, cancer, and infectious disease.

[0120] As described herein, the supramolecular antigenic constructs of the present invention comprise peptides modified to enhance antigenic effect wherein such peptides are modified via pegylation (using polyethylene glycol or modified polyethylene glycol), or modified via other methods such by palmitic acid, poly-amino acids (e.g. poly-glycine, poly-histidine) poly-saccharides (e.g. polygalacturonic acid, polylactic acid, polyglycolide, chitin, chitosan), synthetic polymers (polyamides, polyurethanes, polyesters) or co-polymers (poly(methacrylic acid) and N-(2-hydroxy) propyl methacrylamide) and the like. For therapeutic intervention in neurological disorders such as Alzheimer's disease, the present invention comprises the modification of amyloid beta peptides.

Multidrug resistance 1 (MDR1) in Cancer Cells

[0121] Multidrug resistance 1 in cancer cells is caused by the overexpression of the P-glycoprotein (P₁₇₀) a membrane pump which ejects a large variety of unrelated chemotherapy agents from cancer cells.

[0122] Immunization with palmitoylated extracellular sequences of P₁₇₀, reconstituted in liposomes, led to restoration of the sensitive phenotype *in vitro* in MDR1 L₁₂₁₀ mouse leukemia cells (3). Further results have been obtained *in vivo* (Madoulet, Tosi, Nicolau et al., 2002 - unpublished results) indicating a 70% increase of survival half-life in immunized mice, inoculated with MDR cancer cells, undergoing chemotherapy.

[0123] It is shown herein that an antigen consisting of the P₁₇₀ extracellular sequences 1,4 and 6 constructed, according to the method of the present invention is far more efficient in eliciting antibodies which largely revert the MDR phenotype to the sensitive phenotype *in vitro* and *in vivo*.

[0124] Peptides corresponding to P₁₇₀ extracellular loops 1, 4 and 6 were synthesized and then attached to pegylated lysines - 1 at each end - which in turn were covalently attached to one dioleoyl phosphatylethanolamine molecule at each end. Any fatty acid, myristic, palmitic, stearic or polyunsaturated fatty acids may be used

[0125] These 3 constructs were reconstituted in liposomes consisting of PC-PEA-PG-Cholesterol (or any other phospholipid and cholesterol combination). Lipid A was added at concentration of 40 $\mu\text{g}/\mu\text{mole}$ of phospholipids. The ratio peptide: phospholipid was 1 : 200 (other ratios may be used).

[0126] The length of the polyethyleneglycol chains varied: the longer the peptide sequence, the higher the number of PEG molecules in the chain needs to be. For the 3 sequences used, the PEG chain-length varied from 10 to 5000. Other chain lengths can be used. Figure 3 provides a representative schematic showing a multiple P₁₇₀ antigen.

[0127] IP inoculation of this antigen, followed by three boostings at 2 weeks interval elicited high titres of anti P₁₇₀ antibodies (1:5000-1:10000) capable of blocking the pumping activity of P₁₇₀, *in vitro* and *in vivo*.

Prion Diseases

[0128] Prions cause neurodegenerative diseases such as scrapie in sheep, bovine spongiform encephalopathy in cattle and Creutzfeldt - Jacob- Disease in humans. The only known component of the particle is the scrapie isoform of the protein, PrP^{Sc}. Although prions multiply, there is no evidence that they contain nucleic acid. PrP^{Sc} is derived from the non-infectious, cellular protein PrP^C by a posttranslational process during which PrP^C undergoes a profound conformational change.

[0129] The scrapie protein, PrP^{Sc} has a critical role in neuronal degeneration and during disease development undergoes a three stage transition as follows: (normal cellular isoform of protein) PrP^C - infectious form (scrapie isoform of protein) PrP^{Sc} - protein PrP27-30. Such a cascade of events occurs during the development of Creutzfeldt - Jacob

Disease (CJD), Kuru, Gerstmann - Straussler-Scheinker Syndrome (GSS), fatal familial insomnia in man, scrapie in sheep and goats, encephalopathy in mink and bovine spongiform encephalopathy in cattle.

[0130] The cellular non-toxic protein (PrP^c) is a sialoglycoprotein of MW 33-35 K that is expressed predominantly in neurons. In the diseases mentioned above, PrP^c is converted into an altered form (PrP^{Sc}), which is distinguishable from its normal homologue by its relative resistance to protease digestion. PrP^{Sc} accumulates in the central nervous system of affected animals and individuals and its protease-resistant core aggregates extracellularly. The molecular basis of the pathogenesis is not understood.

[0131] Very interesting observations were made concerning the neurotoxicity of a fragment of the protein, which may have a bearing on the understanding of the mechanism of nerve cell-degeneration occurring in related encephalopathies.

[0132] On the basis of the observation, that the (J-amyloid fragment responsible for the extracellular deposition of amyloid fibrils and plaques in the Alzheimer Disease is neurotoxic, it was hypothesized that neuronal death in related encephalopathies might be due to toxic effects of abnormal extracellular accumulation of PrP^{Sc} and/or its degradation products.

[0133] Synthetic peptides, homologous to different segments of PrP^c were used to investigate their influence on the viability of primary rat hippocampal neurons (Figure 4)

[0134] The present inventors demonstrated that neuronal death occurs from chronic exposure of primary rat hippocampal cultures to micromolar concentrations of a peptide corresponding to residues 106-126 of the amino-acid sequence deduced from human PrP^c cDNA, in a concentration dependent manner (Example 1).

[0135] As detailed in Example 1, the inventors showed that the neuronal death induced by PrP 106-126 occurred by apoptosis in a dose dependent manner. In the terminal stages of subacute encephalopathies, such as scrapie, PrP^{Sc} reaches at whole brain concentrations 10 to 20 times higher than PrP^c, which resembles strikingly the data listed in Table 1 for the 2 concentrations of PrP106-126.

[0136] The process of programmed cell death induced by PrP106-126 is associated, among others with the induction of the testosterone - repressed prostate message - 2 gene (TRPM-2). It is not known whether apoptosis is activated *in vivo* during -related encephalopathies, but the expression of the TRPM-2 mRNA is increased 10-fold in scrapie-infected hamsters.

[0137] It appears from these data, that a neurotoxic mechanism is possibly responsible for neuronal cell loss in related encephalopathies and could also be relevant in Alzheimer's disease.

[0138] The possible mechanism of this neurotoxicity was investigated in a model system aiming at detecting and analyzing ionic channel formations upon the interaction of peptides or proteins with lipid bilayers.

[0139] Low pH, which favors channel formation by PrP106-126, converts also this peptide from ahelical to R-sheet conformation. Whereas peptide mapping of PrP^{Sc} with Edman sequencing and mass spectrometry revealed no differences between its amino acid sequence and that predicted from the PrP^c gene sequence; no chemical modifications were found that might distinguish PrP^{Sc} from PrP^p; Fourier Transform infrared spectroscopy and circular dichroism spectroscopy revealed however a significant conformational difference between PrP^{Sc} and PrP^p.

[0140] PrP^c is essentially α -helical with little or no R-sheet, whereas PrP^{Sc} has a high β -sheet content and less α -helical structure.

[0141] The sequence KTNMKHMAGAAAAGAVVGGLG (PrP106-126) (SEQ ID NO: 6) is not only very hydrophobic but it converts also, at low pH to β -sheet conformation. Moreover, it can convert in solution, other peptides to β -sheet conformation.

[0142] Based upon these observations and upon techniques developed by the inventors, a "vaccine" was developed against diseases by eliciting a strong humoral and cellular immune response in mice to the neurotoxic PrP106-126, and then challenge the immunized mice with brain extracts from scrapie mice.

[0143] As in the previous examples, pegylated lysines were attached covalently at each end of the PrP106-126 sequence. The length of the PEG chain was 12 - 4000. The PEG chains were coupled each other to one molecule of phosphatidyl ethanol amine and reconstituted in PG-PEA-chol liposomes - lipid A.

[0144] Injected into mice these supramolecular antigenic constructs elicited a strong humoral immune response, yielding antibodies with high affinity for the PrP106-126 sequence, and having solubilizing effects within.

[0145] It should be understood that the foregoing relates only to preferred embodiments of the present invention, and that numerous modifications or alterations may be made therein without departing from the scope of the invention as set forth in the appended claims.

EXAMPLE 1

[0146] The present inventors demonstrated that neuronal death occurs from chronic exposure of primary rat hippocampal cultures to micromolar concentrations of a peptide corresponding to residues 106-126 of the amino-acid sequence deduced from human PrP^c cDNA, in a concentration dependent manner. The data are shown in Table 1.

Table 1.

Chronic treatment of hippocampal neurons for 9 days		
Peptide	cell death % 20 μ m	80 μ m
PrP 106-126	18 \pm 8	100 \pm 8
PrP 57-64	0 \pm 5	3 \pm 4
PrP 89-106	5 \pm 2	2 \pm 6
PrP 106-114	0 \pm 3	12 \pm 6
PrP 127-135	3 \pm 6	15 \pm 9
PrP 127-147	1 \pm 7	18 \pm 7
PrP 106-126 scrambled	3 \pm 2	8 \pm 3

[0147] The data are the means \pm s.e. of 6-10 determinations and are normalized to the toxic effect of PrP106-126 (designated 100% response).

[0148] It was shown that the neuronal death induced by PrP 106-126 occurred by apoptosis in a dose dependent manner. In the terminal stages of subacute encephalopathies, such as scrapie, PrP^{Sc} reaches at whole brain concentrations 10 to 20 times higher than PrP^C, which resembles strikingly the data listed in Table 1 for the 2 concentrations of PrP106-126.

[0149] The process of programmed cell death induced by PrP106-126 is associated, among others with the induction of the testosterone - repressed prostate message - 2 gene (TRPM-2). It is not known whether apoptosis is activated *in vivo* during -related encephalopathies, but the expression of the TRPM-2 mRNA is increased 10-fold in scrapie-infected hamsters.

EXAMPLE 2

Methods for Making Supramolecular Antigenic Constructs

[0150] The supramolecular constructs described herein were uniquely synthesized using standard Fmoc/tBu amino acid side-chain protections. Peptides which are modified at both the C- and N-terminus by a PEG-lipid moiety have not previously been reported. Typically, pegylation of peptides results in mixtures of regioisomers. The inventors herein demonstrate a convenient method for the site-specific attachment of a PEG-lipid conjugate to both the C- and N-terminus of A β using partially protected peptides.

[0151] For those peptide sequences containing internal Lys or His residues (4-11, 1-16, 22-35), an orthogonally protected Lys(ivDde) was added to each termini. An additional Gly was added to the C-terminal to facilitate synthesis. The Fmoc group was removed with 20 % piperidine in DMF and N-acetylated using acetic anhydride. Selective cleavage of the ivDde groups was achieved with 3 % hydrazine hydrate in DMF for one hour. The 2-chlorotrityl resin was favored over the more widely used Wang resin since the former proved to be much more resistant to hydrazinolysis. Furthermore, the 2-chlorotrityl resin is extremely acid sensitive and thus, unlike the Wang resin, enables the isolation of protected peptides. Indeed, it was necessary to perform the coupling reaction in the solution phase as coupling of the resin-bound peptide to the pre-activated pegylated lipid reagent DSPE-PEG-SPA did not give rise to any coupling product. Thus selective cleavage from the resin under mild conditions (acetic acid / trifluoroethanol / dichloromethane, 1:1:8, 1h, rt) gave the internally protected peptides (Figure 5).

[0152] Solution-phase couplings were achieved successfully with the peptides derived from sequences A β ₄₋₁₁ (SEQ ID NO: 2), A β ₁₋₁₆ (SEQ ID NO: 5), A β ₂₂₋₃₅ (SEQ ID NO: 3), to DSPE-PEG-SPA in DMSO and excess base (Figure 6). The reactions were then quenched by the addition of excess ethanolamine for 2 h and the solution lyophilized. Purification by HPLC (semi-preparative reverse-phase C₄ column) gave between 50-70 % purity of the N- and C- terminal PEG-lipid conjugates whose identities were confirmed by MALDI (matrix assisted laser desorption ionization). Each sequence showed considerable variation in the ease of the coupling reaction and conditions were adjusted accordingly (temperature, number of molar equivalents DSPE-PEG-SPA, time). Purification by HPLC proved excellent for the separation of excess DSPE-PEG-SPA from the desired product, however since the former shows no affinity to the column, separation of mono-PEG-lipid (both N- and C- terminal) peptide products from the desired product proved difficult. Attempts to separate these products using size-exclusion chromatography also proved unsuccessful, presumably due to their relatively large polydispersities. Nevertheless the present inventors are using cation-exchange chromatography to separate the mono-

and di- coupled products before final side-chain deprotections. Subsequent peptide side-chain deprotections and separation of the excess quenched DSPE-PEG-SPA enables the isolation of the desired conjugates to much higher purity.

EXAMPLE 3

Comparison of Immunogenicity of PEGylated and Palmitoylated antigens, ELISA and Disaggregation assays

[0153] Liposomal antigens were prepared as described above. The sequences PEG-A β_{1-16} , -A β_{4-11} and -A β_{22-35} were reconstituted in a construct consisting of liposomes made of dimyristoyl phosphatidyl choline (DMPC), dimyristoyl phosphatidyl ethanolamine (DMPEA), dimyristoyl phosphatidyl glycerol (DMPG) and cholesterol (0.9: 0.1: 0.1: 0.7 molar ratios) containing monophosphoryl lipid A (40mg/mM phospholipids).

ELISA

[0154] The antigens and palmitoylated A β_{1-16} were used for the immunization in C57BL/6 mice in 2 week intervals. 10-12 animals were immunized with each antigen. Sera were taken 5 days after the boostings and ELISA were conducted with several dilutions of the sera. Comparative results showing the immunogenicity of the different antigens are presented in Figure 7.

[0155] The ELISA data showed that liposomal PEG-A β_{1-16} is significantly more immunogenic than palmitoylated A β_{1-16} . Additional ALUM did not enhance the immunogenicity of PEG-A β_{1-16} in the mice. The antibody response induced by PEG-A β_{4-11} was slower in comparison to PEG-A β_{1-16} .

Disaggregation Assays

[0156] Nine sera (1:100 dilution) from the liposomal-PEG-A β_{4-11} immunized animals were used in an assay where pre-formed A β_{1-42} fibers were incubated with the antisera. The assay was performed as described (Nicolau et al., 2002).

[0157] Solubilization of A β_{1-42} fibers by the different sera was observed with an incubation time of 24 hrs (Figure 8) Some of the sera solubilized the fibers to an extent of 75% (sera from mouse 5 and 6). The spleen cells of these mice were used for the production of monoclonal antibodies.

EXAMPLE 4

Solubilization Assay

[0158] From two palmitoylated A β_{1-16} /liposomes/ lipid A- immunized animals, 25 supernatants were obtained from recently generated hybridoma clones which were shown to be specific for A β_{1-42} specific antibodies. They were tested in a solubilization assay according to methods and protocols described in PNAS 2002, 99, 2332-2337. The results are summarized in Figure 9.

[0159] The supernatants of 5 hybridoma clones were found to be able to solubilize β - amyloid fibers *in vitro* to an extent of up to 75%. The two best clones 15 and 27 were chosen for the purification of monoclonal antibodies. They are being used for further investigations as positive control mAbs *in vivo*.

EXAMPLE 5

Investigation of the β -sheet to α -helix transition of the A β_{1-42} -peptide by solid state NMR spectroscopy

[0160] To avoid loss of ^{13}C -labelled amino acids the synthesis of the A β_{1-42} by Fmoc peptide synthesis was verified by a test-synthesis without labeled amino acids. The identity of the obtained A β_{1-42} peptide could be verified by MALDI mass spectroscopy and a purification procedure using HPLC with a reversed phase column and an ammonia buffered acetonitrile water gradient⁴ could be established.

[0161] The successful setup of a protocol for synthesis and purification of the amyloid β -peptide is followed by the synthesis of the labeled peptide including a ^{13}C labeled valine at position 12 (^{12}val) and a ^{13}C labeled tyrosine at position 10 (^{10}tyr).

[0162] The labeled A β_{1-42} was used to generate fibers by incubating the peptide solution in PBS buffer for one week at 37°C. ^{13}C NMR spectra of the lyophilized fibers confirm the β -sheet structure and are in agreement with published results. Incubation of the fibers with A β_{1-16} specific antibody for 2 days did not show a significant change ^{13}C spectrum. First assessments of NMR measurements indicate a change in the secondary structure (Figure 10).

EXAMPLE 6

Antibodies Elicited by Supramolecular Antigenic Constructs

5 **[0163]** *Manufacturing of the mAbs* : Lipo somal antigens were prepared as described (Nicolau et al., 2002, PNAS, 99, 2332-37). The sequences PEG-A β_{1-15} , -A β_{1-16} , A β_{4-11} -A β_{22-35} and A β_{29-40} were reconstituted in a construct consisting of liposomes made of dimyristoyl phosphatidyl choline (DMPC), dimyristoyl phosphatidyl ethanolamine (DMPEA), dimyristoyl phosphatidyl glycerol (DMPG) and cholesterol (0.9: 0.1: 0.1: 0.7 molar ratios) containing monophosphoryl lipid A (40mg/mM phospholipids). These antigens and palmitoylated A β_{1-16} were used for the immunization in C57BL/6 mice in 2 week intervals. 10-12 animals were immunized with each antigen. After 3 to 6 boostings, mice with therapeutic titers (when a 1:5,000 dilution of the sera were positive in ELISA) were selected for a fusion. The fusion of the mice's B-lymphocytes from the spleens were conducted with the myeloma cell line SP2-0. IgG producing hybridoma clones were selected and tested for their specific binding to the A β_{1-42} peptide by ELISA.

15 **[0164]** *Characterization of the mAb*: Disaggregation assays, NMR-studies, QELS- and SPR- measurements were used for the characterization of the mAbs. The mAbs showed a disaggregation of preformed amyloid fibers of up to 80% (Table 1). A transition of β -sheet to α -helix of amyloid was observed induced by the antibodies (Figure 11, 12). The measurements by Quasi Elastic Light Scattering (QELS) showed that incubation of amyloid fibers with the monoclonal antibodies resulted in fibers with a size of ≤ 800 nm, (40 - 60% of all the fibers present) while amyloid alone gave only very large aggregates ($> 4\mu\text{m}$).

Table 1

mAb obtained by immunization of PEG-A β_{1-15} , -A β_{4-11} and Palin-A β_{1-16}			
Antigen	Mab/hybridoma	dissaggregation [%]	dissociation constants
PEG A β_{1-15}	ET-1H6	75	nd
PEG A β_{1-15}	ET-7E3	55	2×10^{-7}
PEG A β_{4-11}	EN-4H7	80	nd
PEG A β_{4-11}	EN-9H2	35	nd
Palm AP $_{1-16}$	EJ-1A9	55	8×10^{-8}
Palm AP $_{1-16}$	EJ-7H3	60	7×10^{-8}
Palm A β_{1-16}	AN9C-E4	60	nd

20 *NMR studies*: In order to assess the effect of the mAb on fibers a solution of A β -fibers was incubated with the antibody for 12 days. Proton Driven Spin Diffusion measurements (PDSD) were performed to measure a 2D ^{13}C - ^{13}C correlation spectrum. Figures 11-13 provide NMR data and analysis: NMR: ^{13}C - ^{13}C correlation spectrum of Amyloid β -peptide fibers before and after incubation with EN4H7, ET-1H6 or AN9C-E4 for 12 days (A, B). Column (D) and a row (C) extracted of the correlation spectrum. The spectrum (- antibody) shows a spectrum of the pure fibres whereas (+ antibody) shows a spectrum in presence of the mAb.

25 **[0165]** ^{13}C spectra of the lyophilized fibres enables the assignment of the resonances. The chemical shift values of C_α and C_β of Val12 and Tyr10 confirms the β -sheet structure of the pure fibres. (Figure 11) The shifts of the resonances of the C_α and of the C_β nucleus of ^{12}Val clearly indicate a transition from β -sheet to α -helix. The shift of the resonance of C_α and C_β of Tyr 10 on the other hand does not clearly indicate a transition of the secondary structure. The behavior for the Tyr10 resonance can be explained by a model where Tyr10 is at the interface between a β -sheet section and a loop section of the A β -peptide in amyloid fibers.

30 **[0166]** In a second experiment ^{13}C -enriched fibers were incubated with EN4H7 and ET1H6 antibodies (see Table 1) for 12 days. The incubated fibers exhibit shifts of the resonances of the C_β nucleus of ^{12}Val from 32ppm to 28ppm, indicating a transition from β -sheet to α -helix for a significant fraction of the peptide (Figure 12 and 13) for both antibodies. The resonances of C_α and C_β for ^{10}Tyr get broad in the presence of the antibodies. This indicates a transition to a rather unstructured confirmation for the position of ^{10}Tyr .

EXAMPLE 7

Comparison of Immunogenicity of PEGylated and Palmitoylated antigens tested in ELISA

5 **[0167]** ELISA data showed that liposomal PEG-A β_{1-16} is more immunogenic than palmitoylated A β_{1-16} (Figure 14). Additional ALUM did not enhance the immunogenicity of PEG-A β_{1-16} in the mice. With the exception of the antibody response induced by PEG-A β_{4-11} , which was slower in comparison to PEG-A β_{1-16} , in general pegylated peptides appear to be more immunogenic than palmitoylated peptides (Figure 14).

10 EXAMPLE 8

Behavioral Tests for Evaluating Efficacy of Antibodies

15 **[0168]** In order to evaluate the efficacy of antibodies elicited by the methods described herein, namely elicited by use of supramolecular antigenic constructs comprising modified amyloid peptides (such as pegylated amyloid peptides), mice will be treated and then evaluated using the behavior tests outlined below.

Morris Water Maze

20 **[0169]** The pool (a white, circular vessel 1 m in diameter) contains water at 20°C with titanium-dioxide as an odorless, nontoxic additive to hide the escape platform (1 cm beneath the water level). Swimming of each mouse is videotaped and analyzed (Ethovision, Noldus information Technology, Wageningen, the Netherlands). Prior to training, each mouse is placed on top of the platform for 15 seconds. For place navigation tests, mice are trained to locate the hidden platform in five blocks of three trials over three consecutive days. Each trial consists of a forced swim test of maximum 120
25 seconds, followed by 60 seconds of rest. The time each mouse needed for location of the platform is measured. The five consecutive trials result in a learning curve.

[0170] 24 hours after the last training, each animal has a probe trial with the platform removed. Mice are allowed to search for 60 seconds and quadrant search time and crossings of the original platform position is measured.

30 **[0171]** Mice that refuse to swim and search the platform, but instead wait until the performer takes them out of the pool, the so-called "floaters" are to be excluded from analysis.

Open Field

35 **[0172]** A Plexiglas open-field box (52 x 52 x 40 cm) with black vertical walls and a translucent floor, dimly illuminated by a lamp placed underneath the box is used for the test. Different areas are allocated by a computerized system (Ethovision, Noldus information Technology, Wageningen, the Netherlands): the corners (9 x 9 cm), the four sides of the box (9 cm from wall) and the center of the open field box (43 x 43 cm). Each mouse is videotaped and the activity is analyzed (Ethovision) by measuring the distance (cm) of the track, the velocity (cm/sec) of the mouse, the duration/
40 time (sec) spent in the center compared to the border (corners + sides) and the frequency (N) of crossings between both areas. Each mouse is placed in the middle of the box and is allowed to explore the box for ten minutes. In between tests the open-field box is cleaned and dried before the introduction of a new mouse.

Novel Object Recognition Test

45 **[0173]** Mice are familiarized for one hour to a Plexiglas open-field box (52 x 52 x 40 cm) with black vertical walls and a translucent floor, dimly illuminated by a lamp placed underneath the box. The next day the animals are placed in the same box and submitted to a 10 minutes acquisition trial. During this trial mice are placed individually in the open field in the presence of object A (blue ball or red cube, similar sized of ± 4 cm), and the frequency exploring object A (when the animals snout is directed towards the object at a distance of < 1 cm and the mice are actively sniffing in the direction
50 of the object) is recorded (Freq_{AA}). During a 10 minutes retention trial (second trial) which is performed 3 hours later, a novel object (object B, red cube or blue ball) is placed together with the familiar object (object A) into the open field. The frequency the animal explored the two objects is recorded (Freq_A and Freq_B).

55 **[0174]** The recognition index (RI) defined as the ratio of the frequency in which the novel object is explored over the frequency in which both objects were explored $[\text{Freq}_B / (\text{Freq}_A + \text{Freq}_B) \times 100]$ is used to measure non-spatial memory. The frequency object A is explored during the acquisition trial is used to measure curiosity.

References

[0175]

- 5 1. C. Nicolau, R. Greferath, T.S. Balaban, J. Lazarte and R. Hopkins (2002). Proc. Natl. Acad. Sci. USA. 99,2332 - 2337.
2. Fluka AG (2002) Cat. # 79898.
- 10 3. P.-F. Tosi, D. Radu, and C. Nicolau (1995). Biochem. Biophys. Res. Chem. 212, 494-500.
4. Fukuda H, Shimizu T, Nakajima M, Mori H, Shirasawa T. Bioorg. Med. Chem.Lett. 1999; 9: 953-956
- 15 5. Petkova AT, Ishii Y, Balbach JJ, Antzutkin ON, Leapman RD, Delaglio F, Tycko R. Proc. Natl. Acad. Sci. U.S.A 2002; 99: 16742-16747

Claims

- 20 1. A composition comprising an antigenic construct wherein the antigenic construct comprises
- a) an antigenic peptide or active fragment thereof, wherein the antigenic peptide or active fragment thereof is modified to have a polyethylene glycol covalently attached, one at each terminus, and
- b) wherein the antigenic construct is reconstituted in liposomes; to enhance antigenicity.
- 25 2. The composition of Claim 1 comprising an antigenic construct wherein the: antigenic construct comprises
- a) an antigenic peptide or active fragment thereof, wherein the antigenic peptide or active fragment thereof is modified to have a pegylated lysine covalently attached, one at each terminus, and
- b) wherein the antigenic construct is reconstituted in liposomes to enhance antigenicity.
- 30 3. The composition of Claim 1 or 2, wherein the antigenic construct comprises: a peptide sequence, covalently attached to pegylated lysine-one at each terminus; wherein the free PEG terminus is covalently attached to a molecule of phosphatidylethanolamine.
- 35 4. The composition of Claim 1-3, wherein the antigenic construct is reconstituted in liposomes consisting of phospholipids and cholesterol.
5. The composition of any of Claims 1-4, wherein the antigenic peptide comprises an amyloid peptide.
- 40 6. The composition of Claim 5, wherein the amyloid peptide comprises:
- (a) PileArgHisAspSerGlyTyr ;
- (b) PheArgHisAspSerGlyTyrGlu (A β 4-11);
- (c) GluAspValGlySerAsnLysGlyAlalalleGlyLeuMet (A β 22-35);
- (d) GlyAlalalleGlyLeuMetValGlyGlyValVal (A β 29-40); or
- (e) AspAlaGluPheArgHisAspSerGlyTyrGluValHisHisGlnLys (A β 1-16)
- 45 7. The composition of any of Claims 1-4, wherein the antigenic peptide is a prion protein comprising LysThrAsnMetLysHisMetAlaGlyAlaAlaAlaAlaGlyAlaValValGlyGlyLeuGly
- 50 8. The composition of any of Claims 1-7, wherein the polyethylene glycol has a length of n=8-5000.
9. The composition of any of Claims 1-8, further comprising a pharmaceutical carrier.
- 55 10. The composition of any of Claims 1-9 for use in reducing and preventing the occurrence of neurological disorders.
11. The composition of Claim 10, wherein said disorder comprises Alzheimer's disease.

12. The composition of Claim 10, wherein said disorder comprises prion diseases
13. The composition of Claim 1 or 2 for use in the treatment of multidrug resistant cancer.
- 5 14. The composition of any of Claims 1-9 for inducing an immune response in humans or animals.
15. The composition of Claim 14, wherein said immune response is elicited against pathological manifestations in humans or animals comprising abnormalities such as amyloid plaques.
- 10 16. Use of a composition of any of claims 1-9 for the preparation of a medicament for use in reducing and preventing the occurrence of
- a) a neurological disorders; or
- b) disorders comprising Alzheimer's disease, multidrug resistant cancer or prion disease.
- 15

Patentansprüche

- 20 1. Zusammensetzung umfassend ein antigenes Konstrukt, wobei das antigene Konstrukt umfasst:
- a) ein antigenes Peptid oder aktives Fragment davon, wobei das antigene Peptid oder aktive Fragment davon so modifiziert ist, dass es ein Polyethylenglycol kovalent abgehängt enthält, eines an jedem Terminus, und
- b) wobei das antigens Konstrukt in Liposomen rekonstituiert ist;
- 25 zur Verstärkung der Antigenizität.
2. Zusammensetzung gemäß Anspruch 1 umfassend ein antigens Konstrukt wobei das antigene Konstrukt umfasst
- a) ein antigenes Peptid oder aktives Fragment davon, wobei das antigene Peptid oder aktive Fragment davon so modifiziert ist, dass es ein pegyliertes Lysin kovalent angehängt enthält, eines an jedem Terminus, und
- 30 b) wobei das antigene Konstrukt in Liposomen rekonstituiert ist; zur Verstärkung der Antigenizität.
3. Zusammensetzung gemäß Anspruch 1 oder 2, wobei das antigene Konstrukt umfasst: eine Peptidsequenz, kovalent abgehängt an pegyliertes Lysin , eines an jedem Terminus;
- 35 wobei der freie PEG-Terminus kovalent an ein Molekül von Phosphatidylethanolamin abgehängt ist.
4. Zusammensetzung gemäß Anspruch 1 - 3, wobei das antigene Konstrukt in Liposomen bestehend aus Phospholipiden und Cholesterol rekonstituiert ist.
- 40 5. Zusammensetzung gemäß einem der Ansprüche 1 - 4, wobei das antigene Peptid ein amyloides Peptid umfasst.
6. Zusammensetzung gemäß Anspruch 5, wobei das amyloid Peptid umfasst:
- (a) PheArgHisAspSerGlyTyr;
- 45 (b) PheArgHisAspSerGlyTyrGlu (A β 4-11);
- (c) GluAspValGlySerAsnLysGlyAlalleklleGlyLeuMet (A β 22-35);
- (d) GlyAlalleklleGlyLeuMetValGlyGlyValVal (A β 29-40); oder
- (e) AspAlaGluPheArgHisAspSerGlyTyrGluValHisHisGlnLys (A β 1-16).
- 50 7. Zusammensetzung gemäß einem der Ansprüche 1 - 4, wobei das antigene Peptid ein Prionprotein ist umfassend LysThrAsnMetLysHisMetAlaGlyAlaAlaAlaAlaGlyAlaValValGlyGlyLeuGly.
8. Zusammensetzung gemäß einem der Ansprüche 1 - 7, wobei das Polyethylenglycol eine Länge von n = 8 - 5000 hat.
- 55 9. Zusammensetzung gemäß einem der Ansprüche 1 - 8, weiter umfassend einen pharmazeutischen Träger.
10. Zusammensetzung gemäß einem der Ansprüche 1 - 9 zur Verwendung bei der Reduktion und Prävention des Auftretens von neurologischen Erkrankungen.

EP 1 763 364 B9

11. Zusammensetzung gemäß Anspruch 10, wobei die Erkrankung die Alzheimer Krankheit umfasst.
12. Zusammensetzung gemäß Anspruch 10, wobei die Erkrankung Prionkrankheiten umfasst.
- 5 13. Zusammensetzung gemäß Anspruch 1 oder 2 zur Verwendung in der Behandlung von mehrfach Arzneimittel-resistentem Krebs.
14. Zusammensetzung gemäß einem der Ansprüche 1 - 9 zur Induktion einer Immunantwort in Menschen oder Tieren.
- 10 15. Zusammensetzung gemäß Anspruch 14, wobei die Immunantwort gegen pathologische Erscheinungsformen umfassend Anomalien wie amyloide Plaques in Menschen oder Tieren ausgelöst wird.
16. Verwendung von einer Zusammensetzung gemäß einem der Ansprüche 1 - 9 zur Herstellung eines Arzneimittels zur Verwendung bei der Reduktion und Prävention des Auftretens von:
- 15 a) einer neurologischen Erkrankung; oder
b) Erkrankungen umfassend Alzheimer-Krankheit, mehrfach Arzneimittelresistenten Krebs oder Prionkrankheit.

20 Revendications

1. Une composition comprenant une construction antigénique, ladite construction antigénique comprenant
- 25 a) un peptide antigénique ou un de ses fragments actifs, le peptide antigénique ou un de ses fragments actifs étant modifié pour que vienne s'y lier de manière covalente un polyéthylène glycol, un à chaque extrémité; et
b) la construction antigénique étant reconstituée dans des liposomes ;
- pour améliorer l'antigénicité.
- 30 2. La composition selon la revendication 1, comprenant une construction antigénique, ladite construction antigénique comprenant
- a) un peptide antigénique ou un de ses fragments actifs, le peptide antigénique ou un de ses fragments actifs étant modifié pour que vienne s'y lier de manière covalente une lysine pégylée, une à chaque extrémité; et
- 35 b) la construction antigénique étant reconstituée dans des liposomes;
- pour améliorer l'antigénicité.
- 40 3. La composition selon la revendication 1 ou 2, dans laquelle la construction antigénique comprend une séquence peptidique, liée de manière covalente à une lysine pégylée, une à chaque extrémité, l'extrémité PEG libre étant liées de manière covalente à une molécule de phosphatidyléthanolamine.
4. La composition selon les revendications 1 à 3, dans laquelle la construction antigénique est reconstituée dans des liposomes constitués de phospholipides et de cholestérol.
- 45 5. La composition selon l'une quelconque des revendications 1 à 4, dans laquelle le peptide antigénique comprend un peptide amyloïde.
6. La composition selon la revendication 5, dans laquelle le peptide amyloïde comprend:
- 50 (a) PheArgHisAspSerGlyTyr;
(b) PheArgHisAspSerGlyTyrGlu (A β 4-11);
(c) GluAspValGlySerAsnLysGlyAlaIleGlyLeuMet (A β 22-35);
(d) GlyAlaIleGlyLeuMetValGlyGlyValVal (A β 29-40); ou
55 (e) AspAlaGluPheArgHisAspSerGlyTyrGluValHisHisGlnLys (A β 1-16).
7. La composition selon l'une des revendications 1 à 4, dans laquelle le peptide antigénique est une protéine prion comprenant:

EP 1 763 364 B9

LysThrAsnMetLysHisMetAlaGlyAlaAlaAlaAlaGlyAlaValValGlyGlyLeuGly.

- 5
8. La composition selon l'une des revendications 1 à 7, dans laquelle le polyéthylène glycol possède une longueur $n = 8-5000$.
9. La composition selon l'une des revendications 1 à 8, comprenant en outre un véhicule pharmaceutique.
10. La composition selon l'une des revendications 1 à 9, pour une utilisation à but de réduire et de prévenir l'occurrence de maladies neurologiques.
- 10
11. La composition selon la revendication 10, dans laquelle ladite maladie comprend la maladie d'Alzheimer.
12. La composition selon la revendication 10, dans laquelle ladite maladie comprend des maladies à prions.
- 15
13. La composition selon la revendication 1 ou 2, à utiliser dans le traitement d'un cancer manifestant une multirésistance aux médicaments.
14. La composition selon l'une des revendications 1 à 9, destinée à induire une réponse immunitaire dans des êtres humains ou des animaux.
- 20
15. La composition selon la revendication 14, dans laquelle ladite réponse immunitaire est déclenchée en présence de manifestations pathologiques dans des êtres humains ou des animaux, comprenant des anomalies telles que des plaques amyloïdes.
- 25
16. L'utilisation d'une composition selon l'une quelconque des revendications 1 à 9, pour la préparation d'un médicament à utiliser pour la réduction et la prévention de l'occurrence
- 30
- a) des maladies neurologiques ;ou
b) des maladies comprenant la maladie d'Alzheimer, un cancer manifestant une multirésistance aux médicaments ou une maladie à prion.

35

40

45

50

55

Figure 1

Chemically Modified β -amyloid Antigen




Figure 2
Liposome Reconstituted, Chemically Modified Amyloid-Antigen




Figure 3
Multiple P₁₇₀ Antigen




Figure 4
Synthetic Peptides Used For This Study




Fig. 4 : Synthetic peptides used for this study (single-letter amino-acid code)

a. PrP 57-64. WGQPHGGG b. PrP 89-106. WGQGGGTHSQWNKPSKPK: c. PrP 106-140. KTNMKHMAG: d. PrP 106-126. KTNMKHMAGAAAAGAVVGGLG: e. PrP 127-135. GYMLGSAMS: f. PrP 127-147. GYMLGSAMSRPIIHFSGSDYED: g. PrP 106-126 scrambled. NGAKALMGGHGATKVMVGAAA: a-f. amino-acid sequence of peptides homologous to different fragments of the amyloid protein purified from GSS brains (residues 58 to ~150)¹². g. Scrambled version of PrP 106-126. the octapeptide "a" is repeated for 4 or 5 times in the PrP sequence.

A β 4-11

Ac-Lys-Phe-Arg(Pbf)-His(Trt)-Asp(OtBu)-Ser(tBu)-Gly-Tyr(tBu)-Glu(OtBu)-Lys-Gly-OH

A β 1-16

Ac-Lys-Asp(OtBu)-Ala-Glu(OtBu)-Phe-Arg(Pbf)-His(Trt)-Asp(OtBu)-Ser(tBu)-Gly-Tyr(tBu)-Glu(OtBu)-Val-His(Trt)-His(Trt)-Gln(Trt)-Lys(Boc)-Lys-Gly-OH

A β 22-35

Ac-Lys-Glu(OtBu)-Asp(OtBu)-Val-Gly-Ser(tBu)-Asn(Trt)-Lys(Boc)-Gly-Ala-Ile-Ile-Gly-Leu-Met-Lys-Gly-OH

A β 29-40

Ac-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val-Lys-Gly-OH

FIGURE 5




FIGURE 6

FIGURE 7




Figure 7: ELISA conducted with 1:5000 diluted sera from pegylated amyloid/ liposomes/ lipid A immunized C57BL/6 mice. PEG-A β_{1-16} (- -black), PEG-A β_{1-16} + ALUM (- -grey), PEG-A β_{4-11} (—grey). Means of the values of 10 mice per antigen; means of values from 2 mice are shown for A β_{1-16} +ALUM. As a control mean values of 12 palmitoylated A β_{1-16} (- - bright grey) injected animals are shown (published 2002).




Figure 8




Figure 9




Figure 10





Figure 11

	Amyloid β -peptide fibers	Amyloid β -peptide fibers after 12 days of incubation with the antibody
$^{12}\text{Val C}_\alpha$	58 ppm	60 ppm
$^{12}\text{Val C}_\beta$	33 ppm	30 ppm
$^{10}\text{Tyr C}_\alpha$	54 ppm	56 ppm
$^{10}\text{Tyr C}_\beta$	38 ppm	41 ppm

FIGURE 12






Figure 13: pink: Palm.-A β_{1-16} , blue: PEG-A β_{1-16}




Figure 14: pink: Palm.-A β_{1-16} , blue: PEG-A β_{1-16} ,
 turquoise: PEG-A β_{1-15} , violet: PEG-A β_{22-35} ,
 yellow: PEG-A β_{4-11} .

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 9941279 A [0017]

Non-patent literature cited in the description

- **JANSSEN et al.** *International Journal of Pharmaceutics*, 2003, vol. 254, 55-58 [0016]
- **NICOLAU et al.** *PNAS*, 2002, vol. 99 (4), 2332-2337 [0018]
- **ALLXEN et al.** *Biochimica et Biophysica acta*, 1991, vol. 1066, 29-36 [0019]
- **ALVING et al.** *Infect. Immun.*, 1992, vol. 60, 2438-2444 [0083]
- *PNAS*, 2002, vol. 99, 2332-2337 [0158]
- **NICOLAU et al.** *PNAS*, 2002, vol. 99, 2332-37 [0163]
- **C. NICOLAU ; R.GREFERATH ; T.S. BALABAN ; J. LAZARTE ; R. HOPKINS.** *Proc. Natl. Acad. Sci. USA.*, 2002, vol. 99, 2332-2337 [0175]
- **FLUKA AG.** *Cat. # 79898*, 2002 [0175]
- **P.-F. TOSI ; D. RADU ; C.NICOLAU.** *Biochem. Biophys. Res. Chem.*, 1995, vol. 212, 494-500 [0175]
- **FUKUDA H ; SHIMIZU T ; NAKAJIMA M ; MORI H ; SHIRASAWA T.** *Bioorg. Med. Chem.Lett.*, 1999, vol. 9, 953-956 [0175]
- **PETKOVA AT ; ISHII Y ; BALBACH JJ ; ANTZUTKIN ON ; LEAPMAN RD ; DELAGLIO F ; TYCKO R.** *Proc. Natl. Acad. Sci. U.S.A.*, 2002, vol. 99, 16742-16747 [0175]