



(11) **EP 1 781 260 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

- (15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Description Paragraph(s) 14, 19, 28, 34, 36, 38, 39, 43, 44, 49, 54-57, 61, 62, 64, 67, 76, 82, 83
Claims EN 3
- (51) Int Cl.:
A61K 9/20 (2006.01) A61K 9/00 (2006.01)
A61K 31/428 (2006.01) A61K 31/4745 (2006.01)
- (86) International application number:
PCT/EP2005/053602
- (87) International publication number:
WO 2006/015942 (16.02.2006 Gazette 2006/07)
- (48) Corrigendum issued on:
16.02.2011 Bulletin 2011/07
- (45) Date of publication and mention of the grant of the patent:
03.11.2010 Bulletin 2010/44
- (21) Application number: **05777774.0**
- (22) Date of filing: **25.07.2005**

(54) **EXTENDED RELEASE TABLET FORMULATION CONTAINING PRAMIPEXOLE OR A PHARMACEUTICALLY ACCEPTABLE SALT THEREOF, METHOD FOR MANUFACTURING THE SAME AND USE THEREOF**

TABLETTENFORMULIERUNG MIT VERLÄNGERTER FREISETZUNG MIT PRAMIPEXOL ODER EINEM PHARMAZEUTISCH ZULÄSSIGEN SALZ DAVON, HERSTELLUNGSVERFAHREN UND VERWENDUNG DAFÜR

FORMULATION D'UN TABLET AVEC LIBERATION PROLONGUÉE COMPRENNANT PRAMIPEXOLE OU UN SEL PHARMACEUTIQUEMENT ACCEPTABLE, LA METHODE DE LA FABRICATION ET L'UTILISATION DU FORMULATION

- | | |
|--|---|
| <p>(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI SK TR
Designated Extension States:
BA HR YU</p> <p>(30) Priority: 13.08.2004 EP 04019248</p> <p>(43) Date of publication of application:
09.05.2007 Bulletin 2007/19</p> <p>(60) Divisional application:
09166415.1 / 2 135 602</p> | <p>(73) Proprietors:
• Boehringer Ingelheim International GmbH
55216 Ingelheim am Rhein (DE)
Designated Contracting States:
AT BE BG CH CY CZ DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI SK TR
• Boehringer Ingelheim Pharma GmbH & Co. KG
55216 Ingelheim (DE)
Designated Contracting States:
DE</p> <p>(72) Inventors:
• FRIEDL, Thomas
88416 OCHSENHAUSEN (DE)
• EISENREICH, Wolfram
89081 ULM (DE)</p> |
|--|---|

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 781 260 B9

(74) Representative: **Hammann, Heinz et al**
Boehringer Ingelheim GmbH
CD-Patents
Binger Straße 173
55216 Ingelheim am Rhein (DE)

(56) References cited:
WO-A-2004/010997 **WO-A-2004/010998**
WO-A-2004/010999


Description**FIELD OF THE INVENTION**

[0001] The present invention is directed to an extended release tablet formulation containing pramipexole or a pharmaceutically acceptable salt thereof, method for manufacturing the same and use thereof.

BACKGROUND OF THE INVENTION

[0002] Pramipexole is a known dopamine D2 receptor agonist. It is structurally different from the ergot-derived drugs, e.g. bromocriptine or pergolide. It is also pharmacologically unique in that it is a full agonist and has receptor selectivity for the dopamine D2 family of dopamine receptors.

[0003] Pramipexole is designated chemically as (S)-2-Amino-4,5,6,7-tetrahydro-6-(propylamino)benzothiazole and has the molecular formula $C_{10}H_{17}N_3S$ and a relative molecular mass of 211.33. The chemical formula is as follows:



[0004] The salt form commonly used is pramipexole dihydrochloride monohydrate (molecular formula $C_{10}H_{21}Cl_2N_3OS$; relative molecular mass 302.27). Pramipexole dihydrochloride monohydrate is a white to off-white, tasteless, crystalline powder. Melting occurs in the range of 296°C to 301°C, with decomposition. Pramipexole is a chiral compound with one chiral centre. Pure (S)-enantiomer is obtained from the synthetic process by chiral recrystallization of one of the intermediates during synthesis.

[0005] Pramipexole dihydrochloride monohydrate is a highly soluble compound. Water solubility is more than 20 mg/ml and solubility in buffer media is generally above 10 mg/ml between pH2 and pH 7.4. Pramipexole dihydrochloride monohydrate is not hygroscopic, and of highly crystalline nature. Under milling the crystal modification (monohydrate) does not change. Pramipexole is very stable in the solid state, yet in solution it is light sensitive.

[0006] Pramipexole immediate release (IR) tablets were first authorised in the USA in 1997, followed over the course of the next years by marketing authorisations in the European Union (EU), Switzerland, Canada and South America as well as in countries in Eastern Europe, Near East and Asia.

[0007] Pramipexole IR tablets are indicated in the EU and US for the treatment of signs and symptoms of either early parkinson's disease or advanced parkinson's disease in combination with levodopa. The IR tablets have to be taken 3 times a day.

[0008] From the pharmacokinetic point of view pramipexole IR tablets are rapidly and completely absorbed following oral administration. The absolute bioavailability is greater than 90% and the maximum plasma concentration occurs within 1 to 3 hours. The rate of absorption is reduced by food intake but not the overall extent of absorption. Pramipexole shows linear kinetics and a relatively small inter-patient variation of plasma levels. The elimination half-life ($t_{1/2}[h]$) varies from 8 hours in the young to 12 hours in the elderly.

[0009] As commonly known, modified release of active ingredient(s) allows to simplify the patient's administration scheme by reducing the amount of recommended daily intakes, improves patient's compliance, and attenuates adverse events, e.g. related to high plasma peaks. Modified release pharmaceutical preparations regulate the release of the incorporated active ingredient or ingredients over time and comprise formulations with a controlled, a prolonged, a sustained, a delayed, a slow or an extended release, so they accomplish therapeutic or convenience objectives not offered by conventional dosage forms such as solutions or promptly dissolving dosage forms.

[0010] A modified or extended release of active ingredient(s) from a pharmaceutical preparation may be accomplished by homogeneously embedding said active ingredient(s) in a hydrophilic matrix, being a soluble, partially soluble or insoluble network of viscous, hydrophilic polymers, held together by physical or chemical entanglements, by ionic or crystalline interactions, by complex formation, by hydrogen bonds or van der Waals forces. Said hydrophilic matrix swells upon contact with water, thereby creating a protective gellayer from which the active ingredient(s) is (are) slowly, gradually, continuously released in time either by diffusion through the polymeric network, by erosion of the gellayer, by dissolution of the polymer, or by a combination of said release mechanisms.

[0011] However, it has proved difficult to formulate a tablet having a suitable combination of modified, extended or

sustained-release and handling properties, where the drug is one having relatively high solubility, as in the case of pramipexole dihydrochloride.

[0012] There are a number of approaches described in prior art to provide sustained release tablet compositions of pramipexole:

[0013] WO 2004/010997 describes a sustained-release pharmaceutical composition in a form of an orally deliverable tablet comprising a water-soluble salt of pramipexole, dispersed in a matrix comprising a hydrophilic polymer and a starch having a tensile strength of at least about 0.15 kN cm⁻², preferably at least about 0.175 kN cm⁻², and more preferably at least about 0.2 kN cm⁻², at a solid fraction representative of the tablet. The disclosure thereof is concentrated to provide a composition with sufficient hardness yield during a high-speed tableting operation, in particular to resist erosion during application of a coating layer.

[0014] According to a preferred embodiment it is provided a pharmaceutical composition in a form of an orally deliverable tablet having a core comprising pramipexole dihydrochloride monohydrate in an amount of about 0.375, 0.75, 1.5, 3 or 4.5 mg, dispersed in a matrix comprising (a) HPMC type 2208 in an amount of about 35% to about 50% by weight of the tablet and (b) a pregelatinized starch having a tensile strength of at least about 0.15 kN cm⁻² at a solid fraction of 0.8, in an amount of about 45% to about 65% by weight of the tablet; said core being substantially enclosed in a coating that constitutes about 2% to about 7% of the weight of the tablet, said coating comprising an ethylcellulose-based hydrophobic or water-insoluble component and an HPMC-based pore-forming component in an amount of about 10% to about 40% by weight of the ethylcellulose-based component.

[0015] Furthermore, WO 2004/010999 discloses an orally deliverable pharmaceutical composition comprising a therapeutically effective amount of pramipexole or a pharmaceutically acceptable salt thereof and at least one pharmaceutically acceptable excipient, said composition exhibiting at least one of (a) an *in vitro* release profile wherein on average no more than about 20% of the pramipexole is dissolved within 2 hours after placement of the composition in a standard dissolution test; and (b) an *in vivo* pramipexole absorption profile following single dose oral administration to healthy adult humans wherein the time to reach a mean of 20% absorption is greater than about 2 hours and/or the time to reach a mean of 40% absorption is greater than about 4 hours. However, in practical use, it appears that any formulation having an extended or controlled release profile designed for a once daily application would meet the above requirements for which a general teaching how to adjust such a profile is missing.

[0016] WO 2004/010997 and WO 2004/010999 do not disclose the claimed subject matter.

[0017] It is an object of the present invention to provide a controlled release tablet composition of pramipexole or a pharmaceutically acceptable salt thereof that is suitable for once-daily oral administration. It is a further object to provide a tablet composition comprising pramipexole or a pharmaceutically acceptable salt thereof that provides a day-long therapeutic effect and will allow patients to treat their symptoms with a single daily dose, which makes it possible to adjust the release profile of the active ingredient according to a selected release profile dependent or independent from the pH values. Furthermore a method of manufacturing the tablet formulation shall be provided.

DESCRIPTION OF THE INVENTION

[0018] Surprisingly, it has been found that pramipexole or a pharmaceutically acceptable salt thereof may be used in formulations as once daily extended (or slow) release tablets and two alternative formulation principles allow different release rate types dependent or independent from the pH value.

[0019] The present invention relates to an extended release tablet formulation comprising pramipexole or a pharmaceutically acceptable salt thereof in a matrix comprising at least two water swelling polymers other than pregelatinized starch, and wherein at least one of the at least two polymers is an anionic polymer.

[0020] Preferred is an extended release tablet formulation, wherein the anionic polymer is selected from the group of optionally crosslinked acrylic acid polymers, methacrylic acid polymers, alginates, and carboxymethylcellulose.

[0021] Also preferred is an extended release tablet formulation, wherein the anionic polymer is an optionally crosslinked acrylic acid polymer, and wherein the content of the optionally crosslinked acrylic acid polymer in the matrix is from about 0.25 wt.-% to about 25 wt.-%, and preferably from about 0.5 wt.-% to about 15 wt.-%, and preferably from about 1 wt.-% to about 10 wt.-%.

[0022] Also preferred is an extended release tablet formulation, wherein at least one of the at least two polymers is a substantially neutral polymer other than pregelatinized starch.

[0023] Also preferred is an extended release tablet formulation, wherein the substantially neutral polymer is selected from hydroxypropylcellulose and hydroxypropylmethylcellulose.

[0024] Particularly preferred is an extended release tablet formulation, wherein the substantially neutral polymer is hydroxypropyl methylcellulose, and wherein the content of hydroxypropyl methylcellulose in the matrix is from about 10 wt.-% to about 75 wt.-% and preferably from about 25 wt.-% to about 65 wt.-%.

[0025] Particularly preferred is an extended release tablet formulation, wherein the matrix comprises about:

(a)	pramipexole or a salt thereof	0.05 to 5 wt.-%
(b)	anionic water swelling polymer(s)	0.25 to 25 wt.-%
(c)	neutral water swelling polymer(s)	10 to 75 wt.-%
(d)	further excipients	ad 100 wt.-%

[0026] Particularly preferred is an extended release tablet formulation consisting of Pramipexole-dihydrochloride monohydrate, Hypromellose 2208, Corn starch, Carbomer 941. Colloidal silicon dioxide and Magnesium stearate.

[0027] A preferred embodiment of the present invention relates to an extended release tablet formulation comprising pramipexole or a pharmaceutically acceptable salt thereof in a matrix comprising at least one water swelling polymer other than pregelatinized starch preferably a water swelling essentially neutral polymer, a water swelling anionic polymer and optionally excipients, the resulting tablet providing a pH-dependent release characteristic with a faster release characteristic in the range of pH < 4.5, and a slower and further on pH-independent release characteristic in the range from pH 4.5 to 7.5.

[0028] The extended release formulations according to the present invention intended for oral administration allow to select and estimate which *in vitro* release characteristic and timing of a formulation is most suitable to achieve the desired *in vivo* plasma profiles preferably with a once daily application. Therefore, a formulation principle with several variants has been developed for a single unit matrix tablet i.e. formulations having different release rate types are provided and a different pH dependency is available. These alternative formulations are beneficial to patients as the extended release drug delivery will allow patients to treat their symptoms with a single daily dose, thereby increasing patient convenience and compliance.

[0029] The term "*in vitro* release characteristic" as used hereinbefore or hereinafter is directed to a release characteristic as obtained in a kind of normally used liquid medium for *in vitro* experiments wherein the release of active ingredient from the extended release formulation can occur, i.e. for example in *in vitro* dissolution media, but also in body fluids or simulated body fluids, more in particular in the gastro-intestinal fluids.

[0030] In the frame of the present invention the term "extended" release should be understood in contrast to an immediate release, the active ingredient is gradually, continuously liberated over time, sometimes slower or faster, dependent or independent from the pH value. In particular, the term indicates that the formulation does not release the full dose of the active ingredient immediately after oral dosing and that the formulation allows a reduction in dosage frequency, following the definition for extended release, interchangeable with slow release. A slow or extended release, used synonymously with prolonged action, sustained release, or modified release, dosage form is a dosage form that allows a reduction in dosing frequency or a significant increase in patient compliance or therapeutic performance as compared to that presented as a conventional dosage form (e.g. as a solution or an immediate drug-releasing, conventional solid dosage form).

[0031] A release characteristic which is pH-independent indicates that the release characteristic is virtually the same in different pH media.

[0032] According to the teaching of the present invention, extended release tablet formulations are provided with different *in vitro* release profiles.

[0033] The extended release tablets of the present invention are believed to apply a swelling and partly eroding polymer matrix. Based on the assumed mechanisms, the release profile may roughly follow a square root of time to exponential *in vitro* release characteristic. Depending on the particular embodiment the formulation is faster in simulated gastric juice having a pH < 4.5 but is independent from the pH value in the range from 4.5 to 7.5. A faster release in simulated gastric juice versus slower release in the intestinal fluid can be advantageous in cases where a loading dose effect from the dosage form is desired, whereas a widely pH independent release profile can be advantageous to reduce the risk of dose dumping and food effects.

[0034] The water swelling polymer of the present invention represents at least one hydrophilic water swelling polymer constituting the extended release matrix which slowly releases the pramipexole or its salt as active ingredient. The polymer swells upon contact with aqueous fluid following administration, resulting in a viscous, drug release regulating gellayer. The viscosity of the polymer preferably ranges from 150 to 100,000 mPa.s (apparent viscosity of a 2% aqueous solution at 20°C.).

[0035] Examples of such polymers are water swelling substantially neutral polymers or water swelling anionic polymers.

[0036] The term "water swelling substantially neutral polymers" of the present invention comprises alkylcelluloses, such as, methylcellulose; hydroxyalkylcelluloses, for example, hydroxymethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose and hydroxybutylcellulose; hydroxyalkyl alkylcelluloses, such as, hydroxyethyl methylcellulose and hydroxypropyl methylcellulose; carboxyalkylcellulose esters; other natural, semi-synthetic, or synthetic di-, oligo- and polysaccharides such as galactomannans, tragacanth, agar, guar gum, and polyfructans; methacrylate copolymers; polyvinylalcohol; polyvinylpyrrolidone, copolymers of polyvinylpyrrolidone with vinyl acetate; combinations of polyvinyl-

alcohol and polyvinylpyrrolidone; polyalkylene oxides such as polyethylene oxide and polypropylene oxide and copolymers of ethylene oxide and propylene oxide, preferably cellulose ether derivatives such as hydroxypropyl methylcellulose and hydroxypropyl cellulose, most preferred hydroxypropyl methylcellulose.

[0037] The term "water swelling anionic polymer" of the present invention comprises acrylic acid polymerisate, methacrylic acid copolymers, alginates, carrageenans, acacia, xanthan gum, chitin derivatives such as chitosan, carmellose sodium, carmellose calcium, preferably acrylic acid polymerisate.

[0038] Different viscosity grades of hydroxypropyl cellulose and hydroxypropyl methylcellulose are commercially available. Hydroxypropyl methylcellulose (HPMC) preferably used in the present invention has a viscosity grade ranging from about 3,500 mPa.s to about 100,000 mPa.s, in particular ranging from about 4,000 mPa.s to about 20,000 mPa.s and most in particular a viscosity grade of about 6,500 mPa.s to about 15,000 mPa.s (apparent viscosity of a 2% aqueous solution at 20°C.), e.g. hypromellose 2208 or 2206 (DOW, Antwerp, Belgium). HPMC type 2208 contains 19-24% by weight methoxy and 4-12% by weight hydroxypropoxy substituents.

[0039] Hydroxypropyl cellulose having a viscosity higher than 1,500 mPa.s (apparent viscosity of a 1% aqueous solution at 20°C) is preferred, in particular hydroxypropyl cellulose having a viscosity in the range from about 1500 to about 3000 mPa.s, preferably from 4000 to 6500 mPa.s (2% aqueous solutions), e.g. the Klucel series such as Klucel M (Hercules, Wilmington, USA).

[0040] Without wishing to be bound by theory, there are believed to exist three main mechanisms by which pramipexole or a salt thereof can be released from a hydrophilic matrix: dissolution, erosion and diffusion. Pramipexole or its salt will be released by the dissolution mechanism when it is homogeneously dispersed in a matrix network of a soluble polymer. The network will gradually dissolve in the gastrointestinal tract, thereby gradually releasing its load. The matrix polymer can also gradually be eroded from the matrix surface, likewise releasing pramipexole or its salt in time. When pramipexole is processed in a matrix made up of an insoluble polymer, it will be released by diffusion: the gastro-intestinal fluids penetrate the insoluble, sponge-like matrix and diffuse back out loaded with drug.

[0041] Therefore, the water swelling polymers constituting the matrix mainly provide for the controlled pharmacokinetic release profile of the preparation. Depending on the amount of water swelling polymer(s) processed in the preparation, the release profile can be tuned, i.e. larger amounts of swelling polymer lead to a more pronounced sustained release effect and vice versa. Preferably, the amount of water swelling polymer in the present formulation ranges from about 10 to about 80% by weight.

[0042] In addition, when using a combination of polymers, the ratio of said polymers also influences the release profile of the preparation. A combination of different polymers offers the possibility of combining different mechanisms by which pramipexole is released from the matrix. Such combination facilitates control of the pharmacokinetic release profile of the preparation at will. For example, when using one or more water swelling polymers, in particular hydroxypropyl cellulose and hydroxypropyl methylcellulose, the weight percentage of hydroxypropyl methylcellulose preferably ranges from 25 to about 62%; the weight percentage of hydroxypropyl cellulose preferably ranges between 0% and about 16%.

[0043] Release of pramipexole or a salt thereof from a matrix containing hydroxypropyl cellulose and hydroxypropyl methylcellulose occurs by a combined set of release mechanisms. Due to the higher solubility of hydroxypropyl methylcellulose compared with hydroxypropyl cellulose, the former will gradually dissolve and erode from the matrix, whereas the latter will more act as a sponge-like matrix former releasing the active ingredient mainly by diffusion.

[0044] Beside pramipexole or a salt thereof, and the water swelling polymer(s), the formulation of the present invention may also optionally comprise further excipients, i.e. pharmaceutically acceptable formulating agents, in order to promote the manufacture, compressibility, appearance and taste of the preparation. These formulating agents comprise, for example, diluents or fillers, glidants, binding agents, granulating agents, anti-caking agents, lubricants, flavors, dyes and preservatives. Other conventional excipients known in the art can also be included.

[0045] The filler may be selected from soluble fillers, for example, sucrose, lactose, in particular lactose monohydrate, trehalose, maltose, mannitol and sorbitol. Different grades of lactose can be used. One type of lactose preferably used in the present invention is lactose monohydrate 200 mesh (DMV, Veghel, The Netherlands). Another lactose monohydrate, lactose monohydrate of the type DCL 11 (DMV, Veghel, The Netherlands), can also preferably be used. The notation DCL refers to "Direct Compression Lactose". The number 11 is a reference number of the manufacturer. In case of a water soluble active ingredient, like the one described in this invention, more preferably water insoluble fillers, such as starch and starch derivatives other than pregelatinized starch, e.g. corn starch, potato starch, rice starch or wheat starch, microcrystalline cellulose, dibasic calcium phosphate dihydrate and anhydrous dibasic calcium phosphate, preferably corn starch, can be used in addition or instead of the water soluble fillers. The total weight percentage of filler ranges between about 5% and about 75% by weight.

[0046] A glidant can be used to improve powder flow properties prior to and during tableting and to reduce caking. Suitable glidants include colloidal silicon dioxide, magnesium trisilicate, powdered cellulose, talc, tribasic calcium phosphate and the like. Colloidal silicon dioxide is preferably included as a glidant in an amount up to about 2%, preferably about 0.2% to about 0.8%, by weight of the tablet.

[0047] A lubricant can be used to enhance release of a tablet from apparatus on which it is formed, for example by

preventing adherence to the face of an upper punch ("picking") or lower punch ("sticking"). Suitable lubricants include magnesium stearate, calcium stearate, canola oil, glyceryl palmitostearate, hydrogenated vegetable oil, magnesium oxide, mineral oil, poloxamer, polyethylene glycol, polyvinyl alcohol, sodium benzoate, sodium lauryl sulfate, sodium stearyl fumarate, stearic acid, talc, hydrogenated vegetable oil, zinc stearate and the like. In one embodiment, magnesium stearate is included as a lubricant in an amount of about 0.1% to about 1.5%, preferably about 0.3% to about 1%, by weight of the tablet.

[0048] Among the optional formulating agents that further may be comprised in the matrix formulation there may be mentioned agents such as polyvidone; copovidone; starch; acacia; gelatin; seaweed derivatives, e.g. alginic acid, sodium and calcium alginate; cellulose, preferably microcrystalline cellulose, cellulose derivatives, e.g. ethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, having useful dry or wet binding and granulating properties; and anti-adherents such as talc and magnesium stearate.

[0049] In some embodiments of the present invention it is provided a pH-dependent release profile, the release of pramipexole or its salt from the tablet and subsequent the absorption into the blood stream can vary during the passage of the dosage form along the gastro-intestinal tract. Thus, the formulation provides a pH-dependent release characteristic wherein the release characteristic in the range of pH < 4.5 is faster and a slower and further on pH-independent release characteristic in the range from 4.5 = pH = 7.5.

[0050] The above details for the water swelling polymer and selection and type of optional excipients, may apply.

[0051] Moreover, an anionic water swelling polymer, preferably an acrylic acid polymerisate is mandatorily present, which is preferably selected from carbomer or carbopol® series, known acrylic acid polymerisates having high molecular weights. Particularly preferred are for example carbomer 941 (carbopol® 71 G, carbopol® 971) and carbomer 934 (carbopol® 974). The acrylic acid polymerisate is preferably present in the range of 0.25 to 25% by weight, particularly preferred 0.5 to 15% by weight, most preferred 1 to 10% by weight. pH dependency results from the presence of an anionic water swelling polymer, particularly preferred from the presence of acrylic acid polymerisate which intends to swell in a greater extent in the acid pH range above pH 4.5 and in the alkaline pH range.

[0052] An increasing amount of acrylic acid leads to a decrease of the release rate. Therefore, adjusting the amount of acrylic acid polymerisate makes it possible to further tune the dissolution profiles as desired. To adjust the amount of acrylic acid polymerisate in the preferred range from 0.25 to 25 % by weight provides the further advantage that the desired, resp. matching dissolution profiles can be adjusted, resp. maintained, for a variety of formulations composed of different amounts and/or types of gel-forming agents, water swelling polymers, fillers, and dry binders.

[0053] According to a preferred embodiment of the present invention the matrix of the extended release tablet formulation comprises or essentially consists of hydroxypropyl methylcellulose, acrylic acid polymerisate and further excipients. The amount of hydroxypropyl methylcellulose is preferably in the range from 10 to 75%, particularly preferred from 25 to 65%, most preferred from 35 to 55% by weight. The amount of acrylic acid polymerisate is preferably as above-mentioned. The amount of additional excipients is preferably in the range from 90 to 25% particularly preferred from 75 to 35%, most preferred from 65 to 45 % by weight. Optionally carboxymethylcellulose sodium may additionally be present preferably in the range from 5 to 50%, particularly preferred from 10 to 40%, most preferred from 15 to 30% by weight.

[0054] As active ingredient, pramipexole or a pharmaceutically acceptable salt thereof, may be present in any amount suitable for the desired treatment of a patient. A preferred salt of pramipexole is the dihydrochloride salt, most preferably in the form of the monohydrate. Usual amounts are from about 0.1 to about 5 mg pramipexole salt. According to a particularly preferred embodiment e.g. 0.750 mg pramipexole dihydrochloride monohydrate, corresponding to 0.524 mg anhydrous base, is used in the extended release tablet formulation according to the present invention. However, any other amount of active ingredient suitable for treatment may be used with the only proviso that the amount of pramipexole or salt is sufficient to provide a daily dose in one to a small plurality, for example one to about 4, of tablets to be administered at one time. Preferably the full daily dose is delivered in a single tablet. An amount of pramipexole salt, expressed as pramipexole dihydrochloride monohydrate equivalent, of about 0.1 to about 10 mg per tablet, or about 0.05% to about 5% by weight of the composition, will generally be suitable. Preferably an amount of about 0.2 to about 6 mg, more preferably an amount of about 0.3 to about 5 mg, per tablet is present. Specific dosage amounts per tablet c.g. include 0.375, 0.5, 0.75, 1.0, 1.5, 3.0 and 4.5 mg pramipexole dihydrochloride monohydrate. The amount that constitutes a therapeutically effective amount varies according to the condition being treated, the severity of said condition, and the patient being treated.

[0055] An extended release tablet formulation according to the present invention, has preferably the following composition:

pramipexole or a salt thereof	0.05 to 5% by weight
water swelling polymer(s)	10 to 75% by weight
acrylic acid polymerisate	> 0 to 25% by weight
optional further excipient(s)	ad 100% by weight.

Therefore, a particularly preferred extended release tablet formulation of the present invention consists of

0.1 to 2% by weight of pramipexole or a salt thereof;
 25 to 65% by weight of hydroxypropyl methylcellulose;
 > 0 to 40% by weight of carboxymethylcellulose sodium;
 0 to 75% by weight of corn starch other than pregelatinized starch;
 > 0 to 15% by weight of acrylic polymerisate, preferably carbomer 941;
 0.5 to 50% by weight of excipients, preferably selected from the group consisting of colloidal silicon dioxide, magnesium stearate, lactose monohydrate, mannitol, microcrystalline cellulose, dibasic anhydrous calcium phosphate, hydroxypropylcellulose, povidone, copovidone, talc, macrogols, sodium dodecylsulfate, iron oxides and titanium dioxide.

[0056] According to the present invention starch other than pregelatinized starch, preferably corn starch if present, may impart several functions at the same time such as filler, glidant, and the like. However, it may be preferred to exclude starch completely from the tablet formulation according to the present invention, which may be replaced by one or more of the above-mentioned other excipient(s). Furthermore, a starch having a tensile strength of at least about 0.15 kN cm⁻² at a solid fraction representative of the tablet as claimed according to WO 2004/010997 is not required according to the present invention.

[0057] It is preferred that no coating is present on the tablet formulation according to the present invention. However, the extended release tablet of the invention may comprise a nonfunctional coating. A nonfunctional coating can comprise a polymer component, for example HPMC, optionally with other ingredients, for example one or more plasticizers, colorants, etc. The term "nonfunctional" in the present context means having no substantial effect on release properties of the tablet, and the coating serves another useful purpose. For example, such a coating can impart a distinctive appearance to the tablet, provide protection against attrition during packaging and transportation, improve ease of swallowing, and/or have other benefits. A nonfunctional coating should be applied in an amount sufficient to provide complete coverage of the tablet. Typically an amount of about 1 % to about 10%, more typically an amount of about 2% to about 5%, by weight of the tablet as a whole, is suitable.

[0058] The tablets of the present invention can be of any suitable size and shape, for example round, oval, polygonal or pillow-shaped, and optionally bear nonfunctional surface markings. According to the present invention it is preferred that the extended release tablets are white to off-white and of oval or round, biconvex, shape.

[0059] Tablets of the invention can be packaged in a container, accompanied by a package insert providing pertinent information such as, for example, dosage and administration information, contraindications, precautions, drug interactions and adverse reactions.

[0060] The present invention is further directed to the use of the extended release tablet formulation according to the present invention for preparing a medical composition for the treatment of parkinson's disease and complications or disorders associated therewith.

[0061] Furthermore, the present invention is preferably directed to a method of manufacturing the extended release tablet formulations via a direct compression process comprising the steps of

(1) producing an active ingredient trituration wherein the active ingredient is pramipexole or a pharmaceutically acceptable salt thereof by preblending it with a portion of water swelling polymer(s) and/or further excipient(s) in a mixer, wherein pramipexole or the pharmaceutically acceptable salt thereof is milled, preferably peg-milled, prior to use;

(2) premixing the active ingredient trituration of step (1), the main portion of the water swelling polymer(s) and/or excipients in a mixer to obtain a pre-mixture;

(3) optionally dry screening the pre-mixture through a screen in order to segregate cohesive particles and to improve content uniformity;

(4) mixing the pre-mixture of step (2) or (3) in a mixer, optionally by adding remaining excipients to the mixture and continuing mixing; and

(5) tableting the final mixture by compressing it on a suitable tablet press to produce matrix tablets.

[0062] Therefore, the tablets are manufactured via a direct compression process. To achieve adequate content uniformity in this low drug load formulation, the active ingredient is preferably Peg-milled. Preferably the particle size distribution of the peg-milled drug substance, as determined by laser diffractometry using a dry dispensing system, is

characterized by particle fraction of 90 % (V/V) being smaller than 100 μm , most preferably a particle fraction of 90 % (V/V) being smaller than 75 μm in diameter.

[0063] Also other processes can be applied to the manufacturing of Pramipexole extended release tablets, like conventional wet granulation and roller compaction. In case of wet granulation preferably Pramipexole is granulated with suitable fillers, like e.g. starches other than pregelatinized starch, microcrystalline cellulose, lactose monohydrate or anhydrous dibasic calcium phosphate, and wet binding agents, like e.g. hydroxypropylmethylcellulose, hydroxypropylcellulose, povidone, copovidone, and starch paste, leading to a active ingredient concentrate, which after drying and dry screening is mixed with the main fraction of gel forming excipients, like all the above described retarding principles.

[0064] In case of roller compaction, or in other words dry granulation, either a premix of Pramipexole with part of the excipients used in the direct compression process, or the complete mixture containing all excipients, is processed through a conventional roller compactor to form ribbons, which are thereafter screened down to granules which are finally mixed with other excipients, like glidants, lubricants and antiadherents.

BRIEF DESCRIPTION OF THE DRAWINGS

[0065]

Figure 1 is a flow diagram illustrating a preferred embodiment of the direct compression manufacturing process according to the present invention;

Figure 2 is a graph illustrating the dissolution profiles of a matrix tablet formulation according to the present invention which contains 4% by weight carbopol® in 3 different pH media; and

Figure 3 is a graph illustrating the dissolution profiles of 3 matrix tablet formulations which contain 0%, 1% and 4% by weight of carbopol®, respectively.

[0066] Figure 1 illustrates a preferred embodiment of the manufacturing process with reference to a flow diagram wherein the manufacture of the extended release tablets of Examples 1 and 2 are exemplarily shown. Figure 1 shows the detailed process steps and the in process controls performed.

[0067] Process step (1) is directed to the active ingredient trituration, i.e. in the present case a salt of pramipexole, pramipexole dihydrochloride monohydrate, in peg-milled quality, is preblended with a portion of the polymer, in this case hydroxypropyl methylcellulose, in a commonly known mixer. In the flow chart a Turbula free-fall mixer or blender is used. The mixing time is several minutes, in the present case preferably 10 min.

[0068] In process step (2) according to the flow chart a premixing is performed, wherein the active ingredient trituration and the main portion of the water swelling polymer(s) and excipients are premixed for several minutes to obtain a pre-mix. In the present case the main portion of hydroxypropyl methylcellulose (hypromellose), corn starch carbomer 941 and colloidal silicon dioxide are premixed for 5 min. in the above-mentioned Turbula mixer or blender.

[0069] According to the following process step (3) a dry screening may optionally take place. The pre-mixture may be manually screened through a screen, for example a 0.8 mm mesh size screen, in order to segregate cohesive particles and to improve content uniformity.

[0070] In the subsequent process step (4) the main mixing step is performed according to which the components are mixed for several minutes, preferably 5 min. in the Turbula mixer after screening. Optionally further excipients may be added at this time, in the flow chart the component magnesium stearate is added to the main mixture, and further mixing for several minutes, e.g. 3 min., in the Turbula mixer is performed (final mixing) to obtain the final mixture.

[0071] Process step (5) of the process according to the present invention is the tableting. The final mixture is compressed on a suitable tablet press to produce, for example, oblong shaped matrix tablets (ER tablets = extended release tablets). In order to control and maintain the required quality the obtained matrix tablets are subjected to the following in-process controls: tablet mass, hardness, tablet height and friability.

[0072] The obtained pramipexole extended release tablets of the present invention may then be filled, for example, into High Density Polyethylene (HDPE) bottles. The bottles are closed tightly with screw caps and appropriately labelled, whereby all packaging and labelling activities are performed according to cGMP regulations. Alternatively, a blister type packaging can be used, e.g. using aluminium/aluminium foil blisters.

[0073] Figure 2 represents a graph illustrating the dissolution profiles of a matrix tablet formulation according to the present invention. The matrix tablet contains 4% by weight carbopol®, the detailed composition is given in Example 2. The release characteristics of the matrix tablet in 3 different pH media are shown, i.e. in 0.05 M phosphate buffer, pH = 6.8, $n = x$, in simulated gastric juice, pH = 1.2, $n = x$, and in McIlvaine buffer, pH = 4.5, $n = x$; (x ...represents the number of units tested). The value percent of released active ingredient is plotted against the time (hours).

[0074] Figure 3 represents a graph illustrating the dissolution profiles of 3 matrix tablet formulations. The matrix tablets

contain no carbopol®, 1% or 4% by weight carbopol®, respectively, the detailed compositions are given in Examples 1, 2 and 4. The medium is a 0.05 M phosphate buffer, pH = 6.8. The value percent of released active ingredient is plotted against the time (hours).

[0075] Figures 2 and 3 show a pH-independent *in vitro* release characteristic in the range from pH 1 to 7.5 in case carbopol® is not present and a pH-dependent release characteristic wherein the release characteristic in the range of pH < 4.5 is faster in case carbopol® is present. An increase of the amount of carbopol® leads to a decreased releasing rate.

[0076] The advantages of the present invention are manifold:

[0077] According to the present invention, extended release tablets containing pramipexole or its salt are available showing different *in vitro* release profiles. It is possible to select a tailor-made release characteristic for patient's needs, symptoms and clinical picture observed.

[0078] The primary indication for pramipexole, Parkinson's disease, is an affliction that becomes more prevalent with advancing age and is often accompanied by decline in memory. Therefore, the matrix tablets according to the present invention providing an extended or slow release of pramipexole or a salt thereof allows to simplify the patient's administration scheme by reducing the amount of recommended daily intakes and improves patient's compliance, particularly relevant for elderly patients. The inventive extended release tablet formulation provides a daily dose preferably administered at one time.

[0079] Furthermore, the tablets of the present invention may be manufactured via a direct compression, wet or dry granulation process which applies to both types of extended release matrix tablets.

[0080] The invention described will now be illustrated by the Examples which follow various other embodiments and will become apparent to the skilled person from the present specification.

Examples

[0081] According to the present invention pramipexole extended release tablets have been manufactured. The tablets of the Examples are white to off-white, 14 x 6.8 mm oblong shaped, biconvex tablets. The tablets are intended to be administered orally, and shall not be divided into halves. The pramipexole tablets in the Examples contain 0.75 mg of pramipexole dihydrochloride monohydrate, corresponding to 0.524 mg of pramipexole free, anhydrous base.

Example 1

[0082] One embodiment of the qualitative and quantitative composition of pramipexole extended release tablets according to the present invention is shown in TABLE 1.

TABLE 1: Qualitative and quantitative composition of pramipexole extended release tablet

Ingredient	mg per 0.75 mg tablet	Function	Reference to Standards
Pramipexole-dihydrochloride monohydrate, peg-milled	0.750	Active ingredient	Corporate standard
Hypromellose 2208 (Methocel K 15 M)	157.500	Swelling agent	Ph.Eur. / USP
Corn starch	183.700	Filler	Ph.Eur. / NF
Carbomer 941 (Carbopol® 71 G)	3.500	Gelling agent	Ph.Eur. / NF
Colloidal Silicon dioxide	2.800	Glidant	Ph.Eur. / NF
Magnesium stearate	1.750	Lubricant	Ph.Eur. / NF
Total	350.000		

Example 2

[0083] A further embodiment of the qualitative and quantitative composition of pramipexole extended release tablets according to the present invention is shown in TABLE 2.

TABLE 2: Qualitative and quantitative composition of pramipexole extended release tablet

Ingredient	mg per 0.75 mg tablet	Function	Reference to Standards
Pramipexole-dihydrochloride monohydrate, peg-milled	0.750	Active ingredient	Corporate standard
Hypromellose 2208 (Methocel K 15 M)	157.500	Swelling agent	Ph.Eur. / USP
Corn starch	174.600	Filler	Ph.Eur. / NF
Carbomer 941 (Carbopol® 71 G)	14.000	Gelling agent	Ph.Eur. / NF
Colloidal Silicon dioxide	1.400	Glidant	Ph.Eur. / NF
Magnesium stearate	1.750	Lubricant	Ph.Eur. / NF
Total	350.000		

EXAMPLE 3

[0084] The batch formula for the two pramipexole tablet formulations of Example 1 and 2 is shown in Table 3. The batch size of the final mixture corresponds to a batch size of 2000 tablets.

Table 3: Composition per batch of pramipexole 0.75 mg ER tablets

Ingredient	Grams per batch Example 1	Grams per batch Example 2
Pramipexole-dihydrochloride monohydrate, peg-milled	1.500	1.500
Hypromellose 2208	315.000	315.000
Corn starch	367.400	349.200
Carbomer 941	7.000	28.000
Colloidal Silicon dioxide	5.600	2.800
Magnesium stearate	3.500	3.500
Total Mass	700.000	700.000

Example 4 - Reference example see Fig.3

[0085] The following Example shows a pramipexole tablet formulation which provides a release characteristic independent in the pH range of 1 to 7.5.

Table 4

Constituents	mg/tablet
Pramipexole-dihydrochloride monohydrate, peg-milled	0.750
Hypromellose 2208 (Methocel K 100 M)	157.500
Corn starch	187.900
Colloidal silicon dioxide	2.100
Magnesium stearate	1.750
Total weight matrix tablet	350.000

Example 5

[0086] The following Examples 6 to 14 show pramipexole tablet formulations which correspond to a formulation providing a faster release characteristic for pH < 4.5.

EP 1 781 260 B9

Table 5

Constituents	mg/tablet
Pramipexole-dihydrochloride monohydrate, peg-milled	0.750
Hypromellose 2208 (Methocel K 15 M)	175.000
Carboxymethylcellulose sodium	87.500
Lactose monohydrate (200 mesh)	52.500
Microcrystalline cellulose (grade PH 101)	31.100
Colloidal silicon dioxide	1.400
Magnesium stearate	1.750
Total weight matrix tablet	350.000

Example 6

[0087]

Table 6

Constituents	mg/tablet
Pramipexole-dihydrochloride monohydrate, peg-milled	0.750
Hypromellose 2208 (Methocel K 15 M)	175.000
Carboxymethylcellulose sodium	87.500
Lactose monohydrate (200 mesh)	52.500
Microcrystalline cellulose (grade PH 101)	27.600
Carbomer 941 (Carbopol® 71 G)	3.500
Colloidal silicon dioxide	1.400
Magnesium stearate	1.750
Total weight matrix tablet	350.000

Example 7

[0088]

Table 7

Constituents	mg/tablet
Pramipexole-dihydrochloride monohydrate, peg-milled	0.750
Hypromellose 2208 (Methocel K 15 M)	175.000
Carboxymethylcellulose sodium	87.500
Lactose monohydrate (200 mesh)	45.500
Microcrystalline cellulose (grade PH 101)	24.100
Carbomer 941 (Carbopol® 71 G)	14.000
Colloidal silicon dioxide	1.400

EP 1 781 260 B9

(continued)

Constituents	mg/tablet
Magnesium stearate	1.750
Total weight matrix tablet	350.00

[0089] Example 8 - 11 omitted.

Example 12

[0090]

Table 12

Constituents	mg/tablet
Pramipexole-dihydrochloride monohydrate, peg-milled	0.750
Hypromellose 2208 (Methocel K 15 M)	157.500
Corn starch	163.400
Carbomer 941 (Carbopol® 71 G)	24.500
Colloidal silicon dioxide	2.100
Magnesium stearate	1.750
Total weight matrix tablet	350.000

Example 13

[0091]

Table 13

Constituents	mg/tablet
Pramipexole-dihydrochloride monohydrate, peg-milled	0.750
Hypromellose 2910 (Methocel E 5)	0.788
Corn starch	173.812
Hypromellose 2208 (Methocel K 15 M)	157.500
Carbomer 941 (Carbopol® 71 G)	14.000
Colloidal silicon dioxide	1.400
Magnesium stearate	1.750
Total weight matrix tablet	350.000

Example 14

[0092]

Table 14

Constituents	mg/tablet
Pramipexole-dihydrochloride monohydrate, peg milled	0.750

(continued)

Constituents	mg/tablet
Hypromellose 2208 (Methocel K 15 M)	148.500
Corn starch	160.620
Carbomer 941 (Carbopol® 71 G)	16.500
Colloidal silicon dioxide	1.980
Magnesium stearate	1.650
Total weight matrix tablet	330.000

Claims

1. An extended release tablet formulation comprising pramipexole or a pharmaceutically acceptable salt thereof in a matrix comprising at least two water swelling polymers other than pregelatinized starch, and wherein at least one of the at least two polymers is an anionic polymer.
2. The extended release tablet formulation of claim 1, wherein the anionic polymer is selected from the group of optionally crosslinked acrylic acid polymers, methacrylic acid polymers, alginates and carboxymethylcellulose.
3. The extended release tablet formulation of claim 2, wherein the anionic polymer is an optionally crosslinked acrylic acid polymer, and wherein the content of the optionally crosslinked acrylic acid polymer in the matrix is from about 0.25 wt.-% to about 25 wt.-%, and preferably from about 0.5 wt.-% to about 15 wt.-%, and preferably from about 1 wt.-% to about 10 wt.-%.
4. The extended release tablet formulation of claim 1, wherein at least one of the at least two polymers is a substantially neutral polymer other than pregelatinized starch.
5. The extended release tablet formulation of claim 4, wherein the substantially neutral polymer is selected from hydroxypropylcellulose and hydroxypropylmethylcellulose.
6. The extended release tablet formulation of claim 4, wherein the substantially neutral polymer is hydroxypropyl methylcellulose, and wherein the content of hydroxypropyl methylcellulose in the matrix is from about 10 wt.-% to about 75 wt.-%, and preferably from about 25 wt.-% to about 65 wt.-%.
7. The extended release tablet formulation of claim 4, wherein the matrix comprises about:

(a)	pramipexole or a salt thereof	0.05 to 5 wt.-%
(b)	anionic water swelling polymer(s)	0.25 to 25 wt.-%
(c)	neutral water swelling polymer(s)	10 to 75 wt.-%
(d)	further excipients	ad 100 wt.-%
8. The extended release tablet formulation of claim 1 consisting of Pramipexole-dihydrochloride monohydrate, Hypromellose 2208, Corn starch, Carbomer 941, Colloidal silicon dioxide and Magnesium stearate.
9. The extended release tablet formulation according to claim 1, the resulting tablet providing a pH-dependent release characteristic with a faster release characteristic in the range of pH < 4.5, and a slower and further on pH-independent release characteristic in the range from pH 4.5 to 7.5.
10. The extended release tablet formulation according to any of the previous claims, wherein the contained amount of pramipexole or pharmaceutically acceptable salt thereof is sufficient to provide a daily dose administered at one time.

11. Method of manufacturing the extended release tablet formulation according to one of the preceding claims 1 to 10, via a direct compression process comprising the steps of

- (1) producing an active ingredient trituration wherein the active ingredient is pramipexole or a pharmaceutically acceptable salt thereof by preblending it with a portion of water swelling polymer(s) and/or excipient(s) in a mixer, wherein pramipexole or the pharmaceutically acceptable salt thereof is milled, preferably peg-milled, prior to use;
- (2) premixing the active ingredient trituration of step (1), the main portion of the water swelling polymer(s) and/or excipients in a mixer to obtain a pre-mixture;
- (3) optionally dry screening the pre-mixture through a screen in order to segregate cohesive particles and to improve content uniformity;
- (4) mixing the pre-mixture of step (2) or (3) in a mixer, optionally by adding remaining excipients to the mixture and continuing mixing; and
- (5) tableting the final mixture by compressing it on a suitable tablet press to produce matrix tablets.

12. Method of manufacturing the extended release tablet formulation according to one of the preceding claims 1 to 10, via a wet granulation process comprising the steps of

- (1) producing an active ingredient trituration wherein the active ingredient is pramipexole or a pharmaceutically acceptable salt thereof by blending it with a portion of the excipients in a mixer, wherein pramipexole or the pharmaceutically acceptable salt thereof is milled, preferably peg-milled, prior to use;
- (2) granulating the active ingredient trituration of step (1) by adding the granulation liquid, preferably water;
- (3) drying the granules of step (2) in a fluidized bed dryer or a drying oven;
- (4) mixing the dried granules of step (3) with the water swelling polymer(s) and/or excipients in a mixer to obtain the final mixture;
- (5) tableting the final mixture of step (4) by compressing it on a suitable tablet press to produce matrix tablets.

13. Method of manufacturing the extended release tablet formulation according to one of the claims 1 to 10, via a dry granulation process comprising the steps of

- (1) mixing the active ingredient pramipexole or a pharmaceutically acceptable salt thereof with either a portion of the fillers or all the excipients in a mixer, wherein pramipexole or the pharmaceutically acceptable salt thereof is milled, preferably peg-milled, prior to use;
- (2) compaction of the mixture of step (1) on a suitable roller compactor;
- (3) reducing the ribbons obtained during step (1) to small granules by suitable milling or sieving steps;
- (4) optionally mixing the granules of step (3) with the remaining excipients in a mixer to obtain the final mixture;
- (5) tableting the granules of step (3) or the final mixture of step (4) by compressing it on a suitable tablet press to produce matrix tablets.

14. Use of an extended release tablet formulation according to one of the preceding claims 1 to 10 for preparing a medical composition for the treatment of Parkinson's disease and complications or disorders associated therewith.

Patentansprüche

1. Tablettenformulierung mit verzögerter Freisetzung, umfassend Pramipexol oder ein pharmazeutisch akzeptables Salz davon in einer Matrix, umfassend mindestens zwei wasserquellbare Polymere außer vorgelatinisierter Stärke, und wobei mindestens eines der mindestens zwei Polymere ein anionisches Polymer darstellt.
2. Tablettenformulierung mit verzögerter Freisetzung nach Anspruch 1, wobei das anionische Polymer ausgewählt ist aus der Gruppe der gegebenenfalls vernetzten Acrylsäurepolymere, Methacrylsäurepolymere, Alginate und Carboxymethylcellulose.
3. Tablettenformulierung mit verzögerter Freisetzung nach Anspruch 2, wobei das anionische Polymer ein gegebenenfalls vernetztes Acrylsäurepolymer darstellt und wobei der Gehalt des gegebenenfalls vernetzten Acrylsäurepolymeren in der Matrix von etwa 0,25 Gew.-% bis etwa 25 Gew.-% und bevorzugt von etwa 0,5 Gew.-% bis etwa 15 Gew.-% und bevorzugt von etwa 1 Gew.-% bis etwa 10 Gew.-% reicht.

4. Tablettenformulierung mit verzögerter Freisetzung nach Anspruch 1, wobei mindestens eines der mindestens zwei Polymeren ein im wesentlichen neutrales Polymer außer vorgelatinisierter Stärke darstellt.

5. Tablettenformulierung mit verzögerter Freisetzung nach Anspruch 4, wobei das im wesentlichen neutrale Polymer ausgewählt ist aus Hydroxypropylcellulose und Hydroxypropylmethylcellulose.

6. Tablettenformulierung mit verzögerter Freisetzung nach Anspruch 4, wobei das im wesentlichen neutrale Polymer Hydroxypropylmethylcellulose darstellt und wobei der Gehalt der Hydroxypropylmethylcellulose in der Matrix etwa 10 Gew.-% bis etwa 75 Gew.-% und bevorzugt etwa 25 Gew.-% bis etwa 65 Gew.-% beträgt.

7. Tablettenformulierung mit verzögerter Freisetzung nach Anspruch 4, wobei die Matrix etwa umfasst:

- | | | |
|-----|---|--------------------|
| (a) | Pramipexol oder ein Salz davon | 0,05 bis 5 Gew.-% |
| (b) | anionische(s) wasserquellbare(s) Polymer(e) | 0,25 bis 25 Gew.-% |
| (c) | neutrale(s) wasserquellbare(s) Polymer(e) | 10 bis 75 Gew.-% |
| (d) | weitere Hilfsstoffe | auf 100 Gew.-%. |

8. Tablettenformulierung mit verzögerter Freisetzung nach Anspruch 1, bestehend aus Pramipexoldihydrochlorid-Monohydrat, Hypromellose 2208, Maisstärke, Carbomer 941, kolloidalem Siliciumdioxid und Magnesiumstearat.

9. Tablettenformulierung mit verzögerter Freisetzung nach Anspruch 1, wobei die resultierende Tablette eine pH-abhängige Freisetzungseigenschaft mit einer schnelleren Freisetzungseigenschaft im pH-Bereich < 4,5 bereitstellt und einer langsameren und weiterhin pH-unabhängigen Freisetzungseigenschaft im pH-Bereich von 4,5 bis 7,5.

10. Tablettenformulierung mit verzögerter Freisetzung nach irgendeinem der vorhergehenden Ansprüche, wobei die enthaltene Menge an Pramipexol oder des pharmazeutisch akzeptablen Salzes davon ausreichend ist, um eine Tagesdosis bereitzustellen, die auf einmal verabreicht wird.

11. Verfahren zur Herstellung der Tablettenformulierung mit verzögerter Freisetzung nach irgendeinem der vorhergehenden Ansprüche 1 bis 10 über ein direktes Kompressionsverfahren, umfassend die Schritte:

- (1) Herstellen einer Verreibung der aktiven Bestandteile, wobei der aktive Bestandteil Pramipexol oder ein pharmazeutisch akzeptables Salz davon darstellt, durch Vormischen desselben mit einer Portion des (der) wasserquellbaren Polymers(en) und/oder Hilfsstoff(en) in einem Mischer, wobei Pramipexol oder das pharmazeutisch akzeptable Salz davon vor der Verwendung gemahlen wird, insbesondere mit einer Stößmühle;
- (2) Vormischen der Verreibung der aktiven Bestandteile von Schritt (1), der Hauptmenge des (der) wasserquellbaren Polymers(en) und/oder Hilfsstoffen in einem Mischer, um eine Vormischung zu erhalten;
- (3) gegebenenfalls trockenes Sieben der Vormischung durch ein Sieb, um kohäsive Partikel abzuscheiden und die Einheitlichkeit des Inhalts zu verbessern;
- (4) Mischen der Vormischung von Schritt (2) oder (3) in einem Mischer, gegebenenfalls durch Zugabe der verbliebenen Hilfsstoffe zu der Mischung und Fortsetzen des Mischens und
- (5) Tablettieren der Endmischung durch Komprimieren auf einer geeigneten Tablettenpresse, um Matrixtabletten herzustellen.

12. Verfahren zur Herstellung der Tablettenformulierung mit verzögerter Freisetzung nach irgendeinem der vorhergehenden Ansprüche 1 bis 10 über ein Nassgranulationsverfahren, umfassend die Schritte:

- (1) Herstellen einer Verreibung der aktiven Bestandteile, wobei der aktive Bestandteil Pramipexol oder ein pharmazeutisch akzeptables Salz davon darstellt, durch Mischen desselben mit einer Portion der Hilfsstoffe in einem Mischer, wobei Pramipexol oder das pharmazeutisch akzeptable Salz davon vor der Verwendung gemahlen wird, insbesondere mit einer Stößmühle;
- (2) Granulieren der Verreibung der aktiven Bestandteile von Schritt (1) durch Zugabe der Granulierungsflüssigkeit, bevorzugt Wasser;
- (3) Trocknen der Granulate von Schritt (2) in einem Fließbett-Trockner oder einem Trockenofen;
- (4) Mischen der getrockneten Granulate von Schritt (3) mit dem (den) wasserquellbaren Polymer(en) und/oder

Hilfsstoffen in einem Mischer, um die Endmischung zu erhalten;
(5) Tablettieren der Endmischung von Schritt (4) durch Komprimieren auf einer geeigneten Tablettendruckmaschine, um Matrixtabletten herzustellen.

- 5 **13.** Verfahren zur Herstellung der Tablettenformulierung mit verzögerter Freisetzung nach irgendeinem der Ansprüche 1 bis 10 über ein Trockengranulationsverfahren, umfassend die Schritte:

- 10 (1) Mischen des aktiven Bestandteils Pramipexol oder eines pharmazeutisch akzeptablen Salzes davon entweder mit einer Portion Füllstoffe oder sämtlichen Hilfsstoffen in einem Mischer, wobei Pramipexol oder das pharmazeutisch akzeptable Salz davon vor der Verwendung gemahlen wird, insbesondere mit einer Stößmühle;
(2) Kompaktieren der Mischung von Schritt (1) auf einem geeigneten Walzenkompaktor;
(3) Zerkleinern der während des Schritts (1) erhaltenen Schülpen zu kleinen Granulaten durch geeignete Mahl- oder Siebschritte;
(4) gegebenenfalls Mischen der Granulate von Schritt (3) mit den verbliebenen Hilfsstoffen in einem Mischer, um die Endmischung zu erhalten;
15 (5) Tablettieren der Granulate von Schritt (3) oder der Endmischung von Schritt (4) durch Komprimieren auf einer geeigneten Tablettendruckmaschine, um Matrixtabletten zu erzeugen.

- 20 **14.** Verwendung einer Tablettenformulierung mit verzögerter Freisetzung nach irgendeinem der vorhergehenden Ansprüche 1 bis 10 zur Herstellung einer medizinischen Zusammensetzung zur Behandlung der Parkinsonschen Erkrankung und damit verbundenen Komplikationen oder Störungen.

Revendications

- 25 **1.** Formulation pour comprimés à libération prolongée comprenant du pramipexole ou un sel pharmaceutiquement acceptable de celui-ci dans une matrice comprenant au moins deux polymères gonflant dans l'eau différents de l'amidon prégélatinisé et où au moins l'un des au moins deux polymères est un polymère anionique.
- 30 **2.** Formulation pour comprimés à libération prolongée selon la revendication 1 où le polymère anionique est choisi dans le groupe des polymères d'acide acrylique éventuellement réticulés, des polymères d'acide méthacrylique, des alginates et de la carboxyméthylcellulose.
- 35 **3.** Formulation pour comprimés à libération prolongée selon la revendication 2 où le polymère anionique est un polymère d'acide acrylique éventuellement réticulé et où la teneur du polymère d'acide acrylique éventuellement réticulé dans la matrice est d'environ 0,25 % en poids à environ 25 % en poids, et de préférence d'environ 0,5 % en poids à environ 15 % en poids, et de préférence d'environ 1 % en poids à environ 10 % en poids.
- 40 **4.** Formulation pour comprimés à libération prolongée selon la revendication 1 où au moins l'un des au moins deux polymères est un polymère sensiblement neutre différent de l'amidon prégélatinisé.
- 5.** Formulation pour comprimés à libération prolongée selon la revendication 4 où le polymère sensiblement neutre est choisi parmi l'hydroxypropylcellulose et l'hydroxypropylméthylcellulose.
- 45 **6.** Formulation pour comprimés à libération prolongée selon la revendication 4 où le polymère sensiblement neutre est l'hydroxypropylméthylcellulose et où la teneur de l'hydroxypropylméthylcellulose dans la matrice est d'environ 10 % en poids à environ 75 % en poids et de préférence d'environ 25 % en poids à environ 65 % en poids.
- 50 **7.** Formulation pour comprimés à libération prolongée selon la revendication 4 où la matrice comprend environ :
- | | | |
|-----|--|---|
| (a) | du pramipexole ou un sel de celui-ci | 0,05 à 5 % en poids |
| (b) | un ou des polymères anioniques gonflant dans l'eau | 0,25 à 25 % en poids |
| (c) | un ou des polymères neutres gonflant dans l'eau | 10 à 75 % en poids |
| 55 | (d) | d'autres excipients pour 100 % en poids |

- 8.** Formulation pour comprimés à libération prolongée selon la revendication 1 consistant en dichlorhydrate de prami-

pexole monohydraté,
hypromellose 2208, amidon de maïs, carbomère 941, dioxyde de silicium colloïdal et stéarate de magnésium.

9. Formulation pour comprimés à libération prolongée selon la revendication 1, le comprimé résultant fournissant une caractéristique de libération dépendante du pH avec une caractéristique de libération plus rapide dans la plage de pH < 4,5 et une caractéristique de libération plus lente et en outre indépendante du pH dans la plage de pH 4,5 à 7,5.

10. Formulation pour comprimés à libération prolongée selon l'une quelconque des revendications précédentes où la quantité contenue de pramipexole ou de son sel pharmaceutiquement acceptable est suffisante pour fournir une dose quotidienne administrée en une fois.

11. Procédé de fabrication de la formulation pour comprimés à libération prolongée selon l'une des revendications 1 à 10 précédentes via un procédé de compression directe comprenant les étapes de

- (1) production d'une trituration d'ingrédient actif où l'ingrédient actif est le pramipexole ou un sel pharmaceutiquement acceptable de celui-ci par prémélange de celui-ci avec une portion du ou des polymères gonflant dans l'eau et/ou du ou des excipients dans un mélangeur, où le pramipexole ou son sel pharmaceutiquement acceptable est broyé, de préférence broyé dans un broyeur à broches, avant l'utilisation ;
- (2) prémélange de la trituration d'ingrédient actif de l'étape (1), de la portion principale du ou des polymères gonflant dans l'eau et/ou des excipients dans un mélangeur pour obtenir un prémélange ;
- (3) éventuellement tamisage à sec du prémélange sur un tamis pour réaliser une ségrégation des particules cohésives et pour améliorer l'uniformité de la teneur ;
- (4) mélange du prémélange de l'étape (2) ou (3) dans un mélangeur, éventuellement par addition des excipients restants au mélange et continuation du mélange ; et
- (5) mise sous forme de comprimés du mélange final par compression de celui-ci sur une presse à comprimés appropriée pour produire des comprimés à matrice.

12. Procédé de fabrication de la formulation pour comprimés à libération prolongée selon l'une des revendications 1 à 10 précédentes via un procédé de granulation humide comprenant les étapes de

- (1) production d'une trituration d'ingrédient actif où l'ingrédient actif est le pramipexole ou un sel pharmaceutiquement acceptable de celui-ci par mélange de celui-ci avec une portion des excipients dans un mélangeur, où le pramipexole ou son sel pharmaceutiquement acceptable est broyé, de préférence broyé dans un broyeur à broches, avant l'utilisation ;
- (2) granulation de la trituration d'ingrédient actif de l'étape (1) par addition du liquide de granulation, de préférence l'eau ;
- (3) séchage des granules de l'étape (2) dans un sécheur à lit fluidisé ou un four de séchage ;
- (4) mélange des granules séchés de l'étape (3) avec le ou les polymères gonflant dans l'eau et/ou les excipients dans un mélangeur pour obtenir le mélange final ;
- (5) mise sous forme de comprimés du mélange final de l'étape (4) par compression de celui-ci sur une presse à comprimés appropriée pour produire des comprimés à matrice.

13. Procédé de fabrication de la formulation pour comprimés à libération prolongée selon l'une des revendications 1 à 10 via un procédé de granulation sèche comprenant les étapes de

- (1) mélange de l'ingrédient actif pramipexole ou d'un sel pharmaceutiquement acceptable de celui-ci avec une portion des charges ou tous les excipients dans un mélangeur, où le pramipexole ou son sel pharmaceutiquement acceptable est broyé, de préférence broyé dans un broyeur à broches, avant l'utilisation ;
- (2) compactage du mélange de l'étape (1) sur un compacteur à rouleaux approprié ;
- (3) réduction des rubans obtenus pendant l'étape (1) en petits granules par des étapes de broyage ou de tamisage appropriées ;
- (4) éventuellement mélange des granules de l'étape (3) avec les excipients restants dans un mélangeur pour obtenir le mélange final ;
- (5) mise sous forme de comprimés des granules de l'étape (3) ou du mélange final de l'étape (4) par compression de celui-ci sur une presse à comprimés appropriée pour produire des comprimés à matrice.

14. Utilisation d'une formulation pour comprimés à libération prolongée selon l'une des revendications 1 à 10 précédentes pour préparer une composition médicale pour le traitement de la maladie de Parkinson et des complications ou

troubles associés à celle-ci.

5

10

15

20

25

30

35

40

45

50

55

Fig. 1




Fig. 2





Fig. 3



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2004010997 A [0013] [0016] [0056]
- WO 2004010999 A [0015] [0016]